

Sheaf quantization from exact WKB analysis

Tatsuki Kuwagaki

October 2020

Abstract

A sheaf quantization is a sheaf associated to a Lagrangian brane. By using the ideas of exact WKB analysis, spectral networks, and scattering diagrams, we sheaf-quantize spectral curves over the Novikov ring under some assumptions on the behavior of Stokes curves.

For Schrödinger equations, we prove that the local system associated to the sheaf quantization (microlocalization a.k.a. abelianization) over the spectral curve can be identified with the Voros–Iwaki–Nakanishi coordinate.

We expect that these sheaf quantizations are the object-level realizations of the \hbar -enhanced Riemann–Hilbert correspondence.

1 Introduction

A quantization of a Lagrangian submanifold L usually means a module supported on L over a quantized symplectic manifold. For example, a \mathcal{D} -module on a complex manifold is a quantization of its characteristic variety. Via the Riemann–Hilbert correspondence, one can introduce a notion of quantization with more topological flavor: *sheaf quantization*. A sheaf is a sheaf quantization of its microsupport SS [KS94].

Characteristic varieties and microsupports are always conic. To quantize non-conic Lagrangians, it is classical to use \mathcal{D}^h -modules. A \mathcal{D}^h -module \mathcal{E} is a quantization of $\text{supp}(\mathcal{E}/\hbar\mathcal{E})$. Tamarkin introduced an analogue of this trick on the Riemann–Hilbert dual side [Tam18]. Namely, for a real manifold M , consider the product $T^*M \times T_{\tau>0}^*\mathbb{R}_t$ where \mathbb{R}_t is the real line with the standard coordinate t , τ is the dual cotangent coordinate of t , and $T_{\tau>0}^*\mathbb{R}_t$ is the subset of $T^*\mathbb{R}_t$ defined by $\tau > 0$. Let ρ be a map defined by $\rho: T^*M \times T_{\tau>0}^*\mathbb{R}_t \rightarrow T^*M; (x, \xi, t, \tau) \mapsto (x, \xi/\tau)$ where $x \in M$ and $\xi \in T_x^*M$. Roughly speaking, we say that a sheaf \mathcal{E} on $M \times \mathbb{R}_t$ is a *sheaf quantization* of $\mu\text{supp}(\mathcal{E}) := \rho(SS(\mathcal{E}) \cap (T^*M \times T_{\tau>0}^*\mathbb{R}_t))$. This idea is quite useful to study symplectic topology. For example, Tamarkin proved some results on Hamiltonian non-displaceability [Tam18] and Guillermou proved some results on the nearby Lagrangian conjecture [Gui16]. The statements obtained by sheaf quantizations are parallel to those obtained by Floer-theoretic methods. In fact, one of Tamarkin’s first motivations to introduce \mathbb{R}_t is to realize the Novikov ring Λ_0 in sheaf theory. Also, Jin–Treumann [JT] proved that a sheaf quantization is associated to an exact Lagrangian brane, which is nothing but a necessary condition to be an object of Fukaya category.

In the first version of quantization (using \mathcal{D} -modules), via the Riemann–Hilbert correspondence, a sheaf quantization over a complex manifold is equivalent to a quantization by a regular holonomic \mathcal{D} -module. However, in the second version (using \mathcal{D}^{\hbar} -modules), there is no such Riemann–Hilbert correspondence, so the relation is not clear.

One of the purpose of this paper is to reveal a part of the relationship explicitly, using exact WKB analysis. We will explore it more abstractly by constructing a functor relating \mathcal{D}^{\hbar} -modules and sheaf quantizations in the subsequent publication [Kuw22]

Naively speaking, the Riemann–Hilbert functor is about solving differential equations. Similarly, in our case, we have to solve \mathcal{D}^{\hbar} -modules. The appropriate method is *exact WKB analysis* [Vor83, DDP93, KT05]. Classically, the WKB approximation means a semi-classical approximation of quantum mechanics obtained by considering the Planck constant very small. Exact WKB analysis is a method to solve differential equations with \hbar along this line. We solve a given differential equation with special formal power series (WKB solutions) in \hbar and try to sum up the series by Borel’s method.

Exact WKB analysis is quite successful for the second order differential equations over complex 1-dimensional spaces. The result is, roughly speaking, for each $\hbar \in \mathbb{C}_{\hbar}^{\times}$, we have a chamber decomposition of the base space, solutions over each chamber, and connection formulas between solutions in different chambers. For us, these data give a gluing of sheaves.

Theorem 1.1. *Let C be a Riemann surface and \mathcal{M} be a rank 2 flat meromorphic \hbar -connection with Assumption 8.1. For a sufficiently small and generic $\hbar \in \mathbb{C}_{\hbar}^{\times}$, there exists a sheaf quantization $S_{\mathcal{M}}^{\hbar}$ of the spectral curve of \mathcal{M} as an object of $\mathrm{Sh}_{\tau>0}^{\mathbb{R}}(M \times \mathbb{R}_t)$ and its microlocalization is the Voros local system.*

The category $\mathrm{Sh}_{\tau>0}^{\mathbb{R}}(M \times \mathbb{R}_t)$ is the \mathbb{R} -equivariant version of the category $\mathrm{Sh}_{\tau>0}(M \times \mathbb{R}_t)$ introduced by Tamarkin [Tam18]. In the previous works [Gui16, JT], to sheaf-quantize a Lagrangian, we have to assume that the Lagrangian is exact. Since spectral curves are not exact, the previous framework cannot be applied to our situation. One of the technical advance of this paper is to use \mathbb{R} -equivariant sheaves to treat nonexact Lagrangians. This introduction of equivariance naturally induces the Novikov ring action in sheaf theory, which is plausible in comparison with Floer theory. This is the first direct observation of the Novikov ring in sheaf theory, although which has been anticipated since Tamarkin’s work. We will use this Novikov ring to study symplectic topology via sheaf theory in the subsequent work [IK].

In the case of higher order differential equations, there are two major problems in exact WKB analysis:

1. The summability of WKB solutions is not well-established.
2. We cannot obtain a chamber decomposition similarly as in the quadratic case, since there exist collisions of Stokes curves (chamber walls) [BNR82] and the collisions give anomalous monodromies.

In this paper, we will only address the second problem. To cancel the anomalous monodromies, we have to introduce new Stokes curves emanating from the collisions. This

prescription is first introduced by Berk–Nevins–Roberts [BNR82]. Then the new Stokes curves may have more collisions. Hence, in general, we have to add infinitely many new Stokes curves. This phenomenon is recently rediscovered in physics in the work of Gaiotto–Moore–Neitzke as *spectral network* [GMN13].

We will introduce the notion of *inductive tameness*, to have a tame behavior of new Stokes curves in each step. Then we can organize it into an inductive sequence of data. This is an analogue of the notion of *scattering diagram* [KS06, GS11] and we call it a *spectral scattering diagram*. We can mimic the inductive construction of the scattering diagram and obtain the following:

Theorem 1.2. *Let C be a Riemann surface and L be a holomorphic Lagrangian in T^*C with an inductively \hbar -tame initial diagram. Then there exists a sheaf quantization of the spectral curve as an object of $\mathrm{Sh}_{\tau>0}^{\mathbb{R}}(C \times \mathbb{R}_t)$.*

We will discuss the choice of the initial diagram coming from the meromorphic connections at the end of this paper.

Remark 1.3. In the literature of exact WKB analysis, there is another approach to treat collisions of Stokes curves using the notion of *virtual turning points* [AKT91].

As we have already mentioned, we expect our construction is a kind of Riemann–Hilbert correspondence, *\hbar -Riemann–Hilbert correspondence*. In the subsequent work [Kuw22], we construct a functor from the category of \mathcal{D}^{\hbar} -modules to the category of sheaf quantizations, although the coefficients we take is enlarged to obtain the Novikov rings on the side of \mathcal{D}^{\hbar} -modules. For the second order case, our sheaf quantization here can be comparable to the sheaf quantization obtained via the \hbar -RH functor. For the higher order case, we expect that a similar coincidence is possible after exact WKB analysis is settled.

One can also ask relations to other realizations of irregular Riemann–Hilbert correspondence. We will discuss relationships to (1) D’Agnolo–Kashiwara’s holonomic Riemann–Hilbert correspondence [DK16], and (2) Shende–Treumann–Williams–Zaslow’s Legendrian knot description [STWZ19].

Remark 1.4. Microlocal sheaf theory has already met exact WKB analysis in the work of Getmanenko–Tamarkin [GT13]. In contrast to our treatment, they applied microlocal sheaf theory to the problem after the Laplace transform. Hence their solution sheaf corresponds to Borel-transformed solutions. It may be interesting to know the relationship between our sheaves and their sheaves.

Remark 1.5. Recently, Kontsevich–Soibelman [Kon, KS] announced a formulation of Riemann–Hilbert correspondence for deformation quantization. From this point of view, our construction (or our theorem in [Kuw22]) can be considered as an intermediate step. The other step to prove Kontsevich–Soibelman’s conjecture should be a version of Nadler–Zaslow equivalence [NZ09].

Acknowledgment

I would like to thank Kohei Iwaki, Tsukasa Ishibashi, Hiroshi Ohta, and Vivek Shende for many comments and discussions. Especially, Iwaki-san asked me to find relationship between exact WKB analysis and microlocal sheaf theory for some years. I am very happy to write this note as my “homework report” of his lectures. I also would like to thank the members of GHKK reading seminar (Ishibashi, Kano, Mizuno, Oya) where I learned the notion of scattering diagram. This work was partially supported by World Premier International Research Center Initiative (WPI), MEXT, Japan and JSPS KAKENHI Grant Number JP18K13405.

Notation

1. (Base field) Our base field is \mathbb{K} . In a large part of this paper, \mathbb{K} is \mathbb{C} .
2. (Novikov ring) Let R be a commutative ring. Consider $\mathbb{R}_{\geq 0}$ as a semigroup and we denote the group ring by $R[\mathbb{R}_{\geq 0}]$. We denote the indeterminate of $R[\mathbb{R}_{\geq 0}]$ by T^a for $a \in \mathbb{R}_{\geq 0}$. The Novikov ring Λ_0^R is the projective limit

$$\varprojlim_{a \rightarrow \infty} R[\mathbb{R}_{\geq 0}]/T^a R[\mathbb{R}_{\geq 0}]$$

. We denote the fraction field by Λ^R and call it the Novikov field. If the context is clear, we omit the superscript R .

3. (Category) A category usually means a dg-category unless specified. When dg-categories are involved, all the operations are understood to be derived.
4. (Planck) We will use \hbar for a general element or the coordinate function of \mathbb{C}_\hbar^\times and \mathfrak{h} for some fixed element in \mathbb{C}_\hbar .
5. (Sheaf) Let M be a topological space. For us, *sheaf* means *sheaf of \mathbb{K} -vector spaces*. For a locally closed subset Z of M , we use \mathbb{K}_Z to be the constant sheaf of rank 1 supported on Z . If Z is defined by some inequality I , we set $\mathbb{K}_I := \mathbb{K}_Z$.
6. (Complex cotangent bundle) For a complex manifold M , the cotangent bundle T^*M is a holomorphic symplectic manifold by the standard symplectic structure. We denote the projection $T^*M \rightarrow M$ by p_{T^*M} .
On a local coordinate $(U, \{z_i\})$, the standard symplectic form ω can be written as $\sum_i d\zeta_i \wedge dz_i$ where ζ_i is the cotangent coordinate dual to z_i . The form $\sum_i \zeta_i dz_i$ gives a global 1-form λ on T^*M . We call λ the Liouville form.
7. (Real cotangent bundle) Similarly, for a manifold M , the cotangent bundle T^*M is a symplectic manifold by the standard exact symplectic structure. We denote the standard real symplectic form by ω_{st} , and the standard Liouville form by λ_{st} .

2 Microlocal category

Here we will introduce our category which is the living place of our sheaf quantizations. This category is a refined version of the category introduced by Tamarkin [Tam18]. We will generalize the presentation of Guillermou–Schapira [GS11].

2.1 Positively-microsupported sheaves

Let M be a connected differentiable manifold and \mathbb{R}_t be the real line with the standard coordinate t . We denote the derived category of \mathbb{K} -module sheaves by $\mathrm{Sh}(M \times \mathbb{R}_t)$.

We denote the i -th projection by $\mathbb{R}_t \times \mathbb{R}_t \rightarrow \mathbb{R}_t$. We also have the induced map $\mathrm{id}_M \times p_i: M \times \mathbb{R}_t^2 \rightarrow M \times \mathbb{R}_t$ where id_M is the identity map of M . We also denote the addition map $\mathbb{R}_t \times \mathbb{R}_t \rightarrow \mathbb{R}_t$ by m_t .

For objects $\mathcal{E}, \mathcal{F} \in \mathrm{Sh}(M \times \mathbb{R}_t)$, we set

$$\mathcal{E} \star \mathcal{F} := (\mathrm{id}_M \times m_t)_!((\mathrm{id}_M \times p_1)^{-1}\mathcal{E} \otimes (\mathrm{id}_M \times p_2)^{-1}\mathcal{F}). \quad (2.1)$$

We call it the convolution product.

We denote the cotangent bundle of M by T^*M . Let \mathbb{R}_t be the real line with the standard coordinate t . The dual cotangent coordinate of $T^*\mathbb{R}_t$ is denoted by τ . The subset $\{\tau > 0\} := T_{\tau > 0}^*(M \times \mathbb{R}_t) = T^*M \times T_{\tau > 0}^*\mathbb{R}_t$ of $T^*M \times T^*\mathbb{R}_t$ is defined by $\tau > 0$.

We define the full subcategory $\mathrm{Sh}_{\tau \leq 0}(M \times \mathbb{R}_t)$ of $\mathrm{Sh}(M \times \mathbb{R}_t)$ as the full subcategory spanned by the objects satisfying $SS(\mathcal{E}) \subset \{\tau \leq 0\}$.

Lemma 2.1. *For $\mathcal{E} \in \mathrm{Sh}_{\tau \leq 0}(M \times \mathbb{R}_t)$, we have $\mathcal{E} \star \mathbb{K}_{t \geq 0} \simeq 0$.*

Proof. This is a special case of [GS11, Proposition 3.19 (b)]. □

We consider the skyscraper sheaf $\mathbb{K}_{M \times \{t=0\}}$.

Lemma 2.2. *For $\mathcal{E} \in \mathrm{Sh}_{\tau \leq 0}(M \times \mathbb{R}_t)$, we have $\mathcal{E} \star \mathbb{K}_{M \times \{t=0\}} \cong \mathcal{E}$.*

Proof. This is a special case of [GS11, p.25]. □

By the adjunction $\mathrm{Hom}(\mathbb{K}_{t \geq 0}, \mathbb{K}_{t=0}) \cong \mathrm{Hom}(\mathbb{K}_{t=0}, \mathbb{K}_{t=0})$, we have the map $\mathbb{K}_{t \geq 0} \rightarrow \mathbb{K}_{t=0}$ corresponding to the identity. By Lemma 2.2, we have a morphism $\mathcal{E} \star \mathbb{K}_{t \geq 0} \rightarrow \mathcal{E}$ for any $\mathcal{E} \in \mathrm{Sh}(M \times \mathbb{R}_t)$.

Lemma 2.3. *The cone of the above morphism is in ${}^{\perp}\mathrm{Sh}_{\tau \leq 0}(M \times \mathbb{R}_t)$. Moreover, the distinguished triangle gives a semi-orthogonal decomposition:*

$$\mathrm{Sh}(M \times \mathbb{R}_t) \cong \left\langle \mathrm{Sh}_{\tau \leq 0}(M \times \mathbb{R}_t), {}^{\perp}\mathrm{Sh}_{\tau \leq 0}(M \times \mathbb{R}_t) \right\rangle. \quad (2.2)$$

Proof. This is a direct consequence of [GS11, Proposition 3.21]. □

We set

$$\mathrm{Sh}_{\tau>0}(M \times \mathbb{R}_t) := \mathrm{Sh}(M \times \mathbb{R}_t) / \mathrm{Sh}_{\tau \leq 0}(M \times \mathbb{R}_t) \quad (2.3)$$

where $/$ means the Drinfeld–Verdier dg-quotient. We denote the quotient functor $\mathrm{Sh}(M \times \mathbb{R}_t) \rightarrow \mathrm{Sh}_{\tau>0}(M \times \mathbb{R}_t)$ by q .

By Lemma 2.1, the functor $(-)\star \mathbb{K}_{t \geq 0}$ induces a functor $\iota_{>0}: \mathrm{Sh}_{\tau>0}(M \times \mathbb{R}_t) \rightarrow \mathrm{Sh}(M \times \mathbb{R}_t)$.

Lemma 2.4. *The functor $\iota_{>0}$ is fully faithful embedding, and gives an equivalence onto ${}^\perp\mathrm{Sh}_{\tau \leq 0}(M \times \mathbb{R}_t)$.*

Proof. This is a special case of [GS11, Proposition 3.21]. \square

2.2 Equivariant sheaves

For $c \in \mathbb{R}$, we set

$$T_c: M \times \mathbb{R}_t \rightarrow M \times \mathbb{R}_t; x \mapsto x + c. \quad (2.4)$$

The isomorphisms $\{T_c\}_{c \in \mathbb{R}}$ gives an action of \mathbb{R} on $M \times \mathbb{R}_t$. Here we equip the group \mathbb{R} with the discrete topology.

Definition 2.5. An equivariant sheaf on $M \times \mathbb{R}_t$ consists of the following data:

1. a sheaf \mathcal{E} on $M \times \mathbb{R}_t$,
2. an isomorphism $\alpha_c: \mathcal{E} \xrightarrow{\cong} T_c \mathcal{E} := T_{c*} \mathcal{E}$ for any $c \in \mathbb{R}$

such that $T_{c*}(\alpha_{c'}) = \alpha_{c+c'}$ for any $c, c' \in \mathbb{R}$. A morphism between equivariant sheaves is a morphism of sheaves compatible with α_c 's.

Let $\mathrm{Sh}_{\heartsuit}^{\mathbb{R}}(M \times \mathbb{R}_t)$ be the abelian category of \mathbb{R} -equivariant sheaves and $\mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t)$ be the derived dg-category of $\mathrm{Sh}_{\heartsuit}^{\mathbb{R}}(M \times \mathbb{R}_t)$. Note that the equivariant derived category in the sense of Bernstein–Lunts [BL94] coincide with the naive derived category in this case, since \mathbb{R} is a discrete group.

We have the forgetful functor

$$\mathfrak{f}: \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t) \rightarrow \mathrm{Sh}(M \times \mathbb{R}_t). \quad (2.5)$$

Definition 2.6. A functor $F: \mathcal{A} \rightarrow \mathcal{B}$ between additive categories is conservative if $F(a) \simeq 0$ implies that $a \simeq 0$.

Lemma 2.7. *Let $F: \mathcal{A} \rightarrow \mathcal{B}$ be an exact conservative functor between triangulated categories. Then $f: \mathcal{E} \rightarrow \mathcal{F}$ in \mathcal{A} is an isomorphism if $F(f)$ is an isomorphism.*

Proof. Suppose $F(f)$ is an isomorphism. Then $F(\mathrm{Cone}(f)) = \mathrm{Cone}(F(f)) \simeq 0$. By the conservativity, $\mathrm{Cone}(f) \simeq 0$. This completes the proof. \square

Lemma 2.8. *The forgetful functor \mathfrak{f} is conservative.*

Proof. For $\mathcal{E} \in \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t)$, if $\mathfrak{f}(\mathcal{E}) \simeq 0$, then $\mathfrak{f}(\mathcal{E}) \rightarrow 0$ is a quasi-isomorphism. Since this morphism can be lifted to $\mathcal{E} \rightarrow 0$, we obtain the desired result. \square

For an object $\mathcal{E} \in \mathrm{Sh}(M \times \mathbb{R}_t)$, the direct sum $\bigoplus_{c \in \mathbb{R}} T_c \mathcal{E}$ has an obvious equivariant structure

$$\alpha_{c'}: \bigoplus_{c \in \mathbb{R}} T_c \mathcal{E} \xrightarrow{\mathrm{id}} \bigoplus_{c \in \mathbb{R}} T_c \mathcal{E} = T_{c'} \bigoplus_{c \in \mathbb{R}} T_c \mathcal{E}. \quad (2.6)$$

The assignment

$$\mathcal{E} \mapsto \mathfrak{f}^L(\mathcal{E}) := \bigoplus_{c \in \mathbb{R}} T_c \mathcal{E} \quad (2.7)$$

defines a functor $\mathfrak{f}^L: \mathrm{Sh}(M \times \mathbb{R}_t) \rightarrow \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t)$.

Proposition 2.9. *The functor \mathfrak{f}^L is the left adjoint of \mathfrak{f} .*

Proof. Since the both functors are induced from exact functors between abelian categories, it is enough to show that the adjoint holds on the abelian level. We denote the abelian version of the functors by

$$\mathfrak{f}_{\heartsuit}: \mathrm{Sh}_{\heartsuit}^{\mathbb{R}}(M \times \mathbb{R}_t) \xleftarrow{\quad} \mathrm{Sh}_{\heartsuit}(M \times \mathbb{R}_t): \mathfrak{f}_{\heartsuit}^L. \quad (2.8)$$

For objects $\mathcal{E} \in \mathrm{Sh}_{\heartsuit}^{\mathbb{R}}(M \times \mathbb{R}_t)$ and $\mathcal{F} \in \mathrm{Sh}_{\heartsuit}(M \times \mathbb{R}_t)$, we have

$$\mathrm{Hom}_{\mathrm{Sh}_{\heartsuit}^{\mathbb{R}}(M \times \mathbb{R}_t)}(\mathfrak{f}_{\heartsuit}^L \mathcal{F}, \mathcal{E}) \cong \mathrm{Hom}_{\mathrm{Sh}_{\heartsuit}^{\mathbb{R}}(M \times \mathbb{R}_t)}\left(\bigoplus_{c \in \mathbb{R}} T_c \mathcal{F}, \mathcal{E}\right). \quad (2.9)$$

We would like to compute the right hand side. First, we have a morphism

$$\mathrm{Hom}_{\mathrm{Sh}_{\heartsuit}^{\mathbb{R}}(M \times \mathbb{R}_t)}\left(\bigoplus_{c \in \mathbb{R}} T_c \mathcal{F}, \mathcal{E}\right) \rightarrow \mathrm{Hom}_{\mathrm{Sh}_{\heartsuit}(M \times \mathbb{R}_t)}\left(\bigoplus_{c \in \mathbb{R}} T_c \mathcal{F}, \mathfrak{f}(\mathcal{E})\right) \quad (2.10)$$

induced by \mathfrak{f} . Next, we consider the pull-back along the inclusion $\mathcal{F} = T_0 \mathcal{F} \hookrightarrow \bigoplus_{c \in \mathbb{R}} T_c \mathcal{F}$:

$$\mathrm{Hom}_{\mathrm{Sh}_{\heartsuit}(M \times \mathbb{R}_t)}\left(\bigoplus_{c \in \mathbb{R}} T_c \mathcal{F}, \mathfrak{f}(\mathcal{E})\right) \rightarrow \mathrm{Hom}_{\mathrm{Sh}_{\heartsuit}(M \times \mathbb{R}_t)}(\mathcal{F}, \mathfrak{f}(\mathcal{E})). \quad (2.11)$$

Composing these two morphisms, we obtain a morphism

$$C: \mathrm{Hom}_{\mathrm{Sh}_{\heartsuit}^{\mathbb{R}}(M \times \mathbb{R}_t)}\left(\bigoplus_{c \in \mathbb{R}} T_c \mathcal{F}, \mathcal{E}\right) \rightarrow \mathrm{Hom}_{\mathrm{Sh}_{\heartsuit}(M \times \mathbb{R}_t)}(\mathcal{F}, \mathfrak{f}(\mathcal{E})). \quad (2.12)$$

To complete the proof, it is enough to check that this is an isomorphism. For any element $f \in \mathrm{Hom}_{\mathrm{Sh}_{\heartsuit}(M \times \mathbb{R}_t)}(\mathcal{F}, \mathfrak{f}(\mathcal{E}))$, we set

$$f_c := \alpha_c^{-1} \circ T_{c*}(f): T_c \mathcal{F} \rightarrow T_c \mathcal{E} \xrightarrow{\alpha_c^{-1}} \mathcal{E}. \quad (2.13)$$

Then $\prod_c f_c: \bigoplus_{c \in \mathbb{R}} T_c \mathcal{F} \rightarrow \mathcal{E}$ defines an element of $\mathrm{Hom}_{\mathrm{Sh}_{\heartsuit}^{\mathbb{R}}(M \times \mathbb{R}_t)}\left(\bigoplus_{c \in \mathbb{R}} T_c \mathcal{F}, \mathcal{E}\right)$ such that $C(\prod_c f_c) = f$. This construction gives the inverse of C . This completes the proof. \square

For an object $\mathcal{E} \in \mathrm{Sh}(M \times \mathbb{R}_t)$, the microsupport $SS(\mathcal{E})$ is defined by [KS94]. Although, [KS94] defined it for the bounded case, it is well-known that the definition works well for the unbounded case.

Definition 2.10 (Microsupport). For an object $\mathcal{E} \in \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t)$, we set

$$SS(\mathcal{E}) := SS(\mathfrak{f}(\mathcal{E})). \quad (2.14)$$

2.3 Operations for equivariant sheaves

We first recall some basic operations for equivariant sheaves. For details, we refer to [BL94, Gro57].

Let G be a group. Let X_1, X_2 be G -spaces. Let $f: X_1 \rightarrow X_2$ be a G -map. Then we have functors $f_*^G, f_!^G: \mathrm{Sh}^G(X_1) \rightarrow \mathrm{Sh}^G(X_2)$, $f_G^{-1}, f_G^!: \mathrm{Sh}^G(X_2) \rightarrow \mathrm{Sh}^G(X_1)$. Standard adjunctions hold for these functors.

Let G_1, G_2 be groups. Let X_i be G_i -spaces for $i = 1, 2$. We then have the tensor product functor

$$\boxtimes^{G_1 \times G_2}: \mathrm{Sh}^{G_1}(X_1) \times \mathrm{Sh}^{G_2}(X_2) \rightarrow \mathrm{Sh}^{G_1 \times G_2}(X_1 \times X_2). \quad (2.15)$$

We consider $M \times \mathbb{R}_t^2$ on which \mathbb{R}^2 acts by the addition on each component. Through the addition $m: \mathbb{R}^2 \rightarrow \mathbb{R}$, the group \mathbb{R}^2 also acts on $M \times \mathbb{R}_t$. The kernel of the map $m: \mathbb{R}^2 \rightarrow \mathbb{R}$ is the anti-diagonal $\Delta_a := \{(a, -a) \in \mathbb{R} \times \mathbb{R}\}$.

We consider the addition map $\mathrm{id}_M \times m_t: M \times \mathbb{R}_t \times \mathbb{R}_t \rightarrow M \times \mathbb{R}_t$ on \mathbb{R}_t -factors. We then have a functor

$$(\mathrm{id}_M \times m_t)_!^{\mathbb{R}^2}: \mathrm{Sh}^{\mathbb{R}^2}(M \times \mathbb{R}_t^2) \rightarrow \mathrm{Sh}^{\mathbb{R}^2}(M \times \mathbb{R}_t). \quad (2.16)$$

Suppose $\phi: G \rightarrow H$ be a surjective group homomorphism. Let Y be an H -space. Then G acts on Y through ϕ . Let K be the kernel of $G \rightarrow H$. Then we have the invariant functor

$$(-)^K: \mathrm{Sh}^G(Y) \rightarrow \mathrm{Sh}^H(Y) \quad (2.17)$$

and the coinvariant functor

$$(-)_K: \mathrm{Sh}^G(Y) \rightarrow \mathrm{Sh}^H(Y). \quad (2.18)$$

Also, if one has an H -equivariant sheaf on Y , it can also be considered as a G -equivariant sheaf:

$$\iota^\phi: \mathrm{Sh}^H(Y) \rightarrow \mathrm{Sh}^G(Y). \quad (2.19)$$

The following is a standard fact.

Lemma 2.11. ι^ϕ is the right adjoint of $(-)_K$ and the left adjoint of $(-)^K$.

In these notations, we have a functor

$$(-)_{\Delta_a}: \mathrm{Sh}^{\mathbb{R}^2}(M \times \mathbb{R}_t) \rightarrow \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t). \quad (2.20)$$

We set

$$(m_!^{\mathbb{R}^2})_{\Delta_a} := (-)_{\Delta_a} \circ m_!^{\mathbb{R}^2} : \mathrm{Sh}^{\mathbb{R}^2}(M \times \mathbb{R}_t^2) \rightarrow \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t). \quad (2.21)$$

We have the right adjoint of $(m_!^{\mathbb{R}^2})_{\Delta_a}$:

$$(m_!^{\mathbb{R}^2})_{\Delta_a}^R := m_{\mathbb{R}^2}^1 \circ \iota^a : \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t) \rightarrow \mathrm{Sh}^{\mathbb{R}^2}(M \times \mathbb{R}_t^2). \quad (2.22)$$

Let $p_i: M \times \mathbb{R}_t^2 \rightarrow M \times \mathbb{R}_t$ be the i -th projection. We also have the corresponding projection $q_i: \mathbb{R}^2 \rightarrow \mathbb{R}$. We then have

$$(p_{i*}^{\mathbb{R}^2})^{\ker q_i} := (-)^{\ker q_i} \circ p_{i*}^{\mathbb{R}^2} : \mathrm{Sh}^{\mathbb{R}^2}(M \times \mathbb{R}_t^2) \rightarrow \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t) \quad (2.23)$$

and

$$L(p_{i*}^{\mathbb{R}^2})^{\ker q_i} := p_{i, \mathbb{R}^2}^{-1} \circ \iota^{q_i} : \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t) \rightarrow \mathrm{Sh}^{\mathbb{R}^2}(M \times \mathbb{R}_t^2). \quad (2.24)$$

Lemma 2.12. $L(p_{i*}^{\mathbb{R}^2})^{\ker q_i}$ is the left adjoint of $p_{i*}^{\ker q_i}$.

Definition 2.13 (Convolution product). We set, for objects $\mathcal{E}, \mathcal{F} \in \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t)$,

$$\mathcal{E} \star_{\mathbb{R}} \mathcal{F} := (m_!^{\mathbb{R}^2})_{\Delta_a} ((\mathrm{id}_M \times p_1)_{\mathbb{R}}^{-1} \mathcal{E} \otimes^{\mathbb{R} \times \mathbb{R}} (\mathrm{id}_M \times p_2)_{\mathbb{R}}^{-1} \mathcal{F}). \quad (2.25)$$

Definition 2.14 (Convolution product). We set, for objects $\mathcal{E} \in \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t)$ and $\mathcal{F} \in \mathrm{Sh}(M \times \mathbb{R}_t)$,

$$\mathcal{E} \star_R \mathcal{F} := \mathcal{E} \star_{\mathbb{R}} \mathfrak{f}^L(\mathcal{F}) \in \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t). \quad (2.26)$$

Lemma 2.15. 1. For $\mathcal{E}, \mathcal{F}, \mathcal{G} \in \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t)$, we have $(\mathcal{E} \star_{\mathbb{R}} \mathcal{F}) \star_{\mathbb{R}} \mathcal{G} \cong \mathcal{E} \star_{\mathbb{R}} (\mathcal{F} \star_{\mathbb{R}} \mathcal{G})$.

2. For $\mathcal{E}, \mathcal{F} \in \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t)$ and $\mathcal{G} \in \mathrm{Sh}(M \times \mathbb{R}_t)$, we have $(\mathcal{E} \star_{\mathbb{R}} \mathcal{F}) \star_R \mathcal{G} \cong \mathcal{E} \star_{\mathbb{R}} (\mathcal{F} \star_R \mathcal{G})$.

3. For $\mathcal{E} \in \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t)$ and $\mathcal{F}, \mathcal{G} \in \mathrm{Sh}(M \times \mathbb{R}_t)$, we have $(\mathcal{E} \star_R \mathcal{F}) \star_R \mathcal{G} \cong \mathcal{E} \star_R (\mathcal{F} \star \mathcal{G})$.

Proof. The proofs are straightforward. We omit the proof. \square

For \star_R , the underlying sheaf is obtained as the nonequivariant convolution product:

Lemma 2.16. For objects $\mathcal{E} \in \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t)$ and $\mathcal{F} \in \mathrm{Sh}(M \times \mathbb{R}_t)$, we have

$$\mathfrak{f}(\mathcal{E} \star_R \mathcal{F}) \cong \mathfrak{f}(\mathcal{E}) \star \mathcal{F}. \quad (2.27)$$

Proof. We have

$$\mathfrak{f}(\mathcal{E}) \star \mathfrak{f}^L(\mathcal{F}) \cong \bigoplus_{c \in \mathbb{R}} T_c(\mathfrak{f}(\mathcal{E}) \star \mathcal{F}) \cong \bigoplus_{c \in \mathbb{R}} (T_c \mathfrak{f}(\mathcal{E})) \star \mathcal{F} \cong \bigoplus_{c \in \mathbb{R}} \mathfrak{f}(\mathcal{E}) \star \mathcal{F} \quad (2.28)$$

where the last isomorphism is coming from the equivariant structure morphism of \mathcal{E} . Under this identification, the pull back along $a \in \Delta_a$ is $\bigoplus_{c-a \in \mathbb{R}} \mathfrak{f}(\mathcal{E}) \star \mathcal{F}$. By taking the coinvariant, we obtain $\mathfrak{f}(\mathcal{E}) \star \mathcal{F}$. This completes the proof. \square

Note that the equivariant structure of $\mathcal{E} \star \mathcal{F}$ is given by $\mathfrak{f}(\mathcal{E}) \star \mathcal{F} \xrightarrow{\alpha_c \star \text{id}} (T_c \mathfrak{f}(\mathcal{E})) \star \mathcal{F}$ by the proof of the above lemma.

Lemma 2.17. *We have*

$$(-) \star_R \mathbb{K}_{M \times \{t=0\}} \cong \text{id}. \quad (2.29)$$

Proof. By Lemma 2.16 and Lemma 2.2, we have

$$\mathfrak{f}(\mathcal{E} \star_R \mathbb{K}_{M \times \{t=0\}}) \cong \mathfrak{f}(\mathcal{E}) \star \mathbb{K}_{M \times \{t=0\}} \cong \mathfrak{f}(\mathcal{E}). \quad (2.30)$$

By the above remark, the equivariant structures also coincide. \square

2.4 Positively microsupported equivariant sheaves

In this section, we would like to consider the equivariant version of positively microsupported sheaves.

We define the full subcategory $\text{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t)$ of $\text{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t)$ as the full subcategory spanned by the objects satisfying $SS(\mathcal{E}) \subset \{\tau \leq 0\}$.

Lemma 2.18. *The functor \mathfrak{f} can be restricted to $\mathfrak{f}: \text{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t) \rightarrow \text{Sh}_{\tau \leq 0}(M \times \mathbb{R}_t)$.*

Proof. This follows from the definition of SS for equivariant sheaves (2.14). \square

Recall the notation $\mathbb{K}_{t \geq 0} := \mathbb{K}_{M \times \{t \geq 0\}}$.

Lemma 2.19. *For $\mathcal{E} \in \text{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t)$, we have $\mathcal{E} \star_R \mathbb{K}_{t \geq 0} \simeq 0$.*

Proof. For $\mathcal{E} \in \text{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t)$, we have

$$\mathfrak{f}(\mathcal{E} \star_R \mathbb{K}_{t \geq 0}) \cong \mathfrak{f}(\mathcal{E}) \star \mathbb{K}_{t \geq 0} \quad (2.31)$$

by Lemma 2.16. Combining Lemma 2.18 and Lemma 2.1, we can conclude that the right hand side is zero. By Lemma 2.8, this implies that $\mathcal{E} \star_R \mathbb{K}_{t \geq 0}$ is also zero. This completes the proof. \square

By Lemma 2.17, $(-) \star_R \mathbb{K}_{M \times \{t=0\}} \simeq \text{id}$, we have a canonical morphism

$$(-) \star_R \mathbb{K}_{t \geq 0} \rightarrow \text{id} \quad (2.32)$$

induced by the canonical morphism $\mathbb{K}_{t \geq 0} \rightarrow \mathbb{K}_{M \times \{t=0\}}$ corresponding to id under the adjunction isomorphism $\text{Hom}(\mathbb{K}_{t \geq 0}, \mathbb{K}_{M \times \{t=0\}}) \cong \text{Hom}(\mathbb{K}_{M \times \{t=0\}}, \mathbb{K}_{M \times \{t=0\}})$.

In particular, we have a morphism

$$((-) \star_R \mathbb{K}_{t \geq 0}) \star_R \mathbb{K}_{t \geq 0} \rightarrow (-) \star_R \mathbb{K}_{t \geq 0}. \quad (2.33)$$

Lemma 2.20. *The above morphism is an isomorphism*

Proof. By Lemma 2.15, we have

$$((-) \star_R \mathbb{K}_{t \geq 0}) \star_R \mathbb{K}_{t \geq 0} \cong (-) \star_R (\mathbb{K}_{t \geq 0} \star \mathbb{K}_{t \geq 0}) \cong (-) \star \mathbb{K}_{t \geq 0}. \quad (2.34)$$

The latter isomorphism is well-known. \square

We denote the essential image of $(-) \star_R \mathbb{K}_{t \geq 0}$ by $\mathfrak{Im}(\mathbb{K}_{t \geq 0})$.

Lemma 2.21. *The full subcategory $\mathfrak{Im}(\mathbb{K}_{t \geq 0})$ is closed under taking cones.*

Proof. For any morphism $f: \mathcal{E} \rightarrow \mathcal{F}$ in $\mathfrak{Im}(\mathbb{K}_{t \geq 0})$, the natural transformation $(-) \star_R \mathbb{K}_{t \geq 0} \rightarrow \text{id}$ gives a commutative square

$$\begin{array}{ccc} \mathcal{E} & \xrightarrow{f} & \mathcal{F} \\ \downarrow & & \downarrow \\ \mathcal{E} \star_R \mathbb{K}_{t \geq 0} & \xrightarrow{f \star_R \mathbb{K}_{t \geq 0}} & \mathcal{F} \star_R \mathbb{K}_{t \geq 0}. \end{array} \quad (2.35)$$

Since the vertical arrows are isomorphisms by Lemma 2.20, we have

$$\text{Cone}(f) \cong \text{Cone}(f \star_R \mathbb{K}_{t \geq 0}) \cong \text{Cone}(f) \star_R \mathbb{K}_{t \geq 0}. \quad (2.36)$$

This completes the proof. \square

Lemma 2.22. *For any object $\mathcal{E} \in \text{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t)$, we have a distinguished triangle*

$$\mathcal{E}_1 \rightarrow \mathcal{E} \rightarrow \mathcal{E}_2 \xrightarrow{[1]} \quad (2.37)$$

such that $\mathcal{E}_1 \in \mathfrak{Im}(\mathbb{K}_{t \geq 0})$ and $\mathcal{E}_2 \in \text{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t)$.

Proof. We set $\mathcal{E}_1 := \mathcal{E} \star_R \mathbb{K}_{t \geq 0}$. By Lemma 2.16,

$$\mathfrak{f}(\mathcal{E}_2) \cong \text{Cone}(\mathfrak{f}(\mathcal{E}_1) \rightarrow \mathfrak{f}(\mathcal{E})) \cong \text{Cone}(\mathfrak{f}(\mathcal{E}) \star \mathbb{K}_{t \geq 0} \rightarrow \mathfrak{f}(\mathcal{E})). \quad (2.38)$$

The microsupport of the RHS is contained in $\{\tau \leq 0\}$ by Lemma 2.3. Then we have $\mathcal{E}_2 \in \text{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t)$. This completes the proof. \square

Lemma 2.23. *We have a semi-orthogonal decomposition*

$$\text{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t) \cong \left\langle \text{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t), {}^{\perp}\text{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t) \right\rangle. \quad (2.39)$$

Moreover, $\mathfrak{Im}(\star_R \mathbb{K}_{t \geq 0})$ is contained in ${}^{\perp}\text{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t)$, and the inclusion is an equivalence.

Proof. By Lemma 2.20 and Lemma 2.22, $\mathfrak{Im}(\mathbb{K}_{t \geq 0})$ is an admissible subcategory and get a semi-orthogonal decomposition

$$\text{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t) \cong \left\langle \text{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t), \mathfrak{Im}(\star_R \mathbb{K}_{t \geq 0}) \right\rangle. \quad (2.40)$$

This completes the proof. \square

We set

$$\mathrm{Sh}_{\tau>0}^{\mathbb{R}}(M \times \mathbb{R}_t) := \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t) / \mathrm{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t). \quad (2.41)$$

To simplify the notations, we set

$$\mu(T^*M) := \mathrm{Sh}_{\tau>0}^{\mathbb{R}}(M \times \mathbb{R}_t). \quad (2.42)$$

We denote the quotient functor by

$$q_{\mathbb{R}}: \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t) \rightarrow \mu(T^*M). \quad (2.43)$$

Lemma 2.24. *The quotient functor restricts to an equivalence $q_{\mathbb{R}}|_{\perp \mathrm{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t)}: \perp \mathrm{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t) \rightarrow \mu(T^*M)$.*

Proof. This is straightforward from Lemma 2.23. \square

Lemma 2.19 implies that the functor $(-)\star_R \mathbb{K}_{t \geq 0}$ descends to a functor from $\mu(T^*M)$ to $\mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t)$. We denote the induced functor by $\iota_{>0}^{\mathbb{R}}$.

Lemma 2.25. 1. *The functor $\iota_{>0}^{\mathbb{R}}$ is fully faithful.*

2. *The image of the functor $\iota_{>0}^{\mathbb{R}}$ is $\mathfrak{Im}(\star_R \mathbb{K}_{t \geq 0})$. Composing it with the inclusion equivalence $\mathfrak{Im}(\star_R \mathbb{K}_{t \geq 0}) \hookrightarrow \perp \mathrm{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t)$, it gives an inverse of the equivalence of Lemma 2.24.*

3. *Composition of $\iota_{>0}^{\mathbb{R}}: \mu(T^*M) \rightarrow \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t)$ and the quotient functor $q_{\mathbb{R}}: \mathrm{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t) \rightarrow \mu(T^*M)$ is the identity.*

Proof. (1). The composition $\iota_{>0}^{\mathbb{R}} \circ q_{\mathbb{R}}|_{\perp \mathrm{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t)}$ is $(-)\star_R \mathbb{K}_{t \geq 0}$. This is an equivalence onto $\mathfrak{Im}(\star_R \mathbb{K}_{t \geq 0})$ by Lemma 2.20 and Lemma 2.23. Hence $\iota_{>0}^{\mathbb{R}}$ is an equivalence.

(2). Composing further with the inclusion equivalence $\mathfrak{Im}(\star_R \mathbb{K}_{t \geq 0}) \hookrightarrow \perp \mathrm{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t)$, the functor becomes $\star_R \mathbb{K}_{t \geq 0}$. This is id by Lemma 2.20 and Lemma 2.23. Hence $\iota_{>0}^{\mathbb{R}}$ is the inverse of $q_{\mathbb{R}}|_{\perp \mathrm{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t)}$.

(3). By (2), we have

$$q_{\mathbb{R}} \circ \iota_{>0}^{\mathbb{R}} \cong q_{\mathbb{R}}|_{\perp \mathrm{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t)} \circ \iota_{>0}^{\mathbb{R}} \cong \mathrm{id}. \quad (2.44)$$

\square

2.5 Operations for positively microsupported equivariant sheaves

We also would like to introduce the convolution product for positively microsupported sheaves.

We denote the composite functor

$$q_{\mathbb{R}} \circ (\iota_{>0}^{\mathbb{R}}(-)\star_{\mathbb{R}} \iota_{>0}^{\mathbb{R}}(-)): (\mu(T^*M))^{\times 2} \rightarrow \mu(T^*M) \quad (2.45)$$

by $\star_{\mathbb{R}}^{>0}$. If the context is clear, we simply write it by $\star_{\mathbb{R}}$.

We also denote the composition of functors

$$q_{\mathbb{R}} \circ (\iota_{>0}^{\mathbb{R}}(-)\star_R \iota_{>0}^{\mathbb{R}}(-)): \mu(T^*M) \otimes \mathrm{Sh}(M \times \mathbb{R}_t) \rightarrow \mu(T^*M). \quad (2.46)$$

by $\star_R^{>0}$. If the context is clear, we simply write it by \star_R .

2.6 Novikov ring action

We consider $\mathbb{K}_{t \geq 0} \in \text{Sh}(M \times \mathbb{R}_t)$. Then we obtain an object $1_{\mu M} := q_{\mathbb{R}} \circ \mathfrak{f}^L(\mathbb{K}_{t \geq 0})$.

Lemma 2.26.

$$H^0 \text{End}_{\mu(T^*M)}(1_{\mu M}) \cong \Lambda_0. \quad (2.47)$$

Proof. Note that $\mathfrak{f}^L(\mathbb{K}_{t \geq 0}) \in {}^\perp \text{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t)$. We have

$$\begin{aligned} \text{Hom}_{\mu(T^*M)}(1_{\mu M}, 1_{\mu M}) &\cong \text{Hom}_{\perp \text{Sh}_{\tau \leq 0}^{\mathbb{R}}(M \times \mathbb{R}_t)}(\mathfrak{f}^L(\mathbb{K}_{t \geq 0}), \mathfrak{f}^L(\mathbb{K}_{t \geq 0})) \\ &\cong \text{Hom}_{\text{Sh}^{\mathbb{R}}(M \times \mathbb{R}_t)}(\mathfrak{f}^L(\mathbb{K}_{t \geq 0}), \mathfrak{f}^L(\mathbb{K}_{t \geq 0})) \\ &\cong \text{Hom}_{\text{Sh}(M \times \mathbb{R}_t)}(\mathbb{K}_{t \geq 0}, \mathfrak{f}^L(\mathbb{K}_{t \geq 0})) \\ &\cong \text{Hom}_{\text{Sh}(M \times \mathbb{R}_t)}(\mathbb{K}_{t \geq 0}, \bigoplus_{c \in \mathbb{R}} \mathbb{K}_{t \geq c}). \end{aligned} \quad (2.48)$$

We first note that

$$\text{Hom}_{\text{Sh}(M \times \mathbb{R}_t)}(\mathbb{K}_{t \geq 0}, \mathbb{K}_{t \geq c}) = \begin{cases} \mathbb{K} & \text{for } c \geq 0, \\ 0 & \text{otherwise.} \end{cases} \quad (2.49)$$

Hence $\text{Hom}_{\text{Sh}(M \times \mathbb{R}_t)}(\mathbb{K}_{t \geq 0}, \bigoplus_{c \in \mathbb{R}} \mathbb{K}_{t \geq c}) \cong \text{Hom}_{\text{Sh}(M \times \mathbb{R}_t)}(\mathbb{K}_{t \geq 0}, \bigoplus_{c \geq 0} \mathbb{K}_{t \geq c})$. So our computation is reduced to compute the space of sheaf homomorphisms from $\mathbb{K}_{t \geq 0}$ to $\bigoplus_{c \geq 0} \mathbb{K}_{t \geq c}$, which is the same as the global section space of $\bigoplus_{c \geq 0} \mathbb{K}_{t \geq c}$.

For the purpose, we view $\bigoplus_{c \geq 0} \mathbb{K}_{t \geq c}$ as the sheafification of the presheaf

$$U \mapsto \bigoplus_{c \geq 0} (\mathbb{K}_{t \geq c}(U)). \quad (2.50)$$

The presheaf already satisfies the locality condition. Then any global section of the sheafification is given by an open covering $\mathbb{R}_t = \bigcup_{i \in I} U_i$ with compatible sections $\{s_i \in \bigoplus_{c \in \mathbb{R}} \mathbb{K}_{t \geq c}(U_i)\}_{i \in I}$. If $|I|$ is finite, such a section arises from a section of the presheaf. So we consider the case when $|I|$ is not finite. Since \mathbb{R} is paracompact, we can assume $I = \mathbb{N}$. Then, for each $i \in \mathbb{N}$, the set

$$\{c \in \mathbb{R} \mid s_j^c \neq 0, j \leq i\} \quad (2.51)$$

is a finite set where s_j^c is the component of s_j in $\mathbb{K}_{t \geq c}(U_j)$. It implies that there exists a section $s'_i \in \bigoplus_{c \geq 0} (\mathbb{K}_{t \geq c}(\bigcup_{j \leq i} U_j))$ of the presheaf such that $s'_i|_{U_j} = s_j$ for any $j \leq i$.

This observation implies that we have

$$\Gamma(\mathbb{R}_t, \bigoplus_{c \geq 0} \mathbb{K}_{t \geq c}) \cong \lim_{+\infty \leftarrow a} \bigoplus_{c \geq 0} (\mathbb{K}_{t \geq c}((-a, a))). \quad (2.52)$$

We also have an identification of vector spaces $\bigoplus_{c \geq 0} (\mathbb{K}_{t \geq c}((-a, a))) \cong \mathbb{K}[\mathbb{R}_{\geq 0}]/T^a \mathbb{K}[\mathbb{R}_{\geq 0}]$. Hence we have $\Gamma(\mathbb{R}_t, \bigoplus_{c \geq 0} \mathbb{K}_{t \geq c}) \cong \Lambda_0$. It is easy to check that this also gives a ring isomorphism. This completes the proof. \square

We would like to construct the Novikov ring action on the homotopy category of $\mu(T^*M)$.

Lemma 2.27. *The functor $(-)\star_{\mathbb{R}}1_{\mu M}$ is isomorphic to the identity on $\mu(T^*M)$.*

Proof. It is enough to show that $(-)\star_{\mathbb{R}}\mathfrak{f}(\mathbb{K}_{t\geq 0})$ is the identity on ${}^{\perp}\mathrm{Sh}_{\tau\leq 0}^{\mathbb{R}}(M\times\mathbb{R}_t)$. This is already proved in Lemma 2.20 and Lemma 2.23. \square

By the functoriality of $\star_{\mathbb{R}}$, Lemma 2.26, and Lemma 2.27, we get a sequence of morphisms

$$\Lambda_0 \rightarrow \mathrm{End}_{\mu(T^*M)}(1_{\mu M}) \rightarrow \mathrm{End}_{\mu(T^*M)}(\mathcal{E}\star_{\mathbb{R}}1_{\mu M}) \simeq \mathrm{End}_{\mu(T^*M)}(\mathcal{E}) \quad (2.53)$$

This gives a Λ_0 -linear structure of the homotopy category of $\mu(T^*M)$.

Remark 2.28. Since $(-)\star 1_{\mu M}$ is isomorphic to the identity on $\mu(T^*M)$, the image of $(-)\star 1_{\mu M}$ is quasi-equivalent to $\mu(T^*M)$. The former dg-category is enriched over Λ_0 by the above observation, hence the latter category is also enriched over Λ_0 in a homotopical sense.

2.7 Non-conic microsupport

Take an exact symplectic structure ω on T^*M with its primitive λ . Then $(T^*M\times\mathbb{R}_t, dt+\lambda)$ is a contact manifold. We denote the standard symplectic structure of T^*M by ω_{st} with its standard primitive λ_{st} . Then $\omega_{st}+d\tau\wedge dt$ is a symplectic structure of $T^*M\times T_{\tau>0}^*\mathbb{R}_t$. Suppose there exists an $\mathbb{R}_{>0}$ -action on $T^*M\times T_{>0}^*\mathbb{R}_T$ which makes it a homogeneous symplectic manifold and its quotient is the contact manifold $(T^*M\times\mathbb{R}_t, dt+\lambda)$. Then we have projections $T^*M\times T_{>0}^*\mathbb{R}_t\rightarrow T^*M\times\mathbb{R}_t\rightarrow T^*M$ where the second projection is the stupid projection. We denote the composition by ρ . By the uniqueness lemma of Viterbo [Vit], such ρ is unique for a given λ if it exists.

The following is the fundamental example.

Example 2.29. For the standard Liouville structure, $\rho_{st}:=\rho$ can be explicitly written as $\rho_{st}:T^*M\times T_{\tau>0}^*\mathbb{R}_t\rightarrow T^*M$ by $(x,\xi,t,\tau)\mapsto(x,\xi/\tau)$ where $x\in M, \xi\in T_x^*M$.

In the following, we write ρ_{st} by ρ unless specified.

Definition 2.30. For an object $\mathcal{E}\in\mathrm{Sh}_{\tau>0}^{\mathbb{R}}(M\times\mathbb{R}_t)$, we set

$$\mu\mathrm{supp}(\mathcal{E}) := \rho(SS(\mathcal{E})\cap\{\tau>0\}). \quad (2.54)$$

3 Sheaf quantization

In this section, we discuss the microlocalization of sheaves in the equivariant context. The nonequivariant version is discussed in [KS94, Gui16, JT].

3.1 Maslov covering and graded Lagrangian

Let V be a symplectic vector space. The fundamental group of the Lagrangian Grassmannian $LGr(V)$ is isomorphic to \mathbb{Z} . We denote the universal covering by $\pi_V: \widetilde{LGr}(V) \rightarrow LGr(V)$, which is a \mathbb{Z} -covering.

Let X be a symplectic manifold. We denote the Lagrangian Grassmannian bundle by $LGr(X)$.

Definition 3.1 ([Sei00, 2b]). A Maslov covering $\widetilde{LGr}(X)$ is a fiber bundle over X with a morphism $\widetilde{LGr}(X) \rightarrow LGr(X)$ such that it is the universal covering fiberwisely.

Let L be a Lagrangian submanifold of X . By taking its tangent fibers, we get a section of the projection $\lambda_L: LGr(X)|_L \rightarrow L$. We call the section the Lagrangian Gauss map.

Definition 3.2. A graded Lagrangian submanifold is a Lagrangian submanifold L with a choice of a map $L \rightarrow \widetilde{LGr}(X)|_L$ which is a lift of the Lagrangian Gauss map.

When X is the cotangent bundle T^*M , there exists a canonical choice of a Maslov covering [Sei00, 2b] as follows:

Definition 3.3. A Lagrangian distribution over X is a choice of subbundle of TX such that it is a Lagrangian subspace in each fiber.

Let X be a symplectic manifold with a Lagrangian distribution. Then we get a section of $LGr(X) \rightarrow X$. Then we can construct the fiberwise universal covering of $LGr(X)$ by taking the image of $LGr(X)$ as fiberwise base point.

We consider the case when X is a cotangent bundle T^*M . We denote the projection to the base M by π . For any point $p \in T^*M$, the assignment $p \mapsto T_{f,p}T^*M := T_p T_{\pi^{-1}\pi(p)}^*M \subset T_p T^*M$ forms a Lagrangian distribution. Then we can form a fiberwise universal covering as above. In the following, for cotangent bundles, we always use this Maslov covering.

Example 3.4 (cotangent fiber). For a point $x \in M$, the cotangent fiber $L = T_x^*M$ is Lagrangian. The Gauss map $L \rightarrow LGr(T^*M)$ is in the Lagrangian distribution. Hence we can lift it to $L \rightarrow \widetilde{LGr}(T^*M)$ as a fiberwise trivial loop.

Example 3.5 (Graph). The zero section T_M^*M is Lagrangian. We have a splitting $TT^*M|_{T_M^*M} = T_f T^*M \oplus TM$. Each factor is a Lagrangian subspace. Then the rotation action

$$r_\theta := \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} : T_f T^*M \oplus TM \rightarrow T_f T^*M \oplus TM \quad (3.1)$$

acts as a symplectic bundle isomorphism over M . In particular, $r_\theta(T_f T^*M)$ ($\theta \in [0, \pi/2]$) gives a fiberwise path in $LGr(T^*M)$ from the Lagrangian distribution to the Gauss map image of T_M^*M . Hence this gives a grading of T_M^*M . More generally, for a closed 1-form ϕ , the graph $G(\phi)$ of ϕ is a Lagrangian submanifold of T^*M . For $c \in [0, 1]$, the graph of $c \cdot \phi$ gives a Lagrangian isotopy between $G(\phi)$ and T_M^*M . Hence the above grading of T_M^*M induces a grading of $G(\phi)$.

3.2 Relative Pin structure

We also introduce the notion of relative Pin structure. Let L be an n -dimensional manifold. Then the classification map of the tangent bundle is

$$L \rightarrow BO(n). \quad (3.2)$$

The second Stiefel–Whitney class is given by the composition of the above morphism with the universal second SW class $BO(n) \rightarrow B^2\mathbb{Z}/2$. We also have the following homotopy exact sequence:

$$BPin^+(n) \rightarrow BO(n) \rightarrow B^2\mathbb{Z}/2. \quad (3.3)$$

Here $Pin^+(n)$ is a double cover of $O(n)$ with the center $\mathbb{Z}/2 \times \mathbb{Z}/2$.

Definition 3.6. A Pin structure on L is a lift $L \rightarrow BPin^+(n)$ of the map $L \rightarrow BO(n)$.

In other words, it is a null homotopy of the second SW map. From the above homotopy exact sequence, we need the vanishing of the second SW class to obtain a Pin structure.

We can consider a relative version. Fix $w \in H^2(L, \mathbb{Z}/2)$. Then we can twist $L \rightarrow B^2\mathbb{Z}/2$ by adding $-w$.

Definition 3.7. A null-homotopy of the second SW map twisted by w is called a relative Pin structure with the background class w .

In the case of $X = T^*M$, we consider the background class $w_2(T^*M)|_L$.

Example 3.8 (Graph). Suppose X is the cotangent bundle T^*M . We consider the zero section $L = T_M^*M$. Then $w_2(X)|_L = w_2(L)$. Hence the twist gives the trivial map $L \rightarrow B^2\mathbb{Z}/2$. Hence there exists a trivial Pin structure. Similarly, we can equip the graph of closed 1-form with a relative Pin structure.

Example 3.9 (Cotangent fiber). We consider the cotangent fiber $L = T_x^*M$. Then $w_2(X)|_L = 0 = w_2(L)$. Hence the twist gives the trivial map. Hence there exists a trivial Pin structure.

3.3 Microsheaves along Lagrangian

In this subsection, we recall the notion of microsheaves. The content in this subsection is essentially contained in [Jin].

For an open subset $U \subset T^*N$, we set

$$\mathrm{Sh}(U; \mathbb{K}_N) := \mathrm{Sh}(\mathbb{K}_N) / \{ \mathcal{E} \mid \mathrm{SS}(\mathcal{E}) \subset T^*N \setminus U \}. \quad (3.4)$$

The assignment forms a prestack. The stackification is denoted by $\mu\mathrm{Sh}(-)$, called the Kashiwara–Schapira stack.

For a subset $A \subset U$, we can consider the subsheaf spanned by the objects supported in A . We denote it by $\mu\mathrm{Sh}_A$. This can be viewed as a sheaf supported on A .

Let L be a Lagrangian submanifold in the cotangent bundle T^*M . Restricting the Liouville form λ to a contractible open subset U of L , there exists a primitive of $\lambda|_U$ by the Poincare lemma. We fix such a primitive $f_U: U \rightarrow \mathbb{R}$. Then we set

$$L_{f_U} := \{(x, \xi, t, \tau) \in T^*M \times \{\tau > 0\} \mid (x, \xi/\tau) \in U, t = -f_U(x, \xi/\tau)\}. \quad (3.5)$$

Under the twisted projection $\rho := \rho_{st}$, we have $\rho(L_{f_U}) = U$. We can consider the sheaf $\mu\text{Sh}_{L_{f_U}}$ on L_{f_U} . Since $\mu\text{Sh}_{L_{f_U}}$ is conical, it descends to a sheaf on U . Also, the resulting sheaf on U does not depend on the choice of f . Hence we denote the resulting sheaf by μSh_U .

Take an contractible open covering $\{U_i\}_{i \in I}$ of L . For each U_i , put μSh_{U_i} . Again, the sheaf is defined locally on U_i , they can be glued up on the intersections $U_i \cap U_j$. Hence we get a sheaf of categories μSh_L over L . The sheaf is a locally constant sheaf of categories, hence classified by a map

$$\text{KS}: L \rightarrow B\text{Pic}(\text{Mod}(\mathbb{K})). \quad (3.6)$$

where $B\text{Pic}(\text{Mod}(\mathbb{K}))$ is the delooping of the Picard groupoid of $\text{Mod}(\mathbb{K})$. Note that giving a simple global section of μSh_L is equivalent to giving a null homotopy of KS , and further equivalent to giving an equivalence $\mu\text{Sh}_L \cong \text{Loc}_L$ where Loc_L is the sheaf of the local systems over L .

Definition 3.10. A \mathbb{K} -brane structure of L is a null homotopy of KS .

On the other hand, we have the Lagrangian Gauss map $L \rightarrow U/O$ to the stable Lagrangian Grassmannian and the delooped J -homomorphism $B(U/O) \rightarrow \text{Mod}(\mathbb{S})$ where the RHS is the module category of the sphere spectrum.

Theorem 3.11. 1. *The map KS is factored as the composition of the stable Gauss map and the delooping of the J -homomorphism.*

2. *In particular, if $\mathbb{K} = \mathbb{Z}$, the map is given by $L \rightarrow B\mathbb{Z} \times B^2\mathbb{Z}/2\mathbb{Z}$ where the first factor is the Maslov class and the second factor is the relative Stiefel–Whitney class.*

Proof. This is the result of [Jin]. Although the statement in [Jin] is only for exact case, since the statement is essentially local, it works for nonexact case. \square

Since \mathbb{Z} is initial among the rings, we obtain the following.

Corollary 3.12. *For any commutative ring \mathbb{K} , the grading and a relative Pin structure of L gives a \mathbb{K} -brane structure.*

We define a full subcategory as

$$\mu_L(T^*M) := \{\mathcal{E} \in \mu(T^*M) \mid \mu\text{supp}(\mathcal{E}) \subset L\}. \quad (3.7)$$

In the following sections, we will construct the microlocalization functor

$$\mu_L: \mu_L(T^*M) \rightarrow \mu\text{Sh}_L(L). \quad (3.8)$$

3.4 Microlocalization

We consider the Kashiwara–Schapira stack $\mu\mathrm{Sh}(-)$. For a Lagrangian submanifold L , we can consider the substack $\mu\mathrm{Sh}_{\rho^{-1}(L)}(-)$ consisting of objects supported on $\rho^{-1}(L)$.

Take an open contractible covering $\{U_i\}$ of L i.e., U_i is open in T^*M and $U_i \cap L$ is contractible and any intersections of U_i also satisfies the same assumption. For each open subset U_i , we have a presentation $U_i \cap \rho^{-1}(L) = \bigcup_c L_{f+c} \cap U_i$ where f is some primitive of $\lambda|_L$.

Lemma 3.13. *For an object $\mathcal{E} \in \mu\mathrm{Sh}_{\rho^{-1}(L)}(U)$, the restriction $\mathcal{E}|_{L_{f+c}}$ is in $\mu\mathrm{Sh}_{L_f}|_{L_f}(U \cap L_f)$. Here U is some intersection of U_i 's.*

Proof. Then there exists a local symplectomorphism ϕ such that L is mapped to the zero section. By taking the contactization, we get a local contactomorphism (or equivalently, a local homogeneous symplectomorphism) $\tilde{\phi}$ mapping L_{f+c} to $\mathbb{R}_x^n \times \mathbb{R}_{\tau>0} \subset T^*\mathbb{R}_x^n \times \{c\} \times T^*\mathbb{R}_t$. By the quantized contact transformation [KS94], this gives an equivalence $\mu\mathrm{Sh}_{\rho^{-1}(L)}(U) \cong \mu\mathrm{Sh}_{\tilde{\phi}(\rho^{-1}(L))}(\tilde{\phi}(U))$. Then the latter category is given by the sheafification of quotient categories of

$$\{\mathcal{E} \in \mathrm{Sh}_{\tau>0}(M \times \mathbb{R}_t) \mid \mathrm{SS}(\mathcal{E}) \subset \mathbb{R}_x^n \times \mathbb{R}_t \times \mathbb{R}_{\tau>0}\}. \quad (3.9)$$

Since any object of this category is represented by an object of the form $p_M^{-1}\mathcal{E}_0$ where $p_M: \mathbb{R}_x^n \times \mathbb{R}_t \rightarrow \mathbb{R}_t$, the microstalks along $\mathbb{R}_x^n \times \{t=0\} \times \mathbb{R}_{\tau>0}$ is constant. This completes the proof. \square

In the next section, we will prove this lemma by using quantized contact transformation.

Let $C(U)$ be the Cech covering associated to $\{U_i\}$. We have a sequence of functors

$$\mu_L(T^*M) \xrightarrow{\mathfrak{f}} \mathrm{Sh}(M \times \mathbb{R}_t) \rightarrow \prod_{U \in C(U)} \mu\mathrm{Sh}_{\rho^{-1}(L)}(U) \rightarrow \prod_{U \in C(U)} \mu\mathrm{Sh}_{L_f}|_{L_f}(U \cap L_f) \rightarrow \mu\mathrm{Sh}(L) \quad (3.10)$$

where the leftmost morphism is the colimit with respect to the Cech covering. The composition is our desired microlocalization functor, and will be denoted by μ_L .

3.5 Brane structure and microlocalization

Now we can define the notion of Lagrangian brane.

Definition 3.14. A Lagrangian brane $(L, \alpha, b, \mathcal{L})$ is a tuple of the following data:

1. a graded Lagrangian submanifold L with $\alpha: L \rightarrow \widetilde{LGr}(X)|_L$
2. a relative Pin structure b of L
3. a derived local system \mathcal{L} over L .

When \mathcal{L} is the rank 1 constant local system, we simply say that (L, α, b) is a Lagrangian brane.

Remark 3.15. There are further generalizations of the notion of Lagrangian branes. See e.g. [JT].

Given a Lagrangian brane (L, α, b) , we have an equivalence $\mu\text{Sh}_L(L) \cong \text{Loc}(L)$. Composing it with μ_L , we obtain

$$\mu_{(L, \alpha, b)}: \mu_L(T^*M) \rightarrow \text{Loc}(L). \quad (3.11)$$

Definition 3.16. Let $(L, \alpha, b, \mathcal{L})$ be a Lagrangian brane. A sheaf quantization of $(L, \alpha, b, \mathcal{L})$ is an object of $\mu_L(T^*M)$ such that $\mu_{(L, \alpha, b)}(\mathcal{E}) \cong \mathcal{L}$.

More simply,

Definition 3.17. 1. Let L be a Lagrangian submanifold. A sheaf quantization of L is an object of $\mu_L(T^*M)$.

2. Let L be a Lagrangian submanifold. A pure sheaf quantization of L is an object of $\mu_L(T^*M)$ such that $\mu_{(L, \alpha, b)}(\mathcal{E})$ is concentrated in a single degree for some brane structure (L, α, b) .

3. Let L be a Lagrangian submanifold. A simple sheaf quantization of L is a pure sheaf quantization whose microlocalization is rank 1.

We also sometimes use the following terminology.

Definition 3.18. Let (L, α, b) be a Lagrangian brane. The microstalk of an object \mathcal{E} of $\mu_L(T^*M)$ at $(x, \xi) \in L$ is the stalk of $\mu_{(L, \alpha, b)}(\mathcal{E})$ at (x, ξ) . Note that the definition does not depend on b . Moreover if L is connected, the isomorphism type of microstalk does not depend on $(x, \xi) \in L$.

Remark 3.19 (Uniqueness). In the nonexact setting, the present definition of brane structure does not characterize sheaf quantization. To rigidify it, we have to include bounding cochains. We will treat it in another publication.

Remark 3.20 (Sheaf quantization in $\text{Sh}(M)$ and $\text{Sh}_{\tau>0}(M \times \mathbb{R}_t)$). (Version 0). For a conic Lagrangian submanifold L in T^*M , a sheaf quantization \mathcal{E} of L is a constructible sheaf with $SS(\mathcal{E}) = L$.

(Version 1). If L is an exact Lagrangian submanifold, we can take a primitive of $\lambda_{st}|_L$ globally on M . Then we can lift L to $L_f \subset T^*M \times T^*_{>0}\mathbb{R}_t$. An object $\mathcal{E} \in \text{Sh}_{\tau>0}(M \times \mathbb{R}_t)$ on $M \times \mathbb{R}_t$ is said to be a sheaf quantization of L if $SS(\mathcal{E}) \setminus T^*_{M \times \mathbb{R}_t} M \times \mathbb{R}_t = L_f$ and has finite-dimensional pure microstalks. Under certain assumptions on L , Guillermou and Jin–Treumann constructed such sheaves [Gui16, JT]. This notion is a generalization of Version 0 in the following sense: a sheaf quantization \mathcal{E} of a conic Lagrangian L gives a sheaf on $M \times \mathbb{R}_t$ by $\mathcal{E} \boxtimes \mathbb{K}_{[0, \infty)}$, which turns out to be a sheaf quantization (Version 1) of L .

(Version 2). Our concept of sheaf quantization generalizes both. For a sheaf quantization (Version 1) $\mathcal{E} \in \text{Sh}_{\tau>0}(M \times \mathbb{R}_t)$ of an exact Lagrangian L , consider $\bigoplus_{c \in \mathbb{R}} T_c \mathcal{E}$ with the obvious \mathbb{R}_t -equivariant structure. Then this is a sheaf quantization of L in our sense.

Example 3.21 (Graph). Let f be a smooth function $M \rightarrow \mathbb{R}$. We will denote the graph of the differential df by L . It is well-known that the microsupport of the sheaf $\mathbb{K}_{t \geq -f(x)}$ can be computed as

$$SS(\mathbb{K}_{t \geq -f(x)}) \cap \{\tau > 0\} = L_f. \quad (3.12)$$

The direct sum $\mathcal{E}_f := \bigoplus_{c \in \mathbb{R}} \mathbb{K}_{t \geq -f(x)+c} = \mathcal{I}^L(\mathbb{K}_{t \geq -f(x)})$ has $\mu\text{supp}(\mathcal{E}_f) = L$. As we have seen in Example 3.5 and Example 3.8, L is canonically equipped with a brane structure.

We would like to compute the microlocalization. The projection $\pi: L_f \rightarrow M$ is a diffeomorphism. By the Weinstein theorem, we have a symplectomorphism between a neighborhood of L_f and a neighborhood of M induced by the projection. Quantizing this symplectomorphism, \mathcal{E}_f is mapped to $\bigoplus_{c \in \mathbb{R}} \mathbb{K}_{t \geq c}$. Hence the microlocalization is \mathbb{K}_L (possibly with some shift).

4 Meromorphic flat \hbar -connection

4.1 Meromorphic flat \hbar -connection

In this section, we would like to set up our general setting for differential equations.

Let C be a Riemann surface. Let \mathcal{O}_C be the structure sheaf. We consider the subring \mathcal{D}_C^\hbar of the sheaf of $\mathbb{C}[\hbar]$ -linear endomorphisms of $\mathcal{O}_C[\hbar] := \mathcal{O}_C \otimes_{\mathbb{C}} \mathbb{C}[\hbar]$ generated by $\hbar \partial_z$ and \mathcal{O}_C . We would like to consider the modules over \mathcal{D}_C^\hbar .

Our fundamental examples of such modules are given by meromorphic flat \hbar -connections:

Definition 4.1. A meromorphic flat \hbar -connection is given by the following data

1. A meromorphic bundle \mathcal{E} with poles in $M \subset C$ where M is a finite subset of C .
2. A $\mathbb{C}[\hbar]$ -linear morphism $\nabla: \mathcal{E}[\hbar] \rightarrow \mathcal{E}[\hbar] \otimes_{\mathcal{O}_C} K_C$ where $\mathcal{E}[\hbar] := \mathcal{E} \otimes_{\mathbb{C}} \mathbb{C}[\hbar]$ and K_C is the canonical bundle which satisfies the following: For any $f \in \mathcal{O}_C$ and $s \in \mathcal{E}$, we have $\nabla(fs) = s \otimes \hbar df + f \nabla s$.

A meromorphic flat \hbar -connection (\mathcal{E}, ∇) gives a \mathcal{D}^\hbar -module as follows: The underlying $\mathcal{O}_C[\hbar]$ -module is $\mathcal{E}[\hbar]$ and the action of \mathcal{D}^\hbar is defined as follows: For a tangent vector $v \in \mathcal{D}$, we have $\hbar v \in \mathcal{D}^\hbar$. For $s \in \mathcal{E}$, we set

$$(\hbar v) \cdot s := \nabla s(v). \quad (4.1)$$

For $\hbar \in \mathbb{C}^\times$, we define a meromorphic connection $\mathcal{M}^\hbar := (\mathcal{E}, \nabla^\hbar)$ by

$$\nabla^\hbar := \frac{1}{\hbar} \cdot \nabla \otimes_{\mathbb{C}[\hbar]} \mathbb{C} \quad (4.2)$$

where the morphism $\mathbb{C}[\hbar] \rightarrow \mathbb{C}$ of the tensor is given by the evaluation at \hbar .

Remark 4.2. The notion of \hbar -connection is almost the same as Deligne's λ -connections. However, we would like to avoid the terminology in this paper, because of the existence of a non-straight-forward relations between these when one starts from a point in Hitchin's base [Gai, DFK⁺].

4.2 Spectral curve

It is well-known that \mathcal{D}_C^{\hbar} is a deformation quantization of T^*C . Namely, there exists a canonical isomorphism $\mathcal{D}_C^{\hbar}/\hbar\mathcal{D}_C^{\hbar} \cong \mathcal{O}_{T^*C}$. Let \mathcal{M} be a \mathcal{D}_C^{\hbar} -module. Then $\mathcal{M}/\hbar\mathcal{M}$ is a module over \mathcal{O}_{T^*C} . We set

$$SS(\mathcal{M}) := \text{supp}(\mathcal{M}/\hbar\mathcal{M}) \subset T^*C. \quad (4.3)$$

We sometimes call it the *spectral curve* of \mathcal{M} . The spectral curve is holomorphic coisotropic with respect to the standard symplectic structure of T^*C . In the rest of this paper, we will only consider \hbar -connections whose spectral curve is holomorphic Lagrangian.

5 Planck constant in complex and real symplectic geometry

Microlocal sheaf theory is related to real symplectic geometry. On the other hand, the concept of \hbar -connection is related to deformation quantizations of complex symplectic manifolds. In this section, we clarify the relationship.

5.1 Complex symplectic geometry and deformation quantization

Let X be a complex manifold. Then the cotangent bundle T^*X is a complex exact symplectic manifold. The canonical symplectic form ω can be locally written as $\sum d\zeta_i \wedge dz_i$. Here z_i is a local coordinate of X and ζ_i is the corresponding cotangent coordinate. The form ω is exact and has a canonical primitive λ , locally written as $\sum \zeta_i dz_i$.

We will consider the canonical quantization of T^*X from local pieces. The canonical deformation quantization of T^*X is given by the ring $\mathcal{D}^{\hbar} := \mathbb{C}[\hbar][z_i, \hbar\partial_{z_i}]$. In the classical limit, $\hbar\partial_z$ corresponds to ζ .

This gives a hint to make a Lagrangian into a conic Lagrangian: The original coordinate ζ splits into the product $\hbar\partial_z$ in the setting of quantization. Hence we will reconsider \hbar and ∂_z as independent variables. Then a Lagrangian submanifold $L \subset T^*M$ lifts to

$$\tilde{L} := \{(z, p, \hbar) \in T^*X \times \mathbb{C}_\hbar^\times \mid (z, \hbar p) \in L\}. \quad (5.1)$$

For the later use, it is convenient to work with $\eta := \hbar^{-1}$. To get a Lagrangian submanifold, we moreover consider the Fourier dual coordinate y of η . We set

$$L_f^{\mathbb{C}} := \{(z, p, \hbar, y) \in T^*X \times \mathbb{C}_\hbar^\times \times \mathbb{C}_y \mid (z, \hbar p) \in L, y = -f(z, \hbar p)\} \quad (5.2)$$

where f is a primitive of L i.e., $df = \lambda|_L$. We consider $\mathbb{C}_y \times \mathbb{C}_\hbar^\times \cong T_{\eta \neq 0}^* \mathbb{C}_y$. Then $L_f^{\mathbb{C}}$ is a Lagrangian manifold in $T^*X \times T_{\eta \neq 0}^* \mathbb{C}_y$ with respect to the standard symplectic structure and is conic i.e., invariant under the scaling action of \mathbb{C}_\hbar^\times on the fibers of $T^*X \times T_{\eta \neq 0}^* \mathbb{C}_y$.

There exists a projection

$$\rho^{\mathbb{C}}: T^*X \times T_{\eta \neq 0}^* \mathbb{C}_y \rightarrow T^*X; (z, \zeta, y, \eta) \mapsto (z, \zeta/\eta) = (z, \hbar\zeta). \quad (5.3)$$

Then the image of $L_f^{\mathbb{C}}$ is precisely L . Conversely, $L_f^{\mathbb{C}}$ is a Lagrangian leaf of $(\rho^{\mathbb{C}})^{-1}(L)$. This procedure is a classical trick in the theory of deformation quantization (for example, [PS04]).

5.2 From complex symplectic geometry to a family of real symplectic geometries

Considering X as a real manifold and denote it by $X_{\mathbb{R}}$. We have $T^*X_{\mathbb{R}}$ the real cotangent bundle equipped with the standard real symplectic structure ω_{st} .

In a holomorphic local coordinate $\{z_i\}$ of X , we have the associated real coordinate (x_i, y_i) with $z_i = x_i + \sqrt{-1}y_i$. We can further identify the real and complex cotangent bundles via

$$dx_i \mapsto dz_i, dy_i \mapsto -\sqrt{-1}dz_i. \quad (5.4)$$

Hence in the cotangent coordinate, we have $(\xi_i, \eta_i) \mapsto \xi_i - \sqrt{-1}\eta_i = \zeta_i$. This implies that $\Re(\omega) = \omega_{st}$. In the following, we will not distinguish T^*X with $T^*X_{\mathbb{R}}$.

Hence a holomorphic Lagrangian submanifold with respect to ω gives a real Lagrangian submanifold with respect to ω_{st} .

Now we are going to vary the holomorphic symplectic form. For $\hbar \in \mathbb{C}_{\hbar}^{\times}$, we have another holomorphic symplectic form ω/\hbar . This induces a new real symplectic form $\Re(\omega/\hbar)$ on T^*X . A holomorphic Lagrangian submanifold with respect to ω is also a holomorphic Lagrangian with respect to ω/\hbar . Hence it is also a real Lagrangian submanifold with respect to $\Re(\omega/\hbar)$.

If L is a holomorphic exact Lagrangian with respect to λ , then L is also a real exact Lagrangian with respect to $\Re(\lambda/\hbar)$. In this case, a real primitive function $f_{\hbar}: L \rightarrow \mathbb{R}$ of $\Re(\lambda/\hbar)|_L$ can be also considered as a real primitive function of L/\hbar with respect to $\Re(\lambda) = \lambda_{st}$.

Remark 5.1. Since microlocal sheaf theory knows only about the standard symplectic form, to speak about $\Re(\omega/\hbar)$, we will consider the following identification.

$$(T^*X, \Re(\omega/\hbar)) \cong (T^*X, \Re(\omega/\hbar)) \xrightarrow{\cdot(1/\hbar)} (T^*X, \Re \omega) \cong (T^*X, \omega_{st}) \quad (5.5)$$

Hence a sheaf quantization of a real Lagrangian submanifold L on the leftmost side is interpreted as a sheaf quantization of L/\hbar in the rightmost side.

For the Liouville form $\Re(\lambda/\hbar)$, the associated ρ can be written as follows:

$$\rho_{\hbar}: T^*X \times T_{\tau>0}^*\mathbb{R}_t \rightarrow T^*X; (z, \zeta, t, \tau) \mapsto (z, \hbar\zeta/\tau). \quad (5.6)$$

If the context is clear, we will omit the subscript \hbar .

We can compare this picture with the story of deformation quantization. We will consider $\mathbb{R}_{\tau>0}$ is a ray in $\mathbb{C}_{\eta}^{\times}$ defined by $\hbar^{-1}\mathbb{R}_{\tau>0}$. Then we set the following morphism

$$\iota: T^*X \times T_{\tau>0}^*\mathbb{R}_t \rightarrow T^*X \times T_{\eta \neq 0}^*\mathbb{C}_y; (z, \zeta, t, \tau) \mapsto (z, \zeta, \hbar t, \hbar^{-1}\tau). \quad (5.7)$$

Then $\rho^{\mathbb{C}} \circ \iota = \rho_{\hbar}$. This means ρ_{\hbar} precisely corresponds to a rotation of ρ_{st} in Example 2.29 by \hbar .

5.3 Sheaf quantizations in complex cotangent bundles

Now we redefine sheaf quantization at $\hbar \in \mathbb{C}_\hbar^\times$ as follows.

Definition 5.2. Let L be a holomorphic Lagrangian submanifold of T^*X . An object \mathcal{E} of $\mathrm{Sh}_{\tau>0}^{\mathbb{R}}(X \times \mathbb{R}_t)$ is a *sheaf quantization* of L at \hbar if it is a sheaf quantization of L/\hbar .

If one uses ρ_{\hbar} in the definition of $\mu\mathrm{supp}$, we obtain a variant of $\mu\mathrm{supp}$. We denote it by $\mu\mathrm{supp}_{\hbar}$. Then, for a sheaf quantization \mathcal{E} of L at \hbar , we have $\mu\mathrm{supp}_{\hbar}(\mathcal{E}) = L$. We denote the full subcategory of $\mathrm{Sh}_{\tau>0}^{\mathbb{R}}(X \times \mathbb{R}_t)$ spanned by the objects with $\mu\mathrm{supp}_{\hbar}(\mathcal{E}) \subset L$ by $\mu_{L,\hbar}(T^*M)$. Of course, $\mu_{L,\hbar}(T^*M) = \mu_{L/\hbar}(T^*M)$.

Fix a path from 1 to \hbar in \mathbb{C}^\times . It gives a Lagrangian isotopy from L to L/\hbar . Hence, if $(L, \alpha, b, \mathcal{L})$ is a Lagrangian brane, L/\hbar is also canonically equipped with a Lagrangian brane structure $(L, \alpha_{\hbar}, b_{\hbar}, \mathcal{L}_{\hbar})$. Hence we have the microlocalization functor $\mu_{(L, \alpha_{\hbar}, b_{\hbar}, \mathcal{L}_{\hbar})}: \mu_{L,\hbar}(T^*M) \rightarrow \mathrm{Loc}(L/\hbar)$. On the other hand, we have $\mathrm{Loc}(L) \cong \mathrm{Loc}(L/\hbar)$ by the above isotopy. By the composition, we get a functor $\mu_{(L, \alpha, b, \mathcal{L}), \hbar}: \mu_{L,\hbar}(T^*M) \rightarrow \mathrm{Loc}(L)$.

6 Sheaf quantizations associated to meromorphic flat \hbar -connections of rank 0 and 1

In this section, as a warm-up, we would like to construct sheaf quantizations associated to rank 0 and rank 1 connections. We do not need exact WKB analysis here. The material of this section can be applied to any dimensions after minor modifications. In this section, $\mathbb{K} = \mathbb{C}$.

6.1 What will we do?

For a given \mathcal{D}_C^{\hbar} -module \mathcal{M} and $\hbar \in \mathbb{C}_\hbar^\times$, what we will do here is to construct a sheaf quantization $S_{\mathcal{M}}^{\hbar}$ such that

1. $\mu\mathrm{supp}(S_{\mathcal{M}}^{\hbar}) = \mathrm{SS}(\mathcal{M})$, and
2. $S_{\mathcal{M}}^{\hbar}$ is constructed out of the solutions of \mathcal{M} .

The second requirement is not mathematically well-defined, but one can see the meaning from the construction below. The sheaf quantization we will construct below is “correct”, since it is partially recovered from \hbar -Riemann–Hilbert correspondence [Kuw22].

6.2 Rank 0

For a point $z \in C$, the cotangent fiber T_z^*C is a Lagrangian submanifold. We would like to consider deformation quantization and sheaf quantization of T_z^*C . Note that T_z^*C is

canonically equipped with a grading α_{triv} (Example 3.4) and a relative Pin structure b_{triv} (Example 3.9).

Hence, a finite set of points $\{z_i\}_{i \in I} \subset C$, the union $L := \bigcup_{i \in I} T_{z_i}^* C$ is canonically equipped with a grading α_{triv} and a relative Pin structure b_{triv} . With a constant rank 1 local system \mathbb{K}_L , we consider L as a Lagrangian brane. We then have the associated microlocalization functor $\mu_{(L, \alpha_{triv}, b_{triv}, \mathbb{K}_L)}: \mu_L(T^*C) \rightarrow \text{Loc}(L)$. Note that $L = L/\hbar$ in this case.

A connection of rank 0 means a \mathcal{D}_C^{\hbar} -module supported on a 0-dimensional subvariety. Hence it is a direct sum of the form

$$\bigoplus_{i \in I} \mathcal{D}_C^{\hbar}/(z - z_i)^{n_i} \mathcal{D}_C^{\hbar} \cong \bigoplus_{i \in I} (\mathcal{D}_C^{\hbar}/(z - z_i) \mathcal{D}_C^{\hbar})^{\oplus n_i}. \quad (6.1)$$

Here $\{z_i\}_{i \in I}$ is a subset of C .

The ‘‘solution’’ of the equation $\mathcal{D}_C^{\hbar}/(z - z_0) \mathcal{D}_C^{\hbar}$ should be the delta function. Then it is canonical to associate a sheaf

$$\bigoplus_{c \in \mathbb{R}} \mathbb{K}_{z_0} \boxtimes \mathbb{K}_{[c, \infty)} \in \text{Sh}_{\tau > 0}^{\mathbb{R}}(C \times \mathbb{R}_t) \quad (6.2)$$

with a canonical equivariant structure, to $\mathcal{D}_C^{\hbar}/(z - z_0) \mathcal{D}_C^{\hbar}$. Here \mathbb{K}_{z_0} is the skyscraper sheaf on z_0 . It is easy to see that $\mu\text{supp}_{\hbar}(\bigoplus_{c \in \mathbb{R}} \mathbb{K}_{z_0} \boxtimes \mathbb{K}_{[c, \infty)}) = T_{z_0}^* C$ which coincides with $SS(\mathcal{D}_C^{\hbar}/(z - z_0) \mathcal{D}_C^{\hbar}) = T_{z_0}^* C$.

Consequently, we get the bijection of the form

$$\bigoplus_{i \in I} \mathcal{D}_C^{\hbar}/(z - z_i)^{n_i} \mathcal{D}_C^{\hbar} \mapsto \bigoplus_{i \in I} \bigoplus_{c \in \mathbb{R}} \left(\mathbb{K}_{z_i} \boxtimes \mathbb{K}_{[c, \infty)} \in \text{Sh}_{\tau > 0}^{\mathbb{R}}(C \times \mathbb{R}_t) \right)^{\oplus n_i}. \quad (6.3)$$

We can summarize the consequence as follows:

Proposition 6.1. *Fix $\hbar \in \mathbb{C}_{\hbar}$. For any meromorphic \hbar -connection \mathcal{M} of rank 0, there exists a sheaf quantization $\mathcal{S}_{\mathcal{M}}^{\hbar}$ such that*

1. $\mu\text{supp}_{\hbar}(\mathcal{S}_{\mathcal{M}}^{\hbar}) = SS(\mathcal{M})$, and
2. the microlocalization $\mu_{(L, \alpha_{triv}, b_{triv}, \mathbb{K}_L)}(\mathcal{S}_{\mathcal{M}}^{\hbar})$ at z is pure and its rank is the same as the rank of \mathcal{M} at z for any $z \in C$.

Proof. We set

$$\mathcal{S}_{\mathcal{M}}^{\hbar} := \bigoplus_{i \in I} \bigoplus_{c \in \mathbb{R}} \left(\mathbb{K}_{z_i} \boxtimes \mathbb{K}_{[c, \infty)} \in \text{Sh}_{\tau > 0}^{\mathbb{R}}(C \times \mathbb{R}_t) \right)^{\oplus n_i}. \quad (6.4)$$

We have already seen that this satisfies the first condition. It is also easy to see the second condition. \square

Remark 6.2. Upgrading this bijection to an equivalence between categories is a subject of [Kuw22].

6.3 Rank 1

Let \mathcal{M} be a rank 1 meromorphic flat \hbar -connection with poles in $M \subset C$. On a sufficiently small open subset U in C , we can write the equation of flat section of \mathcal{M} as

$$(\hbar\partial - Q(z, \hbar))\psi = 0 \quad (6.5)$$

where $Q(z, \hbar) = \sum_{i=0}^k Q_i(z)\hbar^i$ for some finite k and each Q_i is a meromorphic function. We assume that poles of each $Q_i(z)$ is in M . In T^*U , the spectral curve $SS(\mathcal{M})$ is defined by $\xi = Q_0$. In the following, we set $L := SS(\mathcal{M}) \cap T^*(C \setminus M)$.

The meromorphic function Q_0 is globally a meromorphic 1-form. Hence the spectral curve L is canonically equipped with a grading (Example 3.5) and a relative Pin structure (Example 3.8).

For $\hbar \in \mathbb{C}^\times$, by taking a path from 1 to \hbar , we obtain a Lagrangian isotopy from L to L/\hbar . Hence we obtain a grading α_\hbar and a relative Pin structure b_\hbar on L/\hbar . With a rank 1 constant local system $\mathbb{K}_{L/\hbar}$, we have the associated microlocalization functor $\mu_{(L, \alpha_\hbar, b_\hbar, \mathbb{K}_L), \hbar}: \mu_{L, \hbar}(T^*C) \rightarrow \text{Loc}(L)$.

Take an open covering $\{U_i\}$ of $C \setminus M$ such that each intersection is contractible. Fix $\hbar \in \mathbb{C}^\times$ and a base point z_i in each U_i . We can solve the equation explicitly:

$$\psi(z, \hbar) = \exp\left(\int_{z_i}^z \frac{Q(z, \hbar)}{\hbar} dz\right) \quad (6.6)$$

On each $U_i \times \mathbb{R}_t$, we consider the equivariant sheaf $\bigoplus_{c \in \mathbb{R}} T_c \mathbb{K}_{t \geq -\mathfrak{Re} \int_{z_i}^z \frac{Q_0}{\hbar}}$ where the equivariant structure is the obvious one. We have the following:

Lemma 6.3. $\mu\text{supp}_{\hbar}(\bigoplus_{c \in \mathbb{R}} T_c \mathbb{K}_{t \geq -\mathfrak{Re} \int_{z_i}^z \frac{Q_0}{\hbar}}) = L \cap T^*U_i$. Moreover, the microlocalization along $L \cap T^*U_i$ is the constant sheaf.

Proof. By Example 3.21, we have $\mu\text{supp}_{\hbar}(\bigoplus_{c \in \mathbb{R}} T_c \mathbb{K}_{t \geq -\mathfrak{Re} \int_{z_i}^z \frac{Q_0}{\hbar}}) = (L \cap T^*U_i)/\hbar$ and the microlocalization is the constant sheaf. Multiplying \hbar , we obtain the desired result. \square

On each overlap $U_i \cap U_j$, we consider the isomorphism between $\bigoplus_{c \in \mathbb{R}} T_c \mathbb{K}_{t \geq -\mathfrak{Re} \int_{z_i}^z \frac{Q_0}{\hbar}}$ and $\bigoplus_{c \in \mathbb{R}} T_c \mathbb{K}_{t \geq -\mathfrak{Re} \int_{z_j}^z \frac{Q_0}{\hbar}}$ induced by

$$\exp\left(\int_{z_i}^{z_j} \frac{Q(z, \hbar)}{\hbar} dz\right) \cdot \text{id}: \mathbb{K}_{t \geq -\mathfrak{Re} \int_{z_i}^z \frac{Q_0}{\hbar}} \rightarrow \mathbb{K}_{t \geq -\mathfrak{Re} \int_{z_j}^z \frac{Q_0}{\hbar}} = T_{-\mathfrak{Re} \int_{z_i}^{z_j} \frac{Q_0}{\hbar}} \mathbb{K}_{t \geq -\mathfrak{Re} \int_{z_j}^z \frac{Q_0}{\hbar}}. \quad (6.7)$$

By Lemma 6.3, these gluing-up isomorphisms give a sheaf quantization of L at \hbar .

On the other hand, the localization of \mathcal{M}^{\hbar} on $C \setminus M$ defines a local system $\text{Sol}(\mathcal{M}^{\hbar})$ on $C \setminus M$. Under the projection, we have an equivalence $\pi_*: \text{Loc}(L) \rightarrow \text{Loc}(C \setminus M)$.

Proposition 6.4. Given a rank 1 meromorphic \hbar -connection \mathcal{M} as above, there exists a sheaf quantization $S_{\mathcal{M}}^{\hbar}$ of $SS(\mathcal{M}) \cap T^*(C \setminus M)$ at \hbar such that $\pi_* \mu_{(L, \alpha_\hbar, b_\hbar, \mathbb{K}_L), \hbar}(S_{\mathcal{M}}^{\hbar}) \cong \text{Sol}(\mathcal{M}^{\hbar})$.

Proof. The statement about μsupp immediately follows from Example 3.21. Locally in T^*U_i , the microlocalization is given by $\mathbb{K}_{L \cap T^*U_i}$ from Example 3.21. Each overlapping $L \cap T^*(U_i \cap U_j)$, we have two different identifications of the microlocalization with $\mathbb{K}_{L \cap T^*(U_i \cap U_j)}$. They are related by the gluing morphism: the multiplication by $\exp\left(\int_{z_i}^{z_j} \frac{Q(z, \hbar)}{\hbar} dz\right)$. Hence we get the desired local system. \square

7 Schrödinger equations

We would like to recall some basic facts of exact WKB analysis. We refer to [KT05] and [IN14] for more accounts. In this section, we concentrate on the case arising from Schrödinger equations.

Remark 7.1. Although it is probably not difficult to extend the result to more general rank 2 connections, some necessary results in exact WKB analysis is not available in the literature. Therefore we present our result in the simplest setup, which is enough to present the essence of the construction. The author's most general treatment in the previous version contains some gaps. Another approach can be found in [Kuw22].

7.1 Schrödinger equation and connection

We consider a second order differential operator \mathcal{Q} on C with a parameter \hbar locally written as

$$\mathcal{Q} = (\hbar\partial)^2 - Q(z, \hbar). \quad (7.1)$$

We call this operator *Schrödinger operator*. Here $Q(z, \hbar) = \sum_{i=0}^k Q_i(z)\hbar^i$ for some k and each Q_i is a meromorphic function. We fix a divisor M such that each Q_i is smooth outside M . Then Q_0 defines a global meromorphic quadratic differential on C i.e., a global section of $K^{\otimes 2}(*M)$ for a divisor M .

To lift a Schrödinger operator to a flat \hbar -connection, fix a square root \sqrt{K} of K , which is well-known to be equivalent to choose a spin structure. On $C \setminus M$, we can define a connection by

$$\mathcal{M} := \left(\sqrt{K} \oplus \sqrt{K}^{-1}, \hbar\partial + \begin{pmatrix} 0 & Q \\ 1 & 0 \end{pmatrix} \right). \quad (7.2)$$

We call an \hbar -connection obtained in this way a Schrödinger-type connection. We say $Q_{\mathcal{M}} := Q_0$ is the underlying quadratic differential of \mathcal{M} . We call the equation of the flat sections of this connection the Schrödinger equation of \mathcal{M} .

Remark 7.2. Over each Stokes region (defined below), the bundle \sqrt{K} is trivialized, and a flat section of the connection is equivalent to a solution of the differential operator (7.1).

7.2 WKB solutions

For a Schrödinger operator \mathcal{Q} , consider a formal solution of the form $\psi(z, \hbar) = \exp\left(\int^z \sum_{m=-1}^{\infty} \hbar^m P_m dz\right)$, then one can derive a system of equations governing $P_m(z)$'s:

$$\begin{aligned} P_{-1}^2 &= Q_0 \\ 2P_{-1}P_m + \sum_{\substack{m_1+m_2=m-1 \\ m_1, m_2 \geq 0}} P_{m_1}P_{m_2} + \frac{dP_{m-1}}{dz} &= Q_{m-1}(z). \end{aligned} \quad (7.3)$$

The first equation has two solutions: $P_{-1}^{\pm} = \pm\sqrt{Q_0}$. The second equation has unique solutions for a choice of \pm . We write these two solutions by

$$\begin{aligned} P^{\pm}(z, \hbar) &:= P_{od}^{\pm}(z, \hbar) + P_{ev}(z, \hbar) \\ P_{od}^{\pm} &:= \pm\frac{1}{2}(P^+ - P^-) \\ P_{ev} &:= \frac{1}{2}(P^+ + P^-) \end{aligned} \quad (7.4)$$

Note that P_{ev} does not depend on the choice of \pm , and P_{od}^+ differs from P_{od}^- only by the overall \pm . We also have

$$P_{ev} = -\frac{1}{2} \frac{1}{P_{od}} \frac{dP_{od}}{dz}. \quad (7.5)$$

This means $P_{ev} = \frac{\partial}{\partial z} \log(1/\sqrt{P_{od}})$ formally. Hence we can formally rewrite $\psi(z, \hbar)$ as

$$\begin{aligned} \psi_{\pm}(z, \hbar) &= \exp\left(\pm \int^z P_{od} dz\right) \exp\left(\int^z P_{ev}\right) \\ &= \frac{1}{\sqrt{P_{od}}} \exp\left(\pm \int^z P_{od}\right) \end{aligned} \quad (7.6)$$

These two solutions are called WKB solutions of (7.1).

One can expand the WKB solutions as follows:

$$\psi_{\pm}(z, \hbar) = \exp\left(\int_{z_0}^z \frac{\pm\sqrt{Q_0}}{\hbar} dz\right) \sum_{m=0}^{\infty} \hbar^{m+\frac{1}{2}} \psi_m^{\pm}(z). \quad (7.7)$$

The aim of exact WKB analysis is to lift these formal solutions to analytic solutions.

Definition 7.3 (Resummation). For a point $z \in C$ and $\mathfrak{h} \in \mathbb{C}^{\times}$, a resummation $\Psi_{\pm}(z, \hbar)$ of $\psi_{\pm}(z, \hbar)$ is defined as follows: We set

$$S_{\mathfrak{h}} := \{\hbar \in \mathbb{C}^{\times} \mid |\hbar/\mathfrak{h}| < 2, -\pi/2 < \arg(\hbar/\mathfrak{h}) < \pi/2\}. \quad (7.8)$$

A resummation of Ψ_{\pm} at (z, \mathfrak{h}) is a holomorphic function on $U \times S_{\mathfrak{h}}$ for some open neighborhood U of z such that $\Psi_{\pm}(\hbar, z)$ is asymptotically expanded to $\psi_{\pm}(z, \hbar)$ as $\hbar \rightarrow 0$.

The rest of this section is devoted to explain the known results of resummations of WKB solutions.

7.3 Stokes geometry

Let \mathcal{M} be a meromorphic flat \hbar -connection of Schödinger-type. We introduce the languages of Stokes geometry in this setting. See [IN14, BS15] for the details. We set $L := SS(\mathcal{M})$.

Definition 7.4. We fix $\hbar \in \mathbb{C}^\times$.

1. A turning point of \mathcal{M} is a branched point of L . We denote the set of turning points of \mathcal{M} by $\text{Zero}(\mathcal{M})$. Equivalently, $\text{Zero}(\mathcal{M})$ is the zero set of $Q_{\mathcal{M}}$.
2. For a turning point v of \mathcal{M} , an \hbar -Stokes curve emanating from v is a subset of the closure of

$$\left\{ z \in C \setminus M \mid \Im \left(\int_v^z \left(\frac{\xi_+(z') - \xi_-(z')}{\hbar} \right) dz' \right) = 0 \right\} \quad (7.9)$$

such that the subset is homeomorphic to a connected interval, one of the boundary is v , and the other boundary is another turning point or in M . Here ξ_+ and ξ_- are the restrictions of λ to the sheets of L . Equivalently, $\xi_{\pm} = \pm\sqrt{Q_0}$.

3. The \hbar -Stokes graph $G_{\mathcal{M}}$ of \mathcal{M} is the union of the \hbar -Stokes curves of \mathcal{M} .
4. An \hbar -Stokes region is a connected component of the complement of the closure of the \hbar -Stokes graph in C .
5. A subset of the \hbar -Stokes graph is called an \hbar -Stokes segment if it is diffeomorphic to a closed interval $[a, b] \subset \mathbb{R}$ ($a < b$) and it connects two turning points.

We would like to introduce an easy class of connections.

Definition 7.5 ([BS15]). We say a quadratic differential Q_0 is (complete) GMN if

1. the order of any pole of Q_0 is more than or equal to 2,
2. Q_0 has at least one pole and one zero,
3. every zero of Q_0 is simple.

We say a meromorphic flat \hbar -connection \mathcal{M} is GMN if the underlying quadratic differential $Q_{\mathcal{M}}$ is GMN.

For GMN connections, their Stokes regions are easy:

Theorem 7.6 (Strebel [Str84], Bridgeland–Smith [BS15]). *Suppose \mathcal{M} is GMN and the \hbar -Stokes graph of $Q_{\mathcal{M}}$ does not have any \hbar -Stokes segments. Then an \hbar -Stokes region of \mathcal{M} has one of the following forms:*

1. (*horizontal strip*). a square surrounded by four edges of the \hbar -Stokes graph. Two of four vertices are poles of $Q_{\mathcal{M}}$, and the others are turning points.
2. (*half plane*). a region surrounded by two edges of the \hbar -Stokes graph. One of the vertices is a turning point, and the others are poles of $Q_{\mathcal{M}}$.

Assumption 7.7. *In the rest of this section, we assume that the quadratic differential Q_0 is GMN.*

7.4 WKB regularity

To obtain the resummability of WKB solutions, we have to assume that the higher degree terms with respect to \hbar in the differential operator is tame in some sense.

Locally around a pole p of \mathcal{M} , the equation can be written as

$$((\hbar\partial)^2 - Q(z, \hbar))\psi = 0. \quad (7.10)$$

Assumption 7.8 (WKB-regularity). *We say \mathcal{M} is WKB-regular if, around any pole p , $Q(z, \hbar) = Q_0(z) + \hbar Q_1(z) + \hbar^2 Q_2(z) + \dots$ satisfies the following:*

1. p is a pole of Q_0 .
2. If Q_0 has a pole p of order $m \geq 3$, then the pole order of $Q_i (i \geq 1)$ at p is less than $1 + m/2$.
3. If Q_0 has a pole of order $m = 2$, the following conditions hold: Q_i has at most one simple pole at p for all $i \geq 1$ except for $n = 2$, and $Q_2(z)$ has a double pole at p and satisfies $Q_2(z) = -\frac{1}{4(z-p)^2}(1 + O(z-p))$ as $z \rightarrow p$.

This is defined in [IN14], which attributes it to Koike–Schäfke.

We would like to explain an important property of WKB-regularity.

Definition 7.9. For a singularity z of a meromorphic flat connection \mathcal{N} (especially, of a linear differential equation) on C without \hbar , the formal completion of \mathcal{N} at 0 is, after taking a branched covering $z \mapsto z^k$, isomorphic to the form

$$\bigoplus_i \mathcal{E}^{f_i} \otimes \mathcal{L}_i \quad (7.11)$$

where i runs through finite indices, each \mathcal{L}_i is regular singular, each f_i is Laurent polynomial, and \mathcal{E}^{f_i} is rank 1 connection defined by $d + df_i dz$. This is known as Hukuhara–Levelt–Turrittin theorem. The set of Laurent–Puiseux polynomial $\mathfrak{J}_{\mathcal{N},0} = \{g_i(z) := f_i(z^{1/k})\}_{i \in I}$ is uniquely determined from \mathcal{N} up to the following operations: (i) for some g_i , replace it with $g_i + g$ where g is a regular Puiseux polynomial, and (ii) Galois transformation. We say the class $\mathfrak{J}_{\mathcal{N},0}$ modulo the above operations *the irregularity type of \mathcal{N}* . The rank of the irregularity type $\mathfrak{J}_{\mathcal{N},0}$ is defined to be $\text{rank } \mathcal{N}$. Let r_i be the ramification degree of g_i . Then the rank of the irregularity type is the same as $\sum_{i \in I} r_i$.

For a fixed \hbar , associated to (7.10), we consider the equation

$$\left(\partial^2 - \frac{Q}{\hbar^2} \right) \psi = 0. \quad (7.12)$$

Proposition 7.10. *The irregularity type of (7.12) at a singularity is given by ξ_{\pm}/\hbar for any \hbar .*

Proof. In this proof, we will use the notations in Section 9. In a local coordinate around a pole, Q_0 can be written as $\xi = (\frac{c_1}{z^{n/2}} + \frac{c_2}{z})dz$ with $n \geq 2$ for some $c_1, c_2 \in \mathbb{C}$ [Str84]. Recall (or see Section 9) that the WKB solution can be written as

$$\frac{1}{\sqrt{P_{\text{odd}}}} \exp\left(\int P_{\text{odd}}\right). \quad (7.13)$$

By [IN14, Proposition 2.8], $\int P_{\text{odd},i}$ for $i \geq 0$ does not have a pole. Hence $\exp(\int P_{\text{odd},\geq 0})$ is a (non-Laurent) formal power series.

Also, we have

$$\begin{aligned} \frac{1}{\sqrt{P_{\text{odd}}}} &= \frac{1}{\sqrt{c_1 z^{-n/2} + c_2 z^{-1} + P_{\geq 1, \text{odd}}}} \\ &= \frac{1}{z^{-n/4}} \frac{1}{\sqrt{c_1 + c_2 z^{n/2-1} + z^{n/2} P_{\geq 1, \text{odd}}}} \end{aligned} \quad (7.14)$$

Again, by [IN14, Proposition 2.8], $z^{n/2} P_{\geq 1, \text{odd}}$ is a formal power series. Hence the WKB solution is

$$\exp\left(\int (c_1 z^{-n/2} + c_2 z^{-1})/\hbar\right) F \quad (7.15)$$

where F is a (non-Laurent) Puiseux series. This confirms the claim. \square

7.5 Resummation

To implement the Borel resummation, we first expand (7.6) as

$$\psi_{\pm}(z, \hbar) = \exp\left(\int_{z_0}^z \frac{\xi_{\pm}}{\hbar} dz\right) \sum_{m=0}^{\infty} \hbar^{m+\frac{1}{2}} \psi_m^{\pm}(z). \quad (7.16)$$

The formal Borel transform with respect to \hbar^{-1} is

$$\psi_{\pm, B}(z, y) = \sum_{m=0}^{\infty} \frac{\psi_m^{\pm}(z)}{\Gamma(m + \frac{1}{2})} (y \pm a(z))^{m+\frac{1}{2}} \quad (7.17)$$

where $a(z) := \int_{z_0}^z \sqrt{Q} dz$. This series is convergent. We set

$$\mathcal{S}[\psi_{\pm}](z, \hbar) := \int_{\mp a(z)}^{\infty} e^{-y/\hbar} \psi_{\pm, B}(z, y) dy. \quad (7.18)$$

Theorem 7.11 (Koike–Schäfke, [IN14, Theorem 2.18]). *Suppose \mathcal{M} is GMN, WKB-regular, and of Schrödinger-type. Take $\theta \in \mathbb{R}/2\pi\mathbb{Z}$. Suppose that the $e^{i\theta}$ -Stokes graph of \mathcal{M} does not have any $e^{i\theta}$ -Stokes segments. Fix $z \in C \setminus G_{\mathcal{M}}$. Then there exists $\hbar \in \mathbb{C}_{\hbar}^{\times}$ such that $\arg \hbar = \theta$, $\psi_{\pm, B}(z, y)$ is convergent in a Stokes region D containing z , and $\mathcal{S}[\psi_{\pm}]$ is a resummation of ψ_{\pm} in $D \times \mathcal{S}_{\hbar}$.*

Remark 7.12. The sketch of the proof of the above theorem is available in Takei [Tak17], which is a generalization of the argument in [DLS93].

7.6 Voros' formula

We would like to describe a connection formula known as Voros' formula. Let \mathcal{M} be a meromorphic flat \hbar -connection of Schrödinger-type. We denote the underlying spin structure by \sqrt{K} .

We denote the spectral curve of the connection by L . Considering $Q_0^{1/4}$, as a section of \sqrt{K} , we have a (topological) branched covering $\tilde{\pi}: \tilde{L} \rightarrow L$ branched at the branch points of $\pi: L \rightarrow C$ as the Riemann surface of \sqrt{K} . Since (7.6) involves $Q_0^{1/4}$, to fix a branch of (7.6), it is necessary to designate a sheet of \tilde{L} .

Now we would like to introduce Voros' connection formula. Let D_1, D_2 be adjacent Stokes regions. Let z_0 be a turning point on the Stokes curve separating D_1 and D_2 . We will use the following normalization in the expression (7.6). Fix $z \in D_1$ and $(z, \zeta) \in \pi^{-1}(z) \subset L$. Let us denote the corresponding branch of P_{od} by $P_{od}^{(z, \zeta)}$. Let γ_z be a minimal counterclockwise loop starts from (z, ζ) surrounding z_0 on L . Then we set

$$\int_{z_0}^{(z, \zeta)} P_{od} dz := \frac{1}{2} \int_{\gamma_z} P_{od}^{(z, \zeta)} dz. \quad (7.19)$$

This is called the turning point normalization at z_0 . We also fix a branch of $\sqrt{P_{od}}$, which is given by a choice of $\tilde{z} \in \tilde{\pi}^{-1}(z, \zeta)$. We will write the corresponding resummed solutions on D_1 by $\Psi_{z_0, \tilde{z}, 1}^{D_1}$ i.e.,

$$\Psi_{z_0, \tilde{z}, 1}^{D_1} := \frac{1}{(\sqrt{P_{od}})_{\tilde{z}}} \exp \left(\int_{z_0}^{(z, \zeta)} P_{od} dz \right). \quad (7.20)$$

We also take a lift of γ_z to \tilde{L} starting from \tilde{z} and denote the endpoint by \tilde{z}' . We set

$$\Psi_{z_0, \tilde{z}, 2}^{D_1} := \frac{1}{(\sqrt{P_{od}})_{\tilde{z}'}} \exp \left(\int_{z_0}^{\tilde{\pi}(\tilde{z}')} P_{od} dz \right). \quad (7.21)$$

This is also understood to be resummed. We set $\Psi_{z_0, \tilde{z}}^{D_1} := (\Psi_{z_0, \tilde{z}, 1}^{D_1}, \Psi_{z_0, \tilde{z}, 2}^{D_1})$.

Let γ_{12} be an interval which starts from z , ends at $z_2 \in D_2$, and intersects with the Stokes graph exactly once at a point on the Stokes curve separating D_1 and D_2 . Take the lift $\tilde{\gamma}_{12}$ of γ_{12} to \tilde{L} starting from \tilde{z} . We denote the end point of $\tilde{\gamma}_{12}$ by \tilde{z}_2 . Then we define $\Psi_{z_0, \tilde{z}_2}^{D_2}$ in the same way as $\Psi_{z_0, \tilde{z}}^{D_1}$ replacing \tilde{z} with \tilde{z}_2 .

Theorem 7.13 (Voros [Vor83], Aoki–Kawai–Takei [AKT91]). *The analytic continuation of $\Psi_{z_0, \tilde{z}}^{D_1}$ crossing the Stokes curve is related to $\Psi_{z_0, \tilde{z}_2}^{D_2}$ by*

$$\Psi_{z_0, \tilde{z}}^{D_1} = \Psi_{z_0, \tilde{z}_2}^{D_2} \cdot S \quad (7.22)$$

where

$$S = \begin{cases} \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix} & \text{if } \int_v^z \frac{\sqrt{Q_0}}{\hbar} dz > 0 \text{ on the Stokes curve,} \\ \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} & \text{otherwise.} \end{cases} \quad (7.23)$$

On a horizontal strip D , we have two turning points in its closure. We denote them by v_1, v_2 . Take a path $\gamma_{v_1 v_2}$ in D connects v_1 and v_2 . Take a lift $\widetilde{\gamma}_{v_1 v_2}$ of $\gamma_{v_1 v_2}$ on \widetilde{L} . For $\tilde{z} \in \widetilde{\gamma}$, we have two solutions normalized at v_1 and v_2 and denote them by $\Psi_{v_1, \tilde{z}}^D$ and $\Psi_{v_2, \tilde{z}}^D$. Then we have $\Psi_{v_1, \tilde{z}}^D = \exp(\int_{\widetilde{\gamma}_{v_1 v_2}} P_{od} dz) \Psi_{v_2, \tilde{z}}^D$.

8 Sheaf quantizations associated to meromorphic flat \hbar -connection of rank 2 of Schrödinger type

In this section, \mathcal{M} is a meromorphic flat \hbar -connection satisfying the following assumption:

Assumption 8.1. \mathcal{M} is GMN, WKB-regular, and of Schrödinger-type.

We denote the underlying Schrödinger operator by \mathcal{Q} , and the spin structure by \sqrt{K} . We also set $\mathbb{K} = \mathbb{C}$ in this section.

8.1 Construction

We will first construct our sheaf quantization on a Stoke region D . Then $\widetilde{L}|_D \rightarrow D$ is a trivial covering of degree 4.

For each Stokes edge e , we take a thickening of it D_e . Namely, D_e is a small contractible open neighborhood of the relative interior of e in $C \setminus (M \cup \text{Zero}(\mathcal{M}))$.

Fix a sheet i . We set

$$s_L^v|_{D,i} := \bigoplus_{\pm} \mathbb{K}_{D_{\pm}^v} \quad (8.1)$$

where

$$D_{\pm}^v := \left\{ (z, t) \in D' \times \mathbb{R}_t \mid t \geq \mp \Re \int_v^z (\sqrt{Q_0}/\hbar) dz \right\}, \quad (8.2)$$

v is a turning point in the closure of D , and D' is an open subset which is a union of D and the thickenings of the Stokes edges of D . This object does not depend on i , but we will keep the indices. We set

$$S_L^v|_{D,i} := \bigoplus_{c \in \mathbb{R}} T_c s_L^v|_{D,i}. \quad (8.3)$$

This defines an object of $\text{Sh}_{\tau > 0}^{\mathbb{R}}(D' \times \mathbb{R}_t)$ with an obvious equivariant structure.

We would like to list up the gluing-up isomorphisms:

1. (Change of sheets) For a sheet i , we denote the sheet where we will arrive by one counter clockwise loop around v by i' , Then we define

$$\varphi_{ii'} : S_L^v|_{D,i} \rightarrow S_L^v|_{D,i'} \quad (8.4)$$

by T : namely, the multiplication by $\sqrt{-1}$.

2. (Change of normalizations) For $v_1, v_2 \in D$, the isomorphism $\varphi_{v_1 v_2}: S_L^{v_1}|_{D,i} \rightarrow S_L^{v_2}|_{D,i}$ is induced by the scalar multiplication

$$\exp\left(\int_{\gamma_{v_1, v_2}} P_{od}\right) \cdot (-): T_c \mathbb{K}_{D_\pm^{v_1}} \rightarrow T_{c \mp \Re \int_{\gamma_{v_1 v_2}} (\sqrt{Q_0}/\hbar) dz} \mathbb{K}_{D_\pm^{v_2}} \quad (8.5)$$

where $\int_{\gamma_{v_1, v_2}} P_{od}$ is understood as its resummation.

3. (Change of regions) For adjacent Stokes regions D_1 and D_2 , the Stokes edge e separating two regions satisfies $\Re \int_v^x \sqrt{Q_0}/\hbar dz < 0$ or $\Re \int_v^x \sqrt{Q_0}/\hbar dz > 0$.

We set

$$(D_e)_\pm^v := \left\{ (z, t) \in D_e \times \mathbb{R}_t \mid t \geq \mp \Re \int_v^z (\sqrt{Q_0}/\hbar) dz \right\}, \quad (8.6)$$

which is a subset of $(D_1)_\pm^v \cap (D_2)_\pm^v$.

Then we have

$$\text{End}\left(\bigoplus_{\pm} \mathbb{K}_{(D_e)_\pm^v}\right) = \mathbb{K} \cdot \text{id}_+ \oplus \mathbb{K} \cdot \text{id}_- \oplus \mathbb{K} \cdot e \quad (8.7)$$

where $\text{id}_\pm \in \text{End}\left(\mathbb{K}_{(D_e)_\pm^v}\right)$ are the identities of the components and

$$e \in \begin{cases} \text{Hom}\left(\mathbb{K}_{(D_e)_+^v}, \mathbb{K}_{(D_e)_-^v}\right) & \text{if } \Re \int_v^x \sqrt{Q_0}/\hbar dz > 0 \\ \text{Hom}\left(\mathbb{K}_{(D_e)_-^v}, \mathbb{K}_{(D_e)_+^v}\right) & \text{if } \Re \int_v^x \sqrt{Q_0}/\hbar dz < 0 \end{cases} \quad (8.8)$$

is a canonical nontrivial basis coming from the inclusion of the closed sets. We can reflect the Voros connection formula using this:

$$\varphi_{D_1, D_2} := \text{id}_+ + \text{id}_- - e \in \text{End}\left(\bigoplus_{\pm} \mathbb{K}_{(D_1 \cap D_2)_\pm^v}\right). \quad (8.9)$$

We consider this morphism as an isomorphism $\varphi: S_L^v|_{D_1, i}|_{D_1 \cap D_2} \rightarrow S_L^v|_{D_2, i}|_{D_1 \cap D_2}$.

Lemma 8.2. *The sheaves $S_L^v|_{D, i}$'s can be glued up by φ 's and give an object in $\text{Sh}_{\tau > 0}^{\mathbb{R}}((C \setminus (M \cup \text{Zero}(\mathcal{M}))) \times \mathbb{R}_t)$. We denote this object by $S_{\mathcal{M}}^{\hbar}|_{C \setminus (M \cup \text{Zero}(\mathcal{M}))}$.*

Proof. This is clear from the consistency of the connection formulas. \square

We would like to extend the object to $C \setminus M$. Around a turning point, we have three Stokes regions D_1, D_2, D_3 . Hence $s_L^v|_{D_j, i}$ for $j = 1, 2, 3$ will glue up to a sheaf s_L^v and $S_L^v|_{D_1 \cup D_2 \cup D_3} = \bigoplus_{c \in \mathbb{R}} T_c s_L^v$. Note that there are no monodromies of the solutions, hence we have

$$\text{Ext}^1\left(\bigoplus_{c \in \mathbb{R}} \mathbb{K}_v \boxtimes \mathbb{K}_{[c, \infty)}^{\oplus 2}, \bigoplus_{c \in \mathbb{R}} T_c s_L^v\right) = \Lambda_0^{\oplus 2}. \quad (8.10)$$

where \mathbb{K}_v is the skyscraper sheaf. By using $1 \oplus 1$ in this extension space for each turning point, we get an object on $(C \setminus M) \times \mathbb{R}_t$. By the zero extension, we get an object $S_{\mathcal{M}}^{\hbar} \in \text{Sh}_{\tau > 0}^{\mathbb{R}}(C \times \mathbb{R}_t)$. As a summary, we have the following:

Theorem 8.3. *We have a simple sheaf quantization $S_{\mathcal{M}}^{\hbar} \in \text{Sh}_{\tau>0}^{\mathbb{R}}(C \times \mathbb{R}_t)$ of L at \hbar .*

Proof. The microsupport estimate follows from Lemma 6.3. Outside the turning points, we the object is simple by the computation from Example 3.21. This implies the global simpleness, since the notion is about derived local systems. \square

8.2 Voros symbol

We would like to recall the notion of Voros symbol. We refer to [DDP93] and [IN14] for more accounts.

The “dual” of a segments-free \hbar -Stokes graph gives an ideal triangulation of C with vertices in M . In this situation, $H_1(L, \mathbb{Z})$ is spanned by the following [BS15]:

1. The pull back of cycles in $C \setminus M$.
2. In an \hbar -Stokes region of horizontal strip type, take a path connecting two turning points. This is lifted to a cycle in the spectral curve. We call this cycle, a *Voros edge cycle*.

Definition 8.4. For a cycle $\gamma \in H_1(L, \mathbb{Z})$, the Voros symbol is defined by

$$V_{\gamma}(\hbar) := \oint_{\gamma} P_{od}(\gamma, \hbar). \quad (8.11)$$

The local system on L whose monodromy along γ given by $e^{V_{\gamma}(\hbar)}$ is called the \hbar -Voros local system.

In [IN14], it was proved that the collection $\{e^{V_{\gamma}(\hbar)}\}$ is a set of cluster variables. Regarding this aspect, we sometimes call the set $\{e^{V_{\gamma}(\hbar)}\}$ (or the Voros local system) *Voros–Iwaki–Nakanishi coordinate*.

Note that there is another class of Voros symbols associated to *paths*. We do not recall it here, since we do not use it in this paper.

8.3 Microlocalization

We would like to equip L with a Maslov grading and a relative Pin structure.

We will use the standard Maslov map α on T^*C [NZ09]. Then the zero section is canonically equipped with a Maslov grading $\tilde{\alpha} = 0$. For any Riemann surface, the second Stiefel–Whitney class always vanishes. Hence the notion of relative Pin structures and Spin structures coincide.

Let B be the set of branched points in L . Since we have a trivial grading on the zero section, we can induce a grading on $L \setminus B$. In other words, we have a lift of $L \setminus \pi^{-1}(M) \rightarrow U(1)$ to $L \setminus B \rightarrow \mathbb{R}$. The obstruction class $c \in H^1(L, L \setminus B, \mathbb{Z})$ to extend this lift to L is zero, since the grading can be extended to $B = \text{Zero}(\mathcal{M})$ on the base space. Hence we have a unique extension.

For a spin structure, we use $\sqrt{K}|_{C \setminus M}$ to have a spin structure on $L \setminus B$. We have a diagram

$$\begin{array}{ccc} L \setminus B & \longrightarrow & B\text{Spin}(2) \\ \downarrow & & \downarrow \\ L & \longrightarrow & B\text{SO}(2). \end{array}$$

The obstruction to extend the spin structure lives in $H^2(L, L \setminus B, \mathbb{Z}/2\mathbb{Z})$. Again, the obstruction vanishes. The classes of the extensions is a torsor over $H^1(L, L \setminus B, \mathbb{Z}/2\mathbb{Z})$. We will choose one in the proof of the following corollary.

Corollary 8.5. *The microlocalization of $\mathcal{S}_{\mathcal{M}}^{\hbar}$ along a brane structure is the Voros local system.*

Proof. We explain with Picture 8.1. In this picture, v, w are turning points. Lines emanating from turning points are Stokes lines.

The inverse image of the dotted line connecting v and w along the projection $\pi_L: L \rightarrow C$ is a Voros edge cycle. We would like to compute the monodromy of the microlocalization of $\mathcal{S}_{\mathcal{M}}^{\hbar}$ along the Voros edge cycle.

Take a circle γ' encircling the dotted line. Take a connected component of $\pi_L^{-1}(\gamma')$, and name it γ . The γ is projected down to the circle in Figure 8.2. We denote the sheet of $L|_{D_3}$ above the dotted line over where γ is lying by L_{3+} . Below the dotted line, γ is lying over the other sheet L_{3-} . Running around the tuning points change the sheets.

Since γ is homotopy equivalent to the Voros edge cycle, we compute the monodromy along γ . we can reduce the monodromy computation. Let us first consider over D_3 with




Figure 8.1: Stokes diagram

the normalization v . Namely, we consider the trivialization $\mathcal{S}_{\mathcal{M}}^{\hbar}|_{D_3} \cong \bigoplus_{c \in \mathbb{R}} T_c s_L^v|_{D,i}$. Then we get the rank 1 constant sheaf $\mathbb{K}_{L_{3+}}$ on L_{3+} by the microlocalization as in Example 3.21. Next, we consider over D_3 with the normalization w . We similarly get $\mathbb{K}_{L_{3+}}$ as a result of the microlocalization. By (8.5), these two constant sheaves are glued by the multiplication of $\exp\left(\int_{\gamma_{vw}} P_{od}\right)$.




Figure 8.2: Edge cycle

Similarly, over D_5 and D_4 , we get the rank 1 constant sheaves $\mathbb{K}_{L,4+}$ and $\mathbb{K}_{L,5+}$ under the microlocalization. From the Voros connection formula, they are glued with $\mathbb{K}_{L,3+}$ possibly multiplied with ± 1 .

Returning to v , since we are now going back over the other sheet, the sheaves are glued by the multiplication of $\exp\left(\int_{\gamma_{vw}} P_{od}\right)$. Again, tuning around the turning point possibly multiply ± 1 .

Hence we get the Voros monodromy up to sign. One can fix the sign by retaking the spin structure. \square

9 Irregular Riemann–Hilbert correspondence

In this section, we would like to relate our sheaf quantization to some known formalisms of irregular Riemann–Hilbert correspondence. In the following, we fix a sufficiently small $\hbar \in \mathbb{C}^\times$. In this section, it is convenient to think $S_{\mathcal{M}}^{\hbar}$ as a sheaf quantization of L/\hbar at $\hbar = 1$ with respect to the standard symplectic structure.

9.1 D’Agnolo–Kashiwara’s formalism

We would like to relate our sheaf quantization to D’Agnolo–Kashiwara’s holonomic Riemann–Hilbert correspondence [DK16]. For the details, we refer to [DK16, KS16].

We briefly recall the statement of [DK16]. For a complex manifold X , let us denote the derived category of holonomic \mathcal{D} -modules by $D_{hol}^b(\mathcal{D}_X)$.

For the topological side, we introduce the notion of enhanced ind-sheaves. Let $\overline{\mathbb{R}}$ be the two point compactification of \mathbb{R} i.e., $\mathbb{R} \cong (0, 1) \hookrightarrow [0, 1] = \overline{\mathbb{R}}$. The category of enhanced ind-sheaves is defined in two steps: First, we set

$$D^b(\mathrm{IC}_{X \times (\overline{\mathbb{R}}, \mathbb{R})}) := D^b(\mathrm{IC}_{X \times \overline{\mathbb{R}}}) / D^b(\mathrm{IC}_{X \times \overline{\mathbb{R}} \setminus \mathbb{R}}) \quad (9.1)$$

where $D^b(\mathrm{IC}_X)$ is the bounded derived category of ind-sheaves over X . We set $\mathbb{C}_{t \leq 0} :=$

$\mathbb{C}_{\{(x,t) \in X \times \overline{\mathbb{R}} \mid t \in \mathbb{R}, t \leq 0\}}$. We denote the convolution product along \mathbb{R}_t by $\overset{+}{\otimes}$. We set

$$\mathrm{IC}_{t^*=0} := \left\{ K \mid K \overset{+}{\otimes} \mathbb{C}_{\leq 0} \simeq 0, K \overset{+}{\otimes} \mathbb{C}_{\geq 0} \simeq 0 \right\}. \quad (9.2)$$

The category of enhanced ind-sheaves over X is defined by

$$E^b(\mathrm{IC}_X) := D^b(\mathrm{IC}_{X \times (\overline{\mathbb{R}}, \mathbb{R})}) / \mathrm{IC}_{t^*=0}. \quad (9.3)$$

We set

$$\mathbb{C}_X^E := \text{“lim”}_{a \rightarrow \infty} \mathbb{C}_{t \geq a} \quad (9.4)$$

as an object of $E^b(\mathrm{IC}_X)$. As usual, “lim” means Ind-colimit. An object of $E^b(\mathrm{IC}_X)$ is said to be \mathbb{R} -constructible if there locally exists an \mathbb{R} -constructible sheaf \mathcal{E} and the object is isomorphic to $\mathcal{E} \overset{+}{\otimes} \mathbb{C}_X^E$.

Theorem 9.1 ([DK16]). *There exists a contravariant fully faithful functor $\mathrm{Sol}^E: D_{hol}^b(\mathcal{D}_X) \hookrightarrow E^b(\mathrm{IC}_X)$.*

Let us construct enhanced ind-sheaves by modifying our construction of sheaf quantization.

Recall that our construction of sheaf quantization consists of $S_L^v|_{D,i}$ and gluing morphisms. We replace $S_L^v|_{D,i}$ with $s_L^v|_{D,i} \overset{+}{\otimes} \mathbb{C}_M^E$. We use the same gluing morphisms. We denote the resulting enhanced ind-sheaf by $E(S_{\mathcal{M}}^{\hbar})$.

Proposition 9.2. *We have $\mathrm{Sol}^E(\mathcal{M}^{\hbar}) \cong E(S_{\mathcal{M}}^{\hbar})$.*

Proof. The sheaf is glued up by using the connection matrices of resummed WKB solutions of \mathcal{M}^{\hbar} . Around the singularity, the connection matrices are precisely Stokes matrices, since Proposition 7.10 tells us that the resummed WKB solutions form a Stokes frame. Hence the construction of [DK16] tells us that this is precisely $\mathrm{Sol}^E(\mathcal{M}^{\hbar})$. \square

Remark 9.3. It is desirable to make our construction here functorial.

9.2 Irregularity knots and non-Novikovized quantization

We next compare our sheaf quantization with the Legendrian knot description of Stokes data [STWZ19]. In this section, we set $L_{\mathcal{M}} := L$.

Let us fix a divisor M on C . Fix a positive integer k . For each point $p \in M$, we fix an irregularity type of rank k . We denote this data F_M . Consider meromorphic connections with the given irregularity type F_M .

We first define the notion of irregularity knot. For $p \in M$, we denote the irregularity type of p by $\{g_1, \dots, g_k\}$. Take a small disk D_p around the singularity and fix a coordinate z on the disk. For each g_i , we give an immersion of $\sqcup_k S^1$ in D_p by

$$\bigcup_{i=1}^k \left\{ (\theta, \epsilon \cdot \Re(g_i(e^{\sqrt{-1}\theta}/r))) \mid \theta \in S^1 \right\} \quad (9.5)$$

in the polar coordinate. Here ϵ is taken sufficiently small so that the immersion is into D_p and r is some large positive number. Putting the outward conormals on the immersion, we have a Legendrian link on the cosphere bundle S^*D_p . For a sufficiently large r , the Legendrian isotopy class of the link does not depend on r . We call this link, the irregularity knot and denote it by $L_{\{g_i\}}$. We also identify the Legendrian link with the associated conical Lagrangian in $T^*D_p \subset T^*C$.

To sum up, associated to the given data of irregularity types F_M , we get a conical Lagrangian L_{F_M} contained in $\bigcup_{p \in M} T^*D_p \subset T^*C$.

We denote the following category by $\text{Sh}_{L_{F_M}}^1(C)$: The subcategory of constructible sheaves over C consisting of the objects satisfying the following conditions:

- Assumption 9.4.**
1. *microsupports are contained in $L_{F_M} \cup T_C^*C$,*
 2. *microstalks are rank 1 and concentrated in degree 0,*
 3. *stalks are zero at any $p \in M$.*

Theorem 9.5 ([STWZ19]). *There exists an equivalence between the category of meromorphic connections with the irregularity types of F_M and $\text{Sh}_{L_{F_M}}^1(C)$. We denote the equivalence functor by Sol_{STWZ} .*

We relate this equivalence with our sheaf quantization by slightly modifying the construction of $S_{\mathcal{M}}^{\hbar}$ around the poles.

Let p be a pole of the differential equation and take a local coordinate z around p . Then we consider the following curved cylinder

$$C_p := \{(z, t) \mid |z - p| \leq \eta\} \cup \left\{ (z, t) \mid \begin{array}{l} |z - p| > \eta, 0 < \epsilon(|z - p| - \eta) < \pi, \\ t \geq \tan(-\pi/2 + \epsilon(|z - p| - \eta)) \end{array} \right\} \subset C \times \mathbb{R}_t \quad (9.6)$$

where ϵ (resp. η) is sufficiently large (resp. small) positive number. On each Stokes region D , we take

$$D^{\pm} := \pi^{-1}\pi\left(\left\{(x, t) \in D \times \mathbb{R}_t \mid t \geq -\Re \int \frac{\xi_{\pm}}{\hbar} dz\right\} \setminus \bigcup_{p \in \overline{D}} C_p\right) \cap \left\{(x, t) \in D \times \mathbb{R}_t \mid t \geq -\Re \int \frac{\xi_{\pm}}{\hbar} dz\right\}. \quad (9.7)$$

where $\pi: D \times \mathbb{R} \rightarrow D$ is the projection. Figure 9.1 is an example of the situation where the gray shaded region is an example of D^{\pm} . We set

$$S_{L_c}^D := \bigoplus_{c \in \mathbb{R}} T_c \left(\bigoplus_{\pm} \mathbb{K}_{D^{\pm}} \right). \quad (9.8)$$

We can glue up this by the same rule as in the construction of $S_{\mathcal{M}}^{\hbar}$. We denote the resulting sheaf $S_{\mathcal{M},c}^{\hbar}$ and we set $L_{\mathcal{M},c} := \mu\text{supp}(S_{\mathcal{M},c}^{\hbar})$. Note that $L_{\mathcal{M},c}$ is conical and equal to L_{F_M} outside some neighborhood of the zero section.

Proposition 9.6.

1. *There exists a Lagrangian homotopy between $L_{\mathcal{M}}/\hbar$ and $L_{\mathcal{M},c}$.*




Figure 9.1: An example of D^\pm

2. The microlocalization of $S_{\mathcal{M},c}^{\hbar}$ is the same as the microlocalization of $S_{\mathcal{M}}^{\hbar}$ along the Lagrangian homotopy.

Proof. 1. Since, the both $L_{\mathcal{M}}$ and $L_{\mathcal{M},c}$ are Lagrangian homotopic to $L_{\mathcal{M}} \setminus T^* \pi(\mathcal{C})$ where $\mathcal{C} = \bigcup_p \mathcal{C}_p$.

2. The microlocalization is determined by the microlocalization at $L_{\mathcal{M}} \setminus T^* \pi(\mathcal{C})$. \square

To relate $S_{\mathcal{M},c}^{\hbar}$ to the STWZ functor, we consider the following construction. Let \mathcal{E} be an \mathbb{R} -equivariant sheaf on $C \times \mathbb{R}_t$. For an open subset U of C , we set

$$\mathcal{E}'_{\Lambda}(U) := \text{Hom}_{\text{SH}_{\tau > 0}^{\mathbb{R}}(U \times \mathbb{R}_t)} \left(\bigoplus_{c \in \mathbb{R}} \mathbb{K}_{U \times [c, \infty)}, \mathcal{E}|_U \right) \otimes_{\Lambda_0} \Lambda. \quad (9.9)$$

This assignment defines a presheaf on C valued in Λ -vector spaces. We denote the sheafification of \mathcal{E}'_{Λ} by \mathcal{E}_{Λ} .

Proposition 9.7. *The sheaf $(S_{\mathcal{M},c}^{\hbar})_{\Lambda}$ is isomorphic to $\text{Sol}_{\text{STWZ}}(\mathcal{M}^{\hbar}) \otimes_{\mathbb{C}} \Lambda$.*

Proof. Note that $\pi(\{t = \Re \int \frac{\xi_{\pm}}{\hbar}\} \cap \partial \mathcal{C})$ coincides with the projection of the irregularity knot. The gluing maps are responsible for Stokes phenomena. \square

10 Spectral network and higher order differentials

For higher (≥ 3) order differential equations with \hbar , exact WKB analysis is not so well-established, because of collisions of the Stokes curves. However, the algebraic procedure which we need to construct sheaf quantizations can be also carried out in this case under some hypothesis. The main idea is the use of BPS mass filtration [GMN13] as a grading like in the theory of scattering diagrams [KS06, GS11].

10.1 Stokes geometry of higher order connections

Let us consider a rank n meromorphic flat \hbar -connection \mathcal{M} with $n \geq 3$. We set $L := SS(\mathcal{M})$. Let M be the pole divisor of \mathcal{M} .

We would like to introduce Stokes geometry for higher order differentials.

Definition 10.1. A point z in C is a (*ordinary*) *turning point* if the spectral curve is branched over z .

In the following, we will assume that every branch point of L is a double branch point. We fix $\hbar \in \mathbb{C}_\hbar^\times$. Locally, we name the branches by $1, \dots, n$. Then a turning point is a collision of two sheets. If sheets named by $i, j \in \{1, \dots, n\}$ collide, we say this turning point is of type $\{i, j\}$. A Stokes curve emanating from an $\{i, j\}$ -turning point v can be similarly defined:

Definition 10.2. An \hbar -Stokes curve emanating from v is a subset of the closure of

$$\left\{ z \in C \setminus M \mid \Im \left(\int_v^z \left(\frac{\xi_i(z') - \xi_j(z')}{\hbar} \right) dz' \right) = 0 \right\} \quad (10.1)$$

such that the subset is homeomorphic to a connected interval, one of the boundary is v , and the other boundary is another ordinary turning point or in M . Here ξ_i means the restriction of λ to the i -th branch of L . We say that this is an \hbar -Stokes curve of type (i, j) if $\Re \int_v^z (\zeta_i - \zeta_j) dz > 0$. Otherwise (i.e., $\Re \int_v^z (\zeta_i - \zeta_j) dz < 0$), is of type (j, i) .

Definition 10.3 (Ordered collision). 1. An intersection of two Stokes curves is said to be *ordered* if one of them is of type (i, j) and the other is of type (j, k) .

2. We say a point $p \in C$ is a *cyclically ordered collision* if there exists a set of Stokes curves l_1, l_2, \dots, l_k passing through p and the types of l_1, l_2, \dots, l_k are of the form $(i_1, i_2), (i_2, i_3), \dots, (i_{k-1}, i_k), (i_k, i_1)$.

Note that the above Stokes geometry only depends on the spectral curve L .

Definition 10.4. We say L is \hbar -*tame* if

1. All the turning points are double branches.
2. The set of turning points is finite.
3. Any \hbar -Stokes curve does not have an ordered collision in the set of turning points.
4. There are no cyclically ordered collisions of \hbar -Stokes curves.
5. The complement of the union of \hbar -Stokes curves is open and contractible.
6. The set of ordered collisions of \hbar -Stokes curves is discrete in $C \setminus M$.

10.2 Berk–Nevins–Roberts’ example

We would like to explain a baby version of the construction using Berk–Nevins–Roberts’ example [BNR82], which treats the following equation

$$\left(\hbar^3 \frac{d^3}{dz^3} + 3\hbar \frac{d}{dz} + 2\sqrt{-1}z \right) \psi(z) = 0 \quad (10.2)$$




Figure 10.1: Ordinary Stokes curves of BNR equation

This example has the $\hbar = 1$ -Stokes curve in Figure 10.2. There are only two turning points, the left and right trivalent nodes. Assume that the Voros connection formula holds in this setup for the pair of the resummed WKB solutions corresponding to the Stokes curve. Then Berk–Nevins–Roberts observed nontrivial monodromies around the collision points, which should not occur. Hence the naive application of the Voros formula is wrong in this setup. We explain it in our languages.

Let v_1 be the left turning point, v_2 be the right turning point, and c be the upper collision point c . The Stokes curve emanating from v_1 (resp. v_2) passing through c is of type (21) (resp. (32)).

Let us put $\bigoplus_{c \in \mathbb{R}} T_c \bigoplus_i \mathbb{K}_{t \geq -\Re \int^z \zeta_i}$ on each connected component of the complement of the Stokes curves. We will use the Voros formula to glue up the sheaves as in the case of order 2. Around c , let us compose the gluing morphisms along a counterclockwise loop. We restrict the composition to ζ_3 -component, then we have

$$\begin{aligned} \mathbb{K}_{t \geq -\Re \int_c^z \zeta_3} &\rightarrow T_{-\Re \int_c^{v_2} \zeta_3} \mathbb{K}_{t \geq -\Re \int_{v_2}^z \zeta_3} \\ &\rightarrow T_{-\Re \int_c^{v_2} \zeta_3} (\mathbb{K}_{t \geq -\Re \int_{v_2}^z \zeta_3} \oplus \mathbb{K}_{t \geq -\Re \int_{v_2}^z \zeta_2}) \\ &\rightarrow T_{-\Re \int_c^{v_2} \zeta_3} \mathbb{K}_{t \geq -\Re \int_{v_2}^z \zeta_3} \oplus T_{-\Re \int_c^{v_2} \zeta_3} T_{-\Re \int_{v_2}^{v_1} \zeta_2} \mathbb{K}_{t \geq -\Re \int_{v_1}^z \zeta_2} \\ &\rightarrow T_{-\Re \int_c^{v_2} \zeta_3} \mathbb{K}_{t \geq -\Re \int_{v_2}^z \zeta_3} \oplus T_{-\Re \int_c^{v_2} \zeta_3} T_{-\Re \int_{v_2}^{v_1} \zeta_2} (\mathbb{K}_{t \geq -\Re \int_{v_1}^z \zeta_2} \oplus \mathbb{K}_{t \geq -\Re \int_{v_1}^z \zeta_1}) \\ &\rightarrow \mathbb{K}_{t \geq -\Re \int_c^z \zeta_3} \oplus T_{-\Re \int_c^{v_2} \zeta_3} T_{-\Re \int_{v_2}^{v_1} \zeta_2} T_{-\Re \int_{v_1}^c \zeta_1} \mathbb{K}_{t \geq -\Re \int_c^z \zeta_1}. \end{aligned} \quad (10.3)$$

Each arrow is the following: (1) and (3) These are just different expressions of the same objects. (2) and (4) Voros connection formula. (5) Projection. The projection to the

second component of the last line is nonzero, in particular, the composition is not id. This is the rephrasing of what Berk–Nevins–Roberts observed.

This implies the following. We can construct sheaf quantization like (8.3) on each four region. We can also glue up them by using the connection formula to obtain a sheaf quantization outside the collision. However, we cannot extend the sheaf quantization onto the collision because of the existence of the anomalous monodromy.

To resolve this “paradox”, BNR introduced additional Stokes curves emanating from the collision points (Figure 10.2).




Figure 10.2: New Stokes curves of BNR equation

In our language, it says the following: Add the Stokes curve of (31)-type with the gluing morphism which cancels the morphism $\mathbb{K}_{t \geq -\Re \int_c^z \zeta_3} \rightarrow T_{-\Re \int_c^{v_2} \zeta_3} T_{-\Re \int_{v_2}^{v_1} \zeta_2} T_{-\Re \int_{v_1}^c \zeta_1} \mathbb{K}_{t \geq -\Re \int_c^z \zeta_1}$. Note that

$$-\left(\Re \int_c^{v_2} \zeta_3 + \Re \int_{v_2}^{v_1} \zeta_2 + \Re \int_{v_1}^c \zeta_1 \right) = \Re \int_{v_1}^c (\zeta_2 - \zeta_1) + \Re \int_{v_2}^c (\zeta_3 - \zeta_2) =: w(c). \quad (10.4)$$

Then we can modify our sheaf quantization and the new one can extend over the collision point. We denote the resulting sheaf by S_L .

Without the new Stokes curve, we can still do the following. Instead of considering the local sheaf quantization S_{loc} like (8.3), we use the cone of the endomorphism $T^{w(c)}: S_{loc} \rightarrow S_{loc}$. Again, we can glue up them outside the collision. Moreover, with this replacement, the anomalous monodromy (10.3) vanishes. Hence we can extend the sheaf quantization over the collision. We denote this sheaf quantization S . There exists a canonical morphism $S_L \rightarrow S$, and we view it as an inverse system. We will use this point of view in the general construction.

Remark 10.5. The Stokes curves are expected to correspond to holomorphic disks whose boundaries lying on the spectral curve and the cotangent fibers. Then the integrations like $\Re \int (\zeta_i - \zeta_j)$ are the areas of the holomorphic disks.

10.3 Spectral scattering diagram

To package our construction, we would like to prepare some general notions and their consequences. The package is a variant of the notion of scattering diagram [KS06, GS11].

In the following (until the final section), our fixed data is the following. A finite subset $M \subset C$ (“pole divisor”) and a submanifold $L \subset T^*C$ (“spectral curve”) which is a branched covering of $C \setminus M$ and \mathfrak{h} -tame.

Definition 10.6. For a point $x \in C \setminus M$, an \mathfrak{h} -preStokes curve of type (i, j) emanating from x is a subset of

$$\left\{ z \in C \setminus M \mid \Im \int_x^z (\zeta_i - \zeta_j) dz = 0, \Re \int_x^z (\zeta_i - \zeta_j) dz > 0 \right\} \quad (10.5)$$

such that the subset is homeomorphic to a connected closed subset of \mathbb{R} , and one of the boundary is x .

Note that an \mathfrak{h} -preStokes curve is canonically oriented. The notion of ordered collisions and cyclically ordered collisions for preStokes curves can be defined in the same way as for Stokes curves.

Definition 10.7. Suppose that L is \mathfrak{h} -tame. Let B be a collection of \mathfrak{h} -preStokes curves containing the \mathfrak{h} -Stokes curves. The set $\text{Sing}(B)$ is the set of turning points and ordered intersection points of \mathfrak{h} -preStokes curves in B .

We say B is *tame* if the followings hold:

- (i) Any element of B will end at M .
- (ii) The set $C \setminus (M \cup \bigcup_{l \in B} l)$ is open and each connected component is contractible.
- (iii) Any ordered collisions are not cyclic.
- (iv) The set $\text{Sing}(B)$ is discrete in $C \setminus M$.

Definition 10.8. Let B be a tame collection of \mathfrak{h} -preStokes curves. The union of \mathfrak{h} -preStokes curves forms a graph and we denote the graph by G_B .

1. A connected component of $C \setminus (M \cup \bigcup_{l \in B} l)$ is called a *Stokes region* of B .
2. A *vertex* of B is a node of G_B .
3. For a Stokes region D , let \overline{D} be the closure of a lift of D in the universal covering of C . A vertex of D is a point of \overline{D} which is a lift of a vertex of B .
4. An *edge* of B is an edge of G_B .

Definition 10.9. Let B be a tame collection of \mathfrak{h} -preStokes curves. A *spectral scattering diagram* \mathfrak{S} is a set of pairs (l, ϕ) and a set $\{A_e\}$ where l is an \mathfrak{h} -preStokes curve in B , $\phi \in \Lambda_0$, and $A_e \in GL_n(\mathbb{K})$ is a diagonal matrix parameterized by edges of G_B .

Let \mathfrak{S} be a spectral scattering diagram. For $\mathbf{l} = (l, \phi) \in \mathfrak{S}$ with l of type (i, j) , we denote the starting point of l by $v(l)$. For a point $c \in l \cap \text{Sing}(B)$, the interval $\gamma_c := [v(l), c]$ is an oriented path. We set

$$M_{\mathbf{l}}(c) := \phi T^{\Re \int_{v(l)}^c (\zeta_i - \zeta_j) dz}. \quad (10.6)$$

For $c \in \text{Sing}(B)$, let l_1, \dots, l_n be the counterclockwise ordered collection of \mathfrak{h} -preStokes curves in B passing through c . Let (i_k, j_k) be the type of l_k . We denote the set of elements in \mathfrak{S} supported on l_k by $\{\mathbf{l}_{kl} := (l_k, \phi_{kl})\}_l$. Then l_k also gives an edge containing c . We also denote the corresponding edge by l_k (by abuse of the notation). The monodromy automorphism at c is defined by

$$M(c) := M(\mathfrak{S}, c) := \prod_k A_{l_k} \prod_l (\text{id} + M_{\mathbf{l}_{kl}}(c) E_{i_k j_k}) \in GL_n(\Lambda_0) \quad (10.7)$$

where the product with respect to k is ordered and E_{ij} is the matrix whose (ij) -entry is 1 and the other entries are 0. Note that if $l_i \subset l_j$ then the two \mathfrak{h} -preStokes curves should be of the same type or of separated types by Definition 10.7 (iii), hence the product is well-defined.

Lemma 10.10. *If $i \neq j$ and $M(c)_{ij}$ is nonzero, $M(c)_{ji} = 0$.*

Proof. This is a consequence of the non-existence of cyclically ordered collisions. \square

Definition 10.11. A spectral scattering diagram is *consistent modulo T^w* if $M(\mathfrak{S}, c) = \text{id}$ modulo T^w for any $c \in \text{Sing}(\mathfrak{S}) := \text{Sing}(B)$.

Let \mathfrak{S} be a consistent spectral scattering diagram consistent modulo T^w . We would like to associate the gluing data.

For each Stokes region, let \tilde{D} be a neighborhood of D in $C \setminus (\text{Sing}(B) \cup M)$ such that $\tilde{D} \cap G_B$ is the Stokes edges of D . For each Stokes region and a vertex v of D , we put

$$S_L^v|_D := \bigoplus_{c \in \mathbb{R}} T_c \bigoplus_i \mathbb{K}_{D_i^v}. \quad (10.8)$$

where

$$D_i^v := \left\{ (x, t) \in \tilde{D} \times \mathbb{R}_t \mid t \geq -\Re \int_v^z \zeta_i dz \right\} \quad (10.9)$$

like in the quadratic case. When we go across a Stokes edge, we associate the following gluing-up morphism:

(Change of regions) For adjacent Stokes regions D_1 and D_2 with a common vertex v , let $\mathbf{l}_1, \mathbf{l}_2, \dots, \mathbf{l}_n$ be the subset of \mathfrak{S} supported on the separating edge e . We will use the same notation as in (10.6).

An \mathfrak{h} -preStokes curve of type (i, j) satisfies $\Re \int_v^x (\zeta_i - \zeta_j) dz > 0$. Then we have

$$\text{End} \left(\mathbb{K}_{(D_e)_i^v} \oplus \mathbb{K}_{(D_e)_j^v} \right) = \mathbb{K} \cdot \text{id}_i \oplus \mathbb{K} \cdot \text{id}_j \oplus \mathbb{K} \cdot e_{ij} \quad (10.10)$$

where D_e is a thickening of e and

$$e_{ij} \in \text{Hom} \left(\mathbb{K}_{(D_e)_i^v}, \mathbb{K}_{(D_e)_j^v} \right) \text{ if } \Re \int_v^x (\zeta_i - \zeta_j) dz > 0 \quad (10.11)$$

is a canonical nontrivial basis coming from the inclusion of the closed sets. We set

$$\varphi_{D_1, D_2} := A_e \prod_{\mathbf{l}_k} (\text{id} + M_{\mathbf{l}_k}(c) e_{i_k j_k}) : S_L^v|_{D_1}|_{D_1 \cap D_2} \rightarrow S_L^v|_{D_2}|_{D_1 \cap D_2} \quad (10.12)$$

Proposition 10.12. *Let \mathfrak{S} be a consistent spectral scattering diagram modulo T^w . Then there exists an object with $\mu\text{supp} = L$ in $\text{Sh}_{\tau > 0}^{\mathbb{R}}(C \times \mathbb{R}_t)$.*

Proof. We proceed like in the quadratic case. We can glue up $S_L^v|_D$'s along the gluing morphisms defined above, which give a sheaf quantization outside $C \setminus \text{Sing}(B)$. Instead of using this naive one, we use the same gluing morphisms to glue up $\text{Cone}(T^w: S_L^v|_D \rightarrow S_L^v|_D)$. Then the resulting object does not have any nontrivial monodromy around any point of $\text{Sing}(B)$, because the diagram \mathfrak{S} is consistent modulo T^w . This completes the proof. \square

Remark 10.13. One can say that the notion of spectral scattering diagram is a 2d-version of usual scattering diagrams, and a usual scattering diagram is 4d. Regarding with the aspect of 2d-4d wall-crossing, ultimately, one should consider the notion of "2d-4d scattering diagram" where the consistency will be given by 2d-4d wall-crossing formulas.

10.4 An inductive step

Let \mathfrak{S} be a spectral scattering diagram which is consistent modulo T^w . We would like to construct a new spectral scattering diagram.

The consistency modulo T^w implies that $M(\mathfrak{S}, c) = \text{id}$ modulo T^w for any $c \in \text{Sing}(\mathfrak{S})$. We can write it as $\text{id} + \sum_{i \neq j} c_{ij} T^{a_{ij}} E_{ij}$ and $a_{ij} \geq w$. Here $i \neq j$ comes from the non-existence of cyclic-ordered collision. Let $l_{ij,c}$ be an \mathfrak{h} -preStokes curve of type (i, j) emanating from c if $c_{ij} \neq 0$ and consider the pair $\mathbf{l}_{ij,c} := (l_{ij,c}, -c_{ij} T^{a_{ij}})$. We set $\mathfrak{S}[c] := \{\mathbf{l}_{ij,c}\}_{i,j}$ and $\mathfrak{S}^+ := \mathfrak{S} \cup \bigcup_{c \in \text{Sing}(B)} \mathfrak{S}[c]$. We assume the set $B^+ := B \cup \bigcup_{c, i, j, c_{ij} \neq 0} \{l_{c,ij}\}$ is again a tame collection of \mathfrak{h} -preStokes curves. For A_e 's, if an edge e' of B^+ is a subset of an edge e of B , we set $A_{e'} := A_e$. Otherwise, $A_{e'} = \text{id}$. Then \mathfrak{S}^+ is a spectral scattering diagram.

The following is obvious from the construction.

Lemma 10.14. *$M(\mathfrak{S}^+, c)$ is consistent modulo T^{2w} .*

To speak about the consistency of \mathfrak{S}^+ , we prepare some notions. For a pair $\mathbf{l} := (l, \phi)$ of type (i, j) , its weight is defined by

$$w(\mathbf{l}) := \sup \{w \in \mathbb{R}_{>0} \mid \phi = 0 \text{ modulo } T^w\}, \quad (10.13)$$

namely, the valuation of the Novikov ring. For a point $y \in l$, the \mathbf{l} -weight of y is defined by

$$w_{\mathbf{l}}(y) := w(\mathbf{l}) + \Re \int_{v(l)}^y (\zeta_i - \zeta_j) dz. \quad (10.14)$$

Lemma 10.15. *Let $\mathbf{l}_1, \dots, \mathbf{l}_n$ be a subset of some spectral scattering diagram. For a point $c \in \bigcap l_i$, $M(c)$ is id modulo $T^{\min_{i \neq j} \{w_{\mathbf{l}_i}(c) + w_{\mathbf{l}_j}(c)\}}$. More precisely, $(M(c))_{ij}$, $(i \neq j)$ is zero modulo $T^{w_{ij}}$ where*

$$w_{ij} := \min \{w_{\mathbf{l}_k}(c) + w_{\mathbf{l}_l}(c) \mid k \neq l, \mathbf{l}_k \text{ is of type } (i, m) \text{ and } \mathbf{l}_l \text{ is of type } (m, j) \text{ for some } m\}. \quad (10.15)$$

Proof. This is clear from the definition of the monodromy. \square

An element of $\bigcup_c \mathfrak{S}[c]$ carries the weight greater than or equal to w . By Lemma 10.15, the new diagram \mathfrak{S}^+ is also consistent modulo T^w .

In the following, we would like to show that \mathfrak{S}^+ is consistent modulo $T^{w'}$ for some w' with $w' > w$ for some special \mathfrak{S} .

10.5 A local study of a turning point

We would like to prepare a local notion around a turning point. We assume that L is \mathfrak{h} -tame.

Let v be a turning point (i.e., a double branch). Without loss of generality, we consider the pair (ζ_1, ζ_2) is branched. Let T be the Stokes curve defined by $\Im(\int_v^x (\zeta_1 - \zeta_2)) = 0$. Take a compact neighborhood U_v of x such that

1. By 3 and 6 of Definition 10.4, we can take U_v sufficiently small so that U_v does not contain any other turning points and does not intersect with $(i, 1)$, $(2, i)$ -Stokes curves for any i .
2. $U_v \cap T$ is topologically a trivalent tree and $U_v \cap T$ has three boundaries z_1, z_2, z_3 .

We set

$$d(U_v) := \min_i \left\{ \Re \left(\int_v^{z_i} (\zeta_1 - \zeta_2) dz \right) \right\} > 0 \quad (10.16)$$

We also take a small neighborhood V_v of $\overline{U_v} \cap \overline{T}$ such that the closure of V_v does not contain any other turning points and does not intersect with \mathfrak{h} -Stokes curves of type $(i, 1)$ or $(2, i)$. Consider the set L_v of connected subsets of \mathfrak{h} -preStokes curves of type $(i, 1)$ or $(2, i)$ such that each $l \in L_v$ starts from a point outside V_v and ends at a point in $U_v \cap T$. We set

$$d(V_v) := \inf_{l \in L_v} \left\{ \begin{cases} \Re \left(\int_l (\zeta_i - \zeta_1) dz \right) & \text{if } l \text{ is of type } (i, 1) \\ \Re \left(\int_l (\zeta_2 - \zeta_i) dz \right) & \text{if } l \text{ is of type } (2, i) \end{cases} \right\} > 0. \quad (10.17)$$

Let w be a turning point or an ordered collision of \mathfrak{h} -Stokes curves which is different from v . Let $L_{vw, ij}$ be the set of \mathfrak{h} -Stokes curves of type (ij) emanating from w passing through V_v . We set

$$d_v := \inf_{w \neq v} \inf_{i, j} \inf_{l \in L_{vw, ij}} \inf_{x \in V_v} \left\{ \Re \left(\int_w^x (\zeta_i - \zeta_j) dz \right) \text{ if } l \text{ is of type } (ij) \text{ and the integral is along } l \right\}. \quad (10.18)$$

This is positive, since the set of turning points and the ordered collisions is discrete and w is outside the closure of V_v .

We choose U_v and V_v for each turning point and set

$$w_{\min} := \min_v \{d(U_v), d(V_v), d_v/2\}, \quad (10.19)$$

which is again a positive real number.

10.6 General construction

We would like to construct a sequence of spectral scattering diagrams.

Definition 10.16. An initial spectral scattering diagram \mathfrak{S}_0 of L is the following:

1. $B =: B_0$ is the set of \mathfrak{h} -Stokes curves emanating from the ordinary turning points.
2. For an \mathfrak{h} -Stokes curve l of type (i, j) , we consider the pair (l, ϕ) for some ϕ and A which satisfies the following: For any ordinary turning point c , the equality

$$1 = \prod_k A_k \prod_l (\text{id} + M_{\mathbf{l}_{kl}}(c) E_{i_k j_k}) \in GL_n(\Lambda_0) \quad (10.20)$$

holds where the product runs over the rays emanating from the ordinary turning point.

Fix an initial spectral scattering diagram \mathfrak{S}_0 . We set

$$\mathfrak{S}_n := \mathfrak{S}_0^{+n}, B_n := B_0^{+n}. \quad (10.21)$$

We say an element of $\mathfrak{S}'_{n-1} = \mathfrak{S}_n \setminus \mathfrak{S}_{n-1}$ an n -th Stokes curve.

Definition 10.17. We say \mathfrak{S}_0 is *inductively \mathfrak{h} -tame*, if L is \mathfrak{h} -tame and B_n is tame (Definition 10.7) for any $n \geq 0$.

We expect the inductive \mathfrak{h} -tameness is satisfied by a generic \mathfrak{h} -connection and a generic \mathfrak{h} after imposing some conditions on the order of zeros and poles as in the quadratic case.

Proposition 10.18. *Suppose that \mathfrak{S}_0 is inductively \mathfrak{h} -tame.*

1. For $n \geq 0$, the weight of an n -th Stokes curve is greater than or equal to nw_{min} .
2. The diagram \mathfrak{S}_n is consistent modulo $T^{(n+1)w_{min}}$.

Proof. We will prove by induction. By the definition of w_{min} and Lemma 10.15, the first Stokes curves have weights at least w_{min} . The inductive assumption is as follows: Any m -th Stokes curve ($m \leq n$) has weight at least mw_{min} and the diagram \mathfrak{S}_n is consistent modulo $T^{(n+1)w_{min}}$.

Let \mathbf{l} be an $(n+1)$ -st Stokes curve. By the consistency of \mathfrak{S}_n modulo $T^{(n+1)w_{min}}$, the weight of \mathbf{l} is greater than or equal to $(n+1)w_{min}$. This proves the first part.

We would like to prove the consistency of \mathfrak{S}_{n+1} modulo $T^{(n+2)w_{min}}$. Let c be an ordered collision of \mathbf{l} with another Stokes curve. If a collision consists of m -th Stokes curves with $m \geq 1$, Lemma 10.15 implies that the monodromy is id modulo $T^{(n+1+m)w_{min}}$ and $n+1+m \geq 2$.

Hence we will consider the case when an 0-th Stokes curve \mathbf{l}_0 also has an ordered collision with \mathbf{l} on c . Let v be the starting point of the 0-th Stokes curve \mathbf{l}_0 and we will use the notation from 9.3. Hence \mathbf{l}_0 is of type $(1, 2)$.

If the collision c is not in U_v , the weight $w_{\mathbf{l}_0}(c)$ is greater than or equal to $d(U_v) \geq w_{min}$. Hence the monodromy around the collision is id modulo $(n+2)w_{min}$ by Lemma 10.15.

Hence we can assume that c is in U_v . Suppose \mathbf{l} emanates from a point in $c' \notin V_v$. By Lemma 10.15 and the definition of w_{min} , we have $w_{\mathbf{l}}(c) \geq (n+1)w_{min} + d(V_v) \geq (n+2)w_{min}$. By Lemma 10.15, this case is consistent.

Hence we can assume \mathbf{l} emanates from a point in $c' \in V_v$.

Claim 10.19. *Each element \mathbf{l}' of \mathfrak{S}_n possibly passing through c' satisfies $w_{\mathbf{l}'}(c') \geq 2w_{min}$ except for \mathbf{l}_0 .*

Proof of the claim. For m -th Stokes curves with $m \geq 2$, it is obvious. Any 0-th/first Stokes curve emanates from outside V_v . Hence $w_{\mathbf{l}'}(v) = d_v \geq 2w_{min}$. This completes the proof. \square

By Claim 10.19, any ordered collision of an n -th Stokes curve with an m -th Stokes curve on c' carries the weight $\geq (n+2)w_{min}$. Hence, for the production process of \mathbf{l} , we only have to consider an ordered collision with an n -th Stokes curve and \mathbf{l}_0 . But, this collision only produces the Stokes curve \mathbf{l} of type $(l, 2)$ or $(1, l)$ for some l , which cannot have an ordered collision with \mathbf{l}_0 . \square

Corollary 10.20. *Suppose \mathfrak{S}_0 is inductively \hbar -tame. Then there exists a sheaf quantization of L at \hbar .*

Proof. By Proposition 10.18 and Proposition 10.12, we have a sheaf quantization of L as $S_n \in \text{Sh}_{\tau > 0}^{\mathbb{R}}(C \times \mathbb{R}_t)$ for each n and morphisms $S_{n+1} \rightarrow S_n$. Taking the limit, we obtain the desired object. \square

10.7 Initial data

To conclude the discussion, we would like to present conjectural spectral scattering diagrams associated to meromorphic flat \hbar -connections. We will consider a meromorphic flat \hbar -connection of rank N .

Formal data

We set $\mathbb{K} = \mathbb{C}((\hbar))$. Also, in the higher order cases, WKB solutions have a similar expression (see e.g. [HKT15]):

$$\Psi_i(z, \hbar) := \exp\left(\int^z \frac{P_{-1}^i}{\hbar}\right) \exp\left(\int^z \sum_{j=0} P_j^i \hbar^j\right) \quad (10.22)$$

where i is a label of a sheet of the spectral covering and P_j^i is uniquely determined from P_{-1}^i . Since each component of the complement of the initial spectral scattering diagram is contractible, we can choose one WKB solution for each sheet. Then, on each edge e , we can compare the specified WKB solutions for each sheet, and we obtain the scaling in $\mathbb{C}((\hbar))$ for each sheet. This is our diagonal matrix A_e .

We also have to choose an initial diagram. There exists a (conjecturally) canonical choice for this. For an edge e , we consider the starting turning point $v(e)$. With respect to the turning point normalization, we would like to take ϕ to be -1 . Hence the actual value of ϕ is the ratio between the turning point normalization and our chosen WKB solutions. This choice comes from an ansatz that we can use the Voros connection formula around a simple turning point.

Analytic data

We set $\mathbb{K} := \mathbb{C}$. Conjecturally, one can lift the above initial diagram defined over $\mathbb{C}((\hbar))$ to an analytic data, defined over \mathbb{C} . We would like to state a summability conjecture. Philosophically speaking, the data are ratios of solutions normalized at different points. However, for a point in a Stokes region of \mathfrak{S}_n , we cannot expect the Borel summability of the WKB solutions, since there might exist m -th Stokes curves with $m > n$ passing through the point. Hence we will formulate it as follows. The statements in the following paragraphs are all conjectural.

Again, fix a set of formal WKB solutions on each Stokes region of the initial diagram. On each edge e , the formal WKB solutions on adjacent Stokes regions are Borel-summable using the Laplace transform along any small deformations of $\mathbb{R}_{>0}$ avoiding the singularities of the Borel transformations. Take one of such small deformation of the Laplace path. Consider the resummation of WKB solutions of adjacent Stokes regions along the path. Taking the ratio, we define A_e . This A_e does not depend on the choice of small deformations of $\mathbb{R}_{>0}$.

Next, we would like to define ϕ 's. We assume L is inductively \hbar -tame. Then the initial diagram can be defined in the same way as the formal version. Applying the result in the above section to this initial diagram, we obtain a spectral scattering diagram. The resulting spectral scattering diagram coincide with the following.

For an edge e of \mathfrak{S} , in adjacent regions D_1, D_2 , the resummed WKB solutions $\{\Psi_{1i}\}, \{\Psi_{2i}\}$ should be related by

$$\Psi_1 = A_e \Psi_2 + e^{c_1/\hbar} A_1 \Psi_2 + e^{c_2/\hbar} A_1 \Psi_2 + \dots \quad (10.23)$$

Here $A_i \in M_N(\mathbb{C})$, $c_i/\hbar \in \mathbb{R}_{<0}$. The set $\{(l_i, \phi_i)\}$ supported on e should be related to the above expression by $w(l_i) = c_i$ and ϕ_i are matrix elements of A_i . Namely, the n -th diagram \mathfrak{S}_n only reflects the Stokes–Voros connection automorphisms corresponding to the Borel singularities with $|c_i| < (n+1)w_{min}$.

Remark 10.21. These statements may be stated more elegantly by using the language of Ecalle's resurgence theory.

References

- [AKT91] Takashi Aoki, Takahiro Kawai, and Yoshitsugu Takei. The Bender-Wu analysis and the Voros theory. In *Special functions (Okayama, 1990)*, ICM-90 Satell. Conf. Proc., pages 1–29. Springer, Tokyo, 1991.

- [BL94] Joseph Bernstein and Valery Lunts. *Equivariant sheaves and functors*, volume 1578 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1994.
- [BNR82] H. L. Berk, William McCay Nevins, and K. V. Roberts. New Stokes' line in WKB theory. *J. Math. Phys.*, 23(6):988–1002, 1982.
- [BS15] Tom Bridgeland and Ivan Smith. Quadratic differentials as stability conditions. *Publ. Math. Inst. Hautes Études Sci.*, 121:155–278, 2015.
- [DDP93] É. Delabaere, H. Dillinger, and F. Pham. Résurgence de Voros et périodes des courbes hyperelliptiques. *Ann. Inst. Fourier (Grenoble)*, 43(1):163–199, 1993.
- [DFK⁺] Olivia Dumitrescu, Laura Fredrickson, Georgios Kydonakis, Rafe Mazzeo, Motohico Mulase, and Andrew Neitzke. Opers versus nonabelian hodge. *eprint arXiv:1607.02172*.
- [DK16] Andrea D'Agnolo and Masaki Kashiwara. Riemann-Hilbert correspondence for holonomic D-modules. *Publ. Math. Inst. Hautes Études Sci.*, 123:69–197, 2016.
- [DLS93] T. M. Dunster, D. A. Lutz, and R. Schäfke. Convergent Liouville-Green expansions for second-order linear differential equations, with an application to Bessel functions. *Proc. Roy. Soc. London Ser. A*, 440(1908):37–54, 1993.
- [Gai] Davide Gaiotto. Opers and tba. *eprint arXiv:1403.6137*.
- [GMN13] Davide Gaiotto, Gregory W. Moore, and Andrew Neitzke. Spectral networks. *Ann. Henri Poincaré*, 14(7):1643–1731, 2013.
- [Gro57] Alexander Grothendieck. Sur quelques points d'algèbre homologique. *Tohoku Math. J. (2)*, 9:119–221, 1957.
- [GS11] Mark Gross and Bernd Siebert. From real affine geometry to complex geometry. *Ann. of Math. (2)*, 174(3):1301–1428, 2011.
- [GT13] Alexander Getmanenko and Dmitry Tamarkin. Microlocal properties of sheaves and complex WKB. *Astérisque*, (356):x+111, 2013.
- [Gui16] Stéphane Guillermou. Quantization of exact lagrangian submanifolds in a cotangent bundle, lectures at the 2016 summer school “symplectic topology, sheaves, and mirror symmetry. available at the author's webpage, 2016.
- [HKT15] Naofumi Honda, Takahiro Kawai, and Yoshitsugu Takei. *Virtual turning points*, volume 4 of *SpringerBriefs in Mathematical Physics*. Springer, Tokyo, 2015.
- [IK] Yuichi Ike and Tatsuki Kuwagaki. Microlocal categories over the novikov ring i: cotangent bundles.
- [IN14] Kohei Iwaki and Tomoki Nakanishi. Exact WKB analysis and cluster algebras. *J. Phys. A*, 47(47):474009, 98, 2014.
- [Jin] Xin Jin. Microlocal sheaf categories and the j-homomorphism. *arXiv:2004.14270*.
- [JT] Xin Jin and David Treumann. Brane structures in microlocal sheaf theory. *eprint arXiv:1704.04291*.
- [Kon] Maxim Kontsevich. Resurgence and quantization. *YouTube, IHÉS*.
- [KS] Maxim Kontsevich and Yan Soibelman. Holomorphic floer theory.
- [KS94] Masaki Kashiwara and Pierre Schapira. *Sheaves on manifolds*, volume 292 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1994. With a chapter in French by Christian Houzel, Corrected reprint of the 1990 original.
- [KS06] Maxim Kontsevich and Yan Soibelman. Affine structures and non-Archimedean analytic spaces. In *The unity of mathematics*, volume 244 of *Progr. Math.*, pages 321–385. Birkhäuser Boston, Boston, MA, 2006.

- [KS16] Masaki Kashiwara and Pierre Schapira. *Regular and irregular holonomic D-modules*, volume 433 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2016.
- [KT05] Takahiro Kawai and Yoshitsugu Takei. *Algebraic analysis of singular perturbation theory*, volume 227 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 2005. Translated from the 1998 Japanese original by Goro Kato, Iwanami Series in Modern Mathematics.
- [Kuw22] Tatsuki Kuwagaki. *\hbar -Riemann-Hilbert correspondence*, 2022.
- [NZ09] David Nadler and Eric Zaslow. Constructible sheaves and the Fukaya category. *J. Amer. Math. Soc.*, 22(1):233–286, 2009.
- [PS04] Pietro Polesello and Pierre Schapira. Stacks of quantization-deformation modules on complex symplectic manifolds. *Int. Math. Res. Not.*, (49):2637–2664, 2004.
- [Sei00] Paul Seidel. Graded Lagrangian submanifolds. *Bull. Soc. Math. France*, 128(1):103–149, 2000.
- [Str84] Kurt Strebel. *Quadratic differentials*, volume 5 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1984.
- [STWZ19] Vivek Shende, David Treumann, Harold Williams, and Eric Zaslow. Cluster varieties from Legendrian knots. *Duke Math. J.*, 168(15):2801–2871, 2019.
- [Tak17] Yoshitsugu Takei. WKB analysis and Stokes geometry of differential equations. In *Analytic, algebraic and geometric aspects of differential equations*, Trends Math., pages 263–304. Birkhäuser/Springer, Cham, 2017.
- [Tam18] Dmitry Tamarkin. Microlocal condition for non-displaceability. In *Algebraic and analytic microlocal analysis*, volume 269 of *Springer Proc. Math. Stat.*, pages 99–223. Springer, Cham, 2018.
- [Vit] Claude Viterbo. An introduction to symplectic topology through sheaf theory. <http://www.math.polytechnique.fr/cmat/viterbo/Eilenberg/Eilenberg.pdf>.
- [Vor83] A. Voros. The return of the quartic oscillator: the complex WKB method. *Ann. Inst. H. Poincaré Sect. A (N.S.)*, 39(3):211–338, 1983.

Department of Mathematics, Graduate School of Science, Kyoto University, tatsuki.kuwagaki.a@gmail.com