

# NONCOERCIVE QUASILINEAR ELLIPTIC OPERATORS WITH SINGULAR LOWER ORDER TERMS

FERNANDO FARRONI, LUIGI GRECO,  
GIOCONDA MOSCARIELLO AND GABRIELLA ZECCA

**ABSTRACT.** We consider a family of quasilinear second order elliptic differential operators which are not coercive and are defined by functions in Marcinkiewicz spaces. We prove the existence of a solution to the corresponding Dirichlet problem. The associated obstacle problem is also solved. Finally, we show higher integrability of a solution to the Dirichlet problem when the datum is more regular.

## 1. INTRODUCTION

Given a bounded domain  $\Omega$  of  $\mathbb{R}^N$ ,  $N \geq 2$ , we consider

$$A: (x, u, \xi) \in \Omega \times \mathbb{R} \times \mathbb{R}^N \mapsto \mathbb{R}^N$$

a Carathéodory vector field (i.e. measurable in  $x \in \Omega$  and continuous in  $(u, \xi) \in \mathbb{R} \times \mathbb{R}^N$ ) satisfying for a.e.  $x \in \Omega$  and for every  $u \in \mathbb{R}$  and  $\xi, \eta \in \mathbb{R}^N$  the following structural conditions:

$$(1.1) \quad \langle A(x, u, \xi), \xi \rangle \geq \alpha |\xi|^p - (b(x) |u|)^p - \varphi(x)^p$$

$$(1.2) \quad |A(x, u, \xi)| \leq \beta |\xi|^{p-1} + (b(x) |u|)^{p-1} + \varphi(x)^{p-1}$$

$$(1.3) \quad \langle A(x, u, \xi) - A(x, u, \eta), \xi - \eta \rangle > 0 \quad \text{for } \xi \neq \eta$$

where  $0 < \alpha < \beta$  are positive constants,  $1 < p < N$ , and  $b$  and  $\varphi$  are positive functions verifying  $b \in L^{N,\infty}(\Omega)$  and  $\varphi \in L^p(\Omega)$ . In view of

---

*2020 Mathematics Subject Classification.* 35J25, 35J87.

*Key words and phrases.* Dirichlet problems, Noncoercive elliptic operators, Obstacle problems.

The authors are members of Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of INdAM. The research of G.M. has been partially supported by the National Research Project PRIN "Gradient flows, Optimal Transport and Metric Measure Structures", code 2017TEXA3H.

Sobolev embedding theorem in Lorentz spaces [30, 2, 20], by (1.2) and the assumptions on  $b$  and  $\varphi$ , for each  $u \in W_0^{1,p}(\Omega)$  we have

$$A(x, u, \nabla u) \in L^{p'}(\Omega, \mathbb{R}^N)$$

Hence, we can define the quasilinear elliptic distributional operator

$$(1.4) \quad -\operatorname{div} A(x, u, \nabla u)$$

setting for any  $w \in W_0^{1,p}(\Omega)$

$$(1.5) \quad \langle -\operatorname{div} A(x, u, \nabla u), w \rangle = \int_{\Omega} \langle A(x, u, \nabla u), \nabla w \rangle \, dx.$$

Given  $\Phi \in W^{-1,p'}(\Omega)$ , we study the Dirichlet problem

$$(1.6) \quad \begin{cases} -\operatorname{div} A(x, u, \nabla u) = \Phi \\ u \in W_0^{1,p}(\Omega) \end{cases}$$

By a solution to Problem (1.6) we mean a function  $u \in W_0^{1,p}(\Omega)$  such that

$$(1.7) \quad \int_{\Omega} A(x, u, \nabla u) \nabla w \, dx = \langle \Phi, w \rangle, \quad \forall w \in C_0^\infty(\Omega),$$

where  $\langle \cdot, \cdot \rangle$  denotes the duality product of  $W^{-1,p'}(\Omega)$  and  $W_0^{1,p}(\Omega)$ . Clearly, (1.7) extends to all  $w \in W_0^{1,p}(\Omega)$ .

Our structural conditions allow for a singular lower order term depending on  $u$ . As an example, we consider the following operator

$$(1.8) \quad A(x, u, \xi) := \langle \mathcal{H}(x)\xi, \xi \rangle^{\frac{p-2}{2}} \mathcal{H}(x)\xi + B(x)|u|^{p-2}u$$

with  $1 < p < N$ . Here  $\mathcal{H} = \mathcal{H}(x): \Omega \rightarrow \mathbb{R}^{N \times N}$  is a symmetric, bounded matrix field of  $\mathbb{R}^{N \times N}$  such that

$$\langle \mathcal{H}(x)\xi, \xi \rangle \geq \alpha|\xi|^2 \quad \text{for a.e. } x \in \Omega \text{ and for all } \xi \in \mathbb{R}^N,$$

for a given  $\alpha > 0$ . The vector field  $B: \Omega \rightarrow \mathbb{R}^N$  is a measurable function satisfying  $|B(x)| \leq (b(x))^{p-1}$  for a.e.  $x \in \Omega$  and for some  $b \in L^{N,\infty}(\Omega)$ .

The feature of Problem (1.6) is the lack of coercivity for the operator (1.4) and the singularity in the lower order term due to property of  $b$ . It is well known that, if the operator in (1.4)-(1.5) is coercive, then a solution to problem (1.6) exists. It can for instance be shown by monotone operator theory [27, 6, 7, 4].

On the other hand, the existence of a bounded solution can be expected when  $\Phi$  and  $b$  are sufficiently smooth. For example, in the model case and even for the corresponding linear case, a solution to Problem

(1.6) is bounded whenever  $\Phi$  and  $b$  are in  $W^{-1,p'}(\Omega)$  and  $L^{p'}(\Omega, \mathbb{R}^N)$ , respectively, with  $p' > N/(p-1)$  (see [32, 17]).

The space  $L^{N,\infty}(\Omega)$  is slightly larger than  $L^N(\Omega)$ . Nevertheless, there are essential differences between the case  $b \in L^N(\Omega)$  ([5, 10]), or even  $b \in L^{N,q}(\Omega)$  ([29]) with  $N \leq q < \infty$ , and the case  $b \in L^{N,\infty}(\Omega)$ . In  $L^{N,\infty}(\Omega)$  bounded functions are not dense. Furthermore, in  $L^{N,\infty}(\Omega)$  the norm is not absolutely continuous, namely a function can have large norm even if restricted to a set with small measure.

Our first result is the following

**Theorem 1.1.** *Let  $\Phi \in W^{-1,p'}(\Omega)$ . Under the assumptions (1.1), (1.2) and (1.3), if*

$$(1.9) \quad \text{dist}_{L^{N,\infty}}(b, L^\infty(\Omega)) < \frac{\alpha^{\frac{1}{p}}}{S_{N,p}}$$

*then Problem (1.6) admits a solution.*

Here  $S_{N,p}$  denotes the Sobolev constant of Theorem 2.1 below. As an illustration, we state the following immediate consequence. We denote by  $L_0^{N,\infty}(\Omega)$  the closure of  $L^\infty(\Omega)$  in  $L^{N,\infty}(\Omega)$ .

**Corollary 1.1.** *Assume (1.1), (1.2) and (1.3), with  $b \in L_0^{N,\infty}(\Omega)$ . Then Problem (1.6) admits a solution, for every  $\Phi \in W^{-1,p'}(\Omega)$ .*

The closure  $L_0^{N,\infty}(\Omega)$  contains for example all Lorentz spaces  $L^{N,q}(\Omega)$ , for  $1 < q < +\infty$ , see Subsection 2.1.

It is not clear if the bound in (1.9) is sharp. However, existence of a solution could fail if no restriction on the distance is imposed, even in the linear case, as the Example 1 in Subsection 2.3 shows (see also [19]). Notice that condition (1.9) does not imply the smallness of the norm of  $b$  in  $L^{N,\infty}(\Omega)$  (see [19]).

In the case  $p = 2$  existence results analogous to that of Theorem 1.1 have been proved in [12, 19, 33] and in [9, 31, 34] when the principal part has a coefficient bound in BMO (i.e. the space of functions of bounded mean oscillation). We explicitly remark that in this context the operator (1.4) has the same integrability properties of the principal part (see also [23]). The evolutionary counterpart of Problem (1.6) has been studied in [11]. Other related results can be found in [1, 24, 28].

An additional difficulty in proving Theorem 1.1 lies in the lack of compactness that the operator

$$(1.10) \quad u \in W_0^{1,p}(\Omega) \quad \mapsto \quad (b(x)|u|)^{p-1} \in L^{p'}(\Omega)$$

exhibits in the case  $b \in L^{N,\infty}(\Omega)$ , in contrast with the case  $b \in L^N(\Omega)$  (see Example 3 in Subsection 2.3). In order to overcome this issue, first we consider the case in which  $b \in L^\infty(\Omega)$ . Under this assumption, we deduce the existence of a solution to Problem (1.6) by means of Leray–Schauder fixed point theorem. The a priori estimate required follows from a lemma that could be interesting in itself (see Lemma 3.1 below).

In order to reduce the general case  $b \in L^{N,\infty}(\Omega)$  to the previous one, we consider a sequence of approximating problems, defined essentially by truncating the vector field  $A = A(x, u, \xi)$  in the  $u$ -variable. A bound on the sequence of the solutions is achieved due to the assumption (1.9).

We emphasize that our result is also new when  $b \in L^N(\Omega)$ , in the sense that our approach allows us to treat the general nonlinear operator in (1.6).

Finally, by testing the problems with a suitable admissible test functions, we show that the sequence of solutions to the approximating problems is compact and its limit is a solution to the original problem (1.6).

In Section 5, we show that our approach is robust enough to handle also the corresponding obstacle problem. We prove an existence result in the same spirit of [18] (where the case  $p = 2$  is taken into account).

In Section 6 we present a regularity result. When  $b \in L^N(\Omega)$ , the study of the higher integrability of a solution to (1.6) has been developed in [13, 14] by using the theory of quasiminima. Local summability properties have been recently considered in [8, 22] in the linear case. Here, following [17], we prove higher summability of a solution  $u$  to (1.6) according to that of the data  $\Phi$  and  $\varphi$ .

## 2. PRELIMINARIES AND EXAMPLES

**2.1. Notation and function spaces.** Let  $\Omega$  be a bounded domain in  $\mathbb{R}^N$ . Given  $1 < p < \infty$  and  $1 \leq q < \infty$ , the Lorentz space  $L^{p,q}(\Omega)$  consists of all measurable functions  $f$  defined on  $\Omega$  for which the quantity

$$(2.1) \quad \|f\|_{p,q}^q = p \int_0^{+\infty} |\Omega_t|^{\frac{q}{p}} t^{q-1} dt$$

is finite, where  $\Omega_t = \{x \in \Omega : |f(x)| > t\}$  and  $|\Omega_t|$  is the Lebesgue measure of  $\Omega_t$ , that is,  $\lambda_f(t) = |\Omega_t|$  is the distribution function of  $f$ . Note that  $\|\cdot\|_{p,q}$  is equivalent to a norm and  $L^{p,q}$  becomes a Banach space when endowed with it (see [30, 3, 16]). For  $p = q$ , the Lorentz space

$L^{p,p}(\Omega)$  reduces to the Lebesgue space  $L^p(\Omega)$ . For  $q = \infty$ , the class  $L^{p,\infty}(\Omega)$  consists of all measurable functions  $f$  defined on  $\Omega$  such that

$$\|f\|_{p,\infty}^p = \sup_{t>0} t^p |\Omega_t| < +\infty$$

and it coincides with the Marcinkiewicz class, weak- $L^p(\Omega)$ .

For Lorentz spaces the following inclusions hold

$$L^r(\Omega) \subset L^{p,q}(\Omega) \subset L^{p,r}(\Omega) \subset L^{p,\infty}(\Omega) \subset L^q(\Omega),$$

whenever  $1 \leq q < p < r \leq \infty$ . Moreover, for  $1 < p < \infty$ ,  $1 \leq q \leq \infty$  and  $\frac{1}{p} + \frac{1}{p'} = 1$ ,  $\frac{1}{q} + \frac{1}{q'} = 1$ , if  $f \in L^{p,q}(\Omega)$ ,  $g \in L^{p',q'}(\Omega)$  we have the Hölder-type inequality

$$\int_{\Omega} |f(x)g(x)| \, dx \leq \|f\|_{p,q} \|g\|_{p',q'}.$$

As it is well known,  $L^\infty(\Omega)$  is not dense in  $L^{p,\infty}(\Omega)$ . For a function  $f \in L^{p,\infty}(\Omega)$  we define

$$(2.2) \quad \text{dist}_{L^{p,\infty}(\Omega)}(f, L^\infty(\Omega)) = \inf_{g \in L^\infty(\Omega)} \|f - g\|_{L^{p,\infty}(\Omega)}.$$

In order to characterize the distance in (2.2), we introduce for all  $k > 0$  the truncation operator

$$T_k(y) := \frac{y}{|y|} \min\{|y|, k\}.$$

It is easy to verify that

$$(2.3) \quad \lim_{k \rightarrow \infty} \|f - T_k f\|_{p,\infty} = \text{dist}_{L^{p,\infty}(\Omega)}(f, L^\infty(\Omega)).$$

We denote by  $L_0^{p,\infty}(\Omega)$  the closure of  $L^\infty(\Omega)$ . We have (see [21, Lemma 2.3])

$$(2.4) \quad f \in L_0^{p,\infty}(\Omega) \iff \lim_{t \rightarrow +\infty} t [\lambda_f(t)]^{1/p} = 0.$$

Clearly, for  $1 \leq q < \infty$  we have  $L^{p,q}(\Omega) \subset L_0^{p,\infty}(\Omega)$ , that is, any function in  $L^{p,q}(\Omega)$  has vanishing distance zero to  $L^\infty(\Omega)$ . Indeed,  $L^\infty(\Omega)$  is dense in  $L^{p,q}(\Omega)$ , the latter being continuously embedded into  $L^{p,\infty}(\Omega)$ . Actually, the inclusion also follows from (2.4), since  $\lambda_f(t) = |\Omega_t|$  is decreasing and hence the convergence of the integral at (2.1) implies the condition on the right of (2.4).

Assuming the origin  $0 \in \Omega$ , a typical element of  $L^{N,\infty}(\Omega)$  is  $b(x) = B/|x|$ , with  $B$  a positive constant. An elementary calculation shows that

$$(2.5) \quad \text{dist}_{L^{N,\infty}(\Omega)}(b, L^\infty(\Omega)) = B \omega_N^{1/N}$$

where  $\omega_N$  stands for the Lebesgue measure of the unit ball of  $\mathbb{R}^N$ .

The Sobolev embedding theorem in Lorentz spaces reads as

**Theorem 2.1** ([30, 2, 20]). *Let us assume that  $1 < p < N$ ,  $1 \leq q \leq p$ , then every function  $g \in W_0^{1,1}(\Omega)$  verifying  $|\nabla g| \in L^{p,q}(\Omega)$  actually belongs to  $L^{p^*,q}(\Omega)$ , where  $p^* = \frac{Np}{N-p}$  and*

$$\|g\|_{p^*,q} \leq S_{N,p} \|\nabla g\|_{p,q}$$

where  $S_{N,p}$  is the Sobolev constant.

**2.2. A version of the Leray–Schauder fixed point theorem.** We shall use the well known Leray–Schauder fixed point theorem in the following form (see [15, Theorem 11.3 pg. 280]). A continuous mapping between two Banach spaces is called compact if the images of bounded sets are precompact.

**Theorem 2.2.** *Let  $\mathcal{F}$  be a compact mapping of a Banach space  $X$  into itself, and suppose there exists a constant  $M$  such that  $\|x\|_X < M$  for all  $x \in X$  and  $t \in [0, 1]$  satisfying  $x = t\mathcal{F}(x)$ . Then,  $\mathcal{F}$  has a fixed point.*

**2.3. Critical examples.** Our first example shows that the only assumption that  $b \in L^{N,\infty}(\Omega)$  does not guarantee the existence of a solution to Problem (1.6).

**Example 1.** Let  $\Omega$  be the unit ball. For  $\frac{N}{2} < \gamma + 1 \leq N$ , the problem

$$(2.6) \quad \begin{cases} -\Delta u - \operatorname{div} \left( \gamma u \frac{x}{|x|^2} \right) = -\operatorname{div} \left( \frac{x}{|x|^{N-\gamma}} \right) & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases}$$

does not admit a solution. Assume to the contrary that  $u$  is a solution of (2.6). In the right hand side of the equation we recognize that

$$\frac{x}{|x|^{N-\gamma}} = \nabla v(x),$$

where  $v \in W_0^{1,2}(\Omega)$  is given by

$$v(x) = \begin{cases} \frac{1}{2-N+\gamma} (|x|^{2-N+\gamma} - 1) & \text{for } \gamma \neq N-2 \\ \log |x| & \text{for } \gamma = N-2 \end{cases}$$

Moreover,  $v$  solves the adjoint problem

$$\begin{cases} -\Delta v + \gamma \frac{x}{|x|^2} \cdot \nabla v = 0 & \text{in } \Omega \\ v = 0 & \text{on } \partial\Omega \end{cases}$$

Testing the equation in (2.6) by  $v$  we have

$$\int_{\Omega} |\nabla v|^2 dx = 0.$$

which readily implies  $v \equiv 0$  in  $\Omega$ , which is clearly not the case.  $\square$

Next example shows that for the complete operator

$$\operatorname{div} A(x, u, \nabla u) + B(x, u, \nabla u)$$

in general we do not have existence, even in the linear case.

**Example 2.** Let  $\lambda$  be an eigenvalue of Laplace operator and  $w$  a corresponding eigenfunction

$$\begin{cases} -\Delta w = \lambda w \\ w \in W_0^{1,2}(\Omega) \setminus \{0\} \end{cases}$$

Then the equation

$$-\Delta u - \lambda u = w$$

has no solution of class  $W_0^{1,2}(\Omega)$ .

Our final example shows that compactness of the operator (1.10) in the Introduction could fail.

**Example 3.** Assume  $N \geq 2$  and  $1 < p < N$ . Let  $\Omega$  be the ball of  $\mathbb{R}^N$  centered at the origin of radius 3. Our aim is to construct a sequence of functions  $\{u_n\}_{n \in \mathbb{N}}$  in  $W_0^{1,p}(\Omega)$  and a function  $b \in L^{N,\infty}(\Omega)$  such that  $\{\nabla u_n\}_{n \in \mathbb{N}}$  is bounded in  $L^p(\Omega, \mathbb{R}^N)$ , however it is not possible to extract from  $\{(b|u_n|)^{p-1}\}_{n \in \mathbb{N}}$  any subsequence strongly converging  $L^{p'}(\Omega)$ . To this aim, let

$$b(x) := \frac{1}{|x|}$$

and

$$\gamma := 1 - \frac{N}{p}.$$

We define a sequence  $\{u_n\}_{n \in \mathbb{N}}$  setting for  $x \in \Omega$

$$(2.7) \quad \begin{aligned} u_1(x) &:= \begin{cases} 1 - 2^\gamma & \text{if } |x| < 1 \\ |x|^\gamma - 2^\gamma & \text{if } 1 \leq |x| < 2 \\ 0 & \text{if } |x| \geq 2 \end{cases} \\ u_n(x) &:= n^{-\gamma} u_1(nx) \quad \text{for } n \geq 2 \end{aligned}$$

Observe that  $u_n \in W_0^{1,p}(\Omega)$  since

$$(2.8) \quad |\nabla u_n(x)| = \begin{cases} |\gamma| |x|^{\gamma-1} & \text{if } \frac{1}{n} \leq |x| < \frac{2}{n} \\ 0 & \text{otherwise} \end{cases}$$

and

$$(2.9) \quad \|\nabla u_n\|_{L^p(\Omega)}^p = |\gamma|^p N \omega_N \log 2,$$

where  $\omega_N$  denotes the measure of the unit ball of  $\mathbb{R}^N$ . In particular,  $\|\nabla u_n\|_{L^p(\Omega)}^p$  is independent of  $n$ . On the other hand, a direct calculation shows that

$$(2.10) \quad \begin{aligned} \|(b|u_n|)^{p-1}\|_{L^{p'}(\Omega)}^{p'} &= \int_{|x| < \frac{3}{n}} (b|u_n|)^p dx \\ &= \int_{|x| < \frac{1}{n}} (b|u_n|)^p dx + \int_{\frac{1}{n} \leq |x| < \frac{2}{n}} (b|u_n|)^p dx \\ &= N \omega_N \left[ \frac{(1-2^\gamma)^p}{N-p} + \int_1^2 r^{N-p} (r^\gamma - 2^\gamma)^p \frac{dr}{r} \right] \end{aligned}$$

Hence, we see that the norm of  $(b|u_n|)^{p-1}$  in  $L^{p'}(\Omega)$  is independent of  $n$  as well and strictly positive. On the other hand,  $(b|u_n|)^{p-1} \rightarrow 0$  pointwise in  $\Omega$  and this readily implies that there is no subsequence of  $\{(b|u_n|)^{p-1}\}_{n \in \mathbb{N}}$  strongly converging in  $L^{p'}(\Omega)$ .  $\square$

#### 2.4. An elementary lemma.

**Lemma 2.1.** *Assume  $f_n \rightarrow f$  a.e. Moreover, let  $g_n$ ,  $n \in \mathbb{N}$ , and  $g$  in  $L^q$ ,  $1 \leq q < +\infty$ , verify  $g_n \rightarrow g$  a.e.,  $|f_n| \leq g_n$  a.e.,  $\forall n \in \mathbb{N}$ , and*

$$\int_{\Omega} g_n^q dx \rightarrow \int_{\Omega} g^q dx.$$

*Then  $f_n, f \in L^q$  and*

$$f_n \rightarrow f \text{ in } L^q.$$

It suffices to apply Fatou lemma to the sequence of nonnegative functions

$$F_n = 2^{q-1}(g_n^q + g^q) - |f_n - f|^q.$$

### 3. A WEAK COMPACTNESS RESULT

The aim of this section is to establish a weak compactness criterion in the space  $W_0^{1,p}(\Omega)$  that has an interest by itself.

**Lemma 3.1.** *Let  $\mathcal{B}$  be a nonempty subset of  $W_0^{1,p}(\Omega)$ . Assume that there exists a constant  $C > 0$  such that*

$$(3.1) \quad \|\nabla u\|_{L^p(\Omega \setminus \Omega_\sigma)}^p \leq C \left( 1 + \|u\|_{L^p(\Omega \setminus \Omega_\sigma)}^p \right)$$



for any  $\sigma > 0$  and  $u \in \mathcal{B}$ , where  $\Omega_\sigma := \{x \in \Omega : |u(x)| \geq \sigma\}$ . Then, there exists a constant  $M > 0$  such that

$$(3.2) \quad \|u\|_{W^{1,p}(\Omega)} \leq M$$

for any  $u \in \mathcal{B}$ .

*Proof.* We argue by contradiction and assume  $\mathcal{B}$  unbounded. Then we construct a sequence  $\{u_k\}_k$  in  $\mathcal{B}$  such that

$$\|u_k\| := \|\nabla u_k\|_p \rightarrow \infty$$

as  $k \rightarrow \infty$ . By (3.1) we get, for any  $k \in \mathbb{N}$  and  $\varepsilon > 0$

$$(3.3) \quad \int_{\Omega} |\nabla T_{\varepsilon\|u_k\|} u_k|^p dx \leq C \left( 1 + \int_{\Omega} |u_k|^p \chi_{\{|u_k| < \varepsilon\|u_k\|\}} dx \right)$$

We set

$$v_k = \frac{u_k}{\|u_k\|}.$$

Hence, there exists  $v \in W_0^{1,p}(\Omega)$  such that (up to a subsequence)  $v_k \rightharpoonup v$  weakly in  $W_0^{1,p}$ ,  $v_k \rightarrow v$  strongly in  $L^p$  and  $v_k(x) \rightarrow v(x)$  for a.e.  $x \in \Omega$ . Notice that

$$\frac{T_{\varepsilon\|u_k\|} u_k}{\|u_k\|} = T_{\varepsilon} v_k,$$

thus  $\nabla T_{\varepsilon\|u_k\|} u_k = 0$  on the set  $\{x \in \Omega : |v_k(x)| \geq \varepsilon\}$ . Dividing (3.3) by  $\|u_k\|^p$  we have

$$(3.4) \quad \int_{\Omega} |\nabla T_{\varepsilon} v_k|^p dx \leq C \left( \|u_k\|^{-p} + \int_{\Omega} |v_k|^p \chi_{\{|v_k| < \varepsilon\}} dx \right)$$

Now, we let  $k \rightarrow +\infty$ . To this end, we note that  $T_{\varepsilon} v_k \rightharpoonup T_{\varepsilon} v$  weakly in  $W_0^{1,p}(\Omega)$  and  $T_{\varepsilon} v_k \rightarrow T_{\varepsilon} v$  strongly in  $L^p(\Omega)$ . In the left hand side of (3.4), we use semicontinuity of the norm with respect to weak convergence, while in the right hand side we observe that  $\|u_k\|^{-1} \rightarrow 0$ . Moreover, if

$$(3.5) \quad |\{x \in \Omega : |v(x)| = \varepsilon\}| = 0,$$

then we have  $\chi_{\{|v_k| < \varepsilon\}} \rightarrow \chi_{\{|v| < \varepsilon\}}$  a.e. in  $\Omega$  and hence

$$v_k \chi_{\{|v_k| < \varepsilon\}} \rightarrow v \chi_{\{|v| < \varepsilon\}}$$

strongly in  $L^p$ . Note that the set of values  $\varepsilon > 0$  for which (3.5) fails is at most countable. Thus, we end up with the following estimate

$$(3.6) \quad \int_{\Omega} |\nabla T_{\varepsilon} v|^p dx \leq C \int_{\Omega} |v|^p \chi_{\{|v| < \varepsilon\}} dx$$

Using Poincaré inequality in the left hand side, this yields

$$\varepsilon^p |\{x : |v| \geq \varepsilon\}| \leq C \varepsilon^p |\{x : 0 < |v| < \varepsilon\}|.$$

Passing to the limit as  $\varepsilon \downarrow 0$  (assuming (3.5)), we deduce

$$|\{x : |v| > 0\}| = 0,$$

that is,  $v(x) = 0$  a.e. Once we know that  $v_k \rightharpoonup 0$  weakly in  $W_0^{1,p}(\Omega)$ , the above argument (formally with  $\varepsilon = +\infty$ , i.e. without truncating  $v_k$ ) actually shows that  $v_k \rightarrow 0$  strongly in  $W_0^{1,p}(\Omega)$ , compare with (3.6), and this is not possible, as  $\|v_k\| = 1$ , for all  $k$ .  $\square$

#### 4. PROOF OF THEOREM 1.1

**4.1. The case of bounded coefficient.** In this subsection we assume  $b \in L^\infty(\Omega)$ . For a given function  $v \in L^p(\Omega)$ , we define the vector field on  $\Omega \times \mathbb{R}^N$

$$(4.1) \quad A_v(x, \xi) := A(x, v(x), \xi)$$

which satisfies similar conditions as  $A$ , namely

$$(4.2) \quad \langle A_v(x, \xi), \xi \rangle \geq \alpha |\xi|^p - (b(x) |v|)^p - \varphi(x)^p$$

$$(4.3) \quad |A_v(x, \xi)| \leq \beta |\xi|^{p-1} + (b(x) |v|)^{p-1} + \varphi(x)^{p-1}$$

$$(4.4) \quad \langle A_v(x, \xi) - A_v(x, \eta), \xi - \eta \rangle > 0 \quad \text{for } \xi \neq \eta$$

Hence, we can consider a quasilinear elliptic operator similar to (1.4)

$$(4.5) \quad u \in W_0^{1,p}(\Omega) \quad \mapsto \quad -\operatorname{div} A_v(x, \nabla u) \in W^{-1,p'}(\Omega)$$

defined by the rule

$$(4.6) \quad \langle -\operatorname{div} A_v(x, \nabla u), w \rangle = \int_{\Omega} \langle A(x, v, \nabla u), \nabla w \rangle \, dx$$

for any  $w \in W_0^{1,p}(\Omega)$ . The operator at (4.5) is invertible. Indeed,

**Proposition 4.1.** *For every  $\Phi \in W^{-1,p'}(\Omega)$ , there exists a unique  $u \in W_0^{1,p}(\Omega)$  such that*

$$(4.7) \quad -\operatorname{div} A_v(x, \nabla u) = \Phi$$

Moreover, the mapping

$$(4.8) \quad (v, \Phi) \in L^p(\Omega) \times W^{-1,p'}(\Omega) \quad \mapsto \quad u \in W_0^{1,p}(\Omega)$$

is continuous.

*Proof.* Existence of a solution is classical, see e.g. [27], [7, pg. 27], or [26, Théorème 2.8, pg. 183]. Uniqueness trivially holds by monotonicity.

For the sake of completeness, we prove continuity of the map (4.8). Given  $v_n \rightarrow v$  in  $L^p(\Omega)$  and  $\Phi_n \rightarrow \Phi$  in  $W^{-1,p'}(\Omega)$ , let  $u_n \in W_0^{1,p}(\Omega)$  solve

$$(4.9) \quad -\operatorname{div} A(x, v_n, \nabla u_n) = \Phi_n.$$

The sequence  $\{u_n\}_n$  is clearly bounded, hence we may assume  $u_n \rightharpoonup u$  weakly in  $W_0^{1,p}(\Omega)$ . Moreover, testing equation (4.9) with  $u_n - u$ , we have

$$(4.10) \quad \lim_{n \rightarrow \infty} \int_{\Omega} A(x, v_n, \nabla u_n)(\nabla u_n - \nabla u) \, dx = \lim_{n \rightarrow \infty} \langle \Phi_n, u_n - u \rangle = 0.$$

On the other hand, we easily see that  $A(x, v_n, \nabla u) \rightarrow A(x, v, \nabla u)$  strongly in  $L^{p'}(\Omega, \mathbb{R}^N)$  and thus (4.10) implies

$$(4.11) \quad \lim_{n \rightarrow \infty} \int_{\Omega} [A(x, v_n, \nabla u_n) - A(x, v_n, \nabla u)] \nabla(u_n - u) \, dx = 0.$$

The integrands in (4.11) are nonnegative by monotonicity. Hence, arguing as in the proof of [27, Lemma 3.3], we also get  $\nabla u_n(x) \rightarrow \nabla u(x)$  a.e. in  $\Omega$ , and

$$A(x, v_n, \nabla u_n) \rightharpoonup A(x, v, \nabla u)$$

weakly in  $L^{p'}(\Omega, \mathbb{R}^N)$ . Combining this with (4.10) yields

$$(4.12) \quad \lim_{n \rightarrow \infty} \int_{\Omega} A(x, v_n, \nabla u_n) \nabla u_n \, dx = \int_{\Omega} A(x, v, \nabla u) \nabla u \, dx.$$

By coercivity condition (1.1), we deduce

$$\alpha |\nabla u_n|^p \leq A(x, v_n, \nabla u_n) \nabla u_n + (b|v_n|)^p + \varphi^p$$

Trivially  $\int_{\Omega} (b|v_n|)^p \, dx$  converges to  $\int_{\Omega} (b|v|)^p \, dx$ . In view of (4.12), by Lemma 2.1 we get  $u_n \rightarrow u$  strongly in  $W_0^{1,p}(\Omega)$ , and  $u$  solves the equation

$$-\operatorname{div} A(x, v, \nabla u) = \Phi.$$

□

In view of Rellich Theorem, we have

**Corollary 4.1.** *For fixed  $\Phi \in W^{-1,p'}(\Omega)$ , the mapping*

$$(4.13) \quad \mathcal{F}: v \in W_0^{1,p}(\Omega) \mapsto u \in W_0^{1,p}(\Omega)$$

*which takes  $v$  to the unique solution  $u$  of equation (4.7) is compact.*

Now we state an existence result to Problem (1.6) when  $b \in L^\infty(\Omega)$ .

**Proposition 4.2.** *Let (1.1), (1.2) and (1.3) be in charge with  $b \in L^\infty(\Omega)$ . Then Problem (1.6) has a solution  $u \in W_0^{1,p}(\Omega)$ .*

*Proof.* If  $\mathcal{F}$  is the operator defined in Corollary 4.1, clearly a fixed point of  $\mathcal{F}$  is a solution to Problem (1.6). To apply Leray-Schauder theorem, we need an a priori estimate on the solution  $u \in W_0^{1,p}(\Omega)$  of the equation

$$u = t\mathcal{F}[u]$$

that is

$$(4.14) \quad -\operatorname{div} A\left(x, u, \frac{1}{t} \nabla u\right) = \Phi,$$

as  $t \in ]0, 1]$  varies. By using  $T_\sigma u$  with  $\sigma > 0$  as a test function in (4.14) we get

$$(4.15) \quad \int_{\Omega} \left\langle A\left(x, u, \frac{1}{t} \nabla u\right), \nabla T_\sigma u \right\rangle dx = \langle \Phi, T_\sigma u \rangle$$

Therefore, using the point-wise condition (4.2) we get

$$(4.16) \quad \alpha t^{1-p} \int_{\Omega} |\nabla T_\sigma u|^p dx \leq \|\Phi\| \|\nabla T_\sigma u_k\|_p + \int_{\Omega} \left[ b(x)^p |u|^p \chi_{\{|u| < \sigma\}} + \varphi(x)^p \right] dx$$

As  $0 < t \leq 1$ , by Young inequality (4.16) yields

$$(4.17) \quad \frac{\alpha}{2} \int_{\Omega} |\nabla T_\sigma u|^p dx \leq \|\Phi\|^{p'} + \|b\|_\infty^p \int_{\Omega} |u|^p \chi_{\{|u| \leq \sigma\}} dx + \|\varphi\|_p^p$$

The conclusion follows by Lemma 3.1.  $\square$

**4.2. The approximating problems.** For each  $n \in \mathbb{N}$ , we set

$$(4.18) \quad \vartheta_n(x) = \frac{T_n b(x)}{b(x)}, \quad \text{a.e. } x \in \Omega,$$

and define the vector field

$$(4.19) \quad A_n : (x, u, \xi) \in \Omega \times \mathbb{R} \times \mathbb{R}^N \mapsto \mathbb{R}^N$$

letting

$$(4.20) \quad A_n(x, u, \xi) = A(x, \vartheta_n u, \xi)$$

The vector field  $A_n$  has similar properties as  $A$ , with  $b$  replaced by  $T_n b$ . More precisely,

$$(4.21) \quad \langle A_n(x, u, \xi), \xi \rangle \geq \alpha |\xi|^p - (T_n b(x) |u|)^p - \varphi(x)^p$$

$$(4.22) \quad |A_n(x, u, \xi)| \leq \beta |\xi|^{p-1} + (T_n b(x) |u|)^{p-1} + \varphi(x)^{p-1}$$

$$(4.23) \quad \langle A_n(x, u, \xi) - A_n(x, u, \eta), \xi - \eta \rangle > 0 \quad \text{for } \xi \neq \eta$$

Applying Proposition 4.2 with  $A_n$  in place of  $A$ , fixed  $\Phi \in W^{-1,p'}(\Omega)$ , we find  $u_n \in W_0^{1,p}(\Omega)$  such that

$$(4.24) \quad -\operatorname{div} A_n(x, u_n, \nabla u_n) = \Phi.$$

Notice that we have, for  $\sigma > 0$

$$(4.25) \quad \alpha \int_{\Omega} |\nabla T_{\sigma} u_n|^p dx \leq \|\Phi\| \|\nabla T_{\sigma} u_n\|_p + \int_{\Omega} \left[ (T_n b)^p |u_n|^p \chi_{\{|u_n| < \sigma\}} + \varphi^p \right] dx$$

which implies

$$(4.26) \quad \alpha^{\frac{1}{p}} \|\nabla T_{\sigma} u_n\|_p \leq (\|\Phi\| \|\nabla T_{\sigma} u_n\|_p)^{\frac{1}{p}} + \|(T_n b) u_n \chi_{\{|u_n| < \sigma\}}\|_p + \|\varphi\|_p$$

Our next step consists in showing that the sequence  $\{u_n\}_n$  is bounded in  $W_0^{1,p}(\Omega)$ . Let  $m$  be a positive integer to be chosen later. For  $n \geq m$  we have

$$T_n b \leq T_m b + (b - T_m b)$$

and hence

$$(4.27) \quad \|(T_n b) u_n \chi_{\{|u_n| < \sigma\}}\|_p \leq \|(T_m b) u_n \chi_{\{|u_n| < \sigma\}}\|_p + \|(b - T_m b) u_n \chi_{\{|u_n| < \sigma\}}\|_p$$

Using Hölder and Sobolev inequalities we get

$$\|(b - T_m b) u_n \chi_{\{|u_n| < \sigma\}}\|_p \leq \|b - T_m b\|_{N,\infty} \|T_{\sigma} u_n\|_{p^*,p} \leq S_{N,p} \|b - T_m b\|_{N,\infty} \|\nabla T_{\sigma} u_n\|_p$$

Then (4.26) and (4.27) give

$$(4.28) \quad \alpha^{\frac{1}{p}} \|\nabla T_{\sigma} u_n\|_p \leq (\|\Phi\| \|\nabla T_{\sigma} u_n\|_p)^{\frac{1}{p}} + \|(T_m b) u_n \chi_{\{|u_n| < \sigma\}}\|_p + \|\varphi\|_p + S_{N,p} \|b - T_m b\|_{L^{N,\infty}(\Omega)} \|\nabla T_{\sigma} u_n\|_{L^p(\Omega)}$$

By our assumption (1.9), the level  $m$  can be chosen large enough so that

$$S_{N,p} \|b - T_m b\|_{L^{N,\infty}(\Omega)} < \alpha^{\frac{1}{p}}$$

Then, by absorbing in (4.28) the latter term of the right hand side in the left hand side, we get

$$(4.29) \quad C \int_{\Omega} |\nabla T_{\sigma} u_n|^p dx \leq \|\Phi\| \|\nabla T_{\sigma} u_n\|_p + \int_{\Omega} \left[ (T_m b)^p |T_{\sigma} u_n|^p \chi_{\{|u_n| < \sigma\}} + \varphi^p \right] dx$$

for a positive constant  $C$  which is independent of  $n$ . Now, it is clear that (4.29), via Young inequality, allows us to apply Lemma 3.1, then

$$(4.30) \quad \|u_n\| \leq M$$

for a constant  $M$  independent of  $n$ .

In the model case (1.8), it is easy to show that the operator  $\mathcal{F}$  defined in (4.13) is compact, also for  $b \in L^N(\Omega)$  (see Remark 4.1 below). In the general case, in which  $b \in L^{N,\infty}(\Omega)$  we need more work.

**4.3. Passing to the limit.** Now, we are in a position to conclude the proof of Theorem 1.1. Taking into account estimate (4.30) we may assume

(4.31)

$$\begin{aligned} u_n &\rightharpoonup u && \text{in } W_0^{1,p}(\Omega) \text{ weakly} \\ u_n &\rightarrow u && \text{in } L^q(\Omega) \text{ strongly for any } q < p^*, \text{ and also a.e. in } \Omega \end{aligned}$$

for some  $u \in W_0^{1,p}(\Omega)$ . We shall conclude our proof showing that  $u$  solves Problem (1.6). In the rest of our argument, we let for simplicity  $\gamma(t) := \arctan t$ . Obviously,  $\gamma \in C^1(\mathbb{R})$ ,  $|\gamma(t)| \leq |t|$  and  $0 \leq \gamma'(t) \leq 1$  for all  $t \in \mathbb{R}$ . In particular,  $\gamma$  is Lipschitz continuous in the whole of  $\mathbb{R}$  and therefore

$$u_n, u \in W_0^{1,p}(\Omega) \implies \gamma(u_n - u) \in W_0^{1,p}(\Omega).$$

Moreover, since  $\gamma(0) = 0$  we have

$$(4.32) \quad \gamma(u_n - u) \rightharpoonup 0 \quad \text{in } W_0^{1,p}(\Omega) \text{ weakly}.$$

Testing equation (4.24) with the function  $\gamma(u_n - u)$  we get

$$\int_{\Omega} A_n(x, u_n, \nabla u_n) \nabla \gamma(u_n - u) \, dx = \langle \Phi, \gamma(u_n - u) \rangle$$

where  $\nabla \gamma(u_n - u) = \gamma'(u_n - u)(\nabla u_n - \nabla u)$ . In view of (4.32) we necessarily have

$$(4.33) \quad \lim_{n \rightarrow \infty} \int_{\Omega} A_n(x, u_n, \nabla u_n) \nabla \gamma(u_n - u) \, dx = 0.$$

We claim that

$$(4.34) \quad \lim_{n \rightarrow \infty} \int_{\Omega} A_n(x, u_n, \nabla u) \nabla \gamma(u_n - u) \, dx = 0.$$

In order to prove (4.34), since  $\nabla u_n - \nabla u \rightharpoonup 0$ , it suffices to show that

$$(4.35) \quad \gamma'(u_n - u) A_n(x, u_n, \nabla u) = \frac{A_n(x, u_n, \nabla u)}{1 + |u_n - u|^2} \quad \text{is compact in } L^{p'}.$$

Preliminarily, we observe that combining (4.31) with the property that  $\vartheta_n \rightarrow 1$  as  $n \rightarrow \infty$ , we have

$$\frac{A_n(x, u_n, \nabla u)}{1 + |u_n - u|^2} \rightarrow A(x, u, \nabla u) \quad \text{a.e. in } \Omega.$$

We are going to use Lemma 2.1. To this end, by (4.22) we deduce that

$$\left| \frac{A_n(x, u_n, \nabla u)}{1 + |u_n - u|^2} \right|^{p'} \leq C \left[ |\nabla u|^p + \varphi^p + (b|u|)^p + \frac{(b|u_n - u|)^p}{1 + |u_n - u|^2} \right]$$

for a positive constant  $C = C(p, \beta)$ . Hence, we can pass to the limit if  $1 < p \leq 2$ . For  $p > 2$  we choose  $s$  satisfying

$$\frac{p^*}{p} < s < \frac{p^*}{p-2},$$

so that  $ps' < N$ , and we conclude also in this case, further estimating with the aid of Young inequality

$$\frac{(b|u_n - u|)^p}{1 + |u_n - u|^2} \leq b^{ps'} + |u_n - u|^{(p-2)s}.$$

Now, from (4.33) and (4.34) we get

$$(4.36) \quad \lim_{n \rightarrow \infty} \int_{\Omega} [A_n(x, u_n, \nabla u_n) - A_n(x, u_n, \nabla u)] \nabla \gamma(u_n - u) \, dx = 0.$$

As the integrand is nonnegative, we have (up to a subsequence)

$$[A_n(x, u_n, \nabla u_n) - A_n(x, u_n, \nabla u)] \nabla \gamma(u_n - u) \rightarrow 0$$

a.e. in  $\Omega$ . Moreover, since  $\gamma'(u_n - u) \rightarrow 1$  a.e. in  $\Omega$ , the above in turn implies

$$(4.37) \quad [A_n(x, u_n, \nabla u_n) - A_n(x, u_n, \nabla u)] (\nabla u_n - \nabla u) \rightarrow 0$$

Arguing as in the proof of [27, Lemma 3.3], we see that

$$(4.38) \quad \nabla u_n \rightarrow \nabla u \quad \text{a.e. in } \Omega$$

and

$$(4.39) \quad A_n(x, u_n, \nabla u_n) \rightharpoonup A(x, u, \nabla u) \quad \text{in } L^{p'}(\Omega, \mathbb{R}^N) \text{ weakly}$$

and we conclude that  $u$  is a solution to the original problem (1.6).

**Remark 4.1.** We discuss briefly the particular case in which the operator has the form

$$A(x, v, \xi) = A'(x, \xi) + A''(x, v),$$

with

$$|A''(x, v)| \leq (b(x) |v|)^{p-1} + \varphi(x)^{p-1}.$$

and  $b \in L^N(\Omega)$  (see also [5]). We can easily show that the operator  $\mathcal{F}$  defined in (4.13) is compact, also for  $b \in L^N(\Omega)$ . Indeed, equation (4.7) in this case becomes

$$(4.40) \quad -\operatorname{div} A'(x, \nabla u) = \Phi + \operatorname{div} A''(x, v).$$

Defined  $\vartheta_n$  as in (4.18), each mapping

$$v \in W_0^{1,p}(\Omega) \quad \mapsto \quad A''(x, \vartheta_n v) \in L^{p'}(\Omega, \mathbb{R}^N)$$

is clearly compact. Moreover,

$$(4.41) \quad |A''(x, v) - A''(x, \vartheta_n v)| \leq 2[(b|v|)^{p-1} + \varphi^{p-1}] \chi_{E_n},$$

where

$$E_n = \{x \in \Omega : |b(x)| > n\}.$$

Therefore, as  $n \rightarrow +\infty$  we have

$$A''(x, \vartheta_n v) \rightarrow A''(x, v) \quad \text{strongly in } L^{p'}(\Omega, \mathbb{R}^N),$$

the convergence being uniform when  $v$  varies in a bounded subset of  $W_0^{1,p}(\Omega)$ , and compactness is preserved for the limit mapping.

An a priori bound for solutions of equation

$$u = t \mathcal{F}[u]$$

can be easily obtained as above, splitting  $b \in L^N(\Omega)$  as

$$b = T_m b + (b - T_m b)$$

for a sufficiently large  $m$ . Therefore, in this particular case the existence result of Theorem 1.1 follows simply applying Leray–Schauder fixed point theorem.

## 5. THE OBSTACLE PROBLEM

This section is devoted to the obstacle problem naturally related with problem (1.6) (see [25] for a comprehensive treatment of the topic). We again assume that (1.1), (1.2) and (1.3) are in charge and we let  $\Phi \in W^{-1,p}(\Omega)$ . Given a measurable function  $\psi: \Omega \rightarrow \overline{\mathbb{R}}$ , where  $\overline{\mathbb{R}} := [-\infty, \infty]$ , we consider the convex subset of  $\mathcal{K}_\psi(\Omega)$  of  $W_0^{1,p}(\Omega)$  given by

$$(5.1) \quad \mathcal{K}_\psi(\Omega) := \{w \in W_0^{1,p}(\Omega) : w \geq \psi \text{ a.e. in } \Omega\}.$$

We will assume that  $\mathcal{K}_\psi(\Omega)$  is nonempty. An element  $u \in \mathcal{K}_\psi(\Omega)$  is a solution to the obstacle problem associated with (1.6) if the following variational inequality holds

$$(5.2) \quad \int_{\Omega} \langle A(x, u, \nabla u), \nabla(w - u) \rangle dx \geq \langle \Phi, w - u \rangle, \quad \forall w \in \mathcal{K}_\psi(\Omega).$$

As  $\mathcal{K}_\psi(\Omega) \neq \emptyset$ , we may assume without loss of generality that

$$(5.3) \quad \psi \leq 0 \text{ a.e. in } \Omega.$$

In fact, if  $g \in \mathcal{K}_\psi(\Omega)$ , then one can consider the operator defined by the vector field

$$\tilde{A}(x, u, \xi) := A(x, u + g(x), \xi + \nabla g(x)),$$



satisfying conditions similar to (1.1), (1.2) and (1.3). Now it is clear that, if function  $\tilde{u} \in \mathcal{K}_{\psi-g}(\Omega)$  satisfies the following variational inequality

$$(5.4) \quad \int_{\Omega} \langle \tilde{A}(x, \tilde{u}, \nabla \tilde{u}), \nabla(w - \tilde{u}) \rangle dx \geq \langle \Phi, w - \tilde{u} \rangle \quad \forall w \in \mathcal{K}_{\psi-g}(\Omega)$$

correspondingly  $u = \tilde{u} + g$  is a solution to (5.2). Notice that the obstacle function for problem (5.4) is nonpositive, as we are assuming for the original problem.

**Theorem 5.1.** *Let  $\Phi \in W^{-1,p'}(\Omega)$  and  $\psi : \Omega \rightarrow [-\infty, 0]$  be a measurable function. Under the assumption (1.1), (1.2) and (1.3), if (1.9) holds, then the obstacle problem (5.2) admits a solution.*

*Proof.* We follow closely the arguments of Section 4. For each  $n \in \mathbb{N}$ , we consider the function  $\vartheta_n$  as in (4.18) and define the vector fields  $A_n = A_n(x, u, \xi)$  as in (4.20). We consider a sequence of obstacle problems provided by

$$(5.5) \quad \int_{\Omega} \langle A_n(x, u_n, \nabla u_n), \nabla(w - u_n) \rangle dx \geq \langle \Phi, w - u_n \rangle, \quad \forall w \in \mathcal{K}_{\psi}(\Omega).$$

The existence of a solution  $u_n \in \mathcal{K}_{\psi}(\Omega)$  to (5.5) is proven applying [26, Théorème 8.2, pg. 247] to the operator

$$-\operatorname{div} A_n(x, v, \nabla u),$$

for a fixed  $v \in W_0^{1,p}(\Omega)$ , and then using Leray–Schauder Theorem, arguing as in Subsection 4.1. Due to (5.3), for every  $k > 0$  the function

$$w := u_n - T_k u_n \in \mathcal{K}_{\psi}(\Omega)$$

is a test function for (5.5). Arguing as in Section 4.2 we obtain

$$\|u_n\| \leq M$$

with  $M$  independent of  $n$  (as in (4.30)). Therefore (4.31) holds for some  $u \in W_0^{1,p}(\Omega)$ . It is clear from (4.31) itself that

$$(5.6) \quad u \in \mathcal{K}_{\psi}(\Omega)$$

As for Theorem 1.1, we shall prove that  $u$  is a solution to the original problem (5.2). We proceed as follows. We use

$$(5.7) \quad w := u_n - \gamma(u_n - v)$$

in (5.5), where  $\gamma(s) = \lambda \arctan(s/\lambda)$ , for  $\lambda > 0$ , and  $v \in \mathcal{K}_{\psi}(\Omega)$  is arbitrary. Note that this is a legitimate test function, that is  $w \in \mathcal{K}_{\psi}(\Omega)$ . Indeed, on the set where  $u_n \geq v$  we have  $\gamma(u_n - v) \leq u_n - v$

and so  $w \geq v$ ; on the other hand, on the set where  $u_n \leq v$  we have  $\gamma(u_n - v) \leq 0$  and so  $w \geq u_n$ . Therefore, from (5.5) we get

$$(5.8) \quad \int_{\Omega} A_n(x, u_n, \nabla u_n) \nabla \gamma(u_n - v) \, dx \leq \langle \Phi, \gamma(u_n - v) \rangle .$$

Following the lines of the proof of Theorem 1.1 (where  $\lambda = 1$ ), we get in turn (4.36), (4.38) and finally (4.39). To pass to the limit for fixed general  $\lambda > 0$  in (5.8), we rewrite it as follows:

$$(5.9) \quad \begin{aligned} \int_{\Omega} [A_n(x, u_n, \nabla u_n) - A_n(x, u_n, \nabla v)] \nabla \gamma(u_n - v) \, dx \\ \leq \langle \Phi, \gamma(u_n - v) \rangle - \int_{\Omega} A_n(x, u_n, \nabla v) \nabla \gamma(u_n - v) \, dx . \end{aligned}$$

In the left hand side we use Fatou lemma, as by monotonicity condition (1.3) the integrand is nonnegative. In the right hand side, we note that  $A_n(x, u_n, \nabla v) \gamma'(u_n - v)$  converges to  $A(x, u, \nabla v) \gamma'(u - v)$  in  $L^{p'}$ , compare with (4.35) where we did not use that  $u_n \rightarrow u$ . Hence, we deduce from (5.9)

$$\begin{aligned} \int_{\Omega} [A(x, u, \nabla u) - A(x, u, \nabla v)] \nabla \gamma(u - v) \, dx \\ \leq \langle \Phi, \gamma(u - v) \rangle - \int_{\Omega} A(x, u, \nabla v) \nabla \gamma(u - v) \, dx , \end{aligned}$$

that is

$$(5.10) \quad \int_{\Omega} A(x, u, \nabla u) \nabla \gamma(u - v) \, dx \leq \langle \Phi, \gamma(u - v) \rangle .$$

Now we let  $\lambda \rightarrow \infty$  in (5.10), noting that  $\gamma(u - v) \rightarrow u - v$  strongly in  $W_0^{1,p}(\Omega)$ . Therefore, we get

$$\int_{\Omega} A(x, u, \nabla u) \nabla (u - v) \, dx \leq \langle \Phi, u - v \rangle ,$$

for all  $v \in K_{\psi}(\Omega)$ , which means exactly that  $u$  is a solution to our obstacle problem.  $\square$

**Remark 5.1.** Clearly, Theorem 5.1 is more general than Theorem 1.1 since we are allowed to choose  $\psi \equiv -\infty$ . Indeed, in this case, the obstacle problem (5.2) reduces to (1.6).

## 6. REGULARITY OF THE SOLUTION

In this Section, following [17] we study regularity of the problem (1.6).

**Theorem 6.1.** *Let  $1 < p < r < N$  and  $\Phi \in W^{-1, \frac{r}{p-1}}(\Omega)$ . Assume that (1.1), (1.2) and (1.3) hold with  $\varphi \in L^r(\Omega)$ . Under these hypotheses, there exists  $\delta = \delta(\alpha, N, p, r) > 0$  such that if*

$$(6.1) \quad \text{dist}_{L^{N,\infty}}(b, L^\infty(\Omega)) < \delta,$$

*then any solution  $u \in W_0^{1,p}(\Omega)$  of (1.6) satisfies*

$$(6.2) \quad |u|^{r^*/p^*} \in W_0^{1,p}(\Omega)$$

*In particular  $u \in L^{r^*}(\Omega)$ .*

*Proof.* Let  $u \in W_0^{1,p}(\Omega)$  be a solution of (1.6). We may write  $\Phi \in W^{-1, \frac{r}{p-1}}(\Omega)$  as

$$\Phi = \text{div}(|F|^{p-2}F)$$

for a suitable  $F \in L^r(\Omega, \mathbb{R}^N)$ .

For fixed  $k > 0$ , we use  $v := u - T_k u$  as a test function in (1.7) to get

$$(6.3) \quad \alpha \int_{\Omega_k} |\nabla u|^p dx \leq \int_{\Omega_k} |F|^{p-1} |\nabla u| dx + \int_{\Omega_k} (b^p |u|^p + \varphi^p) dx$$

where  $\Omega_k$  denotes the superlevel set  $\{|u| > k\}$ . For  $0 < \varepsilon < \alpha$ , by Young inequality we get

$$(6.4) \quad (\alpha - \varepsilon) \int_{\Omega_k} |\nabla u|^p dx \leq \int_{\Omega_k} (C |F|^p + b^p |u|^p + \varphi^p) dx$$

with  $C = C(p, \varepsilon) > 0$ . We let

$$(6.5) \quad \lambda = \frac{r^*}{p^*} - 1$$

and multiply both sides of (6.4) by  $k^{p\lambda-1}$  and integrate w.r.t.  $k$  over the interval  $[0, K]$ , for  $K > 0$  fixed. By Fubini theorem we have

$$(6.6) \quad (\alpha - \varepsilon) \int_{\Omega} |\nabla u|^p |T_K u|^{p\lambda} dx \leq \int_{\Omega} (C |F|^p + b^p |u|^p + \varphi^p) |T_K u|^{p\lambda} dx$$

which implies

$$(6.7) \quad (\alpha - \varepsilon)^{\frac{1}{p}} \|\nabla u |T_K u|^\lambda\|_p \leq C \|F |T_K u|^\lambda\|_p + \|b u |T_K u|^\lambda\|_p + \|\varphi |T_K u|^\lambda\|_p$$

For  $M > 0$  we write

$$(6.8) \quad \|b u |T_K u|^\lambda\|_p \leq \|(b - T_M b) u |T_K u|^\lambda\|_p + M \|u |T_K u|^\lambda\|_p$$

By Hölder inequality and Sobolev embedding Theorem 2.1

$$(6.9) \quad \begin{aligned} \|(b - T_M b) u |T_K u|^\lambda\|_p &\leq \|b - T_M b\|_{N,\infty} \|u |T_K u|^\lambda\|_{p^*,p} \\ &\leq \|b - T_M b\|_{N,\infty} S_{N,p} \|\nabla(u |T_K u|^\lambda)\|_p \end{aligned}$$

Moreover,

$$(6.10) \quad |\nabla(u |T_K u|^\lambda)| \leq (1 + \lambda) |\nabla u| |T_K u|^\lambda$$

Therefore

$$(6.11) \quad \|(b - T_M b) u |T_K u|^\lambda\|_{L^p(\Omega)} \leq \|b - T_M b\|_{N,\infty} S_{N,p} (1 + \lambda) \|\nabla u |T_K u|^\lambda\|_p$$

Under the assumption

$$(6.12) \quad \|b - T_M b\|_{N,\infty} S_{N,p} (1 + \lambda) < \alpha^{\frac{1}{p}}$$

choosing  $\varepsilon$  small enough we get from (6.7)

$$(6.13) \quad \|\nabla u |T_K u|^\lambda\|_p \leq C \|G |T_K u|^\lambda\|_p$$

with  $C = C(p, r, M, \alpha) > 0$ , where we set

$$(6.14) \quad G^p = |F|^p + |u|^p + \varphi^p.$$

We first show the claim under the additional assumption  $u \in L^r(\Omega)$ , so that  $G \in L^r(\Omega)$ . By Hölder inequality we have

$$(6.15) \quad \|G |T_K u|^\lambda\|_p \leq \|G\|_r \|T_K u\|_{\lambda \frac{rp}{r-p}}^\lambda$$

From (6.5) we get

$$(6.16) \quad \lambda \frac{rp}{r-p} = r^*.$$

Hence, by Sobolev embedding theorem we have

$$(6.17) \quad \begin{aligned} \|T_K u\|_{\lambda \frac{rp}{r-p}}^\lambda &= \|T_K u\|_{r^*}^\lambda = \| |T_K u|^{\frac{r^*}{p^*}} \|_{p^*}^{\lambda \frac{p^*}{r^*}} \leq C \|\nabla |T_K u|^{\frac{r^*}{p^*}}\|_p^{\lambda \frac{p^*}{r^*}} \\ &\leq C \|\nabla u |T_K u|^\lambda\|_p^{\frac{\lambda}{\lambda+1}} \end{aligned}$$

Then, combining (6.13), (6.15) and (6.17), we get

$$(6.18) \quad \|\nabla u |T_K u|^\lambda\|_p^{\frac{p^*}{r^*}} \leq C \|G\|_r$$

Passing to the limit as  $K \rightarrow +\infty$  and recalling (6.14), we have

$$(6.19) \quad \|\nabla u |u|^\lambda\|_p^{\frac{p^*}{r^*}} \leq C (\|F\|_r + \|\varphi\|_r + \|u\|_r)$$

that is

$$(6.20) \quad \|\nabla |u|^{\frac{r^*}{p^*}}\|_p \leq C (\|F\|_r + \|\varphi\|_r + \|u\|_r)^{\frac{r^*}{p^*}}$$

Hence, (6.2) holds as long as  $u \in L^r(\Omega)$ . At this point we observe that if  $r \leq p^*$ , using the Sobolev embedding theorem,  $u \in L^{p^*}(\Omega)$

and the proof is concluded. In the complementary case  $r > p^*$ , we use a bootstrap approach. Precisely, we repeat the previous argument replacing  $r$  with  $p^*$  to get  $u \in L^{p^{**}}(\Omega)$ . Using this information, if  $r \leq p^{**}$ , there is nothing left to prove. Otherwise we repeat previous argument again. In a finite number of similar steps we can conclude our proof.  $\square$

**Remark 6.1.** In view of (6.12), we may take

$$\delta = \frac{\alpha^{\frac{1}{p}}}{S_{N,p}} \frac{p^*}{r^*}$$

in (6.1). Since  $r \mapsto r^*$  is increasing, a similar condition to (6.12) will hold in all intermediate steps, in case we need the bootstrap argument. Note that  $\delta$  reduces to the distance in (1.9), for  $r = p$ .

## REFERENCES

- [1] Alfano E. A., Monsurr S., *Noncoercive nonlinear Dirichlet problems in unbounded domains*, Nonlinear Analysis 192 (2020) doi/10.1016/j.na.2019.111694
- [2] Alvino A., *Sulla disuguaglianza di Sobolev in Spazi di Lorentz*. Boll. Un. Mat. It. A (5) 14, (1977), 148-156.
- [3] Bennett C., Sharpley R., *Interpolation of operators*. Academic Press (1988)
- [4] Bensoussan, A., Boccardo, L., Murat, F., *On a non linear partial differential equation having natural growth terms and unbounded solution*, Annales de l'I.H.P. Analyse non linaire, Volume 5 (1988) no. 4, 347-364
- [5] Boccardo L., *Finite Energy Solutions of Nonlinear Dirichlet Problems with Discontinuous Coefficients*. Boll. Un. Mat. It. (9) 5, (2012), 357-368.
- [6] Brezis, H., Browder, F. E., *Strongly nonlinear elliptic boundary value problems*. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 5 (1978), no. 3, 587-603.
- [7] Browder F. E., *Existence theorems for nonlinear partial differential equations*, Proceedings of Symposia in Pure Mathematics, Vol. 16, S. S. CHERN and S. SMALE Eds., A.M.S., Providence, 1970, 1-60.
- [8] Cirmi, G.R., D'Asero, S., Leonardi, S., Porzio, M.M., *Local regularity results for solutions of linear elliptic equations with drift term*. Advances in Calculus of Variations. (2019) DOI10.1515/acv-2019-0048.
- [9] Di Gironimo P., Zecca G., *Sobolev-Zygmund solutions for nonlinear elliptic equations with growth coefficients in BMO*, Journal of Elliptic and Parabolic Equations, (2020) DOI : 10.1007/s41808-020-00064-y
- [10] Droniou J., *Non-coercive linear elliptic problems*, Potential Analysis, vol. 17, no. 2, pp. 181-203, 2002.
- [11] Farroni F., Moscariello G., *A nonlinear parabolic equation with drift term*, Nonlinear Analysis, 177, Part B, (2018), 397-412
- [12] Giannetti F., Greco L., Moscariello G., *Linear elliptic equations with lower order terms*, Differential Integral Equations 26, no. 5-6, (2013) 623-638.
- [13] Giaquinta, M., Giusti, E., *Quasiminima*. Ann. Inst. H. Poincaré Anal. Non Linaire 1 (1984), no. 2, 79-107.

- [14] Giusti E., *Direct Methods In The Calculus Of Variations*. World Scientific Publishing Co., Inc., River Edge, NJ, 2003. viii+403 pp. ISBN: 981-238-043-4.
- [15] Gilbarg D., Trudinger N.S., *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, Berlin. (1983).
- [16] Grafakos L. *Classical Fourier analysis*. Third edition. Graduate Texts in Mathematics, 249. Springer, New York, (2014).
- [17] Greco L., Moscariello G., Zecca G. *Regularity for solutions to nonlinear elliptic equations*, Differential and Integral Equations, 26 n. 9-10 (2013) 1105–1113.
- [18] Greco L., Moscariello G., Zecca G., An obstacle problem for noncoercive operators, *Abstract and Applied Analysis*, Article ID 890289, Volume 2015, Article ID 890289 8 pag. <http://dx.doi.org/10.1155/2015/890289>
- [19] Greco L., Moscariello G., Zecca G., Very weak solutions to elliptic equations with singular convection term. *Journal of Mathematical Analysis and Applications*, 457, Issue 2, 15 (2018). 1376–1387.
- [20] Greco L., Moscariello G., *An embedding Theorem in Lorentz-Zygmund spaces*. *Potential Anal.* 5, (1996), 581-590.
- [21] Greco L., Schiattarella R., *An embedding theorem for BV-functions*. *Comm. in Contemporary Math.* Vol. 22, No. 4 (2020) 1950032 (13 pages). DOI: 10.1142/S0219199719500329.
- [22] Kang B., Kim H.,  *$W^{2,p}$ -estimates for elliptic equations with lower order terms*, *Comm. Pure and Appl. Anal.*, Volume 16, Number 3, (2017), 799–821.
- [23] Kim H., Kim Y.-H., *On Weak Solutions of Elliptic Equations with Singular Drifts*, *SIAM J. Math. Anal.*, 47(2), 1271–1290.
- [24] Kim H., Tsai Tai-Peng, *Existence, uniqueness, regularity results for elliptic equations with drift terms in critical weak spaces*, *SIAM J. Math. Anal.*, (2020) Vol. 52, No. 2, pp. 1146–1191.
- [25] Kinderlehrer D., Stampacchia G., *An Introduction to Variational Inequalities and Their Applications*, vol. 88 of Pure and Applied Mathematics, Academic Press, Harcourt Brace Jovanovich, New York, NY, USA, 1980.
- [26] Lions J. L., *Quelques méthodes de résolution des problèmes aux limites non linéaires*. Dunod et Gauthier-Villars, Paris, (1969).
- [27] Leray J., Lions J. L., *Quelques résultats de Visik sur le problèmes elliptiques non linéaires par les méthodes de Minty-Browder*. *Bull. Soc. Math. France*, 93 (1965), 97–107.
- [28] Maz'ya V. G. , Verbitsky I. E., *Accreativity of the General Second order Linear Differential Operators*. *Acta Math. Sinica, English Series*, 35, 6, 199–217 (2019).
- [29] Moscariello G., *Existence and uniqueness for elliptic equations with lower-order terms*, *Adv. Calc. Var.*, 4, no. 4, (2011), 421–444.
- [30] O'Neil R., *Convolutions operators and  $L(p, q)$  spaces*. *Duke Math. J.* 30, (1963) 129–142.
- [31] Radice, T, Zecca, G., Existence and uniqueness for nonlinear elliptic equations with unbounded coefficients, *Ricerche di Matematica*, 63 (2), (2014) 355-367.
- [32] Stampacchia G., *Le problème de Dirichlet pour les équations elliptiques du second ordre à coefficients discontinus*. *Ann. Inst. Fourier* 15, 1, (1965), 189–258.

- [33] Zecca G., *Existence and uniqueness for nonlinear elliptic equations with lower-order terms*. Nonlinear Analysis, Theory, Methods and Applications **75**, no. 2, (2012), 899–912.
- [34] Zecca G., *An optimal control problem for some nonlinear elliptic equations with unbounded coefficients*. Discrete and Continuous Dynamical Systems, Series B, 24 (3) (2019) 1393–1409.

DIPARTIMENTO DI INGEGNERIA ELETTRICA E DELLE TECNOLOGIE DELL'INFORMAZIONE,  
UNIVERSITÀ DEGLI STUDI DI NAPOLI "FEDERICO II", VIA CLAUDIO 21, 80125  
NAPOLI, ITALY

*E-mail address:* luigreco@unina.it

DIPARTIMENTO DI MATEMATICA E APPLICAZIONI "R. CACCIOPPOLI", UNI-  
VERSITÀ DEGLI STUDI DI NAPOLI "FEDERICO II", VIA CINTIA, 80126 NAPOLI,  
ITALY

*E-mail address:* fernando.farroni@unina.it

*E-mail address:* gmoscari@unina.it

*E-mail address:* g.zecca@unina.it