

A Liouville theorem for an integral equation of the Ginzburg-Landau type

YUTIAN LEI AND XIN XU

Abstract In this paper, we are concerned with a Liouville-type result of the nonlinear integral equation

$$u(x) = \vec{l} + C_* \int_{\mathbb{R}^n} \frac{u(1 - |u|^2)}{|x - y|^{n-\alpha}} dy.$$

Here $u : \mathbb{R}^n \rightarrow \mathbb{R}^k$ is a bounded, uniformly continuous and differentiable function with $k \geq 1$ and $1 < \alpha < n$, $\vec{l} \in \mathbb{R}^k$ is a constant vector, and C_* is a real constant. If u is the finite energy solution, we prove that $|\vec{l}| \in \{0, 1\}$. Furthermore, we also give a Liouville type theorem (i.e., $u \equiv \vec{l}$).

Keywords: Ginzburg-Landau equation, Liouville theorem, Riesz potential

MSC2020: 45G05, 45E10, 35Q56, 35R11

1 Introduction

We first recall several Liouville theorems. If a harmonic function u is bounded on \mathbb{R}^n , then $u \equiv \text{Constant}$. When $\alpha \in (0, 2)$, u is a bounded function satisfying $(-\Delta)^{\frac{\alpha}{2}} u = 0$ on \mathbb{R}^n , then $u \equiv \text{Constant}$ (cf. [3], [15]).

In 1994, Brezis, Merle and Rivière [2] studied the quantization effects of the following equation

$$-\Delta u = u(1 - |u|^2) \quad \text{on } \mathbb{R}^2. \quad (1.1)$$

Here $u : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a vector valued function. It is the Euler-Lagrange equation of the Ginzburg-Landau energy

$$E_{GL}(u) = \frac{1}{2} \|\nabla u\|_{L^2(\mathbb{R}^2)}^2 + \frac{1}{4} \|1 - |u|^2\|_{L^2(\mathbb{R}^2)}^2.$$

In particular, they proved that the finite energy solution (i.e., u satisfies $\nabla u \in L^2(\mathbb{R}^2)$) is bounded (see also [7])

$$|u| \leq 1 \quad \text{on } \mathbb{R}^2.$$

Based on this result, they used the Pohozaev identity to obtain a Liouville type theorem for finite energy solutions

Proposition 1.1. *(Theorem 2 in [2]) Let $u : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a classical solution of (1.1). If $\nabla u \in L^2(\mathbb{R}^2)$, then either $u \in L^2(\mathbb{R}^2)$ which implies $u \equiv \vec{0}$, or $1 - |u|^2 \in L^1(\mathbb{R}^2)$ which implies $u \equiv \vec{C}$ with $|\vec{C}| = 1$.*

In addition, for the integral equation

$$u(x) = \int_{\mathbb{R}^n} \frac{u(1 - |u|^2)}{|x - y|^{n-\alpha}} dy, \quad (1.2)$$

there holds the following Liouville theorem.

Proposition 1.2. *(Theorem 1 in [8]) Assume that $u : \mathbb{R}^n \rightarrow \mathbb{R}^k$ is bounded and differentiable, and solves (1.2) with $k \geq 1$ and $\alpha \in (1, n/2)$. If $u \in L^2(\mathbb{R}^n)$, then $u(x) \equiv \vec{0}$.*

For the Lane-Emden type integral equations with the critical case, the finite energy solutions can be classified (cf. [5]). In this paper, we consider finite energy solutions of the integral equation involving the Riesz potential

$$u(x) = \vec{l} + C_* \int_{\mathbb{R}^n} \frac{u(1 - |u|^2)}{|x - y|^{n-\alpha}} dy, \quad (1.3)$$

where $u : \mathbb{R}^n \rightarrow \mathbb{R}^k$ with $k \geq 1$ and $0 < \alpha < n$, $\vec{l} \in \mathbb{R}^k$ is a constant vector, and $C_* \in \mathbb{R}$ is a constant.

Eq. (1.3) is associated with the fractional Ginzburg-Landau equation (cf. [11], [12] and [14])

$$(-\Delta)^{\frac{\alpha}{2}}u = (1 - |u|^2)u \quad \text{on } \mathbb{R}^n. \quad (1.4)$$

Here $u = (u_1, u_2, \dots, u_k) : \mathbb{R}^n \rightarrow \mathbb{R}^k$. Write $u_i^+ = \max\{u_i, 0\}$ and $u_i^- = -\min\{u_i, 0\}$. Then $u_i^+, u_i^- \geq 0$ and $u_i = u_i^+ - u_i^-$.

When $\alpha \geq 2$ is an even number, Theorem 3.21 in [6] (see also Theorem 2.4 in [4]) shows that if u solves (1.4), there exist two constants $l_i^+, l_i^- \in \mathbb{R}$ such that

$$\begin{aligned} u_i^+(x) &= l_i^+ + C_\alpha \int_{\mathbb{R}^n} \frac{u_i^+(y)(1 - |u(y)|^2)}{|x - y|^{n-\alpha}} dy, \\ u_i^-(x) &= l_i^- + C_\alpha \int_{\mathbb{R}^n} \frac{u_i^-(y)(1 - |u(y)|^2)}{|x - y|^{n-\alpha}} dy. \end{aligned} \quad (1.5)$$

Here $C_\alpha = \Gamma(\frac{n-\alpha}{2})[2^\alpha \pi^{n/2} \Gamma(\alpha/2)]^{-1}$. Denote $l_i^+ - l_i^-$ by l_i , then there holds

$$u_i(x) = l_i + C_\alpha \int_{\mathbb{R}^n} \frac{u_i(y)(1 - |u(y)|^2)}{|x - y|^{n-\alpha}} dy,$$

which implies that u also solves (1.3), where $\vec{l} = (l_1, l_2, \dots, l_k)$. For the Lane-Emden equation, Chen, Li and Ou obtained an analogous result (cf. [5]).

When $\alpha \in (0, 2)$, if u solves (1.4) and $|u| \leq 1$ on \mathbb{R}^n , then

$$(-\Delta)^{\frac{\alpha}{2}}u_i^+ = (1 - |u|^2)u_i^+ \quad \text{on } \mathbb{R}^n. \quad (1.6)$$

On the other hand, $C_\alpha|x|^{\alpha-n}$ is a fundamental solution of $(-\Delta)^{\frac{\alpha}{2}}u = 0$ (cf. Chapter 5 in [13]), i.e.,

$$(-\Delta)^{\frac{\alpha}{2}}(C_\alpha|x|^{\alpha-n}) = \delta_o,$$

where δ_o is the Dirac mass at the origin o . Therefore, on \mathbb{R}^n we have

$$\begin{aligned} &(-\Delta)^{\frac{\alpha}{2}}[C_\alpha|x|^{\alpha-n} * (u_i^+(1 - |u|^2))] \\ &= \delta_x * [u_i^+(1 - |u|^2)] = u_i^+(x)(1 - |u(x)|^2). \end{aligned}$$

Combining with (1.6) yields an α -harmonic equation

$$(-\Delta)^{\frac{\alpha}{2}}[u_i^+ - C_\alpha|x|^{\alpha-n} * (u_i^+(1 - |u|^2))] = 0 \quad \text{on } \mathbb{R}^n.$$

Since $u_i^+ \geq 0$ and u is bounded, we can see that the α -harmonic function

$$u_i^+ - C_\alpha|x|^{\alpha-n} * (u_i^+(1 - |u|^2))$$

has an upper bound. And hence it is a constant (cf. [3], [15]), which is denoted by l_i^+ . Thus, (1.5) is also true. By the same argument above, we also see that u solves (1.3).

In this paper, we expect to obtain the analogous results in Propositions 1.1 and 1.2 for the finite energy solutions of (1.3).

First, if the finite energy solution u is bounded, we determine the value of $|\vec{l}|$.

Theorem 1.1. *Assume that a uniformly continuous function $u : \mathbb{R}^n \rightarrow \mathbb{R}^k$ solves (1.3) with $k \geq 1$ and $n \geq 2$. If*

$$|u| \leq 1 \quad \text{on } \mathbb{R}^n; \tag{1.7}$$

and

$$\int_{\mathbb{R}^n} |u|^2(1 - |u|^2)dx < \infty. \tag{1.8}$$

Then, one of the following results holds true

- (Rt-i) $u \in L^2(\mathbb{R}^n)$ and $|\vec{l}| = 0$ if $\alpha \in (0, n/2)$;
- (Rt-ii) $1 - |u|^2 \in L^1(\mathbb{R}^n)$ and $|\vec{l}| = 1$ if $\alpha \in (0, n)$.

Next, we have a Liouville theorem for finite energy solutions.

Theorem 1.2. *(Liouville theorem) Under the same assumption of Theorem 1.1, if u is also differentiable, then*

- (Rt-iii) when (Rt-i) happens and $\alpha \in (1, n/2)$, we have $u \equiv \vec{0}$;
- (Rt-iv) when (Rt-ii) happens and $\alpha \in (1, n)$, we have $u \equiv \vec{l}$.

Finally, we give remarks on conditions (1.7) and (1.8).

Remark 1.1. The condition (1.7) shows the boundedness which is a necessary condition in the Liouville theorem. For (1.4) with $\alpha = 2$, (1.7) was proved firstly by Brezis (cf. [1], where the author used the Keller-Osserman theory via Kato's inequality), and independently by Ma (cf. [9], where the maximum principle was applied). When $\alpha \in (0, 2)$ and $1 - |u|^2 \in L^2(\mathbb{R}^n)$, Ma also proved (1.7) by the Kato inequality (cf. [10]).

Remark 1.2. Sometimes (1.8) is called a finite energy condition. For example, u is a finite energy (i.e., $\nabla u \in L^2(\mathbb{R}^n)$) solution of (1.4) with $\alpha = 2$. Multiplying (1.4) by u and integrating on $B_R := B_R(0)$ yield

$$\int_{B_R} |u|^2(1 - |u|^2)dx = \int_{B_R} |\nabla u|^2 dx - \int_{\partial B_R} u \partial_\nu u ds, \quad (1.9)$$

where ν is the unit outwards norm vector on ∂B_R . The Sobolev inequality implies $\nabla u \in L^2(\mathbb{R}^n) \Rightarrow u \in L^{2^*}(\mathbb{R}^n)$. Therefore,

$$R \int_{\partial B_R} (|u|^{2^*} + |\nabla u|^2) ds \rightarrow 0, \quad \text{when } R = R_j \rightarrow \infty.$$

Thus, by the Hölder inequality, when $R \rightarrow \infty$,

$$\begin{aligned} & \left| \int_{\partial B_R} u \partial_\nu u ds \right| \\ & \leq \left(R \int_{\partial B_R} |u|^{2^*} ds \right)^{\frac{1}{2^*}} \left(R \int_{\partial B_R} |\nabla u|^2 ds \right)^{\frac{1}{2}} |\partial B_R|^{\frac{1}{2} - \frac{1}{2^*}} R^{-\frac{1}{2} - \frac{1}{2^*}} \\ & \rightarrow 0. \end{aligned} \quad (1.10)$$

Inserting this into (1.9) we see that

$$\|\nabla u\|_{L^2(\mathbb{R}^n)}^2 = \int_{\mathbb{R}^n} |u|^2(1 - |u|^2)dx, \quad (1.11)$$

and hence (1.8) holds true.

On the contrary, if (1.8) holds, then $\nabla u \in L^2(\mathbb{R}^n)$. In fact, take ζ_R as the cut-off function satisfying $\zeta_R(x) = 1$ when $|x| \leq R/2$ and $\zeta_R(x) = 0$ when

$|x| \geq R$. Multiplying (1.4) by $u\zeta_R^2$ and integrating on B_R , we get

$$\int_{B_R} \zeta_R^2 |\nabla u|^2 dx = -2 \int_{B_R} (u \nabla u) \cdot (\zeta_R \nabla \zeta_R) dx + \int_{B_R} \zeta_R^2 |u|^2 (1 - |u|^2) dx.$$

Using the Cauchy-Schwarz inequality and (1.8), we have

$$\int_{B_R} \zeta_R^2 |\nabla u|^2 dx \leq C,$$

where $C > 0$ is independent of R . Letting $R \rightarrow \infty$ yields $\nabla u \in L^2(\mathbb{R}^n)$. Moreover, by the same argument above, we also have (1.10), and hence (1.11) still holds true.

2 Proof of Theorem 1.1

Theorem 2.1. *Under the same assumptions of Theorem 1.1, we have*

- (i) either $u \in L^2(\mathbb{R}^n)$ and $\lim_{|x| \rightarrow \infty} |u(x)| = 0$;
- (ii) or $1 - |u|^2 \in L^1(\mathbb{R}^n)$ and $\lim_{|x| \rightarrow \infty} |u(x)| = 1$.

Proof. Here an idea in Section 3.2 of [2] was used.

Set $S_* = \{x \in \mathbb{R}^n; \frac{1}{4} \leq |u| \leq \frac{3}{4}\}$. We claim that there exists suitably large $R_0 > 0$ such that $S_* \subset B_{R_0}(0)$.

Otherwise, we can find a sequence $\{x_j\} \subset S_*$ satisfies $\lim_{j \rightarrow \infty} x_j = \infty$. Since u is uniformly continuous, there exists $\eta \in (0, 1)$ such that

$$\frac{1}{8} \leq |u(x)| \leq \frac{7}{8}, \quad \text{when } |x - x_j| < \eta, \quad \forall j.$$

Choose a subsequence of $\{x_j\}$ (still denoted by itself) satisfying $|x_i - x_j| > 3\eta$ for $i \neq j$. Therefore,

$$\int_{\cup_j B(x_j, \eta)} |u|^2 (1 - |u|^2) dx \geq C |\cup_j B(x_j, \eta)| = \infty,$$

which contradicts with (1.8). The claim is proved.

Since u is uniformly continuous and $\mathbb{R}^n \setminus B_{R_0}(0)$ is connected, either $|u| \leq 1/4$ or $|u| \geq 3/4$ holds true on $\mathbb{R}^n \setminus B_{R_0}(0)$.

When $|u| \leq 1/4$ on $\mathbb{R}^n \setminus B_{R_0}(0)$, it is led to $1 - |u|^2 \geq 15/16$ on $\mathbb{R}^n \setminus B_{R_0}(0)$.

Thus,

$$\begin{aligned} \int_{\mathbb{R}^n} |u|^2 dx &= \int_{B_{R_0}(0)} |u|^2 dx + \int_{\mathbb{R}^n \setminus B_{R_0}(0)} |u|^2 dx \\ &\leq |B_{R_0}| + \frac{16}{15} \int_{\mathbb{R}^n \setminus B_{R_0}(0)} |u|^2 (1 - |u|^2) dx < \infty, \end{aligned}$$

by virtue of (1.7) and (1.8). Namely, $u \in L^2(\mathbb{R}^n)$.

Now, we claim that

$$\lim_{|x| \rightarrow \infty} |u(x)| = 0. \quad (2.1)$$

Otherwise, we can find $\epsilon_0 > 0$ and $x_j \rightarrow \infty$ such that

$$|u(x_j)| \geq 2\epsilon_0.$$

Since u is uniformly continuous, there exists $\eta > 0$, such that

$$|u(x) - u(y)| < \epsilon_0, \quad \text{when } |x - y| < \eta.$$

Therefore, for $x \in B_\eta(x_j)$, $|u(x)| > |u(x_j)| - \epsilon_0 \geq \epsilon_0$. Thus,

$$\int_{B_\eta(x_j)} |u(x)|^2 dx > \epsilon_0^2 |B_\eta|.$$

This contradicts with $u \in L^2(\mathbb{R}^n)$.

When $|u| \geq 3/4$ on $\mathbb{R}^n \setminus B_{R_0}(0)$, by the same argument above, we can see firstly

$$\int_{\mathbb{R}^n} (1 - |u|^2) dx < \infty$$

in view of (1.7) and (1.8). This is $1 - |u|^2 \in L^1(\mathbb{R}^n)$. Second, since u is uniformly continuous, there also holds

$$\lim_{|x| \rightarrow \infty} |u(x)| = 1. \quad (2.2)$$

The proof of Theorem 2.1 is complete. \square

Theorem 2.2. *Under the same assumptions of Theorem 1.1, we have*

(i) *when $u \in L^2(\mathbb{R}^n)$ and $\alpha \in (0, n/2)$, then $|\vec{l}| = 0$;*

(ii) *when $1 - |u|^2 \in L^1(\mathbb{R}^n)$ and $\alpha \in (0, n)$, then $|\vec{l}| = 1$.*

Proof. Set

$$v(x) = \frac{1}{n - \alpha} \int_{\mathbb{R}^n} \frac{|u(y)|(1 - |u(y)|^2)}{|x - y|^{n-\alpha}} dy. \quad (2.3)$$

By exchanging the integral variants, we can get

$$v(x) = \int_0^\infty \frac{\int_{B_t(x)} |u(y)|[1 - |u(y)|^2] dy dt}{t^{n-\alpha}} \frac{1}{t}.$$

Once there holds

$$\lim_{|x| \rightarrow \infty} v(x) = 0, \quad (2.4)$$

from (1.3) it follows that

$$\lim_{|x| \rightarrow \infty} u(x) = \vec{l}.$$

By (2.1) and (2.2), the proof of Theorem 2.2 is complete.

Proof of (2.4). Take $x_0 \in \mathbb{R}^n$. By (1.7), we know that $\forall \varepsilon > 0$, there exists $\delta \in (0, 1)$ such that

$$\int_0^\delta \frac{\int_{B_t(x_0)} |u(z)|[1 - |u(z)|^2] dz dt}{t^{n-\alpha}} \frac{1}{t} \leq C \int_0^\delta t^\alpha \frac{dt}{t} < \varepsilon.$$

When $|x - x_0| < \delta$, $B_t(x_0) \subset B_{t+\delta}(x)$. Therefore,

$$\begin{aligned} & \int_\delta^\infty \frac{\int_{B_t(x_0)} |u(z)|[1 - |u(z)|^2] dz dt}{t^{n-\alpha}} \frac{1}{t} \\ & \leq C \int_\delta^\infty \frac{\int_{B_{t+\delta}(x)} |u(z)|[1 - |u(z)|^2] dz}{(t + \delta)^{n-\alpha}} \left(\frac{t + \delta}{t}\right)^{n-\alpha+1} \frac{d(t + \delta)}{t + \delta} \\ & \leq C 2^{n-\alpha+1} \int_0^\infty \frac{\int_{B_t(x)} |u(z)|[1 - |u(z)|^2] dz dt}{t^{n-\alpha}} \frac{1}{t} \\ & \leq C v(x). \end{aligned}$$

Combining two estimates above, we get

$$v(x_0) < \varepsilon + Cv(x), \quad \text{for } |x - x_0| < \delta.$$

Thus, for any $s > 1$,

$$\begin{aligned} v^s(x_0) &= |B_\delta(x_0)|^{-1} \int_{B_\delta(x_0)} v^s(x) dx \\ &\leq C\varepsilon^s + C|B_\delta(x_0)|^{-1} \int_{B_\delta(x_0)} v^s(x) dx. \end{aligned} \tag{2.5}$$

We claim that

$$v \in L^s(\mathbb{R}^n) \text{ for some } s > 1. \tag{2.6}$$

In fact, using the Hardy-Littlewood-Sobolev inequality to (2.3) yields

$$\|v\|_{L^s(\mathbb{R}^n)} \leq C\|u(1 - |u|^2)\|_{L^{\frac{ns}{n+s\alpha}}(\mathbb{R}^n)}.$$

When $u \in L^2(\mathbb{R}^n)$, we can choose $s = \frac{2n}{n-2\alpha}$ (in view of $0 < \alpha < n/2$). When $1 - |u|^2 \in L^1(\mathbb{R}^n)$, we can choose some $s > \frac{n}{n-\alpha}$. Thus, (2.6) is easy to prove by (1.7).

In view of (2.6), when $|x_0| \rightarrow \infty$,

$$\int_{B_\delta(x_0)} v^s(x) dx \rightarrow 0.$$

Inserting this result into (2.5), we have

$$\lim_{|x_0| \rightarrow \infty} v^s(x_0) = 0.$$

This result means that (2.4) holds. □

3 Proof of Theorem 1.2

Proof. Proof of (Rt-iii). When (Rt-i) holds true, u solves

$$u(x) = C_* \int_{\mathbb{R}^n} \frac{u(1 - |u|^2)}{|x - y|^{n-\alpha}} dy.$$

By the same proof of Proposition 1.2, we also obtain $u \equiv \vec{0}$.

Proof of (Rt-iv).

For convenience, we denote $B_R(0)$ by B_R here.

Step 1. We claim that the improper integral

$$\int_{\mathbb{R}^n} \frac{z \cdot \nabla[u(z)(1 - |u(z)|^2)]}{|x - z|^{n-\alpha}} dz \quad (3.1)$$

is convergent at each $x \in \mathbb{R}^n$.

In fact, since $1 - |u|^2 \in L^1(\mathbb{R}^n)$, we can find $R = R_j \rightarrow \infty$ such that

$$R \int_{\partial B_R} (1 - |u(z)|^2) ds \rightarrow 0. \quad (3.2)$$

By (1.7), we obtain that for sufficiently large R , there holds

$$R \left| \int_{\partial B_R} \frac{u(z)(1 - |u(z)|^2)}{|x - z|^{n-\alpha}} ds \right| \leq CR^{1-n+\alpha} \int_{\partial B_R} (1 - |u(z)|^2) ds.$$

Letting $R = R_j \rightarrow \infty$ and using (3.2) we get

$$R \int_{\partial B_R} \frac{u(z)(1 - |u(z)|^2)}{|x - z|^{n-\alpha}} ds \rightarrow 0 \quad (3.3)$$

when $R = R_j \rightarrow \infty$.

Next, we claim that the improper integral

$$I(\mathbb{R}^n) := \int_{\mathbb{R}^n} \frac{u(z)(1 - |u(z)|^2)(x - z) \cdot z}{|x - z|^{n-\alpha+2}} dz \quad (3.4)$$

absolutely converges for each $x \in \mathbb{R}^n$.

In fact, we observe that the defect points of $I(\mathbb{R}^n)$ are x and ∞ . When z is near ∞ , we have

$$\begin{aligned} |I(\mathbb{R}^n \setminus B_r)| &\leq C \int_{\mathbb{R}^n \setminus B_r} \frac{1 - |u(z)|^2}{|x - z|^{n-\alpha}} dz \\ &\leq C \left(\int_{\mathbb{R}^n} (1 - |u|^2)^s dz \right)^{\frac{1}{s}} \left(\int_r^\infty \rho^{n - \frac{s}{s-1}(n-\alpha)} \frac{d\rho}{\rho} \right)^{1 - \frac{1}{s}}. \end{aligned}$$

Here $s \in (1, n/\alpha)$. In view of $1 - |u|^2 \in L^1(\mathbb{R}^n)$ and (1.7), we get $1 - |u|^2 \in L^s(\mathbb{R}^n)$ for all $s \geq 1$. Therefore,

$$|I(\mathbb{R}^n \setminus B_r)| < \infty. \quad (3.5)$$

When z is near x , by (1.7) and $\alpha > 1$,

$$|I(B_\delta(x))| \leq C \int_{B_\delta(x)} \frac{dz}{|x - z|^{n-\alpha+1}} \leq C \int_0^\delta \rho^{\alpha-1} \frac{d\rho}{\rho} < \infty.$$

Combining this with (3.5), we prove that (3.4) is absolutely convergent.

Finally we prove that (3.1) is convergent. Integrating by parts yields

$$\begin{aligned} & \int_{B_R} \frac{z \cdot \nabla[u(z)(1 - |u(z)|^2)]}{|x - z|^{n-\alpha}} dz \\ &= R \int_{\partial B_R} \frac{u(z)(1 - |u(z)|^2)}{|x - z|^{n-\alpha}} ds \\ & \quad - n \int_{B_R} \frac{u(z)(1 - |u(z)|^2)}{|x - z|^{n-\alpha}} dz \\ & \quad - (n - \alpha) \int_{B_R} \frac{u(z)(1 - |u(z)|^2)(x - z) \cdot z}{|x - z|^{n-\alpha+2}} dz. \end{aligned} \quad (3.6)$$

Letting $R = R_j \rightarrow \infty$ in (3.6) and using (1.3) and (3.3), we can see that

$$\int_{\mathbb{R}^n} \frac{z \cdot \nabla[u(z)(1 - |u(z)|^2)]}{|x - z|^{n-\alpha}} dz = -\frac{n}{C_*} \left(u(x) - \vec{l} \right) + (\alpha - n)I(\mathbb{R}^n),$$

and hence it is convergent at each $x \in \mathbb{R}^n$.

Step 2. Proof of (Rt-iv).

For any $\lambda > 0$, from (1.3) it follows

$$u(\lambda x) = \vec{l} + C_* \lambda^\alpha \int_{\mathbb{R}^n} \frac{u(\lambda z)(1 - |u(\lambda z)|^2)}{|x - z|^{n-\alpha}} dz.$$

Differentiating both sides with respect to λ yields

$$\begin{aligned} & x \cdot \nabla u(\lambda x) \\ &= C_* \alpha \lambda^{\alpha-1} \int_{\mathbb{R}^n} \frac{u(\lambda z)(1 - |u(\lambda z)|^2)}{|x - z|^{n-\alpha}} dz \\ & \quad + C_* \lambda^\alpha \int_{\mathbb{R}^n} \frac{(z \cdot \nabla u(\lambda z))(1 - |u(\lambda z)|^2) + u(\lambda z)[-2u(\lambda z)(z \cdot \nabla u(\lambda z))]}{|x - z|^{n-\alpha}} dz. \end{aligned}$$

Letting $\lambda = 1$ yields

$$x \cdot \nabla u(x) = \alpha \left(u(x) - \vec{l} \right) + C_* \int_{\mathbb{R}^n} \frac{z \cdot \nabla [u(z)(1 - |u(z)|^2)]}{|x - z|^{n-\alpha}} dz. \quad (3.7)$$

Integrating by parts, we get

$$\begin{aligned} & \int_{B_R} u(1 - |u|^2)(x \cdot \nabla u) dx \\ &= -\frac{1}{4} \int_{B_R} x \cdot \nabla [(1 - |u|^2)^2] dx \\ &= \frac{n}{4} \int_{B_R} (1 - |u|^2)^2 dx - \frac{R}{4} \int_{\partial B_R} (1 - |u|^2)^2 ds. \end{aligned}$$

Since (1.7) and (3.2), it follows that

$$R \int_{\partial B_R} (1 - |u|^2)^2 ds \rightarrow 0 \quad (3.8)$$

for some $R = R_j \rightarrow \infty$. Thus, by virtue of (1.7) and $1 - |u|^2 \in L^1(\mathbb{R}^n)$,

$$\int_{\mathbb{R}^n} u(x)(1 - |u(x)|^2)(x \cdot \nabla u(x)) dx = \frac{n}{4} \int_{\mathbb{R}^n} (1 - |u(x)|^2)^2 dx < \infty. \quad (3.9)$$

From (1.7) and $1 - |u|^2 \in L^1(\mathbb{R}^n)$, it also follows that

$$\left| \int_{\mathbb{R}^n} [u(x) \cdot (u(x) - \vec{l})](1 - |u(x)|^2) dx \right| < \infty.$$

Multiply (3.7) by $u(x)(1 - |u(x)|^2)$ and integrate over B_R . Letting $R = R_j \rightarrow \infty$, from the result above and (3.9), we get

$$\left| \int_{\mathbb{R}^n} u(x)(1 - |u(x)|^2) \int_{\mathbb{R}^n} \frac{z \cdot \nabla [u(z)(1 - |u(z)|^2)]}{|x - z|^{n-\alpha}} dz dx \right| < \infty,$$

and

$$\begin{aligned} & \int_{\mathbb{R}^n} u(x)(1 - |u(x)|^2)(x \cdot \nabla u(x)) dx \\ & - \alpha \int_{\mathbb{R}^n} [u(x) \cdot (u(x) - \vec{l})](1 - |u(x)|^2) dx \\ &= C_* \int_{\mathbb{R}^n} u(x)(1 - |u(x)|^2) \int_{\mathbb{R}^n} \frac{z \cdot \nabla [u(z)(1 - |u(z)|^2)]}{|x - z|^{n-\alpha}} dz dx. \end{aligned} \quad (3.10)$$

We use the Fubini theorem and (1.3) to handle the term of the right hand side. Thus,

$$\begin{aligned}
& C_* \int_{\mathbb{R}^n} u(x)(1 - |u(x)|^2) \int_{\mathbb{R}^n} \frac{z \cdot \nabla[u(1 - |u|^2)]}{|x - z|^{n-\alpha}} dz dx \\
&= C_* \int_{\mathbb{R}^n} z \cdot \nabla[u(z)(1 - |u(z)|^2)] \int_{\mathbb{R}^n} \frac{u(x)(1 - |u(x)|^2)}{|x - z|^{n-\alpha}} dx dz \\
&= \int_{\mathbb{R}^n} (x \cdot \nabla[u(x)(1 - |u(x)|^2)])(u(x) - \vec{l}) dx \tag{3.11} \\
&= \int_{\mathbb{R}^n} (u(x) - \vec{l})(1 - |u(x)|^2)(x \cdot \nabla u(x)) dx \\
&\quad + \int_{\mathbb{R}^n} u(x)(u(x) - \vec{l})[x \cdot \nabla(1 - |u(x)|^2)] dx.
\end{aligned}$$

Inserting this result into (3.10), we have

$$\begin{aligned}
0 &= \alpha \int_{\mathbb{R}^n} u(x)(u(x) - \vec{l})(1 - |u(x)|^2) dx \\
&\quad - \int_{\mathbb{R}^n} \vec{l} (1 - |u(x)|^2)(x \cdot \nabla u(x)) dx \\
&\quad - \int_{\mathbb{R}^n} u(x)(u(x) - \vec{l})[x \cdot \nabla(|u(x)|^2 - 1)] dx \\
&:= I + II + III.
\end{aligned}$$

We deal with the first and the third terms of the right hand side.

Noting $|\vec{l}| = 1$, we have

$$\begin{aligned}
I &= \alpha \int_{\mathbb{R}^n} (u + \vec{l})(u - \vec{l})(1 - |u|^2) dx \\
&\quad - \alpha \int_{\mathbb{R}^n} \vec{l} (u - \vec{l})(1 - |u|^2) dx \\
&= -\alpha \int_{\mathbb{R}^n} (|u|^2 - 1)^2 dx + \alpha \int_{\mathbb{R}^n} (1 - |u|^2) dx \\
&\quad - \alpha \int_{\mathbb{R}^n} \vec{l} u (1 - |u|^2) dx.
\end{aligned}$$

Next, integrating by parts, we obtain

$$\begin{aligned}
III &= - \int_{\mathbb{R}^n} (u + \vec{l})(u - \vec{l})[x \cdot \nabla(|u|^2 - 1)]dx \\
&\quad + \int_{\mathbb{R}^n} \vec{l}(u - \vec{l})[x \cdot \nabla(|u|^2 - 1)]dx \\
&= - \int_{\mathbb{R}^n} (|u|^2 - 1)[x \cdot \nabla(|u|^2 - 1)]dx \\
&\quad - \vec{l} \int_{\mathbb{R}^n} (|u|^2 - 1)\nabla \cdot [x(u - \vec{l})]dx \\
&= \frac{n}{2} \int_{\mathbb{R}^n} (|u|^2 - 1)^2 dx + \vec{l} \int_{\mathbb{R}^n} (1 - |u|^2)(x \cdot \nabla u)dx \\
&\quad + n \vec{l} \int_{\mathbb{R}^n} (u - \vec{l})(1 - |u|^2)dx.
\end{aligned}$$

Substituting these results into $I + II + III = 0$, we get

$$\begin{aligned}
&(\alpha - \frac{n}{2}) \int_{\mathbb{R}^n} (|u|^2 - 1)^2 dx \\
&= \int_{\mathbb{R}^n} (-\alpha \vec{l} u + \alpha + n \vec{l} u - n)(1 - |u|^2)dx \\
&= \int_{\mathbb{R}^n} (n - \alpha)(\vec{l} u - 1)(1 - |u|^2)dx \\
&\leq (n - \alpha) \int_{\mathbb{R}^n} (|u| - 1)(1 - |u|^2)dx.
\end{aligned}$$

In view of $|u| - 1 \leq \frac{1}{2}(|u|^2 - 1)$, it follows that

$$(\alpha - \frac{n}{2}) \int_{\mathbb{R}^n} (|u|^2 - 1)^2 dx \leq \frac{n - \alpha}{2} \int_{\mathbb{R}^n} (|u|^2 - 1)(1 - |u|^2)dx,$$

which implies $|u| \equiv 1$ a.e. on \mathbb{R}^n . Inserting this into (1.3), we see that $u \equiv \vec{l}$ and hence (Rt-iv) is proved. \square

Acknowledgements. This research was supported by NNSF (11871278) of China.

References

- [1] H. Brezis, *Comments on two notes by L. Ma and X. Xu*, C. R. Math. Acad. Sci. Paris, **349** (2011), 269–271.
- [2] H. Brezis, F. Merle, T. Rivière, *Quantization effects for $-\Delta u = u(1 - |u|^2)$ in \mathbb{R}^2* , Arch. Rational Mech. Anal., **126** (1994), 35–58.
- [3] K. Bogdan, T. Kulczycki, A. Nowak, *Gradient estimates for harmonic and q -harmonic functions of symmetric stable processes*, Illinois J. Math., **46** (2002) 541-556.
- [4] G. Caristi, L. D’Ambrosio, E. Mitidieri, *Representation formulae for solutions to some classes of higher order systems and related Liouville theorems*, Milan J. Math., **76** (2008), 27–67.
- [5] W. Chen, C. Li, B. Ou, *Classification of solutions for an integral equation*, Comm. Pure Appl. Math., **59** (2006), 330–343.
- [6] W. K. Hayman, P. B. Kennedy, *Subharmonic functions*, Vol. I, Academic Press, London, New York, San Francisco, 1976.
- [7] M. Hervé, R. M. Hervé, *Quelques propriétés des solutions de l’équation de Ginzburg-Landau sur un ouvert de \mathbb{R}^2* , Potential Anal., **5** (1996), 591–609.
- [8] Y. Li, Q. Chen, Y. Lei, *A Liouville theorem of the fractional Ginzburg-Landau equation*, (2020) arXiv:2006.12664. C. R. Math. Acad. Sci. Paris, to appear.
- [9] L. Ma, *Liouville type theorem and uniform bound for the Lichnerowicz equation and the Ginzburg-Landau equation*, C. R. Math. Acad. Sci. Paris, **348** (2010), 993–996.

- [10] L. Ma, *Boundedness of solutions to Ginzburg-Landau fractional Laplacian equation*, Inter. J. Math., **27** (2016), No.5, 1650048.
- [11] V. Millot, Y. Sire, *On a fractional Ginzburg-Landau equation and 1/2-harmonic maps into spheres*, Arch. Rational Mech. Anal., **215** (2015), 125–210.
- [12] A.V. Milovanov, J. Juul Rasmussen, *Fractional generalization of the Ginzburg-Landau equation: an unconventional approach to critical phenomena in complex media*, Phys. Lett. A, **337** (2005), 75–80.
- [13] E. M. Stein, *Singular integrals and differentiability properties of function*, Princeton Math. Series, Vol. 30. Princeton University Press, 1970.
- [14] V. Tarasov, G. Zaslavsky, *Fractional Ginzburg-Landau equation for fractal media*, Phys. A, **354** (2005), 249–261.
- [15] R. Zhuo, W. Chen, X. Cui, Z. Yuan. *Symmetry and non-existence of solutions for a nonlinear system involving the fractional Laplacian*, Discrete Contin. Dyn. Syst., **36** (2016), 1125–1141.

Yutian Lei

Jiangsu Key Laboratory for NSLSCS, School of Mathematical Sciences,
Nanjing Normal University, Nanjing, 210023, China

Xin Xu

Depart. of Math., Southern University of Science and Technology,
Shenzhen, 518055, China

Faculty of Science and Technology, University of Macau, Macau, China