

Full vertex algebra and bootstrap - consistency of four point functions in 2d CFT

Yuto Moriwaki *

Kavli Institute for the Physics and Mathematics of the Universe, Chiba, Japan

Abstract. In physics, it is believed that the consistency of two dimensional conformal field theory follows from the bootstrap equation. In this paper, we introduce the notion of a full vertex algebra by analyzing the bootstrap equation, which is a “real analytic” generalization of a $\mathbb{Z}$ -graded vertex algebra. We also give a mathematical formulation of the consistency of four point correlation functions in two dimensional conformal field theory and prove it for a full vertex algebra with additional assumptions on the conformal symmetry. In particular, we show that the bootstrap equation together with the conformal symmetry implies the consistency of four point correlation functions. As an application, a deformable family of full vertex algebras parametrized by the Grassmanian is constructed, which appears in the toroidal compactification of string theory. This give us examples satisfying the above assumptions.

INTRODUCTION

Quantum field theory has a long history in physics (see e.g. [S]) and has attracted mathematicians in the past several decades due to its unexpected mathematical consequences or conjectures, for example [W, Ma]. One aim of quantum field theory is to calculate n *point correlation functions*, that is, the vacuum expectation value of an interaction of n particles. An interaction of n particles decomposes into subsequent interactions of three particles. Thus, an n point correlation function can be expressed in terms of three point correlation functions, depending on a choice of decompositions. A quantum field theory requires that the resulting n point correlation functions are *independent* of the choice of decompositions. This principle is known as the *consistency of quantum field theory*. Although it is known to be difficult to construct mathematically rigorous quantum field theories, surprisingly many such examples, especially *conformal field theories*, (i.e., quantum field theories with “conformal symmetry”), have been found in two dimension, see [FMS].

In (not necessarily two dimensional) conformal field theories, it is believed in physics, that the whole consistency of n point correlation functions follows from the *bootstrap equations (or hypothesis)*, which are distinguished consistencies of four point correlation functions [FGG, Poly2]. This hypothesis was used successively by Belavin, Polyakov

*email: moriwaki@ms.u-tokyo.ac.jp

and Zamolodchikov in [BPZ] where the modern study of two dimensional conformal field theories was initiated.

In two dimensional conformal field theory, fields are operator valued real analytic functions. It is noteworthy that the subalgebra consisting of holomorphic fields satisfies a purely algebraic axiom, which was introduced by Borchers [B], see also [G]. It is called a vertex algebra or a vertex operator algebra [FLM] and has been studied intensively by many authors, see e.g., [LL, FHL, FB, K]. We note that in the language of a vertex algebra, three point correlation functions correspond to the vertex operator and the bootstrap equations essentially correspond to the Borchers identity, which is an axiom of vertex algebras. In contrast, the axiom of the (non-holomorphic) whole algebra of a conformal field theory needs analytic properties and seems impossible to describe in an algebraic way.

Moore and Seiberg constructed a conformal field theory as an extension of holomorphic and anti-holomorphic vertex operator algebras by their modules [MS1, MS2]. The bootstrap equations in this case are translated as a monodromy invariant property of the four point correlation functions. In the physics literature, this property was reformulated later by Fuchs, Runkel and Schweigert in [FRS], which says that the algebra describing the conformal field theory is a Frobenius algebra object in a braided tensor category constructed from holomorphic and anti-holomorphic vertex operator algebras.

A mathematical approach in this direction is due to Huang and Kong [HK] based on the abstract representation theory of vertex algebras developed by Huang and Lepowsky in a series of papers [HL1, HL2, HL3, H1, H2]. One of the prominent results is obtained by Huang [H3, H4], which states that the representation category of a regular vertex operator algebra (of strong CFT type) inherits a modular tensor category structure. Note that a vertex operator algebra is called regular if every (weak) module is completely reducible.

Then, Huang and Kong [HK] introduced a notion of full field algebras, which is a mathematical axiomatization of the algebras describing two dimensional conformal field theories. Their definition is based upon a part of the consistencies of n point correlation functions. In [HK], they constructed conformal field theories, called *diagonal theories* in physics, as finite module extensions of the tensor product of regular vertex operator algebras in two variables “ $(z, \bar{z})$ ”.

In this paper, we study an axiom of conformal field theory on $\mathbb{CP}^1$ and introduce a notion of *full vertex algebras* starting with the bootstrap equations, which are expected to be sufficient to derive the whole consistency of the theory.

The definition of full vertex algebras is independent of the theory of vertex algebras, which is an essential building block of the approach by Huang and Kong. Rather, the existence of vertex subalgebras as holomorphic and anti-holomorphic part of the algebra is naturally derived from our definition as expected in physics, see below for details.

The notion of full vertex algebras essentially appears in [HK]. There they showed that as long as a full field algebra is an extension of a tensor product of regular vertex

algebras, the notions of full field algebras and full vertex algebras are equivalent [HK, Theorem 2.11].

Before going into the details, let us summarize the key ingredients of the paper:

- to introduce a reasonable space Cor_4 of real analytic functions serving as the four point correlation functions (Section 0.1),
- to define and analyze parenthesized correlation functions S_A (Section 0.2),
- to formulate and prove the consistency of four point functions in the language of full vertex algebras (Section 0.3).

As a byproduct, we give an example of a deformable family of full vertex algebras corresponding to the irrational conformal field theory appearing in the toroidal compactification of string theory [Polc1]. Mathematically, this is a generalization of a lattice vertex algebra [B, FLM]. We remark that a deformation of a theory is one of the most important ingredient of quantum field theory and conformal field theory [IZ]. This kind of deformations will be studied in more general setting in our next paper [Mo2]. The author hopes that our approach will be beneficial in the future investigation on the conformal field theories in higher dimensions.

0.1. Space of four point correlation function. In order to formulate the consistency of conformal field theory on the projective space $\mathbb{CP}^1$, we need to fix a space of functions “consisting of four point correlation functions”. Let $X_4 = \{(z_1, \dots, z_4) \in (\mathbb{CP}^1)^4 \mid z_i \neq z_j \text{ for any } i \neq j\}$ be a space of ordered configurations of four points in $\mathbb{CP}^1$. Then, a four point correlation function is, roughly, a $\mathbb{C}$ -valued real analytic function on X_4 with possible singularities along the diagonals, $\{(z_1, \dots, z_4) \in (\mathbb{CP}^1)^4 \mid z_i = z_j \text{ for } 1 \leq i < j \leq 4\}$. Recall that the automorphism group of $\mathbb{CP}^1$ is $\text{PSL}_2\mathbb{C}$, whose action is called the linear fractional transformation and which is the global conformal symmetry in two dimension. Since the real analytic function

$$\xi : X_4 \rightarrow \mathbb{CP}^1 \setminus \{0, 1, \infty\}, (z_1, z_2, z_3, z_4) \mapsto \frac{(z_1 - z_2)(z_3 - z_4)}{(z_1 - z_3)(z_2 - z_4)}$$

is invariant under the diagonal action of $\text{PSL}_2\mathbb{C}$ on $X_4 \subset (\mathbb{CP}^1)^4$, it gives a homeomorphism from the coset space $\text{PSL}_2\mathbb{C} \backslash X_4$ to $\mathbb{CP}^1 \setminus \{0, 1, \infty\}$. Any four point function of quasi-primary states in 2d conformal field theory possesses the global conformal symmetry. Thus, by the homeomorphism, a four point function can be regarded as a real analytic function on $\mathbb{CP}^1 \setminus \{0, 1, \infty\}$ with possible singularities at $\{0, 1, \infty\}$, which we call *conformal singularities*. We remark that $|z|^r$ is not a real analytic function at 0, where $|z| = \sqrt{z\bar{z}}$, the square of the absolute value, and $r \in \mathbb{R}$. A function with a conformal singularity at 0 has an expansion

$$(0.1) \quad \sum_{r,s \in \mathbb{R}} a_{r,s} z^r \bar{z}^s,$$

where $a_{r,s} \in \mathbb{C}$ and $a_{r,s} = 0$ if $r - s \notin \mathbb{Z}$. This series is absolutely convergent in an annulus $0 < |z| < R$ (for the precise definition, see Section 1.1). By the assumption, $a_{r,s} z^r \bar{z}^s = a_{r,s} z^{r-s} |z|^s$ is a single-valued function around 0. Denote by $F_{0,1,\infty}$ the space of real

analytic functions on $\mathbb{CP}^1 \setminus \{0, 1, \infty\}$ with possible conformal singularities at $\{0, 1, \infty\}$. The space $F_{0,1,\infty}$ contains a monodromy invariant linear combination of solutions of differential equations with possible regular singularities at $\{0, 1, \infty\}$. For example, a combination of hypergeometric functions,

$$(0.2) \quad f_{\text{Ising}}(z) = |1 - \sqrt{1-z}|^{1/2} + |1 + \sqrt{1-z}|^{1/2},$$

has conformal singularities at $\{0, 1, \infty\}$, which appears as a four point function of the two dimensional Ising model. The expansion of f_{Ising} at $z = 0$ is

$$2 + |z|^{1/2}/2 - z/4 - \bar{z}/4 + |z|^{1/2}(z + \bar{z})/16 + z\bar{z}/32 - 5z^2/64 - 5\bar{z}^2/64 + \dots$$

We introduce a space Cor_4 spanned by real analytic functions on X_4 of the form:

$$(0.3) \quad \Pi_{1 \leq i < j \leq 4} (z_i - z_j)^{\alpha_{ij}} (\bar{z}_i - \bar{z}_j)^{\beta_{ij}} f \circ \xi(z_1, z_2, z_3, z_4),$$

where $f \in F_{0,1,\infty}$ and $\alpha_{ij}, \beta_{ij} \in \mathbb{R}$ such that $\alpha_{ij} - \beta_{ij} \in \mathbb{Z}$ for any $1 \leq i < j \leq 4$. Here, we allow these functions to have singularities around $\{z_i = \infty\}_{i=1,2,3,4}$. By the condition $\alpha_{ij} - \beta_{ij} \in \mathbb{Z}$, we have

$$(z_i - z_j)^{\alpha_{ij}} (\bar{z}_i - \bar{z}_j)^{\beta_{ij}} = |z_i - z_j|^{\alpha_{ij}} (\bar{z}_i - \bar{z}_j)^{\beta_{ij} - \alpha_{ij}},$$

which implies that $\Pi_{1 \leq i < j \leq 4} (z_i - z_j)^{\alpha_{ij}} (\bar{z}_i - \bar{z}_j)^{\beta_{ij}}$ is a single valued real analytic function on X_4 . Physically, a four point correlation function of quasi-primary states in compact conformal field theory in two dimensions is of the form (0.3). Thus, we call Cor_4 a space of four point correlation functions. We remark that non-compact conformal field theories, e.g., the Liouville field theory and non-compact WZW conformal field theory are excluded from our consideration, where we need to consider some measure on the space of irreducible representations and integral over there.

0.2. Consistency and parenthesized correlation functions. Roughly speaking, a conformal field theory on $\mathbb{CP}^1$, considered in this paper, consists of an $\mathbb{R}^2$ -graded $\mathbb{C}$ -vector space $F = \bigoplus_{h, \bar{h} \in \mathbb{R}} F_{h, \bar{h}}$ with a distinguished vector $\mathbf{1} \in F$, a symmetric bilinear form $(-, -) : F \times F \rightarrow \mathbb{C}$, which is normalized as $(\mathbf{1}, \mathbf{1}) = 1$, and a linear map

$$Y(-, \underline{z}) : F \rightarrow \text{End}(F)[[z^{\pm}, \bar{z}^{\pm}, |z|^{\mathbb{R}}]], \quad a \mapsto Y(a, \underline{z}) = \sum_{r, s \in \mathbb{R}} a(r, s) z^{-r-1} \bar{z}^{-s-1},$$

where $\underline{z}$ means the pair of the formal variables z and $\bar{z}$. Physically, a vector in $F_{h, \bar{h}}$ is a state with the spin $h - \bar{h}$ and the energy $h + \bar{h}$, $\mathbf{1}$ corresponds to the vacuum state and $Y(a, \underline{z})$ is the field located at z associated with a state $a \in F$, called a *vertex operator*. It is convenient to write the vertex operator $Y(a, \underline{z})$ as $a(\underline{z})$. For example, $(a_1(x_1)a_2)(x_2)$ means $Y(Y(a_1, \underline{x}_1)a_2, \underline{x}_2)$ and $a_1(x_1)a_2(x_2)$ means $Y(a_1, \underline{x}_1)Y(a_2, \underline{x}_2)$. A four point correlation function is a linear map

$$S : F^{\otimes 4} \rightarrow \text{Cor}_4, \quad (a_1, a_2, a_3, a_4) \mapsto S(a_1, a_2, a_3, a_4)(z_1, z_2, z_3, z_4)$$

which physically calculates the vacuum expectation value of states $a_1, a_2, a_3, a_4 \in F$ at $(z_1, z_2, z_3, z_4) \in X_4$. A four point correlation can be calculated in many ways by using the vertex operator, e.g., the formal power series of the variables $(z_1, \bar{z}_1, \dots, z_4, \bar{z}_4)$,

$$(\mathbf{1}, a_1(z_1)a_2(z_2)a_3(z_3)a_4(z_4)\mathbf{1}),$$

is absolutely convergent to $S(a_1, a_2, a_3, a_4)(z_1, z_2, z_3, z_4) \in \text{Cor}_4$ in the region $\{|z_1| \gg |z_2| \gg |z_3| \gg |z_4|\}$. Another choice of compositions of vertex operators is

$$(0.4) \quad (\mathbf{1}, \left((a_4(x_4)a_2)(x_2)a_1 \right)(x_1)(a_3(x_3)\mathbf{1})).$$

Such compositions are described by parenthesized products of symbols $\{1, 2, 3, 4, \star\}$, in this case, $((42)1)(3\star)$. To be more precise, let $n \in \mathbb{Z}_{>0}$ and Q_n be the set of parenthesized products of $n + 1$ elements $1, 2, \dots, n, \star$ with $\star$ at right most, e.g., $((31)6)(24))(5\star) \in Q_6$. We can naturally associate a formal power series to each element in Q_n , which we call a *parenthesized n point correlation function*. To give a simple example, an element $((31)6)(24))(5\star) \in Q_6$ defines the following parenthesized 6 point correlation function:

$$S_{((31)6)(24))(5\star)}(a_1, \dots, a_6; x_1, \dots, x_6) := (\mathbf{1}, \left[\left((a_3(x_3)a_1)(x_1)a_6 \right)(x_6)a_2(x_2)a_4 \right](x_4)a_5(x_5)\mathbf{1}),$$

where $a_1, \dots, a_6 \in F$. For $A \in Q_n$, we combinatorially give a space T_A of formal variables $x_1, \bar{x}_1, \dots, x_n, \bar{x}_n$ which contains the image of all the parenthesized n point functions. Thus, the parenthesized correlation function defines the map, $S_A : F^{\otimes n} \rightarrow T_A$ for each $A \in Q_n$.

The consistency of quantum field theory says that all parenthesized four point correlation functions associated with all the elements in Q_4 and $a_1, a_2, a_3, a_4 \in F$ (or more generally n point correlation functions) are expansions of the same function $S(a_1, a_2, a_3, a_4)(z_1, z_2, z_3, z_4)$ in different regions after taking the change of variables. For example, the parenthesized correlation function $S_{((42)1)(3\star)}(a_1, a_2, a_3, a_4; x_1, x_2, x_3, x_4)$ in (0.4) is the expansion of $S(a_1, a_2, a_3, a_4)(z_1, z_2, z_3, z_4)$ around $\{|x_1| \gg |x_2| \gg |x_4|, |x_3|\}$, where $(x_1, x_2, x_3, x_4) = (z_1, z_2 - z_1, z_3, z_4 - z_2)$. We remark that for a vertex algebra, the composition of vertex operators $Y(Y(a_1, \underline{x}_0)a_2, \underline{x}_2)$ is an analytic continuation of $Y(a_1, \underline{x}_1)Y(a_2, \underline{x}_2)$ after taking the change of variables $x_0 = x_1 - x_2$. Physically, the change of variables comes from the fact that any state in F is located at 0, i.e., $\lim_{z \rightarrow 0} Y(a, \underline{z})\mathbf{1} = a$ (the state-field correspondence). In Appendix, for each $A \in Q_4$, we give a combinatorial rule for the change of variables and the region by using trees and in Section 1.3 we define the power series expansion of a function in Cor_4 , which gives $e_A : \text{Cor}_4 \rightarrow T_A$.

Then, our formulation of the consistency of a four point function is stated as follows: For any $A \in Q_4$, the composition of the four point function $S : F^{\otimes 4} \rightarrow \text{Cor}_4$ and the expansion map $e_A : \text{Cor}_4 \rightarrow T_A$ is equal to the parenthesized correlation function $S_A : F^{\otimes 4} \rightarrow T_A$, that is,

$$(0.5) \quad S_A = e_A \circ S.$$

In physics, the bootstrap hypothesis says that all consistencies follows from the consistency for special parenthesized correlation functions, $S_{(21)(34)}$ (s-channel), $S_{(41)(23)}$ (t-channel) and $S_{(31)(24)}$ (u-channel). The purpose of this paper is to define a full vertex algebra by slightly modifying the bootstrap hypothesis as a generalization of a $\mathbb{Z}$ -graded vertex algebra and to prove the consistency as a consequence of the axiom of a full vertex algebra. In the theory of vertex algebras, it is commonly used the limit of a four point correlation function as $(z_1, z_4) \rightarrow (\infty, 0)$, which we call a *generalized two point function*. Before explaining the definition of a full vertex algebra, we briefly describe the limit $(z_1, z_4) \rightarrow (\infty, 0)$ in relation to the consistency. Let $a_1, a_2, a_3, a_4 \in F$ with $a_1 \in F_{h_1, \bar{h}_1}$. Then, the limit $(z_1, z_4) \rightarrow (\infty, 0)$ of $z_1^{2h_1} \bar{z}_1^{2\bar{h}_1} S(a_1, a_2, a_3, a_4)$ exists and is a linear combination of

$$(0.6) \quad (z_2 - z_3)^{\alpha_{23}} z_2^{\alpha_{24}} z_3^{\alpha_{34}} (\bar{z}_2 - \bar{z}_3)^{\beta_{23}} \bar{z}_2^{\beta_{24}} \bar{z}_3^{\beta_{34}} f(z_3/z_2),$$

which is a real analytic function on $Y_2 = \{(z_2, z_3) \in \mathbb{C}^2 \mid z_2 \neq z_3, z_2 \neq 0, z_3 \neq 0\}$. Denote by GCor_2 the space of real analytic function spanned by (0.6).

Since $(\infty, z_2, z_3, 0) \in X_4$ with $|z_2| > |z_3|$ is in the convergent region of $S_{1(2(3(4\star)))}(a_1, a_2, a_3, a_4)$, which is $\{|z_1| > |z_2| > |z_3| > |z_4|\}$, the limit of $(-1)^{h_1 - \bar{h}_1} z_1^{2h_1} \bar{z}_1^{2\bar{h}_1} S(a_1, a_2, a_3, a_4)$ as $(z_1, z_4) \rightarrow (\infty, 0)$ with $|z_2| > |z_3|$ is equal to the formal limit,

$$\lim_{(z_1, z_4) \rightarrow (\infty, 0)} (-1)^{h_1 - \bar{h}_1} z_1^{2h_1} \bar{z}_1^{2\bar{h}_1} S_{1(2(3(4\star)))}(a_1, a_2, a_3, a_4) = (a_1, Y(a_2, z_2)Y(a_3, z_3)a_4).$$

The convergent region of $S_{1(3(2(4\star)))}(a_1, a_2, a_3, a_4)$ and $S_{1((23)(4\star))}(a_1, a_2, a_3, a_4)$ are $\{|z_1| > |z_3| > |z_2| > |z_4|\}$ and $\{|z_1| > |z_3| > |z_2 - z_3|, |z_4|\}$. Thus, the limits of $(-1)^{h_1 - \bar{h}_1} z_1^{2h_1} \bar{z}_1^{2\bar{h}_1} S(a_1, a_2, a_3, a_4)$ as $(z_1, z_4) \rightarrow (\infty, 0)$ with $|z_3| > |z_2|$ and $|z_3| > |z_2 - z_3|$ are given by $(a_1, Y(a_3, z_3)Y(a_2, z_2)a_4)$ and $(a_1, Y(Y(a_2, z_{23})a_3, z_3)a_4)$, respectively, where $z_{23} = z_2 - z_3$. Furthermore, in the regions, $\{|z_2| > |z_3|\}$, $\{|z_3| > |z_2|\}$ and $\{|z_3| > |z_2 - z_3|\}$, $\frac{z_3}{z_2}$ is close to 0, ∞ and 1, respectively. Thus, the expansions of (0.6) in these regions are determined by the expansion of $f \in F_{0,1,\infty}$ around $\{0, 1, \infty\}$.

0.3. Definition of full vertex algebra and main result. Let us describe the precise definition of a full vertex algebra. A full vertex algebra is an $\mathbb{R}^2$ -graded vector space $F = \bigoplus_{h, \bar{h} \in \mathbb{R}} F_{h, \bar{h}}$ with a distinguished vector $\mathbf{1} \in F_{0,0}$ and a vertex operator

$$Y(-, \underline{z}) : F \rightarrow \text{End } F[[z, \bar{z}, |z|^{\mathbb{R}}]], \quad a \mapsto Y(a, \underline{z}) = \sum_{r, s \in \mathbb{R}} a(r, s) z^{-r-1} \bar{z}^{-s-1}$$

satisfying the following conditions: For any $a, b \in F$, there exists $N \in \mathbb{R}$ such that $a(r, s)b = 0$ unless $r \leq N$ and $s \leq N$; $F_{h, \bar{h}} = 0$ unless $h - \bar{h} \in \mathbb{Z}$; The vacuum vector $\mathbf{1}$ satisfies $Y(\mathbf{1}, \underline{z}) = \text{id}_F$ and $Y(a, \underline{z})\mathbf{1} \in F[[z, \bar{z}]]$ and $\lim_{z \rightarrow 0} Y(a, \underline{z})\mathbf{1} = a$ for any $a \in F$; $F_{h, \bar{h}}(r, s)F_{h', \bar{h}'} \subset F_{h+h'-r-1, \bar{h}+\bar{h}'-s-1}$ for any $h, h', \bar{h}, \bar{h}', r, s \in \mathbb{R}$; For any $a, b, c \in F$ and

$u \in F^\vee = \bigoplus_{h, \bar{h} \in \mathbb{R}} F_{h, \bar{h}}^*$, there exists $\mu(z_1, z_2) \in \text{GCor}_2$ such that

$$(0.7) \quad \begin{aligned} u(Y(a, \underline{z}_1)Y(b, \underline{z}_2)c) &= \mu(z_1, z_2)|_{|z_1| > |z_2|}, \\ u(Y(Y(a, \underline{z}_0)b, \underline{z}_2)c) &= \mu(z_0 + z_2, z_2)|_{|z_2| > |z_0|}, \\ u(Y(b, \underline{z}_2)Y(a, \underline{z}_1)c) &= \mu(z_1, z_2)|_{|z_2| > |z_1|}, \end{aligned}$$

where $F_{h, \bar{h}}^*$ is the dual vector space and $\mu(z_1, z_2)|_{|z_1| > |z_2|}$ is the expansion of μ in $\{|z_1| > |z_2|\}$. For a full vertex algebra F , we define linear maps $D, \bar{D} \in \text{End } F$ by

$$Y(a, \underline{z})\mathbf{1} = a + Daz + \bar{D}a\bar{z} + \dots$$

Then, one can show that $[D, Y(a, \underline{z})] = Y(Da, \underline{z}) = d/dz Y(a, \underline{z})$ and $[\bar{D}, Y(a, \underline{z})] = Y(\bar{D}a, \underline{z}) = d/d\bar{z} Y(a, \underline{z})$ (Proposition 2.1). Thus, if $a, b \in \ker \bar{D}$, then the generalized correlation function (0.7) satisfies $d/d\bar{z}_1 \mu = d/d\bar{z}_2 \mu = 0$, that is, μ is a holomorphic function on Y_2 . We prove that if $\mu \in \text{GCor}_2$ is holomorphic, then $\mu \in \mathbb{C}[z_1^\pm, z_2^\pm, (z_1 - z_2)^\pm]$, which implies that $\ker \bar{D}$ is a vertex algebra (Proposition 2.4). This reflects the fact that a function in $F_{0,1,\infty}$ is holomorphic if and only if $f \in \mathbb{C}[z^\pm, (1 - z)^{-1}]$. Thus, in the definition of a full vertex algebra, we replace the function space of a vertex algebra $\mathbb{C}[z^\pm, (1 - z)^{-1}]$ with $F_{0,1,\infty}$, which implies that the notion of a full vertex algebra is a generalization of the notion of a $\mathbb{Z}$ -graded vertex algebra.

We remark that, for a vertex algebra, the pole of $u(Y(a, z_1)Y(b, z_2)c)$ is $(z_1 - z_2)^{-k}$ for some $k \in \mathbb{Z}$, which implies that $(z_1 - z_2)^N [Y(a, z_1), Y(b, z_2)] = 0$ for a sufficiently large $N \in \mathbb{Z}_{\geq 0}$. However, for a full vertex algebra, the pole of $u(Y(a, \underline{z}_1)Y(b, \underline{z}_2)c)$ is a linear sum of $(z_1 - z_2)^n |z_1 - z_2|^r$ for $r \in \mathbb{R}$ with $n \in \mathbb{Z}$, e.g., $1 + |z_1 - z_2|^{1/2} + \dots$. Hence, we could not simultaneously cancel all poles by multiplication by a polynomial. The number of the index $r \in \mathbb{R}/\mathbb{Z}$ appeared in $(z_1 - z_2)^n |z_1 - z_2|^r$ reflects the number of intermediate states, which is mathematically the number of the irreducible components of the tensor product of modules. In the case of a full vertex algebra, the function space $F_{0,1,\infty}$ is much more complicated, however, we can still describe all the consistencies of a four point function in term of the expansions of the same function $f \in F_{0,1,\infty}$ around $\{0, 1, \infty\}$.

In order to prove the consistency, we need the global conformal symmetry. A full vertex algebra with an energy-momentum tensor, a pair of holomorphic and anti-holomorphic conformal vectors, which we call a full vertex operator algebra (see Section 3.1). A full vertex operator algebra admits an action of the Virasoro algebras, $\text{Vir} \oplus \text{Vir}$ and, in particular, a subalgebra, $\text{sl}_2\mathbb{C} \oplus \text{sl}_2\mathbb{C} = \bigoplus_{i=-1,0,1} L(i) \oplus \bigoplus_{j=-1,0,1} \bar{L}(j)$. A vector $v \in F_{h, \bar{h}}$ is called a *quasi-primary* if $L(1)v = \bar{L}(1)v = 0$ and a full vertex operator algebra is called QP-generated if F is generated by quasi-primary vectors as an $\text{sl}_2\mathbb{C} \oplus \text{sl}_2\mathbb{C}$ -module. A notion of a dual module, introduced in [FHL] for a vertex operator algebra, can be generalized to a full vertex operator algebra and its modules. We give a criterion when a full vertex operator algebra is self-dual and QP-generated, which generalize a result of Li [L1]. The main result of this paper is that for a QP-generated self-dual full vertex operator algebra, all the consistencies of four point correlation functions hold (Theorem 3.2).

0.4. Locality and the proof. One can relax the assumption of a full vertex algebra, (0.7), to the existence of $D, \bar{D} \in \text{End } F$ and the following condition: For $u \in F^\vee$ and $a_1, a_2, a_3 \in F$, there exists $\mu \in \text{GCor}_2$ such that

$$\begin{aligned} u(Y(a, \underline{z}_1)Y(b, \underline{z}_2)c) &= \mu(z_1, z_2)|_{|z_1| > |z_2|}, \\ u(Y(b, \underline{z}_2)Y(a, \underline{z}_1)c) &= \mu(z_1, z_2)|_{|z_2| > |z_1|}. \end{aligned}$$

It is a generalization of the Goddard's axiom of a vertex algebra (Proposition 4.1). By using this description, we construct a QP-generated self-dual full vertex operator algebra associated with an even lattice.

Hereafter, we recall the S_4 -symmetry of the correlation function and then explain the key idea of the proof of the consistency. The symmetric group S_4 acts on $X_4 \subset (\mathbb{CP}^1)^4$ by the permutation, thus, on Cor_4 . As a consequence of the consistency, four point correlation functions $S : F^{\otimes 4} \rightarrow \text{Cor}_4$ posses an S_4 -symmetry,

$$(S_4\text{-symmetry}) \quad \sigma \cdot S(a_1, a_2, a_3, a_4) = S(a_{\sigma^{-1}1}, a_{\sigma^{-1}2}, a_{\sigma^{-1}3}, a_{\sigma^{-1}4}),$$

for any $a_1, a_2, a_3, a_4 \in F$ and $\sigma \in S_4$. Interestingly, in the Goddard's axiom, we only assume that the correlation function is invariant under the permutation $(23) \in S_4$. In contrast, four point correlation functions should be invariant under the S_4 -symmetry. We observe that since the S_4 -action commutes with the diagonal action of $\text{PSL}_2\mathbb{C}$ on $X_4 \subset (\mathbb{CP}^1)^4$, the homeomorphism $\xi : \text{PSL}_2\mathbb{C} \backslash X_4 \rightarrow \mathbb{CP}^1 \setminus \{0, 1, \infty\}$ induces a S_4 -action on $\mathbb{CP}^1 \setminus \{0, 1, \infty\}$, which permutes $\{0, 1, \infty\}$. Thus, the S_4 -action is degenerate to the S_3 -action, that is, the action of the Klein subgroup $C_2 \times C_2 \subset S_4$ is a part of the global conformal symmetry. Since $(34) \cdot \xi = \frac{\xi}{\xi-1}$ and both ξ and $\frac{\xi}{\xi-1}$ go to zero as $\xi \rightarrow 0$, there is a simple relation between the expansion of $f \circ \xi$ in the domain $|z_1| > |z_2| > |z_3| > |z_4|$ and $|z_1| > |z_2| > |z_4| > |z_3|$. This relation gives the skew-symmetry of a full vertex algebra F , i.e.,

$$Y(a, \underline{z})b = \exp(Dz + \bar{D}\bar{z})Y(b, -\underline{z})a$$

for any $a, b \in F_{0,1,\infty}$. Since S_4 is generated by the Klein subgroup together with $(23), (34) \in S_4$, the axiom of a full vertex algebra and the global conformal symmetry implies that the S_4 -symmetry of the correlation function (Theorem 3.1).

Another important point is that in the definition of full vertex algebra, we consider the generalized correlation function, whereas in the consistency, we need the four point correlation functions. This problem is solved by the differential equations derived from the global conformal symmetry.

The bootstrap hypothesis is also proved by using this differential equation. More precisely, consider the following parenthesized correlation functions:

$$\begin{aligned} (\text{s-channel}) \quad & (\mathbf{1}, ((a_2(x_2)a_1)(x_1)a_3(x_3)a_4)(x_4)\mathbf{1})) \\ (\text{t-channel}) \quad & (\mathbf{1}, ((a_4(x_4)a_1)(x_1)a_3(x_3)a_2)(x_2)\mathbf{1})) \\ (\text{u-channel}) \quad & (\mathbf{1}, ((a_3(x_3)a_1)(x_1)a_2(x_2)a_4)(x_4)\mathbf{1})). \end{aligned}$$

The equality of s-channel and t-channel (or s-channel and u-channel) are called a *bootstrap equation*. By using the differential equation, we can relate them with the generalized two point functions in (0.7) and one can show that under some assumptions if a vertex operator Y on F satisfies the bootstrap equation, then F is a full vertex algebra (Proposition 4.7).

0.5. Outline. In Section 1, the definition of the space of correlation functions Cor_4 and the expansions $e_A : \text{Cor}_4 \rightarrow T_A$ associated with parenthesized products are given. Properties of the functions and expansions, e.g., relations among the expansions in different regions and differential equations, holomorphic correlation functions, are studied. In Section 2, we define a full vertex algebra and study the consequence of the axiom by using the results in the previous section. In particular, we study a property of the translation map $D, \bar{D}$ and show that $\ker \bar{D}$ and $\ker D$ is a vertex algebra. A tensor product of full vertex algebras is also constructed here. Section 3 is devoted to studying the parenthesized correlation functions and its consistency. The existence of bilinear form on a full vertex operator algebra are also studied. In Section 4, a full vertex algebra is constructed from the Goddard's axiom and the bootstrap equation and In Section 5, we construct an example of a QP-generated self-dual full vertex operator algebra from an even lattice, which generalize lattice vertex algebras. In Appendix, the combinatorial rule for the expansions is given.

ACKNOWLEDGEMENTS

The author would like to express his gratitude to his supervisor Masahito Yamazaki and Professor Yuji Tachikawa for useful discussions and to Shigenori Nakatsuka for reading the manuscript and valuable comments. This work was supported by World Premier International Research Center Initiative (WPI Initiative), MEXT, Japan. The author was also supported by the Program for Leading Graduate Schools, MEXT, Japan.

CONTENTS

Introduction	1
0.1. Space of four point correlation function	3
0.2. Consistency and parenthesized correlation functions	4
0.3. Definition of full vertex algebra and main result	6
0.4. Locality and the proof	8
0.5. Outline	9
Acknowledgements	9
Preliminaries and Notations	10
1. Correlation functions and formal calculus	12
1.1. Definition of four point functions	12
1.2. S_4 -symmetry	14
1.3. Expansions	16

1.4.	Relations among expansions	22
1.5.	Formal Differential Equation and Convergence	24
1.6.	Holomorphic correlation functions	26
1.7.	Vacuum state and D_4 -symmetry	28
1.8.	Generalized two point Correlation function	30
2.	Full vertex algebra	33
2.1.	Definition of full vertex algebra	33
2.2.	Holomorphic vertex operators	37
2.3.	Tensor product of full vertex algebras	38
3.	Correlation functions and full vertex algebras	39
3.1.	Self-duality	39
3.2.	Quasi-primary vectors	42
3.3.	Parenthesized correlation functions and formal power series	44
3.4.	Consistency of four point functions	46
4.	Construction	52
4.1.	Locality and Associativity I	52
4.2.	Locality and Associativity II	54
4.3.	Bootstrap	57
5.	Example	58
5.1.	Full vertex algebra and AH pair	59
5.2.	Troidal compactification of string theory	63
5.3.	Commutative algebra object in H-Vect	64
6.	Appendix	65
6.1.	Binary tree and expansions	65
	References	67

PRELIMINARIES AND NOTATIONS

We assume that the base field is $\mathbb{C}$ unless otherwise stated. Throughout of this paper, z and $\bar{z}$ are independent formal variables. We will use the notation $\underline{z}$ for the pair $(z, \bar{z})$ and $|z|$ for $z\bar{z}$. For a vector space V , we denote by $V[[z_1, \bar{z}_1, |z_1|^\mathbb{R}, \dots, z_n, \bar{z}_n, |z_n|^\mathbb{R}]]$ the set of formal sums

$$\sum_{s_1, \bar{s}_1, \dots, s_n, \bar{s}_n \in \mathbb{R}} a_{s_1, \bar{s}_1, \dots, s_n, \bar{s}_n} z_1^{s_1} \bar{z}_1^{\bar{s}_1} \dots z_n^{s_n} \bar{z}_n^{\bar{s}_n}$$

such that

- (1) $a_{s_1, \bar{s}_1, \dots, s_n, \bar{s}_n} = 0$ unless $s_i - \bar{s}_i \in \mathbb{Z}$ for all $i = 1, \dots, n$;
- (2) $\{(s_1, \bar{s}_1, \dots, s_n, \bar{s}_n) \in \mathbb{R}^{2n} \mid a_{s_1, \bar{s}_1, \dots, s_n, \bar{s}_n} \neq 0\}$ is a countable set.

We also denote by $V((z_1, \bar{z}_1, |z_1|^\mathbb{R}, \dots, z_n, \bar{z}_n, |z_n|^\mathbb{R}))$ the subspace of $V[[z_1, \bar{z}_1, |z_1|^\mathbb{R}, \dots, z_n, \bar{z}_n, |z_n|^\mathbb{R}]]$ spanned by the series satisfying

- (3) There exists $N \in \mathbb{Z}$ such that $a_{s_1, \bar{s}_1, \dots, s_n, \bar{s}_n} = 0$ unless $s_i, \bar{s}_i \geq N$ for all $i = 1, \dots, n$.

For a series $f(\underline{z}) = \sum_{r,s \in \mathbb{R}} a_{r,s} z^r \bar{z}^s \in V((z, \bar{z}, |z|^{\mathbb{R}}))$, The number of the elements $\{r \in \mathbb{R} \mid a_{r,s} \neq 0\}$ in the image of the natural map $\mathbb{R} \rightarrow \mathbb{R}/\mathbb{Z}$ is called an *exponent of the series*. If the exponent of $f(\underline{z})$ is $l \in \mathbb{Z}_{\geq 0}$, then $f(\underline{z})$ could be written as

$$\sum_{i=1}^l \sum_{n,m=0}^{\infty} a_{n,m}^i z^n \bar{z}^m |z|^{r_i},$$

where $r_i \in \mathbb{R}$ and $r_i - r_j \notin \mathbb{Z}$ for any $i \neq j$.

We will consider the following subspaces of $V[[z, \bar{z}, |z|^{\mathbb{R}}]]$:

$$\begin{aligned} V[[z, \bar{z}]] &= \left\{ \sum_{s, \bar{s} \in \mathbb{Z}_{\geq 0}} a_{s, \bar{s}} z^s \bar{z}^{\bar{s}} \mid a_{s, \bar{s}} \in V \right\}, \\ V[z^{\pm}, \bar{z}^{\pm}] &= \left\{ \sum_{s, \bar{s} \in \mathbb{Z}} a_{s, \bar{s}} z^s \bar{z}^{\bar{s}} \mid a_{s, \bar{s}} \in V, \text{ all but finitely many } a_{s, \bar{s}} = 0 \right\}, \\ V[|z|^{\mathbb{R}}] &= \left\{ \sum_{r \in \mathbb{R}} a_r z^r \bar{z}^r \mid a_r \in V, \text{ all but finitely many } a_r = 0 \right\}. \end{aligned}$$

We will also consider their combinations, e.g., $V((y/x, \bar{y}/\bar{x}, |y/x|^{\mathbb{R}})) [x^{\pm}, \bar{x}^{\pm}, |x|^{\mathbb{R}}]$, which is a subspace of $V[[x, y, \bar{x}, \bar{y}, |x|^{\mathbb{R}}, |y|^{\mathbb{R}}]]$ spanned by

$$\sum_{i=1}^k \sum_{n,m=-l}^l \sum_{r,s \in \mathbb{R}} a_{n,m,r,s}^i x^{n+r_i} \bar{x}^{m+r_i} (y/x)^r (\bar{y}/\bar{x})^s$$

for some $k, l \in \mathbb{Z}_{>0}$ and $r_i \in \mathbb{R}$ and $a_{n,m,r,s}^i \in V$ such that $a_{n,m,r,s}^i = 0$ unless $r - s \in \mathbb{Z}$ and there exists N such that $a_{n,m,r,s}^i = 0$ unless $r \geq N$ and $s \geq N$.

Let $\frac{d}{dz}$ and $\frac{d}{d\bar{z}}$ be formal differential operators acting on $V[[z, \bar{z}, |z|^{\mathbb{R}}]]$ by

$$\begin{aligned} \frac{d}{dz} \sum_{s, \bar{s} \in \mathbb{R}} a_{s, \bar{s}} z^s \bar{z}^{\bar{s}} &= \sum_{s, \bar{s} \in \mathbb{R}} s a_{s, \bar{s}} z^{s-1} \bar{z}^{\bar{s}}, \\ \frac{d}{d\bar{z}} \sum_{s, \bar{s} \in \mathbb{R}} a_{s, \bar{s}} z^s \bar{z}^{\bar{s}} &= \sum_{s, \bar{s} \in \mathbb{R}} \bar{s} a_{s, \bar{s}} z^s \bar{z}^{\bar{s}-1}. \end{aligned}$$

Since $\frac{d}{dz} |z|^s = s |z|^{s-1}$, the differential operators $\frac{d}{dz}$ and $\frac{d}{d\bar{z}}$ acts on all the above vector spaces. Define a linear map $\lim_{z \rightarrow f(w)}$ from a space of the formal variable $(z, \bar{z})$ to a space of the formal variable $(w, \bar{w})$ if the substitution of $(f(w), f(\bar{w}))$ into $(z, \bar{z})$ is well-defined, that is, each coefficient is a finite sum. For example, $\lim_{z \rightarrow -z} V((z, \bar{z}, |z|^{\mathbb{R}})) \rightarrow V((z, \bar{z}, |z|^{\mathbb{R}}))$ is given by

$$\lim_{z \rightarrow -z} \sum_{r,s \in \mathbb{R}} a_{r,s} z^r \bar{z}^s = \sum_{r,s \in \mathbb{R}} (-1)^{r-s} a_{r,s} z^r \bar{z}^s.$$

Since $a_{r,s} = 0$ for any $r - s \notin \mathbb{Z}$, $\lim_{z \rightarrow -z}$ is well-defined. Another example is

$$\lim_{x \rightarrow x+y} : V((y/x, \bar{y}/\bar{x}, |y/x|^{\mathbb{R}})) \rightarrow V((y/x, \bar{y}/\bar{x}, |y/x|^{\mathbb{R}})),$$

which is defined by

$$\lim_{x \rightarrow x+y} \left(\sum_{r,s \in \mathbb{R}} a_{r,s} (y/x)^r (\bar{y}/\bar{x})^s \right) = \sum_{r,s \in \mathbb{R}} \sum_{i,j \in \mathbb{Z}_{\geq 0}} \binom{-r}{i} \binom{-s}{j} a_{r,s} (y/x)^{r+i} (\bar{y}/\bar{x})^{s+j},$$

where we used $(x + y)^s = \sum_{m \geq 0} \binom{s}{m} x^{s-m} y^m$. It is well-defined since $a_{r,s} = 0$ for sufficiently small r or s . We end this section by discussing a convergence of a formal power series in $\mathbb{C}((z_1, \bar{z}_1, |z_1|^\mathbb{R}, \dots, z_n, \bar{z}_n, |z_n|^\mathbb{R}))$. Let $f \in \mathbb{C}((z_1, \bar{z}_1, |z_1|^\mathbb{R}, \dots, z_n, \bar{z}_n, |z_n|^\mathbb{R}))$. Then, there exists $N \in \mathbb{R}$ such that

$$(0.8) \quad |z_1|^N |z_2|^N \cdots |z_n|^N f(z_1, \dots, z_n) = \sum_{\substack{s_1, \bar{s}_1, \dots, s_n, \bar{s}_n \in \mathbb{R} \\ s_1, \bar{s}_1, \dots, s_n, \bar{s}_n \geq 0}} a_{s_1, \bar{s}_1, \dots, s_n, \bar{s}_n} z_1^{s_1} \bar{z}_1^{\bar{s}_1} \cdots z_n^{s_n} \bar{z}_n^{\bar{s}_n}.$$

We say the series f is absolutely convergent around 0 if $|z_1|^N |z_2|^N \cdots |z_n|^N f(z_1, \dots, z_n)$ is absolutely convergent in $\{(z_1, z_2, \dots, z_n) \in \mathbb{C}^n \mid |z_1|, |z_2|, \dots, |z_n| < R\}$ for some $R \in \mathbb{R}_{>0}$. In this case, $f(z_1, \dots, z_n)$ is compactly absolutely-convergent to a continuous function in the annulus $\{(z_1, z_2, \dots, z_n) \in \mathbb{C}^n \mid 0 < |z_1|, |z_2|, \dots, |z_n| < R\}$.

1. CORRELATION FUNCTIONS AND FORMAL CALCULUS

In this section, we define the space of four point correlation functions, Cor_4 and develop formal calculus which we need to study a full vertex algebra. In Section 1.1, we define Cor_4 and in Section 1.2, we study an action of the symmetric group S_4 on Cor_4 . Section 1.3 and 1.7 is devoted to studying series expansions of a function in Cor_4 . We study formal differential equations in Section 1.5 and holomorphic correlation function in Section 1.6. Certain limit of a four point correlation function is studied in Section 1.8. The reader who is only interested in the definition of a full vertex algebra can skip Section 1.3, 1.4, 1.5, 1.6 and 1.7.

1.1. Definition of four point functions. For $n \in \mathbb{Z}_{\geq 0}$, set $X_n = \{(z_1, \dots, z_n) \in (\mathbb{CP}^1)^n \mid z_i \neq z_j \text{ for any } i \neq j\}$, called a space of ordered configurations of n points in the projective space $\mathbb{CP}^1$. In this section, we define and study a space of four point correlation functions in two dimensional conformal field theory, which are $\mathbb{C}$ -valued real analytic functions on X_4 with possible singularity along $\{(z_1, \dots, z_4) \in (\mathbb{CP}^1)^4 \mid z_i = z_j \text{ for } 1 \leq i < j \leq 4\}$. Recall that the automorphism group of $\mathbb{CP}^1$ is $\text{PSL}_2\mathbb{C}$, whose action is called the linear fractional transformation and also called a global conformal symmetry. Define the real analytic function $\xi : X_4 \rightarrow P^1\mathbb{C} \setminus \{0, 1, \infty\}$ by

$$\xi(z_1, z_2, z_3, z_4) = \frac{(z_1 - z_2)(z_3 - z_4)}{(z_1 - z_3)(z_2 - z_4)}.$$

It is easy to show that the map $\xi : X_4 \rightarrow P^1\mathbb{C} \setminus \{0, 1, \infty\}$ is invariant under the diagonal action of $\text{PSL}_2\mathbb{C}$ on $X_4 \subset (\mathbb{CP}^1)^4$, in fact, which gives a homeomorphism $\text{PSL}_2\mathbb{C} \backslash X_4 \rightarrow P^1\mathbb{C} \setminus \{0, 1, \infty\}$ (see, for example, [Y]).

Remark 1.1. *The hypersurfaces in $(\mathbb{CP}^1)^4$,*

$$\{z_1 = z_2\} \cup \{z_3 = z_4\}$$

and

$$\{z_1 = z_4\} \cup \{z_2 = z_3\}$$

and

$$\{z_1 = z_3\} \cup \{z_2 = z_4\}$$

are maps to 0 and 1, ∞ respectively by $\xi : (\mathbb{CP}^1)^4 \rightarrow \mathbb{CP}^1$.

Let $\alpha_1, \dots, \alpha_n \in \mathbb{CP}^1$ and f be a $\mathbb{C}$ -valued real analytic function on $\mathbb{CP}^1 \setminus \{\alpha_1, \dots, \alpha_n\}$.

A chart (χ, α) of $\mathbb{CP}^1$ at a point $\alpha \in \mathbb{CP}^1$ is a biholomorphism χ from an open subset U of $\mathbb{CP}^1$ to an open subset of $\mathbb{C}$ such that $\alpha \in U$ and $\chi(\alpha) = 0$. We say that f has a *conformal singularity* at α_i if for any chart (χ, α_i) of $\mathbb{CP}^1$ at α_i , there exists a formal power series

$$(1.1) \quad \sum_{r,s \in \mathbb{R}} a_{r,s} p^r \bar{p}^s \in \mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}}))$$

such that it is compactly absolutely-convergent to $f \circ \chi^{-1}(p)$ in the annulus $\{p \in \mathbb{C} \mid 0 < |p| < R\}$ for some $R \in \mathbb{R}_{>0}$ (see Section 0.5). It is clear that the above condition is independent of a choice of a chart and the coefficients of the series is uniquely determined by the chart.

Denote by $F_{0,1,\infty}$ the space of real analytic functions on $\mathbb{CP}^1 \setminus \{0, 1, \infty\}$ with possible conformal singularities at $\{0, 1, \infty\}$. Let p be a canonical coordinate of $\mathbb{C} \subset \mathbb{CP}^1$. A non-trivial example of such functions is

$$(1.2) \quad f_{\text{Ising}}(p) = |1 - \sqrt{1-p}|^{1/2} + |1 + \sqrt{1-p}|^{1/2},$$

which appears in a four point function of the 2 dimensional Ising model. We remark that the exponent of the series (1.1) is independent of the choice of the chart. An exponent of a function $f \in F_{0,1,\infty}$ is the maximal number of the exponents of the series expansion of f at $\{0, 1, \infty\}$. For example, the expansion of f_{Ising} for the charts p is

$$(1.3) \quad 2 + |p|^{1/2}/2 - p/4 - \bar{p}/4 + |p|^{1/2}(p + \bar{p})/16 + p\bar{p}/32 - 5p^2/64 - 5\bar{p}^2/64 + \dots$$

Since $f_{\text{Ising}}(p)$ satisfies $f_{\text{Ising}}(p) = f_{\text{Ising}}(1-p) = (z\bar{z})^{1/4} f_{\text{Ising}}(1/p)$, the exponent of f_{Ising} is 2. We remark that for a rational conformal field theory (or more generally quasi-rational conformal field theory), the exponent is always finite.

In this paper, we consider the following special charts of $\{0, 1, \infty\}$:

$$\text{Chart}(0, 1, \infty) = \{p, \frac{p}{p-1}, 1-p, 1-\frac{1}{p}, \frac{1}{p}, \frac{1}{1-p}\}.$$

Recall that the ring of regular functions on the affine scheme $\mathbb{CP}^1 \setminus \{0, 1, \infty\}$ is $\mathbb{C}[p^\pm, (1-p)^\pm]$ and $\text{Chart}(0, 1, \infty)$ is a generator of the ring. It is easy to show that a function in $\mathbb{C}[p^\pm, (1-p)^\pm]$ has conformal singularities at $\{0, 1, \infty\}$. Thus, $\mathbb{C}[p^\pm, (1-p)^\pm] \subset F_{0,1,\infty}$. Conversely, from the existence of the expansion the following proposition follows:

Proposition 1.1. *If $f \in F_{0,1,\infty}$ is a holomorphic function on $\mathbb{CP}^1 \setminus \{0, 1, \infty\}$, then $f \in \mathbb{C}[p^\pm, (1-p)^\pm]$.*

We will consider the real analytic functions on X_4 of the form:

$$(1.4) \quad \prod_{1 \leq i < j \leq 4} (z_i - z_j)^{\alpha_{ij}} (\bar{z}_i - \bar{z}_j)^{\beta_{ij}} f \circ \xi(z_1, z_2, z_3, z_4),$$

where $f \in F_{0,1,\infty}$ and $\alpha_{ij}, \beta_{ij} \in \mathbb{R}$ such that $\alpha_{ij} - \beta_{ij} \in \mathbb{Z}$ for any $1 \leq i < j \leq 4$. In this paper, we always allow to have singularities around $\{z_i = \infty\}_{i=1,2,3,4}$; The asymptotic behavior of (1.4) as $z_1 \rightarrow \infty$ is $z_1^{\alpha_{12} + \alpha_{13} + \alpha_{14}} \bar{z}_1^{\beta_{12} + \beta_{13} + \beta_{14}}$. By the condition $\alpha_{ij} - \beta_{ij} \in \mathbb{Z}$, we have

$$(z_i - z_j)^{\alpha_{ij}} (\bar{z}_i - \bar{z}_j)^{\beta_{ij}} = |z_i - z_j|^{\alpha_{ij}} (\bar{z}_i - \bar{z}_j)^{\beta_{ij} - \alpha_{ij}},$$

which implies that $\prod_{1 \leq i < j \leq 4} (z_i - z_j)^{\alpha_{ij}} (\bar{z}_i - \bar{z}_j)^{\beta_{ij}}$ is a single valued real analytic function on X_4 .

Let Cor_4 be the space of $\mathbb{C}$ -valued real analytic functions spanned by (1.4). We also consider the spaces of functions on X_2 and X_3 and X_4 spanned by

$$\begin{aligned} & (z_1 - z_2)^{\alpha_{12}} (\bar{z}_1 - \bar{z}_2)^{\beta_{12}}, \\ & \prod_{1 \leq i < j \leq 3} (z_i - z_j)^{\alpha_{ij}} (\bar{z}_i - \bar{z}_j)^{\beta_{ij}}, \\ & \prod_{1 \leq i < j \leq 4} (z_i - z_j)^{\alpha_{ij}} (\bar{z}_i - \bar{z}_j)^{\beta_{ij}} \end{aligned}$$

where $\alpha_{ij}, \beta_{ij} \in \mathbb{R}$ such that $\alpha_{ij} - \beta_{ij} \in \mathbb{Z}$ for any $1 \leq i < j \leq 4$. We denote them by $\text{Cor}_2, \text{Cor}_3$ and Cor_4^f , respectively. By the definition, we have:

Lemma 1.1. *A function $\phi \in \text{Cor}_4$ is translation invariant, i.e., $\phi(z_1, z_2, z_3, z_4) = \phi(z_1 + \alpha, z_2 + \alpha, z_3 + \alpha, z_4 + \alpha)$ for any $\alpha \in \mathbb{C}$.*

1.2. S_4 -symmetry. Define the left action of the symmetric group S_4 on $(\mathbb{CP}^1)^4$ by the permutation, $\sigma \cdot (z_1, z_2, z_3, z_4) = (z_{\sigma^{-1}1}, z_{\sigma^{-1}2}, z_{\sigma^{-1}3}, z_{\sigma^{-1}4})$ for $(z_1, z_2, z_3, z_4) \in (\mathbb{CP}^1)^4$ and $\sigma \in S_4$. Since the action commutes with the diagonal action of $\text{PSL}_2\mathbb{C}$ on $X_4 \subset (\mathbb{CP}^1)^4$, we have an S_4 -action on $\mathbb{CP}^1 \setminus \{0, 1, \infty\}$. It is easy to show that each image of $\sigma \in S_4$ in $\text{Aut } \mathbb{CP}^1 \setminus \{0, 1, \infty\}$ is a linear fractional transformation and preserves $\{0, 1, \infty\}$ (see Remark 1.1). Let $\text{Aut}(0, 1, \infty)$ be the subgroup of the linear fractional transformations $\text{PSL}_2\mathbb{C}$ which preserves $\{0, 1, \infty\}$. Since the action of $\text{PSL}_2\mathbb{C}$ on $\mathbb{CP}^1$ is strictly 3-transitive, an element of $\text{Aut}(0, 1, \infty)$ is uniquely determined by the action on the 3 points $\{0, 1, \infty\}$. Thus, $\text{Aut}(0, 1, \infty)$ is isomorphic to the permutation group S_3 . Hence, we have a group homomorphism $t : S_4 \rightarrow \text{Aut}(0, 1, \infty)$ and the kernel of this map is the Klein subgroup $K_4 = \{1, (12)(34), (13)(24), (14)(23)\}$.

The action of S_4 on X_4 induces the dual action on the function space Cor_4 by $\sigma f(-) = f(\sigma^{-1}-)$ for $f \in \text{Cor}_4$ and $\sigma \in S_4$, which also induces the action on the function space of $\mathbb{CP}^1 \setminus \{0, 1, \infty\}, F_{0,1,\infty}$.

Remark 1.2. *For $\sigma \in S_4$ and $\prod_{1 \leq i < j \leq 4} (z_i - z_j)^{\alpha_{ij}} (\bar{z}_i - \bar{z}_j)^{\beta_{ij}} \in \text{Cor}_4^f$,*

$$\sigma \cdot \prod_{1 \leq i < j \leq 4} (z_i - z_j)^{\alpha_{ij}} (\bar{z}_i - \bar{z}_j)^{\beta_{ij}} = \prod_{1 \leq i < j \leq 4} (z_{\sigma i} - z_{\sigma j})^{\alpha_{ij}} (\bar{z}_{\sigma i} - \bar{z}_{\sigma j})^{\beta_{ij}}.$$

Regard the chart $p \in \text{Chart}(0, 1, \infty)$ as a function on $\mathbb{CP}^1 \setminus \{0, 1, \infty\}$, for $\sigma \in S_4$, we have $\sigma \cdot p(-) = p(t_\sigma^{-1} -) = t_\sigma^{-1} \cdot p$, where $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot p$ is defined by $\frac{ap+b}{cp+d} \in \mathbb{C}(p)$ for $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{PSL}_2\mathbb{C}$.

Lemma 1.2. *The explicit expressions of the map $t : S_4 \rightarrow \text{Aut}(0, 1, \infty) \subset \text{PSL}_2\mathbb{C}$ and the orbit $\sigma \cdot p = t_\sigma^{-1} \cdot p$ is*

$$\begin{aligned} K_4 &\mapsto (1) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, & p &\mapsto p, \\ (12)K_4 &\mapsto (1\infty) = \begin{pmatrix} 1 & 0 \\ 1 & -1 \end{pmatrix}, & p &\mapsto \frac{p}{p-1}, \\ (23)K_4 &\mapsto (0\infty) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, & p &\mapsto \frac{1}{p}, \\ (13)K_4 &\mapsto (01) = \begin{pmatrix} -1 & 1 \\ 0 & 1 \end{pmatrix}, & p &\mapsto 1-p, \\ (123)K_4 &\mapsto (01\infty) = \begin{pmatrix} 0 & 1 \\ -1 & 1 \end{pmatrix}, & p &\mapsto \frac{p-1}{p}, \\ (132)K_4 &\mapsto (0\infty 1) = \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix}, & p &\mapsto \frac{1}{1-p}. \end{aligned}$$

Furthermore, $\sigma \cdot \xi = t_\sigma^{-1} \cdot \xi$ for any $\sigma \in S_4$.

Thus, the S_4 -action induces an action on $\text{Chart}(0, 1, \infty)$, i.e., $\sigma \cdot \chi(p) = \chi(p(\sigma^{-1} -)) = \chi(t_\sigma^{-1} \cdot p)$ for $\chi(p) \in \text{Chart}(0, 1, \infty)$ and $\sigma \in S_4$. Hence, we have:

Lemma 1.3. *The function spaces Cor_4 and Cor_4^f and $F_{0,1,\infty}$ are preserved by the action of S_4 .*

By the definition of a conformal singularity, for a chart $\chi(p) \in \text{Chart}(0, 1, \infty)$ and $f \in F_{0,1,\infty}$, we have the expansion of $f \circ \chi^{-1}(p)$ around $p = 0$, which defines a linear map $j(\chi(p), -) : F_{0,1,\infty} \rightarrow \mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}}))$, that is,

$$j(\chi(p), f) = \sum_{r,s \in \mathbb{R}} a_{r,s} p^r \bar{p}^s,$$

where the right-hand-side is compactly absolutely convergent to $f(\chi^{-1}(p))$ around $p = 0$.

For $t \in \text{Aut}(0, 1, \infty)$, $t \cdot p = t^{-1} \cdot p \in \text{Chart}(0, 1, \infty)$ is a chart at $t \cdot 0$ and the inverse $(t \cdot p)^{-1} : \mathbb{CP}^1 \rightarrow \mathbb{CP}^1$ is given by $t^{-1} \cdot p = t \cdot p$, e.g., $\frac{1}{1-p}$ is a chart at $\infty \in \mathbb{CP}^1$ whose inverse is $\frac{p-1}{p}$.

Lemma 1.4. *For $\sigma \in S_4$, $\chi(p) \in \text{Chart}(0, 1, \infty)$ and $f \in F_{0,1,\infty}$, $j(\chi(p), \sigma \cdot f) = j(\sigma^{-1} \cdot \chi(p), f)$.*

Proof. We may assume that $\chi(p) = t.p$ for some $t \in \text{Aut}(0, 1, \infty)$. The series $j(t.p, \sigma f)$ is absolutely convergent to $f(t_\sigma^{-1} t^{-1}.p)$ around $p = 0$. Since $f(t_\sigma^{-1} t^{-1}.p) = f((t t_\sigma)^{-1}.p)$, we have $j(t.p, \sigma f) = j(t t_\sigma.p, f) = j(\sigma^{-1} \cdot t.p, f)$. $\square$

The stabilizer subgroup of a point $0 \in \mathbb{C}P^1$ in S_4 (resp. $\text{Aut}(0, 1, \infty)$) is generated by the Klein subgroup K_4 and (12) (resp. $\{1, (0\infty)\}$), which is isomorphic to the dihedral group D_4 . For $\sigma \in K_4$, the action of σ on $F_{0,1,\infty}$ is trivial and for $\sigma \in (12)K_4$, the action of σ on $F_{0,1,\infty}$ is $(12)K_4 \cdot f(p) = f(\frac{p}{p-1})$. Define the map $\lim_{p \mapsto \frac{p}{p-1}} : \mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}})) \rightarrow \mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}}))$ by substituting $-\sum_{n \geq 1} p^n$ into p . Then, we have:

Lemma 1.5. *For $\sigma \in K_4$, $j(\sigma \cdot p, -) = j(p, -)$ and $j((12) \cdot p, -) = \lim_{p \mapsto \frac{p}{p-1}} j(p, -)$.*

1.3. Expansions. In this section, we consider expansions of a function in Cor_4 . By Lemma 1.1, a function $\phi \in \text{Cor}_4$ can be expanded in three variables, e.g., $z_2 - z_3, z_3 - z_4, z_1 - z_4$. Since ϕ is a multi-variable function, we need to determine the order of the expansion. For example, we expand ϕ around $z_2 = z_3$ first and then around $z_3 = z_4$, and $z_1 = z_4$. By setting $x = z_1 - z_4, y = z_3 - z_4, z = z_2 - z_3$, the resulting series is a formal power series of the variables $(x, \bar{x}, y, \bar{y}, z, \bar{z})$, which is absolutely convergent in $\{|x| > |y| > |z|\}$. All such series expansions are described by parenthesized products of four symbols. The above case corresponds to $1((23)4)$. The innermost product (23) means that we expand ϕ around $z_2 - z_3$ first. Let P_4 be the set of parenthesized products of four elements $1, 2, 3, 4$, e.g., $1(2(34))$. We remark that the permutation group S_4 naturally acts on P_4 and P_4 consists of the permutations of the following elements, called standard elements of P_4 :

$$\begin{array}{ll}
 & (12)(34), \\
 \text{(standard elements of } P_4) & ((12)3)4, \\
 & (1(23))4, \\
 & 1((23)4), \\
 & 1(2(34)).
 \end{array}$$

We introduce formal variables $(x_A, y_A, z_A, \bar{x}_A, \bar{y}_A, \bar{z}_A)$ for each $A \in P_4$. In this section, for each $A \in P_4$, we define a series expansion

$$e_A : \text{Cor}_4 \rightarrow T_A,$$

where T_A is a space of formal variables $(x_A, y_A, z_A, \bar{x}_A, \bar{y}_A, \bar{z}_A)$ defined below and show that the expansions are completely described by $j(\chi(p), -) : F_{0,1,\infty} \rightarrow \mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}}))$ for some $\chi(p) \in \text{Chart}(0, 1, \infty)$.

Remark 1.3. *In introduction 0.2, we consider the expansions in four variables associated with an element in Q_4 . Such expansions are described in Section 1.7 by using the results in this section.*

Set

$$\begin{aligned} T(x, y, z) &= \mathbb{C}[[y/x, \bar{y}/\bar{x}]][(z/y, \bar{z}/\bar{y}, |z/y|^{\mathbb{R}})] [x^{\pm}, y^{\pm}, \bar{x}^{\pm}, \bar{y}^{\pm}, |x|^{\mathbb{R}}, |y|^{\mathbb{R}}], \\ T^f(x, y, z) &= \mathbb{C}[[y/x, \bar{y}/\bar{x}, z/y, \bar{z}/\bar{y}]] [x^{\pm}, y^{\pm}, z^{\pm}, \bar{x}^{\pm}, \bar{y}^{\pm}, \bar{z}^{\pm}, |x|^{\mathbb{R}}, |y|^{\mathbb{R}}, |z|^{\mathbb{R}}], \end{aligned}$$

for the formal variables $x, y, z, \bar{x}, \bar{y}, \bar{z}$.

We first examine the case of $A = 1(2(34))$ in detail. In this case, we first expand a function $\phi \in \text{Cor}_4$ in the variable $z_{1(2(34))} = z_3 - z_4$ (related to the innermost product (34)), then in $y_{1(2(34))} = z_2 - z_4$ (related to $2(34)$) and $x_{1(2(34))} = z_1 - z_4$ (For the general rule for the change of variables, see Appendix). The resulting formal power series is convergent in some open region in $\{|x_{1(2(34))}| > |y_{1(2(34))}| > |z_{1(2(34))}|\}$, which define a linear map

$$e_{1(2(34))} : \text{Cor}_4 \rightarrow T(x_{1(2(34))}, y_{1(2(34))}, z_{1(2(34))})$$

such that for $\phi(z_1, z_2, z_3, z_4) \in \text{Cor}_4$, the formal power series $e_{1(2(34))}(\phi)$ is absolutely convergent to the function $f(z_1, z_2, z_3, z_4)$ by taking the change of variables $(z_1 - z_4, z_2 - z_4, z_3 - z_4) = (x_{1(2(34))}, y_{1(2(34))}, z_{1(2(34))})$.

For a function in $\text{Cor}_4^f \subset \text{Cor}_4$, the map $e_{1(2(34))}$ is defined as a binomial expansion in the domain $|x_{1(2(34))}| \gg |y_{1(2(34))}| \gg |z_{1(2(34))}|$. For example,

$$|z_1 - z_3|^r \mapsto |x_{1(2(34))} - z_{1(2(34))}|^r = |x_{1(2(34))}|^r \sum_{i,j \geq 0} (-1)^i \binom{r}{i} \binom{r}{j} x_{1(2(34))}^{-i} \bar{x}_{1(2(34))}^{-j} z_{1(2(34))}^i \bar{z}_{1(2(34))}^j,$$

where the right hand side is absolutely convergent to $|z_1 - z_3|^r$ in the open domain $|z_1 - z_4| > |z_3 - z_4|$ and is an element of $T^f(x_{1(2(34))}, y_{1(2(34))}, z_{1(2(34))})$. We denote this map by $e_{1(2(34))}^f : \text{Cor}_4^f \rightarrow T^f(x_{1(2(34))}, y_{1(2(34))}, z_{1(2(34))})$. Let $f \in F_{0,1,\infty}$. For $f \circ \xi \in \text{Cor}_4$, expand $\xi = \frac{(z_1 - z_2)(z_3 - z_4)}{(z_1 - z_3)(z_2 - z_4)}$ in $|x_{1(2(34))}| \gg |y_{1(2(34))}| \gg |z_{1(2(34))}|$, e.g.,

$$\begin{aligned} \xi &= \frac{(z_1 - z_2)(z_3 - z_4)}{(z_1 - z_3)(z_2 - z_4)} \mapsto \frac{(x_{1(2(34))} - y_{1(2(34))})z_{1(2(34))}}{y_{1(2(34))}(x_{1(2(34))} - z_{1(2(34))})} = z_{1(2(34))}/y_{1(2(34))}(1 - y_{1(2(34))}/x_{1(2(34))}) \\ &\quad \times \sum_{i \geq 0} (z_{1(2(34))}/x_{1(2(34))})^i, \end{aligned}$$

which is equal to $e_{1(2(34))}^f(\frac{(z_1 - z_2)(z_3 - z_4)}{(z_1 - z_3)(z_2 - z_4)})$, defined above. Define the map $s_{1(2(34))} : \mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}})) \rightarrow T(x_{1(2(34))}, y_{1(2(34))}, z_{1(2(34))})$ by substituting $p = e_{1(2(34))}^f(\frac{(z_1 - z_2)(z_3 - z_4)}{(z_1 - z_3)(z_2 - z_4)})$ into $\sum_{r,s \in \mathbb{R}} a_{r,s} p^r \bar{p}^s$. Since $p = e_{1(2(34))}^f(\frac{(z_1 - z_2)(z_3 - z_4)}{(z_1 - z_3)(z_2 - z_4)})$ is an element of

$$z_{1(2(34))}/y_{1(2(34))} \mathbb{C}[[y_{1(2(34))}/x_{1(2(34))}, z_{1(2(34))}/y_{1(2(34))}]],$$

by the definition of $\mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}}))$, $s_{1(2(34))}$ is well-defined. The following lemma is obvious:

Lemma 1.6. $T(x, y, z)$ is a module over the $\mathbb{C}$ -algebra $T^f(x, y, z)$.

Then, $e_{1(2(34))} : \text{Cor}_4 \rightarrow T(x_{1(2(34))}, y_{1(2(34))}, z_{1(2(34))})$ is defined by

$$e_{1(2(34))}(\phi \cdot f \circ \xi) = e_{1(2(34))}^f(\phi) s_{1(2(34))}(j(p, f))$$

for $\phi \in \text{Cor}_4^f$ and $f \in F_{0,1,\infty}$, which is absolutely convergent to $\phi \cdot f \circ \xi$ in $|x_{1(2(34))}| \gg |y_{1(2(34))}| \gg |z_{1(2(34))}|$. Here, $|x_{1(2(34))}| \gg |y_{1(2(34))}| \gg |w_{1(2(34))}|$ means some non-empty

open domain in $\{(z_1, z_2, z_3, z_4) \mid |z_1 - z_4| > |z_2 - z_4| > |z_3 - z_4|\}$. The map $e_{1((23)4)}$ is independent of a choice of the decomposition $\phi \cdot f \circ \xi \in \text{Cor}_4$, since the coefficient of a convergent series is unique.

Second, we give the definition of e_A in case of $A = 1((23)4)$. We consider the following change of variables;

$$(z_1 - z_4, z_3 - z_4, z_2 - z_3) \mapsto (x_{1((23)4)}, y_{1((23)4)}, z_{1((23)4)}).$$

The map $e_{1((23)4)}^f : \text{Cor}_4^f \rightarrow T^f(x_{1((23)4)}, y_{1((23)4)}, z_{1((23)4)})$ is defined by the binomial expansion in the domain $|x_{1((23)4)}| \gg |y_{1((23)4)}| \gg |z_{1((23)4)}|$ as above. The expansion of $\xi = \frac{(z_1 - z_2)(z_3 - z_4)}{(z_1 - z_3)(z_2 - z_4)}$ is equal to

$$\begin{aligned} \xi &= \frac{(x_{1((23)4)} - y_{1((23)4)} + z_{1((23)4)})y_{1((23)4)}}{(x_{1((23)4)} - y_{1((23)4)})(z_{1((23)4)} + y_{1((23)4)})} = (x_{1((23)4)} - y_{1((23)4)} + z_{1((23)4)})y_{1((23)4)} \\ &\quad \times \sum_{i,j \geq 0} (-1)^j x_{1((23)4)}^{-i-1} y_{1((23)4)}^{i-j-1} z_{1((23)4)}^j. \end{aligned}$$

Importantly, this is not in

$$z_{1((23)4)}/y_{1((23)4)} \mathbb{C}[[y_{1((23)4)}/x_{1((23)4)}, z_{1((23)4)}/y_{1((23)4)}]].$$

Thus, the substitution of this into $j(p, f)$ is not well-defined for $f \in F_{0,1,\infty}$. The reason is that when $z_{1((23)4)} = z_2 - z_3$ goes to zero, ξ goes to 1. Thus, we have to consider the expansion of f around 1 and the expansion of $1 - \xi = \frac{(z_1 - z_4)(z_2 - z_3)}{(z_1 - z_3)(z_2 - z_4)}$. Then, the expansion is

$$\begin{aligned} 1 - \xi &= \frac{x_{1((23)4)} z_{1((23)4)}}{(x_{1((23)4)} - y_{1((23)4)})(y_{1((23)4)} + z_{1((23)4)})} \\ &= z_{1((23)4)}/y_{1((23)4)} \sum_{i,j \geq 0} (-1)^j (y_{1((23)4)}/x_{1((23)4)})^i (z_{1((23)4)}/y_{1((23)4)})^j, \end{aligned}$$

which is an element of $z_{1((23)4)}/y_{1((23)4)} \mathbb{C}[[y_{1((23)4)}/x_{1((23)4)}, z_{1((23)4)}/y_{1((23)4)}]]$. Define the map

$$s_{1((23)4)} : \mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}})) \rightarrow T(x_{1((23)4)}, y_{1((23)4)}, z_{1((23)4)})$$

by substituting $p = e_{1((23)4)}^f(1 - \xi)$ into $\sum_{r,s \in \mathbb{R}} a_{r,s} p^r \bar{p}^s$. Then, $e_{1((23)4)} : \text{Cor}_4 \rightarrow T(x_{1((23)4)}, y_{1((23)4)}, z_{1((23)4)})$ is defined by

$$e_{1((23)4)}(\phi f \circ \xi) = e_{1((23)4)}^f(\phi) s_{1((23)4)}(j(1 - p, f))$$

for $\phi \in \text{Cor}_4^f$ and $f \in F_{0,1,\infty}$. Since $j(1 - p, f)$ is absolutely convergent to $f(1 - p)$, $s_{1((23)4)}(j(1 - p, f))$ is absolutely convergent to $f \circ \xi$ in $|x_{1((23)4)}| \gg |y_{1((23)4)}| \gg |z_{1((23)4)}|$.

Remark 1.4. In the domain $|x_{1((23)4)}| \gg |y_{1((23)4)}| \gg |z_{1((23)4)}|$, the difference $|z_{1((23)4)}| = |z_2 - z_3|$ is assumed to be small. Then, by Remark 1.1, $\xi \mapsto 1$. Thus, we have to consider the expansion of $1 - \xi$ or $1 - \xi^{-1}$. If we choose $1 - \xi^{-1}$, we substitute $e_{1((23)4)}^f(1 - \xi^{-1})$ into $j(\frac{1}{1-p}, f)$. Then, the resulting series is, roughly, convergent to $f(\frac{1}{1-(1-p^{-1})}) = f(p)$. Hence, both of the choices $1 - \xi$ and $1 - \xi^{-1}$ give the same result.

The last example is the case of $A = (12)(34)$, which has a special property. We consider the following change of variables:

$$(z_2 - z_4, z_1 - z_2, z_3 - z_4) \mapsto (x_{(12)(34)}, y_{(12)(34)}, z_{(12)(34)}).$$

Then,

$$z_1 - z_3 \mapsto x_{(12)(34)} + y_{(12)(34)} - z_{(12)(34)},$$

$$z_1 - z_4 \mapsto x_{(12)(34)} + y_{(12)(34)},$$

$$z_2 - z_3 \mapsto x_{(12)(34)} - z_{(12)(34)}.$$

Since there is no $y_{(12)(34)} - z_{(12)(34)}$ or $y_{(12)(34)} + z_{(12)(34)}$ in the above list, the image of $e_{(12)(34)}^f : \text{Cor}_4^f \rightarrow T^f(x_{(12)(34)}, y_{(12)(34)}, z_{(12)(34)})$ is in

$$\mathbb{C}[[y/x, z/x, \bar{y}/\bar{x}, \bar{z}/\bar{x}]] [x^\pm, y^\pm, z^\pm, \bar{x}^\pm, \bar{y}^\pm, \bar{z}^\pm, |x|^\mathbb{R}, |y|^\mathbb{R}, |z|^\mathbb{R}],$$

that is, both $y_{(12)(34)}$ and $z_{(12)(34)}$ are bounded below, which happens only for the S_4 -conjugate elements of $(12)(34)$. Set

$$T_A^f(x, y, z) = \mathbb{C}[[y/x, z/x, \bar{y}/\bar{x}, \bar{z}/\bar{x}]] [x^\pm, y^\pm, z^\pm, \bar{x}^\pm, \bar{y}^\pm, \bar{z}^\pm, |x|^\mathbb{R}, |y|^\mathbb{R}, |z|^\mathbb{R}],$$

for any $A \in P_4$ which is S_4 -conjugate of $(12)(34)$. Then, the binomial expansion defines $e_{(12)(34)} : \text{Cor}_4 \rightarrow T_{(12)(34)}^f(x_{(12)(34)}, y_{(12)(34)}, z_{(12)(34)})$ and

$$\begin{aligned} e_{(12)(34)}(\xi) &= \frac{y_{(12)(34)} z_{(12)(34)}}{x_{(12)(34)} (y_{(12)(34)} + x_{(12)(34)} - z_{(12)(34)})} \\ &= \frac{z_{(12)(34)}}{x_{(12)(34)}} \frac{y_{(12)(34)}}{x_{(12)(34)}} \sum_{n=0}^{\infty} \sum_{i=0}^n (-1)^{n-i} \binom{n}{i} \left(\frac{y_{(12)(34)}}{x_{(12)(34)}} \right)^{n-i} \left(\frac{z_{(12)(34)}}{x_{(12)(34)}} \right)^i. \end{aligned}$$

Let $T'(x, y, z)$ be a subspace of $\mathbb{C}((y/x, z/x, \bar{y}/\bar{x}, \bar{z}/\bar{x}, |y/x|^\mathbb{R}, |z/x|^\mathbb{R}))$ spanned by

$$\sum_{s_1, s_2, \bar{s}_1, \bar{s}_2 \in \mathbb{R}} c_{s_1, \bar{s}_1, s_2, \bar{s}_2} \left(\frac{y}{x} \right)^{s_1} \left(\frac{\bar{y}}{\bar{x}} \right)^{\bar{s}_1} \left(\frac{z}{x} \right)^{s_2} \left(\frac{\bar{z}}{\bar{x}} \right)^{\bar{s}_2}$$

such that:

- (1) $c_{s_1, \bar{s}_1, s_2, \bar{s}_2} = 0$ unless $s_1 - \bar{s}_1, s_2 - \bar{s}_2 \in \mathbb{Z}$;
- (2) There exists $N \in \mathbb{R}$ such that $c_{s_1, \bar{s}_1, s_2, \bar{s}_2} = 0$ unless $s_1, \bar{s}_1, s_2, \bar{s}_2 \geq N$;
- (3) $c_{s_1, \bar{s}_1, s_2, \bar{s}_2} = 0$ unless $s_1 - s_2, \bar{s}_1 - \bar{s}_2 \in \mathbb{Z}$.

Set

$$T_A(x, y, z) = T'(x, y, z) [x^\pm, y^\pm, z^\pm, \bar{x}^\pm, \bar{y}^\pm, \bar{z}^\pm, |x|^\mathbb{R}, |y|^\mathbb{R}, |z|^\mathbb{R}],$$

for any $A \in P_4$ which is S_4 -conjugate to $(12)(34)$. Define the map

$$s_{(12)(34)} : \mathbb{C}((p, \bar{p}, |p|^\mathbb{R})) \rightarrow T_{(12)(34)}(x_{(12)(34)}, y_{(12)(34)}, z_{(12)(34)})$$

by substituting $p = e_{(12)(34)}^f(\xi)$ into $\sum_{r,s \in \mathbb{R}} a_{r,s} p^r \bar{p}^s$. The following lemma is obvious:

Lemma 1.7. *For any $A \in P_4$ which is S_4 -conjugate of $(12)(34)$, $T_A^f(x, y, z)$ is a subspace of $T_A(x, y, z)$ and $T_A(x, y, z)$ is a module over the $\mathbb{C}$ -algebra $T_A^f(x, y, z)$.*

Then, $e_{(12)(34)} : \text{Cor}_4 \rightarrow T_{(12)(34)}(x_{(12)(34)}, y_{(12)(34)}, z_{(12)(34)})$ can be defined as above.

Now, we will consider the general case. For standard elements of P_4 , consider the following change of variables:

$$\begin{aligned}
(12)(34) &: (x_{(12)(34)}, y_{(12)(34)}, z_{(12)(34)}) = (z_2 - z_4, z_1 - z_2, z_3 - z_4), \\
((12)3)4 &: (x_{((12)3)4}, y_{((12)3)4}, z_{((12)3)4}) = (z_3 - z_4, z_2 - z_3, z_1 - z_2), \\
(1(23))4 &: (x_{(1(23))4}, y_{(1(23))4}, z_{(1(23))4}) = (z_3 - z_4, z_1 - z_3, z_2 - z_3), \\
1((23)4) &: (x_{1((23)4)}, y_{1((23)4)}, z_{1((23)4)}) = (z_1 - z_4, z_3 - z_4, z_2 - z_3), \\
1(2(34)) &: (x_{1(2(34))}, y_{1(2(34))}, z_{1(2(34))}) = (z_1 - z_4, z_2 - z_4, z_3 - z_4).
\end{aligned}$$

For general $A \in P_4$, regard the above x_A, y_A, z_A as polynomials of (z_1, z_2, z_3, z_4) . Then, S_4 naturally acts on them. For a standard element A and $\sigma \in S_4$, set $(x_{\sigma A}, y_{\sigma A}, z_{\sigma A}) = (\sigma x_A, \sigma y_A, \sigma z_A)$. For example,

$$(x_{2((43)1)}, y_{2((43)1)}, z_{2((43)1)}) = (124) \cdot (z_1 - z_4, z_3 - z_4, z_2 - z_3) = (z_2 - z_1, z_3 - z_1, z_4 - z_3).$$

For each $A \in P_4$ which is not S_4 -conjugate to $(12)(34)$, set $T_A(x_A, y_A, z_A) = T(x_A, y_A, z_A)$ and $T_A^f(x_A, y_A, z_A) = T^f(x_A, y_A, z_A)$. Then, we have the binomial expansion convergent in $|x_A| \gg |y_A| \gg |z_A|$, $e_A^f : \text{Cor}_4^f \rightarrow T_A^f(x_A, y_A, z_A)$ as above. Define the map $T_\sigma : T(x_A, y_A, z_A) \rightarrow T(x_{\sigma A}, y_{\sigma A}, z_{\sigma A})$ by $(x_A, y_A, z_A, \bar{x}_A, \bar{y}_A, \bar{z}_A) \mapsto (x_{\sigma A}, y_{\sigma A}, z_{\sigma A}, \bar{x}_{\sigma A}, \bar{y}_{\sigma A}, \bar{z}_{\sigma A})$. Then, we have:

Lemma 1.8. *For any $\sigma \in S_4$ and $A \in P_4$ and $\phi \in \text{Cor}_4^f$, $T_\sigma \circ e_A(\phi) = e_{\sigma A}(\sigma\phi)$.*

By Remark 1.4, we need to choose an appropriate choice of $\xi, \frac{\xi}{\xi-1}, 1-\xi, 1-\xi^{-1}, \xi^{-1}, \frac{1}{1-\xi}$ for each $A \in P_4$, when we expand a function $f \in F_{0,1,\infty}$. The map $\tau : P_4 \rightarrow S_4$ is chosen for standard elements as

$$\begin{aligned}
(12)(34) &\mapsto 1, \\
((12)3)4 &\mapsto 1, \\
(1(23))4 &\mapsto (13), \\
1((23)4) &\mapsto (13), \\
1(2(34)) &\mapsto 1,
\end{aligned}$$

and for $\sigma A \in P_4$ as $\tau(\sigma A) = \sigma \cdot \tau(A)$, where $\sigma \in S_4$ and A is a standard element of P_4 . Then, for standard element A , the expansion of $\tau(A) \cdot \xi$ is given by:

$$\begin{aligned}\xi &= \frac{y_{(12)(34)}z_{(12)(34)}}{x_{(12)(34)}(y_{(12)(34)} + x_{(12)(34)} - z_{(12)(34)})} = z_{(12)(34)}y_{(12)(34)} \sum_{n \geq 0} \sum_{i=0}^n (-1)^{n-i} \binom{n}{i} x_{(12)(34)}^{-n-2} y_{(12)(34)}^{n-i} z_{(12)(34)}^i, \\ \xi &= \frac{x_{((12)3)4}z_{((12)3)4}}{(z_{((12)3)4} + y_{((12)3)4})(y_{((12)3)4} + x_{((12)3)4})} = z_{((12)3)4}/y_{(12)(34)} \sum_{i,j \geq 0} (-1)^{i+j} x_{(12)(34)}^{-i} y_{(12)(34)}^{i-j} z_{(12)(34)}^j, \\ 1 - \xi &= \frac{(x_{(1(23))4} + y_{(1(23))4})z_{(1(23))4}}{y_{(1(23))4}(x_{(1(23))4} + z_{(1(23))4})} = z_{(1(23))4}/y_{(1(23))4} (1 - y_{(1(23))4}/x_{(1(23))4}) \sum_{i \geq 0} (-1)^i x_{(1(23))4}^{-i} z_{(1(23))4}^i, \\ 1 - \xi &= \frac{x_{1((23)4)}z_{1((23)4)}}{(x_{1((23)4)} - y_{1((23)4)})(y_{1((23)4)} + z_{1((23)4)})} = z_{1((23)4)}/y_{1((23)4)} \sum_{i,j \geq 0} (-1)^j x_{1((23)4)}^{-i} y_{1((23)4)}^{i-j} z_{1((23)4)}^j, \\ \xi &= \frac{(x_{1(2(34))} - y_{1(2(34))})z_{1(2(34))}}{y_{1(2(34))}(x_{1(2(34))} - z_{1(2(34))})} = z_{1(2(34))}/y_{1(2(34))} (1 - y_{1(2(34))}/x_{1(2(34))}) \sum_{i \geq 0} x_{1(2(34))}^{-i} z_{1(2(34))}^i.\end{aligned}$$

Thus, for $\sigma \in S_4$, by Lemma 1.8, $e_{\sigma A}^f(\tau(\sigma A) \cdot \xi) = e_{\sigma A}^f(\sigma \tau(A) \cdot \xi) = T_\sigma e_A^f(\tau(A) \cdot \xi)$, which implies that $e_{\sigma A}^f(\tau(\sigma A) \cdot \xi) \in z_{\sigma A}/x_{\sigma A}y_{\sigma A}/x_{\sigma A} \mathbb{C}[[y_{\sigma A}/x_{\sigma A}, z_{\sigma A}/x_{\sigma A}]]$ if A is S_4 -conjugate to $(12)(34)$ and $e_{\sigma A}^f(\tau(\sigma A) \cdot \xi) \in z_{\sigma A}/y_{\sigma A} \mathbb{C}[[y_{\sigma A}/x_{\sigma A}, z_{\sigma A}/y_{\sigma A}]]$ otherwise.

Define the map

$$s_A : \mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}})) \rightarrow T_A(x_A, y_A, w_A)$$

by substituting $p = e_A^f(\tau(A) \cdot \xi)$ into $\sum_{r,s \in \mathbb{R}} a_{r,s} p^r \bar{p}^s$. Then, the composition of $j(\tau(A) \cdot p, -) : F_{0,1,\infty} \rightarrow \mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}}))$ and s_A define the map $e_A : \text{Cor}_4 \rightarrow T_A(x_A, y_A, z_A)$. Then, we have:

Proposition 1.2. *For $A \in P_4$ and $\phi \in \text{Cor}_4$, $e_A(\phi) \in T_A(x_A, y_A, z_A)$ is absolutely convergent to ϕ in some non-empty domain in $|x_A| > |y_A| > |z_A|$ and $T_\sigma e_A(\phi) = e_{\sigma A}(\sigma \cdot \phi)$.*

Proof. Let $f \in F_{0,1,\infty}$. Then, $e_A(f \circ \xi) = s_A(j(\tau(A) \cdot p, f)) = s_A(j(t_{\tau(A)}^{-1} \cdot p, f))$ is, by Lemma 1.2, absolutely convergent to

$$f(t_{\tau(A)} \cdot (\tau(A) \cdot \xi)) = f(t_{\tau(A)} \cdot t_{\tau(A)}^{-1} \cdot \xi) = f(\xi).$$

Furthermore, for $\sigma \in S_4$, since $T_\sigma e_A^f(\tau(A) \cdot \xi) = e_{\sigma A} f(\tau(\sigma A) \cdot \xi)$, by Lemma 1.4, we have

$$e_{\sigma A}(\sigma \cdot f \circ \xi) = s_{\sigma A}(j(\sigma \cdot \tau(A) \cdot p, \sigma \cdot f)) = T_\sigma s_A j(\tau(A) \cdot p, f) = T_\sigma e_A(f \circ \xi)$$

□

The symbols e_A and x_A, y_A, z_A is convenient to express the various expansions systematically. However, in practice it is cumbersome to work in this notation. For $A = (1(23))4$, x_A, y_A, z_A represents $z_3 - z_4, z_1 - z_3, z_2 - z_3$. Thus, we introduce new formal variables z_{34}, z_{13}, z_{23} and set $z_{34} = x_{(1(23))4}, z_{13} = y_{(1(23))4}, z_{23} = z_{(1(23))4}$. Then, the map e_A is a map from Cor_4 to $T(z_{34}, z_{13}, z_{23})$. For, $\phi \in \text{Cor}_4$, we also denote $e_{(1(23))4}(\phi)$ by $\phi|_{|z_{34}| > |z_{13}| > |z_{23}|}$. The notations x_A, y_A, z_A and z_{ij} will be used interchangeably in this paper.

We remark that we can define the expansion maps $\text{Cor}_2, \text{Cor}_3$ in similar way, e.g., $\text{Cor}_3 \rightarrow T(z_{13}, z_{23}), \phi \mapsto \phi|_{|z_{13}| > |z_{23}|}$.

1.4. Relations among expansions. In this section, we establish relations among expansions e_A for $A \in P_4$. For $A, B \in P_4$, define the map

$$T'^B_A : T(x_A, y_A, z_A) \rightarrow T(x_B, y_B, z_B)$$

by sending $(x_A, y_A, z_A, \bar{x}_A, \bar{y}_A, \bar{z}_A)$ into $(x_B, y_B, z_B, \bar{x}_B, \bar{y}_B, \bar{z}_B)$.

We first construct a map $T^{1(4(23))}_{1((23)4)} : T(x_{1((23)4)}, y_{1((23)4)}, z_{1((23)4)}) \rightarrow T(x_{1(4(23))}, y_{1(4(23))}, z_{1(4(23))})$ such that $T^{1(4(23))}_{1((23)4)} \circ e_{1((23)4)} = e_{1(4(23))}$. By definition, there should be a relation

$$x_{1(4(23))} = z_1 - z_3 = x_{1((23)4)} - y_{1((23)4)},$$

$$y_{1(4(23))} = z_4 - z_3 = -y_{1((23)4)},$$

$$z_{1(4(23))} = z_2 - z_3 = z_{1((23)4)}.$$

Thus, define $T^{1(4(23))}_{1((23)4)} : T(x_{1((23)4)}, y_{1((23)4)}, z_{1((23)4)}) \rightarrow T(x_{1(4(23))}, y_{1(4(23))}, z_{1(4(23))})$ by

$$T^{1(4(23))}_{1((23)4)} = T'^{1(4(23))}_{1((23)4)} \exp(-y_{1((23)4)} \frac{d}{dx_{1((23)4)}}) \exp(-\bar{y}_{1((23)4)} \frac{d}{d\bar{x}_{1((23)4}}) \lim_{y_{1((23)4)} \mapsto -y_{1((23)4)}}.$$

We remark that the operator $\exp(y \frac{d}{dx}) : \mathbb{C}((y/x)) \rightarrow \mathbb{C}((y/x))$ changes the variable x into $x + y$, that is,

$$\exp(y \frac{d}{dx}) \sum_n a_n x^{-n} y^n = \sum_n \sum_{i \geq 0} \binom{-n}{i} a_n x^{-n-i} y^{n+i}.$$

For $\sigma \in S_4$, define $T^{\sigma 1(4(23))}_{\sigma 1((23)4)} : T(x_{\sigma 1((23)4)}, y_{\sigma 1((23)4)}, z_{\sigma 1((23)4)}) \rightarrow T(x_{\sigma 1(4(23))}, y_{\sigma 1(4(23))}, z_{\sigma 1(4(23))})$ by

$$T^{\sigma 1(4(23))}_{\sigma 1((23)4)} = T_\sigma \circ T^{1(4(23))}_{1((23)4)} \circ T_{\sigma^{-1}}.$$

Then, we have:

Lemma 1.9. For any $\sigma \in S_4$, $T^{\sigma 1(4(23))}_{\sigma 1((23)4)} \circ e_{\sigma 1((23)4)} = e_{\sigma 1(4(23))}$.

Second, we construct $T^{4(1(23))}_{(1(23)4)} : T(x_{(1(23)4)}, y_{(1(23)4)}, z_{(1(23)4)}) \rightarrow T(x_{4(1(23))}, y_{4(1(23))}, z_{4(1(23))})$ and $T^{4(3(12))}_{((12)3)4} : T(x_{((12)3)4}, y_{((12)3)4}, z_{((12)3)4}) \rightarrow T(x_{4(3(12))}, y_{4(3(12))}, z_{4(3(12))})$. Similarly to the above, by the relation

$$x_{4(1(23))} = z_4 - z_3 = -x_{(1(23)4)},$$

$$y_{4(1(23))} = z_1 - z_3 = y_{(1(23)4)},$$

$$z_{4(1(23))} = z_2 - z_3 = z_{(1(23)4)},$$

and

$$x_{4(3(12))} = z_4 - z_2 = -x_{((12)3)4} - y_{((12)3)4},$$

$$y_{4(3(12))} = z_3 - z_2 = -y_{((12)3)4},$$

$$z_{4(3(12))} = z_1 - z_2 = z_{((12)3)4}.$$

Thus, for $\sigma \in S_4$, set

$$T_{(1(23))4}^{4(1(23))} = T_{(1(23))4}'^{4(1(23))} \lim_{x_{(1(23))4} \rightarrow -x_{(1(23))4}}$$

$$T_{((12)3)4}^{4(3(12))} = T_{((12)3)4}'^{4(3(12))} \lim_{(x_{((12)3)4}, y_{((12)3)4}) \rightarrow (-x_{((12)3)4}, -y_{((12)3)4})} \exp y_{((12)3)4} \frac{d}{dx_{((12)3)4}} \exp \bar{y}_{((12)3)4} \frac{d}{d\bar{x}_{((12)3)4}}$$

and

$$T_{\sigma(1(23))4}^{\sigma 4(1(23))} = T_{\sigma} \circ T_{(1(23))4}^{4(1(23))} \circ T_{\sigma^{-1}},$$

$$T_{\sigma((12)3)4}^{\sigma 4(3(12))} = T_{\sigma} \circ T_{((12)3)4}^{4(3(12))} \circ T_{\sigma^{-1}}.$$

Then, we have

Lemma 1.10. *For any $\sigma \in S_4$, $T_{\sigma(1(23))4}^{\sigma 4(1(23))} \circ e_{\sigma(1(23))4} = e_{\sigma 4(1(23))}$ and $T_{\sigma((12)3)4}^{\sigma 4(3(12))} \circ e_{\sigma((12)3)4} = e_{\sigma 4(3(12))}$.*

By the above lemmas, all expansions except for $e_{(12)(34)}$ can be transformed into $e_{1(2(34))}$. For $e_{(12)(34)}$, the relation is

$$x_{2(1(34))} = z_2 - z_4 = x_{(12)(34)},$$

$$y_{2(1(34))} = z_1 - z_4 = y_{(12)(34)} + x_{(12)(34)},$$

$$z_{2(1(34))} = z_3 - z_4 = z_{(12)(34)}.$$

It seems that we only need to consider the operator $\exp(-x_{(12)(34)} \frac{d}{dy_{(12)(34)}})$. However, it is not well-defined, since, informally, $\exp(x \frac{d}{dy})(x - y)^{-1} = -y$ and $\exp(x \frac{d}{dy}) \sum_{n \geq 0} x^{-1-n} y^n = \sum_{n \geq 0} \sum_{k=0}^n x^{-1-n} (y + x)^n$ whose coefficient of x^{-1} is $1 + 1 + \dots$. We will discuss this issue in Section 1.5.

Finally, we transform $e_{1(2(34))}$ into $e_{1(2(43))}$. In this case, the relation is

$$x_{1(2(34))} = z_1 - z_4 = x_{1(2(43))} - z_{1(2(43))},$$

$$y_{1(2(34))} = z_2 - z_4 = y_{1(2(43))} - z_{1(2(43))},$$

$$z_{1(2(34))} = z_3 - z_4 = -z_{1(2(43))}.$$

Set

$$T_{1(2(34))}^{1(2(43))} = T_{1(2(34))}'^{1(2(43))} \lim_{z \rightarrow -z} \exp(z(\frac{d}{dy} + \frac{d}{dx})) \exp(\bar{z}(\frac{d}{d\bar{y}} + \frac{d}{d\bar{x}})).$$

Then,

Lemma 1.11. $T_{1(2(34))}^{1(2(43))} e_{1(2(34))} = e_{1(2(43))}$.

Remark 1.5. *We remark that the existence of this transformation reflects the fact that both $e_{1(2(34))}$ and $e_{1(2(43))}$ correspond to the expansion of a function in $F_{0,1,\infty}$ at the same point, $0 \in \mathbb{CP}^1$. The Borchers identity, an axiom of a vertex algebra, can be derived from a property of holomorphic functions, the Cauchy integral formula, which relate the expansions at the different points. However, for a real analytic function in $F_{0,1,\infty}$, it seems difficult to compare the expansions at the different points.*

1.5. Formal Differential Equation and Convergence. Since $\xi : X_4 \rightarrow \mathbb{C}P^1 \setminus \{0, 1, \infty\}$ is $\mathrm{PSL}_2\mathbb{C}$ invariant, the image of an expansion e_A satisfies differential equations. We first consider the formal differential equations for $e_{1(2(34))}$. We abbreviate $(x_{1(2(34))}, y_{1(2(34))}, z_{1(2(34))})$ to (x, y, z) . Set

$$\begin{aligned} D_1 &= x^2 d/dx + y^2 d/dy + z^2 d/dz, \\ \bar{D}_1 &= \bar{x}^2 d/d\bar{x} + \bar{y}^2 d/d\bar{y} + \bar{z}^2 d/d\bar{z}, \\ D_0 &= x d/dx + y d/dy + z d/dz, \\ \bar{D}_0 &= \bar{x} d/d\bar{x} + \bar{y} d/d\bar{y} + \bar{z} d/d\bar{z}, \end{aligned}$$

which acts on $T(x, y, z)$ and Set $S(x, y, z) = \cap_{i \in \{0, 1\}} (\ker D_i \cap \ker \bar{D}_i)$, which is the space of the formal solutions of the differential equations. The following lemma follows from the $\mathrm{PSL}_2\mathbb{C}$ -invariance of ξ :

Lemma 1.12. *The image of $s_{1(2(34))} : \mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}})) \rightarrow T(x, y, z)$ is in $S(x, y, z)$.*

We will show that $s_{1(2(34))} : \mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}})) \rightarrow S(x, y, z)$ is an isomorphism of vector spaces. Let $f(x, y, z) \in S(x, y, z)$. Since $D_0 f = 0$ and $\bar{D}_0 f = 0$, we can assume that

$$\begin{aligned} f(x, y, z) &= \sum_{n, m, r, s \in \mathbb{R}} a_{n, m, r, s} (y/x)^n (z/y)^m (\bar{y}/\bar{x})^r (\bar{z}/\bar{y})^s \\ &\in \mathbb{C}[[y/x, \bar{y}/\bar{x}]][(z/x, \bar{z}/\bar{x}, |z/x|^{\mathbb{R}})][(y/x)^{\pm}, (\bar{y}/\bar{x})^{\pm}, |y/x|^{\mathbb{R}}]. \end{aligned}$$

By $D_1 f = 0$ and $\bar{D}_1 f = 0$, we have

$$\begin{aligned} (n - m)a_{n, m, r, s} + (m - 1)a_{n, m-1, r, s} - (n + 1)a_{n+1, m, r, s} &= 0, \\ (r - s)a_{n, m, r, s} + (s - 1)a_{n, m, r, s-1} - (r + 1)a_{n, m, r+1, s} &= 0. \end{aligned}$$

By the definition of $T(x, y, z)$, $a_{n, m, r, s} = 0$ if one of n, m, r, s is sufficiently small. Fix $r, s \in \mathbb{R}$ and let $n_0 = \min\{n \mid a_{n, m, r, s} \neq 0 \text{ and } m \in \mathbb{R}\}$. Then, since $n_0 a_{n_0, m, r, s} = (n_0 - 1 - m)a_{n_0-1, m, r, s} + (m - 1)a_{n_0-1, m-1, r, s} = 0$, we have $n_0 = 0$. If there exists $n, m, r, s \in \mathbb{R}$ such that $n \in \mathbb{R} \setminus \mathbb{Z}$ and $a_{n, m, r, s} \neq 0$, then either $a_{n-1, m, r, s}$ or $a_{n-1, m-1, r, s}$ is non-zero, which contradicts the fact that $a_{n, m, r, s} = 0$ for sufficiently small n . Thus, $f(x, y, z) = \sum_{n, r \in \mathbb{Z}_{\geq 0}} \sum_{m, s \in \mathbb{R}} a_{n, m, r, s} (y/x)^n (z/y)^m (\bar{y}/\bar{x})^r (\bar{z}/\bar{y})^s$. Define the map $v_{1(2(34))} : S(x, y, z) \rightarrow \mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}}))$ by

$$f(x, y, z) \mapsto \sum_{m, s \in \mathbb{R}} a_{0, m, 0, s} p^m \bar{p}^s,$$

which is a formal limit of $(x, y, z) \mapsto (\infty, 1, p)$.

Lemma 1.13. *The map $v_{1(2(34))}$ is inverse to $s_{1(2(34))}$.*

Proof. Recall that $e_{1(2(34))}\xi = (1-y/x)z/y \sum_{m \geq 0} (z/x)^m$, which implies that $v_{1(2(34))}(s_{1(2(34))}(p)) = p$. Thus, $v_{1(2(34))}$ is a left inverse of $s_{1(2(34))}$. Hence, it suffices to show that $v_{1(2(34))}$ is injective. Let $f(x, y, z) = \sum_{n, r \in \mathbb{Z}_{\geq 0}} \sum_{m, s \in \mathbb{R}} a_{n, m, r, s} (y/x)^n (z/y)^m (\bar{y}/\bar{x})^r (\bar{z}/\bar{y})^s \in S(x, y, z)$ satisfy

$v_{1(2(34))}(f(x, y, z)) = 0$, that is, $a_{0,m,0,r} = 0$ for any $m, r \in \mathbb{R}$. Then, it is easy to show that $f = 0$ by the above recurrence formula. $\square$

Corollary 1.1. *Let $g \in S(x, y, z)$. If there exists $f \in F_{0,1,\infty}$ such that $v_{1(2(34))}(g) = j(p, f)$, then $g = e_{1(2(34))}(f \circ \xi)$. In particular, g is absolutely convergent to a function in Cor_4 in $|x| \gg |y| \gg |z|$.*

We second consider the formal differential equations for $e_{(12)(34)}$. We abbreviate $(x_{(12)(34)}, y_{(12)(34)}, z_{(12)(34)})$ to (x, y, z) again. Set

$$D'_1 = x^2 d/dx + y^2 d/dy + z^2 d/dz,$$

$$\bar{D}'_1 = \bar{x}^2 d/d\bar{x} + \bar{y}^2 d/d\bar{y} + \bar{z}^2 d/d\bar{z},$$

$$D'_0 = xd/dx + yd/dy + zd/dz,$$

$$\bar{D}'_0 = \bar{x}d/d\bar{x} + \bar{y}d/d\bar{y} + \bar{z}d/d\bar{z},$$

and let $S'(x, y, z)$ be the space of the formal solutions of $D'_0, D'_1, \bar{D}'_0, \bar{D}'_1$ in $T_{(12)(34)}(x, y, z)$. Similarly to the above, we have

$$s_{(12)(34)} : \mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}})) \rightarrow S'(x, y, z).$$

We will show that the map is an isomorphism of vector spaces by constructing the inverse map. By D'_0 and $\bar{D}'_0$, any element of $S'(x, y, z)$ is of the form $\sum_{n,m,r,s \in \mathbb{R}} a_{n,m,r,s} (y/x)^n (z/x)^m (\bar{y}/\bar{x})^r (\bar{z}/\bar{x})^s$. The map $v_{(12)(34)} : S'(x, y, z) \rightarrow \mathbb{C}((p^{1/2}, \bar{p}^{1/2}, |p|^{\mathbb{R}}))$ defined by

$$\sum_{n,m,r,s} a_{n,m,r,s} (y/x)^n (z/x)^m (\bar{y}/\bar{x})^r (\bar{z}/\bar{x})^s \mapsto \sum_{n,m,r,s} a_{n,m,r,s} p^{\frac{n+m}{2}} \bar{p}^{\frac{r+s}{2}}$$

is well-defined by the definition of $T_{(12)(34)}(x, y, z)$. Furthermore, since $e_{(12)(34)}(\xi) = (y/x)(z/x)(1 + (z/x - y/z) + (z/x - y/z)^2 + \dots)$, we have $v_{(12)(34)} \circ s_{(12)(34)}(\xi) = p$, which implies that $e_{(12)(34)}$ is injective. Thus, it suffices to show that $e_{(12)(34)}$ is surjective. Let $g(x, y, z) = \sum_{n,m,r,s} a_{n,m,r,s} (y/x)^n (z/x)^m (\bar{y}/\bar{x})^r (\bar{z}/\bar{x})^s \in S'(x, y, z)$. Since it is in the kernel of D'_1 and $\bar{D}'_1$,

$$(n - m)a_{n,m,r,s} + na_{n-1,m,r,s} + ma_{n,m-1,r,s} = 0,$$

$$(r - s)a_{n,m,r,s} + ra_{n,m,r-1,s} + sa_{n,m,r,s-1} = 0.$$

Lemma 1.14. *For non-zero $g(x, y, z) \in S(x, y, z)$. Set $N_0 = \min\{\frac{n+m}{2} \mid a_{n,m,r,s} \neq 0 \text{ for some } r, s \in \mathbb{R}\}$. and $R_0 = \min\{\frac{r+s}{2} \mid a_{n,2N_0-n,r,s} \neq 0 \text{ for some } n, r, s \in \mathbb{R}\}$. Then, $a_{N_0,N_0,R_0,R_0} \neq 0$ and $a_{n,2N_0-n,r,2R_0-r} = 0$ for any $n \neq N_0$ or $r \neq R_0$. Furthermore, $N_0 - R_0 \in \mathbb{Z}$.*

Proof. Let $n_0, r_0 \in \mathbb{R}$ satisfy $a_{n_0,2N_0-n_0,r_0,2R_0-r_0} \neq 0$. Then, by the above recurrence formula, $2(N_0 - n_0)a_{n_0,2N_0-n_0,r,2R_0-r_0} = n_0a_{n_0-1,2N_0-n_0,r_0,2R_0-r_0} + (2N_0 - n_0)a_{n_0,2N_0-n_0-1,r_0,2R_0-r_0} = 0$. Thus, $n_0 = N_0$ and $r_0 = R_0$. Since $g(x, y, z) \in T_{(12)(34)}(x, y, z)$, $a_{n,m,r,s} = 0$ if $n - r \notin \mathbb{Z}$ or $m - s \notin \mathbb{Z}$, which implies that $N_0 - R_0 \in \mathbb{Z}$. $\square$

By the above lemma, $p^{N_0} \bar{p}^{R_0} \in \mathbb{C}[p, \bar{p}, |p|^{\mathbb{R}}]$. Thus, we can replace $g(x, y, z)$ by $g(x, y, z) - e_{(12)(34)}(a_{N_0,N_0,R_0,R_0} p^{N_0} \bar{p}^{R_0})$, that is, $g(x, y, z) - e_{(12)(34)}(a_{N_0,N_0,R_0,R_0} p^{N_0} \bar{p}^{R_0}) \in S'(x, y, z)$. We repeat this process for R_0 and N_0 . Thus, we have $g(x, y, z) \in \text{Im}e_{(12)(34)}$, which implies that

$e_{(12)(34)}$ is surjective and the image of $v_{(12)(34)}$ is in $\mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}})) \subset \mathbb{C}((p^{1/2}, \bar{p}^{1/2}, |p|^{\mathbb{R}}))$. Hence, we have:

Lemma 1.15. *The map $v_{(12)(34)} : S'(x, y, z) \rightarrow \mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}}))$ is the inverse of $s_{(12)(34)} : \mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}})) \rightarrow S'(x, y, z)$.*

Corollary 1.2. *Let $g' \in S'(x, y, z)$. If there exists $f \in F_{0,1,\infty}$ such that $v_{(12)(34)}(g') = j(p, f)$, then $g' = e_{(12)(34)}(f \circ \xi)$. In particular, g' is absolutely convergent to a function in Cor_4 in $|x| \gg |y| \gg |z|$.*

Finally, we state a similar result on Cor_3 . Set

$$\begin{aligned} D_1'' &= x^2 d/dx + y^2 d/dy, \\ \bar{D}_1'' &= \bar{x}^2 d/d\bar{x} + \bar{y}^2 d/d\bar{y} \\ D_0'' &= x d/dx + y d/dy, \\ \bar{D}_0'' &= \bar{x} d/d\bar{x} + \bar{y} d/d\bar{y}, \end{aligned}$$

and let $S''(x, y)$ be a space of the formal solutions of $D_0', D_1', \bar{D}_0', \bar{D}_1'$ in

$$T(x, y) = \mathbb{C}[[y/x, \bar{y}/\bar{x}][x^\pm, \bar{x}^\pm, |x|, y^\pm, \bar{y}^\pm, |y|].$$

Then, we have:

Lemma 1.16. *The space $S''(x, y)$ consists of the constant functions.*

Proof. Let $f \in S''(x, y)$. By $D_0'' f = \bar{D}_0'' f = 0$, we can assume that $f(x, y) = \sum_{r,s \in \mathbb{R}} a_{r,s} (y/x)^r (\bar{y}/\bar{x})^s$. By $D_1'' f = \bar{D}_1'' f = 0$, $a_{n,\bar{n}}$ satisfies $ra_{r,s} = (r+1)a_{r+1,s}$ and $sa_{r,s} = (s+1)a_{r,s+1}$. Thus, $a_{r,s} = 0$ unless $(r, s) = (0, 0)$. $\square$

1.6. Holomorphic correlation functions. In this section, we consider holomorphic correlation functions. The following proposition is important:

Proposition 1.3. *If $d/d\bar{z}_a \phi = 0$ for $\phi \in \text{Cor}_4$ and $a \in \{1, 2, 3, 4\}$, then ϕ is in Cor_4^f . Furthermore, ϕ is a finite sum of functions $\prod_{1 \leq i < j \leq 4} (z_i - z_j)^{\alpha_{ij}} (\bar{z}_i - \bar{z}_j)^{\beta_{ij}}$ such that $\alpha_{ij} - \beta_{ij} \in \mathbb{Z}$ and $\beta_{ai} = 0$ for $i, j = 1, 2, 3, 4$.*

Proof. By the S_4 -symmetry, we may assume that $a = 2$. By the definition of Cor_4 , we can assume that $\phi(z_1, z_2, z_3, z_4) = \sum_{t=1}^N \phi^t(z_1, z_2, z_3, z_4)$ and

$$(1.5) \quad \phi^t(z_1, z_2, z_3, z_4) = \prod_{1 \leq i < j \leq 4} (z_i - z_j)^{\alpha_{ij}^t} (\bar{z}_i - \bar{z}_j)^{\beta_{ij}^t} f^t\left(\frac{(z_1 - z_2)(z_3 - z_4)}{(z_1 - z_3)(z_2 - z_4)}\right)$$

and $f^t \in F_{0,1,\infty}$ and for $\chi \in \{p, 1 - p, p^{-1}\}$

$$(1.6) \quad j(\chi, f^t) = \sum_{k=1}^{\infty} |p|^{r_k^t \chi} \sum_{n,m \geq 0} a_{n,m}^{t,\chi} p^n \bar{p}^m.$$

We first prove that there exists integers $n_{12}, n_{23}, n_{24}, N_\infty$ such that $d/dz_2^{N_\infty} ((z_1 - z_2)^{-n_{12}+1} (z_2 - z_3)^{-n_{23}+1} (z_2 - z_4)^{-n_{24}+1} \phi) = 0$. Fix z_1, z_3, z_4 . Since $d/d\bar{z}_2 \phi = 0$, ϕ is a holomorphic function for the variable z_2 on $\mathbb{C} \setminus \{z_1, z_3, z_4\}$. Let n_{12} (resp. n_{23}, n_{24}) be the largest integer

smaller than any element of the set $\{\alpha_{12}^t + \beta_{12}^t + r_k^{t,p}\}_{t=1,\dots,N,k=1,2,3,\dots}$ (resp. $\{\alpha_{23}^t + \beta_{23}^t + r_k^{t,1-p}\}_{t=1,\dots,N,k=1,2,3,\dots}$ and $\{\alpha_{24}^t + \beta_{24}^t + r_k^{t,p-1}\}_{t=1,\dots,N,k=1,2,3,\dots}$), which exists since $r_k^{t,\chi} \in \mathbb{R}$ is bounded below. Then, by (1.5) and (1.6), $\lim_{z_2 \rightarrow z_1} |z_2 - z_1|^{\frac{-n_{12}+1}{2}} f = 0$ and $\lim_{z_2 \rightarrow z_3} |z_2 - z_3|^{\frac{-n_{23}+1}{2}} f = 0$ and $\lim_{z_2 \rightarrow z_4} |z_2 - z_4|^{\frac{-n_{24}+1}{2}} f = 0$. Hence, $(z_1 - z_2)^{-n_{12}+1} (z_2 - z_3)^{-n_{23}+1} (z_2 - z_4)^{-n_{24}+1} \phi$ is a holomorphic function on $\mathbb{C}$. Since $d/d\bar{z}_2 (z_1 - z_2)^{-n_{12}+1} (z_2 - z_3)^{-n_{23}+1} (z_2 - z_4)^{-n_{24}+1} \phi = 0$, we may assume without loss of generality that ϕ is a holomorphic function for the variable z_2 on $\mathbb{C}$. Let n_∞ be the largest integer smaller than any element of the set $\{\alpha_{12}^t + \alpha_{23}^t + \alpha_{24}^t + 1\}_{t=1,\dots,N}$. Then, by (1.5), $\lim_{z_2 \rightarrow \infty} |z_2|^{-n_\infty} \phi = 0$. Hence, ϕ is a rational function on the projective plane $\mathbb{CP}^1$ and holomorphic on $\mathbb{C}$. Thus, ϕ is a polynomial whose degree is less than n_∞ , which implies that $d/dz_2^{n_\infty} \phi = 0$.

Now, we consider the expansion map $e_{1(4(23))} : \text{Cor}_4 \rightarrow T(x_{1(4(23))}, y_{1(4(23))}, z_{1(4(23))})$ and set $(x_{1(4(23))}, y_{1(4(23))}, z_{1(4(23))}, \bar{x}_{1(4(23))}, \bar{y}_{1(4(23))}, \bar{z}_{1(4(23))}) = (x, y, z, \bar{x}, \bar{y}, \bar{z})$. Since $T(x, y, z)$ is a completion of

$$\bigoplus_{r,s \in \mathbb{R}} z^r \bar{z}^s \mathbb{C}[[y/x, \bar{y}/\bar{x}]] [x^\pm, y^\pm, \bar{x}^\pm, \bar{y}^\pm, |x|^\mathbb{R}, |y|^\mathbb{R}],$$

by $d/dz_2^{n_\infty} \phi = 0$ and $d/d\bar{z}_2 \phi = 0$,

$$e_{1(4(23))}(\phi) \in \bigoplus_{k=0}^{n_\infty-1} z^k \mathbb{C}[[y/x, \bar{y}/\bar{x}]] [x^\pm, y^\pm, \bar{x}^\pm, \bar{y}^\pm, |x|^\mathbb{R}, |y|^\mathbb{R}].$$

Denote by $pr_{n_\infty} : T(x, y, z) \rightarrow \bigoplus_{k=0}^{n_\infty-1} z^k \mathbb{C}[[y/x, \bar{y}/\bar{x}]] [x^\pm, y^\pm, \bar{x}^\pm, \bar{y}^\pm, |x|^\mathbb{R}, |y|^\mathbb{R}]$ the projection of $T(x, y, z)$ onto the coefficients of $1, z, \dots, z^{n_\infty-1}$. Set $z_{ij} = z_i - z_j$ for $i, j = 1, \dots, 4$. Then,

$$\begin{aligned} e_{1(4(23))}(\phi) &= pr_{n_\infty} e_{1(4(23))}(\phi) \\ &= pr_{n_\infty} \sum_{t=1}^N e_{1(4(23))}^f (\prod_{1 \leq i < j \leq 4} (z_{ij})^{\alpha'_{ij}} (\bar{z}_{ij})^{\beta'_{ij}}) s_{1(4(23))}(j(1-p, f^t), \\ (1.7) \quad &= pr_{n_\infty} \sum_{t=1}^N \sum_{k=1}^{\infty} e_{1(4(23))}^f (\prod_{1 \leq i < j \leq 4} (z_{ij})^{\alpha'_{ij}} (\bar{z}_{ij})^{\beta'_{ij}}) s_{1(4(23))} \left(\sum_{n,m \geq 0} a_{n,m}^{t,1-p} p^n \bar{p}^m |p|^{t,1-p} \right). \end{aligned}$$

Since $e_{1(4(23))}^f (\prod_{1 \leq i < j \leq 4} z_{ij}^{\alpha'_{ij}} \bar{z}_{ij}^{\beta'_{ij}}) \in z^{\alpha'_{23}} \bar{z}^{\beta'_{23}} \mathbb{C}[[y/x, z/y, \bar{y}/\bar{x}, \bar{z}/\bar{y}]] [x^\pm, y^\pm |x|^\mathbb{R}, |y|^\mathbb{R}]$ and $s_{1(4(32))}(p^a \bar{p}^{a'}) \in z^a \bar{z}^{a'} \mathbb{C}[[y/x, z/y, \bar{y}/\bar{x}, \bar{z}/\bar{y}]] [x^\pm, y^\pm |x|^\mathbb{R}, |y|^\mathbb{R}]$ for any $a, a' \in \mathbb{R}$,

$$pr_{n_\infty} e_{1(4(32))}^f (\prod_{1 \leq i < j \leq 4} z_{ij}^{\alpha'_{ij}} \bar{z}_{ij}^{\beta'_{ij}}) s_{1(4(23))}(p^a \bar{p}^{a'})$$

is equal to zero unless $\alpha_{23}^t + a \in \mathbb{Z}_{\leq n_\infty}$ and $\beta_{23}^t + a' \in \mathbb{Z}_{\leq 0}$. Thus, the right hand side of (1.7) is finite sum, that is,

$$\begin{aligned}
&= pr_{n_\infty} \sum_{t=1}^N \sum_{\substack{k=1,2,3,\dots \\ r_k^{t,1-p} + \beta_{23}^t \in \mathbb{Z}_{\leq 0} \\ r_k^{t,1-p} + \alpha_{23}^t \in \mathbb{Z}_{\leq n_\infty}}} e_{1(4(23))}^f (\Pi_{1 \leq i < j \leq 4} (z_{ij})^{\alpha_{ij}^t} (\bar{z}_{ij})^{\beta_{ij}^t}) s_{1(4(23))} \left(\sum_{\substack{n_\infty - \alpha_{23}^t - r_k^{t,1-p} \geq n \geq 0 \\ -\beta_{23}^t - r_k^{t,1-p} \geq m \geq 0}} a_{n,m}^{t,1-p} p^n \bar{p}^m |p|^{r_k^{t,1-p}} \right). \\
&= pr_{n_\infty} \sum_{t=1}^N \sum_{\substack{k=1,2,3,\dots \\ r_k^{t,1-p} + \beta_{23}^t \in \mathbb{Z}_{\leq 0} \\ r_k^{t,1-p} + \alpha_{23}^t \in \mathbb{Z}_{\leq n_\infty}}} \sum_{\substack{n_\infty - \alpha_{23}^t - r_k^{t,1-p} \geq n \geq 0 \\ -\beta_{23}^t - r_k^{t,1-p} \geq m \geq 0}} e_{1(4(32))}^f (\Pi_{1 \leq i < j \leq 4} z_{ij}^{\alpha_{ij}^t} \bar{z}_{ij}^{\beta_{ij}^t} (1 - \xi)^{r_k^{t,1-p} + n} (1 - \bar{\xi})^{r_k^{t,1-p} + m}).
\end{aligned}$$

Thus, the coefficients of $1, z, \dots, z^{n_\infty-1}$ in $e_{1(4(32))}(\phi)$ is an element in Cor_4^f . This finishes the proof, the detailed verification of the assertion being left to the reader. $\square$

Corollary 1.3. *If a function $f \in \text{Cor}_4$ is independent of the variable z_4 , i.e., $d/dz_4 f = d/d\bar{z}_4 f$, then $f \in \text{Cor}_3$.*

1.7. Vacuum state and D_4 -symmetry. In section 1.3, we consider the expansions of a function in Cor_4 in three variables associated with an element in $A \in P_4$. In this section, we consider expansions in four variables. Let Q_4 be the set of parenthesized products of five elements $1, 2, 3, 4, \star$ with $\star$ at right most, e.g., $3((4((12)\star))$ (see Introduction 0.2). The permutation group S_4 acts on Q_4 , which fixes $\star$, and Q_4 consists of the permutations of the following elements, called standard elements of Q_4 :

$$\begin{aligned}
&((12)(34))\star, (12)((34)\star), (12)(3(4\star)), \\
&(\text{standard elements of } Q_4) \quad (((12)3)4)\star, ((12)3)(4\star), \\
&\quad ((1(23))4)\star, (1(23))(4\star), \\
&\quad (1((23)4))\star, 1(((23)4)\star), 1((23)(4\star)), \\
&\quad (1(2(34)))\star, 1((2(34))\star), 1(2((34)\star)), 1(2(3(4\star))).
\end{aligned}$$

The rule for the change of variables is given in Appendix. In this section, we briefly explain that all the expansions associated with Q_4 are given by the expansions associated with P_4 .

We start with the example $1(2(3(4\star))) \in Q_4$. In this case, we consider the expansion of a function Cor_4 in $|z_1| > |z_2| > |z_3| > |z_4|$. Set

$$T^\star(x, y, z, w) = \mathbb{C}[[y/x, w/z, \bar{y}/\bar{x}, \bar{w}/\bar{z}]][(z/y, \bar{z}/\bar{y}, |z/y|^{\mathbb{R}})][x^\pm, y^\pm, z^\pm, \bar{x}^\pm, \bar{y}^\pm, \bar{z}^\pm, |x|^{\mathbb{R}}, |y|^{\mathbb{R}}, |z|^{\mathbb{R}}],$$

for the formal variables $x, y, z, w, \bar{x}, \bar{y}, \bar{z}, \bar{w}$. We recall that the expansion $e_{1(2(34))}$ is given by the change of variables $(z_1 - z_4, z_2 - z_4, z_3 - z_4) \mapsto (z_{14}, z_{24}, z_{34})$ with the expansion in $|z_{14}| > |z_{24}| > |z_{34}|$. Then, let $T_{1(2(34))}^{1(2(3(4\star)))} : T(z_{14}, z_{24}, z_{34}) \rightarrow T^\star(z_1, z_2, z_3, z_4)$ be the linear map defined by

$$T_{1(2(34))}^{1(2(3(4\star)))} = \lim_{(z_{14}, z_{24}, z_{34}) \rightarrow (z_1, z_2, z_3)} \exp(-z_4(d/dz_{14} + d/dz_{24} + d/dz_{34})) \exp(-\bar{z}_4(d/d\bar{z}_{14} + d/d\bar{z}_{24} + d/d\bar{z}_{34}))$$

and set

$$e_{1(2(3(4\star)))} = T_{1(2(34))}^{1(2(3(4\star)))} \circ e_{1(2(34))} : \text{Cor}_4 \rightarrow T^*(z_1, z_2, z_3, z_4).$$

Then, similarly to Lemma 1.9, we have:

Lemma 1.17. *For $\phi \in \text{Cor}_4$, the formal power series $e_{1(2(3(4\star)))}(\phi)$ is absolutely convergent to $\phi(z_1, z_2, z_3, z_4)$ in $|z_1| \gg |z_2| \gg |z_3| \gg |z_4|$.*

For the case of $1((2(34))\star) \in Q_4$, we consider the following expansion with the change of variables

$$(z_1, z_{24}, z_{34}, z_4) = (z_1, z_2 - z_4, z_3 - z_4, z_4), \{ |z_1| \gg |z_{24}| \gg |z_{34}| \gg |z_4| \}.$$

Similarly to the above, such expansion $e_{1((2(34))\star)} : \text{Cor}_4 \rightarrow T^*(z_1, z_{24}, z_{34}, z_4)$ is given by

$$\begin{aligned} e_{1((2(34))\star)} &= T_{1(2(34))}^{1((2(34))\star)} \circ e_{1((2(34)))}, \\ T_{1(2(34))}^{1((2(34))\star)} &= \lim_{z_{14} \rightarrow z_1} \exp(-z_4 d/dz_{14}) \exp(-\bar{z}_4 d/d\bar{z}_{14}). \end{aligned}$$

In this way, we can define expansions and the space of formal power series for all elements in Q_4 by using the expansions defined in the previous section.

Hereafter, we study $e_{1(2(3(4\star)))}$ in more detail. Since the expansions and the derivations are commute with each other, by Lemma 1.1, the image of $e_{1(2(3(4\star)))}$ is in the kernel of $\frac{d}{dz_1} + \frac{d}{dz_2} + \frac{d}{dz_3} + \frac{d}{dz_4}$ and $\frac{d}{d\bar{z}_1} + \frac{d}{d\bar{z}_2} + \frac{d}{d\bar{z}_3} + \frac{d}{d\bar{z}_4}$. Set

$$T_0^*(z_1, z_2, z_3, z_4) = \{ f \in T^*(z_1, z_2, z_3, z_4) \mid \frac{d}{dz_1} + \frac{d}{dz_2} + \frac{d}{dz_3} + \frac{d}{dz_4} f = \frac{d}{dz_1} + \frac{d}{dz_2} + \frac{d}{dz_3} + \frac{d}{dz_4} f = 0 \}.$$

Then, the image of $T_{1(2(34))}^{1(2(3(4\star)))} : T(z_{14}, z_{24}, z_{34}) \rightarrow T^*(z_1, z_2, z_3, z_4)$ is in $T_0^*(z_1, z_2, z_3, z_4)$, since $[\frac{d}{dz_4}, \exp(-z_4(\frac{d}{dz_1} + \frac{d}{dz_2} + \frac{d}{dz_3}))] = -(\frac{d}{dz_1} + \frac{d}{dz_2} + \frac{d}{dz_3})$.

Lemma 1.18. *The above map $T_{1(2(34))}^{1(2(3(4\star)))} : T(z_{14}, z_{24}, z_{34}) \rightarrow T_0^*(z_1, z_2, z_3, z_4)$ is an isomorphism.*

Proof. By substituting $z_4 = \bar{z}_4 = 0$, we obtain the left inverse of the map, which implies that the map is injective. Let $f(z_1, z_2, z_3, z_4) = \sum_{n,m \geq 0} f_{n,m} z_4^n \bar{z}_4^m \in T_0^*(z_1, z_2, z_3, z_4)$. It suffices to show that f is in the image of the map. We may assume that $f_{0,0} = 0$. Then, $n f_{n,m} = (\frac{d}{dz_1} + \frac{d}{dz_2} + \frac{d}{dz_3}) f_{n-1,m}$ and $m f_{n,m} = (\frac{d}{d\bar{z}_1} + \frac{d}{d\bar{z}_2} + \frac{d}{d\bar{z}_3}) f_{n,m-1}$. Thus, $f_{n,m} = 0$ for all $n, m \geq 0$. $\square$

We end this section by studying the relation between $e_{1(2(3(4\star)))}$ and $e_{4(3(2(1\star)))}$. The map $e_{1(2(3(4\star)))}$ expands a function in $|z_1| > |z_2| > |z_3| > |z_4|$, whereas $e_{4(3(2(1\star)))}$ expands in $|z_4| > |z_3| > |z_2| > |z_1|$. We consider the following involution: $X_4 \rightarrow X_4$, $(z_1, z_2, z_3, z_4) \mapsto (z_1^{-1}, z_2^{-1}, z_3^{-1}, z_4^{-1})$ and set

$$T_d^*(x, y, z, w) = \mathbb{C}[[y/x, w/z, \bar{y}/\bar{x}, \bar{w}/\bar{z}]][(z/y, \bar{z}/\bar{y}, |z/y|^{\mathbb{R}})][y^{\pm}, z^{\pm}, \bar{y}^{\pm}, \bar{z}^{\pm}, |y|^{\mathbb{R}}, |z|^{\mathbb{R}}],$$

which is a subspace of $T^\star(x, y, z, w)$. Define the map $I_d : T_d^\star(z_1, z_2, z_3, z_4) \rightarrow T_d^\star(z_4, z_3, z_2, z_1)$ by $(\underline{z}_1, \underline{z}_2, \underline{z}_3, \underline{z}_4) \mapsto (\underline{z}_1^{-1}, \underline{z}_2^{-1}, \underline{z}_3^{-1}, \underline{z}_4^{-1})$. We observe that

$$\begin{aligned} I_d(\bar{z}_1^{-r} e_{1(2(3(4\star)))}(z_1 - z_2)^r) &= I_d\left(\sum_{i \geq 0} (-1)^i \binom{r}{i} \bar{z}_1^{-i} z_2^i\right) \\ &= \sum_{i \geq 0} (-1)^i \binom{r}{i} \bar{z}_1^i z_2^{-i} \\ &= \bar{z}_2^{-r} (-1)^r e_{4(3(2(1\star)))}((z_1 - z_2)^r). \end{aligned}$$

Let

$$\phi(z_1, z_2, z_3, z_4) = \prod_{1 \leq i < j \leq 4} (z_i - z_j)^{\alpha_{ij}} (\bar{z}_i - \bar{z}_j)^{\beta_{ij}} f \circ \xi(z_1, z_2, z_3, z_4) \in \text{Cor}_4,$$

where $f \in F_{0,1,\infty}$ and $\alpha_{ij}, \beta_{ij} \in \mathbb{R}$ satisfy $\alpha_{ij} - \beta_{ij} \in \mathbb{Z}$ for any $1 \leq i < j \leq 4$. Set $P(\alpha, \beta, z) = \prod_{1 \leq i < j \leq 4} (-1)^{\alpha_{ij} - \beta_{ij}} (z_i z_j)^{\alpha_{ij}} (\bar{z}_i \bar{z}_j)^{\beta_{ij}}$. Then, we have:

Lemma 1.19. $I_d(P(\alpha, \beta, z)^{-1} e_{1(2(3(4\star)))}(\phi)) = e_{4(3(2(1\star)))}(\phi)$.

The above transformation corresponds to (14)(23) in S_4 . It is well-known that the axiom of a vertex algebra possesses S_3 -symmetry (see [FHL]). We will later prove that the correlation functions is S_4 -invariant by using the S_3 -symmetry together with (14)(23)-symmetry above (see Proposition 3.7).

1.8. Generalized two point Correlation function. In the theory of a vertex algebra, four point correlation functions in the limit of $(z_1, z_4) \mapsto (\infty, 0)$ are used, which we call generalized two point functions (see Introduction 0.2). This subsection is devoted to studying the property of this generalized two point function. Set

$$U(y, z) = \mathbb{C}((z/y, \bar{z}/\bar{y}, |z/y|^{\mathbb{R}})) [y^\pm, z^\pm, \bar{y}^\pm, \bar{z}^\pm, |y|^{\mathbb{R}}, |z|^{\mathbb{R}}]$$

and

$$Y_2 = \{(z_1, z_2) \in \mathbb{C}^2 \mid z_1 \neq z_2, z_1 \neq 0, z_2 \neq 0\}.$$

Let $\eta(z_1, z_2) : Y_2 \rightarrow \mathbb{C}P^1 \setminus \{0, 1, \infty\}$ be the real analytic function defined by $\eta(z_1, z_2) = \frac{z_2}{z_1}$. For $f \in F_{0,1,\infty}$, $f \circ \eta$ is a real analytic function on Y_2 . Denote by GCor_2 the space of real analytic functions on Y_2 spanned by

$$(1.8) \quad z_1^{\alpha_1} z_2^{\alpha_2} (z_1 - z_2)^{\alpha_{12}} \bar{z}_1^{\beta_1} \bar{z}_2^{\beta_2} (\bar{z}_1 - \bar{z}_2)^{\beta_{12}} f \circ \eta(z_1, z_2),$$

where $f \in F_{0,1,\infty}$ and $\alpha_1, \alpha_2, \alpha_{12}, \beta_1, \beta_2, \beta_{12} \in \mathbb{R}$ satisfy $\alpha_1 - \beta_1, \alpha_2 - \beta_2, \alpha_{12} - \beta_{12} \in \mathbb{Z}$. Define the action of $\text{Aut}(0, 1, \infty)$ on Y_2 by

$$\begin{aligned} (01) \cdot (z_1, z_2) &= (z_1, z_1 - z_2), \\ (0\infty) \cdot (z_1, z_2) &= (z_2, z_1), \\ (1\infty) \cdot (z_1, z_2) &= (z_2 - z_1, z_2), \\ (01\infty) \cdot (z_1, z_2) &= (z_1 - z_2, z_1), \\ (10\infty) \cdot (z_1, z_2) &= (z_2, z_2 - z_1), \end{aligned}$$

for $(z_1, z_2) \in Y_2$. Then, we have:

Lemma 1.20. *The map $\eta : Y_2 \rightarrow \mathbb{CP}^1 \setminus \{0, 1, \infty\}$ commutes with the action of $\text{Aut}(0, 1, \infty)$, i.e., $\eta(t \cdot (z_1, z_2)) = t \cdot \frac{z_2}{z_1}$ for any $(z_1, z_2) \in Y_2$ and $t \in \text{Aut}(0, 1, \infty)$. In particular, for $\mu \in \text{GCor}_2$, $(t \cdot \mu)(-) = \mu(t^{-1} -) \in \text{GCor}_2$ for any $t \in \text{Aut}(0, 1, \infty)$.*

Let $\mu(z_1, z_2) = z_1^{\alpha_1} z_2^{\alpha_2} (z_1 - z_2)^{\alpha_{12}} \bar{z}_1^{\beta_1} \bar{z}_2^{\beta_2} (\bar{z}_1 - \bar{z}_2)^{\beta_{12}} f \circ \eta(z_1, z_2)$ in (1.8). The expansions of μ in $\{|z_1| > |z_2|\}$ and $\{|z_2| > |z_1|\}$ are respectively given by

$$\begin{aligned} & z_1^{\alpha_1 + \alpha_{12}} \bar{z}_1^{\beta_1 + \beta_{12}} z_2^{\alpha_2} \bar{z}_2^{\beta_2} \sum_{i,j \geq 0} \binom{\alpha_{12}}{i} \binom{\beta_{12}}{j} (-z_2/z_1)^i (-\bar{z}_2/\bar{z}_1)^j \lim_{p \rightarrow z_2/z_1} j(p, f) \\ & (-1)^{\alpha_{12} - \beta_{12}} z_1^{\alpha_1} \bar{z}_1^{\beta_1} z_2^{\alpha_2 + \alpha_{12}} \bar{z}_2^{\beta_2 + \beta_{12}} \sum_{i,j \geq 0} \binom{\alpha_{12}}{i} \binom{\beta_{12}}{j} (-z_1/z_2)^i (-\bar{z}_1/\bar{z}_2)^j \lim_{p \rightarrow z_1/z_2} j(p^{-1}, f), \end{aligned}$$

which define maps

$$|_{|z_1| > |z_2|} : \text{GCor}_2 \rightarrow U(z_1, z_2), \mu(z_1, z_2) \mapsto \mu(z_1, z_2)|_{|z_1| > |z_2|}$$

and

$$|_{|z_2| > |z_1|} : \text{GCor}_2 \rightarrow U(z_2, z_1), \mu(z_1, z_2) \mapsto \mu(z_1, z_2)|_{|z_2| > |z_1|}.$$

Since $f(\frac{z_2}{z_1}) = f(\frac{z_2}{z_2 + (z_1 - z_2)})$, the expansions of μ in $\{|z_2| > |z_1 - z_2|\}$ is given by

$$z_2^{\alpha_1 + \alpha_2} \bar{z}_2^{\beta_1 + \beta_2} z_0^{\alpha_{12}} \bar{z}_0^{\beta_{12}} \sum_{i,j \geq 0} \binom{\alpha_{12}}{i} \binom{\beta_{12}}{j} (z_0/z_2)^i (\bar{z}_0/\bar{z}_2)^j \lim_{p \rightarrow -z_0/z_2} j(1 - p^{-1}, f),$$

where $z_0 = z_1 - z_2$. We denote it by

$$|_{|z_2| > |z_1 - z_2|} : \text{GCor}_2 \rightarrow U(z_2, z_0), \mu(z_1, z_2) \mapsto \mu(z_1, z_2)|_{|z_2| > |z_1 - z_2|}.$$

Then, we have:

Lemma 1.21. *For $f \in F_{0,1,\infty}$,*

$$\begin{aligned} f \circ \eta|_{|z_1| > |z_2|} &= \lim_{p \rightarrow z_1/z_2} j(1, f), \\ f \circ \eta|_{|z_2| > |z_1|} &= \lim_{p \rightarrow z_2/z_1} j(p^{-1}, f), \\ f \circ \eta|_{|z_2| > |z_1 - z_2|} &= \lim_{p \rightarrow -z_0/z_2} j(1 - p^{-1}, f). \end{aligned}$$

For $a, a' \in \mathbb{R}$, define a $\mathbb{C}$ -linear map $C_{a,a'}(x) : T(x, y, z) \rightarrow U(y, z)$ by taking the coefficient of $x^a \bar{x}^{a'}$. Then, we have the $\mathbb{C}$ -linear map $C_{a,a'}(x) : T(x, y, z) \rightarrow U(y, z)$.

Proposition 1.4. *For $\phi \in \text{Cor}_4$ and $a, a' \in \mathbb{R}$, there exists $\eta \in \text{GCor}_2$ satisfying the following conditions:*

- (1) $C_{a,a'}(x_{1(2(34))})e_{1(2(34))}\phi = \eta|_{|z_1| > |z_2|}$,
- (2) $C_{a,a'}(x_{1(3(24))})e_{1(3(24))}\phi = \eta|_{|z_2| > |z_1|}$,
- (3) $C_{a,a'}(x_{1((23)4)})e_{1((23)4)}\phi = \eta|_{|z_2| > |z_1 - z_2|}$,

where we identify $(y_{1(2(34))}, z_{1(2(34))})$ and $(y_{1(3(24))}, z_{1(3(24))})$, $(y_{1((23)4)}, z_{1((23)4)})$ as (z_1, z_2) and (z_2, z_1) , (z_2, z_0) , respectively.

Proof. Since the expansion is linear, we may assume that $\phi = \prod_{1 \leq i < j \leq 4} (z_i - z_j)^{\alpha_{ij}} (\bar{z}_i - \bar{z}_j)^{\beta_{ij}} f \circ \xi$, where $f \in F_{0,1,\infty}$ and $\alpha_{ij} - \beta_{ij} \in \mathbb{Z}$. Set

$$T_g(x, y, z) = \mathbb{C}[[y/x, \bar{y}/\bar{x}]][(z/y, \bar{z}/\bar{y}, |z/y|^{\mathbb{R}})] [y^{\pm}, z^{\pm}, \bar{y}^{\pm}, \bar{z}^{\pm}, |y|^{\mathbb{R}}, |z|^{\mathbb{R}}],$$

which involves only $x^{-n} \bar{x}^{-n'}$ with $n, n' \in \mathbb{Z}_{\geq 0}$, and $c = \alpha_{12} + \alpha_{13} + \alpha_{14}$ and $c' = \beta_{12} + \beta_{13} + \beta_{14}$. Let $A = 1(2(34)), 1(3(24)), 1((23)4)$. Then, $e_A(\phi) \subset z_1^c \bar{z}_1^{c'} T_g(x_A, y_A, z_A)$ and, thus, $C_{a,a'}(x_A) e_A(\phi) = 0$ unless $c - a, c' - a' \in \mathbb{Z}_{\geq 0}$. Hence, we may assume that $a = c - n$ and $a' = c' - n'$ for some $n, n' \in \mathbb{Z}_{\geq 0}$. Since $\lim_{z_1 \rightarrow \infty} z_1^{-c} \bar{z}_1^{-c'} \phi$ exists and is equal to

$$\prod_{2 \leq i < j \leq 4} (z_i - z_j)^{\alpha_{ij}} (\bar{z}_i - \bar{z}_j)^{\beta_{ij}} f\left(\frac{z_3 - z_4}{z_2 - z_4}\right),$$

$C_{c-n, c'-n'}(x_A) e_A(\phi)$ is convergent to the real analytic function

$$\eta(z_1, z_2) = \frac{1}{n!n'!} \lim_{(z_1, z_2, z_3, z_4) \rightarrow (\infty, z_1, z_2, 0)} (-z_1^2 d/dz_1)^n (-\bar{z}_1^2 d/d\bar{z}_1)^{n'} (z_1^{-c} \bar{z}_1^{-c'} \phi(z_1, z_2, z_3, z_4)) \in \text{GCor}_2,$$

where we used $-z^2 \frac{d}{dz} z^{-k} = k z^{-(k-1)}$. Since $e_{1((23)4)}$ is the expansion around $|z_1 - z_4| > |z_3 - z_4| > |z_2 - z_3|$, in the limit of $\lim_{(z_1, z_2, z_3, z_4) \rightarrow (\infty, z_1, z_2, 0)}$, it gives the expansion around $|z_2| > |z_1 - z_2|$. $\square$

Similarly to the proof of Proposition 1.3, we have:

Lemma 1.22. *If $\mu \in \text{GCor}_2$ satisfies $\frac{d}{dz_1} \mu = 0$, then $\mu(z_1, z_2) \in \mathbb{C}[z_1^{\pm}, (z_1 - z_2)^{\pm}, z_2^{\pm}, \bar{z}_2^{\pm}, |z_2|^{\mathbb{R}}]$. Furthermore, if $\frac{d}{dz_1} \mu = \frac{d}{dz_2} \mu = 0$, then $\mu(z_1, z_2) \in \mathbb{C}[z_1^{\pm}, z_2^{\pm}, (z_1 - z_2)^{\pm}]$.*

Similarly to Section 1.5, let $S_g(z_1, z_2)$ be the subspace of $U(z_1, z_2)$ consisting of $f \in U(z_1, z_2)$ with $(z_1 d/dz_1 + z_2 d/dz_2) f = (\bar{z}_1 d/d\bar{z}_1 + \bar{z}_2 d/d\bar{z}_2) f = 0$. Clearly, the map $s_g : \mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}})) \rightarrow S_g(z_1, z_2)$ defined by the substitution of z_2/z_1 into p and the map $v_g : S_g(z_1, z_2) \rightarrow \mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}}))$ defined by the limit $(z_1, z_2) \rightarrow (1, p)$ are mutually inverse. Then, we have:

Lemma 1.23. *Let $\mu \in \text{GCor}_2$ satisfy $(z_1 d/dz_1 + z_2 d/dz_2) \mu = (\bar{z}_1 d/d\bar{z}_1 + \bar{z}_2 d/d\bar{z}_2) \mu = 0$. Then, there exists $f \in F_{0,1,\infty}$ such that*

$$\begin{aligned} \mu|_{|z_1| > |z_2|} &= \lim_{p \rightarrow z_2/z_1} j(p, f) \\ \mu|_{|z_2| > |z_1 - z_2|} &= \lim_{p \rightarrow -\bar{z}_0/\bar{z}_1} j(1 - p^{-1}, g) \\ \mu|_{|z_2| > |z_1|} &= \lim_{p \rightarrow z_1/z_2} j(p^{-1}, g). \end{aligned}$$

Proof. Set $f(p) = \mu(1, p)$, which is a real analytic function on $\mathbb{C}P^1 \setminus \{0, 1, \infty\}$. By the existence of the expansion around $\{0, 1, \infty\}$, f has conformal singularities at $\{0, 1, \infty\}$. Thus, the assertion holds. $\square$

We end this section by studying the behavior of the expansion $|_{|z_2| > |z_1 - z_2|}$ of generalized two point functions under the involution $I_Y : Y_2 \rightarrow Y_2, (z_1, z_2) \mapsto (z_1^{-1}, z_2^{-1})$, which is

important to define a dual module for a full vertex algebra. The involution I_Y acts on GCor_2 by $\mu(z_1, z_2) \mapsto \mu(z_1^{-1}, z_2^{-1})$ for $\mu \in \text{GCor}_2$. Define the map

$$\lim_{(z_0, z_2) \mapsto (\frac{-z_0}{z_2(z_2+z_0)}, \frac{1}{z_2})} : U(z_2, z_0) \rightarrow U(z_2, z_0)$$

by substituting $(-z_0/z_2^2 \sum_{i \geq 0} (-1)^i (z_0/z_2)^i, z_2^{-1})$ into z_0 and z_2 . The map is well-defined, since $U(z_2, z_0) = \mathbb{C}((z_0/z_2, \bar{z}_0/\bar{z}_2, |z_0/z_2|)) [z_0^\pm, z_2^\pm, \bar{z}_0^\pm, \bar{z}_2^\pm, |z_0|^\mathbb{R}, |z_2|^\mathbb{R}]$ and $\lim_{(z_0, z_2) \mapsto (\frac{-z_0}{z_2(z_2+z_0)}, \frac{1}{z_2})} z_0/z_2 = -z_0/z_2(1 - z_0/z_2 + (z_0/z_2)^2 - \dots)$.

Lemma 1.24. *For any $\mu \in \text{GCor}_2$,*

$$\lim_{(z_0, z_2) \mapsto (\frac{-z_0}{z_2(z_2+z_0)}, \frac{1}{z_2})} \mu|_{|z_2| > |z_1 - z_2|} = I_Y(\mu)|_{|z_2| > |z_1 - z_2|}.$$

Proof. Set $\mu'(z_0, z_2) = \mu(z_0 + z_2, z_2)$. Since the series $\mu|_{|z_2| > |z_1 - z_2|} \in U(z_2, z_0)$ is absolutely convergent in $|z_2| > |z_0|$, $\lim_{(z_0, z_2) \mapsto (\frac{-z_0}{z_2(z_2+z_0)}, \frac{1}{z_2})} \mu|_{|z_2| > |z_1 - z_2|}$ is also absolutely convergent to $\mu'(\frac{-z_0}{z_2(z_2+z_0)}, \frac{1}{z_2})$ in $|z_2| \gg |z_0|$, which is equal to $\mu(\frac{-z_0}{z_2(z_2+z_0)} + \frac{1}{z_2}, \frac{1}{z_2}) = \mu((z_0 + z_2)^{-1}, z_2^{-1})$. The series $I_Y(\mu)|_{|z_2| > |z_1 - z_2|}$ is also absolutely convergent to $\mu((z_0 + z_2)^{-1}, z_2^{-1})$ in the same domain. Hence, they coincide with each other. $\square$

2. FULL VERTEX ALGEBRA

In this section, we introduce the notion of a full vertex algebra, which is a generalization of a $\mathbb{Z}$ -graded vertex algebra.

2.1. Definition of full vertex algebra. For an $\mathbb{R}^2$ -graded vector space $F = \bigoplus_{h, \bar{h} \in \mathbb{R}^2} F_{h, \bar{h}}$, set $F^\vee = \bigoplus_{h, \bar{h} \in \mathbb{R}^2} F_{h, \bar{h}}^*$, where $F_{h, \bar{h}}^*$ is the dual vector space of $F_{h, \bar{h}}$. A full vertex algebra is an $\mathbb{R}^2$ -graded $\mathbb{C}$ -vector space $F = \bigoplus_{h, \bar{h} \in \mathbb{R}^2} F_{h, \bar{h}}$ equipped with a linear map

$$Y(-, \underline{z}) : F \rightarrow \text{End}(F)[[z^\pm, \bar{z}^\pm, |z|^\mathbb{R}]], \quad a \mapsto Y(a, \underline{z}) = \sum_{r, s \in \mathbb{R}} a(r, s) z^{-r-1} \bar{z}^{-s-1}$$

and an element $\mathbf{1} \in F_{0,0}$ satisfying the following conditions:

- FV1) For any $a, b \in F$, there exists $N \in \mathbb{R}$ such that $a(r, s)b = 0$ for any $r \geq N$ or $s \geq N$, that is, $Y(a, \underline{z})b \in F((z, \bar{z}, |z|^\mathbb{R}))$;
- FV2) $F_{h, \bar{h}} = 0$ unless $h - \bar{h} \in \mathbb{Z}$;
- FV3) For any $a \in F$, $Y(a, \underline{z})\mathbf{1} \in F[[z, \bar{z}]]$ and $\lim_{z \rightarrow 0} Y(a, \underline{z})\mathbf{1} = a(-1, -1)\mathbf{1} = a$.
- FV4) $Y(\mathbf{1}, \underline{z}) = \text{id} \in \text{End} F$;
- FV5) For any $a, b, c \in F$ and $u \in F^\vee$, there exists $\mu(z_1, z_2) \in \text{GCor}_2$ such that

$$\begin{aligned} u(Y(a, \underline{z}_1)Y(b, \underline{z}_2)c) &= \mu|_{|z_1| > |z_2|}, \\ u(Y(Y(a, \underline{z}_0)b, \underline{z}_2)c) &= \mu|_{|z_2| > |z_1 - z_2|}, \\ u(Y(b, \underline{z}_2)Y(a, \underline{z}_1)c) &= \mu|_{|z_2| > |z_1|}, \end{aligned}$$

where $z_0 = z_1 - z_2$.

- FV6) $F_{h, \bar{h}}(r, s)F_{h', \bar{h}'} \subset F_{h+h'-r-1, \bar{h}+\bar{h}'-s-1}$ for any $r, s, h, h', \bar{h}, \bar{h}' \in \mathbb{R}$.

Remark 2.1. Physically, the energy and the spin of a state in $F_{h,\bar{h}}$ are $h + \bar{h}$ and $h - \bar{h}$. Thus, the condition (FV2) implies that we only consider the particles whose spin is an integer, that is, we consider only bosons and not fermions. The notion of a full super vertex algebra can be defined by modifying (FV5) and (FV2).

Remark 2.2. Define the linear map $L(0), \bar{L}(0) \in \text{End } F$ by $L(0)v = hv$ and $\bar{L}(0)a = \bar{h}a$ for any $h, \bar{h} \in \mathbb{R}$ and $a \in F_{h,\bar{h}}$. Then, the condition (FV6) is equivalent to the the following condition: For any $h, \bar{h} \in \mathbb{R}$ and $a \in F_{h,\bar{h}}$,

$$[L(0), Y(a, \underline{z})] = (zd/dz + h)Y(a, \underline{z}),$$

$$[\bar{L}(0), Y(a, \underline{z})] = (\bar{z}d/d\bar{z} + \bar{h})Y(a, \underline{z}).$$

Let $(F^1, Y^1, \mathbf{1}^1)$ and $(F^2, Y^2, \mathbf{1}^2)$ be full vertex algebras. A full vertex algebra homomorphism from F^1 to F^2 is a linear map $f : F^1 \rightarrow F^2$ such that

$$(1) f(\mathbf{1}^1) = \mathbf{1}^2$$

$$(2) f(Y^1(a, \underline{z})-) = Y^2(f(a), \underline{z})f(-) \text{ for any } a \in F^1.$$

The notions of a subalgebra and an ideal are defined in the usual way.

A module of a full vertex algebra F is an $\mathbb{R}^2$ -graded $\mathbb{C}$ -vector space $M = \bigoplus_{h,\bar{h} \in \mathbb{R}^2} M_{h,\bar{h}}$ equipped with a linear map

$$Y_M(-, \underline{z}) : F \rightarrow \text{End}(M)[[z^\pm, \bar{z}^\pm, |z|^\mathbb{R}]], \quad a \mapsto Y_M(a, z) = \sum_{r,s \in \mathbb{R}} a(r, s) z^{-r-1} \bar{z}^{-s-1}$$

satisfying the following conditions:

FM1) For any $a \in F$ and $m \in M$, there exists $N \in \mathbb{R}$ such that $a(r, s)m = 0$ for any $r \geq N$ or $s \geq N$;

FM2) $M_{n,m} = 0$ unless $n - m \in \mathbb{Z}$;

FM3) $Y_M(\mathbf{1}, \underline{z}) = \text{id} \in \text{End } M$;

FM4) For any $a, b \in F$, $m \in M$ and $u \in M^\vee$, there exists $\mu \in \text{GCor}_2$ such that

$$u(Y_M(a, \underline{z}_1)Y_M(b, \underline{z}_2)m) = \mu|_{|z_1| > |z_2|},$$

$$u(Y_M(Y_M(a, \underline{z}_0)b, \underline{z}_2)m) = \mu|_{|z_2| > |z_1 - z_2|},$$

$$u(Y_M(b, \underline{z}_2)Y_M(a, \underline{z}_1)m) = \mu|_{|z_2| > |z_1|};$$

FM5) $F_{h,\bar{h}}(r, s)M_{h',\bar{h}'} \subset M_{h+h'-r-1, \bar{h}+\bar{h}'-s-1}$ for any $r, s, h, h', \bar{h}, \bar{h}' \in \mathbb{R}$.

As a consequence of (FM1) and (FM5), we have:

Lemma 2.1. Let $h_i, \bar{h}_i \in \mathbb{R}$, $a_i \in F_{h_i, \bar{h}_i}$ ($i = 1, 2$), $m \in M_{h_3, \bar{h}_3}$ and $u \in M_{h_0, \bar{h}_0}^*$. Then, $u(Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)m) \in z_1^{h_0-h_1-h_2-h_3} \bar{z}_1^{\bar{h}_0-\bar{h}_1-\bar{h}_2-\bar{h}_3} \mathbb{C}((z_2/z_1, \bar{z}_2/\bar{z}_1, |z_2/z_1|^\mathbb{R}))$.

Proof. Set

$$\sum_{s_1, \bar{s}_1, s_2, \bar{s}_2 \in \mathbb{R}} c_{s_1, \bar{s}_1, s_2, \bar{s}_2} z_1^{s_1} \bar{z}_1^{\bar{s}_1} z_2^{s_2} \bar{z}_2^{\bar{s}_2} = u(Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)m).$$

Then,

$$c_{s_1, \bar{s}_1, s_2, \bar{s}_2} = u(a_1(-s_1 - 1, -\bar{s}_1 - 1)a_2(-s_2 - 1, -\bar{s}_2 - 1)m).$$

By (FM5), $a_1(-s_1 - 1, -\bar{s}_1 - 1)a_2(-s_2 - 1, -\bar{s}_2 - 1)m \in M_{h_1+h_2+h_3+s_1+s_2, \bar{h}_1+\bar{h}_2+\bar{h}_3+\bar{s}_1+\bar{s}_2}$. Hence, $c_{s_1, \bar{s}_1, s_2, \bar{s}_2} = 0$ unless $h_0 = h_1 + h_2 + h_3 + s_1 + s_2$ and $\bar{h}_0 = \bar{h}_1 + \bar{h}_2 + \bar{h}_3 + \bar{s}_1 + \bar{s}_2$. Thus, we have

$$u(Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)m) = z_1^{h_0-h_1-h_2-h_3} \bar{z}_1^{\bar{h}_0-\bar{h}_1-\bar{h}_2-\bar{h}_3} \sum_{s_2, \bar{s}_2 \in \mathbb{R}} c_{s_1, \bar{s}_1, s_2, \bar{s}_2} (z_2/z_1)^{s_2} (\bar{z}_2/\bar{z}_1)^{\bar{s}_2},$$

where $s_1 = h_0 - (h_1 + h_2 + h_3 + s_2)$ and $\bar{s}_1 = \bar{h}_0 - (\bar{h}_1 + \bar{h}_2 + \bar{h}_3 + \bar{s}_2)$. By (FM1), the assertion holds. $\square$

By Lemma 2.1 and Lemma 1.23, we have:

Lemma 2.2. *Let $h_i, \bar{h}_i \in \mathbb{R}$, $a_i \in F_{h_i, \bar{h}_i}$ ($i = 1, 2$), $m \in M_{h_3, \bar{h}_3}$ and $u \in M_{h_0, \bar{h}_0}^*$, there exists $f \in F_{0,1,\infty}$ such that*

$$\begin{aligned} z_2^{-h_0+h_1+h_2+h_3} \bar{z}_2^{-\bar{h}_0+\bar{h}_1+\bar{h}_2+\bar{h}_3} u(Y(a, \underline{z}_1)Y(b, \underline{z}_2)m) &= \lim_{p \rightarrow z_1/z_2} j(p, f), \\ z_2^{-h_0+h_1+h_2+h_3} \bar{z}_2^{-\bar{h}_0+\bar{h}_1+\bar{h}_2+\bar{h}_3} u(Y(Y(a, \underline{z}_0)b, \underline{z}_2)m) &= \lim_{p \rightarrow -z_0/z_2} j(1 - p^{-1}, f), \\ z_2^{-h_0+h_1+h_2+h_3} \bar{z}_2^{-\bar{h}_0+\bar{h}_1+\bar{h}_2+\bar{h}_3} u(Y(b, \underline{z}_2)Y(a, \underline{z}_1)m) &= \lim_{p \rightarrow z_2/z_1} j(1/p, f). \end{aligned}$$

Let M, N be a F -module. A F -module homomorphism from M to N is a linear map $f : M \rightarrow N$ such that $f(Y_M(a, \underline{z})-) = Y_N(a, \underline{z})f(-)$ for any $a \in F$.

Let M be a F -module. According to [L1], a vector $v \in M$ is said to be a vacuum-like vector if $Y(a, \underline{z})v \in M[[z, \bar{z}]]$ for any $a \in F$.

Lemma 2.3. *Let $v \in M$ be a vacuum-like vector and $a, b \in F$ and $u \in M^\vee$ and $\mu \in \text{GCor}_2$ satisfy $u(Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)v) = \mu|_{|z_1| > |z_2|}$. Then, μ is a linear combination of the functions of the form $(z_1 - z_2)^{\alpha_{12}} z_2^{\alpha_2} (\bar{z}_1 - \bar{z}_2)^{\beta_{12}} \bar{z}_2^{\beta_2}$, where $\alpha_2, \beta_2 \in \mathbb{Z}_{\geq 0}$ and $\alpha_{12}, \beta_{12} \in \mathbb{R}$ satisfy $\alpha_{12} - \beta_{12} \in \mathbb{Z}$. Furthermore, the linear function $F_v : F \rightarrow M$ defined by $a \mapsto a(-1, -1)v$ is a F -module homomorphism.*

Proof. By (FM4), $u(Y(Y(a_1, \underline{z}_0)a_2, \underline{z}_2)v) = \mu|_{|z_1| > |z_1 - z_2|}$. Since v is a vacuum like vector, by Lemma 2.1 $p(z_0, z_2) = \mu|_{|z_1| > |z_1 - z_2|} \in \mathbb{C}[z_0^\pm, \bar{z}_0^\pm, |z_0|^\mathbb{R}, z_2, \bar{z}_2] \subset U(z_2, z_0)$, which proves the first part of the lemma. It suffices to show that $F_v(Y(a_1, \underline{z}_0)a_2) = Y(a_1, \underline{z}_0)F_v(a_2)$. Since

$$\begin{aligned} u(Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)v) &= \mu|_{|z_1| > |z_2|} \\ &= \lim_{z_0 \rightarrow (z_1 - z_2)|_{|z_1| > |z_2|}} p(z_0, z_2), \end{aligned}$$

we have

$$(2.1) \quad u(Y(a_1, \underline{z}_0)Y(a_2, \underline{z}_2)v) = \exp(-z_2 d/dz_0 - \bar{z}_2 d/d\bar{z}_0) u(Y(Y(a_1, \underline{z}_0)a_2, \underline{z}_2)v).$$

Thus,

$$\begin{aligned} Y(a_1, \underline{z}_0)F_v(a_2) &= \lim_{z_2 \mapsto 0} u(Y(a_1, \underline{z}_0)Y(a_2, \underline{z}_2)v) \\ &= \lim_{z_2 \mapsto 0} \exp(-z_2 d/dz_0 - \bar{z}_2 d/d\bar{z}_0) u(Y(Y(a_1, \underline{z}_0)a_2, \underline{z}_2)v) \\ &= F_v(Y(a_1, \underline{z}_0)a_2). \end{aligned}$$

□

Let F be a full vertex algebra and D and $\bar{D}$ denote the endomorphism of F defined by $Da = a(-2, -1)\mathbf{1}$ and $\bar{D}a = a(-1, -2)$ for $a \in F$, i.e.,

$$Y(a, z)\mathbf{1} = a + Da z + \bar{D}a \bar{z} + \dots$$

Define $Y(a, -\underline{z})$ by $Y(a, -\underline{z}) = \sum_{r,s} (-1)^{r-s} a(r, s) z^r \bar{z}^s$, where we used $a(r, s) = 0$ for $r - s \notin \mathbb{Z}$, which follows from (FV2) and (FV6).

Proposition 2.1. *For $a \in F$, the following properties hold:*

- (1) $Y(Da, \underline{z}) = d/dz Y(a, \underline{z})$ and $Y(\bar{D}a, \underline{z}) = d/d\bar{z} Y(a, \underline{z})$;
- (2) $D\mathbf{1} = \bar{D}\mathbf{1} = 0$;
- (3) $[D, \bar{D}] = 0$;
- (4) $Y(a, \underline{z})b = \exp(zD + \bar{z}\bar{D})Y(b, -\underline{z})a$;
- (5) $Y(\bar{D}a, \underline{z}) = [\bar{D}, Y(a, \underline{z})]$ and $Y(Da, \underline{z}) = [D, Y(a, \underline{z})]$.

Proof. Let $u \in F^\vee$ and $a, b \in F$ and $\mu_1, \mu_2 \in \text{GCor}_2$ satisfy

$$u(Y(a, \underline{z}_1)Y(\mathbf{1}, \underline{z}_2)b) = \mu_1|_{|z_1| > |z_2|}, u(Y(a, \underline{z}_1)Y(b, \underline{z}_2)\mathbf{1}) = \mu_2|_{|z_1| > |z_2|}.$$

By (FV4) and (FV5), $p_1(z_1) = \mu_1|_{|z_1| > |z_2|} \in \mathbb{C}[z_1^\pm, \bar{z}_1^\pm, |z_1|^\mathbb{R}]$. Then,

$$u(Y(Y(a, \underline{z}_0)\mathbf{1}, \underline{z}_2)b) = \mu_1|_{|z_2| > |z_1 - z_2|} = \lim_{z_1 \rightarrow z_2} \exp(z_0 \frac{d}{dz_1}) \exp(\bar{z}_0 \frac{d}{d\bar{z}_1}) p_1(z_1).$$

Thus, $u(Y(Da, \underline{z}_2)b) = \lim_{z_1 \rightarrow z_2} \frac{d}{dz_1} p_1(z_1) = \frac{d}{dz_2} u(Y(a, z_2)b)$, which implies that $Y(Da, \underline{z}) = \frac{d}{dz} Y(a, \underline{z})$ and similarly $Y(\bar{D}a, \underline{z}) = \frac{d}{d\bar{z}} Y(a, \underline{z})$.

By (FV4), $Y(D\mathbf{1}, \underline{z}) = \frac{d}{dz} Y(\mathbf{1}, \underline{z}) = 0$. Thus, by (FV3), $D\mathbf{1} = \bar{D}\mathbf{1} = 0$. Since $Y(D\bar{D}a, \underline{z}) = \frac{d}{dz} \frac{d}{d\bar{z}} Y(a, \underline{z}) = \frac{d}{d\bar{z}} \frac{d}{dz} Y(a, \underline{z}) = Y(\bar{D}Da, \underline{z})$, we have $[D, \bar{D}] = 0$.

By Lemma 2.3, $\mu_2|_{|z_2| > |z_1 - z_2|} \in \mathbb{C}[z_2, \bar{z}_2][z_0^\pm, \bar{z}_0^\pm, |z_0|^\mathbb{R}]$. Set $p(z_0, z_2) = \mu_2|_{|z_2| > |z_1 - z_2|} = u(Y(Y(a, \underline{z}_0)b, \underline{z}_2)\mathbf{1})$. Since $u(Y(Y(b, -\underline{z}_0)a, \underline{z}_1)\mathbf{1}) = p(z_0, z_1 - z_0)|_{|z_1| > |z_0|}$, we have

$$\begin{aligned} u(Y(a, \underline{z}_0)b) &= p(z_0, 0) = \lim_{z_1 \rightarrow 0} \exp(z_0 \frac{d}{dz_1} + \bar{z}_0 \frac{d}{d\bar{z}_1}) p(z_0, z_1 - z_0) \\ &= \lim_{z_1 \rightarrow 0} \exp(z_0 \frac{d}{dz_1} + \bar{z}_0 \frac{d}{d\bar{z}_1}) u(Y(Y(b, -\underline{z}_0)a, \underline{z}_1)\mathbf{1}) \\ &= \lim_{z_1 \rightarrow 0} u(Y(\exp(z_0 D + \bar{z}_0 \bar{D})Y(b, -\underline{z}_0)a, \underline{z}_1)\mathbf{1}) \\ &= u(\exp(z_0 D + \bar{z}_0 \bar{D})Y(b, -\underline{z}_0)a). \end{aligned}$$

Finally,

$$\begin{aligned} \frac{d}{dz} Y(a, \underline{z})b &= \frac{d}{dz} \exp(Dz + \bar{D}\bar{z})Y(b, -\underline{z})a \\ &= D \exp(Dz + \bar{D}\bar{z})Y(b, -\underline{z})a - \exp(Dz + \bar{D}\bar{z})Y(Db, -\underline{z})a \\ &= DY(a, \underline{z})b - Y(a, \underline{z})Db. \end{aligned}$$

□

Let $(V, Y, \mathbf{1})$ be a $\mathbb{Z}$ -graded vertex algebra. Then, by a standard result of the theory of a vertex algebra (see for example [FLM, FB]), $u(Y(a, z_1)Y(b, z_2)c)$ is an expansion of a rational polynomial in $\mathbb{C}[z_1^\pm, z_2^\pm, (z_1 - z_2)^{-1}] \subset \text{GCor}_2$ in $|z_1| > |z_2|$ for any $u \in V^\vee$ and $a, b, c \in V$. Thus, we have:

Proposition 2.2. *A $\mathbb{Z}$ -graded vertex algebra is a full vertex algebra.*

Let $(F, Y, \mathbf{1})$ be a full vertex algebra. Set $\bar{F} = F$ and $\bar{F}_{h, \bar{h}} = F_{h, h}$ for $h, \bar{h} \in \mathbb{R}$. Define $\bar{Y}(-, \underline{z}) : \bar{F} \rightarrow \text{End}(\bar{F})[[z, \bar{z}, |z|^{\mathbb{R}}]]$ by $\bar{Y}(a, \underline{z}) = \sum_{s, \bar{s} \in \mathbb{R}} a(s, \bar{s}) \bar{z}^{-s-1} z^{-\bar{s}-1}$. Let $C : Y_2 \rightarrow Y_2$ be the conjugate map $(z_1, z_2) \mapsto (\bar{z}_1, \bar{z}_2)$ for $(z_1, z_2) \in Y_2$. For $u \in \bar{F}^\vee$ and $a, b, c \in \bar{F}$, let $\mu \in \text{GCor}_2$ satisfy $u(Y(a, \underline{z}_1)Y(b, \underline{z}_2)c) = \mu(z_1, z_2)|_{|z_1| > |z_2|}$. Then, $u(\bar{Y}(a, \underline{z})\bar{Y}(b, \underline{z})c) = \mu \circ C(z_1, z_2)$. Since $\mu \circ C \in \text{GCor}_2$, we have:

Proposition 2.3. *$(\bar{F}, \bar{Y}, \mathbf{1})$ is a full vertex algebra.*

We call it a conjugate full vertex algebra of $(F, \mathbf{1}, Y)$.

2.2. Holomorphic vertex operators. Let F be a full vertex algebra. A vector $a \in F$ is said to be a holomorphic vector (resp. an anti-holomorphic vector) if $Da = 0$ (resp. $\bar{D}a = 0$). Let $a \in \ker \bar{D}$. Then, since $0 = Y(\bar{D}a, \underline{z}) = d/d\bar{z}Y(a, \underline{z})$, we have $a(r, s) = 0$ unless $s = -1$. Hence, $Y(a, \underline{z}) = \sum_{n \in \mathbb{Z}} a(n, -1)z^{-n-1}$.

Lemma 2.4. *Let $a, b \in F$. If $\bar{D}a = 0$, then for any $n \in \mathbb{Z}$,*

$$[a(n, -1), Y(b, \underline{z})] = \sum_{i \geq 0} \binom{n}{i} Y(a(i, -1)b, \underline{z})z^{n-i},$$

$$Y(a(n, -1)b, \underline{z}) = \sum_{i \geq 0} \binom{n}{i} (-1)^i a(n-i, -1)z^i Y(b, \underline{z}) - Y(b, \underline{z}) \sum_{i \geq 0} \binom{n}{i} (-1)^{i+n} a(i, -1)z^{n-i}.$$

Proof. For any $u \in F^\vee$ and $c \in F$, there exists $\mu \in \text{GCor}_2$ such that (FV5) holds. Since $\bar{D}a = 0$, by Proposition 2.1, $d/d\bar{z}_1\mu(z_1, z_2) = 0$. Then, by Lemma 1.22, $\mu \in \mathbb{C}[z_1^\pm, (z_1 - z_2)^\pm, z_2^\pm, \bar{z}_2^\pm, |z_2|^{\mathbb{R}}]$. Thus, by the Cauchy integral formula, the assertion holds. $\square$

By Proposition 2.1, $\bar{D}Y(a, \underline{z})b = Y(\bar{D}a, \underline{z})b + Y(a, \underline{z})\bar{D}b = 0$. Thus, the restriction of Y on $\ker \bar{D}$ define a linear map $Y(-, z) : \ker \bar{D} \rightarrow \text{End } \ker \bar{D}[[z^\pm]]$. By the above Lemma and Lemma 1.22, we have:

Proposition 2.4. *$\ker \bar{D}$ is a vertex algebra and F is a $\ker \bar{D}$ -module.*

Lemma 2.5. *For a holomorphic vector $a \in F$ and an anti-holomorphic vector $b \in F$, $[Y(a, z), Y(b, \bar{w})] = 0$, that is, $[a(n, -1), b(-1, m)] = 0$ and $a(k, -1)b = 0$ for any $n, m \in \mathbb{Z}$ and $k \in \mathbb{Z}_{\geq 0}$.*

Proof. By Lemma 2.4, it suffices to show that $a(k, -1)b = 0$ for any $k \geq 0$. Since $DY(a, z)b = [D, Y(a, z)]b + Y(a, z)Db = d/dzY(a, z)b$, we have $Da(n, -1)b = -na(n-1, -1)b$ for any $n \in \mathbb{Z}$. Thus, the assertion follows from (FV1). $\square$

2.3. Tensor product of full vertex algebras. In this section, we define a tensor product of full vertex algebras and study the subalgebra of a full vertex algebra generated by holomorphic and anti-holomorphic vectors.

Let F be a full vertex algebra. For $h, \bar{h} \in \mathbb{R}$, the energy and spin of a vector in $F_{h, \bar{h}}$ are $h + \bar{h}$ and $h - \bar{h}$ and the set $\{(h, \bar{h}) \in \mathbb{R}^2 \mid F_{h, \bar{h}} \neq 0\}$ is called a spectrum. The spectrum of F is said to be *bounded below* if there exists $N \in \mathbb{R}$ such that $F_{h, \bar{h}} = 0$ for any $h \leq N$ or $\bar{h} \leq N$ and *discrete* if for any $H \in \mathbb{R}$, $\sum_{h+\bar{h} < H} \dim F_{h, \bar{h}}$ is finite.

Lemma 2.6. *If the spectrum of F is discrete, then for any $N \in \mathbb{Z}_{>0}$ and $a, b \in F$, the number of the set*

$$\{(s, \bar{s}) \in \mathbb{R}^2 \mid a(s, \bar{s})b \neq 0, -N \leq s, \bar{s} \leq N\}$$

is finite.

Proof. We may assume that $a \in F_{h, \bar{h}}$ and $b \in F_{h', \bar{h}'}$. Since $a(s, \bar{s})b \in F_{h+h'-s-1, \bar{h}+\bar{h}'-\bar{s}-1}$, the energies of vectors $\{a(s, \bar{s})b \mid -N \leq s, \bar{s} \leq N\}$ are bounded by $h + h' + \bar{h} + \bar{h}' + 2N - 2$. Thus, the assertion holds. $\square$

Let $(F^1, Y^1, \mathbf{1}^1)$ and $(F^2, Y^2, \mathbf{1}^2)$ be full vertex algebras and assume that the spectrum of F^1 is discrete and the spectrum of F^2 is bounded below. Define the linear map $Y(-, \underline{z}) : F^1 \otimes F^2 \rightarrow \text{End} F^1 \otimes F^2[[z, \bar{z}, |z|^{\mathbb{R}}]]$ by $Y(a \otimes b, \underline{z}) = Y^1(a, \underline{z}) \otimes Y^2(b, \underline{z})$ for $a \in F^1$ and $b \in F^2$. Then, for $a, c \in F^1$ and $b, d \in F^2$,

$$Y(a \otimes b, \underline{z})c \otimes d = \sum_{s, \bar{s}, r, \bar{r} \in \mathbb{R}} a(s, \bar{s})c \otimes b(r, \bar{r})d z^{-s-r-2} \bar{z}^{-\bar{s}-\bar{r}-2}.$$

By (FV1) and the above lemma, the coefficient of $z^k \bar{z}^{\bar{k}}$ is a finite sum for any $k, \bar{k} \in \mathbb{R}$. Thus, $Y(-, \underline{z})$ is well-defined. For any $h_0, \bar{h}_0 \in \mathbb{R}$, set $(F^1 \otimes F^2)_{h_0, \bar{h}_0} = \bigoplus_{a, \bar{a} \in \mathbb{R}} F_{a, \bar{a}}^1 \otimes F_{h_0-a, \bar{h}_0-\bar{a}}^2$. Since the spectrum of F^2 is bounded below, there exists $N \in \mathbb{R}$ such that $(F^1 \otimes F^2)_{h_0, \bar{h}_0} = \bigoplus_{a, \bar{a} \leq N} F_{a, \bar{a}}^1 \otimes F_{h_0-a, \bar{h}_0-\bar{a}}^2$. Since the spectrum of F^1 is discrete, the sum is finite. Thus, $(F^1 \otimes F^2)_{h_0, \bar{h}_0}^* = \bigoplus_{a, \bar{a} \in \mathbb{R}} (F_{a, \bar{a}}^1)^* \otimes (F_{h_0-a, \bar{h}_0-\bar{a}}^2)^*$, which implies that $F^\vee = (F^1)^\vee \otimes (F^2)^\vee$. Let $u_i \in (F^i)^\vee$ and $a_i, b_i, c_i \in F^i$ for $i = 1, 2$. Since

$$u_1 \otimes u_2 (Y(a_1 \otimes a_2, \underline{z}_1) Y(b_1 \otimes b_2, \underline{z}_2) c_1 \otimes c_2) = u_1 (Y(a_1, \underline{z}_1) Y(b_1, \underline{z}_2) c_1) u_2 (Y(a_2, \underline{z}_1) Y(b_2, \underline{z}_2) c_2),$$

we have:

Proposition 2.5. *Let $(F^1, Y^1, \mathbf{1}^1)$ and $(F^2, Y^2, \mathbf{1}^2)$ be full vertex algebras. If the spectrum of F^1 is discrete and the spectrum of F^2 is bounded below, then $(F^1 \otimes F^2, Y^1 \otimes Y^2, \mathbf{1}^1 \otimes \mathbf{1}^2)$ is a full vertex algebra. Furthermore, if the spectrum of F^1 and F^2 are bounded below (resp. discrete), then the spectrum of $F_1 \otimes F^2$ is also bounded below (resp. discrete).*

By Proposition 2.2 and Proposition 2.3, we have:

Corollary 2.1. *Let V, W be a $\mathbb{Z}_{\geq 0}$ -graded vertex algebras such that $\dim V_n$ and $\dim W_n$ are finite for any $n \in \mathbb{Z}_{\geq 0}$. Then, $V \otimes \bar{W}$ is a full vertex algebra with a discrete spectrum, where $\bar{W}$ is the conjugate full vertex algebra.*

Let F be a full vertex algebra. By Proposition 2.4, $\ker \bar{D}$ and $\ker D$ are subalgebras of F . Let $\ker \bar{D} \otimes \ker D$ be the tensor product full vertex algebra. Define the linear map $t : \ker \bar{D} \otimes \ker D \rightarrow F$ by $(a \otimes b) \mapsto a(-1, -1)b$ for $a \in \ker \bar{D}$ and $b \in \ker D$. Then, we have:

Proposition 2.6. *Let F be a full vertex algebra. Then, $t : \ker \bar{D} \otimes \ker D \rightarrow F$ is a full vertex algebra homomorphism.*

Proof. Let $a, c \in \ker \bar{D}$, $b, d \in \ker D$. By Lemma 2.5 and Lemma 2.4,

$$Y(a(-1, -1)b, z) = Y(a, z)Y(b, \bar{z}) = Y(b, \bar{z})Y(a, z).$$

Thus, it suffices to show that $t(a \otimes b(n, m)c \otimes d) = t(a \otimes b)(n, m)t(c \otimes d)$ for any $n, m \in \mathbb{Z}$. By Lemma 2.4

$$\begin{aligned} t(a \otimes b(n, m)c \otimes d) &= t(a(n, -1)c \otimes b(-1, m)d) \\ &= (a(n, -1)c)(-1, -1)b(-1, m)d \\ &= \sum_{i=0}^n \binom{n}{i} (-1)^i (a(n-i, -1)c(-1+i, -1) + c(-1+n-i, -1)a(i, -1))b(-1, m)d. \end{aligned}$$

Since $b(-1, m)d \in \ker D$, by Lemma 2.5, $t(a \otimes b(n, m)c \otimes d) = a(n, -1)c(-1, -1)b(-1, m)d = a(n, -1)b(-1, m)c(-1, -1)d = t(a \otimes b)(n, m)t(c \otimes d)$. Thus, the assertion holds. $\square$

We remark that if $\ker \bar{D} \otimes \ker D$ is simple, then the above map is injective.

3. CORRELATION FUNCTIONS AND FULL VERTEX ALGEBRAS

3.1. Self-duality. A full vertex operator algebra (full VOA) is a full vertex algebra F with a holomorphic vector $\omega \in F$ and an anti-holomorphic vector $\bar{\omega} \in F$ satisfying the following conditions:

FVO1) There exist a pair of scalars $(c, \bar{c}) \in \mathbb{C}$ such that

$$\begin{aligned} [L(n), L(m)] &= (n-m)L(n+m) + \frac{n^3-n}{12}\delta_{n+m,0}c, \\ [\bar{L}(n), \bar{L}(m)] &= (n-m)\bar{L}(n+m) + \frac{n^3-n}{12}\delta_{n+m,0}\bar{c} \end{aligned}$$

holds for any $n, m \in \mathbb{Z}$, where $L(n) = \omega(n+1, -1)$ and $\bar{L}(n) = \bar{\omega}(-1, n+1)$;

FVO2) $D = L(-1)$ and $\bar{D} = \bar{L}(-1)$;

FVO3) $L(0)|_{F_{h,\bar{h}}} = h$ and $\bar{L}(0)|_{F_{h,\bar{h}}} = \bar{h}$ for any $h, \bar{h} \in \mathbb{R}^2$;

FVO4) $F_{h,\bar{h}}$ is a finite dimensional vector space for any $h, \bar{h} \in \mathbb{R}$;

FVO5) The spectrum of F is bounded below, that is, there exists $N \in \mathbb{R}$ such that $F_{h,\bar{h}} = 0$ for any $h \leq N$ or $\bar{h} \leq N$.

The pair of scalars $(c, \bar{c})$ is called a central charge and the pair $(\omega, \bar{\omega})$ is called an *energy-momentum tensor* of the full vertex operator algebra F . A module M of a full vertex algebra F is said to be a module of a full vertex operator algebra if it satisfies

FVOM1) $L(0)|_{M_{h,\bar{h}}} = h$ and $\bar{L}(0)|_{M_{h,\bar{h}}} = \bar{h}$ for any $h, \bar{h} \in \mathbb{R}$;

FVOM2) $M_{h,\bar{h}}$ is a finite dimensional vector space for any $h, \bar{h} \in \mathbb{R}$;

FVOM3) The spectrum of M is bounded below.

Let F be a full VOA and M be a F -module. By Lemma 2.4, we have:

Lemma 3.1. *For $h, \bar{h} \in \mathbb{R}$ and $a \in F_{h,\bar{h}}$,*

$$[L(0), Y(a, \underline{z})] = Y((L(0) + zL(-1))a, \underline{z}) = (zd/dz + h)Y(a, \underline{z}),$$

$$[\bar{L}(0), Y(a, \underline{z})] = Y((\bar{L}(0) + \bar{z}\bar{L}(-1))a, \underline{z}) = (\bar{z}d/d\bar{z} + \bar{h})Y(a, \underline{z}),$$

$$[L(1), Y(a, \underline{z})] = Y((L(1) + 2zL(0) + z^2L(-1))a, \underline{z}) = (z^2d/dz + 2hz)Y(a, \underline{z}) + Y(L(1)a, \underline{z}),$$

$$[\bar{L}(1), Y(a, \underline{z})] = Y((\bar{L}(1) + 2\bar{z}\bar{L}(0) + \bar{z}^2\bar{L}(-1))a, \underline{z}) = (\bar{z}^2d/d\bar{z} + 2\bar{h}\bar{z})Y(a, \underline{z}) + Y(\bar{L}(1)a, \underline{z}).$$

Set $M^\vee = \bigoplus_{h,\bar{h} \in \mathbb{R}} M_{h,\bar{h}}^*$ and let $\langle \cdot, \cdot \rangle: M^\vee \times M \rightarrow \mathbb{C}$ be a canonical pairing. Define $S_z: M \rightarrow M[z^\pm, \bar{z}^\pm, |z|^\mathbb{R}]$, by

$$S_z a = \exp(zL(1) + \bar{z}\bar{L}(1))(-1)^{h-\bar{h}} z^{-2h} \bar{z}^{-2\bar{h}},$$

for $h, \bar{h} \in \mathbb{R}$ and $a \in M_{h,\bar{h}}$, where we used the fact that $L(1)$ and $\bar{L}(1)$ are locally nilpotent by (FVOM3). Define the vertex operator $Y_{M^\vee}(-, \underline{z}): F \rightarrow \text{End} M^\vee[[z^\pm, \bar{z}^\pm, |z|^\mathbb{R}]]$ by

$$\langle Y_{M^\vee}(a, \underline{z})u, v \rangle \equiv \langle u, Y_M(S_z a, \underline{z}^{-1})v \rangle,$$

for $a \in F$ and $v \in M$ and $u \in M^\vee$.

We will prove the following Proposition:

Proposition 3.1. *$(M^\vee, Y_{M^\vee}(-, \underline{z}))$ is a module of the full VOA F .*

Properties of the operators $\{L(i)\}_{i=-1,0,1}$ acting on a vertex operator algebra which satisfy the equations in Lemma 3.1 are studied in [FHL], which can be easily generalized to a full vertex operator algebra.

Lemma 3.2 ([FHL]). *For a full vertex operator algebra F and its module M , the following equations hold:*

$$(1) [L(0), \exp(L(1)z)] = -L(1)z \exp(L(1)z).$$

$$(2) \text{ For any } a \in F, S_{z_2} Y_M(a, \underline{z}_0) = Y_M(S_{z_2+z_0} a, \frac{-z_0}{z_2(z_2+z_0)}) S_{z_2},$$

where $z_2 + z_0$ is expanded in $|z_2| > |z_0|$.

proof of Proposition 3.1. Let $u \in F^\vee$ and $v \in M$ and $a, b \in F$. By Lemma 2.5 and (FVO1), $L(1)\omega = L(1)\bar{\omega} = \bar{L}(1)\omega = \bar{L}(1)\bar{\omega} = 0$. Thus, $\langle Y(\omega, \underline{z})u, v \rangle = \langle u, z^4 Y(\omega, \underline{z}^{-1})v \rangle$, which implies that $\langle L(n)u, v \rangle = \langle u, L(-n)v \rangle$. By Lemma 3.1 and Lemma 3.2, Proposition 2.1,

$$\begin{aligned} & \langle [L(0), Y_{M^\vee}(a, \underline{z})]u, v \rangle = \langle Y_{M^\vee}(L(0)a, \underline{z})u, v \rangle \\ & = \langle u, Y(S_z a, \underline{z}^{-1})L(0)v \rangle = \langle L(0)u, Y(S_z a, \underline{z}^{-1})v \rangle = \langle u, Y(S_z L(0)a, \underline{z}^{-1})v \rangle \\ & = -\langle u, Y((L(0) + z^{-1}L(-1))S_z a, \underline{z}^{-1})v \rangle = -\langle u, Y(S_z L(0)a, \underline{z}^{-1})v \rangle \\ & = \langle u, Y(L(1)z S_z a, \underline{z}^{-1})v \rangle = \langle u, Y(S_z 2L(0)a, \underline{z}^{-1})v \rangle = -z^{-1} \langle u, [L(-1), Y(S_z a, \underline{z}^{-1})]v \rangle. \end{aligned}$$

Since

$$\begin{aligned} z d/dz < u, Y(S_z a, \underline{z}^{-1}) v > &= < u, Y(L(1) z S_z a, \underline{z}^{-1}) v > \\ &- < u, Y(L(1) z S_z 2L(0) a, \underline{z}^{-1}) v > - z^{-1} < u, [L(-1), Y(S_z a, \underline{z}^{-1})] v >, \end{aligned}$$

we have $[L(0), Y_{M^\vee}(a, \underline{z})] = Y_{M^\vee}(L(0)a, \underline{z}) + d/dz Y_{M^\vee}(a, \underline{z})$. Thus, by Remark 2.2, $F_{h, \bar{h}}(r, s) M_{h', \bar{h}'}^\vee \subset M_{h+h'-r-1, \bar{h}+\bar{h}'-s-1}^\vee$. (FM2), (FM3), (FVOM1), (FVOM2) and (FVOM3) clearly follows. (FM1) follows from (FM5) together with (FVOM3). Thus, it suffices to show (FM4). By (FM4), there exists $\mu \in \text{GCor}_2$ such that

$$\begin{aligned} < u, Y(S_{z_2^{-1}} b, \underline{z}_2) Y(S_{z_1^{-1}} a, \underline{z}_1) v > &= \mu|_{|z_2| > |z_1|}, \\ < u, Y(S_{z_1^{-1}} a, \underline{z}_1) Y(S_{z_2^{-1}} b, \underline{z}_2) v > &= \mu|_{|z_1| > |z_2|}, \\ \lim_{z_1 \rightarrow (z_2 + z_0)|_{|z_2| > |z_0|}} < u, Y(Y(S_{z_1^{-1}} a, \underline{z}_0) S_{z_2^{-1}} b, \underline{z}_2) v > &= \mu|_{|z_2| > |z_1 - z_2|}. \end{aligned}$$

Then, $< Y(a, \underline{z}_1) Y(b, \underline{z}_2) u, m > = I_Y(\mu)|_{|z_1| > |z_2|}$ and $< Y(b, \underline{z}_2) Y(a, \underline{z}_1) u, m > = I_Y(\mu)|_{|z_2| > |z_1|}$. To show (FM4), it suffices to show that $I_Y(\mu)|_{|z_2| > |z_1 - z_2|} = < Y(Y(a, \underline{z}_0) b, \underline{z}_2) u, v >$. By Lemma 1.24,

$$\begin{aligned} I_Y(\mu)|_{|z_2| > |z_1 - z_2|} &= \lim_{(z_0, z_2) \mapsto (\frac{-z_0}{z_2(z_2 + z_0)}, \frac{1}{z_2})} \mu|_{|z_2| > |z_1 - z_2|} \\ &= \lim_{(z_0, z_2) \mapsto (\frac{-z_0}{z_2(z_2 + z_0)}, \frac{1}{z_2})} \lim_{z_1 \rightarrow z_2 + z_0} < u, Y(Y(S_{z_1^{-1}} a, \underline{z}_0) S_{z_2^{-1}} b, \underline{z}_2) v > \\ &= < u, Y(Y(S_{z_2 + z_0} a, \frac{-\underline{z}_0}{\underline{z}_2(\underline{z}_2 + \underline{z}_0)}) S_{z_2} b, \underline{z}_2^{-1}) v >. \end{aligned}$$

By Lemma 3.2,

$$\begin{aligned} < Y_M^\vee(Y(a, \underline{z}_0) b, \underline{z}_2) u, v > &= < u, Y(S_{z_2} Y(a, \underline{z}_0) b, \underline{z}_2^{-1}) v > \\ &= < u, Y(Y(S_{z_2 + z_0} a, \frac{-\underline{z}_0}{\underline{z}_2(\underline{z}_2 + \underline{z}_0)}) S_{z_2} b, \underline{z}_2^{-1}) v >. \end{aligned}$$

Thus, the assertion holds. $\square$

An invariant bilinear form on a full vertex operator algebra F is a bilinear form $(-, -) : F \times F \rightarrow \mathbb{C}$ such that

$$(Y(a, \underline{z}) b, c) = (b, Y(S_z a, \underline{z}^{-1}) c)$$

holds for any $a, b, c \in F$. By the proof of Proposition 3.1, $(L(n)a, b) = (a, L(-n)b)$, which implies that $F_{h, \bar{h}}$ and $F_{h', \bar{h}'}$ are orthogonal to each other unless $(h, \bar{h}) = (h', \bar{h}')$.

The following proposition is a straightforward generalization of Li's result on the invariant bilinear form on a vertex operator algebra (see [L1]):

Proposition 3.2. *There is a one-to-one correspondence between $\text{Hom}_{\mathbb{C}}(F_{0,0}/(L(1)F_{1,0} + \bar{L}(1)F_{0,1}), \mathbb{C})$ and the space of invariant bilinear forms on F .*

According to [L1], we will use the following lemma:

Lemma 3.3. *Let M be an F -module. A vector $v \in M$ is a vacuum-like vector if and only if $L(-1)v = \bar{L}(-1)v = 0$.*

Proof. If v is a vacuum-like vector, then $Y(\omega, z)v \in F[[z]]$. Thus, $L(-1)v = 0$ and similarly $\bar{L}(-1)v = 0$. Assume that $L(-1)v = \bar{L}(-1)v = 0$. Then, $L(-1)Y_M(a, \underline{z})v = [L(-1), Y_M(a, \underline{z})]v = d/dz Y_M(a, \underline{z})v$. Thus, similarly to the proof of Lemma 2.5, the assertion holds. $\square$

proof of Proposition 3.2. Let I_F be the space of invariant bilinear forms on F and $(-, -) \in I_F$. Then, $(\mathbf{1}, -) : F \rightarrow \mathbb{C}$, $a \mapsto (\mathbf{1}, a)$ satisfies $(\mathbf{1}, F_{h, \bar{h}}) = 0$ for any $(h, \bar{h}) \neq (0, 0)$. Since $L(-1)\mathbf{1} = \bar{L}(-1)\mathbf{1} = 0$, we have $(\mathbf{1}, L(1)-) = (\mathbf{1}, \bar{L}(1)-) = 0$. Thus, we have a linear map $\rho : I_F \rightarrow \text{Hom}_{\mathbb{C}}(F_{0,0}/(L(1)F_{1,0} + \bar{L}(1)F_{0,1}), \mathbb{C})$. Assume that $(\mathbf{1}, -) = 0$. Then, for any $a, b \in F$, $(a, b) = \lim_{\underline{z} \rightarrow 0} (Y(a, \underline{z})\mathbf{1}, b) = \lim_{\underline{z} \rightarrow 0} (\mathbf{1}, Y(S_z a, \underline{z}^{-1})b) = 0$. Hence, ρ is injective.

Let $u \in \text{Hom}_{\mathbb{C}}(F_{0,0}/(L(1)F_{1,0} + \bar{L}(1)F_{0,1}), \mathbb{C})$. Then, u is an element of $F_{0,0}^* \subset F^\vee$. Since $\langle L(-1)u, a \rangle = \langle u, L(1)a \rangle$ for any $a \in F$, by Lemma 3.3, u is a vacuum-like vector. Hence, by Lemma 2.3, we have a F -module homomorphism $F_u : F \rightarrow F^\vee$, $a \mapsto a(-1, -1)u$. Define the bilinear form $(-, -)_u : F \times F \rightarrow \mathbb{C}$ by $(a, b)_u = \langle F_u(a), b \rangle$ for $a, b \in F$. Then,

$$\begin{aligned} (Y(a, \underline{z})b, c)_u &= \langle F_u(Y(a, \underline{z})b), c \rangle = \langle Y_{F^\vee}(a, \underline{z})F_u(b), c \rangle \\ &= \langle F_u(b), Y(S_z a, \underline{z}^{-1})c \rangle = (b, Y(S_z a, \underline{z}^{-1})c)_u, \end{aligned}$$

which implies that $(-, -)_u$ is an invariant bilinear form. Since $(\mathbf{1}, -)_u = \langle F_u(\mathbf{1}), b \rangle = \langle u, b \rangle$, $\rho((-, -)_u) = u$. Thus, ρ is an isomorphism. $\square$

Similarly to [FHL], we have:

Proposition 3.3. *An invariant bilinear form on F is symmetric.*

Proposition 3.4. *A full vertex operator algebra F is simple if and only if $F^\vee$ is simple.*

A full vertex operator algebra F is said to be self-dual if F is isomorphic to $F^\vee$ as an F -module, or equivalently, there exists a non-degenerate invariant bilinear form on F .

Corollary 3.1. *If F is simple and $F_{0,0} = \mathbb{C}\mathbf{1}$ and $L(1)F_{1,0} = \bar{L}(1)F_{0,1} = 0$, then F is self-dual.*

For a self-dual full vertex operator algebra F and $a \in F$, $(\mathbf{1}, Y(a, \underline{z})\mathbf{1}) = (\mathbf{1}, \exp(Dz + \bar{D}\bar{z})a) = (\exp(L(1)z + \bar{L}(1)\bar{z})\mathbf{1}, a) = (\mathbf{1}, a)$. Thus, we have:

Lemma 3.4. *For $a \in F$, $(\mathbf{1}, Y(a, \underline{z})\mathbf{1})$ is equal to $(\mathbf{1}, a) \in \mathbb{C}$.*

3.2. Quasi-primary vectors. For $h, \bar{h} \in \mathbb{R}$, set

$$QF_{h, \bar{h}} = \{v \in F_{h, \bar{h}} \mid L(1)v = \bar{L}(1)v = 0\}.$$

A homogeneous vector in $QF = \bigoplus_{h, \bar{h} \in \mathbb{R}} QF_{h, \bar{h}}$ is called a quasi-primary vector. Recall that $L(i), \bar{L}(i)$ ($i = -1, 0, 1$) generate the Lie algebra $\mathfrak{sl}_2 \oplus \mathfrak{sl}_2$. Following the terminology of [L1], we call a full vertex operator algebra F *QP-generated* if F is generated by QF as a $\mathfrak{sl}_2 \oplus \mathfrak{sl}_2$ -module, or equivalently, $F = \mathbb{C}[L(-1), \bar{L}(-1)]QF$.

Proposition 3.5. *A self-dual full vertex operator algebra F is QP-generated if and only if the following conditions hold:*

- (1) $F_{h,\bar{h}} = 0$ if $h \in \frac{1}{2}\mathbb{Z}_{<0}$ or $\bar{h} \in \frac{1}{2}\mathbb{Z}_{<0}$.
- (2) $L(1)F_{1,n} = 0$ and $\bar{L}(1)F_{n,1} = 0$ for any $n \in \mathbb{Z}$.
- (3) $L(-1)F_{0,n} = 0$ and $\bar{L}(-1)F_{n,0} = 0$ for any $n \in \mathbb{Z}$.

Lemma 3.5. *If F is self-dual and QP-generated, then the restriction of the invariant bilinear form on QF is non-degenerate. Furthermore, $\text{Im } L(-1) \cap \ker L(1) = 0$ and $\text{Im } \bar{L}(-1) \cap \ker \bar{L}(1) = 0$ hold.*

Proof. Assume that $a \in QF_{h,\bar{h}}$ satisfy $(a, b) = 0$ for any $b \in QF_{h,\bar{h}}$. Since $(a, L(-1)b) = (L(1)a, b) = 0$ and $(a, \bar{L}(-1)b) = (\bar{L}(1)a, b) = 0$ for any $b \in F$, by $F = \mathbb{C}[L(-1), \bar{L}(-1)]QP$, $a = 0$, which proves the first part of the lemma. Set $QF' = \{v \in F \mid L(1)v = 0\}$. Since $F = \mathbb{C}[L(-1)][\bar{L}(-1)]QF$ and $L(1)\mathbb{C}[\bar{L}(-1)]QF = 0$, we have $F = \mathbb{C}[L(-1)]QF'$. Then, by the same proof, the restriction of the invariant bilinear form on QF' is non-degenerate. Since $\text{Im } L(-1) \cap \ker L(1)$ is in the radical of this invariant bilinear form, the assertion holds. $\square$

proof of Proposition 3.5. Suppose that F is QP-generated. Let $0 \neq v \in QF_{-n,m}$ for $n \in \frac{1}{2}\mathbb{Z}_{>0}$. Then, $L(1)L(-1)^{2n+1}v = 0$. By the representation theory of sl_2 and Lemma 3.5, we have $0 \neq L(-1)^{2n}v \in F_{n,m}$ and $L(-1)^{2n+1}v = 0$. Since $L(-1)^{2n+1}v = 0$ implies $L(0)L(-1)^{2n}v = 0$, a contradiction. Hence, $QF_{-n,m} = 0$ for any $n \in \frac{1}{2}\mathbb{Z}_{>0}$.

Let $v \in F_{0,n}$. Since $L(1)L(-1)v = 0$, by Lemma 3.5, $L(-1)v = 0$, which implies (3). Since $F = \mathbb{C}[L(-1), \bar{L}(-1)]QF$, by (1) and (3), $F_{1,n} = \bigoplus_{k \geq 0} \bar{L}(-1)^k QF_{1,n-k}$. Since $L(1)$ commutes with $\bar{L}(-1)$, we have $L(1)F_{1,n} = 0$, thus, (2).

Suppose that F satisfies (1), (2) and (3). Let $a, b \in QF_{n,m}$ with $n, m \in \mathbb{R} \setminus \frac{1}{2}\mathbb{Z}_{<0}$. Since

$$(L(-1)^k \bar{L}(-1)^l a, L(-1)^k \bar{L}(-1)^l b) = k!l! \prod_{\substack{k-1 \geq i \geq 0 \\ l-1 \geq j \geq 0}} (2n+i)(2m+j)(a, b),$$

the restriction of the bilinear form on $L(-1)^k \bar{L}(-1)^l Q_{n,m}$ is non-degenerate if $(-, -)|_{Q_{n,m}}$ is non-degenerate. Let $h, \bar{h} \in \mathbb{R}$ satisfy $F_{h,\bar{h}} \neq 0$ and $F_{h-k, \bar{h}-l} = 0$ for any $k, l \in \mathbb{Z}_{>0}$. Then, $QF_{h,\bar{h}} = F_{h,\bar{h}}$, which implies that $(-, -)|_{QF_{h,\bar{h}}}$ is non-degenerate. Since $QF_{h+1,\bar{h}} = \{v \in F_{h+1,\bar{h}} \mid (v, L(-1)F_{h,\bar{h}}) = 0\}$, $QF_{h+1,\bar{h}}$ is non-degenerate and $F_{h+1,\bar{h}} = QF_{h+1,\bar{h}} \oplus L(-1)Q_{h,\bar{h}}$. Similarly, we have $F_{a+k,b+l} = \bigoplus_{\substack{0 \leq i \leq k \\ 0 \leq j \leq l}} L(-1)^{k-i} \bar{L}(-1)^{l-j} QF_{a+i,b+j}$ by the induction on $k, l \in \mathbb{Z}_{\geq 0}$. $\square$

By Proposition 3.5 and Lemma 3.1, we have:

Corollary 3.2. *Suppose that simple full vertex operator algebra F satisfies the following conditions:*

- PN1) $F_{0,0} = \mathbb{C}\mathbf{1}$;
- PN2) $F_{n,m} = 0$ if $n \in \frac{1}{2}\mathbb{Z}_{<0}$ or $m \in \frac{1}{2}\mathbb{Z}_{<0}$;
- PN3) $L(1)F_{1,n} = 0$ and $\bar{L}(1)F_{n,1} = 0$ for any $n \in \mathbb{Z}$;
- PN4) $L(-1)F_{0,n} = 0$ and $\bar{L}(-1)F_{n,0} = 0$ for any $n \in \mathbb{Z}$.

Then, F is self-dual and QP -generated.

3.3. Parenthesized correlation functions and formal power series. Let F be a self-dual vertex operator algebra and denote $Y(a, \bar{x})$ by $a(x)$ in this section for $a \in F$. For $n \in \mathbb{Z}_{>0}$, let P_n be the set of parenthesized products of n elements $1, 2, \dots, n$, e.g., $((32)1)((47)(65)) \in P_7$ and Q_n be the set of parenthesized products of $n+1$ elements $1, 2, \dots, n, \star$ with $\star$ at right most, e.g., $((32)1)((4(65))\star) \in Q_6$. We can naturally associate a parenthesized n point correlation function to each element in P_n and Q_n (see also Introduction 0.2 and Section 1.3). To give a simple example, an element $((31)6)(24))(57) \in P_7$ and $a_1, \dots, a_7 \in F$ define the following parenthesized 7 point correlation function:

$$S_{((31)6)(24))(57)}(a_1, a_2, a_3, a_4, a_5, a_6, a_7) := \left(\mathbf{1}, \left[\left((a_3(x_3)a_1)(x_1)a_6 \right) (x_6)a_2(x_2)a_4 \right] (x_4)a_5(x_5)a_7 \right).$$

For an element in Q_n and $a_1, \dots, a_n \in F$, we consider parenthesized $n+1$ point correlation for states $a_1, \dots, a_n, \mathbf{1}$ where $\mathbf{1}$ corresponds to $\star$, e.g., for $((31)6)(24))(5\star) \in Q_6$,

$$S_{((31)6)(24))(5\star)}(a_1, a_2, a_3, a_4, a_5, a_6) := \left(\mathbf{1}, \left[\left((a_3(x_3)a_1)(x_1)a_6 \right) (x_6)a_2(x_2)a_4 \right] (x_4)a_5(x_5)\mathbf{1} \right).$$

In this subsection, we study the space of formal variables for each $A \in P_n$ and $A \in Q_n$.

We start from $A = 1(2(\dots(n-1n))\dots) \in P_n$. In this case, for $a_1, \dots, a_n \in F$, the parenthesized correlation function is

$$S_{1(2(\dots(n-1n))\dots)}(a_1, \dots, a_n) = (\mathbf{1}, a_1(x_1)a_2(x_2) \dots a_{n-1}(x_{n-1})a_n).$$

Set

$$\begin{aligned} & T_{1(2(\dots(n-1n))\dots)}(x_1, \dots, x_{n-1}) \\ &= \mathbb{C}[[x_2/x_1, \bar{x}_2/\bar{x}_1]]((x_3/x_2, \bar{x}_3/\bar{x}_2, |x_3/x_2|^{\mathbb{R}}, \dots, x_{n-1}/x_{n-2}, \bar{x}_{n-1}/\bar{x}_{n-2}, |x_{n-1}/x_{n-2}|^{\mathbb{R}})) \\ & \quad [x_1^{\pm}, \bar{x}_1^{\pm}, |x_1|^{\mathbb{R}}, x_2^{\pm}, \bar{x}_2^{\pm}, |x_2|^{\mathbb{R}}]. \end{aligned}$$

For $h, \bar{h}, h', \bar{h}' \in \mathbb{R}$ and $a \in F_{h, \bar{h}}$ and $b \in F_{h', \bar{h}'}$, since

$$\begin{aligned} (\mathbf{1}, Y(a, \underline{z})b) &= (Y(S_{\underline{z}}a, \underline{z}^{-1})\mathbf{1}, b) \\ &= (\exp(D\underline{z}^{-1} + \bar{D}\bar{\underline{z}}^{-1})S_{\underline{z}}a, b) \end{aligned}$$

and $F_{h, \bar{h}}$ and $F_{h', \bar{h}'}$ is orthogonal if $(h, \bar{h}) \neq (h', \bar{h}')$, we have:

Lemma 3.6. For $h, \bar{h}, h', \bar{h}' \in \mathbb{R}$ and $a \in F_{h, \bar{h}}$ and $b \in F_{h', \bar{h}'}$, $(\mathbf{1}, Y(a, \underline{z})b) = 0$ unless $h - h' \in \mathbb{Z}$ and $\bar{h} - \bar{h}' \in \mathbb{Z}$.

Then, we have:

Lemma 3.7. For $a_1, \dots, a_n \in F$,

$$S_{1(2(\dots(n-1n))\dots)}(a_1, \dots, a_n; x_1, \dots, x_{n-1}) \in T_{1(2(\dots(n-1n))\dots)}(x_1, \dots, x_{n-1}).$$

Proof. We may assume that $a_i \in F_{h_i, \bar{h}_i}$ for $h_i, \bar{h}_i \in \mathbb{R}$ and $i = 1, \dots, n$. Set $h = h_1 + \dots + h_n$ and $\bar{h} = \bar{h}_1 + \dots + \bar{h}_n$ and

$$\sum_{s_1, \bar{s}_1, \dots, s_{n-1}, \bar{s}_{n-1} \in \mathbb{R}} c_{s_1, \bar{s}_1, \dots, s_{n-1}, \bar{s}_{n-1}} z_1^{s_1} \bar{z}_1^{\bar{s}_1} \dots z_{n-1}^{s_{n-1}} \bar{z}_{n-1}^{\bar{s}_{n-1}} = (\mathbf{1}, Y(a_1, \underline{z}_1) Y(a_2, \underline{z}_2) \dots Y(a_{n-1}, \underline{z}_{n-1}) a_n).$$

Then,

$$c_{s_1, \bar{s}_1, \dots, s_{n-1}, \bar{s}_{n-1}} = (\mathbf{1}, a_1(-s_1-1, -\bar{s}_1-1) a_2(-s_2-1, -\bar{s}_2-1) \dots a_{n-1}(-s_{n-1}-1, -\bar{s}_{n-1}-1) a_n).$$

By (FVO5), there exists $N \in \mathbb{R}$ such that $F_{h, \bar{h}} = 0$ for any $h \leq N$ or $\bar{h} \leq N$. Since for any $k = 1, 2, \dots, n-1$, $a_k(-s_k-1, -\bar{s}_k-1) a_{k-1}(-s_{k-1}-1, -\bar{s}_{k-1}-1) \dots a_{n-1}(-s_{n-1}-1, -\bar{s}_{n-1}-1) a_n$ is in $F_{h_k + \dots + h_n + s_k + \dots + s_{n-1}, \bar{h}_k + \dots + \bar{h}_n + \bar{s}_k + \dots + \bar{s}_{n-1}}$, by Lemma 3.6 and (FVO5), $c_{s_1, \bar{s}_1, \dots, s_{n-1}, \bar{s}_{n-1}} = 0$ unless $0 = h + s_1 + \dots + s_{n-1}$, $0 = \bar{h} + \bar{s}_1 + \dots + \bar{s}_{n-1}$ and $(h - h_1) + s_2 + \dots + s_{n-1} - h_1 \in \mathbb{Z}$, $(\bar{h} - \bar{h}_1) + \bar{s}_2 + \dots + \bar{s}_{n-1} - \bar{h}_1 \in \mathbb{Z}$ and $h_k + \dots + h_n + s_k + \dots + s_{n-1} \geq N$, $\bar{h}_k + \dots + \bar{h}_n + \bar{s}_k + \dots + \bar{s}_{n-1} \geq N$ for any $k = 1, \dots, n-1$. Thus, we have

$$\begin{aligned} & (\mathbf{1}, Y(a_1, \underline{x}_1) Y(a_2, \underline{x}_2) \dots Y(a_{n-1}, \underline{x}_{n-1}) a_n) \\ &= x_1^{-h} \bar{x}_1^{-\bar{h}} \sum_{\substack{n, m \in \mathbb{Z} \\ s_3, \bar{s}_3, \dots, s_n, \bar{s}_n \in \mathbb{R}}} c_{s_1, \bar{s}_1, \dots, s_{n-1}, \bar{s}_{n-1}} \frac{x_2^{-h+2h_1+n}}{x_1} \frac{\bar{x}_2^{-\bar{h}+2\bar{h}_1+m}}{\bar{x}_1} \dots \frac{x_{n-2}^{s_{n-1}}}{x_{n-2}} \frac{\bar{x}_{n-1}^{\bar{s}_{n-1}}}{\bar{x}_{n-2}}, \end{aligned}$$

where $s_1 = -h - s_2 - \dots - s_{n-1}$, $\bar{s}_1 = -\bar{h} - \bar{s}_2 - \dots - \bar{s}_{n-1}$, $s_2 = -h + 2h_1 - s_3 - \dots - s_{n-1} + n$ and $\bar{s}_2 = -\bar{h} + 2\bar{h}_1 - \bar{s}_3 - \dots - \bar{s}_{n-1} + m$. Thus, the assertion follows. $\square$

We have the embedding $P_n \rightarrow Q_n$ defined by $A \mapsto (A)\star$ for $A \in P_n$ and surjection $d_\star : Q_n \rightarrow P_n$ defined by deleting $\star$ from $A \in Q_n$. By lemma 3.4, we have:

Lemma 3.8. For $A \in P_n$ and $a_1, \dots, a_n$,

$$S_A(a_1, \dots, a_n) = S_{(A)\star}(a_1, \dots, a_n).$$

By Proposition 2.1, $Y(a, \underline{z})\mathbf{1} = \exp(D\underline{z} + \bar{D}\bar{\underline{z}})a$ and $[D, Y(a, \underline{z})] = d/d\underline{z} Y(a, \underline{z})$, any parenthesized n point correlation function for $A \in Q_n$ is an expansion of a parenthesized correlation function for $d_\star(A) \in P_n$.

For example, let us consider the case of $A = 1(2(3(4\star))) \in Q_4$. In this case,

$$S_{1(2(3(4\star)))}(a_1, a_2, a_3, a_4; z_1, z_2, z_3, z_4) = (\mathbf{1}, a_1(z_1) a_2(z_2) a_3(z_3) a_4(z_4) \mathbf{1}).$$

Since

$$\begin{aligned} & (\mathbf{1}, a_1(z_1) a_2(z_2) a_3(z_3) a_4(z_4) \mathbf{1}) \\ &= (\mathbf{1}, a_1(z_1) a_2(z_2) a_3(z_3) \exp(Dz_4 + \bar{D}\bar{z}_4) a_4) \\ &= \exp(-z_4(d/dz_1 + d/dz_2 + d/dz_3)) \exp(-\bar{z}_4(d/d\bar{z}_1 + d/d\bar{z}_2 + d/d\bar{z}_3)) (\mathbf{1}, a_1(z_1) a_2(z_2) a_3(z_3) a_4) \\ &= \exp(-z_4(d/dz_1 + d/dz_2 + d/dz_3)) \exp(-\bar{z}_4(d/d\bar{z}_1 + d/d\bar{z}_2 + d/d\bar{z}_3)) S_{1(2(3(4)))}(a_1, a_2, a_3, a_4; z_1, z_2, z_3), \end{aligned}$$

by Lemma 1.18, we have:

Lemma 3.9. For a_1, a_2, a_3, a_4 ,

$$S_{1(2(3(4\star)))}(a_1, a_2, a_3, a_4; z_1, z_2, z_3, z_4) = T_{1(2(3(4)))}^{1(2(3(4\star)))} S_{1(2(3(4)))}(a_1, a_2, a_3, a_4; z_{14}, z_{24}, z_{34}).$$

Hereafter, we concentrate on the case of $n = 4$ and $A \in P_4$. Similarly to the proof of Lemma 3.7, we can show that $S_A(a_1, a_2, a_3, a_4) \in T_A$, where T_A is the space of formal power series defined in Section 1.3.

For example, for (12)(34), we have:

Lemma 3.10. *For any $a_1, a_2, a_3, a_4 \in F$, $(\mathbf{1}, Y(Y(a_1, \underline{x}_1)a_2, \underline{x}_2)Y(a_3, \underline{x}_3)a_4) \in T_{(12)(34)}(x_2, x_1, x_3)$.*

Proof. We may assume that $a_i \in F_{h_i, \bar{h}_i}$ for $h_i, \bar{h}_i \in \mathbb{R}$ and $i = 1, \dots, 4$. Set $h = h_1 + \dots + h_4$ and $\bar{h} = \bar{h}_1 + \dots + \bar{h}_4$ and

$$\sum_{s_1, \bar{s}_1, \dots, s_3, \bar{s}_3 \in \mathbb{R}} c_{s_1, \bar{s}_1, s_2, \bar{s}_2, s_3, \bar{s}_3} x_1^{s_1} \bar{x}_1^{\bar{s}_1} x_2^{s_2} \bar{x}_2^{\bar{s}_2} x_3^{s_3} \bar{x}_3^{\bar{s}_3} = (\mathbf{1}, Y(Y(a_1, \underline{x}_1)a_2, \underline{x}_2)Y(a_3, \underline{x}_3)a_4).$$

Then,

$$c_{s_1, \bar{s}_1, \dots, s_3, \bar{s}_3} = (\mathbf{1}, (a_1(-s_1 - 1, -\bar{s}_1 - 1)a_2)(-s_2 - 1, -\bar{s}_2 - 1)a_3(-s_3 - 1, -\bar{s}_3 - 1)a_4).$$

By (FVO5), there exists $N \in \mathbb{R}$ such that $F_{h, \bar{h}} = 0$ for any $h \leq N$ or $\bar{h} \leq N$. Thus, similarly to the proof of Lemma 3.7, $c_{s_1, \bar{s}_1, \dots, s_3, \bar{s}_3} = 0$ unless $0 = h + (s_1 + s_2 + s_3)$, $h_1 + h_2 + s_1 - h_3 - h_4 - s_3 \in \mathbb{Z}$ and $h_3 + h_4 + s_3, h_1 + h_2 + s_1 \geq N$.

Thus, there exists $N' \in \mathbb{R}$ such that

$$\begin{aligned} & (\mathbf{1}, Y(Y(a_1, \underline{x}_1)a_2, \underline{x}_2)Y(a_3, \underline{x}_3)a_4) \\ &= x_2^{-h} \bar{x}_2^{-\bar{h}} \left(\frac{x_1}{x_2}\right)^{-h_1-h_2+h_3+h_4} \left(\frac{\bar{x}_1}{\bar{x}_2}\right)^{-\bar{h}_1-\bar{h}_2+\bar{h}_3+\bar{h}_4} \sum_{s'_1, \bar{s}'_1, s_3, \bar{s}_3 \in \mathbb{R}} c_{s_1, \bar{s}_1, \dots, s_3, \bar{s}_3} \left(\frac{x_1}{x_2}\right)^{s'_1} \left(\frac{\bar{x}_1}{\bar{x}_2}\right)^{\bar{s}'_1} \left(\frac{x_3}{x_2}\right)^{s_3} \left(\frac{\bar{x}_3}{\bar{x}_2}\right)^{\bar{s}_3} \end{aligned}$$

where $s_2 = -h - s_1 - s_3$, $\bar{s}_2 = -\bar{h} - \bar{s}_1 - \bar{s}_3$, $s'_1 = h_1 + h_2 - h_3 - h_4 + s_1$ and $\bar{s}'_1 = \bar{h}_1 + \bar{h}_2 - \bar{h}_3 - \bar{h}_4 + \bar{s}_1$ and the sum runs through $s'_1, s_3, \bar{s}'_1, \bar{s}_3 \geq N'$ and $s'_1 - s_3 \in \mathbb{Z}$, $\bar{s}'_1 - \bar{s}_3 \in \mathbb{Z}$. Thus, the assertion holds. $\square$

3.4. Consistency of four point functions. Let F be a self-dual full vertex operator algebra. In this section, we will prove the main result of this paper, the consistency of four point correlation functions. The following lemma follows from Lemma 3.1:

Lemma 3.11. *For $h_i, \bar{h}_i \in \mathbb{R}$ and $a_i \in F_{h_i, \bar{h}_i}$ and $b_i \in QF_{h_i, \bar{h}_i}$,*

(1)

$$\begin{aligned} & \left(\sum_{i=1}^n z_i d/dz_i + h_i\right)(\mathbf{1}, Y(a_1, \underline{z}_1) \dots Y(a_{n-1}, \underline{z}_{n-1})a_n) = 0, \\ & \left(\sum_{i=1}^n \bar{z}_i d/d\bar{z}_i + \bar{h}_i\right)(\mathbf{1}, Y(a_1, \underline{z}_1) \dots Y(a_{n-1}, \underline{z}_{n-1})a_n) = 0; \end{aligned}$$

(2)

$$\begin{aligned} & \left(\sum_{i=1}^n z_i^2 d/dz_i + 2h_i z_i\right)(\mathbf{1}, Y(b_1, \underline{z}_1) \dots Y(b_{n-1}, \underline{z}_{n-1})b_n) = 0, \\ & \left(\sum_{i=1}^n \bar{z}_i^2 d/d\bar{z}_i + 2\bar{h}_i \bar{z}_i\right)(\mathbf{1}, Y(b_1, \underline{z}_1) \dots Y(b_{n-1}, \underline{z}_{n-1})b_n) = 0. \end{aligned}$$

For $h_i, \bar{h}_i \in \mathbb{R}$ with $h_i - \bar{h}_i \in \mathbb{Z}$ ($i = 1, 2, 3, 4$), set

$$Q(\underline{h}, z) = (z_1 - z_2)^{-2h_2} (z_3 - z_4)^{h_1 - h_2 - h_3 - h_4} (z_1 - z_4)^{h_2 - h_1 + h_3 - h_4} (z_1 - z_3)^{h_2 - h_1 - h_3 + h_4} \\ (\bar{z}_1 - \bar{z}_2)^{-2\bar{h}_2} (\bar{z}_3 - \bar{z}_4)^{\bar{h}_1 - \bar{h}_2 - \bar{h}_3 - \bar{h}_4} (\bar{z}_1 - \bar{z}_4)^{\bar{h}_2 - \bar{h}_1 + \bar{h}_3 - \bar{h}_4} (\bar{z}_1 - \bar{z}_3)^{\bar{h}_2 - \bar{h}_1 - \bar{h}_3 + \bar{h}_4} \in \text{Cor}_4^f.$$

Then, by an easy computation, we have:

Lemma 3.12. *For $h_i, \bar{h}_i \in \mathbb{R}$ with $h_i - \bar{h}_i \in \mathbb{Z}$ ($i = 1, 2, 3, 4$), as a function on X_4 , $Q(\underline{h}, z)$ satisfies*

$$\begin{aligned} \left(\sum_{i=1}^3 z_i d/dz_i + h_i \right) Q(\underline{h}, z) &= 0. \\ \left(\sum_{i=1}^3 \bar{z}_i d/d\bar{z}_i + \bar{h}_i \right) Q(\underline{h}, z) &= 0. \\ \left(\sum_{i=1}^4 z_i^2 d/dz_i + 2z_i h_i \right) Q(\underline{h}, z) &= 0. \\ \left(\sum_{i=1}^4 \bar{z}_i^2 d/d\bar{z}_i + 2\bar{z}_i \bar{h}_i \right) Q(\underline{h}, z) &= 0. \end{aligned}$$

Remark 3.1. *We can choose another $Q(\underline{h}, z) \in \text{Cor}_4$ which satisfies the above differential equations. In fact, we can multiply $Q(\underline{h}, z)$ by $\mathbb{C}[\xi^\pm, \bar{\xi}^\pm, |\xi|^\mathbb{R}]$. We also remark that*

$$\prod_{1 \leq i < j \leq 4} (z_i - z_j)^{-h_i - h_j + h/3} (\bar{z}_i - \bar{z}_j)^{-\bar{h}_i - \bar{h}_j + \bar{h}/3},$$

is commonly used in physics, where $h = h_1 + h_2 + h_3 + h_4$ and $\bar{h} = \bar{h}_1 + \bar{h}_2 + \bar{h}_3 + \bar{h}_4$. However, since $(h - \bar{h})/3$ is not always integer, this function is not in Cor_4^f .

Proposition 3.6. *For $h_i, \bar{h}_i \in \mathbb{R}$ and $a_i \in QF_{h_i, \bar{h}_i}$ ($i = 1, 2, 3, 4$), the following conditions hold:*

(1)

$$(\mathbf{1}, Y(a_1, \underline{z}_{12})a_2) = (a_1, a_2)(-1)^{h_1 - \bar{h}_1} z_{12}^{-2h_1} \bar{z}_{12}^{-2\bar{h}_1}.$$

In particular, it is zero unless $h_1 = h_2$ and $\bar{h}_1 = \bar{h}_2$.

(2) *There exists $C_{a_1, a_2, a_3} \in \mathbb{C}$ such that*

$$(\mathbf{1}, Y(a_1, \underline{z}_{13})Y(a_2, \underline{z}_{23})a_3) = C_{a_1, a_2, a_3} \prod_{1 \leq i < j \leq 3} (z_i - z_j)^{h_1 + h_2 + h_3 - 2h_i - 2h_j} (\bar{z}_i - \bar{z}_j)^{\bar{h}_1 + \bar{h}_2 + \bar{h}_3 - 2\bar{h}_i - 2\bar{h}_j} \Big|_{|z_{13}| > |z_{23}|}.$$

(3) *There exists $f \in F_{0,1,\infty}$ such that*

$$(\mathbf{1}, Y(a_1, \underline{z}_{14})Y(a_2, \underline{z}_{24})Y(a_3, \underline{z}_{34})a_4) = \left(Q(\underline{h}, z) f \circ \xi \right) \Big|_{|z_{14}| > |z_{24}| > |z_{34}|}.$$

In particular, this series is absolutely convergent to the function in Cor_4 .

Proof. Since $(\mathbf{1}, Y(a_1, \underline{z}_{12})a_2) = (Y(S_{z_{12}}a_1, \underline{z}_{12}^{-1})\mathbf{1}, a_2) = (-1)^{h_1 - \bar{h}_1} z_{12}^{-2h_1} \bar{z}_{12}^{-2\bar{h}_1} (\exp(L(-1)z_{12}^{-1} + \bar{L}(-1)\bar{z}_{12}^{-1})a_1, a_2) = (-1)^{h_1 - \bar{h}_1} z_{12}^{-2h_1} \bar{z}_{12}^{-2\bar{h}_1} (a_1, \exp(L(1)z_{12}^{-1} + \bar{L}(1)\bar{z}_{12}^{-1})a_2) = (-1)^{h_1 - \bar{h}_1} z_{12}^{-2h_1} \bar{z}_{12}^{-2\bar{h}_1} (a_1, a_2)$, (1) holds.

By Lemma 3.7, the product of the formal power series $\prod_{1 \leq i < j \leq 3} (z_i - z_j)^{-h_1 - h_2 - h_3 + 2h_i + 2h_j} (\bar{z}_i - \bar{z}_j)^{-\bar{h}_1 - \bar{h}_2 - \bar{h}_3 + 2\bar{h}_i + 2\bar{h}_j} |_{|z_{13}| > |z_{23}|}$ and $(\mathbf{1}, Y(a_1, \underline{z}_{13})Y(a_2, \underline{z}_{23})a_3)$ is in

$$\mathbb{C}[[z_{23}/z_{13}, \bar{z}_{23}/\bar{z}_{13}]] [z_{13}^{\pm}, z_{23}^{\pm}, \bar{z}_{13}^{\pm}, \bar{z}_{23}^{\pm}, |z_{13}|^{\mathbb{R}}, |z_{23}|^{\mathbb{R}}]$$

and is in the kernel of the formal differentials $D_0'', \bar{D}_0'', D_1'', \bar{D}_1''$ by Lemma 3.11 and the similar result to Lemma 3.12. Hence, it is constant $C_{a_1, a_2, a_3} \in \mathbb{C}$ by Lemma 1.16, which implies that (2) holds.

Similarly, set

$$\phi = Q(\underline{h}, z)^{-1} |_{|z_{14}| > |z_{24}| > |z_{34}|} (\mathbf{1}, Y(a_1, \underline{z}_{14})Y(a_2, \underline{z}_{24})Y(a_3, \underline{z}_{34})a_4),$$

which is an element of $T(z_{14}, z_{24}, z_{34})$ and in the kernel of the formal differential $D_0, \bar{D}_0, D_1, \bar{D}_1$. Thus, by Lemma 1.13, $v_{1(2(34))}\phi \in \mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}}))$ determines ϕ . Recall that $v_{1(2(34))} : S(z_{14}, z_{24}, z_{34}) \rightarrow \mathbb{C}((p, \bar{p}, |p|^{\mathbb{R}}))$ is defined by the evaluation $(z_{14}, z_{24}, z_{34}) \mapsto (\infty, 1, p)$. Since

$$\begin{aligned} & (\mathbf{1}, Y(a_1, \underline{z}_{14})Y(a_2, \underline{z}_{24})Y(a_3, \underline{z}_{34})a_4) \\ &= (-1)^{h_1 - \bar{h}_1} z_{14}^{-2h_1} \bar{z}_{14}^{-2\bar{h}_1} (\exp(L(-1)z_{14} + \bar{L}(-1)\bar{z}_{14})a_1, Y(a_2, \underline{z}_{24})Y(a_3, \underline{z}_{34})a_4) \end{aligned}$$

and

$$\begin{aligned} Q(\underline{h}, z) &= z_{14}^{-2h_1} z_{34}^{h_1 - h_2 - h_3 - h_4} (1 - z_{24}/z_{14})^{-2h_2} (1 - z_{34}/z_{14})^{h_2 - h_1 - h_3 + h_4} \\ &\quad \bar{z}_{14}^{-2\bar{h}_1} \bar{z}_{34}^{\bar{h}_1 - \bar{h}_2 - \bar{h}_3 - \bar{h}_4} (1 - \bar{z}_{24}/\bar{z}_{14})^{-2\bar{h}_2} (1 - \bar{z}_{34}/\bar{z}_{14})^{\bar{h}_2 - \bar{h}_1 - \bar{h}_3 + \bar{h}_4}. \end{aligned}$$

the formal limit of ϕ as $\underline{z}_{14} \rightarrow \infty$ is

$$(-1)^{h_1 - \bar{h}_1} z_{34}^{-h_1 + h_2 + h_3 + h_4} \bar{z}_{34}^{-\bar{h}_1 + \bar{h}_2 + \bar{h}_3 + \bar{h}_4} (a_1, Y(a_2, \underline{z}_{24})Y(a_3, \underline{z}_{34})a_4)$$

and

$$v_{1(2(34))}(\phi) = (-1)^{h_1 - \bar{h}_1} p^{-h_1 + h_2 + h_3 + h_4} \bar{p}^{-\bar{h}_1 + \bar{h}_2 + \bar{h}_3 + \bar{h}_4} (a_1, Y(a_2, 1)Y(a_3, \underline{p})a_4).$$

By Lemma 2.2, there exists $f \in F_{0,1,\infty}$ such that $v_{1(2(34))}(\phi) = j(p, f)$. By Corollary 1.1, we have $\phi = e_{1(2(34))}(f)$. $\square$

For $h_i, \bar{h}_i \in \mathbb{R}$ and $a_i \in QF_{h_i, \bar{h}_i}$ ($i = 1, 2, 3, 4$), by the above proof,

$$\begin{aligned} & \lim_{\underline{z}_{14} \rightarrow \infty} Q(\underline{h}, z)^{-1} (\mathbf{1}, Y(a_1, \underline{z}_{14})Y(a_2, \underline{z}_{24})Y(a_3, \underline{z}_{34})a_4) \\ &= (-1)^{h_1 - \bar{h}_1} z_{34}^{-h_1 + h_2 + h_3 + h_4} \bar{z}_{34}^{-\bar{h}_1 + \bar{h}_2 + \bar{h}_3 + \bar{h}_4} (a_1, Y(a_2, \underline{z}_{24})Y(a_3, \underline{z}_{34})a_4) \\ & \lim_{\underline{z}_{14} \rightarrow \infty} Q(\underline{h}, z)^{-1} (\mathbf{1}, Y(a_1, \underline{z}_{14})Y(a_3, \underline{z}_{34})Y(a_2, \underline{z}_{24})a_4) \\ &= (-1)^{h_1 - \bar{h}_1} z_{34}^{-h_1 + h_2 + h_3 + h_4} \bar{z}_{34}^{-\bar{h}_1 + \bar{h}_2 + \bar{h}_3 + \bar{h}_4} (a_1, Y(a_3, \underline{z}_{34})Y(a_2, \underline{z}_{24})a_4). \end{aligned}$$

Thus, by Lemma 2.2, we have:

Lemma 3.13. For $h_i, \bar{h}_i \in \mathbb{R}$ and $a_i \in QF_{h_i, \bar{h}_i}$ ($i = 1, 2, 3, 4$), there exists $f \in F_{0,1,\infty}$ such that

$$\begin{aligned} (\mathbf{1}, Y(a_1, \underline{z}_{14})Y(a_2, \underline{z}_{24})Y(a_3, \underline{z}_{34})a_4) &= e_{1(2(34))}(Q(h, z)f \circ \xi) \\ (\mathbf{1}, Y(a_1, \underline{z}_{14})Y(a_3, \underline{z}_{34})Y(a_2, \underline{z}_{24})a_4) &= e_{1(3(24))}(Q(h, z)f \circ \xi). \end{aligned}$$

Proposition 3.7. Let F be a self-dual QP -generated full vertex operator algebra. Then, for any $a_i \in F_i$, there exists $\phi \in \text{Cor}_4$ such that

$$(\mathbf{1}, Y(a_{\sigma 1}, \underline{z}_{\sigma 1}) \dots Y(a_{\sigma 4}, \underline{z}_{\sigma 4})\mathbf{1}) = e_{\sigma 1(2(3(4\star)))}(\phi),$$

for any $\sigma \in S_4$. In particular, this series is absolutely convergent to a function in Cor_4 .

Proof. First, we assume that $a_i \in QF_{h_i, \bar{h}_i}$ for $i = 1, 2, 3, 4$. Set $Q = Q(\underline{h}, z)$ and $h = h_1 + h_2 + h_3 + h_4$ and $\bar{h} = \bar{h}_1 + \bar{h}_2 + \bar{h}_3 + \bar{h}_4$. Then, by Proposition 3.6, there exists $f \in F_{0,1,\infty}$ such that $(\mathbf{1}, Y(a_1, \underline{z}_{14})Y(a_2, \underline{z}_{24})Y(a_3, \underline{z}_{34})a_4) = e_{1(2(34))}(Qf)$. Then, by Lemma 1.18 and Lemma 3.9, we have

$$\begin{aligned} &(\mathbf{1}, Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)Y(a_3, \underline{z}_3)Y(a_4, \underline{z}_4)\mathbf{1}) \\ &= T_{1(2(34))}^{1(2(3(4\star)))}(\mathbf{1}, Y(a_1, \underline{z}_{14})Y(a_2, \underline{z}_{24})Y(a_3, \underline{z}_{34})a_4) \\ &= T_{1(2(34))}^{1(2(3(4\star)))}e_{1(2(34))}(Qf) \\ &= e_{1(2(3(4\star)))}(Qf). \end{aligned}$$

Let $f_1, f_2 \in F_{0,1,\infty}$ satisfy

$$\begin{aligned} e_{1(2(43))}(Qf_1) &= (\mathbf{1}, Y(a_1, \underline{z}_{13})Y(a_2, \underline{z}_{23})Y(a_4, \underline{z}_{43})a_3) \\ e_{4(3(2(10)))}(Qf_2) &= (\mathbf{1}, Y(a_4, \underline{z}_4)Y(a_3, \underline{z}_3)Y(a_2, \underline{z}_2)Y(a_1, \underline{z}_1)\mathbf{1}). \end{aligned}$$

We will show that f and f_1, f_2 are the same function. Then, since $(34), (14)(23), (23) \in S_4$ is a generator of S_4 , by Lemma 3.13, the assertion holds for quasi-primary vectors. By Proposition 2.1 and Lemma 1.11,

$$\begin{aligned} e_{1(2(43))}(Qf_1) &= (\mathbf{1}, Y(a_1, \underline{z}_{13})Y(a_2, \underline{z}_{23})Y(a_4, \underline{z}_{43})a_3) \\ &= (\mathbf{1}, Y(a_1, \underline{z}_{13})Y(a_2, \underline{z}_{23})\exp(\underline{z}_{43}D + \bar{\underline{z}}_{43}\bar{D})Y(a_3, -\underline{z}_{43})a_4) \\ &= \exp(-\underline{z}_{43}(d/d\underline{z}_{13} + d/d\underline{z}_{23}) - \bar{\underline{z}}_{43}(d/d\bar{\underline{z}}_{13} + d/d\bar{\underline{z}}_{23}))(\mathbf{1}, Y(a_1, \underline{z}_{13})Y(a_2, \underline{z}_{23})Y(a_3, -\underline{z}_{43})a_4) \\ &= T_{1(2(34))}^{1(2(43))}e_{1(2(34))}(Qf), \end{aligned}$$

which implies that $f_1 = f$.

Since

$$\begin{aligned} e_{1(2(3(4\star)))}(Qf) &= (\mathbf{1}, Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)Y(a_3, \underline{z}_3)Y(a_4, \underline{z}_4)\mathbf{1}) \\ &= (-1)^{h-\bar{h}}\prod_{1 \leq i \leq 4} \underline{z}_i^{-2h_i} \bar{\underline{z}}_i^{-2\bar{h}_i} (\mathbf{1}, Y(a_4, \underline{z}_4^{-1})Y(a_3, \underline{z}_3^{-1})Y(a_2, \underline{z}_2^{-1})Y(a_1, \underline{z}_1^{-1})\mathbf{1}), \end{aligned}$$

by Lemma 1.19,

$$\begin{aligned}
e_{4(3(2(1\star)))}(Qf_2) &= (\mathbf{1}, Y(a_4, \underline{z}_4)Y(a_3, \underline{z}_3)Y(a_2, \underline{z}_2)Y(a_1, \underline{z}_1)\mathbf{1}) \\
&= I_d\left((-1)^{h-\bar{h}}\prod_{1\leq i\leq 4}z_i^{2h_i}\bar{z}_i^{2\bar{h}_i}(\mathbf{1}, Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)Y(a_3, \underline{z}_3)Y(a_4, \underline{z}_4)\mathbf{1})\right) \\
&= I_d\left((-1)^{h-\bar{h}}\prod_{1\leq i\leq 4}z_i^{2h_i}\bar{z}_i^{2\bar{h}_i}e_{1(2(3(40)))}(Qf)\right) \\
&= e_{4(3(2(1\star)))}(Qf),
\end{aligned}$$

which implies $f_2 = f$.

We now turn to the case that $a_i \in F_{h_i, \bar{h}_i}$. Since F is QP-generated, we may assume that $a_i = D^{n_i} \bar{D}^{\bar{n}_i} b_i$ for some $n_i, \bar{n}_i \in \mathbb{Z}_{\geq 0}$ and $b_i \in QF_{h_i-n_i, \bar{h}_i-\bar{n}_i}$. Since

$$\begin{aligned}
&(\mathbf{1}, Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)Y(a_3, \underline{z}_3)Y(a_4, \underline{z}_4)\mathbf{1}) \\
&= (\mathbf{1}, Y(D^{n_1} \bar{D}^{\bar{n}_1} b_1, \underline{z}_1)Y(D^{n_2} \bar{D}^{\bar{n}_2} b_2, \underline{z}_2)Y(D^{n_3} \bar{D}^{\bar{n}_3} b_3, \underline{z}_3)Y(D^{n_4} \bar{D}^{\bar{n}_4} b_4, \underline{z}_4)\mathbf{1}) \\
&= \prod_{1\leq i\leq 4} d/dz_i^{n_i} d/d\bar{z}_i^{\bar{n}_i} (\mathbf{1}, Y(b_1, \underline{z}_1)Y(b_2, \underline{z}_2)Y(b_3, \underline{z}_3)Y(b_4, \underline{z}_4)\mathbf{1}),
\end{aligned}$$

by Proposition 3.6, there exists $\phi \in \text{Cor}_4$ such that $(\mathbf{1}, Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)Y(a_3, \underline{z}_3)Y(a_4, \underline{z}_4)\mathbf{1}) = e_{1(2(3(4\star)))}(\phi)$. The S_4 -invariance follows from the S_4 -invariance for quasi-primary vectors. $\square$

Hence, we can define the linear map $S : F^{\otimes 4} \rightarrow \text{Cor}_4$ by $e_{1(2(3(4\star)))}(S(a_1, \dots, a_4)) = (\mathbf{1}, Y(a_1, \underline{z}_1) \dots Y(a_4, \underline{z}_4)\mathbf{1})$. We recall that the symmetric group S_4 acts on Cor_4 (see Section 1.2). Then, the S_4 -symmetry of the correlation function follows:

Theorem 3.1. *For a self-dual QP-generated full vertex operator algebra F and $a_1, a_2, a_3, a_4 \in F$ and $\sigma \in S_4$,*

$$\sigma \cdot S(a_1, a_2, a_3, a_4) = S(a_{\sigma^{-1}1}, a_{\sigma^{-1}2}, a_{\sigma^{-1}3}, a_{\sigma^{-1}4}).$$

Proof. By Proposition 1.2 and Proposition 3.7,

$$\begin{aligned}
e_{1(2(3(4\star)))}(\sigma \cdot S(a_1, a_2, a_3, a_4)) &= T_\sigma e_{\sigma^{-1}1(2(3(4\star)))}S(a_1, a_2, a_3, a_4) \\
&= T_\sigma(\mathbf{1}, Y(a_{\sigma^{-1}1}, \underline{z}_{\sigma^{-1}1})Y(a_{\sigma^{-1}2}, \underline{z}_{\sigma^{-1}2})Y(a_{\sigma^{-1}3}, \underline{z}_{\sigma^{-1}3})Y(a_{\sigma^{-1}4}, \underline{z}_{\sigma^{-1}4})\mathbf{1}) \\
&= (\mathbf{1}, Y(a_{\sigma^{-1}1}, \underline{z}_1)Y(a_{\sigma^{-1}2}, \underline{z}_2)Y(a_{\sigma^{-1}3}, \underline{z}_3)Y(a_{\sigma^{-1}4}, \underline{z}_4)\mathbf{1}) \\
&= e_{1(2(3(4\star)))}S(a_{\sigma^{-1}1}, a_{\sigma^{-1}2}, a_{\sigma^{-1}3}, a_{\sigma^{-1}4}).
\end{aligned}$$

$\square$

Now, the result for the consistency of four point correlation functions in conformal field theory on $\mathbb{CP}^1$ can be stated as follows:

Theorem 3.2. *Let F be a self-dual QP-generated full vertex operator algebra. Then, $S_A = e_A \circ S$ for any $A \in Q_4$.*

Lemma 3.14. *For $a_i \in QF_{h_i, \bar{h}_i}$ ($i = 1, 2, 3, 4$),*

$$\begin{aligned}
S_{(12)(34)}(a_1, a_2, a_3, a_4) &= (-1)^{h_2-\bar{h}_2} z_{12}^{-2h_1} \bar{z}_{12}^{-2\bar{h}_1} z_{24}^{-2h_2+2h_1} \bar{z}_{24}^{-2\bar{h}_2+2\bar{h}_1} (1 - z_{34}/z_{24})^{-2h_3} (1 - \bar{z}_{34}/\bar{z}_{24})^{-2\bar{h}_3} \\
&\quad \left(a_2, Y\left(a_1, -\underline{z}_{24}(1 + \underline{z}_{24}/\underline{z}_{12})\right)Y\left(a_3, \frac{\underline{z}_{34}}{1 - \underline{z}_{34}/\underline{z}_{24}}\right)a_4\right)
\end{aligned}$$

Proof. By the invariance,

$$\begin{aligned} (\mathbf{1}, Y(Y(a_1, \underline{z}_{12})a_2, \underline{z}_{24})Y(a_3, \underline{z}_{34})a_4) &= (Y(S_{\underline{z}_{24}}Y(a_1, \underline{z}_{12})a_2, \underline{z}_{24}^{-1})\mathbf{1}, Y(a_3, \underline{z}_{34})a_4) \\ &= (\exp(\underline{z}_{24}^{-1}D + \bar{\underline{z}}_{24}^{-1}\bar{D})S_{\underline{z}_{24}}Y(a_1, \underline{z}_{12})a_2, Y(a_3, \underline{z}_{34})a_4) \\ &= (a_2, Y(S_{\underline{z}_{12}}a_1, \underline{z}_{12}^{-1})S_{\underline{z}_{24}}^\dagger(\exp(\underline{z}_{24}^{-1}\bar{L}(1) + \bar{\underline{z}}_{24}^{-1}\bar{L}(1))Y(a_3, \underline{z}_{34})a_4), \end{aligned}$$

where $S_{\underline{z}_{24}}^\dagger = (-\underline{z}_{24})^{L(0)}(-\bar{\underline{z}}_{24})^{\bar{L}(0)}\exp(L(-1)\underline{z}_{24} + \bar{L}(-1)\bar{\underline{z}}_{24})$. By [FHL],

$$\underline{z}^{L(0)}Y(a, \underline{z}_0)\underline{z}^{-L(0)} = Y(\underline{z}^{L(0)}a, \underline{z}\underline{z}_0)$$

$$\exp(\underline{z}L(1))Y(a, \underline{z}_0)\exp(-\underline{z}L(1)) = Y(\exp(\underline{z}(1 - \underline{z}\underline{z}_0)L(1))(1 - \underline{z}\underline{z}_0)^{-2L(0)}a, \underline{z}_0/(1 - \underline{z}\underline{z}_0)).$$

Thus, the assertion holds. $\square$

proof of Theorem 3.2. We only prove the Theorem for $A \in P_4$. The general result can be obtained by using Proposition 2.1 and the similar argument as Lemma 3.9. For the case of $A = 1((23)4), ((12)3)4$ and $(1(23))4$, the assertion follows from Lemma 1.9 and Lemma 1.10. We will prove the theorem for $A = (12)(34)$. Similarly to the proof of Theorem 3.7, we may assume that $a_i \in QF_{h_i, \bar{h}_i}$ for $i = 1, 2, 3, 4$. Set $Q = Q(\underline{h}, \underline{z})$ and let $f \in F_{0,1,\infty}$ satisfy $S(a_1, a_2, a_3, a_4) = Qf(\xi)$. Then, by Lemma 3.1 and Lemma 3.10,

$$e_{(12)(34)}(Q^{-1})(\mathbf{1}, Y(Y(a_1, \underline{z}_{12})a_2, \underline{z}_{24})Y(a_3, \underline{z}_{34})a_4)$$

is in $S'(\underline{z}_{24}, \underline{z}_{12}, \underline{z}_{34})$. Thus, by Lemma 1.15, $e_{(12)(34)}(Q^{-1})(\mathbf{1}, Y(Y(a_1, \underline{z}_{12})a_2, \underline{z}_{24})Y(a_3, \underline{z}_{34})a_4)$ is uniquely determined by the limit of $(\underline{z}_{24}, \underline{z}_{12}, \underline{z}_{34}) \rightarrow (1, q, q)$. By Lemma 3.14, the limit is equal to

$$\begin{aligned} &(-1)^{h_2 - \bar{h}_2} q^{-3h_1 + 3h_2 + h_3 + h_4} (1 + q)^{h_1 - h_2 - h_3 + h_4} (1 - q)^{-2h_3} \bar{q}^{-3\bar{h}_1 + 3\bar{h}_2 + \bar{h}_3 + \bar{h}_4} (1 + \bar{q})^{\bar{h}_1 - \bar{h}_2 - \bar{h}_3 + \bar{h}_4} (1 - \bar{q})^{-2\bar{h}_3} \\ &(\text{q-series}) \quad \cdot (a_2, Y(a_1, -(1 + \underline{q}^{-1}))Y(a_3, \underline{q}/(1 - \underline{q}))a_4). \end{aligned}$$

Since the formal limit (we omit the anti-holomorphic part)

$$\begin{aligned} &\lim_{(\underline{z}_2, \underline{z}_1, \underline{z}_3, \underline{z}_4) \rightarrow (\infty, -(1+q^{-1}), q/(1-q), 0)} \underline{z}_2^{-2h_2} \bar{\underline{z}}_2^{-2\bar{h}_2} Q(\underline{z})^{-1} \\ &= \frac{q}{1-q}^{-h_1 + h_2 + h_3 + h_4} (-q^{-1}(1+q))^{h_1 - h_2 - h_3 + h_4} (-q(1-q))^{-h_1 + h_2 - h_3 + h_4} \\ &= (-1)^{2h_1 - 2h_2} q^{-3h_1 + 3h_2 + h_3 + h_4} (1+q)^{h_1 - h_2 - h_3 + h_4} (1-q)^{-2h_3} \end{aligned}$$

and $2h_1 - 2h_2 - 2\bar{h}_1 - 2\bar{h}_2 \in 2\mathbb{Z}$, we can verify that the series q-series is equal to the formal limit of

$$Q(\underline{z})^{-1}|_{|\underline{z}_2| > |\underline{z}_1| > |\underline{z}_3| > |\underline{z}_4|}(\mathbf{1}, Y(a_2, \underline{z}_2)Y(a_1, \underline{z}_1)Y(a_3, \underline{z}_3)Y(a_4, \underline{z}_4)\mathbf{1})$$

as $(\underline{z}_2, \underline{z}_1, \underline{z}_3, \underline{z}_4) \rightarrow (\infty, -(1 + q^{-1}), q/(1 - q), 0)$. Since the limit of $\xi(\underline{z}_1, \underline{z}_2, \underline{z}_3, \underline{z}_4)$ as $(\underline{z}_2, \underline{z}_1, \underline{z}_3, \underline{z}_4) \rightarrow (\infty, -(1 + q^{-1}), q/(1 - q), 0)$ is q^2 and

$$Q(\underline{z})^{-1}|_{|\underline{z}_2| > |\underline{z}_1| > |\underline{z}_3| > |\underline{z}_4|}(\mathbf{1}, Y(a_2, \underline{z}_2)Y(a_1, \underline{z}_1)Y(a_3, \underline{z}_3)Y(a_4, \underline{z}_4)\mathbf{1}) = e_{2(1(3(4\star)))}(f(\xi)),$$

$$\nu_{(12)(34)}\left(e_{(12)(34)}(Q^{-1})(\mathbf{1}, Y(Y(a_1, \underline{z}_{12})a_2, \underline{z}_{24})Y(a_3, \underline{z}_{34})a_4)\right) = \lim_{p \rightarrow q^2} j(p, f).$$

Hence, $(\mathbf{1}, Y(Y(a_1, \underline{z}_{12})a_2, \underline{z}_{24})Y(a_3, \underline{z}_{34})a_4) = e_{(12)(34)}(Qf(\xi))$. $\square$

4. CONSTRUCTION

In this section, we generalize the Goddard's axiom for a full vertex algebra (Section 4.1 and Section 4.2). We also show that if a vertex operator $Y(-, \underline{z})$ satisfies the bootstrap equation then it gives a full vertex algebra.

A full prevertex algebra is an $\mathbb{R}^2$ -graded $\mathbb{C}$ -vector space $F = \bigoplus_{h, \bar{h} \in \mathbb{R}^2} F_{h, \bar{h}}$ equipped with a linear map

$$Y(-, \underline{z}) : F \rightarrow \text{End}(F)[[z^\pm, \bar{z}^\pm, |z|^\mathbb{R}]], \quad a \mapsto Y(a, \underline{z}) = \sum_{r, s \in \mathbb{R}} a(r, s) z^{-r-1} \bar{z}^{-s-1}$$

and an element $\mathbf{1} \in F_{0,0}$ such that

PV1) For any $a, b \in F$, there exists $N \in \mathbb{R}$ such that $a(r, s)b = 0$ for any $r \leq N$ or $s \leq N$;

PV2) $F_{h, \bar{h}} = 0$ unless $h - \bar{h} \in \mathbb{Z}$;

PV3) For any $a \in F$, $Y(a, \underline{z})\mathbf{1} \in F[[z, \bar{z}]]$ and $\lim_{z \rightarrow 0} Y(a, z)\mathbf{1} = a(-1, -1)\mathbf{1} = a$;

PV4) $Y(\mathbf{1}, \underline{z}) = \text{id} \in \text{End} F$;

PV5) $F_{h, \bar{h}}(r, s)F_{h', \bar{h}'} \subset F_{h+h'-r-1, \bar{h}+\bar{h}'-s-1}$ for any $r, s, h, \bar{h}, h', \bar{h}' \in \mathbb{R}$.

A full prevertex algebra $(F, Y, \mathbf{1})$ is said to be translation covariant if there exist linear maps $D, \bar{D} \in \text{End } F$ such that

T1) $D\mathbf{1} = \bar{D}\mathbf{1} = 0$;

T2) For any $a \in F$, $[D, Y(a, \underline{z})] = \frac{d}{dz}Y(a, \underline{z})$ and $[\bar{D}, Y(a, \underline{z})] = \frac{d}{d\bar{z}}Y(a, \underline{z})$;

4.1. Locality and Associativity I. Let $(F, Y, \mathbf{1}, D, \bar{D})$ be a translation covariant full prevertex algebra.

In this section, we consider the following conditions:

FL) For any $a_1, a_2, a_3 \in F$ and $u \in F^\vee$, there exists $\mu \in \text{GCor}_2$ such that

$$u(Y(a_2, \underline{z}_2)Y(a_1, \underline{z}_1)a_3) = \mu|_{|z_1| > |z_2|}$$

$$u(Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)a_3) = \mu|_{|z_2| > |z_1|}$$

and

FA) For any $a_1, a_2, a_3 \in F$ and $u \in F$, there exists $\mu \in \text{GCor}_2$ such that

$$u(Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)a_3) = \mu|_{|z_1| > |z_2|}$$

$$u(Y(Y(a_1, \underline{z}_0)a_2, \underline{z}_2)a_3) = \mu|_{|z_2| > |z_1 - z_2|}$$

and

FSS) $Y(a_1, \underline{z})a_2 = e^{D\underline{z} + \bar{D}\bar{z}}Y(a_2, -\underline{z})a_1$ for any $a_1, a_2 \in F$.

The condition (FL) (resp. (FA) and (FSS)) is a generalization of the locality (resp. associativity, skew-symmetry) of a vertex algebra.

Lemma 4.1. *If a translation covariant full prevertex algebra $(F, Y, \mathbf{1}, D, \bar{D})$ satisfies condition (FL), then the skew-symmetry, (FSS), holds.*

Proof. Since $DY(a, z)\mathbf{1} = d/dzY(a, z)\mathbf{1}$, we have $Y(a, z)\mathbf{1} = \exp(Dz + \bar{D}\bar{z})a$, which implies that $DF_{h, \bar{h}} \subset F_{h+1, \bar{h}}$ and $\bar{D}F_{h, \bar{h}} \subset F_{h, \bar{h}+1}$. Let $a_i \in F_{h_i, \bar{h}_i}$ ($i = 1, 2$) and $u \in F_{h_0, \bar{h}_0}^*$. Then,

$$\begin{aligned} u(Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)\mathbf{1}) &= u(Y(a_1, \underline{z}_1)\exp(Dz_2 + \bar{D}\bar{z}_2)a_2) \\ &= \lim_{z_{12} \rightarrow (z_1 - z_2) \parallel_{|z_1| > |z_2|}} u(\exp(Dz_2 + \bar{D}\bar{z}_2)a_2)Y(a_1, \underline{z}_{12})a_2. \end{aligned}$$

Set $h = h_1 + h_2 - h_0$ and $\bar{h} = \bar{h}_1 + \bar{h}_2 - \bar{h}_0$. Then, by (PV5),

$$\begin{aligned} u(\exp(Dz_2 + \bar{D}\bar{z}_2)a_2)Y(a_1, \underline{z}_{12})a_2 &= \sum_{s_1, \bar{s}_1 \in \mathbb{R}} \sum_{n, \bar{n} \in \mathbb{Z}_{\geq 0}} \frac{1}{n!\bar{n}!} u(D^n \bar{D}^{\bar{n}} a_1(s_1, \bar{s}_1)a_2) z_{12}^{-s_1-1} \bar{z}_{12}^{-\bar{s}_1-1} z_2^n \bar{z}_2^{\bar{n}} \\ &= \sum_{n, \bar{n} \in \mathbb{Z}_{\geq 0}} \frac{1}{n!\bar{n}!} u(D^n \bar{D}^{\bar{n}} a_1(h+n-1, \bar{h}+\bar{n}-1)a_2) z_{12}^{-h-n} \bar{z}_{12}^{-\bar{h}-\bar{n}} z_2^n \bar{z}_2^{\bar{n}}. \end{aligned}$$

By (PV1), there exists an integer N such that $a_1(h+n-1, \bar{h}+\bar{n}-1)a_2 = 0$ for any $n \geq N$ or $\bar{n} \geq N$. Thus, $z_{12}^{N+h} \bar{z}_{12}^{\bar{N}+\bar{h}} u(\exp(Dz_2 + \bar{D}\bar{z}_2)a_2)Y(a_1, \underline{z}_{12})a_2 \in \mathbb{C}[z_{12}, z_2, \bar{z}_{12}, \bar{z}_2]$. Hence, by (FL), $\{(z_1 - z_2)^{N+h}(\bar{z}_1 - \bar{z}_2)^{\bar{N}+\bar{h}}\}_{\parallel_{|z_1| > |z_2|}} u(Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)\mathbf{1}) = \{(z_1 - z_2)^{N+h}(\bar{z}_1 - \bar{z}_2)^{\bar{N}+\bar{h}}\}_{\parallel_{|z_2| > |z_1|}} u(Y(a_2, \underline{z}_2)Y(a_1, \underline{z}_1)\mathbf{1})$, which implies

$$\begin{aligned} (-z_2)^{N+h}(-\bar{z}_2)^{\bar{N}+\bar{h}} u(Y(a_2, z_2)a_1) &= \lim_{z_1 \rightarrow 0} \{(z_1 - z_2)^{N+h}(\bar{z}_1 - \bar{z}_2)^{\bar{N}+\bar{h}}\}_{\parallel_{|z_1| > |z_2|}} u(Y(a_1, z_1)Y(a_2, z_2)\mathbf{1}) \\ &= \lim_{z_1 \rightarrow 0} \lim_{z_{12} \rightarrow (z_1 - z_2) \parallel_{|z_1| > |z_2|}} z_{12}^{N+h} \bar{z}_{12}^{\bar{N}+\bar{h}} u(\exp(Dz_2 + \bar{D}\bar{z}_2)a_2)Y(a_1, z_{12})a_2 \\ &= (-z_2)^{N+h}(-\bar{z}_2)^{\bar{N}+\bar{h}} u(\exp(Dz_2 + \bar{D}\bar{z}_2)a_2)Y(a_1, -z_2)a_2. \end{aligned}$$

Hence, the assertion holds. $\square$

Then, we have:

Proposition 4.1. *Assume that a translation covariant full prevertex algebra $(F, Y, \mathbf{1}, D, \bar{D})$ satisfies the condition (FL) and the spectrum of F is bounded below, that is, there exists $N \in \mathbb{R}$ such that $F_{h, \bar{h}} = 0$ for any $h \leq N$ or $\bar{h} \leq N$. Then, $(F, Y, \mathbf{1})$ is a full vertex algebra.*

Proof. Let $a_1, a_2, a_3 \in F$ and $u \in F^\vee$. By (FL), there exists $\mu \in \text{GCor}_2$ such that

$$\begin{aligned} u(Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)a_3) &= \mu_{\parallel_{|z_1| > |z_2|}} \\ u(Y(a_2, \underline{z}_2)Y(a_1, \underline{z}_1)a_3) &= \mu_{\parallel_{|z_2| > |z_1|}}. \end{aligned}$$

By Lemma 4.1,

$$u(Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)a_3) = \lim_{z_{12} \rightarrow (z_1 - z_2) \parallel_{|z_1| > |z_2|}} u(\exp(Dz_2 + \bar{D}\bar{z}_2)Y(a_1, \underline{z}_{12})Y(a_3, -\underline{z}_2)a_2).$$

By the assumption, $u(D^n \bar{D}^{\bar{n}} -) = 0$ for sufficiently large n or $\bar{n}$ in $\mathbb{Z}_{\geq 0}$. Then, by (FL),

$$\begin{aligned} u(\exp(Dz_2 + \bar{D}\bar{z}_2)Y(a_1, \underline{z}_{12})Y(a_3, -\underline{z}_2)a_2) &= \mu_{\parallel_{|z_{12}| > |z_2|}} \\ u(\exp(Dz_2 + \bar{D}\bar{z}_2)Y(a_3, -\underline{z}_2)Y(a_1, \underline{z}_{12})a_2) &= \mu_{\parallel_{|z_2| > |z_{12}|}}. \end{aligned}$$

Since $u(\exp(Dz_2 + \bar{D}\bar{z}_2)Y(a_3, -\underline{z}_2)Y(a_1, \underline{z}_{12})a_2) = u(Y(Y(a_1, \underline{z}_{12})a_2, \underline{z}_2)a_3)$, (FV5) holds. $\square$

In the above proposition, we assume that the spectrum of F is bounded below. However, in the proof, we only use the property that for any $a_1, a_2, a_3 \in F$ and $u \in F^\vee$, there exists N such that $u(D^n \bar{D}^m Y(a, z_1) Y(b, z_2) c) = 0$ for any $n \geq N$ or $m \geq N$. Thus, we have:

Proposition 4.2. *Let $(F, Y, \mathbf{1}, D, \bar{D})$ be a translation covariant full prevertex algebra satisfies the condition (FL) and M be an abelian group. Suppose that there exists an $M \times \mathbb{R}^2$ -grading on F , $F = \bigoplus_{\alpha \in M, h, \bar{h} \in \mathbb{R}} F_{h, \bar{h}}^\alpha$ such that:*

- (1) $F_{h, \bar{h}} = \bigoplus_{\alpha \in M} F_{h, \bar{h}}^\alpha$ for any $h, \bar{h} \in \mathbb{R}$ and $\alpha \in M$
- (2) $F_{h, \bar{h}}^\alpha(r, s) F_{h', \bar{h}'}^{\alpha'} \subset F_{h+h'-r-1, \bar{h}+\bar{h}'-s-1}^{\alpha+\alpha'}$ for any $r, s, h, \bar{h}, h', \bar{h}' \in \mathbb{R}$ and $\alpha, \alpha' \in M$;
- (3) For any α , there exists N such that $F_{h, \bar{h}}^\alpha = 0$ for any $h \geq N$ or $\bar{h} \geq N$.

Then, F is a full vertex algebra.

Proposition 4.3. *Assume that a translation covariant full prevertex algebra $(F, Y, \mathbf{1}, D, \bar{D})$ satisfies the condition (FA) and the skew-symmetry. Then, $(F, Y, \mathbf{1})$ is a full vertex algebra.*

Proof. We first prove that $Y(Da, \underline{z}) = d/dz Y(a, \underline{z})$ for any $a \in F$. For any $a, b \in F$ and $u \in F^\vee$, by (FFA), there exists $\mu \in \text{GCor}_2$ such that

$$\begin{aligned} u(Y(a, \underline{z}_1) b) &= \mu|_{|z_1| > |z_2|} \\ u(Y(Y(a, \underline{z}_0) \mathbf{1}, \underline{z}_2) b) &= \mu|_{|z_2| > |z_1 - z_2|}. \end{aligned}$$

By (PV5), $u(Y(a, \underline{z}_1) b) = p(z_1) \in \mathbb{C}[z_1^\pm, \bar{z}_1^\pm, |z_1|^\mathbb{R}]$. Thus, $\mu|_{|z_2| > |z_1 - z_2|} = \lim_{z_1 \rightarrow (z_2 + z_0)|_{|z_2| > |z_0|}} p(z_1) = \exp(z_0 d/dz_2 + \bar{z}_0 d/d\bar{z}_2) p(z_2)$. Hence,

$$u(Y(Y(a, \underline{z}_0) \mathbf{1}, \underline{z}_2) b) = \exp(z_0 d/dz_2 + \bar{z}_0 d/d\bar{z}_2) u(Y(a, \underline{z}_2) b),$$

which implies that $Y(Da, \underline{z}_2) = d/dz_2 Y(a, \underline{z}_2)$ and similarly $Y(\bar{D}a, \underline{z}_2) = d/d\bar{z}_2 Y(a, \underline{z}_2)$.

Now, we will show the assertion. Let $a_1, a_2, a_3 \in F$ and $u \in F^\vee$. By (FA), there exists $\mu \in \text{GCor}_2$ such that

$$\begin{aligned} u(Y(a_1, \underline{z}_1) Y(a_2, \underline{z}_2) a_3) &= \mu|_{|z_1| > |z_2|} \\ u(Y(Y(a_1, \underline{z}_0) a_2, \underline{z}_2) a_3) &= \mu|_{|z_2| > |z_1 - z_2|}. \end{aligned}$$

By the skew symmetry,

$$\begin{aligned} u(Y(Y(a_1, \underline{z}_0) a_2, \underline{z}_2) a_3) &= u(Y(\exp(Dz_0 + \bar{D}\bar{z}_0) Y(a_2, -\underline{z}_0) a_1, \underline{z}_2) a_3) \\ &= \lim_{\underline{z}_1 \rightarrow (\underline{z}_2 + \underline{z}_0)|_{|z_2| > |z_0|}} u(Y(Y(a_2, -\underline{z}_0) a_1, \underline{z}_1) a_3). \end{aligned}$$

Then, applying (FA) again, (FV5) holds. $\square$

4.2. Locality and Associativity II. In this section, we give a sufficient condition in order to construct a full vertex algebra by using four point functions. We remark that for a self-dual QP-generated full vertex operator algebra F , there is the linear map $(\mathbf{1}, -) : F \rightarrow \mathbb{C}$ such that Proposition 3.7 holds.

In this section, we assume that $(F, Y, \mathbf{1}, D, \bar{D})$ is a translation covariant full prevertex algebra equipped with a linear map $\langle \rangle : F \rightarrow \mathbb{C}$ such that

E1) $F_{h,\bar{h}}$ is a finite dimensional vector space for any $h, \bar{h} \in \mathbb{R}$;

E2) $\langle \mathbf{1} \rangle = 1$ and $\langle Da \rangle = \langle \bar{D}a \rangle = 0$ for any $a \in F$;

E3) $\langle F_{h,\bar{h}} \rangle = 0$ for any $(h, \bar{h}) \neq (0, 0)$;

E4) There exists $N \in \mathbb{R}$ such that $F_{h,\bar{h}} = 0$ for $h \leq N$ or $\bar{h} \leq N$.

We remark that if $F_{0,0} = \mathbb{C}\mathbf{1}$, then $\langle \rangle$ is simply the projection $F \rightarrow F_{0,0}$.

For $(F, Y, \mathbf{1}, D, \bar{D}, \langle \rangle)$, set

$$R_{F,\langle \rangle} = \{b \in F \mid \langle Y(a, \underline{z})b \rangle = 0 \text{ for any } a \in F\}.$$

We say $(F, Y, \mathbf{1}, D, \bar{D}, \langle \rangle)$ is non-degenerate if $R_{F,\langle \rangle} = 0$.

In this section, we consider the following conditions:

FFL) For any $a_1, a_2, a_3, a_4 \in F$, there exists $\phi \in \text{Cor}_4$ such that

$$\begin{aligned} \langle Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)Y(a_3, \underline{z}_3)Y(a_4, \underline{z}_4)\mathbf{1} \rangle &= e_{1(2(3(4\star)))}\phi \\ \langle Y(a_1, \underline{z}_1)Y(a_3, \underline{z}_3)Y(a_2, \underline{z}_2)Y(a_4, \underline{z}_4)\mathbf{1} \rangle &= e_{1(3(2(4\star)))}\phi \end{aligned}$$

and

FFA) For any $a_1, a_2, a_3, a_4 \in F$, there exists $\phi \in \text{Cor}_4$ such that

$$\begin{aligned} \langle Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)Y(a_3, \underline{z}_3)Y(a_4, \underline{z}_4)\mathbf{1} \rangle &= e_{1(2(3(4\star)))}\phi \\ \langle Y(a_1, \underline{z}_1)Y(a_4, \underline{z}_4)Y(a_3, \underline{z}_3)Y(a_2, \underline{z}_2)\mathbf{1} \rangle &= e_{1(4(3(2\star)))}\phi. \end{aligned}$$

By Corollary 1.3 and (PV4), we have:

Lemma 4.2. *Let $(F, Y, \mathbf{1}, D, \bar{D}, \langle \rangle)$ satisfy (FFL) or (FFA). Then, for $a_1, a_2, a_3 \in F$, there exists $\phi \in \text{Cor}_3$ such that for any $\sigma \in S_3$,*

$$\langle Y(a_{\sigma 1}, z_{\sigma 1})Y(a_{\sigma 2}, z_{\sigma 2})Y(a_{\sigma 3}, z_{\sigma 3})\mathbf{1} \rangle = e_{\sigma 1(2(3\star))}\phi.$$

Lemma 4.3. *Let $(F, Y, \mathbf{1}, D, \bar{D}, \langle \rangle)$ satisfy (FFL) or (FFA). Then, for $a_1, a_2, a_3 \in F$,*

$$\langle Y(a_1, \underline{z}_{12})Y(a_3, \underline{z}_{32})a_2 \rangle = \langle Y(a_1, \underline{z}_{12})\exp(Dz_{32} + \bar{D}\bar{z}_{32})Y(a_2, -\underline{z}_{32})a_3 \rangle.$$

Proof. Let $\phi \in \text{Cor}_3$ such that $\langle Y(a_1, \underline{z}_{13})Y(a_2, \underline{z}_{23})a_3 \rangle = e_{1(23)}(\phi)$. Similarly to the proof of Lemma 1.11,

$$e_{1(32)} = \lim_{\substack{\underline{z}_{23} \rightarrow -\underline{z}_{32} \\ \underline{z}_{13} \rightarrow (\underline{z}_{12} - \underline{z}_{32}) \parallel \|\underline{z}_{12}\| > \|\underline{z}_{32}\|}} e_{1(23)}.$$

Thus, by Lemma 4.2 and (E2),

$$\begin{aligned} \langle Y(a_1, \underline{z}_{12})Y(a_3, \underline{z}_{32})a_2 \rangle &= e_{1(32)}(\phi) \\ &= \exp(-z_{32}d/dz_{12} - \bar{z}_{32}d/d\bar{z}_{12}) \langle Y(a_1, \underline{z}_{12})Y(a_2, -\underline{z}_{32})a_3 \rangle \\ &= \langle \exp(-Dz_{32} - \bar{D}\bar{z}_{32})Y(a_1, \underline{z}_{12})\exp(Dz_{32} + \bar{D}\bar{z}_{32})Y(a_2, -\underline{z}_{32})a_3 \rangle \\ &= \langle Y(a_1, \underline{z}_{12})\exp(Dz_{32} + \bar{D}\bar{z}_{32})Y(a_2, -\underline{z}_{32})a_3 \rangle. \end{aligned}$$

□

We remark that we cannot obtain the skew-symmetry from the above lemma, since the skew-symmetry holds only in $\langle -, - \rangle \in F^\vee$. However, if $(F, Y, \mathbf{1}, D, \bar{D}, \langle, \rangle)$ is non-degenerate, then, by (E1) and (PV5), $F^\vee$ is spanned by $\{\langle a(s, \bar{s}) - \rangle\}_{a \in F, s, \bar{s} \in \mathbb{R}}$. Thus, we have:

Lemma 4.4. *If $(F, Y, \mathbf{1}, D, \bar{D}, \langle, \rangle)$ is non-degenerate and satisfy (FFL) or (FFA), then the skew-symmetry holds.*

Proposition 4.4. *If $(F, Y, \mathbf{1}, D, \bar{D}, \langle, \rangle)$ is non-degenerate and satisfy (FFL) or (FFA), then $(F, Y, \mathbf{1})$ is a full vertex algebra.*

Proof. First, we assume that (FFL) holds. Then, by Lemma 1.4, the condition (FL) holds. Thus, by Proposition 4.1, F is a full vertex algebra. Next, we assume that (FFA) holds. Then, for any $a_1, a_2, a_3, a_4 \in F$, there exists $\phi \in \text{Cor}_4$ such that

$$\begin{aligned} \langle Y(a_1, \underline{z}_{14})Y(a_2, \underline{z}_{24})Y(a_3, \underline{z}_{34})a_4 \rangle &= e_{1(2(34))}\phi \\ \langle Y(a_1, \underline{z}_{12})Y(a_4, \underline{z}_{42})Y(a_3, \underline{z}_{32})a_2 \rangle &= e_{1(4(32))}\phi. \end{aligned}$$

Then, by Lemma 4.4 and Lemma 1.11, $\langle Y(a_1, \underline{z}_{12})Y(a_4, \underline{z}_{42})Y(a_2, \underline{z}_{23})a_3 \rangle = e_{1(4(23))}\phi$. By Lemma 4.4 and (E2),

$$\begin{aligned} &\langle Y(a_1, \underline{z}_{12})Y(a_4, \underline{z}_{42})Y(a_2, \underline{z}_{23})a_3 \rangle \\ &= \langle Y(a_1, \underline{z}_{12})\exp(D\underline{z}_{42} + \bar{D}\bar{z}_{42})Y(Y(a_2, \underline{z}_{23})a_3, -\underline{z}_{42})a_4 \rangle \\ &= \exp(\underline{z}_{42}d/d\underline{z}_{12} + \bar{z}_{42}d/d\bar{z}_{12})\langle Y(a_1, \underline{z}_{12})Y(Y(a_2, \underline{z}_{23})a_3, -\underline{z}_{42})a_4 \rangle. \end{aligned}$$

Then, by Lemma 1.9,

$$\langle Y(a_1, \underline{z}_{14})Y(Y(a_2, \underline{z}_{23})a_3, \underline{z}_{24})a_4 \rangle = e_{1((23)4)}\phi.$$

By Lemma 1.4 again, the condition (FA) holds. Thus, by Proposition 4.3, F is a full vertex algebra. $\square$

In the rest of this section, we study the condition that $(F, Y, \mathbf{1}, D, \bar{D}, \langle, \rangle)$ is non-degenerate. A two-sided ideal of $(F, Y, \mathbf{1}, D, \bar{D}, \langle, \rangle)$ is a subspace $I \subset F$ such that $I = \bigoplus_{h, \bar{h} \in \mathbb{R}} I \cap F_{h, \bar{h}}$, i.e., a graded subspace, and for any $v \in I$ and $a \in F$, $Y(a, \underline{z})v, Y(v, \underline{z})a \in I((\underline{z}, \bar{z}, |\underline{z}|^{\mathbb{R}}))$.

Then, we have:

Lemma 4.5. *If $(F, Y, \mathbf{1}, D, \bar{D}, \langle, \rangle)$ satisfy (FFL) or (FFA), the subspace $R_{F, \langle, \rangle}$ is a two-sided ideal of F .*

Proof. By (PV5) and (E3), $R_{F, \langle, \rangle}$ is a graded subspace of F . Let $a_1, a_2 \in F$ and $v \in R_{F, \langle, \rangle}$. It suffices to show that $\langle Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)Y(v, \underline{z}_3)\mathbf{1} \rangle = \langle Y(a_1, \underline{z}_1)Y(v, \underline{z}_3)Y(a_2, \underline{z}_2)\mathbf{1} \rangle = 0$. By Lemma 4.3 and (E2), we have $\langle Y(a, \underline{z})b \rangle = \langle Y(b, -\underline{z})a \rangle$ for any $a, b \in F$. Since $\langle Y(v, \underline{z}_3)Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)\mathbf{1} \rangle = \langle Y(Y(a_1, \underline{z}_1)Y(a_2, \underline{z}_2)\mathbf{1}, -\underline{z}_3)v \rangle = 0$, by Lemma 4.2, the assertion holds. $\square$

By the above lemma, the vertex operator $Y(-, \underline{z}) : F/R_{F, \langle \rangle} \rightarrow \text{End } (F/R_{F, \langle \rangle})[[z, \bar{z}, |z|^{\mathbb{R}}]]$, $D, \bar{D} \in F/R_{F, \langle \rangle}$ and $\langle \rangle : F/R_{F, \langle \rangle} \rightarrow \mathbb{C}$ are induced from $(F, Y, \mathbf{1}, D, \bar{D}, \langle \rangle)$ and are non-degenerate. We remark that by the construction, a parenthesized correlation function of $F/R_{F, \langle \rangle}$ and F are the same. Thus, we have:

Proposition 4.5. *Let $(F, Y, \mathbf{1}, D, \bar{D}, \langle \rangle)$ satisfy (FFL) or (FFA). Then, $F/R_{F, \langle \rangle}$ is a full vertex algebra.*

$(F, Y, \mathbf{1}, D, \bar{D}, \langle \rangle)$ is said to be simple if there is no proper left ideal of F . By the above lemma, we have:

Proposition 4.6. *Let $(F, Y, \mathbf{1}, D, \bar{D}, \langle \rangle)$ satisfy (FFL) or (FFA). If $(F, Y, \mathbf{1}, D, \bar{D}, \langle \rangle)$ is simple, then it is non-degenerate. Conversely, if $(F, Y, \mathbf{1}, D, \bar{D}, \langle \rangle)$ is non-degenerate and $F_{0,0} = \mathbb{C}\mathbf{1}$, then $(F, Y, \mathbf{1}, D, \bar{D}, \langle \rangle)$ is simple.*

Proof. By Lemma 4.5, $R_{F, \langle \rangle}$ is a left ideal. If F is simple, then $R_{F, \langle \rangle} = 0$ by (E2), thus non-degenerate. Suppose that $(F, Y, \mathbf{1}, D, \bar{D}, \langle \rangle)$ is non-degenerate and $F_{0,0} = \mathbb{C}\mathbf{1}$ and let $I \subset F$ be a left ideal which does not contain $\mathbf{1}$. Since $I \cap F_{0,0} = 0$, $\langle Y(a, \underline{z})v \rangle = 0$ for any $a \in F$ and $v \in I$. Hence, $I \subset R_{F, \langle \rangle} = 0$. $\square$

4.3. Bootstrap. In this section, we study the bootstrap equation. Set $L(-1) = D$ and $\bar{L}(-1) = \bar{D}$ and let $L(0), \bar{L}(0) \in \text{End } F$ be linear maps defined by $L(0)|_{F_{h,\bar{h}}} = h$ and $\bar{L}(0)|_{F_{h,\bar{h}}} = \bar{h}$ for any $h, \bar{h} \in \mathbb{R}$. We assume that $(F, Y, \mathbf{1}, D, \bar{D})$ is a translation covariant full prevertex algebra equipped with linear maps $L(1), \bar{L}(1) \in \text{End } F$ and a non-degenerate symmetric bilinear form $(-, -) : F \times F \rightarrow \mathbb{C}$ such that:

- (1) There exists $N \in \mathbb{R}$ such that $F_{h,\bar{h}} = 0$ for any $h \leq N$ or $\bar{h} \leq N$;
- (2) $\dim F_{h,\bar{h}}$ is finite for any $h, \bar{h} \in \mathbb{R}$;
- (3) For any $i, j = -1, 0, 1$,

$$[L(i), L(j)] = (i - j)L(i + j),$$

$$[\bar{L}(i), \bar{L}(j)] = (i - j)\bar{L}(i + j),$$

$$[L(i), \bar{L}(j)] = 0;$$

- (4) For any $i = -1, 0, 1$ and $a, b \in F$,

$$(L(i)a, b) = (a, L(-i)b)$$

$$(\bar{L}(i)a, b) = (a, \bar{L}(-i)b);$$

- (5) For $k = -1, 0, 1$,

$$[L(k), Y(a, \underline{z})] = \sum_{i=0}^{k+1} \binom{k+1}{i+1} Y(L(i+1)a, \underline{z}) z^{-i}$$

$$[\bar{L}(k), Y(a, \underline{z})] = \sum_{i=0}^{k+1} \binom{k+1}{i+1} Y(\bar{L}(i+1)a, \underline{z}) \bar{z}^{-i};$$

- (6) $L(1)\mathbf{1} = \bar{L}(1)\mathbf{1} = 0$;

(7) For any $a, b, c \in F$,

$$(a, Y(b, \underline{z})c) = (Y(S_{\underline{z}}b, \underline{z}^{-1})a, c).$$

Let $(F, Y, \mathbf{1}, L(i), \bar{L}(i), (-, -))$ satisfy the above condition. For any $a_1, a_2, a_3, a_4 \in F$, define s,t,u-channels by

$$\text{(s-channel)} \quad S_{(21)(34)}(a_1, a_2, a_3, a_4) = (\mathbf{1}, ((a_2(z_{21})a_1)(z_{14})a_3(z_{34})a_4))$$

$$\text{(t-channel)} \quad S_{(41)(32)}(a_1, a_2, a_3, a_4) = (\mathbf{1}, ((a_4(z_{41})a_1)(z_{12})a_3(z_{32})a_2))$$

$$\text{(u-channel)} \quad S_{(31)(24)}(a_1, a_2, a_3, a_4) = (\mathbf{1}, ((a_3(z_{31})a_1)(z_{14})a_2(z_{24})a_4)).$$

$(F, Y, \mathbf{1}, L(i), \bar{L}(i), (-, -))$ is said to satisfy a bootstrap equation for the s-channel and t-channel (resp. for the s-channel and u-channel) if for any $a_1, a_2, a_3, a_4 \in F$, there exists $\phi \in \text{Cor}_4$ such that $S_{(21)(34)}(a_1, a_2, a_3, a_4) = e_{(21)(34)}(\phi)$ and $S_{(41)(32)}(a_1, a_2, a_3, a_4) = e_{(41)(32)}(\phi)$ (resp. $S_{(21)(34)}(a_1, a_2, a_3, a_4) = e_{(21)(34)}(\phi)$ and $S_{(31)(24)}(a_1, a_2, a_3, a_4) = e_{(31)(24)}(\phi)$).

The notion, QP-generated, is defined for $(F, Y, \mathbf{1}, L(i), \bar{L}(i), (-, -))$ similarly to the full vertex operator algebra (see Section 3.2).

Let $(F, Y, \mathbf{1}, L(i), \bar{L}(i), (-, -))$ be QP-generated and satisfy the bootstrap equation for the s-channel and t-channel. Set $\langle - \rangle = (\mathbf{1}, -)$. We will show that $(F, Y, \mathbf{1}, D, \bar{D}, \langle \rangle)$ satisfy the assumption of Proposition 4.4. It suffices to show that (FFA) holds. Similarly to the proof of Proposition 3.7, we may assume that all $a_i \in F_{h_i, \bar{h}_i}$ are quasi-primary states. Then, by $\text{PSL}_2\mathbb{C}$ -symmetry, in order to verify (FFA), it suffices to show that there exists $f \in F_{0,1,\infty}$ such that

$$Q(h, z)^{-1}(\mathbf{1}, Y(a_1, z_1)Y(a_2, z_2)Y(a_3, z_3)a_4) = e_{1(2(34))}(f \circ \xi),$$

$$Q(h, z)^{-1}(\mathbf{1}, Y(a_1, z_1)Y(a_4, z_4)Y(a_2, z_2)a_3) = e_{1(4(32))}(f \circ \xi).$$

By Lemma 3.14, we can calculate $v_{1(2(34))}(Q(h, z)^{-1}S_{1(2(34))})$ and $v_{1(4(32))}(Q(h, z)^{-1}S_{1(4(32))})$ similarly to the proof of Theorem 3.2, which are equal to $j(p, f)$ and $j(1 - p, f)$. The bootstrap equation for the s-channel and u-channel implies the same result. The detail verification is left to the reader. Thus, we have:

Proposition 4.7. *If $(F, Y, \mathbf{1}, L(i), \bar{L}(i), (-, -))$ is QP-generated and satisfy the bootstrap equation for the s-channel and t-channel or the s-channel and u-channel, then F is a full vertex algebra.*

5. EXAMPLE

In this section, we construct a full vertex algebras from an AH pair, which is introduced in [Mo1]. One can construct an AH pair from an even lattice, called twisted group algebra [FLM, Mo1]. The full vertex algebra constructed from an even lattice is appeared in the toroidal compactification of string theory and is shown to be a QP-generated self-dual full vertex operator algebra. A conceptual origin of AH pairs is provided in Section 5.3, where we show that an AH pair is a commutative algebra object in some braided tensor category.

5.1. Full vertex algebra and AH pair. We first recall a concept of an AH pair introduced in [Mo1]. Let H be a finite-dimensional vector space over $\mathbb{R}$ equipped with a non-degenerate symmetric bilinear form $(-, -) : H \times H \rightarrow \mathbb{R}$ and A a unital associative algebra over $\mathbb{C}$ with the unity 1. Assume that A is graded by H as $A = \bigoplus_{\alpha \in H} A^\alpha$.

We will say that such a pair (A, H) is an *AH pair* if the following conditions hold:

- AH1) $1 \in A^0$ and $A^\alpha A^\beta \subset A^{\alpha+\beta}$ for any $\alpha, \beta \in H$;
- AH2) $A^\alpha A^\beta \neq 0$ implies $(\alpha, \beta) \in \mathbb{Z}$;
- AH3) For $v \in A^\alpha, w \in A^\beta, vw = (-1)^{(\alpha, \beta)} wv$.

An AH pair (A, H) is called an *even AH pair* if

$$\text{EAH } (\alpha, \alpha) \in 2\mathbb{Z} \text{ for any } \alpha \in H \text{ with } A^\alpha \neq 0.$$

Let $P(H)$ be the set of linear maps $p \in \text{End } H$ such that:

- P1) p is a projection, that is, $p^2 = p$;
- P2) The subspaces $\ker p$ and $\ker(1 - p)$ are orthogonal to each other.

Let $P_>(H)$ be the subset of $P(H)$ such that

- P3) $\ker(1 - p)$ is positive definite and $\ker p$ is negative-definite.

Let (A, H) be an even AH pair and $p \in P(H)$. In this section, we construct a full vertex algebra $F_{A, H, p}$ for the triple (A, H, p) .

Set $\bar{p} = 1 - p$ and $H_l = \ker \bar{p}$ and $H_r = \ker p$. Define the new symmetric bilinear forms $(-, -)_p : H \times H \rightarrow \mathbb{R}$ by

$$(h, h')_p = (ph, ph') - (\bar{p}h, \bar{p}h').$$

By (P1) and (P2), $(-, -)_p$ is non-degenerate. Let $\hat{H}^p = \bigoplus_{n \in \mathbb{Z}} H \otimes \mathbb{C}t^n \oplus \mathbb{C}c$ be the affine Heisenberg Lie algebra associated with $(\hat{H}^p, (-, -)_p)$ and $\hat{H}_{\geq 0}^p = \bigoplus_{n \geq 0} H \otimes \mathbb{C}t^n \oplus \mathbb{C}c$ a subalgebra of $\hat{H}^p$. Define the action of $\hat{H}_{\geq 0}^p$ on A by

$$\begin{aligned} ca &= a \\ h \otimes t^n a &= \begin{cases} 0, & n \geq 1, \\ (h, \alpha)_p a, & n = 0 \end{cases} \end{aligned}$$

for $\alpha \in H$ and $a \in A^\alpha$. Let $F_{A, H, p}$ be the $\hat{H}^p$ -module induced from A and $M_{H, p}$ be the submodule of $F_{A, H, p}$ generated by the unit $1 \in A^0$ as a $\hat{H}^p$ -module. Denote by $h(n)$ the

action of $h \otimes t^n$ on $F_{A,H,p}$ for $n \in \mathbb{Z}$. For $h \in H$, set

$$\begin{aligned} h(z, \bar{z}) &= \sum_{n \in \mathbb{Z}} ((ph)(n)z^{-n-1} + (\bar{p}h)(n)\bar{z}^{-n-1}) \in \text{End} F_{A,H,p}[[z^\pm, \bar{z}^\pm]] \\ h^+(\underline{z}) &= \sum_{n \geq 0} ((ph)(n)z^{-n-1} + (\bar{p}h)(n)\bar{z}^{-n-1}) \\ h^-(\underline{z}) &= \sum_{n \geq 0} ((ph)(-n-1)z^n + (\bar{p}h)(-n-1)\bar{z}^n). \\ E^+(h, \underline{z}) &= \exp\left(-\sum_{n \geq 1} \left(\frac{ph(n)}{n}z^{-n} + \frac{\bar{p}h(n)}{n}\bar{z}^{-n}\right)\right) \\ E^-(h, \underline{z}) &= \exp\left(-\sum_{n \leq -1} \left(\frac{ph(n)}{n}z^{-n} + \frac{\bar{p}h(n)}{n}\bar{z}^{-n}\right)\right). \end{aligned}$$

For $h_r \in H_r$ and $h_l \in H_l$, $h_r(\underline{z})$ and $h_l(\underline{z})$ are denoted by $h_r(z)$ and $h_l(z)$.

Then, similarly to the case of a lattice vertex algebra, we have:

Lemma 5.1. *For any $h_1, h_2 \in H$,*

$$E^+(h_1, \underline{z}_1)E^-(h_2, \underline{z}_2) = \left(\sum_{n, \bar{n} \geq 0} \binom{(ph_1, ph_2)_p}{n} \binom{(\bar{p}h_1, \bar{p}h_2)_p}{\bar{n}} z_2^n z_1^{-n} \bar{z}_2^{\bar{n}} \bar{z}_1^{-\bar{n}}\right) E^-(h_2, \underline{z}_2)E^+(h_1, \underline{z}_1).$$

We remark that the formal power series $\sum_{n, \bar{n} \geq 0} \binom{(ph_1, ph_2)_p}{n} \binom{(\bar{p}h_1, \bar{p}h_2)_p}{\bar{n}} z_2^n z_1^{-n} \bar{z}_2^{\bar{n}} \bar{z}_1^{-\bar{n}}$ is equal to $(1 - z_2/z_1)^{(ph_1, ph_2)_p} (1 - \bar{z}_2/\bar{z}_1)^{(\bar{p}h_1, \bar{p}h_2)_p} |_{|z_1| > |z_2|}$.

Let $\alpha \in H$ and $a \in A^\alpha$. Denote by $l_a \in \text{End} A$ the left multiplication by a and define the linear map $l_a z^{p\alpha} \bar{z}^{\bar{p}\alpha} : A \rightarrow A[z, \bar{z}, |z|^\mathbb{R}]$ by $l_a z^{p\alpha} \bar{z}^{\bar{p}\alpha} b = z^{(p\alpha, p\beta)_p} \bar{z}^{(\bar{p}\alpha, \bar{p}\beta)_p} ab$ for $\beta \in H$ and $b \in A^\beta$. Assume that $ab \neq 0$. Then, $(\alpha, \beta) \in \mathbb{Z}$ and since $z^{(p\alpha, p\beta)_p} \bar{z}^{(\bar{p}\alpha, \bar{p}\beta)_p} = |z|^{(p\alpha, p\beta)_p} \bar{z}^{(\bar{p}\alpha, \bar{p}\beta)_p - (p\alpha, p\beta)_p}$ and $(\bar{p}\alpha, \bar{p}\beta)_p - (p\alpha, p\beta)_p = -(\alpha, \beta) \in \mathbb{Z}$, we have $l_a z^{p\alpha} \bar{z}^{\bar{p}\alpha} b \in A[z, \bar{z}, |z|^\mathbb{R}]$. Then, set

$$a(\underline{z}) = E^-(\alpha, \underline{z})E^+(\alpha, \underline{z})l_a z^{p\alpha} \bar{z}^{\bar{p}\alpha},$$

which is a linear map $F_{A,H,p} \rightarrow F_{A,H,p}[[z, \bar{z}, |z|^\mathbb{R}]]$.

By Poincaré-Birkhoff-Witt theorem, $F_{A,H,p} \cong M_{H,p} \otimes A$, that is, $F_{A,H,p}$ is spanned by

$$\{h_l^1(-n_1 - 1) \dots h_l^l(-n_l - 1) h_r^1(-\bar{n}_1 - 1) \dots h_r^k(-\bar{n}_k - 1) a\},$$

where $h_l^i \in H_l$, $n_i \in \mathbb{Z}_{\geq 0}$ and $h_r^j \in H_r$, $\bar{n}_j \in \mathbb{Z}_{\geq 0}$ for any $1 \leq i \leq l$ and $1 \leq j \leq k$ and $a \in A$. Then, a map $Y : F_{A,H,p} \rightarrow \text{End} F_{A,H,p}[[z, \bar{z}, |z|^\mathbb{R}]]$ is defined inductively as follows: For $a \in A$, define $Y(a, \underline{z})$ by $Y(a, \underline{z}) = a(\underline{z})$. Assume that $Y(v, \underline{z})$ is already defined for $v \in F_{A,H,p}$. Then, for $h_r \in H_r$ and $h_l \in H_l$ and $n, \bar{n} \in \mathbb{Z}_{\geq 0}$, $Y(h_l(-n-1)v, \underline{z})$ and $Y(h_r(-\bar{n}-1)v, \underline{z})$ is defined by

$$\begin{aligned} Y(h_l(-n-1)v, \underline{z}) &= 1/n! \left(\frac{d}{dz}\right)^n h_l^-(z) Y(v, \underline{z}) + Y(v, \underline{z}) 1/n! \left(\frac{d}{dz}\right)^n h_l^+(z) \\ Y(h_r(-\bar{n}-1)v, \underline{z}) &= 1/\bar{n}! \left(\frac{d}{d\bar{z}}\right)^{\bar{n}} h_r^-(\bar{z}) Y(v, \underline{z}) + Y(v, \underline{z}) 1/\bar{n}! \left(\frac{d}{d\bar{z}}\right)^{\bar{n}} h_r^+(\bar{z}). \end{aligned}$$

By the direct calculus, we can easily prove the following lemma:

Lemma 5.2. For any $v \in F_{A,H,p}$ and $h_l \in H_l$ and $h_r \in H_r$,

$$[h_l(n), Y(v, \underline{z})] = \sum_{i \geq 0} \binom{n}{i} Y(h_l(i)v, \underline{z}) z^{n-i},$$

$$[h_r(n), Y(v, \underline{z})] = \sum_{i \geq 0} \binom{n}{i} Y(h_r(i)v, \underline{z}) \bar{z}^{n-i}.$$

By the above lemma,

$$\begin{aligned} [h_l^+(z_1), Y(v, \underline{z}_2)] &= \sum_{i \geq 0, n \geq 0} \binom{n}{i} Y(h_l(i)v, \underline{z}_2) z_1^{-n-1} z_2^{n-i} \\ &= \sum_{i \geq 0} Y(h_l(i)v, \underline{z}_2) \frac{1}{i!} \frac{d^i}{dz_2} \sum_{n \geq 0} z_1^{-n-1} z_2^n \\ &= \sum_{i \geq 0} Y(h_l(i)v, \underline{z}_2) \frac{1}{(z_1 - z_2)^{i+1}} \Big|_{|z_1| > |z_2|}. \end{aligned}$$

Thus, we have:

Lemma 5.3. For any $v \in F_{A,H,p}$ and $h_l \in H_l$,

$$[h_l^+(z_1), Y(v, \underline{z}_2)] = \sum_{i \geq 0} Y(h_l(i)v, \underline{z}_2) \frac{1}{(z_1 - z_2)^{i+1}} \Big|_{|z_1| > |z_2|},$$

$$[Y(v, \underline{z}_1), h_l^-(z_2)] = \sum_{i \geq 0} Y(h_l(i)v, \underline{z}) \frac{(-1)^{i+1}}{(z_1 - z_2)^{i+1}} \Big|_{|z_1| > |z_2|}.$$

Set

$$\begin{aligned} \mathbf{1} &= 1 \otimes 1, \\ \omega &= 1/2 \sum_{i=1}^{\dim H_l} h_l^i(-1) h_l^i, \\ \bar{\omega} &= 1/2 \sum_{j=1}^{\dim H_r} h_r^j(-1) h_r^j, \end{aligned}$$

where h_l^i and h_r^j is an orthonormal basis of $H_l \otimes_{\mathbb{R}} \mathbb{C}$ and $H_r \otimes_{\mathbb{R}} \mathbb{C}$ with respect to the bilinear form $(-, -)_p$.

We will prove the following proposition by using Proposition 4.2:

Proposition 5.1. For an even AH pair (A, H) and $p \in P(H)$, $(F_{A,H,p}, Y, \mathbf{1})$ is a full vertex algebra.

Set $F = F_{A,H,p}$ and $D = \omega(0)$ and $\bar{D} = \bar{\omega}(0)$ and

$$F_{h, \bar{h}}^\alpha = \{v \in F \mid \omega(1)v = hv, \bar{\omega}(1)v = \bar{h}v, h(0)v = (\alpha, h)_p v \text{ for any } h \in H\}$$

and

$$F_{h, \bar{h}} = \bigoplus_{\alpha \in H} F_{h, \bar{h}}^\alpha,$$

for $h, \bar{h} \in \mathbb{R}$ and $\alpha \in H$. Then, $F = \bigoplus_{h, \bar{h} \in \mathbb{R}} \bigoplus_{\alpha \in H} F_{h, \bar{h}}^\alpha$. In order to prove $(F, Y, \mathbf{1})$ is a full vertex algebra, it suffices to show that $(F, Y, \mathbf{1}, D, \bar{D})$ satisfies the condition in Proposition 4.2. An easy computation show that $(F, Y, \mathbf{1}, D, \bar{D})$ is a translation covariant full prevertex algebra and $F = \bigoplus_{h, \bar{h} \in \mathbb{R}} \bigoplus_{\alpha \in H} F_{h, \bar{h}}^\alpha$ satisfies the conditions (1), (2) and (3) in Proposition 4.2. Thus, it suffices to show that the condition (FL).

For $\alpha \in H$ and $n, m \in \mathbb{Z}_{\geq 0}$, it is easy to show that $F_{\frac{1}{2}(p\alpha, p\alpha)_p + n, \frac{1}{2}(\bar{p}\alpha, \bar{p}\alpha)_p + m}^\alpha$ is spanned by $\{h_l^1(-i_1) \dots h_l^k(-i_k) h_r^1(j_1) \dots h_r^l(j_l) a\}$, where $k, l \in \mathbb{Z}_{\geq 0}$, $h_l^a \in H_l$, $h_r^b \in H_r$, $a \in A^\alpha$, $i_a, j_b \in \mathbb{Z}_{\geq 1}$, $i_1 + \dots + i_k = n$ and $j_1 + \dots + j_l = m$ for any $a = 1, \dots, k$ and $b = 1, \dots, l$. Then,

$$F^\alpha = \bigoplus_{n, m \in \mathbb{Z}_{\geq 0}} F_{\frac{1}{2}(p\alpha, p\alpha)_p + n, \frac{1}{2}(\bar{p}\alpha, \bar{p}\alpha)_p + m}^\alpha$$

and

$$F_{\frac{1}{2}(p\alpha, p\alpha)_p, \frac{1}{2}(\bar{p}\alpha, \bar{p}\alpha)_p}^\alpha = A^\alpha.$$

Let $a^* \in A^\vee = \bigoplus_{\alpha \in H} (A^\alpha)^*$ and $\langle a^*, - \rangle$ be the linear map $F \rightarrow \mathbb{C}$ defined by the composition of the projection $F = A \oplus \bigoplus_{\substack{n, m \in \mathbb{Z}_{\geq 0} \\ (n, m) \neq (0, 0)}} F_{\frac{1}{2}(p\alpha, p\alpha)_p + n, \frac{1}{2}(\bar{p}\alpha, \bar{p}\alpha)_p + m}^\alpha \rightarrow A$ and $a^* : A \rightarrow \mathbb{C}$. Then, it is easy to verify $\langle a^*, - \rangle$ is a highest weight vector, that is, $\langle a^*, h(-n) - \rangle = 0$ for any $n \geq 1$ and $h \in H$. Thus, for any $\alpha \in H$, we have:

$$\begin{aligned} E^+(\alpha, \underline{z}) \mathbf{1} &= \mathbf{1}, \\ \langle a^*, E^-(\alpha, \underline{z}) - \rangle &= \langle a^*, - \rangle. \end{aligned}$$

Thus, by using the above fact and Lemma 5.1, we have:

Lemma 5.4. For $\alpha_0, \alpha_1, \alpha_2, \alpha_3 \in H$ and $a_0^* \in (A^{\alpha_0})^*$ and $a_i \in A^{\alpha_i}$ (for $i = 1, 2, 3$),

$$\begin{aligned} &\langle a_0^*, Y(a_1, \underline{z}_1) Y(a_2, \underline{z}_2) a_3 \rangle \\ &= z_1^{(p\alpha_1, p\alpha_3)_p} z_2^{(p\alpha_2, p\alpha_3)_p} (z_1 - z_2)^{(p\alpha_1, p\alpha_2)_p} \Big|_{|z_1| > |z_2|} \\ &\quad \bar{z}_1^{(\bar{p}\alpha_1, \bar{p}\alpha_3)_p} \bar{z}_2^{(\bar{p}\alpha_2, \bar{p}\alpha_3)_p} (\bar{z}_1 - \bar{z}_2)^{(\bar{p}\alpha_1, \bar{p}\alpha_2)_p} \Big|_{|\bar{z}_1| > |\bar{z}_2|} \langle a_0^*, a_1 a_2 a_3 \rangle. \end{aligned}$$

Assume that none of $a_1 a_2, a_1 a_3, a_2 a_3$ is equal to zero. Then, $(\alpha_1, \alpha_2), (\alpha_2, \alpha_3), (\alpha_1, \alpha_3) \in \mathbb{Z}$ by (AH2). Set

$$\phi = z_1^{(p\alpha_1, p\alpha_3)_p} z_2^{(p\alpha_2, p\alpha_3)_p} (z_1 - z_2)^{(p\alpha_1, p\alpha_2)_p} \Big|_{|z_1| > |z_2|} \bar{z}_1^{(\bar{p}\alpha_1, \bar{p}\alpha_3)_p} \bar{z}_2^{(\bar{p}\alpha_2, \bar{p}\alpha_3)_p} (\bar{z}_1 - \bar{z}_2)^{(\bar{p}\alpha_1, \bar{p}\alpha_2)_p} \Big|_{|\bar{z}_1| > |\bar{z}_2|},$$

which is a function in GCor_2 . The above lemma implies that $\langle a_0^*, Y(a_2, \underline{z}_2) Y(a_1, \underline{z}_1) a_3 \rangle = (-1)^{(\alpha_2, \alpha_3)} \phi \Big|_{|z_1| > |z_2|} \langle a_0^*, a_2 a_1 a_3 \rangle$. By the definition of an AH pair, $a_3 a_2 = (-1)^{(\alpha_2, \alpha_3)} a_2 a_3$. Thus, (FL) holds. If one of $a_1 a_2, a_1 a_3, a_2 a_3$ is equal to zero, then both $\langle a_0^*, a_1 a_2 a_3 \rangle$ and $\langle a_0^*, a_2 a_1 a_3 \rangle$ are zero, which implies (FL). For $u \in F^\vee$ and $v_1, v_2, v_3 \in F$, set

$$S(u; v_1, v_2, v_3) = u(Y(v_1, \underline{z}_1) Y(v_2, \underline{z}_2) v_3).$$

We can compute $S(u; v_1, v_2, v_3)$ by using Lemma 5.4 and Lemma 5.3. We observe that $\bigoplus_{h, \bar{h}} (F_{h, \bar{h}}^\alpha)^*$ is spanned by $\{\langle a^*, h_1(i_1) \dots h_k(i_k) - \rangle\}$, where $a^* \in (A^\alpha)^*$ and $i_1, \dots, i_k \in \mathbb{Z}_{\geq 1}$ and

$h_1, \dots, h_k \in H$. Let $u \in F^\vee$ and $v_1, v_2, v_3 \in F$ and $h_l \in H_l$ and $n \in \mathbb{Z}_{\geq 1}$. Then, by Lemma 5.3,

$$\begin{aligned} & S(u(h_l(n)-); v_1, v_2, v_3) \\ &= u(h_l(n)Y(v_1, \underline{z}_1)Y(v_2, \underline{z}_2)v_3) \\ &= S(u; v_1, v_2, h_l(n)v_3) + \sum_{i \geq 0} \binom{n}{i} \left(S(u; h_l(i)v_1, v_2, v_3)z_1^{n-i} + S(u; v_1, h_l(i)v_2, v_3)z_2^{n-i} \right). \end{aligned}$$

Since the above formula is symmetric for v_1 and v_2 , in order to show (FL), we may assume that $u = a^* \in A^\vee$. Similarly, for $n \in \mathbb{Z}_{\geq 0}$,

$$\begin{aligned} a^*(Y(h_l(-n-1)v_1, \underline{z}_1)Y(v_2, \underline{z}_2)v_3) &= \sum_{i \geq 0} \binom{-i-1}{n} \left(\frac{(-1)^n}{(z_1 - z_2)^{n+i+1}} \Big|_{|z_1| > |z_2|} a^*(Y(v_1, \underline{z}_1)Y(h_l(i)v_2, \underline{z}_2)v_3) \right. \\ &\quad \left. + \frac{1}{z_1^{n+i+1}} a^*(Y(v_1, \underline{z}_1)Y(v_2, \underline{z}_2)h_l(i)v_3) \right) \end{aligned}$$

and

$$\begin{aligned} a^*(Y(v_2, \underline{z}_2)Y(h_l(-n-1)v_1, \underline{z}_1)v_3) &= \sum_{i \geq 0} \binom{-i-1}{n} \left(\frac{(-1)^n}{(z_1 - z_2)^{n+i+1}} \Big|_{|z_2| > |z_1|} a^*(Y(h_l(i)v_2, \underline{z}_2)Y(v_1, \underline{z}_1)v_3) \right. \\ &\quad \left. + \frac{1}{z_1^{n+i+1}} a^*(Y(v_2, \underline{z}_2)Y(v_1, \underline{z}_1)h_l(i)v_3) \right), \end{aligned}$$

which implies that $a^*(Y(h_l(-n-1)v_1, \underline{z}_1)Y(v_2, \underline{z}_2)v_3)$ and $a^*(Y(v_2, \underline{z}_2)Y(h_l(-n-1)v_1, \underline{z}_1)v_3)$ are the expansion of the same function in the different region $|z_1| > |z_2|$ and $|z_2| > |z_1|$ if (FL) holds for $(a^*, v_1, h_l(i)v_2, v_3)$ and $(a^*, v_1, v_2, h_l(i)v_3)$ for any $i \geq 0$. In this way, we can inductively show the assumption (FL), by Lemma 5.4 and Lemma 5.3.

5.2. Troidal compactification of string theory. In this section, we construct a full vertex operator algebra associated with an even lattice. An even lattice is a free abelian group L of finite rank equipped with a bilinear map $(-, -) : L \times L \rightarrow \mathbb{Z}$ such that $(\alpha, \alpha) \in 2\mathbb{Z}$ for any $\alpha \in L$. Set $H_L = L \otimes_{\mathbb{Z}} \mathbb{R}$. Then, a bilinear form on H_L is induced by the bilinear form on L , denoted by $(-, -)$ again. An even lattice is called non-degenerate if $(H_L, (-, -))$ is non-degenerate. Let L be a non-degenerate even lattice. In [FLM], they construct an AH pair associated with an even lattice:

Proposition 5.2. [FLM] *There exists a unital associative algebra $\mathbb{C}[\hat{L}] = \bigoplus_{\alpha \in L} \mathbb{C}e_\alpha$ such that:*

- (1) e_0 is the unit;
- (2) $e_\alpha e_\beta \neq 0$ for any $\alpha, \beta \in L$;
- (3) $e_\alpha e_\beta = (-1)^{(\alpha, \beta)} e_\beta e_\alpha$ for any $\alpha, \beta \in L$.

In fact, a unital associative algebra satisfies the conditions in the above proposition is unique up to isomorphism as an AH pair ([FLM, Mo1]). The algebra $\mathbb{C}[\hat{L}]$ is called a twisted group algebra, which is clearly an even AH pair. For $p \in P(H_L)$, set $F_{L, H_L, p} = F_{\mathbb{C}[\hat{L}], H_L, p}$. By Proposition 5.1, we have:

Proposition 5.3. *For a non-degenerate even lattice L and $p \in P(H_L)$, $F_{L,H_L,p}$ is a full vertex algebra. Furthermore, if $p \in P_>(H_L)$, then $(F_{L,H_L,p}, Y, \mathbf{1}, \omega_{H_L}, \bar{\omega}_{H_L})$ is a self-dual QP-generated full vertex operator algebra.*

Proof. Let $p \in P_>(H_L)$ and set $F = F_{L,H_L,p}$. Since the spectrum of $F_{L,H_L,p}$ is $\{(p\alpha, p\alpha)/2 + n, -(\bar{p}\alpha, \bar{p}\alpha)/2 + m\}_{\alpha \in L, n, m \in \mathbb{Z}_{\geq 0}}$ and the bilinear form $(p-, p-)$ on H_l and $-(\bar{p}-, \bar{p}-)$ on H_r is positive definite, the spectrum is discrete, thus, bounded below. It is easy to show that $(F, Y, \mathbf{1}, \omega, \bar{\omega})$ is a simple full vertex operator algebra. To verify the assertion, it suffices to show that F satisfies the assumption of Corollary 3.2. By the positivity of the bilinear forms, (PN1) and (PN2) is clear. Let $0 \neq v \in F_{0,n}^\alpha$ for $\alpha \in L$. Then, $(p\alpha, p\alpha) = 0$ and, thus, $p\alpha = 0$, which implies that $Dv = 0$ and (PN4). Let $v \in F_{1,n}^\alpha$ for $\alpha \in L$. Suppose that $L(1)v \neq 0$. Then, $L(1)v \in F_{0,n}^\alpha$ implies that $p\alpha = 0$, that is, $\alpha \in H_r$. Thus, $F_{1,n}^\alpha$ is spanned by $\{h_l(-1)e_\alpha\}_{h_l \in H_l}$ and $(\alpha, h_l) = 0$. Then, for $h_l \in H_l$, $L(1)h_l(-1)e_\alpha = h_l(0)e_\alpha = (h_l, \alpha)e_\alpha = 0$, a contradiction, which implies (PN3). Hence, the assertion holds. $\square$

Let (n, m) be the signature of H_L . Then, the orthogonal group $O(H_L) \cong O(n, m; \mathbb{R})$ acts on $P_>(H_L)$ by $g \cdot p = g \circ p \circ g^{-1}$ for $p \in P_>(H_L)$ and $g \in O(H_L)$. By the elementary calculus in linear algebra, we can show that the action is transitive. The stabilizer subgroup of p is $O(n; \mathbb{R}) \times O(m; \mathbb{R})$. Thus, $P_>(H_L)$ is homeomorphic to the Grassmannian $O(n, m; \mathbb{R})/O(n; \mathbb{R}) \times O(m; \mathbb{R})$. Hence, we construct a family of full vertex operator algebras which is parametrized by the Grassmannian $O(n, m; \mathbb{R})/O(n; \mathbb{R}) \times O(m; \mathbb{R})$.

Let $\Pi_{1,1}$ be the unique even unimodular lattice with the signature $(1, 1)$, more explicitly, $\Pi_{1,1} = \mathbb{Z}z \oplus \mathbb{Z}w$ and $(z, z) = (w, w) = 0$ and $(z, w) = -1$ and for $N \in \mathbb{Z}_>$, set $\Pi_{N,N} = \Pi_{1,1}^N$. Then, we have a family of full vertex operator algebras of the central charge (N, N) parametrized by $O(N, N; \mathbb{R})/O(N; \mathbb{R}) \times O(N; \mathbb{R})$.

Remark 5.1. *Let $\text{Aut } \Pi_{N,N}$ be the lattice automorphism group of $\Pi_{N,N}$, which is isomorphic to $O(N, N; \mathbb{Z})$, the $\mathbb{Z}$ -value points of the algebraic group $O(N, N)$. Then, we can prove that if $\sigma \in \text{Aut } \Pi_{N,N}$ then $F_{\Pi_{N,N}, H_{\Pi_{N,N}}, p}$ is isomorphic to $F_{\Pi_{N,N}, H_{\Pi_{N,N}}, \sigma p \sigma^{-1}}$ as a full vertex operator algebra. Thus, the true parameter space is*

$$O(N, N; \mathbb{Z}) \backslash O(N, N; \mathbb{R}) / O(N; \mathbb{R}) \times O(N; \mathbb{R}).$$

This statement is proved in [Mo2] in more general setting. We also remark that this conformal field theory and parameter space appear in the toroidal compactification of string theory [Polc1].

5.3. Commutative algebra object in H-Vect. Let H be a finite-dimensional real vector space with a non-degenerate symmetric bilinear form

$$(-, -) : H \times H \rightarrow \mathbb{R}.$$

Let H-Vect be the category whose object is an H -graded vector space over $\mathbb{C}$ and morphism is a grading preserving $\mathbb{C}$ -linear map. Let $V = \bigoplus_{\alpha \in H} V^\alpha$ and $W = \bigoplus_{\beta \in H} W^\beta$ be

H -graded vector spaces. A tensor product of H -graded vector spaces are defined by the tensor product of vector spaces with the H -grading:

$$(V \otimes W)^\alpha = \bigoplus_{\alpha' \in H} V^{\alpha'} \otimes W^{\alpha - \alpha'}.$$

Then, $\underline{H}\text{-Vect}$ is a strict monoidal category whose unit is $\mathbb{C} = \mathbb{C}^0$. Define a braiding $B_{V,W} : V \otimes W \rightarrow W \otimes V$ by $v_\alpha \otimes w_\beta \mapsto \exp((\alpha, \beta)\pi i) w_\beta \otimes v_\alpha$ for $\alpha, \beta \in H$, $v_\alpha \in V^\alpha$ and $w_\beta \in W^\beta$. Then, it is easy to show that B satisfies the hexagon identity, thus $(\underline{H}\text{-Vect}, B)$ is a braided tensor category. Let A be a unital commutative associative algebra object in $\underline{H}\text{-Vect}$ and $a \in A^\alpha$ and $b \in A^\beta$ for $\alpha, \beta \in H$. If $ab \neq 0$, then, by the commutativity, $ab = \exp((\alpha, \beta)\pi i)ba$ and $ba = \exp((\alpha, \beta)\pi i)ab$. Thus, $(\alpha, \beta) \in \mathbb{Z}$ and $ab = (-1)^{(\alpha, \beta)}ba$, which implies that A is an AH pair. Thus, we have:


Proposition 5.4. *A unital commutative associative algebra object in $\underline{H}\text{-Vect}$ is an AH pair, and vice versa.*

6. APPENDIX

6.1. Binary tree and expansions. Let F be a self-dual full vertex operator algebra and denote $Y(a, \underline{x})$ by $a(x)$ for $a \in F$.

For $n \in \mathbb{Z}_{>0}$, let P_n be the set of parenthesized products of n elements $1, 2, \dots, n$, e.g., $((((31)6)(24))(57))$.

We can naturally associate a tree and a parenthesized correlation function (see Section 6.1) to each element in P_n . For example, an element $((((31)6)(24))(57)) \in P_7$ defines the following binary tree:



and the parenthesized correlation function:

$$S_{((((31)6)(24))(57))}(a_1, a_2, a_3, a_4, a_5, a_6, a_7) := \left(1, \left[\left(\left(a_3(x_3)a_1\right)(x_1)a_6\right)(x_6)a_2(x_2)a_4\right](x_4)a_5(x_5)a_7\right), \right.$$

where $a_1, \dots, a_7 \in F$.

We conjecture that $S_{((((31)6)(24))(57))}(a_1, a_2, a_3, a_4, a_5, a_6, a_7)$ is the expansion of a real analytic function $\phi(z_1, \dots, z_7)$ on X_7 in the domain

$$\{(x_1, \dots, x_7) \in \mathbb{C}^7 \mid \{|x_4| > |x_6| > |x_1| > |x_3|\} \cap \{|x_6| > |x_2|\} \cap \{|x_4| > |x_5|\}\}$$

after the change of variables:

$$(x_1, x_2, x_3, x_4, x_5, x_6) = (z_1 - z_6, z_2 - z_4, z_3 - z_1, z_4 - z_7, z_5 - z_7, z_6 - z_4).$$

This conjecture is proved in the case where $n \leq 4$ or $n = 5$ and the right most state is the vacuum state (Theorem 3.2). Here, we give the rule for the change of variables and the convergence region by using trees.

We remark that the tree associated with $A \in P_n$ is a full binary tree, i.e., every node has either 0 or 2 children. A node with no children is called a leaf. We will inductively assign a number to each edge of the tree and a formal variable to a node of the tree. First, the number i is assigned to the unique edge connected to leaf labeled by i . On each node, the number of the above edge is equal to the number of the right below edge, which determine the numbers of the all edges (Fig.1).





Fig.1 Fig.2

Then, we assigned the formal variables $z_i - z_j$ on the node whose left below edge is labeled by i and right below edge is labeled by j and change the variable x_i into $z_i - z_j$ (Fig.2).

For example, for $((((31)6)(24))(57)) \in P_7$, we have:



Thus, the rule for the change of variables are determined.

We now explain how to get the convergence domain. If a node labeled by $z_i - z_j$ is a child of $z_k - z_l$, then $|z_k - z_l| > |z_i - z_j|$ or in the other coordinate $|x_k| > |x_l|$.

We also consider the special case that the right most state is the vacuum state. Let Q_n be the set of parenthesized products of $n + 1$ elements $1, 2, \dots, n, \star$ with $\star$ at the right most, e.g., $2(3(1(4\star))) \in Q_4$. Then, the parenthesized correlation function associated with $A \in Q_n$ is defined by $S_A(a_1, \dots, a_n, \mathbf{1})$, that is, we insert the vacuum state at $\star$. We denote it by $S_A(a_1, \dots, a_n)$ again. For example, for $((((31)6)(24))(5\star)) \in Q_6$ and $a_1, \dots, a_6 \in F$, we have

$$S_{((((31)6)(24))(5\star))}(a_1, a_2, a_3, a_4, a_5, a_6) := \left(\mathbf{1}, \left[\left(\left(a_3(x_3)a_1 \right)(x_1)a_6 \right)(x_6)a_2(x_2)a_4 \right](x_4)a_5(x_5)\mathbf{1} \right),$$


The rule for the change of variable is given by the substitution of $z_\star = 0$ into the rule for P_{n+1} . For $((((31)6)(24))(5\star)) \in P_7$, the rule for the change of variables is:

$$(x_1, x_2, x_3, x_4, x_5, x_6) = (z_1 - z_6, z_2 - z_4, z_3 - z_1, z_4 - z_\star, z_5 - z_\star, z_6 - z_4).$$

Then, the rule for the change of variables for $((((31)6)(24))(5\star)) \in Q_6$ is given by

$$(x_1, x_2, x_3, x_4, x_5, x_6) = (z_1 - z_6, z_2 - z_4, z_3 - z_1, z_4, z_5, z_6 - z_4).$$

We list the trees and the rules for the change of variables for all the standard elements in P_4 :



REFERENCES

- [B] R. Borcherds, Vertex algebras, Kac-Moody algebras, and the Monster, Proc. Nat. Acad. Sci. U.S.A., **83**, 1986, (10), 3068–3071.
- [BPZ] A. A. Belavin, A. M. Polyakov, A. B. Zamolodchikov, Infinite conformal symmetry in two-dimensional quantum field theory, Nuclear Phys. B, **241**, 1984, (2), 333–380.
- [FB] E. Frenkel, D. Ben-Zvi, Vertex algebras and algebraic curves, Mathematical Surveys and Monographs, **88**, Second, American Mathematical Society, Providence, RI, 2004.
- [FGG] S. Ferrara and A.F. Grillo and R. Gatto, Tensor representations of conformal algebra and conformally covariant operator product expansion, Ann. Physics, **76**, 1973, 161–188.
- [FHL] I. Frenkel, Y. Huang and J. Lepowsky, On axiomatic approaches to vertex operator algebras and modules, Mem. Amer. Math. Soc., **104**, 1993, (494).
- [FLM] I. Frenkel, J. Lepowsky, and A. Meurman, Vertex operator algebras and the Monster, Pure and Applied Mathematics, **134**, Academic Press, Inc., Boston, MA, 1988.
- [FMS] P. Di Francesco, P. Mathieu and D. Sénéchal, Conformal field theory, Graduate Texts in Contemporary Physics, Springer-Verlag, New York, 1997.
- [FRS] J. Fuchs, I. Runkel and C. Schweigert, Conformal correlation functions, Frobenius algebras and triangulations, Nucl. Phys. **624** 2002, 452–468.
- [G] P. Goddard, Meromorphic conformal field theory, Infinite-dimensional Lie algebras and groups (Luminy-Marseille, 1988), Adv. Ser. Math. Phys., **7**, 556–587, 1989.
- [H1] Y.-Z. Huang, A theory of tensor products for module categories for a vertex operator algebra, IV, J. Pure Appl. Alg., **100**, (1995), 173–216.
- [H2] Y.-Z. Huang, Virasoro vertex operator algebras, (nonmeromorphic) operator product expansion and the tensor product theory, J. Alg., **182**, (1996), 201–234.
- [H3] Y.-Z. Huang, Vertex operator algebras and the Verlinde conjecture, Comm. Contemp. Math., **10**, 2008, 103–154.
- [H4] Y.-Z. Huang, Rigidity and modularity of vertex tensor categories, Comm. Contemp. Math. **10**, 2008, 871–911.
- [HK] Y.-Z. Huang, L. Kong, Full field algebras, Comm. Math. Phys., **272**, 2007, 2, 345–396.

- [HL1] Y.-Z. Huang and J. Lepowsky, A theory of tensor products for module categories for a vertex operator algebra, I, *Selecta Mathematica (New Series)*, **1**, 1995, 699–756.
- [HL2] Y.-Z. Huang and J. Lepowsky, A theory of tensor products for module categories for a vertex operator algebra, II, *Selecta Mathematica (New Series)*, **1**, 1995, 757–786.
- [HL3] Y.-Z. Huang and J. Lepowsky, A theory of tensor products for module categories for a vertex operator algebra, III, *J. Pure Appl. Alg.*, **100**, 1995, 141–171.
- [IZ] C. Itzykson and J.B. Zuber, *Quantum field theory*, International Series in Pure and Applied Physics, McGraw-Hill International Book Co., New York, 1980.
- [K] V. G. Kac, *Vertex algebras for beginners*, University lecture series, **10**, A.M.S. 1996.
- [L1] H. Li, Symmetric invariant bilinear forms on vertex operator algebras, *J. Pure Appl. Algebra*, **96**, 1994, (3), 279–297.
- [LL] J. Lepowsky and H. Li, *Introduction to vertex operator algebras and their representations*, Progress in Mathematics, **227**, Birkhäuser Boston, Inc., Boston, MA, 2004.
- [Ma] J. Maldacena, The large N limit of superconformal field theories and supergravity, *Adv. Theor. Math. Phys.*, **2**, 1998, 2, 231–252.
- [Mo1] Y. Moriwaki, Genus of vertex algebras and mass formula, arXiv:2004.01441 [q-alg].
- [Mo2] Y. Moriwaki, Full vertex algebra and all order current-current deformation of 2d CFT, preprint.
- [MS1] G. Moore and N. Seiberg, Polynomial equations for rational conformal field theories, *Phys. Lett.* **212**, 1988, 451–460.
- [MS2] G. Moore and N. Seiberg, Classical and quantum conformal field theory, *Comm. Math. Phys.* **123**, 1989, 177–254.
- [Polc1] J. Polchinski, *String theory. Vol. I*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, Cambridge, 1998.
- [Poly2] A. M. Polyakov, Non-Hamiltonian approach to conformal quantum field theory, *Ž. Èksper. Teoret. Fiz.*, **66**, 1974, 23–42.
- [S] M. Srednicki, *Quantum field theory*, Cambridge University Press, Cambridge, 2010.
- [W] E. Witten, Quantum field theory and the Jones polynomial, *Comm. Math. Phys.*, **121**, 1989, (3), 351–399.
- [Y] M. Yoshida, *Hypergeometric functions, my love*, Aspects of Mathematics, E32, Friedr. Vieweg & Sohn, Braunschweig, 1997.