

Near-Optimal SQ Lower Bounds for Agnostically Learning Halfspaces and ReLUs under Gaussian Marginals

Ilias Diakonikolas*
University of Wisconsin-Madison
iliask@cs.wisc.edu

Daniel M. Kane†
University of California, San Diego
dakane@cs.ucsd.edu

Nikos Zarifis‡
University of Wisconsin-Madison
zarifis@wisc.edu

June 30, 2020

Abstract

We study the fundamental problems of agnostically learning halfspaces and ReLUs under Gaussian marginals. In the former problem, given labeled examples (\mathbf{x}, y) from an unknown distribution on $\mathbb{R}^d \times \{\pm 1\}$, whose marginal distribution on \mathbf{x} is the standard Gaussian and the labels y can be arbitrary, the goal is to output a hypothesis with 0-1 loss $\text{OPT} + \epsilon$, where OPT is the 0-1 loss of the best-fitting halfspace. In the latter problem, given labeled examples (\mathbf{x}, y) from an unknown distribution on $\mathbb{R}^d \times \mathbb{R}$, whose marginal distribution on \mathbf{x} is the standard Gaussian and the labels y can be arbitrary, the goal is to output a hypothesis with square loss $\text{OPT} + \epsilon$, where OPT is the square loss of the best-fitting ReLU. We prove Statistical Query (SQ) lower bounds of $d^{\text{poly}(1/\epsilon)}$ for both of these problems. Our SQ lower bounds provide strong evidence that current upper bounds for these tasks are essentially best possible.

*Supported by NSF Award CCF-1652862 (CAREER), a Sloan Research Fellowship, and a DARPA Learning with Less Labels (LwLL) grant.

†Supported by NSF Award CCF-1553288 (CAREER) and a Sloan Research Fellowship.

‡Supported in part by a DARPA Learning with Less Labels (LwLL) grant.

1 Introduction

1.1 Background and Problem Motivation

We study the fundamental problems of agnostically learning halfspaces and ReLU regression in the distribution-specific agnostic PAC model. In both of these problems, we are given i.i.d. samples from a joint distribution \mathcal{D} on labeled examples (\mathbf{x}, y) , where $\mathbf{x} \in \mathbb{R}^d$ is the example and $y \in \mathbb{R}$ is the corresponding label, and the goal is to compute a hypothesis that is competitive with the best-fitting halfspace (with respect to the 0-1 loss) or ReLU (with respect to the square loss) respectively.

A halfspace (or Linear Threshold Function) is any Boolean function $f : \mathbb{R}^d \rightarrow \{\pm 1\}$ of the form $f(\mathbf{x}) = \text{sign}(\langle \mathbf{w}, \mathbf{x} \rangle + \theta)$, where $\mathbf{w} \in \mathbb{R}^d$ is called the weight vector and θ is called the threshold. (The function $\text{sign} : \mathbb{R} \rightarrow \{\pm 1\}$ is defined as $\text{sign}(u) = 1$ if $u \geq 0$ and $\text{sign}(u) = -1$ otherwise.) The task of learning an unknown halfspace from samples is one of the oldest and most well-studied problems in machine learning, starting with the Perceptron algorithm [Ros58] and leading to influential techniques, including SVMs [Vap98] and AdaBoost [FS97]. In the realizable setting [Val84], this learning problem amounts to linear programming and can be solved in polynomial time (see, e.g., [MT94]) without distributional assumptions. In contrast, in the distribution-independent agnostic model [Hau92, KSS94], even *weak* learning is computationally hard [GR06, FGKP06, Dan16].

A line of research [KKMS08, KLS09, ABL17, Dan15, DKS18] has focused on learning halfspaces in the *distribution-specific* agnostic PAC model, where it is assumed that the marginal distribution on the examples is well-behaved. In this paper, we study the important case that the marginal distribution is the standard Gaussian. For concreteness, we formally define this problem.

Problem 1.1 (Agnostically Learning Halfspaces with Gaussian Marginals). *Let \mathcal{C}_{LTF} be the class of halfspaces on \mathbb{R}^d . Given i.i.d. samples (\mathbf{x}, y) from a distribution \mathcal{D} on $\mathbb{R}^d \times \{\pm 1\}$, where the marginal $\mathcal{D}_{\mathbf{x}}$ on \mathbb{R}^d is the standard Gaussian $\mathcal{N}(\mathbf{0}, \mathbf{I})$ and no assumptions are made on the labels y , the goal of the learning algorithm is to output a hypothesis $h : \mathbb{R}^d \rightarrow \{\pm 1\}$ such that with high probability we have $\Pr_{(\mathbf{x}, y) \sim \mathcal{D}}[h(\mathbf{x}) \neq y] \leq \text{OPT} + \epsilon$, where $\text{OPT} = \inf_{f \in \mathcal{C}_{\text{LTF}}} \Pr_{(\mathbf{x}, y) \sim \mathcal{D}}[f(\mathbf{x}) \neq y]$.*

The L_1 -regression algorithm of [KKMS08] solves Problem 1.1 with sample complexity and running time $d^{O(1/\epsilon^2)}$ [DGJ⁺10, DKN10]. This algorithm is also known to succeed for all log-concave distributions and for certain discrete distributions. A related line of work [ABL17, Dan15, DKS18] has given $\text{poly}(d/\epsilon)$ time algorithms with weaker guarantees, specifically with misclassification error $C \cdot \text{OPT} + \epsilon$, for a universal constant $C > 1$. The fastest known algorithm with optimal error is the one from [KKMS08].

A Rectified Linear Unit (ReLU) is any real-valued function $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$ of the form $f(\mathbf{x}) = \text{ReLU}(\langle \mathbf{w}, \mathbf{x} \rangle + \theta)$, where $\mathbf{w} \in \mathbb{R}^d$ is called the weight vector and θ is called the threshold. (The function $\text{ReLU} : \mathbb{R} \rightarrow \mathbb{R}_+$ is defined as $\text{ReLU}(u) = \max\{0, u\}$.) ReLUs are the most commonly used activation functions in modern deep neural networks. Finding the best-fitting ReLU with respect to square-loss is a fundamental primitive in the theory of neural networks. A number of recent works have studied this problem both in terms of finding efficient algorithms and proving hardness results [Sol17, GKK17, MR18, GKK19, DGK⁺20]. Similarly to the case of halfspaces, the problem is efficiently solvable in the realizable setting and computationally hard in the distribution-independent agnostic setting [MR18]. Here we study the case of Gaussian marginals, which we now define.

Problem 1.2 (ReLU Regression with Gaussian Marginals). *Let $\mathcal{C}_{\text{ReLU}}$ be the class of ReLUs on \mathbb{R}^d . Given i.i.d. samples (\mathbf{x}, y) from a distribution \mathcal{D} on $\mathbb{R}^d \times \mathbb{R}$, where the marginal $\mathcal{D}_{\mathbf{x}}$ on \mathbb{R}^d is the standard Gaussian $\mathcal{N}(\mathbf{0}, \mathbf{I})$ and no assumptions are made on the labels y , the goal of the*

learning algorithm is to output a hypothesis $h : \mathbb{R}^d \rightarrow \mathbb{R}$ such that with high probability we have $\mathbf{E}_{(\mathbf{x}, y) \sim \mathcal{D}}[(h(\mathbf{x}) - y)^2] \leq \text{OPT} + \epsilon$, where $\text{OPT} = \inf_{f \in \mathcal{C}_{\text{ReLU}}} \mathbf{E}_{(\mathbf{x}, y) \sim \mathcal{D}}[(f(\mathbf{x}) - y)^2]$.

Recent work [DGK⁺20] gave an algorithm for Problem 1.2 with sample complexity and runtime $d^{\text{poly}(1/\epsilon)}$. While $\text{poly}(d/\epsilon)$ time algorithms are known with weaker guarantees [GKK19, DGK⁺20], the fastest known algorithm with $\text{OPT} + \epsilon$ error is the one of [DGK⁺20].

In terms of computational hardness, prior work has given evidence that no $\text{poly}(d, 1/\epsilon)$ time algorithm exists for either problem. Specifically, [KK14] gave a reduction from the problem of learning sparse parities with noise to Problem 1.1. Based on the presumed computational hardness of the former problem, this reduction implies a computational lower bound of $d^{\Omega(\log(1/\epsilon))}$ for Problem 1.1. More recently, [GKK19] gave a qualitatively similar reduction implying a computational lower bound of $d^{\Omega(\log(1/\epsilon))}$ for Problem 1.2. Interestingly, both of these lower bounds cannot be improved in the sense that the corresponding hard instances can be solved in time $d^{O(\log(1/\epsilon))}$.

In summary, the best known algorithms for Problems 1.1 and 1.2 have running time $d^{\text{poly}(1/\epsilon)}$, while the best known computational hardness results give $d^{\Omega(\log(1/\epsilon))}$ lower bounds. This raises the following natural question:

What is the precise complexity of Problems 1.1 and 1.2?

Given the lower bounds of [KK14, GKK19], it is conceivable that there exist algorithms for these problems running in time $d^{\text{polylog}(1/\epsilon)}$, i.e., quasi-polynomial in $1/\epsilon$. A positive result of this form would represent significant algorithmic progress in the theory of PAC learning.

In this paper, we show that the latter possibility is unlikely. Specifically, we prove Statistical Query (SQ) lower bounds of $d^{\text{poly}(1/\epsilon)}$ for both Problems 1.1 and 1.2. Our SQ lower bounds provide evidence that known algorithms for these problems are essentially best possible.

Before we state our contributions in detail, we give some background on Statistical Query (SQ) algorithms. SQ algorithms are a broad class of algorithms that are only allowed to query expectations of bounded functions of the distribution rather than directly access samples. The SQ model was introduced by Kearns [Kea98] in the context of supervised learning as a natural restriction of the PAC model [Val84]. Subsequently, the SQ model has been extensively studied in a plethora of contexts (see, e.g., [Fel16] and references therein).

Formally, an SQ algorithm has access to the following oracle.

Definition 1.3 (STAT Oracle). Let \mathcal{D} be a distribution over some domain X and let $f : X \rightarrow [-1, 1]$. A statistical query is a function $q : X \times [-1, 1] \rightarrow [-1, 1]$. We define $\text{STAT}(\tau)$ to be the oracle that given a query $q(\cdot, \cdot)$ outputs a value v such that $|v - \mathbf{E}_{\mathbf{x} \sim \mathcal{D}}[q(\mathbf{x}, f(\mathbf{x}))]| \leq \tau$, where $\tau > 0$ is the tolerance parameter of the query.

We note that the class of SQ algorithms is rather general and captures most of the known supervised learning algorithms. More broadly, a wide range of known algorithmic techniques in machine learning are known to be implementable using SQs. These include spectral techniques, moment and tensor methods, local search (e.g., Expectation Maximization), and many others (see, e.g., [FGR⁺13, FGV17]). For the supervised learning problems studied in this paper, all known algorithms with non-trivial performance guarantees are SQ or are easily implementable using SQs.

One can prove lower bounds on the complexity of SQ algorithms via the notion of *Statistical Query (SQ) dimension* [BFJ⁺94, FGR⁺13]. A lower bound on the SQ dimension of a learning problem provides an unconditional lower bound on the complexity of any SQ algorithm for the problem.

1.2 Our Results and Techniques

We are now ready to formally state our main results. For Problem 1.1 we prove:

Theorem 1.4. *Let $d \geq 1$ and $\epsilon \geq d^{-c}$, for some sufficiently small constant $c > 0$. Any SQ algorithm that agnostically learns halfspaces on \mathbb{R}^d under Gaussian marginals within additive error $\epsilon > 0$ requires at least $d^{c/\epsilon}$ many statistical queries to $\text{STAT}(d^{-c/\epsilon})$.*

Intuitively, the above statement says that any SQ algorithm for Problem 1.1 requires time at least $d^{\Omega(1/\epsilon)}$. This comes close to the known upper bound of $d^{O(1/\epsilon^2)}$ [KKMS08] and exponentially improves on the best known lower bound of $d^{\Omega(\log(1/\epsilon))}$ [KK14].

For Problem 1.2 we prove:

Theorem 1.5. *There exist constants $c, c' > 0$ such that the following holds: For $d \geq 1$ and $\epsilon \geq d^{-c}$, any SQ algorithm with excess error ϵ for ReLU regression on \mathbb{R}^d under Gaussian marginals requires at least $d^{c/\epsilon^{c'}}$ many statistical queries to $\text{STAT}(d^{-c/\epsilon^{c'}})$.*

Intuitively, the above statement says that any SQ algorithm for Problem 1.2 requires time at least $d^{(1/\epsilon)^{\Omega(1)}}$. This qualitatively matches the upper bound of $d^{\text{poly}(1/\epsilon)}$ [DGK⁺20], up to the degree of the polynomial, and exponentially improves on the best known lower bound of $d^{\Omega(\log(1/\epsilon))}$ [GKK19].

Discussion. The reduction-based hardness of [KK14, GKK19] imply SQ lower bounds of $d^{\Omega(\log(1/\epsilon))}$ for both problems. Our new SQ lower bounds are qualitatively optimal, nearly matching current algorithms. Interestingly, for both problems, our results show a sharp separation in the complexity of obtaining error $O(\text{OPT}) + \epsilon$ (which is $\text{poly}(d/\epsilon)$) versus optimal error $\text{OPT} + \epsilon$. In particular, our lower bounds suggest that the accuracy-runtime tradeoff of known polynomial time approximation schemes (PTAS) for these problems [Dan15, DGK⁺20] that achieve error $(1 + \gamma)\text{OPT} + \epsilon$, for all $\gamma > 0$, in time $\text{poly}(d^{\text{poly}(1/\gamma)}, 1/\epsilon)$ is qualitatively best possible.

Technical Overview The starting point for our SQ lower bound constructions is the framework of [DKS17]. This work establishes the following: Suppose we have a one-dimensional distribution A that matches its first k moments with the standard one-dimensional Gaussian. Consider the set of distributions $\{\mathbf{P}_{\mathbf{v}}\}$, where \mathbf{v} is any unit vector, such that the projection of $\mathbf{P}_{\mathbf{v}}$ in the \mathbf{v} -direction is equal to A and in the orthogonal complement $\mathbf{P}_{\mathbf{v}}$ is an independent standard Gaussian. Then this set of distributions has SQ dimension $d^{\Omega(k)}$. By known results (see, e.g., [Fel16]) this implies that distinguishing such a distribution from the standard Gaussian or learning a distribution with better than $1/\text{poly}(d^k)$ correlation with such a distribution is hard in the SQ model.

To leverage the aforementioned result, a first hurdle that must be overcome is adapting the results of [DKS17] – which apply to the unsupervised task of learning distributions – to the supervised task of learning functions. For Boolean functions $F : \mathbb{R}^d \rightarrow \{\pm 1\}$, this is relatively straightforward. Essentially, sampling from the distribution $(\mathbf{x}, F(\mathbf{x}))$ is equivalent to sampling from the conditional distributions of \mathbf{x} conditioned on $F(\mathbf{x}) = 1$ and on $F(\mathbf{x}) = -1$ in a way that is easy to make rigorous in the SQ model. To apply the techniques from [DKS17], we want to construct a one-dimensional Boolean-valued function $f : \mathbb{R} \rightarrow \{\pm 1\}$ such that the conditional distributions match moments, or more conveniently so that $\mathbf{E}_{z \sim \mathcal{N}(0,1)}[f(z) z^i] = 0$ for all $0 \leq i \leq k$. Given such a moment matching function, it is not hard to show that the conditional distributions corresponding to the function $F_{\mathbf{v}}(\mathbf{x}) = f(\langle \mathbf{x}, \mathbf{v} \rangle)$, for a unit vector \mathbf{v} , are of the type covered by [DKS17].

Of course, for such a construction to have any implications for the problems of agnostically learning halfspaces/ReLUs, we require an additional key property: We want to find a univariate

Boolean-valued function f that not only has this kind of matching moments property, but also *correlates non-trivially* with a halfspace or ReLU. If we can guarantee non-trivial correlation, agnostically learning the function $F_{\mathbf{v}}$ with respect to this class will require having a *weak learner* for F (and, for example, preventing the learning from just outputting the identically zero function).

To achieve both aforementioned goals, we use analytic tools to show that there exists an $O(k)$ -piecewise constant Boolean-valued function f with k matching moments. Such a function has correlation $\Omega(1/k)$ with some halfspace. For the case of ReLUs, we use an analysis making use of Legendre polynomials to find a function $f' : \mathbb{R} \rightarrow [-1, 1]$ with vanishing first k moments and non-trivial correlation with a ReLU. We then show that this can be rounded to a Boolean-valued function with the same guarantees.

Concurrent Work Concurrent and independent work [GGK20] established qualitatively similar SQ lower bounds for agnostically learning halfspaces and ReLUs. Their techniques are different than ours, building on prior SQ lower bounds for learning depth-2 neural networks [DKKZ20, GGJ⁺20], and employing a reduction-based approach.

2 Preliminaries

Notation For $n \in \mathbb{Z}_+$, we denote $[n] \stackrel{\text{def}}{=} \{1, \dots, n\}$ and $\overline{\mathbb{R}} \stackrel{\text{def}}{=} \mathbb{R} \cup \{\pm\infty\}$. For $\mathbf{x} \in \mathbb{R}^d$, and $i \in [d]$, \mathbf{x}_i denotes the i -th coordinate of \mathbf{x} . We will use $\langle \mathbf{x}, \mathbf{y} \rangle$ for the inner product between $\mathbf{x}, \mathbf{y} \in \mathbb{R}^d$. We will use $\mathbf{E}[X]$ for the expectation of random variable X and $\Pr[\mathcal{E}]$ for the probability of event \mathcal{E} .

Let \mathbf{e}_i be the i -th standard basis vector in \mathbb{R}^d . Let $\mathcal{N}(0, 1)$ denote the standard univariate Gaussian distribution and $\mathcal{N}(\mathbf{0}, \mathbf{I})$ denote the standard multivariate Gaussian distribution. We will use ϕ to denote the pdf of the standard Gaussian.

Correlation and Statistical Query Dimension To bound the complexity of SQ learning a concept class \mathcal{C} , we will use the standard notion of Statistical Query Dimension [BFJ⁺94].

For $f, g : \mathbb{R}^d \rightarrow \mathbb{R}$, we define the correlation between f and g under the distribution \mathcal{D} to be the expectation $\mathbf{E}_{\mathbf{x} \sim \mathcal{D}}[f(\mathbf{x})g(\mathbf{x})]$. To prove that the SQ dimension of \mathcal{C} under the distribution \mathcal{D} is large, we need to find a set of functions in the class that are nearly uncorrelated.

Definition 2.1 (Statistical Query Dimension). For a class of functions \mathcal{C} and distribution \mathcal{D} , $\text{SQ-DIM}(\mathcal{C}, \mathcal{D}) = s$, if s is the largest integer value for which there exist s functions $f_1, f_2, \dots, f_s \in \mathcal{C}$ such that for every $i \neq j$, it holds $|\mathbf{E}_{\mathbf{x} \sim \mathcal{D}}[f_i(\mathbf{x})f_j(\mathbf{x})]| \leq 1/s$.

Our SQ lower bounds will use the following lemma (see, e.g., Theorem 2 of [Szö09]).

Lemma 2.2. *Let \mathcal{C} be a concept class of functions on \mathbb{R}^d and \mathcal{D} be a distribution on \mathbb{R}^d . Let $s = \text{SQ-DIM}(\mathcal{C}, \mathcal{D})$. Any SQ algorithm that outputs a hypothesis with correlation at least $1/s^{1/3}$ from an unknown function in \mathcal{C} requires at least $s^{1/3}/2 - 1$ queries to $\text{STAT}(1/s^{1/3})$.*

We note that the above theorem was initially shown for Boolean-valued functions, but also holds for real-valued functions of bounded norm (see, e.g., [Fel09]).

3 SQ Lower Bound for Agnostically Learning Halfspaces

In this section, we prove Theorem 1.4. To do so, we construct a family \mathcal{F}_k of k -decision lists of halfspaces on \mathbb{R}^d satisfying the following properties: (1) Any SQ algorithm that weakly learns

\mathcal{F}_k requires many high accuracy SQ queries. (2) Each $F \in \mathcal{F}_k$ is non-trivially correlated with a halfspace. Formally, we establish the following statement.

Proposition 3.1. *Assuming d is at least a sufficiently large power of k , there exists a set \mathcal{F}_k of k -decision lists of halfspaces on \mathbb{R}^d such that any SQ algorithm that learns \mathcal{F}_k within 0-1 error $\leq 1/2 - d^{-\Omega(k)}$ with respect to $\mathcal{N}(\mathbf{0}, \mathbf{I})$ needs $d^{\Omega(k)}$ queries to $\text{STAT}(d^{-\Omega(k)})$. Moreover, for any $F \in \mathcal{F}_k$, there is a halfspace σ such that $\mathbf{E}_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})}[F(\mathbf{x})\sigma(\mathbf{x})] \geq 1/(2k)$.*

Given the above statement, Theorem 1.4 follows.

Proof of Theorem 1.4. Let \mathcal{A} be an agnostic SQ learner for halfspaces under Gaussian marginals. We use \mathcal{A} to weakly learn \mathcal{F}_k , for a value of k to be determined. That is, we feed \mathcal{A} a set of i.i.d. labeled examples from an arbitrary function $F \in \mathcal{F}_k$. By definition, algorithm \mathcal{A} computes a hypothesis $h : \mathbb{R}^d \rightarrow \{\pm 1\}$ such that $\Pr_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})}[h(\mathbf{x}) \neq F(\mathbf{x})] \leq \text{OPT} + \epsilon$, for $\epsilon > 0$. By the second statement of Proposition 3.1, it follows that $\text{OPT} \leq 1/2 - 1/(2k)$. Thus, we have that

$$\Pr_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})}[h(\mathbf{x}) \neq F(\mathbf{x})] \leq 1/2 - 1/(2k) + \epsilon.$$

For $\epsilon = 1/(4k)$, Proposition 3.1 gives that \mathcal{A} needs at least $d^{\Omega(1/\epsilon)}$ queries to $\text{STAT}(d^{-\Omega(1/\epsilon)})$. This completes the proof. \square

3.1 Proof of Proposition 3.1

The main idea for our construction of a hard-to-learn family of functions \mathcal{F}_k is the following: We first establish the existence of a one-dimensional Boolean-valued function $f : \mathbb{R} \rightarrow \{\pm 1\}$ whose first k moments match the first k moments of the standard univariate Gaussian distribution (Proposition 3.2). Importantly, this function f is $(k + 1)$ -piecewise constant, i.e., there exists a partition of its domain into $k + 1$ intervals I_1, \dots, I_k such that f is constant within each I_j . The k points $z_1, z_2, \dots, z_k \in \mathbb{R}$ where the function changes value are called breakpoints.

Given our univariate construction, we construct our family of d -dimensional functions by using a copy of this one-dimensional function f oriented in a random direction. More specifically, let S be a set of $2^{d^{\Omega(1)}}$ nearly orthogonal unit vectors on \mathbb{R}^d (Lemma 3.4). Then we define the family $\mathcal{F}_k = \{F_{\mathbf{v}}(\mathbf{x})\}_{\mathbf{v} \in S}$, where $F_{\mathbf{v}}(\mathbf{x}) \stackrel{\text{def}}{=} f(\langle \mathbf{v}, \mathbf{x} \rangle)$ for the univariate function f from Proposition 3.2. Since f is k -piecewise constant, each $F \in \mathcal{F}_k$ is a k -decision list. From this, it follows that each such F is non-trivially correlated with a halfspace (Lemma 3.7).

Figure 1 shows how the function $F_{\mathbf{v}}$ labels the examples in a 2-dimensional projection.




Figure 1: The “red” region is the set of points where $F_{\mathbf{v}}(\mathbf{x}) = -1$ and the “blue” region where $F_{\mathbf{v}}(\mathbf{x}) = 1$.

We show (Proposition 3.1) that any SQ algorithm that can distinguish between an unknown $F_{\mathbf{v}}$ and a function with uniformly random ± 1 labels requires many high accuracy queries.

The key structural result that we require is the following:

Proposition 3.2. *For any $k \geq 1$, there exists an at most $(k + 1)$ -piecewise constant function $f : \mathbb{R} \rightarrow \{\pm 1\}$ such that $\mathbf{E}_{z \sim \mathcal{N}(0,1)}[f(z)z^t] = 0$, for every non-negative integer $t < k$.*

Proposition 3.2 is the most technically involved result of this section. We give its proof in Section 3.2.

In the remainder of this subsection, we prove Proposition 3.1, assuming Proposition 3.2.

We say that a distribution A has k -matching moments with $\mathcal{N}(0, 1)$ if $\mathbf{E}_{z \sim A}[z^t] = \mathbf{E}_{z \sim \mathcal{N}(0,1)}[z^t]$, for all $0 \leq t < k$. Proposition 3.2 implies the following.

Lemma 3.3. *Let $f : \mathbb{R} \rightarrow \{\pm 1\}$ be such that $\mathbf{E}_{z \sim \mathcal{N}(0,1)}[f(z)z^t] = 0$, for every non-negative integer $t < k$. For $z \sim \mathcal{N}(0, 1)$, define $A \stackrel{\text{def}}{=} \mathbb{1}\{f(z) = 1\}$ and $B \stackrel{\text{def}}{=} \mathbb{1}\{f(z) = -1\}$. Then the random variables A and B have k -matching moments with z .*

Proof. We prove the lemma for the random variable A . The proof for B is similar. From the definition of f , we have that $\mathbf{E}_{z \sim \mathcal{N}(0,1)}[f(z)] = 0$, thus

$$\mathbf{E}_{z \sim \mathcal{N}(0,1)}[\mathbb{1}\{f(z) = 1\}] = \mathbf{E}_{z \sim \mathcal{N}(0,1)}[\mathbb{1}\{f(z) = -1\}]$$

or equivalently

$$\mathbf{E}_{z \sim \mathcal{N}(0,1)}[\mathbb{1}\{f(z) = 1\}] = \frac{1}{2}. \quad (1)$$

Similarly, from $\mathbf{E}_{z \sim \mathcal{N}(0,1)}[f(z)z^t] = 0$, we have

$$\mathbf{E}_{z \sim \mathcal{N}(0,1)}[z^t \mathbb{1}\{f(z) = 1\}] = \mathbf{E}_{z \sim \mathcal{N}(0,1)}[z^t \mathbb{1}\{f(z) = -1\}]. \quad (2)$$

Let $\phi(z|f(z) = 1)$ be the probability distribution of z conditional that $f(z) = 1$. We have that

$$\begin{aligned} \mathbf{E}_{z \sim A}[z^t] &= \int_{-\infty}^{\infty} z^t \phi(z|f(z) = 1) dz = \int_{-\infty}^{\infty} z^t \frac{\phi(z)}{\mathbf{Pr}_{z' \sim \mathcal{N}(0,1)}[f(z') = 1]} \mathbb{1}\{f(z) = 1\} dz \\ &= 2 \int_{-\infty}^{\infty} z^t \phi(z) \mathbb{1}\{f(z) = 1\} dz = \int_{-\infty}^{\infty} z^t \phi(z) \mathbb{1}\{f(z) = 1\} dz + \int_{-\infty}^{\infty} z^t \phi(z) \mathbb{1}\{f(z) = -1\} dz \\ &= \int_{-\infty}^{\infty} z^t \phi(z) dz = \mathbf{E}_{z \sim \mathcal{N}(0,1)}[z^t], \end{aligned}$$

where we used Equations (1), (2). □

We will require the following technical lemmas from [DKS17]. The first lemma says that there exists a large set of unit vectors that are pairwise nearly orthogonal.

Lemma 3.4 (Lemma 3.7 of [DKS17]). *For any $0 < c < 1/2$, there exists a set S of $2^{\Omega(d^c)}$ unit vectors in \mathbb{R}^d such that for each pair of distinct $\mathbf{u}, \mathbf{v} \in S$, we have $|\langle \mathbf{u}, \mathbf{v} \rangle| \leq O(d^{c-1/2})$.*

If we define the conditional distribution on the event $F_{\mathbf{v}}(\mathbf{x}) = 1$, we can see that the directions orthogonal to \mathbf{v} follow a standard $(d - 1)$ -dimensional Gaussian distribution. Thus, we define the following distribution for this case.

Definition 3.5 (High Dimensional Hidden Direction Distribution). For a univariate distribution A with probability density function $A(z)$ and a unit vector $\mathbf{u} \in \mathbb{R}^d$, consider the distribution over \mathbb{R}^d with pdf $\mathbf{P}_{\mathbf{u}}(\mathbf{x}) = A(\langle \mathbf{u}, \mathbf{x} \rangle) \exp(-\|\mathbf{x} - \langle \mathbf{u}, \mathbf{x} \rangle \mathbf{u}\|_2^2/2)/(2\pi)^{(d-1)/2}$. That is, $\mathbf{P}_{\mathbf{u}}$ is the product distribution whose orthogonal projection onto the direction \mathbf{u} is A , and onto the subspace perpendicular to \mathbf{u} is the standard $(d-1)$ -dimensional Gaussian distribution.

Let $\mathcal{D}_1, \mathcal{D}_2 : \mathbb{R}^d \rightarrow \mathbb{R}_+$ be probability density functions. The χ^2 -divergence of $\mathcal{D}_1, \mathcal{D}_2$ is defined as $\chi^2(\mathcal{D}_1, \mathcal{D}_2) \stackrel{\text{def}}{=} \int_{\mathbb{R}^d} \mathcal{D}_1(\mathbf{x})^2 / \mathcal{D}_2(\mathbf{x}) d\mathbf{x} - 1$. We also define the correlation between $\mathcal{D}_1, \mathcal{D}_2$ and a reference distribution \mathcal{D} as $\chi_{\mathcal{D}}(\mathcal{D}_1, \mathcal{D}_2) \stackrel{\text{def}}{=} \int_{\mathbb{R}^d} \mathcal{D}_1(\mathbf{x}) \mathcal{D}_2(\mathbf{x}) / \mathcal{D}(\mathbf{x}) d\mathbf{x} - 1$.

The second lemma states that the distributions $\mathbf{P}_{\mathbf{u}}, \mathbf{P}_{\mathbf{v}}$ have correlation depending on the angle of the corresponding vectors.

Lemma 3.6 (Lemma 3.4 of [DKS17]). *For any $\mathbf{u}, \mathbf{v} \in \mathbb{S}^{d-1}$, let $A : \mathbb{R} \rightarrow \mathbb{R}_+$ be the pdf of a distribution that agrees with the first k moments of $\mathcal{N}(0, 1)$. Then, we have that*

$$|\chi_{\mathcal{N}(\mathbf{0}, \mathbf{I})}(\mathbf{P}_{\mathbf{u}}, \mathbf{P}_{\mathbf{v}})| \leq |\langle \mathbf{u}, \mathbf{v} \rangle|^{k+1} \chi^2(A, \mathcal{N}(0, 1)) .$$

The second statement of Proposition 3.1, establishing non-trivial correlation with a halfspace is shown in the lemma below.

Lemma 3.7. *For any $F \in \mathcal{F}_k$, there is a halfspace σ such that $\mathbf{E}_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})}[F(\mathbf{x})\sigma(\mathbf{x})] \geq 1/(2k)$.*

Proof. We start by noting that each $F \in \mathcal{F}_k$ is of the form $F_{\mathbf{v}}(\mathbf{x}) = f(\langle \mathbf{v}, \mathbf{x} \rangle)$, where f is the function from Proposition 3.2 and $\mathbf{v} \in S$. We will take σ to be $\sigma(\mathbf{x}) = \text{sign}(\langle \mathbf{v}, \mathbf{x} \rangle + \beta)$. Let z_1, \dots, z_k be the breakpoints of $f(z)$. We will show that if we set the value of β to a breakpoint, then the result follows.

Let $a_{i+1} = \int_{z_i}^{z_{i+1}} \phi(z) dz$ for $0 < i < k+1$, $a_1 = \int_{-\infty}^{z_1} \phi(z) dz$ and $a_{k+1} = \int_{z_k}^{\infty} \phi(z) dz$. Let $\beta = z_l$, for a breakpoint z_l , and $b = \text{sign}(f((z_l + z_{l+1})/2))$. Then we have that

$$\mathbf{E}_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})}[F_{\mathbf{v}}(\mathbf{x})\sigma(\mathbf{x})] = 2 \int_{z_l}^{\infty} f(z)\phi(z) dz = 2b \sum_{j=l}^{k+1} (-1)^{j-l} a_j ,$$

where the first equality holds because

$$\mathbf{E}_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})}[F_{\mathbf{v}}(\mathbf{x})\mathbf{1}\{\mathbf{x} \in A\}] = - \mathbf{E}_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})}[F_{\mathbf{v}}(\mathbf{x})\mathbf{1}\{\mathbf{x} \in A^c\}] ,$$

for any $A \subseteq \mathbb{R}^d$. From the fact that $\sum_{i=1}^{k+1} a_i = 1$, it follows that there exists an index i such that $a_i \geq 1/(k+1)$. Assume, for the sake of contradiction, that for all $l > i$ we have that $|\sum_{j=l}^{k+1} (-1)^{l-j} a_j| \leq (1/4k)$, since otherwise there exists a breakpoint that satisfies the equation. Then, for $b = \text{sign}(f((z_i + z_{i+1})/2))$, we have that either $2b(\sum_{j=i+1}^{k+1} (-1)^{j-i} a_j + a_i) \geq (1/2k)$ or $2b(\sum_{j=i+1}^{k+1} (-1)^{j-i} a_j + a_i) \leq -(1/2k)$. In the former case, we are done. In the latter case, the halfspace $-\sigma(\mathbf{x})$ satisfies the desired correlation property. \square

We are now ready to prove (the first statement of) Proposition 3.1.

Proof of Proposition 3.1. Let S be the set of nearly orthogonal vectors from Lemma 3.4 and Y be a uniform ± 1 random variable. For each $\mathbf{v} \in S$, let $F_{\mathbf{v}}(\mathbf{x}) = f(\langle \mathbf{v}, \mathbf{x} \rangle)$, where f is the function from Proposition 3.2. Let \mathcal{D}_D be the set that contains the distributions $(X, F_{\mathbf{v}}(X))$ for any $\mathbf{v} \in S$ and $X \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$. We will prove that for any $\mathbf{u}, \mathbf{v} \in S$, $\mathbf{u} \neq \mathbf{v}$, we have

$$\chi_{(X, Y)}((X, F_{\mathbf{u}}(X)), (X, F_{\mathbf{v}}(X))) \leq 2 \cdot |\langle \mathbf{v}, \mathbf{u} \rangle|^{k+1} . \quad (3)$$

To prove Equation (3), for a unit vector \mathbf{v} , denote by $A_{\mathbf{v}}$ the conditional distribution of the event $F_{\mathbf{v}} = 1$, and by $B_{\mathbf{v}}$ the conditional distribution of the event $F_{\mathbf{v}} = -1$. Let $\mathcal{D}_{\mathbf{v}}$ be the probability density function of $(X, F_{\mathbf{v}}(X))$. We then have

$$\begin{aligned}\chi_{(X,Y)}((X, F_{\mathbf{u}}), (X, F_{\mathbf{v}})) &= 2 \int_{\mathbb{R}^d} \frac{\mathcal{D}_{\mathbf{u}}(\mathbf{x}, 1)\mathcal{D}_{\mathbf{v}}(\mathbf{x}, 1)}{\phi(\mathbf{x})} d\mathbf{x} + 2 \int_{\mathbb{R}^d} \frac{\mathcal{D}_{\mathbf{u}}(\mathbf{x}, -1)\mathcal{D}_{\mathbf{v}}(\mathbf{x}, -1)}{\phi(\mathbf{x})} d\mathbf{x} - 1 \\ &= \frac{1}{2} (\chi_X(A_{\mathbf{u}}, A_{\mathbf{v}}) + \chi_X(B_{\mathbf{u}}, B_{\mathbf{v}})) ,\end{aligned}\quad (4)$$

where we used that Y gets each label with probability $1/2$ and that $\Pr_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})}[F_{\mathbf{v}}(\mathbf{x}) = \pm 1] = 1/2$. From Lemma 3.6 and Lemma 3.3, we have

$$(\chi_X(A_{\mathbf{u}}, A_{\mathbf{v}}) + \chi_X(B_{\mathbf{u}}, B_{\mathbf{v}})) \leq \langle \mathbf{u}, \mathbf{v} \rangle^{k+1} (\chi^2(A, \mathcal{N}(0, 1)) + \chi^2(B, \mathcal{N}(0, 1))) . \quad (5)$$

We also have

$$\begin{aligned}\chi^2(A, \mathcal{N}(0, 1)) &= \int_{-\infty}^{\infty} A(z)^2 / \phi(z) dz = \int_{-\infty}^{\infty} \frac{\phi(z)^2}{\phi(z) \Pr_{z' \sim \mathcal{N}(0, 1)}[f(z') = 1]^2} \mathbf{1}\{f(z) = 1\} dz \\ &= 4 \int_{-\infty}^{\infty} \phi(z) \mathbf{1}\{f(z) = 1\} dz = 2 ,\end{aligned}\quad (6)$$

where we used the conditional expectation and Equation (1). Putting Equations (4), (5) and (6) together, we get Equation (3). Using Lemma 3.4, we have that $|\langle \mathbf{v}, \mathbf{u} \rangle| \leq d^{-(1/2-c)}$, thus

$$\chi_{(X,Y)}((X, F_{\mathbf{u}}(X)), (X, F_{\mathbf{v}}(X))) \leq \Omega\left(d^{-(k+1)(1/2-c)}\right) .$$

To finish our argument, for $\mathbf{u} \neq \mathbf{v}$, we have that

$$\chi_{(X,Y)}((X, F_{\mathbf{u}}(X)), (X, F_{\mathbf{v}}(X))) = \mathbf{E}_{X \sim \mathcal{N}(\mathbf{0}, \mathbf{I})}[F_{\mathbf{u}}(X)F_{\mathbf{v}}(X)] \leq \Omega(d^{-(k+1)(1/2-c)}) .$$

Thus, we have that $\text{SQ-DIM}(\mathcal{F}_k, \mathcal{N}(\mathbf{0}, \mathbf{I})) = \min(d^{\Omega(k)}, 2^{d^c}) = d^{\Omega(k)}$, where the last inequality uses the relation between k and d . By Lemma 2.2, any SQ algorithm that finds a function h such that $\Pr_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})}[F_{\mathbf{v}}(\mathbf{x}) \neq h(\mathbf{x})] \leq 1/2 - d^{-\Omega(k)}$ needs at least $d^{\Omega(k)}$ queries to $\text{STAT}(d^{-\Omega(k)})$. This completes the proof. \square

3.2 Proof of Proposition 3.2

The key lemma for the proof is the following.

Lemma 3.8. *Let m and k be positive integers such that $m > k + 1$ and $\epsilon > 0$. If there exists an m -piecewise constant $f : \mathbb{R} \mapsto \{\pm 1\}$ such that $|\mathbf{E}_{z \sim \mathcal{N}(0, 1)}[f(z)z^t]| < \epsilon$ for all non-negative integers $t < k$, then there exists an at most $(m - 1)$ -piecewise constant $g : \mathbb{R} \mapsto \{\pm 1\}$ such that $|\mathbf{E}_{z \sim \mathcal{N}(0, 1)}[g(z)z^t]| < \epsilon$ for all non-negative integers $t < k$.*

Proof. Let $\{b_1, b_2, \dots, b_{m-1}\}$ be the breakpoints of f . Then let $F(z_1, z_2, \dots, z_{m-1}, z) : \overline{\mathbb{R}}^m \mapsto \mathbb{R}$ be an m -piecewise constant function with breakpoints on z_1, \dots, z_{m-1} , where $z_1 < z_2 < \dots < z_{m-1}$ and $F(b_1, b_2, \dots, b_{m-1}, z) = f(z)$. For simplicity, let $\mathbf{z} = (z_1, \dots, z_{m-1})$ and define $M_i(\mathbf{z}) = \mathbf{E}_{z \sim \mathcal{N}(0, 1)}[F(\mathbf{z}, z)z^i]$ and let $\mathbf{M}(\mathbf{z}) = [M_0(\mathbf{z}), M_1(\mathbf{z}), \dots, M_{k-1}(\mathbf{z})]^T$. It is clear from the definition that $M_i(\mathbf{z}) = \sum_{n=0}^{m-1} \int_{z_n}^{z_{n+1}} F(\mathbf{z}, z)z^i \phi(z) dz = \sum_{n=0}^{m-1} a_n \int_{z_n}^{z_{n+1}} z^i \phi(z) dz$, where $z_0 = -\infty$ and $z_m = \infty$ and

a_n is the sign of $F(\mathbf{z}, z)$ in the interval (z_n, z_{n+1}) . Note that $a_n = -a_{n+1}$ for every $0 \leq n < m$. By taking the derivative of M_i in z_j , for $0 < j < m$, we get that

$$\frac{\partial}{\partial z_j} M_i(\mathbf{z}) = 2a_{j-1} z_j^i \phi(z_j) \quad \text{and} \quad \frac{\partial}{\partial z_j} \mathbf{M}(\mathbf{z}) = 2a_{j-1} \phi(z_j) [1, z_j^1, \dots, z_j^{k-1}]^T.$$

We now argue that for any \mathbf{z} with distinct coordinates that there exists a vector $\mathbf{u} \in \mathbb{R}^{m-1}$ such that $\mathbf{u} = (\mathbf{u}_1, \dots, \mathbf{u}_k, 0, 0, \dots, 0, 1)$ and the directional derivative of \mathbf{M} in the \mathbf{u} direction is zero. To prove this, we construct a system of linear equations such that $\nabla_{\mathbf{u}} M_i(\mathbf{z}) = 0$, for all $0 \leq i < k$. Indeed, we have $\sum_{j=1}^k \frac{\partial}{\partial z_j} M_i(\mathbf{z}) \mathbf{u}_j = -\frac{\partial}{\partial z_{m-1}} M_i(\mathbf{z})$ or $\sum_{j=1}^k a_{j-1} z_j^i \phi(z_j) \mathbf{u}_j = -a_{m-2} z_{m-1}^i \phi(z_{m-1})$, which is linear in the variables \mathbf{u}_j . Let $\hat{\mathbf{u}}$ be the vector with the first k variables and let \mathbf{w} be the vector of the right hand side of the system, i.e., $\mathbf{w}_i = -a_{m-2} z_{m-1}^i \phi(z_{m-1})$. Then this system can be written in matrix form as $\mathbf{V} \mathbf{D} \hat{\mathbf{u}} = \mathbf{w}$, where \mathbf{V} is the Vandermonde matrix, i.e., the matrix that is $\mathbf{V}_{i,j} = \alpha_i^{j-1}$, for some values α_i and \mathbf{D} is a diagonal matrix. In our case, $\mathbf{V}_{i,j} = z_i^{j-1}$ and $\mathbf{D}_{j,j} = 2a_{j-1} \phi(z_j)$. It is known that the Vandermonde matrix has full rank iff for all $i \neq j$ we have $\alpha_i \neq \alpha_j$, which holds in our setting. Thus, the matrix $\mathbf{V} \mathbf{D}$ is nonsingular and there exists a solution to the equation. Thus, there exists a vector \mathbf{u} with our desired properties and, moreover, any vector in this direction is a solution of this system of linear equations. Note that the vector \mathbf{u} depends on the value of \mathbf{z} , thus we consider $\mathbf{u}(\mathbf{z})$ be the (continuous) function that returns a vector \mathbf{u} given \mathbf{z} .

We define a differential equation for the function $\mathbf{v} : \overline{\mathbb{R}} \mapsto \overline{\mathbb{R}}^{m-1}$, as follows: $\mathbf{v}(0) = \mathbf{b}$, where $\mathbf{b} = (b_1, \dots, b_{m-1})$, and $\mathbf{v}'(T) = \mathbf{u}(\mathbf{v}(T))$ for all $T \in \overline{\mathbb{R}}$. If \mathbf{v} is a solution to this differential equation, then we have:

$$\frac{d}{dT} \mathbf{M}(\mathbf{v}(T)) = \frac{d}{d\mathbf{v}(T)} \mathbf{M}(\mathbf{v}(T)) \frac{d}{dT} \mathbf{v}(T) = \frac{d}{d\mathbf{v}(T)} \mathbf{M}(\mathbf{v}(T)) \mathbf{u}(\mathbf{v}(T)) = \mathbf{0},$$

where we used the chain rule and that the directional derivative in $\mathbf{u}(\mathbf{v}(T))$ direction is zero. This means that the function $\mathbf{M}(\mathbf{v}(t))$ is constant, and for all $0 \leq j < k$, we have $|M_j| < \epsilon$, because we have that $|\mathbf{E}_{z \sim \mathcal{N}(0,1)} [F(z_1, \dots, z_{m-1}, z) z^t]| < \epsilon$. Furthermore, since $\mathbf{u}(\mathbf{v}(T))$ is continuous in $\mathbf{v}(T)$, this differential equation will be well founded and have a solution up until the point where either two of the z_i approach each other or one of the z_i approaches plus or minus infinity (the solution cannot oscillate, since $\mathbf{v}'_{m-1}(T) = 1$ for all T).

Running the differential equation until we reach such a limit, we find a limiting value \mathbf{v}^* of $\mathbf{v}(T)$ so that either:

1. There is an i such that $\mathbf{v}_i^* = \mathbf{v}_{i+1}^*$, which gives us a function that is at most $(m-2)$ -piecewise constant, i.e., taking $F(\mathbf{v}^*, z)$.
2. Either $\mathbf{v}_{m-1}^* = \infty$ or $\mathbf{v}_1^* = -\infty$, which gives us an at most $(m-1)$ -piecewise constant function, i.e., taking $F(\mathbf{v}^*, z)$. Since when the $\mathbf{v}_{m-1}^* = \infty$, the last breakpoint becomes ∞ , we have one less breakpoint, and if $\mathbf{v}_1^* = -\infty$ we lose the first breakpoint.

Thus, in either case we have a function with at most $m-1$ breakpoints and the same moments. This completes the proof. \square

We also require the following simple fact, establishing the existence of a k' -piecewise constant Boolean-valued function (for some finite k'), satisfying the desired moment conditions.

Fact 3.9. *For any $\epsilon > 0$, there exists a $(k/\epsilon)^{O(k)}$ -piecewise constant function $f : \mathbb{R} \mapsto \{\pm 1\}$ such that $|\mathbf{E}_{z \sim \mathcal{N}(0,1)} [f(z) z^t]| \leq \epsilon$, for every integer $0 \leq t < k$.*

Proof. We define f to take alternative values ± 1 in intervals of length s . Let us denote $I_i = (is, (i+1)s)$ for $-1/(s\epsilon^k) \leq i \leq 1/(s\epsilon^k)$ for an integer i . If $f(z) = 1$ for $z \in I_i$, then we have $f(z) = -1$ for $z \in I_{i+1}$. Moreover, we will have that $f(z) = 1$ for $z \leq -1/\epsilon^k$ and $f(z) = -1$ for $z > 1/\epsilon^k$. We assume that the number of constant pieces is even for simplicity. To prove that for all $0 \leq t < k$, $\mathbf{E}_{z \sim \mathcal{N}(0,1)}[f(z)z^t] < 4\epsilon$, observe that for all even moments the expectation is equal to zero. So it suffices to prove the desired statement for odd moments. Note that $\mathbf{E}_{z \sim \mathcal{N}(0,1)}[z^t f(z) \mathbb{1}\{z \geq 0\}] = \mathbf{E}_{z \sim \mathcal{N}(0,1)}[|z|^t |f(z)| \mathbb{1}\{z < 0\}]$ for odd moments. Thus, we will prove that $\mathbf{E}_{z \sim \mathcal{N}(0,1)}[z^t f(z) \mathbb{1}\{z \geq 0\}] \leq 2\epsilon$. We have that

$$\int_{1/\epsilon^k}^{\infty} z^t \phi(z) dz \leq \epsilon^{k/t},$$

where we used the inequality $\Pr_{z \sim \mathcal{N}(0,1)}[|z|^t \geq y] \leq \frac{1}{\sqrt{2\pi}y^{1/t}} e^{-y^{2/t}} \leq 1/y^{1/t}$. Moreover, we bound from above the absolute ratio between two subsequent regions, i.e., $\left| \frac{\mathbf{E}_{z \sim \mathcal{N}(0,1)}[z^t f(z) \mathbb{1}\{z \in I_i\}]}{\mathbf{E}_{z \sim \mathcal{N}(0,1)}[z^t f(z) \mathbb{1}\{z \in I_{i+1}\}]} \right|$. For $i \geq 0$, we have that

$$\frac{\int_{is}^{(i+1)s} z^t \phi(z) dz}{\int_{(i+1)s}^{(i+2)s} z^t \phi(z) dz} \leq \frac{s((i+1)s)^t \phi(is)}{s((i+1)s)^t \phi((i+2)s)} = e^{2is^2 + 2s^2} \leq 1 + 3is^2 + 9i^2 s^4, \quad (7)$$

where in the first inequality we used the maximum value and the minimum of the integral, and in the second one we used that $e^x \leq 1 + x + x^2$ for $x \leq 1$, which holds for $s < \epsilon^k$. Thus, for two subsequent intervals we have

$$\int_{is}^{(i+1)s} z^t \phi(z) dz - \int_{(i+1)s}^{(i+2)s} z^t \phi(z) dz \leq 4is^2 \int_{(i+1)s}^{(i+2)s} z^t \phi(z) dz \leq 4ks^2 \int_{(i+1)s}^{(i+2)s} z^t \phi(z) dz.$$

On the other direction, from Equation (7) we have that

$$- \int_{is}^{(i+1)s} z^t \phi(z) dz + \int_{(i+1)s}^{(i+2)s} z^t \phi(z) dz \geq -4is^2 \int_{(i+1)s}^{(i+2)s} z^t \phi(z) dz \geq -4ks^2 \int_{(i+1)s}^{(i+2)s} z^t \phi(z) dz.$$

Thus, we have

$$-4ks^2(t-1)!! \leq \sum_{i=0}^{1/(s\epsilon^k)} (-1)^i \int_{is}^{(i+1)s} z^t \phi(z) dz \leq 4ks^2 \int_{-\infty}^{\infty} z^t \phi(z) dz = 4ks^2(t-1)!!.$$

Choosing $s = \epsilon^{(k+1)/2}/k^k$ and setting $\epsilon = \epsilon/2$, the proof follows. \square

Proposition 3.2 follows from the above using a compactness argument.

Proof of Proposition 3.2. For every $\epsilon > 0$, using the function f' from Fact 3.9 and Lemma 3.8, we can obtain a function f_ϵ such that $|\mathbf{E}_{z \sim \mathcal{N}(0,1)}[f_\epsilon(z)z^t]| \leq \epsilon$, for every non-negative integer $t < k$ and the function f_ϵ is at most $(k+1)$ -piecewise constant. Let $\mathbf{M} : \overline{\mathbb{R}^k} \mapsto \mathbb{R}^k$, where $M_i(\mathbf{b}) = \sum_{n=0}^k (-1)^{n+1} \int_{b_n}^{b_{n+1}} z^i \phi(z) dz$ and $b_1 \leq b_2 \leq \dots \leq b_k$, $b_0 = -\infty$ and $b_{k+1} = \infty$. Here we assume without loss of generality that before the first breakpoint the function is negative because we can always set the first breakpoint to be $-\infty$. It is clear that the function \mathbf{M} is a continuous map and $\overline{\mathbb{R}^{k+1}}$ is a compact set, thus $\mathbf{M}(\overline{\mathbb{R}^{k+1}})$ is a compact set. We also have that for every $\epsilon > 0$ there is a point $\mathbf{b} \in \overline{\mathbb{R}^{k+1}}$ such that $|\langle \mathbf{M}(\mathbf{b}), \mathbf{e}_i \rangle| \leq \epsilon$. Thus, from compactness, we have that there exists a point $\mathbf{b}^* \in \overline{\mathbb{R}^{k+1}}$ such that $\mathbf{M}(\mathbf{b}^*) = \mathbf{0}$. This completes the proof. \square

4 SQ Lower Bound for ReLU Regression

In this section, we give the proof of Theorem 1.5. To prove our theorem, we construct a class \mathcal{F} of Boolean-valued functions that is SQ hard to weakly learn. We use the SQ hardness of \mathcal{F} to show that agnostically learning a ReLU with respect the square loss is also SQ hard.

Our proof depends critically on the following technical result.

Proposition 4.1. *For all $k \geq 1$, there exists an $O(k)$ -piecewise constant function $f : \mathbb{R} \rightarrow \{\pm 1\}$ such that $\mathbf{E}_{z \sim \mathcal{N}(0,1)}[f(z)z^t] = 0$ for every non-negative integer $t \leq k$, and $\mathbf{E}_{z \sim \mathcal{N}(0,1)}[f(z)\text{ReLU}(z)] \geq 1/\text{poly}(k)$.*

The proof of Proposition 4.1 is given in Section 4.1.

We start with a brief overview of the proof. To prove Proposition 4.1, we first make essential use of Legendre polynomials to construct an explicit function $f' : \mathbb{R} \rightarrow [-1, 1]$ with the correct properties. By rounding f' to a Boolean-valued function, it is not hard to show that for every $\epsilon > 0$ there is a Boolean-valued function whose first k moments are at most ϵ , and whose correlation with a ReLU is at least $1/\text{poly}(k)$. Using a slight variation of the techniques from Lemma 3.8, we can obtain a function with these moments that is $O(k)$ -piecewise constant. Taking a limit of such functions with ϵ tending to 0 gives the proposition.

The hard family of functions to learn will then be $\mathcal{F} = \{F_{\mathbf{v}}\}_{\mathbf{v} \in S}$, where S is a set of nearly orthogonal unit vectors, and $F_{\mathbf{v}}(\mathbf{x}) = C \cdot f(\langle \mathbf{v}, \mathbf{x} \rangle)$, for $C > 0$ some appropriately chosen number of size polynomial in k . Since $f(\langle \mathbf{v}, \mathbf{x} \rangle)$ correlates with $\text{ReLU}(\langle \mathbf{v}, \mathbf{x} \rangle)$, taking C sufficiently large, we have

$$\mathbf{E} \left[(F_{\mathbf{v}}(\mathbf{x}) - \text{ReLU}(\langle \mathbf{v}, \mathbf{x} \rangle))^2 \right] < \mathbf{E}[F_{\mathbf{v}}(\mathbf{x})^2] (1 - 1/\text{poly}(k)) .$$

If ϵ is a sufficiently small polynomial in $1/k$, any learner would need to return a function g where $\mathbf{E}[(F_{\mathbf{v}}(\mathbf{x}) - g(\mathbf{x}))^2] < \mathbf{E}[F_{\mathbf{v}}(\mathbf{x})^2](1 - 1/\text{poly}(k))$. This implies both that $\mathbf{E}[(1/C^2)g^2]$ is not too large and that g correlates non-trivially with the Boolean-valued function $(1/C)F_{\mathbf{v}}$. However, since the class of functions $\{(1/C)F_{\mathbf{v}}\}_{\mathbf{v} \in S}$ has large SQ-dimension (because of the moment matching property established by Proposition 4.1), Theorem 1.5 follows from Lemma 2.2.

We now give the proof of Theorem 1.5, assuming Proposition 4.1

Proof of Theorem 1.5. The proof follows using the same construction as in Theorem 1.4, but using the $O(k)$ -piecewise constant function f from Proposition 4.1. Let $C(k)$ be a constant that depends on k and \mathcal{F}_k be the family of $O(k)$ -decision lists of halfspaces, where each $F_{\mathbf{v}} \in \mathcal{F}_k$ has the form $F_{\mathbf{v}}(\mathbf{x}) = C(k) \cdot f(\langle \mathbf{v}, \mathbf{x} \rangle)$, for a unit vector $\mathbf{v} \in S$, where we use the set S from Lemma 3.4. Let \mathcal{A} be an agnostic SQ learner for ReLUs under Gaussian marginals. We feed \mathcal{A} a set of i.i.d. labeled examples from an arbitrary function $F_{\mathbf{v}} \in \mathcal{F}_k$. By definition, algorithm \mathcal{A} computes a hypothesis $h : \mathbb{R}^d \mapsto \mathbb{R}$ such that

$$\mathbf{E}_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})} [(h(\mathbf{x}) - F_{\mathbf{v}}(\mathbf{x}))^2] \leq \inf_{f \in \mathcal{C}_{\text{ReLU}}} \mathbf{E}_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})} [(f(\mathbf{x}) - F_{\mathbf{v}}(\mathbf{x}))^2] + \epsilon .$$

We denote $\|g\|_2^2 = \mathbf{E}_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})} [g(\mathbf{x})^2]$ for a function $g : \mathbb{R}^d \mapsto \mathbb{R}$. Let $C(k) = \frac{\|\text{ReLU}\|_2^2}{\mathbf{E}_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})} [f(\langle \mathbf{x}, \mathbf{v} \rangle) \text{ReLU}(\langle \mathbf{x}, \mathbf{v} \rangle)]}$. Then we have that

$$\begin{aligned} \mathbf{E}_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})} [(\text{ReLU}(\langle \mathbf{x}, \mathbf{v} \rangle) - F_{\mathbf{v}}(\mathbf{x}))^2] &= \|F_{\mathbf{v}}\|_2^2 + \|\text{ReLU}\|_2^2 - 2 \mathbf{E}_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})} [F_{\mathbf{v}}(\mathbf{x}) \text{ReLU}(\langle \mathbf{x}, \mathbf{v} \rangle)] \\ &= C^2(k) \|f\|_2^2 - \|\text{ReLU}\|_2^2 . \end{aligned}$$

Furthermore, using that $\|f\|_2^2 = 1$ and $\|\text{ReLU}\|_2^2 = 1/2$, if we choose $\epsilon = o(1/C^2(k))$, the algorithm returns a hypothesis such that

$$\mathbf{E}_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})} [(h(\mathbf{x}) - F_{\mathbf{v}}(\mathbf{x}))^2] \leq C^2(k) (1 - \Omega(1/C^2(k))) .$$

Thus, from the triangle inequality, we have that $\|h/C(k)\|_2^2 \leq 2\|f\|_2^2$, and also

$$2 \mathbf{E}_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})} \left[\frac{h(\mathbf{x})}{C(k)} \frac{F_{\mathbf{v}}(\mathbf{x})}{C(k)} \right] \geq \Omega(1/C^2(k)) + \|h\|_2^2 / C^2(k) \geq \Omega(1/C^2(k)) .$$

Finally,

$$\mathbf{E}_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})} \left[\frac{h(\mathbf{x})}{\|h\|_2} \frac{F_{\mathbf{v}}(\mathbf{x})}{\|F_{\mathbf{v}}\|_2} \right] \geq \frac{1}{2} \mathbf{E}_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})} \left[\frac{h(\mathbf{x})}{C(k)} \frac{F_{\mathbf{v}}(\mathbf{x})}{C(k)} \right] \geq \Omega(1/C^2(k)) .$$

Let $h^*(\mathbf{x}) = \frac{h(\mathbf{x})}{\|h\|_2}$ and $F_{\mathbf{v}}^*(\mathbf{x}) = \frac{F_{\mathbf{v}}(\mathbf{x})}{\|F_{\mathbf{v}}\|_2}$. Then we have that $\mathbf{E}_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})} [h^*(\mathbf{x})F_{\mathbf{v}}^*(\mathbf{x})] \geq \Omega(1/C^2(k))$. Thus, using Proposition 4.1 to bound $C(k)$, we get that

$$\mathbf{E}_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})} [h^*(\mathbf{x})F_{\mathbf{v}}^*(\mathbf{x})] \geq \Omega(1/\text{poly}(k)) .$$

Since the function $F_{\mathbf{v}}$ is an $O(k)$ -decision list of halfspaces, we can apply Proposition 3.1 to get that any SQ algorithm needs $d^{\Omega(k)}$ queries to $\text{STAT}(d^{-\Omega(k)})$ to get $\mathbf{E}_{\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})} [h^*(\mathbf{x})F_{\mathbf{v}}^*(\mathbf{x})] \geq d^{-\Omega(k)}$. Thus, in order to learn with error $\text{OPT} + \epsilon$, for $\epsilon = o(1/\text{poly}(k))$, the algorithm \mathcal{A} needs to use $d^{\Omega((1/\epsilon)^c)}$ queries to $\text{STAT}(d^{-\Omega((1/\epsilon)^c)})$, for a constant $c > 0$. \square

4.1 Proof of Proposition 4.1

To prove Proposition 4.1, we first need to prove that there exists a function that has non-trivial correlation with the ReLU and whose first k moments are zero.

We have the following crucial proposition.

Proposition 4.2. *Let k be a positive integer. There exists a function $f : \mathbb{R} \mapsto [-1, 1]$ such that $\mathbf{E}_{z \sim \mathcal{N}(0,1)} [f(z)z^t] = 0$, for $0 \leq t \leq k$, and $\mathbf{E}_{z \sim \mathcal{N}(0,1)} [f(z)\text{ReLU}(z)] > 1/\text{poly}(k)$.*

The proof of Proposition 4.2 requires analytic properties of the Legendre polynomials and is deferred to Section 4.2. In the main part of this subsection, we prove Proposition 4.1, assuming Proposition 4.2.

In the following lemma, we show that there exists a piecewise constant Boolean-valued function with near-vanishing moments of degree at most k and non-trivial correlation with the ReLU.

Lemma 4.3. *For any $\epsilon > 0$ and any non-negative integer k , there exists a piecewise constant function $G : \mathbb{R} \mapsto \{\pm 1\}$ such that $|\mathbf{E}_{z \sim \mathcal{N}(0,1)} [G(z)z^t]| \leq \epsilon$, for $0 \leq t \leq k$, and $\mathbf{E}_{z \sim \mathcal{N}(0,1)} [G(z)\text{ReLU}(z)] > 1/\text{poly}(k) + O(\epsilon)$.*

Proof. The proof is similar to the proof of Fact 3.9. The main difference here is that we need to construct a function that will also have non-trivial correlation with the ReLU. To do this, we use a probabilistic argument to show that there exists a function that is bounded in the range $[-1, 1]$, that has non trivial correlation, and then we discretize the function as in Fact 3.9. Let f be the function from Proposition 4.2. We split the interval $[-1, 1]$ into sub-intervals of length δ and we define the random piecewise constant function G in each interval $[z_0, z_0 + \delta]$ as $G(z) = 1$ with probability $(1 + \int_{z_0}^{z_0+\delta} f(z)\phi(z)dz / \int_{z_0}^{z_0+\delta} \phi(z)dz) / 2$ and $G(z) = -1$ with probability $(1 -$

$\int_{z_0}^{z_0+\delta} f(z)\phi(z)dz / \int_{z_0}^{z_0+\delta} \phi(z)dz / 2$. Thus, in each interval, we have $\mathbf{E}[G(z)] = \int_{z_0}^{z_0+\delta} f(z)\phi(z)dz / \int_{z_0}^{z_0+\delta} \phi(z)dz$. Then, for any $|z_0| \leq 1 - \delta$, we have that

$$\begin{aligned} \mathbf{E} \left[\int_{z_0}^{z_0+\delta} G(z)z^t \phi(z)dz \right] &= \mathbf{E} \left[\int_{z_0}^{z_0+\delta} G(z)(z_0 + O(\delta))^t \phi(z)dz \right] = \int_{z_0}^{z_0+\delta} f(z)\phi(z)(z_0 + O(\delta))^t dz \\ &= \int_{z_0}^{z_0+\delta} f(z)\phi(z)z^t dz + \int_{z_0}^{z_0+\delta} t \cdot O((|z_0| + \delta)^{t-1}\delta) dz \\ &= \int_{z_0}^{z_0+\delta} f(z)z^t \phi(z)dz + t \cdot O((|z_0| + \delta)^{t-1}\delta^2) , \end{aligned}$$

where we used the Taylor series $z^t = (z_0 + O(\delta))^t + t \cdot O(\delta(|z_0| + \delta)^{t-1})$. Thus, we obtain

$$\mathbf{E} \left[\int_{-1}^1 G(z)z^t \phi(z)dz \right] = \int_{-1}^1 f(z)z^t \phi(z)dz + t \cdot O(\delta) = t \cdot O(\delta) ,$$

where we used that all the moments of f with degree at most k are zero and that $|z_0| + \delta \leq 1$. Moreover, for $0 \leq z_0 \leq 1$, it holds that

$$\mathbf{E} \left[\int_{z_0}^{z_0+\delta} G(z)\text{ReLU}(z)\phi(z)dz \right] = \mathbf{E} \left[\int_{z_0}^{z_0+\delta} G(z)z\phi(z)dz \right] = \int_{z_0}^{z_0+\delta} f(z)\text{ReLU}(z)\phi(z)dz + t \cdot O((|z_0| + \delta)^{t-1}\delta^2) ,$$

where we used the same method as before. Thus, it follows that

$$\mathbf{E} \left[\int_0^1 G(z)\text{ReLU}(z)\phi(z)dz \right] = \int_0^1 f(z)\text{ReLU}(z)\phi(z)dz + t \cdot O(\delta) > 1/\text{poly}(k) + t \cdot O(\delta) .$$

Define the random variable $X_{i,t} = \int_{i\cdot\delta}^{i\cdot\delta+\delta} G(z)z^t \phi(z)dz$ and $X_t = \sum_{i=-1/\delta}^{1/\delta} X_{i,t}$. Using Hoeffding bounds, we have that

$$\Pr[|X_t - \mathbf{E}[X_t]| > \sqrt{\delta} \log(4/(t+1))] \leq 1/(2(t+1)) ,$$

where we used that $|X_{i,t}| \leq \delta$. By the union bound, we get that there is positive probability that all X_t are within $\pm\sqrt{\delta} \log(4/(t+1))$ from the mean value, and thus, from the probabilistic method there is a function with this property. Furthermore, we round the rest of the values of $G(z)$ as in the proof of Fact 3.9 (because $\text{ReLU}(z) = z$ for $z > 0$). Choosing the correct constant value of δ , the result follows. \square

Lemma 4.4. *Let m and k be positive integers such that $m > 2k + 5$ and $\epsilon > 0$. If there exists an m -piecewise constant function $f : \mathbb{R} \mapsto \{\pm 1\}$ such that $|\mathbf{E}_{z \sim \mathcal{N}(0,1)}[f(z)z^t]| < \epsilon$ for all non-negative integers $t \leq k$, and $\mathbf{E}_{z \sim \mathcal{N}(0,1)}[f(z)\text{ReLU}(z)] > 1/\text{poly}(k) + O(\epsilon)$, then there exists an at most $(2k + 5)$ -piecewise constant function $g : \mathbb{R} \mapsto \{\pm 1\}$ such that $|\mathbf{E}_{z \sim \mathcal{N}(0,1)}[g(z)z^t]| < \epsilon$ for all non-negative integers $t \leq k$ and $\mathbf{E}_{z \sim \mathcal{N}(0,1)}[g(z)\text{ReLU}(z)] > 1/\text{poly}(k) + O(\epsilon)$.*

Proof. This proof is similar to the proof of Lemma 3.8. The only difference is that we have to keep also the correlation with the ReLU constant. For completeness, we provide a full proof.

Let $\{b_1, b_2, \dots, b_{m-1}\}$ be the breakpoints of f . Let $F(z_1, z_2, \dots, z_{m-1}, z) : \overline{\mathbb{R}}^m \mapsto \mathbb{R}$ be an m -piecewise constant function with breakpoints on z_1, \dots, z_{m-1} , where $z_1 < z_2 < \dots < z_{m-1}$ and $F(b_1, b_2, \dots, b_{m-1}, z) = f(z)$. For simplicity, let $\mathbf{z} = (z_1, \dots, z_{m-1})$ and define $M_i(\mathbf{z}) =$

$\mathbf{E}_{z \sim \mathcal{N}(0,1)}[F(\mathbf{z}, z)z^i]$, for all $0 \leq i \leq k$ and $M_c(\mathbf{z}) = \mathbf{E}_{z \sim \mathcal{N}(0,1)}[F(\mathbf{z}, z)\text{ReLU}(z)]$. Finally, let $\mathbf{M}(\mathbf{z}) = [M_0(\mathbf{z}), M_1(\mathbf{z}), \dots, M_k(\mathbf{z}), \text{ar}M_c(\mathbf{z})]^T$. It is clear that

$$M_i(\mathbf{z}) = \sum_{n=0}^{m-1} \int_{z_n}^{z_{n+1}} F(\mathbf{z}, z)z^i \phi(z) dz = \sum_{n=0}^{m-1} a_n \int_{z_n}^{z_{n+1}} z^i \phi(z) dz ,$$

and

$$M_c(\mathbf{z}) = \sum_{n=0}^{m-1} \int_{z_n}^{z_{n+1}} F(\mathbf{z}, z)z \mathbb{1}\{z > 0\} \phi(z) dz = \sum_{n=0}^{m-1} a_n \int_{z_n}^{z_{n+1}} z \mathbb{1}\{z > 0\} \phi(z) dz ,$$

where $z_0 = -\infty$, $z_m = \infty$, and a_n is the sign of $F(\mathbf{z}, z)$ in the interval (z_n, z_{n+1}) . Note that $a_n = -a_{n+1}$ for every $0 \leq n < m$. By taking the derivative of M_c and M_i in z_j , for $0 < j < m$, we get that

$$\frac{\partial}{\partial z_j} M_i(\mathbf{z}) = 2a_{j-1}z_j^i \phi(z_j) \quad \text{and} \quad \frac{\partial}{\partial z_j} M_c(\mathbf{z}) = \begin{cases} 2a_{j-1}z_j \phi(z_j), & \text{if } a_j > 0 \\ 0, & \text{if } a_j \leq 0 \end{cases} .$$

Combining the above, we get

$$\frac{\partial}{\partial z_j} \mathbf{M}(\mathbf{z}) = \begin{cases} 2a_{j-1} \phi(z_j) [1, z_j^1, \dots, z_j^k, z_j]^T, & \text{if } z_j > 0 \\ 2a_{j-1} \phi(z_j) [1, z_j^1, \dots, z_j^k, 0]^T, & \text{if } z_j \leq 0 . \end{cases}$$

We first work with the positive breakpoints. Let i_0 be the index of the first positive breakpoint and assume that the positive breakpoints are $m' > k + 2$. We argue that there exists a vector $\mathbf{u} \in \mathbb{R}^{m-1}$ such that $\mathbf{u} = (0, \dots, 0, \mathbf{u}_{i_0+1}, \dots, \mathbf{u}_{i_0+k+2}, 0, 0, \dots, 0, 1)$ and the directional derivative of \mathbf{M} in \mathbf{u} is zero. To prove this, we construct a system of linear equations, such that $\nabla_{\mathbf{u}} M_i(\mathbf{z}) = 0$ for all $0 \leq i \leq k$ and $\nabla_{\mathbf{u}} M_c(\mathbf{z}) = 0$. Indeed, we have $\sum_{j=1}^k \frac{\partial}{\partial z_j} M_i(\mathbf{z}) \mathbf{u}_j = -\frac{\partial}{\partial z_{m-1}} M_i(\mathbf{z})$ or $\sum_{j=1}^k a_{j-1} z_j^i \phi(z_j) \mathbf{u}_j = -a_{m-2} z_{m-1}^i \phi(z_{m-1})$ and $\sum_{j=1}^k a_{j-1} z_j \phi(z_j) \mathbf{u}_j \mathbb{1}\{z_j \geq 0\} = -a_{m-2} z_{m-1} \phi(z_{m-1}) \mathbb{1}\{z_{m-1} \geq 0\}$, which is linear in the variables \mathbf{u}_j . Note that the last equation is the same equation as the $\nabla_{\mathbf{u}} M_1(\mathbf{z}) = 0$, because we have positive breakpoints only. Let $\hat{\mathbf{u}}$ be the vector with the variables from index $i_0 + 1$ to $i_0 + k + 2$, and let \mathbf{w} be the vector of the right hand side of the system, i.e., $\mathbf{w}_i = -a_{m-2} z_{m-1}^i \phi(z_{m-1})$. Then this system can be written in matrix form as $\mathbf{V} \mathbf{D} \hat{\mathbf{u}} = \mathbf{w}$, where \mathbf{V} is the Vandermonde matrix, i.e., the matrix that is $\mathbf{V}_{i,j} = \alpha_i^{j-1}$, for some values α_i and \mathbf{D} is a diagonal matrix. In our case, $\mathbf{V}_{i,j} = z_i^{j-1}$ and $\mathbf{D}_{j,j} = 2a_{j-1} \phi(z_j)$. It is known that the Vandermonde matrix has full rank iff for all $i \neq j$ we have $\alpha_i \neq \alpha_j$, which holds in our setting. Thus, the matrix $\mathbf{V} \mathbf{D}$ is nonsingular and there exists a solution to the equation. Thus, there exists a vector \mathbf{u} with our desired properties and, moreover, any vector in this direction is a solution to this system of linear equations. Note that the vector \mathbf{u} depends on the value of \mathbf{z} , thus we consider $\mathbf{u}(\mathbf{z})$ be the (continuous) function that returns a vector \mathbf{u} given \mathbf{z} .

We define a differential equation for the function $\mathbf{v} : \overline{\mathbb{R}} \mapsto \overline{\mathbb{R}}^{m-1}$, as follows: $\mathbf{v}(0) = \mathbf{b}$, where $\mathbf{b} = (b_1, \dots, b_{m-1})$, and $\mathbf{v}'(T) = \mathbf{u}(\mathbf{v}(T))$ for all $T \in \overline{\mathbb{R}}$. If \mathbf{v} is a solution to this differential equation, then we have:

$$\frac{d}{dT} \mathbf{M}(\mathbf{v}(T)) = \frac{d}{d\mathbf{v}(T)} \mathbf{M}(\mathbf{v}(T)) \frac{d}{dT} \mathbf{v}(T) = \frac{d}{d\mathbf{v}(T)} \mathbf{M}(\mathbf{v}(T)) \mathbf{u}(\mathbf{v}(T)) = \mathbf{0} ,$$

where we used the chain rule and that the directional derivative in the $\mathbf{u}(\mathbf{v}(T))$ direction is zero. This means that the function $\mathbf{M}(\mathbf{v}(t))$ is constant and, for all $0 \leq j < k$, we have $|M_j| < \epsilon$, because we have that $|\mathbf{E}_{z \sim \mathcal{N}(0,1)}[F(z_1, \dots, z_{m-1}, z)z^t]| < \epsilon$. Furthermore, since $\mathbf{u}(\mathbf{v}(T))$ is continuous in

$\mathbf{v}(T)$, this differential equation will be well founded and have a solution up until the point where either two of the z_i approach each other or one of the z_i approaches plus or to zero (the solution cannot oscillate, since $\mathbf{v}'_{m-1}(T) = 1$ for all T).

Running the differential equation until we reach such a limit, we find a limiting value \mathbf{v}^* of $\mathbf{v}(T)$ so that either:

1. There is an i such that $\mathbf{v}_i^* = \mathbf{v}_{i+1}^*$, which gives us a function that is at most $(m-2)$ -piecewise constant, i.e., taking $F(\mathbf{v}^*, z)$.
2. $\mathbf{v}_{m-1}^* = \infty$, which gives us an at most $(m-1)$ -piecewise constant function, i.e., taking $F(\mathbf{v}^*, z)$. Since when the $\mathbf{v}_{m-1}^* = \infty$, the last breakpoint becomes ∞ , we have one less breakpoint.
3. $\mathbf{v}_{i_0+1}^* = 0$, which gives us one less positive breakpoint.

By iterating this method, we can get a function f' that has at most $k+2$ positive breakpoints. For the negative breakpoints, we work in a similar way, with the only difference that $\frac{\partial}{\partial z_j} M_c(\mathbf{z}) = 0$, for all the negative breakpoints, and that the direction we increase has the form $\mathbf{u} = (-1, \mathbf{u}_1, \dots, 0, \mathbf{u}_{k+2}, 0, \dots, 0)$. Thus, we get a function g that has at most $2k+5$ breakpoints, where we can get an extra breakpoint if 0 is a breakpoint. \square

Proof of Proposition 4.1. For every $\epsilon > 0$, using the function f' from Lemma 4.3 in Lemma 4.4, we can obtain a function f_ϵ such that $|\mathbf{E}_{z \sim \mathcal{N}(0,1)}[f_\epsilon(z)z^t]| \leq \epsilon$, for every non-negative integer $t \leq k$ and $\mathbf{E}_{z \sim \mathcal{N}(0,1)}[f_\epsilon(z)\text{ReLU}(z)] > 1/\text{poly}(k) + O(\epsilon)$. Moreover, the function f_ϵ is at most $(2k+5)$ -piecewise constant.

Let $\mathbf{M} : \overline{\mathbb{R}}^{2k+5} \mapsto \mathbb{R}^{k+2}$, where $M_i(\mathbf{b}) = \sum_{n=0}^{2k+5} (-1)^{n+1} \int_{b_n}^{b_{n+1}} z^i \phi(z) dz$, for $0 \leq i < k+2$, and $M_{k+2}(\mathbf{b}) = \sum_{n=0}^{2k+5} (-1)^{n+1} \int_{b_n}^{b_{n+1}} \text{ReLU}(z) \phi(z) dz$, where $b_0 \leq b_1 \dots \leq b_{2k+5}$, $b_0 = -\infty$ and $b_{2k+6} = \infty$. Here we assume without loss of generality that before the first breakpoint the function is negative, because we can always set the first breakpoint to be $-\infty$. It is clear that the function \mathbf{M} is a continuous map and $\overline{\mathbb{R}}^{2k+5}$ is a compact set, thus $\mathbf{M}(\overline{\mathbb{R}}^{2k+5})$ is a compact set. We also have that for every $\epsilon > 0$, there is a point $\mathbf{b} \in \overline{\mathbb{R}}^{2k+5}$ such that $|\langle \mathbf{M}(\mathbf{b}), \mathbf{e}_i \rangle| \leq \epsilon$, for $0 \leq i < k+2$, and $\langle \mathbf{M}(\mathbf{b}), \mathbf{e}_{k+2} \rangle > 1/\text{poly}(k) + O(\epsilon)$. Thus, from compactness, we have that there exists a point $\mathbf{b}^* \in \overline{\mathbb{R}}^{2k+5}$ such that $|\langle \mathbf{M}(\mathbf{b}^*), \mathbf{e}_i \rangle| = 0$ for $0 \leq i < k+2$, and $\langle \mathbf{M}(\mathbf{b}^*), \mathbf{e}_{k+2} \rangle > 1/\text{poly}(k)$. \square

4.2 Proof of Proposition 4.2

Below we state some important properties of the Legendre polynomials that we use in our proofs.

Fact 4.5 (see, e.g., [Sze89]). *The Legendre polynomials, $P_n(z)$, for n non-negative integer, satisfy the following properties:*

- (i) $P_n(z)$ is a degree- n univariate polynomial, with $P_0(z) = 1$ and $P_1(z) = z$.
- (ii) $\int_{-1}^1 P_i(z)P_j(z)dz = \delta_{ij} \frac{2}{2i+1}$, for all i, j non-negative integers (orthogonality).
- (iii) $|P_n(z)| \leq 1$, for all $|z| \leq 1$ (bounded).
- (iv) $P'_n(z) = \sum_{t=0}^n \frac{2t+1}{2} P_t(z)$ (closed form of derivative).

Using the Legendre polynomials, we can construct a function for which the first $k+1$ moments are zero and which has non-trivial correlation with the ReLU function.

Proof of Proposition 4.2. Define $f(z) = c \frac{\text{ReLU}(z) - p(z)}{\phi(z)} \mathbb{1}\{z \in [-1, 1]\}$, for a degree- k polynomial $p(z)$ and a constant $c > 0$. Then, we have

$$\mathbf{E}_{z \sim \mathcal{N}(0,1)} [f(z)z^t] = c \int_{-1}^1 (\text{ReLU}(z) - p(z))z^t dz .$$

We want $\mathbf{E}_{z \sim \mathcal{N}(0,1)} [f(z)z^t] = 0$, thus we want to find a polynomial $p(z)$ such that

$$\int_{-1}^1 \text{ReLU}(z)z^t dz = \int_{-1}^1 p(z)z^t dz . \quad (8)$$

Equation (8) is equivalent to saying that for all $0 \leq t < k$, it holds

$$\int_{-1}^1 \text{ReLU}(z)P_t(z)dz = \int_{-1}^1 p(z)P_t(z)dz , \quad (9)$$

because the Legendre polynomials of degree at most k span the space of polynomials of degree at most k . Using Fact 4.5 (ii) and a standard computation involving orthogonal polynomials, gives that for $p(z) = \sum_{t=0}^k \frac{2t+1}{2} P_t(z) \int_{-1}^1 \text{ReLU}(z)P_t(z)dz$, Equation (9) and Equation (8) hold. We want the function f to take values inside the interval $[-1, 1]$. To achieve this, we bound from above the constant c . It holds that $\int_{-1}^1 \text{ReLU}(z)P_t(z)dz \leq 2$, where we used Fact 4.5 (iii) and $|\text{ReLU}(z)| \leq 1$ for $|z| \leq 1$. Moreover, we get that

$$|p(z)| \leq 2 \sum_{t=0}^k \frac{2t+1}{2} |P_t(z)| \leq k^2 + 2k \leq 2k^2 ,$$

for all $|z| \leq 1$. Thus, it must hold that $c \leq g(1)/(2k^2 + 1)$, and by taking $c = g(1)/(2k^2 + 1)$, we get that $|f(z)| \leq 1$.

Next we prove that $\mathbf{E}_{z \sim \mathcal{N}(0,1)} [f(z)\text{ReLU}(z)] > 1/\text{poly}(k)$. We have that

$$\mathbf{E}_{z \sim \mathcal{N}(0,1)} [f(z)\text{ReLU}(z)] = c \int_{-1}^1 \text{ReLU}(z)(\text{ReLU}(z) - p(z))dz = c \int_{-1}^1 (\text{ReLU}(z) - p(z))^2 dz ,$$

where we used that $\int_{-1}^1 q(z)(\text{ReLU}(z) - p(z))dz = 0$, for any polynomial q of degree at most k , and thus it holds for $q(z) = p(z)$. Note that $|p'(z)| \leq 5k^4$ and $|p''(z)| \leq 7k^6 =: N$, because from Fact 4.5 (iv), we have that $|P'_n(z)| \leq 2n^2$ and $|P''_n(z)| \leq 4n^4$, for all $|z| \leq 1$. For $\epsilon > 0$ sufficiently small, we then have

$$\int_{-1}^1 (\text{ReLU}(z) - p(z))^2 dz \geq \int_{-\epsilon}^{\epsilon} (\text{ReLU}(z) - p(z))^2 dz .$$

Using the Taylor expansion of p , we get that there exists a linear function L , such that $p(z) = L(z) + O(N\epsilon^2)$, for $|z| \leq \epsilon$. We thus have that

$$\int_{-\epsilon}^{\epsilon} (\text{ReLU}(z) - p(z))^2 dz = \int_{-\epsilon}^{\epsilon} (\text{ReLU}(z) - L(z) + O(N\epsilon^2))^2 dz .$$

Note that every function can be written as $G(z) = G_{\text{even}}(z) + G_{\text{odd}}(z)$, where $G_{\text{even}}(z)$ is the even part of G and $G_{\text{odd}}(z)$ is the odd part. For $\ell > 0$, it holds that

$$\int_{-\ell}^{\ell} G^2(z)dz = \int_{-\ell}^{\ell} (G_{\text{even}}^2(z) + G_{\text{odd}}^2(z) + 2G_{\text{even}}(z)G_{\text{odd}}(z)) dz \geq \int_{-\ell}^{\ell} G_{\text{even}}^2(z)dz ,$$

where we used that $\int_{-\epsilon}^{\epsilon} G_{\text{even}}(z)G_{\text{odd}}(z) = 0$. Using that $\text{ReLU}(z) = |z|/2 + z/2$, it holds

$$\int_{-\epsilon}^{\epsilon} (\text{ReLU}(z) - L(z) + O(N\epsilon^2))^2 dz \geq \int_{-\epsilon}^{\epsilon} (|z|/2 - L(0) + O(N\epsilon^2))^2 dz ,$$

where we used that L is linear, thus the even part is $L(0)$. Choosing ϵ such that $N < \epsilon^{-1}/C$ for a large enough $C > 0$, we have that $||z|/2 - L(0)| \geq \epsilon/8$ for at least half of the interval $[-\epsilon, \epsilon]$. To prove this, note that we have two cases. First, if $L(0) > \epsilon/2$ or $L(0) \leq 0$, this holds trivially. Again in the other case trivially in half the points we have $||z|/2 - L(0)| \geq \epsilon/4$. Moreover, from the choice of ϵ , we have that $N\epsilon^2 \leq \epsilon/C$, thus $||z|/2 - L(0) + O(N\epsilon^2)| \geq ||z|/2 - L(0)| - |O(N\epsilon^2)| \geq \epsilon/8$ for at least half of the interval. Therefore, we have

$$\int_{-\epsilon}^{\epsilon} (|z|/2 - L(0) + O(N\epsilon^2))^2 dz \geq \Omega(\epsilon^3) .$$

By our choice of ϵ , we have

$$c \int_{-1}^1 (\text{ReLU}(z) - p(z))^2 dz \geq c \cdot \Omega(\epsilon^3) \geq c \cdot \Omega(N^{-3}) \geq \Omega(1/k^{20}) .$$

This completes the proof. □

References

- [ABL17] P. Awasthi, M. F. Balcan, and P. M. Long. The power of localization for efficiently learning linear separators with noise. *J. ACM*, 63(6):50:1–50:27, 2017.
- [BFJ⁺94] A. Blum, M. Furst, J. Jackson, M. Kearns, Y. Mansour, and S. Rudich. Weakly learning DNF and characterizing statistical query learning using Fourier analysis. In *Proceedings of the Twenty-Sixth Annual Symposium on Theory of Computing*, pages 253–262, 1994.
- [Dan15] A. Daniely. A PTAS for agnostically learning halfspaces. In *Proceedings of The 28th Conference on Learning Theory, COLT 2015*, pages 484–502, 2015.
- [Dan16] A. Daniely. Complexity theoretic limitations on learning halfspaces. In *Proceedings of the 48th Annual Symposium on Theory of Computing, STOC 2016*, pages 105–117, 2016.
- [DGJ⁺10] I. Diakonikolas, P. Gopalan, R. Jaiswal, R. Servedio, and E. Viola. Bounded independence fools halfspaces. *SIAM Journal on Computing*, 39(8):3441–3462, 2010.
- [DGK⁺20] I. Diakonikolas, S. Goel, S. Karmalkar, A. Klivans, and M. Soltanolkotabi. Approximation schemes for relu regression. In *COLT 2020, to appear*, 2020. Available at <https://arxiv.org/abs/2005.12844>.
- [DKKZ20] I. Diakonikolas, D. M. Kane, V. Kontonis, and N. Zarifis. Algorithms and SQ lower bounds for PAC learning one-hidden-layer relu networks. *CoRR*, abs/2006.12476, 2020. To appear in COLT 2020.
- [DKN10] I. Diakonikolas, D. M. Kane, and J. Nelson. Bounded independence fools degree-2 threshold functions. In *51th Annual IEEE Symposium on Foundations of Computer Science, FOCS 2010*, pages 11–20. IEEE Computer Society, 2010.

- [DKS17] I. Diakonikolas, D. M. Kane, and A. Stewart. Statistical query lower bounds for robust estimation of high-dimensional Gaussians and Gaussian mixtures. In *FOCS*, pages 73–84, 2017.
- [DKS18] I. Diakonikolas, D. M. Kane, and A. Stewart. Learning geometric concepts with nasty noise. In *Proceedings of the 50th Annual ACM SIGACT Symposium on Theory of Computing, STOC 2018*, pages 1061–1073, 2018.
- [Fel09] V. Feldman. A complete characterization of statistical query learning with applications to evolvability. In *Proc. 50th Symposium on Foundations of Computer Science (FOCS)*, pages 375–384, 2009.
- [Fel16] V. Feldman. Statistical query learning. In *Encyclopedia of Algorithms*, pages 2090–2095, 2016.
- [FGKP06] V. Feldman, P. Gopalan, S. Khot, and A. Ponnuswami. New results for learning noisy parities and halfspaces. In *Proc. FOCS*, pages 563–576, 2006.
- [FGR⁺13] V. Feldman, E. Grigorescu, L. Reyzin, S. Vempala, and Y. Xiao. Statistical algorithms and a lower bound for detecting planted cliques. In *Proceedings of STOC’13*, pages 655–664, 2013.
- [FGV17] V. Feldman, C. Guzman, and S. S. Vempala. Statistical query algorithms for mean vector estimation and stochastic convex optimization. In *Proceedings of the Twenty-Eighth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2017*, pages 1265–1277. SIAM, 2017.
- [FS97] Y. Freund and R. Schapire. A decision-theoretic generalization of on-line learning and an application to boosting. *Journal of Computer and System Sciences*, 55(1):119–139, 1997.
- [GGJ⁺20] S. Goel, A. Gollakota, Z. Jin, S. Karmalkar, and A. R. Klivans. Superpolynomial lower bounds for learning one-layer neural networks using gradient descent. *CoRR*, abs/2006.12011, 2020. To appear in ICML 2020.
- [GGK20] S. Goel, A. Gollakota, and A. Klivans. Statistical-query lower bounds via functional gradients. Manuscript, 2020.
- [GKK19] S. Goel, S. Karmalkar, and A. R. Klivans. Time/accuracy tradeoffs for learning a relu with respect to gaussian marginals. In *Advances in Neural Information Processing Systems 32: Annual Conference on Neural Information Processing Systems 2019, NeurIPS 2019*, pages 8582–8591, 2019.
- [GKKT17] S. Goel, V. Kanade, A. R. Klivans, and J. Thaler. Reliably learning the relu in polynomial time. In *Proceedings of the 30th Conference on Learning Theory, COLT 2017*, volume 65 of *Proceedings of Machine Learning Research*, pages 1004–1042. PMLR, 2017.
- [GR06] V. Guruswami and P. Raghavendra. Hardness of learning halfspaces with noise. In *Proc. 47th IEEE Symposium on Foundations of Computer Science (FOCS)*, pages 543–552. IEEE Computer Society, 2006.
- [Hau92] D. Haussler. Decision theoretic generalizations of the PAC model for neural net and other learning applications. *Information and Computation*, 100:78–150, 1992.

- [Kea98] M. J. Kearns. Efficient noise-tolerant learning from statistical queries. *Journal of the ACM*, 45(6):983–1006, 1998.
- [KK14] A. R. Klivans and P. Kothari. Embedding hard learning problems into gaussian space. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques, APPROX/RANDOM 2014*, pages 793–809, 2014.
- [KKMS08] A. Kalai, A. Klivans, Y. Mansour, and R. Servedio. Agnostically learning halfspaces. *SIAM Journal on Computing*, 37(6):1777–1805, 2008.
- [KLS09] A. Klivans, P. Long, and R. Servedio. Learning halfspaces with malicious noise. To appear in *Proc. 17th Internat. Colloq. on Algorithms, Languages and Programming (ICALP)*, 2009.
- [KSS94] M. Kearns, R. Schapire, and L. Sellie. Toward Efficient Agnostic Learning. *Machine Learning*, 17(2/3):115–141, 1994.
- [MR18] P. Manurangsi and D. Reichman. The computational complexity of training relu (s). *arXiv preprint arXiv:1810.04207*, 2018.
- [MT94] W. Maass and G. Turan. How fast can a threshold gate learn? In S. Hanson, G. Drastal, and R. Rivest, editors, *Computational Learning Theory and Natural Learning Systems*, pages 381–414. MIT Press, 1994.
- [Ros58] F. Rosenblatt. The Perceptron: a probabilistic model for information storage and organization in the brain. *Psychological Review*, 65:386–407, 1958.
- [Sol17] M. Soltanolkotabi. Learning relus via gradient descent. In *Advances in neural information processing systems*, pages 2007–2017, 2017.
- [Sze89] G. Szegő. *Orthogonal Polynomials*, volume XXIII of *American Mathematical Society Colloquium Publications*. A.M.S, Providence, 1989.
- [Szö09] B. Szörényi. Characterizing statistical query learning: simplified notions and proofs. In *International Conference on Algorithmic Learning Theory*, pages 186–200. Springer, 2009.
- [Val84] L. Valiant. A theory of the learnable. *Communications of the ACM*, 27(11):1134–1142, 1984.
- [Vap98] V. Vapnik. *Statistical Learning Theory*. Wiley-Interscience, New York, 1998.