arXiv:2006.16436v2 [math.SG] 20 Jan 2023

AUGMENTATIONS AND IMMERSED LAGRANGIAN FILLINGS

YU PAN AND DAN RUTHERFORD

ABSTRACT. For a Legendrian link $\Lambda \subset J^1M$ with $M = \mathbb{R}$ or S^1, immersed exact Lagrangian fillings $L \subset \text{Symp}(J^1M) \cong T^*(\mathbb{R}_{>0} \times M)$ of Λ can be lifted to conical Legendrian fillings $\Sigma \subset J^1(\mathbb{R}_{>0} \times M)$ of Λ. When Σ is embedded, using the version of functoriality for Legendrian contact homology (LCH) from [32], for each augmentation $\alpha : \mathcal{A}(\Sigma) \to \mathbb{Z}/2$ of the LCH algebra of Σ, there is an induced augmentation $\epsilon_{(\Sigma,\alpha)} : \mathcal{A}(\Lambda) \to \mathbb{Z}/2$. With Σ fixed, the set of homotopy classes of all such induced augmentations, $I_\Sigma \subset Aug(\Lambda)/\sim$, is a Legendrian isotopy invariant of Σ. We establish methods to compute I_Σ based on the correspondence between Morse complex families and augmentations. This includes developing a functoriality for the cellular DGA from [33] with respect to Legendrian cobordisms, and proving its equivalence to the functoriality for LCH. For arbitrary $n \geq 1$, we give examples of Legendrian torus knots with $2n$ distinct conical Legendrian fillings distinguished by their induced augmentation sets. We prove that when $\rho \neq 1$ and $\Lambda \subset J^1\mathbb{R}$ *every* ρ-graded augmentation of Λ can be induced in this manner by an immersed Lagrangian filling. Alternatively, this is viewed as a computation of cobordism classes for an appropriate notion of ρ-graded augmented Legendrian cobordism.

CONTENTS

1. Introduction 1
2. Immersed DGA maps and cobordisms 5
3. Immersed DGA maps and augmentations 15
4. Immersed maps and the cellular DGA 19
5. Computations via Morse complex families 29
6. Examples 34
7. Every augmentation is induced by an immersed filling 38
References 50

1. INTRODUCTION

A fundamental holomorphic curve invariant of a Legendrian submanifold, Λ, is the Legendrian contact homology (LCH) dg-algebra (DGA), denoted $\mathcal{A}(\Lambda)$. As part of the symplectic field theory package, the LCH algebra is functorial for an appropriate class of cobordisms. In this article we consider 1-dimensional Legendrian links in the 1-jet spaces, J^1M with $M = \mathbb{R}$ or S^1, and exact Lagrangian cobordisms in the symplectization, $Symp(J^1M) = \mathbb{R} \times J^1M$; throughout, our coefficient field is $\mathbb{Z}/2$. For $\Lambda_-, \Lambda_+ \subset J^1M$, such a cobordism, $L : \Lambda_- \to \Lambda_+$, cylindrical over Λ_- and Λ_+ at the negative and positive ends of $Symp(J^1M)$, equipped with a $\mathbb{Z}/\rho$-valued Maslov potential induces a $\mathbb{Z}/\rho$-graded DGA map $f_L : \mathcal{A}(\Lambda_+) \to \mathcal{A}(\Lambda_-)$, cf. [13, 16]. In particular, when L is an exact Lagrangian filling, i.e. a cobordism $L : \emptyset \to \Lambda$, the induced map

$$\epsilon_L : \mathcal{A}(\Lambda) \to \mathbb{Z}/2$$

is a **ρ-graded augmentation** which by definition is a unital ring homomorphism that satisfies $\epsilon_L \circ \partial = 0$ and preserves a $\mathbb{Z}/\rho$-grading on $\mathcal{A}(\Lambda)$.

A natural question is:

Question 1.1. Which augmentations come from exact Lagrangian fillings?

While orientable exact Lagrangian fillings have been constructed for several classes of Legendrian knots [16, 22, 38, 40], there are many augmentations that cannot be induced by any orientable filling as obstructions to such fillings arise from the Thurston-Bennequin number of Λ and from the linearized homology of the augmentation; see [7, 13, 11]. The main result of this article shows that if one extends the setting to allow immersed cobordisms with double points then the algebra more closely matches the geometry. Indeed we prove that when ρ is even, *every ρ-graded augmentation can be induced by an orientable immersed exact Lagrangian filling.*

An extension of the functoriality for LCH to immersed Lagrangian cobordisms is implemented in [32] by working with a class of Legendrian cobordisms as follows. Applying a symplectomorphism $Symp(J^1M) \cong T^*(\mathbb{R}_{>0} \times M)$ an exact immersed Lagrangian cobordism, $L \subset Symp(J^1M)$, can be lifted to a Legendrian $\Sigma \subset J^1(\mathbb{R}_{>0} \times M)$ with the cylindrical ends of L translating to conical ends for Σ and double points of L becoming Reeb chords of Σ. See Section 2.4. When Σ is embedded and equipped with a $\mathbb{Z}/\rho$-valued Maslov potential, such a **conical Legendrian cobordism**, $\Sigma : \Lambda_- \to \Lambda_+$, induces a diagram of ρ-graded DGA maps

$$\mathcal{A}(\Lambda_+) \xrightarrow{f_\Sigma} \mathcal{A}(\Sigma) \xleftarrow{i_\Sigma} \mathcal{A}(\Lambda_-) \tag{1.1}$$

where $\mathcal{A}(\Sigma)$ is generated by $\mathcal{A}(\Lambda_-)$ and the Reeb chords of Σ. Diagrams of the above form are referred to in [32] as **immersed DGA maps**. There, a notion of homotopy for immersed DGA maps is introduced, and the homotopy type of the immersed map (1.1) is shown to be an invariant of the conical Legendrian isotopy type of Σ. When $\Sigma : \emptyset \to \Lambda$ is a conical Legendrian filling equipped with a choice of ρ-graded augmentation, $\alpha : \mathcal{A}(\Sigma) \to \mathbb{Z}/2$, we can then define an **induced augmentation** $\epsilon_{(\Sigma,\alpha)} : \mathcal{A}(\Lambda) \to \mathbb{Z}/2$ as the composition $\epsilon_{(\Sigma,\alpha)} = \alpha \circ f_\Sigma$. This generalizes the construction of induced augmentations from embedded Lagrangian fillings.

We can now state our first main result, where in the following we write $\epsilon \simeq \epsilon'$ to indicate that two augmentations are DGA homotopic.

Theorem 1.2. *Let $\Lambda \subset J^1\mathbb{R}$ have the $\mathbb{Z}/\rho$-valued Maslov potential μ where $\rho \geq 0$, and let $\epsilon : \mathcal{A}(\Lambda) \to \mathbb{Z}/2$ be any ρ-graded augmentation.*

1. *If $\rho \neq 1$, there exists a conical Legendrian filling Σ of Λ with $\mathbb{Z}/\rho$-valued Maslov potential extending μ together with a ρ-graded augmentation $\alpha : \mathcal{A}(\Sigma) \to \mathbb{Z}/2$ such that $\epsilon \simeq \epsilon_{(\Sigma,\alpha)}$. Moreover, if ρ is even, then Σ is orientable.*
2. *If $\rho = 1$, then there exists a conical Legendrian cobordism $\Sigma : U \to \Lambda$ where U is the standard Legendrian unknot with $tb(U) = -1$ together with a 1-graded augmentation $\alpha : \mathcal{A}(\Sigma) \to \mathbb{Z}/2$ such that $\epsilon \simeq \alpha \circ f_\Sigma$.*

The algebra of the standard Legendrian unknot, U, is generated by a single Reeb chord b of degree 1. In the case where $\rho = 1$, if the restriction of α to $\mathcal{A}(U) \subset \mathcal{A}(\Sigma)$ sends b to 0, then by concatenating with the standard filling of U we see that ϵ can be induced by a pair (Σ', α') where Σ' is a conical Legendrian filling. However, in the case that α restricts to the augmentation of $\mathcal{A}(U)$ that maps b to 1, we are not sure whether ϵ can be induced by a Legendrian filling. In fact, we conjecture that this is not possible.

Conjecture 1.3. *There is no conical Legendrian filling Σ of the Legendrian unknot U with 1-graded augmentation $\alpha : \mathcal{A}(\Sigma) \to \mathbb{Z}/2$ such that the induced augmentation $\epsilon_{(\Sigma,\alpha)}$ maps the unique Reeb chord b to 1.*

For the case of embedded Lagrangians, induced augmentations provide an effective means of distinguishing Lagrangian fillings of a given Legendrian knot. The results of [32] show that these induced augmentations are actually invariants of the associated conical Legendrian fillings. (Note that conical Legendrian isotopy appears to be a much less restrictive notion of equivalence than isotopy of the corresponding exact Lagrangians, since during the course of a Legendrian isotopy any number of double points can be added and removed from the Lagrangian projections.)

A key new feature in the immersed case is that a single Legendrian filling, Σ, can induce more than one augmentation of Λ due to the dependence of the construction on the choice of augmentation α of $\mathcal{A}(\Sigma)$. As a result, to obtain an invariant of Σ we should consider the *set* of (DGA homotopy classes of) induced augmentations $I_\Sigma \subset Aug_\rho(\Lambda)/\sim$. After developing methods to compute this invariant **induced augmentation set**, we demonstrate that I_Σ can be effective for distinguishing conical Legendrian fillings in the following theorem.

Theorem 1.4. *For each $n \geq 1$, there exists $2n$ distinct conical Legendrian fillings, $\Sigma_1, \dots, \Sigma_{2n}$ of the max-tb Legendrian torus knot $T(2, 2n+1)$ such that*

(i) *the Σ_i are all orientable with genus $n-1$ and have $\mathbb{Z}$-valued Maslov potentials,*
(ii) *each Σ_i has a single Reeb chord of degree 0,*
(iii) *and the induced augmentation sets satisfy $I_{\Sigma_i} \neq I_{\Sigma_j}$ when $i \neq j$.*

Note that the Lagrangian projections $L_1, \dots, L_n$ are immersed Lagrangian fillings of $T(2, 2n+1)$ with a single double point. Moreover, the dg-algebras $\mathcal{A}(\Sigma_i)$ are all isomorphic to one another, and thus do not distinguish the Σ_i on their own.

1.1. **Cobordism classes of augmented Legendrians.** In the context of relative symplectic field theory (SFT) type invariants [20], it is natural to consider Lagrangian cobordisms between Legendrian submanifolds in the symplectization of a contact manifold (or more generally in symplectic cobordisms with concave and convex ends modeled on the negative and positive ends of symplectizations). In the following discussion, we refer to exact Lagrangian cobordisms in $Symp(J^1M)$ with cylindrical ends as **SFT-cobordisms**. The relation defined by embedded SFT-cobordisms, or even SFT-concordances, is not symmetric [8, 29], and hence does not define an equivalence relation on Legendrian links in J^1M. In fact, it is a major open question [10] in the field whether or not SFT-cobordisms define a partial order on the set of Legendrian isotopy classes in $J^1\mathbb{R}$.

The lack of a readily available symmetry is visible after lifting an SFT-cobordism to a conical Legendrian cobordism $\Sigma \subset J^1(\mathbb{R}_{>0} \times M) = T^*(\mathbb{R}_{>0} \times M) \times \mathbb{R}_z$ in the difference in behavior at the two ends of Σ: as Σ approaches 0 (resp. $+\infty$) along the $\mathbb{R}_{>0}$ factor the cotangent coordinates and the z-coordinates of Σ appear as those of Λ_- (resp. Λ_+) but shrinking (resp. expanding). However, when one allows for SFT-cobordisms to be immersed it becomes possible to reverse their direction as an expanding end can be modified to a shrinking end (and vice-versa) at the expense of creating some additional Reeb chords. Thus, the relation of conical Legendrian cobordism is equivalent to another standard notion of Legendrian cobordism in 1-jet spaces introduced by Arnold, cf. [1, 2, 3]. In Arnold's definition, a Legendrian cobordism between two Legendrians $\Lambda_0, \Lambda_1 \subset J^1M$ is a compact Legendrian, $\Sigma \subset J^1([0,1] \times M)$, whose restriction to $J^1(\{i\} \times M)$ is Λ_i. Seminal results in this theory of Legendrian (and also Lagrangian) cobordisms were achieved by Audin, Eliashberg, and Vassiliev in the 1980's, including homotopy theoretic characterizations of various Lagrange and Legendre cobordism groups. See [18, 4, 41].

For an alternate perspective on our main theorem, we can incorporate augmentations into an Arnold-type cobordism theory. Define a **ρ-graded augmented Legendrian** to be a Legendrian submanifold $\Lambda \subset J^1M$ equipped with a $\mathbb{Z}/\rho$-valued Maslov potential and a ρ-graded augmentation $\epsilon : \mathcal{A}(\Lambda) \to \mathbb{Z}/2$. For compact Legendrian cobordisms $\Sigma \subset J^1([0,1] \times M)$ satisfying a suitable Morse minimum boundary condition (see Section 2.5), the LCH dg-algebra of Σ is defined as in [17], and contains $\mathcal{A}(\Lambda_0)$ and $\mathcal{A}(\Lambda_1)$ as sub-dg-algebras. We then declare two ρ-graded augmentation Legendrians, (Λ_0, ϵ_0) and (Λ_1, ϵ_1), to be cobordant if there exists a pair (Σ, α) consisting of a Legendrian cobordism equipped with a ρ-graded augmentation of $\mathcal{A}(\Sigma)$ whose restriction to $\mathcal{A}(\Lambda_i)$ is homotopic to ϵ_i for $i = 0, 1$. As a variant on Theorem 1.2 we obtain the following.

Theorem 1.5. *Let $\rho \geq 0$ be a non-negative integer, and let (Λ, ϵ) be a ρ-graded augmented Legendrian in $J^1\mathbb{R}$.*

(1) *If* $\rho \neq 1$, *then* (Λ, ϵ) *is cobordant to* $\emptyset$.
(2) *If* $\rho = 1$, *then* (Λ, ϵ) *is either cobordant to* $\emptyset$ *or* (U, ϵ_1) *where* U *is the standard Legendrian unknot and* $\epsilon_1 : \mathcal{A}(U) \to \mathbb{Z}/2$ *satisfies* $\epsilon_1(b) = 1$ *on the unique Reeb chord of* U.

As in Conjecture 1.3, we expect that (U, ϵ_1) is not null-cobordant, so that there should be exactly one cobordism class when $\rho \neq 1$ and exactly two cobordism classes when $\rho = 1$.

In a further article [31], we give a complete classification of cobordism classes of augmented Legendrians in J^1S^1. Interestingly, in J^1S^1 it is often the case that cobordant Legendrians may become non-cobordant once they are equipped with augmentations. We note that for long Legendrian knots in $\mathbb{R}^3$ a concordance group of similar spirit, incorporating quadratic at infinity generating families as additional equipment rather than augmentations, is considered recently in the work of Limouzineau, [26].

1.2. **Methods and outline.** Our approach is based on working with certain algebraic/combinatorial structures equivalent to augmentations called Morse complex families (MCFs) that were introduced by Pushkar (unpublished) and studied by Henry in [23]. MCFs can be viewed as combinatorial approximations to generating families of functions, as they consist of formal Morse complexes and handleslide data assigned to a Legendrian submanifold. For 1-dimensional Legendrians, the work of Henry [23, 25] establishes a bijection between equivalence classes of MCFs and homotopy classes of augmentations. More recently, a correspondence between MCFs and augmentations for 2-dimensional Legendrians was obtained in [36] using the cellular DGA which is a cellular model for LCH developed in the work of the second author and Sullivan [33, 34, 35]. The strategy of the present article is to extend these methods to the case of Legendrian cobordisms, and then apply Morse complex families to compute induced augmentation sets. In more detail, to prove Theorem 1.2 we accomplish the following tasks (1)-(3). Note that (1) and (2) may be of some independent interest.

(1) We develop a functoriality of the cellular DGA for Legendrian cobordisms, and we extend the equivalence with LCH from [33, 34, 35] to an isomorphism between the immersed LCH functor from [32] and its cellular analog; see Proposition 4.9.

Although it seems crucial for this construction to be carried out in the broader setting of Legendrian cobordisms and immersed DGA maps, this allows the cellular DGA to be applied just as well for working with embedded SFT-cobordisms.

(2) We extend the correspondence between MCFs and augmentations to the case of 2-dimensional Legendrian cobordisms. In particular, we give a method based on MCFs for computing the induced augmentation sets of Legendrian cobordisms and fillings; see Proposition 5.9.

In Section 6, we illustrate this method with examples and use it to prove Theorem 1.4. Making use of (1) and (2), Theorem 1.2 is then reduced to the following construction.

(3) Given a 1-dimensional Legendrian knot, $\Lambda \subset J^1\mathbb{R}$, with a ρ-graded MCF $\mathcal{C}_\Lambda$ with $\rho \neq 1$ (resp. $\rho = 1$), we produce a Legendrian filling Σ of Λ (resp. a Legendrian cobordism $\Sigma : U \to \Lambda$) with a 2-dimensional ρ-graded MCF $\mathcal{C}_\Sigma$ extending $\mathcal{C}_\Lambda$.

In Section 7 we provide such a construction via an induction on the complexity of the front projection of Λ. The resulting cobordism is built up out of elementary building blocks that include those used in constructing decomposable, embedded Lagrangian cobordisms (Legendrian isotopies, pinch and unknot moves) as well as clasp moves that produce double points in the Lagrangian projection. Care is taken to ensure that at each step the MCF can be extended over the cobordism, and these considerations are simplified by making use of a standard form for MCFs (the "SR-form") introduced by Henry.

Remark 1.6. For embedded, decomposable SFT cobordisms, explicit computations of induced DGA maps and in particular induced augmentations are possible via the results of [16], and this approach has been commonly used in the literature. A possible alternate route to proving Theorem 1.2 could be to extend the arsenal from [16] to include computations of the immersed DGA map associated to the Clasp Move and attempt a similar inductive construction in this context. Such a computation would be useful more generally for handling immersed Lagrangian cobordisms within the framework of [16]. We leave this as an interesting direction for possible future research, while noting that the approach taken in the current article and [32] is independent of [16] in both the foundations (based on [14, 12]) and computational methods (based on [33, 36]).

The remainder of the paper is organized as follows. In Section 2 we collect background from [32] on immersed DGA maps, conical Legendrian cobordisms, and the immersed LCH functor, F. In addition, we discuss Morse minimum cobordisms and show that Theorem 1.5 follows from Theorem 1.2. In Section 3, we establish some basic properties of induced augmentation sets associated to immersed DGA maps including a composition formula and Legendrian isotopy invariance. In Section 4, we review the cellular DGA and define a cellular LCH functor, F_{cell}. We state an isomorphism between the cellular LCH functor and the immersed LCH functor. The proof of the isomorphism is an extension of [34, 35] that is sketched here with a more detailed argument appearing in the preprint version of this article [30]. In Section 5, after reviewing the definition of Morse complex families, we extend the correspondence between MCFs and augmentations of the cellular DGA to the cobordism case and establish an MCF characterization of the induced augmentation set. In Section 6, we make use of A-form MCFs to compute the induced augmentation sets for several examples of Legendrian fillings, and we prove Theorem 1.4. In particular, we give explicit examples of augmentations that can be induced by Legendrian fillings but cannot be induced by any embedded fillings. Finally, the article concludes with Section 7 that, after recalling Henry's SR-form MCFs, provides the proof of Theorem 1.2. The reader that is content to blackbox the characterization of the induced augmentation set in terms of MCFs can more quickly arrive at the proof of Theorem 1.2 by omitting Sections 4, 5.3, and 6.

Remark 1.7. In the present article we restrict considerations to $\mathbb{Z}/2$ valued augmentations. This is mainly because the isomorphism between the cellular DGA and LCH DGA proven in [33, 34, 35] is only established with $\mathbb{Z}/2$ coefficients. We expect that a version of this isomorphism should extend to a more general coefficient ring, eg. to the setting of the LCH DGA with fully non-commutative $\mathbb{Z}[\pi_1(\Lambda)]$ coefficients which is appropriate for considering augmentations to general rings. Assuming this point, the proof of Theorem 1.2 should be extendable to apply to augmentations with values in an arbitrary *field*; see also the discussion on coefficients in [31, Section 3.3]. The proof breaks down in the case of augmentations to a general ring, as the ability to take multiplicative inverses of handleslide coefficients is crucial in extending the arguments in Section 7.

1.3. **Acknowledgements.** DR is partially supported by grant 429536 from the Simons Foundation. YP is partially supported by the NSF Grant (DMS-1510305).

2. Immersed DGA maps and cobordisms

In this section we review the algebraic and geometric setup from [32], and discuss two classes of Legendrian cobordisms. Section 2.1 recalls a class of DGAs relevant for Legendrian contact homology, and records in Proposition 2.1 a method for producing stable tame isomorphisms. In Section 2.2 we review the concept of immersed DGA maps and immersed homotopy then discuss two equivalent characterizations of composition. In addition, we recall the category,

$\mathfrak{DGA}^{\rho}_{im}$, of $\mathbb{Z}/\rho$-graded DGAs with immersed maps constructed in [32], and its connection with the ordinary homotopy category of DGAs, $\mathfrak{DGA}^{\rho}_{hom}$; see Proposition 2.7.

Next, we turn to the geometric side with a brief review of Legendrians and LCH in the setting of 1-jet spaces in 2.3. Section 2.4 discusses conical Legendrian cobordisms and, after observing their connection with immersed Lagrangian cobordisms, recalls the construction of immersed DGA maps from conical Legendrian cobordisms as in [32]. The construction is nicely encoded in a functor, F, called the *immersed LCH functor* that plays a central role in this article. Finally, Section 2.5 discusses Morse minimum Legendrian cobordisms, connects them with conical Legendrian cobordisms, and provides a proposition that allows for Morse minimum cobordisms to be used for computations with the immersed LCH functor. The section concludes by defining the equivalence relation of ρ-graded augmented Legendrian cobordism and showing that Theorem 1.5 follows from Theorem 1.2.

2.1. **Differential graded algebras.** We work in the algebraic context of [32, Sections 2 and 3] that we will now briefly review. In this article, **differential graded algebras** (abbr. **DGAs**), $(\mathcal{A}, \partial)$, are defined over $\mathbb{Z}/2$ and are graded by $\mathbb{Z}/\rho$ for some fixed $\rho \geq 0$. The differential, ∂, has degree -1 (mod ρ). We restrict attention to **based DGAs** where the $\mathbb{Z}/2$-algebra $\mathcal{A} = \mathbb{Z}/2\langle x_1, \ldots, x_n\rangle$ is free associative (non-commutative) with identity element and is equipped with a choice of (finite) free generating set $\{x_1, \ldots, x_n\}$; generators have degrees $|x_i| \in \mathbb{Z}/\rho$. Subalgebras (resp. 2-sided ideals) generated by a subset $Y \subset \mathcal{A}$ are notated as $\mathbb{Z}/2\langle Y\rangle$ (resp. $\mathcal{I}(Y)$). Such a DGA $(\mathcal{A}, \partial)$ is **triangular** if (with respect to some ordering of the generating set) we have $\partial x_i \in \mathbb{Z}/2\langle x_1, \ldots, x_{i-1}\rangle$ for all $1 \leq i \leq n$. The coproduct of based DGAs, $\mathcal{A} * \mathcal{B}$, is the based DGA whose generating set is the union of the generating sets of $\mathcal{A}$ and $\mathcal{B}$ and whose differential extends the differentials of $\mathcal{A}$ and $\mathcal{B}$. A **stabilization** of a DGA $\mathcal{A}$ is a DGA of the form $\mathcal{A} * S$ where S has generating set of the form $\{a_1, b_1, \ldots, a_r, b_r\}$ with differential $\partial a_i = b_i$, $1 \leq i \leq r$. **DGA morphisms** are unital, algebra homomorphisms that preserve the $\mathbb{Z}/\rho$-grading and commute with differentials. A **stable tame isomorphism** from $\mathcal{A}$ to $\mathcal{B}$ is a DGA isomorphism $\varphi : \mathcal{A} * S \to \mathcal{B} * S'$ between stabilizations of $\mathcal{A}$ and $\mathcal{B}$ that is tame, i.e. it is a composition of isomorphisms that have a certain form on generators; see [32, Section 2.2]. We say DGAs are **equivalent** if they are stable tame isomorphic. Two DGA maps $f, g : \mathcal{A} \to \mathcal{B}$ are **DGA homotopic** if they satisfy

$$f - g = \partial_{\mathcal{B}} \circ K + K \circ \partial_{\mathcal{A}}$$

for some (f, g)-derivation, $K : \mathcal{A} \to \mathcal{B}$, where an (f, g)**-derivation** is a degree 1 (mod ρ) linear map satisfying $K(xy) = K(x)g(y) + (-1)^{|x|}f(x)K(y)$. When $\varphi : \mathcal{A} * S \to \mathcal{B} * S'$ is a stable isomorphism there is an **associated DGA homotopy equivalence** $h : \mathcal{A} \to \mathcal{B}$ given by $h = \pi' \circ h \circ \iota$ where $\iota : \mathcal{A} \to \mathcal{A} * S$ and $\pi' : \mathcal{B} * S' \to \mathcal{B}$ are inclusion and projection.

The following proposition is contained in [32, Propositions 2.3 and 2.5].

Proposition 2.1. *Let $(\mathcal{A}, \partial)$ be a DGA that is triangular with respect to the ordered generating set $\{x_1, \ldots, x_n\}$, and suppose that $\partial x_i = x_j + w$ where $w \in \mathbb{Z}/2\langle x_1, \ldots, x_{j-1}\rangle$.*

(1) *Then, $\mathcal{A}/I$ where $I = \mathcal{I}(x_i, \partial x_i)$ is a triangular DGA with respect to the generating set $\{\overline{x_1}, \ldots, \widehat{\overline{j}}, \ldots, \widehat{\overline{i}}, \ldots, \overline{x_n}\}$, and there is a stable tame isomorphism*

$$\varphi : \mathcal{A}/I * S \to \mathcal{A}$$

with $S = \mathbb{Z}/2\langle y, z\rangle$. Moreover, $\varphi = g * h$ where $h(y) = x_i$, $h(z) = \partial x_i$, and $g : \mathcal{A}/I \to \mathcal{A}$ is DGA homotopy inverse to the quotient map $p : \mathcal{A} \to \mathcal{A}/I$ via

$$g \circ p - id_{\mathcal{A}} = \partial \circ H + H \circ \partial$$

where $H : \mathcal{A} \to \mathcal{A}$ is the $(g \circ p, id_{\mathcal{A}})$-derivation satisfying $H(x_j) = x_i$ and $H(x_l) = 0$ for $l \neq j$.

(2) *If $\mathcal{B} \subset \mathcal{A}$ is a based sub-DGA generated by $Y \subset \{x_1, \dots, x_n\}$ and $x_i, x_j \notin Y$, then $\varphi(\overline{x_k}) = x_k$ for all $x_k \in Y$.*

(3) *For φ^{-1}, the associated DGA homotopy equivalence $\mathcal{A} \xrightarrow{\varphi^{-1}} \mathcal{A}/I * S \xrightarrow{\pi} \mathcal{A}/I$ has $\pi \circ \varphi^{-1} = p$ where $p : \mathcal{A} \to \mathcal{A}/I$ is the quotient map.*

Definition 2.2. A **ρ-graded augmentation** to $\mathbb{Z}/2$ of a $\mathbb{Z}/\rho$-graded DGA, $(\mathcal{A}, \partial)$, is a DGA morphism $\epsilon : (\mathcal{A}, \partial) \to (\mathbb{Z}/2, 0)$ where the grading on $\mathbb{Z}/2$ is concentrated in degree 0 mod ρ. I.e., ϵ is a $\mathbb{Z}/2$-algebra homomorphism from $\mathcal{A}$ to $\mathbb{Z}/2$ that satisfies

$$\epsilon \circ \partial = 0, \quad \epsilon(1) = 1,$$

and $\epsilon(x) \neq 0$ implies that $|x| = 0 \in \mathbb{Z}/\rho$.

2.2. **Immersed DGA maps.** The main construction of [32] extends the functoriality of the Legendrian contact homology DGA to the case where the domain category consists of Legendrians with a class of *immersed* Lagrangian cobordisms. To accomplish this it is natural to also enlarge the class of morphisms in the target category of DGAs, and this is done by introducing *immersed* DGA maps with a suitable notion of homotopy. Here, we recall these notions which will be central in the remainder of the article.

Definition 2.3. Let $(\mathcal{A}_1, \partial_1)$ and $(\mathcal{A}_2, \partial_2)$ be triangular DGAs. An **immersed DGA map**, M, from $\mathcal{A}_1$ to $\mathcal{A}_2$ is a diagram of DGA maps


$$M = \left(\mathcal{A}_1 \xrightarrow{f} \mathcal{B} \xhookleftarrow{i} \mathcal{A}_2\right)$$

where $\mathcal{B}$ is a triangular DGA and f and i are DGA maps such that i is an inclusion induced by an inclusion of the generating set of $\mathcal{A}_2$ into the generating set of $\mathcal{B}$.

Two immersed DGA maps $M = \left(\mathcal{A}_1 \xrightarrow{f} \mathcal{B} \xhookleftarrow{i} \mathcal{A}_2\right)$ and $M' = \left(\mathcal{A}_1 \xrightarrow{f'} \mathcal{B}' \xhookleftarrow{i'} \mathcal{A}_2\right)$ are **immersed homotopic** if there exists a stable tame isomorphism $\varphi : \mathcal{B} * S \to \mathcal{B} * S'$ such that

- $\varphi \circ i = i'$ and
- $\varphi \circ f \simeq f'$ (DGA homotopy).


That is, the left (resp. right) half of the diagram



is commutative up to DGA homotopy (resp. fully commutative).

Definition 2.4. The **composition** $M_2 \circ M_1$ of two immersed DGA maps $M_k = \left(\mathcal{A}_k \xrightarrow{f_k} \mathcal{B}_k \xhookleftarrow{i_k} \mathcal{A}_{k+1}\right)$, for $k = 1, 2$ is given by $M_2 \circ M_1 = \left(\mathcal{A}_1 \xrightarrow{f} \mathcal{B} \xhookleftarrow{i} \mathcal{A}_3\right)$, where $\mathcal{B} = \mathcal{B}_1 * \mathcal{B}_2 / \mathcal{I}(\{i_1(x) - f_2(x) \,|\, x \in \mathcal{A}_2\})$ is the categorical push out of i_1 and f_2 and $f = p_1 \circ f_1$, $i = p_2 \circ i_2$ with $p_k : \mathcal{B}_k \to \mathcal{B}$ the projection maps. The algebra $\mathcal{B}$ is triangular with respect to the generating set obtained as the union of the generators of $\mathcal{B}_1$ and $\mathcal{B}_2$ with the generators of $\mathcal{A}_2$ removed.

This is summarized by the diagram:



There is an alternate characterization of composition of immersed maps, up to immersed homotopy, as follows. Again, let $M_k = \left(\mathcal{A}_k \overset{f_k}{\to} \mathcal{B}_k \overset{i_k}{\hookleftarrow} \mathcal{A}_{k+1}\right)$, $k = 1, 2$ be immersed DGA maps. Let $a_1, \ldots, a_m$ denote the generators of $\mathcal{A}_2$, and let $\widehat{\mathcal{A}}_2$ have free generating set $\widehat{a}_1, \ldots, \widehat{a}_m$ with degree shift $|\widehat{a}_i| = |a_i| + 1$.

Proposition 2.5 (Proposition 3.6 of [32])**.** *Suppose that $(\mathcal{D}, \partial)$ is a triangular DGA such that*

- $\mathcal{D} = \mathcal{B}_1 * \widehat{\mathcal{A}}_2 * \mathcal{B}_2$,
- *for $k = 1, 2$, $\partial|_{\mathcal{B}_k} = \partial_{\mathcal{B}_k}$, and*
- *for $1 \leq i \leq m$,*

$$\partial \widehat{a}_i = i_1(a_i) + f_2(a_i) + \gamma_i \tag{2.1}$$

where $\gamma_i \in \mathcal{I}(\widehat{a}_1, \ldots, \widehat{a}_m)$.

Then, there is an immersed homotopy

$$M_2 \circ M_1 \simeq \left(\mathcal{A}_1 \overset{j_1 \circ f_1}{\to} \mathcal{D} \overset{j_2 \circ i_2}{\hookleftarrow} \mathcal{A}_3\right)$$

where $j_k : \mathcal{B}_k \to \mathcal{B}_1 * \widehat{\mathcal{A}}_2 * \mathcal{B}_2$ is the inclusion.

Note that differentials of the form (2.1) always exist on $\mathcal{D}$. For instance, one can define

$$\partial \widehat{a}_i = i_1(a_i) + f_2(a_i) + \Gamma \circ \partial a_i \tag{2.2}$$

where $\Gamma : \mathcal{A}_2 \to \mathcal{D}$ is the unique (i_1, f_2)-derivation satisfying $\Gamma(a_i) = \widehat{a}_i$.

Definition 2.6. For $\rho \in \mathbb{Z}_{\geq 0}$ fixed, we define a category $\mathfrak{DGA}^{\rho}_{im}$ whose objects are triangular DGAs graded by $\mathbb{Z}/\rho$ and whose morphisms are immersed homotopy classes of immersed DGA maps. Define a related category $\mathfrak{DGA}^{\rho}_{hom}$ to have the same objects as $\mathfrak{DGA}^{\rho}_{im}$ but with morphisms given by DGA homotopy classes of ordinary DGA maps.

Proposition 2.7 (Propositions 3.9 and 3.10 of [32])**.** *For any $\rho \in \mathbb{Z}_{\geq 0}$, $\mathfrak{DGA}^{\rho}_{im}$ is a category with identity morphisms given by the homotopy classes of the immersed maps*

$$\left(\mathcal{A} \overset{id}{\to} \mathcal{A} \overset{id}{\hookleftarrow} \mathcal{A}\right).$$

Moreover, there is a functor $I : \mathfrak{DGA}^{\rho}_{hom} \to \mathfrak{DGA}^{\rho}_{im}$ that is the identity on objects and has

$$I\left([\mathcal{A}_1 \overset{f}{\to} \mathcal{A}_2]\right) = \left[\left(\mathcal{A}_1 \overset{f}{\to} \mathcal{A}_2 \overset{id}{\hookleftarrow} \mathcal{A}_2\right)\right],$$

and I is injective on all hom-spaces.

We emphasize that the triangularity condition is used crucially in [32] in establishing the well-definedness of compositions in $\mathfrak{DGA}^{\rho}_{im}$.

2.3. **The Legendrian contact homology DGA and exact Lagrangian cobordisms.** Recall that the 1-jet space, J^1E, of an n-dimensional manifold, E, has its standard contact structure $\xi = \ker \alpha$ where $\alpha = dz - \sum y_i dx_i$ in coordinates $(x_1, \dots, x_n, y_1, \dots, y_n, z) \in T^*E \times \mathbb{R} = J^1E$ arising from local coordinates $(x_1, \dots, x_n)$ on E. A **Legendrian submanifold**, Λ, is an n-dimensional submanifold that is tangent to ξ everywhere. In this article, we will only need to consider Legendrian submanifolds of dimension $n = 1$ or 2, i.e. Legendrian knots and surfaces. We use the notations $\pi_x : J^1E \to E$ and $\pi_{xz} : J^1E \to J^0E = E \times \mathbb{R}$ for the **base** and **front projections**. Generically, outside of a codimension 1 subset $\Lambda_{cusp} \subset \Lambda$ consisting of cusp points (resp. cusp edge and swallowtail points) the front projection, $\pi_{xz}|_\Lambda$, of a Legendrian knot (resp. surface) is an immersion. Moreover, any $p \in \Lambda \setminus \Lambda_{cusp}$ has a neighborhood $W \subset \Lambda$ that is the 1-jet, $j^1 f$, of some **local defining function** $f : U \to \mathbb{R}$ defined in a neighborhood $U \subset E$ of $\pi_x(p)$. In particular, $\pi_{xz}(W)$ agrees with the graph of f. Suppose that E has boundary and $N \subset \partial E$ is a boundary component. If Λ intersects $\pi_x^{-1}(N)$ transversally (this is equivalent to the map $\pi_x|_\Lambda$ being transverse to N), then the **restriction** of Λ to J^1N is the Legendrian submanifold $\Lambda|_N \subset J^1N$ that is the image of $\Lambda \cap \pi_x^{-1}(N)$ under the restriction map $\pi_x^{-1}(N) \to J^1N$. The front projection of $\Lambda|_N$ is the intersection of $\pi_{xz}(\Lambda) \subset E \times \mathbb{R}$ with $N \times \mathbb{R}$.

The standard Reeb vector field on J^1E is $\frac{\partial}{\partial z}$, and Reeb chords of Λ, i.e. trajectories of $\frac{\partial}{\partial z}$ having endpoints on Λ, correspond to critical points of **local difference functions**, $f_{i,j} = f_i - f_j$, where f_i, f_j are local defining functions for Λ with $f_i > f_j$. For $\rho \in \mathbb{Z}_{\geq 0}$, a $\mathbb{Z}/\rho$-valued **Maslov potential** μ for a 1 or 2 dimensional Legendrian Λ is a locally constant map $\mu : \Lambda \backslash \Lambda_{cusp} \to \mathbb{Z}/\rho$, such that near each cusp point or edge, the value of μ at the upper sheet of the cusp is one more than the value of μ at the lower sheet; such a Maslov potential exists if and only if ρ is a divisor of the Maslov number of Λ which is $2rot(\Lambda)$ when Λ is 1-dimensional, cf. [32, Section 4]. When Λ is equipped with a choice of $\mathbb{Z}/\rho$-valued Maslov potential, each Reeb chord, c, is assigned a $\mathbb{Z}/\rho$-grading by

$$|c| = \mu(c_u) - \mu(c_l) + ind(f_{i,j}) - 1 \in \mathbb{Z}/\rho$$

where c_u and c_l are the upper and lower endpoints of c and $ind(f_{i,j})$ is the Morse index of c when viewed as a critical point of the local difference function $f_{i,j} = f_i - f_j$ (where f_i, f_j are defining functions for Λ near c_u and c_l.) In the case that Λ is connected, the grading of Reeb chords is independent of the choice of Maslov potential, but this is not true in the multi-component case.

In order to have a well defined grading of Reeb chords (and also of the Legendrian contact homology algebra) we will always work with Legendrians equipped with a choice of Maslov potential.

Definition 2.8. For $\rho \in \mathbb{Z}_{\geq 0}$, a **$\rho$-graded Legendrian** is a pair (Λ, μ) consisting of a Legendrian submanifold $\Lambda \subset J^1E$ together with a choice of $\mathbb{Z}/\rho$-valued Maslov potential, μ.

Remark 2.9. When the base space E is oriented a $\mathbb{Z}/2$-valued Maslov potential is equivalent to a choice of orientation for Λ. [Indeed, given an orientation for Λ one defines a $\mathbb{Z}/2$-valued Maslov potential, $\mu : \Lambda \setminus \Lambda_{cusp} \to \mathbb{Z}/2$, to be 0 (resp. 1) at points where the base projection of Λ to E is orientation preserving (resp. reversing). Moreover, this procedure can be reversed to produce an orientation from a $\mathbb{Z}/2$-valued Maslov potential.] Thus, $\mathbb{Z}/2$-graded Legendrians are the same as oriented Legendrians, and any $\mathbb{Z}/\rho$-graded Legendrian with ρ even has a well defined orientation (from reducing the Maslov potential mod 2).

For a ρ-graded Legendrian submanifold Λ in J^1E, symplectic field theory gives a Floer type invariant called the **Legendrian contact homology DGA** (aka. the Chekanov-Eliashberg algebra) [9, 19, 15] that we will denote by $(\mathcal{A}(\Lambda), \partial)$. It is a $\mathbb{Z}/\rho$-graded triangular, based DGA over $\mathbb{Z}/2$ generated by Reeb chords of Λ with $\mathbb{Z}/\rho$-grading arising from the choice of Maslov potential, μ. The differential ∂ is defined by counting holomorphic disks either in T^*M, cf. [15], or $Symp(J^1M)$, cf. [11], with boundary on either the cotangent projection of Λ or the Lagrangian cylinder $\mathbb{R} \times \Lambda$. According to [12], one can alternatively compute the differential by

counting **gradient flow trees** (abbrv. **GFTs**) which are certain trees whose edges parametrize flowlines of gradients of local difference functions, $-\nabla f_{i,j}$, with $f_{i,j} > 0$; see also [34]. The $\mathbb{Z}/\rho$-graded DGA $(\mathcal{A}(\Lambda), \partial)$ is an invariant of (Λ, μ) up to stable tame isomorphism.

The LCH DGA is functorial for exact Lagrangian cobordisms, and we will restrict our attention to the case of cobordisms in the symplectization of J^1M. An **exact Lagrangian cobordism** L from Λ_- to Λ_+ is an embedded surface in $Symp(J^1M) := (\mathbb{R}_t \times J^1M, d(e^t\alpha))$ (as shown in Figure 1) that

- agrees with the cylinder $\mathbb{R} \times \Lambda_+$ (resp. $\mathbb{R} \times \Lambda_-$) when t is very positive (resp. very negative; and
- there is a function $g : L \to \mathbb{R}$ such that $(e^t\alpha)|_L = dg$ and g is constant for t near $\pm\infty$. Such a function g is called a **primitive**.

When Λ_- is empty, we say that L is an **exact Lagrangian filling** of L.




FIGURE 1. An exact Lagrangian cobordism from Λ_- to Λ_+.

According to [16, 13, 17], an exact Lagrangian cobordism L from Λ_- to Λ_+ induces a DGA map f_L from $\mathcal{A}(\Lambda_+)$ to $\mathcal{A}(\Lambda_-)$. (For now, we suppress discussion of the grading.) When two such cobordisms are isotopic through exact Lagrangian cobordisms, their induced DGA maps are DGA homotopic. Moreover, when two exact Lagrangian cobordisms L_i, $i = 1, 2$ from Λ_i to Λ_{i+1} are concatenated together to form $L_2 \circ L_1$ (by translating L_2 in the positive t-direction, truncating the two cobordisms, and gluing along a region of the form $(a, b) \times \Lambda_1$), the induced DGA map $f_{L_2 \circ L_1}$ is DGA homotopic to $f_{L_1} \circ f_{L_2}$. In summary, we have a functor between a suitably defined category of Legendrian knots with morphisms exact Lagrangian cobordisms and the homotopy category of DGAs.

2.4. **Conical Legendrian cobordisms.** In [32], functoriality for the LCH DGA was generalized to a class of *immersed* exact Lagrangian cobordisms by working with conical Legendrian cobordisms. We now review relevant results from [32].

Let M be a 1-manifold. In $J^1(\mathbb{R}_{>0} \times M)$, we denote the $\mathbb{R}_{>0}$ coordinate by s.

Definition 2.10. Let Λ be a Legendrian link in J^1M that is parametrized by $\theta \mapsto (x(\theta), y(\theta), z(\theta))$, let $f : I \to \mathbb{R}_{>0}$ where $I \subset \mathbb{R}_{>0}$ is an interval, and let $A \in \mathbb{R}$ be a constant. Define

$$j^1(f(s) \cdot \Lambda + A) \subset J^1(I \times M)$$

to be the Legendrian that is parametrized by

$$\begin{aligned}(s, \theta) \mapsto &(s, x(\theta), f'(s)z(\theta), f(s)y(\theta), f(s)z(\theta) + A)\\ &= (s, x, u, y, z) \in J^1(I \times M)\end{aligned}$$

where (s, x) are the coordinates on $I \times M$, (u, y) are the coordinates on the cotangent fibers, and z is the $\mathbb{R}$ coordinate on $J^1(I \times M) = T^*(I \times M) \times \mathbb{R}$.

Note that the front projection of $j^1(f(s)\cdot\Lambda + A)$ is obtained from $\pi_{xz}(\Lambda)$ by forming the cylinder in the s-direction, multiplying z-coordinates by $f(s)$, and then shifting them by A.

Definition 2.11. A **conical Legendrian cobordism** Σ from Λ_- to Λ_+ is an embedded Legendrian surface in $J^1(\mathbb{R}_{>0}\times M)$ (see Figure 2) such that

- Σ has conical ends, i.e, when $s > s_+$ (resp. $s < s_-$) for some positive number $s_\pm$, the Legendrian surface Σ is $j^1(s\cdot\Lambda_\pm + A_\pm)$ for some constant $A_\pm$; and
- the intersection $\Sigma \cap J^1([s_-, s_+]\times M)$ is compact.

Two conical Legendrian cobordisms from Λ_- to Λ_+ are **conical Legendrian isotopic** if they are isotopic through conical Legendrian cobordisms from Λ_- to Λ_+.




FIGURE 2. A sketch of a conical Legendrian cobordism in the front projection to $J^0(\mathbb{R}_{>0}\times M)$.

Note that there is a contactomorphism between $J^1(\mathbb{R}_{>0}\times M)$ and $(Symp(J^1M)\times\mathbb{R}_w, dw+e^t\alpha)$ (see [32] for details). Consider the image of Σ in $Symp(J^1M)\times\mathbb{R}$ and then take the Lagrangian projection to $Symp(J^1M)$. The resulting surface L is an (immersed) exact Lagrangian surface in $Symp(J^1M)$, and we call surfaces obtained in this manner **good Lagrangian cobordisms** from Λ_- to Λ_+. Note that the conical ends condition on Σ implies that L has cylindrical ends and that the primitive is constant on top and bottom cylinders. As long as the immersed Lagrangians are equipped with such primitives, the construction is reversible, so we have a bijection between conical Legendrian cobordisms and good Lagrangian cobordisms; see eg. [32, Proposition 4.9]. Thus, conical Legendrian cobordisms generalize exact Lagrangian cobordisms from the embedded case to the immersed case. Conical Legendrian cobordisms can be concatenated in a way that generalizes the concatenation of exact Lagrangian cobordisms; see [32, Section 4.4].

The functoriality of the LCH DGA extends to conical Legendrian cobordisms provided the induced maps are allowed to be immersed DGA maps.

Theorem 2.12 ([32])**.** *A conical Legendrian cobordism Σ from Λ_- to Λ_+ induces an immersed DGA map*

$$M_\Sigma = \big(\mathcal{A}(\Lambda_+) \xrightarrow{f} \mathcal{A}(\Sigma) \xleftarrow{i} \mathcal{A}(\Lambda_-)\big),$$

satisfying:

1. *The DGA $\mathcal{A}(\Sigma)$ is generated by Reeb chords of Σ and Reeb chords of Λ_-.*
2. *When $\Lambda_\pm$ and Σ are ρ-graded such that the $\mathbb{Z}/\rho$-valued Maslov potential of Σ restricts to the Maslov potentials on $\Lambda_\pm$, all the DGAs $\mathcal{A}(\Lambda_+), \mathcal{A}(\Sigma)$, and $\mathcal{A}(\Lambda_-)$ inherit $\mathbb{Z}/\rho$-gradings that are preserved by f and i.*
3. *When two conical Legendrian cobordisms Σ and Σ' from Λ_- to Λ_+ are conical Legendrian isotopic, their induced immersed DGA maps are immersed homotopic.*
4. *When $\Sigma_1 : \Lambda_1 \to \Lambda_2$ and $\Sigma_2 : \Lambda_2 \to \Lambda_3$ are concatenated the immersed maps satisfy*

$$M_{\Sigma_2\circ\Sigma_1} \simeq M_{\Sigma_1}\circ M_{\Sigma_2} \quad \textit{(immersed homotopy).}$$

(5) *When Σ has no Reeb chords, i.e., when it corresponds to an embedded exact Lagrangian cobordism L, we have that $\mathcal{A}(\Sigma) = \mathcal{A}(\Lambda_-)$ and the f map is DGA homotopic to the induced map f_L from* [16].

The construction of the immersed DGA map is summarized as follows. Given a conical Legendrian cobordism Σ from Λ_- to Λ_+, we construct a Morse cobordism $\widetilde{\Sigma}$ (as in [16]) by replacing the conical ends $j^1(s\cdot\Lambda_\pm + A_\pm)$ with standard Morse ends $j^1(h_\pm(s)\cdot\Lambda_\pm + B_\pm)$, where h_- (resp. h_+) are positive Morse functions with a single minimum (resp. maximum). The set of Reeb chords of $\widetilde{\Sigma}$ is in bijection with the set of Reeb chords of Σ and $\Lambda_\pm$. The DGA $\mathcal{A}(\Sigma)$ is generated by the Reeb chords of $\widetilde{\Sigma}$ that correspond to Reeb chords of Σ and Λ_-, and the differential is defined by the usual count of GFTs with respect to a suitable choice of metric g on $\mathbb{R}_{>0}\times M$ having the form $g_{\mathbb{R}}\times g_\pm$ near the critical points of $h_\pm$ where $g_{\mathbb{R}}$ is the Euclidean metric on $\mathbb{R}_{>0}$ and $g_\pm$ are regular[1] metrics used for computing $\mathcal{A}(\Lambda_\pm)$. Moreover, the DGA $\mathcal{A}(\Lambda_-)$ is a sub-DGA of $\mathcal{A}(\Sigma)$ and the map i is the natural DGA inclusion map. Finally, the DGA map f is defined by counting GFTs with positive puncture at one of the Λ_+ Reeb chords of $\widetilde{\Sigma}$ and with image to the left of the local maximum of h_+.

In [32, Section 6.3], the construction of Theorem 2.12 is formulated as a functor

$$F : \mathfrak{Leg}^\rho_{im} \to \mathfrak{DGA}^\rho_{im}$$

called the **immersed LCH functor**. The category $\mathfrak{Leg}^\rho_{im}$ has objects (Λ, g, μ) consisting of an (embedded) ρ-graded Legendrian knot, $\Lambda\subset J^1M$, with Maslov potential, μ, equipped with a choice of regular metric, g, on M. Morphisms from (Λ_-, g_-, μ_-) to (Λ_+, g_+, μ_+) are conical Legendrian isotopy classes of ρ-graded conical Legendrian cobordisms, $\Sigma : \Lambda_- \to \Lambda_+$, with Maslov potentials, μ, extending the Maslov potentials μ_- and μ_+.

Corollary 2.13. *The correspondence*

$$\begin{array}{rcl}(\Lambda, g, \mu) & \mapsto & \mathcal{A}(\Lambda, g, \mu),\\ (\Sigma, \mu) & \mapsto & [M_\Sigma]\end{array}$$

where $[M_\Sigma]$ denotes the immersed homotopy class of the immersed map M_Σ from Theorem 2.12 defines a contravariant functor $F : \mathfrak{Leg}^\rho_{im} \to \mathfrak{DGA}^\rho_{im}$.

2.5. **Morse minimum cobordisms.** Let $[a_-, a_+]\subset\mathbb{R}$ be a closed interval. A compact Legendrian $\Sigma_{min}\subset J^1([a_-, a_+]\times M)$ is called a **Morse minimum cobordism** from Λ_- to Λ_+ if there are neighborhoods $U_- = [a_-, a_- + \delta)$ and $U_+ = (a_+ - \delta, a_+]$, for some $\delta > 0$, such that Σ_{min} has the form $j^1(h_\pm(s)\cdot\Lambda_\pm + B_\pm)$ in $J^1(U_\pm\times M)$ where $h_\pm : U_\pm \to \mathbb{R}_{>0}$ are positive Morse functions with unique critical points that are non-degenerate local minima at $a_\pm$.

As in [17], the LCH DGA of a Morse minimum cobordism $\mathcal{A}(\Sigma_{min})$ is well-defined and can be computed using GFTs with respect to a regular metric g on $[a_-, a_+]\times M$ having the form $g_{\mathbb{R}}\times g_\pm$ in a neighborhood of $\{a_\pm\}\times M$ where $g_\pm$ are regular metrics for $\Lambda_\pm$. Moreover, there are DGA inclusions

$$i_\pm : \mathcal{A}(\Lambda_\pm)\hookrightarrow\mathcal{A}(\Sigma_{min})$$

obtained via identifying Reeb chords of $\Lambda_\pm$ with the Reeb chords of Σ_{min} located above $\{a_\pm\}\times M$. See also [32, Section 5.3].

Construction 2.14. Given Σ_{min} one can form an associated conical Legendrian cobordism $\Sigma_{conic}\subset J^1(\mathbb{R}_{>0}\times M)$ by shifting the interval $[a_-, a_+]$ into $[a_- + s_0, a_+ + s_0]\subset\mathbb{R}_{>0}$, and modifying the ends to have the form $j^1(\widehat{h}_\pm(s)\cdot\Lambda_\pm + B_\pm)$ with

$$\widehat{h}_- : (0, a_- + s_0 + \delta)\to\mathbb{R}_{>0}, \quad \text{and} \quad \widehat{h}_+ : (a_+ + s_0 - \delta, +\infty)\to\mathbb{R}_{>0}$$

chosen as follows:

[1]Here, **regular** means that (i) there are no GFTs for Λ of negative formal dimension and (ii) all 0-dimensional GFTs for Λ are transversally cut out. See [12].

- The function $\widehat{h}_-$ has no critical points, is increasing on $(0, a_- + s_0 + \delta)$, and satisfies
$$\begin{aligned} \widehat{h}_-(s) &= s, && \text{for } s \in (0, a_- + s_0], \\ \widehat{h}_-(s) &= h_-(s - s_0), && \text{near } s = a_- + s_0 + \delta. \end{aligned}$$
- The function $\widehat{h}_+$ has a unique critical point that is a non-degenerate local minimum at $s = a_+ + s_0$, and satisfies
$$\begin{aligned} \widehat{h}_+(s) &= h_+(s - s_0), && \text{for } s \in (a_+ + s_0 - \delta, a_+ + s_0], \\ \widehat{h}_+(s) &= s, && \text{for } s \gg 0. \end{aligned}$$

Moreover, from a regular metric g_{min} for Σ_{min} defined on $[a_-, a_+] \times M$ and having the form $g_{\mathbb{R}} \times g_{\pm}$ near $\{a_\pm\} \times M$ we construct a metric g_{conic} on $\mathbb{R}_{>0} \times M$ by shifting g_{min} by s_0 in the $\mathbb{R}_{>0}$-direction and then extending to agree with $g_{\mathbb{R}} \times g_-$ on $(0, a_- + s_0] \times M$ and $g_{\mathbb{R}} \times g_+$ on $[a_+ + s_0, +\infty) \times M$.

Note that the conditions on $\widehat{h}_-$ can be arranged with an appropriate choice of s_0. See Figure 3.




FIGURE 3. A Morse minimum cobordism, an associated conical cobordism, and the cobordism $\widetilde{\Sigma}_{conic}$ used in the proof of Proposition 2.15.

Proposition 2.15. *When $\Sigma_{min} \subset J^1([a_-, a_+] \times M)$ is a Morse minimum cobordism from Λ_- to Λ_+ and Σ_{conic} is an associated conical cobordism as in Construction 2.14. Then, the immersed LCH functor satisfies*
$$F(\Sigma_{conic}) = \left[\mathcal{A}(\Lambda_+) \xrightarrow{i_+} \mathcal{A}(\Sigma_{min}) \xleftarrow{i_-} \mathcal{A}(\Lambda_-) \right].$$

Proof. A Morse cobordism $\widetilde{\Sigma}_{conic}$ (with a Morse minimum at the negative end and a Morse maximum at the positive end) that can be used to compute the induced immersed map
$$M_{\Sigma_{conic}} = \left(\mathcal{A}(\Lambda_+) \xrightarrow{f} \mathcal{A}(\Sigma_{conic}) \xleftarrow{i} \mathcal{A}(\Lambda_-) \right)$$
from Theorem 2.12 is obtained as follows:

(1) At the negative end of Σ_{conic}, replace $j^1(\widehat{h}_-(s) \cdot \Lambda_- + B_-)$ with $j^1(h_-(s - s_0) \cdot \Lambda_- + B_-)$. This matches the negative end of Σ_{min} shifted by s_0.
(2) At the postive end, replace the function $\widehat{h}_+$ with some $\widetilde{h}_+ : (a_+ + s_0 - \delta, s_1 + \delta] \to \mathbb{R}_{>0}$. Here, s_1 is chosen large enough so that $\widehat{h}_+(s) = s$ for $s \geq s_1 - \delta$ and we require
   - $\widetilde{h}_+(s) = \widehat{h}_+(s)$ for $s \in (a_+ + s_0 - \delta, s_1]$,
   - $\widetilde{h}_+$ is increasing on $[s_1, s_1 + \delta]$ and has a single critical point on this interval that is a non-degenerate local maximum at $s_1 + \delta$.

   See Figure 3.

By definition, the DGA $\mathcal{A}(\Sigma_{conic})$ is generated by those Reeb chords of $\widetilde{\Sigma}_{conic}$ that appear in the region where $s < s_1 + \delta$. These Reeb chords are the same as the Reeb chords of Σ_{min} but shifted by s_0 in the s direction. Moreover, using metrics of the form g_{min} and g_{conic} as in Construction 2.14 to compute GFTs, because of the Morse minima all of the GFTs with positive punctures at these chords are contained in the region $[a_- + s_0, a_+ + s_0] \times M$ and therefore coincide (up to the shift in the s direction) with the GFTs of Σ_{min}; see eg. [17]. Thus, $\mathcal{A}(\Sigma_{conic}) = \mathcal{A}(\Sigma_{min})$ and the map i agrees with i_- by definition. The GFTs that define the map f (by definition) have their unique positive punctures at the Reeb chords located at $\{s_1 + \delta\} \times M$, and (because of the local minimum of $\widetilde{h}_+$ at $s = a_+ + s_0$) have there images contained in $(a_+ + s_0, s_1 + \delta) \times M$. In this region, $\widetilde{\Sigma}_{conic} = j^1(\widetilde{h}_+(s) \cdot \Lambda_+ + B_+)$ and $\widetilde{h}_+$ strictly increases, so the computation of such GFTs is as in the case of the identity cobordism from Λ_+ to itself found in [32, Proposition 6.15]. For each Reeb chord, b, of Λ_+ there is a single gradient trajectory that connects the Reeb chords of $\widetilde{\Sigma}_{conic}$ at $s = s_1 + \delta$ corresponding to b to the Reeb chord of $\widetilde{\Sigma}_{conic}$ at $s = a_+ + s_0$ corresponding to b, and these are the only rigid GFTs. It follows that $f = i_+$. □

As discussed in the introduction, Morse minimum cobordisms may be used to define an equivalence relation on augmented Legendrians. Refer to a triple (Λ, μ, ϵ) consisting of a ρ-graded Legendrian $\Lambda \subset J^1M$ with Maslov potential, μ, and a ρ-graded augmentation, $\epsilon : \mathcal{A}(\Lambda) \to \mathbb{Z}/2$, as a **$\rho$-graded augmented Legendrian**.

Definition 2.16. Two ρ-graded augmented Legendrians $(\Lambda_i, \mu_i, \epsilon_i)$, $i = 0, 1$, in J^1M are **cobordant** if there exists a triple (Σ, μ, α) consisting of a Morse minimum cobordism $\Sigma \subset J^1([0,1] \times M)$ from Λ_0 to Λ_1 together with a Maslov potential μ extending the μ_i and a ρ-graded augmentation $\alpha : \mathcal{A}(\Sigma) \to \mathbb{Z}/2$ satisfying $\alpha|_{\mathcal{A}(\Lambda_i)} \simeq \epsilon_i$ (DGA homotopy).


It is straightforward to see that cobordism defines an equivalence relation on ρ-graded augmented Legendrians.

We can now show that Theorem 1.2 implies Theorem 1.5 from the introduction.

Proof of Theorem 1.5. Given a ρ-graded augmented Legendrian (Λ, μ, ϵ), assuming Theorem 1.2, there exists a conical Legendrian cobordism with $\mathbb{Z}/\rho$-valued Maslov potential,

$$\Sigma : \emptyset \to \Lambda \quad (\text{if } \rho \neq 1) \quad \text{or} \quad \Sigma : U \to \Lambda \quad (\text{if } \rho = 1),$$

together with a ρ-graded augmentation $\alpha : \mathcal{A}(\Sigma) \to \mathbb{Z}/2$ such that $\epsilon \simeq \alpha \circ f_\Sigma$. Now, a Legendrian isotopy that is compactly supported in the conical ends of Σ modifies Σ to have the form Σ_{conic} for some Morse minimum cobordism Σ_{min} as in Construction 2.14. Then, from Theorem 2.12 there is an immersed DGA homotopy $M_\Sigma \simeq M_{\Sigma_{conic}}$ that (after using Proposition 2.15 to evaluate $M_{\Sigma_{conic}}$ and replacing the stable tame isomorphism $\varphi : \mathcal{A}(\Sigma_{min}) * S \to \mathcal{A}(\Sigma_{min}) * S'$ with its associated homotopy equivalence) gives rise to a DGA homotopy commutative diagram:



Then, we can compute

$$\epsilon \simeq \alpha \circ f_\Sigma \simeq (\alpha \circ h) \circ i_+,$$

so that $(\Sigma_{min}, \alpha \circ h)$ provides the cobordism of ρ-graded augmented Legendrians from $\emptyset$ (if $\rho \neq 1$) or (U, ϵ_1) (if $\rho = 1$) to (Λ, ϵ) as in the statement of Theorem 1.5 where $\epsilon_1 = \alpha \circ h \circ i_-$. □

Remark 2.17. (1) When ρ is even, the cobordism Σ and Legendrians are canonically oriented by the Maslov potential μ. If ρ is odd, Σ may be orientable or not. In the odd case, a refined relation of oriented cobordism for ρ-graded augmentation Legendrians arises from requiring that the Λ_i and Σ are additionally equipped with orientations. We leave the computation of such oriented, odd-graded cobordism classes of augmented Legendrians in $J^1\mathbb{R}$ as an open problem.

(2) Without augmentations, Legendrian cobordism classes in $J^1\mathbb{R}$ are computed as follows; see eg. [3, Section 5.1]. Two Legendrians in $J^1\mathbb{R}$ are oriented cobordant if and only if they have the same rotation number, while any two Legendrians in $J^1\mathbb{R}$ are non-oriented cobordant. In particular, Theorem 1.5 implies the well known result of Sabloff, see [37], that if $\Lambda \subset J^1\mathbb{R}$ has an ρ-graded augmentation with ρ even, then $rot(\Lambda) = 0$.

(3) As Legendrians that admit augmentations exhibit significantly more rigid behavior than general Legendrians, we do not see any *a priori* reason to expect that cobordism classes of ρ-graded augmented Legendrians should closely match the classical cobordism classes of Legendrians. In fact, in the case of J^1S^1 we will show in [31] that there are many examples of non-cobordant augmented Legendrians that become cobordant if one ignores the augmentations.

3. Immersed DGA maps and augmentations

A DGA morphism $f : \mathcal{A} \to \mathcal{B}$ contravariantly induces a map on homotopy classes of augmentations, $f^* : Aug(\mathcal{B})/\sim \to Aug(\mathcal{A})/\sim$. In Section 3.1, we consider analogous constructions for immersed DGA maps focusing on the *induced augmentation set* of an immersed DGA map, M, that is a subset $I(M) \subset Aug(\mathcal{B})/\sim \times Aug(\mathcal{A})/\sim$. We show that the induced augmentation set induced by a conical Legendrian cobordism Σ is a Legendrian invariant of Σ. In addition, we make some observations about the form of the augmentation set in the case of Legendrian fillings and embedded Legendrian cobordisms. In Section 3.2 we show that immersed augmentation sets compose as relations, and we record the effect of concatenating a conical Legendrian cobordism with the (invertible) Legendrian cobordism arising from a Legendrian isotopy.

3.1. **Induced augmentation sets.** We work with $\mathbb{Z}/\rho$-graded DGAs, *with $\rho \geq 0$ understood to be fixed*. As such, when the grading does not need to be emphasized we may refer to ρ-graded augmentations simply as augmentations. We denote by $Aug_\rho(\mathcal{A}) = Aug(\mathcal{A})$ the set of all (ρ-graded) augmentations of $\mathcal{A}$ to $\mathbb{Z}/2$, and we write $Aug(\mathcal{A})/\sim$ for the set of all DGA homotopy classes of augmentations. In the case that $\mathcal{A} = \mathcal{A}(\Lambda)$ is the DGA of some ρ-graded Legendrian knot or cobordism, we may shorten these notations to $Aug(\Lambda)$ and $Aug(\Lambda)/\sim$. A DGA map $f : \mathcal{A} \to \mathcal{B}$ induces a pullback map

$$f^* : Aug(\mathcal{B}) \to Aug(\mathcal{A}), \quad f^*\epsilon = \epsilon \circ f,$$

and this gives a well-defined map on DGA homotopy classes also denoted as

$$f^* : Aug(\mathcal{B})/\sim \to Aug(\mathcal{A})/\sim.$$

Let $M = \left(\mathcal{A}_1 \xrightarrow{f} \mathcal{B} \xleftarrow{i} \mathcal{A}_2\right)$ be an immersed DGA map. Then, the pullback construction results in maps

$$Aug(\mathcal{A}_1) \xleftarrow{f^*} Aug(\mathcal{B}) \xrightarrow{i^*} Aug(\mathcal{A}_2) \quad \text{and} \quad Aug(\mathcal{A}_1)/\sim \xleftarrow{f^*} Aug(\mathcal{B})/\sim \xrightarrow{i^*} Aug(\mathcal{A}_2)/\sim.$$

The latter diagram is equivalent to the map

$$i^* \times f^* : Aug(\mathcal{B})/\sim \to (Aug(\mathcal{A}_2)/\sim) \times (Aug(\mathcal{A}_1)/\sim)$$

that we call the **augmentation map** induced by M.


Definition 3.1. The **induced augmentation set**, $I(M)$, of an immersed DGA map $M = \left(\mathcal{A}_1 \xrightarrow{f} \mathcal{B} \xhookleftarrow{i} \mathcal{A}_2\right)$ is the image of the augmentation map,

$$I(M) = \mathrm{Im}(i^* \times f^*) \subset Aug(\mathcal{A}_2)/\sim \times Aug(\mathcal{A}_1)/\sim.$$

The induced augmentation set is an invariant of the immersed homotopy class of M.

Proposition 3.2. *Suppose that $M = \left(\mathcal{A}_1 \xrightarrow{f} \mathcal{B} \xhookleftarrow{i} \mathcal{A}_2\right)$ and $M' = \left(\mathcal{A}_1 \xrightarrow{f'} \mathcal{B}' \xhookleftarrow{i'} \mathcal{A}_2\right)$ are immersed DGA maps that are immersed homotopic. Then, there is a bijection $h^* : Aug(\mathcal{B}')/\sim \xrightarrow{\cong} Aug(\mathcal{B})/\sim$ fitting into a commutative diagram*


(3.1)



In particular, the induced augmentation sets satisfy $I(M) = I(M')$.

Proof. There exists a diagram

(3.2)



where φ is a DGA isomorphism such that

$$\varphi \circ f \simeq f' \quad \text{and} \quad \varphi \circ i = i'.$$

Let $h = \pi' \circ \varphi \circ \iota$, where $\iota : \mathcal{B} \to \mathcal{B} * S$ and $\pi' : \mathcal{B}' * S' \to \mathcal{B}'$ are the inclusions and projections, be the associated homotopy equivalence from $\mathcal{B}$ to $\mathcal{B}'$. Then, since the maps such as $f : \mathcal{A}_1 \to \mathcal{B} * S$ in (3.2) are implicitly understood to mean $\iota \circ f$, the diagram (3.2) leads to a similar homotopy commutative diagram (3.2)' with the vertical map replaced with $h : \mathcal{B} \to \mathcal{B}'$.


Now, the association

$$\begin{array}{lcl} \mathcal{A} & \rightsquigarrow & Aug(\mathcal{A})/\sim, \\ g : \mathcal{A} \to \mathcal{B} & \rightsquigarrow & g^* : Aug(\mathcal{B})/\sim \to Aug(\mathcal{A})/\sim \end{array}$$

gives a well-defined contravariant functor from the category $\mathfrak{DGA}^{\rho}_{hom}$ (where morphisms are DGA homotopy classes of maps) to the category of sets. In particular, since h is a homotopy equivalence, h^* is a bijection, and since (3.2)' is homotopy commutative, the diagram (3.1) is indeed fully commutative. □

When $\Sigma : \Lambda_- \to \Lambda_+$ is a conical Legendrian cobordism with immersed DGA map, M_Σ, as in Theorem 2.12, we write $I_\Sigma = I(M_\Sigma) \subset Aug(\Lambda_-)/\sim \times Aug(\Lambda_+)/\sim$ and refer to I_Σ as the **induced augmentation set** of Σ.

Corollary 3.3. *Suppose that $\Sigma, \Sigma' : \Lambda_- \to \Lambda_+$ are conical Legendrian cobordisms related by a conical Legendrian isotopy. Then, there is a commutative diagram*

(3.3)


i.e., h^* is a bijection and $i^*_{\Sigma'} \times f^*_{\Sigma'} = (i^*_\Sigma \times f^*_\Sigma) \circ h^*$.

In particular, the induced augmentation set

$$I_\Sigma = Im(i^*_\Sigma \times f^*_\Sigma) \subset Aug(\Lambda_-)/\sim \times Aug(\Lambda_+)/\sim,$$

is an invariant of Σ.

Remark 3.4.
(1) To provide a more refined invariant of Σ, one can take multiplicities into account when considering I_Σ. Eg., the function
$$Aug(\Lambda_-)/\sim \times Aug(\Lambda_+)/\sim \to \mathbb{Z}_{\geq 0}, \quad ([\epsilon_-],[\epsilon_+]) \mapsto \left|(i^*_\Sigma \times f^*_\Sigma)^{-1}\big(([\epsilon_-],[\epsilon_+])\big)\right|$$
is a conical Legendrian invariant of Σ. A similar invariant (using a normalized count of augmentations rather than homotopy classes) is studied for 1-dimensional Σ in [39].
(2) The set $Aug(\Lambda_\pm)$ can be equipped with the additional structure of an A_∞-category whose moduli space of objects is $Aug(\Lambda_\pm)/\sim$, cf. [5, 27], and one could hope for a further refinement of the augmentation map $i^*_\Sigma \times f^*_\Sigma$ to take this structure into account. In the case of embedded cobordisms, such a refinement is made in [29].

3.1.1. *The embedded case.* When the conical Legendrian cobordism $\Sigma : \Lambda_- \to \Lambda_+$ corresponds to an embedded Lagrangian cobordism, L, i.e. when Σ has no Reeb chords, $\mathcal{A}(\Sigma) = \mathcal{A}(\Lambda_-)$ and $i_\Sigma = id_{\mathcal{A}(\Lambda_-)}$. Therefore, in this case the augmentation map $i^*_\Sigma \times f^*_\Sigma$ is determined by the induced augmentation set

$$I_\Sigma = \{([\epsilon], f^*_\Sigma[\epsilon]) \,|\, [\epsilon] \in Aug(\Lambda_-)/\sim\} \subset Aug(\Lambda_-)/\sim \times Aug(\Lambda_+)/\sim \tag{3.4}$$

which is simply the function f^*_Σ viewed as a relation.

More generally, when $M = (\mathcal{A}_1 \xrightarrow{f} \mathcal{A}_2 \xleftarrow{id} \mathcal{A}_2)$ is the image of an ordinary DGA map $f : \mathcal{A}_1 \to \mathcal{A}_2$ under the functor from Proposition 2.7, the induced augmentation set for M is just the graph of f^*.

3.1.2. *Induced augmentations and immersed fillings.* Let Σ be a conical Legendrian filling, or equivalently a good immersed Lagrangian filling. Since $\Lambda_- = \emptyset$, we have $\mathcal{A}(\Lambda_-) = \mathbb{Z}/2$. Thus, $Aug(\Lambda_-) = Aug(\Lambda_-)/\sim$ consists of a single element, so that we can view I_Σ as a subset of $Aug(\Lambda_+)$. To emphasize the analogy with the case of embedded Lagrangian fillings, given an augmentation $\alpha : \mathcal{A}(\Sigma) \to \mathbb{Z}/2$, we use the notation

$$\epsilon_{(\Sigma,\alpha)} : \mathcal{A}(\Lambda_+) \to \mathbb{Z}/2, \quad \epsilon_{(\Sigma,\alpha)} = \alpha \circ f_\Sigma$$

and refer to $\epsilon_{(\Sigma,\alpha)}$ as the **augmentation induced by Σ via α**. Thus, the induced augmentation set $I_\Sigma \subset Aug(\Lambda_+)$ consists of those augmentations of Λ_+ that can be induced by some choice of augmentation for Σ.

3.2. **Concatenation and induced augmentation sets.** Let $\mathfrak{Rel}$ denote the category whose objects are sets and morphisms are relations, i.e. a morphism $R \subset Hom_{\mathfrak{Rel}}(X,Y)$ is just a subset $R \subset X \times Y$. Given relations $R \subset X \times Y$ and $S \subset Y \times Z$, there composition is

$$S \circ R = \{(x,z) \in X \times Z \,|\, \exists y \in Y \text{ such that } (x,y) \in R \text{ and } (y,z) \in S\}.$$

Observation 3.5. (1) Any function $f : X \to Y$ defines a relation $\Gamma_f = \{(x, f(x)) \,|\, x \in X\} \subset X \times Y$.

(2) Given any $R \subset X \times Y$ and $g : Y \to Z$, we have

$$\Gamma_g \circ R = \{(x, g(y)) \,|\, (x,y) \in R\} = (id \times g)(R).$$

(3) If $f : X \to Y$ is a bijection, and $S \subset Y \times Z$,then

$$S \circ \Gamma_f = (f^{-1} \times id)(S).$$

Proposition 3.6. *The induced augmentation set construction gives a contravariant functor* $\mathfrak{DGA}^{\rho}_{im} \to \mathfrak{Rel}$,

$$\begin{array}{ccc} \mathcal{A} & \rightsquigarrow & Aug(\mathcal{A})/\sim \\ [M] & \rightsquigarrow & I(M). \end{array}$$

In particular, the induced augmentation sets for a pair of conical Legendrian cobordisms, $\Sigma_i : \Lambda_{i+1} \to \Lambda_i$, $i = 1, 2$, *satisfy*

$$I_{\Sigma_1 \circ \Sigma_2} = I_{\Sigma_1} \circ I_{\Sigma_2}. \tag{3.5}$$

Proof. Let $M_1 = \left(\mathcal{A}_1 \xrightarrow{f_1} \mathcal{B}_1 \xleftarrow{j_1} \mathcal{A}_2\right)$ and $M_2 = \left(\mathcal{A}_2 \xrightarrow{f_2} \mathcal{B}_2 \xleftarrow{j_2} \mathcal{A}_3\right)$ be immersed maps. The set $I(M_2 \circ M_1)$ only depends on the immersed homotopy class of $M_2 \circ M_1$. Thus, we can compute it using the immersed map $\mathcal{A}_1 \xrightarrow{f_1} \mathcal{D} \xleftarrow{j_2} \mathcal{A}_3$ as in Proposition 2.5 where $\mathcal{D} = \mathcal{B}_1 * \widehat{\mathcal{A}}_2 * \mathcal{B}_2$ with differential as in (2.2). That $I(M_2 \circ M_1) = I(M_1) \circ I(M_2)$ then follows from:

Claim: Let X be the set of triples (α_1, α_2, K) such that $\alpha_i \in Aug(\mathcal{B}_i)$ and $K : \mathcal{A}_2 \to \mathbb{Z}/2$ is a DGA homotopy operator from $i_1^*\alpha_1$ to $f_2^*\alpha_2$, i.e. a $(i_1^*\alpha_1, f_2^*\alpha_2)$-derivation with $i_1^*\alpha_1 - f_2^*\alpha_2 = K \circ \partial$. There is a bijection

$$X \to Aug(\mathcal{D}), \quad (\alpha_1, \alpha_2, K) \mapsto \alpha$$

where $\alpha : \mathcal{D} \to \mathbb{Z}/2$ satisfies $\alpha|_{\mathcal{B}_i} = \alpha_i$ and $\alpha(\widehat{a}_i) = K(a_i)$ for all generators $a_i \in \mathcal{A}_2$.

To verify the claim note that $(\alpha \circ \partial)|_{\mathcal{B}_i} = 0$ if and only if $\mathcal{B}_i$ is an augmentation. In addition, since $\partial(\widehat{a}_i) = i_1(a_i) + f_2(a_i) + \Gamma \circ \partial a_i$ where $\Gamma(a_i) = \widehat{a}_i$ is an (i_1, f_2)-derivation, the equation $\alpha \circ \partial(\widehat{a}_i) = 0$ is equivalent to $i_1^*\alpha_1(a_i) + f_2^*\alpha_2(a_i) = K \circ \partial a_i$. □

In the case that the corresponding exact Lagrangian cobordism is embedded, we get a simpler statement.

Proposition 3.7. *Suppose* $\Sigma : \Lambda_- \to \Lambda_+$ *is a conical Legendrian cobordism with embedded Lagrangian projection, and let* $\Sigma' : \Lambda'_- \to \Lambda_-$ *and* $\Sigma'' : \Lambda_+ \to \Lambda''_+$.

(1) *Then,*

$$I_{\Sigma \circ \Sigma'} = (id \times f_\Sigma^*)(I_{\Sigma'})$$

(2) *If* $f_\Sigma^* : Aug(\Lambda_-)/\sim \to Aug(\Lambda_+)/\sim$ *is a bijection, then*

$$I_{\Sigma'' \circ \Sigma} = ((f_\Sigma^*)^{-1} \times id)(I_{\Sigma''}).$$

Proof. As discussed in 3.1.1, when Σ is embedded I_Σ is the relation $\Gamma_{f_\Sigma^*}$ associated to the function f_Σ^*. Thus, the formulas from Observation 3.5 can be applied.

□

An important case where (2) of Proposition 3.7 applies is when Σ is induced by a Legendrian isotopy. We now briefly review a version of this construction.

Let $\Phi = \{\Lambda_s\}_{s\in\mathbb{R}_{>0}}$ be a Legendrian isotopy from Λ_- to Λ_+ so that $\Lambda_s : \sqcup_{i=1}^c S^1 \to J^1M$ for $s \in \mathbb{R}_{>0}$ is a Legendrian embedding satisfying $\Lambda_s = \Lambda_-$ for $0 < s \leq 1$ and $\Lambda_s = \Lambda_+$ for $s \gg 0$. Writing

$$\Lambda_s(\theta) = (x(s,\theta),\, y(s,\theta),\, z(s,\theta))$$

there is a conical Legendrian cobordism

$$\Sigma_\Phi : \mathbb{R}_{>0} \times (\sqcup_{i=1}^c S^1) \to J^1(\mathbb{R}_{>0} \times M) = \{(s,x,u,y,z)\}$$

from Λ_- to Λ_+ associated to Φ that is parametrized by

$$\Sigma_\Phi(s,\theta) = \left(s,\, x(s,\theta),\, z(s,\theta) + s\cdot\frac{\partial z}{\partial s}(s,\theta) - s\cdot y(s,\theta)\cdot\frac{\partial x}{\partial s}(s,\theta),\, s\cdot y(s,\theta),\, s\cdot z(s,\theta)\right).$$

It can be shown that after reparametrizing by an appropriate orientation preserving diffeomorphism, $\alpha : \mathbb{R}_{>0} \to \mathbb{R}_{>0}$, the conical Legendrian cobordism corresponding to $\Phi' = \{\Lambda_{\alpha(s)}\}$ will not have Reeb chords. Indeed, with our setup, one can take $\alpha(s) = s^a$ with $a > 0$ suitably small. See [16, Section 6.1] for a version of this construction for exact Lagrangian cobordisms in $Symp(J^1M)$.

Corollary 3.8. *Let $\Phi = \{\Lambda_s\}_{s\in\mathbb{R}_{>0}}$ be a Legendrian isotopy parametrized so that the conical Legendrian cobordism Σ_Φ does not have Reeb chords. Then, $f^*_{\Sigma_\Phi} : Aug(\Lambda_-)/\sim \to Aug(\Lambda_+)/\sim$ is a bijection. In particular, we have*

$$I_{\Sigma_\Phi\circ\Sigma'} = (id \times f^*_{\Sigma_\Phi})(I_{\Sigma'}) \quad \text{and} \quad I_{\Sigma''\circ\Sigma_\Phi} = ((f^*_{\Sigma_\Phi})^{-1} \times id)(I_{\Sigma''}).$$

whenever Σ' and Σ'' are composable with Σ_Φ.

Proof. Let Φ^{-1} be be the inverse Legendrian isotopy $\{\Lambda_{1/s}\}_{s\in\mathbb{R}_{>0}}$ reparametrized, if necessary, to ensure that $\Sigma_{\Phi^{-1}}$ has no Reeb chords. Then, the immersed LCH functor satisfies

$$F(\Sigma_\Phi) = [M_{\Sigma_\Phi}] = I(f_{\Sigma_\Phi}) \quad \text{and} \quad F(\Sigma_{\Phi^{-1}}) = [M_{\Sigma_{\Phi^{-1}}}] = I(f_{\Sigma_{\Phi^{-1}}})$$

where $I : \mathfrak{DGA}^\rho_{hom} \to \mathfrak{DGA}^\rho_{im}$ is the functor from Proposition 2.7. There is a clear conical Legendrian isotopy between $\Sigma_\Phi \circ \Sigma_{\Phi^{-1}}$ (resp. $\Sigma_{\Phi^{-1}} \circ \Sigma_\Phi$) and the identity cobordism $j^1(s\cdot\Lambda_+)$ (resp. $j^1(s\cdot\Lambda_-)$). From functoriality, it follows that $I(f_{\Sigma_\Phi} \circ f_{\Sigma_{\Phi^{-1}}})$ and $I(f_{\Sigma_{\Phi^{-1}}} \circ f_{\Sigma_\Phi})$ are identity morphisms in $\mathfrak{DGA}^\rho_{im}$, and since I is injective on hom-sets we conclude that f_{Σ_Φ} and $f_{\Sigma_{\Phi^{-1}}}$ are homotopy inverses. □

4. Immersed maps and the cellular DGA

In [33, 34, 35], a cellular version of Legendrian contact homology is introduced and shown to be equivalent to the usual LCH DGA in the case of closed Legendrian surfaces. In Sections 4.1 and 4.2, we briefly review the cellular DGA and extend its definition to the case of (compact) Legendrian cobordisms. The DGA of the identity cobordisms is computed in Section 4.3 and shown to be a mapping cylinder DGA. In Section 4.4 a cellular version, F_{cell}, of the immersed LCH functor is defined with a domain category consisting of ρ-graded Legendrians equipped with some additional data and with compact Legendrian cobordisms as morphisms. Finally, in Section 4.5 we state in Proposition 4.9 a precise relationship between the immersed and cellular LCH functors, F and F_{cell}, that will allow us to work with F_{cell} in place of F when considering induced augmentation sets; see Corollary 4.12. Specifically, after unifying the domain categories by precomposing with suitable functors, F and F_{cell} become isomorphic. This isomorphism is an extension of the isomorphism between the cellular and LCH DGAs for closed Legendrian surfaces from [33, 34, 35], and is presented here in a condensed manner. A more detailed presentation appears in Appendices A and B of the preprint version of this article [30].

4.1. **Review of the cellular DGA.** The cellular DGA construction requires as input a Legendrian knot or surface equipped with a suitable polygonal decomposition of its base projection. Recall that for a generic 1-dimensional Legendrian knot $\Lambda \subset J^1M$ the singularities of the front projection, $\pi_{xz}(\Lambda) \subset M \times \mathbb{R}$, are cusp points and crossings points (i.e. transverse double points). Generically, front projections of Legendrian surfaces have crossing arcs and cusp edges as codimension 1 singularities and triple points (intersection of three smooth sheets of Λ), cusp-sheet intersections (where a smooth sheet intersects a cusp edge), and swallowtail points as codimension 2 singularities. See eg. [3]; the front singularities for surfaces are illustrated in [33, Section 2.2]. We say that Λ has **generic front and base projections** if the front singularities are generic, and the base projections (to M) of the different classes of front singularities are all self transverse and transverse to one another.

Definition 4.1. Let $\Lambda \subset J^1M$ be a closed Legendrian submanifold with $\dim \Lambda = 1$ or 2 having generic front and base projections. A **compatible polygonal decomposition** for Λ is a polygonal[2] decomposition, $\mathcal{E} = \{e^d_\alpha\}$, of the base projection of Λ,

$$\sqcup^2_{d=0} \sqcup_\alpha e^d_\alpha = \pi_x(\Lambda) \subset M,$$

(where the superscript $0 \leq d \leq 2$ denotes the dimension of a cell) such that the base projection of the singular set of $\pi_{xz}(\Lambda)$ (crossings, cusps, swallow tail points, etc.) is contained in the 1-skeleton of $\mathcal{E}$. In addition, we require that:

(1) Each 1-cell is assigned an orientation.
(2) Each 2-cell is assigned an initial and terminal vertex, v_0 and v_1. If $v_0 = v_1$, then a prefered direction around the boundary of the 2-cell is also chosen.
(3) At each swallowtail point, s_0, the two polygonal corners that border the crossing arc near its endpoint at s_0 are labelled as S and T.

Let $(\Lambda, \mathcal{E})$ be a pair consisting of a closed Legendrian knot or surface, $\Lambda \subset J^1M$, together with a choice of compatible polygonal decomposition. The **cellular DGA** of $(\Lambda, \mathcal{E})$ will be denoted $\mathcal{A}(\Lambda, \mathcal{E})$ or $\mathcal{A}_{cell}(\Lambda, \mathcal{E})$, and is defined as follows.

Algebra: Given a cell, e^d_α, denote by $\{S^\alpha_p\}$ the set of **sheets** of Λ above e^d_α. By definition, sheets above e^d_α are those components of $\Lambda \cap \pi_x^{-1}(e^d_\alpha)$ not contained in any cusp edge. (Note: (i) Sheets are subsets of Λ, not $\pi_{xz}(\Lambda)$, so that, eg., a crossing arc of $\pi_{xz}(\Lambda)$ above a 1-cell corresponds to two sheets. (ii) A swallowtail point *is* considered to be a sheet above a 0-cell.) The algebra $\mathcal{A}(\Lambda, \mathcal{E})$ is the free unital, associative $\mathbb{Z}/2$-algebra whose generators are in bijection with triples $(e^d_\alpha, S^\alpha_i, S^\alpha_j)$ where S^α_i and S^α_j are sheets above e^d_α such that the inequality $z(S^\alpha_i) > z(S^\alpha_j)$ holds pointwise above e^d_α. We denote the generator associated to $(e^d_\alpha, S^\alpha_i, S^\alpha_j)$ as

$$a^\alpha_{i,j}, \quad b^\alpha_{i,j}, \quad \text{or} \quad c^\alpha_{i,j}$$

when the dimension of e^d_α is 0, 1, or 2 respectively.

Grading: A $\mathbb{Z}/\rho$-grading on $\mathcal{A}(\Lambda, \mathcal{E})$ arises from a choice of $\mathbb{Z}/\rho$-valued Maslov potential, μ, for Λ. The $\mathbb{Z}/\rho$-grading of generators is

$$|a^\alpha_{i,j}| = \mu(S^\alpha_i) - \mu(S^\alpha_j) - 1, \quad |b^\alpha_{i,j}| = \mu(S^\alpha_i) - \mu(S^\alpha_j), \quad |c^\alpha_{i,j}| = \mu(S^\alpha_i) - \mu(S^\alpha_j) + 1.$$

(If S^α_i is a swallowtail point above a 0-cell, then take $\mu(S^\alpha_i)$ to be the value of μ on the two sheets that cross near S^α_i.)

Differential: The differential $\partial : \mathcal{A}(\Lambda, \mathcal{E}) \to \mathcal{A}(\Lambda, \mathcal{E})$ is characterized on the generators of $\mathcal{A}(\Lambda, \mathcal{E})$ by matrix formulas whose precise form depends on the dimension of the associated cell $e^d_\alpha \in \mathcal{E}$. We review these formulas here *in the case that* Λ *does not have swallowtail points.* See also [33, 36].

[2]By polygonal decomposition, we mean a CW-complex decomposition such that the boundary of each 2-cell consists of a sequence of vertices and edges (with repeats allowed).

- *For* 0*-cells:* We choose a linear ordering of the sheets of Λ above e^0_α so that the z-coordinates appear in non-increasing order
$$z(S^\alpha_{\iota(1)}) \geq z(S^\alpha_{\iota(2)}) \geq \cdots \geq z(S^\alpha_{\iota(n)})$$
and use it to place the generators, $a^\alpha_{p,q}$, into a matrix, A, whose (i,j)-entry is $a^\alpha_{\iota(i),\iota(j)}$ when $z(S^\alpha_{\iota(i)}) > z(S^\alpha_{\iota(j)})$ and is 0 otherwise. When ∂ is applied entry-by-entry to A we have
$$\partial A = A^2.$$
- *For* 1*-cells:* After a choice of linear ordering of sheets as above, we place the generators, $b^\alpha_{p,q}$, associated to a 1-cell, e^1_α, into an $n \times n$ matrix, B. In addition, we form $n \times n$ **boundary matrices**, A_- and A_+, associated to the initial and terminal vertices, e^0_- and e^0_+, of e^1_α. The sheets above $e^0_\pm$ are identified with a subset of the sheets above e^1_α (since every sheet of $e^0_\pm$ belongs to the closure of a unique sheet of e^1_α). Using this identification, we place the $e^0_\pm$ generators, $a^\pm_{p,q}$, into the corresponding rows and columns of the $n \times n$ matrices $A_\pm$. Whenever two sheets of e^1_α, S^α_k and $S^\alpha_{k'}$, meet at a cusp point above $e^0_\pm$, we place a 2×2 block of the form
$$N = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$
on the diagonal at the location of the two (possibly non-consecutive) rows and columns of $A_\pm$ that correspond to S^α_k and $S^\alpha_{k'}$ with respect to the linear ordering of sheets of e^1_α. All other entries of $A_\pm$ are 0. The differential on the $b^\alpha_{p,q}$ satisfies
$$\partial B = A_+(I+B) + (I+B)A_-.$$
- *For* 2*-cells:* The sheets above a 2-cell, e^2_α, are already linearly ordered by descending z-coordinate, and using this ordering we place the generators $c^\alpha_{i,j}$ into an $n \times n$-matrix, C. We form $n \times n$ boundary matrices, A_{v_0} and A_{v_1}, associated to the initial and terminal vertices, v_0 and v_1, for e^2_α following the same procedure as in the 1-cell case. Additional boundary matrices $B_1, \ldots, B_l$ are associated to the 1-cells that appear around the boundary of C. We assume that the numbering is such that $B_1, \ldots, B_j$ (resp. $B_{j+1}, \ldots, B_l$) are the boundary matrices for the sequence of edges that appear along the path γ_+ (resp. γ_-) where $\gamma_\pm$ are the two paths that travel around the boundary of e^2_α (in the domain of the characteristic map) from v_0 to v_1. When $v_0 = v_1$ one of $\gamma_\pm$ is constant (as specified by the choice of preferred direction around ∂e^2_α). The B_i are formed using the same procedure as for the boundary matrices associated to vertices except that the 2×2 N blocks that correspond to pairs of cusp sheets are replaced with 2×2 blocks of 0. The differential satisfies
$$\partial C = A_{v_1}C + CA_{v_0} + (I+B_j)^{\eta_j} \cdots (I+B_1)^{\eta_1} + (I+B_l)^{\eta_l} \cdots (I+B_{j+1})^{\eta_{j+1}}$$
where η_i is $+1$ (resp. -1) when the orientation of the B_i edge agrees (resp. disagrees) with the orientation (from v_0 to v_1) of the corresponding path $\gamma_\pm$. Note that since the B_i are strictly upper triangular, $(I+B_i)^{-1} = I + B_i + B_i^2 + \cdots + B_i^{n-1}$.

Remark 4.2. To define ∂ in the case that Λ has swallowtail points, the following additions should be made.

- The definition of the boundary matrices $A_\pm$, A_{v_0}, and A_{v_1} needs to be adjusted when the vertex is a swallowtail point of Λ.
- Whenever a 2-cell contains one of the corners labeled S or T at a swallowtail point, an additional matrix need to be inserted into the product $(I+B_j)\cdots(I+B_1)$ or $(I+B_l)\cdots(I+B_{j+1})$.

These details may be found in [33] or [36] and are not needed for the arguments that follow.




FIGURE 4. Notation for cells in $\mathcal{E}$ (left) and $\mathcal{E}'$ (right).

Observation 4.3. Whenever $\mathcal{E}' \subset \mathcal{E}$ is a (CW) sub-complex, the collection of generators associated to cells of $\mathcal{E}'$ form a sub DGA of $\mathcal{A}_{cell}(\Sigma, \mathcal{E})$. Moreover, if $\mathcal{E}'$ is a decomposition of a curve or curve segment $C \subset M$ such that the restriction of Σ to C (as in Section 2.3) is a 1-dimensional Legendrian knot $\Lambda \subset J^1C$, then this sub-DGA is precisely $\mathcal{A}_{cell}(\Lambda, \mathcal{E}')$.

4.2. **The cellular DGA for compact Legendrian cobordisms.** In extending the definition of the cellular DGA to Legendrian cobordisms, it is most natural to consider compact cobordisms rather than conical cobordisms. The following definition coincides precisely with a standard notion of Legendrian cobordism introduced by Arnold, cf. [1, 2, 3].

Definition 4.4. Given an interval $I \subset [0,1]$ and $\Lambda \subset J^1M$, we write $j^1(1_I \cdot \Lambda)$ for the **Legendrian cylinder** on Λ in $J^1(I \times M)$. It is defined to be the product of $\Lambda \subset J^1M$ with the 0-section in T^*I, i.e.

$$j^1(1_I \cdot \Lambda) = 0_{T^*I} \times \Lambda \subset T^*I \times J^1M \cong J^1(I \times M). \tag{4.1}$$

The notation is chosen to be consistent with that of Definition 2.10 with $1_I : I \to \mathbb{R}_{>0}$ denoting the constant function 1.

Let $\Lambda_0, \Lambda_1 \subset J^1M$. A **compact Legendrian cobordism** from Λ_0 to Λ_1, written $\Sigma : \Lambda_0 \to \Lambda_1$, is a Legendrian surface $\Sigma \subset J^1([0,1] \times M)$ that, for some $\epsilon > 0$, Σ agrees with the Legendrian cylinder $j^1(1_{[0,\epsilon]} \cdot \Lambda_0)$ in $J^1([0,\epsilon] \times M)$ and agrees with $j^1(1_{[1-\epsilon,1]} \cdot \Lambda_1)$ in $J^1([1-\epsilon, 1], \times M)$.

When $\Sigma : \Lambda_0 \to \Lambda_1$ is a compact Legendrian cobordism, we modify the definition of compatible polygonal decomposition for Σ to require that $\pi_x(\Sigma) \cap (\{0\} \times M)$ and $\pi_x(\Sigma) \cap (\{1\} \times M)$ are (CW) sub-complexes of $\mathcal{E}$, that we denote $\mathcal{E}_0$ and $\mathcal{E}_1$. Then, the definition of the cellular DGA extends immediately to give DGAs $\mathcal{A}(\Sigma, \mathcal{E})$ when Σ is a compact Lagrangian cobordism. Moreover, since $\mathcal{E}_0$ and $\mathcal{E}_1$ may be viewed as polygonal decompositions for Λ_0 and Λ_1, as in the Observation 4.3, we have inclusion maps

$$i_0 : \mathcal{A}(\Lambda_0, \mathcal{E}_0) \to \mathcal{A}(\Sigma, \mathcal{E}) \quad \text{and} \quad i_1 : \mathcal{A}(\Lambda_1, \mathcal{E}_1) \to \mathcal{A}(\Sigma, \mathcal{E}). \tag{4.2}$$

4.3. **The DGA of a product cobordism.** Given a 1-dimensional Legendrian with compatible polygonal decomposition, $(\Lambda, \mathcal{E})$, we now compute the DGA of the product cobordism $\Sigma = j^1(1_{[0,1]} \cdot \Lambda) \subset J^1([0,1] \times M)$. This will be a useful ingredient in a few later arguments.

As a preliminary, use $\mathcal{E}$ to form the **product decomposition**, $\mathcal{E}'$, for $\pi_x(\Sigma) = [0,1] \times \pi_x(\Lambda)$ as follows: For each d-cell $e^d_\alpha \in \mathcal{E}$ (here, $d = 0$ or 1), decompose

$$[0,1] \times e^d_\alpha = \{0\} \times e^d_\alpha \sqcup (0,1) \times e^d_\alpha \sqcup \{1\} \times e^d_\alpha$$

so that $\{k\} \times e^d_\alpha$ are d-cells in $\mathcal{E}'$, for $k = 0, 1$, and $(0,1) \times e^d_\alpha$ is a $(d+1)$-cell in $\mathcal{E}'$. Orient 1-cells of the form $\{k\} \times e^1_\beta$ in the same direction as e^1_β, and those of the form $(0,1) \times e^0_\alpha$ using the standard orientation of $(0,1)$. For each 2-cell, $(0,1) \times e^1_\beta$ choose the initial and terminal vertices

v_0 and v_1 to be $\{0\} \times e^0_-$ and $\{1\} \times e^0_+$ where e^0_- and e^0_+ are the initial and terminal vertices of e^1_β. See Figure 4.

Let us fix notation for the generators of the cellular DGA $\mathcal{A}(\Sigma, \mathcal{E}')$. Observe that for any $e^d_\alpha \in \mathcal{E}$, the sheets of Λ above e^d_α are in bijection with the sheets of Σ above any of the $\{k\} \times e^d_\alpha$ or $(0,1) \times e^d_\alpha$ cells of $\mathcal{E}'$, so the generators associated to these cells are also in bijection with those of e^d_α.

- For a 0-cell $e^0_\alpha \in \mathcal{E}$ with corresponding generators $a^\alpha_{i,j}$ we notate the generators corresponding to $\{0\} \times e^0_\alpha$, $\{1\} \times e^0_\alpha$, and $(0,1) \times e^0_\alpha$ as $i_0(a^\alpha_{i,j})$, $i_1(a^\alpha_{i,j})$, and $b^\alpha_{i,j}$.
- For a 1-cell $e^1_\beta \in \mathcal{E}$ with corresponding generators $b^\beta_{i,j}$ we notate the generators corresponding to $\{0\} \times e^1_\beta$, $\{1\} \times e^1_\beta$, and $(0,1) \times e^1_\beta$ as $i_0(b^\beta_{i,j})$, $i_1(b^\beta_{i,j})$, and $c^\beta_{i,j}$.

In Proposition 4.5, the DGA of the product cobordism Σ is described as a mapping cylinder DGA. We now briefly review the relevant definitions, referring the reader to [32, Section 2] for a more thorough treatment in the present algebraic setting of triangular DGAs over $\mathbb{Z}/2$. Let $f : (\mathcal{A}, \partial_\mathcal{A}) \to (\mathcal{B}, \partial_\mathcal{B})$ be a DGA map between based DGAs. The **standard mapping cylinder DGA** of f is $(\mathcal{A} * \widehat{\mathcal{A}} * \mathcal{B}, \partial_\Gamma)$ where $\widehat{\mathcal{A}}$ has generators $\{\widehat{a_i}\}$ in correspondence with the generators $\{a_i\}$ of $\mathcal{A}$ but with the degree shift $|\widehat{a_i}| = |a_i| + 1$. The differential, ∂_Γ, satisfies

$$\partial_\Gamma|_\mathcal{A} = \partial_\mathcal{A}, \quad \partial_\Gamma|_\mathcal{B} = \partial_\mathcal{B}, \quad \text{and} \quad \partial_\Gamma(\widehat{a_i}) = f(a_i) + a_i + \Gamma \circ \partial_\mathcal{A}(a_i)$$

where $\Gamma : \mathcal{A} \to \mathcal{A} * \widehat{\mathcal{A}} * \mathcal{B}$ is the unique $(f, id_\mathcal{A})$-derivation satisfying $\Gamma(a_i) = \widehat{a_i}$.

Proposition 4.5. *Given $(\Lambda, \mathcal{E})$ with $\Lambda \subset J^1M$ and $\dim \Lambda = 1$, let $(\Sigma, \mathcal{E}')$ be the product cobordism, $\Sigma = j^1(1_{[0,1]} \cdot \Lambda) \subset J^1([0,1] \times M)$ equipped with the product decomposition, $\mathcal{E}'$.*

1. *For $k = 0, 1$, the maps $i_k : \mathcal{A}(\Lambda, \mathcal{E}) \to \mathcal{A}(\Sigma, \mathcal{E}')$ that extend the correspondence of generators are DGA isomorphisms from $\mathcal{A}(\Lambda, \mathcal{E})$ onto the sub-DGAs $\mathcal{A}_k \subset \mathcal{A}(\Sigma, \mathcal{E}')$ associated to the subcomplexes $\{k\} \times \mathcal{E} \subset \mathcal{E}'$.*
2. *Identifying the sub-algebra of $\mathcal{A}(\Sigma, \mathcal{E}')$ generated by the $b^\alpha_{i,j}$ and $c^\beta_{i,j}$ with $\widehat{\mathcal{A}_0}$ using the grading preserving bijection $\widehat{i_0(a^\alpha_{i,j})} \leftrightarrow b^\alpha_{i,j}$ and $\widehat{i_0(b^\beta_{i,j})} \leftrightarrow c^\beta_{i,j}$ gives a DGA isomorphism*

$$\varphi : (\mathcal{A}(\Sigma, \mathcal{E}'), \partial) \xrightarrow{\cong} (\mathcal{A}_0 * \widehat{\mathcal{A}_0} * \mathcal{A}_1, \partial_\Gamma)$$

with the standard mapping cylinder DGA of the map $i_1 \circ i_0^{-1} : \mathcal{A}_0 \to \mathcal{A}_1$.

Proof. (1) is obvious. For (2), φ is clearly an algebra isomorphism which is the identity on the $\mathcal{A}_k$, so we just to check that $\partial_\Gamma \circ \varphi(x) = \varphi \circ \partial(x)$ when $x = b^\alpha_{i,j}$ or $c^\beta_{i,j}$. From the definition of ∂_Γ, we have

$$\partial_\Gamma \widehat{i_0(a^\alpha_{i,j})} = i_1(a^\alpha_{i,j}) + i_0(a^\alpha_{i,j}) + \Gamma \circ \partial_\Lambda a^\alpha_{i,j} \tag{4.3}$$

$$\partial_\Gamma \widehat{i_0(b^\beta_{i,j})} = i_1(b^\beta_{i,j}) + i_0(b^\beta_{i,j}) + \Gamma \circ \partial_\Lambda b^\beta_{i,j} \tag{4.4}$$

where ∂_Λ is the differential on $\mathcal{A}(\Lambda, \mathcal{E})$ and $\Gamma : \mathcal{A}(\Lambda, \mathcal{E}) \to \mathcal{A}_1 * \widehat{\mathcal{A}_0} * \mathcal{A}_0$ is the (i_1, i_0)-derivation satisfying $\Gamma(x) = \widehat{i_0(x)}$ on generators.

Using matrix notation, this allows us to compute

$$\begin{aligned}
\varphi \circ \partial(B_\alpha) &= \varphi\left(i_1(A_\alpha)(I + B_\alpha) + (I + B_\alpha) i_0(A_\alpha)\right) \\
&= i_1(A_\alpha) + i_1(A_\alpha)\widehat{i_0(A_\alpha)} + i_0(A_\alpha) + \widehat{i_0(A_\alpha)} i_0(A_\alpha) \\
&= i_1(A_\alpha) + i_0(A_\alpha) + \Gamma(A^2_\alpha) \\
&= \partial_\Gamma(\widehat{i_0(A_\alpha)}) = \partial_\Gamma \circ \varphi(B_\alpha),
\end{aligned}$$

and

$$\begin{aligned}
\partial_\Gamma \circ \varphi(C_\beta) &= \partial_\Gamma(\widehat{i_0(B_\beta)}) = i_1(B_\beta) + i_0(B_\beta) + \Gamma(\partial B_\beta) \\
&= i_1(B_\beta) + i_0(B_\beta) + \Gamma(A_+(I + B_\beta) + (I + B_\beta)A_-) \\
&= i_1(B_\beta) + i_0(B_\beta) + \Gamma(A_+)(I + i_0(B_\beta)) + i_1(A_+)\widehat{i_0(B_\beta)} \\
&\quad + \widehat{i_0(B_\beta)}i_0(A_-) + (I + i_1(B_\beta))\Gamma(A_-) \\
&= \varphi\Big(i_1(B_\beta) + i_0(B_\beta) + B_+(I + i_0(B_\beta)) + i_1(A_+)C_\beta \\
&\quad + C_\beta i_0(A_-) + (I + i_1(B_\beta))B_- \Big) \\
&= \varphi\Big(i_1(A_+)C_\beta + C_\beta i_0(A_-) + (I + B_+)(I + i_0(B_\beta)) + (I + i_1(B_\beta))(I + B_-) \Big) \\
&= \varphi \circ \partial(C_\beta).
\end{aligned}$$

At the fourth equality, it should be observed that when A_- and A_+ are the boundary matrices for e^1_β associated to the vertices e^0_- and e^0_+, the boundary matrices B_- and B_+ for $(0,1) \times e^1_\beta$ associated to the edges $(0,1) \times e^0_-$ and $(0,1) \times e^0_+$ indeed satisfy $\varphi(B_\pm) = \Gamma(A_\pm)$. [Note that when a $\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ block appears on the diagonal of $A_\pm$ due to two sheets of e^1_β meeting at a cusp above $e^0_\pm$, since $\Gamma(1) = 0$ (any derivation has this property), the appropriate $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ block will appear in $\Gamma(A_\pm)$.] □

4.4. **Immersed DGA maps from cobordisms via the cellular DGA.** Our aim is to now define a cellular version of the immersed LCH functor $F : \mathfrak{Leg}^\rho_{im} \to \mathfrak{DGA}^\rho_{im}$ from Corollary 2.13. Recall that the domain category for F has ρ-graded Legendrians in J^1M equipped with regular metrics as objects and has (conical Legendrian isotopy classes of) conical Legendrian cobordisms in $J^1(\mathbb{R}_{>0} \times M)$ (equivalently, good immersed Lagrangian cobordisms in $Symp(J^1M)$) as morphisms. For the cellular construction we instead work with a category of compact cobordisms.

With $M = \mathbb{R}$ or S^1 and $\rho \geq 0$ fixed, define a **cellular Legendrian cobordism category**, denoted $\mathfrak{Leg}^\rho_{cell}$, whose objects are triples $(\Lambda, \mathcal{E}, \mu)$ consisting of a 1-dimensional Legendrian link, $\Lambda \subset J^1M$, together with a choice, $\mathcal{E}$, of compatible polygonal decomposition, and a choice, μ, of $\mathbb{Z}/\rho$-valued Maslov potential. Morphisms from $(\Lambda_0, \mathcal{E}_0, \mu_0)$ to $(\Lambda_1, \mathcal{E}_1, \mu_1)$ are equivalence classes of compact Legendrian cobordisms $\Sigma \subset J^1([0,1] \times M)$ from Λ_- to Λ_+ having generic base and front projection and equipped with a $\mathbb{Z}/\rho$-valued Maslov potential extending μ_0 and μ_1. Here, two cobordisms are considered equivalent if their front and base projections are **combinatorially equivalent**. That is, Σ and Σ' are equivalent if there is a homeomorphism $\phi : ([0,1] \times M) \times \mathbb{R} \to ([0,1] \times M) \times \mathbb{R}$ with $\phi(\pi_{xz}(\Sigma)) = \phi(\pi_{xz}(\Sigma'))$ that is a composition of (i) a homeomorphism, ϕ_1, of the $[0,1] \times M$ factor, and (ii) a homeomorphism, ϕ_2, that preserves the $[0,1] \times M$ factor. Moreover, ϕ_1 and ϕ_2 should be isotopic to the identity and equal to the identity in a neighborhood of the boundary.

Remark 4.6.
(1) As with the category $\mathfrak{Leg}^\rho_{im}$, in the definition of morphisms Σ is not equipped with any additional structure beyond a choice of Maslov potential, eg. Σ is not equipped with a polygonal decomposition.
(2) In contrast to $\mathfrak{Leg}^\rho_{im}$, we do NOT allow general Legendrian isotopies of Σ in the equivalence relation used to define morphisms. This is because from the initial definition of our cellular LCH functor we will not check directly that the assignment of immersed DGA maps to Legendrian cobordisms factors through general Legendrian isotopies. However,

it will later be established in Corollary 4.11 that the induced immersed DGA maps (considered up to immersed homotopy) are indeed Legendrian invariants of Σ.

The next proposition defines the **cellular LCH functor**, F_{cell}.

Proposition 4.7. *There is a well-defined contravariant functor $F_{cell} : \mathfrak{Leg}^{\rho}_{cell} \to \mathfrak{DGA}^{\rho}_{im}$ given by*

$$\begin{aligned} (\Lambda, \mathcal{E}) &\rightsquigarrow \mathcal{A}_{cell}(\Lambda, \mathcal{E}), \\ \Sigma : (\Lambda_0, \mathcal{E}_0) \to (\Lambda_1, \mathcal{E}_1) &\rightsquigarrow \left[\mathcal{A}_{cell}(\Lambda_1, \mathcal{E}_1) \xrightarrow{i_1} \mathcal{A}_{cell}(\Sigma, \mathcal{E}') \xleftarrow{i_0} \mathcal{A}_{cell}(\Lambda_0, \mathcal{E}_0)\right] \end{aligned}$$

where $\mathcal{E}'$ is any choice of compatible polygonal decomposition for Σ that restricts to $\mathcal{E}_k$ on $\{k\} \times M$, $k = 0, 1$, and the maps i_0 and i_1 are as in (4.2).

Proof. To see that F_{cell} is well-defined it is enough to verify independence of the choice of $\mathcal{E}'$. (Since modifying $(\Sigma, \mathcal{E}')$ in a manner that preserves the combinatorics of the front and base projections does not affect the cellular DGA.)

In [33, Section 4.2-4.4], it is shown in the case when Σ is a closed surface that any compatible polygonal decompositions $\mathcal{E}'$ and $\mathcal{E}''$ for Σ can be made the same after some sequence of the following modifications and their inverses (see [33] for the precise meanings).

(1) Subdivide a 1-cell.
(2) Subdivide a 2-cell.
(3) Delete a 1-valent edge.
(4) Switch the S and T decorations at a swallowtail point.

The proof extends with minor adjustments to the case of a compact cobordism. Moreover, if $\mathcal{E}'$ and $\mathcal{E}''$ agree above $\partial([0,1] \times M)$, then the polygonal decomposition of $\partial([0,1] \times M)$ can be left unchanged throughout the sequence of modifications. [To see this, follow the proof of [33, Theorem 4.1], but treat all 0-cells (resp. 1-cells) in $\partial([0,1] \times M)$ as if they belong to what is notated there as Σ_2 (resp. Σ_1).]

If $\mathcal{E}'$ and $\mathcal{E}''$ are related by one of the modifications (1)-(4), then [33, Theorems 4.2-4.5] provide stable tame isomorphisms $\varphi : \mathcal{A}_{cell}(\Sigma, \mathcal{E}') * S' \to \mathcal{A}_{cell}(\Sigma, \mathcal{E}'') * S''$. Moreover, in all cases φ can be seen to restrict to the identity on the sub-algebras $\mathcal{A}_{cell}(\Lambda_k, \mathcal{E}_k)$, $k = 0, 1$. [For (1)-(3), φ is defined using the construction of Proposition 2.1, and generators from $\mathcal{A}_{cell}(\Lambda_k, \mathcal{E}_k)$ are never among the generators that are canceled. For (4), φ is the identity on all generators except for 1-cells with endpoints at swallowtail points.] Thus, we will have identities $\varphi \circ i_k = i_k$, $k = 0, 1$, so that φ provides the required immersed DGA homotopy.

To verify that F_{cell} preserves composition, suppose that $\mathcal{E}'$ and $\mathcal{E}''$ are compatible polygonal decompositions for a pair of composable (compact) cobordisms, $\Sigma' : (\Lambda_0, \mathcal{E}_0) \to (\Lambda_1, \mathcal{E}_1)$ and $\Sigma'' : (\Lambda_0, \mathcal{E}_0) \to (\Lambda_1, \mathcal{E}_1)$. Since $\mathcal{E}'$ and $\mathcal{E}''$ agree with $\mathcal{E}_1$ along Λ_1, they can be glued to form a well-defined compatible polygonal decomposition, $\mathcal{E}' \cup \mathcal{E}''$, on $\Sigma'' \circ \Sigma'$. Moreover, when this decomposition is used for computing it is readily verified from Definition 2.4 that $F_{cell}(\Sigma'' \circ \Sigma') = F_{cell}(\Sigma') \circ F_{cell}(\Sigma'')$.

Finally, we check that F_{cell} preserves identities. The identity morphism in $Hom_{\mathfrak{Leg}^{\rho}_{cell}}((\Lambda, \mathcal{E}), (\Lambda, \mathcal{E}))$ is the product cobordism, $\Sigma = j^1(1_{[0,1]} \cdot \Lambda)$, and for computing $F_{cell}(\Sigma)$ we equip it with the product decomposition, $\mathcal{E}'$. The computation from Proposition 4.5 shows that the morphism $F_{cell}(\Sigma)$ is the immersed homotopy class of the immersed DGA map

$$D = \left(\mathcal{A}(\Lambda, \mathcal{E}) \xrightarrow{i_1} (\mathcal{A}_0 * \widehat{\mathcal{A}_0} * \mathcal{A}_1, \partial_\Gamma) \xleftarrow{i_0} \mathcal{A}(\Lambda, \mathcal{E})\right)$$

where $\mathcal{A}_0$ and $\mathcal{A}_1$ are two copies of $\mathcal{A}(\Lambda, \mathcal{E})$ and $(\mathcal{A}_0 * \widehat{\mathcal{A}_0} * \mathcal{A}_1, \partial_\Gamma)$ is the standard mapping cylinder DGA for the identity map on $\mathcal{A}(\Lambda, \mathcal{E})$. On the other hand, the identity morphism in $\mathfrak{DGA}^{\rho}_{im}$ is just $[I]$ where I is the immersed DGA map

$$I = \left(\mathcal{A}(\Lambda, \mathcal{E}) \xrightarrow{id} \mathcal{A}(\Lambda, \mathcal{E}) \xleftarrow{id} \mathcal{A}(\Lambda, \mathcal{E})\right).$$

Note that the immersed map D is obtained from two copies of I precisely as in the statement of Proposition 2.5, which therefore shows that D is immersed homotopic to $I \circ I = I$ as required. □

4.5. **Isomorphism between the cellular and immersed LCH functors.** In this section, we state a precise relationship between the immersed and cellular LCH functors, F and F_{cell}.

The functors $F : \mathfrak{Leg}^{\rho}_{im} \to \mathfrak{DGA}^{\rho}_{im}$ and $F_{cell} : \mathfrak{Leg}^{\rho}_{cell} \to \mathfrak{DGA}^{\rho}_{im}$ have slightly different domain categories. In $\mathfrak{Leg}^{\rho}_{im}$ (resp. $\mathfrak{Leg}^{\rho}_{cell}$), the objects are (closed, 1-dimensional) ρ-graded Legendrians in J^1M equipped with a suitable choice of metric (resp. polygonal decomposition), while morphisms are equivalence classes of conical (resp. compact) ρ-graded Legendrian cobordisms. For comparing the two functors, it is convenient to work with a third domain category whose objects again are 1-dimensional Legendrians equipped with some additional data: an *admissible transverse decomposition*, $\mathcal{E}_{\pitchfork}$. The role of $\mathcal{E}_{\pitchfork}$ is to simultaneously specify a compatible polygonal decomposition of Λ, $\mathcal{E}_{||}$, and a modification of Λ by a Legendrian isotopy to $\widetilde{\Lambda}$ so that the cellular DGA of $(\Lambda, \mathcal{E}_{||})$ and the LCH DGA of $\widetilde{\Lambda}$ are related by a canonical homotopy equivalence (in fact, a stable tame isomorphism). This is precisely the role of the *transverse square decompositions* from [34, Section 3] in the case of closed 2-dimensional Legendrians.

Definition 4.8. With $\rho \in \mathbb{Z}_{\geq 0}$ and 1-dimensional M fixed, define the **transverse Legendrian category**, $\mathfrak{Leg}^{\rho}_{\pitchfork}$, to have objects $(\Lambda, \mathcal{E}_{\pitchfork}, \mu)$ where $\Lambda \subset J^1M$ is Legendrian, μ is a $\mathbb{Z}/\rho$-valued Maslov potential, and $\mathcal{E}_{\pitchfork}$ is an **admissible transverse decomposition** for Λ as defined in [30, Appendix A, Definition A.1]. Morphisms are compact ρ-graded Legendrian cobordisms up to combinatorial equivalence of front and base projections (as in Section 4.4).

An admissible transverse decomposition of Λ is a cellular decomposition of a neighborhood of the base projection, $\pi_x(\Lambda) \subset M$, that has cells transverse to the projection of the singular set of $\pi_{xz}(\Lambda)$ and is subject to some technical restrictions that are useful for the detailed proof of Proposition 4.9 but are not relevant when applying the proposition. For use of Proposition 4.9 in the remainder of the article, it will be sufficient to note that any 1-dimensional Legendrian $\Lambda \subset J^1M$ with generic front and base projections admits an admissible transverse decomposition.

In [30, Appendix A], two functors are constructed

$$\begin{array}{c} G_{std} : \mathfrak{Leg}^{\rho}_{\pitchfork} \to \mathfrak{Leg}^{\rho}_{im}, \\ (\Lambda, \mathcal{E}_{\pitchfork}) \mapsto (\widetilde{\Lambda}, \widetilde{g}) \end{array} \qquad \text{and} \qquad \begin{array}{c} G_{||} : \mathfrak{Leg}^{\rho}_{\pitchfork} \to \mathfrak{Leg}^{\rho}_{cell}, \\ (\Lambda, \mathcal{E}_{\pitchfork}) \mapsto (\Lambda, \mathcal{E}_{||}) \end{array}$$

where $\widetilde{\Lambda}$, called the **standard geometric model** for Λ with respect to $\mathcal{E}_{\pitchfork}$, is Legendrian isotopic to Λ with front and base projections combinatorially equivalent to those of Λ. The actions on morphisms are as follows:

- The morphism spaces in $\mathfrak{Leg}^{\rho}_{\pitchfork}$ and $\mathfrak{Leg}^{\rho}_{cell}$ only depend on Λ and are the same in both categories, and $G_{||}$ acts as the identity on morphisms.
- Given a compact cobordism Σ from Λ_0 to Λ_1, we can choose any conical Legendrian cobordism $\Sigma' \subset J^1(\mathbb{R}_{>0} \times M)$ from $G_{std}(\Lambda_0)$ to $G_{std}(\Lambda_1)$ whose front and base projections are combinatorially equivalent to Σ (after truncating the conical ends). The conical Legendrian isotopy type of Σ' depends only on Σ, and we have $G_{std}([\Sigma]) = [\Sigma']$.

The standard geometric model, $G_{std}(\Lambda, \mathcal{E}_{\pitchfork}) = (\widetilde{\Lambda}, \widetilde{g})$, is constructed so that the Reeb chords of $\widetilde{\Lambda}$ closely match the generators of the cellular DGA, $\mathcal{A}(\Lambda, \mathcal{E}_{||})$. In fact, $\mathcal{A}(\widetilde{\Lambda}, \widetilde{g})$ is isomorphic to a stabilization of $\mathcal{A}(\Lambda, \mathcal{E}_{||})$, and an explicitly defined homotopy equivalence

$$q_{\Lambda} : \mathcal{A}(\widetilde{\Lambda}, \widetilde{g}) \to \mathcal{A}(\Lambda, \mathcal{E}_{||})$$

that we call the **canonical quotient map** is identified in [30, Section A.4]. As q_{Λ} is invertible in the DGA homotopy category $\mathfrak{DGA}^{\rho}_{hom}$, it follows that the immersed map

$$Q_{\Lambda} = \left[\mathcal{A}(\widetilde{\Lambda}, \widetilde{g}) \xrightarrow{q_{\Lambda}} \mathcal{A}(\Lambda, \mathcal{E}_{||}) \xleftarrow{id} \mathcal{A}(\Lambda, \mathcal{E}_{||})\right] \in \mathfrak{DGA}^{\rho}_{im}$$

(that is the image of q_Λ under the functor from Proposition 2.7) is an isomorphism in $\mathfrak{DGA}^\rho_{im}$.

Proposition 4.9. *The canonical quotient map construction,*

$$(\Lambda, \mathcal{E}_{\pitchfork}) \in \mathfrak{Leg}^\rho_{\pitchfork} \mapsto Q_\Lambda \in Hom_{\mathfrak{DGA}^\rho_{im}}(\mathcal{A}(\widetilde{\Lambda}, \widetilde{g}), \mathcal{A}(\Lambda, \mathcal{E}_{||}))$$

gives an isomorphism (invertible natural transformation) from the functor that is the composition

$$\mathfrak{Leg}^\rho_{\pitchfork} \overset{G_{std}}{\to} \mathfrak{Leg}^\rho_{im} \overset{F}{\to} \mathfrak{DGA}^\rho_{im}.$$


to the functor

$$\mathfrak{Leg}^\rho_{\pitchfork} \overset{G_{||}}{\to} \mathfrak{Leg}^\rho_{cell} \overset{F_{cell}}{\to} \mathfrak{DGA}^\rho_{im}.$$

That is, the Q_Λ are invertible, and for any compact Legendrian cobordism $\Sigma : (\Lambda_-, \mathcal{E}^{\pitchfork}_-) \to (\Lambda_+, \mathcal{E}^{\pitchfork}_+)$ with generic base and front projection we have a commutative diagram in $\mathfrak{DGA}^\rho_{im}$,

$$\begin{array}{ccc} \mathcal{A}(\widetilde{\Lambda}_+, \widetilde{g}_+) & \xrightarrow{F\circ G_{std}(\Sigma)} & \mathcal{A}(\widetilde{\Lambda}_-, \widetilde{g}_-) \\ \downarrow{\scriptstyle Q_{\Lambda_+}} & & \downarrow{\scriptstyle Q_{\Lambda_-}} \\ \mathcal{A}_{cell}(\Lambda_+, \mathcal{E}^{||}_+) & \xrightarrow[F_{cell}\circ G_{||}(\Sigma)]{} & \mathcal{A}_{cell}(\Lambda_-, \mathcal{E}^{||}_-) \end{array} . \tag{4.5}$$

In summary, the following diagram is commutative up to a canonical isomorphism of functors:



A detailed proof of Proposition 4.9 is presented over the course of [30, Appendices A and B]. It is an extension of the isomorphism from [34, 35] for closed surfaces, and we provide here a summary of the key points.

Sketch of Proof. The technical requirements in the definition of admissible transverse decomposition are designed so that, with arbitrary $(\Lambda_-, \mathcal{E}^{\pitchfork}_-), (\Lambda_+, \mathcal{E}^{\pitchfork}_+) \in \mathfrak{Leg}^\rho_{\pitchfork}$ fixed, for any (compact) $\Sigma : \Lambda_- \to \Lambda_+$ the construction of [34, Section 3] for closed surfaces generalizes to produce a transverse square decomposition, $\mathcal{E}^{\pitchfork}_\Sigma$, that in a closed collar neighborhood of $\partial\Sigma = \Lambda_- \cup \Lambda_+$ agrees with the product of $\mathcal{E}^{\pitchfork}_-$ and $\mathcal{E}^{\pitchfork}_+$ with the standard CW decomposition of a closed interval. Applying the square-by-square coordinate construction from [35] using $\mathcal{E}^{\pitchfork}_\Sigma$ produces an initial geometric model for Σ, $(\overline{\Sigma}, \overline{g})$. The restriction of $(\overline{\Sigma}, \overline{g})$ to the negative boundary only depends on $(\Lambda_-, \mathcal{E}^{\pitchfork}_-)$, and this restriction is the definition of $G_{std}(\Lambda_-, \mathcal{E}^{\pitchfork}_-) = (\widetilde{\Lambda}_-, \widetilde{g}_-)$. Similarly, $G_{std}(\Lambda_+, \mathcal{E}^{\pitchfork}_+)$ is the positive boundary of $(\overline{\Sigma}, \overline{g})$. In addition, [34, Section 3] constructs from $\mathcal{E}^{\pitchfork}_\Sigma$ a compatible decomposition $\mathcal{E}^{||}_\Sigma$ whose restriction to the boundary is independent of Σ and matches the $\mathcal{E}^{||}_\pm$ that appear in $G_{||}(\Lambda_\pm, \mathcal{E}^{\pitchfork}_\pm) = (\Lambda_\pm, \mathcal{E}^{||}_\pm)$. A technical point is that for computing the immersed DGA map $F \circ G_{std}(\Sigma) = [\mathcal{A}(\widetilde{\Lambda}_+, \widetilde{g}_+) \to \mathcal{A}(\widetilde{\Sigma}, \widetilde{g}) \hookleftarrow \mathcal{A}(\widetilde{\Lambda}_+, \widetilde{g}_+)]$ we need to pick a specific $(\widetilde{\Sigma}, \widetilde{g})$ that is a conical version of Σ. This is done [30, Appendix B] by carefully modifying $(\overline{\Sigma}, \overline{g})$ in the collar neighborhood of the boundary to be a Morse minimum cobordism and then applying the Construction 2.14. As verified in [30, Appendix B], the square-by-square computation of LCH from [34] still applies to produce an explicit stable tame isomorphism relating $\mathcal{A}(\widetilde{\Sigma}, \widetilde{g})$ and $\mathcal{A}(\Sigma, \mathcal{E}^{||}_\Sigma)$. Moreover, [30, Appendix A.6] uses this stable tame isomorphism to verify the commutativity of (4.5). □

Remark 4.10. An arbitrary Legendrian $\Lambda \subset J^1M$ is related to one in the image of G_{std} by a Legendrian isotopy (that preserves the combinatorial appearance of the front projection). Thus Proposition 4.9 allows the cellular DGA to be used for computing immersed maps $F(\Sigma)$ induced by a conical Legendrian cobordism $\Sigma : \Lambda_- \to \Lambda_+$ after possibly changing the Legendrians at the ends from $\Lambda_\pm$ to $\Lambda'_\pm$ by Legendrian isotopy. Composing Σ with cobordisms $\Sigma_+ : \Lambda_+ \to \Lambda'_+$ and $\Sigma_- : \Lambda'_- \to \Lambda_-$ induced by the isotopies modifies Σ to a cobordism $\Sigma' : \Lambda'_- \to \Lambda'_+$ to which the cellular computation of Proposition 4.9 can be applied directly. Moreover, if one wants to work with the original $\Lambda_\pm$, then the maps $F(\Sigma_-)$ and $F(\Sigma_+)$ can be explicitly computed as in [16, Section 6].

The following corollary of Proposition 4.9 shows that the immersed DGA maps associated to a compact cobordism Σ by the cellular functor, F_{cell}, only depend on the Legendrian isotopy type of Σ.

Corollary 4.11. *Given $(\Lambda_0, \mathcal{E}_0), (\Lambda_1, \mathcal{E}_1) \in \mathfrak{Leg}^\rho_{cell}$ and compact cobordisms Σ and Σ'. If Σ and Σ' are Legendrian isotopic rel. boundary, then $F_{cell}(\Sigma) = F_{cell}(\Sigma')$ in $\mathfrak{DGA}^\rho_{im}$.*

Proof. **Case 1.** Suppose $(\Lambda_k, \mathcal{E}_k) = G_{||}(\Lambda_k, \mathcal{E}_k^\pitchfork)$ for some admissible transverse decompositions $\mathcal{E}_k^\pitchfork$ for Λ_k, $k = 0, 1$.

In this case, Proposition 4.9 allows us to compute in $\mathfrak{DGA}^\rho_{im}$,

$$F_{cell}(\Sigma) = Q_{\Lambda_1} \circ F(G_{std}(\Sigma)) \circ Q^{-1}_{\Lambda_0} = Q_{\Lambda_1} \circ F(G_{std}(\Sigma')) \circ Q^{-1}_{\Lambda_0} = F_{cell}(\Sigma').$$

[At the second equality we used that $F(G_{std}(\Sigma))$ only depends on the conical Legendrian isotopy type of $G_{std}(\Sigma)$.]

Case 2. For general $\mathcal{E}_0$ and $\mathcal{E}_1$.

In this case, fix some choice of admissible transverse decompositions $\mathcal{E}_k^\pitchfork$ for Λ_k, $k = 0, 1$. Let I_k denote the product cobordism $j^1(1_{[0,1]} \cdot \Lambda_k)$ viewed as a morphism in $Hom_{\mathfrak{Leg}^\rho_{cell}}((\Lambda_k, \mathcal{E}_k), (\Lambda_k, G_{||}(\mathcal{E}_k^\pitchfork))$, and let $\Sigma_\pitchfork$ and $\Sigma'_\pitchfork$ denote Σ and Σ' viewed as morphisms in $Hom_{\mathfrak{Leg}^\rho_{cell}}((\Lambda_0, G_{||}(\mathcal{E}_0^\pitchfork)), (\Lambda_1, G_{||}(\mathcal{E}_1^\pitchfork))$. Using Case 1, we compute

$$\begin{aligned} F_{cell}(\Sigma) = F_{cell}(I_1^{-1} \circ \Sigma_\pitchfork \circ I_0) &= F_{cell}(I_1^{-1}) \circ F_{cell}(\Sigma_\pitchfork) \circ F_{cell}(I_0) \\ &= F_{cell}(I_1^{-1}) \circ F_{cell}(\Sigma'_\pitchfork) \circ F_{cell}(I_0) = F_{cell}(\Sigma'). \end{aligned}$$

□

As in Section 3.1, given an immersed DGA map $M = \left(\mathcal{A}_1 \xrightarrow{f} \mathcal{B} \xleftarrow{i} \mathcal{A}_2\right)$ we write $I(M) = Im(i^* \times f^*) \subset Aug_\rho(\mathcal{A}_1)/\sim \times Aug_\rho(\mathcal{A}_2)/\sim$ for the *induced augmentation set* of M. According to Proposition 3.2, $I(M)$ only depends on the immersed homotopy class of M.

Corollary 4.12. *Let $\Sigma \subset J^1([0,1] \times M)$ be a compact Legendrian cobordism from Λ_0 to Λ_1. Let $\mathcal{E}_k^\pitchfork$ be an admissible transverse decomposition for Λ_k, $k = 0, 1$, and write $G_{std}(\Lambda_k, \mathcal{E}_k^\pitchfork) = (\widetilde{\Lambda}_k, \widetilde{g}_k)$ and $G_{||}(\Lambda_k, \mathcal{E}_k^\pitchfork) = (\Lambda_k, \mathcal{E}_k^{||})$. Then, the induced augmentation sets $I_\Sigma := I(F(G_{std}(\Sigma)))$ and $I_\Sigma^{cell} := I(F_{cell}(G_{||}(\Sigma)))$ are related by*

$$(q^*_{\Lambda_0} \times q^*_{\Lambda_1})\big(I_\Sigma^{cell}\big) = I_\Sigma \tag{4.6}$$

where

$$q^*_{\Lambda_0} \times q^*_{\Lambda_1} : Aug_\rho(\Lambda_0, \mathcal{E}_0^{||})/\sim \times Aug_\rho(\Lambda_1, \mathcal{E}_1^{||})/\sim \xrightarrow{\cong} Aug_\rho(\widetilde{\Lambda}_0, \widetilde{g}_0)/\sim \times Aug_\rho(\widetilde{\Lambda}_1, \widetilde{g}_1)/\sim$$

is the bijection induced by the canonical quotient maps, $q_{\Lambda_k} : \mathcal{A}(\widetilde{\Lambda}_k, \widetilde{g}_k) \to \mathcal{A}(\Lambda_k, \mathcal{E}_k^{||})$.

Proof. Applying the functor from Proposition 3.6 to (4.5) produces the diagram of relations

$$\begin{array}{ccc} Aug(\widetilde{\Lambda}_1), \widetilde{g}_1)/\sim & \xleftarrow{I_\Sigma} & Aug(\widetilde{\Lambda}_0, \widetilde{g}_0)/\sim \\ {\scriptstyle I(Q_{\Lambda_1})}\uparrow & & \uparrow{\scriptstyle I(Q_{\Lambda_0})} \\ Aug(\Lambda_1, \mathcal{E}_1^{||})/\sim & \xleftarrow{I_\Sigma^{cell}} & Aug(\Lambda_0, \mathcal{E}_0)/\sim \end{array} \tag{4.7}$$

As noted in 3.1.1, the induced augmentation sets associated to Q_{Λ_k} are just the graphs of $q^*_{\Lambda_k}$, so Observation 3.5 (2) and (3) can be applied to arrive at (4.6). □

5. Computations via Morse complex families

With Corollary 4.12 we have seen that for a compact Legendrian cobordism, Σ, the two versions of the induced augmentation sets, I_Σ^{cell} and I_Σ, defined via the cellular and ordinary versions of the LCH functor, F_{cell} and F, are equivalent invariants of Σ (in a canonical way). In this section, we focus on obtaining methods for computing I_Σ^{cell} based on an extension of correspondences between augmentations and Morse complex families (abbr. MCFs), cf. [23, 25, 36], to the case of 2-dimensional Legendrian cobordisms. In Section 5.1, we review the definition of MCFs and include the present context of 2-dimensional Legendrian cobordisms. In Section 5.2, we make use of a bijection between the set of equivalence classes of MCFs and DGA homotopy classes of augmentions of the cellular DGA for 1-dimensional Legendrians to establish a characterization of induced augmentation sets in terms of MCFs; see Proposition 5.9. The section concludes in Section 5.3 with a discussion of A-form MCFs, as defined by Henry [23], that are particularly convenient for computing induced augmentation sets in explicit examples.

5.1. Morse complex families.

We now review the definition of Morse complex families for Legendrian knots and surfaces, as in [23, 36], allowing for the case of Legendrian cobordisms.

5.1.1. *The 1-dimensional case.* Let $(\Lambda, \mu) \subset J^1M$ be a 1-dimensional ρ-graded Legendrian knot. Recall that we use the respective notations $\pi_x : J^1M \to M$ and $\pi_{xz} : J^1M \to M \times \mathbb{R}$ for the base and front projections.

Definition 5.1. A **ρ-graded Morse complex family** (abbr. **MCF**) for (Λ, μ) is a pair $\mathcal{C} = (H, \{d_\nu\})$ consisting of the following items:

(1) A **handleslide set** H which is a finite collection of points in M such that each $p \in H$ is equipped with lifts $u(p), l(p) \in \Lambda$ not belonging to cusp edges and satisfying $\pi_x(u(p)) = \pi_x(l(p)) = p$, $z(u(p)) > z(l(p))$, and $\mu(u(p)) = \mu(l(p))$. We refer to $p \in H$ as a (formal) handleslide of $\mathcal{C}$ with upper and lower endpoints at $u(p)$ and $l(p)$.

Denote by Λ_{sing} the base projection of the singular set of $\pi_{xz}(\Lambda)$ (crossings and cusps), and let $\{R_\nu\}$ be the collection of path components of $M \setminus (H \cup \Lambda_{sing})$. Note that above each R_ν, the sheets of Λ are totally ordered by their z-coordinates, so they can be enumerated as

$$S_1^\nu, \dots, S_{n_\nu}^\nu \quad \text{with } z(S_1^\nu) > \cdots > z(S_{n_\nu}^\nu).$$

We then have $\mathbb{Z}/\rho$-graded vector spaces $V(R_\nu) = \mathrm{Span}_{\mathbb{Z}/2}\{S_i^\nu\}$ where the grading is via the Maslov potential $|S_i^\nu| = \mu(S_i^\nu)$. We will sometimes omit the superscript ν in the notation for sheets.

(2) The second piece of information defining an MCF is a collection of strictly upper triangular differentials $\{d_\nu\}$ of degree $+1$ mod ρ making each $(V(R_\nu), d_\nu)$ into a $\mathbb{Z}/\rho$-graded cochain complex. Here, the strictly upper triangular condition is that the matrix coefficient $\langle d_\nu S_q^\nu, S_p^\nu \rangle$ is 0 unless $z(S_p^\nu) > z(S_q^\nu)$.

Moreover, H and $\{d_\nu\}$ are required to satisfy the following Axiom 5.2.

Axiom 5.2. Let R_0 and R_1 be connected components of $\pi_x(\Lambda) \setminus (H \cup \Lambda_{sing})$ that border each other at a point $p \in H \cup \Lambda_{sing}$.

(1) If $p \in H$ is a *handleslide* with upper endpoint $u(p) \in S_u$ and $l(p) \in S_l$, then the handleslide map
$$h_{u,l}(S_i) = \begin{cases} S_i, & i \neq l, \\ S_u + S_l, & i = l, \end{cases}$$
gives a chain isomorphism $h_{u,l} : (V(R_0), d_0) \xrightarrow{\cong} (V(R_1), d_1)$.

(2) If p is the base projection of a *crossing* between sheets S_k and S_{k+1}, then the permutation map
$$Q(S_k) = S_{k+1},\ Q(S_{k+1}) = S_k,\ Q(S_i) = S_i, i \notin \{k, k+1\}$$
is a chain isomorphism, $Q : (V(R_0), d_0) \xrightarrow{\cong} (V(R_1), d_1)$.

(3) Suppose that p is a *cusp* point such that the two sheets of the cusp, S^1_k and S^1_{k+1}, exist above R_1 but not above R_0. Then, identifying sheets of R_0 and R_1 whose closures intersect at p gives an inclusion $\iota : V(R_0) \hookrightarrow V(R_1)$ with
$$V(R_1) = \iota(V(R_0)) \oplus V_{cusp}, \quad V_{cusp} = \mathrm{Span}_{\mathbb{Z}/2}\{S^1_k, S^1_{k+1}\}.$$
We require that $\iota \circ d_0 = d_1 \circ \iota$ and $d_1 S^1_{k+1} = S^1_k$, i.e. as a complex $(V(R_1), d_1)$ is the split extension of $(V(R_0), d_0)$ by (V_{cusp}, d') where d' maps the lower sheet of the cusp to the upper sheet of the cusp.

Remark 5.3. (1) The requirement in Axiom 5.2 (3) used here is stronger than in some other versions of the definition, eg. in [23, 24, 25]. In the terminology of [23], with our definition we only consider Morse complex families with "simple births and deaths". The collection of equivalence classes of MCFs is not affected by making this restriction; see [23, Proposition 3.17].

(2) If $\Lambda \subset J^1\mathbb{R}$, then the collection of differentials $\{d_\nu\}$ is uniquely determined by Λ and H, since one can work from left to right and apply Axiom 5.2 to determine the $\{d_\nu\}$ inductively. However, it should be noted that not every handleslide set H for Λ will produce an MCF. For example, consider the case when R_0 and R_1 are seperated by a crossing point, and d_0 has already been determined. If $\langle d_0 S_{k+1}, S_k\rangle \neq 0$, then the differential $d_1 = Q \circ d_0 \circ Q^{-1}$ required by Axiom 5.2 (2) will not be strictly upper triangular.

In figures, we picture handleslides as vertical segments connecting $u(p)$ and $l(p)$ on the front projection of Λ. When convenient, differentials d_ν may be indicated above R_ν by drawing a dotted arrow from S_i to S_j whenever $\langle d_\nu S_j, S_i\rangle = 1$. See for example Figures 5 and 9 below.

5.1.2. *The 2-dimensional case.* Consider a 2-dimensional ρ-graded compact Legendrian surface, $(\Sigma, \mu) \subset J^1M$ where $\dim M = 2$. We allow for $\partial M \neq \emptyset$, but in this case require that Σ is properly embedded, eg. Σ could be a compact Legendrian cobordism in $J^1([0,1] \times \mathbb{R})$ or $J^1([0,1] \times S^1)$. We again use the notation $\Sigma_{sing} \subset M$ for the base projection of the singular set of $\pi_{xz}(\Sigma)$ (which now includes crossings, cusps, and swallowtail points).

Definition 5.4. A **ρ-graded MCF** for Σ is a triple $(H_0, H_{-1}, \{d_\nu\})$ consisting of:

(1) A set of **super-handleslide points**, $H_{-1} \subset Int(M) \setminus \Sigma_{sing}$, equipped with upper and lower endpoint lifts to Σ satisfying $\mu(u(p)) = \mu(l(p)) - 1$ for all $p \in H_{-1}$.

(2) A collection of **handleslide arcs** which is an immersion $H_0 : X \to M$ of a compact 1-manifold, transverse to ∂M. Aside from the exceptions at boundary points specified in Axiom 5.5 below, we require that H_0 is transverse to $H_0 \cup \Sigma_{sing}$ and that the only self intersections are at transverse double points in $Int(M)$. In addition, H_0 should be equipped with upper and lower endpoint lifts defined on the interior of X
$$u, l : Int(X) \to \Sigma, \quad \pi_x \circ u = \pi_x \circ l = H_0,$$

satisfying

$$u(s) > l(s) \quad \text{and} \quad \mu(u(s)) = \mu(l(s)) \quad \forall s \in Int(X).$$

Endpoints of handleslide arcs must satisfy Axiom 5.5 below.

(3) A collection of differentials $\{d_\nu\}$ of degree $+1$ mod ρ making each $(V(R_\nu), d_\nu)$ into a $\mathbb{Z}/\rho$-graded cochain complex where $\{R_\nu\}$ is the collection of path components of $M \setminus (H_{-1} \cup H_0(X) \cup \Sigma_{sing})$. Note that $H_{-1} \cup H_0(X) \cup \Sigma_{sing}$ is a stratified subset of M whose 1-dimensional strata are *handleslide arcs*, *crossing arcs*, and *cusp arcs*. For each such 1-dimensional stratum, we require that the complexes $(V(R_0), d_0)$ and $(V(R_1), d_1)$ that appear adjacent to the stratum are related as in Axiom 5.2 (1)-(3).

Before stating Axiom 5.5 we introduce some convenient terminology. Above $M \setminus \Sigma_{sing}$ sheets of Σ are locally labelled $S_1, \dots, S_n$ with descending z-coordinate. We will refer to a handleslide arc that has its upper and lower lift on sheets S_i and S_j as an (i,j)**-handleslide**. We caution that this terminology only applies locally since the labeling of upper and lower sheets of a handleslide arc may change when the arc passes through Σ_{sing}. We use a similar terminology for super-handleslide points.

Axiom 5.5. Endpoints of handleslide arcs can occur at transverse intersections of H_0 and ∂M. All other endpoints of handleslide arcs occurring in the interior of M are as specified in the following:

(1) Suppose that $p \in M$ is a transverse double point for $H_0|_{Int(X)}$ where a (u_1, l_1)-handleslide arc crosses a (u_2, l_2)-handleslide arc. If $l_1 = u_2$, then at p there is a single endpoint of a (u_1, l_2)-handleslide arc.

(2) Let $p \in H_0$ be a (u,l)-super-handleslide point, and fix some region R_ν bordering p.
- For every $i < u$ such that $\langle d_\nu S_u, S_i \rangle = 1$, there is a single (i,l)-handleslide arc with an endpoint at p.
- For every $l < j$ such that $\langle d_\nu S_j, S_l \rangle = 1$, there is a single (u,j)-handleslide arc with an endpoint at p.

(3) Let $p \in \Sigma_{sing}$ be the base projection of an upward swallowtail point, i.e. one where in the front projection the crossing arc appears below the third sheet that that limits to the swallowtail point. The two cusp arcs in Σ_{sing} that meet at p divide a small neighborhood $N \subset M$ of p into halfs N_- and N_+ above which Σ has $(n-2)$ or n sheets respectively. Let k be such that above N_- (resp. above N_+) the sheets that border the swallowtail point are numbered S_k (resp. S_k, S_{k+1}, S_{k+2}) with S_{k+1} and S_{k+2} meeting along the crossing arc, c, that ends at p and is contained in N_+. Let $(V(R_0), d_0)$ be the complex associated to the region, R_0, that borders p on the same side as N_-.
- There are two $(k+1, k+2)$-handleslide arcs contained in N_+ and ending at p, one on each side of c.
- For each $i < k$ such that $\langle d_0 S_k, S_i \rangle = 1$ there is a (i,k)-handle slide arc in N_+ with endpoint at p.

When p is a downward swallowtail, the requirement is symmetric. The sheets S_k in N_- (resp. S_k, S_{k+1}, S_{k+2} in N_+) correspond now to sheets S_{l-2} in N_- (resp. S_l, S_{l-1}, S_{l-2}). There are now two $(l-2, l-1)$-handleslide arcs in N_+ on opposite sides of c with endpoints at p, and for each $l-2 < j$ with $\langle d_0 S_j, S_{l-2} \rangle = 1$ there is an $(l, j+2)$-handleslide arc in N_+ with endpoint at p.

The three types of endpoints for handleslide arcs are pictured in Figure 5.

Remark 5.6.
(1) In Axiom 5.5 (2), if the condition holds for some region R_ν bordering p, then it holds for every region that borders p.
(2) In Axiom 5.5 (3), in a neighborhood of a swallowtail point p, assume that a differential d_0 has been assigned to the region R_0 that borders p in N_-. Then, as long as handleslide




FIGURE 5. The three possible endpoints for handleslide arcs specified by Axiom 5.5 (1)-(3). The left column depicts the base projection (to M) of H_0 (in red), H_{-1} (a green star) and Σ_{sing}, (in blue). The center and right column depict slices of the front projection, $\pi_{xz}(\Sigma)$, at $x_2 = a$ and $x_2 = b$ with dotted black arrows from S_i to S_j used to indicate when $\langle dS_j, S_i\rangle = 1$. In (3), a downward swallowtail point is pictured.

arcs are placed in N_+ as specified by the axiom, there is always a unique way to assign differentials to regions in N_+ to produce an MCF near p.

(3) See [36, Section 6.1] for a further discussion of constructions of MCFs.

Note that when Σ and M are 2-dimensional and have boundary, an MCF $\mathcal{C}$ for Σ restricts to an MCF for the 1-dimensional Legendrian $\Sigma|_{\partial M} \subset J^1(\partial M)$.

Definition 5.7. Let $\mathcal{C}_0$ and $\mathcal{C}_1$ be two MCFs for a 1-dimensional Legendrian knot $\Lambda \subset J^1M$. We say that $\mathcal{C}_0$ and $\mathcal{C}_1$ are **equivalent** if there exists an MCF $\mathcal{C}$ for the product cobordism $j^1(1_{[0,1]} \cdot \Lambda) \subset J^1([0,1] \times M)$ such that $\mathcal{C}|_{\{0\}\times M} = \mathcal{C}_0$ and $\mathcal{C}|_{\{1\}\times M} = \mathcal{C}_1$.

Remark 5.8. The statement of Definition 5.7 is slightly different than the definition of equivalence in [23, 24, 25]. These earlier works did not consider MCFs for 2-dimensional Legendrians and instead present a collection of elementary moves on 1-dimensional MCFs with two MCFs $\mathcal{C}_0$ and $\mathcal{C}_1$ declared to be equivalent if they can be related by a sequence of elementary moves. The two versions of the definitions are equivalent, since the elementary moves simply describe the bifurcations that occur along the slices, $\mathcal{C}|_{\{t\}\times M}$, of a generic 2-dimensional MCF $\mathcal{C}$ for $j^1(1_{[0,1]} \cdot \Lambda)$ as t increases from 0 to 1. For instance, as numbered in [24, Section 4.3], Move 1 corresponds to passing a critical point of $t \circ H_0$; Moves 2-6 correspond to a transverse self intersection of H_0 with Move 6 demonstrating the handleslide endpoints required by Axiom 5.5 (1); Moves 7, 8, and 10 are transverse intersections of H_0 with the base projection of a crossing arc of $j^1(1_{[0,1]} \cdot \Lambda)$;

Moves 9 and 11 are transverse intersections of H_0 and the base projection of a cusp arc; and Move 15 corresponds to passing a super-handleslide point. The Moves 12-14 do not occur in our case since we only consider MCFs with "simple births and deaths", and this does not affect the resulting equivalence classes. Indeed, when two such MCFs with only "simple births and deaths" are related by a sequence of the Moves 1-15, possibly passing through some MCFs with non-simple births or deaths, we can always arrive at an alternate sequence of moves avoiding non-simple cusps by treating the implicit handleslides discussed in [24] as explicit handleslides located near a cusp. See [23, Proposition 3.17].

5.2. **MCFs and augmentations.** The following proposition characterizes the (cellular) induced augmentation set of a compact Legendrian cobordism (see Section 3.2 for the definition) in terms of MCFs.

Proposition 5.9. (1) *For any $(\Lambda, \mathcal{E}, \mu)$ with $\Lambda \subset J^1M$ and $\dim \Lambda = 1$, there exists a canonical bijection*

$$\widehat{\Phi} : MCF_\rho(\Lambda)/\sim \to Aug_\rho(\Lambda, \mathcal{E})/\sim$$

between equivalence classes of ρ-graded MCFs for (Λ, μ) and DGA homotopy classes of ρ-graded augmentations of the cellular DGA $\mathcal{A}(\Lambda, \mathcal{E})$.

(2) *Let $\Sigma \subset J^1([0,1] \times M)$ be a compact ρ-graded Legendrian cobordism from $(\Lambda_0, \mu_0, \mathcal{E}_0)$ to $(\Lambda_1, \mu_1, \mathcal{E}_1)$. Using the canonical bijections $\widehat{\Phi} : MCF_\rho(\Lambda_i)/\sim \xrightarrow{\cong} Aug_\rho(\Lambda_i, \mathcal{E}_i)/\sim$, the induced augmentation set*

$$I_\Sigma^{cell} \subset Aug_\rho(\Lambda_0, \mathcal{E}_0)/\sim \times Aug_\rho(\Lambda_1, \mathcal{E}_1)/\sim$$

viewed as a subset

$$I_\Sigma^{MCF} \subset MCF_\rho(\Lambda_0, \mathcal{E}_0)/\sim \times MCF_\rho(\Lambda_1, \mathcal{E}_1)/\sim$$

satisfies

$$I_\Sigma^{MCF} = \{([\mathcal{C}|_{\Lambda_0}], [\mathcal{C}|_{\Lambda_1}]) \,|\, \mathcal{C} \in MCF_\rho(\Sigma)\}\,.$$

A detailed proof of this, including review of the relevant parts of [36], appears in the preprint version of this article; see [30, Proposition 5.17 and 5.19]. We provide here a sketch.

Sketch of proof. Both (1) and (2) are consequences of a straightforward extension of the correspondences between MCFs and augmentations from [36, Propositions 5.5 and 6.4] for closed surfaces to the case of 2-dimensional cobordisms. These results produce augmentations of $\mathcal{A}(\Lambda, \mathcal{E})$ (that are viewed in [36] as *Chain Homotopy Diagrams*) from MCFs that are suitably transverse to $\mathcal{E}$ (aka *nice* MCFs), and vice-versa.

In particular, in the 1-dimensional case we have a map Φ from nice MCFs to augmentations of $\mathcal{A}(\Lambda, \mathcal{E})$. That Φ induces a bijection $\widehat{\Phi}$ as in (1) is seen from applying the MCF/augmentation correspondences to the product cobordism $j^1(1_{[0,1]} \cdot \Lambda)$. Indeed, the above Proposition 4.5 shows that two augmentations of $\mathcal{A}(\Lambda, \mathcal{E})$ are homotopic if and only if they extend to an augmentation of $j^1(1_{[0,1]} \cdot \Lambda)$, and this characterization matches the definition of equivalence for MCFs. The second statement (2) follows since in the cobordism case the correspondences can be made to commute with the restrictions to the boundary components.

□

5.3. **A-form MCFs.** From Proposition 5.9 and Corollary 4.11, we see that I_Σ^{MCF} is invariant of Σ up to Legendrian isotopy rel. $\partial\Sigma$. Moreover, I_Σ^{MCF} is equivalent in a canonical way to both the cellular induced augmentation set I_Σ^{cell} and, via Corollary 4.12, to the induced augmentation set I_Σ of the conical version of Σ, $G_{std}(\Sigma)$. For computing I_Σ^{MCF} in explicit examples, it is convenient to have an efficient characterization of the sets, $MCF_\rho(\Lambda)/\sim$, not requiring the cellular DGA which often has a large number of generators. For Legendrian links in $J^1\mathbb{R}$, work of Henry [23, 25], which we now review, gives an explicit bijection $MCF_\rho(\Lambda)/\sim \leftrightarrow Aug_\rho(\Lambda_{res})/\sim$ where




FIGURE 6. The allowed locations for handleslides in an A-form MCF.

Λ_{res} is the Ng resolution of Λ. Recall that the DGA $\mathcal{A}(\Lambda_{res})$ has generators in bijection with the crossings and right cusps of the front projection of Λ; see [28, Section 2].

Definition 5.10 ([23])**.** A ρ-graded MCF $\mathcal{C}$ for a Legendrian link $\Lambda \subset J^1\mathbb{R}$ is in **A-form** if its handleslides occur only in the following locations:

(1) Handleslides connecting the two crossing strands appear immediately to the left of *some subset* of the crossings of $\pi_{xz}(\Lambda)$.
(2) In the case $\rho = 1$, handleslides connecting the two strands that meet at a right cusp appear immediately to the left of *some subset* of the right cusps of $\pi_{xz}(\Lambda)$.
(3) Near a right cusp, any number of additional handleslides may appear connecting the upper (resp. lower) strand of the right cusp to a strand above (resp. below) the right cusp. These handleslides are located to the right of the handleslide from (2), if it exists.

See Figure 6.

Remark 5.11. The definition appears slightly different from that found in [23, 25] as the handleslides from (3) are implicit handleslides in these references; see Remarks 5.3 (1) and 5.8. Moreover, an A-form MCS is uniquely determined by its handleslides of type (1) and (2), since the handleslides of type (3) are determined by the form of the complexes (R_ν, d_ν) near right cusps.

Given an A-form MCF, $\mathcal{C}$, for $\Lambda \subset J^1\mathbb{R}$ we can define an algebra homomorphism

$$\epsilon_{\mathcal{C}} : \mathcal{A}(\Lambda_{res}) \to \mathbb{Z}/2$$

where for each crossing or right cusp $b_i \in \mathcal{A}(\Lambda_{res})$ we set $\epsilon_{\mathcal{C}}(b_i)$ to be 1 (resp. 0) if a handleslide connecting the two strands of the crossing or cusp appears (resp. does not appear) to the left of b_i.

Proposition 5.12 ([23, 25])**.** *For any $\Lambda \subset J^1\mathbb{R}$, the correspondence $\mathcal{C} \mapsto \epsilon_{\mathcal{C}}$ defines a surjective map from the set of ρ-graded A-form MCFs for Λ to $Aug_\rho(\Lambda_{res})$. Moreover, this induces a bijection*

$$MCF_\rho(\Lambda)/\sim \xrightarrow{\cong} Aug_\rho(\Lambda_{res})/\sim.$$

Proof. In [23], a map $\widehat{\Psi} : MCF_\rho(\Lambda)/\sim \to Aug_\rho(\Lambda_{res})/\sim$ is defined and shown to be surjective. Moreover, this map has the property that when $\mathcal{C}$ is an A-form MCF we have $\widehat{\Psi}([\mathcal{C}]) = [\epsilon_{\mathcal{C}}]$, and it is shown that every MCF is equivalent to an A-form MCF. Finally, in [25] it is shown that $\widehat{\Psi}$ is injective. □

6. Examples

In this section, we apply the methods developed in Section 5 to compute induced augmentations and augmentation sets for some particular Legendrian fillings. In Section 6.1, we give examples of oriented Legendrian fillings inducing augmentations that cannot be induced by any oriented embedded Lagrangian fillings. In Section 6.2, we prove Theorem 1.4 by constructing $2n$ Legendrian fillings of the Legendrian torus knot, $T(2, 2n+1)$, each with a single degree 0 Reeb chord, and distinguished by their induced augmentation sets.

6.1. **Initial examples.** Before launching into examples, we summarize the method that we use for computing induced augmentations and induced augmentation sets arising from a (compact) Legendrian filling, Σ. For a 1-dimensional Legendrian, Λ, with admissible transverse decomposition $\mathcal{E}_{\pitchfork}$, Proposition 5.9 and Corollary 4.12 provide a commutative diagram of bijections and inclusions

$$\begin{array}{ccccc} MCF_\rho(\Lambda)/\sim & \cong & Aug_\rho(\Lambda, \mathcal{E}_{||})/\sim & \cong & Aug_\rho(\widetilde{\Lambda})/\sim \\ \cup & & \cup & & \cup \\ I_\Sigma^{MCF} & \cong & I_\Sigma^{cell} & \cong & I_{\widetilde{\Sigma}} \end{array} \tag{6.1}$$

where, in the top row, from left to right the sets are the equivalence classes of MCFs of Λ; homotopy classes of augmentations of the cellular DGA of $(\Lambda, \mathcal{E}_{||})$; and homotopy classes of augmentations of the LCH DGA $\mathcal{A}(\widetilde{\Lambda})$ where $\widetilde{\Lambda}$ is Legendrian isotopic to Λ. (It is the standard geometric model from Section 4.5.) The bottom row consists of the (equivalent) versions of the induced augmentation set in these three different settings (where $\widetilde{\Sigma} = G_{std}(\Sigma)$ is Σ with its positive end extended to become a conical cobordism). The MCFs in I_Σ^{MCF} are those that arise from restricting a 2-dimensional MCF, $\mathcal{C}_\Sigma$, for Σ to Λ. Thus, elements of I_Σ^{MCF} (and so also induced augmentations in $I_{\widetilde{\Sigma}}$) may be constructed by specifying a 2-dimensional MCF for Σ, and then observing its restriction to Λ. To keep track of the different MCFs of Λ we use the further bijection from Proposition 5.12 with augmentations of the Ng resolution,

$$MCF_\rho(\Lambda)/\sim \ \cong \ Aug(\Lambda_{res})/\sim$$

following Section 5.3. That is, once a 2-dimensional MCF has been restricted to the Legendrian boundary Λ, if necessary, we transform it by an equivalence into A-form to arrive at an augmentation of Λ_{res}.

For the rest of the section, all of the augmentations we talk about are of Λ_{res}, and we will refer to Reeb chords of Λ_{res} as crossings and right cusps of the front projection.

Example 6.1. An augmentation that can be induced by a Legendrian filling for a knot that does not have any oriented embedded Lagrangian fillings.

The work [32, Section 8.3] gives an infinite family of such knots without explicitly computing the induced augmentations. One of them is the knot 4_1 shown in Figure 7 where a conical Legendrian filling is constructed by a clasp move (see Figure 17 below for a discussion of local moves for constructing Legendrian cobordisms) followed by Reidemeister I moves and an unknot move; the pictures are slices of the front projection of the corresponding compact filling, $\Sigma \subset J^1([0,1] \times M)$, with $t \in [0,1]$ constant. As shown in [32, Section 8.3], the clasp move can be realized by a conical Legendrian cobordism with a single Reeb chord, so that a conical version of Σ (after an appropriate Legendrian isotopy) is a disk with a single double point in the Lagrangian projection. Figure 7 indicates the MCF for Σ as a movie of handleslides. Note that the handleslides that appear after the Reidemeister I moves are as required in Axiom 5.5 (3). Using the correspondence between A-form MCFs and augmentations of Λ_{res}, the induced augmentation ϵ_1 sends the crossings b_1 and b_2 to 1 and all other Reeb chords to 0. Note that a $\mathbb{Z}$-valued Maslov potential for 4_1 extends (uniquely) over Σ, and the MCF for Σ is 0-graded. Correspondingly, ϵ_1 is the unique 0-graded augmentation of Λ_{res}. As discussed in [32, Section 8.3], the 4_1 knot does not have any oriented embedded Lagrangian fillings due to a restriction on the Thurston-Bennequin number.

We note that 4_1 does admit *non-orientable* embedded Lagrangian fillings. Such a filling, constructed via applying two pinch moves (see Figure 17) and then an unknot move, is pictured in Figure 8. It induces the 1-graded augmentation ϵ_2 that sends b_1, b_2, a, and c to 1 and other Reeb chords to 0.

Question 6.2. Can the augmentation ϵ_1 induced by the Legendrian filling from Figure 7 also be induced by a non-orientable embedded Lagrangian filling?




FIGURE 7. A conical Legendrian filling of Legendrian 4_1 constructed by a movie. The arrow indicates the negative t direction. Red vertical line segments indicates handle slides in the MCF.

We suspect that the answer is no, but we do not know of any obstructions to such a non-orientable filling that are currently available in the literature.




FIGURE 8. A non-orientable conical Legendrian filling of Legendrian 4_1 with embedded Lagrangian projection. A 1-graded MCF inducing the augmentation ϵ_2 is pictured.

Example 6.3. Two augmentations for the same knot, one can be induced by an embedded Lagrangian filling and one cannot.

Figure 9 shows a Legendrian of knot type $m(8_{21})$. The knot has a genus one embedded Lagrangian filling constructed by doing three pinch moves (see Figure 17) and closing up the two disjoint unknots (see the first row in Figure 9). The associated MCF induces the augmentation ϵ_1 that sends b_1, b_4, b_5 to 1 and others to 0.

One the other hand, one can construct a conical Legendrian filling through doing three clasp moves and then close up the unknot as shown in the movie in the second row of Figure 9. This is a Legendrian disk with three Reeb chords. The associated MCF as shown in Figure 9 induces the augmentation ϵ_2 that sends b_1, b_2, b_4 and b_5 to 1 and other Reeb chords to 0. This augmentation cannot be induced by an embedded orientable Lagrangian filling since its linearized homology has Poincare polynomial $t^{-1} + 4 + 2t$. By Seidel's Isomorphism [12, 11], if an augmentation was induced by an embedded orientable genus g filling, using the $\mathbb{Z}/2$-grading on linearized homology arising from a $\mathbb{Z}/2$-valued Maslov potential on the filling (such a Maslov potential exists by orientability, see Remark 2.9), its Poincare polynomial with $\mathbb{Z}/2$-grading should be $t + 2g$.

6.2. **Proof of Theorem 1.4.** The theorem provides examples of knots with an arbitrary (finite) number of immersed fillings, all having isomorphic DGAs, but distinguished by induced augmentation sets.

Theorem 1.4. *For each $n \geq 1$, there exists $2n$ distinct conical Legendrian fillings, $\Sigma_1, \ldots, \Sigma_{2n}$ of the max-tb Legendrian torus knot $T(2, 2n+1)$ such that*




FIGURE 9. The first row gives an embedded Lagrangian filling of $m(8_{21})$ constructed by a movie. The second row gives a conical Legendrian filling of $m(8_{21})$ through a movie. The arrow indicates negative t direction.

(i) *the Σ_i are all orientable with genus $n-1$ and have $\mathbb{Z}$-valued Maslov potentials,*
(ii) *each Σ_i has a single Reeb chord of degree 0,*
(iii) *and the induced augmentation sets satisfy $I_{\Sigma_i} \neq I_{\Sigma_j}$ when $i \neq j$.*




FIGURE 10. Part (a) is a Legendrian $(2, 2n+1)$ torus knot and (b) is a Legendrian $(2, 2n-1)$ torus knot. Part (c) is the unknot got after doing $2n-2$ pinch moves to $T(2, 2n-1)$.

Note that each of the DGAs $\mathcal{A}(\Sigma_i)$ is $\mathbb{Z}$-graded with a single generator in degree 0 and the differential is necessarily zero for grading reasons. Thus, they are all isomorphic.

Proof. Consider the Legendrian torus knot $T(2, 2n+1)$ as shown in Figure 10 part (a). There are $2n$ "eye shapes" that one can do a clasp move on to get a $(2, 2n-1)$ Legendrian torus knot as shown in part (b). One can then do $2n-2$ pinch moves to the $T(2, 2n-1)$ knot and close the unknot in part (c). Let Σ_i be the conical Legendrian filling of $T(2, 2n+1)$ that is constructed

by performing the first clasp move at the "eye shape" between b_i and b_{i+1}. According to the MCF shown in Figure 10 part (b) and (c), the pinching procedure on $T(2, 2n-1)$ induces an augmentation of $T(2, 2n-1)$ that only sends c_1 to 1 and all others to 0.

The clasp move produces a degree 0 Reeb chord, a, (see Proposition 8.3 of [32] for the degree computation) of the resulting conical Legendrian surface Σ_i, and the rest of the cobordism can be constructed without Reeb chords. Thus, $\mathcal{A}(\Sigma_i)$ has two augmentations, $a \mapsto 0$ and $a \mapsto 1$, and it follows that I_{Σ_i} can have at most two elements. In fact, I_{Σ_i} does have two elements induced by the two MCFs of the conical Legendrian surface shown in Figure 11. For a subset I of the set $\{1, \cdots 2n+1\}$, denote by ϵ_I the algebra map from $\mathcal{A}(T(2, 2n+1))$ to $\mathbb{Z}/2$ that only sends $\{b_i | i \in I\}$ to 1 and all other Reeb chords to 0. Then, Figure 11 shows

$$I_{\Sigma_i} = \begin{cases} \{\epsilon_{\{1\}}, \epsilon_{\{3\}}\}, & \text{if } i = 1 \\ \{\epsilon_{\{1\}}, \epsilon_{\{1,i,i+2\}}\}, & \text{if } 1 < i < 2n \\ \{\epsilon_{\{1\}}, \epsilon_{\{1,2n\}}\}, & \text{if } i = 2n. \end{cases}$$




FIGURE 11. Two MCFs for the clasp move constructing Σ_i pictured in the base projection. These MCFs extend the MCF from Figure 10 (b) and (c) to all of Σ_i. Columns $(a), (b), (c)$ are for $i = 1$, $1 < i < 2n$ and $i = 2n$, respectively. The black dotted lines represent double points (in the front projection) and red lines represent handleslides.

□

7. Every augmentation is induced by an immersed filling

With the characterization of induced augmentation sets for Legendrian cobordisms in terms of MCFs now in place, the present section establishes Theorem 1.2 from the introduction. For convenience, we repeat the statement.

Theorem 1.2. *Let $\Lambda \subset J^1\mathbb{R}$ have the $\mathbb{Z}/\rho$-valued Maslov potential μ where $\rho \geq 0$, and let $\epsilon : \mathcal{A}(\Lambda) \to \mathbb{Z}/2$ be any ρ-graded augmentation.*

(1) *If $\rho \neq 1$, there exists a conical Legendrian filling Σ of Λ with $\mathbb{Z}/\rho$-valued Maslov potential extending μ together with a ρ-graded augmentation $\alpha : \mathcal{A}(\Sigma) \to \mathbb{Z}/2$ such that $\epsilon \simeq \epsilon_{(\Sigma,\alpha)}$. Moreover, if ρ is even, then Σ is orientable.*

(2) *If $\rho = 1$, then there exists a conical Legendrian cobordism $\Sigma : U \to \Lambda$ where U is the standard Legendrian unknot with $tb(U) = -1$ together with an augmentation $\alpha : \mathcal{A}(\Sigma) \to \mathbb{Z}/2$ such that $\epsilon \simeq f_\Sigma^* \alpha$.*

Proof of Theorem 1.2. Note that the orientability of Σ when ρ is even follows from the existence of the $\mathbb{Z}/\rho$-valued Maslov potential on Σ. See Remark 2.9.

In view of Corollary 3.8, if Λ_1 and Λ_2 are Legendrian isotopic and the statement holds for Λ_1, then it also holds for Λ_2. Therefore, we can replace Λ with its standard geometric model $G_{std}(\Lambda, \mathcal{E}_{\pitchfork})$ with respect to some admissible transverse decomposition (see Section 4.5), so that Corollary 4.12 and Proposition 5.9 apply to reduce Theorem 1.2 to Proposition 7.1 below. □

Proposition 7.1. *Let $\mathcal{C}_\Lambda$ be any ρ-graded MCF for $\Lambda \subset J^1\mathbb{R}$ with respect to the Maslov potential μ.*

(1) *If $\rho \neq 1$, then there exists a compact Legendrian filling, $\Sigma \subset J^1([0,1] \times \mathbb{R})$, of Λ with a Maslov potential extending μ such that $\mathcal{C}_\Lambda$ can be extended to a ρ-graded MCF, $\mathcal{C}_\Sigma$, on Σ.*

(2) *If $\rho = 1$, then the same statement holds except that Σ is a compact Legendrian cobordism from the Legendrian unknot U to Λ.*

We will prove Proposition 7.1 at the end of this section after establishing some preliminaries about extending MCFs over various elementary cobordisms.

7.1. **Normal rulings and SR-form MCFs.** In [23], Henry introduced a class of MCFs (the "SR-form MCFs") for 1-dimensional Legendrian links that are in a clear many-to-one correspondence with the normal rulings of Λ. Moreover, Henry proved that any MCF is equivalent to an SR-form MCF. It will be convenient to make use of this result in proving Proposition 7.1, so we briefly discuss the relevant definitions.

See any of [24, 21, 37] for a detailed definition of **normal rulings** for Legendrian links in $J^1\mathbb{R}$. Here, we recall that a normal ruling, σ, for $\Lambda \subset J^1\mathbb{R}$ (where Λ has generic front and base projections) may be viewed as a continuous family of fixed point free involutions, $\sigma_{x_0} : \Lambda \cap \{x = x_0\} \to \Lambda \cap \{x = x_0\}$, defined for $x_0 \in \mathbb{R} \setminus \Lambda_{sing}$, i.e. for all values of x_0 such that $\pi_{xz}(\Lambda)$ has no crossing or cusp along the vertical line $x = x_0$. For each connected component $R_\nu \subset \mathbb{R} \setminus \Lambda_{sing}$ there is a resulting involution on the sheets of Λ above R_ν, notated simply as

$$\sigma : \{S_i^\nu\} \to \{S_i^\nu\},$$

which partitions $\{S_i^\nu\}$ into a collection of disjoint pairs (because of the fixed point free condition). Moreover, if two regions R_i and R_{i+1} share a border at a cusp or crossing, then the pairings of sheets above R_i and R_{i+1} are required to be related in a standard way. In particular, sheets that meet at a cusp point are paired near the cusp point, and every crossing of $\pi_{xz}(\Lambda)$ is either a **departure**, a **return**, or a **switch** for σ as in Figure 12. The **degree** of a crossing of $\pi_{xz}(\Lambda)$ with respect to a $\mathbb{Z}/\rho$-valued Maslov potential μ is defined to be the difference $\mu(u) - \mu(l) \in \mathbb{Z}/\rho$ where u and l denote the upper and lower strands at the left side of the crossing. We say that σ is **ρ-graded** with respect to μ if all switches of σ have degree 0 mod ρ. On each region $R_\nu \subset \mathbb{R} \setminus \Lambda_{sing}$, we can use σ to define a **standard ruling differential**

$$d_\nu^\sigma : V(R_\nu) \to V(R_\nu), \quad d_\nu^\sigma S_j^\nu = \begin{cases} S_i^\nu, & \sigma(S_j^\nu) = S_i^\nu \text{ and } z(S_i^\nu) > z(S_j^\nu), \\ 0, & \text{else.} \end{cases} \tag{7.1}$$

Definition 7.2. A ρ-graded MCF $\mathcal{C} = (H, \{d_\nu\})$ for a 1-dimensional Legendrian link Λ is in **SR-form** with respect to a ρ-graded normal ruling σ for Λ if the only handleslides of $\mathcal{C}$ are as follows:

(1) At *every* switch of σ, handleslides are placed as specified in Figure 13.




FIGURE 12. Near a crossing the two sheets that cross cannot be paired with one another by a normal ruling σ. The figure indicates the allowed configurations for the crossing sheets and their companion sheets. (There may be other sheets of Λ in between those pictured.)




FIGURE 13. The placement of handleslides near switches and some subset of the degree 0 returns of σ for an SR-form MCF. When $\rho = 1$, handleslides are also placed next to some collection of right cusps as indicated.

(2) At *some subset* of the degree 0 (mod ρ) returns of σ, handleslides are placed as in Figure 13.
(3) In the 1-graded case, at *some subset* of the right cusps of Λ, a single handeslide connects the two cusp strands near the cusp.

Remark 7.3. (1) It can be shown that for any ρ-graded normal ruling σ of Λ and any chosen subset of the ρ-graded returns of Λ (and also of the right cusps of Λ in the 1-graded case), there is an SR-form MCF with handleslide set as in Figure 13.
(2) Whenever $\mathcal{C} = (H, \{d_\nu\})$ is an SR-form MCF and $R_\nu \subset \mathbb{R}\setminus(H \cup \Lambda_{sing})$ is a region outside of the collection of handleslides near switches, returns, and cusps, the differential d_ν agrees with the standard ruling differential d_ν^σ defined in (7.1).

The following proposition is Theorem 6.17 from [23].

Proposition 7.4. *Any ρ-graded MCF on a Legendrian link $\Lambda \subset J^1\mathbb{R}$ is equivalent to a SR-form MCF with respect to some ρ-graded normal ruling of Λ.*

7.2. **Construction of MCFs.** The following results from [23] and [36] are useful for constructing MCFs.

Proposition 7.5. *Let $\Lambda \subset J^1M$ be a 1-dimensional Legendrian, and let $\mathcal{C}_0 = (H, \{d_\nu\})$ be an ρ-graded MCF for Λ.*

1. *Suppose that H' is a handleslide set for the product cobordism $j^1(1_{[0,1]} \cdot \Lambda) \subset J^1([0,1] \times M)$ as in Definition 5.4 (2) (and without super-handleslide points) that agrees with H when restricted to $\{0\} \times \Lambda$ and satisfies Axiom 5.5. Then, there is a unique ρ-graded MCF for $j^1(1_{[0,1]} \cdot \Lambda)$ that agrees with $\mathcal{C}_0$ on $\{0\} \times \Lambda$ and has handleslide set H'.*
2. *Construct a handleslide set H' for $j^1(1_{[0,1]} \cdot \Lambda) \subset J^1([0,1] \times M)$ by extending each handleslide point p of $\mathcal{C}_0$ to a handleslide arc along $[0,1] \times \{p\}$, and for some region $R_\nu \subset M \setminus (\Lambda_{sing} \cap H)$ and some $i < j$ such that $\mu(S_i^\nu) = \mu(S_j^\nu) - 1$ placing a single (i,j)-super handleslide point, q, in the interior of $[0,1] \times R_\nu$ and adding handleslide arcs in $[0,1] \times R_\nu$ as specified by Axiom 5.5 (2) that connect q to $\{1\} \times R_\nu$ with monotonically increasing $[0,1]$ component. Then, there is a unique ρ-graded MCF for $j^1(1_{[0,1]} \cdot \Lambda)$ that agrees with $\mathcal{C}_0$ on $\{0\} \times \Lambda$ and has handleslide set H'.*

Proof. In all cases, it needs to be shown that it is possible to produce the required collection of differentials $\{d_\nu\}$ extending the given ones and so that Axiom 5.2 is satisfied. (1) and (2) are both consequences of Proposition 3.8 from [23] which shows that the differentials $\{d_\nu\}$ can be extended from $[0, t_0 - \epsilon] \times M$ to $[0, t_0 + \epsilon] \times M$ when a generic bifurcation of the t-slices of H occurs at $t = t_0$. (2) corresponds to Henry's Move 17 in [23]. Alternatively, see Proposition 6.3 of [36]. □

Next, we make a sequence of observations (B1)-(B6) about extending MCFs along various types of elementary cobordisms that will form the building blocks for the proof of Proposition 7.1. In all cases it should be observed that the Maslov potential for the given 1-dimensional Legendrian extends in a unique way over the elementary cobordism, though we mostly leave this implicit. Moreover, although we omit the terminology, all MCFs under consideration are ρ-graded.

(B1) *If $\mathcal{C}_0$ and $\mathcal{C}_1$ are equivalent MCFs on $\Lambda \subset J^1\mathbb{R}$, then there exists a MCF $\mathcal{C}$ on $j^1(1_{[0,1]} \cdot \Lambda) \subset J^1([0,1] \times \mathbb{R})$ that restricts to $\mathcal{C}_i$ above $\{i\} \times \mathbb{R}$, for $i = 0, 1$.*

Proof. This is the definition of equivalence. □

(B2) *Let Λ_t, $0 \leq t \leq 1$ be a Legendrian isotopy, and consider the Legendrian surface $\Sigma \subset J^1([0,1] \times \mathbb{R})$ such that the $\{t\} \times \mathbb{R}$ slices of the front projection of Σ are the front projections of the Λ_t. Any MCF on Λ_0 extends over Σ.*

Proof. It suffices to consider the case where the isotopy contains a single bifurcation of the front or base projection, i.e. a single Reidemeister move or an instance at which two singularities (crossings or cusps) at different locations in the front diagram share the same x-coordinate. See Figure 14.

Before applying the move, we apply (B1) together with Henry's result (Proposition 7.4) to assume that the given MCF $\mathcal{C}_0$ is in SR-form with respect to some normal ruling σ.

- For a Type III move or any of the "Other" moves (both directions): Use Proposition 7.5 to first move all of the handleslides outside of the x-interval that contains the pictured part of the diagram where the move occurs. Then, we apply the move, and check that there is a unique way to assign differentials to the new region(s) that arise from the move so that Axiom 5.2 is satisfied.




FIGURE 14. Generic bifurcations of the front and base projections during a Legendrian isotopy include the Reidemeister moves I-III, and instances where two front singularities share the same x-coordinate (marked "Other" in the figure). Vertical and horizontal reflections of all moves are allowed.




FIGURE 15. Labeling of the regions in the base projection near a triple point.

This is a straightforward case-by-case check that we illustrate by considering the Type III move in detail. (Other cases are left to the reader.) The t-slices before (resp. after) the move occurs intersect a sequence of 4 regions of $\pi_x(\Sigma) \setminus (\Sigma_{sing} \cup H)$, which we label R_A, R_B, R_C, and R_D (resp. R_A, $R_{B'}$, $R_{C'}$, and R_D); see Figure 15. The differentials on R_A, R_B, R_C, and R_D are already specified by $\mathcal{C}_0$. Suppose that the three sheets that intersect at the triple point are numbered as $S^{\nu}_k, S^{\nu}_{k+1}, S^{\nu}_{k+2}$ above these regions. (Recall that above any particular region we label sheets with decreasing z-coordinate, eg. the sheet labeled S^A_k becomes S^B_{k+1} when it passes through the crossing locus.) Write $Q_{i\,i+1}$ for a linear map that interchanges the i and $i+1$ sheets. According to Axiom 5.2 (2) we must define the differentials on $R_{B'}$ and $R_{C'}$ so that the following maps are chain isomorphisms,

$$Q_{k+1\,k+2} : (V(R_A), d_A) \xrightarrow{\cong} (V(R_{B'}), d_{B'}), \quad \text{and} \quad Q_{k\,k+1} : (V(R_{B'}), d_{B'}) \xrightarrow{\cong} (V(R_{C'}), d_{C'}). \tag{7.2}$$

In order for this to define a valid MCF we only need to check that Axiom 5.2 (2) is satisfied along the border between $R_{C'}$ and R_D, i.e., we need to show that

$$Q_{k+1\,k+2} : (V(R_{C'}), d_{C'}) \to (V(R_D), d_D)$$

is a chain map. In view of (7.2), this is equivalent to the composition $Q_{k+1\,k+2} \circ Q_{k\,k+1} \circ Q_{k+1\,k+2} : (V(R_A), d_A) \to (V(R_D), d_D)$ being a chain isomorphism. Since we know that Axiom 5.2 is satisfied by $\mathcal{C}_0$, when we pass from R_A to R_D by way of R_B and R_C we see that $Q_{k\,k+1} \circ Q_{k+1\,k+2} \circ Q_{k\,k+1} : (V(R_A), d_A) \to (V(R_D), d_D)$ is a chain isomorphism, so we can just note the braid relation $Q_{k\,k+1} \circ Q_{k+1\,k+2} \circ Q_{k\,k+1} = Q_{k+1\,k+2} \circ Q_{k\,k+1} \circ Q_{k+1\,k+2}$.

- For a Type I move ($\leftarrow$ direction): A crossing, b, a left cusp, c_l, and a right cusp, c_r, of Λ_0 all vanish at a swallowtail point, s, during the move. The crossing, b, must be a switch for σ, so under the SR-form assumption, when $\rho \neq 1$, the handleslides appearing near the crossing are exactly those required to have endpoints at the swallowtail point by Axiom 5.5 (3). Extend these to handleslide arcs with endpoints at s. Let ∂_{ν_l} (resp.




FIGURE 16. Removing a handleslide at a return to prepare for a Type II move. The top row illustrates the MCF on a sequence of t-slices with t increasing. The bottom row depicts the MCF in the base projection. If the return is as in the 2nd or 3rd column of Figure 12, there will be a second handleslide arc with endpoint at the super-handleslide point. Neither of its endpoints are on the cusp sheets, so it can be moved away from the pictured part of the front projection.

∂_{ν_r}) denote the differential from $\mathcal{C}_0$ assigned to the region R_{ν_l} (resp. R_{ν_r}) that borders c_l on the left (resp. c_r on the right). After the Type I move, R_{ν_l} and R_{ν_r} merge to become a single region, so it is important to note that ∂_{ν_l} and ∂_{ν_r} agree. This is the case since by Remark 7.3 (2), they both agree with the standard ruling differential for σ.

When $\rho = 1$, the SR-form may also have a handleslide connecting the strands of c_r, and this requires a preliminary step to remove the handleslide. This is done using Proposition 7.5 to alter the handleslide set via the sequence of adjustments pictured in [31, Figure 16].

- For a Type I move ($\rightarrow$ direction): To extend $\mathcal{C}_0$, we just add additional handleslide arcs with end points at the swallowtail point as required by Axiom 5.5. Differentials for the new regions can be defined since the handleslide set on the slices after the swallow tail point occurs is in SR-form for the ruling σ' obtained from σ by making the new crossing into a switch. Alternatively, see Proposition 6.2 of [36].
- For a Type II move ($\leftarrow$ direction): Once again, the strategy is to use an equivalence to move all of the handleslides out of the x-inteval, I, where the front diagram is pictured and then perform the move. It is then routine to check that the required differentials can be defined to complete the extension of $\mathcal{C}_0$ to $\mathcal{C}$. The two pictured crossings of Λ_0 are a departure followed by a return, so (since $\mathcal{C}_0$ is in SR-form) the only handleslides that need to be moved are those that may appear near the return. Figure 16 illustrates a 2-dimensional MCF on $j^1(1_{[0,1]} \cdot \Lambda)$ that will remove all handleslides from I as required. This MCF involves a single super-handleslide point, so Proposition 7.5 (1) and (2) produce the required differentials. (A similar procedure applies for the horizontally and/or vertically reflected versions of Move II.)
- For a Type II move ($\rightarrow$ direction): There are no handleslides in the interval where the move occurs. We perform the move and then check that the differentials may be extended.

□

Remark 7.6. An alternate approach to establishing (B2) is made possible by the correspondence between MCFs and augmentations. When Φ is a Legendrian isotopy from Λ_- to Λ_+, as in Section 3.2, there is an invertible conical Legendrian cobordism, Σ_Φ, with embedded Lagrangian projection. Thus, because the induced augmentation set I_Σ has the form found in equation (3.4)




FIGURE 17. The Clasp, Pinch, and Unknot Moves, pictured as front projection slices (left) and in the base projection of the corresponding 2-dimensional cobordism (right). Shading in the base projection indicates the region where the two cusp sheets exist.




FIGURE 18. The Cusp Tangency Move refers to the interchange of the far left and far right diagrams. This is accomplished by a Type II Move and a Clasp Move, as pictured.




FIGURE 19. The requirement on handleslides from (B3) that allows for MCFs to be extended through a clasp move (left) or cusp tangency (right).

the calculation of I_Σ in terms of MCFs via Proposition 5.9 and Corollary 4.12 implies that any MCF for Λ_- can be extended over (a compact version of) Σ.

Along with moves associated to 1-dimensional Legendrian isotopies, the generic front bifurcations of the t-slices of a Legendrian cobordism include the **Clasp Move**, the **Pinch Move**, and the **Unknot Move** as pictured in Figure 17. These moves correspond to a local maximum or minimum in the t-direction for the crossing locus (in the case of the Clasp Move) or the cusp locus (for the Pinch and Unknot Move). It will also be convenient to consider a (non-generic) **Cusp Tangency Move** pictured in Figure 18 which can be realized by a combination of the Clasp Move with a Legendrian isotopy.

(B3) (a) *An MCF can be extended along the $\to$ direction of the Clasp Move if there is no handleslide that connects the crossing sheets and has its x-coordinate between the two crossings.*

(b) *An MCF can be extended along the Cusp Tangency Move (either direction) provided there is no handleslide that connects the crossing strands and has its x-coordinate to the left of the crossing.*

See Figure 19.

Proof. Let Σ be the Legendrian surface corresponding to the Clasp Move ($\rightarrow$), with Λ_0 and Λ_1 denoting the 1-dimensional slices before and after the move. Let $\mathcal{C}_0$ be an MCF for Λ_0 with no handleslides between the crossings. Then, there is a sequence of three adjacent regions R_A, R_B, R_C for $\mathcal{C}_0$ where R_B is between the crossings and R_A and R_C are to the left and to the right. Since there are no handleslides between the crossings, we can define a MCF for Σ by extending all handleslide points of $\mathcal{C}_0$ along straight line segments in the t-direction. Axiom 5.2 (2) for $\mathcal{C}_0$ shows that the differentials on the two regions R_A and R_B agree, so that there is a well defined differential on the common region for Σ that contains them.

The Cusp Tangency Move follows from the first case of (B3) since the hypothesis restricting the location of handleslides before the move implies that the move can be realized (in either direction) by a Type II Move followed by a Clasp Move ($\rightarrow$ direction) in such a way that there are no handleslides between the crossings when the Clasp Move is applied. □

(B4) *An SR-form MCF for Λ associated to a ρ-graded normal ruling σ can be extended to a ρ-graded MCF on the elementary cobordism Σ arising from applying a Pinch Move ($\rightarrow$ direction) to adjacent sheets of Λ, S_k^ν and S_{k+1}^ν, above a region where they are paired by σ.*

Note that the $\mathbb{Z}/\rho$-valued Maslov potential μ for Λ extends over Σ since the fact that σ is ρ-graded implies that $\mu(S_k^\nu) = \mu(S_{k+1}^\nu) + 1 \bmod \rho$.

Proof. At the location of the pinch move, the differential for the SR-form MCF, d_ν, is the standard ruling differential for σ. (See Remark 7.3 (2).) Thus, $d_\nu S_{k+1} = S_k$, and S_k and S_{k+1} do not appear in the differentials of the other generators. That is, the complex for $(V(R_\nu), d_\nu)$ splits as a direct sum $(V(R_\nu), d_\nu) = (C_1, d_1) \oplus (C_2, d_2)$ with C_2 spanned by $\{S_k, S_{k+1}\}$ and C_1 spanned by the rest of the sheets. Therefore, when we extend the MCF over the surface by using (C_1, d_1) in the region where S_k and S_{k+1} do not exist, the Axiom 5.2 is satisfied. □

(B5) *An MCF $\mathcal{C}$ can be extended along the Unknot Move ($\leftarrow$ direction), provided there is no handleslide connecting the two sheets of the unknot. (Note that such a handleslide cannot exist if $\mathcal{C}$ is ρ-graded and $\rho \neq 1$.)*

Proof. Using (B1) we can assume $\mathcal{C}$ is in SR-Form on the slice that precedes the unknot move. It then follows that (i) there are no handleslides with endpoints on the unknot sheets, and (ii) in the region where the unknot exists $(V(R_\nu), d_\nu) = (C_1, d_1) \oplus (C_2, d_2)$ where C_1 is spanned by the unknot sheets, S_k^ν and S_{k+1}^ν, and $d_\nu S_{k+1}^\nu = S_k$. Moreover, by Axiom 5.2 the differentials on the two regions adjacent to the unknot both agree with d_2. Thus, $\mathcal{C}$ extends in an obvious way over the surface. □

(B6) *Any MCF $\mathcal{C}$ can be extended along the $\leftarrow$ direction of the Pinch Move.*

Proof. Again using (B1) we can assume $\mathcal{C}$ has no handleslides near the two cusps that are joined by the Pinch Move. From Axiom 5.2 the differentials agree in the two regions where the cusp sheets exist, so that $\mathcal{C}$ extends over the surface. □

Remark 7.7. Although we will not need to use them in our proof of Theorem 1.2, it is also easy to give necessary and sufficient conditions for an MCF to extend over the remaining moves.

(1) Any MCF can be extended over the Unknot Move ($\rightarrow$ direction). (This is easy to see directly, but also follows from the fact that there are conical Legendrian cobordisms realizing these moves that have embedded Lagrangian projections; cf. [16, 6]. The same comment applies to the $\leftarrow$ Pinch Move.)

(2) An MCF can be extended over the Clasp Move ($\leftarrow$ direction) if and only if $\langle d_\nu S_{k+1}, S_k \rangle = 0$ where R_ν is the region where the crossings will appear and S_k and S_{k+1} are the two crossing sheets.




FIGURE 20. The elementary tangles $l_k^{n-2,n}, r_k^{n,n-2}$, and $\sigma_k^{n,n}$. The picture illustrates $l_3^{5,7}$, $r_4^{7,5}$, and $\sigma_4^{6,6}$ from left to right.




FIGURE 21. The front projection corresponding to the word $w_\Lambda = Xl_k\sigma_{k-1}\sigma_{k-2}\cdots\sigma_{k-s}\sigma_{k+1}\sigma_{k+2}\cdots\sigma_{k+t}Y$ when $s = 3$ and $t = 2$.

7.3. **Construction of the Legendrian filling.** The front projection of a generic Legendrian $\Lambda \subset J^1\mathbb{R}$ can be represented as a word w_Λ that is a product (left to right concatenation) of elementary Legendrian tangles, each one of which contains a single crossing or cusp. We notate these **elementary tangles** as $l_k^{n-2,n}, r_k^{n,n-2}$, and $\sigma_k^{n,n}$ in the case of a tangle with a left cusp, right cusp, or crossing respectively, where $n \geq 2$ and $1 \leq k \leq n-1$. The superscripts indicate the number of strands of the tangle at its left and right vertical boundaries, while k is the numbering of the upper of the two strands that is involved with the crossing or cusp when we number strands as 1 to n from top to bottom. See Figure 20. In the following, we suppress the superscripts from notation.

Proof of Proposition 7.1. Assume $\mathcal{C}_\Lambda$ is an ρ-graded MCF for $\Lambda \subset J^1\mathbb{R}$. We first prove the proposition *assuming* $\rho \neq 1$, and then close by indicating the minor modifications to the proof when $\rho = 1$.

Let w_Λ be the word representing the front projection of Λ, and note that if $\Lambda \neq \emptyset$ then w_Λ can be written in the form

$$w_\Lambda = Xl_k\sigma_{k-1}\sigma_{k-2}\cdots\sigma_{k-s}\sigma_{k+1}\sigma_{k+2}\cdots\sigma_{k+t}Y \tag{7.3}$$

where $s, t \geq 0$ and Y contains no left cusps. (For example, just take the l_k term to be the left cusp of Λ with largest x-coordinate, and put $s = t = 0$.) See Figure 21.

We prove the following statement by a nested induction. The outer induction is on c and the inner induction is on n.

Inductive Statement: Suppose that w_Λ has c left cusps, and if $c \geq 1$ assume that w_Λ can be written in the form (7.3) such that the length of Y is $|Y| = n$. Then, there exists a compact Legendrian filling $\Sigma \subset J^1([0,1] \times \mathbb{R})$ with a ρ-graded MCF $\mathcal{C}$ such that $\mathcal{C}|_\Lambda = \mathcal{C}_\Lambda$.

The case $c = 0$ is trivial since $\Lambda = \emptyset$. For fixed $c \geq 1$, assuming the statement holds for smaller values of c, we establish the statement via induction on n. The case $n = 0$ is vacuously true, since Y must have at least one right cusp. Assuming $n \geq 1$, we write $Y = zY'$ (allowing the possibility that Y' is the empty word) where z is an elementary tangle that is necessarily a

right cusp or a crossing, $z = r_j$ or $z = \sigma_j$. We consider cases depending on the vertical location of this right cusp (in Cases 1-5) or crossing (in Cases 6-10). By symmetry we can assume $j \leq k$. In addition, using (B1) and Proposition 7.4 we can assume that $\mathcal{C}_\Lambda$ is in SR-form with respect to a ρ-graded normal ruling σ.

Case 1: $z = r_j$ with $j \leq k - s - 2$. Then, z is a right cusp above all the sheets with cusps or crossings in the product $l_k\sigma_{k-1}\sigma_{k-2}\cdots\sigma_{k-s}\sigma_{k+1}\sigma_{k+2}\cdots\sigma_{k+t}$. So, a Legendrian isotopy of Λ modifies the front diagram by

$$Xl_k\sigma_{k-1}\sigma_{k-2}\cdots\sigma_{k-s}\sigma_{k+1}\sigma_{k+2}\cdots\sigma_{k+t}(r_jY') \to (Xr_j)l_{k-2}\sigma_{k-3}\sigma_{k-4}\cdots\sigma_{k-2-s}\sigma_{k-1}\sigma_k\cdots\sigma_{k-2+t}Y'.$$

Use (B2) to extend $\mathcal{C}_\Lambda$ over the Legendrian isotopy. Then, the inductive hypothesis on n applies to complete the construction of $(\Sigma, \mathcal{C})$.

Case 2: $z = r_{k-s-1}$. See Figure 22 for a summary of this case. Using a Legendrian isotopy, we have

$$w_\Lambda \to Xl_k\sigma_{k-1}\sigma_{k-2}\cdots\sigma_{k-s}r_{k-s-1}\sigma_{k-1}\sigma_k\cdots\sigma_{k-2+t}Y'.$$

Note that we must have $s \geq 1$, since otherwise a "zig-zag" l_kr_{k-1} would occur in Λ. This is not possible since MCFs do not exist for stabilized Legendrian links. The idea is to try to move the cusp l_k up next to the r_{k-s-1}. Consider the first crossing to the right of l_k, $x = \sigma_{k-1}$.




FIGURE 22. The first line outlines the cobordism used in Case 2, Subcase A and the second line outlines the cobordism for Subcase B. Subcase A will necessarily occur before the front at the far right of (B) is reached.

Subcase A: x is a switch for σ. Since $\mathcal{C}$ is in SR-form for σ, we apply a $\to$ Pinch Move to the right of x (extending the MCF using (B4)) followed by a Type I Reidemeister Move to produce an MCF on a surface with the slices

$$\cdots l_k\sigma_{k-1}\cdots \overset{(B4)}{\to} \cdots l_k\sigma_{k-1}r_kl_k\cdots \to \cdots l_k\sigma_{k-2}\cdots\sigma_{k-s}r_{k-s-1}\sigma_{k-1}\sigma_k\cdots\sigma_{k-2+t}Y'$$
$$\to \cdots\sigma_{k-2}\cdots\sigma_{k-s}r_{k-s-1}l_{k-2}\sigma_{k-1}\sigma_k\cdots\sigma_{k-2+t}Y'.$$

Then, the inductive hypothesis on n applies.

Subcase B: x is not a switch for σ. Then, it is a departure, so since $\mathcal{C}_\Lambda$ is in SR-form, it has no handleslides connecting the crossing strands to the left of $x = \sigma_{k-1}$. Then, we can apply (B3) to extend past the Cusp Tangency Move:

$$Xl_k\sigma_{k-1}\sigma_{k-2}\cdots\sigma_{k-s}r_{k-s-1}\sigma_{k-1}\sigma_k\cdots\sigma_{k-2+t}Y' \overset{(B3)}{\to} Xl_{k-1}\sigma_k\sigma_{k-2}\cdots\sigma_{k-s}r_{k-s-1}\sigma_{k-1}\sigma_k\cdots\sigma_{k-2+t}Y'$$
$$\to Xl_{k-1}\sigma_{k-2}\cdots\sigma_{k-s}r_{k-s-1}\sigma_{k-2}\sigma_{k-1}\sigma_k\cdots\sigma_{k-2+t}Y'.$$

We then repeat this argument with $x = \sigma_{k-2}$. Continuing in this manner, we must encounter the switch case at some point; if not, the left cusp would eventually be moved next to the right cusp, and we would have an MCF for a stabilized link which is impossible. (The front would have a "zig-zag", $l_{k-s}r_{k-s-1}$.)

Case 3: $s \geq 1$ and $z = r_{k-s}$. Then, the presence of the term $\sigma_{k-s}r_{k-s}$ shows that Λ is stabilized which is impossible. (There is a standard Legendrian isotopy that will turn the "fish tail" from the $\sigma_{k-s}r_{k-s}$ product into a "zig-zag".)

Case 4: $z = r_j$ with $k - s < j < k$. Then, a Type II move can be applied to produce a Legendrian isotopy

$$Xl_k\sigma_{k-1}\sigma_{k-2}\cdots\sigma_{k-s}\sigma_{k+1}\sigma_{k+2}\cdots\sigma_{k+t}(r_jY') \to (Xr_{j-1})l_{k-2}\sigma_{k-3}\cdots\sigma_{k-s}\sigma_{k-1}\sigma_k\cdots\sigma_{k-2+t}Y',$$

where the product $\sigma_{k-3}\cdots\sigma_{k-s}$ just means the identity tangle if $s = 2$. Again, we can apply (B2) and then the inductive hypothesis on n to produce $(\Sigma, \mathcal{C})$.

Case 5: $z = r_k$.

Subcase A: $s = t = 0$. Then, we have a standard Legendrian unknot in the middle of the diagram. *As long as* $\rho \neq 1$, there can be no handleslides connecting the strands of this unknot, so we apply (B5) to extend $\mathcal{C}$ as we remove the unknot. This decreases c, so that the inductive hypothesis applies.

Subcase B: Exactly one of $s > 0$ or $t > 0$. Then, we can apply a Type I Reidemeister move to reduce the number of cusps. The $(\Sigma, \mathcal{C})$ is then constructed via (B2) and the inductive hypothesis on c.

Subcase C: $s > 0$ and $t > 0$. If the crossing σ_{k-1} is a switch for the normal ruling σ associated to $\mathcal{C}_\Lambda$, then we can apply a pinch move, as in Case 2, followed by two Type I moves to decrease c while extending $\mathcal{C}$ via (B4) and (B2). If σ_{k-1} is not a switch, then we apply a Cusp Tangency Move as in (B3) to arrive at

$$Xl_{k-1}\sigma_{k-2}\cdots\sigma_{k-s}\sigma_k\sigma_{k+1}\sigma_{k+2}\cdots\sigma_{k+t}r_kY'.$$

At this point, Case 4 (reflected vertically) applies.

Case 6: $z = \sigma_j$ with $j \leq k - s - 2$. As in Case 1, move z to the left of l_k and then apply the inductive hypothesis on n.

Case 7: $z = \sigma_{k-s-1}$. Using a Legendrian isotopy, we can group σ_{k-s-1} into the product

$$X(l_k\sigma_{k-1}\sigma_{k-2}\cdots\sigma_{k-s}\sigma_{k+1}\sigma_{k+2}\cdots\sigma_{k+t})(\sigma_{k-s-1}Y') \to X(l_k\sigma_{k-1}\sigma_{k-2}\cdots\sigma_{k-s}\sigma_{k-s-1}\sigma_{k+1}\sigma_{k+2}\cdots\sigma_{k+t})Y'$$

and apply the inductive hypothesis on n.

Case 8: $s \geq 1$ and $z = \sigma_{k-s}$. Using an argument similar to Case 2, we attempt to apply cusp tangency moves to move the left cusp directly next to z. Initially, assume $k - 1 > k - s$. If the crossing $x = \sigma_{k-1}$ directly to the right of l_k is a switch, we can apply a Pinch Move followed by a Type I Reidemeister Move to go from

$$\begin{aligned}\cdots l_k\sigma_{k-1}\sigma_{k-2}\cdots \overset{(B4)}{\to} \cdots l_k\sigma_{k-1}r_kl_k\sigma_{k-2}\cdots \to \cdots l_k\sigma_{k-2}\cdots\sigma_{k-s}\sigma_{k-s}\sigma_{k+1}\sigma_{k+2}\cdots\sigma_{k+t}Y'\\ \to \cdots\sigma_{k-2}\cdots\sigma_{k-s}\sigma_{k-s}l_k\sigma_{k+1}\sigma_{k+2}\cdots\sigma_{k+t}Y'\end{aligned}$$

and then apply induction on n.

If $x = \sigma_{k-1}$ is not a switch, then it is a departure. Then, we apply (B3) to extend $\mathcal{C}$ during the sequence

$$\begin{aligned}Xl_k\sigma_{k-1}\sigma_{k-2}\cdots\sigma_{k-s}\sigma_{k+1}\sigma_{k+2}\cdots\sigma_{k+t}(\sigma_{k-s}Y') \overset{(B3)}{\to} Xl_{k-1}\sigma_k\sigma_{k-2}\cdots\sigma_{k-s}\sigma_{k-s}\sigma_{k+1}\sigma_{k+2}\cdots\sigma_{k+t}Y'\\ \to Xl_{k-1}\sigma_{k-2}\cdots\sigma_{k-s}\sigma_{k-s}\sigma_k\sigma_{k+1}\sigma_{k+2}\cdots\sigma_{k+t}Y'.\end{aligned}$$

We repeat this argument until we either find a switch or arrive at a word of the form

$$Xl_{k-s+1}\sigma_{k-s}\sigma_{k-s}\sigma_{k-s+2}\cdots\sigma_{k+t}Y'$$

If the first σ_{k-s} is a switch, then the usual combination of Pinch Move and Type I Reidemeister Move produces

$$Xl_{k-s+1}\sigma_{k-s}\sigma_{k-s+2}\cdots\sigma_{k+t}Y'$$

which has the form (7.3) with $|Y'| = n-1$ so that induction applies. If instead σ_{k-s} is a departure, then we can extend $\mathcal{C}$ over another Cusp Tangency Move and then apply a Reidemeister II move:

$$\begin{aligned} Xl_{k-s+1}\sigma_{k-s}\sigma_{k-s}\sigma_{k-s+2}\cdots\sigma_{k+t}Y' &\overset{(B3)}{\to} Xl_{k-s}\sigma_{k-s+1}\sigma_{k-s}\sigma_{k-s+2}\cdots\sigma_{k+t}Y' \\ &\to Xl_{k-s+1}\sigma_{k-s+2}\cdots\sigma_{k+t}Y'. \end{aligned}$$

Again, we are able to use the inductive hypothesis on n.

Case 9: $z = \sigma_j$ with $k-s < j < k$. Then, a Type III Reidemeister move can be applied:

$$Xl_k\sigma_{k-1}\sigma_{k-2}\cdots\sigma_{k-s}\sigma_{k+1}\sigma_{k+2}\cdots\sigma_{k+t}(\sigma_jY') \to (X\sigma_{j-1})l_k\sigma_{k-1}\sigma_{k-2}\cdots\sigma_{k-s}\sigma_{k+1}\sigma_{k+2}\cdots\sigma_{k+t}Y'.$$

We apply (B2) and then the inductive hypothesis on n to produce $(\Sigma, \mathcal{C})$.

Case 10: $z = \sigma_k$.

Subcase A: $s = t = 0$. This case cannot occur since the appearance of the product $l_k\sigma_k$ would show that Λ is stabilized.

Subcase B: Exactly one of $s > 0$ or $t > 0$. Then, a Type II Move allows us to apply induction on n. For instance, when $s > 0$ and $t = 0$,

$$\begin{aligned} Xl_k\sigma_{k-1}\sigma_{k-2}\cdots\sigma_{k-s}(\sigma_kY') &\to Xl_k\sigma_{k-1}\sigma_k\sigma_{k-2}\cdots\sigma_{k-s}Y' \\ &\to Xl_{k-1}\sigma_{k-2}\cdots\sigma_{k-s}Y'. \end{aligned}$$

Subcase C: Both $s > 0$ and $t > 0$. We have

$$Xl_k\sigma_{k-1}\cdots\sigma_{k-s}\sigma_{k+1}\cdots\sigma_{k+t}(\sigma_kY') \to Xl_k\sigma_{k-1}\sigma_{k+1}\sigma_k(\sigma_{k-2}\cdots\sigma_{k-s})(\sigma_{k+2}\cdots\sigma_{k+t})Y'.$$

If the σ_{k-1} is a switch, then we apply (B4) to do a Pinch Move followed by a Type I and Type II Move:

$$\begin{aligned} Xl_k\sigma_{k-1}\sigma_{k+1}\sigma_k\cdots &\overset{(B4)}{\to} Xl_k\sigma_{k-1}r_kl_k\sigma_{k+1}\sigma_k\cdots \\ &\to Xl_k\sigma_{k+1}\sigma_k\cdots \\ &\to Xl_{k+1}(\sigma_{k-2}\cdots\sigma_{k-s})(\sigma_{k+2}\cdots\sigma_{k+t})Y' \\ &\to (X\sigma_{k-2}\cdots\sigma_{k-s})(l_{k+1}\sigma_{k+2}\cdots\sigma_{k+t})Y'. \end{aligned}$$

Finally, if σ_{k-1} is not a switch, we use (B3) and apply a Cusp Tangency Move followed by a Type III Move:

$$\begin{aligned} Xl_k\sigma_{k-1}\sigma_{k+1}\sigma_k\cdots &\overset{(B3)}{\to} Xl_{k-1}\sigma_k\sigma_{k+1}\sigma_k\cdots \\ &\to Xl_{k-1}\sigma_{k+1}\sigma_k\sigma_{k+1}(\sigma_{k-2}\cdots\sigma_{k-s})(\sigma_{k+2}\cdots\sigma_{k+t})Y' \\ &\to (X\sigma_{k+1})l_{k-1}(\sigma_{k-2}\cdots\sigma_{k-s})(\sigma_k\sigma_{k+1}\sigma_{k+2}\cdots\sigma_{k+t})Y'. \end{aligned}$$

Then, induction on n applies.

This completes the proof when $\rho \neq 1$. The only place where the hypothesis $\rho \neq 1$ was used in the above induction was Subcase A of Case 5 where (B5) was applied to remove a Legendrian unknot component. When $\rho = 1$, instead of applying (B5) for this subcase, we can simply apply a Legendrian isotopy to move the unknot component to the left of the rest of the front. In this manner, the inductive argument produces a cobordism Σ from a disjoint union of standard Legendrian unknots (each with 2 cusps and no crossings) to Λ together with an MCF $\mathcal{C}$ on Σ

extending $\mathcal{C}_\Lambda$. Finally, we can apply the (B6) to use the ($\leftarrow$) direction of the Pinch Move to join all of the unknots into a single unknot via a cobordism.

□

References

[1] V. I. Arnol'd. Lagrange and Legendre cobordisms. I. *Funktsional. Anal. i Prilozhen.*, 14(3):1–13, 96, 1980.
[2] V. I. Arnol'd. Lagrange and Legendre cobordisms. II. *Funktsional. Anal. i Prilozhen.*, 14(4):8–17, 95, 1980.
[3] V. I. Arnol'd. *Singularities of caustics and wave fronts*, volume 62 of *Mathematics and its Applications (Soviet Series)*. Kluwer Academic Publishers Group, Dordrecht, 1990.
[4] M. Audin. *Cobordismes d'immersions lagrangiennes et legendriennes*, volume 20 of *Travaux en Cours [Works in Progress]*. Hermann, Paris, 1987.
[5] F. Bourgeois and B. Chantraine. Bilinearized Legendrian contact homology and the augmentation category. *J. Symplectic Geom.*, 12(3):553–583, 2014.
[6] F. Bourgeois, J. M. Sabloff, and L. Traynor. Lagrangian cobordisms via generating families: construction and geography. *Algebr. Geom. Topol.*, 15(4):2439–2477, 2015.
[7] B. Chantraine. Lagrangian concordance of Legendrian knots. *Algebr. Geom. Topol.*, 10(1):63–85, 2010.
[8] B. Chantraine. Lagrangian concordance is not a symmetric relation. *Quantum Topol.*, 6(3):451–474, 2015.
[9] Y. Chekanov. Differential algebra of Legendrian links. *Invent. Math.*, 150(3):441–483, 2002.
[10] C. Cornwell, L. Ng, and S. Sivek. Obstructions to Lagrangian concordance. *Algebr. Geom. Topol.*, 16(2):797–824, 2016.
[11] G. Dimitroglou Rizell. Lifting pseudo-holomorphic polygons to the symplectisation of $P\times\mathbb{R}$ and applications. *Quantum Topol.*, 7(1):29–105, 2016.
[12] T. Ekholm. Morse flow trees and Legendrian contact homology in 1-jet spaces. *Geom. Topol.*, 11:1083–1224, 2007.
[13] T. Ekholm. Rational symplectic field theory over $\mathbb{Z}_2$ for exact Lagrangian cobordisms. *J. Eur. Math. Soc. (JEMS)*, 10(3):641–704, 2008.
[14] T. Ekholm, J. Etnyre, and M. Sullivan. Legendrian contact homology in $P \times \mathbb{R}$. *Trans. Amer. Math. Soc.*, 359(7):3301–3335, 2007.
[15] T. Ekholm, J. Etnyre, and M. Sullivan. Legendrian contact homology in $P \times \mathbb{R}$. *Trans. Amer. Math. Soc.*, 359(7):3301–3335, 2007.
[16] T. Ekholm, K. Honda, and T. Kálmán. Legendrian knots and exact Lagrangian cobordisms. *J. Eur. Math. Soc. (JEMS)*, 18(11):2627–2689, 2016.
[17] T. Ekholm and T. Kálmán. Isotopies of Legendrian 1-knots and Legendrian 2-tori. *J. Symplectic Geom.*, 6(4):407–460, 2008.
[18] J. Eliashberg. Cobordisme des solutions de relations différentielles. In *South Rhone seminar on geometry, I (Lyon, 1983)*, Travaux en Cours, pages 17–31. Hermann, Paris, 1984.
[19] Y. Eliashberg. Invariants in contact topology. In *Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998)*, number Extra Vol. II, pages 327–338, 1998.
[20] Y. Eliashberg, A. Givental, and H. Hofer. Introduction to symplectic field theory. Number Special Volume, Part II, pages 560–673. 2000. GAFA 2000 (Tel Aviv, 1999).
[21] D. Fuchs. Chekanov-Eliashberg invariant of Legendrian knots: existence of augmentations. *J. Geom. Phys.*, 47(1):43–65, 2003.
[22] K. Hayden and J. M. Sabloff. Positive knots and Lagrangian fillability. *Proc. Amer. Math. Soc.*, 143(4):1813–1821, 2015.
[23] M. B. Henry. Connections between Floer-type invariants and Morse-type invariants of Legendrian knots. *Pacific J. Math.*, 249(1):77–133, 2011.
[24] M. B. Henry and D. Rutherford. A combinatorial DGA for Legendrian knots from generating families. *Commun. Contemp. Math.*, 15(2):1250059, 60, 2013.
[25] M. B. Henry and D. Rutherford. Equivalence classes of augmentations and Morse complex sequences of Legendrian knots. *Algebr. Geom. Topol.*, 15(6):3323–3353, 2015.
[26] M. Limouzineau. On Legendrian cobordisms and generating functions. *J. Knot Theory Ramifications*, 29(3):2050008, 17, 2020.
[27] L. Ng, D. Rutherford, V. Shende, S. Sivek, and E. Zaslow. Augmentations are sheaves. *Geom. Topol., to appear*.
[28] L. L. Ng. Computable Legendrian invariants. *Topology*, 42(1):55–82, 2003.
[29] Y. Pan. The augmentation category map induced by exact Lagrangian cobordisms. *Algebr. Geom. Topol.*, 17(3):1813–1870, 2017.
[30] Y. Pan and D. Rutherford. Augmentations and immersed lagrangian fillings. *Preprint, arXiv:2006.16436*.

[31] Y. Pan and D. Rutherford. Augmented Legendrian cobordism in J^1S^1. *Preprint, arXiv:2111.10065v2.*
[32] Y. Pan and D. Rutherford. Functorial LCH for immersed Lagrangian cobordisms. *Preprint.*
[33] D. Rutherford and M. Sullivan. Cellular Legendrian contact homology for surfaces, part I. *Preprint.*
[34] D. Rutherford and M. Sullivan. Cellular Legendrian contact homology for surfaces, part II. *Internat. J. Math.*, 30(7):1950036, 135, 2019.
[35] D. Rutherford and M. Sullivan. Cellular Legendrian contact homology for surfaces, part III. *Internat. J. Math.*, 30(7):1950037, 111, 2019.
[36] D. Rutherford and M. G. Sullivan. Generating families and augmentations for Legendrian surfaces. *Algebr. Geom. Topol.*, 18(3):1675–1731, 2018.
[37] J. M. Sabloff. Augmentations and rulings of Legendrian knots. *Int. Math. Res. Not.*, (19):1157–1180, 2005.
[38] V. Shende, D. Treumann, H. Williams, and E. Zaslow. Cluster varieties from Legendrian knots. *Duke Math. J.*, 168(15):2801–2871, 2019.
[39] T. Su. Ruling polynomials and augmentations for Legendrian tangles. *Preprint.*
[40] K. Tagami. On the Lagrangian fillability of almost positive links. *J. Korean Math. Soc.*, 56(3):789–804, 2019.
[41] V. A. Vassilyev. *Lagrange and Legendre characteristic classes*, volume 3 of *Advanced Studies in Contemporary Mathematics*. Gordon and Breach Science Publishers, New York, 1988. Translated from the Russian.

Tianjin University
Email address: ypan@tju.edu.cn

Ball State University
Email address: rutherford@bsu.edu