Guarantees for Tuning the Step Size using a Learning-to-Learn Approach

Xiang Wang* Shuai Yuan[†] Chenwei Wu[‡] Rong Ge[§]

Abstract

Choosing the right parameters for optimization algorithms is often the key to their success in practice. Solving this problem using a learning-to-learn approach—using meta-gradient descent on a meta-objective based on the trajectory that the optimizer generates—was recently shown to be effective. However, the meta-optimization problem is difficult. In particular, the meta-gradient can often explode/vanish, and the learned optimizer may not have good generalization performance if the meta-objective is not chosen carefully. In this paper we give meta-optimization guarantees for the learning-to-learn approach on a simple problem of tuning the step size for quadratic loss. Our results show that the naïve objective suffers from meta-gradient explosion/vanishing problem. Although there is a way to design the meta-objective so that the meta-gradient remains polynomially bounded, computing the meta-gradient directly using backpropagation leads to numerical issues. We also characterize when it is necessary to compute the meta-objective on a separate validation set to ensure the generalization performance of the learned optimizer. Finally, we verify our results empirically and show that a similar phenomenon appears even for more complicated learned optimizers parametrized by neural networks.

1 Introduction

Choosing the right optimization algorithm and related hyper-parameters is important for training a deep neural network. Even for simple algorithms like gradient descent and stochastic gradient descent, choosing a good step size can be important to the convergence speed and generalization performance. Empirically, the parameters are often chosen based on past experiences or grid search. Recently, Maclaurin et al. (2015) considered the idea of tuning these parameters by optimization—that is, consider a meta-optimization problem where the goal is to find the best parameters for an optimizer. A series of works (e.g., Andrychowicz et al. (2016); Wichrowska et al. (2017)) extended such ideas and parametrized the set of optimizers by neural networks.

Although this approach has shown empirical success, there are very few theoretical guarantees for learned optimizers. Gupta & Roughgarden (2017) gave sample complexity bounds for tuning the step size, but they did not address how one can find the learned optimizer efficiently. In practice, the meta-optimization problem is often solved by meta-gradient descent—define a meta-objective function based on the trajectory that the optimizer generates, and then compute the meta-gradient using back-propagation (Franceschi et al., 2017). The optimization for meta-parameters is usually a nonconvex problem, therefore it is unclear why simple meta-gradient descent would find an optimal solution.

In this paper we consider using learning-to-learn approach to tune the step size of standard gradient descent/stochastic gradient descent algorithm. Even in this simple setting, many of the challenges still remain and we can get better learned optimizers by choosing the right meta-objective function. Though our results are proved only in the simple setting, we empirically verify the results using complicated learned optimizers with neural network parametrizations.

^{*}Duke University. Email: xwang@cs.duke.edu

[†]Duke University. Email: shuai@cs.duke.edu

[‡]Duke University. Email: chenwei.wu592@duke.edu

[§]Duke University. Email: rongge@cs.duke.edu

1.1 Our Results

In this paper we focus on two basic questions on learning-to-learn for gradient descent optimizer. First, will the metagradient explode/vanish and is there a way to fix the problem? Second, how could we guarantee that the learned optimizer has good generalization properties?

Our first result shows that meta-gradient can explode/vanish even for tuning the step size for gradient descent on a simple quadratic objective. In this setting, we show that there is a unique local and global minimizer for the step size, and we also give a simple way to get rid of the gradient explosion/vanishing problem.

Theorem 1 (Informal version of Theorem 3 and Theorem 4). For tuning the step size of gradient descent on a quadratic objective, if the meta-objective is the loss of the last iteration, then the meta-gradient will explode/vanish. If the meta-objective is the log of the loss of the last iteration, then the meta-gradient is polynomially bounded. Further, doing meta-gradient descent with a meta step size of $1/\sqrt{k}$ (where k is the number of meta-gradient steps) provably converges to the optimal step size for the inner-optimizer.

Surprisingly, even though taking the log of the objective solves the meta-gradient explosion/vanishing problem, one cannot simply implement such an algorithm using back-propagation (which is standard in auto-differentiation tools such as those used in TensorFlow (Abadi et al., 2016)). The reason is that even though the meta-gradient is polynomially bounded, back-propagation algorithm will compute the meta-gradient as the ratio of two exponentially large/small numbers, which causes numerical issues. Detailed discussion for the first result appears in Section 3.

Our second result shows that defining meta-objective on the same training set (later referred to as the "train-by-train" approach) could lead to overfitting; while defining meta-objective on a separate validation set ("train-by-validation", see Metz et al. (2019)) can solve this issue. We consider a simple least squares setting where $y = \langle w^*, x \rangle + \xi$ and $\xi \sim \mathcal{N}(0, \sigma^2)$. We show that when the number of samples is small and the noise is large, it is important to use train-by-validation; while when the number of samples is much larger train-by-train can also learn a good optimizer.

Theorem 2 (Informal version of Theorem 5 and Theorem 6). For a least squares problem in d dimensions, if the number of samples n is a constant fraction of d (e.g., d/2), and the samples have large noise, then the train-by-train approach performs much worse than train-by-validation. On the other hand, when the number of samples n is large, train-by-train can get close to error $d\sigma^2/n$, which is optimal.

We discuss the details in Section 4. In Section 5 we show that such observations also hold empirically for more complicated learned optimizers—an optimizer parametrized by a neural network.

1.2 Related Work

Learned optimizer The idea of learning an optimizer has appeared in early works decades ago (Bengio et al., 1990, 1992; Hochreiter et al., 2001). Recently, with the rise of deep learning, researchers started to consider more complex optimizers on more challenging tasks. One line of research views the optimizer as a policy and apply reinforcement learning techniques to train it (Li & Malik, 2016, 2017; Bello et al., 2017). The other line of papers use gradient descent on the meta-objective to update the optimizer parameters (Maclaurin et al., 2015; Andrychowicz et al., 2016; Lv et al., 2017; Wichrowska et al., 2017; Metz et al., 2019).

Mostly relevant to our work, Metz et al. (2019) highlighted several challenges in the meta-optimization for learning-to-learn approach. First, they observed the meta-gradient exploding/vanishing issue and proposed to use a gradient estimator for a variational meta-objective. They also observed that train-by-train approach can overfit the training tasks while train-by-validation generalizes well.

Data-driven algorithm design In data-driven algorithm design, we aim to find an algorithm that works well on a particular distribution of tasks. Gupta & Roughgarden (2017) first modeled this algorithm-selection process as a statistical learning problem. In particular, they analyzed the sample complexity of choosing the step size for gradient descent. But they didn't consider the meta-optimization problem. They also restricted the step size into a small range so that gradient descent is guaranteed to converge on every task. We don't have such a restriction and allow the meta-learning to choose a more aggressive step size.

Following the work by Gupta & Roughgarden (2017), data-driven algorithms have been studied in many problems, including partitioning and clustering (Balcan et al., 2016a), tree search (Balcan et al., 2018a), pruning (Alabi et al., 2019) and machanism design (Morgenstern & Roughgarden, 2015, 2016; Balcan et al., 2016b, 2018b).

Step size schedule for GD/SGD Shamir & Zhang (2013) showed that SGD with polynomial step size scheduling can almost match the minimax rate in convex non-smooth settings, which was later tightened by Harvey et al. (2018) for standard step size scheduling. Assuming that the number of training steps is known to the algorithm, the information-theoretically optimal bound in convex non-smooth setting was later achieved by Jain et al. (2019) which used another step size schedule, and Ge et al. (2019) showed that exponentially decaying step size scheduling can achieve near optimal rate for least squares regression.

A closely related paper that appeared later than our work also studied the comparison between train-by-train and train-by-validation (Bai et al., 2020). They considered a very different meta-learning problem, where the goal is to find the best common initialization for adapting to a linear predictor on each task. They proved train-by-train can work better than train-by-validation in the noiseless setting.

2 Preliminaries

In this section, we first introduce some notations, then formulate the learning-to-learn framework.

2.1 Notations

For any integer n, we use [n] to denote $\{1, 2, \cdots, n\}$. We use $\|\cdot\|$ to denote the ℓ_2 norm for a vector and the spectral norm for a matrix. We use $\langle \cdot, \cdot \rangle$ to denote the inner product of two vectors. For a symmetric matrix $A \in \mathbb{R}^{d \times d}$, we denote its eigenvalues as $\lambda_1(A) \geq \cdots \geq \lambda_d(A)$. We denote the d-dimensional identity matrix as I_d or simply as I when the dimension is clear. We use $O(\cdot), \Omega(\cdot), \Theta(\cdot)$ to hide constant factor dependencies. We use $\operatorname{poly}(\cdot)$ to represent a polynomial on the relevant parameters with constant degree.

2.2 Learning-to-learn Framework

We consider the learning-to-learn approach applied to training a distribution of learning tasks. Each task is specified by a tuple $(\mathcal{D}, S_{\text{train}}, S_{\text{valid}}, \ell)$. Here \mathcal{D} is a distribution of samples in $X \times Y$, where X is the domain for the sample and Y is the domain for the label/value. The sets S_{train} and S_{valid} are samples generated independently from \mathcal{D} , which serve as the training and validation set (the validation set is optional). The learning task looks to find a parameter $w \in W$ that minimizes the loss function $\ell(w, x, y) : W \times X \times Y \to \mathbb{R}$, which gives the loss of the parameter w for sample (x, y). The training loss for this task is

$$\hat{f}(w) := \frac{1}{|S_{\text{train}}|} \sum_{(x, y) \in S_{\text{train}}} \ell(w, x, y),$$

while the population loss is $f(w) := \mathbb{E}_{(x,y) \sim \mathcal{D}}[\ell(w,x,y)].$

The goal of inner-optimization is to minimize the population loss f(w). For the learned optimizer, we consider it as an update rule $u(\cdot)$ on weight w. The update rule is a parameterized function that maps the weight at step τ and its history to the step $\tau+1: w_{\tau+1}=u(w_{\tau},\nabla \hat{f}(w_{\tau}),\nabla \hat{f}(w_{\tau-1}),\cdots;\theta)$. In most parts of this paper, we consider the update rule u as gradient descent mapping with step size as the trainable parameter (here $\theta=\eta$ which is the step size for gradient descent). That is, $u(w;\eta)=w-\eta\nabla \hat{f}(w)$ for gradient descent and $u(w;\eta)=w-\eta\nabla_w\ell(w,x,y)$ for stochastic gradient descent where (x,y) is a sample randomly chosen from the training set S_{train} .

In the outer (meta) level, we consider a distribution \mathcal{T} of tasks. For each task $P \sim \mathcal{T}$, we can define a meta-loss function $\Delta(\theta, P)$. The meta-loss function measures the performance of the optimizer on this learning task. The meta-objective, for example, can be chosen as the target training loss \hat{f} at the last iteration (train-by-train), or the loss on the validation set (train-by-validation).

The training loss for the meta-level is the average of the meta-loss across m different specific tasks $P_1, P_2, ..., P_m$, that is,

$$\hat{F}(\theta) = \frac{1}{m} \sum_{i=1}^{m} \Delta(\theta, P_k).$$

The population loss for the meta-level is the expectation over all the possible specific tasks $F(\theta) = \mathbb{E}_{P \sim \mathcal{T}}[\Delta(\theta, P)]$.

In order to train an optimizer by gradient descent, we need to compute the gradient of meta-objective \hat{F} in terms of meta parameters θ . The meta parameter is updated once after applying the optimizer on the inner objective t times to generate the trajectory $w_0, w_1, ..., w_t$. The meta-gradient is then computed by unrolling the optimization process and back-propagating through the t applications of the optimizer.

3 Alleviating Gradient Explosion/Vanishing Problems

First we consider the meta-gradient explosion/vanishing problem. More precisely, we say the meta-gradient explodes/vanishes if it is exponentially large/small with respect to the number of steps t of the inner-optimizer.

In this section, we consider a simple instance of the learning-to-learn approach, where the distribution \mathcal{T} only contains a single task P, and the task also just defines a single loss function f^1 . Therefore, in this section $\hat{F}(\eta) = F(\eta) = \Delta(\eta, P)$. We will simplify notation and only use $\hat{F}(\eta)$.

The inner task P is a simple quadratic problem, where the starting point is fixed at w_0 with unit norm, and the loss function is $f(w) = \frac{1}{2} w^{\top} H w$ for some fixed positive definite matrix $H \in \mathbb{R}^{d \times d}$.

Let $\{w_{\tau,\eta}\}_{\tau=0}^t$ be the GD sequence running on f(w) starting from w_0 with step size η . We consider two ways of defining meta-objective: using the loss of the last point directly or using the log of this value. We first show that although choosing $\hat{F}(\eta) = f(w_{t,\eta})$ does not have any bad local optimal solution, it has the meta-gradient explosion/vanishing problem. We use $\hat{F}'(\eta)$ to denote the derivative of \hat{F} in η .

In the analysis, we use eigen-decomposition to transform H into a diagonal matrix. We introduce related notations here: suppose the eigenvalue decomposition of H is $\sum_{i=1}^d \lambda_i u_i u_i^{\top}$. We denote $L:=\lambda_1(H)$ and $\alpha:=\lambda_d(H)$ as the largest and smallest eigenvalues of H. For each $i\in [d]$, let c_i be $\langle w_0,u_i\rangle$ and let c_{\min} be $\min(|c_1|,|c_d|)$. We assume $c_{\min}>0$ and $L>\alpha$ for simplicity².

Theorem 3. Let the meta-objective be $\hat{F}(\eta) = f(w_{t,\eta})$, we know $\hat{F}(\eta)$ is a strictly convex function in η with an unique minimizer. However, for any step size $0 < \eta < 2/L$,

$$|\hat{F}'(\eta)| \le tL^2 \max(|1 - \eta \alpha|^{2t-1}, |1 - \eta L|^{2t-1});$$

for any step size $\eta > 2/L$,

$$|\hat{F}'(\eta)| \ge c_1^2 L^2 t (\eta L - 1)^{2t - 1} - L^2 t.$$

Note that in Theorem 3, when $0 < \eta < 2/L$, $|\hat{F}'(\eta)|$ is exponentially small because $|1 - \eta \alpha|$, $|1 - \eta L| < 1$; when $\eta > 2/L$, $|\hat{F}'(\eta)|$ is exponentially large because $\eta L - 1 > 1$. The strict convexity of $\hat{F}(\eta)$ is proved by showing the second order derivative of $\hat{F}(\eta)$ is positive; the upper and lower bounds of $\hat{F}'(\eta)$ follows from direct calculation.

Intuitively, gradient explosion/vanishing happens because the meta-objective becomes too small or too large. A natural idea to fix the problem is to take the log of the meta-objective to reduce its range. If we choose $\hat{F}(\eta) = \frac{1}{t} \log f(w_{t,\eta})$, we have

Theorem 4. Let the meta-objective be $\hat{F}(\eta) = \frac{1}{t} \log f(w_{t,\eta})$. We know $\hat{F}(\eta)$ has a unique minimizer η^* and $\hat{F}'(\eta) = O\left(\frac{L^3}{c_{\min}^2\alpha(L-\alpha)}\right)$ for all $\eta \geq 0$. Let $\{\eta_k\}$ be the GD sequence running on \hat{F} with meta step size $\mu_k = 1/\sqrt{k}$. Suppose the starting step size $\eta_0 \leq M$. Given any $1/L > \epsilon > 0$, there exists $k' = \frac{M^6}{\epsilon^2} poly(\frac{1}{c_{\min}}, L, \frac{1}{\alpha}, \frac{1}{L-\alpha})$ such that for all $k \geq k', |\eta_k - \eta^*| \leq \epsilon$.

In the notation of Section 2, one can think that \mathcal{D} contains a single point (0,0) and the loss function $f(w) = \ell(w,0,0)$.

²If w_0 is uniformly sampled from the unit sphere, with high probability c_{\min} is at least $\Omega(1/\sqrt{d})$; if H is XX^{\top} with $X \in \mathbb{R}^{d \times 2d}$ as a random Gaussian matrix, with constant probability, both α and $L - \alpha$ are at least $\Omega(d)$.

For convenience, in the above algorithmic result, we reset η to zero once η goes negative (this corresponds to doing a projected gradient descent on η under constraint $\eta \ge 0$). We give a proof sketch of Theorem 4 in Section 3.1.

Surprisingly, even though we showed that the meta-gradient is well-behaved, it cannot be effectively computed by doing back-propagation due to numerical issues. More precisely:

Corollary 1. If we choose the meta-objective as $\hat{F}(\eta) = \frac{1}{t} \log f(w_{t,\eta})$, when computing the meta-gradient using back-propagation, there are intermediate results that are exponentially large/small in number of inner-steps t.

If we use back-propagation to compute $\hat{F}'(\eta)$, we need to separately compute the numerator and denominator in Eqn. (1), which are exponentially large or small as we showed in Theorem 3. Indeed, in Section 5 we empirically verify that standard auto-differentiation tools can fail in this setting. In contrast, the meta training succeeds if we use the formula derived in Section 3.1 (Eqn. (2)). This suggests that one should be more careful about using standard back-propagation in the learning-to-learn approach. The proofs of the results in this section are deferred into Appendix A.

3.1 Proof Sketch of Theorem 4

Throughout the proof, we work in the eigenspace of H which reduces the problem to having a diagonal matrix H. The proof goes in three steps:

- Claim 1 shows that the meta-objective \hat{F} has a unique minimizer η^* and the minus meta-gradient always points to the minimizer.
- Claim 2 shows meta-gradient $\hat{F}'(\eta)$ never explodes.
- Claim 3 shows meta-gradient is large when η is far from η^* .

Claim 1. The meta-objective \hat{F} has only one stationary point that is also its unique minimizer η^* . For any $\eta \in [0, \eta^*)$, $\hat{F}'(\eta) < 0$ and for any $\eta \in (\eta^*, \infty)$, $\hat{F}'(\eta) > 0$.

The lemma follows from a direct calculation $\hat{F}'(\eta)$:

$$\hat{F}'(\eta) = \frac{-2\sum_{i=1}^{d} c_i^2 \lambda_i^2 (1 - \eta \lambda_i)^{2t-1}}{\sum_{i=1}^{d} c_i^2 \lambda_i (1 - \eta \lambda_i)^{2t}}.$$
(1)

Claim 1 is proved by noticing that the denominator in $\hat{F}'(\eta)$ is always positive and the numerator is strictly increasing in η . Next, we show the meta derivative is polynomially upper bounded.

Claim 2. For any
$$\eta \in [0, \infty)$$
, we have $|\hat{F}'(\eta)| \leq \frac{4L^3}{c_{\min}^2 \alpha(L-\alpha)}$.

To prove this claim we observe that the numerator and denominator are both polynomially bounded once we divide them by a common factor, which is $(1-\eta\alpha)^{2t}$ when $\eta\in[0,\frac{2}{\alpha+L}]$. More precisely we have when $\eta\in[0,\frac{2}{\alpha+L}]$

$$\left| \hat{F}'(\eta) \right| = 2 \frac{\left| \sum_{i=1}^{d} \frac{c_i^2 \lambda_i^2}{1 - \eta \alpha} (\frac{1 - \eta \lambda_i}{1 - \eta \alpha})^{2t - 1} \right|}{c_d^2 \alpha + \sum_{i=1}^{d-1} c_i^2 \lambda_i (\frac{1 - \eta \lambda_i}{1 - \eta \alpha})^{2t}} \le \frac{2 \sum_{i=1}^{d} c_i^2 \lambda_i^2}{c_d^2 \alpha (1 - \eta \alpha)}. \tag{2}$$

This leads to the claimed bounds based on our assumptions. The case when η is large is similar. Finally, we show the meta-gradient is lower bounded if η is away from η^* and is not too large. The proof follows from a similar calculation as above.

Claim 3. Given
$$\hat{M} \geq 2/\alpha$$
 and $1/L > \epsilon > 0$, for any $\eta \in [0, \eta^* - \epsilon] \cup [\eta^* + \epsilon, \hat{M}]$, we have $|F'(\eta)| \geq 2\epsilon c_{\min}^2 \min\left(\frac{\alpha^3}{L}, \frac{1}{\hat{M}^2}\right)$.

With the above three claims, we are ready to sketch the proof of Theorem 4. Due to Claim 1, we know the minus meta-gradient always points to the minimizer η^* . This alone is not sufficient to prove the convergence result because the iterates might significantly overshoot the minimizer if $|\hat{F}'|$ is too large or the iterates might converge very slowly if $|\hat{F}'|$ is too small. Fortunately, these two problematic cases can be excluded by Claim 2 and Claim 3.

4 Generalization for Trained Optimizer

Next we consider the generalization ability of simple trained optimizers. In this section we consider a simple family of least squares problems. Let \mathcal{T} be a distribution of tasks where every task $(\mathcal{D}(w^*), S_{\text{train}}, S_{\text{valid}}, \ell)$ is determined by a parameter $w^* \in \mathbb{R}^d$ that is sampled uniformly at random from the unit sphere. For each individual task, $(x,y) \sim \mathcal{D}(w^*)$ is generated by first choosing $x \sim \mathcal{N}(0, I_d)$ and then computing $y = \langle w^*, x \rangle + \xi$ where $\xi \sim \mathcal{N}(0, \sigma^2)$ with $\sigma \geq 1$. The loss function $\ell(w, x, y)$ is just the squared loss $\ell(w, x, y) = \frac{1}{2}(y - \langle w, x \rangle)^2$. That is, the tasks are just standard least-squares problems with ground-truth equal to w^* and noise level σ^2 .

We consider two different ways to define the meta-objective.

Train-by-train: In the train-by-train setting, the training set S_{train} contains n independent samples, and the metaloss function is chosen to be the training loss. That is, in each task P, we first choose w^* uniformly at random, then generate $(x_1, y_1), ..., (x_n, y_n)$ as the training set S_{train} . The meta-loss function $\Delta_{TbT(n)}(\eta, P)$ is defined to be

$$\Delta_{TbT(n)}(\eta, P) = \frac{1}{2n} \sum_{i=1}^{n} (y_i - \langle w_{t,\eta}, x_i \rangle)^2.$$

Here $w_{t,\eta}$ is the result of running t iterations of gradient descent starting from point 0 with step size η . Note we truncate a sequence and declare the meta loss is high once the weight norm exceeds certain threshold³. We can safely do this because we assume the ground truth weight w^* has unit norm, so if the weight norm is too high, it means the inner training has diverged and the step size is too large.

As before, the empirical meta-objective in train-by-train setting is the average of the meta-loss across m different specific tasks $P_1, P_2, ..., P_m$, that is,

$$\hat{F}_{TbT(n)}(\eta) = \frac{1}{m} \sum_{k=1}^{m} \Delta_{TbT(n)}(\eta, P_k).$$
(3)

Train-by-validation: In the train-by-validation setting, the specific tasks are generated by sampling n_1 training samples and n_2 validation samples for each task, and the meta-loss function is the validation loss. That is, in each specific task P, we first choose w^* uniformly at random, then generate $(x_1, y_1), ..., (x_{n_1}, y_{n_1})$ as the training set S_{train} and $(x'_1, y'_1), ..., (x'_{n_2}, y'_{n_2})$ as the validation set S_{valid} . The meta-loss function $\Delta_{TbV(n_1, n_2)}(\eta, P)$ is defined to be

$$\Delta_{TbV(n_1,n_2)}(\eta,P) = \frac{1}{2n_2} \sum_{i=1}^{n_2} (y_i' - \langle w_{t,\eta}, x_i' \rangle)^2.$$

Here again $w_{t,\eta}$ is the result of running t iterations of the gradient descent on the training set starting from point 0, and we use the same truncation as before. The empirical meta-objective is defined as

$$\hat{F}_{TbV(n_1,n_2)}(\eta) = \frac{1}{m} \sum_{k=1}^{m} \Delta_{TbV(n_1,n_2)}(\eta, P_k), \tag{4}$$

where each P_k is independently sampled according to the described procedure.

We first show that when the number of samples is small (in particular n < d) and the noise is a large enough constant, train-by-train can be much worse than train-by-validation, even when $n_1 + n_2 = n$ (the total number of samples used in train-by-validation is the same as in train-by-train)

Theorem 5. Let $\hat{F}_{TbT(n)}(\eta)$ and $\hat{F}_{TbV(n_1,n_2)}(\eta)$ be as defined in Equation (3) and Equation (4) respectively. Assume $n, n_1, n_2 \in [d/4, 3d/4]$. Assume noise level σ is a large constant c_1 . Assume unroll length $t \geq c_2$, number of training

³Specifically, if at the τ -th step $||w_{\tau,\eta}|| \ge 40\sigma$, we freeze the training on this task and set $w_{\tau',\eta} = 40\sigma u$ for all $\tau \le \tau' \le t$, for some arbitrary vector u with unit norm. Setting the weight to a large vector is just one way to declare the loss is high.

tasks $m \ge c_3 \log(mt)$ and dimension $d \ge c_4 \log(mt)$ for certain constants c_2, c_3, c_4 . With probability at least 0.99 in the sampling of training tasks, we have

$$\eta_{train}^* = \Theta(1)$$
 and $\mathbb{E} \left\| w_{t,\eta_{train}^*} - w^* \right\|^2 = \Omega(1)\sigma^2$,

for all $\eta_{train}^* \in \arg\min_{\eta \geq 0} \hat{F}_{TbT(n)}(\eta);$

$$\eta_{\textit{valid}}^* = \Theta(1/t) \ \textit{and} \ \mathbb{E} \left\| w_{t,\eta_{\textit{valid}}^*} - w^* \right\|^2 = \left\| w^* \right\|^2 - \Omega(1)$$

for all $\eta_{valid}^* \in \arg\min_{\eta \geq 0} \hat{F}_{TbV(n_1,n_2)}(\eta)$. In both equations the expectation is taken over new tasks.

In Theorem 5, $w_{t,\eta_{\mathrm{rain}}^*}$ and $w_{t,\eta_{\mathrm{valid}}^*}$ are the results obtained on the new task and w^* is the ground truth of the new task. If σ is a large enough constant, we know $\mathbb{E} \left\| w_{t,\eta_{\mathrm{train}}^*} - w^* \right\|^2$ is larger than $\mathbb{E} \left\| w_{t,\eta_{\mathrm{valid}}^*} - w^* \right\|^2$ by some constant. The probability 0.99 is an arbitrary number, which can be replaced by any constant smaller than 1.

Note that in this case, the number of samples n is smaller than d, so the least square problem is under-determined and the optimal training loss would go to 0 (there is always a way to simultaneously satisfy all n equations). This is exactly what train-by-train would do—it will choose a large constant learning rate which guarantees the optimizer converges exponentially to the empirical risk minimizer $(ERM)^4$. However, when the noise is large making the training loss go to 0 will overfit to the noise and hurt the generalization performance. In contrast, train-by-validation will choose a smaller learning rate which allows it to leverage the signal in the training samples without overfitting to noise.

We separately give a proof sketch for the train-by-train setting and train-by-validation setting in Section 4.1 and Section 4.2, respectively. The detailed proof of Theorem 5 is deferred to Appendix B. We also prove similar results for SGD in Appendix D

We emphasize that neural networks are often over-parameterized, which corresponds to the case when d > n. Therefore in order to train neural networks, it is usually better to use train-by-validation. On the other hand, we show when the number of samples is large $(n \gg d)$, train-by-train can also perform well.

Theorem 6. Let $\hat{F}_{TbT(n)}(\eta)$ be as defined in Equation (3). Assume noise level is a constant c_1 . Given any $1 > \epsilon > 0$, assume training set size $n \ge \frac{cd}{\epsilon^2} \log(\frac{nm}{\epsilon d})$, unroll length $t \ge c_2 \log(\frac{n}{\epsilon d})$, number of training tasks $m \ge \frac{c_3 n^2}{\epsilon^4 d^2} \log(\frac{tnm}{\epsilon d})$ and dimension $d \ge c_4$ for certain constants c, c_2, c_3, c_4 . With probability at least 0.99 in the sampling of training tasks, we have

$$\mathbb{E} \left\| w_{t,\eta_{\text{train}}}^* - w^* \right\|^2 \le (1 + \epsilon) \frac{d\sigma^2}{n},$$

for all $\eta_{train}^* \in \arg\min_{\eta \geq 0} \hat{F}_{TbT(n)}(\eta)$, where the expectation is taken over new tasks.

Therefore if the learning-to-learn approach is applied to a traditional optimization problem that is not over-parameterized, train-by-train can work well. In this case, the empirical risk minimizer (ERM) already has good generalization performance, and train-by-train optimizes the convergence towards the ERM. We defer the proof of Theorem 6 into Appendix C.

4.1 Proof Sketch for Train-by-train

In this section, we will give a proof sketch for the first half of Theorem 5 (train-by-train with small number of samples). At the end of this section, we will briefly discuss the proof of Theorem 6 (train-by-train with large number of samples). For convenience, we denote \hat{F}_{TbT} as the empirical meta-objective and F_{TbT} as the population meta-objective. We implicit assume the conditions in Theorem 5 hold in the following lemmas.

Our meta-optimization problem works on a distribution of tasks. Since different tasks can have different smoothness condition, it's possible that under the same step size, the inner training converges on some tasks, but diverges on others. One way to avoid this issue is to restrict the step size into a small range under which the inner training converges on all tasks (Gupta & Roughgarden, 2017). But this is too conservative and may lead to suboptimal step

⁴In an under-determined problem, there are actually multiple ERM solutions. Here, we focus on the unique ERM solution in the span of training data. This is also the solution that GD converges to when the initialization is 0.

size. Instead, we allow any positive step size and truncate the inner training if the weight norm goes too large. This approach resolves the diverging issues and also allow the meta-learning algorithm to choose a more aggressive step size. As we explain later, this brings some technical challenges into our proof.

In order to prove $\mathbb{E} \left\| w_{t,\eta_{\text{train}}^*} - w^* \right\|^2$ is large, we only need to show the population meta-objective $F_{TbT}(\eta_{\text{train}}^*)$ is small. This is because $F_{TbT}(\eta_{\text{train}}^*)$ measures the distance between $w_{t,\eta_{\text{train}}^*}$ and the ERM solution while ERM solution is far from w^* . Since η_{train}^* minimizes the empirical meta-objective, we know $\hat{F}_{TbT}(\eta_{\text{train}}^*)$ is small. Thus we only need to show F_{TbT} and \hat{F}_{TbT} are similar. This is easy to prove for small step sizes when the inner training always converges, but is difficult when the inner training can diverge and gets truncated. To address this problem we break the step size into three intervals separated by 1/L and $\hat{\eta}$ (L is a large constant that bounds the smoothness on all tasks). Intuitively, when $\eta \leq 1/L$ almost all inner training converges and larger step size leads to faster convergence and smaller \hat{F}_{TbT} ; on the other hand, when $\eta > \tilde{\eta}$, we show $\hat{F}_{TbT}(\eta)$ is always large so the minimizer of \hat{F}_{TbT} cannot be in this region. Therefore, the optimal step size must be in $[1/L, \tilde{\eta}]$. We only need to prove in the interval $[1/L, \tilde{\eta}]$ the empirical meta-objective \hat{F}_{TbT} is close to the population meta-objective F_{TbT} . This proof is still nontrivial since the inner training can still diverge on a small fraction of sampled tasks.

We first show that for $\eta \in [0, 1/L]$, the empirical meta-objective \hat{F}_{TbT} strictly decreases as η increases and \hat{F}_{TbT} is exponentially small in t at step size 1/L.

Lemma 1. With probability at least $1 - m \exp(-\Omega(d))$, $\hat{F}_{TbT}(\eta)$ is monotonically decreasing in [0, 1/L] and

$$\hat{F}_{TbT}(1/L) \le 2L^2\sigma^2\left(1 - \frac{1}{L^2}\right)^t.$$

Next we show that the minimizer cannot be larger than $\tilde{\eta}$ for suitably chosen $\tilde{\eta}$ (see the precise definition in the appendix). Intuitively, this is because when η is too large the inner-optimizer would diverge on a significant fraction of the sampled tasks.

Lemma 2. With probability at least $1 - \exp(-\Omega(m))$,

$$\hat{F}_{TbT}(\eta) \ge \frac{\sigma^2}{10L^8}$$

for all $\eta > \tilde{\eta}$.

By Lemma 1 and Lemma 2, we know when t is large enough, the optimal step size η^*_{train} must lie in $[1/L, \tilde{\eta}]$. We can also show $1/L < \tilde{\eta} < 3/L$, so η^*_{train} is a constant. To relate the empirical loss at η^*_{train} to the population loss, we prove the following uniform convergence result when $\eta \in [1/L, \tilde{\eta}]$.

Lemma 3. With probability at least $1 - m \exp(-\Omega(d)) - O(t + m) \exp(-\Omega(m))$,

$$|F_{TbT}(\eta) - \hat{F}_{TbT}(\eta)| \le \frac{\sigma^2}{L^3},$$

for all $\eta \in [1/L, \tilde{\eta}]$.

The proof of this Lemma involves constructing special ϵ -nets for F_{TbT} and \hat{F}_{TbT} and showing that for each fixed η , $|F_{TbT}(\eta) - \hat{F}_{TbT}(\eta)|$ is small with high probability using concentration inequalities.

Combining the above lemmas, we know the population meta-objective F_{TbT} is small at η_{train}^* , which means $w_{t,\eta_{\text{train}}^*}$ is close to the ERM solution. Since the ERM solution overfits to the noise in the training samples, we know $\mathbb{E} \|w_{t,\eta_{\text{train}}^*} - w^*\|$ has to be large.

Train-by-train with large number of samples: The proof of Theorem 6 follows the same strategy as above. We prove that under the optimal step size η_{train}^* , $w_{t,\eta_{\text{train}}^*}$ converges to the ERM solution. But with more samples, the ERM solution w_{ERM} becomes closer to the ground truth w^* . More precisely, we can prove $\mathbb{E} \|w_{\text{ERM}} - w^*\|^2$ is roughly $\frac{d\sigma^2}{n}$, which leads to the bound in Theorem 6.

4.2 Proof Sketch for Train-by-Validation

In this section, we give a proof sketch for the second half of Theorem 5. We denote \hat{F}_{TbV} as the empirical meta-objective and F_{TbV} as the population meta-objective.

The overall proof strategy is similar as before: we will show the empirical meta-objective is high when the step size is beyond certain threshold, and only prove generalization result for step sizes below this threshold. Under the train-by-validation meta-objective, the optimal step size η_{valid}^* is in order $\Theta(1/t)$. So we will choose a smaller threshold step size to be 1/L.

When $\eta < 1/L$, we show that the learned signal is linear in ηt while the fitted noise is quadratic in ηt . So there exists certain step size in the order $\Theta(1/t)$ such that our model can leverage the signal in the training set without overfitting the noise. More precisely, we prove the following lemma.

Lemma 4. There exist $\eta_1, \eta_2, \eta_3 = \Theta(1/t)$ with $\eta_1 < \eta_2 < \eta_3$ such that

$$F_{TbV}(\eta_2) \le \frac{1}{2} \|w^*\|^2 - \frac{9}{10}C + \frac{\sigma^2}{2}$$

$$F_{TbV}(\eta) \ge \frac{1}{2} \|w^*\|^2 - \frac{6}{10}C + \frac{\sigma^2}{2}, \forall \eta \in [0, \eta_1] \cup [\eta_3, 1/L]$$

where C is a positive constant.

We then show whenever η is large, either the gradient descent diverges and the sequence gets truncated or it converges and overfits the noise. In both cases, the meta-objective must be high.

Lemma 5. With probability at least $1 - \exp(-\Omega(m))$,

$$\hat{F}_{TbV}(\eta) \ge C'\sigma^2 + \frac{1}{2}\sigma^2,$$

for all $\eta \geq 1/L$, where C' is a positive constant independent with σ .

To relate the behavior of F_{TbV} to the behavior of \hat{F}_{TbV} , we prove the following uniform convergence result for step sizes in [0, 1/L]. The proof is similar as in Lemma 3.

Lemma 6. With probability at least $1 - O(1/\epsilon) \exp(-\Omega(\epsilon^2 m))$,

$$|\hat{F}_{TbV}(\eta) - F_{TbV}(\eta)| \le \epsilon,$$

for all $\eta \in [0, 1/L]$.

By choosing a small enough ϵ in Lemma 6, we ensure that the behavior of \hat{F}_{TbV} is similar as that of F_{TbV} in Lemma 4. Combing with Lemma 5, we know $\eta_{\text{valid}}^* = \Theta(1/t)$ and $F_{TbV}(\eta_{\text{valid}}^*) \leq \frac{1}{2} \|w^*\|^2 + \frac{1}{2}\sigma^2 - \Omega(1)$. This concludes our proof since $F_{TbV}(\eta) = \frac{1}{2} \mathbb{E} \|w_{t,\eta} - w^*\|^2 + \frac{1}{2}\sigma^2$.

5 Experiments

In this section, we give experiment results on both synthetic data and realistic data to verify our theory.⁵

Optimizing step size for quadratic objective We first validate the results in Section 3. We fixed a 20-dimensional quadratic objective as the inner problem and vary the number of inner steps t and initial value η_0 . We compute the meta-gradient directly using the formula in Eqn. (2). In this way, we avoid the computation of exponentially small/large intermediate terms. We use the algorithm suggested in Theorem 4, except we choose the meta-step size to be $1/(100\sqrt{k})$ as the constants in the theorem were not optimized.

⁵Our code is available at https://github.com/Kolin96/learning-to-learn.





Figure 1: Meta training trajectory for η ($t = 80, \eta_0 = 0.1$).

An example training curve of η for t=80 and $\eta_0=0.1$ is shown in Figure 1, and we can see that η converges quickly within 300 steps. Similar convergence also holds for larger t or larger initial η_0 . In contrast, we also implemented the meta-training with Tensorflow, where the code was adapted from the previous work of Wichrowska et al. (2017). Experiments show that in many settings (especially with large t and large t are the initial value throughout the meta training because the meta-gradient explodes and gives NaN value. More details can be found in Appendix F.

Train-by-train vs. train-by-validation, synthetic data Here we validate our theoretical results in Section 4 using the least-squares model defined there. We fix the input dimension d to be 1000.

In the first experiment, we fix the size of the data (n=500 for train-by-train, $n_1=n_2=250$ for train-by-validation). Under different noise levels, we find the optimal η^* by a grid search on its meta-objective for train-by-train and train-by-validation settings respectively. We then use the optimal η^* found in each of these two settings to test on 10 new least-squares problem. The mean RMSE, as well as its range over the 10 test cases, are shown in Figure 2. We can see that for all of these cases, the train-by-train model overfits easily, while the train-by-validation model performs much better and does not overfit. Also, when the noise becomes larger, the difference between these two settings becomes more significant.







Figure 2: Training and testing RMSE for different σ values (500 samples)

In the next experiment, we fix $\sigma=1$ and change the sample size. For train-by-validation, we always split the samples evenly into training and validation set. From Figure 3, we can see that the gap between these two settings is decreasing as we use more data, as expected by Theorem 6.

Train-by-train vs. train-by-validation, MLP optimizer on MNIST Here we consider the more interesting case of a multi-layer perceptron (MLP) optimizer on MNIST data set. We use the same MLP optimizer as in Metz et al. (2019), and details of this optimizer is discussed in Appendix F. As the inner problem, we use a two-layer fully-connected network of 100 and 20 hidden units with ReLU activations. The inner objective is the classic 10-class cross entropy loss, and we use mini-batches of 32 samples at inner training. In all the following experiments, we use SGD as a






Figure 3: Training and testing RMSE for different samples sizes ($\sigma = 1$)

baseline with step size tuned by grid search against validation loss. For each optimizer, we run 5 independent tests and collect training accuracy and test accuracy for evaluation. The plots show the mean of the 5 tests⁶.




Figure 4: The test accuracy of different optimizers in various settings. Comparison between (a) and (b) shows that the advantage of train-by-validation over train-by-train increases when the samples have more noise; comparison between (a) and (c) shows that when the number of samples increases, train-by-train gets comparable performance as train-by-validation.

In Figure 4, we show the test accuracy for different optimizers for different sample size and noise level. In this figure, "TbTx" represents train-by-train approach with x training samples; "TbVx + y" represents train-by-validation approach with x training samples and y validation samples. In Figure 4(a) the optimizer is applied to 1000 randomly sub-sampled data (split between training and validation for train-by-validation); in Figure 4(b) we use the same amount of data, except we add 20% label noise; in Figure 4(c) we use the whole MNIST dataset without label noise. Comparing

⁶We didn't show the measure of the spread because the results of these 5 tests are so close to each other, such that the range or standard deviation marks will not be readable in the plots.

Figure 4(a) and (b), we see that when the noise is large train-by-validation significantly outperforms train-by-train. Figure 5 gives the training accuracy in the same setting as Figure 4(b), which clearly shows that train-by-validation can avoid overfitting to noisy labels. Comparing Figure 4(a) and (c), we see that when the number of samples is large enough there is no significant difference between train-by-train and train-by-validation.




Figure 5: Training accuracy for 1000 samples and 20% noise (same setting as in Figure 4(b))

6 Conclusions

In this paper, we have proved optimization and generalization guarantees for tuning the step size for quadratic loss. From the optimization perspective, we considered a simple task whose objective is a quadratic function. We proved that the meta-gradient can explode/vanish if the meta-objective is simply the loss of the last iteration; we then showed that the log-transformed meta-objective has polynomially bounded meta-gradient and can be successfully optimized. To study the generalization issues, we considered the least squares problem—when the number of samples is small and the noise is large, train-by-validation approach generalizes better than train-by-train; while when the number of samples is large, train-by-train can also work well.

Although our theoretical results are proved for quadratic loss, this simple setting already yields interesting phenomenons and requires non-trivial techniques to analyze. We have also verified our theoretical results on an optimizer parameterized by neural networks and on MNIST dataset. There are still many open problems, including extending similar analysis to more complicated optimizers, or generalizing the idea to prevent numerical issues to neural network optimizers. We hope our work can lead to more theoretical understanding of the learning-to-learn approach.

Acknowledgements

Rong Ge, Xiang Wang and Chenwei Wu are supported in part by NSF Award CCF-1704656, CCF-1845171 (CA-REER), CCF-1934964 (Tripods), a Sloan Research Fellowship, and a Google Faculty Research Award. Part of the work was done when Rong Ge and Xiang Wang were visiting Instituted for Advanced Studies for "Special Year on Optimization, Statistics, and Theoretical Machine Learning" program. We acknowledge the valuable early discussions with Yatharth Dubey.

References

Abadi, M., Barham, P., Chen, J., Chen, Z., Davis, A., Dean, J., Devin, M., Ghemawat, S., Irving, G., Isard, M., et al. Tensorflow: A system for large-scale machine learning. In *12th USENIX Symposium on Operating Systems Design and Implementation (OSDI 16)*, pp. 265–283, 2016.

Alabi, D., Kalai, A. T., Ligett, K., Musco, C., Tzamos, C., and Vitercik, E. Learning to prune: Speeding up repeated computations. *arXiv preprint arXiv:1904.11875*, 2019.

- Andrychowicz, M., Denil, M., Gomez, S., Hoffman, M. W., Pfau, D., Schaul, T., Shillingford, B., and De Freitas, N. Learning to learn by gradient descent by gradient descent. In *Advances in neural information processing systems*, pp. 3981–3989, 2016.
- Bai, Y., Chen, M., Zhou, P., Zhao, T., Lee, J. D., Kakade, S., Wang, H., and Xiong, C. How important is the train-validation split in meta-learning? *arXiv preprint arXiv:2010.05843*, 2020.
- Balcan, M.-F., Nagarajan, V., Vitercik, E., and White, C. Learning-theoretic foundations of algorithm configuration for combinatorial partitioning problems. *arXiv preprint arXiv:1611.04535*, 2016a.
- Balcan, M.-F., Sandholm, T., and Vitercik, E. Sample complexity of automated mechanism design. *arXiv preprint* arXiv:1606.04145, 2016b.
- Balcan, M.-F., Dick, T., Sandholm, T., and Vitercik, E. Learning to branch. arXiv preprint arXiv:1803.10150, 2018a.
- Balcan, M.-F., Sandholm, T., and Vitercik, E. A general theory of sample complexity for multi-item profit maximization. In *Proceedings of the 2018 ACM Conference on Economics and Computation*, pp. 173–174, 2018b.
- Bello, I., Zoph, B., Vasudevan, V., and Le, Q. V. Neural optimizer search with reinforcement learning. In *Proceedings* of the 34th International Conference on Machine Learning-Volume 70, pp. 459–468. JMLR. org, 2017.
- Bengio, S., Bengio, Y., Cloutier, J., and Gecsei, J. On the optimization of a synaptic learning rule. In *Preprints Conf. Optimality in Artificial and Biological Neural Networks*, volume 2, 1992.
- Bengio, Y., Bengio, S., and Cloutier, J. Learning a synaptic learning rule. Citeseer, 1990.
- Franceschi, L., Donini, M., Frasconi, P., and Pontil, M. Forward and reverse gradient-based hyperparameter optimization. *arXiv preprint arXiv:1703.01785*, 2017.
- Ge, R., Kakade, S. M., Kidambi, R., and Netrapalli, P. The step decay schedule: A near optimal, geometrically decaying learning rate procedure for least squares. In *Advances in Neural Information Processing Systems*, pp. 14951–14962, 2019.
- Gupta, R. and Roughgarden, T. A pac approach to application-specific algorithm selection. *SIAM Journal on Computing*, 46(3):992–1017, 2017.
- Harvey, N. J., Liaw, C., Plan, Y., and Randhawa, S. Tight analyses for non-smooth stochastic gradient descent. *arXiv* preprint arXiv:1812.05217, 2018.
- Hochreiter, S., Younger, A. S., and Conwell, P. R. Learning to learn using gradient descent. In *International Conference on Artificial Neural Networks*, pp. 87–94. Springer, 2001.
- Jain, P., Nagaraj, D., and Netrapalli, P. Making the last iterate of sgd information theoretically optimal. *arXiv* preprint arXiv:1904.12443, 2019.
- Johnson, W. B. and Lindenstrauss, J. Extensions of lipschitz mappings into a hilbert space. *Contemporary mathematics*, 26(189-206):1, 1984.
- Li, K. and Malik, J. Learning to optimize. arXiv preprint arXiv:1606.01885, 2016.
- Li, K. and Malik, J. Learning to optimize neural nets. arXiv preprint arXiv:1703.00441, 2017.
- Lv, K., Jiang, S., and Li, J. Learning gradient descent: Better generalization and longer horizons. In *International Conference on Machine Learning*, pp. 2247–2255. PMLR, 2017.
- Maclaurin, D., Duvenaud, D., and Adams, R. Gradient-based hyperparameter optimization through reversible learning. In *International Conference on Machine Learning*, pp. 2113–2122, 2015.

- Metz, L., Maheswaranathan, N., Nixon, J., Freeman, D., and Sohl-Dickstein, J. Understanding and correcting pathologies in the training of learned optimizers. In *International Conference on Machine Learning*, pp. 4556–4565, 2019.
- Morgenstern, J. and Roughgarden, T. Learning simple auctions. In *Conference on Learning Theory*, pp. 1298–1318. PMLR, 2016.
- Morgenstern, J. H. and Roughgarden, T. On the pseudo-dimension of nearly optimal auctions advances in neural information processing systems. 136–144. *Google Scholar Google Scholar Digital Library Digital Library*, 2015.
- Shamir, O. and Zhang, T. Stochastic gradient descent for non-smooth optimization: Convergence results and optimal averaging schemes. In *International conference on machine learning*, pp. 71–79, 2013.
- Vershynin, R. Introduction to the non-asymptotic analysis of random matrices. arXiv preprint arXiv:1011.3027, 2010.
- Vershynin, R. *High-dimensional probability: An introduction with applications in data science*, volume 47. Cambridge university press, 2018.
- Wichrowska, O., Maheswaranathan, N., Hoffman, M. W., Colmenarejo, S. G., Denil, M., de Freitas, N., and Sohl-Dickstein, J. Learned optimizers that scale and generalize. In *Proceedings of the 34th International Conference on Machine Learning-Volume 70*, pp. 3751–3760. JMLR. org, 2017.

In the appendix, we first give the missing proofs for the theorems in the main paper. Later in Appendix F we give details for the experiments.

Notations: Besides the notations defined in Section 2, we define more notations that will be used in the proofs.

For a matrix $X \in \mathbb{R}^{n \times d}$ with $n \leq d$, we denote its singular values as $\sigma_1(X) \geq \cdots \geq \sigma_n(X)$.

For a positive semi-definite matrix $A \in \mathbb{R}^{d \times d}$, we denote $u^{\top}Au$ as $\|u\|_A^2$. For a matrix $X \in \mathbb{R}^{d \times n}$, let $\operatorname{Proj}_X \in \mathbb{R}^{d \times d}$ be the projection matrix onto the column span of X. That means, $\operatorname{Proj}_X = SS^{\top}$, where the columns of S form an orthonormal basis for the column span of X.

For any event \mathcal{E} , we use $\mathbb{1}\{\mathcal{E}\}$ to denote its indicator function: $\mathbb{1}\{\mathcal{E}\}$ equals 1 when \mathcal{E} holds and equals 0 otherwise. We use $\bar{\mathcal{E}}$ to denote the complementary event of \mathcal{E} .

A Proofs for Section 3 – alleviating gradient explosion/vanishing problem for quadratic objective

In this section, we prove the results in Section 3. Recall the meta learning problem as follows:

The inner task is a fixed quadratic problem, where the starting point is fixed at w_0 , and the loss function is $f(w) = \frac{1}{2} w^\top H w$ for some fixed positive definite matrix $H \in \mathbb{R}^{d \times d}$. Suppose the eigenvalue decomposition of H is $\sum_{i=1}^d \lambda_i u_i u_i^\top$. In this section, we assume $L = \lambda_1(H)$ and $\alpha = \lambda_d(H)$ are the largest and smallest eigenvalues of H with $L > \alpha$. We assume the starting point w_0 has unit ℓ_2 norm. For each $i \in [d]$, let c_i be $\langle w_0, u_i \rangle$ and let $c_{\min} = \min(|c_1|, |c_d|)$. We assume $c_{\min} > 0$ for simplicity, which is satisfied if w_0 is chosen randomly from the unit sphere.

Let $\{w_{\tau,\eta}\}$ be the GD sequence running on f(w) starting from w_0 with step size η . For the meta-objective, we consider using the loss of the last point directly, or using the log of this value. In Section A.1, we first show that although choosing $\hat{F}(\eta) = f(w_{t,\eta})$ does not have any bad local optimal solution, it has the gradient explosion/vanishing problem (Theorem 3). Then, in Section A.2, we show choosing $\hat{F}(\eta) = \frac{1}{t}\log f(w_{t,\eta})$ leads to polynomially bounded meta-gradient and further show meta-gradient descent converges to the optimal step size (Theorem 4). Although the meta-gradient is polynomially bounded, if we simply use back-propogation to compute the meta-gradient, the intermediate results can still be exponentially large/small (Corollary 1). This is also proved in Section A.2.

A.1 Meta-gradient vanishing/explosion

In this section, we show although choosing $\hat{F}(\eta) = f(w_{t,\eta})$ does not have any bad local optimal solution, it has the meta-gradient explosion/vanishing problem. Recall Theorem 3 as follows.

Theorem 3. Let the meta-objective be $\hat{F}(\eta) = f(w_{t,\eta})$, we know $\hat{F}(\eta)$ is a strictly convex function in η with an unique minimizer. However, for any step size $0 < \eta < 2/L$,

$$|\hat{F}'(\eta)| \le tL^2 \max(|1 - \eta \alpha|^{2t-1}, |1 - \eta L|^{2t-1});$$

for any step size $\eta > 2/L$,

$$|\hat{F}'(\eta)| \ge c_1^2 L^2 t (\eta L - 1)^{2t - 1} - L^2 t.$$

Intuitively, if we write $w_{t,\eta}$ in the basis of the eigen-decomposition of H, then each coordinate evolve exponentially in t. The gradient of the standard objective is therefore also exponential in t.

Proof of Theorem 3. According to the gradient descent iterations, we have

$$w_{t,\eta} = w_{t-1,\eta} - \eta \nabla f(w_{t-1,\eta}) = w_{t-1,\eta} - \eta H w_{t-1,\eta} = (I - \eta H) w_{t-1,\eta} = (I - \eta H)^t w_0.$$

Therefore, $\hat{F}(\eta) := f(w_{t,\eta}) = \frac{1}{2} w_0^\top (I - \eta H)^{2t} H w_0$. Taking the derivative of $\hat{F}(\eta)$,

$$\hat{F}'(\eta) = -tw_0^{\top} (I - \eta H)^{2t-1} H^2 w_0 = -t \sum_{i=1}^d c_i^2 \lambda_i^2 (1 - \eta \lambda_i)^{2t-1},$$

where $c_i = \langle w_0, u_i \rangle$. Taking the second derivative of $F(\eta)$,

$$F''(\eta) = t(2t-1)w_0^{\top}(I-\eta H)^{2t-2}H^3w_0 = t(2t-1)\sum_{i=1}^d c_i^2 \lambda_i^3 (1-\eta \lambda_i)^{2t-2}.$$

Since $L > \alpha$, we have $\hat{F}''(\eta) > 0$ for any η . That means $\hat{F}(\eta)$ is a strictly convex function in η with a unique minimizer.

For any fixed $\eta \in (0, 2/L)$ we know $|1 - \eta \lambda_i| < 1$ for all $i \in [d]$. We have

$$\begin{split} \left| \hat{F}'(\eta) \right| &\leq t \sum_{i=1}^{d} c_i^2 \lambda_i^2 |1 - \eta \lambda_i|^{2t-1} \\ &\leq t \sum_{i=1}^{d} c_i^2 \max_{i \in [d]} \left(\lambda_i^2 |1 - \eta \lambda_i|^{2t-1} \right) \\ &\leq t L^2 \max \left(|1 - \eta \alpha|^{2t-1}, |1 - \eta L|^{2t-1} \right), \end{split}$$

where the last inequality uses $\sum_{i=1}^{d} c_i^2 = 1$. Note for $\eta \in (0, 2/L)$, it's guaranteed that $|1 - \eta \lambda_i|$ takes the maximum at $|1 - \eta \alpha|$ or $|1 - \eta L|$.

For any fixed $\eta \in (2/L, \infty)$, we know $\eta L - 1 > 1$. We have

$$\begin{split} \hat{F}'(\eta) &= -tc_1^2L^2(1-\eta L)^{2t-1} - t\sum_{i\neq 1: (1-\eta\lambda_i)\leq 0} c_i^2\lambda_i^2(1-\eta\lambda_i)^{2t-1} - t\sum_{i\neq 1: (1-\eta\lambda_i)> 0} c_i^2\lambda_i^2(1-\eta\lambda_i)^{2t-1} \\ &\geq tc_1^2L^2(\eta L-1)^{2t-1} - t\sum_{i=1}^d c_i^2\lambda_i^2 \geq tc_1^2L^2(\eta L-1)^{2t-1} - L^2t, \end{split}$$

where the last inequality uses $\sum_{i=1}^{d} c_i^2 = 1$.

A.2 Alleviating meta-gradient vanishing/explosion

We prove when the meta objective is chosen as $\frac{1}{t} \log f(w_{t,\eta})$, the meta-gradient is polynomially bounded. Furthermore, we show meta-gradient descent can converge to the optimal step size within polynomial iterations. Recall Theorem 4 as follows.

Theorem 4. Let the meta-objective be $\hat{F}(\eta) = \frac{1}{t} \log f(w_{t,\eta})$. We know $\hat{F}(\eta)$ has a unique minimizer η^* and $\hat{F}'(\eta) = O\left(\frac{L^3}{c_{\min}^2\alpha(L-\alpha)}\right)$ for all $\eta \geq 0$. Let $\{\eta_k\}$ be the GD sequence running on \hat{F} with meta step size $\mu_k = 1/\sqrt{k}$. Suppose the starting step size $\eta_0 \leq M$. Given any $1/L > \epsilon > 0$, there exists $k' = \frac{M^6}{\epsilon^2} poly(\frac{1}{c_{\min}}, L, \frac{1}{\alpha}, \frac{1}{L-\alpha})$ such that for all $k \geq k', |\eta_k - \eta^*| \leq \epsilon$.

When we take the log of the function value, the derivative of the function value with respect to η becomes much more stable. We will first show some structural result on $\hat{F}(\eta)$ – it has a unquie minimizer and the gradient is polynomially bounded. Further the gradient is only close to 0 when the point η is close to the unique minimizer. Then using such structural result we prove that meta-gradient descent converges.

Proof of Theorem 4. The proof consists of three claims. In the first claim, we show that \hat{F} has a unique minimizer and the minus meta derivative always points to the minimizer. In the second claim, we show that \hat{F} has bounded derivative. In the last claim, we show that for any η that is outside the ϵ -neighborhood of η^* , $|\hat{F}'(\eta)|$ is lower bounded. Finally, we combine these three claims to finish the proof.

Claim 4. The meta objective \hat{F} has only one stationary point that is also its unique minimizer η^* . For any $\eta \in [0, \eta^*)$, $\hat{F}'(\eta) < 0$ and for any $\eta \in (\eta^*, \infty)$, $\hat{F}'(\eta) > 0$. Furthermore, we know $\eta^* \in [1/L, 1/\alpha]$.

We can compute the derivative of \hat{F} in η as follows,

$$\hat{F}'(\eta) = \frac{-2w_0^\top (I - \eta H)^{2t-1} H^2 w_0}{w_0^\top (I - \eta H)^{2t} H w_0} = \frac{-2\sum_{i=1}^d c_i^2 \lambda_i^2 (1 - \eta \lambda_i)^{2t-1}}{\sum_{i=1}^d c_i^2 \lambda_i (1 - \eta \lambda_i)^{2t}}.$$
 (5)

It's not hard to verify that the denominator $\sum_{i=1}^d c_i^2 \lambda_i (1-\eta\lambda_i)^{2t}$ is always positive. Denote the numerator $-2\sum_{i=1}^d c_i^2 \lambda_i^2 (1-\eta\lambda_i)^{2t-1}$ as $g(\eta)$. Since $g'(\eta)>0$ for any $\eta\in[0,\infty)$, we know $g(\eta)$ is strictly increasing in η . Combing with the fact that g(0) < 0 and $g(\infty) > 0$, we know there is a unique point (denoted as η^*) where $g(\eta^*) = 0$ and $g(\eta) < 0$ for all $\eta \in [0, \eta^*)$ and $g(\eta) > 0$ for all $\eta \in (\eta^*, \infty)$. Since the denominator in $\hat{F}'(\eta)$ is always positive and the numerator equals $g(\eta)$, we know there is a unique point η^* where $\hat{F}'(\eta^*) = 0$ and $\hat{F}'(\eta) < 0$ for all $\eta \in [0, \eta^*)$ and $\hat{F}'(\eta) > 0$ for all $\eta \in (\eta^*, \infty)$. It's clear that η^* is the minimizer of \hat{F} .

Also, it's not hard to verify that for any $\eta \in [0, 1/L)$, $\hat{F}'(\eta) < 0$ and for any $\eta \in (1/\alpha, \infty)$, $\hat{F}'(\eta) > 0$. This implies that $\eta^* \in [1/L, 1/\alpha]$.

Claim 5. For any $\eta \in [0, \infty)$, we have

$$|\hat{F}'(\eta)| \le \frac{4L^3}{c_{\min}^2 \alpha (L - \alpha)} := D_{\max}.$$

For any $\eta \in [0, \frac{2}{\alpha + L}]$, we have $|1 - \eta \lambda_i| \le 1 - \eta \alpha$ for all *i*. Dividing the numerator and denominator in $\hat{F}'(\eta)$ by $(1-\eta\alpha)^{2t}$, we have

$$\left| \hat{F}'(\eta) \right| = 2 \frac{\left| \sum_{i=1}^{d} \frac{c_i^2 \lambda_i^2}{1 - \eta \alpha} (\frac{1 - \eta \lambda_i}{1 - \eta \alpha})^{2t - 1} \right|}{c_d^2 \alpha + \sum_{i=1}^{d-1} c_i^2 \lambda_i (\frac{1 - \eta \lambda_i}{1 - \eta \alpha})^{2t}} \le \frac{2 \sum_{i=1}^{d} c_i^2 \lambda_i^2}{c_d^2 \alpha (1 - \eta \alpha)} \le \frac{2(\alpha + L) \sum_{i=1}^{d} c_i^2 \lambda_i^2}{c_d^2 \alpha (L - \alpha)} \le \frac{4L^3}{c_d^2 \alpha (L - \alpha)},$$

where the second last inequality uses $\eta \leq \frac{2}{\alpha + L}$. Similarly for any $\eta \in (\frac{2}{\alpha + L}, \infty)$, we have $|1 - \eta \lambda_i| \leq \eta L - 1$ for all i. Dividing the numerator and denominator in $\hat{F}'(\eta)$ by $(\eta L - 1)^{2t}$, we have

$$\hat{F}'(\eta) = 2 \frac{\left| \sum_{i=1}^{d} \frac{c_i^2 \lambda_i^2}{\eta L - 1} (\frac{1 - \eta \lambda_i}{\eta L - 1})^{2t - 1} \right|}{c_1^2 L + \sum_{i=2}^{d} c_i^2 \lambda_i (\frac{1 - \eta \lambda_i}{\eta L - 1})^{2t}} \le \frac{2 \sum_{i=1}^{d} c_i^2 \lambda_i^2}{c_1^2 L (\eta L - 1)} \le \frac{2(\alpha + L) \sum_{i=1}^{d} c_i^2 \lambda_i^2}{c_1^2 L (L - \alpha)} \le \frac{4L^3}{c_1^2 L (L - \alpha)}$$

where the last inequality uses $\eta \ge \frac{2}{\alpha + L}$. Overall, we know for any $\eta \ge 0$,

$$|\hat{F}'(\eta)| \le \frac{4L^3}{L - \alpha} \max\left(\frac{1}{c_d^2 \alpha}, \frac{1}{c_1^2 L}\right) \le \frac{4L^3}{c_{\min}^2 \alpha (L - \alpha)}.$$

Claim 6. Given $\hat{M} \geq 2/\alpha$ and $1/L > \epsilon > 0$, for any $\eta \in [0, \eta^* - \epsilon] \cup [\eta^* + \epsilon, \hat{M}]$, we have

$$|F'(\eta)| \ge \min\left(\frac{2\epsilon c_d^2 \alpha^3}{L}, \frac{2\epsilon c_1^2 L^2}{(\hat{M}L - 1)^2}\right) \ge 2\epsilon c_{\min}^2 \min\left(\frac{\alpha^3}{L}, \frac{1}{\hat{M}^2}\right) := D_{\min}(\hat{M}).$$

If $\eta \in [0, \eta^* - \epsilon]$ and $\eta \leq \frac{2}{\alpha + L}$, we have

$$\hat{F}'(\eta) = -2 \frac{\sum_{i=1}^{d} c_i^2 \lambda_i^2 (1 - \eta \lambda_i)^{2t-1}}{\sum_{i=1}^{d} c_i^2 \lambda_i (1 - \eta \lambda_i)^{2t}} = -2 \frac{\sum_{i=1}^{d} c_i^2 \lambda_i^2 (1 - \eta \lambda_i)^{2t-1} - \sum_{i=1}^{d} c_i^2 \lambda_i^2 (1 - \eta^* \lambda_i)^{2t-1}}{\sum_{i=1}^{d} c_i^2 \lambda_i (1 - \eta \lambda_i)^{2t}},$$

where the second equality holds because $\sum_{i=1}^d c_i^2 \lambda_i^2 (1-\eta^*\lambda_i)^{2t-1} = 0$. For the numerator, we have

$$\sum_{i=1}^{d} c_i^2 \lambda_i^2 (1 - \eta \lambda_i)^{2t-1} - \sum_{i=1}^{d} c_i^2 \lambda_i^2 (1 - \eta^* \lambda_i)^{2t-1} \ge c_d^2 \alpha^2 \left((1 - \eta \alpha)^{2t-1} - (1 - \eta^* \alpha)^{2t-1} \right) \\ \ge c_d^2 \alpha^2 \left((1 - \eta \alpha)^{2t-1} - (1 - \eta \alpha - \epsilon \alpha)^{2t-1} \right);$$

for the denominator, we have

$$\sum_{i=1}^{d} c_i^2 \lambda_i (1 - \eta \lambda_i)^{2t} \le \left(\sum_{i=1}^{d} c_i^2 \lambda_i\right) (1 - \eta \alpha)^{2t},$$

where the second inequality holds because $|1 - \eta \lambda_i| \le 1 - \eta \alpha$ for all i. Overall, we have when $\eta \in [0, \eta^* - \epsilon]$ and $\eta \le \frac{2}{\alpha + L}$,

$$\begin{split} \left| \hat{F}'(\eta) \right| \geq & 2 \frac{c_d^2 \alpha^2 \left((1 - \eta \alpha)^{2t - 1} - (1 - \eta \alpha - \epsilon \alpha)^{2t - 1} \right)}{\left(\sum_{i=1}^d c_i^2 \lambda_i \right) (1 - \eta \alpha)^{2t}} \\ \geq & \frac{2\epsilon c_d^2 \alpha^3}{\left(\sum_{i=1}^d c_i^2 \lambda_i \right) (1 - \eta \alpha)} \geq \frac{2\epsilon c_d^2 \alpha^3}{L}, \end{split}$$

where the last inequality holds because $(1-\eta\alpha)\leq 1$ and $\sum_i^d c_i^2\lambda_i\leq L$. Similarly, if $\eta\in[0,\eta^*-\epsilon]$ and $\eta\geq\frac{2}{\alpha+L}$, we have

$$\begin{split} \left| \hat{F}'(\eta) \right| &\geq & 2 \frac{c_1^2 L^2 \left((1 - \eta L)^{2t - 1} - (1 - \eta L - \epsilon L)^{2t - 1} \right)}{\left(\sum_{i = 1}^d c_i^2 \lambda_i \right) (1 - \eta L)^{2t}} \\ &= & 2 \frac{c_1^2 L^2 \left((\eta L + \epsilon L - 1)^{2t - 1} - (\eta L - 1)^{2t - 1} \right)}{\left(\sum_{i = 1}^d c_i^2 \lambda_i \right) (\eta L - 1)^{2t}} \\ &\geq & \frac{2\epsilon c_1^2 L^3}{\left(\sum_{i = 1}^d c_i^2 \lambda_i \right) (\eta L - 1)^2} \geq \frac{2\epsilon c_1^2 \alpha^2 L^2}{(L - \alpha)^2}, \end{split}$$

where the last inequality holds because $\eta \leq \eta^* - \epsilon \leq 1/\alpha$ and $\sum_i^d c_i^2 \lambda_i \leq L$. If $\eta \in [\eta^* + \epsilon, \infty)$ and $\eta \leq \frac{2}{\alpha + L}$, we have

$$\begin{aligned} \left| \hat{F}'(\eta) \right| &\geq 2 \frac{c_d^2 \alpha^2 \left((1 - \eta \alpha + \epsilon \alpha)^{2t - 1} - (1 - \eta \alpha)^{2t - 1} \right)}{\left(\sum_{i=1}^d c_i^2 \lambda_i \right) (1 - \eta \alpha)^{2t}} \\ &\geq \frac{2\epsilon c_d^2 \alpha^3}{L}, \end{aligned}$$

If $\eta \in [\eta^* + \epsilon, \infty)$ and $\eta \ge \frac{2}{\alpha + L}$, we have

$$\begin{split} \left| \hat{F}'(\eta) \right| \geq & 2 \frac{c_1^2 L^2 \left((1 - \eta L + \eta \epsilon)^{2t - 1} - (1 - \eta L)^{2t - 1} \right)}{\left(\sum_{i=1}^d c_i^2 \lambda_i \right) (1 - \eta L)^{2t}} \\ \geq & \frac{2 \epsilon c_1^2 L^3}{\left(\sum_{i=1}^d c_i^2 \lambda_i \right) (\eta L - 1)^2} \geq \frac{2 \epsilon c_1^2 L^2}{(\hat{M} L - 1)^2}, \end{split}$$

where the last inequality uses the assumption that $\eta \leq \hat{M}$.

With the above three claims, we are ready to prove the optimization result. By Claim 4, we know $F'(\eta) < 0$ for any $\eta \in [0, \eta^*)$ and $F'(\eta) > 0$ for any $\eta \in (\eta^*, \infty)$. So the opposite gradient descent always points to the minimizer.

Since $\mu_k = 1/\sqrt{k}$, when $k \ge k_1 := \frac{D_{\max}^2}{\epsilon^2}$ we know $\mu_k \le \frac{\epsilon}{D_{\max}}$. By Claim 5, we know $|\hat{F}'(\eta)| \le D_{\max}$ for all $\eta \ge 0$, which implies $|\mu_k \hat{F}'(\eta)| \le \epsilon$ for all $k \ge k_1$. That means, meta gradient descent will never overshoot the minimizer by more than ϵ when $k \ge k_1$. In other words, after k_1 meta iterations, once η enters the ϵ -neighborhood of η^* , it will never leave this neighborhood.

We also know that at meta iteration k_1 , we have $\eta_{k_1} \leq \max(1/\alpha + D_{\max}, M) := \hat{M}$. Here, $1/\alpha + D_{\max}$ comes from the case that the eta starts from the left of η^* and overshoot to the right of η^* by D_{\max} . Since $\eta^* \in [1/L, 1/\alpha]$, we have $|\eta_{k_1} - \eta^*| \leq \max(1/\alpha, 1/\alpha + D_{\max} - 1/L, M - 1/L) := R$. By Claim 6, we know that $|\hat{F}'(\eta)| \geq D_{\min}(\hat{M})$ for any $\eta \in [0, \eta^* - \epsilon] \cup [\eta^* + \epsilon, \hat{M}]$. Choosing some k_2 satisfying $\sum_{k=k_1}^{k_2} 1/\sqrt{k} \geq \frac{R}{D_{\min}}$, we know for any $k \geq k_2$, $|\eta_k - \eta^*| \leq \epsilon$. Plugging in all the bounds for D_{\min}, D_{\max} from Claim 6 and Claim 5, we know there exists $k_1 = \frac{1}{\epsilon^2} \mathrm{poly}(\frac{1}{c_{\min}}, L, \frac{1}{\lambda}, \frac{1}{L-\alpha})$, $k_2 = \frac{M^6}{\epsilon^2} \mathrm{poly}(\frac{1}{c_{\min}}, L, \frac{1}{\alpha}, \frac{1}{L-\alpha})$ satisfying these conditions. \square Next, we show although the meta-gradient is polynomially bounded, the intermediate results can still vanish or

Next, we show although the meta-gradient is polynomailly bounded, the intermediate results can still vanish or explode if we use back-propagation to compute the meta-gradient.

Corollary 1. If we choose the meta-objective as $\hat{F}(\eta) = \frac{1}{t} \log f(w_{t,\eta})$, when computing the meta-gradient using back-propagation, there are intermediate results that are exponentially large/small in number of inner-steps t.

Proof of Corollary 1. This is done by direct calculation. If we use back-propagation to compute the derivative of $\frac{1}{t}\log(f(w_{t,\eta}))$, we need to first compute $\frac{\partial f(w_{t,\eta})}{\partial t}\frac{1}{t}\log(f(w_{t,\eta}))$ that equals $\frac{1}{tf(w_{t,\eta})}$. Same as the analysis in Theorem 3, we can show $\frac{1}{tf(w_{t,\eta})}$ is exponentially large when $\eta < 2/L$ and is exponentially small when $\eta > 2/L$. \square

B Proofs of train-by-train v.s. train-by-validation (GD)

In this section, we show when the number of samples is small and when the noise level is a large constant, train-by-train overfits to the noise in training tasks while train-by-validation generalizes well. We separately prove the results for train-by-train and train-by-validation in Theorem 7 and Theorem 8, respectively. Then, Theorem 5 is simply a combination of Theorem 7 and Theorem 8.

Recall that in the train-by-train setting, each task P contains a training set S_{train} with n samples. The inner objective is defined as $\hat{f}(w) = \frac{1}{2n} \sum_{(x,y) \in S_{\text{train}}} (\langle w, x \rangle - y)^2$. Let $\{w_{\tau,\eta}\}$ be the GD sequence running on $\hat{f}(w)$ from initialization 0 (with truncation). The meta-loss on task P is defined as the inner objective of the last point, $\Delta_{TbT(n)}(\eta,P) = \hat{f}(w_{t,\eta}) = \frac{1}{2n} \sum_{(x,y) \in S_{\text{train}}} (\langle w_{t,\eta}, x \rangle - y)^2$. The empirical meta objective $\hat{F}_{TbT(n)}(\eta)$ is the average of the meta-loss across m different tasks. We show that under $\hat{F}_{TbT(n)}(\eta)$, the optimal step size is a constant and the learned weight is far from ground truth w^* on new tasks. We prove Theorem 7 in Section B.2.

Theorem 7. Let the meta objective $\hat{F}_{TbT(n)}(\eta)$ be as defined in Equation 3 with $n \in [d/4, 3d/4]$. Assume noise level σ is a large constant c_1 . Assume unroll length $t \ge c_2$, number of training tasks $m \ge c_3 \log(mt)$ and dimension $d \ge c_4 \log(m)$ for certain constants c_2, c_3, c_4 . With probability at least 0.99 in the sampling of the training tasks, we have

$$\eta_{train}^* = \Theta(1)$$
 and $\mathbb{E} \left\| w_{t,\eta_{min}^*} - w^* \right\|^2 = \Omega(1)\sigma^2$,

for all $\eta_{train}^* \in \arg\min_{n>0} \hat{F}_{TbT(n)}(\eta)$, where the expectation is taken over new tasks.

In Theorem 7, $\Omega(1)$ is an absolute constant independent with σ . Intuitively, the reason that train-by-train performs badly in this setting is because there is a way to set the step size to a constant such that gradient descent converges very quickly to the empirical risk minimizer, therefore making the train-by-train objective very small. However, when the noise is large and the number of samples is smaller than the dimension, the empirical risk minimizer (ERM) overfits to the noise and is not the best solution.

In the train-by-validation setting, each task P contains a training set S_{train} with n_1 samples and a validation set with n_2 samples. The inner objective is defined as $\hat{f}(w) = \frac{1}{2n_1} \sum_{(x,y) \in S_{\text{train}}} \left(\langle w, x \rangle - y \rangle^2 \right)$. Let $\{w_{\tau,\eta}\}$ be the GD sequence running on $\hat{f}(w)$ from initialization 0 (with truncation). For each task P, the meta-loss $\Delta_{TbV(n_1,n_2)}(\eta,P)$ is defined as the loss of the last point $w_{t,\eta}$ evaluated on the validation set S_{valid} . That is, $\Delta_{TbV(n_1,n_2)}(\eta,P) = \frac{1}{2n_2} \sum_{(x,y) \in S_{\text{valid}}} \left(\langle w_{t,\eta}, x \rangle - y \right)^2$. The empirical meta objective $\hat{F}_{TbV(n_1,n_2)}(\eta)$ is the average of the meta-loss across m different tasks $P_1, P_2, ..., P_m$. We show that under $\hat{F}_{TbV(n_1,n_2)}(\eta)$, the optimal step size is $\Theta(1/t)$ and the learned weight is better than initialization 0 by a constant on new tasks. Theorem 8 is proved in Section B.3.

Theorem 8. Let the meta objective $\hat{F}_{TbV(n_1,n_2)}(\eta)$ be as defined in Equation 4 with $n_1, n_2 \in [d/4, 3d/4]$. Assume noise level σ is a large constant c_1 . Assume unroll length $t \geq c_2$, number of training tasks $m \geq c_3$ and dimension $d \ge c_4 \log(t)$ for certain constants c_2, c_3, c_4 . With probability at least 0.99 in the sampling of training tasks, we have

$$\eta_{valid}^* = \Theta(1/t) \text{ and } \mathbb{E} \|w_{t,\eta_{valid}^*} - w^*\|^2 = \|w^*\|^2 - \Omega(1)$$

for all $\eta_{valid}^* \in \arg\min_{\eta \geq 0} \hat{F}_{TbV(n_1,n_2)}(\eta)$, where the expectation is taken over new tasks.

Intuitively, train-by-validation is optimizing the right objective. As long as the meta-training problem has good generalization performance (that is, good performance on a few tasks implies good performance on the distribution of tasks), then train-by-validation should be able to choose the optimal learning rate. The step size of $\Theta(1/t)$ here serves as regularization similar to early-stopping, which allows gradient descent algorithm to achieve better error on test data.

Notations We define more quantities that are useful in the analysis. In the train by train setting, given a task $P_k := (\mathcal{D}(w_k^*), S_{\text{train}}^{(k)}, \ell)$. The training set $S_{\text{train}}^{(k)}$ contains n samples $\{x_i^{(k)}, y_i^{(k)}\}_{i=1}^n$ with $y_i^{(k)} = \left\langle w_k^*, x_i^{(k)} \right\rangle + \xi_i^{(k)}$.

Let $X_{\text{train}}^{(k)}$ be an $n \times d$ matrix with its i-th row as $(x_i^{(k)})^{\top}$. Let $H_{\text{train}}^{(k)} := \frac{1}{n}(X_{\text{train}}^{(k)})^{\top}X_{\text{train}}^{(k)}$ be the covariance matrix of the inputs in $S_{\text{train}}^{(k)}$. Let $\xi_{\text{train}}^{(k)}$ be an n-dimensional column vector with its i-th entry equal to $\xi_i^{(k)}$. Since $n \leq d$, with probability 1, we know $X_{\text{train}}^{(k)}$ is full row rank. Therefore, $X_{\text{train}}^{(k)}$ has pseudo-inverse $(X_{\text{train}}^{(k)})^{\dagger}$ such that $X_{\text{train}}^{(k)}(X_{\text{train}}^{(k)})^{\dagger} = I_n$. It's not hard to verify that there exists $w_{\text{train}}^{(k)} = \text{Proj}_{(X_{\text{train}}^{(k)})^{\top}} w_k^* + (X_{\text{train}}^{(k)})^{\dagger} \xi_{\text{train}}^{(k)}$ such that $y_i^{(k)} = \left\langle w_{\text{train}}^{(k)}, x_i^{(k)} \right\rangle \text{ for every } (x_i^{(k)}, y_i^{(k)}) \in S_{\text{train}}^{(k)}. \text{ Here, } \operatorname{Proj}_{(X_{\text{train}}^{(k)})^\top} \text{ is the projection matrix onto the column span } \operatorname{of}(X_{\text{train}}^{(k)})^\top. \text{ We also denote } \operatorname{Proj}_{(X_{\text{train}}^{(k)})^\top} w_k^* \text{ as } (w_{\text{train}}^{(k)})^*. \text{ We use } B_{t,\eta}^{(k)} \text{ to denote } (I - (I - \eta H_{\text{train}}^{(k)})^t). \text{ Let } w_{t,\eta}^{(k)} \text{ be the } W_{t,\eta}^{(k)} \text{ to denote } (I - (I - \eta H_{\text{train}}^{(k)})^t). \text{ The substitution of } W_{t,\eta}^{(k)} \text{ to denote } W_{t,\eta}^{(k$ weight obtained by running GD on $S_{\mathrm{train}}^{^{(\mathrm{ran})}}$ with step size η (with truncation).

With the above notations, it's not hard to verify that for task P_k , the inner objective $\hat{f}(w) = \frac{1}{2} \left\| w - w_{\text{train}}^{(k)} \right\|_{L^{(k)}}^2$

The meta-loss on task
$$P_k$$
 is just $\Delta_{TbT(n)}(\eta,P_k)=\frac{1}{2}\left\|w_{t,\eta}-w_{\mathrm{train}}^{(k)}\right\|_{H_{\mathrm{train}}^{(k)}}^2$

In the train-by-validation setting, each task P_k contains a training set $S_{\text{train}}^{(k)}$ with n_1 samples and a validation set $S_{\text{valid}}^{(k)}$ with n_2 samples. Similar as above, for the training set $S_{\text{train}}^{(k)}$, we can define $\xi_{\text{train}}^{(k)}$, $X_{\text{train}}^{(k)}$, $X_{\text{trai$

We also use these notations without index k to refer to the quantities defined on task P. In the proofs, we ignore the subsripts on n, n_1, n_2 and simply write $\Delta_{TbT}(\eta, P_k), \Delta_{TbV}(\eta, P_k), \hat{F}_{TbT}, \hat{F}_{TbV}, F_{TbT}, F_{TbV}$.

Overall Proof Strategy B.1

In this section (and the next), we follow similar proof strategies that consists of three steps.

Step 1: First, we show for both train-by-train and train-by-validation, there is a good step size that achieves small empirical meta-objective (however the step sizes and the empirical meta-objective they achieve are different in the two settings). This does not necessarily mean that the actual optimal step size is exactly the good step size that we propose, but it gives an upperbound on the empirical meta-objective for the optimal step size.

Step 2: Second, we define a threshold step size such that for any step size larger than it, the empirical meta-objective must be higher than what was achieved at the good step size in Step 1. This immediately implies that the optimal step size cannot exceed this threshold step size.

Step 3: Third, we show the meta-learning problem has good generalization performance, that is, if a learning rate η performs well on the training tasks, it must also perform well on the task distribution, and vice versa. Thanks to Step 1 and Step 2, we know the optimal step size cannot exceed certain threshold and then only need to prove generalization result within this range. The generalization result is not surprising as we only have a single trainable parameter η , however we also emphasize that this is non-trivial as we will not restrict the step size η to be small enough that the algorithms do not diverge. Instead we use a truncation to alleviate the diverging problem (this allows us to run the algorithm on distribution of data whose largest possible learning rate is unknown).

Combing Step 1, 2, 3, we know the population meta-objective has to be small at the optimal step size. Finally, we show that as long as the population meta-objective is small, the performance of the algorithms satisfy what we stated in Theorem 5. The last step is easier for the train-by-validation setting, because its meta-objective is exactly the correct measure that we are looking at; for the train-by-train setting we instead look at the property of empirical risk minimizer (ERM), and show that anything close to the ERM is going to behave similarly.

B.2 Train-by-train (GD)

Recall Theorem 7 as follows.

Theorem 7. Let the meta objective $\hat{F}_{TbT(n)}(\eta)$ be as defined in Equation 3 with $n \in [d/4, 3d/4]$. Assume noise level σ is a large constant c_1 . Assume unroll length $t \geq c_2$, number of training tasks $m \geq c_3 \log(mt)$ and dimension $d \geq c_4 \log(m)$ for certain constants c_2, c_3, c_4 . With probability at least 0.99 in the sampling of the training tasks, we have

$$\eta_{train}^* = \Theta(1)$$
 and $\mathbb{E} \left\| w_{t,\eta_{train}^*} - w^* \right\|^2 = \Omega(1)\sigma^2$,

for all $\eta_{train}^* \in \arg\min_{\eta > 0} \hat{F}_{TbT(\eta)}(\eta)$, where the expectation is taken over new tasks.

According to the data distribution, we know X_{train} is an $n \times d$ random matrix with each entry i.i.d. sampled from standard Gaussian distribution. In the following lemma, we show that the covariance matrix H_{train} is approximately isotropic when $d/4 \le n \le 3d/4$. Specifically, we show $\frac{\sqrt{d}}{\sqrt{L}} \le \sigma_i(X_{\text{train}}) \le \sqrt{Ld}$ and $\frac{1}{L} \le \lambda_i(H_{\text{train}}) \le L$ for all $i \in [n]$ with L = 100. We use letter L to denote the upper bound of $\|H_{\text{train}}\|$ to emphasize that this bounds the smoothness of the inner objective. Throughout this section, we use letter L to denote constant 100. The proof of Lemma 7 follows from random matrix theory. We defer its proof into Section B.2.4.

Lemma 7. Let $X \in \mathbb{R}^{n \times d}$ be a random matrix with each entry i.i.d. sampled from standard Gaussian distribution. Let $H := 1/nX^{\top}X$. Assume n = cd with $c \in [\frac{1}{4}, \frac{3}{4}]$. Then, with probability at least $1 - \exp(-\Omega(d))$, there exists constant L = 100 such that

$$\frac{\sqrt{d}}{\sqrt{L}} \le \sigma_i(X) \le \sqrt{Ld} \text{ and } \frac{1}{L} \le \lambda_i(H) \le L,$$

for all $i \in [n]$.

In this section, we always assume the size of each training set is within [d/4, 3d/4] so Lemma 7 holds. Since $\|H_{\text{train}}\|$ is upper bounded by L with high probability, we know the GD sequence converges to w_{train} for $\eta \in [0, 1/L]$. In Lemma 1, we prove that the empirical meta objective \hat{F}_{TbT} monotonically decreases as η increases until 1/L. Also, we show \hat{F}_{TbT} is exponentially small in t at step size 1/L. This serves as step 1 in Section B.1. The proof is deferred into Section B.2.1.

Lemma 1. With probability at least $1 - m \exp(-\Omega(d))$, $\hat{F}_{TbT}(\eta)$ is monotonically decreasing in [0, 1/L] and

$$\hat{F}_{TbT}(1/L) \le 2L^2\sigma^2\left(1 - \frac{1}{L^2}\right)^t.$$

When the step size is larger than 1/L, the GD sequence can diverge, which incurs a high loss in meta objective. Later in Definition 1, we define a step size $\tilde{\eta}$ such that the GD sequence gets truncated with descent probability for any step size that is larger than $\tilde{\eta}$. In Lemma 2, we show with high probability, the empirical meta objective is high for all $\eta > \tilde{\eta}$. This serves as step 2 in the proof strategy described in Section B.1. The proof is deferred into Section B.2.2.

Lemma 2. With probability at least $1 - \exp(-\Omega(m))$,

$$\hat{F}_{TbT}(\eta) \ge \frac{\sigma^2}{10L^8}$$

for all $\eta > \tilde{\eta}$.

By Lemma 1 and Lemma 2, we know the optimal step size must lie in $[1/L, \tilde{\eta}]$. We can also show $1/L < \tilde{\eta} < 3L$, so η^*_{train} is a constant. To relate the empirical loss at η^*_{train} to the population loss. We prove a generalization result for step sizes within $[1/L, \tilde{\eta}]$. The following lemma is a formal version of Lemma 3. This serves as step 3 in Section B.1. The proof is deferred into Section B.2.3.

Lemma 8. Suppose σ is a large constant c_1 . Assume $t \geq c_2, d \geq c_4$ for certain constants c_2, c_4 . With probability at least $1 - m \exp(-\Omega(d)) - O(t + m) \exp(-\Omega(m))$,

$$|F_{TbT}(\eta) - \hat{F}_{TbT}(\eta)| \le \frac{\sigma^2}{L^3},$$

for all $\eta \in [1/L, \tilde{\eta}]$,

Combining the above lemmas, we know the population meta objective F_{TbT} is small at η^*_{train} , which means $w_{t,\eta^*_{\text{train}}}$ is close to the ERM solution. Since the ERM solution overfits to the noise in training tasks, we know $\|w_{t,\eta^*_{\text{train}}} - w^*\|$ has to be large. We present the proof of Theorem 7 as follows.

Proof of Theorem 7. We assume σ is a large constant in this proof. According to Lemma 1, we know with probability at least $1-m\exp(-\Omega(d))$, $\hat{F}_{TbT}(\eta)$ is monotonically decreasing in [0,1/L] and $\hat{F}_{TbT}(1/L) \leq 2L^2\sigma^2(1-1/L^2)^t$. This implies that the optimal step size $\eta^*_{\text{train}} \geq 1/L$ and $\hat{F}_{TbT}(\eta^*_{\text{train}}) \leq 2L^2\sigma^2(1-1/L^2)^t$. By Lemma 2, we know with probability at least $1-\exp(-\Omega(m))$, $\hat{F}_{TbT}(\eta) \geq \frac{\sigma^2}{10L^8}$ for all $\eta > \tilde{\eta}$, where $\tilde{\eta}$ is defined in Definition 1. As long as $t \geq c_2$ for certain constant c_2 , we know $\frac{\sigma^2}{10L^8} > 2L^2\sigma^2(1-1/L^2)^t$, which then implies that the optimal step size η^*_{train} lies in $[1/L, \tilde{\eta}]$. According to Lemma 10, we know $\tilde{\eta} \in (1/L, 3L)$. Therefore η^*_{train} is a constant.

According to Lemma 8, we know with probability at least $1-m\exp(-\Omega(d))-O(t+m)\exp(-\Omega(m))$, $|F_{TbT}(\eta)-D(t+m)|^2$

According to Lemma 8, we know with probability at least $1-m\exp(-\Omega(d))-O(t+m)\exp(-\Omega(m)), |F_{TbT}(\eta)-\hat{F}_{TbT}(\eta)| \leq \frac{\sigma^2}{L^3}$, for all $\eta \in [1/L, \tilde{\eta}]$. As long as t is larger than some constant, we have $\hat{F}_{TbT}(\eta^*_{\text{train}}) \leq \frac{\sigma^2}{L^3}$. Combing with the generalization result, we have $F_{TbT}(\eta^*_{\text{train}}) \leq \frac{2\sigma^2}{L^3}$. Next, we show that under a small population loss, $\mathbb{E} \|w_{t,\eta^*_{\text{train}}} - w^*\|^2$ has to be large.

Let \mathcal{E}_1 be the event that $\sqrt{d}/\sqrt{L} \leq \sigma_i(X_{\text{train}}) \leq \sqrt{Ld}$ and $1/L \leq \lambda_i(H_{\text{train}}) \leq L$ for all $i \in [n]$ and $\sqrt{d}\sigma/4 \leq \|\xi_{\text{train}}\| \leq \sqrt{d}\sigma$. We have

$$\begin{split} \mathbb{E} \left\| w_{t,\eta_{\text{train}}^*} - w_{\text{train}} \right\|_{H_{\text{train}}}^2 &\geq \frac{1}{L} \mathbb{E} \left\| w_{t,\eta_{\text{train}}^*} - w_{\text{train}} \right\|^2 \mathbb{1} \left\{ \mathcal{E}_1 \right\} \\ &\geq \frac{1}{L} \left(\mathbb{E} \left\| w_{t,\eta_{\text{train}}^*} - w_{\text{train}}^* - (X_{\text{train}})^\dagger \xi_{\text{train}} \right\| \mathbb{1} \left\{ \mathcal{E}_1 \right\} \right)^2 \\ &\geq \frac{1}{L} \left(\mathbb{E} \left\| (X_{\text{train}})^\dagger \xi_{\text{train}} \right\| \mathbb{1} \left\{ \mathcal{E}_1 \right\} - \mathbb{E} \left\| w_{t,\eta_{\text{train}}^*} - w_{\text{train}}^* \right\| \mathbb{1} \left\{ \mathcal{E}_1 \right\} \right)^2. \end{split}$$

Since $\mathbb{E} \left\| w_{t,\eta_{\text{train}}^*} - w_{\text{train}} \right\|_{H_{\text{train}}}^2 \leq \frac{4\sigma^2}{L^3}$, this then implies

$$\mathbb{E} \left\| (X_{\text{train}})^{\dagger} \xi_{\text{train}} \right\| \mathbb{1} \left\{ \mathcal{E}_{1} \right\} - \mathbb{E} \left\| w_{t, \eta_{\text{train}}^{*}} - w_{\text{train}}^{*} \right\| \mathbb{1} \left\{ \mathcal{E}_{1} \right\} \leq \sqrt{L \frac{4\sigma^{2}}{L^{3}}} = \frac{2\sigma}{L}.$$

Conditioning on \mathcal{E}_1 , we can lower bound $\|(X_{\text{train}})^{\dagger}\xi_{\text{train}}\|$ by $\frac{\sigma}{4\sqrt{L}}$. According to Lemma 7 and Lemma 49, we know $\Pr[\mathcal{E}_1] \geq 1 - \exp(-\Omega(d))$. As long as d is at least certain constant, we have $\Pr[\mathcal{E}_1] \geq 0.9$. This then implies $\mathbb{E}\|(X_{\text{train}})^{\dagger}\xi_{\text{train}}\|\mathbb{1}\{\mathcal{E}_1\} \geq \frac{9\sigma}{40\sqrt{L}}$. Therefore, we have

$$\mathbb{E}\left\|w_{t,\eta_{\text{train}}^*} - w_{\text{train}}^*\right\| \mathbb{1}\left\{\mathcal{E}_1\right\} \geq \frac{9\sigma}{40\sqrt{L}} - \frac{2\sigma}{L} = \frac{9\sigma}{4L} - \frac{2\sigma}{L} = \frac{\sigma}{4L},$$

where the first equality uses L = 100. Then, we have

$$\mathbb{E}\left\|w_{t,\eta_{\text{train}}^*} - w^*\right\|^2 \ge \mathbb{E}\left\|w_{t,\eta_{\text{train}}^*} - w_{\text{train}}^*\right\|^2 \mathbb{1}\left\{\mathcal{E}_1\right\} \ge \left(\mathbb{E}\left\|w_{t,\eta_{\text{train}}^*} - w_{\text{train}}^*\right\| \mathbb{1}\left\{\mathcal{E}_1\right\}\right)^2 \ge \frac{\sigma^2}{16L^2},$$

where the first inequality holds because for any S_{train} , w_{train}^* is the projection of w^* on the subspace of S_{train} and $w_{t,\eta_{\text{train}}^*}$ is also in this subspace. Taking a union bound for all the bad events, we know this result holds with probability at least 0.99 as long as σ is a large constant c_1 and $t \ge c_2$, $m \ge c_3 \log(mt)$ and $d \ge c_4 \log(m)$ for certain constants c_2 , c_3 , c_4 .

B.2.1 Behavior of \hat{F}_{TbT} for $\eta \in [0, 1/L]$

In this section, we prove the empirical meta objective \hat{F}_{TbT} is monotonically decreasing in [0, 1/L]. Furthermore, we show $\hat{F}_{TbT}(1/L)$ is exponentially small in t.

Lemma 1. With probability at least $1 - m \exp(-\Omega(d))$, $\hat{F}_{TbT}(\eta)$ is monotonically decreasing in [0, 1/L] and

$$\hat{F}_{TbT}(1/L) \le 2L^2\sigma^2 \left(1 - \frac{1}{L^2}\right)^t.$$

Proof of Lemma 1. For each $k \in [m]$, let \mathcal{E}_k be the event that $\sqrt{d}/\sqrt{L} \leq \sigma_i(X_{\text{train}}) \leq \sqrt{Ld}$ and $1/L \leq \lambda_i(H_{\text{train}}) \leq L$ for all $i \in [n]$ and $\sqrt{d}\sigma/4 \leq \|\xi_{\text{train}}\| \leq \sqrt{d}\sigma$. Here, L is constant 100 from Lemma 7. According to Lemma 7 and Lemma 49, we know for each $k \in [m]$, \mathcal{E}_k happens with probability at least $1 - \exp(-\Omega(d))$. Taking a union bound over all $k \in [m]$, we know $\cap_{k \in [m]} \mathcal{E}_k$ holds with probability at least $1 - m \exp(-\Omega(d))$. From now on, we assume $\cap_{k \in [m]} \mathcal{E}_k$ holds.

Let's first consider each individual loss function $\Delta_{TbT}(\eta, P_k)$. Let $\{\hat{w}_{\tau,\eta}^{(k)}\}$ be the GD sequence without truncation. We have

$$\begin{split} \hat{w}_{\tau,\eta}^{(k)} - w_{\text{train}}^{(k)} = & \hat{w}_{\tau-1,\eta}^{(k)} - w_{\text{train}}^{(k)} - \eta H_{\text{train}}^{(k)} (\hat{w}_{\tau-1,\eta}^{(k)} - w_{\text{train}}^{(k)}) \\ = & (I - \eta H_{\text{train}}^{(k)}) (\hat{w}_{\tau-1,\eta}^{(k)} - w_{\text{train}}^{(k)}) = - (I - \eta H_{\text{train}}^{(k)})^t w_{\text{train}}^{(k)} \end{split}$$

For any $\eta \in [0,1/L]$, we have $\left\|\hat{w}_{ au,\eta}^{(k)}\right\| \leq \left\|w_{ ext{train}}^{(k)}\right\| = \left\|(w_{ ext{train}}^{(k)})^* + (X_{ ext{train}}^{(k)})^\dagger \xi_{ ext{train}}^{(k)}\right\| \leq 2\sqrt{L}\sigma$ for any τ . Therefore, $\left\|w_{t,\eta}^{(k)}\right\|$ never exceeds the norm threshold and never gets truncated.

Noticing that $\Delta_{TbT}(\eta, P_k) = \frac{1}{2} (w_{t,\eta}^{(k)} - w_{\text{train}}^{(k)})^{\top} H_{\text{train}}^{(k)}(w_{t,\eta}^{(k)} - w_{\text{train}}^{(k)})$, we have

$$\Delta_{TbT}(\eta, P_k) = \frac{1}{2} (w_{\text{train}}^{(k)})^\top H_{\text{train}}^{(k)} (I - \eta H_{\text{train}}^{(k)})^{2t} w_{\text{train}}^{(k)}.$$

Taking the derivative of $\Delta_{TbT}(\eta, P_k)$ in η , we have

$$\frac{\partial}{\partial \eta} \Delta_{TbT}(\eta, P_k) = -t(w_{\text{train}}^{(k)})^{\top} (H_{\text{train}}^{(k)})^2 (I - \eta H_{\text{train}}^{(k)})^{2t-1} w_{\text{train}}^{(k)}.$$

Conditioning on \mathcal{E}_k , we know $1/L \leq \lambda_i(H_{\text{train}}^{(k)}) \leq L$ for all $i \in [n]$ and $H_{\text{train}}^{(k)}$ is full rank in the row span of $X_{\text{train}}^{(k)}$. Therefore, we know $\frac{\partial}{\partial \eta} \Delta_{TbT}(\eta, P_k) < 0$ for all $\eta \in [0, 1/L)$. Here, we assume $\left\| w_{\text{train}}^{(k)} \right\| > 0$, which happens with probability 1.

Overall, we know that conditioning on $\bigcap_{k \in [m]} \mathcal{E}_k$, every $\Delta_{TbT}(\eta, P_k)$ is strictly decreasing for $\eta \in [0, 1/L]$. Since $\hat{F}_{TbT}(\eta) := \frac{1}{m} \sum_{k=1}^m \Delta_{TbT}(\eta, P_k)$, we know $\hat{F}_{TbT}(\eta)$ is strictly decreasing when $\eta \in [0, 1/L]$. At step size $\eta = 1/L$, we have

$$\Delta_{TbT}(\eta, P_k) = \frac{1}{2} (w_{\text{train}}^{(k)})^{\top} H_{\text{train}}^{(k)} (I - \eta H_{\text{train}}^{(k)})^{2t} w_{\text{train}}^{(k)}$$

$$\leq \frac{1}{2} L \left(1 - \frac{1}{L^2} \right)^t \left\| w_{\text{train}}^{(k)} \right\|^2 \leq 2L^2 \sigma^2 \left(1 - \frac{1}{L^2} \right)^t,$$

where we upper bound $\left\|w_{\text{train}}^{(k)}\right\|^2$ by $4L\sigma^2$ at the last step. Therefore, we have $\hat{F}_{TbT}(1/L) \leq 2L^2\sigma^2(1-\frac{1}{L^2})^t$.

B.2.2 Lower bounding \hat{F}_{ThT} for $\eta \in (\tilde{\eta}, \infty)$

In this section, we prove that the empirical meta objective is lower bounded by $\Omega(\sigma^2)$ with high probability for $\eta \in (\tilde{\eta}, \infty)$. Step size $\tilde{\eta}$ is defined such that there is a descent probability of diverging for any step size larger than $\tilde{\eta}$. Then, we show the contribution from these truncated sequence will be enough to provide an $\Omega(\sigma^2)$ lower bound for \tilde{F}_{TbT} . The proof of Lemma 2 is given at the end of this section.

Lemma 2. With probability at least $1 - \exp(-\Omega(m))$,

$$\hat{F}_{TbT}(\eta) \ge \frac{\sigma^2}{10L^8}$$

for all $\eta > \tilde{\eta}$.

We define $\tilde{\eta}$ as the smallest step size such that the contribution from the truncated sequence in the population meta objective exceeds certain threshold. The precise definition is as follows.

Definition 1. Given a training task P, let \mathcal{E}_1 be the event that $\sqrt{d}/\sqrt{L} \le \sigma_i(X_{train}) \le \sqrt{Ld}$ and $1/L \le \lambda_i(H_{train}) \le L$ for all $i\in[n]$ and $\sqrt{d}\sigma/4\le\|\xi_{train}\|\le\sqrt{d}\sigma$. Let $\bar{\mathcal{E}}_2(\eta)$ be the event that the GD sequence is truncated with step size η . Define $\tilde{\eta}$ as follows,

$$\tilde{\eta} = \inf \left\{ \eta \geq 0 \middle| \mathbb{E}\frac{1}{2} \left\| w_{t,\eta} - w_{\textit{train}} \right\|_{H_{\textit{train}}}^2 \mathbb{1} \left\{ \mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\eta) \right\} \geq \frac{\sigma^2}{L^6} \right\}.$$

In the next lemma, we prove that for any fixed training set, $\mathbb{1}\left\{\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\eta')\right\} \geq \mathbb{1}\left\{\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\eta)\right\}$ for any $\eta' \geq \eta$.

This immediately implies that $\Pr[\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\eta)]$ and $\mathbb{E}_{\frac{1}{2}} \| w_{t,\eta} - w_{\text{train}} \|_{H_{\text{train}}}^2 \mathbb{1} \left\{ \mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\eta) \right\}$ is non-decreasing in η . Basically we need to show, conditioning on \mathcal{E}_1 , if a GD sequence gets truncated at step size η , it must be also truncated for larger step sizes. Let $\{w'_{\tau,\eta}\}$ be the GD sequence without truncation. We only need to show that for any τ , if $\|w'_{\tau,\eta}\|$ exceeds the norm threshold, $\|w'_{\tau,\eta'}\|$ must also exceed the norm threshold for any $\eta' \geq \eta$. This is easy to prove if τ is odd because in this case $\|w'_{\tau,\eta}\|$ is always non-decreasing in η . The case when τ is even is trickier because there indeed exists certain range of η such that $\|w'_{\tau,\eta}\|$ is decreasing in η . We manage to prove that this problematic case cannot happen when $||w'_{\tau,n}||$ is at least $4\sqrt{L}\sigma$. The full proof of Lemma 9 is deferred into Section B.2.4.

Lemma 9. Fixing a task P, let \mathcal{E}_1 and $\bar{\mathcal{E}}_2(\eta)$ be as defined in Definition 1. We have

$$\mathbb{1}\left\{\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\eta')\right\} \ge \mathbb{1}\left\{\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\eta)\right\},\,$$

for any $\eta' \geq \eta$.

In the next Lemma, we prove that $\tilde{\eta}$ must lie within (1/L, 3L). We prove this by showing that the GD sequence never gets truncated for $\eta \in [0, 2/L]$ and almost always gets truncated for $\eta \in [2.5L, \infty)$. The proof is deferred into Section B.2.4.

Lemma 10. Let $\tilde{\eta}$ be as defined in Definition 1. Suppose σ is a large constant c_1 . Assume $t \geq c_2, d \geq c_4$ for some constants c_2, c_4 . We have

$$1/L < \tilde{\eta} < 3L$$
.

Now, we are ready to give the proof of Lemma 2.

Proof of Lemma 2. Let \mathcal{E}_1 and $\bar{\mathcal{E}}_2(\eta)$ be as defined in Definition 1. For the simplicity of the proof, we assume $\mathbb{E}\frac{1}{2}\|w_{t,\tilde{\eta}}-w_{ ext{train}}\|_{H_{ ext{train}}}^2\mathbb{1}\left\{\mathcal{E}_1\cap\bar{\mathcal{E}}_2(\tilde{\eta})\right\}\geq \frac{\sigma^2}{L^6}$. We will discuss the proof for the other case at the end, which is very

Conditioning on \mathcal{E}_1 , we know $\frac{1}{2} \|w_{t,\tilde{\eta}} - w_{\text{train}}\|_{H_{\text{train}}}^2 \leq 18L^2\sigma^2$. Therefore, we know $\Pr[\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\tilde{\eta})] \geq \frac{1}{18L^8}$. For each task P_k , define $\mathcal{E}_1^{(k)}$ and $\bar{\mathcal{E}}_2^{(k)}(\eta)$ as the corresponding events on training set $S_{\text{train}}^{(k)}$. By Hoeffding's inequality, we know with probability at least $1 - \exp(-\Omega(m))$,

$$\frac{1}{m}\sum_{k=1}^{m}\mathbb{1}\left\{\mathcal{E}_{1}^{(k)}\cap\bar{\mathcal{E}}_{2}^{(k)}(\tilde{\eta})\right\}\geq\frac{1}{20L^{8}}.$$

By Lemma 9, we know $\mathbb{1}\left\{\mathcal{E}_1^{(k)}\cap\bar{\mathcal{E}}_2^{(k)}(\eta)\right\}\geq\mathbb{1}\left\{\mathcal{E}_1^{(k)}\cap\bar{\mathcal{E}}_2^{(k)}(\tilde{\eta})\right\}$ for any $\eta\geq\tilde{\eta}$. Then, we can lower bound \hat{F}_{TbT} for any $\eta > \tilde{\eta}$ as follows,

$$\begin{split} \hat{F}_{TbT}(\eta) &= \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{\text{train}}^{(k)} \right\|_{H_{\text{train}}}^{2} \geq \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{\text{train}}^{(k)} \right\|_{H_{\text{train}}}^{2} \mathbbm{1} \left\{ \mathcal{E}_{1}^{(k)} \cap \bar{\mathcal{E}}_{2}^{(k)}(\eta) \right\} \\ &\geq 2\sigma^{2} \frac{1}{m} \sum_{k=1}^{m} \mathbbm{1} \left\{ \mathcal{E}_{1}^{(k)} \cap \bar{\mathcal{E}}_{2}^{(k)}(\eta) \right\} \\ &\geq 2\sigma^{2} \frac{1}{m} \sum_{k=1}^{m} \mathbbm{1} \left\{ \mathcal{E}_{1}^{(k)} \cap \bar{\mathcal{E}}_{2}^{(k)}(\tilde{\eta}) \right\} \geq \frac{\sigma^{2}}{10L^{8}}, \end{split}$$

where the second inequality lower bounds the loss for one task by $2\sigma^2$ when the sequence gets truncated.

We have assumed $\mathbb{E}^1_2 \| w_{t,\tilde{\eta}} - w_{\text{train}} \|^2_{H_{\text{train}}} \mathbb{1} \left\{ \mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\tilde{\eta}) \right\} \geq \frac{\sigma^2}{L^6}$ in the proof. Now, we show the proof also works when $\mathbb{E}^1_2 \| w_{t,\tilde{\eta}} - w_{\text{train}} \|^2_{H_{\text{train}}} \mathbb{1} \left\{ \mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\tilde{\eta}) \right\} < \frac{\sigma^2}{L^6}$ with slight changes. According to the definition and Lemma 9, we know $\mathbb{E}^1_2 \| w_{t,\tilde{\eta}} - w_{\text{train}} \|^2_{H_{\text{train}}} \mathbb{1} \left\{ \mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\tilde{\eta}) \right\} > \frac{\sigma^2}{L^6}$ for all $\eta > \tilde{\eta}$. At each training set S_{train} , we can define $\mathbb{1} \left\{ \mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\tilde{\eta}') \right\}$ as $\lim_{\eta \to \tilde{\eta}^+} \mathbb{1} \left\{ \mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\eta) \right\}$. We also have $\Pr[\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\tilde{\eta}')] \geq \frac{1}{18L^8}$. The remaining proof is the same as before as we substitute $\mathbb{1} \left\{ \mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\tilde{\eta}) \right\}$. same as before as we substitute $\mathbb{1}\left\{\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\tilde{\eta})\right\}$ by $\mathbb{1}\left\{\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\tilde{\eta}')\right\}$.

B.2.3 Generalization for $\eta \in [1/L, \tilde{\eta}]$

In this section, we show empirical meta objective \hat{F}_{TbT} is point-wise close to population meta objective F_{TbT} for all

Lemma 8. Suppose σ is a large constant c_1 . Assume $t \geq c_2$, $d \geq c_4$ for certain constants c_2 , c_4 . With probability at least $1 - m \exp(-\Omega(d)) - O(t + m) \exp(-\Omega(m))$,

$$|F_{TbT}(\eta) - \hat{F}_{TbT}(\eta)| \le \frac{\sigma^2}{L^3},$$

for all $\eta \in [1/L, \tilde{\eta}]$,

In this section, we first show \hat{F}_{TbT} concentrates on F_{TbT} for any fixed η and then construct ϵ -net for \hat{F}_{TbT} and F_{TbT} for $\eta \in [1/L, \tilde{\eta}]$. We give the proof of Lemma 8 at the end.

We first show that for a fixed η , $\hat{F}_{TbT}(\eta)$ is close to $F_{TbT}(\eta)$ with high probability. We prove the meta-loss on each task $\Delta_{TbT}(\eta, P_k)$ is O(1)-subexponential. Then we apply Bernstein's inequality to get the result. The proof is deferred into Section B.2.4. We will assume σ is a large constant and $t \ge c_2, d \ge c_4$ for some constants c_2, c_4 so that Lemma 10 holds and $\tilde{\eta}$ is a constant.

Lemma 11. Suppose σ is a constant. For any fixed η and any $1 > \epsilon > 0$, with probability at least $1 - \exp(-\Omega(\epsilon^2 m))$,

$$\left| \hat{F}_{TbT}(\eta) - F_{TbT}(\eta) \right| \le \epsilon.$$

Next, we construct an ϵ -net for F_{TbT} . By the definition of $\tilde{\eta}$, we know for any $\eta \leq \tilde{\eta}$, the contribution from truncated sequences in $F_{TbT}(\eta)$ is small. We can show the contribution from the un-truncated sequences is O(t)-

Lemma 12. Suppose σ is a large constant c_1 . Assume $t \geq c_2, d \geq c_4$ for some constant c_2, c_4 . There exists an $\frac{11\sigma^2}{L^4}$ -net $N \subset [1/L, \tilde{\eta}]$ for F_{TbT} with |N| = O(t). That means, for any $\eta \in [1/L, \tilde{\eta}]$,

$$|F_{TbT}(\eta) - F_{TbT}(\eta')| \le \frac{11\sigma^2}{L^4},$$

for $\eta' = \arg\min_{\eta'' \in N, \eta'' < \eta} (\eta - \eta'')$.

Proof of Lemma 12. Let \mathcal{E}_1 and $\bar{\mathcal{E}}_2(\eta)$ be as defined in Definition 1. For the simplicity of the proof, we assume $\mathbb{E}\frac{1}{2}\|w_{t,\tilde{\eta}}-w_{\text{train}}\|_{H_{\text{train}}}^2\mathbb{1}\left\{\mathcal{E}_1\cap\bar{\mathcal{E}}_2(\tilde{\eta})\right\}\leq \frac{\sigma^2}{L^6}$. We will discuss the proof for the other case at the end, which is very

We can divide $\mathbb{E}^{\frac{1}{2}} \| w_{t,\eta} - w_{\text{train}} \|_{H_{\text{train}}}^2$ as follows,

$$\begin{split} & \mathbb{E}\frac{1}{2}\left\|w_{t,\eta} - w_{\text{train}}\right\|_{H_{\text{train}}}^2 \\ = & \mathbb{E}\frac{1}{2}\left\|w_{t,\eta} - w_{\text{train}}\right\|_{H_{\text{train}}}^2 \mathbb{1}\left\{\mathcal{E}_1 \cap \mathcal{E}_2(\tilde{\eta})\right\} + \mathbb{E}\frac{1}{2}\left\|w_{t,\eta} - w_{\text{train}}\right\|_{H_{\text{train}}}^2 \mathbb{1}\left\{\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\tilde{\eta})\right\} \\ & + \mathbb{E}\frac{1}{2}\left\|w_{t,\eta} - w_{\text{train}}\right\|_{H_{\text{train}}}^2 \mathbb{1}\left\{\bar{\mathcal{E}}_1\right\}. \end{split}$$

We will construct an ϵ -net for the first term and show the other two terms are small. Let's first consider the third term. Since $\frac{1}{2} \|w_{t,\eta} - w_{\text{train}}\|_{H_{\text{train}}}^2$ is O(1)-subexponential and $\Pr[\bar{\mathcal{E}}_1] \leq \exp(-\Omega(d))$, we have $\mathbb{E}\frac{1}{2} \|w_{t,\eta} - w_{\text{train}}\|_{H_{\text{train}}}^2 \mathbb{1} \left\{ \bar{\mathcal{E}}_1 \right\} = 0$ $O(1) \exp(-\Omega(d)). \text{ Choosing } d \text{ to be at least certain constant, we know } \frac{1}{2} \|w_{t,\eta} - w_{\text{train}}\|_{H_{\text{train}}}^2 \mathbb{1} \left\{ \bar{\mathcal{E}}_1 \right\} \leq \sigma^2 / L^4.$ Then we upper bound the second term. Since $\mathbb{E} \frac{1}{2} \|w_{t,\tilde{\eta}} - w_{\text{train}}\|_{H_{\text{train}}}^2 \mathbb{1} \left\{ \mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\tilde{\eta}) \right\} \leq \frac{\sigma^2}{L^6}$ and

 $\frac{1}{2} \|w_{t,\tilde{\eta}} - w_{\mathrm{train}}\|_{H_{\mathrm{train}}}^2 \geq 2\sigma^2$ when $w_{t,\tilde{\eta}}$ diverges, we know $\Pr[\mathcal{E}_1 \cap \widetilde{\mathcal{E}}_2(\tilde{\eta})] \leq \frac{1}{2L^6}$. Then, we can upper bound the

$$\mathbb{E}\frac{1}{2}\left\|w_{t,\eta} - w_{\text{train}}\right\|_{H_{\text{train}}}^2 \mathbb{1}\left\{\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\tilde{\eta})\right\} \leq 18L^2\sigma^2 \frac{1}{2L^6} = \frac{9\sigma^2}{L^4}$$

Next, we show the first term $\frac{1}{2} \|w_{t,\eta} - w_{\text{train}}\|_{H_{\text{train}}}^2 \mathbbm{1} \{\mathcal{E}_1 \cap \mathcal{E}_2(\tilde{\eta})\}$ has desirable Lipschitz condition. According to Lemma 9, we know $\mathbbm{1} \{\mathcal{E}_1 \cap \mathcal{E}_2(\eta)\} \geq \mathbbm{1} \{\mathcal{E}_1 \cap \mathcal{E}_2(\tilde{\eta})\}$ for any $\eta \leq \tilde{\eta}$. Therefore, conditioning on $\mathcal{E}_1 \cap \mathcal{E}_2(\tilde{\eta})$, we know $w_{t,\eta}$ never gets truncated for any $\eta \leq \tilde{\eta}$. This means $w_{t,\eta} = B_{t,\eta} w_{\text{train}}$ with $B_{t,\eta} = (I - (I - \eta H_{\text{train}})^t)$. We can compute the derivative of $\frac{1}{2} \|w_{t,\eta} - w_{\text{train}}\|_{H_{\text{train}}}^2$ as follows,

$$\frac{\partial}{\partial n} \frac{1}{2} \left\| w_{t,\eta} - w_{\text{train}} \right\|_{H_{\text{train}}}^2 = \left\langle t H_{\text{train}} (I - \eta H_{\text{train}})^{t-1} w_{\text{train}}, H_{\text{train}} (w_{t,\eta} - w_{\text{train}}) \right\rangle.$$

Since $\|w_{t,\eta}\| = \|(I - (I - \eta H_{\mathrm{train}})^t)w_{\mathrm{train}}\| \le 4\sqrt{L}\sigma$ and $\|w_{\mathrm{train}}\| \le 2\sqrt{L}\sigma$, we have $\|(I - \eta H_{\mathrm{train}})^tw_{\mathrm{train}}\| \le 4\sqrt{L}\sigma$ $6\sqrt{L}\sigma$. We can bound $\|(I-\eta H_{\text{train}})^{t-1}w_{\text{train}}\|$ with $\|(I-\eta H_{\text{train}})^t w_{\text{train}}\| + \|w_{\text{train}}\|$ by bounding the expanding directions using $\|(I - \eta H_{\text{train}})^t w_{\text{train}}\|$ and bounding the shrinking directions using $\|w_{\text{train}}\|$. Therefore, we can bound the derivative as follows,

$$\left|\frac{\partial}{\partial \eta}\frac{1}{2}\left\|w_{t,\eta}-w_{\mathrm{train}}\right\|_{H_{\mathrm{train}}}^{2}\right|\leq tL\times 8\sqrt{L}\sigma\times 6L\sqrt{L}\sigma=48L^{3}\sigma^{2}t.$$

Suppose σ is a constant, we know $\mathbb{E}\frac{1}{2}\|w_{t,\eta}-w_{\text{train}}\|_{H_{\text{train}}}^2\mathbb{1}\{\mathcal{E}_1\cap\mathcal{E}_2(\tilde{\eta})\}$ is O(t)-lipschitz. Therefore, there exists an $\frac{\sigma^2}{L^4}$ -net N for $\mathbb{E} \frac{1}{2} \|w_{t,\eta} - w_{ ext{train}}\|_{H_{ ext{train}}}^2 \mathbb{1} \{\mathcal{E}_1 \cap \mathcal{E}_2(\tilde{\eta})\}$ with size O(t). That means, for any $\eta \in [1/L, \tilde{\eta}]$,

$$\left| \mathbb{E}\frac{1}{2} \left\| w_{t,\eta} - w_{\text{train}} \right\|_{H_{\text{train}}}^2 \mathbb{1} \left\{ \mathcal{E}_1 \cap \mathcal{E}_2(\tilde{\eta}) \right\} - \mathbb{E}\frac{1}{2} \left\| w_{t,\eta'} - w_{\text{train}} \right\|_{H_{\text{train}}}^2 \mathbb{1} \left\{ \mathcal{E}_1 \cap \mathcal{E}_2(\tilde{\eta}) \right\} \right| \leq \frac{\sigma^2}{L^4}$$

for $\eta' = \arg\min_{\eta'' \in N, \eta'' \leq \eta} (\eta - \eta'')$. Note we construct the ϵ -net in a particular way such that η' is chosen as the largest step size in N that is at most η .

Combing with the upper bounds on the second term and the third term, we have for any $\eta \in [1/L, \tilde{\eta}]$,

$$|F_{TbT}(\eta) - F_{TbT}(\eta')| \le \frac{11\sigma^2}{L^4}$$

for $\eta' = \arg\min_{\eta'' \in N, \eta'' < \eta} (\eta - \eta'')$.

In the above analysis, we have assumed $\mathbb{E}\frac{1}{2}\|w_{t,\tilde{\eta}}-w_{\text{train}}\|_{H_{\text{train}}}^2\mathbb{I}\left\{\mathcal{E}_1\cap\bar{\mathcal{E}}_2(\tilde{\eta})\right\}\leq \frac{\sigma^2}{L^6}$. The proof can be easily generalized to the other case. We can define $\mathbb{I}\left\{\mathcal{E}_1\cap\bar{\mathcal{E}}_2(\tilde{\eta}')\right\}$ as $\lim_{\eta\to\tilde{\eta}^-}\mathbb{I}\left\{\mathcal{E}_1\cap\bar{\mathcal{E}}_2(\eta)\right\}$. Then the proof works as long as we substitute $\mathbb{I}\left\{\mathcal{E}_1\cap\bar{\mathcal{E}}_2(\tilde{\eta})\right\}$ by $\mathbb{I}\left\{\mathcal{E}_1\cap\bar{\mathcal{E}}_2(\tilde{\eta}')\right\}$. We will also add $\tilde{\eta}$ into the ϵ -net.

In order to prove F_{TbT} is close to \hat{F}_{TbT} point-wise in $[1/L, \tilde{\eta}]$, we still need to construct an ϵ -net for the empirical meta objective \hat{F}_{TbT} .

Lemma 13. Suppose σ is a large constant c_1 . Assume $t \geq c_2, d \geq c_4$ for certain constants c_2, c_4 . With probability at least $1 - m \exp(-\Omega(d))$, there exists an $\frac{\sigma^2}{L^4}$ -net $N' \subset [1/L, \tilde{\eta}]$ for \hat{F}_{TbT} with |N| = O(t+m). That means, for any $\eta \in [1/L, \tilde{\eta}]$,

 $|\hat{F}_{TbT}(\eta) - \hat{F}_{TbT}(\eta')| \le \frac{\sigma^2}{L^4},$

for $\eta' = \arg\min_{\eta'' \in N', \eta'' < \eta} (\eta - \eta'')$.

Proof of Lemma 13. For each $k \in [m]$, let $\mathcal{E}_{1,k}$ be the event that $\sqrt{d}/\sqrt{L} \leq \sigma_i(X_{\text{train}}^{(k)}) \leq \sqrt{Ld}$ and $1/L \leq \lambda_i(H_{\text{train}}^{(k)}) \leq L$ for all $i \in [n]$ and $\sqrt{d}\sigma/4 \leq \left\|\xi_{\text{train}}^{(k)}\right\| \leq \sqrt{d}\sigma$. According to Lemma 7 and Lemma 49, we know with probability at least $1 - m \exp(-\Omega(d))$, $\mathcal{E}_{1,k}$'s hold for all $k \in [m]$. From now on, we assume all these events hold. Recall that the empirical meta objective as follows,

$$\hat{F}_{TbT}(\eta) := \frac{1}{m} \sum_{k=1}^{m} \Delta_{TbT}(\eta, P_k).$$

For any $k \in [m]$, let $\eta_{c,k}$ be the smallest step size such that $w_{t,\eta}^{(k)}$ gets truncated. If $\eta_{c,k} > \hat{\eta}$, by similar argument as in Lemma 12, we know $\Delta_{TbT}(\eta,P_k)$ is O(t)-Lipschitz in $[1/L,\hat{\eta}]$ as long as σ is a constant. If $\eta_{c,k} \leq \hat{\eta}$, by Lemma 9 we know $w_{t,\eta}^{(k)}$ gets truncated for any $\eta \geq \eta_{c,k}$. This then implies that $\Delta_{TbT}(\eta,P_k)$ is a constant function for $\eta \in [\eta_{c,k},\hat{\eta}]$. We can also show that $\Delta_{TbT}(\eta,P_k)$ is O(t)-Lipschitz in $[1/L,\eta_{c,k})$. There might be a discontinuity in function value at $\eta_{c,k}$, so we need to add $\eta_{c,k}$ into the ϵ -net.

Overall, we know there exists an $\frac{\sigma^2}{L^4}$ -net N' with |N'| = O(t+m) for \hat{F}_{TbT} . That means, for any $\eta \in [1/L, \tilde{\eta}]$,

$$\left| \hat{F}_{TbT}(\eta) - \hat{F}_{TbT}(\eta') \right| \le \frac{\sigma^2}{L^4}$$

for $\eta' = \arg\min_{\eta'' \in N', \eta'' \le \eta} (\eta - \eta'')$.

Finally, we combine Lemma 11, Lemma 12 and Lemma 13 to prove that \hat{F}_{TbT} is point-wise close to F_{TbT} for $\eta \in [1/L, \tilde{\eta}]$.

Proof of Lemma 8. We assume σ as a constant in this proof. By Lemma 11, we know with probability at least $1-\exp(-\Omega(\epsilon^2m)), \ \left|\hat{F}_{TbT}(\eta)-F_{TbT}(\eta)\right| \leq \epsilon$ for any fixed η . By Lemma 12, we know there exists an $\frac{11\sigma^2}{L^4}$ -net N for F_{TbT} with size O(t). By Lemma 13, we know with probability at least $1-m\exp(-\Omega(d))$, there exists an $\frac{\sigma^2}{L^4}$ -net N' for \hat{F}_{TbT} with size O(t+m). According to the proofs of Lemma 12 and Lemma 13, it's not hard to verify that $N \cup N'$ is still an $\frac{11\sigma^2}{L^4}$ -net for \hat{F}_{TbT} and F_{TbT} . That means, for any $\eta \in [1/L, \tilde{\eta}]$, we have

$$|F_{TbT}(\eta) - F_{TbT}(\eta')|, |\hat{F}_{TbT}(\eta) - \hat{F}_{TbT}(\eta')| \le \frac{11\sigma^2}{L^4},$$

for $\eta' = \arg\min_{\eta'' \in N \cup N', \eta'' \leq \eta} (\eta - \eta'')$.

Taking a union bound over $N \cup N'$, we have with probability at least $1 - O(t + m) \exp(-\Omega(m))$,

$$\left| \hat{F}_{TbT}(\eta) - F_{TbT}(\eta) \right| \le \frac{\sigma^2}{L^4}$$

for all $\eta \in N \cup N'$.

Overall, we know with probability at least $1 - m \exp(-\Omega(d)) - O(t + m) \exp(-\Omega(m))$, for all $\eta \in [1/L, \tilde{\eta}]$,

$$|F_{TbT}(\eta) - \hat{F}_{TbT}(\eta)|$$

$$\leq |F_{TbT}(\eta) - F_{TbT}(\eta')| + |\hat{F}_{TbT}(\eta) - \hat{F}_{TbT}(\eta')| + |\hat{F}_{TbT}(\eta') - F_{TbT}(\eta')|$$

$$\leq \frac{23\sigma^2}{L^4} \leq \frac{\sigma^2}{L^3},$$

where $\eta' = \arg\min_{\eta'' \in N \cup N', \eta'' < \eta} (\eta - \eta'')$. We use the fact that L = 100 in the last inequality.

B.2.4 Proofs of Technical Lemmas

Proof of Lemma 7. Recall that X_{train} is an $n \times d$ matrix with n = cd where $c \in [1/4, 3/4]$. According to Lemma 52, with probability at least $1 - 2\exp(-t^2/2)$, we have

$$\sqrt{d} - \sqrt{cd} - t \le \sigma_i(X_{\text{train}}) \le \sqrt{d} + \sqrt{cd} + t,$$

for all $i \in [n]$.

Since $H_{\text{train}} = 1/nX_{\text{train}}^{\top}X_{\text{train}}$, we know $\lambda_i(H_{\text{train}}) = 1/n\sigma_i^2(X_{\text{train}})$. Since $c \in [\frac{1}{4}, \frac{3}{4}]$, we have $\frac{1}{cd}(\sqrt{d} + \sqrt{cd})^2 \leq 100 - c'$ and $\frac{1}{cd}(\sqrt{d} - \sqrt{cd})^2 \geq \frac{1}{100} + c'$, for some constant c'. Therefore, we know with probability at least $1 - \exp(-\Omega(d))$,

$$\frac{1}{100} \le \lambda_i(H_{\text{train}}) \le 100.$$

for all $i \in [n]$.

Similarly, since there exists constant c'' such that $\sqrt{d} + \sqrt{cd} \le (10 - c'')\sqrt{d}$ and $\sqrt{d} - \sqrt{cd} \ge (1/10 + c'')\sqrt{d}$, we know with probability at least $1 - \exp(-\Omega(d))$,

$$\frac{1}{10}\sqrt{d} \le \sigma_i(X_{\text{train}}) \le 10\sqrt{d},$$

for all $i \in [n]$. Choosing L = 100 finishes the proof.

Proof of Lemma 9. We prove that for any training set S_{train} , $\mathbbm{1}\left\{\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\eta')\right\} \geq \mathbbm{1}\left\{\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\eta')\right\}$ for any $\eta' > \eta$. This is trivially true if \mathcal{E}_1 is false on S_{train} . Therefore, we focus on the case when \mathcal{E}_1 holds for S_{train} . Suppose η_c is the smallest step size such that the GD sequence gets truncated. Let $\{w'_{\tau,\eta_c}\}$ be the GD sequence without truncation. There must exists $\tau \leq t$ such that $\|w'_{\tau,\eta_c}\| \geq 4\sqrt{L}\sigma$. We only need to prove that $\|w'_{\tau,\eta}\| \geq 4\sqrt{L}\sigma$ for any $\eta \geq \eta_c$. We prove this by showing the derivative of $\|w'_{\tau,\eta}\|^2$ in η is non-negative assuming $\|w'_{\tau,\eta}\|^2 \geq 4\sqrt{L}\sigma$.

Recall the recursion of $w'_{\tau,\eta}$ as $w'_{\tau,\eta} = w_{\text{train}} - (I - \eta H_{\text{train}})^{\tau} w_{\text{train}}$. If τ is an odd number, it's clear that $\frac{\partial}{\partial \eta} \|w'_{\tau,\eta}\|^2$ is non-negative at any $\eta \geq 0$. From now on, we assume τ is an even number. Actually in this case, $\frac{\partial}{\partial \eta} \|w'_{\tau,\eta}\|^2$ can be negative for some η . However, we can prove the derivative must be non-negative assuming $\|w'_{\tau,\eta}\|^2 \geq 4\sqrt{L}\sigma$.

Suppose the eigenvalue decomposition of H_{train} is $\sum_{i=1}^{n} \lambda_i u_i u_i^{\top}$ with $\lambda_1 \geq \cdots \lambda_n$. Denote c_i as $\langle w_{\text{train}}, u_i \rangle$. Let λ_j be the smallest eigenvalue such that $(1 - \eta \lambda_j) \leq -1$. This implies $\lambda_i \leq 2/\eta$ for any $i \geq j+1$. We can write down $\|w_{\tau,n}'\|^2$ as follows

$$\begin{aligned} \left\| w_{\tau,\eta}' \right\|^2 &= \sum_{i=1}^j \left(1 - (1 - \eta \lambda_i)^t \right)^2 c_i^2 + \sum_{i=j+1}^n \left(1 - (1 - \eta \lambda_i)^t \right)^2 c_i^2 \\ &\leq \sum_{i=1}^j \left(1 - (1 - \eta \lambda_i)^t \right)^2 c_i^2 + \left\| w_{\text{train}} \right\|^2. \end{aligned}$$

Since \mathcal{E}_1 holds, we know $\|w_{\text{train}}\|^2 \leq 4L\sigma^2$. Combining with $\|w'_{\tau,\eta}\|^2 \geq 16L\sigma^2$, we have $\sum_{i=1}^{j} (1-(1-\eta\lambda_i)^t)^2 c_i^2 \geq 12L\sigma^2$. We can lower bound the derivative as follows,

$$\frac{\partial}{\partial \eta} \|w_{\tau,\eta}\|^{2} = \sum_{i=1}^{j} 2t\lambda_{i}(1 - \eta\lambda_{i})^{t-1} \left(1 - (1 - \eta\lambda_{i})^{t}\right) c_{i}^{2} + \sum_{i=j+1}^{n} 2t\lambda_{i}(1 - \eta\lambda_{i})^{t-1} \left(1 - (1 - \eta\lambda_{i})^{t}\right) c_{i}^{2}
\geq 2t \sum_{i=1}^{j} \lambda_{i}(1 - \eta\lambda_{i})^{t-1} \left(1 - (1 - \eta\lambda_{i})^{t}\right) c_{i}^{2} - 2t \frac{2}{\eta} \sum_{i=j+1}^{n} c_{i}^{2}
\geq 2t \sum_{i=1}^{j} \lambda_{i}(1 - \eta\lambda_{i})^{t-1} \left(1 - (1 - \eta\lambda_{i})^{t}\right) c_{i}^{2} - 2t \times 8L\sigma^{2}/\eta.$$

Then, we only need to show that $\sum_{i=1}^{j} \lambda_i (1-\eta\lambda_i)^{t-1} (1-(1-\eta\lambda_i)^t) c_i^2$ is larger than $8L\sigma^2/\eta$. We have

$$\begin{split} \sum_{i=1}^{j} \lambda_{i} (1 - \eta \lambda_{i})^{t-1} \left(1 - (1 - \eta \lambda_{i})^{t} \right) c_{i}^{2} &= \sum_{i=1}^{j} \lambda_{i} \frac{(1 - \eta \lambda_{i})^{t-1}}{1 - (1 - \eta \lambda_{i})^{t}} \left(1 - (1 - \eta \lambda_{i})^{t} \right)^{2} c_{i}^{2} \\ &= \sum_{i=1}^{j} \lambda_{i} \frac{(\eta \lambda_{i} - 1)^{t-1}}{(\eta \lambda_{i} - 1)^{t} - 1} \left(1 - (1 - \eta \lambda_{i})^{t} \right)^{2} c_{i}^{2} \\ &= \sum_{i=1}^{j} \lambda_{i} \frac{(\eta \lambda_{i} - 1)^{t}}{(\eta \lambda_{i} - 1)^{t} - 1} \frac{1}{\eta \lambda_{i} - 1} \left(1 - (1 - \eta \lambda_{i})^{t} \right)^{2} c_{i}^{2} \\ &\geq \sum_{i=1}^{j} \frac{1}{\eta} \left(1 - (1 - \eta \lambda_{i})^{t} \right)^{2} c_{i}^{2} \geq 12L\sigma^{2}/\eta > 8L\sigma^{2}/\eta. \end{split}$$

Proof of Lemma 10. Similar as the analysis in Lemma 1, conditioning on \mathcal{E}_1 , we know the GD sequence never exceeds the norm threshold for any $\eta \in [0, 2/L]$. This then implies

$$\mathbb{E}\frac{1}{2}\left\|w_{t,\eta}-w_{\mathrm{train}}\right\|_{H_{\mathrm{train}}}^2\mathbb{1}\left\{\mathcal{E}_1\cap\bar{\mathcal{E}}_2(\eta)\right\}=0.$$

for all $\eta \in [0, 2/L]$.

Let $\{w'_{\tau,\eta}\}$ be the GD sequence without truncation. For any step size $\eta \in [2.5L, \infty]$, conditioning on \mathcal{E}_1 , we have

$$\|w'_{t,\eta}\| \ge \left((\eta/L - 1)^t - 1 \right) \|w_{\text{train}}\| \ge \left(1.5^t - 1 \right) \left(\frac{\sigma}{4\sqrt{L}} - 1 \right) \ge 4\sqrt{L}\sigma,$$

where the last inequality holds as long as $\sigma \geq 5\sqrt{L}, t \geq c_2$ for some constant c_2 . Therefore, we know when $\eta \in [2.5L,\infty), \mathbbm{1}\left\{\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\eta)\right\} = \mathbbm{1}\left\{\mathcal{E}_1\right\}$. Then, we have for any $\eta \geq 2.5L$,

$$\mathbb{E}\frac{1}{2} \|w_{t,\eta} - w_{\text{train}}\|_{H_{\text{train}}}^2 \mathbb{1} \left\{ \mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\eta) \right\} \ge \frac{1}{2L} \left(4\sqrt{L}\sigma - 2\sqrt{L}\sigma \right)^2 \Pr[\mathcal{E}_1] \ge 2\sigma^2 \Pr[\mathcal{E}_1] \ge \frac{\sigma^2}{L^3}$$

where the last inequality uses $\Pr[\mathcal{E}_1] \geq 1 - \exp(-\Omega(d))$ and assume $d \geq c_4$ for some constant c_4 .

Overall, we know $\mathbb{E}\frac{1}{2}\|w_{t,\eta}-w_{\mathrm{train}}\|_{H_{\mathrm{train}}}^2\mathbb{1}\left\{\mathcal{E}_1\cap\bar{\mathcal{E}}_2(\eta)\right\}$ equals zero for all $\eta\in[0,2/L]$ and is at least $\frac{\sigma^2}{L^3}$ for all $\eta\in[2.5L,\infty)$. By definition, we know $\hat{\eta}\in(1/L,3L)$.

Proof of Lemma 11. Recall that $\hat{F}_{TbT}(\eta) := \frac{1}{m} \sum_{k=1}^{m} \Delta_{TbT}(\eta, P_k)$. We prove that each $\Delta_{TbT}(\eta, P_k)$ is O(1)-subexponential. We can further write $\Delta_{TbT}(\eta, P_k)$ as follows,

$$\begin{split} \Delta_{TbT}(\eta, P_k) = & \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_k^* - (X_{\text{train}}^{(k)})^\dagger \xi_{\text{train}}^{(k)} \right\|_{H_{\text{train}}^{(k)}}^2 \\ \leq & \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_k^* \right\|^2 \left\| H_{\text{train}}^{(k)} \right\| + \frac{1}{2n} \left\| \xi_{\text{train}}^{(k)} \right\|^2 + \left\| w_{t,\eta}^{(k)} - w_k^* \right\| \left(\frac{1}{\sqrt{n}} \left\| \xi_{\text{train}}^{(k)} \right\| \right) \left(\frac{1}{\sqrt{n}} \left\| X_{\text{train}}^{(k)} \right\| \right). \end{split}$$

We can write $\left\|H_{\text{train}}^{(k)}\right\|$ as $\sigma_{\max}^2(\frac{1}{\sqrt{n}}X_{\text{train}}^{(k)})$. According to Lemma 51, we know $\sigma_{\max}(X_{\text{train}}^{(k)}) - \mathbb{E}\sigma_{\max}(X_{\text{train}}^{(k)})$ is O(1)-subgaussian, which implies that $\sigma_{\max}(\frac{1}{\sqrt{n}}X_{\text{train}}^{(k)}) - \mathbb{E}\sigma_{\max}(\frac{1}{\sqrt{n}}X_{\text{train}}^{(k)})$ is $O(1/\sqrt{d})$ -subgaussian. Since $\mathbb{E}\sigma_{\max}(\frac{1}{\sqrt{n}}X_{\text{train}}^{(k)})$ is a constant, we know $\sigma_{\max}(\frac{1}{\sqrt{n}}X_{\text{train}}^{(k)})$ is O(1)-subgaussian and $\sigma_{\max}^2(\frac{1}{\sqrt{n}}X_{\text{train}}^{(k)})$ is O(1)-subexponential. Similarly, we know both $\frac{1}{2n}\left\|\xi_{\text{train}}^{(k)}\right\|^2$ and $\left(\frac{1}{\sqrt{n}}\left\|X_{\text{train}}^{(k)}\right\|\right)\left(\frac{1}{\sqrt{n}}\left\|\xi_{\text{train}}^{(k)}\right\|\right)$ are O(1)-subexponential.

Suppose σ is a constant, we know $\left\|w_{t,\eta}^{(k)} - w_k^*\right\|$ is upper bounded by a constant. Then, we know $\Delta_{TbT}(\eta, P_k)$ is O(1)-subexponential. Therefore, $\hat{F}_{TbT}(\eta)$ is the average of m i.i.d. O(1)-subexponential random variables. By standard concentration inequality, we know for any $1 > \epsilon > 0$, with probability at least $1 - \exp(-\Omega(\epsilon^2 m))$,

$$\left|\hat{F}_{TbT}(\eta) - F_{TbT}(\eta)\right| \le \epsilon.$$

B.3 Train-by-validation (GD)

In this section, we show that the optimal step size under \hat{F}_{TbV} is $\Theta(1/t)$. Furthermore, we show under this optimal step size, GD sequence makes constant progress towards the ground truth. Precisely, we prove the following theorem.

Theorem 8. Let the meta objective $\hat{F}_{TbV(n_1,n_2)}(\eta)$ be as defined in Equation 4 with $n_1,n_2 \in [d/4,3d/4]$. Assume noise level σ is a large constant c_1 . Assume unroll length $t \geq c_2$, number of training tasks $m \geq c_3$ and dimension $d \geq c_4 \log(t)$ for certain constants c_2, c_3, c_4 . With probability at least 0.99 in the sampling of training tasks, we have

$$\eta_{\textit{valid}}^* = \Theta(1/t) \ \textit{and} \ \mathbb{E} \left\| w_{t,\eta_{\textit{valid}}^*} - w^* \right\|^2 = \left\| w^* \right\|^2 - \Omega(1)$$

for all $\eta_{valid}^* \in \arg\min_{\eta \geq 0} \hat{F}_{TbV(n_1,n_2)}(\eta)$, where the expectation is taken over new tasks.

In this section, we still use L to denote constant 100. We start from analyzing the behavior of the population meta-objective F_{TbV} for step sizes within [0,1/L]. We show the optimal step size within this range is $\Theta(1/t)$ and GD sequence moves towards w^* under the optimal step size. The following lemma is a formal version of Lemma 4. This serves as step 1 in Section B.1. We defer the proof of Lemma 14 into Section B.3.1.

Lemma 14. Suppose noise level σ is a large enough constant c_1 . Assume unroll length $t \ge c_2$ and dimension $d \ge c_4$ for some constants c_2, c_4 . There exist $\eta_1, \eta_2, \eta_3 = \Theta(1/t)$ with $\eta_1 < \eta_2 < \eta_3$ such that

$$F_{TbV}(\eta_2) \le \frac{1}{2} \|w^*\|^2 - \frac{9}{10}C + \frac{\sigma^2}{2}$$
$$F_{TbV}(\eta) \ge \frac{1}{2} \|w^*\|^2 - \frac{6}{10}C + \frac{\sigma^2}{2}, \forall \eta \in [0, \eta_1] \cup [\eta_3, 1/L]$$

where C is a positive constant.

To relate the behavior of F_{TbV} to the behavior of \hat{F}_{TbV} , we prove the following generalization result for step sizes in [0, 1/L]. The following lemma is a formal version of Lemma 6. This serves as step 3 in Section B.1. The proof is deferred into Section B.3.2.

Lemma 15. For any $1 > \epsilon > 0$, assume $d \ge c_4 \log(1/\epsilon)$ for some constant c_4 . With probability at least $1 - O(1/\epsilon) \exp(-\Omega(\epsilon^2 m))$,

$$|\hat{F}_{TbV}(\eta) - F_{TbV}(\eta)| \le \epsilon,$$

for all $\eta \in [0, 1/L]$.

In Lemma 16, we show the empirical meta objective \hat{F}_{TbV} is high for all step size larger than 1/L, which then implies $\eta_{\text{valid}}^* \in [0, 1/L]$. The following lemma is a formal version of Lemma 5. This serves as step 2 in Section B.1. We prove this lemma in Section B.3.3.

Lemma 16. Suppose σ is a large constant. Assume $t \geq c_2$, $d \geq c_4 \log(t)$ for some constants c_2 , c_4 . With probability at least $1 - \exp(-\Omega(m))$,

$$\hat{F}_{TbV}(\eta) \ge C'\sigma^2 + \frac{1}{2}\sigma^2,$$

for all $\eta \geq 1/L$, where C' is a positive constant independent with σ .

Combining Lemma 14, Lemma 15 and Lemma 16, we give the proof of Theorem 8.

Proof of Theorem 8. According to Lemma 14, we know as long as d and t are larger than certain constants, there exists $\eta_1, \eta_2, \eta_3 = \Theta(1/t)$ with $\eta_1 < \eta_2 < \eta_3$ such that

$$F_{TbV}(\eta_2) \le \frac{1}{2} \|w^*\|^2 - \frac{9}{10}C + \sigma^2/2$$

$$F_{TbV}(\eta) \ge \frac{1}{2} \|w^*\|^2 - \frac{6}{10}C + \sigma^2/2, \forall \eta \in [0, \eta_1] \cup [\eta_3, 1/L],$$

for some positive constant C.

Choosing $\epsilon = \min(1, C/10)$ in Lemma 15, we know as long as d is larger than certain constant, with probability at least $1 - \exp(-\Omega(m))$,

$$|\hat{F}_{TbV}(\eta) - F_{TbV}(\eta)| \le C/10,$$

for all $\eta \in [0, 1/L]$.

Therefore,

$$\hat{F}_{TbV}(\eta_2) \le \frac{1}{2} \|w^*\|^2 - \frac{8}{10}C + \sigma^2/2$$

$$\hat{F}_{TbV}(\eta) \ge \frac{1}{2} \|w^*\|^2 - \frac{7}{10}C + \sigma^2/2, \forall \eta \in [0, \eta_1] \cup [\eta_3, 1/L].$$

By Lemma 16, we know as long as $t \ge c_2, d \ge c_4 \log(t)$ for some constants c_2, c_4 , with probability at least $1 - \exp(-\Omega(m)),$

$$\hat{F}_{TbV}(\eta) \ge C'\sigma^2 + \frac{1}{2}\sigma^2,$$

for all $\eta \geq 1/L$. As long as $\sigma \geq 1/\sqrt{C'}$, we have $\hat{F}_{TbV}(\eta) \geq 1 + \frac{1}{2}\sigma^2$ for all $\eta \geq 1/L$. Combining with $\hat{F}_{TbV}(\eta_2) \leq 1$ $\frac{1}{2}\|w^*\|^2 - \frac{8}{10}C + \sigma^2/2$, we know $\eta_{\text{valid}}^* \in [0, 1/L]$. Furthermore, since $\hat{F}_{TbV}(\eta) \geq \frac{1}{2}\|w^*\|^2 - \frac{7}{10}C + \sigma^2/2$, $\forall \eta \in [0, 1/L]$ $[0,\eta_1] \cup [\eta_3,1/L]$, we have $\eta_1 \leq \eta_{\text{valid}}^* \leq \eta_3$. Recall that $\eta_1,\eta_3 = \Theta(1/t)$, we know $\eta_{\text{valid}}^* = \Theta(1/t)$. At the optimal step size, we have

$$F_{TbV}(\eta_{\text{valid}}^*) \le \hat{F}_{TbV}(\eta_{\text{valid}}^*) + C/10 \le \hat{F}_{TbV}(\eta_2) + C/10 \le \frac{1}{2} \|w^*\|^2 - \frac{7}{10}C + \sigma^2/2.$$

Since $F_{TbV}(\eta_{\text{valid}}^*) = \mathbb{E}\frac{1}{2} \|w_{t,\eta_{\text{valid}}^*} - w^*\|^2 + \sigma^2/2$, we have

$$\mathbb{E} \| w_{t,\eta_{\text{valid}}^*} - w^* \|^2 \le \| w^* \|^2 - \frac{7}{5} C.$$

Choosing m to be at least certain constant, this holds with probability at least 0.99.

B.3.1 Behavior of F_{TbV} for $\eta \in [0, 1/L]$

In this section, we study the behavior of F_{TbV} when $\eta \in [0, 1/L]$. We prove the following Lemma.

Lemma 14. Suppose noise level σ is a large enough constant c_1 . Assume unroll length $t \ge c_2$ and dimension $d \ge c_4$ for some constants c_2, c_4 . There exist $\eta_1, \eta_2, \eta_3 = \Theta(1/t)$ with $\eta_1 < \eta_2 < \eta_3$ such that

$$F_{TbV}(\eta_2) \le \frac{1}{2} \|w^*\|^2 - \frac{9}{10}C + \frac{\sigma^2}{2}$$

$$F_{TbV}(\eta) \ge \frac{1}{2} \|w^*\|^2 - \frac{6}{10}C + \frac{\sigma^2}{2}, \forall \eta \in [0, \eta_1] \cup [\eta_3, 1/L]$$

where C is a positive constant.

It's not hard to verify that $F_{TbV}(\eta) = \mathbb{E}1/2 \|w_{t,\eta} - w^*\|^2 + \sigma^2/2$. For convenience, denote $Q(\eta) := 1/2 \|w_{t,\eta} - w^*\|^2$. In order to prove Lemma 14, we only need to show that $\mathbb{E}Q(\eta_2) \leq \frac{1}{2} \|w^*\|^2 - \frac{9}{10}C$ and $\mathbb{E}Q(\eta) \geq \frac{1}{2} \|w^*\|^2 - \frac{6}{10}C$ for all $\eta \in [0, \eta_1] \cup [\eta_3, 1/L]$. In Lemma 17, we first show that this happens with high probability over the sampling of tasks.

Lemma 17. Suppose noise level σ is a large enough constant c_1 . Assume unroll length $t \geq c_2$ for certain constant c_2 . Then, with probability at least $1 - \exp(-\Omega(d))$ over the sampling of tasks, there exists $\eta_1, \eta_2, \eta_3 = \Theta(1/t)$ with $\eta_1 < \eta_2 < \eta_3$ such that

$$Q(\eta_2) := \frac{1}{2} \|w_{t,\eta_2} - w^*\|^2 \le \frac{1}{2} \|w^*\|^2 - C$$

$$Q(\eta) := \frac{1}{2} \|w_{t,\eta} - w^*\|^2 \ge \frac{1}{2} \|w^*\|^2 - \frac{C}{2}, \forall \eta \in [0, \eta_1] \cup [\eta_3, 1/L]$$

where C is a positive constant.

Since we are in the small step size regime, we know the GD sequence converges with high probability and will not be truncated. For now, let's assume $w_{t,\eta} = B_{t,\eta} w_{\text{train}}^* + B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}}$, where $B_{t,\eta} = I - (I - \eta H_{\text{train}})^t$. We have

$$\begin{split} Q(\eta) = & \frac{1}{2} \left\| B_{t,\eta} w_{\text{train}}^* + B_{t,\eta} (X_{\text{train}})^\dagger \xi_{\text{train}} - w^* \right\|^2 \\ = & \frac{1}{2} \left\| B_{t,\eta} w_{\text{train}}^* - w^* \right\|^2 + \frac{1}{2} \left\| B_{t,\eta} (X_{\text{train}})^\dagger \xi_{\text{train}} \right\|^2 \\ & + \left\langle B_{t,\eta} w_{\text{train}}^* - w^*, B_{t,\eta} (X_{\text{train}})^\dagger \xi_{\text{train}} \right\rangle \\ = & \frac{1}{2} \left\| w^* \right\|^2 + \frac{1}{2} \left\| B_{t,\eta} w_{\text{train}}^* \right\|^2 + \frac{1}{2} \left\| B_{t,\eta} (X_{\text{train}})^\dagger \xi_{\text{train}} \right\|^2 - \left\langle B_{t,\eta} w_{\text{train}}^*, w^* \right\rangle \\ & + \left\langle B_{t,\eta} w_{\text{train}}^* - w^*, B_{t,\eta} (X_{\text{train}})^\dagger \xi_{\text{train}} \right\rangle. \end{split}$$

In Lemma 18, we show that with high probability the crossing term $\langle B_{t,\eta} w_{\text{train}}^* - w^*, B_{t,\eta}(X_{\text{train}})^\dagger \xi_{\text{train}} \rangle$ is negligible for all $\eta \in [0,1/L]$. By Hoeffding's inequality, we know the crossing term is small for any fixed η . Constructing an ϵ -net for the crossing term in η , we can take a union bound and show it's small for all $\eta \in [0,1/L]$. We defer the proof of Lemma 18 to Section B.3.4.

Lemma 18. Assume σ is a constant. For any $1 > \epsilon > 0$, we know with probability at least $1 - O(1/\epsilon) \exp(-\Omega(\epsilon^2 d))$,

$$\left|\left\langle B_{t,\eta}w_{\textit{train}}^* - w^*, B_{t,\eta}(X_{\textit{train}})^{\dagger} \xi_{\textit{train}} \right\rangle\right| \leq \epsilon,$$

for all $\eta \in [0, 1/L]$.

Denote

$$G(\eta) := \frac{1}{2} \|w^*\|^2 + \frac{1}{2} \|B_{t,\eta} w_{\text{train}}^*\|^2 + \frac{1}{2} \|B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}}\|^2 - \langle B_{t,\eta} w_{\text{train}}^*, w^* \rangle.$$

Choosing $\epsilon = C/4$ in Lemma 18, we only need to show $G(\eta_2) \leq \|w^*\|^2 - 5C/4$ and $G(\eta) \geq \|w^*\|^2 - C/4$ for all $\eta \in [0, \eta_1] \cup [\eta_3, 1/L]$.

We first show that there exists $\eta_2 = \Theta(1/t)$ such that $G(\eta_2) \leq \frac{1}{2} \|w^*\|^2 - 5C/4$ for some constant C. It's not hard to show that $\frac{1}{2} \|B_{t,\eta} w^*_{\text{train}}\|^2 + \frac{1}{2} \|B_{t,\eta} (X_{\text{train}})^\dagger \xi_{\text{train}}\|^2 = O(\eta^2 t^2 \sigma^2)$. In Lemma 19, we show that the improvement $\langle B_{t,\eta} w^*_{\text{train}}, w^* \rangle = \Omega(\eta t)$ is linear in η . Therefore there exists $\eta_2 = \Theta(1/t)$ such that $G(\eta_2) \leq \frac{1}{2} \|w^*\|^2 - 5C/4$ for some constant C. We defer the proof of Lemma 19 to Section B.3.4.

Lemma 19. For any fixed $\eta \in [0, L/t]$ with probability at least $1 - \exp(-\Omega(d))$,

$$\langle B_{t,\eta} w_{train}^*, w^* \rangle \ge \frac{\eta t}{16L}.$$

To lower bound $G(\eta)$ for small η , we notice

$$G(\eta) \geq \frac{1}{2} \|w^*\|^2 - \langle B_{t,\eta} w_{\text{train}}^*, w^* \rangle.$$

We can show that $\langle B_{t,\eta} w_{\text{train}}^*, w^* \rangle = O(\eta t)$. Therefore, there exists $\eta_1 = \Theta(1/t)$ such that $\langle B_{t,\eta} w_{\text{train}}^*, w^* \rangle \leq C/4$ for all $\eta \in [0, \eta_1]$.

To lower bound $G(\eta)$ for large η , we lower bound $G(\eta)$ using the noise square term,

$$G(\eta) \geq \frac{1}{2} \left\| B_{t,\eta}(X_{\text{train}})^{\dagger} \xi_{\text{train}} \right\|^{2}.$$

We show that with high probability $\|B_{t,\eta}(X_{\text{train}})^{\dagger}\xi_{\text{train}}\|^2 = \Omega(\sigma^2)$ for all $\eta \in [\log(2)L/t, 1/L]$. Therefore, as long as σ is larger than some constant, there exists $\eta_3 = \Theta(1/t)$ such that $G(\eta) \geq \frac{1}{2} \|w^*\|^2$ for all $\eta \in [\eta_3, 1/L]$.

Combing Lemma 18 and Lemma 19, we give a complete proof for Lemma 17.

Proof of Lemma 17. Recall that

$$Q(\eta) = \frac{1}{2} \|B_{t,\eta} w_{\text{train}}^* - w^*\|^2 + \frac{1}{2} \|B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}}\|^2$$

$$+ \langle B_{t,\eta} w_{\text{train}}^* - w^*, B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}} \rangle$$

$$= G(\eta) + \langle B_{t,\eta} w_{\text{train}}^* - w^*, B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}} \rangle$$

We first show that with probability at least $1 - \exp(-\Omega(d))$, there exist $\eta_1, \eta_2, \eta_3 = \Theta(1/t)$ with $\eta_1 < \eta_2 < \eta_3$ such that $G(\eta_2) \le 1/2 \|w^*\|^2 - 5C/4$ and $G(\eta) \ge 1/2 \|w^*\|^2 - C/4$ for all $\eta \in [0, \eta_1] \cup [\eta_3, 1/L]$.

According to Lemma 7, we know with probability at least $1 - \exp(-\Omega(d))$, $\sqrt{d}/\sqrt{L} \le \sigma_i(X_{\text{train}}) \le \sqrt{Ld}$ and $1/L \le \lambda_i(H_{\text{train}}) \le L$ for all $i \in [n]$ with L = 100.

Upper bounding $G(\eta_2)$: We can expand $G(\eta)$ as follows:

$$\begin{split} G(\eta) := & \frac{1}{2} \left\| B_{t,\eta} w_{\text{train}}^* - w^* \right\|^2 + \frac{1}{2} \left\| B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}} \right\|^2 \\ = & \frac{1}{2} \left\| w^* \right\|^2 + \frac{1}{2} \left\| B_{t,\eta} w_{\text{train}}^* \right\|^2 + \frac{1}{2} \left\| B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}} \right\|^2 - \left\langle B_{t,\eta} w_{\text{train}}^*, w^* \right\rangle. \end{split}$$

Recall that $B_{t,\eta} = I - (I - \eta H_{\text{train}})^t$, for any vector w in the span of H_{train} ,

$$||B_{t,\eta}w|| = ||(I - (I - \eta H_{\text{train}})^t)w|| \le L\eta t ||w||.$$

According to Lemma 49, we know with probability at least $1 - \exp(-\Omega(d))$, $\|\xi_{\text{train}}\| \leq \sqrt{d}\sigma$. Therefore, we have

$$\frac{1}{2} \left\| B_{t,\eta} w_{\text{train}}^* \right\|^2 + \frac{1}{2} \left\| B_{t,\eta} (X_{\text{train}})^\dagger \xi_{\text{train}} \right\|^2 \le L^2 \eta^2 t^2 / 2 + L^3 \eta^2 t^2 \sigma^2 / 2 \le L^3 \eta^2 t^2 \sigma^2,$$

where the second inequality uses $\sigma, L \geq 1$. According to Lemma 19, for any fixed $\eta \in [0, L/t]$, with probability at least $1 - \exp(-\Omega(d))$, $\langle B_{t,\eta} w^*_{\text{train}}, w^* \rangle \geq \frac{\eta t}{16L}$. Therefore,

$$G(\eta) \le \frac{1}{2} \|w^*\|^2 + L^3 \eta^2 t^2 \sigma^2 - \frac{\eta t}{16L} \le \frac{1}{2} \|w^*\|^2 - \frac{\eta t}{32L}$$

where the second inequality holds as long as $\eta \leq \frac{1}{32L^4\sigma^2t}$. Choosing $\eta_2 := \frac{1}{32L^4\sigma^2t}$, we have

$$G(\eta_2) \le \frac{1}{2} \|w^*\|^2 - \frac{1}{1024L^5\sigma^2} = \frac{1}{2} \|w^*\|^2 - \frac{5C}{4}$$

where $C = \frac{1}{819.2L^5\sigma^2}$. Note C is a constant as σ, L are constants.

Lower bounding $G(\eta)$ for $\eta \in [0, \eta_1]$: Now, we prove that there exists $\eta_1 = \Theta(1/t)$ with $\eta_1 < \eta_2$ such that for any $\eta \in [0, \eta_1]$, $G(\eta) \ge \frac{1}{2} \|w^*\|^2 - \frac{C}{4}$. Recall that

$$\begin{split} G(\eta) = & \frac{1}{2} \left\| w^* \right\|^2 + \frac{1}{2} \left\| B_{t,\eta} w_{\text{train}}^* \right\|^2 + \frac{1}{2} \left\| B_{t,\eta} (X_{\text{train}})^\dagger \xi_{\text{train}} \right\|^2 - \left\langle B_{t,\eta} w_{\text{train}}^*, w^* \right\rangle. \\ \geq & \frac{1}{2} \left\| w^* \right\|^2 - \left\langle B_{t,\eta} w_{\text{train}}^*, w^* \right\rangle. \end{split}$$

Since $|\langle B_{t,\eta} w_{\text{train}}^*, w^* \rangle| \leq L \eta t$, we know for any $\eta \in [0, \eta_1]$,

$$G(\eta) \ge \frac{1}{2} \|w^*\|^2 - L\eta_1 t.$$

Choosing $\eta_1 = \frac{C}{4Lt}$, we have for any $\eta \in [0, \eta_1]$,

$$G(\eta) \ge \frac{1}{2} \|w^*\|^2 - \frac{C}{4}.$$

Lower bounding $G(\eta)$ for $\eta \in [\eta_3, 1/L]$: Now, we prove that there exists $\eta_3 = \Theta(1/t)$ with $\eta_3 > \eta_2$ such that for all $\eta \in [\eta_3, 1/L]$,

$$G(\eta) \ge \frac{1}{2} \|w^*\|^2 - \frac{C}{4}.$$

Recall that

$$G(\eta) = \frac{1}{2} \|B_{t,\eta} w_{\text{train}}^* - w^*\|^2 + \frac{1}{2} \|B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}}\|^2 \ge \frac{1}{2} \|B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}}\|^2.$$

According to Lemma 49, we know with probability at least $1 - \exp(-\Omega(d))$, $\frac{\sqrt{d}\sigma}{2\sqrt{2}} \le \|\xi_{\text{train}}\|$. Therefore,

$$\left\|B_{t,\eta}(X_{\text{train}})^{\dagger} \xi_{\text{train}}\right\|^{2} \ge \left(1 - e^{-\eta t/L}\right)^{2} \frac{\sigma^{2}}{8L} \ge \frac{\sigma^{2}}{32L},$$

where the last inequality assumes $\eta \ge \log(2)L/t$. As long as $t \ge \log(2)L^2$, we have $\log(2)L/t \le 1/L$. Choosing $\eta_3 = \log(2)L/t$, we know for all $\eta \in [\eta_3, 1/L]$,

$$G(\eta) \ge \frac{1}{2} \left\| B_{t,\eta}(X_{\text{train}})^{\dagger} \xi_{\text{train}} \right\|^2 \ge \frac{\sigma^2}{64L}.$$

Note that $\frac{1}{2} \|w^*\|^2 = 1/2$. Therefore, as long as $\sigma \geq 8\sqrt{L}$, we have

$$G(\eta) \ge \frac{1}{2} \left\| w^* \right\|^2$$

for all $\eta \in [\eta_3, 1/L]$.

Overall, we have shown that there exist $\eta_1, \eta_2, \eta_3 = \Theta(1/t)$ with $\eta_1 < \eta_2 < \eta_3$ such that $G(\eta_2) \le 1/2 \|w^*\|^2 - 5C/4$ and $G(\eta) \ge 1/2 \|w^*\|^2 - C/4$ for all $\eta \in [0, \eta_1] \cup [\eta_3, 1/L]$. Recall that $Q(\eta) = G(\eta) + \langle B_{t,\eta} w_{\text{train}}^* - w^*, B_{t,\eta}(X_{\text{train}})^\dagger \xi_{\text{train}} \rangle$. Choosing $\epsilon = C/4$ in Lemma 18, we know with probability at least $1 - \exp(-\Omega(d)), |\langle B_{t,\eta} w_{\text{train}}^* - w^*, B_{t,\eta}(X_{\text{train}})^\dagger \xi_{\text{train}} \rangle| \le C/4$ for all $\eta \in [0, 1/L]$. Therefore, we know $Q(\eta_2) \le 1/2 \|w^*\|^2 - C$ and $Q(\eta) \ge 1/2 \|w^*\|^2 - C/2$ for all $\eta \in [0, \eta_1] \cup [\eta_3, 1/L]$.

Next, we give the proof of Lemma 14.

Proof of Lemma 14. Recall that $F_{TbV}(\eta) = \mathbb{E}1/2 \|w_{t,\eta} - w^*\|^2 + \frac{\sigma^2}{2}$. For convenience, denote $Q(\eta) := 1/2 \|w_{t,\eta} - w^*\|^2$. In order to prove Lemma 14, we only need to show that $\mathbb{E}Q(\eta_2) \leq \frac{1}{2} \|w^*\|^2 - \frac{9}{10}C$ and $\mathbb{E}Q(\eta) \geq \frac{1}{2} \|w^*\|^2 - \frac{6}{10}C$ for all $\eta \in [0, \eta_1] \cup [\eta_3, 1/L]$.

According to Lemma 17, as long as σ is a large enough constant c_1 and t is at least certain constant c_2 , with probability at least $1 - \exp(-\Omega(d))$ over the sampling of S_{train} , there exists $\eta_1, \eta_2, \eta_3 = \Theta(1/t)$ with $\eta_1 < \eta_2 < \eta_3$ such that

$$Q(\eta_2) := 1/2 \|w_{t,\eta_2} - w^*\|^2 \le \frac{1}{2} \|w^*\|^2 - C$$

$$Q(\eta) := 1/2 \|w_{t,\eta} - w^*\|^2 \ge \frac{1}{2} \|w^*\|^2 - \frac{C}{2}, \forall \eta \in [0, \eta_1] \cup [\eta_3, 1/L]$$

where C is a positive constant. Call this event \mathcal{E} . Suppose the probability that \mathcal{E} happens is $1 - \delta$. We can write $\mathbb{E}Q(\eta)$ as follows,

$$\mathbb{E}Q(\eta) = \mathbb{E}[Q(\eta)|\mathcal{E}] \Pr[\mathcal{E}] + \mathbb{E}[Q(\eta)|\bar{\mathcal{E}}] \Pr[\bar{\mathcal{E}}].$$

According to the algorithm, we know $||w_{t,\eta}||$ is always bounded by $4\sqrt{L}\sigma$. Therefore, $Q(\eta) := 1/2 ||w_{t,\eta} - w^*||^2 \le$ $13L\sigma^2$. When $\eta = \eta_2$, we have

$$\mathbb{E}Q(\eta_2) \le \left(\frac{1}{2} \|w^*\|^2 - C\right) (1 - \delta) + 13L\sigma^2 \delta$$

$$= \frac{1}{2} \|w^*\|^2 - \frac{\delta}{2} - C + (C + 13L\sigma^2) \delta$$

$$\le \frac{1}{2} \|w^*\|^2 - \frac{9C}{10},$$

where the last inequality assumes $\delta \leq \frac{C}{10C + 130L\sigma^2}$

When $\eta \in [0, \eta_1] \cup [\eta_3, 1/L]$, we have

$$\mathbb{E}Q(\eta_2) \ge \left(\frac{1}{2} \|w^*\|^2 - \frac{C}{2}\right) (1 - \delta) - 13L\sigma^2 \delta$$

$$= \frac{1}{2} \|w^*\|^2 - \frac{\delta}{2} - (1 - \delta)\frac{C}{2} - 13L\sigma^2 \delta$$

$$\ge \frac{1}{2} \|w^*\|^2 - \frac{C}{2} - (1/2 + 13L\sigma^2) \delta$$

$$\ge \frac{1}{2} \|w^*\|^2 - \frac{6C}{10},$$

where the last inequality holds as long as $\delta \leq \frac{C}{5C+130L\sigma^2}$. According to Lemma 17, we know $\delta \leq \exp(-\Omega(d))$. Therefore, the conditions for δ can be satisfied as long as dis larger than certain constant.

B.3.2 Generalization for $\eta \in [0, 1/L]$

In this section, we show \hat{F}_{TbV} is point-wise close to F_{TbV} for all $\eta \in [0, 1/L]$. Recall Lemma 15 as follows.

Lemma 15. For any $1 > \epsilon > 0$, assume $d \ge c_4 \log(1/\epsilon)$ for some constant c_4 . With probability at least $1 - \epsilon$ $O(1/\epsilon) \exp(-\Omega(\epsilon^2 m))$

$$|\hat{F}_{TbV}(\eta) - F_{TbV}(\eta)| < \epsilon$$

for all $\eta \in [0, 1/L]$.

In order to prove Lemma 15, let's first show that for a fixed η with high probability $F_{TbV}(\eta)$ is close to $F_{TbV}(\eta)$. Similar as in Lemma 11, we show each $\Delta_{TbV}(\eta, P_k)$ is O(1)-subexponential. We defer its proof to Section B.3.4.

Lemma 20. Suppose σ is a constant. For any fixed $\eta \in [0, 1/L]$ and any $1 > \epsilon > 0$, with probability at least $1 - \exp(-\Omega(\epsilon^2 m)),$

$$\left|\hat{F}_{TbV}(\eta) - F_{TbV}(\eta)\right| \le \epsilon.$$

Next, we show that there exists an ϵ -net for F_{TbV} with size $O(1/\epsilon)$. By ϵ -net, we mean there exists a finite set N_{ϵ} of step size such that $|F_{TbV}(\eta) - F_{TbV}(\eta')| \le \epsilon$ for any $\eta \in [0, 1/L]$ and $\eta' \in \arg\min_{\eta \in N_{\epsilon}} |\eta - \eta'|$. We defer the proof of Lemma 21 to Section B.3.4.

Lemma 21. Suppose σ is a constant. For any $1 > \epsilon > 0$, assume $d \ge c_4 \log(1/\epsilon)$ for constant c_4 . There exists an ϵ -net N_{ϵ} for F_{TbV} with $|N_{\epsilon}| = O(1/\epsilon)$. That means, for any $\eta \in [0, 1/L]$,

$$|F_{TbV}(\eta) - F_{TbV}(\eta')| \le \epsilon,$$

for $\eta' \in \arg\min_{\eta \in N_{\epsilon}} |\eta - \eta'|$.

Next, we show that with high probability, there also exists an ϵ -net for \hat{F}_{TbV} with size $O(1/\epsilon)$.

Lemma 22. Suppose σ is a constant. For any $1 > \epsilon > 0$, assume $d \ge c_4 \log(1/\epsilon)$ for constant c_4 . With probability at least $1 - \exp(-\Omega(\epsilon^2 m))$, there exists an ϵ -net N'_{ϵ} for \hat{F}_{TbV} with $|N_{\epsilon}| = O(1/\epsilon)$. That means, for any $\eta \in [0, 1/L]$,

$$|\hat{F}_{TbV}(\eta) - \hat{F}_{TbV}(\eta')| \le \epsilon,$$

for $\eta' \in \arg\min_{\eta \in N_{\epsilon}} |\eta - \eta'|$.

Combing Lemma 20, Lemma 21 and Lemma 22, now we give the proof of Lemma 15.

Proof of Lemma 15. The proof is very similar as in Lemma 8. By Lemma 20, we know with probability at least $1 - \exp(-\Omega(\epsilon^2 m))$, $\left|\hat{F}_{TbV}(\eta) - F_{TbV}(\eta)\right| \le \epsilon$ for any fixed η . By Lemma 21 and Lemma 22, we know as long as $d = \Omega(\log(1/\epsilon))$, with probability at least $1 - \exp(-\Omega(\epsilon^2 m))$, there exists ϵ -net N_ϵ and N'_ϵ for F_{TbV} and \hat{F}_{TbV} respectively. Here, both of N_ϵ and N'_ϵ have size $O(1/\epsilon)$. According to the proofs of Lemma 21 and Lemma 22, it's not hard to verify that $N_\epsilon \cup N'_\epsilon$ is still an ϵ -net for \hat{F}_{TbV} and F_{TbV} . That means, for any $\eta \in [0, 1/L]$, we have

$$|F_{TbV}(\eta) - F_{TbV}(\eta')|, |\hat{F}_{TbV}(\eta) - \hat{F}_{TbV}(\eta')| \le \epsilon,$$

for $\eta' \in \arg\min_{\eta \in N_{\epsilon} \cup N'_{\epsilon}} |\eta - \eta'|$.

Taking a union bound over $N_{\epsilon} \cup N'_{\epsilon}$, we have with probability at least $1 - O(1/\epsilon) \exp(-\Omega(\epsilon^2 m))$,

$$\left| \hat{F}_{TbV}(\eta) - F_{TbV}(\eta) \right| \le \epsilon$$

for any $\eta \in N_{\epsilon} \cup N'_{\epsilon}$.

Overall, we know with probability at least $1 - O(1/\epsilon) \exp(-\Omega(\epsilon^2 m))$, for all $\eta \in [0, 1/L]$,

$$|F_{TbV}(\eta) - \hat{F}_{TbV}(\eta)|$$

$$\leq |F_{TbV}(\eta) - F_{TbV}(\eta')| + |\hat{F}_{TbV}(\eta) - \hat{F}_{TbV}(\eta')| + |\hat{F}_{TbV}(\eta') - F_{TbV}(\eta')|$$

$$\leq 3\epsilon.$$

where $\eta' \in \arg\min_{\eta \in N_{\epsilon} \cup N'_{\epsilon}} |\eta - \eta'|$. Changing ϵ to $\epsilon'/3$ finishes the proof.

B.3.3 Lower bounding \hat{F}_{TbV} for $\eta \in [1/L, \infty)$

In this section, we prove \hat{F}_{TbV} is large for any step size $\eta \geq 1/L$. Therefore, the optimal step size η^*_{valid} must be smaller than \hat{F}_{TbV} .

Lemma 16. Suppose σ is a large constant. Assume $t \ge c_2, d \ge c_4 \log(t)$ for some constants c_2, c_4 . With probability at least $1 - \exp(-\Omega(m))$,

$$\hat{F}_{TbV}(\eta) \ge C' \sigma^2 + \frac{1}{2} \sigma^2,$$

for all $\eta \geq 1/L$, where C' is a positive constant independent with σ .

When the step size is very large (larger than 3L), we know the GD sequence gets truncated with high probability, which immediately implies the loss is high. The proof of Lemma 23 is deferred into Section B.3.4.

Lemma 23. Assume $t \ge c_2$, $d \ge c_4$ for some constants c_2 , c_4 . With probability at least $1 - \exp(-\Omega(m))$,

$$\hat{F}_{TbV}(\eta) \geq \sigma^2$$
,

for all $\eta \in [3L, \infty)$

The case for step size within [1/L, 3L] requires more efforts. We give the proof of Lemma 24 in this section later.

Lemma 24. Suppose σ is a large constant. Assume $t \ge c_2, d \ge c_4 \log(t)$ for some constants c_2, c_4 . With probability at least $1 - \exp(-\Omega(m))$,

$$\hat{F}_{TbV}(\eta) \ge C_4 \sigma^2 + \frac{1}{2} \sigma^2,$$

for all $\eta \in [1/L, 3L]$, where C_4 is a positive constant independent with σ .

With the above two lemmas, Lemma 16 is just a combination of them.

Proof of Lemma 16. The result follows by taking a union bound and choosing $C' = \min(C_4, 1/2)$.

In the remaining of this section, we give the proof of Lemma 24. When the step size is between 1/L and 3L, if the GD sequence has a reasonable probability of diverging, we can still show the loss is high similar as before. If not, we need to show the GD sequence overfits the noise in the training set, which incurs a high loss.

Recall that the noise term is roughly $\frac{1}{2} \left\| (I - (I - \eta H_{\text{train}})^t) (X_{\text{train}})^\dagger \xi_{\text{train}} \right\|^2$. When $\eta \in [1/L, 3L]$, the eigenvalues of $I - \eta H_{\text{train}}$ in S_{train} subspace can be negative. If all the non-zero n eigenvalues of H_{train} have the same value, there exists a step size such that the eigenvalues of $I - \eta H_{\text{train}}$ in subspace S_{train} is $S_{\text{train}} = 1$. If $S_{\text{train}} = 1$ is even, the eigenvalues of $I - \eta H_{\text{train}}$ in $S_{\text{train}} = 1$ is $S_{\text{train}} = 1$.

Notice that the above problematic case cannot happen when the eigenvalues of H_{train} are spread out. Basically, when there are two different eigenvalues, there won't exist any large η that can cancel both directions at the same time. In Lemma 25, we show with constant probability, the eigenvalues of H_{train} are indeed spread out. The proof is deferred into Section B.3.4.

Lemma 25. Let the top n eigenvalues of H_{train} be $\lambda_1 \ge \cdots \ge \lambda_n$. Assume dimension $d \ge c_4$ for certain constant c_4 . There exist positive constants μ, μ', μ'' such that with probability at least μ ,

$$\lambda_{\mu'n} - \lambda_{n-\mu'n+1} \ge \mu''$$
.

Next, we utilize this variance in eigenvalues to prove that the GD sequence has to learn a constant fraction of the noise in training set.

Lemma 26. Suppose noise level σ is a large enough constant c_1 . Assume unroll length $t \ge c_2$ and dimension $d \ge c_4$ for some constants c_2, c_4 . Then, with probability at least C_1

$$\left\|B_{t,\eta}w_{train} - w^*\right\|_{H_{train}}^2 \ge C_2\sigma^2,$$

for all $\eta \in [1/L, 3L]$, where C_1, C_2 are positive constants.

Proof of Lemma 26. Let \mathcal{E}_1 be the event that $\sqrt{d}/\sqrt{L} \leq \sigma_i(X_{\text{train}}) \leq \sqrt{Ld}$ and $1/L \leq \lambda_i(H_{\text{train}}) \leq L$ for all $i \in [n]$ and $\sqrt{d}\sigma/4 \leq \|\xi_{\text{train}}\| \leq \sqrt{d}\sigma$. Let \mathcal{E}_3 be the event that $\sqrt{d}/\sqrt{L} \leq \sigma_i(X_{\text{valid}}) \leq \sqrt{Ld}$ and $1/L \leq \lambda_i(H_{\text{valid}}) \leq L$ for all $i \in [n]$ and $\sqrt{d}\sigma/4 \leq \|\xi_{\text{valid}}\| \leq \sqrt{d}\sigma$. According to Lemma 7 and Lemma 49, we know both \mathcal{E}_1 and \mathcal{E}_3 hold with probability at least $1 - \exp(-\Omega(d))$.

Let the top n eigenvalues of H_{train} be $\lambda_1 \geq \cdots \geq \lambda_n$. According to Lemma 25, assuming d is larger than certain constant, we know there exist positive constants μ_1, μ_2, μ_3 such that with probability at least $\mu_1, \lambda_{\mu_2 n} - \lambda_{n-\mu_2 n+1} \geq \mu_3$. Call this event \mathcal{E}_2 .

Let S_1 and S_2 be the span of the bottom and top $\mu_2 n$ eigenvectors of H_{train} respectively. According to Lemma 49, we know $\|\xi_{\text{train}}\| \geq \frac{\sqrt{d}}{4}\sigma$ with probability at least $1 - \exp(-\Omega(d))$. Let $P_1 \in \mathbb{R}^{n \times n}$ be a rank- $\mu_2 n$ projection matrix such that the column span of $(X_{\text{train}})^{\dagger} P_1$ is S_1 . By Johnson-Lindenstrauss Lemma, we know with probability at least $1 - \exp(-\Omega(d))$, $\|\operatorname{Proj}_{P_1}\xi_{\text{train}}\| \geq \frac{\sqrt{\mu_2}}{2}\|\xi_{\text{train}}\|$. Taking a union bound, with probability at least $1 - \exp(-\Omega(d))$, $\|\operatorname{Proj}_{P_1}\xi_{\text{train}}\| \geq \frac{\sqrt{\mu_2}d\sigma}{8}$. Similarly, we can define P_2 for the S_2 subspace and show with probability at least $1 - \exp(-\Omega(d))$, $\|\operatorname{Proj}_{P_2}\xi_{\text{train}}\| \geq \frac{\sqrt{\mu_2}d\sigma}{8}$. Call the intersection of both events as \mathcal{E}_4 , which happens with with probability at least $1 - \exp(-\Omega(d))$.

Taking a union bound, we know $\mathcal{E}_1 \cap \mathcal{E}_2 \cap \mathcal{E}_3 \cap \mathcal{E}_4$ holds with probability at least $\mu_1/2$ as long as d is larger than certain constant. Through the proof, we assume $\mathcal{E}_1 \cap \mathcal{E}_2 \cap \mathcal{E}_3 \cap \mathcal{E}_4$ holds.

Let's first lower bound $||B_{t,\eta}w_{\text{train}} - w_{\text{train}}^*||$ as follows,

$$||B_{t,\eta}w_{\text{train}} - w_{\text{train}}^*|| = ||B_{t,\eta}\left(w_{\text{train}}^* + (X_{\text{train}})^{\dagger}\xi_{\text{train}}\right) - w_{\text{train}}^*|$$

$$\geq \left(||B_{t,\eta}\left(w_{\text{train}}^* + (X_{\text{train}})^{\dagger}\xi_{\text{train}}\right)|| - 1\right)$$

Recall that we define S_1 and S_2 as the span of the bottom and top $\mu_2 n$ eigenvectors of H_{train} respectively. We rely on S_1 to lower bound $\|w_{t,\eta} - w^*\|$ when η is small and rely on S_2 when η is large.

Case 1: Let $\sigma_{\min}^{S_1}(B_{t,\eta})$ be the smallest singular value of $B_{t,\eta}$ within S_1 subspace. If $\eta \lambda_{n-\mu_2 n+1} \leq 2 - \mu_3/(2L)$, we have

$$\sigma_{\min}^{S_1}(B_{t,\eta}) \ge \min\left(1 - \left(1 - \frac{1}{L^2}\right)^t, 1 - \left(1 - \frac{\mu_3}{2L}\right)^t\right) \ge \frac{1}{2},$$

where the second inequality assumes $t \ge \max(L^2, 2L/\mu_3) \log 2$. Then, we have

$$||w_{t,\eta} - w^*|| \ge \left(\sigma_{\min}^{S_1}(B_{t,\eta}) \left(\left\| \operatorname{Proj}_{S_1}(X_{\operatorname{train}})^{\dagger} \xi_{\operatorname{train}} \right\| - 1 \right) - 1 \right)$$

$$\ge \left(\frac{1}{2} \left(\frac{\sqrt{\mu_2}\sigma}{8\sqrt{L}} - 1 \right) - 1 \right) \ge \frac{\sqrt{\mu_2}\sigma}{32\sqrt{L}},$$

where the second inequality uses $\|\operatorname{Proj}_{P_1}\xi_{\operatorname{train}}\| \geq \frac{\sqrt{\mu_2 d\sigma}}{8}$ and the last inequality assumes $\sigma \geq \frac{48\sqrt{L}}{\sqrt{\mu_2}}$.

Case 2: If $\eta \lambda_{n-\mu_2 n+1} > 2 - \mu_3/(2L)$, we have $\eta \lambda_{\mu_2 n} \ge 2 + \mu_3/(2L)$ since $\lambda_{\mu_2 n} - \lambda_{n-\mu_2 n+1} \ge \mu_3$ and $\eta \ge 1/L$. Let $\sigma_{\min}^{S_2}(B_{t,\eta})$ be the smallest singular value of $B_{t,\eta}$ within S_2 subspace. We have

$$\sigma_{\min}^{S_2}(B_{t,\eta}) \ge \left(\left(1 + \frac{\mu_3}{2L}\right)^t - 1\right) \ge \frac{1}{2},$$

where the last inequality assumes $t \ge 4L/\mu_3$. Then, similar as in Case 1, we can also prove $||w_{t,\eta} - w^*|| \ge \frac{\sqrt{\mu_2}\sigma}{32\sqrt{L}}$. Therefore, we have

$$\|B_{t,\eta}w_{\text{train}} - w^*\|_{H_{\text{train}}}^2 = \|B_{t,\eta}w_{\text{train}} - w_{\text{train}}^*\|_{H_{\text{train}}}^2 \ge \frac{1}{L}\|B_{t,\eta}w_{\text{train}} - w_{\text{train}}^*\|^2 \ge \frac{\mu_2\sigma^2}{1024L^2},$$

for all $\eta \in [1/L, 3L]$. We denote $C_1 := \mu_1/2$ and $C_2 = \frac{\mu_2}{1024L^2}$.

Before we present the proof of Lemma 24, we still need a technical lemma that shows the noise in S_{valid} concentrates at its mean. The proof of Lemma 27 is deferred into Section B.3.4.

Lemma 27. Suppose σ is constant. For any $1 > \epsilon > 0$, with probability at least $1 - O(t/\epsilon) \exp(-\Omega(\epsilon^2 d))$, $\lambda_n(H_{valid}) \ge 1/L$ and

$$\|w_{t,\eta} - w_{valid}\|_{H_{valid}}^2 \ge \|w_{t,\eta} - w^*\|_{H_{valid}}^2 + (1 - \epsilon)\sigma^2,$$

for all $\eta \in [1/L, 3L]$.

Combing the above lemmas, we give the proof of Lemma 24.

Proof of Lemma 24. According to Lemma 27, we know given $1 > \epsilon > 0$, with probability at least $1 - O(t/\epsilon) \exp(-\Omega(\epsilon^2 d))$, $\lambda_n(H_{\text{valid}}) \ge 1/L$ and $\|w_{t,\eta} - w_{\text{valid}}\|_{H_{\text{valid}}}^2 \ge \|w_{t,\eta} - w^*\|_{H_{\text{valid}}}^2 + (1 - \epsilon)\sigma^2$ for all $\eta \in [1/L, 3L]$. Call this event \mathcal{E}_1 . Suppose $\Pr[\mathcal{E}_1] \ge 1 - \delta/2$, where δ will be specifies later. For each training set $S_{\text{train}}^{(k)}$, we also define $\mathcal{E}_1^{(k)}$. By concentration, we know with probability at least $1 - \exp(-\Omega(\delta^2 m))$, $1/m \sum_{k=1}^m \mathbbm{1}\left\{\mathcal{E}_1^{(k)}\right\} \ge 1 - \delta$.

According to Lemma 26, we know there exist constants C_1, C_2 such that with probability at least C_1 , $\|B_{t,\eta}w_{\text{train}} - w^*\|_{H_{\text{train}}}^2 \geq C_2\sigma^2$ for all $\eta \in [1/L, 3L]$. Call this event \mathcal{E}_2 . For each training set $S_{\text{train}}^{(k)}$, we also define $\mathcal{E}_2^{(k)}$. By concentration, we know with probability at least $1 - \exp(-\Omega(m))$, $1/m \sum_{k=1}^m \mathbb{1}\left\{\mathcal{E}_2^{(k)}\right\} \geq C_1/2$.

For any step size $\eta \in [1/L, 3L]$, we can lower bound $\hat{F}_{TbV}(\eta)$ as follows,

$$\begin{split} \hat{F}_{TbV}(\eta) = & \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{\text{valid}}^{(k)} \right\|_{H_{\text{valid}}^{(k)}}^{2} \\ \geq & \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{\text{valid}}^{(k)} \right\|_{H_{\text{valid}}^{(k)}}^{2} \mathbbm{1} \left\{ \mathcal{E}_{1}^{(k)} \right\} \\ \geq & \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{k}^{*} \right\|_{H_{\text{valid}}}^{2} \mathbbm{1} \left\{ \mathcal{E}_{1}^{(k)} \right\} + \frac{1}{2} (1 - \epsilon) (1 - \delta) \sigma^{2} \\ \geq & \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{k}^{*} \right\|_{H_{\text{valid}}}^{2} \mathbbm{1} \left\{ \mathcal{E}_{1}^{(k)} \cap \mathcal{E}_{2}^{(k)} \right\} + \frac{1}{2} (1 - \epsilon) (1 - \delta) \sigma^{2}. \end{split}$$

As long as $\delta \leq C_1/4$, we know $\frac{1}{m} \sum_{k=1}^m \mathbb{1}\left\{\mathcal{E}_1^{(k)} \cap \mathcal{E}_2^{(k)}\right\} \geq C_1/4$. Let $\bar{\mathcal{E}}_3(\eta)$ be the event that $w_{t,\eta}^{(k)}$ gets truncated with step size η . We have

$$\begin{split} &\frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{k}^{*} \right\|_{H_{\text{valid}}}^{2} \mathbb{1} \left\{ \mathcal{E}_{1}^{(k)} \cap \mathcal{E}_{2}^{(k)} \right\} \\ = &\frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{k}^{*} \right\|_{H_{\text{valid}}}^{2} \mathbb{1} \left\{ \mathcal{E}_{1}^{(k)} \cap \mathcal{E}_{2}^{(k)} \cap \mathcal{E}_{3}^{(k)} \right\} \\ &+ \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{k}^{*} \right\|_{H_{\text{valid}}}^{2} \mathbb{1} \left\{ \mathcal{E}_{1}^{(k)} \cap \mathcal{E}_{2}^{(k)} \cap \bar{\mathcal{E}}_{3}^{(k)} \right\}. \end{split}$$

If $\frac{1}{m} \sum_{k=1}^m \mathbb{1}\left\{\mathcal{E}_1^{(k)} \cap \mathcal{E}_2^{(k)} \cap \bar{\mathcal{E}}_3^{(k)}\right\} \ge C_1/8$, we have

$$\frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{k}^{*} \right\|_{H_{\text{valid}}}^{2} \mathbb{1} \left\{ \mathcal{E}_{1}^{(k)} \cap \mathcal{E}_{2}^{(k)} \right\} \ge \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{k}^{*} \right\|_{H_{\text{valid}}}^{2} \mathbb{1} \left\{ \mathcal{E}_{1}^{(k)} \cap \mathcal{E}_{2}^{(k)} \cap \bar{\mathcal{E}}_{3}^{(k)} \right\} \\
\ge \frac{C_{1}}{8} \times \frac{9\sigma^{2}}{2} = \frac{9C_{1}\sigma^{2}}{16}.$$

Here, we lower bound $\left\|w_{t,\eta}^{(k)}-w_k^*\right\|_{H_{\text{min}}}^2$ by $9\sigma^2$ when the sequence gets truncated.

If
$$\frac{1}{m} \sum_{k=1}^{m} \mathbb{1} \left\{ \mathcal{E}_{1}^{(k)} \cap \mathcal{E}_{2}^{(k)} \cap \bar{\mathcal{E}}_{3}^{(k)} \right\} < C_{1}/8$$
, we know $\frac{1}{m} \sum_{k=1}^{m} \mathbb{1} \left\{ \mathcal{E}_{1}^{(k)} \cap \mathcal{E}_{2}^{(k)} \cap \mathcal{E}_{3}^{(k)} \right\} \ge C_{1}/8$. Then, we have

$$\begin{split} \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{k}^{*} \right\|_{H_{\text{valid}}}^{2} \mathbbm{1} \left\{ \mathcal{E}_{1}^{(k)} \cap \mathcal{E}_{2}^{(k)} \right\} \geq & \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| B_{t,\eta}^{(k)} w_{\text{train}} - w_{k}^{*} \right\|_{H_{\text{valid}}}^{2} \mathbbm{1} \left\{ \mathcal{E}_{1}^{(k)} \cap \mathcal{E}_{2}^{(k)} \cap \mathcal{E}_{3}^{(k)} \right\} \\ \geq & \frac{C_{1}}{8} \times \frac{C_{2}\sigma^{2}}{2} = \frac{C_{1}C_{2}\sigma^{2}}{16} \end{split}$$

Letting $C_3 = \min(\frac{9C_1}{16}, \frac{C_1C_2}{16})$, we then have

$$\hat{F}_{TbV}(\eta) \ge C_3 \sigma^2 + \frac{1}{2} (1 - \epsilon)(1 - \delta)\sigma^2 \ge \frac{C_3 \sigma^2}{2} + \frac{1}{2} \sigma^2,$$

where the last inequality chooses $\delta = \epsilon = C_3/2$. In order for $\Pr[\mathcal{E}_1] \ge 1 - \delta/2$, we only need $d \ge c_4 \log(t)$ for some constant c_4 . Replacing $C_3/2$ by C_4 finishes the proof.

B.3.4 Proofs of Technical Lemmas

Proof of Lemma 18. We first show that for a fixed $\eta \in [0, 1/L]$, the crossing term $|\langle B_{t,\eta} w_{\text{train}}^* - w^*, B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}} \rangle|$ is small with high probability. We can write down the crossing term as follows:

$$\left\langle B_{t,\eta} w_{\text{train}}^* - w^*, B_{t,\eta} (X_{\text{train}})^\dagger \xi_{\text{train}} \right\rangle = \left\langle [(X_{\text{train}})^\dagger]^\top B_{t,\eta} (B_{t,\eta} w_{\text{train}}^* - w^*), \xi_{\text{train}} \right\rangle.$$

Noticing that ξ_{train} is independent with $[(X_{\text{train}})^{\dagger}]^{\top}B_{t,\eta}(B_{t,\eta}w_{\text{train}}^* - w^*)$, we will use Hoeffding's inequality to bound $|\langle B_{t,\eta}w_{\text{train}}^* - w^*, B_{t,\eta}(X_{\text{train}})^{\dagger}\xi_{\text{train}}\rangle|$. According to Lemma 7, we know with probability at least $1 - \exp(-\Omega(d))$, $\sqrt{d}/\sqrt{L} \leq \sigma_i(X_{\text{train}}) \leq \sqrt{Ld}$ and $1/L \leq \lambda_i(H_{\text{train}}) \leq L$ for all $i \in [n]$ with L = 100. Since $\eta \leq 1/L$, we know $\|B_{t,\eta}\| = \|I - (I - \eta H_{\text{train}})^t\| \leq 1$. Therefore, we have

$$\left\| [(X_{\text{train}})^{\dagger}]^{\top} B_{t,\eta} (B_{t,\eta} w_{\text{train}}^* - w^*) \right\| \leq \frac{2\sqrt{L}}{\sqrt{d}},$$

for any $\eta \in [0, 1/L]$. Then, for any $\epsilon > 0$, by Hoeffding's inequality, with probability at least $1 - \exp(-\Omega(\epsilon^2 d))$,

$$\left|\left\langle B_{t,\eta}w_{\text{train}}^* - w^*, B_{t,\eta}(X_{\text{train}})^{\dagger} \xi_{\text{train}}\right\rangle\right| \leq \epsilon.$$

Next, we construct an ϵ -net on η and show the crossing term is small for all $\eta \in [0, 1/L]$. Let

$$g(\eta) := \langle B_{t,\eta} w_{\text{train}}^* - w^*, B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}} \rangle.$$

We compute the derivative of $g(\eta)$ as follows:

$$\begin{split} g'(\eta) = & \left\langle t H_{\text{train}} (I - \eta H_{\text{train}})^{t-1} w_{\text{train}}^*, B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}} \right\rangle \\ & + \left\langle B_{t,\eta} w_{\text{train}}^* - w^*, t H_{\text{train}} (I - \eta H_{\text{train}})^{t-1} (X_{\text{train}})^{\dagger} \xi_{\text{train}} \right\rangle \end{split}$$

By Lemma 49, we know with probability at least $1 - \exp(-\Omega(d))$, $\|\xi_{\text{train}}\| \leq \sqrt{d}\sigma$. Therefore,

$$|g'(\eta)| \le L^{1.5} t \left(1 - \frac{\eta}{L}\right)^{t-1} \sigma + 2L^{1.5} t \left(1 - \frac{\eta}{L}\right)^{t-1} \sigma = 3L^{1.5} t \left(1 - \frac{\eta}{L}\right)^{t-1} \sigma.$$

We can control $|g'(\eta)|$ in different regimes:

- For $\eta \in [0, \frac{L}{t-1}]$, we have $|g'(\eta)| \leq 3L^{1.5}t\sigma$.
- Given any $1 \le i \le \log t 1$, for any $\eta \in (\frac{iL}{t-1}, \frac{(i+1)L}{t-1}]$, we have $|g'(\eta)| \le \frac{3L^{1.5}t\sigma}{e^i}$.
- For any $\eta \in (\frac{L \log t}{t-1}, 1/L]$, we have $|g'(\eta)| \leq 3L^{1.5}\sigma$.

Fix any $\epsilon > 0$, we know there exists an ϵ -net N_{ϵ} with size

$$|N_{\epsilon}| = \frac{1}{\epsilon} \left(\frac{L}{t-1} \sum_{i=0}^{\log t - 1} \frac{3L^{1.5}t\sigma}{e^i} + \left(\frac{1}{L} - \frac{L\log t}{t-1} \right) 3L^{1.5}\sigma \right)$$

$$\leq \frac{1}{\epsilon} \left(\frac{3eL^{2.5}t\sigma}{t-1} + 3\sqrt{L}\sigma \right) = O(\frac{1}{\epsilon})$$

such that for any $\eta \in [0, 1/L]$, there exists $\eta' \in N_{\epsilon}$ with $|g(\eta) - g(\eta')| \le \epsilon$. Note that L = 100 and σ is a constant. Taking a union bound over N_{ϵ} and all the other bad events, we have with probability at least $1 - \exp(-\Omega(d)) - O(1/\epsilon) \exp(-\Omega(\epsilon^2 d))$, for all $\eta \in [0, 1/L]$,

$$\left|\left\langle B_{t,\eta} w_{\text{train}}^* - w^*, B_{t,\eta}(X_{\text{train}})^{\dagger} \xi_{\text{train}} \right\rangle\right| \leq \epsilon + \epsilon = 2\epsilon.$$

As long as $1 > \epsilon > 0$, this happens with probability at least $1 - O(1/\epsilon) \exp(-\Omega(\epsilon^2 d))$. Replacing ϵ by $\epsilon'/2$ finishes the proof.

Proof of Lemma 19. According to Lemma 7, we know with probability at least $1 - \exp(-\Omega(d))$, $1/L \le \lambda_i(H_{\text{train}}) \le L$ for all $i \in [n]$ with L = 100. We can lower bound $\langle B_{t,\eta} w_{\text{train}}^*, w^* \rangle$ as follows,

$$\langle B_{t,\eta} w_{\text{train}}^*, w^* \rangle = \langle \left(I - (I - \eta H_{\text{train}})^t \right) w_{\text{train}}^*, w_{\text{train}}^* \rangle$$

$$\geq \lambda_{\min} \left(I - (I - \eta H_{\text{train}})^t \right) \|w_{\text{train}}^*\|^2$$

$$\geq \left(1 - \exp\left(-\frac{\eta t}{L} \right) \right) \|w_{\text{train}}^*\|^2.$$

By Johnson-Lindenstrauss lemma (Lemma 53), we know with probability at least $1 - 2\exp(-c\epsilon^2 d/4)$,

$$||w_{\text{train}}^*|| \ge \frac{1}{2}(1-\epsilon) ||w^*|| = \frac{1}{2}(1-\epsilon).$$

Then, we know with probability at least $1 - 2\exp(-c\epsilon^2 d/4) - \exp(-\Omega(d))$,

$$\langle B_{t,\eta} w_{\text{train}}^*, w^* \rangle \ge \left(1 - \exp\left(-\frac{\eta t}{L} \right) \right) \|w_{\text{train}}^*\|^2$$

$$\ge \left(1 - \exp\left(-\frac{\eta t}{L} \right) \right) \frac{1}{4} (1 - \epsilon)^2$$

$$\ge \frac{1 - 2\epsilon}{4} \left(1 - \exp\left(-\frac{\eta t}{L} \right) \right)$$

Since $e^x \le 1 - x + x^2/2$ for any $x \le 0$, we know $\exp(-\eta t/L) \le 1 - \eta t/L + \eta^2 t^2/(2L^2)$. For any $\eta \le L/t$, we have $\exp(-\eta t/L) \le 1 - \eta t/(2L)$. Then with probability at least $1 - 2\exp(-c\epsilon^2 d/4) - \exp(-\Omega(d))$,

$$\langle B_{t,\eta} w_{\text{train}}^*, w^* \rangle \ge \frac{1 - 2\epsilon}{4} \frac{\eta t}{2L}$$

 $\ge \frac{\eta t}{16L},$

where the second inequality holds by choosing $\epsilon = 1/4$.

Proof of Lemma 20. Recall that

$$\hat{F}_{TbV}(\eta) := \frac{1}{m} \sum_{k=1}^{m} \Delta_{TbV}(\eta, P_k)$$

For each individual loss function $\Delta_{TbV}(\eta, P_k)$, we have

$$\begin{split} \Delta_{TbV}(\eta, P_k) &= \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w^* - (X_{\text{valid}}^{(k)})^{\dagger} \xi_{\text{valid}}^{(k)} \right\|_{H_{\text{valid}}^{(k)}}^2 \\ &= \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w^* \right\|_{H_{\text{valid}}^{(k)}}^2 + \frac{1}{2n} \left\| \xi_{\text{valid}}^{(k)} \right\|^2 + \left\langle w_{t,\eta}^{(k)} - w^*, \frac{1}{n} (X_{\text{valid}}^{(k)})^{\top} \xi_{\text{valid}}^{(k)} \right\rangle \\ &\leq \frac{25L\sigma^2}{2} \left\| H_{\text{valid}}^{(k)} \right\| + \frac{1}{2n} \left\| \xi_{\text{valid}}^{(k)} \right\|^2 + 5\sqrt{L}\sigma \left(\frac{1}{\sqrt{n}} \left\| X_{\text{valid}}^{(k)} \right\| \right) \left(\frac{1}{\sqrt{n}} \left\| \xi_{\text{valid}}^{(k)} \right\| \right) \end{split}$$

We can write $\left\|H_{\mathrm{valid}}^{(k)}\right\|$ as $\sigma_{\mathrm{max}}^2(\frac{1}{\sqrt{n}}X_{\mathrm{valid}}^{(k)})$. According to Lemma 51, we know $\sigma_{\mathrm{max}}(X_{\mathrm{valid}}^{(k)}) - \mathbb{E}\sigma_{\mathrm{max}}(X_{\mathrm{valid}}^{(k)})$ is O(1)-subgaussian, which implies that $\sigma_{\mathrm{max}}(\frac{1}{\sqrt{n}}X_{\mathrm{valid}}^{(k)}) - \mathbb{E}\sigma_{\mathrm{max}}(\frac{1}{\sqrt{n}}X_{\mathrm{valid}}^{(k)})$ is $O(1/\sqrt{d})$ -subgaussian. Since $\mathbb{E}\sigma_{\mathrm{max}}(\frac{1}{\sqrt{n}}X_{\mathrm{valid}}^{(k)})$ is a constant, we know $\sigma_{\mathrm{max}}(\frac{1}{\sqrt{n}}X_{\mathrm{valid}}^{(k)})$ is O(1)-subgaussian and $\sigma_{\mathrm{max}}^2(\frac{1}{\sqrt{n}}X_{\mathrm{valid}}^{(k)})$ is O(1)-subexponential. Similarly, we know both $\frac{1}{2n}\left\|\xi_{\mathrm{valid}}^{(k)}\right\|^2$ and $\left(\frac{1}{\sqrt{n}}\left\|X_{\mathrm{valid}}^{(k)}\right\|\right)\left(\frac{1}{\sqrt{n}}\left\|\xi_{\mathrm{valid}}^{(k)}\right\|\right)$ are O(1)-subexponential. This further implies that $\Delta_{TbV}(\eta, P_k)$ is O(1)-subexponential. Therefore, \hat{F}_{TbV} is the average of m i.i.d. O(1)-subexponential random variables. By standard concentration inequality, we know for any $1 > \epsilon > 0$, with probability at least $1 - \exp(-\Omega(\epsilon^2 m))$,

$$\left|\hat{F}_{TbV}(\eta) - F_{TbV}(\eta)\right| \le \epsilon.$$

Proof of Lemma 21. Recall that

 $F_{TbV}(\eta) = \mathbb{E}\frac{1}{2} \|w_{t,\eta} - w^*\|^2 + \sigma^2/2.$

We only need to construct an ϵ -net for $\mathbb{E}\frac{1}{2}\|w_{t,\eta}-w^*\|^2$. Let \mathcal{E} be the event that $\sqrt{d}/\sqrt{L} \leq \sigma_i(X_{\text{train}}) \leq \sqrt{Ld}$ and $1/L \leq \lambda_i(H_{\text{train}}) \leq L$ for all $i \in [n]$ and $\|\xi_{\text{train}}\| \leq \sqrt{d}\sigma$. We have

$$\mathbb{E}\frac{1}{2}\left\|w_{t,\eta} - w^*\right\|^2 = \mathbb{E}\left[\frac{1}{2}\left\|w_{t,\eta} - w^*\right\|^2 |\mathcal{E}\right] \Pr[\mathcal{E}] + \mathbb{E}\left[\frac{1}{2}\left\|w_{t,\eta} - w^*\right\|^2 |\bar{\mathcal{E}}\right] \Pr[\bar{\mathcal{E}}]$$

We first construct an ϵ -net for $\mathbb{E}\left[\frac{1}{2}\|w_{t,\eta}-w^*\|^2|\mathcal{E}\right]\Pr[\mathcal{E}]$. Let $Q(\eta):=\frac{1}{2}\|w_{t,\eta}-w^*\|^2$. Fix a training set S_{train} under which event \mathcal{E} holds. We show that $Q(\eta)$ has desirable lipschitz property.

The derivative of $Q(\eta)$ can be computed as follows,

$$Q'(\eta) = \left\langle t H_{\text{train}} (I - \eta H_{\text{train}})^{t-1} w_{\text{train}}, w_{t,\eta} - w^* \right\rangle.$$

Conditioning on \mathcal{E} , we have

$$|Q'(\eta)| = O(1)t(1 - \frac{\eta}{L})^{t-1}.$$

Therefore, we have

$$\left| \frac{\partial}{\partial \eta} \mathbb{E} \left[\frac{1}{2} \| w_{t,\eta} - w^* \|^2 | \mathcal{E} \right] \Pr[\mathcal{E}] \right| = O(1)t(1 - \frac{\eta}{L})^{t-1}.$$

Similar as in Lemma 18, for any $\epsilon > 0$, we know there exists an ϵ -net N_{ϵ} with size $O(1/\epsilon)$ such that for any $\eta \in [0, 1/L]$,

$$\left| \mathbb{E}\left[\frac{1}{2} \| w_{t,\eta} - w^* \|^2 | \mathcal{E} \right] \Pr[\mathcal{E}] - \mathbb{E}\left[\frac{1}{2} \| w_{t,\eta'} - w^* \|^2 | \mathcal{E} \right] \Pr[\mathcal{E}] \right| \le \epsilon$$

for $\eta' \in \arg\min_{\eta \in N_{\epsilon}} |\eta - \eta'|$.

Suppose the probability of $\bar{\mathcal{E}}$ is δ . We have

$$\mathbb{E}\left[\frac{1}{2}\|w_{t,\eta} - w^*\|^2 |\bar{\mathcal{E}}\right] \Pr[\bar{\mathcal{E}}] \le \frac{25L\sigma^2}{2}\delta \le \epsilon,$$

where the last inequality assumes $\delta \leq \frac{2\epsilon}{25L\sigma^2}$. According to Lemma 7 and Lemma 49, we know $\delta := \Pr[\bar{\mathcal{E}}] \leq \exp(-\Omega(d))$. Therefore, given any $\epsilon > 0$, there exists constant c_4 such that $\delta \leq \frac{2\epsilon}{25L\sigma^2}$ as long as $d \geq c_4 \log(1/\epsilon)$.

Overall, for any $\epsilon > 0$, as long as $d = \Omega(\log(1/\epsilon))$, there exists N_{ϵ} with size $O(1/\epsilon)$ such that for any $\eta \in [0, 1/L]$, $|F_{TbV}(\eta) - F_{TbV}(\eta')| \leq 3\epsilon$ for $\eta' \in \arg\min_{\eta \in N_{\epsilon}} |\eta - \eta'|$. Changing ϵ to $\epsilon'/3$ finishes the proof.

Proof of Lemma 22. For each $k \in [m]$, let \mathcal{E}_k be the event that $\sqrt{d}/\sqrt{L} \leq \sigma_i(X_{\text{train}}^{(k)}) \leq \sqrt{Ld}$ for any $i \in [n]$ and $\left\|\xi_{\text{train}}^{(k)}\right\| \leq \sqrt{d}\sigma$. Then, we can write the empirical meta objective as follows,

$$\hat{F}_{TbV}(\eta) := \frac{1}{m} \sum_{k=1}^{m} \Delta_{TbT}(\eta, P_k) \mathbb{1}_{\mathcal{E}_k} + \frac{1}{m} \sum_{k=1}^{m} \Delta_{TbT}(\eta, P_k) \mathbb{1}_{\bar{\mathcal{E}}_k}.$$

Similar as Lemma 21, we will show that the first term has desirable Lipschitz property and the second term is small. Now, let's focus on the first term $\frac{1}{m}\sum_{k=1}^m \Delta_{TbT}(\eta,P_k)\mathbb{1}_{\mathcal{E}_k}$. Recall that

$$\begin{split} \Delta_{TbT}(\eta, P_k) = & \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{\text{valid}}^{(k)} \right\|_{H_{\text{valid}}^{(k)}}^2 \\ = & \frac{1}{2} \left\| B_{t,\eta}^{(k)} w_{\text{train}}^{(k)} - w^* - (X_{\text{valid}}^{(k)})^\dagger \xi_{\text{valid}}^{(k)} \right\|_{H_{\text{valid}}^{(k)}}^2. \end{split}$$

Computing the derivative of $\Delta_{TbT}(\eta, P_k)$ in terms of η , we have

$$\frac{\partial}{\partial \eta} \Delta_{TbT}(\eta, P_k) = \left\langle tH_{\text{train}}^{(k)} (I - \eta H_{\text{train}}^{(k)})^{t-1} w_{\text{train}}^{(k)}, H_{\text{valid}}^{(k)} \left(w_{t,\eta}^{(k)} - w^* - (X_{\text{valid}}^{(k)})^\dagger \xi_{\text{valid}}^{(k)} \right) \right\rangle$$

Conditioning on \mathcal{E}_k , we can bound the derivative,

$$\left| \frac{\partial}{\partial \eta} \Delta_{TbT}(\eta, P_k) \right| = O(1)t \left(1 - \frac{\eta}{L} \right)^{t-1} \left(\left\| H_{\text{valid}}^{(k)} \right\| + \left(\frac{1}{\sqrt{d}} \left\| X_{\text{valid}}^{(k)} \right\| \right) \left(\frac{1}{\sqrt{d}} \left\| \xi_{\text{valid}}^{(k)} \right\| \right) \right).$$

Therefore, we have

$$\left| \frac{1}{m} \sum_{k=1}^{m} \frac{\partial}{\partial \eta} \Delta_{TbT}(\eta, P_k) \mathbb{1}_{\mathcal{E}_k} \right| = O(1)t \left(1 - \frac{\eta}{L} \right)^{t-1} \frac{1}{m} \sum_{k=1}^{m} \left(\left\| H_{\text{valid}}^{(k)} \right\| + \left(\frac{1}{\sqrt{d}} \left\| X_{\text{valid}}^{(k)} \right\| \right) \left(\frac{1}{\sqrt{d}} \left\| \xi_{\text{valid}}^{(k)} \right\| \right) \right).$$

Similar as in Lemma 20, we know both $\left\|H_{\mathrm{valid}}^{(k)}\right\|$ and $\left(\frac{1}{\sqrt{d}}\left\|X_{\mathrm{valid}}^{(k)}\right\|\right)\left(\frac{1}{\sqrt{d}}\left\|\xi_{\mathrm{valid}}^{(k)}\right\|\right)$ are O(1)-subexponential. Therefore, we know with probability at least $1-\exp(-\Omega(m))$, $\frac{1}{m}\sum_{k=1}^{m}\left(\left\|H_{\mathrm{valid}}^{(k)}\right\|+\left(\frac{1}{\sqrt{d}}\left\|X_{\mathrm{valid}}^{(k)}\right\|\right)\left(\frac{1}{\sqrt{d}}\left\|\xi_{\mathrm{valid}}^{(k)}\right\|\right)\right)=O(1)$. This further shows that with probability at least $1-\exp(-\Omega(m))$,

$$\left| \frac{1}{m} \sum_{k=1}^{m} \frac{\partial}{\partial \eta} \Delta_{TbT}(\eta, P_k) \mathbb{1}_{\mathcal{E}_k} \right| = O(1)t \left(1 - \frac{\eta}{L} \right)^{t-1}.$$

Similar as in Lemma 18, we can show that for any $\epsilon > 0$, there exists an ϵ -net with size $O(1/\epsilon)$ for $\frac{1}{m} \sum_{k=1}^m \Delta_{TbT}(\eta, P_k) \mathbb{1}_{\mathcal{E}_k}$. Next, we show that the second term $\frac{1}{m} \sum_{k=1}^m \Delta_{TbT}(\eta, P_k) \mathbb{1}_{\bar{\mathcal{E}}_k}$ is small with high probability. According to the proof in Lemma 20, we know

$$\Delta_{TbT}(\eta, P_k) = O(1) \left(\left\| H_{\text{valid}}^{(k)} \right\| + \frac{1}{d} \left\| \xi_{\text{valid}}^{(k)} \right\|^2 + \left(\frac{1}{\sqrt{d}} \left\| X_{\text{valid}}^{(k)} \right\| \right) \left(\frac{1}{\sqrt{d}} \left\| \xi_{\text{valid}}^{(k)} \right\| \right) \right)$$

Therefore, there exists constant C such that

$$\frac{1}{m} \sum_{k=1}^{m} \Delta_{TbT}(\eta, P_k) \mathbb{1}_{\bar{\mathcal{E}}_k} \leq C \frac{1}{m} \sum_{k=1}^{m} \left(\left\| H_{\text{valid}}^{(k)} \right\| + \frac{1}{d} \left\| \xi_{\text{valid}}^{(k)} \right\|^2 + \left(\frac{1}{\sqrt{d}} \left\| X_{\text{valid}}^{(k)} \right\| \right) \left(\frac{1}{\sqrt{d}} \left\| \xi_{\text{valid}}^{(k)} \right\| \right) \right) \mathbb{1}_{\bar{\mathcal{E}}_k}.$$

It's not hard to verify that $\left(\left\|H_{\mathrm{valid}}^{(k)}\right\| + \frac{1}{d}\left\|\xi_{\mathrm{valid}}^{(k)}\right\|^2 + \left(\frac{1}{\sqrt{d}}\left\|X_{\mathrm{valid}}^{(k)}\right\|\right)\left(\frac{1}{\sqrt{d}}\left\|\xi_{\mathrm{valid}}^{(k)}\right\|\right)\right)\mathbb{1}_{\bar{\mathcal{E}}_k}$ is O(1)-subexponential. Suppose the expectation of $\left(\left\|H_{\mathrm{valid}}^{(k)}\right\| + \frac{1}{d}\left\|\xi_{\mathrm{valid}}^{(k)}\right\|^2 + \left(\frac{1}{\sqrt{d}}\left\|X_{\mathrm{valid}}^{(k)}\right\|\right)\left(\frac{1}{\sqrt{d}}\left\|\xi_{\mathrm{valid}}^{(k)}\right\|\right)\right)$ is μ , which is a constant. Suppose

the probability of $\bar{\mathcal{E}}_k$ be δ . We know the expectation of $\left(\left\|H_{\text{valid}}^{(k)}\right\| + \frac{1}{d}\left\|\xi_{\text{valid}}^{(k)}\right\|^2 + \left(\frac{1}{\sqrt{d}}\left\|X_{\text{valid}}^{(k)}\right\|\right)\left(\frac{1}{\sqrt{d}}\left\|\xi_{\text{valid}}^{(k)}\right\|\right)\right)\mathbb{1}_{\bar{\mathcal{E}}_k}$ is $\mu\delta$ due to independence. By standard concentration inequality, for any $1 > \epsilon > 0$, with probability at least $1 - \exp(-\Omega(\epsilon^2 m))$,

$$C\frac{1}{m}\sum_{k=1}^{m} \left(\left\| H_{\text{valid}}^{(k)} \right\| + \frac{1}{d} \left\| \xi_{\text{valid}}^{(k)} \right\|^2 + \left(\frac{1}{\sqrt{d}} \left\| X_{\text{valid}}^{(k)} \right\| \right) \left(\frac{1}{\sqrt{d}} \left\| \xi_{\text{valid}}^{(k)} \right\| \right) \right) \mathbb{1}_{\bar{\mathcal{E}}_k} \leq C\mu\delta + C\epsilon \leq (C+1)\epsilon,$$

where the second inequality assumes $\delta \le \epsilon/(C\mu)$. By Lemma 7 and Lemma 49, we know $\delta \le \exp(-\Omega(d))$. Therefore, as long as $d \ge c_4 \log(1/\epsilon)$ for some constant c_4 , we have $\delta \le \epsilon/(C\mu)$.

Overall, we know that as long as $d \ge c_4 \log(1/\epsilon)$, with probability at least $1 - \exp(-\Omega(\epsilon^2 m))$, there exists N'_{ϵ} with $|N'_{\epsilon}| = O(1/\epsilon)$ such that for any $\eta \in [0, 1/L]$,

$$|\hat{F}_{TbV}(\eta) - \hat{F}_{TbV}(\eta')| \le (2C+3)\epsilon,$$

for $\eta' \in \arg\min_{\eta \in N_{\epsilon}} |\eta - \eta'|$. Changing ϵ to $\epsilon'/(2C+3)$ finishes the proof.

Proof of Lemma 23. Let \mathcal{E}_1 be the event that $\sqrt{d}/\sqrt{L} \leq \sigma_i(X_{\text{train}}) \leq \sqrt{Ld}$ and $1/L \leq \lambda_i(H_{\text{train}}) \leq L$ for all $i \in [n]$ and $\sqrt{d}\sigma/4 \leq \|\xi_{\text{train}}\| \leq \sqrt{d}\sigma$. Let \mathcal{E}_2 be the event that $\sqrt{d}/\sqrt{L} \leq \sigma_i(X_{\text{valid}}) \leq \sqrt{Ld}$ and $1/L \leq \lambda_i(H_{\text{valid}}) \leq L$ for all $i \in [n]$ and $\sqrt{d}\sigma/4 \leq \|\xi_{\text{valid}}\| \leq \sqrt{d}\sigma$. According to Lemma 7 and Lemma 49, we know both \mathcal{E}_1 and \mathcal{E}_2 hold with probability at least $1 - \exp(-\Omega(d))$. Assuming $d \geq c_4$ for certain constant c_4 , we know $\Pr[\mathcal{E}_1 \cap \mathcal{E}_2] \geq 2/3$. Also define $\mathcal{E}_1^{(k)}$ and $\mathcal{E}_2^{(k)}$ on each training set $S_{\text{train}}^{(k)}$. By concentration, we know with probability at least $1 - \exp(-\Omega(m))$,

$$\frac{1}{m} \sum_{k=1}^{m} \mathbb{1}\left\{\mathcal{E}_1^{(k)} \cap \mathcal{E}_2^{(k)}\right\} \ge \frac{1}{2}.$$

It's easy to verify that conditioning on \mathcal{E}_1 , the GD sequence always exceeds the norm threshold and gets truncated for $\eta \geq 3L$ as long as t is larger than certain constant. We can lower bound \hat{F}_{TbV} for any $\eta \geq 3L$ as follows,

$$\begin{split} \hat{F}_{TbV}(\eta) = & \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{\text{valid}}^{(k)} \right\|_{H_{\text{valid}}^{(k)}}^{2} \\ \geq & \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{\text{valid}}^{(k)} \right\|_{H_{\text{valid}}^{(k)}}^{2} \mathbb{1} \left\{ \mathcal{E}_{1} \cap \mathcal{E}_{2} \right\} \geq 2\sigma^{2} \frac{1}{2} = \sigma^{2}, \end{split}$$

where the last inequality lower bounds $\left\|w_{t,\eta}^{(k)}-w_{\mathrm{valid}}^{(k)}\right\|_{H^{(k)}}^2$ by $2\sigma^2$ when $w_{t,\eta}^{(k)}$ gets truncated.

Proof of Lemma 25. We first show that with constant probability in X_{train} , the variance of the eigenvalues of H_{train} is lower bounded by a constant. Let $\bar{\lambda}$ be $1/n\sum_{i=1}^n \lambda_i$. Specifically, we show $1/n\sum_{i=1}^n \lambda_i^2 - \bar{\lambda}^2$ is lower bounded by a constant

Let's first compute the variance of the eigenvalues in expectation. Let the *i*-th row of X_{train} be x_i^{\top} . We have,

$$\mathbb{E}\left[\bar{\lambda}^{2}\right] = \frac{1}{n^{2}} \mathbb{E}\left[\left(\operatorname{tr}\left(\frac{1}{n}X_{\operatorname{train}}^{\top}X_{\operatorname{train}}\right)\right)^{2}\right] = \frac{1}{n^{4}} \mathbb{E}\left[\left(\sum_{i=1}^{n}\|x_{i}\|^{2}\right)^{2}\right]$$

$$= \frac{1}{n^{4}} \sum_{i=1}^{n} \mathbb{E}\|x_{i}\|^{4} + \frac{1}{n^{4}} \sum_{1 \leq i \neq j \leq n} \mathbb{E}\|x_{i}\|^{2} \|x_{j}\|^{2}$$

$$= \frac{1}{n^{4}} \left(nd(d+2) + n(n-1)d^{2}\right) = \frac{d^{2}}{n^{2}} + \frac{2d}{n^{3}}.$$

Similarly, we compute $\mathbb{E}\left[1/n\sum_{i=1}^n \lambda_i^2\right]$ as follows,

$$\mathbb{E}\left[\frac{1}{n}\sum_{i=1}^{n}\lambda_{i}^{2}\right] = \frac{1}{n^{3}}\mathbb{E}\left[\operatorname{tr}\left(X_{\operatorname{train}}^{\top}X_{\operatorname{train}}X_{\operatorname{train}}^{\top}X_{\operatorname{train}}\right)\right]$$

$$= \frac{1}{n^{3}}\sum_{i=1}^{n}\mathbb{E}\left\|x_{i}\right\|^{4} + \frac{1}{n^{3}}\sum_{1\leq i\neq j\leq n}\mathbb{E}\left\langle x_{i}, x_{j}\right\rangle^{2}$$

$$= \frac{1}{n^{3}}\left(nd(d+2) + n(n-1)d\right) = \frac{d^{2}}{n^{2}} + \frac{d}{n} + \frac{d}{n^{2}}$$

Therefore, we have

$$\mathbb{E}\left[\frac{1}{n}\sum_{i=1}^n \lambda_i^2 - \bar{\lambda}^2\right] = \frac{d}{n} + \frac{d}{n^2} - \frac{2d}{n^3} \ge \frac{d}{n} \ge \frac{4}{3},$$

where the first inequality assumes $n \geq 2$ and the last inequality uses $n \leq \frac{3d}{4}$. Since $n \geq \frac{1}{4}d$, we know $n \geq 2$ as long as $d \geq 8$.

Let \mathcal{E} be the event that $\sqrt{d}/\sqrt{L} \leq \sigma_i(X_{\text{train}}) \leq \sqrt{Ld}$ and $1/L \leq \lambda_i(H_{\text{train}}) \leq L$ for $i \in [n]$ with L = 100. According to Lemma 7, we know \mathcal{E} happens with probability at least $1 - \exp(-\Omega(d))$. Let $\mathbb{1}\{\mathcal{E}\}$ be the indicator

function for event \mathcal{E} . Next we show that $\mathbb{E}[1/n\sum_{i=1}^n(\lambda_i-\bar{\lambda})^2\mathbb{1}\{\mathcal{E}\}]$ is also lower bounded. It's clear that $\mathbb{E}\left[\bar{\lambda}^2\mathbb{1}\{\mathcal{E}\}\right]$ is upper bounded by $\mathbb{E}\left[\bar{\lambda}^2\right]$. In order to lower bound $\mathbb{E}\left[\frac{1}{n}\sum_{i=1}^n\lambda_i^2\mathbb{1}\{\mathcal{E}\}\right]$, we first show that $\mathbb{E}\left[\frac{1}{n}\sum_{i=1}^n\lambda_i^2\mathbb{1}\{\bar{\mathcal{E}}\}\right]$ is small. We can decompose $\mathbb{E}\left[\frac{1}{n}\sum_{i=1}^n\lambda_i^2\mathbb{1}\{\bar{\mathcal{E}}\}\right]$ into two parts,

$$\mathbb{E}\left[\frac{1}{n}\sum_{i=1}^n\lambda_i^2\mathbbm{1}\left\{\bar{\mathcal{E}}\right\}\right] = \mathbb{E}\left[\frac{1}{n}\sum_{i=1}^n\lambda_i^2\mathbbm{1}\left\{\bar{\mathcal{E}} \text{ and } \lambda_1 \leq L\right\}\right] + \mathbb{E}\left[\frac{1}{n}\sum_{i=1}^n\lambda_i^2\mathbbm{1}\left\{\lambda_1 > L\right\}\right].$$

The first term can be bounded by $L^2 \Pr[\bar{\mathcal{E}}]$. Since $\Pr[\bar{\mathcal{E}}] \leq \exp(-\Omega(d))$, we know the first term is at most 1/6 as long as d is larger than certain constant. The second term can be bounded by $\mathbb{E}[\lambda_1^2 \mathbb{I}\{\lambda_1 > L\}]$. According to Lemma 52, we know $\Pr[\lambda_1 \geq L + t] \leq \exp(-\Omega(dt))$. Then, it's not hard to verify that $\mathbb{E}\left[\lambda_1^2\mathbbm{1}\left\{\lambda_1 > L\right\}\right] = O(1/d)$ that is bounded by 1/6 as long as d is larger than certain constant. Overall, we know $\mathbb{E}\left[\frac{1}{n}\sum_{i=1}^n \lambda_i^2\mathbbm{1}\left\{\mathcal{E}\right\}\right] \geq \mathbb{E}\left[\frac{1}{n}\sum_{i=1}^n \lambda_i^2\right] - 1/3$. Combing with the upper bounds on $\mathbb{E}\left[\bar{\lambda}^2\mathbbm{1}\left\{\mathcal{E}\right\}\right]$, we have $\mathbb{E}\left[\frac{1}{n}\sum_{i=1}^n (\lambda_i - \bar{\lambda})^2\mathbbm{1}\left\{\mathcal{E}\right\}\right] \geq 1$.

Since conditioning on \mathcal{E} , λ_i is bounded by L for all $i \in [n]$. In order to make $\mathbb{E}\left[\frac{1}{n}\sum_{i=1}^n(\lambda_i-\bar{\lambda})^2\mathbb{1}\left\{\mathcal{E}\right\}\right]$ lower bounded by one, there must exist positive constants μ_1, μ_2 such that with probability at least μ_1, \mathcal{E} holds and $\frac{\frac{1}{n}\sum_{i=1}^n(\lambda_i-\bar{\lambda})^2\geq\mu_2.}{\text{Since }\frac{1}{n}\sum_{i=1}^n(\lambda_i-\bar{\lambda})^2\geq\mu_2\text{ and }\lambda_i\leq L\text{ for all }i\in[n],\text{ we know there exists a subset of eigenvalues }S\subset\{\lambda_i\}_1^n$

with size $\mu_3 n$ such that $|\lambda_i - \bar{\lambda}| \ge \mu_4$ for all $\lambda_i \in S$, where μ_3, μ_4 are both positive constants.

If at least half of eigenvalues in S are larger than $\bar{\lambda}$, we know at least $\frac{\mu_3 \mu_4 n}{2L}$ number of eigenvalues are smaller than $\bar{\lambda}$. Otherwise, the expectation of the eigenvalues will be larger than $\bar{\lambda}$, which contradicts the definition of $\bar{\lambda}$. Similarly, if at least half of eigenvalues in S are smaller than $\bar{\lambda}$, we know at least $\frac{\mu_3 \mu_4 n}{2L}$ number of eigenvalues are larger than $\bar{\lambda}$. Denote $\mu_5:=\frac{\mu_3\mu_4}{2L}$. We know $\lambda_{\mu_5n}-\lambda_{n-\mu_5n+1}\geq \mu_4$.

Proof of Lemma 27. Let $\mathcal{E}_{\underline{1}}$ be the event that $\sqrt{d}/\sqrt{L} \leq \underline{\sigma}_i(X_{\text{train}}) \leq \sqrt{Ld}$ and $1/\underline{L} \leq \lambda_i(H_{\text{train}}) \leq L$ for all $i \in [n]$ and $\sqrt{d}\sigma/4 \le \|\xi_{\text{train}}\| \le \sqrt{d}\sigma$. Let \mathcal{E}_3 be the event that $\sqrt{d}/\sqrt{L} \le \sigma_i(X_{\text{valid}}) \le \sqrt{Ld}$ and $1/L \le \lambda_i(H_{\text{valid}}) \le L$ for all $i \in [n]$ and $\sqrt[n]{d\sigma/4} \le \|\xi_{\text{valid}}\| \le \sqrt[n]{d\sigma}$. According to Lemma 7 and Lemma 49, we know both \mathcal{E}_1 and \mathcal{E}_3 hold with probability at least $1 - \exp(-\Omega(d))$. In this proof, we assume both properties hold and take a union bound at the end.

We can lower bound $\|w_{t,\eta} - w_{\text{valid}}\|_{H_{\text{valid}}}^2$ as follows,

$$\begin{split} \left\| w_{t,\eta} - w_{\text{valid}} \right\|_{H_{\text{valid}}}^2 &= \left\| w_{t,\eta} - w^* - (X_{\text{valid}})^\dagger \xi_{\text{valid}} \right\|_{H_{\text{valid}}}^2 \\ &\geq \left\| w_{t,\eta} - w^* \right\|_{H_{\text{valid}}}^2 + \frac{1}{n} \left\| \xi_{\text{valid}} \right\|^2 - 2 \left| \left\langle w_{t,\eta} - w^*, H_{\text{valid}}(X_{\text{valid}})^\dagger \xi_{\text{valid}} \right\rangle \right|. \end{split}$$

For the second term, by Lemma 49, we know for any $1 > \epsilon > 0$, with probability at least $1 - \exp(-\Omega(\epsilon^2 d))$,

$$\frac{1}{n} \|\xi_{\text{valid}}\|^2 \ge (1 - \epsilon)\sigma^2.$$

We can write down the third term as $\langle [(X_{\text{valid}})^{\dagger}]^{\top} H_{\text{valid}}(w_{t,\eta} - w^*), \xi_{\text{valid}} \rangle$. Suppose σ is a constant, we know $\|[(X_{\text{valid}})^{\dagger}]^{\top} H_{\text{valid}}(w_{t,\eta} - w^*)\| = O(1/\sqrt{d})$. Therefore, for a fixed $\eta \in [1/L, 3L]$, we have with probability at least $1 - \exp(-\Omega(\epsilon^2 d))$,

$$\left|\left\langle w_{t,\eta} - w^*, H_{\text{valid}}(X_{\text{valid}})^{\dagger} \xi_{\text{valid}}\right\rangle\right| \leq \epsilon.$$

To prove this crossing term is small for all $\eta \in [1/L, 3L]$, we need to construct an ϵ -net for the crossing term. Similar as in Lemma 13, we can show there exists an ϵ -net for the crossing term with size $O(t/\epsilon)$. Taking a union bound over this ϵ -net, we are able to show with probability at least $1 - O(t/\epsilon) \exp(-\Omega(\epsilon^2 d))$,

$$\left|\left\langle w_{t,\eta} - w^*, H_{\text{valid}}(X_{\text{valid}})^{\dagger} \xi_{\text{valid}}\right\rangle\right| \leq \epsilon,$$

for all $\eta \in [1/L, 3L]$.

Overall, we have with probability at least $1 - O(t/\epsilon) \exp(-\Omega(\epsilon^2 d))$,

$$||w_{t,\eta} - w_{\text{valid}}||_{H_{\text{valid}}}^2 \ge ||w_{t,\eta} - w^*||_{H_{\text{valid}}}^2 + \frac{1}{n} ||\xi_{\text{valid}}||^2 - 2 |\langle w_{t,\eta} - w^*, H_{\text{valid}}(X_{\text{valid}})^{\dagger} \xi_{\text{valid}} \rangle|$$

$$\ge ||w_{t,\eta} - w^*||_{H_{\text{valid}}}^2 + (1 - \epsilon)\sigma^2 - 2\epsilon \ge (1 - 3\epsilon)\sigma^2,$$

for all $\eta \in [1/L, 3L]$, where the last inequality uses $\sigma \geq 1$. The proof finishes as we change 3ϵ to ϵ' .

C Proofs of train-by-train with large number of samples (GD)

In this section, we give the proof of Theorem 6. We show when the size of each training set n and the number of training tasks m are large enough, train-by-train also performs well. Recall Theorem 6 as follows.

Theorem 6. Let $\hat{F}_{TbT(n)}(\eta)$ be as defined in Equation (3). Assume noise level is a constant c_1 . Given any $1 > \epsilon > 0$, assume training set size $n \geq \frac{cd}{\epsilon^2}\log(\frac{nm}{\epsilon d})$, unroll length $t \geq c_2\log(\frac{n}{\epsilon d})$, number of training tasks $m \geq \frac{c_3n^2}{\epsilon^4 d^2}\log(\frac{tnm}{\epsilon d})$ and dimension $d \geq c_4$ for certain constants c, c_2, c_3, c_4 . With probability at least 0.99 in the sampling of training tasks, we have

$$\mathbb{E} \left\| w_{t,\eta_{train}^*} - w^* \right\|^2 \le (1+\epsilon) \frac{d\sigma^2}{n},$$

for all $\eta_{train}^* \in \arg\min_{\eta \geq 0} \hat{F}_{TbT(n)}(\eta)$, where the expectation is taken over new tasks.

In the proof, we use the same notations defined in Section B. On each training task P, in Lemma 28 we show the meta-loss can be decomposed into two terms:

$$\Delta_{TbT}(\eta, P) = \frac{1}{2} \left\| w_{t,\eta} - w_{\text{train}} \right\|_{H_{\text{train}}}^2 + \frac{1}{2n} \left\| (I_n - \text{Proj}_{X_{\text{train}}}) \xi_{\text{train}} \right\|^2$$

where $w_{\text{train}} = w^* + (X_{\text{train}})^\dagger \xi_{\text{train}}$. Recall that X_{train} is a $n \times d$ matrix with its i-th row as x_i^\top . The pseudo-inverse $(X_{\text{train}})^\dagger$ has dimension $d \times n$ satisfying $X_{\text{train}}^\dagger X_{\text{train}} = I_d$. Here, $\text{Proj}_{X_{\text{train}}} \in \mathbb{R}^{n \times n}$ is a projection matrix onto the column span of X_{train} .

In Lemma 28, we show with a constant step size, the first term in $\Delta_{TbT}(\eta, P)$ is exponentially small. The second term is basically the projection of the noise on the orthogonal subspace of the data span. We show this term concentrates well on its mean. This lemma servers as step 1 in Section B.1. The proof of Lemma 28 is deferred into Section C.1.

Lemma 28. Assume $n \ge 40d$. Given any $1 > \epsilon > 0$, with probability at least $1 - m \exp(-\Omega(n)) - \exp(-\Omega(\epsilon^4 m d/n))$,

$$\hat{F}_{TbT}(2/3) \le 20(1 - \frac{1}{3})^{2t}\sigma^2 + \frac{n - d}{2n}\sigma^2 + \frac{\epsilon^2 d\sigma^2}{20n}.$$

In the next lemma, we show the empirical meta objective is large when η exceeds certain threshold. We define this threshold $\hat{\eta}$ such that for any step size larger than $\hat{\eta}$ the GD sequence has reasonable probability being truncated. In the proof, we rely on the truncated sequences to argue the meta-objective must be high. The precise definition of $\hat{\eta}$ is in Definition 2. This lemma serves as step 2 in Section B.1. We leave the proof of Lemma 29 into Section C.2.

Lemma 29. Let $\hat{\eta}$ be as defined in Definition 2 with $1 > \epsilon > 0$. Assume $n \ge cd, t \ge c_2, d \ge c_4$ for some constants c, c_2, c_4 . With probability at least $1 - \exp(-\Omega(\epsilon^4 m d^2/n^2))$.

$$\hat{F}_{TbT}(\eta) \ge \frac{\epsilon^2 d\sigma^2}{8n} + \frac{n-d}{2n}\sigma^2 - \frac{\epsilon^2 d\sigma^2}{20n}$$

for all $\eta > \hat{\eta}$.

By Lemma 28 and Lemma 29, we know when t is reasonably large, $\hat{F}_{TbT}(\eta)$ is larger than $\hat{F}_{TbT}(2/3)$ for all step sizes $\eta > \hat{\eta}$. This means the optimal step size $\hat{\eta}$ must lie in $[0, \hat{\eta}]$. In Lemma 30, we show a generalization result for $\eta \in [0, \hat{\eta}]$. This serves as step 3 in Section B.1. We prove this lemma in Section C.3.

Lemma 30. Let $\hat{\eta}$ be as defined in Definition 2 with $1 > \epsilon > 0$. Suppose σ is a constant. Assume $n \ge c \log(\frac{n}{\epsilon d})d$, $t \ge c \log(\frac{n}{\epsilon d})d$ $c_2, d \geq c_4$ for some constants c, c_2, c_4 . With probability at least $1 - m \exp(-\Omega(n)) - O(\frac{tn}{\epsilon^2 d} + m) \exp(-\Omega(m\epsilon^4 d^2/n^2))$,

$$|F_{TbT}(\eta) - \hat{F}_{TbT}(\eta)| \le \frac{17\epsilon^2 d\sigma^2}{n},$$

for all $\eta \in [0, \hat{\eta}],$

Combining Lemma 28, Lemma 29 and Lemma 30, we present the proof of Theorem 6 as follows.

Proof of Theorem 6. According to Lemma 28, assuming $n \geq 40d$, given any $1/2 > \epsilon > 0$, with probability at least $1 - m \exp(-\Omega(n)) - \exp(-\Omega(\epsilon^4 m d/n))$, $\hat{F}_{TbT}(2/3) \leq 20(1 - \frac{1}{3})^{2t} \sigma^2 + \frac{n-d}{2n} \sigma^2 + \frac{\epsilon^2 d \sigma^2}{20n}$. As long as $t \geq c_2 \log(\frac{n}{\epsilon d})$ for certain constant c_2 , we have

$$\hat{F}_{TbT}(2/3) \le \frac{n-d}{2n}\sigma^2 + \frac{7\epsilon^2 d\sigma^2}{100n}.$$

Let $\hat{\eta}$ be as defined in Definition 2 with the same ϵ . According to Lemma 29, as long as $n \geq cd, t \geq c_2, d \geq c_4$ with probability at least $1 - \exp(-\Omega(\epsilon^4 m d^2/n^2))$,

$$\hat{F}_{TbT}(\eta) \ge \frac{\epsilon^2 d\sigma^2}{8n} + \frac{n-d}{2n}\sigma^2 - \frac{\epsilon^2 d\sigma^2}{20n} = \frac{n-d}{2n}\sigma^2 + \frac{7.5\epsilon^2 d\sigma^2}{100n}$$

for all $\eta > \hat{\eta}$. We have $\hat{F}_{TbT}(\eta) > \hat{F}_{TbT}(2/3)$ for all $\eta \geq \hat{\eta}$. This implies that η^*_{train} is within $[0,\hat{\eta}]$ and $\hat{F}_{TbT}(\eta^*_{\text{train}}) \leq \hat{F}_{TbT}(2/3) \leq \frac{n-d}{2n}\sigma^2 + \frac{7\epsilon^2 d\sigma^2}{100n}$.

By Lemma 30, assuming σ is a constant and assuming $n \geq c\log(\frac{n}{\epsilon d})d$ for some constant c, we have with probability at least $1 - m\exp(-\Omega(n)) - O(\frac{tn}{\epsilon^2 d} + m)\exp(-\Omega(m\epsilon^4 d^2/n^2))$,

$$|F_{TbT}(\eta) - \hat{F}_{TbT}(\eta)| \le \frac{17\epsilon^2 d\sigma^2}{n},$$

for all $\eta \in [0, \hat{\eta}]$. This then implies

$$F_{TbT}(\eta_{\text{train}}^*) \leq \hat{F}_{TbT}(\eta_{\text{train}}^*) + \frac{17\epsilon^2 d\sigma^2}{n} \leq \frac{n-d}{2n}\sigma^2 + \frac{24\epsilon^2 d\sigma^2}{n}.$$

By the analysis in Lemma 28, we have

$$\begin{split} F_{TbT}(\eta_{\text{train}}^*) = & \mathbb{E}\frac{1}{2} \left\| w_{t,\eta_{\text{train}}^*} - w_{\text{train}} \right\|_{H_{\text{train}}}^2 + \mathbb{E}\frac{1}{2n} \left\| (I_n - \text{Proj}_{X_{\text{train}}}) \xi_{\text{train}} \right\|^2 \\ = & \mathbb{E}\frac{1}{2} \left\| w_{t,\eta_{\text{train}}^*} - w_{\text{train}} \right\|_{H_{\text{train}}}^2 + \frac{n-d}{2n} \sigma^2. \end{split}$$

Therefore, we know $\mathbb{E}\frac{1}{2}\left\|w_{t,\eta_{\text{train}}^*}-w_{\text{train}}\right\|_{H_{\text{train}}}^2 \leq \frac{24\epsilon^2d\sigma^2}{n}$. Next, we show this implies $\mathbb{E}\left\|w_{t,\eta_{\text{train}}^*}-w^*\right\|^2$ is small. Let \mathcal{E} be the event that $1-\epsilon \leq \lambda_i(H_{\text{train}}) \leq 1+\epsilon$ for all $i\in[d]$. According to Lemma 31, we know $\Pr[\mathcal{E}] \geq 1$ $1 - \exp(-\Omega(\epsilon^2 n))$ as long as $n \ge 10d/\epsilon^2$. Then, we can decompose $\mathbb{E} \|w_{t,\eta_{\text{train}}^*} - w^*\|^2$ as follows,

$$\mathbb{E}\left\|w_{t,\eta_{\text{train}}^*}-w^*\right\|^2 = \mathbb{E}\left\|w_{t,\eta_{\text{train}}^*}-w^*\right\|^2\mathbb{1}\left\{\mathcal{E}\right\} + \mathbb{E}\left\|w_{t,\eta_{\text{train}}^*}-w^*\right\|^2\mathbb{1}\left\{\bar{\mathcal{E}}\right\}.$$

Let's first show the second term is small. Due to the truncation in our algorithm, we know $\|w_{t,\eta_{\text{train}}^*} - w^*\|^2 \le 41^2\sigma^2$, which then implies $\mathbb{E}\|w_{t,\eta_{\text{train}}^*} - w^*\|^2 \mathbb{1}\{\bar{\mathcal{E}}\} \le 41^2\sigma^2 \exp(-\Omega(\epsilon^2 n))$. As long as $n \ge \frac{c}{\epsilon^2}\log(\frac{n}{\epsilon d})$ for some constant c, we have $\mathbb{E}\|w_{t,\eta_{\text{train}}^*} - w^*\|^2 \mathbb{1}\{\bar{\mathcal{E}}\} \le \frac{\epsilon d\sigma^2}{n}$.

We can upper bound the first term by Young's inequality,

$$\mathbb{E}\left\|w_{t,\eta_{\text{train}}^*} - w^*\right\|^2 \mathbb{1}\left\{\mathcal{E}\right\} \leq \left(1 + \frac{1}{\epsilon}\right) \mathbb{E}\left\|w_{t,\eta_{\text{train}}^*} - w_{\text{train}}\right\|^2 \mathbb{1}\left\{\mathcal{E}\right\} + \left(1 + \epsilon\right) \mathbb{E}\left\|w_{\text{train}} - w^*\right\|^2 \mathbb{1}\left\{\mathcal{E}\right\}.$$

Conditioning on \mathcal{E} , we have $\left\|w_{t,\eta_{\text{train}}^*} - w_{\text{train}}\right\|_{H_{\text{train}}}^2 \geq (1-\epsilon) \left\|w_{t,\eta_{\text{train}}^*} - w_{\text{train}}\right\|^2$ which implies $\left\|w_{t,\eta_{\text{train}}^*} - w_{\text{train}}\right\|^2 \leq (1+2\epsilon) \left\|w_{t,\eta_{\text{train}}^*} - w_{\text{train}}\right\|_{H_{\text{train}}}^2$ as long as $\epsilon \leq 1/2$. Similarly, we also have $\left\|w_{\text{train}} - w^*\right\|^2 \leq (1+2\epsilon) \left\|w_{\text{train}} - w^*\right\|_{H_{\text{train}}}^2$. Then, we have

$$\begin{split} & \mathbb{E} \left\| w_{t,\eta_{\text{train}}^*} - w^* \right\|^2 \mathbb{1} \left\{ \mathcal{E} \right\} \\ & \leq (1 + \frac{1}{\epsilon})(1 + 2\epsilon) \mathbb{E} \left\| w_{t,\eta_{\text{train}}^*} - w_{\text{train}} \right\|_{H_{\text{train}}}^2 \mathbb{1} \left\{ \mathcal{E} \right\} + (1 + \epsilon)(1 + 2\epsilon) \mathbb{E} \left\| w_{\text{train}} - w^* \right\|_{H_{\text{train}}}^2 \mathbb{1} \left\{ \mathcal{E} \right\} \\ & \leq (5 + \frac{1}{\epsilon}) \mathbb{E} \left\| w_{t,\eta_{\text{train}}^*} - w_{\text{train}} \right\|_{H_{\text{train}}}^2 + (1 + 5\epsilon) \mathbb{E} \left\| w_{\text{train}} - w^* \right\|_{H_{\text{train}}}^2 \\ & \leq (5 + \frac{1}{\epsilon}) \frac{48\epsilon^2 d\sigma^2}{n} + (1 + 5\epsilon) \frac{d\sigma^2}{n} \leq (1 + 293\epsilon) \frac{d\sigma^2}{n}. \end{split}$$

Overall, we have $\mathbb{E}\left\|w_{t,\eta_{\text{train}}^*}-w^*\right\|^2 \leq (1+293\epsilon)\frac{d\sigma^2}{n}+\frac{\epsilon d\sigma^2}{n}=(1+294\epsilon)\frac{d\sigma^2}{n}.$ Combining all the conditions, we know this holds with probability at least 0.99 as long as σ is a constant $c_1, n \geq \frac{cd}{\epsilon^2}\log(\frac{nm}{\epsilon d}), t \geq c_2\log(\frac{n}{\epsilon d}), m \geq \frac{c_3n^2}{\epsilon^4d^2}\log(\frac{tnm}{\epsilon d}), d \geq c_4$ for some constants c, c_2, c_3, c_4 . We finish the proof by choosing $\epsilon = \epsilon'/294$.

C.1 Upper bounding $\hat{F}_{TbT}(2/3)$

In this section, we show there exists a step size that achieves small empirical meta objective. On each training task P, we show the meta-loss can be decomposed into two terms:

$$\Delta_{TbT}(\eta, P) = \frac{1}{2n} \sum_{i=1}^{n} \left(\langle w_{t,\eta} - w_{\text{train}}, x_i \rangle - \left(\xi_i - x_i^{\top} X_{\text{train}}^{\dagger} \xi_{\text{train}} \right) \right)^2$$

$$= \frac{1}{2} \left\| w_{t,\eta} - w_{\text{train}} \right\|_{H_{\text{train}}}^2 + \frac{1}{2n} \left\| (I_n - \text{Proj}_{X_{\text{train}}}) \xi_{\text{train}} \right\|^2,$$

where $w_{\text{train}} = w^* + (X_{\text{train}})^{\dagger} \xi_{\text{train}}$. In Lemma 28, we show with a constant step size, the first term is exponentially small and the second term concentrates on its mean.

Lemma 28. Assume $n \ge 40d$. Given any $1 > \epsilon > 0$, with probability at least $1 - m \exp(-\Omega(n)) - \exp(-\Omega(\epsilon^4 m d/n))$,

$$\hat{F}_{TbT}(2/3) \le 20(1 - \frac{1}{3})^{2t}\sigma^2 + \frac{n-d}{2n}\sigma^2 + \frac{\epsilon^2 d\sigma^2}{20n}.$$

Before we go to the proof of Lemma 28, let's first show the covariance matrix H_{train} is very close to identity when n is much larger than d. The proof follows from the concentration of singular values of random Gaussian matrix (Lemma 52). We leave the proof into Section C.4.

Lemma 31. Given $1 > \epsilon > 0$, assume $n \ge 10d/\epsilon^2$. With probability at least $1 - \exp(-\Omega(\epsilon^2 n))$,

$$(1 - \epsilon)\sqrt{n} \le \sigma_i(X_{\textit{train}}) \le (1 + \epsilon)\sqrt{n} \ \textit{and} \ 1 - \epsilon \le \lambda_i(H_{\textit{train}}) \le 1 + \epsilon,$$

for all $i \in [d]$.

Now, we are ready to present the proof of Lemma 28.

Proof of Lemma 28. Let's first look at one training set S_{train} , in which $y_i = \langle w^*, x_i \rangle + \xi_i$ for each sample. Recall the meta-loss as

$$\Delta_{TbT}(\eta, P) = \frac{1}{2n} \sum_{i=1}^{n} \left(\langle w_{t,\eta}, x_i \rangle - \langle w^*, x_i \rangle - \xi_i \right)^2.$$

Recall that X_{train} is an $n \times d$ matrix with its i-th row as x_i^{\top} . With probability 1, we know X_{train} is full column rank. Denote the pseudo-inverse of X_{train} as $X_{\text{train}}^{\dagger} \in \mathbb{R}^{d \times n}$ that satisfies $X_{\text{train}}^{\dagger} X_{\text{train}} = I_d$ and $X_{\text{train}} X_{\text{train}}^{\dagger} = \text{Proj}_{X_{\text{train}}}$, where $\text{Proj}_{X_{\text{train}}} \in \mathbb{R}^{n \times n}$ is a projection matrix onto the column span of X_{train} .

Let w_{train} be $w^* + X_{\text{train}}^{\dagger} \xi_{\text{train}}$, where ξ_{train} is an *n*-dimensional vector with its *i*-th entry as ξ_i . We have,

$$\begin{split} &\Delta_{TbT}(\eta, P) \\ &= \frac{1}{2n} \sum_{i=1}^{n} \left(\langle w_{t,\eta} - w_{\text{train}}, x_i \rangle - \left(\xi_i - x_i^\top X_{\text{train}}^\dagger \xi_{\text{train}} \right) \right)^2 \\ &= \frac{1}{2} \left\| w_{t,\eta} - w_{\text{train}} \right\|_{H_{\text{train}}}^2 + \frac{1}{2n} \left\| (I_n - \text{Proj}_{X_{\text{train}}}) \xi_{\text{train}} \right\|^2 - \frac{1}{n} \sum_{i=1}^{n} \left\langle w_{t,\eta} - w_{\text{train}}, x_i \xi_i - x_i x_i^\top X_{\text{train}}^\dagger \xi_{\text{train}} \right\rangle. \end{split}$$

We first show the crossing term is actually zero. We have,

$$\begin{split} \frac{1}{n} \sum_{i=1}^{n} \left\langle w_{t,\eta} - w_{\text{train}}, x_{i} \xi_{i} - x_{i} x_{i}^{\intercal} X_{\text{train}}^{\dagger} \xi_{\text{train}} \right\rangle &= \frac{1}{n} \left\langle w_{t,\eta} - w_{\text{train}}, \sum_{i=1}^{n} x_{i} \xi_{i} - \sum_{i=1}^{n} x_{i} x_{i}^{\intercal} X_{\text{train}}^{\dagger} \xi_{\text{train}} \right\rangle \\ &= \frac{1}{n} \left\langle w_{t,\eta} - w_{\text{train}}, X_{\text{train}}^{\intercal} \xi_{\text{train}} - X_{\text{train}}^{\intercal} X_{\text{train}} X_{\text{train}}^{\dagger} \xi_{\text{train}} \right\rangle \\ &= \frac{1}{n} \left\langle w_{t,\eta} - w_{\text{train}}, X_{\text{train}}^{\intercal} \xi_{\text{train}} - X_{\text{train}}^{\intercal} \xi_{\text{train}} \right\rangle = 0, \end{split}$$

where the second last equality holds because $X_{\text{train}}X_{\text{train}}^{\dagger} = \text{Proj}_{X_{\text{train}}}$. We can define $w_{\text{train}}^{(k)}$ as $w_k^* + (X_{\text{train}}^{(k)})^{\dagger}\xi_{\text{train}}^{(k)}$ for every training set $S_{\text{train}}^{(k)}$. Then, we have

$$\hat{F}_{TbT}(\eta) = \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{\text{train}}^{(k)} \right\|_{H_{\text{train}}^{(k)}}^2 + \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2n} \left\| (I_n - \text{Proj}_{X_{\text{train}}^{(k)}}) \xi_{\text{train}}^{(k)} \right\|^2$$

We first prove that the second term concentrates on its mean. We can concatenate m noise vectors $\xi_{\text{train}}^{(k)}$ into a single noise vector $\bar{\xi}_{\text{train}}$ with dimension nm. We can also construct a data matrix $\bar{X}_{\text{train}} \in \mathbb{R}^{nm \times dm}$ that consists of $X_{\text{train}}^{(k)}$ as diagonal blocks. Then the second term can be written as

$$\frac{1}{2} \left\| \frac{1}{\sqrt{nm}} (I_{nm} - \operatorname{Proj}_{\bar{X}_{\text{train}}}) \bar{\xi}_{\text{train}} \right\|^2.$$

According to Lemma 49, with probability at least $1 - \exp(-\Omega(\epsilon^4 m d^2/n))$,

$$\left(1 - \frac{\epsilon^2 d}{n}\right) \sigma \leq \frac{1}{\sqrt{nm}} \left\|\bar{\xi}_{\text{train}}\right\| \leq \left(1 + \frac{\epsilon^2 d}{n}\right) \sigma.$$

By Johnson-Lindenstrauss Lemma (Lemma 53), we know with probability at least $1 - \exp(-\Omega(\epsilon^4 m d))$.

$$\frac{1}{\sqrt{nm}} \left\| \operatorname{Proj}_{\bar{X}_{\operatorname{train}}} \bar{\xi}_{\operatorname{train}} \right\| \geq (1 - \epsilon^2) \frac{\sqrt{md}}{\sqrt{mn}} \frac{1}{\sqrt{nm}} \left\| \bar{\xi}_{\operatorname{train}} \right\| \geq (1 - \epsilon^2) \sqrt{\frac{d}{n}} (1 - \frac{\epsilon^2 d}{n}) \sigma.$$

Therefore, we have $\left\|\frac{1}{\sqrt{nm}}\bar{\xi}_{\text{train}}\right\|^2 \leq (1+\frac{3\epsilon^2d}{n})\sigma^2$ and $\left\|\frac{1}{\sqrt{nm}}\text{Proj}_{\bar{X}_{\text{train}}}\bar{\xi}_{\text{train}}\right\|^2 \geq (1-2\epsilon^2)\frac{d}{n}\sigma^2$. Overall, we know with probability at least $1-\exp(-\Omega(\epsilon^4md/n))$,

$$\frac{1}{2} \left\| \frac{1}{\sqrt{nm}} (I_{nm} - \operatorname{Proj}_{\bar{X}_{\text{train}}}) \bar{\xi}_{\text{train}} \right\|^2 \leq \frac{n - d}{2n} \sigma^2 + \frac{5\epsilon^2 d\sigma^2}{2n}.$$

Now, we show the first term in meta objective is small when we choose a right step size. According to Lemma 31, we know as long as $n \geq 40d$, with probability at least $1 - \exp(-\Omega(n))$, $\sqrt{n}/2 \leq \sigma_i(X_{\text{train}}^{(k)}) \leq 3\sqrt{n}/2$ and $1/2 \leq \lambda_i(H_{\text{train}}^{(k)}) \leq 3/2$, for all $i \in [d]$. According to Lemma 49, we know with probability at least $1 - \exp(-\Omega(n))$, $\left\|\xi_{\text{train}}^{(k)}\right\| \leq 2\sqrt{n}\sigma$. Taking a union bound on m tasks, we know all these events hold with probability at least $1 - m \exp(-\Omega(n))$.

For each $k \in [m]$, we have $\left\| w_{\text{train}}^{(k)} \right\| \le 1 + \frac{2}{\sqrt{n}} 2\sqrt{n}\sigma \le 5\sigma$. It's easy to verify that for any step size at most 2/3, the GD sequence will not be truncated since we choose the threshold norm as 40σ . Then, for any step size $\eta \le 2/3$, we have

$$\begin{split} \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{\text{train}}^{(k)} \right\|_{H_{\text{train}}^{(k)}}^{2} &= \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| (I - \eta H_{\text{train}}^{(k)})^{t} w_{\text{train}}^{(k)} \right\|_{H_{\text{train}}^{(k)}}^{2} \\ &\leq \frac{3}{4} (1 - \frac{\eta}{2})^{2t} 25 \sigma^{2} \leq 20 (1 - \frac{1}{3})^{2t} \sigma^{2}, \end{split}$$

where the last inequality chooses η as 2/3.

Overall, we know with probability at least $1 - m \exp(-\Omega(n)) - \exp(-\Omega(\epsilon^4 m d/n))$,

$$\hat{F}_{TbT}(2/3) \le 20(1 - \frac{1}{3})^{2t}\sigma^2 + \frac{n - d}{2n}\sigma^2 + \frac{5\epsilon^2 d\sigma^2}{2n}.$$

We finish the proof by changing $\frac{5\epsilon^2}{2}$ by $(\epsilon')^2/20$.

C.2 Lower bounding \hat{F}_{TbT} for $\eta \in (\hat{\eta}, \infty)$

In this section, we show the empirical meta objective is large when the step size exceeds certain threshold. Recall Lemma 29 as follows.

Lemma 29. Let $\hat{\eta}$ be as defined in Definition 2 with $1 > \epsilon > 0$. Assume $n \ge cd, t \ge c_2, d \ge c_4$ for some constants c, c_2, c_4 . With probability at least $1 - \exp(-\Omega(\epsilon^4 m d^2/n^2))$,

$$\hat{F}_{TbT}(\eta) \ge \frac{\epsilon^2 d\sigma^2}{8n} + \frac{n-d}{2n}\sigma^2 - \frac{\epsilon^2 d\sigma^2}{20n},$$

for all $\eta > \hat{\eta}$.

Roughly speaking, we define $\hat{\eta}$ such that for any step size larger than $\hat{\eta}$ the GD sequence has a reasonable probability being truncated. The definition is very similar as $\hat{\eta}$ in Definition 1.

Definition 2. Given a training task P, let \mathcal{E}_1 be the event that $\sqrt{n}/2 \le \sigma_i(X_{train}) \le 3\sqrt{n}/2$ and $1/2 \le \lambda_i(H_{train}) \le 3/2$ for all $i \in [d]$ and $\sqrt{n}\sigma/2 \le \|\xi_{train}\| \le 2\sqrt{n}\sigma$. Let $\bar{\mathcal{E}}_2(\eta)$ be the event that the GD sequence is truncated with step size η . Given $1 > \epsilon > 0$, define $\hat{\eta}$ as follows,

$$\hat{\eta} = \inf \left\{ \eta \geq 0 \middle| \mathbb{E}\frac{1}{2} \left\| w_{t,\eta} - w_{\textit{train}} \right\|_{H_{\textit{train}}}^2 \mathbb{1} \left\{ \mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\eta) \right\} \geq \frac{\epsilon^2 d\sigma^2}{n} \right\}.$$

Similar as in Lemma 9, we show $\mathbb{I}\left\{\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\eta')\right\} \geq \mathbb{I}\left\{\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\eta)\right\}$ for any $\eta' \geq \eta$. This means conditioning on \mathcal{E}_1 , if a GD sequence gets truncated with step size η , it has to be truncated with any step size $\eta' \geq \eta$. The proof is deferred into Section C.4.

Lemma 32. Fixing a training set S_{train} , let \mathcal{E}_1 and $\bar{\mathcal{E}}_2(\eta)$ be as defined in Definition 2. We have

$$\mathbb{1}\left\{\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\eta')\right\} \ge \mathbb{1}\left\{\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\eta)\right\},\,$$

for any $\eta' \geq \eta$.

Next, we show $\hat{\eta}$ does exist and is a constant. Similar as in Lemma 10, we show that the GD sequence almost never diverges when η is small and diverges with high probability when η is large. The proof is left in Section C.4.

Lemma 33. Let $\hat{\eta}$ be as defined in Definition 2. Suppose σ is a constant. Assume $n \geq cd, t \geq c_2, d \geq c_4$ for some constants c, c_2, c_4 . We have

$$\frac{4}{3} < \tilde{\eta} < 6.$$

Next, we show the empirical loss is large for any η larger than $\tilde{\eta}$. The proof is very similar as the proof of Lemma 2. **Proof of Lemma 29.** By Lemma 33, we know $\hat{\eta}$ is a constant as long as $n \geq cd, t \geq c_2, d \geq c_4$ for some constants c, c_2, c_4 . Let \mathcal{E}_1 and $\bar{\mathcal{E}}_2(\eta)$ be as defined in Definition 2. For the simplicity of the proof, we assume $\mathbb{E}\frac{1}{2}\|w_{t,\hat{\eta}}-w_{\text{train}}\|_{H_{\text{train}}}^2\mathbb{1}\left\{\mathcal{E}_1\cap\bar{\mathcal{E}}_2(\hat{\eta})\right\}\geq \frac{\epsilon^2d\sigma^2}{n}$. The other case can be resolved using same techniques in Lemma 2

Conditioning on \mathcal{E}_1 , we know $\frac{1}{2}\|w_{t,\hat{\eta}} - w_{\text{train}}\|_{H_{\text{train}}}^2 \leq \frac{3}{4}45^2\sigma^2$. Therefore, we know $\Pr[\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\hat{\eta})] \geq \frac{4\epsilon^2d}{3\times 45^2n}$. For each task k, define $\mathcal{E}_1^{(k)}$ and $\bar{\mathcal{E}}_2^{(k)}(\eta)$ as the corresponding events on training set $S_{\text{train}}^{(k)}$. By Hoeffding's inequality, we know with probability at least $1 - \exp(-\Omega(\epsilon^4md^2/n^2))$,

$$\frac{1}{m} \sum_{k=1}^{m} \mathbb{1}\left\{\mathcal{E}_1^{(k)} \cap \bar{\mathcal{E}}_2^{(k)}(\hat{\eta})\right\} \ge \frac{\epsilon^2 d}{45^2 n}.$$

By Lemma 32, we know $\mathbb{1}\left\{\mathcal{E}_1^{(k)}\cap\bar{\mathcal{E}}_2^{(k)}(\eta)\right\}\geq\mathbb{1}\left\{\mathcal{E}_1^{(k)}\cap\bar{\mathcal{E}}_2^{(k)}(\hat{\eta})\right\}$ for any $\eta\geq\hat{\eta}$. Recall that

$$\hat{F}_{TbT}(\eta) = \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{\text{train}}^{(k)} \right\|_{H_{\text{train}}^{(k)}}^2 + \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2n} \left\| (I_n - \text{Proj}_{X_{\text{train}}^{(k)}}) \xi_{\text{train}}^{(k)} \right\|^2.$$

We can lower bound the first term for any $\eta > \hat{\eta}$ as follows,

$$\begin{split} \hat{F}_{TbT}(\eta) &= \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{\text{train}}^{(k)} \right\|_{H_{\text{train}}}^{2} \geq \frac{1}{m} \sum_{k=1}^{m} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{\text{train}}^{(k)} \right\|_{H_{\text{train}}}^{2} \mathbbm{1} \left\{ \mathcal{E}_{1}^{(k)} \cap \bar{\mathcal{E}}_{2}^{(k)}(\eta) \right\} \\ &\geq \frac{35^{2} \sigma^{2}}{4} \frac{1}{m} \sum_{k=1}^{m} \mathbbm{1} \left\{ \mathcal{E}_{1}^{(k)} \cap \bar{\mathcal{E}}_{2}^{(k)}(\eta) \right\} \\ &\geq \frac{35^{2} \sigma^{2}}{4} \frac{1}{m} \sum_{k=1}^{m} \mathbbm{1} \left\{ \mathcal{E}_{1}^{(k)} \cap \bar{\mathcal{E}}_{2}^{(k)}(\hat{\eta}) \right\} \geq \frac{\epsilon^{2} d \sigma^{2}}{8n}, \end{split}$$

where the second inequality lower bounds the loss for one task by $35^2\sigma^2$ when the sequence gets truncated. For the second term, according to the analysis in Lemma 28, with probability at least $1 - \exp(-\Omega(\epsilon^4 md/n))$,

$$\frac{1}{m}\sum_{k=1}^{m}\frac{1}{2n}\left\|(I_n-\operatorname{Proj}_{X_{\operatorname{train}}^{(k)}})\xi_{\operatorname{train}}^{(k)}\right\|^2\geq \frac{n-d}{2n}\sigma^2-\frac{\epsilon^2d\sigma^2}{20n}.$$

Overall, with probability at least $1 - \exp(-\Omega(\epsilon^4 m d^2/n^2))$,

$$\hat{F}_{TbT}(\eta) \ge \frac{\epsilon^2 d\sigma^2}{8n} + \frac{n-d}{2n}\sigma^2 - \frac{\epsilon^2 d\sigma^2}{20n},$$

for all $\eta > \hat{\eta}$.

C.3 Generalization for $\eta \in [0, \hat{\eta}]$

Combing Lemma 28 and Lemma 29, it's not hard to see that the optimal step size η_{train}^* lies in $[0, \hat{\eta}]$. In this section, we show a generalization result for step sizes in $[0, \hat{\eta}]$. The proof of Lemma 30 is given at the end of this section.

Lemma 30. Let $\hat{\eta}$ be as defined in Definition 2 with $1 > \epsilon > 0$. Suppose σ is a constant. Assume $n \ge c \log(\frac{n}{\epsilon d})d$, $t \ge c_2$, $d \ge c_4$ for some constants c, c_2 , c_4 . With probability at least $1 - m \exp(-\Omega(n)) - O(\frac{tn}{\epsilon^2 d} + m) \exp(-\Omega(m\epsilon^4 d^2/n^2))$,

$$|F_{TbT}(\eta) - \hat{F}_{TbT}(\eta)| \le \frac{17\epsilon^2 d\sigma^2}{n},$$

for all $\eta \in [0, \hat{\eta}]$,

In Lemma 34, we show \hat{F}_{TbT} concentrates on F_{TbT} at any fixed step size. The proof is almost the same as Lemma 11. We omit its proof.

Lemma 34. Suppose σ is a constant. For any fixed η and any $1 > \epsilon > 0$, with probability at least $1 - \exp(-\Omega(\epsilon^2 m))$,

$$\left| \hat{F}_{TbT}(\eta) - F_{TbT}(\eta) \right| \le \epsilon.$$

Next, we construct an ϵ -net for F_{TbT} in $[0, \hat{\eta}]$. The proof is very similar as in Lemma 12. We defer its proof into Section C.4.

Lemma 35. Let $\hat{\eta}$ be as defined in Definition 2 with $1 > \epsilon > 0$. Assume the conditions in Lemma 33 hold. Assume $n \ge c\log(\frac{n}{\epsilon d})d$ for some constant c. There exists an $\frac{8\epsilon^2 d\sigma^2}{n}$ -net $N \subset [0,\hat{\eta}]$ for F_{TbT} with $|N| = O(\frac{tn}{\epsilon^2 d})$. That means, for any $\eta \in [0,\hat{\eta}]$,

$$|F_{TbT}(\eta) - F_{TbT}(\eta')| \le \frac{8\epsilon^2 d\sigma^2}{n},$$

for $\eta' = \arg\min_{\eta'' \in N, \eta'' \leq \eta} (\eta - \eta'')$.

We also construct an ϵ -net for the empirical meta objective. The proof is very similar as in Lemma 13. We leave its proof into Section C.4.

Lemma 36. Let $\hat{\eta}$ be as defined in Definition 2 with $1 > \epsilon > 0$. Assume the conditions in Lemma 33 hold. Assume $n \ge 40d$. With probability at least $1 - m \exp(-\Omega(n))$, there exists an $\frac{\epsilon^2 d\sigma^2}{n}$ -net $N' \subset [0, \hat{\eta}]$ for \hat{F}_{TbT} with $|N'| = O(\frac{tn}{\epsilon^2 d} + m)$. That means, for any $\eta \in [0, \hat{\eta}]$,

$$|\hat{F}_{TbT}(\eta) - \hat{F}_{TbT}(\eta')| \le \frac{\epsilon^2 d\sigma^2}{n},$$

for $\eta' = \arg\min_{\eta'' \in N', \eta'' < \eta} (\eta - \eta'')$.

Combing the above three lemmas, we give the proof of Lemma 30.

Proof of Lemma 30. We assume σ as a constant in this proof. By Lemma 34, we know with probability at least $1-\exp(-\Omega(m\epsilon^4d^2/n^2)), \ \left|\hat{F}_{TbT}(\eta)-F_{TbT}(\eta)\right| \leq \frac{\epsilon^2d\sigma^2}{n}$ for any fixed η . By Lemma 35, we know as long as $n\geq c\log(\frac{n}{\epsilon d})d$ for some constant c, there exists an $\frac{8\epsilon^2d\sigma^2}{n}$ -net N for F_{TbT} with size $O(\frac{tn}{\epsilon^2d})$. By Lemma 36, we know with probability at least $1-m\exp(-\Omega(n))$, there exists an $\frac{\epsilon^2d\sigma^2}{n}$ -net N' for \hat{F}_{TbT} with size $O(\frac{tn}{\epsilon^2d}+m)$. It's not hard to verify that $N\cup N'$ is still an $\frac{8\epsilon^2d\sigma^2}{n}$ -net for \hat{F}_{TbV} and F_{TbV} . That means, for any $\eta\in[0,\hat{\eta}]$, we have

$$|F_{TbT}(\eta) - F_{TbT}(\eta')|, |\hat{F}_{TbT}(\eta) - \hat{F}_{TbT}(\eta')| \le \frac{8\epsilon^2 d\sigma^2}{n},$$

for $\eta' = \arg\min_{\eta'' \in N \cup N', \eta'' \le \eta} (\eta - \eta'')$.

Taking a union bound over $N \cup N'$, we have with probability at least $1 - O(\frac{tn}{\epsilon^2 d} + m) \exp(-\Omega(m\epsilon^4 d^2/n^2))$,

$$\left| \hat{F}_{TbT}(\eta) - F_{TbT}(\eta) \right| \le \frac{\epsilon^2 d\sigma^2}{n}$$

for all $\eta \in N \cup N'$.

Overall, we know with probability at least $1 - m \exp(-\Omega(n)) - O(\frac{tn}{\epsilon^2 d} + m) \exp(-\Omega(m\epsilon^4 d^2/n^2))$, for all $\eta \in [0, \hat{\eta}]$,

$$|F_{TbT}(\eta) - \hat{F}_{TbT}(\eta)| \le |F_{TbT}(\eta) - F_{TbT}(\eta')| + |\hat{F}_{TbT}(\eta) - \hat{F}_{TbT}(\eta')| + |\hat{F}_{TbT}(\eta') - F_{TbT}(\eta')| \le \frac{17\epsilon^2 d\sigma^2}{n},$$

where $\eta' = \arg\min_{\eta'' \in N \cup N', \eta'' \leq \eta} (\eta - \eta'')$.

C.4 Proofs of Technical Lemmas

Proof of Lemma 31. According to Lemma 52, we know with probability at least $1 - 2 \exp(-t^2/2)$,

$$\sqrt{n} - \sqrt{d} - t \le \sigma_i(X_{\text{train}}) \le \sqrt{n} + \sqrt{d} + t$$

for all $i \in [d]$. Since $d \le \frac{\epsilon^2 n}{10}$, we have $\sqrt{n} - \frac{\epsilon \sqrt{n}}{\sqrt{10}} - t \le \sigma_i(X_{\text{train}}) \le \sqrt{n} + \frac{\epsilon \sqrt{n}}{\sqrt{10}} + t$. Choosing $t = (\frac{1}{3} - \frac{1}{\sqrt{10}})\epsilon \sqrt{n}$, we have with probability at least $1 - \exp(-\Omega(\epsilon^2 n))$,

$$(1 - \frac{\epsilon}{3})\sqrt{n} \le \sigma_i(X_{\text{train}}) \le (1 + \frac{\epsilon}{3})\sqrt{n}.$$

Since $\lambda_i(H_{\text{train}}) = 1/n\sigma_i^2(X_{\text{train}})$, we have $1 - \epsilon \le \lambda_i(H_{\text{train}}) \le 1 + \epsilon$.

Proof of Lemma 32. The proof is almost the same as in Lemma 9. We omit the details here. Basically, in Lemma 9, the only property we rely on is that the norm threshold is larger than $2 \| w_{\text{train}} \|$ conditioning on \mathcal{E}_1 . Conditioning on \mathcal{E}_1 , we know $\| w_{\text{train}} \| \le 5\sigma$. Recall that the norm threshold is still set as 40σ . So this property is preserved and the previous proof works.

Proof of Lemma 33. The proof is very similar as in Lemma 10. Conditioning on \mathcal{E}_1 , we know $||H_{\text{train}}|| \leq 3/2$ and $||w_{\text{train}}|| \leq 5\sigma$. So the GD sequence never exceeds the norm threshold 40σ for any $\eta \leq 4/3$. That means,

$$\mathbb{E}\frac{1}{2}\left\|w_{t,\eta} - w_{\text{train}}\right\|_{H_{\text{train}}}^2 \mathbb{1}\left\{\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\eta)\right\} = 0$$

for all $\eta < 4/3$.

To lower bound the loss for large step size, we need to first lower bound $\|w_{\text{train}}\|$. Recall that $w_{\text{train}} = w^* + (X_{\text{train}})^\dagger \xi_{\text{train}}$. Conditioning on \mathcal{E}_1 , we know $\|\xi_{\text{train}}\| \le 2\sqrt{n}\sigma$ and $\sigma_d(X_{\text{train}}) \ge \sqrt{n}/2$, which implies $\|(X_{\text{train}})^\dagger\| \le 2/\sqrt{n}$. By Johnson-Lindenstrauss Lemma (Lemma 53), we have $\|\text{Proj}_{X_{\text{train}}}\xi_{\text{train}}\| \le \frac{3}{2}\sqrt{d/n} \|\xi_{\text{train}}\|$ with probability at least $1 - \exp(-\Omega(d))$. Call this event \mathcal{E}_3 . Conditioning on $\mathcal{E}_1 \cap \mathcal{E}_3$, we have

$$\left\|(X_{\text{train}})^{\dagger}\xi_{\text{train}}\right\| \leq 2\sqrt{n}\sigma\frac{2}{\sqrt{n}}\frac{3}{2}\sqrt{\frac{d}{n}} \leq 6\sqrt{\frac{d}{n}}\sigma,$$

which is smaller than 1/2 as long as $n \ge 12^2 d\sigma^2$. Note that we assume σ is a constant. This then implies $||w_{\text{train}}|| \ge 1/2$.

Let $\{w'_{\tau,\eta}\}$ be the GD sequence without truncation. For any step size $\eta \in [6,\infty]$, conditioning on $\mathcal{E}_1 \cap \mathcal{E}_3$, we have

$$\|w'_{t,\eta}\| \ge \left((6 \times \frac{1}{2} - 1)^t - 1 \right) \|w_{\text{train}}\| \ge \left(2^t - 1 \right) \frac{1}{2} \ge 40\sigma,$$

where the last inequality holds as long as $t \geq c_2$ for some constant c_2 . Therefore, we know when $\eta \in [6, \infty)$, $\mathbb{1} \{\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\eta)\} = \mathbb{1} \{\mathcal{E}_1 \cap \mathcal{E}_3\}$. Assuming $n \geq 40d$, we know \mathcal{E}_1 holds with probability at least $1 - \exp(-\Omega(n))$. Then, we have for any $\eta \geq 6$,

$$\mathbb{E}\frac{1}{2}\left\|w_{t,\eta} - w_{\text{train}}\right\|_{H_{\text{train}}}^{2} \mathbb{1}\left\{\mathcal{E}_{1} \cap \bar{\mathcal{E}}_{2}(\eta)\right\} \geq \frac{1}{4}\left(40\sigma - 5\sigma\right)^{2} \Pr[\mathcal{E}_{1} \cap \mathcal{E}_{3}] \geq \frac{\epsilon^{2}d\sigma^{2}}{n},$$

where the last inequality assumes $n \ge c, d \ge c_4$ for some constant c, c_4 .

Overall, we know $\mathbb{E}\frac{1}{2}\|w_{t,\eta}-w_{\text{train}}\|_{H_{\text{train}}}^2\mathbb{1}\left\{\mathcal{E}_1\cap\bar{\mathcal{E}}_2(\eta)\right\}$ equals zero for all $\eta\in[0,4/3]$ and is at least $\frac{\epsilon^2d\sigma^2}{n}$ for all $\eta\in[6,\infty)$. By definition, we know $\hat{\eta}\in(4/3,6)$.

Proof of Lemma 35. By Lemma 33, we know $\hat{\eta}$ is a constant. The proof is very similar as in Lemma 12. Let \mathcal{E}_1 and $\bar{\mathcal{E}}_2(\eta)$ be as defined in Definition 2. For the simplicity of the proof, we assume $\mathbb{E}\frac{1}{2}\|w_{t,\hat{\eta}}-w_{\text{train}}\|_{H_{\text{train}}}^2\mathbb{I}\left\{\mathcal{E}_1\cap\bar{\mathcal{E}}_2(\hat{\eta})\right\}\leq \frac{\epsilon^2d\sigma^2}{n}$. The other case can be resolved using techniques in the proof of Lemma 12.

Recall the population meta objective

$$F_{TbT}(\eta) = \mathbb{E}\frac{1}{2} \left\| w_{t,\eta} - w_{\text{train}} \right\|_{H_{\text{train}}}^2 + \frac{n-d}{2n} \sigma^2.$$

Therefore, we only need to construct an ϵ -net for the first term.

We can divide $\mathbb{E}_{\frac{1}{2}} \| w_{t,\eta} - w_{\text{train}} \|_{H_{\text{train}}}^2$ as follows,

$$\begin{split} & \mathbb{E}\frac{1}{2}\left\|w_{t,\eta} - w_{\text{train}}\right\|_{H_{\text{train}}}^2 \\ = & \mathbb{E}\frac{1}{2}\left\|w_{t,\eta} - w_{\text{train}}\right\|_{H_{\text{train}}}^2 \mathbb{1}\left\{\mathcal{E}_1 \cap \mathcal{E}_2(\hat{\eta})\right\} + \mathbb{E}\frac{1}{2}\left\|w_{t,\eta} - w_{\text{train}}\right\|_{H_{\text{train}}}^2 \mathbb{1}\left\{\mathcal{E}_1 \cap \bar{\mathcal{E}}_2(\hat{\eta})\right\} \\ & + \mathbb{E}\frac{1}{2}\left\|w_{t,\eta} - w_{\text{train}}\right\|_{H_{\text{train}}}^2 \mathbb{1}\left\{\bar{\mathcal{E}}_1\right\}. \end{split}$$

We will construct an ϵ -net for the first term and show the other two terms are small. Let's first consider the third term. Assuming $n \geq 40d$, we know $\Pr[\mathcal{E}_1] \leq \exp(-\Omega(n))$. Since $\frac{1}{2} \|w_{t,\eta} - w_{\text{train}}\|_{H_{\text{train}}}^2$ is O(1)-subexponential, by Cauchy-Schwarz inequality, we have $\mathbb{E}\frac{1}{2} \|w_{t,\eta} - w_{\text{train}}\|_{H_{\text{train}}}^2 \mathbb{E}\left\{\bar{\mathcal{E}}_1\right\} = O(1) \exp(-\Omega(n))$. Choosing $n \geq c \log(n/(\epsilon d))$ for some constant c, we know $\frac{1}{2} \|w_{t,\hat{\eta}} - w_{\text{train}}\|_{H_{\text{train}}}^2 \mathbb{E}\left\{\bar{\mathcal{E}}_1\right\} \leq \frac{\epsilon^2 d\sigma^2}{n}$.

Then we upper bound the second term. Since $\mathbb{E}\frac{1}{2}\|w_{t,\hat{\eta}}-w_{\text{train}}\|_{H_{\text{train}}}^2\mathbb{1}\left\{\mathcal{E}_1\cap\bar{\mathcal{E}}_2(\hat{\eta})\right\}\leq \frac{\epsilon^2d\sigma^2}{n}$ and $\frac{1}{2}\|w_{t,\hat{\eta}}-w_{\text{train}}\|_{H_{\text{train}}}^2\geq \frac{35^2\sigma^2}{4}$ when $w_{t,\hat{\eta}}$ diverges, we know $\Pr[\mathcal{E}_1\cap\bar{\mathcal{E}}_2(\hat{\eta})]\leq \frac{4\epsilon^2d}{35^2n}$. Then, we can upper bound the second term as follows,

$$\mathbb{E}\frac{1}{2}\left\|w_{t,\eta} - w_{\text{train}}\right\|_{H_{\text{train}}}^{2} \mathbb{1}\left\{\mathcal{E}_{1} \cap \bar{\mathcal{E}}_{2}(\hat{\eta})\right\} \leq \frac{3 \times 45^{2}\sigma^{2}}{4} \frac{4\epsilon^{2}d}{35^{2}n} \leq \frac{6\epsilon^{2}d\sigma^{2}}{n}$$

Next, similar as in Lemma 12, we can show the first term $\frac{1}{2}\|w_{t,\eta}-w_{\text{train}}\|_{H_{\text{train}}}^2\mathbbm{1}\{\mathcal{E}_1\cap\mathcal{E}_2(\hat{\eta})\}$ is O(t)-lipschitz. Therefore, there exists an $\frac{\epsilon^2 d\sigma^2}{n}$ -net N for $\mathbb{E}\frac{1}{2}\|w_{t,\eta}-w_{\text{train}}\|_{H_{\text{train}}}^2\mathbbm{1}\{\mathcal{E}_1\cap\mathcal{E}_2(\hat{\eta})\}$ with size $O(\frac{tn}{\epsilon^2 d})$. That means, for any $\eta\in[0,\hat{\eta}]$,

$$\left| \mathbb{E}\frac{1}{2} \left\| w_{t,\eta} - w_{\text{train}} \right\|_{H_{\text{train}}}^2 \mathbb{1} \left\{ \mathcal{E}_1 \cap \mathcal{E}_2(\hat{\eta}) \right\} - \mathbb{E}\frac{1}{2} \left\| w_{t,\eta'} - w_{\text{train}} \right\|_{H_{\text{train}}}^2 \mathbb{1} \left\{ \mathcal{E}_1 \cap \mathcal{E}_2(\hat{\eta}) \right\} \right| \leq \frac{\epsilon^2 d\sigma^2}{n}$$

for $\eta' = \arg\min_{\eta'' \in N, \eta'' < \eta} (\eta - \eta'')$.

Combing with the upper bounds on the second term and the third term, we have for any $\eta \in [0, \hat{\eta}]$,

$$|F_{TbT}(\eta) - F_{TbT}(\eta')| \le \frac{8\epsilon^2 d\sigma^2}{n}$$

for $\eta' = \arg\min_{\eta'' \in N, \eta'' < \eta} (\eta - \eta'')$.

Proof of Lemma 36. By Lemma 33, we know $\hat{\eta}$ is a constant. For each $k \in [m]$, let $\mathcal{E}_{1,k}$ be the event that $\sqrt{n}/2 \leq \sigma_i(X_{\text{train}}^{(k)}) \leq 3\sqrt{n}/2$ and $1/2 \leq \lambda_i(H_{\text{train}}^{(k)}) \leq 3/2$ for all $i \in [d]$ and $\sqrt{n}\sigma/2 \leq \left\|\xi_{\text{train}}^{(k)}\right\| \leq 2\sqrt{n}\sigma$. Assuming $n \geq 40d$, by Lemma 31, we know with probability at least $1 - m \exp(-\Omega(n))$, $\mathcal{E}_{1,k}$'s hold for all $k \in [m]$.

Then, similar as in Lemma 13, there exists an $\frac{\epsilon^2 d\sigma^2}{n}$ -net N' with $|N'| = O(\frac{nt}{\epsilon^2 d} + m)$ for \hat{F}_{TbT} . That means, for any $\eta \in [0, \hat{\eta}]$,

$$\left| \hat{F}_{TbT}(\eta) - \hat{F}_{TbT}(\eta') \right| \le \frac{\epsilon^2 d\sigma^2}{n}$$

for $\eta' = \arg\min_{\eta'' \in N', \eta'' \leq \eta} (\eta - \eta'')$.

D Proofs of train-by-train v.s. train-by-validation (SGD)

Previously, we have shown that train-by-validation generalizes better than train-by-train when the tasks are trained by GD and when the number of samples is small. In this section, we show a similar phenomenon also appears in the SGD setting.

In the train-by-train setting, each task P contains a training set $S_{\text{train}} = \{(x_i, y_i)\}_{i=1}^n$. The inner objective is defined as $\hat{f}(w) = \frac{1}{2n} \sum_{(x,y) \in S_{\text{train}}} (\langle w, x \rangle - y)^2$. Let $\{w_{\tau,\eta}\}$ be the SGD sequence running on $\hat{f}(w)$ from initialization 0 (without truncation). That means, $w_{\tau,\eta} = w_{\tau-1,\eta} - \eta \hat{\nabla} \hat{f}(w_{\tau-1,\eta})$, where $\hat{\nabla} \hat{f}(w_{\tau-1,\eta}) = \left(\left\langle w_{\tau-1,\eta}, x_{i(\tau-1)} \right\rangle - y_{i(\tau-1)} \right) x_{i(\tau-1)}$. Here index $i(\tau-1)$ is independently and uniformly sampled from [n]. We denote the SGD noise as $n_{\tau-1,\eta} := \hat{\nabla} \hat{f}(w_{\tau-1,\eta}) - \nabla \hat{f}(w_{\tau-1,\eta})$. The meta-loss on task P is defined as follows,

$$\Delta_{TbT(n)}(\eta, P) = \mathbb{E}_{SGD}\hat{f}(w_{t,\eta}) = \mathbb{E}_{SGD}\frac{1}{2n} \sum_{(x,y) \in S_{train}} \left(\langle w_{t,\eta}, x \rangle - y \right)^2,$$

where the expectation is taken over the SGD noise. Note $w_{t,\eta}$ depends on the SGD noise along the trajectory. Then, the empirical meta objective $\hat{F}_{TbT(n)}(\eta)$ is the average of the meta-loss across m different specific tasks

$$\hat{F}_{TbT(n)}(\eta) = \frac{1}{m} \sum_{k=1}^{m} \Delta_{TbT(n)}(\eta, P_k).$$
(6)

In order to control the SGD noise in expectation, we restrict the feasible set of step sizes into O(1/d). We show within this range, the optimal step size under $\hat{F}_{TbT(n)}$ is $\Omega(1/d)$ and the learned weight is far from ground truth w^* on new tasks. We prove Theorem 9 in Section D.1.

Theorem 9. Let the meta objective $\hat{F}_{TbT(n)}$ be as defined in Equation 6 with $n \in [d/4, 3d/4]$. Suppose σ is a constant. Assume unroll length $t \ge c_2d$ and dimension $d \ge c_4 \log(m)$ for certain constants c_2, c_4 . Then, with probability at least 0.99 in the sampling of training tasks P_1, \dots, P_m and test task P,

$$\eta_{train}^* = \Omega(1/d)$$
 and $\mathbb{E}_{SGD} \left\| w_{t,\eta_{rain}^*} - w^* \right\|^2 = \Omega(\sigma^2),$

for all $\eta_{train}^* \in \arg\min_{0 \le \eta \le \frac{1}{2L^3d}} \hat{F}_{TbT(n)}(\eta)$, where L = 100 and w_{t,η_{train}^*} is trained by running SGD on test task P.

In the train-by-validation setting, each task P contains a training set S_{train} with n_1 samples and a validation set with n_2 samples. The inner objective is defined as $\hat{f}(w) = \frac{1}{2n_1} \sum_{(x,y) \in S_{\text{train}}} (\langle w,x \rangle - y)^2$. Let $\{w_{\tau,\eta}\}$ be the SGD sequence running on $\hat{f}(w)$ from initialization 0 (with the same truncation defined in Section 4). For each task P, the meta-loss $\Delta_{TbV(n_1,n_2)}(\eta,P)$ is defined as

$$\Delta_{TbV(n_1,n_2)}(\eta,P) = \mathbb{E}_{SGD} \frac{1}{2n_2} \sum_{(x,y) \in S_{valid}} \left(\langle w_{t,\eta}, x \rangle - y \right)^2.$$

The empirical meta objective $\hat{F}_{TbV(n_1,n_2)}(\eta)$ is the average of the meta-loss across m different tasks $P_1,P_2,...,P_m$,

$$\hat{F}_{TbV(n_1,n_2)}(\eta) = \frac{1}{m} \sum_{k=1}^{m} \Delta_{TbV(n_1,n_2)}(\eta, P_k). \tag{7}$$

In order to bound the SGD noise with high probability, we restrict the feasible set of the step sizes into $O(\frac{1}{d^2 \log^2 d})$. Within this range, we prove the optimal step size under $\hat{F}_{TbV(n_1,n_2)}$ is $\Theta(1/t)$ and the learned weight is better than initialization 0 by a constant on new tasks. Theorem 10 is proved in Section D.2.

Theorem 10. Let the meta objective $\hat{F}_{TbV(n_1,n_2)}$ be as defined in Equation 7 with $n_1, n_2 \in [d/4, 3d/4]$. Assume noise level σ is a large constant c_1 . Assume unroll length $t \geq c_2 d^2 \log^2(d)$, number of training tasks $m \geq c_3$ and dimension $d \geq c_4$ for certain constants c_2, c_3, c_4 . There exists constant c_5 such that with probability at least 0.99 in the sampling of training tasks, we have

$$\eta_{valid}^* = \Theta(1/t) \text{ and } \mathbb{E} \left\| w_{t,\eta_{valid}^*} - w^* \right\|^2 = \left\| w^* \right\|^2 - \Omega(1)$$

for all $\eta_{valid}^* \in \arg\min_{0 \le \eta \le \frac{1}{c_5 d^2 \log^2(d)}} \hat{F}_{TbV(n_1, n_2)}(\eta)$, where the expectation is taken over the new tasks and SGD noise.

Notations: In the following proofs, we use the same set of notations defined in Appendix B. We use $\mathbb{E}_{P \sim T}$ to denote the expectation over the sampling of tasks and use \mathbb{E}_{SGD} to denote the expectation over the SGD noise. We use \mathbb{E} to denote $\mathbb{E}_{P \sim T} \mathbb{E}_{SGD}$. Same as in Appendix B, we use letter L to denote constant 100, which upper bounds $\|H_{train}\|$ with high probability.

D.1 Train-by-train (SGD)

Recall Theorem 9 as follows.

Theorem 9. Let the meta objective $\hat{F}_{TbT(n)}$ be as defined in Equation 6 with $n \in [d/4, 3d/4]$. Suppose σ is a constant. Assume unroll length $t \geq c_2d$ and dimension $d \geq c_4\log(m)$ for certain constants c_2, c_4 . Then, with probability at least 0.99 in the sampling of training tasks P_1, \dots, P_m and test task P,

$$\eta_{train}^* = \Omega(1/d) \text{ and } \mathbb{E}_{SGD} \left\| w_{t,\eta_{train}^*} - w^* \right\|^2 = \Omega(\sigma^2),$$

for all $\eta_{train}^* \in \arg\min_{0 \leq \eta \leq \frac{1}{2L^3d}} \hat{F}_{TbT(n)}(\eta)$, where L = 100 and w_{t,η_{train}^*} is trained by running SGD on test task P.

In order to prove Theorem 9, we first show that η^*_{train} is $\Omega(1/d)$ in Lemma 37. The proof is similar as in the GD setting. As long as $\eta = O(1/d)$, the SGD noise is dominated by the full gradient. Then, we can show that $\Delta_{TbT}(\eta, P)$ is roughly $(1 - \Theta(1)\eta)^t$, which implies that $\eta^*_{\text{train}} = \Omega(1/d)$. We leave the proof of Lemma 37 into Section D.1.1.

Lemma 37. Assume $t \ge c_2 d$ with certain constant c_2 . With probability at least $1 - m \exp(-\Omega(d))$ in the sampling of m training tasks,

$$\eta_{train}^* \ge \frac{1}{6L^5d},$$

for all $\eta_{train}^* \in \arg\min_{0 \le \eta \le \frac{1}{2L^3d}} \hat{F}_{TbT}(\eta)$.

Let $P = (\mathcal{D}(w^*), S_{\text{train}}, \ell)$ be an independently sampled test task with $|S_{\text{train}}| = n \in [d/4, 3d/4]$. For any step size $\eta \in [\frac{1}{6L^5d}, \frac{1}{2L^3d}]$, let $w_{t,\eta}$ be the weight obtained by running SGD on $\hat{f}(w)$ for t steps. Next, we show $\mathbb{E}_{\text{SGD}} \|w_{t,\eta} - w^*\|^2 = \Omega(\sigma^2)$ with high probability in the sampling of P.

Lemma 38. Suppose σ is a constant. Assume unroll length $t \ge c_2 d$ for some constant c_2 . With probability at least $1 - \exp(-\Omega(d))$ in the sampling of test task P,

$$\mathbb{E}_{SGD} \left\| w_{t,\eta} - w^* \right\|^2 \ge \frac{\sigma^2}{128L},$$

for all $\eta \in [\frac{1}{6L^5d}, \frac{1}{2L^3d}]$, where $w_{t,\eta}$ is obtained by running SGD on task P for t iterations.

With Lemma Lemma 37 and Lemma 38, the proof of Theorem 9 is straightforward.

Proof of Theorem 9. Combing Lemma 37 and Lemma 38, we know as long as σ is a constant, $t \geq c_2 d$, $d \geq c_4 \log(m)$, with probability at least 0.99, $\eta^*_{\text{train}} = \Omega(1/d)$ and $\mathbb{E}_{\text{SGD}} \|w_{t,\eta^*_{\text{train}}} - w^*\|^2 = \Omega(\sigma^2)$, for all $\eta^*_{\text{train}} \in \arg\min_{0 \leq \eta \leq \frac{1}{2L^3d}} \hat{F}_{TbT}(\eta)$.

D.1.1 Detailed Proofs

Proof of Lemma 37. The proof is very similar to the proof of Lemma 1 except that we need to bound the SGD noise term. For each $k \in [m]$, let \mathcal{E}_k be the event that $\sqrt{d}/\sqrt{L} \leq \sigma_i(X_{\text{train}}) \leq \sqrt{Ld}$ and $1/L \leq \lambda_i(H_{\text{train}}) \leq L$ for all $i \in [n]$ and $\sqrt{d}\sigma/4 \leq \|\xi_{\text{train}}\| \leq \sqrt{d}\sigma$. According to Lemma 7 and Lemma 49, we know for each $k \in [m]$, \mathcal{E}_k happens with probability at least $1 - \exp(-\Omega(d))$. Taking a union bound over all $k \in [m]$, we know $\cap_{k \in [m]} \mathcal{E}_k$ holds with probability at least $1 - m \exp(-\Omega(d))$. From now on, we assume $\cap_{k \in [m]} \mathcal{E}_k$ holds.

For each $k \in [m]$, we have

$$\Delta_{TbT}(\eta, P_k) := \frac{1}{2} \mathbb{E}_{\text{SGD}} \left\| w_{t,\eta}^{(k)} - w_{\text{train}}^{(k)} \right\|_{H_{\text{train}}^{(k)}}^2$$

Since $1/L \le \lambda_i(H_{\text{train}}^{(k)}) \le L$ and $(w_{t,\eta}^{(k)} - w_{\text{train}}^{(k)})$ is in the span of $H_{\text{train}}^{(k)}$, we have

$$\frac{1}{2L} \mathbb{E}_{\text{SGD}} \left\| w_{t,\eta}^{(k)} - w_{\text{train}}^{(k)} \right\|^2 \leq \Delta_{TbT}(\eta, P_k) \leq \frac{L}{2} \mathbb{E}_{\text{SGD}} \left\| w_{t,\eta}^{(k)} - w_{\text{train}}^{(k)} \right\|^2.$$

Recall the updates of stochastic gradient descent,

$$w_{t,\eta}^{(k)} - w_{\text{train}}^{(k)} = (I - \eta H_{\text{train}}^{(k)})(w_{t-1,\eta}^{(k)} - w_{\text{train}}^{(k)}) - \eta n_{t-1,\eta}^{(k)}.$$

Therefore,

$$\mathbb{E}_{\text{SGD}}\left[\left\|w_{t,\eta}^{(k)} - w_{\text{train}}^{(k)}\right\|^2 \left|w_{t-1,\eta}^{(k)}\right] = \left\|(I - \eta H_{\text{train}}^{(k)})(w_{t-1,\eta}^{(k)} - w_{\text{train}}^{(k)})\right\|^2 + \eta^2 \mathbb{E}_{\text{SGD}}\left[\left\|n_{t-1,\eta}^{(k)}\right\|^2 \left|w_{t-1,\eta}^{(k)}\right|\right].$$

We know for any $\eta \leq 1/L$,

$$(1-2\eta L) \left\| w_{t-1,\eta}^{(k)} - w_{\text{train}}^{(k)} \right\|^2 \leq \left\| (I-\eta H_{\text{train}}^{(k)}) (w_{t-1,\eta}^{(k)} - w_{\text{train}}^{(k)}) \right\|^2 \leq (1-\frac{\eta}{L}) \left\| w_{t-1,\eta}^{(k)} - w_{\text{train}}^{(k)} \right\|^2.$$

The noise can be bounded as follows,

$$\begin{split} & \eta^{2}\mathbb{E}_{\text{SGD}}\left[\left\|n_{t-1,\eta}^{(k)}\right\|^{2}|w_{t-1,\eta}^{(k)}\right] \\ = & \eta^{2}\mathbb{E}_{\text{SGD}}\left[\left\|x_{i(t-1)}x_{i(t-1)}^{\intercal}(w_{t-1,\eta}^{(k)} - w_{\text{train}}^{(k)}) - H_{\text{train}}^{(k)}(w_{t-1,\eta}^{(k)} - w_{\text{train}}^{(k)})\right\|^{2}|w_{t-1,\eta}^{(k)}| \\ \leq & \eta^{2}\mathbb{E}_{\text{SGD}}\left[\left\|x_{i(t-1)}x_{i(t-1)}^{\intercal}(w_{t-1,\eta}^{(k)} - w_{\text{train}}^{(k)})\right\|^{2}|w_{t-1,\eta}^{(k)}| \\ \leq & \eta^{2}\max_{i(t-1)}\left\|x_{i(t-1)}\right\|^{2}\left\|w_{t-1,\eta}^{(k)} - w_{\text{train}}^{(k)}\right\|_{H_{\text{train}}^{(k)}}^{2}. \end{split}$$

Since $||X_{\text{train}}|| \leq \sqrt{L}\sqrt{d}$, we immediately know $\max_{i(t-1)} ||x_{i(t-1)}|| \leq \sqrt{L}\sqrt{d}$. Therefore, we can bound the noise as follows,

$$\eta^{2} \mathbb{E}_{SGD} \left[\left\| n_{t-1,\eta}^{(k)} \right\|^{2} |w_{t-1,\eta}^{(k)}| \right] \leq \eta^{2} \max_{i(t-1)} \left\| x_{i(t-1)} \right\|^{2} \left\| w_{t-1,\eta}^{(k)} - w_{\text{train}}^{(k)} \right\|_{H_{\text{train}}^{(k)}}^{2}$$
$$\leq L^{2} \eta^{2} d \left\| w_{t-1,\eta}^{(k)} - w_{\text{train}}^{(k)} \right\|^{2}.$$

As long as $\eta \leq \frac{1}{2L^3d}$, we have

$$(1 - \eta L) \left\| w_{t-1,\eta}^{(k)} - w_{\text{train}}^{(k)} \right\|^2 \leq \mathbb{E}_{\text{SGD}} \left[\left\| w_{t,\eta}^{(k)} - w_{\text{train}}^{(k)} \right\|^2 | w_{t-1,\eta}^{(k)} \right] \leq (1 - \frac{\eta}{2L}) \left\| w_{t-1,\eta}^{(k)} - w_{\text{train}}^{(k)} \right\|^2.$$

This further implies

$$(1 - \eta L)^t \|w_{\text{train}}\|^2 \le \mathbb{E}_{\text{SGD}} \|w_{t,\eta}^{(k)} - w_{\text{train}}^{(k)}\|^2 \le (1 - \frac{\eta}{2L})^t \|w_{\text{train}}\|^2.$$

Let $\eta_2 := \frac{1}{2L^3d}$, we have

$$\Delta_{TbT}(\eta, P_k) \le \frac{L}{2} \left(1 - \frac{1}{4L^4d}\right)^t \left\| w_{\text{train}} \right\|^2$$

Let $\eta_1 := \frac{1}{6L^5d}$, for all $\eta \in [0, \eta_1]$ we have

$$\Delta_{TbT}(\eta, P_k) \ge \frac{1}{2L} (1 - \frac{1}{6L^4d})^t \|w_{\text{train}}\|^2$$

As long as $t \ge c_2 d$ for certain constant c_2 , we know

$$\frac{1}{2L}(1 - \frac{1}{6L^4d})^t \left\| w_{\text{train}} \right\|^2 > \frac{L}{2}(1 - \frac{1}{4L^4d})^t \left\| w_{\text{train}} \right\|^2.$$

As this holds for all $k \in [m]$ and $\hat{F}_{TbT} = 1/m \sum_{i=1}^{m} \Delta_{TbT}(\eta, P_k)$, we know the optimal step size η^*_{train} is within $[\frac{1}{6L^5d},\frac{1}{2L^3d}].$ We rely the following technical lemma to prove Lemma 38.

Lemma 39. Suppose σ is a constant. Given any $\epsilon > 0$, with probability at least $1 - O(1/\epsilon) \exp(-\Omega(\epsilon^2 d))$,

$$\left|\left\langle B_{t,\eta}w_{train}^* - w^*, B_{t,\eta}(X_{train})^{\dagger} \xi_{train}\right\rangle\right| \leq \epsilon,$$

for all $\eta \in [0, \frac{1}{2L^3d}]$.

Proof of Lemma 39. By Lemma 7, with probability at least $1 - \exp(-\Omega(d))$, $\sqrt{d}/\sqrt{L} \le \sigma_i(X_{\text{train}}) \le \sqrt{Ld}$ and $1/L \le \lambda_i(H_{\text{train}}) \le L$ for all $i \in [n]$. Therefore $\|[(X_{\text{train}})^\dagger]^\top B_{t,\eta}(B_{t,\eta}w_{\text{train}}^* - w^*)\| \le 2\sqrt{L}/\sqrt{d}$. Notice that ξ_{train} is independent with $[(X_{\text{train}})^\dagger]^\top B_{t,\eta}(B_{t,\eta}w_{\text{train}}^* - w^*)$. By Hoeffding's inequality, with probability at least $1 - \exp(-\Omega(\epsilon^2 d)),$

$$\left|\left\langle [(X_{\mathrm{train}})^{\dagger}]^{\top} B_{t,\eta}(B_{t,\eta} w_{\mathrm{train}}^* - w^*), \xi_{\mathrm{train}} \right\rangle \right| \leq \epsilon.$$

Next, we construct an ϵ -net for η and show the crossing term is small for all $\eta \in [0, \frac{1}{2L^3d}]$. For simplicity, denote $g(\eta) := \langle B_{t,\eta} w_{\text{train}}^* - w^*, B_{t,\eta}(X_{\text{train}})^{\dagger} \xi_{\text{train}} \rangle$. Taking the derivative of $g(\eta)$, we have

$$\begin{split} g'(\eta) = & t \left\langle H_{\text{train}} (I - \eta H_{\text{train}})^{t-1} w_{\text{train}}^*, B_{t,\eta} (X_{\text{train}})^\dagger \xi_{\text{train}} \right\rangle \\ & + t \left\langle B_{t,\eta} w_{\text{train}}^* - w^*, H_{\text{train}} (I - \eta H_{\text{train}})^{t-1} (X_{\text{train}})^\dagger \xi_{\text{train}} \right\rangle \end{split}$$

According to Lemma 49, we know with probability at least $1 - \exp(-\Omega(d))$, $\|\xi_{\text{train}}\| \le \sqrt{d\sigma}$. Therefore, the derivative $q'(\eta)$ can be bounded as follows,

$$|g'(\eta)| = O(1)t(1 - \frac{\eta}{L})^{t-1}$$

Similar as in Lemma 18, there exists an ϵ -net N_{ϵ} with size $O(1/\epsilon)$ such that for any $\eta \in [0, \frac{1}{3L^3d}]$, there exists $\eta' \in N_{\epsilon}$ with $|g(\eta) - g(\eta')| \le \epsilon$. Taking a union bound over N_{ϵ} , we have with probability at least $1 - O(1/\epsilon) \exp(-\Omega(\epsilon^2 d))$, for every $\eta \in N_{\epsilon}$,

$$\left|\left\langle B_{t,\eta} w_{\text{train}}^* - w^*, B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}} \right\rangle\right| \leq \epsilon.$$

which implies for every $\eta \in [0, \frac{1}{3L^3d}]$

$$\left|\left\langle B_{t,\eta} w_{\text{train}}^* - w^*, B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}} \right\rangle\right| \leq 2\epsilon.$$

Changing ϵ to $\epsilon'/2$ finishes the proof.

Proof of Lemma 38. According to Lemma 7 and Lemma 49, we know with probability at least $1 - \exp(-\Omega(d))$, $\sqrt{d}/\sqrt{L} \le \sigma_i(X_{\text{train}}) \le \sqrt{Ld}$ and $1/L \le \lambda_i(H_{\text{train}}) \le L$ for all $i \in [n]$ and $\sqrt{d}\sigma/4 \le \|\xi_{\text{train}}\| \le \sqrt{d}\sigma$. We assume these properties hold in the proof and take a union bound at the end.

Recall that $\mathbb{E}_{SGD} \|w_{t,\eta} - w^*\|^2$ can be lower bounded as follows,

$$\mathbb{E}_{\text{SGD}} \| w_{t,\eta} - w^* \|^2 = \mathbb{E}_{\text{SGD}} \left\| B_{t,\eta} (w_{\text{train}}^* + (X_{\text{train}})^{\dagger} \xi_{\text{train}}) - \eta \sum_{\tau=0}^{t-1} (I - \eta H_{\text{train}})^{t-1-\tau} n_{\tau,\eta} - w^* \right\|^2$$

$$\geq \| B_{t,\eta} (w_{\text{train}}^* + (X_{\text{train}})^{\dagger} \xi_{\text{train}}) - w^* \|^2$$

$$\geq \| B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}} \|^2 + 2 \left\langle B_{t,\eta} w_{\text{train}}^* - w^*, B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}} \right\rangle$$

For any $\eta \in \left[\frac{1}{6L^5d}, \frac{1}{2L^3d}\right]$, we can lower bound the first term as follows,

$$\begin{aligned} \left\| B_{t,\eta}(X_{\text{train}})^{\dagger} \xi_{\text{train}} \right\|^{2} &\geq \left(1 - \exp\left(-\frac{\eta t}{L} \right) \right)^{2} \frac{\sigma^{2}}{16L} \\ &\geq \left(1 - \exp\left(-\frac{t}{6L^{6}d} \right) \right)^{2} \frac{\sigma^{2}}{16L} \\ &\geq \frac{\sigma^{2}}{64L}, \end{aligned}$$

where the last inequality holds as long as $t \ge c_2 d$ for certain constant c_2 .

Choosing $\epsilon = \frac{\sigma^2}{256L}$ in Lemma 39, we know with probability at least $1 - \exp(-\Omega(d))$,

$$\left|\left\langle B_{t,\eta}w_{\text{train}}^* - w^*, B_{t,\eta}(X_{\text{train}})^{\dagger} \xi_{\text{train}} \right\rangle\right| \leq \frac{\sigma^2}{256L}$$

for all $\eta \in [0, \frac{1}{2L^3d}]$.

Overall, we have $\mathbb{E}_{SGD} \|w_{t,\eta} - w^*\|^2 \ge \frac{\sigma^2}{128L}$. Taking a union bound over all the bad events, we know this happens with probability at least $1 - \exp(-\Omega(d))$.

D.2 Train-by-validation (SGD)

Recall Theorem 10 as follows.

Theorem 10. Let the meta objective $\hat{F}_{TbV(n_1,n_2)}$ be as defined in Equation 7 with $n_1, n_2 \in [d/4, 3d/4]$. Assume noise level σ is a large constant c_1 . Assume unroll length $t \geq c_2 d^2 \log^2(d)$, number of training tasks $m \geq c_3$ and dimension $d \geq c_4$ for certain constants c_2, c_3, c_4 . There exists constant c_5 such that with probability at least 0.99 in the sampling of training tasks, we have

$$\eta_{\textit{valid}}^* = \Theta(1/t) \textit{ and } \mathbb{E} \left\| w_{t,\eta_{\textit{valid}}^*} - w^* \right\|^2 = \left\| w^* \right\|^2 - \Omega(1)$$

for all $\eta_{valid}^* \in \arg\min_{0 \le \eta \le \frac{1}{c_5 d^2 \log^2(d)}} \hat{F}_{TbV(n_1, n_2)}(\eta)$, where the expectation is taken over the new tasks and SGD noise.

To prove Theorem 10, we first study the behavior of the population meta objective F_{TbV} . That is,

$$F_{TbV}(\eta) := \mathbb{E}_{P \sim \mathcal{T}} \Delta_{TbV}(\eta, P) = \mathbb{E}_{P \sim \mathcal{T}} \mathbb{E}_{SGD} \frac{1}{2} \left\| w_{t,\eta} - w^* - (X_{\text{valid}})^{\dagger} \xi_{\text{valid}} \right\|_{H_{\text{valid}}}^{2}$$
$$= \mathbb{E}_{P \sim \mathcal{T}} \mathbb{E}_{SGD} \frac{1}{2} \left\| w_{t,\eta} - w^* \right\|^{2} + \frac{\sigma^{2}}{2}.$$

We show that the optimal step size for the population meta objective F_{TbV} is $\Theta(1/t)$ and $\mathbb{E}_{P \sim \mathcal{T}} \mathbb{E}_{SGD} \| w_{t,\eta} - w^* \|^2 = \| w^* \|^2 - \Omega(1)$ under the optimal step size.

Lemma 40. Suppose σ is a large constant c_1 . Assume $t \ge c_2 d^2 \log^2(d)$, $d \ge c_4$ for some constants c_2 , c_4 . There exist $\eta_1, \eta_2, \eta_3 = \Theta(1/t)$ with $\eta_1 < \eta_2 < \eta_3$ and constant c_5 such that

$$F_{TbV}(\eta_2) \le \frac{1}{2} \|w^*\|^2 - \frac{9}{10}C + \frac{\sigma^2}{2}$$

$$F_{TbV}(\eta) \ge \frac{1}{2} \|w^*\|^2 - \frac{6}{10}C + \frac{\sigma^2}{2}, \forall \eta \in [0, \eta_1] \cup [\eta_3, \frac{1}{c_5 d^2 \log^2(d)}]$$

where C is a positive constant.

In order to relate the behavior of F_{TbV} to \hat{F}_{TbV} , we show a generalization result from \hat{F}_{TbV} to F_{TbV} for $\eta \in [0, \frac{1}{c_5 d^2 \log^2(d/\epsilon)}]$.

Lemma 41. For any $1 > \epsilon > 0$, assume σ is a constant and $d \ge c_4 \log(1/\epsilon)$ for some constant c_4 . There exists constant c_5 such that with probability at least $1 - O(1/\epsilon) \exp(-\Omega(\epsilon^2 m))$,

$$|\hat{F}_{TbV}(\eta) - F_{TbV}(\eta)| \le \epsilon,$$

for all $\eta \in [0, \frac{1}{c_5 d^2 \log^2(d/\epsilon)}]$.

Combining Lemma 40 and Lemma 41, we give the proof of Theorem 10.

Proof of Theorem 10. The proof is almost the same as in the GD setting (Theorem 8). We omit the details here.

D.2.1 Behavior of F_{TbV} for $\eta \in [0, \frac{1}{c_5 d^2 \log^2 d}]$

In this section, we give the proof of Lemma 40. Recall the lemma as follows,

Lemma 40. Suppose σ is a large constant c_1 . Assume $t \ge c_2 d^2 \log^2(d)$, $d \ge c_4$ for some constants c_2 , c_4 . There exist $\eta_1, \eta_2, \eta_3 = \Theta(1/t)$ with $\eta_1 < \eta_2 < \eta_3$ and constant c_5 such that

$$F_{TbV}(\eta_2) \le \frac{1}{2} \|w^*\|^2 - \frac{9}{10}C + \frac{\sigma^2}{2}$$

$$F_{TbV}(\eta) \ge \frac{1}{2} \|w^*\|^2 - \frac{6}{10}C + \frac{\sigma^2}{2}, \forall \eta \in [0, \eta_1] \cup [\eta_3, \frac{1}{c_5 d^2 \log^2(d)}]$$

where C is a positive constant.

Recall that $F_{TbV}(\eta) = \mathbb{E}_{P \sim \mathcal{T}} \mathbb{E}_{SGD} 1/2 \|w_{t,\eta} - w^*\|^2 + \sigma^2/2$. Denote $Q(\eta) := \mathbb{E}_{SGD} 1/2 \|w_{t,\eta} - w^*\|^2$. Recall that we truncate the SGD sequence once the weight norm exceeds $4\sqrt{L}\sigma$. Due to the truncation, the expectation of $1/2 \|w_{t,\eta} - w^*\|^2$ over SGD noise is very tricky to analyze.

Instead, we define an auxiliary sequence $\{w'_{\tau,\eta}\}$ that is obtained by running SGD on task P without truncation and we first study $Q'(\eta) := 1/2\mathbb{E}_{\text{SGD}} \|w'_{t,\eta} - w^*\|^2$. In Lemma 42, we show that with high probability in the sampling of task P, the minimizer of $Q'(\eta)$ is $\Theta(1/t)$. The proof is very similar as the proof of Lemma 17 except that we need to bound the SGD noise at step size η_2 . We defer the proof into Section D.2.3.

Lemma 42. Given a task P, let $\{w'_{\tau,\eta}\}$ be the weight obtained by running SGD on task P without truncation. Choose σ as a large constant c_1 . Assume unroll length $t \geq c_2d$ for some constant c_2 . With probability at least $1 - \exp(-\Omega(d))$ over the sampling of task $P, \sqrt{d}/\sqrt{L} \le \sigma_i(X_{train}) \le \sqrt{Ld}$ and $1/L \le \lambda_i(H_{train}) \le L$ for all $i \in [n]$ and $\sqrt{d\sigma/4} \le 1$ $\|\xi_{train}\| \leq \sqrt{d}\sigma$ and there exists $\eta_1, \eta_2, \eta_3 = \Theta(1/t)$ with $\eta_1 < \eta_2 < \eta_3$ such that

$$Q'(\eta_2) := 1/2\mathbb{E}_{SGD} \|w'_{t,\eta_2} - w^*\|^2 \le \frac{1}{2} \|w^*\|^2 - C$$

$$Q'(\eta) := 1/2\mathbb{E}_{SGD} \|w'_{t,\eta} - w^*\|^2 \ge \frac{1}{2} \|w^*\|^2 - \frac{C}{2}, \forall \eta \in [0, \eta_1] \cup [\eta_3, 1/L]$$

where C is a positive constant.

To relate the behavior of $Q'(\eta)$ defined on $\{w'_{\tau,\eta}\}$ to the behavior of $Q(\eta)$ defined on $\{w_{\tau,\eta}\}$. We show when the step size is small enough, the SGD sequence gets truncated with very small probability so that sequence $\{w_{\tau,\eta}\}$ almost always coincides with sequence $\{w'_{\tau,n}\}$. The proof of Lemma 43 is deferred into Section D.2.3.

Lemma 43. Given a task P, assume $\sqrt{d}/\sqrt{L} \le \sigma_i(X_{train}) \le \sqrt{Ld}$ and $1/L \le \lambda_i(H_{train}) \le L$ for all $i \in [n]$ and $\sqrt{d}\sigma/4 \le \|\xi_{train}\| \le \sqrt{d}\sigma$. Given any $\epsilon > 0$, suppose $\eta \le \frac{1}{c_5d^2\log^2(d/\epsilon)}$ for some constant c_5 , we have

$$|Q(\eta) - Q'(\eta)| \le \epsilon.$$

Combining Lemma 42 and Lemma 43, we give the proof of lemma 40.

Proof of Lemma 40. Recall that we define $Q(\eta) := 1/2\mathbb{E}_{\text{SGD}} \|w_{t,\eta} - w^*\|^2$ and $Q'(\eta) = 1/2\mathbb{E}_{\text{SGD}} \|w'_{t,\eta} - w^*\|^2$. Here, $\{w'_{\tau,\eta}\}$ is a SGD sequence running on task P without truncation.

According to Lemma 42, with probability at least $1 - \exp(-\Omega(d))$ over the sampling of task $P, \sqrt{d}/\sqrt{L} \le$ $\sigma_i(X_{\text{train}}) \leq \sqrt{Ld}$ and $1/L \leq \lambda_i(H_{\text{train}}) \leq L$ for all $i \in [n]$ and $\sqrt{d}\sigma/4 \leq \|\xi_{\text{train}}\| \leq \sqrt{d}\sigma$ and there exists $\eta_1, \eta_2, \eta_3 = \Theta(1/t)$ with $\eta_1 < \eta_2 < \eta_3$ such that

$$Q'(\eta_2) \le \frac{1}{2} \|w^*\|^2 - C$$

$$Q'(\eta) \ge \frac{1}{2} \|w^*\|^2 - \frac{C}{2}, \forall \eta \in [0, \eta_1] \cup [\eta_3, 1/L]$$

where C is a positive constant. Call this event \mathcal{E} . Suppose the probability that \mathcal{E} happens is $1-\delta$. We can write $\mathbb{E}_{P \sim \mathcal{T}} Q(\eta)$ as follows,

$$\mathbb{E}_{P \sim \mathcal{T}} Q(\eta) = \mathbb{E}_{P \sim \mathcal{T}} [Q(\eta) | \mathcal{E}] \Pr[\mathcal{E}] + \mathbb{E}_{P \sim \mathcal{T}} [Q(\eta) | \bar{\mathcal{E}}] \Pr[\bar{\mathcal{E}}].$$

According to the algorithm, we know $\|w_{t,\eta}\|$ is always bounded by $4\sqrt{L}\sigma$. Therefore, $Q(\eta) := 1/2 \|w_{t,\eta} - w^*\|^2 \le 13L\sigma^2$. By Lemma 43, we know conditioning on \mathcal{E} , $|Q(\eta) - Q'(\eta)| \le \epsilon$ for any $\eta \le \frac{1}{c_5 d^2 \log^2(d/\epsilon)}$. As long as $t \ge c_2 d^2 \log^2(d/\epsilon)$ for certain constant c_2 , we know $\eta_3 \le \frac{1}{c_5 d^2 \log^2(d/\epsilon)}$

When $\eta = \eta_2$, we have

$$\mathbb{E}_{P \sim \mathcal{T}} Q(\eta_2) \le (Q'(\eta_2) + \epsilon) (1 - \delta) + 13L\sigma^2 \delta$$

$$\le \left(\frac{1}{2} \|w^*\|^2 - C + \epsilon\right) (1 - \delta) + 13L\sigma^2 \delta$$

$$\le \frac{1}{2} \|w^*\|^2 - C + 13L\sigma^2 \delta + \epsilon \le \frac{1}{2} \|w^*\|^2 - \frac{9C}{10},$$

where the last inequality assumes $\delta \leq \frac{C}{260L\sigma^2}$ and $\epsilon \leq \frac{C}{20}$. When $\eta \in [0, \eta_1] \cup [\eta_3, \frac{1}{c_5d^2\log^2(d/\epsilon)}]$, we have

$$\mathbb{E}_{P \sim \mathcal{T}} Q(\eta_2) \ge (Q'(\eta) - \epsilon) (1 - \delta) - 13L\sigma^2 \delta$$

$$\ge \left(\frac{1}{2} \|w^*\|^2 - \frac{C}{2} - \epsilon\right) (1 - \delta) - 13L\sigma^2 \delta$$

$$\ge \frac{1}{2} \|w^*\|^2 - \frac{C}{2} - \frac{\delta}{2} - 13L\sigma^2 \delta - \epsilon \ge \frac{1}{2} \|w^*\|^2 - \frac{6C}{10},$$

where the last inequality holds as long as $\delta \leq \frac{C}{280L\sigma^2}$ and $\epsilon \leq \frac{C}{20}$. According to Lemma 42, we know $\delta \leq \exp(-\Omega(d))$. Therefore, the conditions for δ can be satisfied as long as dis larger than certain constant. The condition on ϵ can be satisfied as long as $\eta \leq \frac{1}{c_5 d^2 \log^2(d)}$ for some constant c_5 . \square

D.2.2 Generalization for $\eta \in [0, \frac{1}{c_5 d^2 \log^2 d}]$

In this section, we prove Lemma 41 by showing that $\hat{F}_{TbV}(\eta)$ is point-wise close to $F_{TbV}(\eta)$ for all $\eta \in [0, \frac{1}{c_5 d^2 \log^2(d/\epsilon)}]$. Recall Lemma 41 as follows.

Lemma 41. For any $1 > \epsilon > 0$, assume σ is a constant and $d \geq c_4 \log(1/\epsilon)$ for some constant c_4 . There exists constant c_5 such that with probability at least $1 - O(1/\epsilon) \exp(-\Omega(\epsilon^2 m))$,

$$|\hat{F}_{TbV}(\eta) - F_{TbV}(\eta)| \le \epsilon,$$

for all
$$\eta \in [0, \frac{1}{c_5 d^2 \log^2(d/\epsilon)}]$$
.

In order to prove Lemma 41, we first show that for a fixed η with high probability $F_{TbV}(\eta)$ is close to $F_{TbV}(\eta)$. Similar as in Lemma 20, we can still show that each $\Delta_{TbV}(\eta, P)$ is O(1)-subexponential. The proof is deferred into Section D.2.3.

Lemma 44. Suppose σ is a constant. Given any $1 > \epsilon > 0$, for any fixed η with probability at least $1 - \exp(-\Omega(\epsilon^2 m))$,

$$\left|\hat{F}_{TbV}(\eta) - F_{TbV}(\eta)\right| \le \epsilon.$$

Next, we show that there exists an ϵ -net for F_{TbV} with size $O(1/\epsilon)$. By ϵ -net, we mean there exists a finite set N_{ϵ} of step sizes such that $|F_{TbV}(\eta) - F_{TbV}(\eta')| \le \epsilon$ for any η and $\eta' \in \arg\min_{\eta \in N_{\epsilon}} |\eta - \eta'|$. The proof is very similar as in Lemma 21. We defer the proof of Lemma 45 into Section D.2.3.

Lemma 45. Suppose σ is a constant. For any $1 > \epsilon > 0$, assume $d \ge c_4 \log(1/\epsilon)$ for some c_4 . There exists constant c_5 and an ϵ -net $N_{\epsilon} \subset [0, \frac{1}{c_5 d^2 \log^2(d/\epsilon)}]$ for F_{TbV} with $|N_{\epsilon}| = O(1/\epsilon)$. That means, for any $\eta \in [0, \frac{1}{c_5 d^2 \log^2(d/\epsilon)}]$,

$$|F_{TbV}(\eta) - F_{TbV}(\eta')| < \epsilon$$
,

for $\eta' \in \arg\min_{\eta \in N_{\epsilon}} |\eta - \eta'|$.

Next, we show that with high probability, there also exists an ϵ -net for \hat{F}_{TbV} with size $O(1/\epsilon)$. The proof is very similar as the proof of Lemma 22. We defer the proof into Section D.2.3.

Lemma 46. Suppose σ is a constant. For any $1 > \epsilon > 0$, assume $d \ge c_4 \log(1/\epsilon)$ for some c_4 . With probability at least $1 - \exp(-\Omega(\epsilon^2 m))$, there exists constant c_5 and an ϵ -net $N'_{\epsilon} \subset [0, \frac{1}{c_5 d^2 \log^2(d/\epsilon)}]$ for \hat{F}_{TbV} with $|N_{\epsilon}| = O(1/\epsilon)$. That means, for any $\eta \in [0, \frac{1}{c_5 d^2 \log^2(d/\epsilon)}],$

$$|\hat{F}_{TbV}(\eta) - \hat{F}_{TbV}(\eta')| \le \epsilon,$$

for $\eta' \in \arg\min_{\eta \in N_{\epsilon}} |\eta - \eta'|$.

Combing Lemma 44, Lemma 45 and Lemma 46, now we give the proof of Lemma 41.

Proof of Lemma 41. The proof is almost the same as the proof of Lemma 15. We omit the details here.

D.2.3 Proofs of Technical Lemmas

In Lemma 47, we show when the step size is small, the expected SGD noise square is well bounded. The proof follows from the analysis in Lemma 37.

Lemma 47. Let $\{w'_{\tau,\eta}\}$ be an SGD sequence running on task P without truncation. Let $n'_{\tau,\eta}$ be the SGD noise at $w'_{\tau,\eta}$. Assume $\sqrt{d}/\sqrt{L} \le \sigma_i(X_{train}) \le \sqrt{L}\sqrt{\sigma}$ for all $i \in [n]$ and $\|\xi_{train}\| \le \sqrt{d}\sigma$. Suppose $\eta \in [0, \frac{1}{2L^3d}]$, we have

$$\mathbb{E}_{SGD} \left\| n_{\tau,\eta}' \right\|^2 \le 4L^3 \sigma^2 d$$

for all $\tau \leq t$.

Proof of Lemma 47. Similar as the analysis in Lemma 37, for $\eta \leq \frac{1}{2L^3d}$, we have

$$\mathbb{E}_{\text{SGD}}\left[\left\|n_{\tau,\eta}'\right\|^{2}\left|w_{\tau-1,\eta}'\right| \leq L^{2}d\left\|w_{\tau-1,\eta}'-w_{\text{train}}\right\|^{2}.$$

and

$$\mathbb{E}_{\mathrm{SGD}} \left\| w_{\tau-1,\eta}' - w_{\mathrm{train}} \right\|^2 \leq (1 - \frac{\eta}{2L})^{\tau-1} \left\| w_{\mathrm{train}} \right\|^2 \leq \left\| w_{\mathrm{train}}^* + (X_{\mathrm{train}})^\dagger \xi_{\mathrm{train}} \right\|^2 \leq 4L\sigma^2.$$

Therefore, we have

$$\mathbb{E}_{\mathrm{SGD}} \left\| n_{\tau,\eta}' \right\|^2 \leq L^2 d \mathbb{E}_{\mathrm{SGD}} \left\| w_{\tau,\eta}' - w_{\mathrm{train}} \right\|^2 \leq 4 L^3 \sigma^2 d.$$

Proof of Lemma 42. We can expand $Q'(\eta)$ as follows,

$$\begin{split} Q'(\eta) := & \frac{1}{2} \mathbb{E}_{\text{SGD}} \left\| w'_{t,\eta} - w^* \right\|^2 \\ = & \frac{1}{2} \mathbb{E}_{\text{SGD}} \left\| B_{t,\eta} w^*_{\text{train}} + B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}} - \eta \sum_{\tau=0}^{t-1} (I - \eta H_{\text{train}})^{t-1-\tau} n'_{\tau,\eta} - w^* \right\|^2 \\ = & \frac{1}{2} \left\| B_{t,\eta} w^*_{\text{train}} - w^* \right\|^2 + \frac{1}{2} \left\| B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}} \right\|^2 + \frac{\eta^2}{2} \mathbb{E}_{\text{SGD}} \left\| \sum_{\tau=0}^{t-1} (I - \eta H_{\text{train}})^{t-1-\tau} n'_{\tau,\eta} \right\|^2 \\ + & \left\langle B_{t,\eta} w^*_{\text{train}} - w^*, B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}} \right\rangle \end{split}$$

Denote

$$G(\eta) := \frac{1}{2} \left\| B_{t,\eta} w_{\text{train}}^* - w^* \right\|^2 + \frac{1}{2} \left\| B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}} \right\|^2 + \frac{\eta^2}{2} \mathbb{E}_{\text{SGD}} \left\| \sum_{\tau=0}^{t-1} (I - \eta H_{\text{train}})^{t-1-\tau} n_{\tau,\eta}' \right\|^2.$$

According to Lemma 7, we know with probability at least $1 - \exp(-\Omega(d))$, $\sqrt{d}/\sqrt{L} \le \sigma_i(X_{\text{train}}) \le \sqrt{L}\sqrt{d}$ and $1/L \le \lambda_i(H_{\text{train}}) \le L$ for all $i \in [n]$. According to Lemma 49, we know with probability at least $1 - \exp(-\Omega(d))$, $\sqrt{d}\sigma/4 \le \|\xi_{\text{train}}\| \le \sqrt{d}\sigma$.

Upper bounding $G(\eta_2)$: We can expand $G(\eta)$ as follows:

$$G(\eta) := \frac{1}{2} \|B_{t,\eta} w_{\text{train}}^* - w^*\|^2 + \frac{1}{2} \|B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}}\|^2 + \frac{\eta^2}{2} \mathbb{E}_{\text{SGD}} \left\| \sum_{\tau=0}^{t-1} (I - \eta H_{\text{train}})^{t-1-\tau} n_{\tau,\eta}' \right\|^2$$

$$= \frac{1}{2} \|w^*\|^2 + \frac{1}{2} \|B_{t,\eta} w_{\text{train}}^*\|^2 + \frac{1}{2} \|B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}}\|^2 + \frac{\eta^2}{2} \mathbb{E}_{\text{SGD}} \left\| \sum_{\tau=0}^{t-1} (I - \eta H_{\text{train}})^{t-1-\tau} n_{\tau,\eta}' \right\|^2$$

$$- \langle B_{t,\eta} w_{\text{train}}^*, w^* \rangle.$$

Same as in Lemma 17, we know $\frac{1}{2} \|B_{t,\eta} w_{\text{train}}^*\|^2 + \frac{1}{2} \|B_{t,\eta} (X_{\text{train}})^\dagger \xi_{\text{train}}\|^2 \leq L^3 \eta^2 t^2 \sigma^2$. For the SGD noise, by Lemma 47 we know $\mathbb{E}_{\text{SGD}} \|n_{\tau,\eta}'\|^2 \leq 4L^3 \sigma^2 d$ for all $\tau \leq t$ as long as $\eta \leq \frac{1}{2L^3 d}$. Therefore,

$$\frac{\eta^2}{2} \mathbb{E}_{\text{SGD}} \left\| \sum_{\tau=0}^{t-1} (I - \eta H_{\text{train}})^{t-1-\tau} n_{\tau,\eta}' \right\|^2 \leq \frac{\eta^2}{2} \sum_{\tau=0}^{t-1} \mathbb{E}_{\text{SGD}} \left\| n_{\tau,\eta}' \right\|^2 \leq 2L^3 \eta^2 \sigma^2 dt \leq 2L^3 \eta^2 \sigma^2 t^2,$$

where the last inequality assumes $t \ge d$. According to Lemma 19, for any fixed $\eta \in [0, L/t]$, with probability at least $1 - \exp(-\Omega(d))$ over X_{train} ,

$$\langle B_{t,\eta} w_{\text{train}}^*, w^* \rangle \ge \frac{\eta t}{16L}.$$

Therefore, for any step size $\eta \leq \frac{1}{2L^3d}$:

$$G(\eta) \le \frac{1}{2} \|w^*\|^2 + 3L^3 \eta^2 \sigma^2 t^2 - \frac{\eta t}{16L} \le \frac{1}{2} \|w^*\|^2 - \frac{\eta t}{32L}$$

where the second inequality holds as long as $\eta \leq \frac{1}{96L^4\sigma^2t}$. Choosing $\eta_2 := \frac{1}{96L^4\sigma^2t}$ that is smaller than $\frac{1}{2L^3d}$ assuming $t \geq d$. Then, we have

$$G(\eta_2) \le \frac{1}{2} \|w^*\|^2 - \frac{5C}{4},$$

where constant $C = \frac{1}{3072L^5\sigma^2}$.

Lower bounding $G(\eta)$ for $\eta \in [0, \eta_1]$: Now, we prove that there exists $\eta_1 = \Theta(1/t)$ with $\eta_1 < \eta_2$ such that for any $\eta \in [0, \eta_1], G(\eta) \geq \frac{1}{2} \|w^*\|^2 - \frac{C}{4}$. Recall that

$$G(\eta) = \frac{1}{2} \|w^*\|^2 + \frac{1}{2} \|B_{t,\eta} w_{\text{train}}^*\|^2 + \frac{1}{2} \|B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}}\|^2 + \frac{\eta^2}{2} \mathbb{E}_{\text{SGD}} \left\| \sum_{\tau=0}^{t-1} (I - \eta H_{\text{train}})^{t-1-\tau} n_{\tau,\eta}' \right\|^2 - \langle B_{t,\eta} w_{\text{train}}^*, w^* \rangle.$$

$$\geq \frac{1}{2} \|w^*\|^2 - \langle B_{t,\eta} w_{\text{train}}^*, w^* \rangle.$$

Same as in Lemma 17, by choosing $\eta_1 = \frac{C}{4Lt}$, we have for any $\eta \in [0, \eta_1]$,

$$G(\eta) \ge \frac{1}{2} \|w^*\|^2 - \frac{C}{4}.$$

Lower bounding $G(\eta)$ for $\eta \in [\eta_3, 1/L]$: Now, we prove that there exists $\eta_3 = \Theta(1/t)$ with $\eta_3 > \eta_2$ such that for all $\eta \in [\eta_3, 1/L]$,

$$G(\eta) \ge \frac{1}{2} \|w^*\|^2 - \frac{C}{4}.$$

Recall that

$$G(\eta) = \frac{1}{2} \|B_{t,\eta} w_{\text{train}}^* - w^*\|^2 + \frac{1}{2} \|B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}}\|^2 + \frac{\eta^2}{2} \mathbb{E}_{\text{SGD}} \left\| \sum_{\tau=0}^{t-1} (I - \eta H_{\text{train}})^{t-1-\tau} n_{\tau,\eta}' \right\|^2$$
$$\geq \frac{1}{2} \|B_{t,\eta} (X_{\text{train}})^{\dagger} \xi_{\text{train}}\|^2.$$

Same as in Lemma 17, by choosing $\eta_3 = \log(2)L/t$, as long as $\sigma \ge 8\sqrt{L}$, we have

$$G(\eta) \ge \frac{1}{2} \left\| w^* \right\|^2$$

for all $\eta \in [\eta_3, 1/L]$. Note $\eta_3 \le 1/L$ as long as $t \ge \log(2)L^2$.

Overall, we have shown that there exist $\eta_1, \eta_2, \eta_3 = \Theta(1/t)$ with $\eta_1 < \eta_2 < \eta_3$ such that $G(\eta_2) \le 1/2 \|w^*\|^2 - 5C/4$ and $G(\eta) \ge 1/2 \|w^*\|^2 - C/4$ for all $\eta \in [0, \eta_1] \cup [\eta_3, 1/L]$. Recall that $Q'(\eta) = G(\eta) + \left\langle B_{t,\eta} w_{\text{train}}^* - w^*, B_{t,\eta}(X_{\text{train}})^\dagger \xi_{\text{train}} \right\rangle$. Choosing $\epsilon = C/4$ in Lemma 18, we know with probability at least $1 - \exp(-\Omega(d)), \left| \left\langle B_{t,\eta} w_{\text{train}}^* - w^*, B_{t,\eta}(X_{\text{train}})^\dagger \xi_{\text{train}} \right\rangle \right| \le C/4$ for all $\eta \in [0, 1/L]$. Therefore, we know $Q'(\eta_2) \le 1/2 \|w^*\|^2 - C$ and $Q'(\eta) \ge 1/2 \|w^*\|^2 - C/2$ for all $\eta \in [0, \eta_1] \cup [\eta_3, 1/L]$.

In order to prove Lemma 43, we first construct a super-martingale to show that as long as task P is well behaved, with high probability in SGD noise, the weight norm along the trajectory never exceeds $4\sqrt{L}\sigma$.

Lemma 48. Assume $\sqrt{d}/\sqrt{L} \le \sigma_i(X_{train}) \le \sqrt{Ld}$ and $1/L \le \lambda_i(H_{train}) \le L$ for all $i \in [n]$ and $\sqrt{d}\sigma/4 \le \|\xi_{train}\| \le \sqrt{d}\sigma$. Given any $1 > \delta > 0$, suppose $\eta \le \frac{1}{c_5d^2\log^2(d/\delta)}$ for some constant c_5 , with probability at least $1 - \delta$ in the SGD noise,

$$\|w'_{\tau,\eta}\| < 4\sqrt{L}\sigma$$

for all $\tau \leq t$.

Proof of Lemma 48. According to the proofs of Lemma 47, as long as $\eta \leq \frac{1}{2L^3d}$, we have

$$\mathbb{E}_{\mathrm{SGD}}\left[\left\|w_{t,\eta}'-w_{\mathrm{train}}\right\|^{2}|w_{t-1,\eta}']\leq\left(1-\frac{\eta}{2L}\right)\left\|w_{t-1,\eta}'-w_{\mathrm{train}}\right\|^{2}.$$

Since log is a concave function, by Jenson's inequality, we know

$$\begin{split} & \mathbb{E}_{\text{SGD}}\left[\log\left\|w_{t,\eta}' - w_{\text{train}}\right\|^{2} | w_{t-1,\eta}' \right] \\ & \leq \log\mathbb{E}_{\text{SGD}}\left[\left\|w_{t,\eta}' - w_{\text{train}}\right\|^{2} | w_{t-1,\eta}' \right] \leq \log\left\|w_{t-1,\eta}' - w_{\text{train}}\right\|^{2} + \log(1 - \frac{\eta}{2L}). \end{split}$$

Defining $G_t = \log \left\| w'_{t,\eta} - w_{\text{train}} \right\|^2 - t \log(1 - \frac{\eta}{2L})$, we know G_t is a super-martingale. Next, we bound the martingale differences.

We can bound $|G_t - \mathbb{E}_{SGD}[G_t|w'_{t-1,\eta}]|$ as follows,

$$|G_{t} - \mathbb{E}_{SGD}[G_{t}|w'_{t-1,\eta}]| \leq \max_{n'_{t-1,\eta}, n''_{t-1,\eta}} \log \left(\frac{\left\| (I - \eta H_{\text{train}})(w'_{t-1,\eta} - w_{\text{train}}) - \eta n'_{t-1,\eta} \right\|^{2}}{\left\| (I - \eta H_{\text{train}})(w'_{t-1,\eta} - w_{\text{train}}) - \eta n''_{t-1,\eta} \right\|^{2}} \right)$$

We can expand $\left\| (I - \eta H_{\text{train}})(w'_{t-1,\eta} - w_{\text{train}}) - \eta n'_{t-1,\eta} \right\|^2$ as follows,

$$\begin{split} & \left\| (I - \eta H_{\text{train}})(w'_{t-1,\eta} - w_{\text{train}}) - \eta n'_{t-1,\eta} \right\|^2 \\ & = \left\| (I - \eta H_{\text{train}})(w'_{t-1,\eta} - w_{\text{train}}) \right\|^2 - 2\eta \left\langle n'_{t-1,\eta}, (I - \eta H_{\text{train}})(w'_{t-1,\eta} - w_{\text{train}}) \right\rangle + \eta^2 \left\| n'_{t-1,\eta} \right\|^2 \end{split}$$

We can bound the norm of the noise as follows,

$$\begin{aligned} \|n'_{t-1,\eta}\| &= \left\| x_{i(t-1)} x_{i(t-1)}^{\top} (w'_{t-1,\eta} - w_{\text{train}}) - H_{\text{train}} (w'_{t-1,\eta} - w_{\text{train}}) \right\| \\ &\leq \left\| x_{i(t-1)} x_{i(t-1)}^{\top} (w'_{t-1,\eta} - w_{\text{train}}) \right\| + \left\| H_{\text{train}} (w'_{t-1,\eta} - w_{\text{train}}) \right\| \\ &\leq (Ld+L) \left\| w'_{t-1,\eta} - w_{\text{train}} \right\| \leq 2Ld \left\| w'_{t-1,\eta} - w_{\text{train}} \right\|, \end{aligned}$$

where the second inequality uses $||x_{i(t-1)}|| \leq \sqrt{Ld}$. Therefore, we have

$$\begin{aligned} & \left| 2\eta \left\langle n'_{t-1,\eta}, (I - \eta H_{\text{train}})(w'_{t-1,\eta} - w_{\text{train}}) \right\rangle \right| \leq 4L\eta d \left\| w'_{t-1,\eta} - w_{\text{train}} \right\|^2, \\ & \eta^2 \left\| n'_{t-1,\eta} \right\|^2 \leq 4L^2 \eta^2 d^2 \left\| w'_{t-1,\eta} - w_{\text{train}} \right\|^2. \end{aligned}$$

This further implies,

$$\begin{split} &|G_{t} - \mathbb{E}_{\text{SGD}}[G_{t}|w'_{t-1,\eta}]| \\ &\leq \log \left(\frac{\left\| (I - \eta H_{\text{train}})(w'_{t-1,\eta} - w_{\text{train}}) \right\|^{2} + \left(4L\eta d + 4L^{2}\eta^{2}d^{2} \right) \left\| w'_{t-1,\eta} - w_{\text{train}} \right\|^{2}}{\left\| (I - \eta H_{\text{train}})(w'_{t-1,\eta} - w_{\text{train}}) \right\|^{2} - 4L\eta d \left\| w'_{t-1,\eta} - w_{\text{train}} \right\|^{2}} \right) \\ &\leq \log \left(1 + \frac{8L\eta d + 4L^{2}\eta^{2}d^{2}}{(1 - 2L\eta - 4L\eta d)} \right) \leq 16L\eta d + 8L^{2}\eta^{2}d^{2}, \end{split}$$

where the second inequality uses $\|(I-\eta H_{\text{train}})(w'_{t-1,\eta}-w_{\text{train}})\|^2 \geq (1-2L\eta)\|w'_{t-1,\eta}-w_{\text{train}}\|^2$. The last inequality assumes $\eta \leq \frac{1}{12Ld}$ and uses numerical inequality $\log(1+x) \leq x$. Assuming $\eta \leq 1/(Ld)$, we further have $|G_t - \mathbb{E}_{\text{SGD}}[G_t|w'_{t-1,\eta}]| \leq L^2\eta d$.

By Azuma's inequality, we know with probability at least $1 - \delta/t$,

$$G_t \le G_0 + L^2 \sqrt{2t} \eta d \log(t/\delta).$$

Plugging in $G_t = \log \|w'_{t,n} - w_{\text{train}}\|^2 - t \log(1 - \frac{\eta}{2L})$ and $G_0 = \log \|w_0 - w_{\text{train}}\|^2 = \log \|w_{\text{train}}\|^2$, we have

$$\begin{split} \log \left\| w_{t,\eta}' - w_{\text{train}} \right\|^2 & \leq \log \left\| w_{\text{train}} \right\|^2 + t \log (1 - \frac{\eta}{2L}) + L^2 \sqrt{2t} \eta d \log(t/\delta) \\ & \leq \log \left\| w_{\text{train}} \right\|^2 - \frac{\eta}{2L} t + L^2 \sqrt{2t} \eta d \log(t/\delta). \end{split}$$

This implies,

$$\begin{aligned} \left\| w'_{t,\eta} - w_{\text{train}} \right\|^2 &\leq \left\| w_{\text{train}} \right\|^2 \exp\left(\eta \left(-\frac{1}{2L} t + L^2 \sqrt{2} \log(t/\delta) d\sqrt{t} \right) \right) \\ &= \left\| w_{\text{train}} \right\|^2 \exp\left(O(d^2 \log^2(d/\delta)) \eta \right) \\ &\leq \left\| w_{\text{train}} \right\|^2 \exp\left(2/3 \right), \end{aligned}$$

where the second inequality assumes $\eta \leq \frac{1}{c_5 d^2 log^2 (d/\delta)}$ for some constant c_5 . Furthermore, since $\|w_{\text{train}}\| \leq (1+\sqrt{L})\sigma$, we have $\|w'_{t,\eta}\| \leq (1+e^{1/3}) \|w_{\text{train}}\| < 4\sqrt{L}\sigma$.

Overall, we know as long as $\eta \leq \frac{1}{c_5 d^2 log^2 (d/\delta)}$, with probability at least $1 - \delta/t$, $\|w'_{t,\eta}\| \leq 4\sqrt{L}\sigma$. Since this analysis also applies to any $\tau \leq t$, we know for any τ , with probability at least $1 - \delta/t$, $\|w'_{\tau,\eta}\| < 4\sqrt{L}\sigma$. Taking a union bound over $\tau \leq t$, we have with probability at least $1 - \delta$, $\|w'_{\tau,\eta}\| < 4\sqrt{L}\sigma$ for all $\tau \leq t$.

Proof of Lemma 43. Let \mathcal{E} be the event that $\|w'_{\tau,\eta}\| < 4\sqrt{L}\sigma$ for all $\tau \leq t$. We first show that $\mathbb{E}_{\text{SGD}} \|w_{t,\eta} - w^*\|^2$ is close to $\mathbb{E}_{\text{SGD}} \|w'_{t,\eta} - w^*\|^2 \mathbb{1} \{\mathcal{E}\}$. It's not hard to verify that

$$\mathbb{E}_{\text{SGD}} \|w_{t,\eta} - w^*\|^2 = \mathbb{E}_{\text{SGD}} \|w'_{t,\eta} - w^*\|^2 \mathbb{1} \{\mathcal{E}\} + \|u - w^*\|^2 \Pr[\bar{\mathcal{E}}],$$

where u is a fixed vector with norm $4\sqrt{L}\sigma$. By Lemma 48, we know $\Pr[\bar{\mathcal{E}}] \leq \epsilon/(25L\sigma^2)$ as long as $\eta \leq \frac{1}{c_5d^2\log^2(d/\epsilon)}$ for some constant c_5 . Therefore, we have

$$\left| \mathbb{E}_{\text{SGD}} \left\| w_{t,\eta} - w^* \right\|^2 - \mathbb{E}_{\text{SGD}} \left\| w_{t,\eta}' - w^* \right\|^2 \mathbb{1} \left\{ \mathcal{E} \right\} \right| \leq \epsilon.$$

Next, we show that $\mathbb{E}_{SGD} \|w'_{t,\eta} - w^*\|^2 \mathbb{1} \{\mathcal{E}\}$ is close to $\mathbb{E}_{SGD} \|w'_{t,\eta} - w^*\|^2$. For any $1 \leq \tau \leq t$, let \mathcal{E}_{τ} be the event that $\|w'_{\tau,\eta}\| \geq 4\sqrt{L}\sigma$ and $\|w'_{\tau',\eta}\| < 4\sqrt{L}\sigma$ for all $\tau' < \tau$. Basically \mathcal{E}_{τ} means the weight norm exceeds the threshold at step τ for the first time. It's easy to see that $\bigcup_{\tau=1}^t \mathcal{E}_{\tau} = \bar{\mathcal{E}}$. Therefore, we have

$$\mathbb{E}_{\text{SGD}} \| w'_{t,\eta} - w^* \|^2 = \mathbb{E}_{\text{SGD}} \| w'_{t,\eta} - w^* \|^2 \mathbb{1} \{ \mathcal{E} \} + \sum_{\tau=1}^t \mathbb{E}_{\text{SGD}} \| w'_{t,\eta} - w^* \|^2 \mathbb{1} \{ \mathcal{E}_{\tau} \}.$$

Conditioning on \mathcal{E}_{τ} , we know $\left\|w'_{\tau-1,\eta}\right\| < 4\sqrt{L}\sigma$. Since we assume $\frac{\sqrt{d}}{\sqrt{L}} \leq \sigma_i(X_{\text{train}}) \leq \sqrt{L}\sqrt{d}$ for all $i \in [n]$ and $\xi_{\text{train}} \leq \sqrt{d}\sigma$, we know $\left\|w_{\text{train}}\right\| \leq 2\sqrt{L}\sigma$. Therefore, we have $\left\|w'_{\tau-1,\eta} - w_{\text{train}}\right\| \leq 6\sqrt{L}\sigma$. Recall the SGD updates,

$$w'_{\tau,\eta} - w_{\text{train}} = (I - \eta H_{\text{train}})(w'_{\tau-1,\eta} - w_{\text{train}}) - \eta n'_{\tau-1,\eta}.$$

For the noise term, we have $\eta \left\| n'_{\tau-1,\eta} \right\| \leq 2\eta L d \left\| w'_{\tau-1,\eta} - w_{\mathrm{train}} \right\|$ that is at most $\left\| w'_{\tau-1,\eta} - w_{\mathrm{train}} \right\|$ assuming $\eta \leq \frac{1}{2Ld}$. Therefore, we have $\left\| w'_{\tau,\eta} - w_{\mathrm{train}} \right\| \leq 2 \left\| w'_{\tau-1,\eta} - w_{\mathrm{train}} \right\| \leq 12\sqrt{L}\sigma$. Note that event \mathcal{E}_{τ} is independent with the SGD noises after step τ . Therefore, according to the previous analysis, we know as long as $\eta \leq \frac{1}{2L^3d}$,

$$\mathbb{E}_{\mathrm{SGD}}\left[\left\|w_{t,\eta}'-w_{\mathrm{train}}\right\|^{2}|\mathcal{E}_{\tau}\right]\leq\left\|w_{\tau,\eta}'-w_{\mathrm{train}}\right\|^{2}\leq2L^{2}\sigma^{2}.$$

Then, we can bound $\mathbb{E}_{\mathrm{SGD}}\left[\left\|w_{t,\eta}'-w^*\right\|^2|\mathcal{E}_{ au}
ight]$ as follows,

$$\mathbb{E}_{\text{SGD}} \left[\left\| w'_{t,\eta} - w^* \right\|^2 | \mathcal{E}_{\tau} \right]$$

$$= \mathbb{E}_{\text{SGD}} \left[\left\| w'_{t,\eta} - w_{\text{train}} + w_{\text{train}} - w^* \right\|^2 | \mathcal{E}_{\tau} \right]$$

$$\leq \mathbb{E}_{\text{SGD}} \left[\left\| w'_{t,\eta} - w_{\text{train}} \right\|^2 | \mathcal{E}_{\tau} \right] + 2 \mathbb{E}_{\text{SGD}} \left[\left\| w'_{t,\eta} - w_{\text{train}} \right\| | \mathcal{E}_{\tau} \right] \left\| w_{\text{train}} - w^* \right\| + \left\| w_{\text{train}} - w^* \right\|^2$$

$$\leq 2L^2 \sigma^2 + 2 \cdot 2L\sigma \cdot 3\sqrt{L}\sigma + 9L\sigma^2 \leq 3L^2 \sigma^2.$$

Therefore, we have

$$\sum_{\tau=1}^{t} \mathbb{E}_{SGD} \|w'_{t,\eta} - w^*\|^2 \mathbb{1} \{\mathcal{E}_{\tau}\} = \sum_{\tau=1}^{t} \mathbb{E}_{SGD} \left[\|w'_{t,\eta} - w^*\|^2 |\mathcal{E}_{\tau} \right] \Pr[\mathcal{E}_{\tau}]$$

$$\leq 3L^2 \sigma^2 \sum_{\tau=1}^{t} \Pr[\mathcal{E}_{\tau}] = 3L^2 \sigma^2 \Pr[\bar{\mathcal{E}}] \leq 3L^2 \sigma^2 \epsilon.$$

This then implies that $\left\|\mathbb{E}_{\mathrm{SGD}}\left\|w_{t,\eta}'-w^*\right\|^2-\mathbb{E}_{\mathrm{SGD}}\left\|w_{t,\eta}'-w^*\right\|^2\mathbb{1}\left\{\mathcal{E}\right\}\right|\leq 3L^2\sigma^2\epsilon.$ Finally, we have

$$\begin{aligned} & \left| \mathbb{E}_{\text{SGD}} \left\| w_{t,\eta} - w^* \right\|^2 - \mathbb{E}_{\text{SGD}} \left\| w'_{t,\eta} - w^* \right\|^2 \right| \\ & \leq \left| \mathbb{E}_{\text{SGD}} \left\| w_{t,\eta} - w^* \right\|^2 - \mathbb{E}_{\text{SGD}} \left\| w'_{t,\eta} - w^* \right\|^2 \mathbb{1} \left\{ \mathcal{E} \right\} \right| + \left| \mathbb{E}_{\text{SGD}} \left\| w'_{t,\eta} - w^* \right\|^2 - \mathbb{E}_{\text{SGD}} \left\| w'_{t,\eta} - w^* \right\|^2 \mathbb{1} \left\{ \mathcal{E} \right\} \right| \\ & \leq \left(3L^2 \sigma^2 + 1 \right) \epsilon \end{aligned}$$

as long as $\eta \leq \frac{1}{c_5 d^2 \log^2(d/\epsilon)}$. Therefore, $|Q(\eta) - Q'(\eta)| \leq \left(3L^2\sigma^2 + 1\right)\epsilon/2$. Choosing $\epsilon' = \frac{2\epsilon}{(3L^2\sigma^2 + 1)}$ finishes the proof.

Proof of Lemma 44. Recall that

$$\hat{F}_{TbV}(\eta) := \frac{1}{m} \sum_{k=1}^{m} \Delta_{TbV}(\eta, P) = \frac{1}{m} \sum_{k=1}^{m} \mathbb{E}_{\text{SGD}} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{\text{valid}}^{(k)} \right\|_{H_{\text{valid}}^{(k)}}^{2}.$$

Similar as in Lemma 15, we can show $\frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{\text{valid}}^{(k)} \right\|_{H_{\text{valid}}^{(k)}}^2$ is O(1)-subexponential, which implies

 $\mathbb{E}_{\text{SGD}} \frac{1}{2} \left\| w_{t,\eta}^{(k)} - w_{\text{valid}}^{(k)} \right\|_{H_{\text{valid}}^{(k)}}^2 \text{ is } O(1) \text{-subexponential. Therefore, } \hat{F}_{TbV}(\eta) \text{ is the average of } m \text{ i.i.d. } O(1) \text{-subexponential random variables.}$ By standard concentration inequality, we know for any $1 > \epsilon > 0$, with probability at least $1 - \exp(-\Omega(\epsilon^2 m))$,

$$\left| \hat{F}_{TbV}(\eta) - F_{TbV}(\eta) \right| \le \epsilon.$$

Proof of Lemma 45. Recall that

$$F_{TbV}(\eta) = \mathbb{E}_{P \sim \mathcal{T}} \mathbb{E}_{SGD} \frac{1}{2} \left\| w_{t,\eta} - w^* \right\|^2 + \sigma^2 / 2$$

We only need to construct an ϵ -net for $\mathbb{E}_{P \sim \mathcal{T}} \mathbb{E}_{SGD} \frac{1}{2} \|w_{t,\eta} - w^*\|^2$. Let \mathcal{E} be the event that $\sqrt{d}/\sqrt{L} \leq \sigma_i(X_{train}) \leq \sigma_i(X_{train})$ \sqrt{Ld} and $1/L \le \lambda_i(H_{\text{train}}) \le L$ for all $i \in [n]$ and $\sqrt{d\sigma}/4 \le \|\xi_{\text{train}}\| \le \sqrt{d\sigma}$ We have

$$\mathbb{E}_{P \sim \mathcal{T}} \mathbb{E}_{\text{SGD}} \frac{1}{2} \| w_{t,\eta} - w^* \|^2$$

$$= \mathbb{E}_{P \sim \mathcal{T}} \left[\frac{1}{2} \mathbb{E}_{\text{SGD}} \| w_{t,\eta} - w^* \|^2 |\mathcal{E}| \right] \Pr[\mathcal{E}] + \mathbb{E}_{P \sim \mathcal{T}} \left[\frac{1}{2} \mathbb{E}_{\text{SGD}} \| w_{t,\eta} - w^* \|^2 |\bar{\mathcal{E}}| \right] \Pr[\bar{\mathcal{E}}]$$

According to Lemma 43, we know conditioning on \mathcal{E} ,

$$\left| \frac{1}{2} \mathbb{E}_{SGD} \| w_{t,\eta} - w^* \|^2 - \frac{1}{2} \mathbb{E}_{SGD} \| w'_{t,\eta} - w^* \|^2 \right| \le \epsilon,$$

as long as $\eta \leq \frac{1}{c_5 d^2 \log^2(d/\epsilon)}$. Note $\{w'_{\tau,\eta}\}$ is the SGD sequence without truncation. For the second term, we have

$$\mathbb{E}_{P \sim \mathcal{T}} \left[\frac{1}{2} \mathbb{E}_{\text{SGD}} \| w_{t,\eta} - w^* \|^2 | \bar{\mathcal{E}} \right] \Pr[\bar{\mathcal{E}}] \le 13L\sigma^2 \Pr[\bar{\mathcal{E}}] \le \epsilon,$$

where the last inequality assumes $\Pr[\bar{\mathcal{E}}] \leq \frac{\epsilon}{13L\sigma^2}$. According to Lemma 7 and Lemma 49, we know $\Pr[\bar{\mathcal{E}}] \leq \exp(-\Omega(d))$. Therefore, given any $\epsilon > 0$, we have $\Pr[\bar{\mathcal{E}}] \leq \frac{\epsilon}{13L\sigma^2}$ as long as $d \geq c_4 \log(1/\epsilon)$ for some constant

Then, we only need to construct an ϵ -net for $\mathbb{E}_{P \sim \mathcal{T}}\left[\frac{1}{2}\mathbb{E}_{\text{SGD}} \left\|w'_{t,\eta} - w^*\right\|^2 |\mathcal{E}|\right] \Pr[\mathcal{E}]$. By the analysis in Lemma 37, it's not hard to prove

$$\left|\frac{\partial}{\partial \eta} \mathbb{E}_{P \sim \mathcal{T}} \left[\frac{1}{2} \mathbb{E}_{\text{SGD}} \left\| w_{t,\eta}' - w^* \right\|^2 |\mathcal{E}\right] \Pr[\mathcal{E}] \right| = O(1) t (1 - \frac{\eta}{2L})^{t-1},$$

for all $\eta \in [0, \frac{1}{c_5 d^2 \log^2(d/\epsilon)}]$. Similar as in Lemma 18, for any $\epsilon > 0$, we know there exists an ϵ -net N_{ϵ} with size $O(1/\epsilon)$ such that for any $\eta \in [0, \frac{1}{c_5 d^2 \log^2(d/\epsilon)}]$

$$\left| \mathbb{E}_{P \sim \mathcal{T}} \left[\frac{1}{2} \mathbb{E}_{\text{SGD}} \left\| w'_{t,\eta} - w^* \right\|^2 | \mathcal{E} \right] \Pr[\mathcal{E}] - \mathbb{E}_{P \sim \mathcal{T}} \left[\frac{1}{2} \mathbb{E}_{\text{SGD}} \left\| w'_{t,\eta'} - w^* \right\|^2 | \mathcal{E} \right] \Pr[\mathcal{E}] \right| \le \epsilon$$

for $\eta' \in \arg\min_{\eta \in N_{\epsilon}} |\eta - \eta'|$.

Combing with the bounds on $\left|\frac{1}{2}\mathbb{E}_{\text{SGD}}\left\|w_{t,\eta}-w^*\right\|^2\mathbb{1}\left\{\mathcal{E}\right\}-\frac{1}{2}\mathbb{E}_{\text{SGD}}\left\|w_{t,\eta}'-w^*\right\|^2\mathbb{1}\left\{\mathcal{E}\right\}\right|$ and $\mathbb{E}_{P \sim \mathcal{T}} \left[\frac{1}{2} \mathbb{E}_{\text{SGD}} \left\| w_{t,\eta} - w^* \right\|^2 |\bar{\mathcal{E}}| \right] \Pr[\bar{\mathcal{E}}], \text{ we have for any } \eta \in [0, \frac{1}{c_5 d^2 \log^2(d/\epsilon)}],$

$$F_{TbV}(\eta) - F_{TbV}(\eta') \le 4\epsilon$$

for $\eta' \in \arg\min_{\eta \in N_{\epsilon}} |\eta - \eta'|$. We finish the proof by replacing 4ϵ by ϵ' .

Proof of Lemma 46. The proof is very similar as the proof of Lemma 22. The only difference is that we need to first relate the SGD sequence with truncation to the SGD sequence without truncation and then bound the Lipschitzness on the SGD sequence without truncation (as we did in Lemma 45). We omit the details here.

E Tools

E.1 Norm of random vectors

We use the following lemma to bound the noise in least squares model.

Lemma 49 (Theorem 3.1.1 in Vershynin (2018)). Let $X = (X_1, X_2, \dots, X_n) \in \mathbb{R}^n$ be a random vector with each entry independently sampled from $\mathcal{N}(0,1)$. Then

$$\Pr[||x|| - \sqrt{n}| \ge t] \le 2\exp(-t^2/C^2),$$

where C is an absolute constant.

E.2 Singular values of Gaussian matrices

Given a random Gaussian matrix, in expectation its smallest and largest singular value can be bounded as follows.

Lemma 50 (Theorem 5.32 in Vershynin (2010)). Let A be an $N \times n$ matrix whose entries are independent standard normal random variables. Then

$$\sqrt{N} - \sqrt{n} \le \mathbb{E}s_{\min}(A) \le \mathbb{E}s_{\max}(A) \le \sqrt{N} + \sqrt{n}$$

Lemma 51 shows a lipchitz function over i.i.d. Gaussian variables concentrate well on its mean. We use this lemma to argue for any fixed step size, the empirical meta objective concentrates on the population meta objective.

Lemma 51 (Proposition 5.34 in Vershynin (2010)). Let f be a real valued Lipschitz function on \mathbb{R}^n with Lipschitz constant K. Let X be the standard normal random vector in \mathbb{R}^n . Then for every t > 0 one has

$$\Pr[f(X) - \mathbb{E}f(X) \ge t] \le \exp(-\frac{t^2}{2K^2}).$$

The following lemma shows a tall random Gaussian matrix is well-conditioned with high probability. The proof follows from Lemma 50 and Lemma 51. We use Lemma 52 to show the covariance matrix is well conditioned in the least squares model.

Lemma 52 (Corollary 5.35 in Vershynin (2010)). Let A be an $N \times n$ matrix whose entries are independent standard normal random variables. Then for every $t \ge 0$ with probability at least $1 - 2 \exp(-t^2/2)$ one has

$$\sqrt{N} - \sqrt{n} - t \le s_{\min}(A) \le s_{\max}(A) \le \sqrt{N} + \sqrt{n} + t$$

E.3 Johnson-Lindenstrauss lemma

We also use Johnson-Lindenstrauss Lemma in some of the lemmas. Johnson-Lindenstrauss Lemma tells us the projection of a fixed vector on a random subspace concentrates well as long as the subspace is reasonably large.

Lemma 53 (Johnson & Lindenstrauss (1984)). Let P be a projection in \mathbb{R}^d onto a random n-dimensional subspace uniformly distributed in $G_{d,n}$. Let $z \in \mathbb{R}^d$ be a fixed point and $\epsilon > 0$, then with probability at least $1 - 2\exp(-c\epsilon^2 n)$,

$$(1 - \epsilon)\sqrt{\frac{n}{d}} \|z\| \le \|Pz\| \le (1 + \epsilon)\sqrt{\frac{n}{d}} \|z\|.$$

F Experiment details

We describe the detailed settings of our experiments in Section F.1 and give more experimental results in Section F.2.

F.1 Experiment settings

Optimizing step size for quadratic objective In this experiment, we meta-train a learning rate for gradient descent on a fixed quadratic objective. Our goal is to show that the autograd module in popular deep learning softwares, such as Tensorflow, can have numerical issues when using the log-transformed meta objective. Therefore, we first implement the meta-training process with Tensorflow to see the results. We then re-implement the meta-training using the hand-derived meta-gradient (see Eqn 5) to compare the result.

A general setting for both implementations is as follows. The inner problem is fixed as a 20-dimensional quadratic objective as described in Section 3, and we use the log-transformed meta objective for training. The positive semi-definite matrix H is generated by first sampling a 20×20 matrix X with all entries drawn from the standard normal distribution and then setting $H = X^T X$. The initial point w_0 is drawn from standard normal as well. Note that we use the same quadratic problem (i.e., the same H and w_0) throughout the meta-training. We do 1000 meta-training iterations, and collect results for different settings of the initial learning rate η_0 and the unroll length t.

We first implement the meta-training code with Tensorflow. Our code is adapted from Wichrowska et al. (2017) ⁷. We use their global learning rate optimizer and specify the problem set to have only one quadratic objective instance. We implemented the quadratic objective class ourselves (the "MyQuadratic" class). We also turned off multiple advanced features in the original code, such as attention and second derivatives, by assigning their flags as false. This ensures that the experiments have exactly the same settings as we described. The meta-training learning rate is set to be 0.001, which is of similar scale as our next experiment. We also try RMSProp as the meta optimizer, which alleviates some of the numerical issues as it renormalizes the gradient, but our experiments show that even RMSProp is still much worse than our implementation.

We then implement the meta-training by hand to show the accurate training results that avoid numerical issues. Specifically, we compute the meta-gradient using Eq (5), where we also scaled the numerator and denominator as described in Claim 5 to avoid numerical issues. We use the algorithm suggested in Theorem 4, except we choose the meta-step size to be $1/(100\sqrt{k})$ as the constants in Theorem 4 were not optimized.

Train-by-train vs. train-by-validation, synthetic data In this experiment, we find the optimal learning rate η^* for least-squares problems trained in train-by-train and train-by-validation settings and then see how the learning rate works on new tasks.

Specifically, we generate 300 different 1000-dimensional least-squares tasks with noise as defined in Section 4 for inner-training and then use the meta-objectives defined in Eq (3) and (4) to find the optimal learning rate. The inner-training number of steps t is set as 40. We try different sample sizes and different noise levels for comparison. Subsequently, in order to test how the two η^* (for train-by-train and train-by-validation respectively) work, we use them on 10 test tasks (the same setting as the inner-training problem) and compute training and testing root mean squared error (RMSE).

Note that since we only need the final optimal η^* found under the two meta-objective settings (regardless of how we find it), we do not need to actually do the meta-training. Instead, we do a grid search on the interval $[10^{-6},1]$, which is divided log-linearly to 25 candidate points. For both the train-by-train and train-by-validation settings, we average the meta-objectives over the 300 inner problems and see which η minimizes this averaged meta-objective. The results are shown in Appendix F.2.

Train-by-train vs. train-by-validation, MLP optimizer on MNIST To observe the trade-off between train-by-train and train-by-validation in a broader and more realistic case, we also do experiments to meta-train an MLP optimizer as in Metz et al. (2019) to solve the MNIST classification problem. We use part of their code ⁸ to integrate with our code in the first experiment, and we use exactly the same default setting as theirs, which is summarized below.

The MLP optimizer is a trainable optimizer that works on each parameter separately. When doing inner-training, for each parameter, we first compute some statistics of that parameter (explained below), which are combined into a feature vector, and then feed that feature vector to a Muti-Layer Perceptron (MLP) with ReLU activations, which

 $^{^7} Their open source code is available at \verb|https://github.com/tensorflow/models/tree/master/research/learned_optimizer| and the control of the control of$

⁸ Their code is available at https://github.com/google-research/google-research/tree/master/task_specific_learned_opt

Table 1: Whether the implementation converges for different t (fixed $\eta_0 = 0.1$)

t	10	20	40	80
Ours	√	√	√	√
Tensorflow GD	×	×	×	×
Tensorflow RMSProp	✓	✓	×	×

outputs two scalars, the update direction and magnitude. The update is computed as the direction times the exponential of the magnitude. The feature vector is 31-dimensional, which includes gradient, parameter value, first-order moving averages (5-dim), second-order moving averages (5-dim), normalized gradient (5-dim), reciprocal of square root second-order moving averages (5-dim) and a step embedding (9-dim). All moving averages are computed using 5 different decay rates (0.5, 0.9, 0.99, 0.999, 0.9999), and the step embedding is tanh distortion of the current number of steps divided by 9 different scales (3, 10, 30, 100, 300, 1000, 3000, 10000, 300000). After expanding the 31-dimensional feature vector for each parameter, we also normalize the set of vectors dimension-wise across all the parameters to have mean 0 and standard deviation 1 (except for the step embedding part). More details can be found in their original paper and original implementation.

The inner-training problem is defined as using a two-layer fully connected network (i.e., another "MLP") with ReLU activations to solve the classic MNIST 10-class classification problem. We use a very small network for computational efficiency, and the two layers have 100 and 20 neurons. We fix the cross-entropy loss as the inner-objective and use mini-batches of 32 samples when inner-training.

When we meta-train the MLP optimizer, we use exactly the same process as fixed in experiments by Wichrowska et al. (2017). We use 100 different inner problems by shuffling the 10 classes and also sampling a new subset of data if we do not use the complete MNIST data set. We run each of the problems with three inner-training trajectories starting with different initialization. Each inner-training trajectory is divided into a certain number of unrolled segments, where we compute the meta-objective and update the meta-optimizer after each segment. The number of unrolled segments in each trajectory is sampled from 10 + Exp(30), and the length of each segment is sampled from 50 + Exp(100), where $\text{Exp}(\cdot)$ denotes the exponential distribution. Note that the meta-objective computed after each segment is defined as the average of all the inner-objectives (evaluated on the train/validation set for train-by-train/train-by-val) within that segment for a better convergence. We also do not need to log-transform the inner-objective this time because the cross entropy loss has a log operator itself. The meta-training, i.e. training the parameters of the MLP in the MLP optimizer, is completed using a classic RMSProp optimizer with meta learning rate 0.01.

For each settings of sample sizes and noise levels, we train two MLP optimizer: one for train-by-train, and one for train-by-validation. When we test the learned MLP optimizer, we use similar settings as the inner-training problem, and we run the trajectories longer for full convergence (4000 steps for small data sets; 40000 steps for the complete data set). We run 5 independent tests and collect training accuracy and test accuracy for evaluation. The plots show the mean of the 5 tests. We have also tuned a SGD optimizer (with the same mini-batch size) by doing a grid-search of the learning rate as baseline.

F.2 Additional results

Optimizing step size for quadratic objective We try experiments for the same settings of the initial η_0 and inner training length t for all of three implementations (our hand-derived GD version, Tensorflow GD version and the Tensorflow RMSProp version). We do 1000 meta-training steps for all the experiments.

For both Tensorflow versions, we always see infinite meta-objectives if η_0 is large or t is large, whose meta-gradient is usually treated as zero, so the training get stuck and never converge. Even for the case that both η_0 and t is small, it still has very large meta-objectives (the scale of a few hundreds), and that is why we also try RMSProp, which should be more robust against the gradient scales. Our hand-derived version, however, does not have the numerical issues and can always converge to the optimal η^* . The detailed convergence is summarized in Tab 1 and Tab 2. Note that the optimal η^* is usually around 0.03 under our settings.

Table 2: Whether the implementation converges for different η_0 (fixed t=40)

η_0	0.001	0.01	0.1	1
Ours	√	✓	√	√
Tensorflow GD	×	×	×	×
Tensorflow RMSProp	✓	\checkmark	×	×

Train-by-train vs. train-by-validation, MLP optimizer on MNIST We also do additional experiments on training an MLP optimizer on the MNIST classification problem. We first try using all samples under the 20% noised setting. The results are shown in Fig 6. The train-by-train setting can perform well if we have a large data set, but since there is also noise in the data, the train-by-train model still overfits and is slightly worse than the train-by-validation model.




Figure 6: Training and testing accuracy for different models (all samples, 20% noise)

We then try an intermediate sample size 12000. The results are shown in Fig 7 (no noise) and Fig 8 (20% noise). We can see that as the theory predicts, as the amount of data increases (from 1000 samples to 12000 samples and then to 60000 samples) the gap between train-by-train and train-by-validation decreases. Also, when we condition on the same number of samples, having additional label noise always makes train-by-train model much worse compared to train-by-validation.




Figure 7: Training and testing accuracy for different models (12000 samples, no noise)




Figure 8: Training and testing accuracy for different models (12000 samples, 20% noise)