

HAUSDORFF DIMENSION OF JULIA SETS OF UNICRITICAL CORRESPONDENCES

CARLOS SIQUEIRA

MARCH 6, 2024

ABSTRACT. We show that if $\beta > 1$ is a rational number and the Julia set J of the holomorphic correspondence $z^\beta + c$ is a locally eventually onto hyperbolic repeller, then the Hausdorff dimension of J is bounded from above by the zero of the associated pressure function. As a consequence, we conclude that the Julia set of the correspondence has zero Lebesgue measure for parameters close to zero, whenever $q^2 < p$ and $\beta = p/q$ in lowest terms.

MSC-class 2020: 37F05 (Primary) 37D35 (Secondary).

1. INTRODUCTION

This paper is concerned with the oneparameter family of holomorphic correspondences acting on the complex plane

$$(1.1) \quad (w - c)^q = z^p,$$

where $p > q \geq 2$ are integers. A holomorphic correspondence, as defined in this paper, is a relation $z \mapsto w$ determined by a polynomial equation $P(z, w) = 0$ in two complex variables. The family (1.1) has bidegree (p, q) ; this means that (1.1) defines a multifunction that maps every $z \neq 0$ to q different values of w and its inverse maps every $w \neq c$ to p values of z . We shall establish an upper bound for the Hausdorff dimension of certain Julia sets in this family (Theorem A), thereby showing that the Julia set of (1.1) has zero Lebesgue measure for parameters c close to zero when $q^2 < p$ (Theorem B).

The family (1.1) can be regarded as a generalisation of the quadratic family $z^2 + c$, for if $\beta = p/q$ in lowest terms, then (1.1) is the multifunction $\mathbf{f}_c(z) = z^\beta + c$, in the sense that $\mathbf{f}_c(z)$ is the set of all w satisfying (1.1). There is another way of generalising the quadratic family in the context of holomorphic correspondences using matings. Indeed, the space of holomorphic correspondences includes all possible Kleinian groups and rational maps, and some correspondences concentrate in a single expression $P(z, w) = 0$ the dynamics of a polynomial map and a Kleinian group. This is the case of the oneparameter family of holomorphic correspondences \mathcal{F}_a introduced by Bullett and Penrose nearly thirty years ago in [6], when they discovered that if $4 \leq a \leq 7$, then the correspondence \mathcal{F}_a is a mating between some quadratic map $z^2 + c$ and the modular group $\mathrm{PSL}(2, \mathbb{Z})$. In the same paper [6], the authors conjectured that the connectedness locus of the family \mathcal{F}_a is homeomorphic to the Mandelbrot set. Recently, Bullett and Lomonaco [2] have shown that \mathcal{F}_a is a mating between some parabolic map and the modular group, for every parameter a in the connectedness locus of the family \mathcal{F}_a . Moreover, they have developed a strategy to prove that the connectedness locus of the family of matings is indeed homeomorphic to the Mandelbrot set, see [3, page 4] and [4]. Holomorphic correspondences also appear in many other contexts; indeed, Lee, Lyubich, Makarov and Mukherjee [9] have investigated the dynamics oneparameter families of Schwarz reflections which give rise to anti-holomorphic correspondences that are, in a suitable sense, matings of anti-rational maps with the abstract modular group $\mathbb{Z}_2 * \mathbb{Z}_3$.

1.1. Hausdorff dimension. The Julia set $J(\mathbf{f}_c)$ of $\mathbf{f}_c(z) = z^\beta + c$, when $\beta > 1$, is defined as the closure of all repelling cycles (see section 2); it is always the projection of a solenoid in \mathbb{C}^2 when the parameter c is close to zero, as described by Siqueira and Smalnia in [16]. Simon [13] has derived an explicit formula for the Hausdorff dimension of the Smale-Williams solenoid which relies on the zero of the pressure function, see [13, page 1224]. Before Simon, the pioneer work of Bowen [1] on quasi-Fuchsian groups was the first to establish the formula $P(t\phi) = 0$, relating the Hausdorff dimension to the unique zero of the pressure function. Similarly, in the early eighties Ruelle [12] proved that if the Julia set $J(f)$ of a rational function f is hyperbolic, then the Hausdorff dimension of $J(f)$ depends real analytically on f . The strategy used by Ruelle consists of:

(I) showing that the Hausdorff dimension is given by the *Bowen's formula*: $P(t\phi) = 0$, where $t = \dim_H J$ and $\phi(z) = -\log |f'(z)|$ is the geometric potential; and

(II) proving that $f \mapsto t(f)$ is real analytic, where $t = t(f)$ is implicitly given by $P(t\phi) = 0$.

In the context of holomorphic correspondences, we have the following result.

Theorem A (Theorem 4.1). *Suppose $J(\mathbf{f}_c)$ is a locally eventually onto hyperbolic repeller and let t_c be the unique zero of the pressure function. Then $\dim_H J(\mathbf{f}_c) \leq t_c$.*

The Bowen parameter t_c in Theorem A comes from an expanding and topologically mixing map $f_c : J(f_c) \rightarrow J(f_c)$ acting on a ‘Julia set’ $J(f_c) \subset \mathbb{C}^2$ – see Theorem 3.3. The shape of $J(f_c)$ is similar to that of the Smale-Williams solenoid for parameters close to zero, and the projection of $J(f_c)$ is always the Julia set $J(\mathbf{f}_c)$ in the plane [16]. The sets $J(f_c)$ are related by a holomorphic motion in \mathbb{C}^2 (section 2.2). Since $J(f_c) \subset \mathbb{C}^2$ moves holomorphically, we believe that the Hausdorff dimension of $J(f_c)$ depends real analytically, or at least continuously on c . However, this problem is still unsolved. (It should be noticed that even though t_c comes from the dynamics of f_c on $J(f_c) \subset \mathbb{C}^2$, in this paper the parameter t_c is used to estimate the Hausdorff dimension of the Julia set $J(\mathbf{f}_c)$ in the plane).

The estimate provided by Theorem A can be used to derive the following result.

Theorem B (Corollary 4.1.3). *If c is sufficiently close to zero and $q^2 < p$, then the solenoidal Julia set of $z^{p/q} + c$ has zero Lebesgue measure.*

See [16, page 3106] for a typical figure of the solenoidal Julia set of Theorem B.

1.2. Hyperbolic components. The *connectedness locus* of the family $z^\beta + c$, denoted by \mathcal{M}_β (for $\beta > 1$ and $\beta \in \mathbb{Q}$), is the set of all parameters c for which the Julia set of $z^\beta + c$ is connected. We define $\mathcal{M}_{\beta,0}$ as the set of c for which zero has at least one bounded forward orbit under $z^\beta + c$. Siqueira has shown [15, Theorem 2.3] that \mathcal{M}_β contains $\mathcal{M}_{\beta,0}$. Therefore, the Julia set of $z^\beta + c$ is connected whenever the critical point has at least one bounded forward orbit under \mathbf{f}_c .

For the quadratic family we have $\beta = 2$ and the definitions of \mathcal{M}_2 and $\mathcal{M}_{2,0}$ coincide with the Mandelbrot set. For some non-integer values of β , the parameter space $\mathcal{M}_\beta - \mathcal{M}_{\beta,0}$ is known to be nonempty and generates an intriguing class of Julia sets named Carpets: they are hyperbolic, connected, and have infinitely many holes. In spite of being hyperbolic, Carpets seem to have positive area. See [5, section 3] for more details and figures.

A special version of the Fatou conjecture for polynomial maps states that the interior of \mathcal{M}_d consists of hyperbolic parameters. One implication of this conjecture is: *the Julia set of every polynomial $z^d + c$ has zero Lebesgue measure if c is in the interior of \mathcal{M}_d .* The Hausdorff dimension $d(c)$ of the Julia set of the polynomial $z^d + c$ is real analytic on every hyperbolic component; in particular, it is real analytic on $\mathbb{C} - \mathcal{M}_d$. In the boundary of \mathcal{M}_d , the Hausdorff dimension $d(c)$ is not even continuous at semihyperbolic parameters.

However, Rivera-Letelier [11] has established some sort of continuity of $d(c)$ at semihyperbolic parameters c_0 in the boundary of the multibrot set \mathcal{M}_d , proving that $d(c_n) \rightarrow d(c_0)$, whenever c_n converges to c_0 in an appropriate way (see [11] for more details).

Little is known about the connectedness locus \mathcal{M}_β of the family $z^\beta + c$. One possible way to start the investigation has been presented in [15]: study the dynamics of $z^\beta + c$ when c is close to a centre. Recall from Douady and Hubbard [7, 8] that the Mandelbrot set has infinitely many hyperbolic components U , each of which encoded by a centre $c \in U$; the centre is the only parameter in U for which the orbit of the critical point of $z^2 + c$ is a cycle. For the correspondence $z^\beta + c$, the parameter a is a *simple centre* if only one forward orbit of zero $(0, z_1, z_2, \dots)$ is periodic, and any other orbit $(0, w_1, w_2, \dots)$ diverges to infinity (see Definition 2.5; it is not necessary to compute all orbits to test if a is a simple centre. Indeed, the basin of attraction of ∞ contains a forward invariant disk $|z| > R$, and therefore we have to check only finitely many iterates).

There exists an open set H_β containing the complement of $\mathcal{M}_{\beta,0}$ and every simple centre such that, for any c in H_β , $z^\beta + c$ is hyperbolic and its Julia set is stable by means of branched holomorphic motions; moreover, $J(\mathbf{f}_c)$ is a locally eventually onto (LEO) hyperbolic repeller, whenever $c \in H_\beta$. (See Definition 4.2, Corollary 5.7.1 and Theorems 5.8 and 5.10 of [15]).

Combining these facts with Theorem A we have the following result.

Theorem C (Corollaries 4.1.1 and 4.1.4). *If c is sufficiently close to a simple centre, or if c belongs to the complement of $\mathcal{M}_{\beta,0}$, then*

$$\dim_H J(\mathbf{f}_c) \leq t_c.$$

2. HYPERBOLIC SETS

Consider the holomorphic correspondence (1.1), i.e., the relation $z \mapsto w$ determined by the polynomial equation $(w - c)^q = z^p$. This correspondence shall be denoted by \mathbf{f}_c . The integers $p > q \geq 2$ are fixed. Therefore, $(\mathbf{f}_c)_{c \in \mathbb{C}}$ is a oneparameter family of holomorphic correspondences.

2.1. Preliminary definitions. If z and w are related by (1.1), then we say that w is an *image* of z and write $z \mapsto w$. Every sequence $(z_i)_0^\infty$ of the plane where z_{i+1} is an image of z_i is an *orbit of the correspondence*. An orbit is a *cycle of period n* if $z_n = z_0$, and $z_i \neq z_0$ if $0 < i < n$.

The set of images of a point z under the correspondence (1.1) is denoted by $\mathbf{f}_c(z)$. Similarly, $\mathbf{f}_c(A)$ is the union of all $\mathbf{f}_c(z)$ when z belongs A . The inverse image set $\mathbf{f}_c^{-1}(A)$ is defined by the set of all z which has at least one image in A .

If z_1 is an image of z_0 under the correspondence (1.1) and $z_0 \neq 0$, then there exists a unique univalent function $w = g(z)$ defined on a neighbourhood U of z_0 , implicitly defined by the equation (1.1) and $g(z_0) = z_1$. If $(z_i)_0^n$ is a finite orbit not containing zero, we denote

$$(2.1) \quad g_{z_0, z_1}(z) = g(z) \text{ and } g_{z_0, z_1, \dots, z_n}(z) = (g_{z_{n-1}, z_n} \circ \dots \circ g_{z_0, z_1})(z).$$

The domain of g_{z_0, z_1, \dots, z_n} is an unspecified neighbourhood of z_0 . A cycle $(z_i)_0^n$ with period n is *repelling* if the absolute value of the derivative of g_{z_0, z_1, \dots, z_n} at z_0 is strictly greater than 1. Therefore, no point of a repelling cycle is allowed to be zero.

Definition 2.1 (Julia set). *The closure of the union of all repelling cycles is the Julia set $J(\mathbf{f}_c)$ of the correspondence \mathbf{f}_c .*

Definition 2.2 (Hyperbolic repeller). *A compact set $\Lambda \subset \mathbb{C}^*$ is a hyperbolic repeller if $\mathbf{f}_c^{-1}(\Lambda) = \Lambda$ and there exist $C > 0$ and $\lambda > 1$ such that*

$$(2.2) \quad |g'_{z_0, \dots, z_n}(z_0)| \geq C\lambda^n,$$

for every finite orbit $(z_i)_0^n$ contained in Λ .

Since every hyperbolic repeller Λ is backward invariant, its complement is forward invariant. We define $B_r(a) = \{z \in \mathbb{C} : |z - a| < r\}$.

Lemma 2.1 (Koebe's distortion). *Let \mathcal{F} denote the family of all univalent functions $\varphi : B_r(a) \rightarrow \mathbb{C}$, where $r > 0$ and $a \in \mathbb{C}$ are allowed to vary arbitrarily. There exists a constant $K > 0$ such that, for any φ in \mathcal{F} , we have*

$$(2.3) \quad K^{-1}|\varphi'(a)| \cdot |z - w| \leq |\varphi(z) - \varphi(w)| \leq K|\varphi'(a)| \cdot |z - w|,$$

whenever z and w belong to $B_{r/4}(a)$, where r is the radius of the domain $B_r(a)$ of φ . (Note that K depends neither on r nor a).

Definition 2.3 (Expansive constant for correspondences). Let $\omega \in \mathbb{N}_0 \cup \{\infty\}$. A pair of orbits $(z_i)_0^\omega$ and $(w_i)_0^\omega$ is said to be ϵ -close if $|z_i - w_i| < \epsilon$ for $0 \leq i \leq \omega$. We say that $\epsilon > 0$ is an *expansive constant* for a hyperbolic repeller Λ if every pair of ϵ -close orbits $(z_i)_0^\omega$ and $(w_i)_0^\omega$ in Λ must actually coincide.

Lemma 2.2. *Every hyperbolic repeller Λ of \mathbf{f}_c has an expansive constant $\epsilon > 0$ satisfying the following property: if $(z_i)_0^n$ and $(w_i)_0^n$ is a pair of finite ϵ -close orbits contained in Λ , then both maps $(g_{z_0 \dots z_n})^{-1}$ and $(g_{w_0 \dots w_n})^{-1}$ are well defined and coincide on $B_\epsilon(z_n)$. In particular, $(g_{z_0 \dots z_n})^{-1}$ maps w_n to w_0 .*

Proof. Let $X_{n,c}$ denote the space of all finite orbits $(z_i)_0^n$ contained in Λ . We know that $X_{n,c}$ is a closed subset of the product space Λ^n , and therefore $X_{n,c}$ is a compact metric space with the induced metric from Λ^n . (We have used this fact repeatedly in [16]; see, for example, the beginning of [16, proof of Lemma 2.3]).

The proof is divided into four steps.

Claim A: *given a finite orbit $(z_i)_0^n \in X_{n,c}$ there exist $\epsilon > 0$ and ρ with $\epsilon < \rho$ such that for any $(w_i)_0^n$ in the open set $V_0 \subset X_{n,c}$ consisting of all finite orbits which are ϵ -close to $(z_i)_0^n$, we have:*

- (i) *the univalent branches $\varphi = g_{z_0 \dots z_n}$ and $\psi = g_{w_0 \dots w_n}$ coincide on $B_\rho(z_0)$; and*
- (ii) *the image of $B_\rho(z_0)$ under φ contains $B_\epsilon(z_n)$. Consequently, φ^{-1} and ψ^{-1} coincide on $B_\epsilon(z_n)$ and both send w_n to w_0 .*

Let us prove the case $n = 1$. Recall that if $z_0 \neq 0$, then there exist q univalent branches $g_i : U \rightarrow \mathbb{C}$ defined on a neighbourhood of z_0 such that $\mathbf{f}_c(U)$ is completely determined by the union of all $g_i(U)$. We may assume, without loss of generality, that for some $\delta > 0$, the diameter $|g_i(U)|$ is less than δ and $|g_i(z_0) - g_j(z_0)| > 9\delta$, whenever $i \neq j$. (Roughly speaking, the sets $g_i(U)$ are very small and away from each other). Hence, for some $0 < \epsilon_1 < \delta$, if $w_0 \mapsto w_1$ is ϵ_1 -close to $z_0 \mapsto z_1$, it follows that g_{z_0, z_1} and g_{w_0, w_1} must be determined by the same univalent branch $g_{i_0} : U \rightarrow \mathbb{C}$, otherwise z_1 and w_1 would not belong to the same $g_{i_0}(U)$, and consequently, $|z_1 - w_1| > \epsilon_1$, which is impossible, since we have a pair of ϵ_1 -close orbits. There exists $\rho > 0$ such that $B_\rho(z_0) \subset U$. Hence $g_{z_0, z_1}|_{B_\rho(z_0)} = g_{w_0, w_1}|_{B_\rho(z_0)} = g_{i_0}|_{B_\rho(z_0)}$, and it follows that z_1 belongs to $g_{i_0}(B_\rho(z_0))$. There exists $\epsilon > 0$ such that $B_\epsilon(z_1) \subset g_{i_0}(B_\rho(z_0))$, with $\epsilon < \epsilon_1$ and $\epsilon < \rho$. Then (i) and (ii) of Claim A follow with this choice of ϵ and ρ . This proves Claim A for $n = 1$. The general case $n \geq 1$ follows by induction.

Our next step is to eliminate the dependence of ϵ on $(z_i)_0^n$.

Claim B: *for a fixed $n \geq 1$, there exists $0 < \epsilon < 1$ such that for any pair of ϵ -close orbits $(z_i)_0^n$ and $(w_i)_0^n$ in $X_{n,c}$, both maps $(g_{z_0 \dots z_n})^{-1}$ and $(g_{w_0 \dots w_n})^{-1}$ are well defined and coincide on $B_\epsilon(z_n)$.*

Using Claim A, we construct a finite covering $\{V_i\}$ of $X_{n,c}$ and finitely many $\rho_i > \epsilon_i > 0$ for which (i) and (ii) of Claim A hold, whenever $(z_i)_0^n$ and $(w_i)_0^n$ are in V_i . Then chose $0 < \epsilon < 1$ as being a Lebesgue number of the covering such that $\epsilon < \epsilon_i$, for every i . Any pair of ϵ -close orbits are within the same V_i ; by Claim A (ii) the corresponding inverses coincide on $B_{\epsilon_i}(z_n)$. Since $\epsilon < \epsilon_i$, they coincide on $B_\epsilon(z_n)$.

(The following ϵ does not depend on n).

Claim C: *there exists $0 < \epsilon < 1$ such that, for every $n \geq 1$ and every $(z_i)_0^n$ in $X_{n,c}$, the domain of $(g_{z_0 \dots z_n})^{-1}$ contains $B_\epsilon(z_n)$. Moreover, if $(w_i)_0^n$ is ϵ -close to $(z_i)_0^n$ then $(g_{w_0 \dots w_n})^{-1}$ is defined on $B_\epsilon(z_n)$ and coincide with $(g_{z_0 \dots z_n})^{-1}$ on $B_\epsilon(z_n)$.*

We shall postpone the proof of this claim.

Claim D: $\epsilon_0 = \epsilon/4$ is an expansive constant, where ϵ is given by Claim C.

(Therefore, Lemma 2.2 follows from Claim D with ϵ replaced by $\epsilon/4$.)

Proof of Claim D. Suppose $(z_i)_0^\infty$ is ϵ_0 -close to $(w_i)_0^\infty$. We need to show that $z_n = w_n$, for every n . Fix n and let $\{\varphi_m\}$ denote the family of univalent maps $(g_{z_n \dots z_m})^{-1}$; by Claim C they are all defined on $B_\epsilon(z_m)$, for any $m > n$, and each φ_m maps w_m to w_n . By (2.2), the absolute value of the derivative of $(g_{z_n \dots z_m})^{-1}$ at z_m is at most $C^{-1}\lambda^{-(m-n)}$, where $C > 0$ and $\lambda > 1$ are the constants of (2.2). Using (2.2) and Koebe's Lemma 2.1 applied to the family of all $\varphi_m : B_\epsilon(z_m) \rightarrow \mathbb{C}$ we have

$$(2.4) \quad \begin{aligned} |z_n - w_n| &= |\varphi_m(z_m) - \varphi_m(w_m)| \leq KC^{-1}\lambda^{-(m-n)}|z_m - w_m| \\ &\leq KC^{-1}\lambda^{-(m-n)}2\epsilon_0 \rightarrow 0, \end{aligned}$$

as $m \rightarrow \infty$. Notice that K is the constant of Koebe's Lemma 2.1 and we have used the fact $|z_m - w_m| < \epsilon/4$, which is fundamental for the application of Lemma 2.1. We conclude that $z_n = w_n$, for every n .

Proof of Claim C. Let K be the constant of Lemma 2.1; let $C > 0$ and $\lambda > 1$ as in (2.2). There exists $n > 0$ such that $KC^{-1}\lambda^{-n} < 1/2$. Keep n fixed and find a corresponding $\epsilon' > 0$ satisfying the properties of Claim B. Let $k > 0$ be an integer. Suppose $(w_i)_0^{nk}$ and $(z_i)_0^{nk}$ are finite orbits contained in Λ which are ϵ' -close to one another. By Claim B, for any $0 < i \leq k$, the maps $\varphi_i = (g_{z_{n(i-1)} \dots z_{ni}})^{-1}$ and $\psi_i = (g_{w_{n(i-1)} \dots w_{ni}})^{-1}$ are well defined and coincide on $B_{\epsilon'}(z_{ni})$. Using the same argument of (2.4) in the proof of Claim D, we conclude that the maps φ_i and ψ_i are well defined contractions by factor $1/2$ on $B_{\epsilon'/4}(z_{ni})$ and $B_{\epsilon'/4}(w_{ni})$, respectively. Moreover, they define the same inverse branch on $B_{\epsilon'/4}(z_{ni})$ and φ_i maps $B_{\epsilon'/4}(z_{ni})$ into $B_{\epsilon'/8}(z_{n(i-1)})$. This ball of radius $\epsilon'/8$ is obviously contained in the domain $B_{\epsilon'}(z_{n(i-1)})$ of $(g_{z_{n(i-2)} \dots z_{n(i-1)}})^{-1}$; so that we are allowed to perform the compositions:

$$\begin{aligned} (g_{z_0 \dots z_{nk}})^{-1} &= (g_{z_0 \dots z_n})^{-1} \circ \dots \circ (g_{z_{n(k-1)} \dots z_{nk}})^{-1}, \\ (g_{w_0 \dots w_{nk}})^{-1} &= (g_{w_0 \dots w_n})^{-1} \circ \dots \circ (g_{w_{n(k-1)} \dots w_{nk}})^{-1}, \end{aligned}$$

thereby showing that $(g_{w_0 \dots w_{nk}})^{-1}$ and $(g_{z_0 \dots z_{nk}})^{-1}$ are well defined and coincide on $B_{\epsilon'/4}(z_{nk})$. Since $k > 0$ is arbitrary, Claim C follows with $\epsilon = \epsilon'/4$.

This completes the proof. \square

Definition 2.4 (LEO hyperbolic repellers). *The correspondence \mathbf{f}_c is called locally eventually onto (LEO) on a hyperbolic repeller Λ if for every relatively open subset U of Λ there exists $n > 0$ such that $\mathbf{f}_c^n(U)$ contains Λ .*

According to Theorem 4.3 of [16], every LEO hyperbolic repeller Λ of \mathbf{f}_c is contained in the Julia set $J(\mathbf{f}_c)$.

Definition 2.5 (Simple centre). *The critical point $z = 0$ has infinitely many forward orbits under \mathbf{f}_c . If precisely one forward orbit of the critical point of \mathbf{f}_c is periodic and the others diverge to ∞ , we say that c is a simple centre.*

The following result is stated as Theorem 5.4 in [15].

Theorem 2.1. *If a is a simple centre for the family of holomorphic correspondences \mathbf{f}_c , then $J(\mathbf{f}_c)$ is a LEO hyperbolic repeller for every c in a neighbourhood of a .*

We shall now summarise some results of [16] concerning \mathbb{C}^2 -extensions.

2.2. The \mathbb{C}^2 -extension. Suppose $J(\mathbf{f}_{c_0})$ is a LEO hyperbolic repeller for the correspondence \mathbf{f}_{c_0} . There exists a family of holomorphic maps

$$(2.5) \quad f_c : V \rightarrow \mathbb{C}^2$$

defined on a open subset V of \mathbb{C}^2 and parameterised on a neighbourhood U of c_0 , such that the closure of the periodic points of f_c , denoted by $J(f_c)$, is a subset of \mathbb{C}^2 which is completely invariant under f_c . (See [16, Lemma 2.3]). Moreover, f_c is a p -to-1 map on $J(f_c)$. The notation $J(f_c)$ suggests the definition of a Julia set, or that every periodic point of f_c is repelling. As a matter of fact, the Jacobian determinant of f_c^n at every periodic point of period n is strictly greater than 1. This fact is explained in Remark 2.4 of [16] and somehow justifies the notation $J(f_c)$. In spite of this analogy, for technical reasons $J(f_c)$ shall not be referred to as the Julia set of f_c .

The family (2.5) is defined for parameters in a neighbourhood of c_0 and enjoys some important properties.

- (A) There exists a holomorphic motion $h_c : J(\mathbf{f}_{c_0}) \rightarrow J(f_c)$ parameterised on U and based at c_0 , given by a family of conjugacies h_c from $J(\mathbf{f}_{c_0})$ to $J(f_c)$. (See Theorems B and C of [16]).
- (B) The projection $\pi(z, w) = z$ establishes a semiconjugacy $\pi : J(f_c) \rightarrow J(\mathbf{f}_c)$ between f_c and \mathbf{f}_c , in the sense that $J(\mathbf{f}_c) = \pi J(f_c)$ is also a LEO hyperbolic repeller for \mathbf{f}_c and $\pi f_c(x)$ is an image of $\pi(x)$ under \mathbf{f}_c , for every x in $J(f_c)$. Hence

$$\pi(x) \rightarrow \pi f(x) \rightarrow \pi f^2(x) \rightarrow \dots$$

is a forward orbit under \mathbf{f}_c . This is, by definition, the *projected orbit* of x . (See Lemma 2.3 of [16]).

3. THE BOWEN PARAMETER

Recall that a continuous surjective map $f : X \rightarrow X$ of a compact metric space is *expanding* if there exists $\ell > 1$ such that every point in X has a neighbourhood V such that $f^{-1}(V)$ is a finite union of disjoint open sets U_j , each of which is mapped homeomorphically onto V and

$$d(f(x), f(y)) > \ell d(x, y)$$

for x and y in U_j .

The LEO property may be defined for every $f : X \rightarrow X$. It means that every nonempty open set of X is eventually mapped onto X . The following result is stated as Theorem 4.3 in [16].

Theorem 3.1. *Let c_0 be a parameter in \mathbb{C} such that $J(\mathbf{f}_{c_0})$ is a LEO hyperbolic repeller for \mathbf{f}_{c_0} . For every c in a neighbourhood of c_0 , the map f_c of the family (2.5) is LEO on $J(f_c)$ and expanding with respect to the metric*

$$d_s(x, y) = \sum_{n=0}^{\infty} s^{-n} |\pi f_c^n(x) - \pi f_c^n(y)|$$

where $s > 1$ is arbitrary and x, y belong to $J(f_c)$.

The *dynamic ball* of radius ϵ , time n and centre x is defined by

$$B(x, n, \epsilon) = \{y \in X : d(f^j x, f^j y) < \epsilon, 0 \leq j \leq n\}.$$

Every expanding map $f : X \rightarrow X$ has an *expansive constant* $\epsilon > 0$ characterised by the fact that if $d(f^n x, f^n y) < \epsilon$ for every $n \geq 0$, then $x = y$. (It is clear that any other positive real number $\epsilon_0 < \epsilon$ is also an expansive constant.)

A continuous map $f : X \rightarrow X$ is *topologically mixing* if for every pair of nonempty open subsets U and V there exists n_0 such that $f^n(U) \cap V$ is nonempty, for every $n \geq n_0$. LEO maps are topologically mixing.

Suppose f is an expanding map of a compact metric space (X, d) , and $\phi : X \rightarrow \mathbb{R}$ is a potential (i.e., a real valued continuous function). The topological pressure of ϕ with respect to the system f is denoted by $P(\phi)$, and $S_n \phi(x)$ denotes a Birkhoff sum. Since f is expanding, there exists an expansive constant $\epsilon > 0$. If μ is a probability measure on X and there exists $C_\epsilon > 0$ such that

$$C_\epsilon^{-1} \leq \frac{\mu(B(x, n, \epsilon))}{\exp(S_n \phi(x) - nP(\phi))} \leq C_\epsilon,$$

for every x in X and $n > 0$, then μ is a *Gibbs measure* of f and ϕ . (See [10, Chap. 4]).

3.1. Transfer operators. Let $f : X \rightarrow X$ be an expanding map of a compact metric space (X, d) . The *transfer operator* $\mathcal{L}_\phi : C(X) \rightarrow C(X)$ with potential ϕ acts on the space $C(X)$ of continuous complex valued functions $g : X \rightarrow \mathbb{C}$, and is defined by

$$\mathcal{L}_\phi g(x) = \sum_{f(y)=x} e^{\phi(y)} g(y).$$

(See [10, Chap. 4]). The iterate \mathcal{L}_ϕ^n is precisely the transfer operator $\mathcal{L}_{S_n \phi}$ with respect to $f^n : X \rightarrow X$.

The dual operator $\mathcal{L}_\phi^* : M(X) \rightarrow M(X)$ acts on the space of complex measures defined on the Borel σ -algebra of X . It is defined by $\langle \mathcal{L}_\phi^* \mu, g \rangle = \langle \mu, \mathcal{L}_\phi g \rangle$, for every μ in $M(X)$ and g in $C(X)$, where $\langle \mu, g \rangle = \int_X g d\mu$. The following theorem summarises some well known results concerning the transfer operator, and the first three sentences are often referred to as the *Ruelle-Perron-Frobenius theorem* [10, Chap. 4].

Theorem 3.2. Suppose $f : X \rightarrow X$ is an expanding and topologically mixing map of a compact metric space X . If $\phi : X \rightarrow \mathbb{R}$ is Hölder continuous and $\lambda_\phi = \exp P(\phi)$, then:

- (a) there exists a unique probability measure ν on X such that $\mathcal{L}_\phi^*(\nu) = \lambda_\phi \nu$;
- (b) there exists a unique real valued continuous function $h > 0$ on X such that $\mathcal{L}_\phi(h) = \lambda_\phi h$ and $\int_X h d\nu = 1$;
- (c) for any continuous real valued function g on X ,

$$\lambda_\phi^{-n} \mathcal{L}_\phi^n(g) \rightarrow h \int_X g d\nu,$$

as $n \rightarrow \infty$, and the convergence is uniform on X ;

- (d) the measure $\mu(A) = \int_A h d\nu$ is the unique invariant Gibbs measure of f and ϕ .

Remark 3.1. The number $P(0)$ is also known as the topological entropy of f . Suppose that $f : X \rightarrow X$ is a d -to-1 map (i.e., every point has exactly d preimages). Then f^n is d^n -to-1. By Theorem 3.2,

$$\lambda_\phi^{-n} \mathcal{L}_0^n(1) = \lambda_\phi^{-n} \sum_{f^n(y)=x} e^{S_n \phi(y)} \cdot 1 = \lambda_\phi^{-n} d^n$$

converges to $h > 0$, where $\phi = 0$. It follows that $\lambda_\phi = d$, and the topological entropy of f must be $\log(d)$.

The following result introduces the Bowen parameter t_c .

Let $z_0(x)$ and $z_1(x)$ denote the first two elements of the projected orbit of an element x of $J(f_c)$. We define $\varphi_c : J(f_c) \rightarrow \mathbb{R}$ by $\varphi_c(x) = -\log |g'_{z_0, z_1}(z_0)|$, where z_0 and z_1 are $z_0(x)$ and $z_1(x)$, respectively. The map φ_c is therefore a potential on $J(f_c)$, and the zero of the pressure function $t \mapsto P(t\varphi_c)$ is defined as the *Bowen parameter* t_c . However, we need to show that there exists only one zero of the pressure function. This is established in the following theorem. Note that $\varphi_c(x)$ is defined directly by (3.1), for then $g'_{z_0, z_1}(z_0)$ equals $p(z_1 - z_0)/(qz_0)$.

Theorem 3.3 (Bowen parameter). *If the Julia set $J(f_c) \subset \mathbb{C}$ of the holomorphic correspondence $(w - c)^q = z^p$ is a LEO hyperbolic repeller, then the topological entropy of the \mathbb{C}^2 -extension*

$$f_c : J(f_c) \rightarrow J(f_c)$$

is strictly positive. Moreover, the potential

$$(3.1) \quad \varphi_c(x) = -\log |p(z_1 - z_0)/(qz_0)|$$

is Hölder continuous with respect to the metric d_s on $J(f_c)$, and there exists a unique zero $t_c > 0$ of the pressure function $t \mapsto P(t\varphi_c)$ defined on $[0, \infty)$.

Proof. According to [1, Chap. 2B], the topological pressure of a potential ϕ can be calculated by

$$(3.2) \quad P(\phi) = \lim_{|\mathcal{C}| \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{n} \log \left(\inf_{\mathcal{C}_n} \sum_{U \in \mathcal{C}_n} \exp S_n \phi(U) \right),$$

where \mathcal{C} is any finite covering of $J(f_c)$; $|\mathcal{C}|$ is the greatest diameter of an element of \mathcal{C} ; and $S_n \phi(U)$ is given by the supremum of all $S_n \phi(x)$ when $x \in U$. The infimum in (3.2) is taken over all possible covers \mathcal{C}_n of $J(f_c)$ whose elements can be written as

$$U = \{x \in J(f_c) : f_c^j(x) \in U_j, 0 \leq j \leq n\},$$

for some finite sequence $(U_j)_1^n$ of elements of \mathcal{C} .

Since $t \mapsto P(t\varphi_c)$ is continuous, in order to prove the existence of a unique zero it suffices to show that the pressure function is strictly decreasing with $P(0) > 0$ and $P(t\varphi_c) \rightarrow -\infty$ as $t \rightarrow \infty$.

Indeed, $J(f_c)$ is a hyperbolic repeller, and therefore $|g'_{z_0 \dots z_n}(z_0)| \geq C\lambda^n$, where $\lambda > 1$ and $(z_j)_0^\infty$ denotes the projected orbit of any x . By Theorem 3.3 and a simple computation involving the logarithm,

$$(3.3) \quad \exp S_n(t\varphi_c)(x) = |g'_{z_0 \dots z_n}(z_0)|^{-t}.$$

Hence,

$$\begin{aligned} P((t + \tau)\varphi_c) &= \lim_{|\mathcal{C}| \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{n} \log \left(\inf_{\mathcal{C}_n} \sum_{U \in \mathcal{C}_n} \exp S_n(t\varphi_c)(U) \cdot \exp S_n(\tau\varphi_c)(U) \right) \\ &= \lim_{|\mathcal{C}| \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{n} \log \left(\inf_{\mathcal{C}_n} \sum_{U \in \mathcal{C}_n} \sup_{x \in U} |g'_{z_0 \dots z_n}(z_0)|^{-t} \cdot \exp S_n(\tau\varphi_c)(U) \right) \\ &\leq \lim_{|\mathcal{C}| \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{n} \log (C^{-t} \lambda^{-nt} \inf_{\mathcal{C}_n} \sum_{U \in \mathcal{C}_n} \exp S_n(\tau\varphi_c)(U)) \\ &= -t \log(\lambda) + P(\tau\varphi_c). \end{aligned}$$

Since $\log(\lambda) > 0$, we conclude that the pressure function is indeed strictly decreasing and $P(t\varphi_c) \rightarrow -\infty$ as $t \rightarrow \infty$. Since f_c is a p -to-one map, it follows from Remark 3.1 that $P(0) = \log(p) > 0$. Hence, there exists a unique zero t_c of the pressure function in $(0, \infty)$.

The proof of the Hölder continuity of the potential with respect to the metric d_s is a straightforward application of the mean value theorem. \square

Remark 3.2. *The value of t_c does not depend on the particular choice of f_c .*

In fact, if we consider the space X_c of all forward orbits $(z_i)_0^\infty$ under f_c which are contained in $J(f_c)$, then $g_c(x) = (\pi f_c^n(x))_{n=0}^\infty$ defines a homeomorphism from $J(f_c)$ onto X_c which conjugates the dynamics of f_c to the dynamics of the (left) one-sided shift σ on X_c [16, pages 3110-3112]. Since the pressure is a topological invariant, this is sufficient to show that t_c depends only on the dynamics of the shift map on the space of orbits.

Indeed,

$$(3.4) \quad P(t\varphi_c; f_c) = P(t\varphi_c \circ g_c^{-1}; \sigma).$$

The notation $P(\phi; f)$ makes explicit the dependence of the pressure on the dynamics of f . By definition, g_c^{-1} maps every $(z_i)_0^\infty$ in X_c to the unique $x \in J(f_c)$ such that $\pi f_c^n(x) = z_n$, for every n . Since φ_c is defined by (3.1) and $(z_i)_0^\infty$ is the projected orbit of x ,

$$(t\varphi_c \circ g_c^{-1})((z_i)_0^\infty) = -\log |p(z_1 - z_0)/(qz_0)|.$$

This shows that the potential on the right side of (3.4) does not depend on g_c or f_c . Hence the same is true for the unique zero of the pressure function, as desired.

4. HAUSDORFF DIMENSION

The s -dimensional Hausdorff outer measure \mathcal{H}^s of a set $\Lambda \subset \mathbb{C}$ is defined by

$$\mathcal{H}^s(\Lambda) = \liminf_{\delta \rightarrow 0} \sum_{i=1}^{\infty} |U_i|^s,$$

where \inf is taken over all countable coverings $\{U_i\}_0^\infty$ of Λ with diameter $|U_i| \leq \delta$. There exists a unique nonnegative real number d characterised by the following properties: $\mathcal{H}^s(\Lambda) = 0$ if $s > d$ and $\mathcal{H}^s(\Lambda) = \infty$ if $0 \leq s < d$. The number d is, by definition, the *Hausdorff dimension* of Λ , denoted by $\dim_H \Lambda$.

Theorem 4.1. *Suppose $J(f_c)$ is a LEO hyperbolic repeller and let t_c be the unique zero of the associated pressure function. Then $\mathcal{H}^{t_c}(J(f_c))$ is finite, and therefore*

$$\dim_H J(f_c) \leq t_c.$$

Proof. Fix a parameter c for which $\Lambda = J(f_c)$ is a hyperbolic repeller. By Lemma 2.2, Λ has an expansive constant $\epsilon > 0$. Cover $J(f_c) \subset \mathbb{C}^2$ with finitely many dynamic balls of fixed time n and radius ϵ :

$$B(x_i, n, \epsilon) = \{y \in J(f_c) : d_s(f_c^j(x_i), f_c^j(y)) < \epsilon, 0 \leq j \leq n\}.$$

The finite covering is possible because $J(f_c)$ is compact. The covering is called *minimal* if $x_j \notin B(x_i, n, \epsilon)$ whenever $i \neq j$. We may assume that the covering is minimal by removing $B(x_i, n, \epsilon)$ if x_i is contained in some other $B(x_j, n, \epsilon)$. Since the covering is minimal, the corresponding dynamic balls with radius $\epsilon/2$ are pairwise disjoint, for if $B(x_i, n, \epsilon/2)$ intersects $B(x_j, n, \epsilon/2)$, then by the triangle inequality x_j must belong to $B(x_i, n, \epsilon)$. But x_j should not belong to $B(x_i, n, \epsilon)$ when $i \neq j$ (the covering is minimal).

It will be convenient to denote such covering of $J(f_c)$ by \mathcal{V}_n , and its elements by $V_i = B(x_i, n, \epsilon)$. Now we are going to construct a covering of $J(f_c)$.

Claim: the family of all sets $U_i = (g_{z_0 \dots z_n})^{-1}(B_\epsilon(z_n))$, where $(z_k)_0^\infty$ is the projected orbit of the centre x_i of V_i , is a covering of $J(\mathbf{f}_c)$. We shall denote this covering by \mathcal{U}_n .

By Lemma 2.2, each set U_i is well defined since the domain of $(g_{z_0 \dots z_n})^{-1}$ contains $B_\epsilon(z_n)$. As we shall see, the sets U_i do in fact cover $J(\mathbf{f}_c)$ because each U_i contains πV_i , $J(\mathbf{f}_c)$ is the projection of $J(f_c)$, and $\{V_i\}$ is a covering of $J(f_c)$. In order to check that πV_i is contained in U_i it suffices to show that $\pi(y)$ is in U_i , whenever $y \in V_i$. Since y is in $B(x_i, n, \epsilon)$, by definition the distance with respect to the metric d_s between $f_c^j(x_i)$ and $f_c^j(y)$ is strictly less than ϵ , for $0 \leq j \leq n$. This means

$$\sum_{k=0}^{\infty} s^{-k} |\pi f_c^{k+j}(x_i) - \pi f_c^{k+j}(y)| < \epsilon,$$

for $0 \leq j \leq n$. In particular, the first term of the above series is less than ϵ , and it follows that $|z_j - w_j| < \epsilon$ for $0 \leq j \leq n$, where $(z_j)_0^\infty$ and $(w_j)_0^\infty$ are the projected orbits of x_i and y , respectively. By Lemma 2.2, $(g_{z_0 \dots z_n})^{-1}$ maps w_n to w_0 . Hence $w_0 = \pi(y)$ belongs to U_i , as desired.

The diameter of \mathcal{U}_n is defined as the supremum of all diameters $|U_i|$.

Claim: the diameter of \mathcal{U}_n tends to zero as $n \rightarrow \infty$.

By Definition 2.2 and Koebe's Lemma 2.1, for every element U_i of \mathcal{U}_n we have

$$|U_i| = |(g_{z_0 \dots z_n})^{-1}(B_\epsilon(z_n))| \leq K |g'_{z_0 \dots z_n}(z_0)|^{-1} 2\epsilon \leq (2\epsilon K C^{-1}) \lambda^{-n} \rightarrow 0,$$

as $n \rightarrow \infty$. The claim is proved.

By Theorem 3.1, $f_c : J(f_c) \rightarrow J(f_c)$ is LEO and expanding. Therefore, for every Hölder continuous potential there corresponds a unique invariant Gibbs measure. Let $\phi = t_c \varphi_c$. Since $P(\phi) = 0$ and $\epsilon/2$ is also an expansive constant, the corresponding Gibbs measure μ for the system satisfies:

$$(4.1) \quad C_{\epsilon/2}^{-1} \leq \frac{\mu(B(x, n, \epsilon/2))}{\exp S_n \phi(x)} \leq C_{\epsilon/2},$$

for any x in $J(f_c)$ and n , where $C_{\epsilon/2}$ is a constant independent of x and n . By Koebe's Lemma 2.1, (3.3) and (4.1) we have

$$(4.2) \quad \begin{aligned} |U_i|^{t_c} &= |(g_{z_0 \dots z_n})^{-1}(B_\epsilon(z_n))|^{t_c} \leq (K |g'_{z_0 \dots z_n}(z_0)|^{-1} 2\epsilon)^{t_c} \\ &\leq (2\epsilon K)^{t_c} C_{\epsilon/2} \mu(B(x_i, n, \epsilon/2)), \end{aligned}$$

for every element U_i of \mathcal{U}_n . Since μ is a probability measure and the dynamic balls $B(x_i, n, \epsilon/2)$ are pairwise disjoint, we conclude that $\sum_{i=0}^{\infty} |U_i|^{t_c} \leq (2\epsilon K)^{t_c} C_{\epsilon/2}$. Since \mathcal{U}_n is a covering of $J(\mathbf{f}_c)$ whose diameter tends to zero as $n \rightarrow \infty$, the t_c -dimensional Hausdorff measure of $J(\mathbf{f}_c)$ is finite. Hence, $\dim_H J(\mathbf{f}_c) \leq t_c$. \square

Corollary 4.1.1. For any parameter c sufficiently close to a simple centre,

$$\dim_H J(\mathbf{f}_c) \leq t_c.$$

Proof. By Theorem 5.4 of [15], the Julia set $J(\mathbf{f}_c)$ is a LEO hyperbolic repeller, for every c in a certain subset H_β of the parameter space. By definition 4.1 of [15], the set H_β includes a sufficiently small neighbourhood of every simple centre. \square

Corollary 4.1.2. Suppose $J(\mathbf{f}_c)$ is a LEO hyperbolic repeller and $|g'_{z,w}(z)| \geq \kappa > 1$, for every z, w in J_c such that w is an image of z under \mathbf{f}_c . Then

$$(4.3) \quad \dim_H J(\mathbf{f}_c) \leq (\log p)/(\log \kappa).$$

Proof. Let $\phi = t_c \varphi_c$. Since $P(\phi) = 0$, by the Ruelle-Perron-Frobenius Theorem 3.2, $\mathcal{L}_\phi^n(1) \rightarrow h$ uniformly on $J(f_c)$, where h is a continuous function from $J(f_c)$ to $(0, \infty)$. By (3.3), we have

$$(4.4) \quad \mathcal{L}_\phi^n(1)(x) = \sum_{f_c^n(y)=x} e^{S_n \phi(y)} \cdot 1 = \sum_{f_c^n(y)=x} |g'_{w_0 \dots w_n}(w_0)|^{-t_c} \leq p^n \kappa^{-nt_c},$$

where $(w_i)_0^\infty$ is the projected orbit of y . In the above estimate we have used the hypothesis and the fact that f_c^n is a p^n -to-1 map. Since $\mathcal{L}_\phi^n(1)$ converges to a positive function, we have $p\kappa^{-t_c} \geq 1$, otherwise $p^n \kappa^{-nt_c}$ would converge to zero. Solving the inequality $p\kappa^{-t_c} \geq 1$ for t_c yields (4.3). \square

Corollary 4.1.3 (Zero area). *If $q^2 < p$ and c is sufficiently close to zero, then*

$$(4.5) \quad \dim_H J(\mathbf{f}_c) < 2.$$

Proof. The derivative of a branch $g(z)$ of \mathbf{f}_c is given by $(p/q)(w-c)/z$ where $w = g(z)$. The set function $c \mapsto J(\mathbf{f}_c)$ is continuous with respect to the Hausdorff topology for parameters c in an open set H_β which contains zero and every simple centre (see Theorem 5.11 of [15]). Since $J(\mathbf{f}_0)$ is the unit circle \mathbb{S}^1 , we conclude that $|g'_{z,w}(z)| \geq \kappa$ for z and w in $J(\mathbf{f}_c)$, where $\kappa > 1$ can be chosen arbitrarily close to p/q as c tends to zero. It is easy to check that $\log(p)/\log(p/q) < 2$ if, and only if, $q^2 < p$. In particular, $\log(p)/\log(\kappa) < 2$ for κ close to p/q . From Corollary 4.1.2 we conclude (4.5), for every c sufficiently close to zero. \square

Remark 4.1. *For arbitrary integers $p > q \geq 2$, we have $t_0 > 1$, and therefore the estimate of Theorem 4.1 is not sharp if $c = 0$.*

Proof. Indeed, in (4.4) $|g'_{w_0 \dots w_n}(w_0)|$ equals $(p/q)^n$ when all the elements of the projected orbit $(w_i)_0^\infty$ are contained in \mathbb{S}^1 . It follows from (4.4) that $\mathcal{L}_\phi^n(1)$ is simply $(p/q)^{-nt_0} p^n$ when ϕ is the potential $t_0 \varphi_0$. In particular, $\mathcal{L}_\phi(1)$ is a constant function $\omega \cdot 1$, where ω is a real number and 1 is the function which is constantly one on $J(f_0)$. Since $\mathcal{L}_\phi^n(1)$ is $\omega^n \cdot 1$, which converges to a positive function $h > 0$ on $J(f_0)$, it follows that $\omega = 1$ and consequently $h = 1$. Since $\mathcal{L}_\phi(1)$ is $(p/q)^{-t_0} p$ and 1 is a fixed point of \mathcal{L}_ϕ , it follows that $(p/q)^{-t_0} p = 1$. Since $p > q \geq 2$, we have $t_0 > 1$.

Corollary 4.1.4. *If every orbit of zero under \mathbf{f}_c diverges to infinity, then*

$$\dim_H J(\mathbf{f}_c) \leq t_c.$$

Proof. According to [15, Theorem 5.4], if every orbit of zero under \mathbf{f}_c diverges to infinity, then $J(\mathbf{f}_c)$ is a LEO hyperbolic repeller. It follows from Theorem 4.1 that $\dim_H J(\mathbf{f}_c) \leq t_c$. \square

Acknowledgments. Research partially supported by the grants 2016/16012-6 São Paulo Research Foundation and CNPq 232706/2014-0. The author would like to thank Daniel Smania for many discussions and key insights which led to some preliminary results of this paper, already described in the author's PhD thesis [14].

REFERENCES

1. Rufus Bowen, *Hausdorff dimension of quasi-circles*, Publications Mathématiques de l'Institut des Hautes Études Scientifiques **50** (1979), no. 1, 11–25.
2. S. Bullett and L. Lomonaco, *Mating quadratic maps with the modular group II.*, Invent. Math. **220** (2020), 185–210.
3. Shaun Bullett and Luna Lomonaco, *Dynamics of modular matings*, arXiv preprint: <https://arxiv.org/abs/1707.04764>, 2017.
4. ———, *Mating quadratic maps with the modular group III: the modular Mandelbrot set*, arXiv preprint: <https://arxiv.org/pdf/2010.04273.pdf>, 2020.

5. Shaun Bullett, Luna Lomonaco, and Carlos Siqueira, *Correspondences in complex dynamics*, New Trends in One-Dimensional Dynamics (Cham) (Maria José Pacifico and Pablo Guarino, eds.), Springer International Publishing, 2019, pp. 51–75.
6. Shaun Bullett and Christopher Penrose, *Mating quadratic maps with the modular group*, *Inventiones mathematicae* **115** (1994), no. 1, 483–511.
7. Adrien Douady and John H. Hubbard, *Étude dynamique des polynômes complexes. Partie I.*, Publications Mathématiques d’Orsay **84** (1984), no. 2, 75.
8. ———, *Étude dynamique des polynômes complexes. Partie II. (with the collaboration of P. Lavaurs, Tan Lei and P. Sentenac).*, Publications Mathématiques d’Orsay **85** (1985), no. 4, v–154.
9. Seung-Yeop Lee, Mikhail Lyubich, Nikolai G. Makarov, and Sabyasachi Mukherjee, *Schwarz reflections and anti-holomorphic correspondences*, *Advances in Mathematics* **385** (2021), 107766.
10. Feliks Przytycki and Mariusz Urbanski, *Conformal fractals: Ergodic theory methods*, 1st ed., Cambridge University Press, New York, NY, USA, 2010.
11. Juan Rivera-Letelier, *On the continuity of Hausdorff dimension of Julia sets and similarity between the Mandelbrot set and Julia sets*, *Fund. Math.* **170** (2001), no. 3, 287–317.
12. David Ruelle, *Repellers for real analytic maps*, *Ergodic Theory and Dynamical Systems* **2** (1982), 99–107 (eng).
13. Károly Simon, *The Hausdorff dimension of the Smale-Williams solenoid with different contraction coefficients*, *Proceedings of the American Mathematical Society* **125** (1997), no. 4, 1221–1228 (eng).
14. Carlos Siqueira, *Dynamics of holomorphic correspondences*, Ph.D. thesis, University of São Paulo, ICMC, Digital library USP, 8 2015.
15. Carlos Siqueira, *Dynamics of hyperbolic correspondences*, *Ergodic Theory and Dynamical Systems* (2021), first view 1–32.
16. Carlos Siqueira and Daniel Smania, *Holomorphic motions for unicritical correspondences*, *Nonlinearity* **30** (2017), no. 8, 3104.

DEPARTAMENTO DE MATEMÁTICA, IME-UFBA. SALVADOR – BA, BRAZIL. CEP 40170-115
 Email address, C. Siqueira: carlos.siqueira@ufba.br