

$\Lambda\Lambda N - \Xi NN$ S wave resonance

H. Garcilazo^{1,*} and A. Valcarce^{2,†}

¹*Escuela Superior de Física y Matemáticas,
Instituto Politécnico Nacional, Edificio 9, 07738 México D.F., Mexico*

²*Departamento de Física Fundamental,
Universidad de Salamanca, 37008 Salamanca, Spain*

(Dated: January 18, 2022)

Abstract

We use an existing model of the $\Lambda\Lambda N - \Xi NN$ three-body system based in two-body separable interactions to study the $(I, J^P) = (1/2, 1/2^+)$ three-body channel. For the $\Lambda\Lambda$, ΞN , and $\Lambda\Lambda - \Xi N$ amplitudes we have constructed separable potentials based on the most recent results of the HAL QCD Collaboration. They are characterized by the existence of a resonance just below or above the ΞN threshold in the so-called H -dibaryon channel, $(i, j^p) = (0, 0^+)$. A three-body resonance appears 2.3 MeV above the Ξd threshold. We show that if the $\Lambda\Lambda - \Xi N$ H -dibaryon channel is not considered, the $\Lambda\Lambda N - \Xi NN$ S wave resonance disappears. Thus, the possible existence of a $\Lambda\Lambda N - \Xi NN$ resonance would be sensitive to the $\Lambda\Lambda - \Xi N$ interaction. The existence or nonexistence of this resonance could be evidenced by measuring, for example, the Ξd cross section.

PACS numbers: 21.45.-v, 25.10.+s, 11.80.Jy

Keywords: baryon-baryon interactions, Faddeev equations

*Electronic address: humberto@esfm.ipn.mx

†Electronic address: valcarce@usal.es

The $\Lambda\Lambda - \Xi N$ system in a pure S wave configuration has quantum numbers $(i, j^p) = (0, 0^+)$ so that adding one more nucleon, the $\Lambda\Lambda N - \Xi NN$ system has necessarily quantum numbers $(I, J^P) = (1/2, 1/2^+)$. In a series of works based on a chiral constituent quark model [1–3] this system was studied under the assumption that the H dibaryon [4] has the lower limit mass determined by the E373 experiment at KEK [5] from the observation of a ${}^6_{\Lambda\Lambda}\text{He}$ double hypernucleus. Despite the large amount of experimental and theoretical efforts, the existence of the H dibaryon remains inconclusive, see Ref. [6] for a recent update. Experimental evidence disfavors large binding energies [7], as predicted in Ref. [4], and the high statistics study of Υ decays at Belle [8] found no indication of an H dibaryon with a mass near the $\Lambda\Lambda$ threshold. Recently, the HAL QCD Collaboration [9] has published a $N_f = 2 + 1$ study of coupled channel ($\Lambda\Lambda$ and ΞN) baryon-baryon interactions with near-physical quark masses, namely $m_\pi = 146$ MeV, concluding that the H dibaryon could be a $\Lambda\Lambda$ resonance just below or above the ΞN threshold. Similar results were obtained in a low-energy effective field theory study of the H dibaryon in $\Lambda\Lambda$ scattering [10].

The HAL QCD results are being used as input for the study of strangeness -2 baryon-baryon interactions, as recently done in relativistic chiral effective field theory studies [11]. The HAL QCD ΞN interactions have been also recently used to study the possible existence of ΞNN bound states in Ref. [12], with negative results for the $(I, J^P) = (1/2, 1/2^+)$ channel. For the NN interaction they used the AV8 potential [13]. As the coupling between $\Lambda\Lambda$ and ΞN is found to be weak in Ref. [9], they used an effective single-channel ΞN potential in which the coupling to $\Lambda\Lambda$ in ${}^{11}S_0$ was renormalized into a single Gaussian form chosen to reproduce the ΞN phase shift obtained with channel coupling. The three-body ΞNN problem is solved in the real axis by means of a variational method with Gaussian bases, the Gaussian Expansion Method [14, 15]. The full coupling between the ΞNN and $\Lambda\Lambda N$ channels was not explicitly considered. A similar calculation based on the Nijmegen ESC08c potentials [16–18] was presented in Ref. [19] also with negative results for the $(I, J^P) = (1/2, 1/2^+)$ channel, see Fig. 2(a) of Ref. [19].

Unlike the calculation in Ref. [12], we developed in Ref. [20] a model of the $\Lambda\Lambda N - \Xi NN$ three-body system which allowed us to look for possible three-body resonances. Using separable two-body interactions fitted to the low-energy data of the Nijmegen S wave baryon-baryon amplitudes [16–18], we found a resonance just below the Ξd threshold with a very

small width of only 0.09 MeV¹. Qualitatively similar results have been obtained in Ref. [12], although as stressed in this manuscript they are numerically different due to a different NN potential and different treatment of ESC08c Nijmegen S wave baryon-baryon interactions. Such dependencies on the models and parametrizations of the two-body interactions make this three-body system ideally suited for testing the different models for the two-body interactions.

However, contrary to the recent results of the HAL QCD Collaboration, the Nijmegen baryon-baryon interactions gave no indication of either a bound state or a resonance in the $\Lambda\Lambda - \Xi N$ ($0, 0^+$) two-body channel, the H -dibaryon channel. It is thus interesting to see if the existence of a resonance just below or above the ΞN threshold, as it has been found by the HAL QCD Collaboration [9] and low-energy effective field theory studies [10], may affect the position of the three-body S wave ($1/2, 1/2^+$) $\Lambda\Lambda N - \Xi NN$ resonance found in Ref. [20]. For this purpose, we have now constructed separable potential models of the $\Lambda\Lambda$, ΞN , and $\Lambda\Lambda - \Xi N$ amplitudes reproducing the behavior of the HAL QCD Collaboration results [9] and we have performed a full fledged coupled-channel study of the $\Lambda\Lambda N - \Xi NN$ three-body system.

We use rank-one separable potentials for all the uncoupled two-body channels, that is, for all channels except the $\Lambda\Lambda - \Xi N$ ($0, 0^+$) interaction. They are as follows,

$$V_i^\rho = g_i^\rho \lambda \langle g_i^\rho, \quad (1)$$

such that the two-body t-matrices are

$$t_i^\rho = g_i^\rho \tau_i^\rho \langle g_i^\rho, \quad (2)$$

with

$$\tau_i^\rho = \frac{\lambda}{1 - \lambda \langle g_i^\rho | G_0(i) | g_i^\rho \rangle}, \quad (3)$$

where $G_0(i) = 1/(E - K_i + i\epsilon)$ and K_i is the kinetic energy operator of channel i . We use Yamaguchi form factors [22] for the separable potentials of Eq. (1), i.e.,

$$g_i^\rho(p) = \frac{1}{\alpha^2 + p^2}. \quad (4)$$

¹ It is worth to note that the results for the ΞNN system with maximal isospin have been independently reproduced within the integral Faddeev equation formalism [21] in agreement with high accuracy.

TABLE I: Parameters α and β (in fm^{-1}), λ_{11} , λ_{33} , and λ_{13} (in fm^{-2}) of the separable-potential model of the coupled $(i, j^p) = (0, 0^+)$ $\Lambda\Lambda - \Xi N$ two-body system.

| α | β | λ_{11} | λ_{33} | λ_{13} |
|----------|---------|----------------|----------------|----------------|
| 1.3465 | 1.1460 | -0.1390 | -0.3171 | 0.0977 |

TABLE II: Parameters α (in fm^{-1}) and λ (in fm^{-2}) of the separable-potential model of the uncoupled (i, j^p) ΞN two-body channels.

| Channel | α | λ |
|------------|----------|-----------|
| $(0, 1^+)$ | 1.41 | -0.117 |
| $(1, 0^+)$ | 7.333 | 22.97 |
| $(1, 1^+)$ | 0.803 | -0.016 |

In the case of the coupled $(0, 0^+)$ $\Lambda\Lambda - \Xi N$ channel we use a rank-two separable potential of the form [23]

$$V_{ij}^{\rho\sigma} = g_i^\rho \lambda_{ij} \langle g_j^\sigma, \quad (5)$$

such that

$$t_{ij}^{\rho-\sigma} = g_1^\rho \tau_{ij}^{\rho-\sigma} \langle g_j^\sigma, \quad (6)$$

with

$$\begin{aligned} \tau_{11}^{\Lambda\Lambda-\Lambda\Lambda} &= \frac{-\lambda_{13}^2 G^{\Xi N} - \lambda_{11}(1 - \lambda_{33} G^{\Xi N})}{\lambda_{13}^2 G^{\Lambda\Lambda} G^{\Xi N} - (1 - \lambda_{11} G^{\Lambda\Lambda})(1 - \lambda_{33} G^{\Xi N})}, \\ \tau_{33}^{\Xi N-\Xi N} &= \frac{-\lambda_{13}^2 G^{\Lambda\Lambda} - \lambda_{33}(1 - \lambda_{11} G^{\Lambda\Lambda})}{\lambda_{13}^2 G^{\Lambda\Lambda} G^{\Xi N} - (1 - \lambda_{11} G^{\Lambda\Lambda})(1 - \lambda_{33} G^{\Xi N})}, \\ \tau_{13}^{\Lambda\Lambda-\Xi N} &= \tau_{31}^{\Xi N-\Lambda\Lambda} = \frac{-\lambda_{13}}{\lambda_{13}^2 G^{\Lambda\Lambda} G^{\Xi N} - (1 - \lambda_{11} G^{\Lambda\Lambda})(1 - \lambda_{33} G^{\Xi N})}, \end{aligned} \quad (7)$$

and

$$\begin{aligned} G^{\Lambda\Lambda} &= \langle g_1^{\Lambda\Lambda} | G_0(1) | g_1^{\Lambda\Lambda} \rangle, \\ G^{\Xi N} &= \langle g_3^{\Xi N} | G_0(3) | g_3^{\Xi N} \rangle. \end{aligned} \quad (8)$$

In this case we also use Yamaguchi-type form factors as

$$\begin{aligned} g_1^{\Lambda\Lambda}(p) &= \frac{1}{\alpha^2 + p^2}, \\ g_3^{\Xi N}(p) &= \frac{1}{\beta^2 + p^2}. \end{aligned} \quad (9)$$




FIG. 1: (a) $\Lambda\Lambda$ scattering phase shifts, (b) $\Lambda\Lambda$ inelasticity, and (c) $N\Xi$ scattering phase shifts in the $(i, j^p) = (0, 0^+)$ channel.

The parameters of the $\Lambda\Lambda - \Xi N$ model based on the latest HAL QCD potentials are given in Table I. In Figs. 1(a), (b), and (c) we show the predictions for the $\Lambda\Lambda$ and ΞN phase shifts as well as the inelasticity, which are rather similar to those of model $t/a = 12$ of the HAL QCD Collaboration presented in Fig. 4 of Ref. [9]. The corresponding parameters of the uncoupled ΞN models are given in Table II. Let us note that our results have been obtained by taking the nucleon mass as the average of the proton and neutron masses and the Ξ mass as the average of Ξ^0 and Ξ^- masses. Thus, the ΞN and ΞNN thresholds are 25.6 MeV above the $\Lambda\Lambda$ and $\Lambda\Lambda N$ thresholds, respectively. However, this threshold is 32 MeV for the HAL QCD results [9], since they use for the baryon masses the values obtained from their lattice QCD study. Therefore, to compare our phase shifts with those of Ref. [9] one should keep in mind that the energy scale of Ref. [9] corresponds to those of Fig. 1 multiplied by 1.25. The models of the NN and ΛN subsystems are the same of Ref. [20].

The coupled $\Lambda\Lambda N - \Xi NN$ three-body system presents the special characteristic that each three-body component consists of two identical fermions and a third one that is different. The homogeneous integral equations of this system appropriate for the search of bound and resonant states were derived in Ref. [3] using a graphical method. Using the new separable models of Tables I and II based in the HAL QCD interactions we did not find any bound state below the Ξd threshold, in agreement with the results of Ref. [12]. Therefore, we investigate the possible existence of a resonance above the Ξd threshold by calculating the Ξd scattering amplitude.

We adopt the same convention as in Refs. [3, 20], i.e., particles 2 and 3 are identical and particle 1 is the different one in each three-body component. After the reduction for identical particles the inhomogeneous integral equations appropriate for Ξd elastic scattering take the following form

$$\begin{aligned}
\langle 1|T_1|\phi_0\rangle &= 2\langle 1|t_1^{\Lambda\Lambda}|1\rangle\langle 1|3\rangle G_0(3)\langle 3|T_3|\phi_0\rangle + \langle 1|t_{13}^{\Lambda\Lambda-N\Xi}|3\rangle\langle 3|1\rangle G_0(1)\langle 1|U_1|\phi_0\rangle \\
&\quad - \langle 1|t_{13}^{\Lambda\Lambda-N\Xi}|3\rangle\langle 2|3\rangle G_0(3)\langle 3|U_3|\phi_0\rangle, \\
\langle 3|T_3|\phi_0\rangle &= -\langle 3|t_3^{N\Lambda}|3\rangle\langle 2|3\rangle G_0(3)\langle 3|T_3|\phi_0\rangle + \langle 3|t_3^{N\Lambda}|3\rangle\langle 3|1\rangle G_0(1)\langle 1|T_1|\phi_0\rangle, \\
\langle 1|U_1|\phi_0\rangle &= 2\langle 1|t_1^{NN}|1\rangle\langle 1|3\rangle G_0(3)\langle 3|U_3|\phi_0\rangle, \\
\langle 3|U_3|\phi_0\rangle &= 2\langle 3|t_3^{N\Xi}|\phi_0\rangle \\
&\quad - \langle 3|t_3^{N\Xi}|3\rangle\langle 2|3\rangle G_0(3)\langle 3|U_3|\phi_0\rangle + \langle 3|t_3^{N\Xi}|3\rangle\langle 3|1\rangle G_0(1)\langle 1|U_1|\phi_0\rangle \\
&\quad + 2\langle 3|t_{31}^{N\Xi-\Lambda\Lambda}|1\rangle\langle 1|3\rangle G_0(3)\langle 3|T_3|\phi_0\rangle,
\end{aligned} \tag{10}$$

where $|\phi_0\rangle$ is the initial state consisting of the deuteron wave function times a Ξ plane wave.

Using Eqs. (2) and (6) into the integral equations (10) and introducing the transformations $\langle i|T_i|\phi_0\rangle = \langle i|g_i^{\alpha_i}\rangle\langle i|X_i|\phi_0\rangle$ and $\langle i|U_i|\phi_0\rangle = \langle i|g_i^{\beta_i}\rangle\langle i|Y_i|\phi_0\rangle$ one obtains the inhomogeneous one-dimensional integral equations

$$\begin{aligned}
\langle 1|X_1|\phi_0\rangle &= 2\tau_1^{\Lambda\Lambda}\langle g_1^{\Lambda\Lambda}|1\rangle\langle 1|3\rangle G_0(3)\langle 3|g_3^{N\Lambda}\rangle\langle 3|X_3|\phi_0\rangle \\
&\quad + \tau_{13}^{\Lambda\Lambda-N\Xi}\langle g_3^{N\Xi}|3\rangle\langle 3|1\rangle G_0(1)\langle 1|g_1^{NN}\rangle\langle 1|Y_1|\phi_0\rangle \\
&\quad - \tau_{13}^{\Lambda\Lambda-N\Xi}\langle g_3^{N\Xi}|3\rangle\langle 2|3\rangle G_0(3)\langle 3|g_3^{N\Xi}\rangle\langle 3|Y_3|\phi_0\rangle, \\
\langle 3|X_3|\phi_0\rangle &= -\tau_3^{N\Lambda}\langle g_3^{N\Lambda}|3\rangle\langle 2|3\rangle G_0(3)\langle 3|g_3^{N\Lambda}\rangle\langle 3|X_3|\phi_0\rangle \\
&\quad + \tau_3^{N\Lambda}\langle g_3^{N\Lambda}|3\rangle\langle 3|1\rangle G_0(1)\langle 1|g_1^{\Lambda\Lambda}\rangle\langle 1|X_1|\phi_0\rangle,
\end{aligned}$$

$$\begin{aligned}
\langle 1|Y_1|\phi_0\rangle &= 2\tau_1^{NN}\langle g_1^{NN}|1\rangle\langle 1|3\rangle G_0(3)\langle 3|g_3^{N\Xi}\rangle\langle 3|Y_3|\phi_0\rangle, \\
\langle 3|Y_3|\phi_0\rangle &= 2\tau_3^{N\Xi}\langle g_3^{N\Xi}|\phi_0\rangle \\
&\quad -\tau_3^{N\Xi}\langle g_3^{N\Xi}|3\rangle\langle 2|3\rangle G_0(3)\langle 3|g_3^{N\Xi}\rangle\langle 3|Y_3|\phi_0\rangle \\
&\quad +\tau_3^{N\Xi}\langle g_3^{N\Xi}|3\rangle\langle 3|1\rangle G_0(1)\langle 1|g_1^{NN}\rangle\langle 1|Y_1|\phi_0\rangle \\
&\quad +2\tau_{31}^{N\Xi-\Lambda\Lambda}\langle g_1^{\Lambda\Lambda}|1\rangle\langle 1|3\rangle G_0(3)\langle 3|g_3^{N\Lambda}\rangle\langle 3|X_3|\phi_0\rangle.
\end{aligned} \tag{11}$$

If one neglects the inhomogeneous terms in Eqs. (10) and (11) they become identical to Eqs. (14) and (15) of Ref. [20]. The Ξd scattering amplitude normalized as in the Argand diagram is given by

$$F = -\pi q_0 \nu \langle \phi_0 | U_3 | \phi_0 \rangle, \tag{12}$$

where q_0 and ν are, respectively, the Ξd on-shell momentum and reduced mass. We solved the integral equations (11) using the standard method [24], where the momentum variables are rotated into the complex plane as $q_i \rightarrow q_i e^{-i\phi}$ and checked that the results are independent of the rotation angle ϕ . If the resonance lies below the Ξd threshold, as it was the case in Ref. [20], the contour rotation method allows to take simultaneously both the momentum variables and the energy variable as complex so that one can determine the position of the pole in the complex plane. However, if the resonance lies above the Ξd threshold, like in the present case, the contour rotation method works only if one takes the momentum variables complex but leaves the energy variable real so that one can not determine the position of the pole in the complex plane.

We show in Fig. 2 the Argand diagram of the Ξd system between 0 and 10 MeV above the Ξd threshold where one sees the typical counterclockwise behavior of a resonant amplitude. If one neglects the $(i, j^P) = (0, 0^+)$ channel, the counterclockwise behavior disappears which shows that the H -dibaryon channel $(i, j^P) = (0, 0^+)$ is basic for the existence of the three-body $\Lambda\Lambda N - \Xi NN$ S wave resonance.

As already mentioned in the introduction, the HAL QCD ΞN interactions have been recently used to study the possible existence of ΞNN bound states in Ref. [12] with negative results for the $(I, J^P) = (1/2, 1/2^+)$ channel, in agreement with our findings in spite of using a different NN interaction and a different method.

We have finally evaluated the Ξd elastic cross section as a function of energy where we have included not only the $(I, J^P) = (1/2, 1/2^+)$ Ξd amplitude but also the $(1/2, 3/2^+)$ amplitude, which is very small. The result is shown in Fig. 3. As one can see, the resonance




FIG. 2: Argand diagram of the Ξd system between 0 and 10 MeV above the Ξd threshold. Some relevant energies, in MeV, are indicated.

shows up as a change of slope of the cross section at an energy around 2.3 MeV, i.e., close to the Ξd breakup threshold $\sqrt{S} = 2m_N + m_{\Xi}$. The bump in the cross section would become larger for a stronger $(i, j^p) = (0, 0^+)$ transition potential, and, as said above, it would disappear if the $(i, j^p) = (0, 0^+)$ channel is not considered or the two-body resonance in the H -dibaryon channel would not exist. The Ξd cross section would allow to discriminate among the different models for the strangeness -2 two-baryon interactions. It could be studied




FIG. 3: Ξd elastic cross section.

through the quasifree Ξ^- production in the (K^-, K^+) reaction on a deuteron target [25, 26].

Let us finally note that if one drops the coupling to the $\Lambda\Lambda N$ channel Fig. 2 changes by about 10 % while keeping its shape rotating slightly to the right; similarly, in Fig. 3 the cross section at $E = 2.3$ MeV changes from 55 mb to 58 mb.

It is interesting to compare this resonance with the nucleon-nucleon 1D_2 Hoshizaki resonance [27] which has a mass close to $\sqrt{S} = m_N + m_\Delta$ since it arises due to the process $NN \rightarrow \pi NN$ which is driven by the pion-nucleon Δ resonance [28]. The resonance we are studying here is driven by the $\Lambda\Lambda - \Xi N$ H -dibaryon resonance which appears either just below or just above the $N\Xi$ threshold [9] so that it has a mass $m_H = m_N + m_\Xi$. Following the comparison with the Hoshizaki state one expects that the $\Lambda\Lambda N - \Xi NN$ resonance will have a mass close to $\sqrt{S} = m_N + m_H = 2m_N + m_\Xi$ which is precisely the Ξd threshold in agreement with Figs. 2 and 3.

In brief, we have shown that the possible existence of a $\Lambda\Lambda N - \Xi NN$ resonance would be highly sensitive to the $\Lambda\Lambda - \Xi N$ interaction. In particular, by using a separable potential based on the most recent results of the HAL QCD Collaboration, characterized by the existence of a resonance just below or above the ΞN threshold in the so-called H -dibaryon channel, $(i, j^p) = (0, 0^+)$, a three-body resonance appears 2.3 MeV above the Ξd threshold. A theoretical and experimental effort to constrain the $\Lambda\Lambda - \Xi N$ interaction is a basic ingredient to progress in our knowledge of the strangeness -2 sector.

Acknowledgments

This work has been partially funded by COFAA-IPN (México) and by Ministerio de Ciencia e Innovación and EU FEDER under Contracts No. FPA2016-77177-C2-2-P, PID2019-105439GB-C22 and RED2018-102572-T.

-
- [1] H. Garcilazo and A. Valcarce, Phys. Rev. Lett. **110**, 012503 (2013); A. Gal, Phys. Rev. Lett. **110**, 179201 (2013); H. Garcilazo and A. Valcarce, Phys. Rev. Lett. **110**, 179202 (2013).
 - [2] H. Garcilazo, A. Valcarce, and T. Fernández-Caramés, J. Phys. G **41**, 095103 (2014).
 - [3] H. Garcilazo, A. Valcarce, and T. Fernández-Caramés, J. Phys. G **42**, 025103 (2015).
 - [4] R. L. Jaffe, Phys. Rev. Lett. **38**, 195 (1977); **38**, 617(E) (1977).

- [5] H. Takahashi *et al.*, Phys. Rev. Lett. **87**, 212502 (2001).
- [6] A. Francis, J. R. Green, P. M. Junnarkar, Ch. Miao, T. D. Rae, and H. Wittig, Phys. Rev. D. **99**, 074505 (2019).
- [7] A. Gal, E. V. Hungerford, and D. J. Millener, Rev. Mod. Phys. **88**, 035004 (2016).
- [8] B. H. Kim *et al.* [Belle Collaboration], Phys. Rev. Lett. **110**, 222002 (2013).
- [9] K. Sasaki, S. Aoki, T. Doi, S. Gongyo, T. Hatsuda, Y. Ikeda, T. Inoue, T. Iritani, N. Ishii, K. Murano, and T. Miyamoto [HAL QCD Collaboration], Nucl. Phys. A **998**, 121737 (2020).
- [10] Y. Yamaguchi and T. Hyodo, Phys. Rev. C **94**, 065207 (2016).
- [11] K. -W. Li, T. Hyodo, and L. -S. Geng, Phys. Rev. C **98**, 065203 (2018).
- [12] E. Hiyama, K. Sasaki, T. Miyamoto, T. Doi, T. Hatsuda, Y. Yamamoto, and Th. A. Rijken, Phys. Rev. Lett. **124**, 092501 (2020).
- [13] R. B. Wiringa, R. A. Smith, and T. L. Ainsworth, Phys. Rev. C **29**, 1207 (1984).
- [14] E. Hiyama, Y. Kino, and M. Kamimura, Prog. Theor. Nucl. Phys. **51**, 223 (2003).
- [15] E. Hiyama, PTEP **2012**, 01A204 (2012).
- [16] M. M. Nagels, Th. A. Rijken, and Y. Yamamoto, Phys. Rev. C **99**, 044002 (2019).
- [17] M. M. Nagels, Th. A. Rijken, and Y. Yamamoto, Phys. Rev. C **99**, 044003 (2019).
- [18] M. M. Nagels, Th. A. Rijken, and Y. Yamamoto, arXiv:1504.02634.
- [19] H. Garcilazo and A. Valcarce, Phys. Rev. C **93**, 064003 (2016).
- [20] H. Garcilazo, Phys. Rev. C **93**, 024001 (2016).
- [21] I. Filikhin, V. M. Suslov, and B. Vlahovic, Math. Model. Geom. **5**, 1 (2017).
- [22] Y. Yamaguchi, Phys. Rev. **95**, 1628 (1954).
- [23] S. B. Carr, I. R. Afnan, and B. F. Gibson, Nucl. Phys. A **625**, 143 (1997).
- [24] B. C. Pearce and I. R. Afnan, Phys. Rev. C **30**, 2022 (1984).
- [25] T. Tamagawa *et al.*, Nucl. Phys. A **691**, 234c (2001).
- [26] Y. Yamamoto, T. Tamagawa, T. Fukuda, and T. Motoba, Prog. Theor. Phys. **106**, 363 (2001).
- [27] N. Hoshizaki, Phys. Rev. C **45**, R1424 (1992).
- [28] A. Gal and H. Garcilazo, Nucl. Phys. A **928**, (2014) 73.