

A Robust Adaptive Modified Maximum Likelihood Estimator for the Linear Regression Model

S. Acitas^a; Peter Filzmoser^b Birdal Senoglu^c

^aEskisehir Technical University, Department of Statistics, Eskisehir, Turkey; ^bVienna University of Technology, Vienna, Austria; ^cAnkara University, Department of Statistics, Ankara, Turkey

ARTICLE HISTORY

Compiled February 26, 2022

ABSTRACT

In linear regression, the least squares (LS) estimator has certain optimality properties if the errors are normally distributed. This assumption is often violated in practice, partly caused by data outliers. Robust estimators can cope with this situation and thus they are widely used in practice. One example of robust estimators for regression are adaptive modified maximum likelihood (AMML) estimators [1]. However, they are not robust to x outliers, so-called leverage points. In this study, we propose a new regression estimator by employing an appropriate weighting scheme in the AMML estimation method. The resulting estimator is called robust AMML (RAMML) since it is not only robust to y outliers but also to x outliers. A simulation study is carried out to compare the performance of the RAMML estimator with some existing robust estimators such as MM, least trimmed squares (LTS) and S. The results show that the RAMML estimator is preferable in most settings according to the mean squared error (MSE) criterion. Two data sets taken from the literature are also analyzed to show the implementation of the RAMML estimation methodology.

KEYWORDS

Adaptive modified likelihood; efficiency; leverage point; regression; robustness

1. Introduction

Regression analysis is a widely used statistical tool to analyze the relationships between a dependent variable and predictors. The least squares (LS) method is usually utilized to estimate the unknown model parameters. The LS estimator is popular because of its simple computation and good statistical properties, i.e. it is the best linear unbiased estimator under the normality assumption.

The LS estimator loses its efficiency when the normality assumption is not satisfied. Therefore, some alternative methods are employed to accommodate the problems arising in case of non-normality which is caused by outliers. Indeed, there are two types of outliers in the context of regression analysis: (i) vertical outliers, and (ii) leverage points. These outliers refer to outlyingness in the space of the response and the predictors, respectively. The LS estimator is sensitive to both types of outliers, and bad leverage points can even distort the estimator and lead to a non-sense model

[2]. For this reason, many robust estimators which reduce the effects of outliers have been proposed in the literature. For example, the M estimator [3], the S estimator [4], the least trimmed squares (LTS) and the least median of squares (LMS) of Rousseeuw [5], and the MM estimator [6] are well-known and widely used in the context of robust regression. We refer to Maronna et al. [7] or Gschwandtner and Filzmoser [8] in which descriptions of these estimators and related literature information are briefly given. See also Yu and Yao [9] for an extensive review of robust linear regression estimators.

In the literature, the modified maximum likelihood (MML) estimator is also used to reduce the effect of outlying observations. As its name indicates, the MML estimator is obtained applying a particular modification of the maximum likelihood (ML) method [10,11]. This modification allows to obtain explicit forms of the estimator under the assumption of a non-normal error distribution. A long-tailed symmetric distribution is frequently used in this context since it is useful for modelling outliers occurring in the direction of long tails [12]. For example, Tiku et al. [13] obtain the MML estimator of the unknown parameters of the simple linear regression model when the distribution of the error terms is long-tailed symmetric. The MML estimator is also obtained for the multiple linear regression model [14] with long-tailed symmetric error distributions. The MML estimator has good properties, and it is easy to compute. Furthermore, it is not only robust to outliers but also asymptotically equivalent to the ML estimator. However, it has two drawbacks: (i) the shape parameter of the long-tailed symmetric distribution is assumed to be known and (ii) it is only robust to vertical outliers but not to leverage points.

The first drawback is not a new issue in the related literature. For example, Lucas [15] consider the Student's t distribution as an alternative to normal distribution for estimating the location parameter under the assumption of known degrees of freedom ν . This is because of the fact that the ML estimator of the other parameters is no longer robust when it is estimated jointly with ν . Therefore, it is treated as a robustness tuning constant—see also Arslan and Genc [16,17], Acitas et al. [18,19] in which similar discussions are available. Tiku and Surucu [20] also consider this issue and propose a new version of the MML estimator. This version is then called adaptive MML (AMML) by Donmez [1]. In AMML, the main idea of the MML methodology is preserved but the assumption of a known shape parameter on the estimation process is weakened.

To the best of our knowledge, the second drawback is still an open problem. In other words, there is no previous study on the MML and AMML estimators dealing with leverage points as far as we know. Therefore, the motivation of this study is to suggest a new regression estimator which is robust to both types of outliers based on the AMML estimators. This estimator is called robust AMML (RAMML) and obtained by introducing new weights assigned to the predictors in the AMML estimation process [21]. The RAMML is not only robust to both types of outliers but also efficient under the normality assumption. A further advantage of the RAMML estimator is that it is easy to compute since the explicit form of the estimator is available.

The rest of the paper is organized as follows. Section 2 reviews the AMML methodology. Section 3 is reserved to the RAMML estimator. The results of a Monte-Carlo simulation study are given in Section 4. The paper is finalized with some concluding remarks.

2. Review of the AMML estimator

We consider a multiple linear regression model with n observations and m predictors,

$$y_i = \beta_0 + \mathbf{x}_i' \boldsymbol{\beta} + \varepsilon_i, \quad i = 1, 2, \dots, n \quad (1)$$

where y_i is the dependent variable, β_0 is the intercept, $\boldsymbol{\beta} = (\beta_1 \ \beta_2 \ \dots \ \beta_m)'$ is the vector of unknown regression coefficients, $\mathbf{x}_i$ is the vector of predictors defined by $\mathbf{x}_i = (x_{i1} \ x_{i2} \ \dots \ x_{im})'$ and ε_i is the random error term.

Assume that the distribution of the error terms in model (1) is long-tailed symmetric with the probability density function (pdf) given by

$$f(e; p, \sigma) = \frac{1}{\sqrt{q} B(0.5, p-0.5) \sigma} \left(1 + \frac{e^2}{q\sigma^2} \right)^{-p}, \quad q = 2p-3, \quad -\infty < e < \infty \quad (2)$$

where p is the shape parameter which is assumed to be $p \geq 2$, σ is the scale parameter and $B(\cdot, \cdot)$ is the beta function. The kurtosis value of the long-tailed distribution is greater than 3. It tends to normal distribution as p tends to ∞ . Therefore, it has been considered as a good alternative to the normal distribution [12].

The ML estimators of the model parameters are the solutions of the following likelihood equations:

$$\frac{\partial \log L}{\partial \beta_0} = \frac{2p}{q\sigma} \sum_{i=1}^n g(z_i) = 0, \quad (3)$$

$$\frac{\partial \log L}{\partial \beta_j} = \frac{2p}{q\sigma} \sum_{i=1}^n g(z_i) x_{ij} = 0, \quad j = 1, 2, \dots, m \quad (4)$$

$$\frac{\partial \log L}{\partial \sigma} = -\frac{n}{\sigma} + \frac{2p}{q\sigma} \sum_{i=1}^n g(z_i) z_i = 0, \quad (5)$$

where

$$g(z_i) = \frac{z_i}{1 + \frac{z_i^2}{q}}, \quad z_i = \frac{y_i - \beta_0 - \mathbf{x}_i' \boldsymbol{\beta}}{\sigma}, \quad i = 1, 2, \dots, n.$$

Since $g(\cdot)$ is a nonlinear function of the unknown model parameters, the explicit solutions of equations (3)-(5) cannot be obtained, and numerical methods need to be used instead. However, using numerical methods has some drawbacks such as wrong convergency, multiple roots and non-convergence of iterations, see e.g. Puthenpura and Sinha [22], Vaughan [23]. The modified maximum likelihood (MML) methodology proposed in Tiku [10,11] overcomes these problems and allows to obtain the explicit forms of the estimators. The idea underlying the MML methodology is linearization of the intractable term (i.e., the function $g(\cdot)$) in the likelihood equations. This is done using the first two terms of a Taylor series expansion. For more details we refer to Islam and Tiku [14] in which MML estimators for the multiple linear regression model are obtained under a long-tailed symmetric error distribution assumption.

The MML estimators are explicitly formulated, thus no computational efforts are

required. They have also good statistical properties, i.e. they are asymptotically equivalent to the ML estimators and therefore they are minimum variance bound estimators [24]. However, the MML methodology works under the assumption of a known shape parameter p . In some studies, p is identified using the profile likelihood method, see for example Acitas et al. [18,19]. Tiku and Surucu [20] suggested a new version of the MML method in which the known shape parameter assumption is weakened. They estimate the location and the scale parameter using this new version and obtain estimators which are as good as or better than M-estimators. Donmez [1] generalized this idea to the simple and multiple linear regression model and calls the method AMML methodology. In the following, the steps of the AMML are explained based on Donmez [1]:

Step 1. The function $g(\cdot)$ is linearized around

$$t_i = \frac{y_i - \beta_0 - \mathbf{x}_i' \boldsymbol{\beta}}{\sigma}, \quad i = 1, 2, \dots, n \quad (6)$$

using the first two terms of a Taylor series expansion,

$$g(z_i) \cong \alpha_i + \delta_i z_i, \quad i = 1, 2, \dots, n \quad (7)$$

where

$$\alpha_i = \frac{(1/q)t_i}{(1 + (1/q)t_i^2)^2}, \quad \delta_i = \frac{1}{(1 + (1/q)t_i^2)^2}, \quad i = 1, 2, \dots, n. \quad (8)$$

Step 2. The linearized version of the function $g(z_{(i)})$ is incorporated into the likelihood equation and the so-called modified likelihood equations are obtained as follows:

$$\frac{\partial \log L^*}{\partial \beta_0} = \frac{2p}{q\sigma} \sum_{i=1}^n (\alpha_i + \delta_i z_i) = 0 \quad (9)$$

$$\frac{\partial \log L^*}{\partial \beta_j} = \frac{2p}{q\sigma} \sum_{i=1}^n (\alpha_i + \delta_i z_i) x_{ij} = 0, \quad j = 1, 2, \dots, m \quad (10)$$

$$\frac{\partial \log L^*}{\partial \sigma} = -\frac{n}{\sigma} + \frac{2p}{q\sigma} \sum_{i=1}^n (\alpha_i + \delta_i z_i) z_i = 0, \quad (11)$$

where $\log L^*$ stands for the modified log-likelihood function.

Step 3. The solutions of the Equations (9)-(11) are the AMML estimators. They are formulated as follows:

$$\hat{\beta}_0 = \bar{y}_{[\cdot]} - \bar{\mathbf{x}}'_{[\cdot]} \hat{\boldsymbol{\beta}} \quad (12)$$

$$\hat{\boldsymbol{\beta}} = \mathbf{K} + \mathbf{L} \hat{\sigma}, \quad (13)$$

$$\hat{\sigma} = \frac{B + \sqrt{B^2 + 4nC}}{2\sqrt{n(n-m-1)}} \quad (14)$$

where

$$\begin{aligned}\mathbf{K} &= (\mathbf{X}'\boldsymbol{\delta}\mathbf{X})^{-1}(\mathbf{X}'\boldsymbol{\delta}\mathbf{y}), \quad \mathbf{L} = (\mathbf{X}'\boldsymbol{\delta}\mathbf{X})^{-1}(\mathbf{X}'\boldsymbol{\alpha}\mathbf{1}) \\ B &= \frac{2p}{q}(\mathbf{y} - \mathbf{X}\mathbf{K})'\boldsymbol{\alpha}\mathbf{1}, \quad C = \frac{2p}{q}(\mathbf{y} - \mathbf{X}\mathbf{K})'\boldsymbol{\delta}(\mathbf{y} - \mathbf{X}\mathbf{K})\end{aligned}$$

$$\bar{y}_{[\cdot]} = \frac{\sum_{i=1}^n \delta_i y_i}{\sum_{i=1}^n \delta_i}, \quad \bar{\mathbf{x}}_{[\cdot]j} = \frac{\sum_{i=1}^n \delta_i x_{ij}}{\sum_{i=1}^n \delta_i}, \quad \bar{\mathbf{x}}'_{[\cdot]} = [\bar{\mathbf{x}}_{[\cdot]1} \quad \bar{\mathbf{x}}_{[\cdot]2} \quad \cdots \quad \bar{\mathbf{x}}_{[\cdot]m}],$$

$$\boldsymbol{\delta} = \text{diag}(\delta_i) \quad \text{and} \quad \boldsymbol{\alpha} = \text{diag}(\alpha_i).$$

Here, $\mathbf{X}$ is the $n \times m$ matrix of predictors, and $\mathbf{1}$ stands for the vector of ones with n entries.

It is clear that AMML estimators are also explicitly formulated. They preserve the properties of the MML estimators [1]. It can easily be shown that the AMML estimators are regression, scale and affine equivariant. The proof is not given here for the sake brevity, but it can be provided upon request.

As it is clear, the values t_i should be computed at the beginning of the methodology using some initial estimates. The computational details are explained in the following subsection.

2.1. Computation of the AMML estimators

The computation of the AMML estimators is obtained in two iterations. An initial estimate of t_i is required in the first iteration. The initial estimate of t_i , proposed by [1], is obtained based on a reparametrization of the model (1). The reparametrized model is constructed assuming that all β_j coefficients are equal in model (1). This is because of the fact that there is no reason to believe that one predictor is more important than others at the beginning. The reparametrized model is formulated by

$$y_i = \beta_0 + \theta v_i + \varepsilon_i, \quad v_i = \sum_{j=1}^m x_{ij}. \quad i = 1, 2, \dots, n. \quad (15)$$

Then, the initial estimate of t_i is given as

$$\tilde{t}_i = \frac{y_i - T_0 - T_1 v_i}{S_0} \quad (16)$$

where the initial estimators of β_0 , θ and σ are given by

$$T_0 = \text{median}\{y_i - T_1 v_i\}, \quad T_1 = \text{median}\left\{\frac{y_{\ell+1} - y_\ell}{v_{\ell+1} - v_\ell}\right\} \quad (17)$$

and

$$S_0 = 1.483 \cdot \text{median}\{|y_i - T_0 - T_1 v_i|\}, i = 1, 2, \dots, n, \quad (18)$$

respectively. After getting the initial estimate of t_i , initial AMML estimators denoted by $\tilde{\beta}_0$, $\tilde{\beta}_j$ and $\tilde{\sigma}$ are obtained following steps 1-3 given in Section 2.

In the second iteration, t_i is updated using the initial AMML estimators ($\tilde{\beta}_0$, $\tilde{\beta}_j$ and $\tilde{\sigma}$) as follows:

$$t_i = \frac{y_i - \tilde{\beta}_0 - \sum_{j=1}^n x_{ij} \tilde{\beta}_j}{\tilde{\sigma}} \quad (19)$$

and steps 1-3 are utilized one more time. The final AMML estimators are obtained at the end of step 3 using Equations (12)-(14).

It is clear that the shape parameter of the long-tailed symmetric distribution p appears in the formulas. It is considered as robustness tuning constant and taken to be 16.5 so that the resulting AMML estimators are efficient when the distribution of the error terms is normal [1,20].

Using different simulation schemes, Donmez [1] shows that AMML estimators are robust to y -outliers. On the other hand, there are no available results whether they are resistant to x -outliers. Indeed, the simulation schemes used in Donmez [1] do not include any x -outliers. It should also be mentioned that Donmez [1] point out to use standardized versions of predictors to reduce the effect of x -outliers. However, the standardization is done using mean and standard deviation which are not robust to outliers. Standardization is indeed required when working with the reparametrized model (see Equation (15)) in which the sum of the predictors is used. If the predictors are measured under different scales and/or units, this sum is not remarkable. In our proposed methodology, we do not need any standardization for the predictors and the reparametrized model. Therefore, these AMML estimators are not considered in the rest of the paper.

3. Robust AMML estimators

In this section, we propose RAMML estimators which are robust to x -outliers unlike the AMML. This is done by downweighting the x -outliers using extra weights for predictors. The steps for the RAMML methodology are similar to those given for AMML, i.e. step 1 is the same while steps 2 and 3 have to be revised. New versions of step 2 and 3 are shown by Step 2* and 3*, respectively, and explained below.

Step 2*. Consider the modified likelihood Equations (9)-(11). We introduce new weights, denoted by δ_i^x , for the predictors to accommodate the effects of x -

outliers in these equations:

$$\frac{\partial \log L^*}{\partial \beta_0} = \frac{2p}{q\sigma} \sum_{i=1}^n \delta_i^x (\alpha_i + \delta_i z_i) = 0 \quad (20)$$

$$\frac{\partial \log L^*}{\partial \beta_j} = \frac{2p}{q\sigma} \sum_{i=1}^n \delta_i^x (\alpha_i + \delta_i z_i) x_{ij} = 0, \quad j = 1, 2, \dots, m \quad (21)$$

$$\frac{\partial \log L^*}{\partial \sigma} = -\frac{n}{\sigma} + \frac{2p}{q\sigma} \sum_{i=1}^n \delta_i^x (\alpha_i + \delta_i z_i) z_i = 0, \quad (22)$$

where

$$\delta_i^x = \frac{1}{(1 + (1/q)\tilde{x}_i^2)^4} \quad \text{and} \quad \tilde{x}_i = \frac{\|\mathbf{x}_i - \text{med}_{L_1}(\mathbf{X})\|}{\text{median}_i \|\mathbf{x}_i - \text{med}_{L_1}(\mathbf{X})\|}, \quad i = 1, 2, \dots, n. \quad (23)$$

Here, $\text{med}_{L_1}(\cdot)$ denotes the L_1 -median which is also known as geometric median or spatial median. It is a highly robust estimator for the multivariate location, i.e. its breakdown point is 0.5 [25]. We refer to Croux et al. [26] and Fritz et. al [27] in which detailed information is available about L_1 -median.

Step 3*. The RAMML estimators are obtained as solutions of Equations (20)–(22). They are formulated as given in Equations (12)–(14). However, there are some modifications due to the new weighting scheme. They are given as follows:

$$\hat{\beta}_0 = \bar{y}_{[\cdot]} - \bar{\mathbf{x}}'_{[\cdot]} \hat{\boldsymbol{\beta}} + \frac{\Delta}{w} \hat{\sigma} \quad (24)$$

where

$$\bar{y}_{[\cdot]} = \frac{\sum_{i=1}^n \delta_i \delta_i^x y_i}{w}, \quad \bar{\mathbf{x}}_{[\cdot]j} = \frac{\sum_{i=1}^n \delta_i \delta_i^x x_{ij}}{w}, \quad \bar{\mathbf{x}}'_{[\cdot]} = [\bar{\mathbf{x}}_{[\cdot]1} \quad \bar{\mathbf{x}}_{[\cdot]2} \quad \cdots \quad \bar{\mathbf{x}}_{[\cdot]m}],$$

$$w = \sum_{i=1}^n \delta_i \delta_i^x, \quad \Delta = \sum_{i=1}^n \alpha_i \delta_i^x.$$

$$\boldsymbol{\delta} = \text{diag}(\delta_i \delta_i^x) \quad \text{and} \quad \boldsymbol{\alpha} = \text{diag}(\alpha_i \delta_i^x), \quad i = 1, 2, \dots, n.$$

The remaining expressions, i.e. $\hat{\boldsymbol{\beta}}$, $\hat{\sigma}$, $\mathbf{K}$, $\mathbf{L}$, B and C are formulated in the same way as given in the previous section, but using the weights and formulations as introduced above.

It should be noticed that RAMML estimators are defined similar to generalized M (GM) estimators which are robust to both types of outliers, see e.g. Maronna et al. [7] for further details.

3.1. Computation of RAMML estimators

The RAMML estimators are obtained after two iterations. In the first iteration, an initial estimate of t_i is required. Unlike Donmez [1], we do not consider a reparametrization of model (1). We continue with model (1) and suggest to use well-known robust regression estimators such as LTS or S to initialize the iterations. In other words, the initial estimate of t_i is proposed as

$$t_i = \frac{y_i - T_0 - \sum_{j=1}^n x_{ij}T_j}{S_0}, \quad (25)$$

where T_0 , T_j ($j = 1, 2, \dots, m$) and S_0 are obtained from LTS or S regression. After getting the initial estimate of t_i , initial RAMML estimators denoted by $\tilde{\beta}_0$, $\tilde{\beta}_j$ and $\tilde{\sigma}$ are obtained following Steps 1, 2* and 3*.

In the second iteration, t_i is updated with the initial RAMML estimators obtained in the previous iteration as follows:

$$t_i = \frac{y_i - \tilde{\beta}_0 - \sum_{j=1}^n x_{ij}\tilde{\beta}_j}{\tilde{\sigma}} \quad (26)$$

and Steps 1, 2* and 3* are utilized. The final RAMML estimators are obtained at the end of Step 3* using Equations (13), (14) and (24).

The resulting estimators obtained using LTS and S regression are denoted by RAMML₁ and RAMML₂, respectively. It is clear that RAMML estimators are also explicitly formulated and they can be computed after two simple iterations. Therefore, they are computationally straightforward.

If the weights δ_i^x are removed from the estimating equations, RAMML₁ and RAMML₂ will reduce to the AMML₁ and AMML₂, respectively. It should be noted that these AMML₁ and AMML₂ estimators differ from that of Donmez (2010) since the initial t_i s are defined differently.

It should be noted that different initial estimates can be preferred for finding initial t_i s. The important issue here is that the initial estimators should be robust so that the resulting estimator is robust. In this study, we consider two well-known robust estimators as initials, i.e. LTS and S. We also consider the LMS estimator as an initial estimator and obtain more or less the same results, see Acitas et al. [21]. However, we do not consider it here because it is not computationally efficient.

4. Simulation study

In this section, we conduct a simulation study to compare the performances of the proposed estimators RAMML₁ and RAMML₂ along with some known robust estimators such as MM, LTS and S estimators under different simulation scenarios. AMML₁ and AMML₂ are also taken into account during the performance comparisons.

The simulation setup and contamination schemes are considered similar to those used by Gschwandtner and Filzmoser [8]. Model (1) is used during the simulation study where β_0 is assumed to be 0 without loss of generality. The data are generated

as follows:

$$\begin{aligned} \mathbf{x}_i &\sim \mathbf{N}_m(\mathbf{0}, \mathbf{I}_m), \quad i = 1, 2, \dots, n \\ \beta_j &= 1/\sqrt{m}, \quad j = 1, 2, \dots, m \quad \text{and thus} \quad \|\beta\| = 1. \end{aligned}$$

Here, $\mathbf{N}_m$ denotes the m -dimensional multivariate normal distribution, $\mathbf{0}$ stands for the vector of zeros and $\mathbf{I}_m$ is the identity matrix.

Five different error distributions are considered in model (1): (i) $N(0, 1)$, (ii) Laplace, (iii) t_5 , (iv) t_1 (Cauchy) and (v) slash distribution. The leverage points are generated replacing the first n_{out} observations with the following versions:

$$\mathbf{x}_i = [\ell \quad \ell \quad \dots \quad \ell]' \quad \text{and} \quad y_i = \mathbf{x}_i' \mathbf{a} \quad \text{for} \quad i = 1, 2, \dots, n$$

where $\mathbf{a}$ is a normalized unit vector defined by $\mathbf{a} = \boldsymbol{\nu} - (\boldsymbol{\nu}'\beta)\beta$. Here, $\boldsymbol{\nu}$ is a vector having entries $(-1)^j$. If $m = 1$, a is taken as -1. We consider two different versions for leverage points, i.e. $\ell = 5$ and $\ell = 10$. This is because of the fact that we would like to see how the performances of the estimators are affected as the magnitude of the leverage point increases. The sample size is taken to be $n = 50$ and 200 . The number of the predictors m taken from the set $\{1, 5, 10, 20\}$. n_{out} is the number of leverage points and determined according to two contamination levels, i.e. 10% and 20%.

The mean squared error (MSE) formulated as

$$MSE = \frac{1}{n_{rep}} \sum_{r=1}^{n_{rep}} \left\| \hat{\beta}^r - \beta \right\|^2 \quad (27)$$

is calculated to evaluate the performances of the estimators based on $n_{rep} = 500$ Monte-Carlo runs. All computations are carried out using the software environment R. The results are tabulated in Tables 1 and 2. It should be mentioned that LS estimators are not considered in these tables since they are not robust to outliers.

[Table 1 is here]

It can be seen from Table 1 that the MSEs of the RAMML₁ and RAMML₂ are close to each other and smaller than those of MM, LTS, AMML₁, S and AMML₂ for both 10% and 20% contamination levels when the error distribution is normal, Laplace, t_5 , t_1 and slash. The RAMML₂ estimator is almost the best since it has the smallest MSE. However, the RAMML₁ also performs better than the others in some cases, i.e. if the contamination level is 10%, $m = 1, 5$ and 10 , when the error distribution is normal. It is also the best estimator for simple linear regression with normal, Laplace and t_5 error distributions when the contamination level is 20%. In the remaining cases, the RAMML₂ is preferable to other estimators. It should also be mentioned that the AMML₂ performs better than MM, LTS, AMML₁, RAMML₁, S and RAMML₂ when the number of predictors is 20, $\ell = 5$ and the contamination level is 20%. The MM estimator has a remarkable performance but it is not sufficient to be the best. In short, the RAMML₂ is slightly better than its competitors in most cases considered in Table 1.

[Table 2 is here]

When the sample size is increased to 200, see Table 2, we can draw similar conclusions to those obtained from Table 1. However, here the MM estimator gains efficiency

and performs better than the other estimators for some cases. For example, for simple linear regression with normal, Laplace and t_5 error distributions for 10% level of contamination and $\ell = 5$, it has the minimum MSE value, followed by the RAMML₂. If the distribution of the error is heavy-tailed, i.e. t_1 and slash, the RAMML₂ is the best. This conclusion is also true for $\ell = 5$. If m is increased from 5 to 20, the RAMML₂ is mostly the best estimator when the contamination level is 10%. In some cases such as $\ell = 5$, $m = 10$, t_1 error distribution and $\ell = 10$, $m = 20$, normal distribution, the RAMML₁ outperforms the other estimators. When the level of contamination is 20%, the RAMML₁ and RAMML₂ outperform the others. For the simple linear regression case with all error distributions, the RAMML₂ is better than MM, LTS, AMML₁, RAMML₁, S and AMML₂. For $m \in \{5, 10, 20\}$, the RAMML₁ is more preferable for normal, Laplace and t_5 error distributions. However, if the error distribution is heavy-tailed, RAMML₂ is the best regardless of the value of ℓ and contamination level. It can be concluded that RAMML₂ is the best estimator for almost all cases considered in Table 2.

In this part of the simulation study, we just focus on the MSEs of the estimators for the scale parameter σ when the distribution of the error terms is $N(0, 1)$. The MSE of $\hat{\sigma}$ is calculated using (27) in which the symbols β are replaced by σ . The results are tabulated in Table 3. The MSEs of RAMML₁ and RAMML₂ are more or less the same. RAMML₁ and/or RAMML₂ has the minimum MSE value in most of the cases.

[Table 3 is here]

It is well-known that the trade-off between robustness and efficiency is important. We therefore explore the efficiencies of RAMML₁ and RAMML₂ under normality in this part of the simulation study. For this purpose, the performances of all estimators considered in this study are compared with the LS estimators when the error distribution is $N(0, 1)$ with zero contamination. The results are given in Table 4. It is clear from this table that the MSE values of RAMML₁ and RAMML₂ are close to those of LS. This conclusion implies that RAMML₁ and RAMML₂ are not only robust to both types of outliers but also efficient under normality.

[Table 4 is here]

Finally, we explore the performances of the estimators when the predictors are correlated. Therefore, we make a simple modification in the simulation scenario given at the beginning of this section. Here, the $\mathbf{x}_i$ s are generated from $\mathbf{N}_m(\mathbf{0}, \mathbf{V})$ where $\mathbf{V} = [v_{jk}]$ with $v_{jj} = 1$ and $v_{jk} = \rho$ for $j \neq k$ and $k = 1, 2, \dots, m$. In Table 5, we report the MSEs of the estimators for $\rho = 0.90$, $\ell = 10$ and contamination levels 10%. This table shows that the MM estimator is the best one in almost all settings, mostly followed by RAMML₁ and RAMML₂.

Overall, the MSEs of RAMML₁ and RAMML₂ are close to each other and they outperform their competitors when there are x -outliers. Among these two estimators, we see that RAMML₂ is preferable.

[Table 5 is here]

5. Application

In this part of the study, we provide two data sets which are widely used in the literature to show the implementation of the proposed estimators. The full data sets

can be found in the R package “robustbase”.

5.1. *Hertzsprung-Russell Diagram Data of Star Cluster CYG OB1*

This data set is shortly denoted as starsCYC and includes 47 observations with two variables named $\log.Te$ (x) and $\log.light$ (y). The former variable is the logarithm of the effective temperature at the surface of the star while the latter one is the logarithm of its light intensity, see Rousseeuw and Leroy [2]. It is clear from Figure 1 that the starsCYC data contains outliers, i.e. four stars in the left upper corner are far away from bulk of the data, and thus they are leverage points. Rousseeuw and Leroy [2] employ the LMS method and obtain more reliable results than for the LS method. Different from them, in this study, we use the RAMML₁ and RAMML₂ estimation methods along with the other methods considered in the previous section to estimate the regression coefficients. To evaluate the performances of the methods, we use the standard error of prediction (SEP) criterion formulated as

$$SEP = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (y_i - \hat{y}_i - \text{bias})^2}, \quad \text{bias} = \frac{1}{n} \sum_{i=1}^n (y_i - \hat{y}_i)$$

where, $\hat{y}_i$ denotes the fitted value for the i -th observation. It should be noted that using SEP may give misleading results in case of outliers since each observation has the same contribution in the formula of SEP. Therefore, a robust version of the SEP is required. We therefore compute the trimmed version of SEP where the trimming value is 10%. The robust performance criterion will lead to a contrary picture since the influence of the largest contributions is reduced. There is no general rule for the trimming value and thus it is subjective. One can take lower or higher values according to the data quality, see for example Liebmann et al. [28], Filzmoser and Todorov [29].

[Table 6 is here]

The estimated values of the regression coefficients and the scale parameter and also SEP and SEP_{trim} values are tabulated in Table 6. The SEP_{trim} values suggest that RAMML₁ and RAMML₂ are slightly preferable to MM, LTS, AMML₁, S and AMML₂. Furthermore, the scale estimates obtained based on RAMML₁ and RAMML₂ are much smaller than those of other estimation methods. This conclusion implies that estimates obtained by RAMML₁ and RAMML₂ give more reliable results. The scatter plot of the starsCYG data and the fitted regression lines are also illustrated in Figure 1. The fitted regression lines obtained from the RAMML₁ and RAMML₂ estimates are also satisfactory.

The weights obtained at the final step of the robust estimation procedures MM, RAMML₁ and RAMML₂ are plotted against the scaled residuals in Figure 2. The weights obtained from RAMML₁ and RAMML₂ are smaller than those of MM. This is because the weights given by the RAMML methodology allow more flexibility to the data compared to those from the MM, see also Figure 1.

[Figure 1 is here]

[Figure 2 is here]

5.2. Aircraft data

This data set consisting of 23 observations on single-engine aircrafts is obtained from the Office of Naval Research over the years 1947-1979, see Gray [30] and Rousseeuw and Leroy [2]. It contains four predictors and a response variable whose descriptions are given below:

$y = \text{cost (in units of \$100,000)}, \quad x_1 = \text{aspect ratio}, \quad x_2 = \text{lift-to-drag ratio},$

$x_3 = \text{weight of plane (in pounds)} \quad \text{and} \quad x_4 = \text{maximal thrust}.$

We use the methods MM, LTS, AMML₁, RAMML₁, S, AMML₂, and RAMML₂ to estimate the unknown parameters. Robust diagnostics obtained based on LTS show that there are two outliers in the data. The estimated values of the regression coefficients and the scale parameter along with performance criteria are shown in Table 7. The minimum SEP_{trim} value is obtained for LTS which is followed by MM, RAMML₁ and RAMML₂. However, the scale estimates obtained from RAMML₁ and RAMML₂ are smaller than those of LTS and MM. Therefore, RAMML₁ and RAMML₂ are also promising and can be considered as good alternative estimators.

In Figure 3, the weights obtained at the last step of the robust estimation procedures MM, RAMML₁ and RAMML₂ are plotted against the scaled residuals. Also in this example, the RAMML weights are smaller than those of the MM estimator; this is because the RAMML weights are more flexible.

[Table 7 is here]

[Figure 3 is here]

6. Conclusions

In this study, we propose RAMML estimators which are robust to both y and x outliers. The RAMML estimators are obtained by downweighting x -outliers using an extra weight function in the AMML estimation method.

The RAMML estimators have some advantage over existing robust estimators: they are (i) explicitly formulated and thus easy to compute, (ii) efficient under the normality assumption and (iii) more preferable according to the MSE and SEP criteria. These properties make them good alternatives to existing robust estimators. The simulation studies have also shown that the RAMML estimators are improving over the other considered robust estimators if the contamination level increases, in case of heavy-tailed error distributions, and if the number of explanatory variables increases. Thus, in more difficult data situations there the advantage gets more pronounced. Another interesting feature of the RAMML estimators is that the estimated residual variance is in many situation smaller than for the alternatives. Together with the other properties, this is a useful feature which is based on a more flexible weighting scheme within the computation of the estimator.

Although the new robust estimators show good performance, there are still some points that should be considered. One of them is the choice of the weights δ_i^x assigned to the predictors. Different choices of δ_i^x will affects the efficiencies of the RAMML estimators. In this study, the weights δ_i^x are obtained similar to the weights δ_i . In the

future studies, alternative weight functions such as Huber, Fair, etc. will be considered.

Acknowledgements

This study is supported by “The Scientific and Technological Research Council of Turkey (TUBITAK)” as part of the “2219 – International Postdoctoral Research Scholarship Programme”.

References

- [1] Donmez A. Adaptive estimation and hypothesis testing methods. METU; 2010.
- [2] Rousseeuw PJ, Leroy AM. Robust Regression and Outlier Detection. Wiley; 1987.
- [3] Huber PJ. Robust regression: Asymptotics, conjectures, and monte carlo. *Ann. Stat.* 1973;1:799-821.
- [4] Rousseeuw P, Yohai V. Robust Regression by Means of S-Estimators. *Robust nonlinear time Ser. Anal.* Springer; 1984. p. 256-272.
- [5] Rousseeuw PJ. Least median of squares regression. *J. Am. Stat. Assoc.* 1984;79:871-880.
- [6] Yohai VJ. High Breakdown-Point and High Efficiency Robust Estimates for Regression. *Ann. Stat.* 1987;15:642-656.
- [7] Maronna RA, Martin RD, Yohai VJ. Robust Statistics: Theory and Methods. *Robust Stat. Theory Methods.* Wiley; 2006.
- [8] Gschwandtner M, Filzmoser P. Computing Robust Regression Estimators: Developments since Duttor 1977. *Austrian J. Stat.* 2012;41:45-58.
- [9] Yu C, Yao W. Robust linear regression: A review and comparison. *Commun. Stat. Simul. Comput.* Taylor and Francis Inc.; 2017. p. 6261-6282.
- [10] Tiku ML. Estimating the mean and standard deviation from a censored normal sample. *Biometrika.* 1967;54:155-1665.
- [11] Tiku ML. Estimating the parameters of normal and logistic distributions from a censored normal sample. *Aust. J. Stat.* 1968;10:64-74.
- [12] Tiku ML, Suresh RP. A new method of estimation for location and scale parameters. *J. Stat. Plan. Inference.* 1992;30:281-292.
- [13] Tiku ML, Islam MQ, Selcuk AS. Nonnormal regression. II. Symmetric distributions. *Commun. Stat. - Theory Methods.* 2001;30:1021-1045.
- [14] Islam MQ, Tiku ML. Multiple Linear Regression Model Under Nonnormality. *Commun. Stat. - Theory Methods.* 2005;33:2443-2467.
- [15] Lucas A. Robustness of the student t based M-estimator. *Commun. Stat. - Theory Methods.* 1997;26:1165-1182.
- [16] Arslan O, Genc AI. Robust location and scale estimation based on the univariate generalized t (GT) distribution. *Commun. Stat. Methods.* 2003;32:1505-1525.
- [17] Arslan O, Genc AI. The skew generalized t distribution as the scale mixture of a skew exponential power distribution and its applications in robust estimation. *Statistics (Ber).* 2009;43:481-498.
- [18] Acitas S, Kasap P, Senoglu B, et al. Robust estimation with the skew t2 distribution. *Pakistan J. Stat.* 2013;29:409-430.
- [19] Acitas S, Kasap P, Senoglu B, et al. One-step M-estimators: Jones and Faddys skewed t-distribution. *J. Appl. Stat.* 2013;40:1545-1560.
- [20] [1] Tiku ML, Src B. MMLEs are as good as M-estimators or better. *Stat. Probab. Lett.* 2009;79:984-989.
- [21] Acitas S, Filzmoser P, Senoglu B. A Robust Adaptive Modified Maximum Likelihood Estimator for the Linear Regression Model. *Olomoucian Days Appl. Math.* 2019. Olomouc, Czech Republic; 2019.

- [22] Puthenpura S, Sinha NK. Modified maximum likelihood method for the robust estimation of system parameters from very noisy data. *Automatica*. 1986;22:231-235.
- [23] Vaughan DC. On the Tiku–Suresh method of estimation. *Commun. Stat. - Theory Methods*. 1992;21:329-340.
- [24] Bhattacharyya GK. The asymptotics of maximum likelihood and related estimators based on type ii censored data. *J. Am. Stat. Assoc.* 1985;80:398-404.
- [25] Lopuhaa HP, Rousseeuw PJ. Breakdown points of affine equivariant estimators of multivariate location and covariance matrices. *Ann. Stat.* 1991;19:229-248.
- [26] Croux C, Filzmoser P, Oliveira MR. Algorithms for Projection-Pursuit robust principal component analysis. *Chemom. Intell. Lab. Syst.* 2007;87:218-225.
- [27] Fritz, H., Filzmoser P, Croux. C. A comparison of algorithms for the multivariate L_1 -median. *Comput. Stat.* 2012;27:393-410.
- [28] Liebmann B, Filzmoser P, Varmuza K. Robust and classical PLS regression compared. *J. Chemom.* 2010;24:111-120.
- [29] Filzmoser P, Todorov V. Review of robust multivariate statistical methods in high dimension. *Anal. Chim. Acta.* 2011;705:2-14.
- [30] Gray JB. Graphics for Regression Diagnostics. *Am. Stat. Assoc. Proc. Stat. Comput. Sect. ASA. Washington, D.C., U.S.A.*; 1985.

Table 1. The simulated MSEs of MM, LMS, AMML₁, RAMML₁, LTS, AMML₂, RAMML₂ estimators for regression coefficients: $n = 50$.

Estimator	Normal	Laplace	t_5	t_1	Slash	Normal	Laplace	t_5	t_1	Slash
Contamination level = 10%										
$m = 1$										
$\ell = 5$										
MM	0.0474	0.0831	0.0519	1.1207	2.2697	0.0397	0.0774	0.0422	0.4349	1.4712
LTS	0.1111	0.1423	0.1665	0.3463	1.2311	0.1181	0.1363	0.1534	0.3616	1.2682
AMML ₁	0.8011	1.2815	1.1698	2.1221	2.2956	0.2737	0.8501	0.6461	2.6837	3.3277
RAMML ₁	0.0323	0.0476	0.0437	0.1808	0.3142	0.0295	0.0419	0.0355	0.1640	0.2536
S	0.0856	0.1003	0.0803	0.3766	1.7704	0.0800	0.0902	0.0702	0.4027	1.5348
AMML ₂	0.6775	1.0634	1.0226	2.0572	2.2894	0.1907	0.5905	0.4307	2.4178	3.2486
RAMML ₂	0.0321	0.0463	0.0430	0.1687	0.3048	0.0296	0.0415	0.0354	0.1556	0.2419
$m = 5$										
MM	0.6763	0.8421	0.7994	1.3103	1.8802	0.7433	0.8264	0.8419	1.3939	1.8336
LTS	1.1991	1.2970	1.3264	1.6641	2.4065	1.1903	1.3648	1.3543	1.7104	2.4053
AMML ₁	0.9573	1.1174	1.0607	1.5667	2.1197	0.9339	1.0755	1.0617	1.6799	2.1338
RAMML ₁	0.3569	0.4778	0.4295	0.9849	1.5785	0.1352	0.2245	0.1906	0.7523	1.2911
S	0.9484	1.2203	1.2053	1.7732	2.9104	1.0298	1.1326	1.2092	1.8727	2.6679
AMML ₂	0.8496	1.0715	0.9965	1.5387	2.0872	0.7432	0.9161	0.8933	1.6366	2.0738
RAMML ₂	0.2978	0.4413	0.3946	0.9560	1.5511	0.1367	0.2235	0.1961	0.7183	1.2272
$m = 10$										
MM	1.2647	1.6644	1.5661	2.5889	3.798	1.2768	1.6596	1.6618	2.7727	3.6821
LTS	2.5168	3.1116	2.9643	4.1231	7.4949	2.4722	3.0758	2.9561	4.0753	6.1214
AMML ₁	1.3476	1.6566	1.5553	2.9153	4.5046	1.3567	1.6161	1.5779	3.1962	4.1157
RAMML ₁	0.9400	1.218	1.1349	2.4783	4.0340	0.3106	0.5558	0.4720	2.0974	3.1041
S	2.0672	2.8438	2.6515	4.3687	6.9140	2.1380	2.7441	2.6572	4.1898	6.5537
AMML ₂	1.1733	1.5749	1.4535	2.8347	4.2468	1.0396	1.4531	1.3917	3.0774	3.9912
RAMML ₂	0.7825	1.1355	1.0536	2.4038	3.7700	0.3146	0.5553	0.4759	1.9827	2.9669
$m = 20$										
MM	3.8055	5.4072	5.1708	11.4061	19.3850	3.7672	5.3245	5.1794	12.7201	19.0190
LTS	7.1310	9.4504	8.9798	20.3109	35.0990	7.5826	9.4996	9.6198	21.3224	33.3410
AMML ₁	2.8865	3.7237	3.4436	9.6187	16.4300	3.0323	3.9743	3.6027	9.9341	16.4600
RAMML ₁	2.7306	3.6004	3.2992	9.3477	16.1550	1.2295	2.3787	1.9306	8.6733	15.1420
S	6.0778	8.2255	7.5602	20.5418	35.2120	5.9019	8.7704	7.7262	20.0038	29.1150
AMML ₂	2.4717	3.3454	3.1723	10.0115	15.4510	2.4700	3.4223	3.0899	9.6530	15.5550
RAMML ₂	2.2657	3.1453	2.9610	9.6906	15.0810	0.9957	1.9028	1.5561	8.3624	14.3660
Contamination level = 20%										
$m = 1$										
MM	0.9063	1.5115	1.3348	2.8098	3.1718	1.1198	1.6951	1.6449	3.1743	3.6786
LTS	1.9771	2.1900	2.1504	2.9506	3.4027	2.2838	2.3322	2.6162	3.1222	3.7683
AMML ₁	2.6602	2.8145	2.7665	2.9762	3.0118	2.6405	3.1454	3.1654	3.6689	3.6903
RAMML ₁	0.2286	0.3897	0.3260	0.7351	0.9377	0.0328	0.0530	0.0481	0.1758	0.2441
S	1.0514	1.7672	1.5666	3.1428	3.6572	1.1219	1.7432	1.7081	3.2864	3.8538
AMML ₂	2.4319	2.7249	2.6339	2.9805	3.0240	1.8558	2.7676	2.6542	3.6578	3.6942
RAMML ₂	0.1452	0.3293	0.2638	0.7275	0.9437	0.0333	0.0532	0.0488	0.1690	0.2341
$m = 5$										
MM	1.3353	1.4435	1.3949	1.7396	2.0893	1.3779	1.4804	1.4346	1.6947	2.0291
LTS	2.3538	2.6105	2.4865	3.2080	4.4527	2.4167	2.6958	2.5875	3.2793	4.4298
AMML ₁	1.2048	1.2656	1.2523	1.7383	2.1198	1.2499	1.3365	1.2883	1.7571	2.1580
RAMML ₁	0.9746	1.0474	1.0370	1.5205	1.8879	0.1904	0.3055	0.2579	0.8240	1.3294
S	2.1677	2.4305	2.3244	2.9160	3.8286	2.2236	2.4929	2.3276	2.8058	3.8126
AMML ₂	1.1977	1.2625	1.2452	1.7209	2.0833	1.2289	1.3164	1.2828	1.7381	2.1254
RAMML ₂	0.9624	1.0412	1.0248	1.5008	1.8485	0.1847	0.2953	0.2536	0.8018	1.2893
$m = 10$										
MM	2.7202	3.3625	3.3854	4.6944	6.0067	2.9763	3.4669	3.1312	4.4531	5.9192
LTS	4.3832	5.7078	5.3309	8.7250	12.4536	4.3648	5.6491	5.2871	9.4753	14.6904
AMML ₁	1.8994	2.2494	2.1421	3.6946	5.0060	1.9994	2.2793	2.2528	3.8856	5.301
RAMML ₁	1.8389	2.1956	2.0839	3.6169	4.9443	0.9558	1.3027	1.2283	3.0656	4.6646
S	3.8241	4.7879	4.6691	7.5793	10.5919	4.1517	5.1859	4.5923	8.2151	11.6736
AMML ₂	1.6618	1.9274	1.8949	3.3243	4.5066	1.7220	2.0002	1.9096	3.3479	4.4427
RAMML ₂	1.5641	1.8257	1.7968	3.1857	4.3612	0.6103	0.9395	0.7995	2.4694	3.7191
$m = 20$										
MM	10.6851	13.8939	13.6416	50.8300	87.5050	11.3025	14.3318	13.6001	52.1760	77.6950
LTS	13.5703	19.2541	19.0750	89.8350	126.6330	15.2065	20.6194	18.4261	83.5840	123.8700
AMML ₁	12.7179	17.6868	17.7259	80.4410	112.0310	14.2021	19.1793	17.0929	74.2650	107.8330
RAMML ₁	12.7966	17.8217	17.8598	81.1750	113.1480	14.2039	19.2348	17.1026	74.7500	108.9300
S	14.2006	20.8200	18.3638	143.607	143.8900	13.5973	19.8771	18.3664	95.4700	115.5300
AMML ₂	5.0160	7.5507	5.4790	53.6650	40.8890	5.5229	7.1065	7.4644	32.7620	41.8120
RAMML ₂	5.1211	7.7963	5.6209	54.7630	42.1430	4.9299	6.6159	6.9766	33.1670	42.4460

Table 2. The simulated MSEs of MM, LMS, AMML₁, RAMML₁, LTS, AMML₂, RAMML₂ estimators for regression coefficients: $n = 200$.

Estimator	Normal	Laplace	t_5	t_1	Slash	Normal	Laplace	t_5	t_1	Slash
Contamination level = 10%										
$m = 1$										
$\ell = 5$						$\ell = 10$				
MM	0.0060	0.0085	0.0078	0.5136	2.4011	0.3472	0.4862	0.5819	1.1543	1.3631
LTS	0.0069	0.0087	0.0090	0.0257	0.9422	0.8945	0.7963	1.0423	1.2114	1.5357
AMML ₁	0.6964	1.2441	1.1109	2.1133	2.2443	0.9002	1.0152	1.0219	1.2870	1.4867
RAMML ₁	0.0071	0.0128	0.0100	0.0561	0.1066	0.4557	0.4614	0.5554	0.9008	1.1409
S	0.0147	0.0095	0.0156	0.0327	1.1916	0.5249	0.6682	0.8161	1.4541	1.8625
AMML ₂	0.5764	0.9927	0.9086	2.0420	2.2402	0.6607	0.9147	0.8968	1.2699	1.4605
RAMML ₂	0.0071	0.0122	0.0098	0.0493	0.1023	0.2258	0.3525	0.3854	0.9031	1.1201
$m = 5$										
MM	0.2626	0.3484	0.4125	0.9128	1.0750	0.2958	0.3375	0.4387	0.9928	1.1468
LTS	0.7012	0.5478	0.7708	0.8773	1.1372	0.7343	0.5465	0.8001	0.9557	1.2046
AMML ₁	0.8465	0.9088	0.9050	1.0321	1.1099	0.7398	0.8282	0.8656	1.1169	1.2055
RAMML ₁	0.1948	0.2184	0.2537	0.4840	0.6080	0.0296	0.0467	0.0407	0.1427	0.2510
S	0.2876	0.3808	0.4624	1.0463	1.2826	0.3312	0.3912	0.4801	1.0471	1.3217
AMML ₂	0.7168	0.8596	0.8426	1.0253	1.1002	0.4003	0.6415	0.6182	1.1031	1.1935
RAMML ₂	0.0900	0.1616	0.1655	0.4853	0.6009	0.0298	0.0464	0.0409	0.1329	0.2359
$m = 10$										
MM	0.3472	0.4862	0.5819	1.1543	1.3631	0.4094	0.5243	0.5526	1.1824	1.4381
LTS	0.8945	0.7963	1.0423	1.2114	1.5357	0.9454	0.8472	1.0492	1.2255	1.5759
AMML ₁	0.9002	1.0152	1.0219	1.2870	1.4867	0.8160	0.8524	0.9379	1.3207	1.5742
RAMML ₁	0.4557	0.4614	0.5554	0.9008	1.1409	0.0639	0.1018	0.0851	0.3059	0.5717
S	0.5249	0.6682	0.8161	1.4541	1.8625	0.5275	0.6597	0.7939	1.4660	1.8753
AMML ₂	0.6607	0.9147	0.8968	1.2699	1.4605	0.3772	0.6000	0.6214	1.2966	1.5422
RAMML ₂	0.2258	0.3525	0.3854	0.9031	1.1201	0.0642	0.1008	0.0847	0.2860	0.5390
$m = 20$										
MM	0.5684	0.9978	0.9497	1.6374	1.9992	0.5490	1.0318	0.8953	1.6088	2.0397
LTS	1.3601	1.4243	1.5033	1.9556	2.5446	1.3503	1.4359	1.5317	1.8563	2.5387
AMML ₁	1.0659	1.2139	1.2080	1.8028	2.2552	1.0480	1.1482	1.1866	1.7621	2.2977
RAMML ₁	0.8478	0.9343	0.9641	1.5636	2.0235	0.1317	0.2179	0.1891	0.6689	1.2338
S	1.1751	1.8051	1.7165	2.4816	3.3106	1.1005	1.7525	1.6309	2.4502	3.3417
AMML ₂	0.7525	1.1542	1.0870	1.7551	2.1846	0.6228	1.0910	0.9737	1.7253	2.2274
RAMML ₂	0.5599	0.9062	0.8535	1.5359	1.9591	0.1334	0.2177	0.1893	0.6220	1.1623
Contamination level = 20%										
$m = 1$										
$\ell = 5$						$\ell = 10$				
MM	0.4461	0.9931	1.0734	3.0839	3.1299	0.3609	1.0414	0.9840	3.6322	3.7580
LTS	2.1464	1.7778	2.5918	3.1700	3.3282	2.3985	2.1247	2.7607	3.5390	3.8195
AMML ₁	2.7559	2.8206	2.8515	2.9719	2.9715	2.8591	3.2785	3.3429	3.6997	3.6998
RAMML ₁	0.1690	0.2314	0.2571	0.6988	0.8005	0.0073	0.0119	0.0100	0.0418	0.0688
S	0.4982	1.1499	1.2401	3.5014	3.5937	0.3781	1.1068	1.0245	3.7578	3.9076
AMML ₂	2.4541	2.7179	2.6921	2.9850	2.9847	1.4806	2.6798	2.5184	3.7018	3.7049
RAMML ₂	0.0543	0.1662	0.1431	0.6877	0.7970	0.0070	0.0119	0.0097	0.0398	0.0656
$m = 5$										
MM	1.0223	1.0363	1.0332	1.0759	1.1504	1.0541	1.0680	1.0679	1.1116	1.1818
LTS	1.2814	1.2889	1.3084	1.2890	1.4834	1.3104	1.3122	1.3382	1.3256	1.4875
AMML ₁	0.9921	1.0100	1.0050	1.0892	1.1774	1.0375	1.0525	1.0507	1.1288	1.2108
RAMML ₁	0.7619	0.7933	0.7815	0.8871	0.9778	0.0405	0.0654	0.0573	0.1956	0.3214
S	1.3082	1.3170	1.3401	1.3314	1.5485	1.3187	1.3456	1.3467	1.3639	1.5658
AMML ₂	0.9932	1.0109	1.0062	1.0835	1.1666	1.0382	1.0525	1.0513	1.1226	1.2012
RAMML ₂	0.7670	0.7988	0.7873	0.8863	0.9722	0.0422	0.0669	0.0590	0.1870	0.3108
$m = 10$										
MM	1.1484	1.1931	1.1811	1.2888	1.4655	1.1622	1.2124	1.1898	1.3184	1.4636
LTS	1.5638	1.5978	1.6118	1.7976	2.2175	1.5719	1.6271	1.6265	1.8212	2.2973
AMML ₁	1.0943	1.1339	1.1165	1.3262	1.5427	1.1134	1.1600	1.1415	1.3402	1.5488
RAMML ₁	0.9697	1.0133	0.9941	1.2034	1.4133	0.0925	0.1651	0.1362	0.4595	0.7448
S	1.6621	1.7289	1.7171	1.8785	2.4519	1.6672	1.7437	1.7504	1.9083	2.4676
AMML ₂	1.0964	1.1357	1.1185	1.3084	1.5171	1.1153	1.1609	1.1433	1.3237	1.5173
RAMML ₂	0.9760	1.0190	0.9998	1.1905	1.3934	0.0991	0.1732	0.1446	0.4428	0.7166
$m = 20$										
MM	1.5137	1.5901	1.5740	1.7791	2.2969	1.4988	1.5764	1.5847	1.7845	2.2317
LTS	2.0562	2.2847	2.1948	3.0003	4.2073	2.0834	2.2013	2.2635	3.0191	4.2844
AMML ₁	1.2698	1.3726	1.3359	1.8244	2.2852	1.2867	1.3659	1.3439	1.8106	2.2819
RAMML ₁	1.2068	1.3077	1.2721	1.7406	2.1988	0.2928	0.4745	0.4030	1.1256	1.6735
S	2.2856	2.4812	2.4787	3.1068	4.5225	2.2711	2.4297	2.4752	3.1939	4.4752
AMML ₂	1.2800	1.3773	1.3452	1.7765	2.2147	1.2944	1.3700	1.3521	1.7605	2.1994
RAMML ₂	1.2214	1.3168	1.2862	1.7005	2.1384	0.3352	0.5164	0.4477	1.0955	1.6096

Table 3. The simulated MSEs of MM, LMS, AMML₁, RAMML₁, LTS, AMML₂, RAMML₂ estimators for scale parameter when the error distribution is $N(0, 1)$.

	$\ell = 5$				$\ell = 10$			
	$m = 1$	$m = 5$	$m = 10$	$m = 25$	$m = 1$	$m = 5$	$m = 10$	$m = 25$
$n = 50$								
Contamination level = 10%								
MM	0.0460	0.0233	0.0308	0.0860	0.0435	0.0238	0.0284	0.0866
LTS	0.0778	0.0619	0.0576	0.1296	0.0760	0.0702	0.0609	0.1243
AMML ₁	0.3110	0.0777	0.0819	0.0689	0.0837	0.0706	0.0852	0.0655
RAMML ₁	0.0382	0.0194	0.0301	0.0451	0.0406	0.0317	0.0325	0.0614
S	0.0460	0.0252	0.0329	0.0713	0.0435	0.0262	0.0298	0.0734
AMML ₂	0.2623	0.0660	0.0626	0.0673	0.0621	0.0497	0.0505	0.0645
RAMML ₂	0.0390	0.0251	0.0283	0.0378	0.0410	0.0334	0.0353	0.0502
Contamination level = 20%								
MM	0.1366	0.0618	0.1897	0.3920	0.1466	0.0577	0.1875	0.4297
LTS	0.3193	0.0786	0.2786	0.7529	0.3655	0.0689	0.2716	0.7505
AMML ₁	0.6533	0.0418	0.0421	0.6658	0.5751	0.0484	0.0472	0.6669
RAMML ₁	0.0411	0.0183	0.0503	0.6979	0.0537	0.0491	0.0865	0.6993
S	0.1366	0.0570	0.1498	0.2782	0.1467	0.0544	0.1479	0.3162
AMML ₂	0.6240	0.0433	0.0303	0.0890	0.4141	0.0478	0.0321	0.1075
RAMML ₂	0.0367	0.0188	0.0231	0.1093	0.0555	0.0481	0.0556	0.1378
$n = 200$								
Contamination level = 10%								
MM	0.0282	0.0211	0.0181	0.0129	0.0181	0.0225	0.0187	0.0132
LTS	0.0571	0.0751	0.0717	0.0565	0.0717	0.0801	0.0750	0.0599
AMML ₁	0.3020	0.0771	0.0704	0.0689	0.0704	0.0613	0.0593	0.0721
RAMML ₁	0.0294	0.0093	0.0175	0.0256	0.0175	0.0235	0.0224	0.0204
S	0.0282	0.0219	0.0186	0.0113	0.0186	0.0229	0.0196	0.0124
AMML ₂	0.2522	0.0686	0.0500	0.0389	0.0500	0.0291	0.0201	0.0342
RAMML ₂	0.0303	0.0160	0.0196	0.0227	0.0196	0.0251	0.0249	0.0240
Contamination level = 20%								
MM	0.1393	0.0115	0.0236	0.0768	0.1364	0.0119	0.0239	0.0728
LTS	0.4019	0.0182	0.0110	0.0295	0.4286	0.0212	0.0115	0.0260
AMML ₁	0.6431	0.0375	0.0381	0.0310	0.6230	0.0413	0.0400	0.0320
RAMML ₁	0.0133	0.0071	0.0106	0.0093	0.0495	0.0352	0.0292	0.0157
S	0.1393	0.0115	0.0230	0.0753	0.1364	0.0118	0.0237	0.0730
AMML ₂	0.6317	0.0339	0.0331	0.0238	0.3894	0.0375	0.0347	0.0247
RAMML ₂	0.0164	0.0061	0.0086	0.0067	0.0490	0.0368	0.0308	0.0168

Table 4. The simulated MSEs of MM, LMS, AMML₁, RAMML₁, LTS, AMML₂, RAMML₂ estimators for regression coefficients when the distribution of error terms is $N(0, 1)$ with zero contamination.

Estimator	$n = 50$				$n = 200$			
	$m = 1$	$m = 5$	$m = 10$	$m = 20$	$m = 1$	$m = 5$	$m = 10$	$m = 20$
LS	0.0222	0.1145	0.2600	0.7357	0.0051	0.0251	0.0531	0.1112
MM	0.0231	0.1249	0.2922	1.0148	0.0053	0.0270	0.0590	0.1287
LTS	0.0344	0.2306	0.5866	1.8022	0.0074	0.0376	0.0803	0.1793
AMML ₁	0.0224	0.1166	0.2636	0.7574	0.0051	0.0254	0.0542	0.1135
RAMML ₁	0.0273	0.1180	0.2654	0.7635	0.0065	0.0255	0.0546	0.1143
S	0.0663	0.4081	0.9027	2.2393	0.0162	0.0955	0.2187	0.4537
AMML ₂	0.0224	0.1172	0.2653	0.7628	0.0051	0.0255	0.0544	0.1141
RAMML ₂	0.0274	0.1187	0.2677	0.7702	0.0065	0.0256	0.0548	0.1149

Table 5. The simulated MSEs of MM, LMS, AMML₁, RAMML₁, LTS, AMML₂, RAMML₂ estimators for regression coefficients with correlated predictors: $n = 200$, $\ell = 10$ and contamination level = 10%.

Estimator	Normal	Laplace	t_5	t_1	Slash
$m = 5$					
MM	0.2415	0.3600	0.3311	0.8236	2.0186
LTS	0.2822	0.3797	0.3697	0.7139	2.6958
AMML ₁	0.2590	0.5424	0.4623	2.9306	4.1066
RAMML ₁	0.2447	0.3780	0.3460	1.2012	2.1839
S	0.6538	0.4250	0.6039	0.5792	2.9211
AMML ₂	0.2540	0.4548	0.4121	2.5534	3.9090
RAMML ₂	0.2455	0.3723	0.3440	1.1143	2.0515
$m = 10$					
MM	0.5865	1.1378	0.8459	8.5211	14.4712
LTS	1.8531	5.5060	5.7911	27.7480	44.4990
AMML ₁	0.9999	2.8892	2.4072	10.8070	13.7108
RAMML ₁	0.6063	1.0319	0.8598	3.4661	5.9173
S	2.2357	2.7599	2.5881	34.5933	49.7293
AMML ₂	0.8313	1.8118	1.4004	10.5471	13.3625
RAMML ₂	0.6079	1.0161	0.8540	3.2843	5.5056
$m = 20$					
MM	1.2732	1.9421	1.7584	4.8325	9.6923
LTS	1.4942	2.0848	2.0331	4.3235	9.7048
AMML ₁	1.1957	2.0302	1.7448	8.5954	22.3637
RAMML ₁	1.2593	2.0304	1.7781	6.6322	12.7007
S	4.3629	3.4980	6.5179	21.3058	48.5781
AMML ₂	1.3343	2.1102	2.1823	11.3807	25.6140
RAMML ₂	1.2640	1.9994	1.7705	6.6776	12.9944

Table 6. The estimated values of the regression coefficients and the scale parameter and also SEP and SEP_{trim} values for the starsCYG data.

Method	β_0	β_1	σ	SEP	SEP _{trim}
MM	-4.9694	2.2532	0.4715	0.9556	0.3282
LTS	-8.5001	3.0462	0.4562	1.1507	0.3266
AMML ₁	3.4697	0.3409	0.5239	0.6000	0.3994
RAMML ₁	-8.0822	2.9523	0.3249	1.1269	0.3252
S	-9.5708	3.2904	0.4715	1.2133	0.3325
AMML ₂	3.3016	0.3790	0.5214	0.6041	0.3972
RAMML ₂	-8.0907	2.9553	0.3249	1.1277	0.3252

Table 7. The estimated regression coefficients, scale parameter and values of the performance criteria for the Aircraft data.

Method	β_0	β_1	β_2	β_3	β_4	σ	SEP	SEP _{trim}
MM	6.1396	-3.2306	1.6713	0.0019	-0.0009	5.8932	10.5866	3.4636
LTS	9.5007	-3.0488	1.2100	0.0014	-0.0006	5.6927	12.4080	3.4110
AMML ₁	2.2997	-3.4300	2.0052	0.0025	-0.0013	6.5780	8.9124	3.8884
RAMML ₁	7.0264	-3.2411	1.6622	0.0018	-0.0009	3.9639	11.1015	3.4651
S	13.3733	-4.0220	1.5413	0.0017	-0.0010	5.8932	11.8666	3.5001
AMML ₂	1.8030	-3.4535	2.0451	0.0026	-0.0014	6.7367	8.7372	3.9778
RAMML ₂	6.9195	-3.2381	1.6674	0.0018	-0.0009	3.9821	11.0630	3.4699




Figure 1. Fitted regression lines to the starsCYG data.




Figure 2. Weights versus scaled residuals for the estimators MM, RAMML₁ and RAMML₂, based on the starsCYG data.




Figure 3. Weights versus scaled residuals for the Aircraft data.

