

THE DYNAMICAL APPROACH TO THE CONJUGACY IN GROUPS

IGOR PROTASOV, KSENIA PROTASOVA

Abstract. Given a discrete group G , we identify the Stone-Čech compactification βG with the set of all ultrafilters on G and put $G^* = \beta G \setminus G$. The action of G on G by the conjugations $(g, x) \mapsto g^{-1}xg$ induces the action of G on G^* by $(g, p) \mapsto p^g$, $p^g = \{g^{-1}Pg : P \in p\}$. We study interplays between the algebraic properties of G and the dynamical properties of (G, G^*) . In particular, we show that p^G is finite for each $p \in G^*$ if and only if the commutant of G is finite.

1991 MSC: 20E45, 54D80.

Keywords: space G^* of ultrafilters on a group G , the action of G on G^* by the conjugations, ultracenter of G^* .

1. INTRODUCTION AND PRELIMINARIES

Let a group G acts on a set X by

$$G \times X \longrightarrow X : (g, x) \mapsto gx.$$

We endow X with the discrete topology, identify the Stone-Čech extension βX of X with the set of all ultrafilters on X and put $X^* = \beta X \setminus X$. By the universal property of βX , every mapping $f : X \longrightarrow K$, where K is a compact Hausdorff space, gives up to the continuous mapping $f^\beta : \beta X \longrightarrow K$. Using this property, we extend the action of G on X to the action

$$G \times \beta G \longrightarrow \beta G : (g, p) \mapsto gp,$$

where gp is the ultrafilter $\{gP : P \in p\}$. We note that $gp \in X^*$ for each $g \in G$, $p \in X^*$. Thus, we get the *dynamical system* (G, X^*) .

In the case $X = G$ and gx is the product of g and x in G , (G, G^*) is studied in [2]. The general case of (G, X^*) aroused in *Asymptology*, see the survey [4] and [1, Section 5].

In this paper, $X = G$, but G acts on X by the conjugations $(g, x) \mapsto g^x$, $g^x = x^{-1}gx$.

In what follows, all groups under consideration are suppose to be infinite. If $g \in G$ and $p \in G^*$ then

$$p^g = \{g^{-1}Pg : P \in p\}, \quad p^G = \{p^g : g \in G\}$$

and p^G is called the G -orbit of p .

We note that the closure clp^G of p^G coincides with the set $\{p^q : q \in \beta G\}$, where $p^q = q\text{-}\lim\{p^x : x \in G\}$. To describe p^q more explicitly, we take $Q \in q$ and, for each $x \in Q$, pick $P_x \in p$. Then $\bigcup\{x^{-1}P_x x : x \in Q\} \in p^q$ and these subsets form a base of p^q .

For a group G and $p \in G^*$, we denote

$$St(p) = \{g \in G : p^g = p\}, \quad Z_G = \{g \in G : g^x = g \text{ for each } x \in G\},$$

$$UZ_G = \{p \in G^* : St(p) = G\}.$$

and say that UZ_G is the *ultracenter* of G^* .

We note that $g \in St(p)$ if and only if there exists $P \in p$ such that $gx = xp$ for each $x \in P$, and $Z_G^* \subseteq UZ_G$. We use also the following observation: $|p^G| = |G : St(p)|$.

Proposition 1.1. *A group G is Abelian if and only if $UZ_G = G^*$.*

Proof. If G is Abelian then $St(p) = G$ for each $p \in G^*$ so $G^* \subseteq UZ_G$.

We suppose that $G^* = UZ_G$. If $g, x \in G$, $p \in G^*$ then $(xp)^g = xp$ and $(xp)^g = x^g p^g = x^g p$ so $x^g = x$ and G is Abelian. \square

The multiplication on a discrete group G can be extended (see [2]) on βG so that, for each $g \in G$, the mapping

$$\beta G \longrightarrow \beta G : p \mapsto gp$$

is continuous, and for each $p \in \beta G$, the mapping

$$\beta G \longrightarrow \beta G : q \mapsto qp$$

is continuous. Then, βG is a right topological semigroup and G^* is a subsemigroup of βG .

For $q, p \in \beta G$, we choose $Q \in q$ and, for each $g \in Q$, pick $P_g \in p$. Then $\bigcup\{xP_x : x \in Q\} \in qp$ and subsets of this form are a base for qp .

Proposition 1.2. *For a group G , either UZ_G is empty or UZ_G is a subsemigroup of G^* .*

Proof. It suffices to note that $(qp)^g = q^g p^g$ for all $g \in G$ and $q, p \in G^*$. \square

In Section 2, we show that, for every group G , G^* has a plenty of discrete orbits p^G and if G is countable then the set $\{p \in G^* : p^G \text{ is discrete}\}$ is dense in G^* . We give some sufficient conditions on a subset A of G under which some orbit p^G , $p \in A^*$ or each orbit p^G , $p \in A^*$ is infinite.

In Section 3, we characterize groups G such that each orbit p^G , $p \in G^*$ is finite.

In Section 4, we show that every dynamical system (H, X^*) can be embedded into (G, G^*) for an appropriate choice of G . This means that the dynamical systems of the form (G, G^*) could be very complicated.

2. PLENTY OF ORBITS

Theorem 2.1. *For every subset A of a group G such that $|A| = |G|$, we have*

$$|\{p \in G^* : A \in p \text{ and } p^G \text{ is discrete}\}| = 2^{2^{|A|}}.$$

Proof. We enumerate $G = \{g_\alpha : \alpha < |G|\}$, put $K_\alpha = \{g_\beta : \beta < \alpha\}$ and choose inductively $\{a_\alpha : \alpha < |G|\}$ such that

$$a_\alpha^{K_\alpha} \bigcap a_\beta^{K_\beta} = \emptyset$$

for all $\alpha < \beta$. If $g \in G$, $p \in G^*$, $\{a_\alpha : \alpha < |G|\} \in p$ and $|P| = |G|$ for each $P \in p$ then either $p^g = p$ or $(G \setminus \{a_\alpha : \alpha < |G|\}) \in p^G$. It suffices to note that we have $2^{2^{|G|}}$ possibilities to choose p with these properties. \square

Theorem 2.2. *For a countable group G , the following statements hold*

- (i) *the set $\{p \in G^* : p^G \text{ is discrete}\}$ is dense in G^* ;*
- (ii) *if $cl\ p^G \cap cl\ q^G \neq \emptyset$ then either $cl\ p^G \subseteq cl\ q^G$ or $cl\ q^G \subseteq cl\ p^G$;*
- (iii) *there exist an infinite subset $A \subseteq G$ such that $cl\ p^G \cap cl\ q^G = \emptyset$ for all distinct $p, q \in A^*$.*

Proof. (i) Apply Theorem 2.1.

(ii) Apply the Frolik's lemma [2, Corollary 3.42];

(iii) We take a subset A from the proof of Theorem 2.1. Let $p, q \in A^*$, $p \neq q$ but $cl\ p^G \cap cl\ q^G \neq \emptyset$. By (ii), either $p = cl\ q^G$ or $q = cl\ p^G$. In both cases, there exists $r \in A^*$, $g \in G$ such that $r^g \neq r$ and $r^g \in A^*$, contradicting the choice of A . \square

Theorem 2.3. *If some conjugate class a^G of a group G is infinite then there exists $p \in G^*$ such that p^G is infinite.*

Proof. We choose a sequence $(g_n)_{n \in \omega}$ in G such that $a^{g_n} \neq a^{g_m}$ for all distinct m, n . Let $q \in G^*$. If there exists an infinite subset $I \subseteq \omega$ such that $q^{g_m} \neq q^{g_n}$ for all distinct m, n then we put $p = q$.

Otherwise, there exists an infinite subset $J \subseteq \omega$ such that $q^{g_n} = q^{g_m}$ for all distinct $m, n \in J$. Then $(aq)^{g_n} = a^{g_n}q$, $(aq)^{g_m} = a^{g_m}q$. Since $a^{g_n} \neq a^{g_m}$, we have $(aq)^{g_n} \neq (aq)^{g_m}$, put $p = aq$. \square

For a subset A of a group G and $p \in G^*$, we denote $\Delta_p(A) = p^G \cap A^*$.

Theorem 2.4. *Let A be a subset of a group G such that $\Delta_p(A)$ is infinite for each $p \in A^*$. Then there exists $q \in A^*$ such that q^G is not discrete.*

Proof. We use the Zorn's lemma and compactness of G^* to choose the minimal by inclusion, closed subset S of G^* such that $S \cap Y^* \neq \emptyset$ and $g^{-1}Sg \subseteq S$ for each $g \in G$. Let $q \in S \cap Y^*$. Since $q^G \cap Y^*$ is infinite, there exists a limit point $r \in Y$ of q^G . By the minimality of S , q is a limit point of r^G . It follows that q is not isolated in q^G and q^G is not discrete. \square

Let $(y_n)_{n \in \omega}, (b_n)_{n \in \omega}$ be sequences in a group G . Adopting the dynamical terminology from [5] to the action of G on G by conjugations, we say that

$$Y = \left\{ b_n^{y_n^{\varepsilon_n} \dots y_0^{\varepsilon_0}} : n < \omega, \varepsilon_i \in \{0, 1\} \right\}$$

is a *piece-wise shifted FP-set* if

$$(*) \quad b_n^{y_n^{\varepsilon_n} \dots y_0^{\varepsilon_0}} = b_m^{y_m^{\delta_m} \dots y_0^{\delta_0}}, \quad \varepsilon_i, \delta_j \in \{0, 1\}$$

implies $n = m$ and $\varepsilon_0 = \delta_0, \dots, \varepsilon_n = \delta_n$.

Theorem 2.5. *If Y is a piece-wise shifted FP-set in a group G and $p \in Y^*$ then the orbit p^G is infinite.*

Proof. For $n < \omega$, we put

$$Y_m = \left\{ b_n^{y_n^{\varepsilon_n} \dots y_0^{\varepsilon_0}} : \varepsilon_0 = \dots = \varepsilon_{m-1} = 0, \varepsilon_m = 1 \right\}$$

and note that $Y_m \cap Y_{m'} = \emptyset$ if $m \neq m'$.

If $Y_m \not\subseteq p$ for each $m < \omega$ then $p^{y_n} \neq p^{y_k}$ for all distinct n, k .

If $Y_m \subseteq p$ then $p^g \neq p$ for $g = y_m$ and we repeat the arguments for p^g in place of p . \square

3. FINITE ORBITS

We recall that G is an FC-group if a^G is finite for $a \in G$.

Theorem 3.1. *For a group G , the following conditions are equivalent*

- (i) p^G is finite for each $p \in G^*$;
- (ii) there exists a natural number n such that $|p^G| < n$ for each $p \in G^*$;
- (iii) there exists a natural number n such that $|a^G| < n$ for each $a \in G$;

(iv) the commutant of G is finite.

Proof. The equivalence (iii) \Leftrightarrow (iv) is proved in [3], (ii) \Rightarrow (i) is evident.

(i) \Rightarrow (iii). If some class a^G is infinite then, by Theorem 2.3, some orbit p^G is infinite. We suppose that G is an FC-group, but for every natural number n , there exists $a \in G$ such that $|a^G| > n$. In view of Theorem 2.5, it suffices to construct a piece-wise shifted FP-set Y in G .

We choose $c_0, y_0 \in G$ such that $c_0^{y_0} \neq c_0$, put $b_0 = c_0$. Suppose that we have chosen y_0, \dots, y_k and c_0, \dots, c_k such that, for $b_i = c_0, \dots, c_i$, (*) holds for all $n, m \leq k$. We denote by Z the intersection of the centralizers of $c_0, \dots, c_k, y_0, \dots, y_k$. Since G is an FC, $|G : Z|$ is finite. It follows that Z has an arbitrary large class of conjugate elements in Z . Using this observation, we can choose $c_{k+1}, y_{k+1} \in Z$ such that, for $b_{k+1} = b_k c_{k+1}$, (*) holds for all $n, m \leq k+1$.

After ω steps, we get the desired piece-wise shifted FC-group

$$Y = \{b_n^{y_n^{\varepsilon_n}, \dots, y_0^{\varepsilon_0}} : n < \omega, \varepsilon_i \in \{0, 1\}\}.$$

(iii) \Rightarrow (ii). We suppose that there exists $p \in G^*$ such that $|p^G| \geq m$. We choose g_1, \dots, g_m in G such that $p^{g_i} \neq p^{g_j}$ for all $i < j \leq m$. Then there exists $P \in p$ such that $P^{g_i} \cap P^{g_j} = \emptyset$ for all $i < j \leq m$. If $a \in P$ then $a^{g_i} \neq a^{g_j}$ for $i < j \leq m$, contradicting (iii). \square

Example 3.2. We define a transitive group S of permutations of ω such that each orbit $Sp, p \in \omega^*$ is finite, and for every natural number n there exists $q \in \omega^*$ such that $|Sq| = n$.

First, partition ω infinite subsets $\{W_n : n \in \omega\}$. Second, partition each W_n into infinite subsets $\{W_{ni} : i < n\}$. Third, for each n , defines a bijection b_n of ω such that $f_n(W_{n0}) = W_{n1}$, $f_n(W_{n1}) = W_{n2}, \dots$, $f_n(W_{nn-1}) = W_0$, and $f_n(x) = x$ for each $x \in \omega \setminus W_n$. At last, take a group of permutations of ω generated by $\{f_n : n < \omega\}$ and all permutations with finite support. If $p \in \omega^*$ and $W_n \in p$ then $|Sp| = n$, otherwise, $|Sp| = 1$.

Question 3.3. Let G be a group such that p^G is discrete for each $p \in G^*$. Is p^G finite for each $p \in G^*$?

In light of Theorems 2.4 and 2.5, to answer this question in affirmative, it suffices to show that if some orbit p^G is infinite then G contains a piece-wise shifted FP-set.

4. DYNAMICAL EMBEDDINGS

Let $(H, Y^*), (G, X^*)$ be dynamical systems. We suppose that there exists an isomorphic embedding $f : H \longrightarrow G$ and an injective mapping $\phi : Y \longrightarrow X$ such

that, for each $(h, p) \in (H, Y^*)$, we have $\phi^\beta(hp) = f(h)\phi^\beta(p)$. We say that the pair (f, ϕ) is a dynamical embedding of (H, Y^*) into (G, X^*) .

Theorem 4.1. *Every dynamical system (H, Y) admits a dynamical embeddings into (G, G^*) for an appropriate choice of a group G .*

Proof. We consider $\{0, 1\}^Y$ as a group with point-wise addition *mod* 2. For $h \in H$ and $\chi \in \{0, 1\}^Y$, χ_h is defined by $\chi_h(y) = \chi(h^{-1}y)$. Then we define a semidirect product $G = \{0, 1\}^Y \rtimes H$ by

$$(\chi, h)(\chi', h') = (\chi + \chi'_h, hh').$$

Let $f : H \rightarrow G$ be the natural isomorphic embedding and $\phi : Y \rightarrow G$ is defined by $\phi(y) = (\psi_y, id)$, ψ_y is the characteristic function of $\{y\}$, id is the identity of H . Clearly, $\phi^\beta(hp) = (\phi^\beta(p))^{f(h)}$ so (f, ϕ) is the desired dynamical embedding. \square

Example 4.2. Let S be the group of all permutations of $\omega = \{0, 1, \dots\}$. Every orbit Sp , $p \in \omega^*$ is not discrete. Applying Theorem 4.2, we conclude that Theorem 2.2(i) needs not to be true for an uncountable group.

Example 4.3. Let F be the group of all permutations of ω with finite supports. Then $Fp = \{p\}$ for each $p \in \omega^*$. We use G given by Theorem 4.1 for (F, ω^*) . Then $\phi^\beta(\omega^*) \subseteq UZ_G$.

REFERENCES

- [1] T. Banach, I. Protasov, *Set-theoretical problems in Asymptology*, preprint, arXiv: 2004.01979.
- [2] N. Hindman, D. Strauss, *Algebra in the Stone-Ćech Compactification*, de Grueter, Berlin, New York, 1998.
- [3] B.H. Neuman, *Groups covered by permutable subsets*, J. London Math. Soc. **29**, (1954), 236-248.
- [4] I. Protasov, *Coarse spaces, ultrafilters and dynamical systems*, Topology Proceedings (to appear), preprint arXiv: 2003.12014.
- [5] I. Protasov, S. Slobodianiuk, *On the subset combinatorics of G -spaces*, Algebra Discrete Math. **17**: 1 (2014), 98-109.

CONTACT INFORMATION

I. Protasov:
 Faculty of Computer Science and Cybernetics
 Kyiv University by Taras Shevchenko
 Academic Glushkov pr. 4d
 03680 Kyiv, Ukraine
 i.v.protasov@gmail.com

K. Protasova:
Faculty of Computer Science and Cybernetics
Kyiv University
Academic Glushkov pr. 4d
03680 Kyiv, Ukraine
k.d.ushakova@gmail.com