

Spinning Black Holes Fall in Love

Alexandre Le Tiec^{1,2} and Marc Casals^{2,3}

¹*Laboratoire Univers et Théories, Observatoire de Paris, CNRS,
Université PSL, Université de Paris, 92190 Meudon, France*

²*Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, CEP 22290-180, Brazil*

³*School of Mathematics and Statistics, University College Dublin, Belfield, Dublin 4, Ireland*

The open question of whether a black hole can become tidally deformed by an external gravitational field has profound implications for fundamental physics, astrophysics and gravitational-wave astronomy. Love numbers characterize the tidal deformability of compact objects such as astrophysical (Kerr) black holes. We prove that all Love numbers vanish identically for a Kerr black hole in the nonspinning limit or for an axisymmetric tidal perturbation. In contrast to this result, we show that Love numbers are generically *nonzero* for a spinning black hole. Specifically, to linear order in the black hole spin and the weak perturbing tidal field, we compute in closed form the Love numbers that couple the mass-type and current-type quadrupole moments to the electric-type and magnetic-type quadrupolar tidal fields. This tidal deformability is potentially observationally important through its contribution to the accumulated gravitational-wave phase of an inspiralling stellar-mass compact object into a massive black hole. We show that for a dimensionless black hole spin ~ 0.1 , the nonvanishing quadrupolar Love numbers are $\sim 2 \times 10^{-3}$. This indicates that, despite black holes being particularly “stiff” compact objects, their nonvanishing tidal deformability could be detected by the future gravitational-wave interferometer LISA.

Introduction.—The deformability of a self-gravitating body under the effect of an external tidal field is a question of central interest in gravitational physics. Such tidal deformability can be characterized by a discrete set of *tidal Love numbers* (TLNs) [1], the gravitational analogue of the electric susceptibility in electrodynamics. Importantly, the TLNs of a self-gravitating body encode information about its internal structure, such as its composition or equation of state [2–4]. Those numbers were first introduced by Love [5, 6], in the context of Newtonian gravitation, to describe the Earth’s ocean tides due to its gravitational interaction with the Moon. They now play an important role in understanding the internal structure of the planets of the Solar System [7, 8], and even of exoplanets such as WASP-103b [9] and the TRAPPIST-1 system [10, 11].

In the context of relativistic gravitation, current and future gravitational-wave measurements of TLNs in binary inspirals provide a novel way of testing the inspiralling compact objects (neutron stars or black holes) and general relativity in the regime of strong gravitational fields [12]. In events that lead to the coalescence of two neutron stars, such as GW170817 [13] and GW190425 [14], the tidal effects become important for gravitational-wave frequencies of around 600 Hz, by accelerating the coalescence and affecting the gravitational-wave phase. Those two events have been used to set upper bounds on the tidal deformability of neutron stars, thereby constraining their radii and equation of state at supranuclear densities [14–18]. Some universal (i.e., equation-of-state independent) I-Love-Q [19] and I-Love-C [20] relations between the neutron star moment of inertia, quadrupolar TLN, quadrupole moment and compactness can be used to lift degeneracies among parameters in gravitational-wave signals, enhancing the measurability of the tidal effects. Over the coming decades, the observation by the planned LISA mission [21] of the gravitational-wave signals generated by the inspiral of stellar mass compact objects into massive black holes might place constraints on the TLNs of the central body that are roughly eight orders of mag-

nitude more stringent than current ones on neutron stars [22].

It is widely accepted that all astrophysical black holes are rotating and are thus described by the Kerr family [23, 24] of solutions of the Einstein equation. Previous works on the tidal deformability of black holes in general relativity have shown that, differently from the Newtonian case, the tidal field can be decomposed into two sectors, according to their parity, often called electric and magnetic, and, importantly, that the TLNs of nonrotating black holes all vanish [25–29]. This conclusion was extended to slowly rotating black holes, perturbatively in the spin, for a weak quadrupolar tidal field: to quadratic order for an axisymmetric quadrupole of electric-type [30], and to linear order for a generic quadrupole [31]. Given those remarkable results, there appears to be a widespread expectation that the vanishing of black hole TLNs extends to a generic rotating Kerr black hole in a generic multipolar tidal environment, e.g. in [22, 32–38]. In this Letter we will show that, on the contrary, the TLNs of a Kerr black hole do *not* vanish in general. The details will be given in a companion paper.

Throughout this Letter we use units such that $G = c = 1$, an overbar denotes the complex conjugation, we use the shorthand $\sum_{\ell m} \equiv \sum_{\ell=2}^{\infty} \sum_{m=-\ell}^{\ell}$ where ℓ is the multipolar index, as well as the notation $L \equiv i_1 \cdots i_{\ell}$ for a multi-index made of ℓ spatial indices.

Newtonian Love.—Consider an isolated, spherical, Newtonian body of mass M and equilibrium radius R . For a weak and slowly varying external tidal field, the induced mass multipole moments $I_L(t)$ are proportional to the applied tidal moments $\mathcal{E}_L(t)$,

$$I_L = \lambda_{\ell} \mathcal{E}_L, \quad (1)$$

where λ_{ℓ} is a constant. For instance, at quadrupolar order the induced mass quadrupole I_{ij} is proportional to the quadrupolar tidal field \mathcal{E}_{ij} . This adiabatic approximation holds as long as the typical timescales of the physical processes responsible for adjusting the matter distribution are much shorter than the typ-

ical timescale of variation of the tidal environment itself. The tidal deformability parameter λ_ℓ in (1) depends exclusively on the internal structure of the body and scales as $R^{2\ell+1}$. Introducing the dimensionless TLN k_ℓ associated with λ_ℓ , defined via

$$k_\ell \equiv -\frac{(2\ell-1)!!}{2(\ell-2)!} \frac{\lambda_\ell}{R^{2\ell+1}}, \quad (2)$$

the gravitational potential of the tidally-perturbed Newtonian body can be expanded over spherical harmonics $Y_{\ell m}(\theta, \phi)$ according to [1]

$$U = \frac{M}{r} - \sum_{\ell m} \frac{(\ell-2)!}{\ell!} \mathcal{E}_{\ell m} r^\ell \left[1 + 2k_\ell \left(\frac{R}{r} \right)^{2\ell+1} \right] Y_{\ell m}, \quad (3)$$

where r is the Euclidean distance to the center of mass, and the $2\ell+1$ coefficients $\mathcal{E}_{\ell m}$ are the spherical-harmonic modes of the tidal moment \mathcal{E}_L . The growing term $O(r^\ell)$ corresponds to the external 2^ℓ -polar tidal perturbation, and the decaying term $O(r^{-\ell-1})$ to the body's response, proportional to the TLN k_ℓ .

Einsteinian Love.—In general relativity, the tidal environment of a body is fully characterized by *two* families of tidal moments: the electric-type and magnetic-type tidal fields \mathcal{E}_L and \mathcal{B}_L . The former are the relativistic analogues of the Newtonian tidal moments introduced above, while the latter have no counterpart in Newtonian gravity. Similarly, the multipolar structure of that body is now characterized by *two* families of multipole moments: the mass-type and current-type multipole moments M_L and S_L , which are defined in a coordinate-independent manner for any asymptotically flat, stationary solutions of the vacuum Einstein equation [39, 40].

From now on we consider a weak, stationary tidal perturbation of a given compact body. The associated perturbed metric is $\hat{g}_{\alpha\beta} + h_{\alpha\beta}$, where the background $\hat{g}_{\alpha\beta}$ is an exact solution of the Einstein equation and the linear metric perturbation can be decomposed according to

$$h_{\alpha\beta} = h_{\alpha\beta}^{\text{tidal}} + h_{\alpha\beta}^{\text{resp}}. \quad (4)$$

Here, $h_{\alpha\beta}^{\text{tidal}}$ and $h_{\alpha\beta}^{\text{resp}}$ are *uniquely* specified as the linearly independent solutions of the linearized Einstein equation in vacuum (outside the body) that have the appropriate asymptotic behavior at large distances. In particular, the growing solution $h_{\alpha\beta}^{\text{tidal}}$ is unambiguously associated with the perturbing tidal field, while the decaying solution $h_{\alpha\beta}^{\text{resp}}$ is unambiguously associated with the corresponding linear response of the body (see the Supplemental Material). The perturbed metric $\hat{g}_{\alpha\beta} + h_{\alpha\beta}^{\text{resp}}$ is an asymptotically flat, stationary solution of the linearized Einstein equation in vacuum, and the corresponding multipole moments are

$$M_L = \mathring{M}_L + \delta M_L, \quad (5a)$$

$$S_L = \mathring{S}_L + \delta S_L, \quad (5b)$$

where a circle over a quantity indicates it is associated with the background $\hat{g}_{\alpha\beta}$ and a δ preceding that it is associated with the linear response $h_{\alpha\beta}^{\text{resp}}$.

For a nonspinning compact object, the background metric is spherically symmetric. By conservation of parity, the body's linear response contribution to the mass-type (resp. current-type) multipole moments can only couple to the electric-type (resp. magnetic-type) tidal moments,

$$\delta M_L = \lambda_\ell^{\text{el}} \mathcal{E}_L \quad \text{and} \quad \delta S_L = \lambda_\ell^{\text{mag}} \mathcal{B}_L, \quad (6)$$

where the tidal deformability parameters λ_ℓ^{el} and $\lambda_\ell^{\text{mag}}$ are constant. If R denotes the areal radius of the central body, then its dimensionless gravito-electric and gravito-magnetic TLNs k_ℓ^{el} and k_ℓ^{mag} are defined as per the formula (2) above. If the central object is spinning, however, then the spherical symmetry of the background metric is broken. Consequently, (i) the multipoles $(\delta M_L, \delta S_L)$ and the tidal moments $(\mathcal{E}_L, \mathcal{B}_L)$ cannot obey simple proportionality relationships akin to Eq. (6), (ii) the degeneracy of the azimuthal number m is lifted, (iii) fields with different parity can now mix, and (iv) the spherical-harmonic modes $M_{\ell m}$ and $S_{\ell m}$ of M_L and S_L can couple to modes $\mathcal{E}_{\ell' m'}$ and $\mathcal{B}_{\ell' m'}$ of \mathcal{E}_L and \mathcal{B}_L with $(\ell', m') \neq (\ell, m)$.

Quadrupole moments.—We consider a Kerr black hole of mass M and spin angular momentum per unit mass a embedded in a weak and stationary, but otherwise completely generic tidal environment. Hence, we work to linear order in the weak tidal perturbation, so that the TLNs are constants. Denoting by $\mathring{M}_{\ell m} = \mathring{M}_\ell \delta_{m0}$ and $\mathring{S}_{\ell m} = \mathring{S}_\ell \delta_{m0}$ the modes of, respectively, the mass-type and current-type multipole moments of the axisymmetric Kerr background spacetime, the multipole moments of the perturbed Kerr geometry read as

$$M_{\ell m} = \mathring{M}_{\ell m} + \lambda_{\ell m}^{M\mathcal{E}} \mathcal{E}_{\ell m} + \lambda_{\ell m}^{M\mathcal{B}} \mathcal{B}_{\ell m}, \quad (7a)$$

$$S_{\ell m} = \mathring{S}_{\ell m} + \lambda_{\ell m}^{S\mathcal{E}} \mathcal{E}_{\ell m} + \lambda_{\ell m}^{S\mathcal{B}} \mathcal{B}_{\ell m}, \quad (7b)$$

where $\lambda_{\ell m}^{M\mathcal{E}}$, $\lambda_{\ell m}^{M\mathcal{B}}$, $\lambda_{\ell m}^{S\mathcal{E}}$ and $\lambda_{\ell m}^{S\mathcal{B}}$ are the four families of Kerr TLNs, which are complex-valued. In here, we do not include couplings between different (ℓ, m) modes because, as we shall show, such couplings are absent in our results.

We computed (see Supplemental Material) the quadrupolar Kerr TLNs explicitly up to *linear* order in the dimensionless spin parameter $\chi \equiv a/M$. For convenience, we introduce the symbol “ \doteq ” for an equality that holds to that order. For any azimuthal number $|m| \leq 2$, the result simply reads

$$\lambda_{2m}^{M\mathcal{E}} \doteq \lambda_{2m}^{S\mathcal{B}} \doteq \frac{im\chi}{180} (2M)^5 \quad \text{and} \quad \lambda_{2m}^{M\mathcal{B}} \doteq \lambda_{2m}^{S\mathcal{E}} \doteq 0. \quad (8)$$

The mass-type (resp. current-type) quadrupole moment couples *only* to the electric-type (resp. magnetic-type) quadrupolar tidal perturbation. The coupling between $(\mathcal{E}_{2m}, \mathcal{B}_{2m})$ and (M_{2m}, S_{2m}) which arises from (7) and (8) is akin to a Zeeman-like splitting proportional to the azimuthal number m [41]. We have thus shown that a spinning black hole becomes tidally deformed under the effect of a weak and nonaxisymmetric tidal field. In particular, while in a binary system, a *spinning black hole falls in Love with its companion*. In the nonspinning limit ($\chi = 0$) or for an axisymmetric tidal field ($m = 0$), however, the TLNs in (8) vanish, in agreement with the results in [25–30].

In fact, we have extended those results to an arbitrarily spinning black hole in a generic multipolar tidal environment, as shown in the Supplemental Material. Our results agree with the previous ones but there is an apparent disagreement with [31], who found vanishing black hole TLNs for a generic (nonaxisymmetric) quadrupolar tidal perturbation. This disagreement is likely a consequence of the use of different tidal/response splits of the full physical solution.

The event horizon “radius” of a slowly spinning Kerr black hole is $2M(1 + O(\chi^2))$. Therefore, by analogy with the TLNs (2) introduced above for a spherical Newtonian body, we define the dimensionless black hole TLNs according to

$$k_{\ell m}^{M\mathcal{E}} \equiv -\frac{(2\ell-1)!!}{2(\ell-2)!} \frac{\lambda_{\ell m}^{M\mathcal{E}}}{(2M)^{2\ell+1}}, \quad (9)$$

and similarly for the $M\mathcal{B}$, $S\mathcal{E}$ and $S\mathcal{B}$ couplings. These TLNs generalize to slowly spinning black holes those for nonspinning compact objects [25, 26]. For a Kerr black hole with spin $\chi \sim 0.1$, Eqs. (8) and (9) imply $|k_{2,\pm 2}^{M\mathcal{E}}| = |k_{2,\pm 2}^{S\mathcal{B}}| \sim 2 \times 10^{-3}$. This small number could be compared, for instance, to the values $k_2^{\text{el}} \sim 0.05 - 0.15$ and $|k_2^{\text{mag}}| \lesssim 6 \times 10^{-4}$ of the gravito-electric and gravito-magnetic quadrupolar TLNs of a *nonspinning* neutron star, depending on the equation of state [25, 26]. Hence, while spinning black holes do deform like any self-gravitating body, they are particularly “stiff” compact objects.

Tidal Love tensor.—Having related the spherical-harmonic modes of the quadrupole moments to those of the quadrupolar tidal moments, we now relate δM_{ij} and δS_{ij} to \mathcal{E}_{ij} and \mathcal{B}_{ij} themselves. Multiplying Eq. (7) for $\ell = 2$ by Y_{2m} , using Eq. (8), and summing over modes, we obtain $\hat{r}^i \hat{r}^j \delta M_{ij} \propto \hat{r}^i \hat{\phi}^j \mathcal{E}_{ij}$ and $\hat{r}^i \hat{r}^j \delta S_{ij} \propto \hat{r}^i \hat{\phi}^j \mathcal{B}_{ij}$, where the unit angular vector $\hat{\phi}$ is orthogonal to the unit radial vector $\hat{\mathbf{r}}$. Therefore, δM_{ij} and δS_{ij} cannot be simply proportional to \mathcal{E}_{ij} and \mathcal{B}_{ij} , respectively. Rather, they must obey more general tensorial relations of the form

$$\delta M_{ij} = \sum_{k,l} \lambda_{ijkl} \mathcal{E}_{kl} \quad \text{and} \quad \delta S_{ij} = \sum_{k,l} \lambda_{ijkl} \mathcal{B}_{kl}, \quad (10)$$

where the constant tensor $\lambda_{ijkl} = O(\chi)$ is the quadrupolar *tidal Love tensor* (TLT) of the Kerr black hole. Such a complication with respect to the nonspinning case [25, 26] stems from the fact that the black hole spin breaks the spherical symmetry of the background spacetime. Using Eq. (8), an explicit calculation shows that the quadrupolar Kerr black hole TLT is given by

$$(\lambda_{ijkl}) \doteq \frac{\chi}{180} (2M)^5 \begin{pmatrix} \mathbf{I}_{11} & \mathbf{I}_{12} & \mathbf{I}_{13} \\ \mathbf{I}_{12} & -\mathbf{I}_{11} & \mathbf{I}_{23} \\ \mathbf{I}_{13} & \mathbf{I}_{23} & \mathbf{0} \end{pmatrix}, \quad (11)$$

where we introduced the four symmetric and trace-free matri-

ces

$$\begin{aligned} \mathbf{I}_{11} &\equiv \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, & \mathbf{I}_{12} &\equiv \begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \\ \mathbf{I}_{13} &\equiv \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & \frac{1}{2} \\ 0 & \frac{1}{2} & 0 \end{pmatrix}, & \mathbf{I}_{23} &\equiv \begin{pmatrix} 0 & 0 & -\frac{1}{2} \\ 0 & 0 & 0 \\ -\frac{1}{2} & 0 & 0 \end{pmatrix}, \end{aligned} \quad (12)$$

and with the understanding that the second pair of indices in λ_{ijkl} indicates one of these 3×3 matrices and the first pair to an element within it. For instance, if the quadrupolar tidal field \mathcal{E}_{ij} is sourced by a static particle of mass $\mu \ll M$ a distance $r \gg M$ away from the black hole, in the direction $\hat{\mathbf{r}}$, then, in the Newtonian limit, the formulas (10)–(12) imply

$$(\delta M_{ij}) \doteq \frac{\chi}{60} (2M)^5 \frac{\mu}{r^3} [\hat{\mathbf{r}} \otimes (\hat{\mathbf{z}} \times \hat{\mathbf{r}}) + (\hat{\mathbf{z}} \times \hat{\mathbf{r}}) \otimes \hat{\mathbf{r}}], \quad (13)$$

where $\hat{\mathbf{z}}$ is the direction of the Kerr black hole spin, \times denotes the cross product and \otimes the tensor product. If the particle lies along the axisymmetry axis of the background Kerr geometry (above one of the poles), then $\hat{\mathbf{z}} \times \hat{\mathbf{r}} = \mathbf{0}$ and δM_{ij} vanishes, in agreement with the vanishing of the TLNs (8) for an axisymmetric tidal perturbation. If the particle lies on the equatorial plane, then $(\delta M_{ij}) \propto \hat{\mathbf{r}} \otimes \hat{\phi} + \hat{\phi} \otimes \hat{\mathbf{r}}$, which is the gravitational analogue of the quadrupole moment tensor obtained by setting four electric charges with alternating signs at the corners of a square centered at the origin and whose four sides are tangent to the directions $\hat{\mathbf{r}}$ and $\hat{\phi}$. Interestingly, the purely imaginary TLNs in (8) and the induced mass quadrupole moment (13) suggest that the black hole tidal bulge is rotated by 45° with respect to the quadrupolar tidal perturbation.

Effective action.—It is well-known in effective field theory (EFT) that finite-size effects correspond to augmenting the effective (spinning) point particle action by nonminimal world-line couplings to curvature [42, 43]. Specifically, the tidally-induced quadrupole moments (10) are expected to correspond to the nonminimal couplings

$$\sum_{i,j,k,l} \int ds \lambda_{ijkl} \mathcal{E}_{ij} \mathcal{E}_{kl} \quad \text{and} \quad \sum_{i,j,k,l} \int ds \lambda_{ijkl} \mathcal{B}_{ij} \mathcal{B}_{kl}. \quad (14)$$

Given the expression (11)–(12) of the quadrupolar TLT, this corresponds to a fifth post-Newtonian (5PN) effect [1] beyond the leading Newtonian motion. The identification of the Kerr TLNs or TLT computed here with analogous quantities at the level of an effective action such as the couplings (14) requires a proper matching calculation. Such a matching was performed in the case of nonspinning black holes [27, 28, 33], but remains to be done for rotating black holes. Notice that vanishing black hole TLNs would be at odds with the naturalness dogma: since there is no apparent enhanced symmetry in the effective action describing black holes with vanishing TLNs, this would imply an unnatural fine-tuning from the EFT point of view [32, 33].

Prospects.—There is the exciting prospect that the planned space-based gravitational-wave observatory LISA [21] might

be able to detect the quadrupolar tidal deformability of spinning black holes which we have here uncovered. One of the main sources for LISA is the radiation-reaction driven inspiral of a stellar-mass compact object of mass μ into a massive black hole of mass $M \gg \mu$. An order-of-magnitude *estimate* of the contribution of the black hole quadrupolar tidal deformability to the total accumulated gravitational-wave phase in such an inspiral is obtained by considering a low velocity approximation (Einstein's quadrupole formula) for the quasi-circular inspiral down to the innermost stable circular orbit of a nonspinning, massive, compact body, yielding [22]

$$\Phi_{\text{tidal}} = -\frac{k_2 q}{96\sqrt{6}} \simeq -4 \times 10^{-3} k_2 q, \quad (15)$$

where $q \equiv M/\mu$ is the large mass ratio of the binary system, and k_2 is the quadrupolar TLN of the massive compact body. For a slowly rotating Kerr black hole, we may use instead the typical value $|k_{22}| \sim \chi/60$ derived from Eqs. (8)–(9). For instance, for a mass ratio $q = 10^7$ and a Kerr black hole spin $\chi = 0.1$, this yields $\Phi_{\text{tidal}} \simeq -80$ rad. By requiring a detectability threshold of $|\Phi_{\text{tidal}}| > 1$ rad, this effect could be detected down to a mass ratio $q \sim 10^5$. It would be interesting to investigate if the tidal effect (15) could be used to perform a null test of the Kerr-ness nature of the massive black holes at the center of galaxies.

We are particularly grateful to Edgardo Franzin for significant contributions to this work. We are grateful to Eric Poisson, Rafael Porto, Adrian Ottewill and Sam Gralla for helpful correspondence, as well as to the whole Capra 23 community for valuable feedback. ALT acknowledges the financial support of the Action Fédératrice PhyFOG and of the Scientific Council of the Paris Observatory. Both authors acknowledge partial financial support by CNPq (Brazil), process no. 310200/2017-2 and 312917/2019-8.

SUPPLEMENTAL MATERIAL

Curvature scalar.—Contracting the Weyl tensor $C_{\alpha\beta\gamma\delta}$ with the vectors ℓ^α and m^α of a Newman-Penrose null tetrad [44] yields the curvature scalar $\psi_0 \equiv C_{\alpha\beta\gamma\delta} \ell^\alpha m^\beta \ell^\gamma m^\delta$. By considering the relation between the gravitational potential (3) and the Weyl tensor in the Newtonian limit where $c \rightarrow \infty$, we find for the Newtonian curvature scalar:

$$\lim_{c \rightarrow \infty} \psi_0 = \sum_{\ell m} \alpha_{\ell m}(t) r^{\ell-2} \left[1 + 2k_\ell \left(\frac{R}{r} \right)^{2\ell+1} \right] {}_2Y_{\ell m}(\theta, \phi), \quad (16)$$

where $\alpha_{\ell m} \equiv [(\ell+2)(\ell+1)/(\ell(\ell-1))]^{1/2} \mathcal{E}_{\ell m}$ and ${}_2Y_{\ell m}(\theta, \phi)$ is a spin-2 spherical harmonic [45]. Similarly to Eq. (3), the term $O(r^{\ell-2})$ in (16) corresponds to the external 2^ℓ -polar tidal perturbation, and the term $O(r^{-\ell-3})$ to the body's response. The formula (16) can easily be generalized to an axisymmetric, spinning Newtonian body, for which $k_\ell \rightarrow k_{\ell m}$.

From now on we shall consider a Kerr black hole of mass M and spin angular momentum per unit mass a . Using advanced

Kerr coordinates $(x^\alpha) = (v, r, \theta, \phi)$, the Kerr metric $\hat{g}_{\alpha\beta}$ is given by [23, 24]

$$\begin{aligned} \hat{g}_{\alpha\beta} dx^\alpha dx^\beta = & - \left(1 - \frac{2Mr}{\Sigma} \right) dv^2 + 2dvdr \\ & - \frac{4Mr}{\Sigma} a \sin^2 \theta dv d\phi + \Sigma d\theta^2 - 2a \sin^2 \theta dr d\phi \\ & + \left(r^2 + a^2 + \frac{2Mr}{\Sigma} a^2 \sin^2 \theta \right) \sin^2 \theta d\phi^2, \end{aligned} \quad (17)$$

where $\Sigma \equiv r^2 + a^2 \cos^2 \theta$. The coordinate radii of the event (+) and Cauchy (−) horizons are $r_\pm = M \pm \sqrt{M^2 - a^2}$. This black hole is embedded in a weak and slowly varying, but otherwise completely generic tidal environment. Slowly varying means that we can neglect time derivatives, but we emphasize that it is nonstatic as it has a *parametric* dependence on v . The tidal field being weak means that we can use the framework of the linearized Einstein equation. This tidal environment is fully characterized by two sets of symmetric and trace-free tensors (angle brackets notation), both defined out of the Weyl tensor $C_{\alpha\beta\gamma\delta}$ and its covariant derivatives (semicolon notation), and evaluated in the black hole's local asymptotic rest frame: the electric-type and magnetic-type tidal fields [25, 30, 46]

$$\mathcal{E}_L(v) \equiv [(\ell-2)!]^{-1} C_{0(i_1|0|i_2|i_3 \dots i_\ell)}(v), \quad (18a)$$

$$\mathcal{B}_L(v) \equiv \frac{3}{2} [(\ell+1)(\ell-2)!]^{-1} \epsilon_{jk(i_1} C_{i_2|0|j k|i_3 \dots i_\ell)}(v), \quad (18b)$$

where ϵ_{ijk} is the totally antisymmetric permutation symbol. The electric-type tidal tensors (18a) are the general relativistic analogues of the Newtonian tidal tensors. The magnetic-type tidal tensors (18b) have no counterpart in Newtonian gravity.

While the curvature Weyl scalar vanishes for the Kerr background (17), Teukolsky [47] managed to derive a decoupled equation for the *perturbed* ψ_0 . He did so with the choice of the Kinnersley tetrad [48] and showed that, furthermore, the solution admits separation by variables:

$$\psi_0 = \sum_{\ell m} z_{\ell m}(v) R_{\ell m}(r) {}_2Y_{\ell m}(\theta, \phi), \quad (19)$$

for zero frequency. The coefficients $z_{\ell m} = \alpha_{\ell m} + i\beta_{\ell m}$ are slowly varying functions of time v , where $\alpha_{\ell m}$ and $\beta_{\ell m}$ are related to the spherical-harmonic modes $\mathcal{E}_{\ell m}$ and $\mathcal{B}_{\ell m}$ of the electric-type and magnetic-type tidal moments (18) [49]. For instance, at the leading quadrupolar order $\alpha_{2m} = \sqrt{6} \mathcal{E}_{2m}$ and $\beta_{2m} = \sqrt{6} \mathcal{B}_{2m}$. The radial factor $R_{\ell m}$ in the mode decomposition (19) satisfies the zero-frequency radial Teukolsky equation [50]

$$\begin{aligned} x(x+1)R''_{\ell m}(x) + (6x+3+2im\gamma)R'_{\ell m}(x) \\ + \left[4im\gamma \frac{2x+1}{x(x+1)} - (\ell+3)(\ell-2) \right] R_{\ell m}(x) = 0, \end{aligned} \quad (20)$$

where $x \equiv (r-r_+)/ (r_+ - r_-)$ and $\gamma \equiv a/(r_+ - r_-)$. While the physical regime requires $\ell \in \mathbb{L} \equiv \mathbb{Z}^+ \setminus \{1\}$, we shall temporarily carry out the calculations for a *generic* $\ell \in \mathbb{R}$, for the following reason. In the Newtonian limit, the identification of the

growing tidal perturbation and the associated decaying body's linear response in the gravitational potential U in (3) is unambiguous. This is thanks to the linearity of the Poisson equation satisfied by U and the Euclidean nature of 3-space, which provides a unique and unambiguous notion of distance. In general relativity, however, the identification of the tidal and response contributions is quite subtle. In particular, Gralla [51] pointed out that the value of the TLNs depends on the choice of radial coordinate. Moreover, given a choice of coordinates and thus of response contribution, the tidal contribution remains ambiguous for $\ell \in \mathbb{L}$ because one can always add a multiple of the decaying (subdominant) response contribution to the growing (dominant) tidal contribution [25, 41]. We henceforth consider Eq. (20) with $\ell \in \mathbb{R}$, since this allows us to naturally resolve the two ambiguities, as we shall next see.

Tidal/response solutions.—We obtain the specific physical solution by requiring that the curvature scalar ψ_0 is smooth at the event horizon $r = r_+$ when written in the Hartle-Hawking tetrad [52], which is smooth there. The corresponding physical radial factor $R_{\ell m}(x)$, which is a solution of Eq. (20), is thus uniquely determined (up to a normalization) and can be expressed as

$$R_{\ell m}(x) = (r_+ - r_-)^{\ell-2} [\hat{R}_{\ell m}^{\text{tidal}}(x) + 2k_{\ell m} \hat{R}_{\ell m}^{\text{resp}}(x)], \quad (21)$$

where

$$\hat{R}_{\ell m}^{\text{tidal}} \equiv x^{\ell-2} \frac{F(-\ell-2, -\ell-2im\gamma, -2\ell; -1/x)}{(1+1/x)^2}, \quad (22a)$$

$$\hat{R}_{\ell m}^{\text{resp}} \equiv x^{-\ell-3} \frac{F(\ell-1, \ell+1-2im\gamma, 2\ell+2; -1/x)}{(1+1/x)^2}, \quad (22b)$$

with $F(a, b, c; z)$ the Gaussian hypergeometric function and

$$k_{\ell m} = -\frac{i}{4\pi} \sinh(2\pi m\gamma) |\Gamma(\ell+1+2im\gamma)|^2 \frac{(\ell-2)!(\ell+2)!}{(2\ell)!(2\ell+1)!}. \quad (23)$$

Here, $\hat{R}_{\ell m}^{\text{tidal}}$ and $\hat{R}_{\ell m}^{\text{resp}}$ are two linearly independent solutions of the radial Teukolsky equation (20), which are exchanged under the substitution $\ell \rightarrow -\ell-1$. While, as said, the physical ψ_0 (constructed from $R_{\ell m}$) in the Hartle-Hawking tetrad should be—and is—smooth at the Kerr horizon, the contributions to it constructed from $\hat{R}_{\ell m}^{\text{tidal}}$ and $\hat{R}_{\ell m}^{\text{resp}}$ are both regular but, for $m\gamma \neq 0$, nonsmooth there. They need not be smooth on the horizon since neither of them is separately physically accessible.

The solutions $\hat{R}_{\ell m}^{\text{tidal}}$ and $\hat{R}_{\ell m}^{\text{resp}}$ in the formulas (21) and (22) are specified uniquely and unambiguously by the fact that they carry separately the two linearly independent asymptotic behaviours $x^{\ell-2}$ and $x^{-\ell-3}$ as $r \rightarrow \infty$ for $\ell \in \mathbb{R}$. Crucially, the *analytic continuation* to $\ell \in \mathbb{R}$ that we have carried out allows us to ascertain whether or not a given solution which contains the asymptotic behaviour $x^{\ell-2}$ also contains the asymptotic behaviour $x^{-\ell-3}$ within it since, for $\ell \in \mathbb{R}$, neither asymptotic behaviour can be considered to be dominant/subdominant (this is further exemplified below (27) in the case of the Hertz potential). For $\ell \in \mathbb{L}$, the asymptotic behaviours $x^{\ell-2}$ and $x^{-\ell-3}$ physically correspond to the Newtonian tidal field and body's linear response, respectively.

As pointed out in [41, 53] and shown in [28], analytic continuation is a natural method to obtain a unique and unambiguous tidal/response split that is the general relativistic extension of the unique and unambiguous Newtonian tidal/response split in the rotating version of (3) and (16). More precisely, the radial factor (21) reads

$$R_{\ell m} = r^{\ell-2} (1 + \dots) + 2k_{\ell m} (r_+ - r_-)^{2\ell+1} r^{-\ell-3} (1 + \dots), \quad (24)$$

where the dots denote relativistic corrections to the rotating version of the Newtonian (16). We determined the overall normalization of $R_{\ell m}$ so that the coefficient of the asymptotic term $r^{\ell-2}$ as $r \rightarrow \infty$ times $z_{\ell m}$ is in agreement with the asymptotic form of ψ_0 in terms of the spherical-harmonic modes $\mathcal{E}_{\ell m}$ and $\mathcal{B}_{\ell m}$ of the tidal moments [49].

Comparing the relativistic formula (24) with the Newtonian formula (16), the constant coefficients (23) can be interpreted as the Newtonian TLNs of a Kerr black hole. In the particular cases of a Schwarzschild black hole ($a = 0$) and axisymmetric ($m = 0$) tidal perturbations of Kerr, those coefficients are identically zero for $\ell \in \mathbb{L}$, and so the response vanishes, yielding zero relativistic black hole TLNs (9) in these cases. This is in agreement with results in the literature [25–30]. However, the response $2k_{\ell m} \hat{R}_{\ell m}^{\text{resp}}$ is *not* zero in general in Kerr. In particular, for a small spin $\chi = a/M$, the quadrupolar coefficients (23) reduce to $k_{2m} \doteq -im\chi/120$, in agreement with the nonvanishing quadrupolar TLNs (8)–(9).

Hertz potential.—To derive the response part of the metric perturbation from the Weyl scalar (19) with Eqs. (21)–(23), we apply the Hertz potential formalism [54–58]. The first step consists in obtaining a Hertz potential Ψ by solving a fourth-order linear partial differential equation with the Weyl scalar ψ_0 as the source. This can be done mode by mode [59],

$$\bar{\Psi} = \sum_{\ell m} z_{\ell m}(v) G_{\ell m}(r) {}_2Y_{\ell m}(\theta, \phi), \quad (25)$$

where the radial function of the (complex-conjugated) Hertz potential inherits the tidal-response split of the Weyl scalar, according to

$$G_{\ell m}(x) = g_{\ell} (r_+ - r_-)^{\ell+2} [\hat{G}_{\ell m}^{\text{tidal}}(x) + 2k_{\ell m} \hat{G}_{\ell m}^{\text{resp}}(x)], \quad (26)$$

with $g_{\ell} \equiv 2\Gamma(\ell+3)/(\Gamma(\ell+1)[(\ell+2)(\ell+1)\ell(\ell-1)]^2$ and

$$\hat{G}_{\ell m}^{\text{tidal}} = x^{\ell} (1+x)^2 F(-\ell+2, -\ell-2im\gamma, -2\ell; -1/x), \quad (27a)$$

$$\hat{G}_{\ell m}^{\text{resp}} = \frac{(1+x)^2}{x^{\ell+1}} F(\ell+3, \ell+1-2im\gamma, 2\ell+2; -1/x). \quad (27b)$$

The decomposition (26) into tidal and response contributions in the (complex-conjugated) Hertz potential inherited from the Weyl scalar in Eq. (21) has a delicate structure: As one takes $\ell \in \mathbb{L}$, $G_{\ell m}$ becomes a polynomial in r of order $\ell+2$ and so one might be tempted to conclude that the response vanishes in the physical regime $\ell \in \mathbb{L}$. However, analytic continuation to $\ell \in \mathbb{R}$ shows that this is a mere illusion since such polynomial behaviour arises from the fact that $\hat{G}_{\ell m}^{\text{tidal}}$ contains this polynomial plus an infinite series in $1/x$ which exactly cancels out with $2k_{\ell m} \hat{G}_{\ell m}^{\text{resp}}$ when taking $\ell \in \mathbb{L}$.

Metric reconstruction.—Once the response contribution to the Hertz potential is known, one can obtain the corresponding metric perturbation $h_{\alpha\beta}^{\text{resp}}$ by applying a certain second order differential operator to it, e.g. [54, 60, 61]. We carried out the calculation explicitly up to *linear* order in the dimensionless spin parameter $\chi = a/M$. In the so-called ingoing radiation gauge, specified by the conditions $\ell^\alpha h_{\alpha\beta}^{\text{resp}} = 0$ and $\hat{g}^{\alpha\beta} h_{\alpha\beta}^{\text{resp}} = 0$, we obtain

$$h_{\alpha\beta}^{\text{resp}} = \ell_\alpha \ell_\beta h_{nn}^{\text{resp}} + 2\Re [m_\alpha m_\beta h_{\bar{m}\bar{m}}^{\text{resp}} - 2\ell_{(\alpha} m_{\beta)} h_{n\bar{m}}^{\text{resp}}], \quad (28)$$

with

$$h_{nn}^{\text{resp}} \doteq -2\chi \sum_{\ell m} C_\ell Q_\ell \Re(im \bar{z}_{\ell m} \bar{Y}_{\ell m}), \quad (29a)$$

$$h_{n\bar{m}}^{\text{resp}} \doteq -\sqrt{2}\chi \sum_{\ell m} im C_\ell \sqrt{\frac{(\ell-1)!}{(\ell+1)!}} r Q'_\ell \bar{z}_{\ell m} \bar{Y}_{\ell m}, \quad (29b)$$

$$h_{\bar{m}\bar{m}}^{\text{resp}} \doteq -2\chi \sum_{\ell m} im C_\ell \sqrt{\frac{(\ell-2)!}{(\ell+2)!}} (r^2 Q''_\ell + 2r Q'_\ell - 2Q_\ell) \bar{z}_{\ell m} \bar{Y}_{\ell m}, \quad (29c)$$

where we used $C_\ell \equiv \ell![(\ell-2)!]^{3/2}/((2\ell)![(\ell+2)!]^{1/2})(2M)^\ell$ and $Q_\ell(r) \equiv (r/M-2)^2 Q''_\ell(r/M-1)$, with Q_ℓ the Legendre function of the second kind. The ℓ modes of the tetrad components (29) scale as $O(r^{-\ell-1})$.

Multipole moments.—All our results so far were valid for a tidal field which is slowly varying but is allowed to possess a parametric dependence on time ν . From now on we neglect the slow parametric evolution of the coefficients $z_{\ell m}(\nu)$ in the metric perturbation (29). Hence, the metric $g_{\alpha\beta} = \hat{g}_{\alpha\beta} + h_{\alpha\beta}^{\text{resp}}$ is a stationary, asymptotically flat, vacuum solution of the linearized Einstein equation, and its Geroch-Hansen [39, 40] multipole moments are well defined.

More precisely, we compute the norm λ and scalar twist ω associated with the timelike Killing field ξ^α of the perturbed Kerr geometry $g_{\alpha\beta}$, and the 3-metric $\gamma_{\alpha\beta} = \lambda g_{\alpha\beta} + \xi_\alpha \xi_\beta$ on any spacelike hypersurface orthogonal to ξ^α . We then perform a conformal transformation of $\gamma_{\alpha\beta}$, with a choice of conformal factor Ω that ensures that the mass dipole moment vanishes, thus taking the multipole moments about the center-of-mass of the perturbed black hole. The mass-type and current-type quadrupole moments M_{ij} and S_{ij} are given by the formula

$$M_{ij} + iS_{ij} = \frac{1}{3} \left(\tilde{D}_{\langle i} \tilde{D}_{j \rangle} \tilde{\Phi} - \frac{1}{2} \tilde{R}_{\langle ij \rangle} \tilde{\Phi} \right) \Big|_\Lambda, \quad (30)$$

where the scalar field $\tilde{\Phi} \equiv (\Phi_M + i\Phi_S)/\sqrt{\Omega}$ is a conformally rescaled linear combination of the mass-type and current-type potentials $\Phi_M \doteq 1/(4\lambda) - \lambda/4$ and $\Phi_S = \omega/(2\lambda)$, while \tilde{D}_α is the covariant derivative compatible with the conformal metric $\tilde{\gamma}_{\alpha\beta} = \Omega^2 \gamma_{\alpha\beta}$ and $\tilde{R}_{\alpha\beta}$ is the associated Ricci tensor. The right-hand side of Eq. (30) is evaluated at spatial infinity, which corresponds to the point Λ of the extended 3-manifold. The results (8) are obtained by taking the spherical-harmonic modes of (30) and comparing to the formulas (7).

-
- [1] E. Poisson and C. M. Will, *Gravity: Newtonian, post-Newtonian, relativistic* (Cambridge University Press, Cambridge, 2014).
 - [2] T. Hinderer, *Astrophys. J.* **677**, 1216 (2008), *Erratum: Astrophys. J.* **697**, 964 (2009), [arXiv:0711.2420 \[astro-ph\]](#).
 - [3] T. Hinderer, B. D. Lackey, R. N. Lang, and J. S. Read, *Phys. Rev. D* **81**, 123016 (2010), [arXiv:0911.3535 \[astro-ph.HE\]](#).
 - [4] S. Postnikov, M. Prakash, and J. M. Lattimer, *Phys. Rev. D* **82**, 024016 (2010), [arXiv:1004.5098 \[astro-ph.SR\]](#).
 - [5] A. E. H. Love, *Nature* **80**, 252 (1909).
 - [6] A. E. H. Love, *Some problems of geodynamics* (Cornell University Library, Ithaca, 1911).
 - [7] L. Iess *et al.*, *Science* **337**, 457 (2012).
 - [8] V. Lainey, *Celest. Mech. Dyn. Astr.* **126**, 145 (2016), [arXiv:1604.04184 \[astro-ph.EP\]](#).
 - [9] B. Akisnami *et al.*, *Astron. Astrophys.* **621**, A117 (2019), [arXiv:1812.04538 \[astro-ph.EP\]](#).
 - [10] P. Auclair-Desrotour, J. Leconte, E. Bolmont, and S. Mathis, *Astron. Astrophys.* **629**, A132 (2019), [arXiv:1907.06451 \[astro-ph.EP\]](#).
 - [11] E. Bolmont *et al.*, *Astron. Astrophys.* **635**, A117 (2020), [arXiv:2002.02015 \[astro-ph.EP\]](#).
 - [12] V. Cardoso, E. Franzin, A. Maselli, P. Pani, and G. Raposo, *Phys. Rev. D* **95**, 084014 (2017), [arXiv:1701.01116 \[gr-qc\]](#).
 - [13] B. P. Abbott *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), *Phys. Rev. Lett.* **119**, 161101 (2017), [arXiv:1710.05832 \[gr-qc\]](#).
 - [14] B. P. Abbott *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), *Astrophys. J. Lett.* **892**, L3 (2020), [arXiv:2001.01761 \[astro-ph.HE\]](#).
 - [15] B. P. Abbott *et al.* (LIGO Scientific Collaboration and Virgo Collaboration), *Phys. Rev. Lett.* **121**, 161101 (2018), [arXiv:1805.11581 \[gr-qc\]](#).
 - [16] E. R. Most, L. R. Weih, L. Rezzolla, and J. Schaffner-Bielich, *Phys. Rev. Lett.* **120**, 261103 (2018), [arXiv:1803.00549 \[gr-qc\]](#).
 - [17] S. De *et al.*, *Phys. Rev. Lett.* **121**, 091102 (2018), [arXiv:1804.08583 \[astro-ph.HE\]](#).
 - [18] K. Chatziioannou, (2020), [arXiv:2006.03168 \[gr-qc\]](#).
 - [19] N. Yunes and K. Yagi, *Science* **341**, 365 (2013), [arXiv:1302.4499 \[gr-qc\]](#).
 - [20] T. K. Chan, A. P. O. Chan, and P. T. Leung, *Phys. Rev. D* **91**, 044017 (2015), [arXiv:1411.7141 \[astro-ph.SR\]](#).
 - [21] P. Amaro-Seoane *et al.*, “Laser Interferometer Space Antenna,” (2017), [arXiv:1702.00786 \[astro-ph.IM\]](#).
 - [22] P. Pani and A. Maselli, *Int. J. Mod. Phys. D* **28**, 1944001 (2019), [arXiv:1905.03947 \[gr-qc\]](#).
 - [23] R. P. Kerr, *Phys. Rev. Lett.* **11**, 237 (1963).
 - [24] S. Chandrasekhar, *The mathematical theory of black holes* (Oxford University Press, Oxford, 1983).
 - [25] T. Binnington and E. Poisson, *Phys. Rev. D* **80**, 084018 (2009), [arXiv:0906.1366 \[gr-qc\]](#).
 - [26] T. Damour and A. Nagar, *Phys. Rev. D* **80**, 084035 (2009), [arXiv:0906.0096 \[gr-qc\]](#).
 - [27] B. Kol and M. Smolkin, *JHEP* **1202**, 010 (2012), [arXiv:1110.3764 \[hep-th\]](#).
 - [28] S. Chakrabarti, T. Delsate, and J. Steinhoff, (2013), [arXiv:1304.2228 \[gr-qc\]](#).
 - [29] N. Gürlebeck, *Phys. Rev. Lett.* **114**, 151102 (2015), [arXiv:1503.03240 \[gr-qc\]](#).
 - [30] P. Pani, L. Gualtieri, A. Maselli, and V. Ferrari, *Phys. Rev. D* **92**, 024010 (2015), [arXiv:1503.07365 \[gr-qc\]](#).

- [31] P. Landry and E. Poisson, Phys. Rev. D **91**, 104018 (2015), [arXiv:1503.07366 \[gr-qc\]](#).
- [32] R. A. Porto, Phys. Rept. **663**, 1 (2016), [arXiv:1601.04914 \[gr-qc\]](#).
- [33] R. A. Porto, Fortschr. Phys. **64**, 723 (2016), [arXiv:1606.08895 \[gr-qc\]](#).
- [34] S. Cai and K.-D. Wang, (2019), [arXiv:1906.06850 \[hep-th\]](#).
- [35] V. Cardoso, L. Gualtieri, and C. J. Moore, Phys. Rev. D **100**, 124037 (2019), [arXiv:1910.09557 \[gr-qc\]](#).
- [36] V. Cardoso and F. Duque, Phys. Rev. D **101**, 064028 (2020), [arXiv:1912.07616 \[gr-qc\]](#).
- [37] H. S. Tan, (2020), [arXiv:2001.00403 \[gr-qc\]](#).
- [38] C. Chirenti, C. Posada, and V. Guedes, (2020), [arXiv:2005.10794 \[gr-qc\]](#).
- [39] R. P. Geroch, J. Math. Phys. **11**, 2580 (1970).
- [40] R. O. Hansen, J. Math. Phys. **15**, 46 (1974).
- [41] P. Pani, L. Gualtieri, and V. Ferrari, Phys. Rev. D **92**, 124003 (2015), [arXiv:1509.02171 \[gr-qc\]](#).
- [42] T. Damour and G. Esposito-Farèse, Phys. Rev. D **53**, 5541 (1996), [arXiv:gr-qc/9506063](#).
- [43] W. D. Goldberger and I. Z. Rothstein, Phys. Rev. D **73**, 104029 (2006), [arXiv:hep-th/0409156](#).
- [44] E. Newman and R. Penrose, J. Math. Phys. **3**, 566 (1962), *Erratum*: J. Math. Phys. **4**, 998 (1963).
- [45] E. T. Newman and R. Penrose, J. Math. Phys. **7**, 863 (1966).
- [46] X.-H. Zhang, Phys. Rev. D **34**, 991 (1986).
- [47] S. A. Teukolsky, Astrophys. J. **185**, 635 (1973).
- [48] W. Kinnersley, J. Math. Phys. **10**, 1195 (1969).
- [49] K. Chatziioannou, E. Poisson, and N. Yunes, Phys. Rev. D **87**, 044022 (2013), [arXiv:1211.1686 \[gr-qc\]](#).
- [50] S. A. Teukolsky and W. H. Press, Astrophys. J. **193**, 443 (1974).
- [51] S. E. Gralla, Class. Quant. Grav. **35**, 085002 (2018), [arXiv:1710.11096 \[gr-qc\]](#).
- [52] S. W. Hawking and J. B. Hartle, Commun. Math. Phys. **27**, 283 (1972).
- [53] J. Steinhoff, Fund. Theor. Phys. **179**, 615 (2015), [arXiv:1412.3251 \[gr-qc\]](#).
- [54] P. L. Chrzanowski, Phys. Rev. D **11**, 2042 (1975).
- [55] J. M. Cohen and L. S. Kegeles, Phys. Rev. D **10**, 1070 (1974).
- [56] L. S. Kegeles and J. M. Cohen, Phys. Rev. D **19**, 1641 (1979).
- [57] J. M. Stewart, Proc. R. Soc. Lond. A **367**, 527 (1979).
- [58] R. M. Wald, Phys. Rev. Lett. **41**, 203 (1978).
- [59] A. Ori, Phys. Rev. D **67**, 124010 (2003), [arXiv:gr-qc/0207045](#).
- [60] C. Lousto and B. Whiting, Phys. Rev. D **66**, 024026 (2002), [arXiv:gr-qc/0203061](#).
- [61] N. Yunes and J. A. González, Phys. Rev. D **73**, 024010 (2006), *Erratum*: Phys. Rev. D **89**, 089902 (2014), [arXiv:gr-qc/0510076](#).