

SOME HOMOLOGICAL PROPERTIES OF CATEGORY $\mathcal{O}$, V

HANKYUNG KO, VOLODYMYR MAZORCHUK AND RAFAEL MRĐEN

ABSTRACT. We compute projective dimension of translated simple modules in the regular block of the BGG category $\mathcal{O}$ in terms of Kazhdan-Lusztig combinatorics. This allows us to determine which projectives can appear at the last step of a minimal projective resolution for a translated simple module, confirming a conjecture by Johan Kåhrström. We also derive some inequalities, in terms of Lusztig's a -function, for possible degrees in which the top (or socle) of a translated simple module can live. Finally, we relate Kostant's problem with decomposability and isomorphism of translated simple modules, addressing yet another conjecture by Johan Kåhrström.

CONTENTS

1. Introduction, motivation and description of the results	1
2. A zoo of questions about $\mathcal{O}_0$	4
3. Kazhdan-Lusztig combinatorics	5
4. Proof of Theorem 2.2	8
5. KMM vs KM	9
6. Extensions between translated simple modules	10
7. K and KMM	13
8. K vs. KM and Kh	16
9. Further discussion and speculation on KMM	25
10. Small rank results	27
References	28

1. INTRODUCTION, MOTIVATION AND DESCRIPTION OF THE RESULTS

1.1. **Setup.** Let $\mathfrak{g}$ be a semi-simple complex Lie algebra with a fixed triangular decomposition

$$\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$$

and $U(\mathfrak{g})$ be the universal enveloping algebra of $\mathfrak{g}$. Let $\mathcal{O}$ denote the associated Bernstein-Gelfand-Gelfand (BGG) category $\mathcal{O}$, see [BGG, Hu]. Let, further, $\mathcal{O}_0$ denote the *principal block* $\mathcal{O}$, that is, the indecomposable direct summand containing the trivial $\mathfrak{g}$ -module.

Let W be the Weyl group of $\mathfrak{g}$. It acts naturally on $\mathfrak{h}^*$ via $(w, \lambda) \mapsto w(\lambda)$. We consider the *dot-action* $(w, \lambda) \mapsto w \cdot \lambda$ of W which is obtained by shifting the usual action by the half of the sum of all positive roots.

The category $\mathcal{O}_0$ is equivalent to $A\text{-mod}$, for some finite-dimensional associative basic algebra A , unique up to isomorphism. Simple objects in $\mathcal{O}_0$ are exactly the (pairwise non-isomorphic) simple highest weight modules $L_w := L(w \cdot 0)$ of highest weight $w \cdot 0$, for $w \in W$. The category $\mathcal{O}_0$ is equipped with the action of the monoidal category $\mathcal{P}$ of *projective functors*, as defined in [BG]. Up to isomorphism, indecomposable projective functors are also in bijection with the elements in W , where the indecomposable projective functor θ_w , for $w \in W$, is normalized such that it sends the projective cover P_e of L_e to the projective cover P_w of L_w .

The algebra A is Koszul by [So1], in particular, it admits a positive $\mathbb{Z}$ -grading. Denote by $\mathcal{O}_0^{\mathbb{Z}}$ the category of finite dimensional $\mathbb{Z}$ -graded A -modules with morphisms being homogeneous homomorphisms of degree zero, see e.g., [St].

1.2. Motivation. The first major motivation for the present paper is the following:

Conjecture 1.1 ([KM]). *Assume that $\mathfrak{g}$ is of type A . Then, for $x, y \in W$, the module $\theta_x L_y$ is either indecomposable or zero.*

Various approaches to Conjecture 1.1 were considered in [KM] and [CMZ]. In the latter paper, the conjecture was confirmed in the cases $\mathfrak{g} = \mathfrak{sl}_n$, where $n = 2, 3, 4, 5, 6$. For other values of n , a number of special results are obtained in [KM] and [CMZ].

The second major motivation for the present paper is the so-called *Kostant's problem*, as popularized in [Jo], the content of which is to determine all $w \in W$ for which the universal enveloping algebra surjects onto the space of linear endomorphism of L_w that are locally finite with respect to the adjoint action (we will denote this property by $\mathbf{K}(w)$, see § 2.3 for details). This problem was studied in [Ma1, Ma3, MS2, Kh, KhM], see also the references therein. In March 2019, the second author received an email from Johan K  hrstr  m with the following conjecture (based on extensive computer computation).

Conjecture 1.2 (J. K  hrstr  m). *For a Duflo element $d \in W$, the following assertions are equivalent:*

- (a) $\mathbf{K}(d)$.
- (b) $\theta_x L_d \not\cong \theta_y L_d$, for all $x \neq y \in W$ such that $\theta_x L_d \neq 0$ and $\theta_y L_d \neq 0$.
- (c) For all $x \neq y \in W$ such that $\theta_x L_d \neq 0$ and $\theta_y L_d \neq 0$, there exists $z \in W$ and $i \in \mathbb{Z}$ such that $[\theta_x L_d : L_z \langle i \rangle] \neq [\theta_y L_d : L_z \langle i \rangle]$ in $\mathcal{O}_0^{\mathbb{Z}}$.
- (d) For all $x \neq y \in W$ such that $\theta_x L_d \neq 0$ and $\theta_y L_d \neq 0$, there is $z \in W$ such that $[\theta_x L_d : L_z] \neq [\theta_y L_d : L_z]$ in $\mathcal{O}_0$.

The final piece of motivation for the present paper is the problem to determine the projective dimension for all modules of the form $\theta_x L_y \in \mathcal{O}_0$, as formulated in [Ma3, Problem 24]. In connection to this problem, the email from Johan K  hrstr  m mentioned above contained the following conjecture (also based on extensive computer computation):

Conjecture 1.3 (J. K  hrstr  m). *Let $x, y \in W$ and k be the projective dimension of $\theta_x L_y$. Assume that $z \in W$ is such that $\text{Ext}^k(\theta_x L_y, L_z) \neq 0$. Then z and x belong to the same Kazhdan-Lusztig left cell.*

1.3. Description of the results. The first main result of the present paper, see Proposition 6.2, Formula (21), Theorem 6.1 and Corollary 6.3, is:

Theorem A. *Conjecture 1.3 is true. Moreover, the projective dimension of $\theta_x L_y$, for $x, y \in W$ such that $\theta_x L_y \neq 0$, is given by*

$$\mathbf{a}(w_0 x) + \mathbf{b}(y^{-1} w_0, w_0 x^{-1}),$$

where w_0 is the longest element in W , $\mathbf{a}$ denotes Lusztig's $\mathbf{a}$ -function, and $\mathbf{b}(u, v)$ is defined as the maximal degree shift of the composition factors in $\theta_u L_v$ (see Subsection 6.3).

Note that both $\mathbf{a}(w_0 x)$ and $\mathbf{b}(y^{-1} w_0, w_0 x^{-1})$ are “combinatorial” in the sense that they are defined explicitly in terms of Kazhdan-Lusztig polynomials.

Our second main result is the following statement, see Theorem 8.21, which combines ingredients of Conjecture 1.1 with parts of Conjecture 1.2:

Theorem B. *For $y \in W$, the assertion $\mathbf{K}(y)$ is true if and only if both of the following conditions hold.*

- (i) *for all $x \neq z$, we have $\theta_x L_y \not\cong \theta_z L_y$ whenever nonzero (we refer to this property as $\mathbf{Kh}(y)$);*
- (ii) *for each $x \in W$, the module $\theta_x L_y$ is either indecomposable or zero (we refer to this property as $\mathbf{KM}(*, y)$).*

In particular, if Conjecture 1.1 is true, then (a) $\Leftrightarrow$ (b) in Conjecture 1.2 holds in type A.

In Theorem 2.2 we obtain some bounds (given in terms of the $\mathbf{a}$ -function) on the degrees of simple constituents in the top (or socle) of $\theta_x L_y$. When the bound prescribed by Theorem 2.2 is achieved, an interesting question is the multiplicity of the corresponding simple constituent in the top. We formulate a condition on this multiplicity that we call $\mathbf{KMM}$, see (3), which would imply Conjecture 1.1 in type A.

We also relate Kostant's problem for different simple highest weight modules from the same Kazhdan-Lusztig left cell, see Corollary 8.19 and Proposition 8.26 for the condition (e):

Theorem C. *Let $d \in W$ be a Duflo element and let $\mathcal{L}$ be the Kazhdan-Lusztig left cell containing d .*

- (i) *If $\mathbf{K}(d)$ is not true, then $\mathbf{K}(y)$ is not true, for all $y \in \mathcal{L}$;*
- (ii) *If $\mathbf{K}(d)$ is true, then, for each $y \in \mathcal{L}$, the following conditions are equivalent:*
 - (a) $\mathbf{K}(y)$;
 - (b) $\theta_y L_{y^{-1}} \cong \theta_d L_d$;
 - (c) $t_y t_{y^{-1}} = t_d$ in the asymptotic ring for W (see §3.4);
 - (d) $\mathbf{KM}(y, y^{-1})$;
 - (e) *if, furthermore, $\mathfrak{g}$ is of classical type: the $\mathbf{H}$ -cell of y and the $\mathbf{H}$ -cell of d contain the same number of elements (here, an $\mathbf{H}$ -cell is the intersection of a Kazhdan-Lusztig right cell and a Kazhdan-Lusztig left cells, see Subsection 3.3).*

Theorem B and Theorem C give a conjectural answer to Kostant's problem in terms of Kazhdan-Lusztig combinatorics as follows (see Corollary 8.23).

Corollary 1.4. *Suppose (a) $\Leftrightarrow$ (c) in Conjecture 1.2 is true for W . Then, for each $y \in W$, the condition $\mathbf{K}(y)$ is equivalent to the conjunction of the following conditions:*

- (i) $t_y t_{y^{-1}} = t_d$ in the asymptotic ring for W ;
- (ii) *For the Duflo element $d \sim_L y$ and $x, x' \leq_R d$, if $h_{z,x,d} = h_{z,x',d}$, for all $z \in W$, then $x = x'$.*

Here $h_{z,x,d} \in \mathbb{Z}[v, v^{-1}]$ is the structure coefficient for the Kazhdan-Lusztig basis (see (6)).

In the last section, we use Theorem B, Theorem C, Kazhdan-Lusztig combinatorics, and some other results in the paper to determine $\mathbf{K}(y)$, as well as $\mathbf{KM}(y)$ and $\mathbf{Kh}(y)$, for all $y \in W$ in a number of small rank cases. The result confirms Conjecture 1.2 in type A_n for $n \leq 5$ and in type BCD in rank ≤ 4 . In particular, we completely solve Kostant's problem in type A_5 , see Corollary 10.1. This question was considered before in [KhM, Kh], where it was solved completely for A_n , where $n \leq 4$, and a partial answer for A_5 was given.

Acknowledgments. This research was partially supported by the Swedish Research Council, Göran Gustafsson Stiftelse and Vergstiftelsen. The third author was also partially supported by the QuantiXLie Center of Excellence grant no. KK.01.1.1.01.0004 funded by the European Regional Development Fund.

We are especially indebted to Johan Kåhrström who shared with us his ideas which started the work on this paper.

2. A ZOO OF QUESTIONS ABOUT $\mathcal{O}_0$

In this sections we both recall some classical open problems and questions about $\mathcal{O}_0$ and propose some new ones. In the rest of the paper we look deeper into connection between these problems and questions.

For any function $\mathbf{F} : W \rightarrow \{\text{false}, \text{true}\}$, we write $\mathbf{F}(*)$ for the conjunction of all $\mathbf{F}(w)$, where $w \in W$, and similarly for functions of several variables. We also write $\mathbf{F}(*_d)$ for the conjunction of all $\mathbf{F}(w)$, where $w \in W$ is a Duflo element.

2.1. Indecomposability of translation of simple modules. For $x, y \in W$, we denote by $\mathbf{KM}(x, y)$ the statement “the module $\theta_x L_y$ is either indecomposable or zero”. The following problem is still open:

Problem 2.1. *Determine all $x, y \in W$, for which $\mathbf{KM}(x, y)$ is true.*

Conjecture 1.1 asserts that $\mathbf{KM}(x, y)$ is always true in type A . We note that $\theta_x L_y$ is non-zero if and only if $x^{-1} \leq_L y$, where $\leq_L$ denotes the Kazhdan-Lusztig left order on W from [KL], see, for example, [KM, Formula (1)] and the references preceding this formula. If $\mathfrak{g}$ is of type B_2 and $1, 2$ are the two simple reflections in W , then $\mathbf{KM}(12, 21)$ is known to be false, see [KM, Subsection 5.1].

2.2. Graded simple tops of translated simple modules. Denote by $\langle 1 \rangle$ the grading shift on $\mathcal{O}_0^{\mathbb{Z}}$ normalized such that it maps degree one to degree zero. Fix standard graded lifts L_w , where $w \in W$, of simple modules concentrated in degree zero. Fix standard graded lifts P_w , where $w \in W$, of indecomposable projective modules such that their tops are concentrated in degree zero. According to [St], each θ_w , where $w \in W$, also has a graded lift (unique up to isomorphism and shift of grading), which we normalize such that $\theta_w P_e \cong P_w$ holds as graded modules. Morphisms in $\mathcal{O}^{\mathbb{Z}}$ is denoted by hom .

We will use Lusztig's $\mathbf{a}$ -function $\mathbf{a} : W \rightarrow \mathbb{Z}_{\geq 0}$ from [Lu1, Lu2] (see §3.3 for details). One of our principal observations in this paper is the following:

Theorem 2.2. *For $x, y, z \in W$ and $i \in \mathbb{Z}$, the condition*

$$(1) \quad \text{hom}(\theta_x L_y, L_z \langle i \rangle) \neq 0 \quad \text{implies} \quad i \geq \mathbf{a}(x).$$

Theorem 2.2 is proved in Section 4.

Recall that each Kazhdan-Lusztig left (and right) cell contains a unique distinguished involution, called the *Duflo element*. In type A , all involutions are Duflo elements. We propose the following:

Conjecture 2.3. *Let $d, y \in W$ be such that d is a Duflo element. Let, further, M be an indecomposable summand of $\theta_d L_y$. Then*

$$(2) \quad \dim \operatorname{hom}(M, L_y \langle \mathbf{a}(d) \rangle) = 1.$$

Let $d \in W$ be a Duflo element. Then there is a (unique up to a non-zero scalar and homogeneous of degree zero) non-zero natural transformation $\zeta : \theta_d \rightarrow \theta_e \langle \mathbf{a}(d) \rangle$, see [MM3, Section 7]. If $y \in W$ is such that $\theta_d L_y \neq 0$, then ζ_{L_y} is non-zero because the cokernel of ζ is killed by θ_d , see [MM1, Proposition 17], and hence this cokernel must annihilate L_y . Therefore, $\dim \operatorname{hom}(\theta_d L_y, L_y \langle \mathbf{a}(d) \rangle) \geq 1$.

For fixed $x, y \in W$, we denote by $\mathbf{KMM}(x, y)$ the property

$$(3) \quad \dim \operatorname{hom}(\theta_x L_y, L_y \langle \mathbf{a}(x) \rangle) \leq 1.$$

2.3. Kostant's problem. For any $\mathfrak{g}$ -modules M and N , the vector space $\operatorname{Hom}_{\mathbb{C}}(M, N)$ has the natural structure of a $U(\mathfrak{g})$ - $U(\mathfrak{g})$ -bimodule. The subspace $\mathcal{L}(M, N)$ consisting of all vectors of $\operatorname{Hom}_{\mathbb{C}}(M, N)$, the adjoint action of $\mathfrak{g}$ on which is locally finite, is a $U(\mathfrak{g})$ - $U(\mathfrak{g})$ -subbimodule. If $M = N$, then the image of $U(\mathfrak{g})$ in $\operatorname{Hom}_{\mathbb{C}}(M, M)$ belongs to $\mathcal{L}(M, M)$ and this $U(\mathfrak{g})$ - $U(\mathfrak{g})$ -bimodule map is also an algebra map. The following is known as *Kostant's problem*, see [Jo]:

Problem 2.4. *For which $w \in W$, the image of $U(\mathfrak{g})$ in $\operatorname{Hom}_{\mathbb{C}}(L_w, L_w)$ coincides with $\mathcal{L}(L_w, L_w)$?*

This problem was studied, for example, in [GJ, MS1, MS2, Ma1, Ma3, KhM, Kh], where several partial results were obtained. However, the general case is very much open. As already mentioned, we write $\mathbf{K}(w)$ for the statement “the image of $U(\mathfrak{g})$ in $\operatorname{Hom}_{\mathbb{C}}(L_w, L_w)$ coincides with $\mathcal{L}(L_w, L_w)$ ”.

2.4. K  hrstr  m's conditions. For $y \in W$, we write $\mathbf{Kh}(y)$ for the statement “for all $x \neq z \in W$ such that $\theta_x L_y \neq 0$ and $\theta_z L_y \neq 0$, we have $\theta_x L_y \not\cong \theta_z L_y$ ” and write $[\mathbf{Kh}](y)$ for the statement “for all $x \neq z \in W$ such that $\theta_x L_y \neq 0$ and $\theta_z L_y \neq 0$, we have $[\theta_x L_y] \neq [\theta_z L_y]$ in the Grothendieck group $\operatorname{Gr}(\mathcal{O}_0^{\mathbb{Z}})$ ”.

In particular, the equivalence of Conjecture 1.2(a) and Conjecture 1.2(b) can be expressed as $\mathbf{K}(d) \Leftrightarrow \mathbf{Kh}(d)$, and the equivalence of Conjecture 1.2(b) and Conjecture 1.2(c) can be expressed as $\mathbf{Kh}(d) \Leftrightarrow [\mathbf{Kh}](d)$, for a Duflo element $d \in W$.

3. KAZHDAN-LUSZTIG COMBINATORICS

The Kazhdan-Lusztig conjecture from [KL], proved in [BB, BK] (see also [EW]), tells us that a large amount of information on $\mathcal{O}_0^{\mathbb{Z}}$ is encoded in the Hecke algebra of W and thus can be computed combinatorially. We recall in this section some well-known constructions and facts around the Hecke algebra and their relation to $\mathcal{O}_0^{\mathbb{Z}}$.

3.1. Hecke algebra and Kazhdan-Lusztig basis. Let $S \subset W$ be the set of simple reflections. Then (W, S) is a Coxeter group. The Hecke algebra $H(W, S)$ associated to (W, S) is a $\mathbb{Z}[v, v^{-1}]$ -algebra generated by H_s , for $s \in S$, which satisfy the (Coxeter) braid relations and the quadratic relation

$$(H_s + v)(H_s - v^{-1}) = 0,$$

for all $s \in S$. Given a reduced expression $w = st \cdots u$ of $w \in W$, we let $H_w = H_s H_t \cdots H_u$. The element H_w is, in fact, independent of the choice of the reduced expression, and $\{H_w\}_{w \in W}$ is a $(\mathbb{Z}[v, v^{-1}])$ -basis of $H(W, S)$ called the *standard basis*. Now consider the $(\mathbb{Z}$ -algebra-)involution

$$\overline{} : H(W, S) \rightarrow H(W, S)$$

determined by $\overline{v} = v^{-1}$ and $\overline{H_s} = H_s^{-1}$. Then there is a unique element $\underline{H}_w$ in $H(W, S)$ such that $\overline{\underline{H}_w} = \underline{H}_w$ and

$$\underline{H}_w = H_w + \sum_y p_{y,w} H_y,$$

for some $p_{y,w} \in v\mathbb{Z}[v]$. The elements $\underline{H}_w$, where $w \in W$, form a basis of $H(W, S)$ called the *Kazhdan-Lusztig (KL) basis*. We refer to [KL], or [Lu3, §3-5] for details (if referring to [Lu3], note that we are in the special case $L(s) = 1$ for all $s \in S$ and that our v is denoted by v^{-1} , H_w is denoted by T_w , and our $\underline{H}_w$ is denoted by c_w in [Lu3]).

We denote by μ the Kazhdan-Lusztig μ -function.

3.2. Kazhdan-Lusztig theory. Let $\mathcal{P}$ be the monoidal category of (graded) projective functors on $\mathcal{O}_0^{\mathbb{Z}}$ where indecomposables θ_w , for $w \in W$, are normalized such that $\theta_w P_e \cong P_w$ holds in $\mathcal{O}_0^{\mathbb{Z}}$. Then, for the split Grothendieck ring $\text{Gr}_{\oplus}(\mathcal{P})$, we have

$$(4) \quad \begin{aligned} \text{Gr}_{\oplus}(\mathcal{P})^{\text{op}} &\xrightarrow{\sim} H(W, S) \\ [\theta_w] &\mapsto \underline{H}_w \end{aligned}$$

as $\mathbb{Z}[v, v^{-1}]$ -algebras so that $\langle 1 \rangle$ on the left corresponds to v on the right. For example, $[\theta_w \langle m \rangle] \mapsto v^m \underline{H}_w$. Furthermore,

$$(5) \quad \begin{aligned} \text{Gr}(\mathcal{O}_0^{\mathbb{Z}}) &\xrightarrow{\sim} H(W, S) \\ [P_w] &\mapsto \underline{H}_w \end{aligned}$$

as (right) modules over $\text{Gr}_{\oplus}(\mathcal{P})^{\text{op}} \cong H(W, S)$ so that $\langle 1 \rangle$ on the left hand side corresponds to v on the right hand side. We note that, for $w \in W$, we have $[\Delta_w] \mapsto H_w$, where Δ_w denotes the (graded) Verma module with simple top L_w .

Using (5), various multiplicities in $\mathcal{O}_0^{\mathbb{Z}}$ can be described in terms of the Kazhdan-Lusztig polynomials from [KL]. In particular, the graded composition multiplicities in P_e are given by the corresponding coefficients of the Kazhdan-Lusztig polynomials.

3.3. Kazhdan-Lusztig cells and the a-function. For $x, y \in W$, write

$$(6) \quad \underline{H}_x \underline{H}_y = \sum_z h_{x,y,z} \underline{H}_z,$$

where $h_{x,y,z} \in \mathbb{Z}[v^{\pm 1}]$. In fact, we have $h_{x,y,z} \in \mathbb{Z}_{\geq 0}[v^{\pm 1}] \cap \mathbb{Z}[v + v^{-1}]$ by [KL].

Given $x, y \in W$, we say $x \leq_L y$ if there exists $z \in W$ such that $\underline{H}_y$ appears with a nonzero coefficient in $\underline{H}_z \underline{H}_x$, that is, $h_{z,x,y} \neq 0$. This defines the equivalence relation $\sim_L$ on W . A (Kazhdan-Lusztig) left cell is an equivalence class for the relation $\sim_L$. Similarly, define the right preorder $\leq_R$ and (Kazhdan-Lusztig) right cells using multiplication on the right. Finally, we define the two-sided preorder $\leq_J$ and (Kazhdan-Lusztig) two-sided cells using multiplication on both sides. The equivalence relation $\sim_J$ is the minimum equivalence relation containing both $\sim_L$ and $\sim_R$. We define the equivalence relation $\sim_H$ as the intersection of $\sim_L$ and $\sim_R$. Equivalence classes for $\sim_H$ are called H-cells. A two-sided cell is called *strongly regular* if the intersection of each left and each right cell inside this two-sided cell is a singleton. A left (right) cell is *strongly regular* if it belongs to a strongly regular two-sided cell.

We note that the left and right orders are the opposite of that in [Lu3]. Our conventions are consistent with the previous papers [Ma2, Ma3, CM1, CM2] of the series.

Lusztig's a-function $\mathbf{a} : W \rightarrow \mathbb{Z}_{\geq 0}$ is defined as follows:

$$(7) \quad \mathbf{a}(z) := \max_{x,y \in W} \{\deg h_{x,y,z}\}.$$

The following facts can be found in [Lu3] (note that the conjectures P1-15 in [Lu3, §13] are proved in [Lu3, §14-15] in our setting). Let w_0 denote the longest element in W .

Proposition 3.1. *Let $x, y \in W$.*

$$(i) \quad x \leq_L y \iff x^{-1} \leq_R y^{-1}.$$

- (ii) $x \leq_L y \iff w_0 x \geq_L w_0 y$. Furthermore, $x \leq_R y \iff x w_0 \geq_R y w_0$.
- (iii) Let $X \in \{L, R, J\}$. If $x \leq_X y$, then $\mathbf{a}(x) \leq \mathbf{a}(y)$. If $x <_X y$, then $\mathbf{a}(x) < \mathbf{a}(y)$. In particular, the $\mathbf{a}$ -function is J -cell invariant.
- (iv) If $\mathbf{a}(x) = \mathbf{a}(y)$, then $x \leq_L y \Rightarrow x \sim_L y$. Furthermore, if $\mathbf{a}(x) = \mathbf{a}(y)$, then $x \leq_R y \Rightarrow x \sim_R y$.

Recall from [Lu1, Lu2], that the $\mathbf{a}$ -function can be also defined as the maximal (or minimal, depending on the normalization) possible degree of a Kazhdan-Lusztig polynomial between the identity and an element of a given left (or right) cell. As a consequence of this definition, for any $w \in W$, we have

$$(8) \quad [P_e : L_w \langle i \rangle] \neq 0 \quad \text{implies} \quad -\mathbf{a}(w) \geq i \geq -\ell(w),$$

where $\ell(w)$ denotes the length of w , moreover, the inequality $-\mathbf{a}(w) \geq i$ is strict unless w is a Duflo element and in the latter case

$$(9) \quad [P_e : L_w \langle -\mathbf{a}(w) \rangle] = 1.$$

For each right cell $\mathcal{R}$ in W , we have the Serre subcategory $\mathcal{O}_0^{\hat{\mathcal{R}}}$ of $\mathcal{O}_0$ whose simples are L_w , for all $w \in W$ such that $w \leq_R \mathcal{R}$, see [MS1].

3.4. Asymptotic rings. We introduce the asymptotic ring $A(W) = A(W, S)$ defined by Lusztig (and called “the ring J ” in [Lu3]). It has a $(\mathbb{Z})$ -basis $\{t_w\}_{w \in W}$ whose multiplication is defined as

$$(10) \quad t_x t_y = \sum_{z \in W} \gamma_{x,y,z^{-1}} t_z,$$

where $\gamma_{y,x,z^{-1}} \in \mathbb{Z}_{\geq 0}$ is the coefficient of $\theta_z \langle \mathbf{a}(z) \rangle$ in the decomposition of $\theta_x \theta_y$, i.e., the top degree coefficient in $h_{y,x,z}$ (see (6)). One can check that $\gamma_{x,y,z^{-1}} = 0$ unless y and x^{-1} belong to the same right cell. The basis elements t_d corresponding to Duflo elements $d \in W$ are local identities in $A(W)$ in the following sense:

Lemma 3.2. *Let $d \in W$ be a Duflo element in $\mathcal{H} = \mathcal{L} \cap \mathcal{L}^{-1}$. Here, $\mathcal{L}$ is the left cell containing d and $\mathcal{L}^{-1}$ is the right cell containing $d^{-1} = d$. We have*

$$(11) \quad t_d t_x = t_x \quad \text{and} \quad t_y t_d = t_y.$$

for each $y \in \mathcal{L}$ and $x \in \mathcal{L}^{-1}$ and

$$(12) \quad t_{x^{-1}} t_x = t_d + \sum_{d \neq z \in \mathcal{L} \cap \mathcal{L}^{-1}} \gamma_{x^{-1},x,z^{-1}} t_z.$$

Proof. This follows from P5, P7, P8, P13 in [Lu3, §14.1] and positivity of $h_{x,y,z}$. □

3.5. Graded composition multiplicities in $\mathcal{O}_0^{\mathbb{Z}}$. By §3.2 (and [KL, BB, BK]), composition multiplicities of many important objects in $\mathcal{O}_0^{\mathbb{Z}}$ can be computed purely inside the Hecke algebra $H(W, S)$.

Proposition 3.3. *For $x, y, w \in W$, we have*

$$(13) \quad [\theta_x L_y : L_z] = h_{z, x^{-1}, y},$$

where $[\theta_x L_y : L_z]$ denotes the graded composition multiplicity viewed as an element in $\mathbb{Z}[v, v^{-1}]$. In particular, for $i \in \mathbb{Z}$, we have

$$(14) \quad [\theta_x L_y : L_z \langle i \rangle] = [\theta_z L_{y^{-1}} : L_x \langle i \rangle].$$

Proof. Formula (13) follows, by adjunction, from the observation that the multiplicity of $L_z\langle i \rangle$ in $\theta_x L_y$ equals

$$\dim \operatorname{hom}(\theta_z P_e \langle i \rangle, \theta_x L_y) = \dim \operatorname{hom}(\theta_{x^{-1}} \theta_z P_e \langle i \rangle, L_y).$$

By a similar adjunction, (14) is equivalent to

$$\operatorname{hom}(\theta_{x^{-1}} \theta_z L_{y^{-1}}, I_e \langle i \rangle) \cong \operatorname{hom}(\theta_{z^{-1}} \theta_x L_y, I_e \langle i \rangle),$$

where I_e is the indecomposable injective envelope of L_e with socle concentrated in degree zero. Also, by adjunction, for $a, b \in W$ and $j \in \mathbb{Z}$, we have

$$\dim \operatorname{hom}(\theta_{a^{-1}} L_b, I_e \langle j \rangle) = \delta_{a,b} \delta_{j,0}.$$

As $\theta_{x^{-1}} \theta_z$ is adjoint to $\theta_{z^{-1}} \theta_x$ and θ_y is adjoint to $\theta_{y^{-1}}$, the multiplicity of θ_y in $\theta_{x^{-1}} \theta_z$ coincides with the multiplicity of $\theta_{y^{-1}}$ in $\theta_{z^{-1}} \theta_x$. The claim follows. $\square$

4. PROOF OF THEOREM 2.2

For $w \in W$, we denote by T_w the indecomposable tilting module in $\mathcal{O}_0$ with highest weight $w \cdot 0$.

We record a lemma which follows directly from, for example, [MM1, Lemma 13(a)].

Lemma 4.1. *Let θ be a projective functor on $\mathcal{O}_0$ and $M \in \mathcal{O}_0$. If $y \in W$ is such that $[\theta M : L_y] \neq 0$, then there exists $x \in W$ such that $[M : L_x] \neq 0$ and $y \leq_R x$.*

We also need the following statement.

Lemma 4.2. *Assume $M \in \mathcal{O}_0$. Let $i \in \mathbb{Z}$ and $w \in W$ be such that $\operatorname{Ext}_{\mathcal{O}}^i(M, T_w) \neq 0$. Then we have*

$$(15) \quad i \geq \min\{\mathbf{a}(x^{-1}w_0) : x \in W \text{ such that } [M : L_x] \neq 0\}.$$

Proof. As $T_w = \theta_{w_0} T_{w_0}$, by adjunction, we have

$$\operatorname{Ext}_{\mathcal{O}}^i(M, T_w) \cong \operatorname{Ext}_{\mathcal{O}}^i(\theta_{w^{-1}w_0} M, T_{w_0}).$$

By Lemma 4.1, any $y \in W$ such that $[\theta_{w^{-1}w_0} M : L_y] \neq 0$ satisfies $y \leq_R x$, for some $x \in W$ such that $[M : L_x] \neq 0$. For such x and y , we have $w_0 y \geq_R w_0 x$ and thus $\mathbf{a}(w_0 y) \geq \mathbf{a}(w_0 x)$ (see Proposition 3.1). In particular,

$$\begin{aligned} \min\{\mathbf{a}(x^{-1}w_0) : x \in W \text{ such that } [M : L_x] \neq 0\} &= \\ \min\{\mathbf{a}(x^{-1}w_0) : x \in W \text{ such that } [\theta_{w^{-1}w_0} M : L_x] \neq 0\}. \end{aligned}$$

Therefore, it is enough to prove the claim for $w = w_0$, in which case $T_w = T_{w_0} = L_{w_0}$.

Let $\mathcal{I}_{w_0}^\bullet$ be the minimal injective resolution of L_{w_0} . The assumption implies that there is a nonzero morphism from M to $\mathcal{I}_{w_0}^i$, i.e., there is $x \in W$ such that $[M : L_x] \neq 0$ and $I_x \subseteq \mathcal{I}_{w_0}^i$. Recall from [So1] that $\mathcal{O}_0$ is Koszul self-dual and this self-duality maps L_w to $P_{w^{-1}w_0}$ and I_w to $L_{w^{-1}w_0}$. Consequently, the minimal injective resolution $\mathcal{I}_{w_0}^\bullet$ of L_{w_0} corresponds to the dominant projective module P_e in the Koszul dual picture. Thus, for $x \in W$ as above, I_x appearing as a summand of $\mathcal{I}_{w_0}^i$ implies that $L_{x^{-1}w_0}$ appears as a composition subquotient of P_e in degree i . By (8), we have $\mathbf{a}(x^{-1}w_0) \leq i \leq \ell(x^{-1}w_0)$, for such x . This implies the inequality in (15) and completes the proof. $\square$

Proof of Theorem 2.2. Let $x, y, z \in W$. Koszul-Ringel self-duality of $\{\theta_a L_b : a, b \in W\}$ from [Ma3, Theorem 16] maps a non-zero homomorphism from $\theta_x L_y$ to $L_z \langle i \rangle$ to a non-zero element in

$$\operatorname{Ext}_{\mathcal{O}}^i(\theta_{y^{-1}w_0} L_{w_0 x^{-1}}, T_{w_0 z^{-1}w_0}).$$

By Lemma 4.1, $\theta_{y^{-1}w_0} L_{w_0 x^{-1}}$ consists of composition factors isomorphic to L_z with $z \leq_R w_0 x^{-1}$. Equivalently, we have $z^{-1} \leq_L x w_0$ yielding $x \leq_L z^{-1} w_0$ by Proposition 3.1(ii). For all such z , we have $\mathbf{a}(z^{-1}w_0) \geq \mathbf{a}(x)$ by Proposition 3.1(iii). Now the claim of Theorem 2.2 follows from Lemma 4.2. $\square$

5. KMM vs KM

In this section we establish a connection between **KMM** and **KM**.

5.1. The graded endomorphism algebra of $\theta_x L_y$. For $x, y \in W$, we consider the module $\theta_x L_y \in \mathcal{O}_0^{\mathbb{Z}}$ and its endomorphism algebra $\text{End}(\theta_x L_y)$ which is naturally $\mathbb{Z}$ -graded.

Lemma 5.1. *The natural $\mathbb{Z}$ -grading on $\text{End}(\theta_x L_y)$ is non-negative in the sense that all components with negative degrees are zero.*

Proof. By [Ma3, Theorem 16], the module $\theta_x L_y$ is Koszul-Ringel dual to $\theta_{y^{-1}w_0} L_{w_0 x^{-1}}$, where w_0 is the longest element of W . Under this duality, the endomorphism algebra of $\theta_x L_y$ is mapped to the algebra of diagonal self-extensions for $\theta_{y^{-1}w_0} L_{w_0 x^{-1}}$. The grading of the latter algebra is manifestly non-negative, which implies the claim. $\square$

Corollary 5.2. *If $\dim \text{End}(\theta_x L_y)_0 = 1$, then **KM**(x, y) is true.*

Proof. Every idempotent of the non-negatively graded algebra $\text{End}(\theta_x L_y)$ must be homogeneous of degree zero. Therefore the assumption $\dim \text{End}(\theta_x L_y)_0 = 1$ implies that $\text{End}(\theta_x L_y)_0 \cong \mathbb{C}$, that is, the only idempotents of $\text{End}(\theta_x L_y)$ are 0 and 1. This means that $\text{End}(\theta_x L_y)$ is local and hence $\theta_x L_y$ is indecomposable. $\square$

5.2. **KMM($*_d, y$) implies **KM**($*, y$) in type A.** Assume now that $\mathfrak{g}$ is of type A. Then the cell structure of W is especially nice. In particular, if $\mathcal{L}$ and $\mathcal{R}$ are a left and a right cell inside the same two-sided cell, then $|\mathcal{L} \cap \mathcal{R}| = 1$. Moreover, if $\mathcal{R} = \{w^{-1} : w \in \mathcal{L}\}$, then $\mathcal{L} \cap \mathcal{R} = \{d\}$, where d is a Duflo element.

Lemma 5.3. *For $x \in W$ in type A and d a Duflo element such that $x \sim_R d$, we have*

$$\theta_{x^{-1}} \theta_x \cong \theta_d \langle -\mathbf{a}(d) \rangle \oplus \bigoplus_{w \in W} \bigoplus_{i > -\mathbf{a}(w)} \theta_w \langle i \rangle^{\oplus m_{w,i}}.$$

Proof. Since $\mathcal{R} \cap \mathcal{R}^{-1} = \{d\}$, where $x \in \mathcal{R}$, the claim follows from (12) in Lemma 3.2. $\square$

Proposition 5.4. *Assume that $\mathfrak{g}$ is of type A and $x, y \in W$. Then **KMM**($*_d, y$) implies **KM**(x, y).*

Proof. By adjunction, we have

$$\text{hom}(\theta_x L_y, \theta_x L_y) \cong \text{hom}(\theta_{x^{-1}} \theta_x L_y, L_y).$$

Given Theorem 2.2, from Lemma 5.3 it follows that the only term in the decomposition of $\theta_{x^{-1}} \theta_x$ which can contribute to a non-zero element of $\text{hom}(\theta_{x^{-1}} \theta_x L_y, L_y)$ is $\theta_{d'} \langle -\mathbf{a}(x) \rangle$, where $d' \sim_R x$ is a Duflo element. Therefore, we just need to show that

$$\dim \text{hom}(\theta_{d'} L_y, L_y \langle \mathbf{a}(x) \rangle) = 1.$$

If $\theta_{d'} L_y \neq 0$, this is guaranteed by **KMM**(d', y). $\square$

6. EXTENSIONS BETWEEN TRANSLATED SIMPLE MODULES

6.1. Monotonicity of projective dimension. One of our main observations in this section is the following statement in the spirit of [Ma3] and [CM2].

Theorem 6.1. *Let $x, x', y \in W$ be such that $x >_R x'$ and $\theta_{x'} L_y \neq 0$. Then*

$$\text{proj.dim}(\theta_x L_y) < \text{proj.dim}(\theta_{x'} L_y).$$

Note that, in the setup of Theorem 6.1, the inequality $\text{proj.dim}(\theta_x L_y) \leq \text{proj.dim}(\theta_{x'} L_y)$ follows easily applying projective functors to a minimal projective resolution of $\theta_{x'} L_y$, where $\theta_x L_y$ can be eventually found as a direct summand of the homology in the homological position zero, since $x >_R x'$.

We note that the main result of [Ma3] provides an explicit formula for $\text{proj.dim}(\theta_y L_{w_0})$ in terms of the $\mathbf{a}$ -function and thus implies Theorem 6.1 in the case $y = w_0$ by since the $\mathbf{a}$ -function is strictly monotone along any of the Kazhdan-Lusztig orders (see Proposition 3.1 (ii)). More generally, in the case $y = w_0^{\mathfrak{p}} w_0$, where $\mathfrak{p}$ is a parabolic subalgebra of $\mathfrak{g}$, Theorem 6.1 follows from [CM2, Table 2 and Theorem 4.1(i)].

Let M be a $\mathbb{Z}$ -graded module. Then the *graded length* of M is defined as the difference between the maximal and the minimal degrees of non-zero components of M . If $M = 0$, then the graded length is, by convention, $-\infty$. For modules concentrated in a single degree, the graded length is 0. For example, the projective dimension of M is the same as the graded length of a minimal projective resolution of M consider as a module in an appropriate graded category with a $\mathbb{Z}$ -grading given by the homological degree.

For $x, y \in W$, denote by $g_{x,y}$ the graded length of $\theta_x L_y$, viewed as an object of the category of linear complexes of tilting modules in $\mathcal{O}$. From [Ma3, Theorem 16], it follows that $g_{x,y}$ is always even. Applying projective functors, it is easy to see that

$$(16) \quad g_{x,y} \leq g_{x',y} \quad \text{if} \quad x \geq_R x'.$$

Proposition 6.2. *Consider $x, y \in W$ such that $\theta_x L_y \neq 0$ and let $k := \text{proj.dim}(\theta_x L_y)$. Then we have:*

$$(i) \quad k = \mathbf{a}(w_0 x) + \frac{1}{2} g_{x,y} =: k'.$$

$$(ii) \quad \text{For any } z \in W \text{ such that } z \not\sim_L x, \text{ we have } \text{Ext}^k(\theta_x L_y, L_z) = 0.$$

Proof. Let $z \in W$ be such that $\text{Ext}^k(\theta_x L_y, L_z) \neq 0$. Let d be the Duflo element in the left cell of x . Then $\theta_d \theta_x = \theta_x \oplus \theta$, for some projective functor θ , in particular, $\text{Ext}^k(\theta_d \theta_x L_y, L_z) \neq 0$. By adjunction, we obtain $\text{Ext}^k(\theta_x L_y, \theta_d L_z) \neq 0$. We use the Koszul-Ringel duality of [Ma3, Theorem 16] and represent both $\theta_x L_y$ and $\theta_d L_z$ as complexes of tilting modules. We call these complexes $\mathcal{X}^\bullet$ and $\mathcal{Y}^\bullet$, respectively. The minimal non-zero position of $\mathcal{X}^\bullet$ is $-\frac{1}{2} g_{x,y}$. The maximal non-zero position of $\mathcal{Y}^\bullet$ is $\frac{1}{2} g_{d,z}$. As $d \sim_J x$, from [Ma3, Proposition 1] we get the inequality $\frac{1}{2} g_{d,z} \leq \mathbf{a}(w_0 d) = \mathbf{a}(w_0 x^{-1}) = \mathbf{a}(w_0 x)$, moreover, $\frac{1}{2} g_{d,z} = \mathbf{a}(w_0 x)$ if and only if $w_0 z^{-1} \sim_L w_0 d$. Note that $w_0 z \sim_L w_0 d$ is equivalent to $z \sim_L d$ in which case we have $z \sim_L x$. This implies that $k \leq k'$ and that $k = k'$ holds exactly when $z \sim_L x$.

Therefore, we just need to find some $z \in W$ such that $z \sim_L x$ and

$$\text{Ext}^{k'}(\theta_x L_y, L_z) \neq 0.$$

Let L_u be some simple module which appears in the graded degree $-\frac{1}{2} g_{x,y}$, which is the extreme degree, of the module $\theta_{y^{-1} w_0} L_{w_0 x^{-1}}$. By Lemma 4.1 u belongs to the right cell $\bar{\mathcal{R}}$ of $w_0 x^{-1}$. Since all indecomposable projective-injective modules in $\mathcal{O}^{\bar{\mathcal{R}}}$ are of the form $\theta_{z^{-1} w_0} L_{w_0 d}$ for $z \sim_L x$ (see [Ma3]), there exists $z \sim_L x$ such that $\theta_{z^{-1} w_0} L_{w_0 d}$ is the injective envelope of L_u in $\mathcal{O}^{\bar{\mathcal{R}}}$. For this z , the factor L_u appears in the graded degree $\frac{1}{2} g_{d,z}$

of the module $\theta_{z^{-1}w_0}L_{w_0d}$. Since any isomorphism between the corresponding summands of $\mathcal{X}^{-\frac{1}{2}g_{x,y}}$ and $\mathcal{Y}^{\frac{1}{2}g_{d,z}}$ gives rise a non-zero extension of degree k' between $\theta_x L_y$ and $\theta_d L_z$ (e.g., by the argument from [MO, Theorem 1], a), the claim follows. $\square$

Proof of Theorem 6.1. By Proposition 6.2(i) we need to prove that $\mathbf{a}(w_0x) + \frac{1}{2}g_{x,y}$ is strictly monotone along the right Kazhdan-Lusztig order with respect to x . The term $\mathbf{a}(w_0x)$ is strictly monotone by Proposition 3.1 (ii),(iii). The claim follows by (16). $\square$

Directly from Proposition 6.2(ii), we obtain:

Corollary 6.3. *Conjecture 1.3 is true.*

We record a special case of Proposition 6.2(i).

Corollary 6.4. *Let $x, y \in W$ be such that $\theta_x L_y \neq 0$ and $x \sim_J y$. Then*

$$\text{proj.dim}(\theta_x L_y) = 2\mathbf{a}(w_0x) = \mathbf{a}(w_0x) + \mathbf{a}(y^{-1}w_0).$$

Proof. This follows from Proposition 6.2 (i) and [Ma3, Proposition 1]. $\square$

6.2. Theorem 6.1 does not naively extend to the Koszul-Ringel dual. An alternative description of $g_{x,y}$ is given by:

Proposition 6.5. *For $x, y \in W$ such that $\theta_x L_y \neq 0$, we have*

$$g_{x,y} = \max\{i : \text{Ext}^i(\theta_x L_y, \theta_x L_y) \neq 0\}.$$

Proof. By representing $\theta_x L_y$ as a complex $\mathcal{X}^\bullet$ of tilting modules, we can compute $\text{Ext}^i(\theta_x L_y, \theta_x L_y)$ via homomorphisms from $\mathcal{X}^\bullet$ to $\mathcal{X}^\bullet[i]$ in the homotopy category of tilting modules. If $i > g_{x,y}$, then the homomorphism space is obviously zero as non-zero components of $\mathcal{X}^\bullet$ and $\mathcal{X}^\bullet[i]$ never match.

Because of the self-duality of $\theta_x L_y$ combined with the Koszul-Ringel self-duality of $\{\theta_u L_v : u, v \in W\}$ from [Ma3, Theorem 16], the maximal and the minimal non-zero components of $\mathcal{X}^\bullet$ are isomorphic as objects in $\mathcal{O}$. Therefore, when $i = g_{x,y}$, the argument from the proof of [MO, Theorem 1] shows that an isomorphism from the minimal component in $\mathcal{X}^\bullet$ to the maximal component in $\mathcal{X}^\bullet[i]$ induces a non-zero homomorphism between complexes in the homotopy category of complexes of tilting modules. The claim follows. $\square$

In contrast to Theorem 6.1, the function $x \mapsto g_{x,y}$, for y fixed, does not have to be strictly monotone with respect to the right order on W . Indeed, if $u \in W$ is the longest element of some parabolic subgroup, then the graded length of each non-zero $\theta_u L_v$, where $v \in W$, equals $2\ell(u)$. By Koszul-Ringel duality, this gives, in the case $y = w_0 u^{-1}$, that $g_{x,y} = 2\ell(u)$, for any x for which $g_{x,y} \neq -\infty$. In other words, in this case, the function $x \mapsto g_{x,y}$ is, in fact, constant. This example and the discussion above imply the following corollary:

Corollary 6.6. *Let $\mathfrak{p}$ be a parabolic subalgebra of $\mathfrak{g}$ and T any non-zero tilting module in the parabolic category $\mathcal{O}_0^{\mathfrak{p}}$, then*

$$\max\{i : \text{Ext}_{\mathcal{O}}^i(T, T) \neq 0\} = 2\ell(w_0^{\mathfrak{p}}).$$

Proof. This follows from the above discussion noting that all indecomposable tilting modules in $\mathcal{O}_0^{\mathfrak{p}}$ are of the form $\theta_x L_{w_0^{\mathfrak{p}' w_0}}$, for some x , where $\mathfrak{p}'$ is the parabolic subalgebra obtained from $\mathfrak{p}$ using w_0 . $\square$

We emphasize that the extensions in the formulation of Corollary 6.6 are taken in $\mathcal{O}$ and not in $\mathcal{O}^p$. Indeed, T is assumed to be a tilting module in $\mathcal{O}^p$ and, as such, is ext-self-orthogonal in $\mathcal{O}^p$.

6.3. The $\mathbf{b}$ -function. We look further into the number $g_{x,y}$ appear in Proposition 6.5. We introduce a function which refines the $\mathbf{a}$ -function. For $x, y \in W$, consider the non-negative integer

$$(17) \quad \mathbf{b}(x, y) := \max\{\deg h_{z, x^{-1}, y} : z \in W\}$$

which we take as the definition of the function $\mathbf{b} : W \times W \rightarrow \mathbb{N} \sqcup \{-\infty\}$ (by our convention the degree of the zero polynomial is $-\infty$).

By (13), $\mathbf{b}(x, y)$ gives the height (the highest degree of the composition factors) of the module $\theta_x L_y$. Since the latter is self-dual, we have

$$(18) \quad 2\mathbf{b}(x, y) = \text{the graded length of } \theta_x L_y.$$

Using the Koszul-Ringel duality from [Ma3], we also have

$$(19) \quad 2\mathbf{b}(x, y) = \text{tilt. dim } \theta_{y^{-1}w_0} L_{w_0x^{-1}}$$

where $\text{tilt. dim } M$ denotes the length of a minimal complex of tilting modules representing M . Consequently, we have

$$(20) \quad 2\mathbf{b}(x, y) = g_{y^{-1}w_0, w_0x^{-1}}.$$

Thus we can rewrite Proposition 6.2(i) as

$$(21) \quad \text{proj. dim}(\theta_x L_y) = \mathbf{a}(w_0x) + \mathbf{b}(y^{-1}w_0, w_0x^{-1}).$$

On the one hand, this provides a formula for the projective dimension of $\theta_x L_y$ purely in terms of Kazhdan-Lusztig combinatorics. On the other hand, this gives another characterization of the $\mathbf{b}$ -function.

Proof. This follows directly from (21) and Proposition 6.8(i). $\square$

Proposition 6.7. *The value $\mathbf{b}(x, y)$ is $-\infty$ if and only if $x^{-1} \not\leq_L y$. Moreover, if $y \sim_L y'$, then $\mathbf{b}(x, y) = \mathbf{b}(x, y')$.*

Proof. The first statement follows from (18). If $y \sim_L y'$, then $y^{-1}w_0 \sim_R (y')^{-1}w_0$. The second statement follows from (19). Alternatively, given a tilting resolution $T_\bullet$ of $\theta_{y^{-1}w_0} L$, a tilting resolution of $\theta_{(y')^{-1}w_0} L$ is obtained as a direct summand of $\theta T_\bullet$ for some projective functor θ . This implies $\mathbf{b}(x, y) \geq \mathbf{b}(x, y')$ and the desired equality holds by symmetry. $\square$

Proposition 6.8. *Let $x, y \in W$ be such that $x^{-1} \leq_L y$.*

- (i) *If $x \sim_J y$ (i.e., $x^{-1} \sim_L y$), then we have $\mathbf{a}(x) = \mathbf{b}(x, y) = \mathbf{a}(y)$;*
- (ii) *If $x \not\sim_J y$ (i.e., $x^{-1} <_L y$), then $\mathbf{a}(x) \leq \mathbf{b}(x, y) < \mathbf{a}(y)$;*
- (iii) *$\mathbf{b}(x, y) \leq \ell(x)$;*
- (iv) *If $x = w_0^{\mathfrak{p}}$, for some parabolic subalgebra $\mathfrak{p}$, then $\mathbf{a}(x) = \mathbf{b}(x, y) = \ell(x)$;*
- (v) *If $y = w_0 w_0^{\mathfrak{p}}$, for some parabolic subalgebra $\mathfrak{p}$, then $\mathbf{b}(x, y) = s_{\lambda_{\mathfrak{p}w_0}}(w_0x^{-1}) - \mathbf{a}(y)$, where $s_{\lambda_{\mathfrak{p}w_0}}(w)$ denotes the projective dimension of $L(w \cdot \lambda_{\mathfrak{p}w_0})$ for a singular weight $\lambda_{\mathfrak{p}w_0}$ with stabilizer given by the w_0 -conjugate of the parabolic subgroup of W corresponding to $\mathfrak{p}$, cf. [CM2].*

Proof. By (18), we have $\mathbf{a}(x) \leq \mathbf{b}(x, y) \leq \ell(x)$ and $\mathbf{b}(x, y) \leq \mathbf{a}(y)$ is clear from the definition. The latter is equality if and only if $x \sim_J y$ by Proposition 6.4. This proves claims (i), (ii) and (iii). Claim (iv) follows from claims (i), (ii) and (iii) since $\mathbf{a}(w_0^p) = \ell(w_0^p)$.

Claim (v) follows from Proposition 6.2 and (20), after two observations. The first observation is that $L(w_0 x^{-1} \cdot \lambda)$ has a linear projective resolution in its singular block $\mathcal{O}_\lambda$ and this resolution is mapped to a linear projective resolution of $\theta_{w_0^p} L_{w_0 x^{-1}}$ by translating out of the wall from $\mathcal{O}_\lambda$ to $\mathcal{O}_0$. The second observation is that $\mathbf{a}(y) = \mathbf{a}(w_0 y^{-1} w_0)$. $\square$

Remark 6.9. In the case $y = w_0^p w_0$, we alternatively have $\theta_x L_y = \theta_x L_{w_0^p w_0}$ is the indecomposable tilting module with highest weight $w_0^p w_0 x$ in the principal block $\mathcal{O}_0^p$ of the parabolic category $\mathcal{O}^p$. According to [CM2, Table 2], the graded length of this parabolic tilting module is given by $2(\mathbf{s}_\lambda((w_0^p w_0 x)^{-1} w_0^p) - \mathbf{a}(w_0 w_0^p))$. Therefore we have $\mathbf{b}(x, y) = \mathbf{s}_\lambda(w_0(w_0 x^{-1}) w_0) - \mathbf{a}(w_0 y w_0)$ which agrees with Proposition 6.8(v).

Note that Proposition 6.8(v) provides an explicit description of the function $\mathbf{s}_\lambda$ in terms of Kazhdan-Lusztig combinatorics.

7. K AND KMM

7.1. Category $\mathcal{O}$ via Harish-Chandra bimodules. Denote by $\mathcal{H}$ the category of Harish-Chandra $U(\mathfrak{g})$ - $U(\mathfrak{g})$ -bimodules, that is, finitely generated $U(\mathfrak{g})$ - $U(\mathfrak{g})$ -bimodules on which the adjoint action of $\mathfrak{g}$ is locally finite and has finite multiplicities, see [Ja, Kapitel 6]. Note that $\mathcal{H}$ is equipped with a tensor product given by tensoring over $U = U(\mathfrak{g})$ (however, $\mathcal{H}$ is not monoidal as the regular bimodule $U(\mathfrak{g})$ is not a Harish-Chandra bimodule).

Let $\mathbf{m}$ denote the maximal ideal of $Z(\mathfrak{g})$ which annihilates the trivial $\mathfrak{g}$ -module. Let I be the kernel of the surjection from $Z(\mathfrak{g})$ onto $\text{End}(P_{w_0})$, see Endomorphismensatz in [So1]. Let ${}^\infty\mathcal{H}_0^\infty$ denote the full subcategory of $\mathcal{H}$ consisting of all bimodules X such that $X\mathbf{m}^i = 0$ and $\mathbf{m}^i X = 0$, for $i \gg 0$. Similarly let ${}^\infty\mathcal{H}_0^1$ be the full subcategory of $\mathcal{H}$ (or of ${}^\infty\mathcal{H}_0^\infty$) consisting of bimodules X such that $X\mathbf{m} = 0$ and $\mathbf{m}^i X = 0$ for $i \gg 0$. Let ${}^\infty\mathcal{H}_0^I$ be the full subcategory of ${}^\infty\mathcal{H}_0^\infty$ consisting of bimodules X such that $XI = 0$ and $\mathbf{m}^i X = 0$ for $i \gg 0$. By [BG], we have mutually inverse equivalences of categories as follows:

$$(22) \quad \mathcal{O}_0 \begin{array}{c} \xrightarrow{\mathcal{L}(P_e, -)} \\ \xleftarrow{- \otimes_{U(\mathfrak{g})} P_e} \end{array} {}^\infty\mathcal{H}_0^1$$

Similarly, we have a monoidal equivalence

$$(23) \quad (\overline{\mathcal{P}}, \circ) \cong ({}^\infty\mathcal{H}_0^I, \otimes_U),$$

where $\overline{\mathcal{P}}$ denotes the projective abelianization of $\mathcal{P}$, see [MM1, § 3.5].

7.2. Internal hom of projective functors. The monoidal category $({}^\infty\mathcal{H}_0^I, \otimes_U)$ acts (in the sense of [EGNO, Section 7]) on the left on $\mathcal{O}_0$ in the obvious way, and we can apply the theory of internal hom as in [Os] (see [EGNO, §7.8-7.10]). This action restricts to the projective objects, that is, $\text{proj}({}^\infty\mathcal{H}_0^I) \cong \mathcal{P}$ acts on $\text{proj}(\mathcal{O}_0)$ as well as on $\mathcal{O}_0$. The action of $\mathcal{P}$ on $\mathcal{O}_0$ is given by exact functors, and the action of ${}^\infty\mathcal{H}_0^I$ on $\mathcal{O}_0$ is given by right exact functors. For $M \in \mathcal{O}_0$, the internal hom functor $[M, -] : \mathcal{O}_0 \rightarrow {}^\infty\mathcal{H}_0^I$ is defined as the right adjoint of the right exact functor $- \otimes_U M : {}^\infty\mathcal{H}_0^I \rightarrow \mathcal{O}_0$. The object $[M, N]$, for $M, N \in \mathcal{O}_0$, agrees with the subspace $\mathcal{L}(M, N)$ of $\text{Hom}_{\mathbb{C}}(M, N)$ where the action of $\mathfrak{g}$ is locally finite, as defined in §2.3. The following adjunction confirms this fact.

Proposition 7.1. *For all $M, N \in \mathcal{O}_0$ and $X \in {}^\infty\mathcal{H}_0^I$, we have:*

$$\text{Hom}_{\mathcal{O}}(X \otimes_U M, N) \cong \text{Hom}_{{}^\infty\mathcal{H}_0^I}(X, \mathcal{L}(M, N)).$$

Proof. The Hom-tensor adjunction gives

$$\mathrm{Hom}_{\mathcal{O}}(X \otimes_U M, N) = \mathrm{Hom}_{U-}(X \otimes_U M, N) \cong \mathrm{Hom}_{U-U}(X, \mathrm{Hom}_{\mathbb{C}}(M, N)).$$

Now, since $X \in {}^\infty\mathcal{H}_0^I$, any $U - U$ -bimodule map $X \rightarrow \mathrm{Hom}_{\mathbb{C}}(M, N)$ factors through $\mathcal{L}(M, N)$, that is, we have

$$\mathrm{Hom}_{\mathcal{O}}(X \otimes_U M, N) \cong \mathrm{Hom}_{{}^\infty\mathcal{H}_0^I}(X, \mathcal{L}(M, N)). \quad \square$$

Note that $\mathcal{L}(\Delta_e, \Delta_e)$, as well as $\mathcal{L}(M, M)$, for any $M \in \mathcal{O}_0$, is an algebra in ${}^\infty\mathcal{H}_0^I$. The multiplication coming from the internal hom construction and the multiplication restricted from $\mathrm{Hom}_{\mathbb{C}}(M, M)$ coincide. We denote by $\mathrm{mod}_{{}^\infty\mathcal{H}_0^I}(\mathcal{L}(M, M))$ the category of (right) $\mathcal{L}(M, M)$ -modules in ${}^\infty\mathcal{H}_0^I$. We denote by $\mathrm{proj}_{{}^\infty\mathcal{H}_0^I}(\mathcal{L}(M, M))$ the category of projective objects in $\mathrm{mod}_{{}^\infty\mathcal{H}_0^I}(\mathcal{L}(M, M))$. The morphism spaces in both categories are denoted $\mathrm{Hom}_{\mathcal{L}(M, M)}(-, -)$. There is a natural action of ${}^\infty\mathcal{H}_0^I$ on $\mathrm{mod}_{{}^\infty\mathcal{H}_0^I}(\mathcal{L}(M, M))$ on the left given by the monoidal structure of ${}^\infty\mathcal{H}_0^I$. We refer to [EGNO] for further details on module categories in monoidal categories.

Proposition 7.2. *For $M \in \mathcal{O}_0$, we have*

$$\mathrm{add}_{\mathcal{O}_0}\{\theta_x M \mid x \in W\} \cong \mathrm{proj}_{{}^\infty\mathcal{H}_0^I}(\mathcal{L}(M, M))$$

as module categories over $\mathcal{P}$. Moreover, we have

$$\mathrm{add}_{\mathcal{O}_0}\{FM \mid F \in {}^\infty\mathcal{H}_0^I\} \cong \mathrm{mod}_{{}^\infty\mathcal{H}_0^I}(\mathcal{L}(M, M))$$

as module categories over ${}^\infty\mathcal{H}_0^I$.

Proof. Since $\mathcal{P} \cong \mathrm{proj}_{{}^\infty\mathcal{H}_0^I}$ has monoidal duals (i.e., rigid in the sense of [EGNO]; it is also fiat in the sense of [MM1]), the internal hom theory as in [EGNO, Section 7] and [MMMT] applies. Consider the functor

$$\mathcal{L}(M, -) : \mathcal{O}_0 \rightarrow {}^\infty\mathcal{H}_0^I.$$

Objects of the form $\mathcal{L}(M, N)$ can be viewed as objects in $\mathrm{mod}_{{}^\infty\mathcal{H}_0^I}(\mathcal{L}(M, M))$ since composition defines an action $\mathcal{L}(M, N) \otimes \mathcal{L}(M, M) \rightarrow \mathcal{L}(M, N)$. Now, consider the restriction $\overline{\Phi}$ of $\mathcal{L}(M, -)$ as follows:

$$\overline{\Phi} : \mathrm{add}_{\mathcal{O}_0}\{FM \mid F \in {}^\infty\mathcal{H}_0^I\} \rightarrow \mathrm{mod}_{{}^\infty\mathcal{H}_0^I}(\mathcal{L}(M, M)).$$

Then we have that

- (i) $\overline{\Phi}$ is a $\mathcal{P}$ -module functor;
- (ii) $\mathcal{L}(M, M)$ is projective in $\mathrm{mod}_{{}^\infty\mathcal{H}_0^I}(\mathcal{L}(M, M))$.

This yields a $\mathcal{P}$ -module functor

$$\Phi : \mathrm{add}_{\mathcal{O}_0}\{\theta M \mid \theta \in \mathcal{P}\} \rightarrow \mathrm{proj}_{{}^\infty\mathcal{H}_0^I}(\mathcal{L}(M, M))$$

(see the proofs in [MMMT, §4.2] for the dual version). Since $\mathrm{add}_{\mathcal{O}_0}\{\theta M \mid \theta \in \mathcal{P}\}$ is, by definition, a transitive representation of $\mathcal{P}$, an algebra analog of [MMMT, Theorem 4.7] gives the two equivalences in the statement. $\square$

The following is a slight variation of (22).

Proposition 7.3. *We have $\mathrm{mod}_{{}^\infty\mathcal{H}_0^I}(\mathcal{L}(\Delta_e, \Delta_e)) \cong \mathcal{O}_0$.*

Proof. By Proposition 7.2, we have equivalences

$$\mathrm{add}_{\mathcal{O}_0}\{\theta_x \Delta_e \mid x \in W\} \cong \mathrm{proj}_{{}^\infty\mathcal{H}_0^I}(\mathcal{L}(\Delta_e, \Delta_e))$$

and

$$(24) \quad \mathrm{add}_{\mathcal{O}_0}\{F \otimes_U \Delta_e \mid F \in {}^\infty\mathcal{H}_0^I\} \cong \mathrm{mod}_{{}^\infty\mathcal{H}_0^I}(\mathcal{L}(\Delta_e, \Delta_e)).$$

But $\theta_x \Delta_e \cong P_x$, and therefore $\text{add}_{\mathcal{O}_0} \{\theta_x \Delta_e \mid x \in W\} \cong \text{proj } \mathcal{O}_0$. The claim follows. $\square$

7.3. The image of $\mathcal{L}(L_w, L_w)$ in $\mathcal{O}_0$. Graded characters of the projective modules P_w , $w \in W$, can be computed using Kazhdan-Lusztig combinatorics. Recall from Subsection 3.3 the left $\leq_L$, right $\leq_R$ and the two-sided order $\leq_J$ on W .

Note that $\text{Ann}(L_x) = \text{Ann}(L_y)$ if and only if $x \sim_L y$, and moreover, the inclusion order of primitive ideals for simple modules in $\mathcal{O}_0$ is the opposite of $\leq_L$, see [BV1, BV2]. In particular, the image of $U(\mathfrak{g})$ in $\mathcal{L}(L_w, L_w)$ depends (up to isomorphism) only on the left cell of w .

Let $d \in W$ be a Duflo element. The module $\theta_d L_d$ is indecomposable with simple top $L_d \langle \mathbf{a}(d) \rangle$ and simple socle $L_d \langle -\mathbf{a}(d) \rangle$. It has a unique simple subquotient isomorphic to the trivial $\mathfrak{g}$ -module, and this subquotient is in degree zero. Therefore there is a unique, up to scalar, non-zero homomorphism from P_e to $\theta_d L_d$. We denote by D_d the image of this homomorphism. The module D_d is indecomposable with simple top L_e and simple socle $L_d \langle -\mathbf{a}(d) \rangle$. All other simple subquotients of D_d have the form $L_w \langle i \rangle$, where $-\mathbf{a}(d) < i < 0$ and $x <_R d$ (and hence also $x <_L d$ since d is an involution). We refer to [Ma3, Section 3] for details.

Proposition 7.4. *We have*

$$(U(\mathfrak{g})/\text{Ann}(L_d)) \otimes_{U(\mathfrak{g})} P_e \cong D_d.$$

Proof. The equivalence (22) sends, by construction, $U(\mathfrak{g})/(U(\mathfrak{g})\mathbf{m})$ to P_e inducing a bijection between subobjects. Note that subobjects of the former are exactly the two-sided ideals and that D_d is, by definition, a quotient of P_e . Therefore, it is enough to argue that $\text{Ann}(D_d) = \text{Ann}(L_d)$. This follows from [KhM, Lemma 6]. $\square$

Denote by $\overline{D}_d$ the intersection of the kernels of all possible homomorphisms

$$\varphi : \theta_d L_d \rightarrow \theta_w L_d \langle i \rangle, \quad w \sim_R d, \quad i \in \mathbb{Z},$$

satisfying $\varphi(D_d) = 0$. By construction, $D_d \subset \overline{D}_d$. In particular, L_d is the simple socle of $\overline{D}_d$ and all other composition factors of $\overline{D}_d$ are of the form $L_w \langle i \rangle$, where $-\mathbf{a}(d) < i$ and $w <_R d$ (and also $w <_L d$).

Proposition 7.5. *We have*

$$\mathcal{L}(L_d, L_d) \otimes_{U(\mathfrak{g})} P_e \cong \overline{D}_d.$$

Proof. By (22), it is enough to prove that

$$(25) \quad \mathcal{L}(P_e, \overline{D}_d) \cong \mathcal{L}(L_d, L_d).$$

The natural projection $P_e \twoheadrightarrow D_d$ induces an embedding

$$(26) \quad \mathcal{L}(D_d, \overline{D}_d) \subset \mathcal{L}(P_e, \overline{D}_d).$$

Let K be the kernel of $P_e \twoheadrightarrow D_d$. From [Ma3, Corollary 3] it follows that θK has no non-zero homomorphisms to $\overline{D}_d$, for any projective functor θ . Therefore the inclusion (26) is, in fact, an isomorphism. Consequently, we have $\mathcal{L}(P_e, \overline{D}_d) \cong \mathcal{L}(D_d, \overline{D}_d)$. Now (25) follows from [KhM, Lemma 11] and [KhM, Lemma 12]. $\square$

As an immediate corollary from Propositions 7.4 and 7.5, we have the following statement (which is a reformulation of the main result of [KhM]):

Corollary 7.6. *For $d \in W$ Duflo element, $\mathbf{K}(d)$ is equivalent to $D_d = \overline{D}_d$.*

Example 7.7. In type A_3 with Dynkin diagram $\overset{1}{\circ} \text{---} \overset{2}{\circ} \text{---} \overset{3}{\circ}$, where 1, 2, 3 are simple reflections, it is known from [MS2] that $\mathbf{K}(13)$ is false. In this case one can compute that the graded composition multiplicities of D_{13} and $\overline{D}_{13}$ are, respectively, as follows (here e is in degree zero, and we also abbreviate L_w by w , for simplicity):

$$(27) \quad \begin{array}{ccc} & e & \\ 1 & & 3 \\ & 13 & \end{array} \qquad \begin{array}{ccccc} & & e & & \\ & & 1 & & 3 \\ 123 & & & & 321 \\ & & 13 & & \end{array}$$

7.4. $\mathbf{K}(d)$ implies $\mathbf{KMM}(*, d)$.

Proposition 7.8. *Let $d \in W$ be a Duflo element. Then $\mathbf{K}(d)$ implies $\mathbf{KMM}(*, d)$.*

Proof. By Proposition 7.5, the claim $\mathbf{K}(d)$ is equivalent to $D_d = \overline{D}_d$. Note that D_d is a quotient of P_e , and hence (8) and (9) implies that $[D_d : L_x\langle -\mathbf{a}(x) \rangle] \leq 1$. This implies $\mathbf{KMM}(*, d)$. $\square$

From Example 7.7 we see that $\mathbf{K}$ is strictly stronger than $\mathbf{KMM}$. Indeed, the two additional elements on the right picture in (27) (compared with the left picture) both have the $\mathbf{a}$ -value 1 and have multiplicity one (they are not Duflo elements either). Therefore, in this case, we have that $\mathbf{K}(13)$ is false while $\mathbf{KMM}(*, 13)$ is true.

8. $\mathbf{K}$ VS. $\mathbf{KM}$ AND $\mathbf{Kh}$

8.1. Some computations in $\mathcal{O}^{\hat{\mathcal{R}}}$.

Lemma 8.1. *Let $\mathcal{L}$ be a left cell in W and $d \in \mathcal{H} = \mathcal{L} \cap \mathcal{L}^{-1}$ the corresponding Duflo element. For each $y \in \mathcal{L}$ and $y' \in \mathcal{L}^{-1}$, we have*

$$(28) \quad \theta_d \theta_y = \theta_y \langle \mathbf{a}(d) \rangle \oplus \theta, \quad \theta_{y'} \theta_d = \theta_{y'} \langle \mathbf{a}(d) \rangle \oplus \theta',$$

where $\theta, \theta' \in \mathcal{P}$ are (possibly empty) direct sums of (some) $\theta_z \langle a \rangle$ with $z \not\leq_J d$ or $a < \mathbf{a}(d)$.

Proof. This follows from Lemma 3.2. $\square$

The following lemma slightly extends [Ma3, Theorem 6].

Lemma 8.2. *Let $\mathcal{R} \subseteq W$ be a right cell, $y \in \mathcal{R}$ and $x \in W$ be such that $x \sim_J y$. Then $\theta_x L_y$ is a projective-injective object in $\mathcal{O}^{\hat{\mathcal{R}}}$.*

Proof. If $y = d$ is Duflo, the claim follows directly from [Ma3, Theorem 6]. Moreover, the objects $\theta_x L_d$, where $x^{-1} \in \mathcal{R}$, give a complete and irredundant list of indecomposable projective-injective modules in $\mathcal{O}^{\hat{\mathcal{R}}}$. Set $a := \mathbf{a}(d)$. Each of $\theta_x L_d$ has simple top $L_x \langle a \rangle$ and simple socle $L_x \langle -a \rangle$ and no other subquotients in these extreme non-zero components.

For an arbitrary y , by [Ma3, Proposition 1], the module $\theta_x L_y$ is either zero (in which case the claim is obvious) or the top of $\theta_x L_y$ is concentrated in degree $-a$ and the socle of $\theta_x L_y$ is concentrated in degree a , moreover, these top and socle have the same length due to self-duality of $\theta_x L_y$. Therefore, from the previous paragraph it follows that the minimal projective cover of $\theta_x L_y$ is an isomorphism. The claim of the lemma follows. We note that, alternatively, the claim of the lemma follows from [KMMZ, Theorem 2]. $\square$

Proposition 8.3. *Let $\mathcal{L}$ be a left cell and $y, d \in \mathcal{L}$ be such that d is Duflo. Let d' be the Duflo element in the right cell $\mathcal{R}$ of y . Then $\theta_d L_y \cong \theta_y L_{d'} \cong P_y^{\hat{\mathcal{R}}} \langle \mathbf{a}(d) \rangle$, where $P_y^{\hat{\mathcal{R}}}$ denotes the projective cover of L_y in $\mathcal{O}_0^{\hat{\mathcal{R}}}$.*

Proof. By Lemma 8.1, we have

$$\theta_d P_y \cong \theta_d \theta_y P_e \cong P_y \langle \mathbf{a}(d) \rangle \oplus P,$$

where P is a direct sum of (some) $P_z \langle a \rangle$ with either $z \not\leq_J d$ or $a < \mathbf{a}(d)$. Applying θ_d to the canonical map $P_y \twoheadrightarrow L_y$, we get

$$(29) \quad P_y \langle \mathbf{a}(d) \rangle \oplus P \cong \theta_d P_y \twoheadrightarrow \theta_d L_y.$$

By [Ma3, Proposition 1], the top of $\theta_d L_y$ is concentrated in degree $\mathbf{a}(d) = \mathbf{b}(d, y) = \mathbf{a}(y)$. Therefore, this top is a direct sum of $L_w \langle \mathbf{a}(d) \rangle$, for $w \sim_R y$. It follows that the image of any possible map from P to $\theta_d L_y$ does not contain any top component of $\theta_d L_y$, showing that (29) restricts to $P_y \langle \mathbf{a}(d) \rangle \twoheadrightarrow \theta_d L_y$. Now the claim of the proposition follows combining the facts that $\theta_d L_y$ is projective in $\mathcal{O}_0^{\hat{\mathcal{R}}}$, see Lemma 8.2, and that $\theta_y L_{d'}$ is the projective cover of $L_y \langle \mathbf{a}(d) \rangle$ in $\mathcal{O}_0^{\hat{\mathcal{R}}}$, see [Ma3, Theorem 6]. $\square$

Proposition 8.4. *Let $\mathcal{R} \subseteq W$ be a right cell and $d, y \in \mathcal{R}$ with d Duflo. Let $\mathcal{L}$ be the left cell of y and $x \in W$ be such that $x \sim_J y$. Then $\theta_x L_y = 0$ if $x^{-1} \notin \mathcal{L}$ and, in case $x^{-1} \in \mathcal{L}$, we have*

$$(30) \quad \theta_x L_y \cong \bigoplus_{z \in \mathcal{L} \cap \mathcal{R}} \theta_z L_d^{\oplus \gamma_{y,x,z^{-1}}}.$$

Proof. That $\theta_x L_y = 0$, for all $x \in W$ such that $x \sim_J y$ and $x^{-1} \notin \mathcal{L}$ follows from § 2.1. From [Ma3, Proposition 1 and Theorem 6] it follows that $\theta_x L_y$ is a direct sum of $\theta_z L_d$, where $z \in \mathcal{L} \cap \mathcal{R}$, with some multiplicities which we denote by $c_z \in \mathbb{Z}_{\geq 0}$. Applying θ_x to $\theta_y L_d \langle -\mathbf{a}(d) \rangle \twoheadrightarrow L_y$ and using [Ma3, Proposition 1] and § 3.4, we obtain $c_z = \gamma_{y,x,z^{-1}}$. $\square$

8.2. Computation with simple bimodules. Denote by β_x the simple Harish-Chandra bimodule corresponding to $x \in W$, that is, β_x is the simple top of θ_x in ${}^\infty_0 \mathcal{H}_0^I$. Denote by $\star : {}^\infty_0 \mathcal{H}_0^I \rightarrow {}^\infty_0 \mathcal{H}_0^I$ the simple preserving duality satisfying $(\beta \langle a \rangle)^* = \beta^* \langle -a \rangle$, see [MM, § 4.1]. This duality restricts to $\mathcal{O}_0$ and is compatible with the action in the sense that, for $M \in \mathcal{O}_0$ and $\theta \in {}^\infty_0 \mathcal{H}_0^I$, we have a natural isomorphism $\theta^* M^* \cong (\theta M)^*$. The following statement provides a different interpretation of the internal hom.

Proposition 8.5. *For any $y \in W$, we have $\mathcal{L}(L_y, (-)^* I_e)^* \cong - \otimes_U \beta_{y^{-1}}$.*

Proof. Consider the functor $- \otimes_U \beta_{y^{-1}} : {}^\infty_0 \mathcal{H}_0^I \rightarrow {}^\infty_0 \mathcal{H}_0^I$. It naturally commutes (in the sense of [Kho]) with the left action of ${}^\infty_0 \mathcal{H}_0^I$ on itself. Therefore, by the main result of [Kho], it is determined, up to isomorphism, by its value at θ_e . Clearly, this value is $\theta_e \otimes_U \beta_{y^{-1}} = \beta_{y^{-1}}$.

By construction, $\mathcal{L}(L_y, (-)^* I_e)^*$ is also right exact and naturally commutes with the left action of ${}^\infty_0 \mathcal{H}_0^I$ by [Ja, § 6.8]. The value

$$\mathcal{L}(L_y, (\theta_e)^* I_e)^* \cong \mathcal{L}(L_y, (\theta_e P_e)^*)^* \cong \mathcal{L}(L_y, I_e)^*$$

at θ_e can be identified studying homomorphisms from projective objects:

$$\mathrm{Hom}_{{}^\infty_0 \mathcal{H}_0^I}(\theta_w, \mathcal{L}(L_y, I_e)) \cong \mathrm{Hom}_{\mathcal{O}}(\theta_w L_y, I_e) \cong \mathrm{Hom}_{\mathcal{O}}(L_y, \theta_{w^{-1}} I_e) \cong \mathrm{Hom}_{\mathcal{O}}(L_y, I_{w^{-1}}) = \delta_{y, w^{-1}} \mathbb{C}.$$

It follows that $\mathcal{L}(L_y, I_e) \cong \mathcal{L}(L_y, I_e)^* \cong \beta_{y^{-1}}$. The claim follows. $\square$

Let $y, z \in W$. Evaluating both sides of Proposition 8.5 at β_z and noticing that all simples are self-dual, we obtain

$$(31) \quad \mathcal{L}(L_y, L_z) \cong (\beta_z \beta_{y^{-1}})^*.$$

Evaluating both sides of Proposition 8.5 at θ_z , we obtain

$$(32) \quad \mathcal{L}(L_y, I_z) \cong (\theta_z \beta_{y^{-1}})^*.$$

Corollary 8.6. *We have the canonical inclusion $\mathcal{L}(L_y, L_z) \otimes_U P_e \hookrightarrow \theta_z L_{y^{-1}}$, for $y, z \in W$.*

Proof. Applying the left exact functor $\mathcal{L}(L_y, -)$ to the canonical inclusion $L_z \hookrightarrow I_z$, we obtain

$$\mathcal{L}(L_y, L_z) \hookrightarrow \mathcal{L}(L_y, I_z)$$

Applying now the equivalence $- \otimes_U P_e$, the right term becomes $(\theta_z L_{y^{-1}})^* \cong \theta_z L_{y^{-1}}$. $\square$

Note that $\mathcal{L}(L_y, L_z) = 0$ unless $y \sim_R z$. Proposition 8.5 provides the following description of $\mathcal{L}(L_y, L_z)$ in the case when it is nonzero:

Proposition 8.7. *Let $y, z \in W$ be such that $y \sim_R z$. Then the object $\mathcal{L}(L_y, L_z) \otimes_U P_e$ admits a copresentation*

$$(33) \quad 0 \rightarrow \mathcal{L}(L_y, L_z) \otimes_U P_e \rightarrow \theta_z L_{y^{-1}} \rightarrow \bigoplus_{w \in W} \mu(z, w) \theta_w L_{y^{-1}} \langle 1 \rangle.$$

Proof. The simple object $L_z \in \mathcal{O}$ admits an injective copresentation

$$(34) \quad L_z \hookrightarrow I_z \rightarrow \bigoplus_{w \in W} \mu(z, w) I_w \langle 1 \rangle.$$

Applying the left exact functor $\mathcal{L}(L_y, -) \otimes_U P_e$ to (34), we obtain (33) using (32) by the same arguments as in Corollary 8.6. $\square$

Next lemma give us additional information on the copresentation in (33).

Lemma 8.8. *For $y, z \in W$ such that $y \sim_R z$, we have $\text{soc } \mathcal{L}(L_y, L_z) = \text{soc } \mathcal{L}(L_y, I_z)$.*

Proof. We prove the equivalent statement in $\mathcal{O}$, namely, $\text{soc } \mathcal{L}(L_y, L_z) \otimes_U P_e = \text{soc } \mathcal{L}(L_y, I_z) \otimes_U P_e$. Since $y \sim_R z$, the socle of $\mathcal{L}(L_y, I_z) \otimes_U P_e \cong \theta_z L_{y^{-1}}$ is concentrated in degree $\mathbf{a}(z)$. By Proposition 8.7, it is thus enough to note that each $\theta_w L_{y^{-1}} \langle 1 \rangle$ has extreme degree $\mathbf{b}(w, y^{-1}) - 1 \leq \mathbf{a}(y^{-1}) - 1 < \mathbf{a}(z)$, see Proposition 6.8. $\square$

From Proposition 8.7 and Theorem 2.2, we have that, for $y, z \in W$ such that $y \sim_R z$, the assumption

$$[\mathcal{L}(L_y, L_z) \otimes_U P_e : L_w \langle -a \rangle] \neq 0,$$

for some $w \in W$ and $a \in \mathbb{Z}$, implies $\mathbf{a}(w) \leq a \leq \mathbf{a}(y)$.

The following statement generalizes [KhM, Lemma 8(i)].

Corollary 8.9. *Let $y, z \in W$ be such that $y \sim_R z$, and $\mathcal{R}$ be the right cell of y^{-1} . Then $\mathcal{L}(L_y, L_z) \otimes_U P_e$ is isomorphic to the largest submodule M of $\theta_z L_{y^{-1}} \in \mathcal{O}_0^{\hat{\mathcal{R}}}$ such that any simple subquotient of $M / \text{soc } M$ is, up to shift of grading, of the form L_w , for some $w \notin \mathcal{R}$.*

Proof. We start by showing that, for $w \in \mathcal{R}$, the module L_w can only appear in the socle of $\mathcal{L}(L_y, L_z) \otimes_U P_e$, that is in degree $\mathbf{a}(w)$. This follows directly from [Ma3, Proposition 1] and the adjunction

$$\mathrm{Hom}_{\mathcal{O}}(\theta_w L_y, L_z) \cong \mathrm{Hom}_{\infty \mathcal{H}_0^I}(\theta_w, \mathcal{L}(L_y, L_z)),$$

given by Proposition 7.1. Therefore $\mathcal{L}(L_y, L_z) \otimes_U P_e \subset M$ by Corollary 8.6. Now, the necessary isomorphism $\mathcal{L}(L_y, L_z) \otimes_U P_e = M$ follows from Proposition 8.7 since the socle of each summand in the second term of the injective (in $\mathcal{O}_0^{\hat{\mathcal{R}}}$) copresentation of $\mathcal{L}(L_y, L_z) \otimes_U P_e$ is, up to shift of grading, of the form L_w , for some $w \in \mathcal{R}$. $\square$

For a Duflo element $d \in W$, let $\mathcal{R}$ be the right cell of d . Denote by $\Psi : \mathcal{O}_0^{\hat{\mathcal{R}}} \rightarrow \mathcal{O}_0^{\hat{\mathcal{R}}}$ the functor of partial approximation with respect to projective-injective modules in $\mathcal{O}_0^{\hat{\mathcal{R}}}$, see [KM, § 2.4]. The easiest way to define this functor is as follows: we let $\mathcal{X}$ denote the Serre subcategory of $\mathcal{O}_0^{\hat{\mathcal{R}}}$ generated by all simple which do not appear in the socle of projective-injective objects. Then Ψ is the composition of the (exact) natural projection $\mathcal{O}_0^{\hat{\mathcal{R}}} \rightarrow \mathcal{O}_0^{\hat{\mathcal{R}}}/\mathcal{X}$ followed by the right adjoint of this projection. In particular, Ψ is left exact and is equipped with a natural transformation η from the identity to it which is non-zero exactly on those simple modules which appear in the socle of a projective-injective module. By [Ma3, Theorem 6], Corollary 8.9 says precisely that $\mathcal{L}(L_y, L_z) \otimes_U P_e$ is isomorphic to $\Psi(\mathrm{soc}(\theta_z L_{y^{-1}}))$.

We can now relate our discussion closer to the results of [KhM].

Proposition 8.10. *Let $y \in W$ and $\mathcal{R}$ be the right cell containing y^{-1} . Then the left exact functor*

$$\mathcal{L}(L_y, -) \otimes_U P_e : \mathcal{O}_0 \rightarrow \mathcal{O}_0$$

has image in $\mathcal{O}_0^{\hat{\mathcal{R}}}$ and maps $I_z \in \mathcal{O}_0$, for $w \in W$, to zero unless $z \leq_{\mathcal{R}} y$.

Proof. Note that, for any $M \in \mathcal{O}_0$, for L_x to appear in the composition series of $\mathcal{L}(L_y, M) \otimes_U P_e$, the space

$$(35) \quad \mathrm{Hom}_{\mathcal{O}}(P_x, \mathcal{L}(L_y, M) \otimes_U P_e) \cong \mathrm{Hom}_{\infty \mathcal{H}_0^I}(\theta_x, \mathcal{L}(L_y, M)) \cong \mathrm{Hom}_{\mathcal{O}}(\theta_x L_y, M)$$

should be nonzero, which requires $\theta_x L_y \neq 0$ and thus $x \leq_{\mathcal{R}} y^{-1}$. This proves the first claim.

Note that all simple subquotients of $\theta_x L_y$ have the form L_z , where $z \leq_{\mathcal{R}} y$. This implies the second claim. $\square$

For $y \in W$, denote by $\mathcal{R}$ the right cell of y and by $\mathcal{R}'$ the right cell of y^{-1} . Proposition 8.10 says that the functor $\mathcal{L}(L_y, -) \otimes_U P_e$ restricts to the functor

$$(36) \quad \Psi_y : \mathcal{O}_0^{\hat{\mathcal{R}}} \rightarrow \mathcal{O}_0^{\hat{\mathcal{R}'}}.$$

If y is a Duflo element, then $\mathcal{R} = \mathcal{R}'$ and the functor Ψ_y agrees with the functor of partial coapproximation with respect to the projective-injective modules in $\mathcal{O}_0^{\hat{\mathcal{R}}}$, cf. [KhM, Corollary 7.21].

8.3. Kostant's problem via internal hom. By [Ja, § 6.9], we have $U/\mathrm{Ann}(P_e) \cong \mathcal{L}(P_e, P_e)$ as algebras in $\infty \mathcal{H}_0^I$. Recall that $\mathrm{Ann}(P_e) = UI$. More generally, for every $M \in \mathcal{O}_0$, there is an injective map of algebras $\phi_M : U/\mathrm{Ann}(M) \rightarrow \mathcal{L}(M, M)$. In the case $M = L(w)$, since $U/\mathrm{Ann}(P_e) \twoheadrightarrow U/\mathrm{Ann}(L_w)$, we have an algebra homomorphism

$$(37) \quad \mathcal{L}(P_e, P_e) \rightarrow \mathcal{L}(L_w, L_w)$$

which is surjective if and only if $\mathbf{K}(w)$ holds.

Fix $y \in W$, let $\mathcal{L}$ be the left cell of y and $\mathcal{R} = \mathcal{L}^{-1}$, the right cell of y^{-1} . For simplicity, we write $U_y := U/\mathrm{Ann}(L_y)$ and $A_y := \mathcal{L}(L_y, L_y)$, and let $\phi_y := \phi_{L_y} : U_y \hookrightarrow A_y$. This induces the restriction functor

$$\mathrm{Res}_{U_y}^{A_y} : \mathrm{mod}_{\infty \mathcal{H}_0^I}(A_y) \rightarrow \mathrm{mod}_{\infty \mathcal{H}_0^I}(U_y).$$

Proposition 8.11. *The category $\text{mod}_{\infty \mathcal{H}_0^I}(U_y)$ can be identified with ${}_{\infty \mathcal{H}_0^{\text{Ann}(L_y)}}^{\infty}$, i.e., the category of Harish-Chandra bimodules that are annihilated by $\text{Ann}(L_y)$ on the right.*

Proof. Note that $- \otimes_U U_y$ is the identity functor on the full subcategory of ${}_{\infty \mathcal{H}_0^I}$ consisting of all M such that $M \cdot \text{Ann}(L_y) = 0$. In particular, we have $\mu : U_y \otimes_U U_y \cong U_y$ and this, together with the unit map $\epsilon : U/UI \twoheadrightarrow U_y$, makes U_y an algebra object in ${}_{\infty \mathcal{H}_0^I}$. For this algebra object, the corresponding module category $\text{mod}_{\infty \mathcal{H}_0^I}(U_y)$ is a full subcategory of ${}_{\infty \mathcal{H}_0^I}$. The objects in $\text{mod}_{\infty \mathcal{H}_0^I}(U_y)$ are exactly those $M \in {}_{\infty \mathcal{H}_0^I}$ for which $M \otimes_U U_y \cong M$, i.e., $M \cdot \text{Ann}(L_y) = 0$. In fact, if $M \otimes_U U_y \cong M$, then $1_M \otimes_U \mu$ defines a U_y -module structure on M , and, if we have a U_y -modules structure $m : M \otimes_U U_y \rightarrow M$, then the unit law gives $1_M = m \circ (1_M \otimes_U \epsilon)$, which implies that m is an isomorphism since $1_M \otimes_U \epsilon$ is surjective. The claim follows. $\square$

Recall that, for an additive category $\mathcal{A}$, its projective abelianization is denoted $\overline{\mathcal{A}}$, see [MM1, § 3.5].

Proposition 8.12. *We have $\text{mod}_{\infty \mathcal{H}_0^I}(U_y) \cong \overline{\text{add}_{\infty \mathcal{H}_0^I}\{\theta_x \otimes_U U_y : x \in W\}} \cong \mathcal{O}^{\hat{\mathcal{R}}}$.*

Proof. Since β_x , for $x \in W$ such that $x \leq_R y^{-1}$, are exactly the simples that are killed by $\text{Ann}(L_y)$ on the right, the equivalence $- \otimes_U P_e$ given by (22) restricts to the desired equivalence via Proposition 8.11. $\square$

Corollary 8.13. *The indecomposable projectives in $\text{mod}_{\infty \mathcal{H}_0^I}(U_y)$ are exactly $\theta_x \otimes_U U_y$, for $x \in \hat{\mathcal{R}}$.*

Proof. This follows from the fact that the nonzero $\theta_x \otimes_U U_y$ corresponds in $\mathcal{O}_0$ to a quotient of P_x . Therefore such modules are indecomposable and mutually non-isomorphic. $\square$

Proposition 8.14. *For $y \in W$, we have that $\mathbf{K}(y)$ implies both $\mathbf{Kh}(y)$ and $\mathbf{KM}(*, y)$.*

Proof. Suppose $\mathbf{K}(y)$ is true. Then we have

$$(38) \quad \text{proj}_{\infty \mathcal{H}_0^I}(A_y) \cong \text{proj}_{\infty \mathcal{H}_0^I}(U_y),$$

where $\theta_x \otimes_U A_y$ corresponds to $\theta_x \otimes_U U_y$. By Proposition 7.2, we also have

$$(39) \quad \text{add}_{\mathcal{O}_0}\{\theta_x L_y : x \in W\} \cong \text{proj}_{\infty \mathcal{H}_0^I}(A_y),$$

where $\theta_x L_y$ corresponds to $\theta_x A_y$. Combining (38) and (39), we get an equivalence that identifies $\theta_x L_y$ with $\theta_x \otimes_U U_y \in {}_{\infty \mathcal{H}_0^I}$. Thus, by Corollary 8.13, $\theta_x L_y$ are indecomposable and mutually non-isomorphic. $\square$

Proposition 8.15. *For a Duflo element $d \in W$, if both $\mathbf{Kh}(d)$ and $\mathbf{KM}(*, d)$ are true, then $\text{Res}_{U_d}^{A_d}$ is an equivalence.*

Proof. Let $\mathcal{R}$ be the right cell of d . The equivalence in Proposition 8.12 restricts to the equivalence

$$(40) \quad \text{proj}(\mathcal{O}^{\hat{\mathcal{R}}}) \cong \text{proj}_{\infty \mathcal{H}_0^I}(U_d),$$

which sends $P_w^{\hat{\mathcal{R}}}$, for $w \in W$, to $\theta_w \otimes_U U_d$. By [Ma3, Corollary 3], we have $L_d \hookrightarrow P_e^{\hat{\mathcal{R}}}$ and any simple subquotient of the cokernel of this inclusion is of the form L_x , for $x <_R d$, up to shift of grading. For $w \in W$, applying θ_w gives the inclusion $\theta_w L_d \hookrightarrow P_w^{\hat{\mathcal{R}}}$ such that any simple subquotient of the cokernel of this inclusion is of the form L_x , for $x <_R d$, up to shift of grading. In particular, there are no homomorphisms from any such cokernel to any object in $\text{proj}(\mathcal{O}^{\hat{\mathcal{R}}})$. This means that, for $w, u \in W$, any non-zero homomorphism from $P_w^{\hat{\mathcal{R}}}$ to $P_u^{\hat{\mathcal{R}}}$ restricts to a non-zero homomorphism from $\theta_w L_d$ to $\theta_u L_d$. Using (39) and (40), this gives a faithful functor

$$\Upsilon : \text{proj}_{\infty \mathcal{H}_0^I}(U_d) \rightarrow \text{proj}_{\infty \mathcal{H}_0^I}(A_d).$$

Because of $\mathbf{Kh}(d)$ and $\mathbf{KM}(*, d)$, this functor sends (pairwise non-isomorphic) indecomposable objects to (pairwise non-isomorphic) indecomposable objects.

Using (39), Proposition 8.12 and Proposition 7.2, the functor $\text{Res}_{U_d}^{A_d}$ gives rise to the faithful functor

$$\Theta : \text{add}_{\mathcal{O}}\{\theta_x L_d \mid x \in \hat{\mathcal{R}}\} \rightarrow \text{add}_{\infty \mathcal{H}_0^I}\{\theta_x \otimes_U U_d \mid x \in \hat{\mathcal{R}}\}$$

which sends $\theta_x L_d$ to $\theta_x \otimes_U U_d$. The combination of $\mathbf{Kh}(d)$ and $\mathbf{KM}(*, d)$ implies that Θ sends (pairwise non-isomorphic) indecomposable objects to (pairwise non-isomorphic) indecomposable objects. Putting the faithful functors Υ and Θ together, we thus conclude that they are equivalences. This means that $\text{Res}_{U_d}^{A_d}$ is an equivalence when restricted to projective objects and hence is an equivalence. This completes the proof. $\square$

Theorem 8.16. *Let $d \in W$ be a Duflo element. Then $\mathbf{K}(d)$ is true if and only if both $\mathbf{Kh}(d)$ and $\mathbf{KM}(*, d)$ are true.*

Proof. The “only if” direction is Proposition 8.14. For the “if” direction, assume that both $\mathbf{Kh}(d)$ and $\mathbf{KM}(*, d)$ are true. Then, by Proposition 8.15, $\text{Res}_{U_d}^{A_d}$ is an equivalence sending A_d to U_d . Therefore A_d and U_d are isomorphic as objects in $\infty \mathcal{H}_0^I$, which is exactly the claim $\mathbf{K}(d)$. $\square$

8.4. More on $\mathbf{K}$, $\mathbf{Kh}$, and $\mathbf{KM}$. For $y \in W$, we set $E_y := \mathcal{L}(L_y, L_y) \otimes_U P_e \cong \overline{D_y}$ and $D_y := U_y \otimes_U P_e \in \mathcal{O}$. Note that D_y depends only on the left cell of y . From $U_y \rightarrow \mathcal{L}(L_y, L_y)$, we have $D_y \hookrightarrow E_y$.

Proposition 8.17. *Let $y \in W$ and d be the Duflo element in the left cell of y . Then E_d is a summand of E_y .*

Proof. Let $\mathcal{R}$ be the right cell of d . By adjunction, we have

$$\text{Hom}(L_d, \theta_y L_{y^{-1}}) = \text{Hom}(\theta_{y^{-1}} L_d, L_{y^{-1}}).$$

The right hand side has dimension one by [Ma3, Theorem 6]. Therefore $\theta_d L_d$, which is the indecomposable injective envelope of L_d in $\mathcal{O}_0^{\hat{\mathcal{R}}}$, is a summand of the injective module $\theta_y L_{y^{-1}}$ in $\mathcal{O}_0^{\hat{\mathcal{R}}}$ with multiplicity one.

Now we can use Proposition 8.7 and Corollary 8.9. The socle of the module E_d coincides with the socle of $\theta_d L_d$ and $E_d \cong \Psi(\text{Soc}(\theta_d L_d))$. The socle of E_y coincides with the socle of $\theta_y L_{y^{-1}}$ and we have $E_y \cong \Psi(\text{Soc}(\theta_y L_{y^{-1}}))$. Now the claim of the proposition follows from the additivity of Ψ by applying the latter to the unique up to a scalar split injection of $\text{Soc}(\theta_d L_d)$ in $\text{Soc}(\theta_y L_{y^{-1}})$ given by the previous paragraph. $\square$

We summarize the above discussions in the following statement.

Theorem 8.18. *Let $y \in W$ and d be the Duflo element in the left cell of y , and denote by $\mathcal{H}$ the $\mathbb{H}$ -cell of y . Then we have a commutative diagram*

$$(41) \quad \begin{array}{ccccc} D_y & \hookrightarrow & E_y & \hookrightarrow & \theta_y L_{y^{-1}} \cong \bigoplus_{h \in \mathcal{H}} \theta_h L_{y^{-1}} \oplus \gamma_{y^{-1}, y, h^{-1}} \\ \parallel & & \uparrow & & \uparrow \\ D_d & \hookrightarrow & E_d & \hookrightarrow & \theta_d L_d, \end{array}$$

whose vertical maps are split. Moreover, the diagram (41) restricts to

$$(42) \quad \begin{array}{ccccc} L_d = \text{soc } D_y & \hookrightarrow & \text{soc } E_y & \xlongequal{\quad} & \text{soc } \theta_y L_{y^{-1}} \\ \parallel & & \uparrow & & \uparrow \\ L_d = \text{soc } D_d & \xlongequal{\quad} & \text{soc } E_d & \xlongequal{\quad} & \text{soc } \theta_d L_d \end{array}$$

and the object E_y is determined by its socle.

Proof. The first claim is given by Proposition 8.4, Corollary 8.6, and Proposition 8.17. The second claim is given by Lemma 8.8 and Corollary 8.9. $\square$

Corollary 8.19. *Let $d \in W$ be a Duflo element and $\mathcal{L}$ its left cell.*

- (i) *If $\mathbf{K}(d)$ is not true, then $\mathbf{K}(y)$ is not true, for all $y \in \mathcal{L}$.*
- (ii) *If $\mathbf{K}(d)$ is true, then $\mathbf{K}(y)$ is equivalent to $\mathbf{KM}(y, y^{-1})$, for $y \in \mathcal{L}$.*

Proof. This is a direct consequence of Proposition 8.17 and Theorem 8.16. $\square$

Proposition 8.20. *Let d be the Duflo element in a left cell $\mathcal{L}$ in W and $y, z \in \mathcal{L}$. If $t_y t_{y^{-1}} = t_z t_{z^{-1}}$, then, for each $x \in W$, we have $\mathbf{KM}(x, y) \Leftrightarrow \mathbf{KM}(x, z)$ and $\mathbf{Kh}(y) \Leftrightarrow \mathbf{Kh}(z)$. In particular, both $\mathbf{KM}(*, -)$ and $\mathbf{Kh}(-)$ are constant on strongly regular left cells.*

Proof. By Proposition 8.17, the assumption $t_y t_{y^{-1}} = t_z t_{z^{-1}}$ implies $E_y = E_z$. Thus, by Proposition 7.5 and Proposition 7.2, we have equivalences

$$\text{add}\{\theta_x L_y : x \in W\} \cong \text{proj}_{\infty \mathcal{H}_0^I}(A_y) = \text{proj}_{\infty \mathcal{H}_0^I}(A_z) \cong \text{add}\{\theta_x L_z : x \in W\},$$

under which $\theta_x L_y \mapsto \theta_x L_z$, for each x . The claim follows. $\square$

At this point we can strengthen Theorem 8.16.

Theorem 8.21. *Let $y \in W$. Then $\mathbf{K}(y)$ is true if and only if both $\mathbf{Kh}(y)$ and $\mathbf{KM}(*, y)$ are true.*

Proof. The “only if” direction is Proposition 8.14. For the “if” direction, assume that both $\mathbf{Kh}(y)$ and $\mathbf{KM}(*, y)$ are true. Let d' be the Duflo element in the right cell of y and d be the Duflo element in the left cell of y . We will first prove that $\mathbf{K}(d) \Leftrightarrow \mathbf{K}(y)$.

Because of $\mathbf{KM}(*, y)$, the module $\theta_{y^{-1}} L_y$ is indecomposable and hence is isomorphic to $\theta_{d'} L_{d'}$. In particular, $t_{y^{-1}} t_y = t_{d'}$ in the asymptotic ring $A(W)$, see (30). We claim that this implies $t_y t_{y^{-1}} = t_d = t_d t_d$. Indeed, since $t_{d'}$ is a local identity, $t_y t_{y^{-1}}$ is an idempotent of $A(W)$. Using (12), we can write $t_y t_{y^{-1}}$ as $t_d + x$, where x is a linear combination of various t_u with non-negative integer coefficients. Then

$$t_d + x = (t_d + x)(t_d + x) = t_d + t_d x + x t_d + x^2 = t_d + 2x + x^2$$

by Lemma 3.2, and thus $0 = x + x^2$. Since the structure coefficients with respect to the t_u 's are non-negative, we conclude $x = 0$ and $t_y t_{y^{-1}} = t_d$. By Proposition 8.17, this implies $E_y = E_d$, and hence we have $\mathbf{K}(d) \Leftrightarrow \mathbf{K}(y)$.

Now, by Proposition 8.20, $\mathbf{Kh}(y)$ implies $\mathbf{Kh}(d)$, and $\mathbf{KM}(*, y)$ implies $\mathbf{KM}(*, d)$. Therefore $\mathbf{K}(d)$ holds by Theorem 8.16 and we are done. $\square$

In most small rank examples, we have $\mathbf{K}(y)$ if and only if $\mathbf{Kh}(y)$. However, the situation is slightly more complicated in type G_2 .

Example 8.22. Let $\mathfrak{g}$ be of type G_2 , and let (W, S) be its Weyl group. Write $S = \{1, 2\}$. The two-sided cells in W are given by

e	1 121 12121	21 2121
	12 1212	2 212 21212
		212121

where the rows are left, and the columns are right cells. For the right cell $\mathcal{R}$ containing 1, the indecomposables in $\text{add}\{\theta_x L_y : x^{-1}, y \in R\}$ are exactly

$$\theta_1 L_1, \theta_1 L_{12}, \theta_1 L_{121}, \theta_1 L_{1212}, \theta_1 L_{12121}.$$

We describe the situation for $y \in \mathcal{R}$ in Table 1, the rest is either trivial or symmetric to what is given. Note that, for dihedral types, $[\theta_x L_y]$ determines the isomorphism class of $\theta_x L_y$.

y	$\mathbf{K}(y)$	$\mathbf{KM}(*, y)$	$\mathbf{Kh}(y)$	$\theta_{y^{-1}} L_y$	E_y
1	True	True	True	$\theta_1 L_1$	D_1
12	False	False	True	$\theta_1 L_1 \oplus \theta_1 L_{121}$	$D_2 \oplus L_{212}\langle -1 \rangle$
121	False	False	False	$\theta_1 L_1 \oplus \theta_1 L_{121} \oplus \theta_1 L_{12121}$	$D_1 \oplus L_{121}\langle -1 \rangle \oplus L_{12121}\langle -1 \rangle$
1212	False	False	True	$\theta_1 L_1 \oplus \theta_1 L_{121}$	$D_2 \oplus L_{212}\langle -1 \rangle$
12121	True	True	True	$\theta_1 L_1$	D_1

TABLE 1. $\mathbf{K}$, $\mathbf{KM}$ and $\mathbf{Kh}$ for G_2

8.5. A combinatorial statement. Although we reduce $\mathbf{K}(y)$ to $\mathbf{Kh}(y)$ and $\mathbf{KM}(*, y)$, it is not easy to determine $\mathbf{Kh}(y)$ and $\mathbf{KM}(*, y)$ in general. But the combinatorial version $[\mathbf{Kh}](y)$ of $\mathbf{Kh}(y)$ can be determined by computing $h_{x,y,z}$ (see (13)). Recall that Conjecture 1.2 claims $[\mathbf{Kh}](d) = \mathbf{K}(d)$ for Duflo elements $d \in W$. When this is true, the problem $\mathbf{K}(y)$ has a purely combinatorial solution as follows.

Corollary 8.23. *Let d be a Duflo element and let $y \sim_L d$. Suppose $[\mathbf{Kh}(d)] = \mathbf{K}(d)$. Then the condition $\mathbf{K}(y)$ is equivalent to the conjunction of the following conditions:*

- (i) $t_y t_{y^{-1}} = t_d$ in $A(W)$;
- (ii) For $x, x' \leq_R d$, if $h_{z,x,d} = h_{z,x',d}$, for all $z \in W$, then $x = x'$.

Proof. Suppose $\mathbf{K}(y)$ is true. Then Theorem 8.18 says that $E_y = E_d$, Condition (i) is true, and $\mathbf{K}(d)$ is true. By assumption, we also have $[\mathbf{Kh}(d)]$ which is equivalent to Condition (ii) by (13).

For the other direction, suppose Conditions (i), (ii) are true. Theorem 8.18 and Condition (i) implies $\mathbf{K}(d) = \mathbf{K}(y)$. Since Condition (ii) implies $\mathbf{K}(d)$ by assumption, it follows that $\mathbf{K}(y)$ is true. $\square$

8.6. Extra results in classical types. In this subsection, unless explicitly stated otherwise, we assume $\mathfrak{g}$ to be of (classical) type A , B , C , or D .

The asymptotic algebra endows each diagonal H-cell $\mathcal{H}$ with the structure of an abelian group. This group depends only on the two-sided cell containing $\mathcal{H}$ and is isomorphic to $(\mathbb{Z}/2\mathbb{Z})^k$, for some $k \in \mathbb{N}$, see [Lu3]. The off-diagonal H-cells in the same two-sided cell have order 2^l , for $l \leq k$ (see [MMMTZ, Section 7] for more information). If $\mathcal{H}$

is an $\mathbb{H}$ -cell, $\mathcal{H}_l$ is the diagonal $\mathbb{H}$ -cell in the left cell of $\mathcal{H}$, and $\mathcal{H}_r$ is the diagonal $\mathbb{H}$ -cell in the right cell of $\mathcal{H}$, then the asymptotic algebra endows $\mathcal{H}$ with the structure of a transitive $\mathcal{H}_l$ -set via right multiplication, and with the structure of a transitive $\mathcal{H}_r$ -set via left multiplication. In particular, we can speak about stabilizers of elements from $\mathcal{H}$ in both, $\mathcal{H}_l$ and $\mathcal{H}_r$. This combinatorics allows us to formulate a necessary condition for $\mathbf{Kh}(y)$, and thus also for $\mathbf{K}(y)$.

Proposition 8.24. *Let $\mathfrak{g}$ be of classical type and $y \in W$. Assume that the $\mathbb{H}$ -cell of y does not have the maximal cardinality among the $\mathbb{H}$ -cells inside the two-sided cell of y . Then both $\mathbf{Kh}(y)$ and $\mathbf{K}(y)$ are false.*

Proof. Let $y \in \mathcal{H}$ and $\mathcal{H}_l$ and $\mathcal{H}_r$ be as above, in particular, $|\mathcal{H}| < |\mathcal{H}_l| = |\mathcal{H}_r|$. Let $d \in \mathcal{H}_l$ be a Duflo element. As $\mathbf{K}(y)$ implies $\mathbf{Kh}(y)$ by Proposition 8.14, we only need to show that $\mathbf{Kh}(y)$ is false. Taking (30) into account, it is enough to find $z \in \mathcal{H}$ such that $z \neq d$ and $t_y t_d = t_y t_z$. As $|\mathcal{H}| < |\mathcal{H}_l|$, the stabilizer of y in $\mathcal{H}_l$ is non-trivial and hence we can take as z any element from this stabilizer different from d . The claim follows. $\square$

Lemma 8.25. *Suppose W is of classical type, $\mathcal{L}$ is a left cell in W and $y \in \mathcal{L}$. Set $\mathcal{H} = \mathcal{L} \cap \mathcal{L}^{-1}$. Then*

$$\theta_{y^{-1}} \theta_y \cong \bigoplus_{z \in \text{Stab}_{\mathcal{H}}(y)} \theta_z \langle \mathbf{a}(y) \rangle \oplus \theta,$$

where each summand of θ is of the form $\theta_w \langle a \rangle$, where $a < \mathbf{a}(w)$.

Proof. The element $t_y t_{y^{-1}}$ of the asymptotic ring is a linear combination of t_u , for $u \in \mathcal{H}$, and has the same left stabilizer as y . Therefore $t_y t_{y^{-1}}$ must be a scalar multiple of $\sum_{z \in \text{Stab}_{\mathcal{H}}(y)} t_z$. But the multiplicity of θ_d , where $d \in \mathcal{H}$ is the Duflo element, in $\theta_{y^{-1}} \theta_y$ is one. Therefore the scalar in question is one which implies our claim. $\square$

Lemma 8.26. *Suppose W is of classical type, $\mathcal{L}$ is a left cell in W and $y \in \mathcal{L}$. Set $\mathcal{H} = \mathcal{L} \cap \mathcal{L}^{-1}$ and let $d \in \mathcal{H}$ be the Duflo element. Then*

$$\theta_{y^{-1}} L_y \cong \bigoplus_{z \in \text{Stab}_{\mathcal{H}}(y)} \theta_z L_d.$$

In particular, $\theta_{y^{-1}} L_y$ is indecomposable if and only if $\mathcal{H}$ is of maximal cardinality in its two-sided cell.

Proof. This follows from Proposition 8.4 and Lemma 8.25. $\square$

Lemma 8.27. *Let W be of any type and $x, y \in W$. Then we have*

$$\text{hom}(\theta_x L_y, \theta_x L_y) = \bigoplus_{w \in \mathcal{L} \cap \mathcal{L}^{-1}} \text{hom}(\theta_w L_y, L_y \langle \mathbf{a}(x) \rangle)^{\oplus \gamma_{x, x^{-1}, w^{-1}}},$$

where $\mathcal{L}$ is the left cell of x .

Proof. This follows from Theorem 2.2 and the definition of the asymptotic ring by adjunction. $\square$

Lemma 8.28. *Let W be of any type, $x, y \in W$ and d the Duflo element such that $d \sim_{\mathbb{R}} x$. Assume that $\mathbf{KMM}(d, y)$ is true. Then $\mathbf{KM}(x^{-1}, x)$ implies $\mathbf{KM}(x, y)$.*

Proof. If $\theta_{x^{-1}} L_x$ is indecomposable, then Proposition 8.4 implies $t_x t_{x^{-1}} = t_d$. Hence, by Lemma 8.27, $\text{end}(\theta_x L_y)$ is isomorphic to $\text{hom}(\theta_d L_y, L_y \langle \mathbf{a}(x) \rangle)$ and thus has dimension one by $\mathbf{KMM}(d, y)$. This implies the claim. $\square$

Lemma 8.29. *Let W be of any type, $y, z \in W$ and d be the Duflo element such that $d \sim_{\mathbb{R}} w_0 y^{-1}$. Suppose $\mathbf{KMM}(d, z)$ and $\mathbf{KM}(y^{-1}, y)$ are true. Then $\mathbf{KM}(w_0 z^{-1}, y)$ is true.*

degree: composition factors in this degree:

- −3: 123121, 23123121, **231232**^{⊕2}
- −2: 12321, 23123^{⊕2}, 121, 2312312^{⊕2}, 2312321^{⊕4}, 23121, 12312^{⊕2}
- −1: 23123121^{⊕4}, 1231^{⊕3}, 2312^{⊕2}, 2321, 12, 231231^{⊕3}, 123121^{⊕4}, 231232^{⊕4}, 21
- 0: 12321^{⊕3}, 23123^{⊕4}, 121^{⊕2}, 1, 123, 2312312^{⊕4}, 2312321^{⊕8}, 231^{⊕2}, 23121^{⊕2}, 12312^{⊕4}
- 1: 23123121^{⊕4}, 1231^{⊕3}, 2312^{⊕2}, 2321, 12, 231231^{⊕3}, 123121^{⊕4}, 231232^{⊕4}, 21
- 2: 12321, 23123^{⊕2}, 121, 2312312^{⊕2}, 2312321^{⊕4}, 23121, 12312^{⊕2}
- 3: 123121, 23123121, 231232^{⊕2}

FIGURE 1. The composition factors of $\theta_x L_y$ from Subsection 9.1

Proof. The indecomposability of $\theta_{y^{-1}} L_y$ is, by Proposition 8.4, equivalent to $t_y t_{y^{-1}} = t_{d'}$, where d' is an appropriate Duflo element. As we saw in the proof of Theorem 8.21, this, in turn, is equivalent to $t_{y^{-1}} t_y = t_{d''}$, where d'' is an appropriate Duflo element. The latter is equivalent to the indecomposability of $\theta_y L_{y^{-1}}$. By Koszul-Ringel duality, we get that $\theta_{yw_0} L_{w_0 y^{-1}}$ is indecomposable. Then **KMM**(d, z) and Lemma 8.28 implies that $\theta_{w_0 y^{-1}} L_z$ is indecomposable. Then, by Koszul-Ringel duality again, $\theta_{z^{-1} w_0} L_{w_0 y w_0}$ is indecomposable. Since conjugation by w_0 corresponds to an automorphism of the Dynkin diagram, we obtain that $\theta_{w_0 z^{-1}} L_y$ is indecomposable. $\square$

We can now establish a general sufficient condition for equivalence of **K**(y) is equivalent to **Kh**(y).

Proposition 8.30. *Let $\mathfrak{g}$ be of classical type, $y \in W$ and d be the Duflo element such that $d \sim_{\mathbb{R}} w_0 y^{-1}$. Assume that **KMM**(d, z) is true, for all $z \in W$. Then **K**(y) is equivalent to **Kh**(y).*

Proof. By Theorem 8.21, it is enough to show that **Kh**(y) implies **KM**($*, y$). The latter assertion would follow from Lemma 8.29 provided that we can show that $\theta_{y^{-1}} L_y$ is indecomposable. Given **Kh**(y), from Proposition 8.24 it follows that the H -cell of y is of maximal cardinality inside the J -cell of y . Hence indecomposability of $\theta_{y^{-1}} L_y$ follows from Lemma 8.26. $\square$

In this subsection, we do need the assumption on $\mathfrak{g}$ to be of classical type (compare with Example 8.22).

9. FURTHER DISCUSSION AND SPECULATION ON **KMM**

9.1. A counterexample. In this subsection we give an example in which **KMM** is false.

Let (W, S) be of type B_3 . We label the simple reflections in the following way: $\overset{1}{\circ} \text{---} \overset{2}{\circ} \text{---} \overset{3}{\circ}$. Consider $y = 231232$ and a Duflo element $x = 2312312$. We have $\mathbf{a}(x) = 3$. One can check (using a computer) that the graded composition factors of $\theta_x L_y$ is as given in Figure 1 and directly see that $\dim \text{hom}(\theta_x L_y, L_y \langle \mathbf{a}(x) \rangle) = 2$. We note also that we have a decomposition $\theta_x L_y \cong \theta_{12312} L_{23123121} \oplus \theta_{12312} L_{123121}$, supporting Conjecture 2.3.

9.2. A homological approach to **KMM.** Here is a general criterion for **KMM** given in homological terms.

Proposition 9.1. *Let $x, y \in W$ and $\mathcal{H} = \mathcal{L} \cap \mathcal{L}^{-1}$, where $\mathcal{L}$ is the left cell of $w_0 y$. Then **KMM**(x, y) is equivalent to*

$$(43) \quad \sum_{z \in \mathcal{H}} \gamma_{y^{-1} w_0, w_0 y, z^{-1}} \cdot \dim \text{ext}^{\mathbf{a}(x)}(\theta_z L_{w_0 x^{-1}}, L_{w_0} \langle \mathbf{a}(y^{-1} w_0) - \mathbf{a}(x) \rangle) \leq 1.$$

We note that, in type A , the formula (43) simplifies to

$$(44) \quad \dim \operatorname{ext}^{\mathbf{a}(x)}(\theta_d L_{w_0 x^{-1}}, L_{w_0} \langle \mathbf{a}(y^{-1} w_0) - \mathbf{a}(x) \rangle) \leq 1,$$

where $d \in \mathcal{H}$ is the Duflo element (in fact, $\mathcal{H} = \{d\}$ in this case). In classical types, the formula (43) reads

$$\sum_{z \in \operatorname{Stab}_{\mathcal{H}}(y^{-1} w_0)} \dim \operatorname{ext}^{\mathbf{a}(x)}(\theta_d L_{w_0 x^{-1}}, L_{w_0} \langle \mathbf{a}(y^{-1} w_0) - \mathbf{a}(x) \rangle) \leq 1.$$

Proof of Proposition 9.1. We realize $\theta_x L_y$ as a linear complex of tilting modules. Using Koszul-Ringel duality and adjunction, we have

$$(45) \quad \begin{aligned} \operatorname{hom}(\theta_x L_y, L_y \langle \mathbf{a}(x) \rangle) &\cong \operatorname{ext}^{\mathbf{a}(x)}(\theta_{y^{-1} w_0} L_{w_0 x^{-1}}, \theta_{y^{-1} w_0} L_{w_0} \langle -\mathbf{a}(x) \rangle) \\ &\cong \operatorname{ext}^{\mathbf{a}(x)}(\theta_{w_0 y} \theta_{y^{-1} w_0} L_{w_0 x^{-1}}, L_{w_0} \langle -\mathbf{a}(x) \rangle). \end{aligned}$$

For $z \in \mathcal{H}$, the composition $\theta_{w_0 y} \theta_{y^{-1} w_0}$ contains $\theta_z \langle -\mathbf{a}(y^{-1} w_0) \rangle$ with multiplicity $\gamma_{y^{-1} w_0, w_0 y, z^{-1}}$. All other summands of this composition are of the form $\theta_w \langle -a \rangle$, where $w \geq_J d$ and $a < \mathbf{a}(w)$. By (43), the contribution of all $\theta_z \langle -\mathbf{a}(y^{-1} w_0) \rangle$ to the right hand side of (45) is at most one-dimensional. So, to complete the proof we only need to show that the contribution of all other summands of $\theta_{w_0 y} \theta_{y^{-1} w_0}$ to the right hand side of (45) is zero.

To this end, fix $\theta_w \langle -a \rangle$ as above and consider the linear complex $\mathcal{T}_\bullet$ of tilting modules representing $\theta_w L_{w_0 x^{-1}}$. By Koszul-Ringel duality, each indecomposable summand of $\mathcal{T}_{-\mathbf{a}(x)}$ is of the form $T_u \langle -\mathbf{a}(x) \rangle$, for some $u \in W$ such that $u \leq_J w_0 w \leq_J w_0 d$.

For such u , we claim that the condition $\operatorname{hom}(T_u, L_{w_0} \langle b \rangle) \neq 0$ necessarily implies $b \geq \mathbf{a}(w_0 u) = \mathbf{a}(d)$. Indeed, we have $\operatorname{hom}(T_u, L_{w_0} \langle b \rangle) = \operatorname{hom}(T_u, T_{w_0} \langle b \rangle)$ which, in turn, equals $\operatorname{hom}(P_{w_0 u}, P_e \langle b \rangle)$ by Soergel's character formula for tilting modules, see [So2, Theorem 6.7]. Now our $b \geq \mathbf{a}(w_0 u)$ follows from (8).

For such b , we have $b - a > \mathbf{a}(y^{-1} w_0) - \mathbf{a}(x)$ since $-a > -\mathbf{a}(w)$. This implies that the contribution of $\theta_w \langle -a \rangle$ to the right hand side of (45) is zero and completes the proof. $\square$

9.3. KMM via twisting functors. Consider the full twisting functor $\mathbf{T}_{w_0}$, see [AS]. The following statement provides a reformulation of **KMM** in terms of the cohomology of the derived twisting functor $\mathcal{L}\mathbf{T}_{w_0}$ evaluated at a simple module.

Proposition 9.2. *Let $x, y \in W$ and $\mathcal{H} = \mathcal{L} \cap \mathcal{L}^{-1}$, where $\mathcal{L}$ is the left cell of $w_0 y$. Then **KMM**(x, y) is equivalent to*

$$(46) \quad \sum_{z \in \mathcal{H}} \gamma_{y^{-1} w_0, w_0 y, z^{-1}} \cdot [\mathcal{L}_{\mathbf{a}(x)} \mathbf{T}_{w_0}(L_{w_0 x^{-1}}) : L_{z^{-1}} \langle \mathbf{a}(y^{-1} w_0) - \mathbf{a}(x) \rangle] \leq 1.$$

In type A , the formula (46) simplifies to

$$[\mathcal{L}_{\mathbf{a}(x)} \mathbf{T}_{w_0}(L_{w_0 x^{-1}}) : L_d \langle \mathbf{a}(y^{-1} w_0) - \mathbf{a}(x) \rangle] \leq 1,$$

where d is the Duflo element in $\mathcal{H}$. In classical types, the formula (46) reads

$$\sum_{z \in \operatorname{Stab}_{\mathcal{H}}(y^{-1} w_0)} [\mathcal{L}_{\mathbf{a}(x)} \mathbf{T}_{w_0}(L_{w_0 x^{-1}}) : L_{z^{-1}} \langle \mathbf{a}(y^{-1} w_0) - \mathbf{a}(x) \rangle] \leq 1.$$

Proof of Proposition 9.2. In this proof we use standard properties of twisting functors, see [AS] for details. To start with, we note that $\mathcal{L}\mathbf{T}_{w_0}$ is an auto-equivalence of the bounded derived category of $\mathcal{O}_0$. Let us apply this auto-equivalence to the left hand side of (43). In the second argument of the extension space, we get $\mathcal{L}\mathbf{T}_{w_0}(T_{w_0}) = I_e$. In the first argument of the extension space, we note that $\mathcal{L}\mathbf{T}_{w_0}$ commutes with projective functors, so we can move θ_z out. By adjunction, we move θ_z over to the second argument obtaining $\theta_{z^{-1}} I_e \cong I_{z^{-1}}$. The leftover in the first

argument is $\mathcal{L}T_{w_0}(L_{w_0x^{-1}})$. Evaluating at the correct degree of the extension and noting that homomorphisms to $I_{z^{-1}}$ give exactly the composition multiplicity of $L_{z^{-1}}$ for the homology, we obtain (46). $\square$

10. SMALL RANK RESULTS

The results in this paper, together with (computer-assisted computations for) Kazhdan-Lusztig combinatorics, enable us to determine $\mathbf{K}(y)$, $\mathbf{KM}(*, y)$, and $\mathbf{Kh}(y)$ in many cases. We present some of the results in this section.

In computer-assisted calculations, SageMath v.9.0 has been used.

10.1. Type A. It is verified in [CMZ] that $\mathbf{KM}(x, y)$ is true for all x, y in type A_n for $n \leq 5$. Therefore, in this case we only need to determine either $\mathbf{Kh}(y)$ or $\mathbf{K}(y)$, which are equivalent by Theorem 8.21.

In [KhM, Kh], Kostant's problem is solved for simple highest weight modules in $\mathcal{O}_0$, for $\mathfrak{sl}_n$, $n \leq 5$. For $\mathfrak{sl}_6$, i.e., type A_5 , out of 76 Duflo elements, 47 have positive answer to Kostant's problem, 20 have negative, and the following 9 Duflo elements were left as an open problem (the notation for simple reflections is analogous to the one in Example 7.7):

$$(47) \quad \begin{array}{ll} 23432, & 4523412, \\ 234312, & 12342321, \\ 452342, & 23454232, \\ 234512342312, & 3451234231, \\ & 2345412312. \end{array}$$

We can now solve these remaining cases.

For all y in the first column in (47), one can check (by computer) that $[\mathbf{Kh}](y)$ holds, and therefore $\mathbf{Kh}(y)$ holds. From [CMZ] and Theorem 8.21 it follows that $\mathbf{K}(y)$ is true.

For the second column in (47), it is enough to consider only $x := 4523412$, $y := 12342321$ and $z := 3451234231$, because of the symmetry of the Dynkin diagram. By [CMZ, Proposition 46.b)], we have

$$\theta_{23}L_x \cong \theta_3L_{x3} \cong \theta_{43}L_x \quad \text{and} \quad \theta_{32}L_z \cong \theta_2L_{z2} \cong \theta_{12}L_z.$$

Therefore both $\mathbf{Kh}(x)$ and $\mathbf{Kh}(z)$ fail. So again, using [CMZ] and Theorem 8.21, we conclude that neither $\mathbf{K}(x)$ nor $\mathbf{K}(z)$ holds.

A calculation shows that $\theta_{45342} \cong \theta_{342}\theta_{45}$. One can check that $\theta_{45}L_y$ has height 1. Moreover, its top and socle are simple, consisting of $L_{y'}$ with $y' := 123452321$, and L_y in the middle. Since $342 \not\leq_R (y')^{-1}$, it follows that

$$\theta_{45342}L_y \cong \theta_{342}L_y.$$

Therefore $\mathbf{Kh}(y)$ does not hold, and so $\mathbf{K}(y)$ also does not hold. Since the property $\mathbf{K}$ in type A is invariant for the left cells, we have:

Corollary 10.1. *Kostant's problem has a positive solution, for a simple highest weight module L_w for $\mathfrak{sl}_6$, if and only if w does not belong to the left cells containing one of the following 25 Duflo elements:*

$$\begin{array}{lllll} 31, & 531, & 45341, & 512321, & 23454232, \\ 42, & 3431, & 52312, & 3453431, & 34541231, \\ 53, & 4121, & 234232, & 4523412, & 2345412312, \\ 232, & 4542, & 345431, & 5123121, & 3451234231, \\ 343, & 5232, & 454121, & 12342321, & 12345343121. \end{array}$$

10.2. **Types BCD .** We completely determine $\mathbf{K}(w)$, $\mathbf{KM}(*, w)$ and $\mathbf{Kh}(w)$, for each $w \in W$ in types B_3 and D_4 . We also determine $\mathbf{K}(w)$ and $\mathbf{Kh}(w)$ completely in type B_4 . In all above examples, we have $\mathbf{Kh}(w) \Leftrightarrow \mathbf{KM}(*, w)$, supporting Conjecture 1.2. We provide below the results without details for type B_3 and D_4 .

A good way to present these results is to mark the H-cells in W in the following way (note that, since W is of classical type, by Proposition 8.20 and Lemma 8.26, the function $\mathbf{K}(w)$ (resp., $\mathbf{KM}(*, w)$ and $\mathbf{Kh}(w)$) has the same value for w in the same H-cell) :

The elements w in the H-cells colored in

- **this** color satisfy $\mathbf{K}(w)$, $\mathbf{KM}(*, w)$ and $\mathbf{Kh}(w)$,
- **this** color satisfy $\mathbf{KM}(*, w)$ and do not satisfy $\mathbf{Kh}(w)$, thus do not satisfy $\mathbf{K}(w)$,
- white do not satisfy any of the above properties.

Figure 2 presents type B_3 results and Figure 3 gives type D_4 results.

e	2312	312	32312
	231	31	3231
	23123	3123	323123

121	23121	3121
12312	2312312	312312
1231	231231	31231

1	21	321
12321	2321	
12	2	32
1232	232	
123	23	3
		323

23123121	3123121	323121
123121		
2312321	312321	32321
	32312321	3231231
231232	31232	3232
	3231232	32312312

323123121

FIGURE 2. Cells in type B_3 with the labeling $\overset{1}{\circ} \text{---} \overset{2}{\circ} \text{---} \overset{3}{\circ}$. Rows are left, and columns are right cells. Duflo elements are the top elements in the diagonal blocks.

The following example is potentially related to Conjecture 1.2 (b) $\Leftrightarrow$ (c):

Example 10.2. Let W be of type B_3 as in Figure 2. In $\text{Gr}(\mathcal{O}_0^{\mathbb{Z}})$ we have the following equalities:

$$\begin{aligned} [\theta_{123}L_{121}] &= [\theta_3L_{231}] + [\theta_{3231}L_{1231}] \\ &= [\theta_3L_{23123}] + [\theta_{323123}L_{1231}]. \end{aligned}$$

However, as shown in Figure 2, the object $\theta_{123}L_{121}$ is indecomposable.

REFERENCES

- [AS] Andersen, H.; Stroppel, C. Twisting functors on $\mathcal{O}$. Represent. Theory **7** (2003), 681–699.
- [BV1] Barbasch, D.; Vogan, D. Primitive ideals and orbital integrals in complex classical groups. Math. Ann. **259** (1982), no. 2, 153–199.
- [BV2] Barbasch, D.; Vogan, D. Primitive ideals and orbital integrals in complex exceptional groups. J. Algebra **80** (1983), no. 2, 350–382.
- [BB] Beilinson, A.; Bernstein, J. Localisation de $\mathfrak{g}$ -modules. C. R. Acad. Sci. Paris Ser. I Math. **292** (1981), no. 1, 15–18.
- [BG] Bernstein, I. N.; Gelfand, S. I. Tensor products of finite- and infinite-dimensional representations of semisimple Lie algebras. Compositio Math. **41** (1980), no. 2, 245–285.
- [BGG] Bernstein, I. N.; Gelfand, I. M.; Gelfand, S. I. A certain category of $\mathfrak{g}$ -modules. (Russian) Funkcional. Anal. i Priložen. **10** (1976), no. 2, 1–8.

e	1	21	321	421			
	12	2	32	42	2312	312	42312
	123	23	3	423	231	31	4231
	124	24	324	4	23124	3124	423124


2412	32412	412	124321	24321	4321
24123	324123	4123	12432	2432	432
241	3241	41	1243	243	43

121	3121	4121	23121	24121	423121	243121	43121
1243121	31243121	12423121	3243121	2423121	324121		
1231	12321	41231	2321	241231	42321	2412321	412321
12412321	312412321	1242321	32412321	242321	3241231		
1241	31241	12421	324321	2421	32421	2312421	312421
23124321	3124321	423124321	231241	42312421	4231241		
12312	1232	124232	232	24232	4232	241232	41232
1241232	31241232	412312	3241232	2412312	32412312		
12412	312432	1242	32432	242	3242	231242	31242
2312432	312412	42312432	2312412	4231242	42312412		
124123	31243	12423	3243	2423	32423	2312423	312423
231243	3124123	4231243	23124123	42312423	423124123		
124312	3124312	1242312	324312	242312	3242312	24312	4312
						231242312	31242312
12431	312431	124231	32431	24231	324231	2431	431
						23124231	3124231

123121	24123121	4123121	23124123121	324123121	3124123121	124123121
23124312	4231242312	423124312	231241232	42312412312	4231241232	423124232
2312431	423124231	42312431	2312412321	4231241231	42312412321	4231242321
			231243121	423124121	4231243121	42312423121

2312423121	312423121	32423121	124121	23124121	3124121
231242321	31242321	3242321	12412312	2312412312	312412312
23124232	3124232	324232	1241231	231241231	31241231

423124123121

FIGURE 3. Cells in type D_4 with labeling . Rows are left, and columns are right cells. Duflo elements are the top elements in the diagonal blocks.

- [BK] Brylinski, J.-L.; Kashiwara, M. Kazhdan-Lusztig conjecture and holonomic systems. *Invent. Math.* **64** (1981), no. 3, 387–410.
- [CM1] Coulembier, K.; Mazorchuk, V. Some homological properties of category $\mathcal{O}$. III. *Adv. Math.* **283** (2015), 204–231.
- [CM2] Coulembier, K.; Mazorchuk, V. Some homological properties of category $\mathcal{O}$. IV. *Forum Math.* **29** (2017), no. 5, 1083–1124.
- [CMZ] Coulembier, K.; Mazorchuk, V.; Zhang, X. Indecomposable manipulations with simple modules in category $\mathcal{O}$. *Math. Res. Lett.* **26** (2019), no. 2, 447–499.
- [EGNO] Etingof, P.; Gelaki, S.; Nikshych, D.; Ostrik, V. *Tensor categories*. Mathematical Surveys and Monographs, **205**. American Mathematical Society, Providence, RI, 2015. xvi+343 pp.

- [EW] Elias, B., Williamson, G. The Hodge theory of Soergel bimodules. *Ann. of Math. (2)* **180** (2014), no. 3, 1089–1136.
- [GJ] Gabber, O.; Joseph, A. On the Bernstein-Gelfand-Gelfand resolution and the Duflo sum formula. *Compositio Math.* **43** (1981), no. 1, 107–131.
- [Hu] Humphreys, J. E. Representations of semisimple Lie algebras in the BGG category $\mathcal{O}$. *Graduate Studies in Mathematics*, **94**. American Mathematical Society, Providence, RI, 2008. xvi+289 pp.
- [Ja] Jantzen, J. C. *Einhüllende Algebren halbeinfacher Lie-Algebren*. *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*, **3**. Springer-Verlag, Berlin, 1983. ii+298 pp.
- [Jo] Joseph, A. Kostant's problem, Goldie rank and the Gel'fand-Kirillov conjecture. *Invent. Math.* **56** (1980), no. 3, 191–213.
- [KL] Kazhdan, D.; Lusztig, G. Representations of Coxeter groups and Hecke algebras. *Invent. Math.* **53** (1979), no. 2, 165–184.
- [Kho] Khomenko, O. Categories with projective functors. *Proc. London Math. Soc. (3)* **90** (2005), no. 3, 711–737.
- [KMMZ] Kildetoft, T.; Mackaay, M.; Mazorchuk, V.; Zimmermann, J. Simple transitive 2-representations of small quotients of Soergel bimodules. *Trans. Amer. Math. Soc.* **371** (2019), no. 8, 5551–5590.
- [KM] Kildetoft, T.; Mazorchuk, V. Parabolic projective functors in type A . *Adv. Math.* **301** (2016), 785–803.
- [Kh] Kåhrström, J. Kostant's problem and parabolic subgroups. *Glasg. Math. J.* **52** (2010), no. 1, 19–32.
- [KhM] Kåhrström, J.; Mazorchuk, V. A new approach to Kostant's problem. *Algebra Number Theory* **4** (2010), no. 3, 231–254.
- [Lu1] Lusztig, G. Cells in affine Weyl groups, in: *Algebraic Groups and Related Topics*, in: *Adv. Stud. Pure Math.*, vol. **6**, North-Holland, Amsterdam, 1985, pp. 255–287.
- [Lu2] Lusztig, G. Cells in affine Weyl groups. II, *J. Algebra* **109** (2) (1987) 536–548.
- [Lu3] Lusztig, G. Hecke algebras with unequal parameters, *CRM Monograph Series*, **18**. American Mathematical Society, Providence, RI (2003), vi+136 pp.
- [Ma1] Mazorchuk, V. A twisted approach to Kostant's problem. *Glasg. Math. J.* **47** (2005), no. 3, 549–561.
- [Ma2] Mazorchuk, V. Some homological properties of the category $\mathcal{O}$. *Pacific J. Math.* **232** (2) (2007) 313–341.
- [Ma3] Mazorchuk, V. Some homological properties of the category $\mathcal{O}$. II. *Represent. Theory* **14** (2010), 249–263.
- [MM] Mazorchuk, V.; Miemietz, V. Serre functors for Lie algebras and superalgebras. *Ann. Inst. Fourier (Grenoble)* **62** (2012), no. 1, 47–75.
- [MM1] Mazorchuk, V.; Miemietz, V. Cell 2-representations of finitary 2-categories. *Compos. Math.* **147** (2011), no. 5, 1519–1545.
- [MM3] Mazorchuk, V.; Miemietz, V. Endomorphisms of cell 2-representations. *Int. Math. Res. Not. IMRN* **2016**, no. 24, 7471–7498.
- [MMMT] Mackaay, M.; Mazorchuk, V.; Miemietz, V.; Tubbenhauer, D. Simple transitive 2-representations via (co-)algebra 1-morphisms. *Indiana Univ. Math. J.* **68** (2019), no. 1, 1–33.
- [MMMTZ] M. Mackaay, V. Mazorchuk, V. Miemietz, D. Tubbenhauer, and X. Zhang. 2-representations of Soergel bimodules. *arXiv:1906.11468*
- [MO] Mazorchuk, V.; Ovsienko, S. Finitistic dimension of properly stratified algebras. *Adv. Math.* **186** (2004), no. 1, 251–265.
- [MS1] Mazorchuk, V.; Stroppel, C. Categorification of (induced) cell modules and the rough structure of generalised Verma modules. *Adv. Math.* **219** (2008), no. 4, 1363–1426.
- [MS2] Mazorchuk, V.; Stroppel, C. Categorification of Wedderburn's basis for $\mathbb{C}[S_n]$. *Arch. Math. (Basel)* **91** (2008), no. 1, 1–11.
- [Os] Ostrik, V. Module categories, weak Hopf algebras and modular invariants. *Transform. Groups* **8** (2003), 177–206.
- [So1] Soergel, W. Kategorie $\mathcal{O}$, perverse Garben und Moduln über den Koinvarianten zur Weylgruppe. *J. Amer. Math. Soc.* **3** (1990), no. 2, 421–445.
- [So2] Soergel, W. Character formulas for tilting modules over Kac-Moody algebras. *Represent. Theory* **2** (1998), 432–448.
- [St] Stroppel, C. Category $\mathcal{O}$: gradings and translation functors. *J. Algebra* **268** (2003), no. 1, 301–326.

H. K.: Department of Mathematics, Uppsala University, Box. 480, SE-75106, Uppsala, SWEDEN, email: hankyung.ko@math.uu.se

V. M.: Department of Mathematics, Uppsala University, Box. 480, SE-75106, Uppsala, SWEDEN, email: mazor@math.uu.se

R. M.: Department of Mathematics, Uppsala University, Box. 480, SE-75106, Uppsala, SWEDEN, email: rafael.mrden@math.uu.se

(On leave from: Faculty of Civil Engineering, University of Zagreb, Fra Andrije Kačića-Miošića 26, 10000 Zagreb, CROATIA)