QUANTITATIVE INDUCTIVE ESTIMATES FOR GREEN'S FUNCTIONS OF NON-SELF-ADJOINT MATRICES

WENCAI LIU

ABSTRACT. We provide quantitative inductive estimates for Green's functions of matrices with (sub)expoentially decaying off diagonal entries in higher dimensions. Together with Cartan's estimates and discrepancy estimates, we establish explicit bounds for the large deviation theorem for non-self-adjoint Toeplitz operators. As applications, we obtain the modulus of continuity of the integrated density of states with explicit bounds and the pure point spectrum property for analytic quasi-periodic operators. Moreover, our inductions are self-improved and work for perturbations with low complexity interactions.

1. INTRODUCTION

The dynamics and spectral theory for quasi-periodic operators have been made significant progress in the last 40 years, through earlier perturbative methods [28, 31, 33, 34, 64, 69], and then non-perturbative methods by controlling Green's functions/transfer matrices [15, 18, 19, 21, 54, 55] or by reducibility [4, 43]. The case of one dimensional lattice and one frequency potential, has been well understood for both small and large coupling constants, with the recent discovery of global theory [2] and universal structure [49, 50]. In particular, remarkable developments have been achieved for several models motivated by physics: the almost Mathieu operator (the Harper's model), the extended Harper's model and the Maryland model [1, 3, 5, 8, 9, 41, 44–50, 52, 53, 55, 57, 58, 58–60, 68]. We refer readers to [62, 74] and references therein for more details.

Problems are known to be much more complicated if one increases the underlying dimension b of the torus or the dimension d of the lattice. The higher dimension picture is still far from clear. For the one dimensional lattice d = 1and multi-frequencies $b \ge 1$, some special cases have been studied by transfer matrices or Schrödinger cocycles [15, 26, 30, 33, 35, 38, 40]. The first multidimensional localization result was obtained by perturbative (KAM) methods by Chulaevsky-Dinaburg for operators on lattices \mathbb{Z}^d and torus \mathbb{T} for arbitrary d [27]. Bourgain-Goldstein-Schlag developed a celebrated method in the spirit

²⁰¹⁰ Mathematics Subject Classification. 81Q10 (primary); 82B44, 37C55 (secondary).

Key words and phrases. Multi-scale analysis, large deviation theorem, discrepancy, semialgebraic sets, Cartan's techniques, Anderson localization.

of non-perturbative approaches from Bourgain-Goldstein [18] to handle the twodimension and two-frequency case [20] (b = d = 2) and established the Anderson localization for large coupling constants. This is the first higher dimension lattice and multi-frequency result. Moreover, the large deviation theorem in [20], which is a key ingredient to prove the Anderson localization, is purely arithmetic in the sense that removed sets of frequencies are independent of the potential. Roughly speaking, by imposing some purely arithmetic condition on $(\omega_1, \omega_2) \in \mathbb{R}^2$, for any algebraic curve $\Gamma \subset [0, 1]^2$ with degree at most N^C , the number of lattice points

(1)
$$\{(n_1, n_2) \in \mathbb{Z}^2 : |n_1| \le N, |n_2| \le N, (n_1\omega_1, n_2\omega_2) \mod \mathbb{Z}^2 \in \Gamma_\tau \}$$

is bounded by $N^{1-\delta}$ for some $\delta > 0$, where Γ_{τ} is the $e^{-N^{\tau}}$ neighborhood of Γ . The quantity $N^{1-\delta}$ is referred to as the sublinear bound. It is still open whether the analogy for $d \geq 3$ is true or not.

In [20], Bourgain developed a new scheme to prove the large deviation theorem for arbitrary b = d [17] by a delicate study of the semi-algebraic sets. Jitomirskaya-Liu-Shi extended Bourgain's result to the case of arbitrary b and d[51]. However, the removed set of frequencies in [17, 51] depends on the potential.

Bourgain, Goldstein and Schlag [20] mentioned that the sub-linear bound (1) is the only obstruction to establish an arithmetic version of the large deviation theorem in higher dimensions. However, there is no detailed proof available yet. Our first goal of this paper is to provide such a proof. Moreover, we are going to establish the quantitative version of the main results in [20] with generalizations, in particular it can be applied to quasi-periodic operators on arbitrary lattices \mathbb{Z}^d driven by any dynamics on tori \mathbb{Z}^b under the assumption on sub-linear bounds.

Instead of Laplacians or long range operators, we will study Toeplitz matrices with (sub)exponentially decaying off diagonal entries. Among all the motivations of our generalizations, we want to highlight one. Anderson localization receives a lot of attentions from both mathematics and physics. The approach to establish Anderson localization for quasi-periodic operators with analytic potentials turns out to be a breakthrough component to construct quasi-periodic solutions for nonlinear Schrödinger equations and nonlinear wave equations [15, 72]. It is known that the quasi-periodic solutions in PDEs are only sub-exponentially, not exponentially decaying [12, 15, 72]. Therefore, the (sub)exponentially decaying matrices are more natural settings in PDEs.

In our arguments, the matrices are not necessarily self-adjoint and every entry of the matrices is allowed to be a function. For $d \ge 2$, this is the first time to study operators that beyonds long range cases. For d = 1, our assumptions are weaker than Bourgain's [14]. See Remark 3 for details. Moreover, our arguments hold under perturbations with low complexity.

Our proof is definitely inspired by [20]. However, there are a lot of important ingredients being added into the arguments to make it quantitative in our more

general settings. Moreover, we significantly simplify the arguments even for the case appearing in [20]. The analysis of [20] required dealing with many different types of elementary regions, say rectangles and *L*-shapes in \mathbb{Z}^2 . We largely reduced the elementary regions to be square related. See Fig.1. Two novelties are added here. Firstly, we introduce the concept of width of subsets of lattices. In our augments, we always keep the involved regions Λ having large width so that every lattice point in Λ can be covered by a square related elementary region with presetting size contained in Λ . For example, the region like Fig.2 was not allowed because the width determined by the distance between *B* and *C* is too small. Secondly, we reconstruct the exhaustion of *x* in every elementary region. In our new construction, the annuli with small width are absorbed into bigger ones. See Fig.3.

There are several other technical improvements in this paper, which we believe to be of independent interest. For example, we establish the Cartan's estimates for non-self-adjoint matrices.

We will prove a quantitatively inductive theorem about the Green's functions in higher dimensions as stated in Theorem 2.1. This is a deterministic statement, which can be applied to study operators even without dynamics. Based on matrix-valued Cartan-type theorem (estimates on subharmonic functions) in [20] with further developments in [15, 36, 51], we will establish the measure estimates in Theorem 2.2. Imposing proper dynamics on tori, the quantitative inductive estimate for Green's functions is obtained (Theorem 2.3). Moreover, the relation among all constants and parameters is displayed clearly so that the whole picture becomes extremely transparent. We will see how arithmetic conditions on frequencies effect the discrepancy, how structures of semi-algebraic sets effect the number of bad Green's functions, and how the dimensions of lattices and frequencies contribute to bounds.

Finally, we want to talk about the applications. As far as we know, there is no explicit bound yet for the large deviation theorem except for the case d = 1and b = 1, 2. Our approaches (Theorems 2.1, 2.2 and 2.3) are the first time to establish the explicit bounds in higher dimensions and multi-frequencies. We show that in the arithmetic sense, for d = 1 and any b, the bound is arbitrarily close to $\frac{1}{b^3}$ for shift dynamics and $\frac{1}{4^{b-1}b^3}$ for skew-shift dynamics. For b = 1 and arbitrary d, we show that the bound is arbitrarily close to 1.

Another application we want to mention is the regularity of the integrated density of states (IDS) of quasi-periodic operators. The log-Hölder continuity of the integrated density of states is quite general [23, 29]. The Hölder continuity in one dimensional settings was well established [6, 7, 13, 15, 26, 35, 36, 40, 42, 61, 75] for both large and small coupling constants. What we will investigate in this paper is the modulus of continuity $f(x) = e^{-\kappa |\log x|^{\tau}}$. Unfortunately, like the large deviation theorem, except for the case d = 1 and b = 1, 2, there

are no explicit bounds of τ in the region of large coupling constants. Based on the ingredients from [13, 65] and the large deviation theorem, the modulus of continuity of the integrated density of states with explicit estimates will be obtained in Theorem 2.5.

2. Main results

Let A be a (operator) matrix on $\ell^2(\mathbb{Z}^d)$ satisfying,

(2)
$$|A(n,n')| \le K e^{-c_1 |n-n'|^{\tilde{\sigma}}}, K > 0, c_1 > 0, 0 < \tilde{\sigma} \le 1,$$

where $|n| := \max_{1 \le i \le d} |n_i|$ for $n = (n_1, n_2, \cdots, n_d) \in \mathbb{Z}^d$. We say that the off diagonal entries of A are subexponentially decaying if A satisfies (2). Sometimes, we just say A is subexponentially decaying for simplicity.

For d = 1, the elementary region of size N centered at 0 is given by

$$Q_N = [-N, N].$$

For $d \geq 2$, denote by Q_N an elementary region of size N centered at 0, which is one of the following regions,

$$Q_N = [-N, N]^d$$

or

$$Q_N = [-N, N]^d \setminus \{ n \in \mathbb{Z}^d : n_i \varsigma_i 0, 1 \le i \le d \},\$$

where for $i = 1, 2, \dots, d$, $\varsigma_i \in \{<, >, \emptyset\}^d$ and at least two ς_i are not \emptyset .

Denote by \mathcal{E}_N^0 the set of all elementary regions of size N centered at 0. Let \mathcal{E}_N be the set of all translates of elementary regions with center at 0, namely,

$$\mathcal{E}_N := \{ n + Q_N : n \in \mathbb{Z}^d, Q_N \in \mathcal{E}_N^0 \}.$$

We call elements in \mathcal{E}_N elementary regions.

Example 1: For d = 2, there are five types of elementary regions.




Fig.1: elementary regions in \mathbb{Z}^2

The width of a subset $\Lambda \subset \mathbb{Z}^d$, is defined by maximum $M \in \mathbb{N}$ such that for any $n \in \Lambda$, there exists $\hat{M} \in \mathcal{E}_M$ such that

$$n\in \hat{M}\subset \Lambda$$

and

dist
$$(n, \Lambda \setminus \hat{M}) \ge M/2$$
.

Example 2: In Fig.2, the width of Λ is determined by the distance between B and C.




Fig.2: a region with small width

A generalized elementary region is defined to be a subset $\Lambda \subset \mathbb{Z}^d$ of the form

$$\Lambda := R \backslash (R+z),$$

where $z \in \mathbb{Z}^d$ is arbitrary and R is a rectangle,

$$R = \{n = (n_1, n_2, \cdots, n_d) \in \mathbb{Z}^d : |n_1 - n_1'| \le M_1, \cdots, |n_d - n_d'| \le M_d\}.$$

For $\Lambda \subset \mathbb{Z}^d$, we introduce its diameter,

diam(
$$\Lambda$$
) = sup $_{n,n'\in\Lambda} |n - n'|$.

Denote by \mathcal{R}_N all generalized elementary regions with diameter less than or equal to N. Denote by \mathcal{R}_N^M all generalized elementary regions in \mathcal{R}_N with width larger than or equal to M. For $\Lambda \subset \mathbb{Z}^d$, let R_Λ be the restriction operator, i.e., $(R_\Lambda u)(n) = u(n)$ for $n \in \Lambda$, and $(R_\Lambda u)(n) = 0$ for $n \notin \Lambda$.

We say an elementary region $\Lambda \in \mathcal{E}_{N'}$ is in class G (Good) if

(3)
$$|(R_{\Lambda}AR_{\Lambda})^{-1}(n,n')| \le e^{-c_2|n-n'|^{\tilde{\sigma}}}, \text{ for } |n-n'| \ge \frac{N'}{10},$$

where $0 < c_2 \leq \frac{5^{\tilde{\sigma}}-1}{5^{\tilde{\sigma}}}c_1$ and $0 < \tilde{\sigma} \leq 1$. We mentioned that the upper bound $\frac{5^{\tilde{\sigma}}-1}{5^{\tilde{\sigma}}}c_1$ is chosen for technical convenience. See (72) for the explanation.

Denote by $\lfloor x \rfloor$ the largest integer smaller than or equal to x.

Theorem 2.1. Assume A satisfies (2). Let $\varsigma, \sigma, \xi \in (0, 1)$ and $\sigma < \tilde{\sigma} \leq 1$. Let $\tilde{\Lambda}_0 \in \mathcal{E}_N$ be an elementary region with the property that for all $\Lambda \subset \tilde{\Lambda}_0$, $\Lambda \in \mathcal{R}_L^{N^{\xi}}$ with $N^{\xi} \leq L \leq N$, the Green's function $(R_{\Lambda}AR_{\Lambda})^{-1}$ satisfies

(4)
$$||(R_{\Lambda}AR_{\Lambda})^{-1}|| \le e^{L^{\sigma}}.$$

Assume that for any family \mathcal{F} of pairwise disjoint elementary regions in $\tilde{\Lambda}_0$ with size $M = \lfloor N^{\xi} \rfloor$,

(5)
$$\#\{\Lambda \in \mathcal{F} : \Lambda \text{ is not in class } G\} \leq \frac{N^{\varsigma}}{N^{\xi}}.$$

Then for large N (depending on $K, \varsigma, \sigma, \tilde{\sigma}, \xi, c_1$ and the lower bound of c_2),

(6)
$$|(R_{\tilde{\Lambda}_0}AR_{\tilde{\Lambda}_0})^{-1}(n,n')| \le e^{-(c_2-N^{-\vartheta})|n-n'|^{\tilde{\sigma}}}, \text{ for } |n-n'| \ge \frac{N}{10},$$

where $\vartheta = \vartheta(\sigma, \tilde{\sigma}, \xi, \varsigma) > 0.$

Here are several comments about Theorem 2.1.

- **Remark 1.** (1) For d = 1 and $\tilde{\sigma} = 1$, a similar statement was proved by Bourgain [14]. For d = 2 and $\tilde{\sigma} = 1$, a similar statement was proved for the particular case where A is given by the discrete Laplacian [20].
 - (2) The statement in Theorem 2.1 is a robust approach to deal with the spectral theory for quasi-periodic operators and also the construction of quasi-periodic solutions for nonlinear Schrödinger/wave equations. See [14, 19, 20, 24, 25] for applications. Some particular cases of Theorem 2.1 have been used as ingredients to construct quasi-periodic solutions for PDEs and have been stated in [24, 25, 72] without detailed proof. There are no explicit bound estimates in their arguments either.
 - (3) In applications, ς is chosen to be arbitrarily close to 1, namely $\varsigma = 1 \varepsilon$ with arbitrarily small $\varepsilon > 0$. Then the upper bound in (5) equals $N^{1-\xi-\varepsilon}$. Theorem 2.1 says that the "goodness" of Green's functions at small size N^{ξ} will ensure the "goodness" of Green's functions at larger size N under the following two conditions:
 - The number of bad Green's functions of size N^{ξ} in $[-N, N]^d$ is less than $N^{1-\xi-\varepsilon}$ (referred to as the sub-linear bound).
 - The Green's functions can not be "super bad" in the sense that they are controlled by (4). The upper bound $e^{L^{\sigma}}$ with $\sigma < 1$ is referred to as the sub-exponential bound.

Let $b = \sum_{i=1}^{k} b_i$, where $b_i \in \mathbb{N}$. Let $x = (x_1, x_2, \cdots, x_k)$, where $x_i \in \mathbb{T}^{b_i} = (\mathbb{R}/\mathbb{Z})^{b_i}$, $i = 1, 2, \cdots, k$. For any $x \in \mathbb{T}^b$ and $1 \le i \le k$, let

$$x_i^{\neg} = (x_1, \cdots, x_{i-1}, x_{i+1}, \cdots, x_k) \in \mathbb{T}^{b-b_j}.$$

For any $y \in \mathbb{T}^{d_1}$ and $X \subset \mathbb{T}^{d_1+d_2}$, denote the y-section of X:

$$X(y) := \{ z \in \mathbb{T}^{d_2} : (y, z) \in X \}.$$

Write Leb(S) for the Lebesgue measure.

Assume each element of the operator A is a function on \mathbb{T}^b . Sometimes, we indicate the dependence and denote by the element A(x; n, n'). Assume every element A(z; n, n') is analytic in the strip $\{z \in \mathbb{C}^b : |\Im z| \leq \rho\}, \rho > 0$, and satisfies for any $n, n' \in \mathbb{Z}^d$ and $x \in \mathbb{T}^b$,

(7)
$$|A(x;n,n')| \le K e^{-c_1 |n-n'|^{\tilde{\sigma}}}, K > 0, c_1 > 0, 0 < \tilde{\sigma} \le 1.$$

Assume that there exists $K_1 > 1$ such that for any $x \in \mathbb{T}^b$ and $z \in \{z \in \mathbb{C}^b : |\Im z| \le \rho\}$ with $||x - z|| \le e^{-(\log(|n| + |n'| + 2))^{K_1}}$,

(8)
$$|A(x;n,n') - A(z;n,n')| \le K ||x - z||^{\gamma},$$

where $||z|| = \operatorname{dist}(z, \mathbb{Z}^b)$.

Example 3. If A satisfies (7) and for any $n, n' \in \mathbb{Z}^d$, A(x; n, n') is a trigonometric polynomial of degree at most $e^{(\log(|n|+|n'|+2))^{K_1}}$, then (8) holds.

We say an elementary region $\Lambda \in \mathcal{E}_N$ is in class SG_N (strongly good with size N) if

(9)
$$||R_{\Lambda}AR_{\Lambda})^{-1}|| \le e^{N^{\sigma}}$$

and

(10)
$$|(R_{\Lambda}AR_{\Lambda})^{-1}(n,n')| \le e^{-c_2|n-n'|^{\check{\sigma}}}, \text{ for } |n-n'| \ge \frac{N}{10},$$

where $0 < c_2 \leq \frac{5\tilde{\sigma}-1}{5\tilde{\sigma}}c_1$ and $0 < \sigma < \tilde{\sigma} \leq 1$. When there is no confusion, we drop the dependence of N from the notation SG_N .

Theorem 2.2. Assume A satisfies (7) and (8). Fix $\sigma, \delta, \tilde{\sigma}, \zeta \in (0, 1)$ and $\mu \in (1 - \delta, 1), \sigma < \tilde{\sigma}$. Suppose $\mathcal{R} \subset [-N_3, N_3]^d$ has width at least N_2 . For $x \in \mathbb{T}^b$, define $\mathcal{B}_{\mathcal{R}}(x)$ as

$$\mathcal{B}_{\mathcal{R}}(x) = \{ n \in \mathcal{R} : \text{ there exists } Q_{N_1} \in \mathcal{E}_{N_1}^0 \text{ such that } n + Q_{N_1} \notin SG_{N_1} \}$$

Assume that for any $x \in \mathbb{T}^b$,

(11)
$$\#\mathcal{B}_{\mathcal{R}}(x) \le L^{1-\delta}.$$

Assume that there exists a subset $X_{N_2} \subset \mathbb{T}^b$, such that

(12)
$$\sup_{1 \le i \le k, x_i^{\neg} \in \mathbb{T}^{b-b_i}} \operatorname{Leb}(X_{N_2}(x_i^{\neg})) \le e^{-N_2^{\zeta}}$$

and for any $Q_{N_2} \in \mathcal{E}_{N_2}^0$, $x \notin X_{N_2}$ and $n \in \mathcal{R}$, the region $n + Q_{N_2}$ is in class SG_{N_2} . Let

$$\tilde{X}_{\mathcal{R}}(x) = \{x \in \mathbb{T}^b : ||(R_{\mathcal{R}}A(x)R_{\mathcal{R}})^{-1}|| \ge e^{L^{\mu}}\}.$$

Suppose $N_3 \leq e^{N_1^{\frac{1}{2K_1}}}$, $N_2 \geq N_1^{\frac{2}{\zeta}}$ and $L \geq N_2^{\frac{2d+b+2}{\mu-1+\delta}}$. Then there exists $N_0 = N_0(K_1, K, c_1, c_2, \tilde{\sigma}, \sigma, \delta, \gamma, \rho, \mu)^1$ such that for any $N_1 \geq N_0$ and $i = 1, 2, \cdots, k$,

(13)
$$\sup_{x_i^{\neg} \in \mathbb{T}^{b-b_i}} \operatorname{Leb}(\tilde{X}_{\mathcal{R}}(x_i^{\neg})) \le e^{-\left(\frac{L^{\mu-1+\delta}}{N_2^{2d+b+2}}\right)^{1/b_i}}$$

¹It depends on the lower bound of c_2 .

Let f be a function from $\mathbb{Z}^d \times \mathbb{T}^b$ to \mathbb{T}^b . Assume for any $m_1, m_2, \cdots, m_d \in \mathbb{Z}^d$ and $n_1, n_2, \cdots, n_d \in \mathbb{Z}^d$,

$$f(m_1 + n_1, m_2 + n_2, \cdots, m_d + n_d, x) = f(m_1, m_2, \cdots, m_d, f(n_1, n_2, \cdots, n_d, x)).$$

Sometimes, we write down $f^n(x)$ for f(n, x) for convenience, where $n \in \mathbb{Z}^d$ and $x \in \mathbb{T}^b$. We say A is a Toeplitz (operator) matrix on $\ell^2(\mathbb{Z}^d)$ with respect to f, if

(14)
$$A(x; n+k, n'+k) = A(f^k(x); n, n'),$$

for any $n \in \mathbb{Z}^d$, $n' \in \mathbb{Z}^d$ and $k \in \mathbb{Z}^d$. We note that A is not necessarily self-adjoint.

We say the Green's function of an operator A(x) satisfies property P with parameters (μ, ζ, c_2) at size N if the following statement is true: there exists a subset $X_N \subset \mathbb{T}^b$ such that

$$\sup_{1 \le i \le k, x_i^{\neg} \in \mathbb{T}^{b-b_i}} \operatorname{Leb}(X_N(x_i^{\neg})) \le e^{-N^{\zeta}},$$

and for any $x \notin X_N \mod \mathbb{Z}^b$ and $Q_N \in \mathcal{E}_N^0$,

$$\begin{aligned} ||(R_{Q_N}A(x)R_{Q_N})^{-1}|| &\leq e^{N^{\mu}}, \\ |(R_{Q_N}AR_{Q_N})^{-1}(x;n,n')| &\leq e^{-c_2|n-n'|^{\tilde{\sigma}}}, \text{ for } |n-n'| \geq \frac{N}{10}. \end{aligned}$$

Theorem 2.3. Assume A(x) satisfies (7), (8) and (14), and

$$0 < c_2 < (1 - 5^{-\tilde{\sigma}})c_1, 1 - \delta < \sigma < \tilde{\sigma} \le 1, \delta > \iota > 0, \text{ and } 0 < \mu < \tilde{\sigma}.$$

Let $c = \frac{1}{2} \min\{\frac{1}{K_1}, \tilde{\sigma}\}$. Fix any sufficiently small $\varepsilon > 0$. There exists a large constant C depending on all parameters such that the following statements are true. Let N_1 be sufficiently large, $N_2 \in [N_1^C, e^{N_1^{c/2}}]$ and $N_3 \in [N_2^C, e^{N_1^c}]$. Assume that the Green's function satisfies the property P with parameters (μ, ζ, c_2) at sizes N_1 and N_2 . Assume for any $L \in [N_3^{\delta-\iota}, N_3]$ and any $x \in \mathbb{T}^b$,

(15)
$$\#\{n \in \mathbb{Z}^d : |n| \le L, f(n, x) \in X_{N_1} \mod \mathbb{Z}^b\} \le L^{1-\delta}$$

Then there exists $X_{N_3} \subset \mathbb{T}^b$ such that

(16)
$$\sup_{1 \le i \le k, x_i^{\neg} \in \mathbb{T}^{b-b_i}} \operatorname{Leb}(X_{N_3}(x_i^{\neg})) \le e^{-N_3 \frac{\sigma-1}{b_i} \delta + \frac{\delta^2}{b_i} - \varepsilon},$$

and for any $x \notin X_{N_3}$ and $Q_{N_3} \in \mathcal{E}_{N_3}^0$,

(17)
$$||(R_{Q_{N_3}}A(x)R_{Q_{N_3}})^{-1}|| \leq e^{N_3^{\sigma}}$$

and for $|n - n'| \ge \frac{N_3}{10}$,

(18)
$$|(R_{Q_{N_3}}AR_{Q_{N_3}})^{-1}(x;n,n')| \leq e^{-(c_2 - 2N_1^{-\nu_1} - N_3^{-\nu_2})|n-n'|^{\sigma}},$$

where $\vartheta_1 = \vartheta_1(\tilde{\sigma}, \mu, c)$ and $\vartheta_2 = \vartheta_2(\tilde{\sigma}, \sigma, \delta, \varepsilon)$.

Our Theorems work for Topelitz matrices with low complexity interactions. Let U be an operator on $\ell^2(\mathbb{Z}^d)$ satisfying

$$|U(n,n')| \le K e^{-c_1 |n-n'|^{\tilde{\sigma}}}$$

Given $m \in \mathbb{Z}^d$, define the operator U^m by

$$U^m(n,n') = U(m+n,m+n'), n \in \mathbb{Z}^d, n' \in \mathbb{Z}^d$$

We say U has low complexity if there exists 0 < a < 1 such that for any N > 1,

(19)
$$\#\{R_{Q_N}U^m R_{Q_N} : m \in \mathbb{Z}^d, Q_N \in \mathcal{E}_N^0\} \le K e^{N^a}.$$

For any $m \in \mathbb{Z}^d$, denote by

(20)
$$\tilde{A}^{m}(x;n,n') = A(x;n,n') + U^{m}(n,n')$$

We say that the Green's function of an operator A(x) satisfies property \tilde{P} with parameters (μ, ζ, c_2) at size N if the following statement is true: there exists a set $X_N \subset \mathbb{T}^b$ such that

$$\sup_{1 \le i \le k, x_i^{\sim} \in \mathbb{T}^{b-b_i}} \operatorname{Leb}(X_N(x_i^{\sim})) \le e^{-N^{\zeta}},$$

and for any $x \notin X_N \mod \mathbb{Z}^b$, $m \in \mathbb{Z}^d$, and $Q_N \in \mathcal{E}_N^0$

$$\begin{aligned} ||(R_{Q_N}\tilde{A}^m(x)R_{Q_N})^{-1}|| &\leq e^{N^{\mu}}, \\ |(R_{Q_N}\tilde{A}^mR_{Q_N})^{-1}(x;n,n')| &\leq e^{-c_2|n-n'|^{\tilde{\sigma}}}, \text{ for } |n-n'| \geq \frac{N}{10}. \end{aligned}$$

We have

Theorem 2.4. Assume A(x) satisfies (7), (8) and (14), U has low complexity,

$$0 < c_2 < (1 - 5^{-\tilde{\sigma}})c_1, 1 - \delta < \sigma < \tilde{\sigma} \le 1, \delta > \iota > 0, 0 < \mu < \tilde{\sigma},$$

and

(21)
$$a \le \frac{1}{2} \min_{i} \left\{ \frac{\sigma - 1}{b_i} \delta + \frac{\delta^2}{b_i} \right\}.$$

Let \tilde{A}^m be given by (20) and $c = \frac{1}{2} \min\{\frac{1}{K_1}, \tilde{\sigma}\}$. Fix any sufficiently small $\varepsilon > 0$. Then there exists a large constant C depending on all parameters such that the following statements are true. Let N_1 be sufficiently large, $N_2 \in [N_1^C, e^{N_1^{c/2}}]$ and $N_3 \in [N_2^C, e^{N_1^c}]$. Assume the Green's function satisfies the property \tilde{P} with parameters (μ, ζ, c_2) at sizes N_1 and N_2 . Assume for any $L \in [N_3^{\delta-\iota}, N_3]$ and any $x \in \mathbb{T}^b$,

$$#\{n \in \mathbb{Z}^d : |n| \le L, f(n, x) \in X_{N_1} \mod \mathbb{Z}^b\} \le L^{1-\delta}.$$

Then there exists a subset $X_{N_3} \subset \mathbb{T}^b$ such that

$$\sup_{1 \le i \le k, x_i^{\sim} \in \mathbb{T}^{b-b_i}} \operatorname{Leb}(X_{N_3}(x_i^{\sim})) \le e^{-N_3 \frac{\sigma-1}{b_i}\delta + \frac{\delta^2}{b_i} - \varepsilon}.$$

and for any $x \notin X_{N_3}$, $m \in \mathbb{Z}^d$ and $Q_{N_3} \in \mathcal{E}^0_{N_3}$,

$$||(R_{Q_{N_3}}\tilde{A}^m(x)R_{Q_{N_3}})^{-1}|| \leq e^{N_3^{\sigma}},$$

and for $|n - n'| \ge \frac{N_3}{10}$,

$$|(R_{Q_{N_3}}\tilde{A}^m R_{Q_{N_3}})^{-1}(x;n,n')| \leq e^{-(c_2 - N_1^{-\vartheta_1} - N_3^{-\vartheta_2})|n-n'|^{\tilde{\sigma}}},$$

where $\vartheta_1 = \vartheta_1(\tilde{\sigma}, \mu, c)$ and $\vartheta_2 = \vartheta_2(\tilde{\sigma}, \sigma, \delta, \varepsilon)$.

Remark 2. (1) Theorem 2.3 improves the parameters from (μ, ζ, c_2) to

$$(\sigma, \frac{\sigma-1}{b_i}\delta + \frac{\delta^2}{b_i} - \varepsilon, c_2 - N_1^{-\vartheta_1} - N_3^{-\vartheta_2}).$$

Theorem 2.3 gives us opportunities to combine perturbative approaches with non-perturbative approaches. After establishing the property P for initial scales by non-perturbative methods, we can adapt the parameters to establish property P with explicit bounds for larger scales. See Theorems 3.1, 3.2 and 3.3, and Corollaries 3.4, 3.5 and 3.6 for examples.

(2) Roughly speaking Theorem 2.3 says that under the assumption on the sublinear bound, the large deviation theorem at sizes $N = N_1$ and $N = N_2$ will ensure the large deviation theorem at size $N = N_3$.

We are going to discuss the modulus of continuity of the integrated density of states (IDS). In order to make it as general as possible, we do not require the existence of the integrated density of states first. Let $E_1 < E_2$ and define (22)

$$k(x, E_1, E_2) = \limsup_{N \to \infty} \frac{1}{(2N+1)^d} \# \{ \text{ eigenvalues of } R_{[-N,N]^d} A(x) R_{[-N,N]^d} \text{ in } [E_1, E_2] \}.$$

Fix $x \in \mathbb{T}^b$. Assume for any measurable set $\mathcal{S} \subset \mathbb{T}^b$, we have (23)

$$\limsup_{N \to \infty} \frac{1}{(2N+1)^d} \# \{ n \in \mathbb{Z}^d : |n| \le N, f(n_1, n_2, \cdots, n_d, x) \in \mathcal{S} \} \le \operatorname{Leb}(\mathcal{S}).$$

For an operator A(x) on $\ell^2(\mathbb{Z}^d)$, denote by the energy dependent Green's functions

(24)
$$G_{\Lambda}(E,x) = (R_{\Lambda}(A(x) - E)R_{\Lambda})^{-1}.$$

Instead of $G_{\Lambda}(E, x)$, we will write G_{Λ} , $G_{\Lambda}(E)$, or $G_{\Lambda}(x)$ when there is no ambiguity. We will write $G_{\Lambda}(n, n')$, $G_{\Lambda}(E; n, n')$, $G_{\Lambda}(x; n, n')$, or $G_{\Lambda}(E, x; n, n')$ for the element of matrices.

Theorem 2.5. Assume A(x) is a Toeplitz (operator) matrix on $\ell^2(\mathbb{Z}^d)$ with respect to f in the sense of (14). Let $\zeta \in (0, 1)$ and $0 < \sigma < \tilde{\sigma} \leq 1$. Assume for any $E \in \mathbb{R}$, there exists a set $X_N \subset \mathbb{T}^b$ such that

$$\operatorname{Leb}(X_N) \le e^{-N^{\alpha}}$$

and for any $x \notin X_N$ and any $Q_N \in \mathcal{E}_N^0$,

$$||G_{Q_N}(E, x)|| \leq e^{N^{\sigma}} |G_{Q_N}(E, x; n, n')| \leq e^{-c|n-n'|^{\tilde{\sigma}}} \text{ for } |n-n'| \geq \frac{N}{10}$$

where c > 0. Assume (23) holds for some $x_0 \in \mathbb{T}^b$. Then for any $\varepsilon > 0$, we have

$$|k(x_0, E_1, E_2)| \le e^{-|\log|E_1 - E_2||\frac{\zeta}{\sigma} - \varepsilon},$$

provided that $|E_1 - E_2|$ is sufficiently small.

The rest of this paper is organized as follows. Except for some statements in applications (Section 3), this paper is entirely self contained. We will introduce many applications to quasi-periodic operators in Section 3. Sections 4, 5, 6, 7 are devoted to prove Theorems 2.1, 2.2, 2.3, 2.4 and 2.5. We will introduce the discrepancy for semi-algebraic sets in Section 8. In Section 9, we will give the proof for all the results in Section 3.

3. Applications

Let S be a Toeplitz (operator) matrix on $\ell^2(\mathbb{Z}^d)$ with respect to f, namely,

(25)
$$S(x; n+k, n'+k) = S(f^k(x); n, n'),$$

for any $n \in \mathbb{Z}^d$, $n' \in \mathbb{Z}^d$ and $k \in \mathbb{Z}^d$. Assume every element S(z; n, n'), $n, n' \in \mathbb{Z}^d$, is analytic in a strip $\{z : |\Im z| \le \rho\}$ with $\rho > 0$ and satisfies for any $x \in \mathbb{R}$ and $n, n' \in \mathbb{Z}^d$,

(26)
$$|S(x;n,n')| \le K e^{-c_1|n-n'|}, K > 0, c_1 > 0.$$

Assume that there exists $K_1 > 1$ such that for any $x \in \mathbb{T}^b$ and $z \in \{z \in \mathbb{C}^b : |\Im z| \le \rho\}$ with $||x - z|| \le e^{-(\log(|n| + |n'| + 2))^{K_1}}$,

(27)
$$|A(x;n,n') - A(z;n,n')| \le K ||x-z||^{\gamma}.$$

Assume for any N > 1, $n, n' \in \mathbb{Z}^d$ with $|n| \leq N$ and $|n'| \leq N$, there exists a trigonometric polynomial $\tilde{S}(x; n, n')$ of degree less than $e^{(\log N)^{K_1}}$ such that

(28)
$$\sup_{x \in \mathbb{T}^b} |S(x; n, n') - \tilde{S}(x; n, n')| \le K e^{-N^2}.$$

Define a family of operators H(x) on $\ell^2(\mathbb{Z}^d)$:

(29)
$$H(x) = \lambda^{-1}S + v(f(n,x))\delta_{nn'},$$

where v is an analytic function on \mathbb{T}^{b} .

In this section, we always assume

- v is non-constant,
- f is a frequency shift or skew-shift,
- except for subsection 3.6, S is a Toeplitz (operator) matrix on $\ell^2(\mathbb{Z}^d)$ with respect to f and satisfies (25)-(28).

Example 4:

• If S is a long range operator, namely, S does not depend on x and

$$S(n,n') \le K e^{-c_1|n-n'|}, n, n' \in \mathbb{Z}^d,$$

then (26), (27) and (28) hold.

• Let $\phi_k(x)$, $k \in \mathbb{Z}$, be a trigonometric polynomial on \mathbb{T}^b of degree less than $e^{(\log(1+|k|))^{K_1}}$ satisfying

$$\sup_{x \in \mathbb{T}^b} |\phi_k(x)| \le K e^{-c_1|k|}.$$

Let

$$S(x; n, n') = \phi_{n-n'}(f(n, x)) + \phi_{n'-n}(f(n', x)).$$

Then (26), (27) and (28) hold.

Remark 3. For $d \ge 2$, our settings (25)-(28) is the first time to allow every entry of S to depend on x, which beyonds the long range operators. For d = 1, Bourgain [14] studied the case in Example 4 under the assumption that $\phi_k(x)$ is a trigonometric polynomial of degree at most N^C .

We will apply Theorems 2.1, 2.2, 2.3 and 2.5 to operators

$$A(x) = H(x) = \lambda^{-1}S + v(f(n,x))\delta_{nn'}.$$

In this section, the Green's functions always depend on energy E. See (24).

The IDS appearing in applications is always existed, namely, the following limit

$$k(x,E) = \lim_{N \to \infty} \frac{1}{(2N+1)^d} \#\{ \text{ eigenvalues of } R_{[-N,N]^d} A(x) R_{[-N,N]^d} \text{ smaller than } E\},$$

converges to k(E) for almost every x. We write k(E) for the IDS when it exists.

For the large deviation theorem, S is not necessarily self-adjoint. However, in order to establish pure point spectrum property, self-adjointness is necessary because of the energy elimination.

3.1. Shifts: d = 1, arbitrary b. Denote by Δ the discrete Laplacian on $\ell^2(\mathbb{Z})$, that is, for $\{u(n)\} \in \ell^2(\mathbb{Z})$,

$$(\Delta u)(n) = \sum_{|n-n'|=1} u(n').$$

13

We say that $\omega = (\omega_1, \omega_2, \cdots, \omega_b)$ satisfies Diophantine condition $DC(\kappa, \tau)$, if

(30)
$$||k\omega|| \ge \frac{\tau}{|k|^{\kappa}}, k \in \mathbb{Z}^b \setminus \{(0, 0, \cdots, 0)\}$$

By the Dirichlet principle, one has $\kappa \geq b$. When $\kappa > b$, $\cup_{\tau>0} DC(\kappa, \tau)$ has full Lebesgue measure.

We say that $\omega \in \mathbb{R}$ satisfies strong Diophantine conditions if there exist $\kappa > 1$ and $\tau > 0$ such that

(31)
$$||k\omega|| \ge \frac{\tau}{k(1+\log k)^{\kappa}} \text{ for all } k \in \mathbb{N}.$$

It is easy to see that almost every ω satisfies strong Diophantine conditions. Let

$$f^n(x) = x + n\omega = (x_1 + n\omega_1, x_2 + n\omega_2, \cdots, x_b + n\omega_b) \mod \mathbb{Z}^b$$

where $x = (x_1, x_2, \dots, x_b) \in \mathbb{T}^b$, $n \in \mathbb{Z}$ and $\omega = (\omega_1, \omega_2, \dots, \omega_b) \in \mathbb{R}^b$. Let H(x) on $\ell^2(\mathbb{Z})$ be given by

(32)
$$H(x) = \Delta + v(f^n(x)) = \Delta + v(x_1 + n\omega_1, x_2 + n\omega_2, \cdots, x_b + n\omega_b)\delta_{nn'},$$

where $n, n' \in \mathbb{Z}$.

0

Let

(33)
$$A_k^E(x) = \prod_{j=k-1}^0 A^E(x+j\omega) = A^E(x+(k-1)\omega)A^E(x+(k-2)\omega)\cdots A^E(x)$$

and

(34)
$$A_{-k}^{E}(x) = (A_{k}^{E}(x-k\omega))^{-1}$$

for $k \ge 1$, where $A^E(x) = \begin{pmatrix} E - v(x) & -1 \\ 1 & 0 \end{pmatrix}$. A^E_k is called the (k-step) transfer matrix. The Lyapunov exponent is given by

(35)
$$L(E) = \lim_{k \to \infty} \frac{1}{k} \int_{\mathbb{T}^b} \ln \|A_k^E(x)\| dx.$$

Theorem 3.1. Let $\omega \in DC(\kappa, \tau)$ and $1 - \frac{1}{b\kappa} < \sigma < 1$. Let H(x) be given by (32). Assume the Lyapunov exponent L(E) is positive. Then for any $\varepsilon > 0$ and large N, there exists a subset $X_N \subset \mathbb{T}^b$ such that

$$\operatorname{Leb}(X_N) \le e^{-N^{\frac{\sigma-1}{b^2\kappa} + \frac{1}{b^3\kappa^2} - \varepsilon}},$$

and for any $x \notin X_N$, we have

$$||G_{[-N,N]}(E,x)|| \le e^{N^{\sigma}},$$

and

$$|G_{[N,-N]}(E,x;n,n')| \le e^{-(L(E)-\varepsilon)|n-n'|}$$
 for $|n-n'| \ge N/10$.

Theorem 3.2. Let $\omega \in DC(\kappa, \tau)$ and H(x) be given by (32). Suppose the Lyapunov exponent L(E) > 0 for every E in an interval I. Then for any $\varepsilon > 0$,

$$|k(E_1) - k(E_2)| \le e^{-\left(\log \frac{1}{|E_1 - E_2|}\right)^{\frac{1}{b^3 \kappa^2}}}$$

provided that $|E_1 - E_2|$ is sufficiently small and $E_1, E_2 \in I$.

Theorem 3.3. Let H(x) be given by (32). Then the following statement is true for almost every ω . Assume the Lyapunov exponent L(E) > 0 for every E in an interval I. Then for any $\varepsilon > 0$,

$$|k(E_1) - k(E_2)| \le e^{-\left(\log \frac{1}{|E_1 - E_2|}\right)^{\frac{1}{b^3} - \varepsilon}},$$

provided that $|E_1 - E_2|$ is sufficiently small and $E_1, E_2 \in I$.

Remark 4. Under the same assumptions, the large deviation theorem and the modulus of continuity of the IDS were shown in [35] (also see [15]). When b = 2, a better bound b = 1/3 was obtained in [35]. However, there are no explicit bounds in [15, 35] when $b \ge 3$.

Putting a coupling constant λ^{-1} in front of the Laplacian Δ , the operator given by (32) becomes

(36)
$$H(x) = \lambda^{-1} \Delta + v(x + n\omega) \delta_{nn'}.$$

For large λ only depending on the potential v, the Lyapunov exponent L(E) is positive for every E [16]. Therefore, we have the following three corollaries

Corollary 3.4. Assume $\omega \in DC(\kappa, \tau)$ and $1 - \frac{1}{b\kappa} < \sigma < 1$. Let H(x) be given by (36). Then there exists $\lambda_0 = \lambda_0(v)$ such that for any $\varepsilon > 0$, $\lambda > \lambda_0$ and large N, there exists $X_N \subset \mathbb{T}^b$ such that

(37)
$$\operatorname{Leb}(X_N) \le e^{-N\frac{\sigma-1}{b^2\kappa} + \frac{1}{b^3\kappa^2} - \varepsilon}.$$

and for any $x \notin X_N$, we have

$$||G_{[-N,N]}(E,x)|| \le e^{N^{\sigma}},$$

and

$$|G_{[N,-N]}(E,x;n,n')| \le e^{-(L(E)-\varepsilon)|n-n'|}$$
 for $|n-n'| \ge N/10$.

Corollary 3.5. Let $\omega \in DC(\kappa, \tau)$ and H(x) be given by (36). Then there exists $\lambda_0 = \lambda_0(v)$ such that for any $\varepsilon > 0$ and $\lambda > \lambda_0$,

$$|k(E_1) - k(E_2)| \le e^{-\left(\log \frac{1}{|E_1 - E_2|}\right)^{\frac{1}{b^3 \kappa^2} - \varepsilon}}$$

provided that $|E_1 - E_2|$ is sufficiently small.

Corollary 3.6. Let H(x) be given by (36). Then there exists $\lambda_0 = \lambda_0(v)$ such that the following statement is true for almost every ω . For any $\varepsilon > 0$ and $\lambda > \lambda_0$,

$$|k(E_1) - k(E_2)| \le e^{-\left(\log \frac{1}{|E_1 - E_2|}\right)^{\frac{1}{b^3} - \varepsilon}},$$

provided that $|E_1 - E_2|$ is sufficiently small.

Let H(x) on $\ell^2(\mathbb{Z})$ be given by

(38)
$$H(x) = \lambda^{-1}S + v(f^n(x)) = \lambda^{-1}S + v(x + n\omega)\delta_{nn'},$$

where $x, \omega \in \mathbb{R}^b$.

Theorem 3.7. Let H(x) be given by (38). Assume $\omega \in DC(\kappa, \tau)$ and $1 - \frac{1}{b\kappa} < \sigma < 1$. Then for any $\varepsilon > 0$, there exists

$$\lambda_0 = \lambda_0(\varepsilon, \kappa, \tau, \rho, \sigma, \gamma, K, K_1, c_1, v)$$

such that for any $\lambda > \lambda_0$ and any N, there exists $X_N \subset \mathbb{T}^b$ such that

$$\operatorname{Leb}(X_N) \le e^{-N\frac{\sigma-1}{b^2\kappa} + \frac{1}{b^3\kappa^2} - \varepsilon},$$

and for any $x \notin X_N$, we have

$$||G_{[-N,N]}(E,x)|| \le e^{N^{\sigma}},$$

and

$$|G_{[N,-N]}(E,x;n,n')| \le e^{-\frac{1}{2}c_1|n-n'|}$$
 for $|n-n'| \ge N/10$.

Theorem 3.8. Assume S is self-adjoint and $\omega \in DC(\kappa, \tau)$. Let H(x) be given by (38). Then for any $\varepsilon > 0$, there exists

$$\lambda_0 = \lambda_0(\varepsilon, \kappa, \tau, \rho, \gamma, K, K_1, c_1, v)$$

such that for any $\lambda > \lambda_0$,

$$|k(E_1) - k(E_2)| \le e^{-\left(\log \frac{1}{|E_1 - E_2|}\right)^{\frac{1}{b^3 \kappa^2} - \varepsilon}},$$

provided that $|E_1 - E_2|$ is sufficiently small.

Theorem 3.9. Assume S is self-adjoint. Let H(x) be given by (38). Then for almost every $\omega \in \mathbb{R}^b$ the following is true. For any $\varepsilon > 0$, there exists

$$\lambda_0 = \lambda_0(\varepsilon, \omega, \rho, \gamma, K, K_1, c_1, v)$$

such that for any $\lambda > \lambda_0$,

$$|k(E_1) - k(E_2)| \le e^{-\left(\log \frac{1}{|E_1 - E_2|}\right)^{\frac{1}{b^3} - \varepsilon}},$$

provided that $|E_1 - E_2|$ is sufficiently small.

Theorem 3.10. Let H(x) be given by (38). Then for any $\rho > 0$, there is $\lambda_0 = \lambda_0(\rho, \rho, \gamma, K, K_1, c_1, v) > 0$ such that the following statement holds. For any $\lambda > \lambda_0$ and any $x \in \mathbb{T}$, there exists $\Omega = \Omega(x, \lambda, S, v, \rho) \subset \mathbb{T}^b$ with $\text{Leb}(\mathbb{T}^b \setminus \Omega) \leq \rho$ such that for any $\omega \in \Omega$, H(x) satisfies Anderson localization.

3.2. Shifts: b = 1, arbitrary d. Let v be analytic on T. Let

$$f^n(x) = x + n\omega = x + n_1\omega_1 + n_2\omega_2 + \dots + n_d\omega_d \mod \mathbb{Z},$$

where $n = (n_1, n_2, \cdots, n_d) \in \mathbb{Z}^d$ and $x \in \mathbb{T}$. Let H(x) on $\ell^2(\mathbb{Z}^d)$ be given by

(39)
$$H(x) = \lambda^{-1}S + v(f^n(x))\delta_{nn'} = \lambda^{-1}S + v(x + n_1\omega_1 + n_2\omega_2 + \dots + n_d\omega_d)\delta_{nn'}.$$

Theorem 3.11. Let $\omega \in DC(\kappa, \tau)$ and H(x) be given by (39). Then for any $\varepsilon > 0$, there exists $\lambda_0 = \lambda_0(\varepsilon, \kappa, \tau, \rho, \sigma, \gamma, K, K_1, c_1, v)$ such that for any $\lambda > \lambda_0$ and any N, there exists $X_N \subset \mathbb{T}$ such that

(40)
$$\operatorname{Leb}(X_N) \le e^{-N^{\sigma-\varepsilon}},$$

and for any $x \notin X_N$ and any $Q_N \in \mathcal{E}_N^0$, we have

$$||G_{Q_N}(E,x)|| \le e^{N^{\sigma}},$$

and

(41)
$$|G_{Q_N}(E,x;n,n')| \le e^{-\frac{1}{2}c_1|n-n'|} \text{ for } |n-n'| \ge N/10.$$

Theorem 3.12. Assume S is self-adjoint and $\omega \in DC(\kappa, \tau)$. Let H(x) be given by (39). Then for any $\varepsilon > 0$, there exists

$$\lambda_0 = \lambda_0(\varepsilon, \kappa, \tau, \rho, \gamma, K, K_1, c_1, v)$$

such that for any $\lambda > \lambda_0$,

$$|k(E_1) - k(E_2)| \le e^{-\left(\log \frac{1}{|E_1 - E_2|}\right)^{1-\varepsilon}},$$

provided that $|E_1 - E_2|$ is sufficiently small.

Theorem 3.13. Assume S is self-adjoint. Let H(x) be given by (39). Then for any $\rho > 0$, there is $\lambda_0 = \lambda_0(\rho, \rho, \gamma, K, K_1, c_1, v) > 0$ such that the following statement holds. For any $\lambda > \lambda_0$ and any $x \in \mathbb{T}$, there exists $\Omega = \Omega(x, \lambda, S, v, \rho) \subset \mathbb{T}^d$ with Leb $(\mathbb{T}^d \setminus \Omega) \leq \rho$ such that for any $\omega \in \Omega$, H(x) satisfies Anderson localization.

Remark 5. Theorem 3.13 is a generalization of Theorem 2 in p.138 of [15] and main result in [27].

3.3. Skew-shifts: d = 1, arbitrary b. Let $f: \mathbb{T}^b \to \mathbb{T}^b$ be the skew-shift defined as follows

(42)
$$f(x_1, x_2, ..., x_b) = (x_1 + \omega, x_2 + x_1, ..., x_b + x_{b-1}).$$

Let H(x) on $\ell^2(\mathbb{Z})$ be given by

(43)
$$H(x) = \lambda^{-1}S(x) + v(f^n(x))\delta_{nn'},$$

v is analytic on \mathbb{T}^b .

Theorem 3.14. Let H(x) be given by (43). Assume $\omega \in DC(\kappa, \tau)$ and $1 - \frac{1}{2^{b-1}b\kappa} < \sigma < 1$. Then for any $\varepsilon > 0$, there exists $\lambda_0 = \lambda_0(\varepsilon, \kappa, \tau, \rho, \sigma, \gamma, K, K_1, c_1, v)$ such that for any $\lambda > \lambda_0$ and any N, there exists $X_N \subset \mathbb{T}^b$ such that

$$\text{Leb}(X_N) \le e^{-N^{\frac{\sigma-1}{2^{b-1}b^{2_{\kappa}}} + \frac{1}{4^{b-1}b^{3_{\kappa}2}} - \varepsilon}}$$

and for any $x \notin X_N$, we have

$$||G_{[-N,N]}(E,x)|| \le e^{N^{\sigma}}$$

and

$$|G_{[N,-N]}(E,x;n,n')| \le e^{-\frac{1}{2}c_1|n-n'|}$$
 for $|n-n'| \ge N/10$.

Remark 6. Under stronger assumptions that $\omega \in DC(2, \tau)$, v and each element of S are nonconstant trigonometric polynomials, the large deviation theorem appearing in Theorem 3.14 without explicit bounds was proved for d = 2 [15] and arbitrary d [67].

Theorem 3.15. Assume S is self-adjoint and $\omega \in DC(\kappa, \tau)$. Let H(x) be given by (43). Then for any $\varepsilon > 0$, there exists

$$\lambda_0 = \lambda_0(\varepsilon, \kappa, \tau, \rho, \gamma, K, K_1, c_1, v)$$

such that for any $\lambda > \lambda_0$, we have

$$|k(E_1) - k(E_2)| \le e^{-\left(\log \frac{1}{|E_1 - E_2|}\right)^{\frac{1}{4b-1}b^3\kappa^2 - \varepsilon}},$$

provided that $|E_1 - E_2|$ is sufficiently small.

Corollary 3.16. Assume S is self-adjoint. Let H(x) be given by (43). Then for almost every $\omega \in \mathbb{R}$ the following is true. For any $\varepsilon > 0$, there exists $\lambda_0 = \lambda_0(\varepsilon, \omega, \rho, \gamma, K, K_1, c_1, v)$ such that for any $\lambda > \lambda_0$,

$$|k(E_1) - k(E_2)| \le e^{-\left(\log \frac{1}{|E_1 - E_2|}\right)^{\frac{1}{4^{b-1}b^3} - \varepsilon}}$$

provided that $|E_1 - E_2|$ is sufficiently small.

Assume S is taken the particular case, i.e., $S = \Delta$. Let b = 2. In this case, by Corollary 3.16, $\frac{1}{4^{b-1}b^3} = \frac{1}{32}$. A bound $\frac{1}{24}$ was shown by Bourgain, Goldstein and Schlag [19]. By combining the arguments in Bourgain, Goldstein and Schlag [19] with the proof of Corollary 3.16, we are able to improve the bound.

Corollary 3.17. Assume S is self-adjoint. Let b = 2 and H(x) be given by (43). Then for almost every $\omega \in \mathbb{R}$ the following is true. For any $\varepsilon > 0$, there exists $\lambda_0 = \lambda_0(\varepsilon, \omega, \rho, \gamma, K, K_1, c_1, v)$ such that for any $\lambda > \lambda_0$,

$$|k(E_1) - k(E_2)| \le e^{-\left(\log \frac{1}{|E_1 - E_2|}\right)^{\frac{1}{18} - \varepsilon}},$$

provided that $|E_1 - E_2|$ is sufficiently small.

3.4. Skew-shifts: d = b = 1. Let P_b be the projection on the *b*th coordinate of \mathbb{T}^b , namely, $P_b(x_1, x_2, \cdots, x_b) = x_b$, where $(x_1, x_2, \cdots, x_b) \in \mathbb{R}^b$. Define H(x) on $\ell^2(\mathbb{Z})$,

(44)
$$H(x) = \lambda^{-1} \Delta + v(P_b(f^n(x)))\delta_{nn'},$$

where v is analytic on \mathbb{T} and f is the skew-shift on \mathbb{T}^b .

Theorem 3.18. Let H(x) be given by (44). Assume ω is strong Diophantine and $1 - \frac{1}{2^{b-1}b} < \sigma < 1$. Then there exists $\lambda_0 = \lambda_0(v)$ such that for any $\varepsilon > 0$, $\lambda > \lambda_0$ and large N, there exists $X_N \subset \mathbb{T}^b$ such that

Leb
$$(X_N) \le e^{-N^{\frac{\sigma-1}{2^{b-1}} + \frac{1}{4^{b-1}} - \varepsilon}}$$

and for any $x \notin X_N$, we have

$$||G_{[-N,N]}(E,x)|| \le e^{N}$$

and

$$|G_{[N,-N]}(E,x;n,n')| \le e^{-\frac{1}{2}c_1|n-n'|}$$
 for $|n-n'| \ge N/10$.

Theorem 3.19. Let ω be strong Diophantine and H(x) be given by (44). Then there exists $\lambda_0 = \lambda_0(v)$ such that for any $\varepsilon > 0$ and $\lambda > \lambda_0$,

(45)
$$|k(E_1) - k(E_2)| \le e^{-\left(\log \frac{1}{|E_1 - E_2|}\right)^{\frac{1}{4^{b-1}} - \varepsilon}}$$

provided that $|E_1 - E_2|$ is sufficiently small.

- **Remark 7.** Comparing to Theorems 3.14 and 3.15, there is no dimension (b^3) loss in the bounds of Theorems 3.18 and 3.19. This is because the potential v is defined on \mathbb{T} .
 - The large deviation theorem and the modulus of continuity of Lyapunov exponents (the IDS) without explicit bounds was obtained in [70].
 - Let b = 2. The constant in (45) becomes $\frac{1}{4^{b-1}} = \frac{1}{4}$. It is possible to improve the bound from 1/4 to 1/3 by incorporating the arguments in [19]. A weaker result was proved by Tao [71], where a constant $\frac{1}{30}$ was obtained.

3.5. Shifts: d = b = 2. Assume v is analytic on $\mathbb{T}^2 = (\mathbb{R}/\mathbb{Z})^2$. Let

$$f^n(x) = (x_1 + n_1\omega_1, x_2 + n_2\omega_2) \mod \mathbb{Z}^2,$$

where $n = (n_1, n_2) \in \mathbb{Z}^2$, $\omega = (\omega_1, \omega_2) \in \mathbb{R}^2$ and $x = (x_1, x_2) \in \mathbb{T}^2$. Let H(x) on $\ell^2(\mathbb{Z}^2)$ be given by (46) $H(x) = \lambda^{-1}S(x) + v(f^n(x))\delta_{nn'} = \lambda^{-1}S(x_1, x_2) + v(x_1 + n_1\omega_1, x_2 + n_2\omega_2)\delta_{nn'}.$

Theorem 3.20. Let H(x) be given by (46). Suppose v is nonconstant on any line segment contained in $[0,1)^2$, $\omega_1 \in DC(\kappa,\tau)$ and $\omega_2 \in DC(\kappa,\tau)$ with $1 \le \kappa < \frac{13}{12}$. Assume

$$3\kappa - \frac{9}{4} < \sigma < 1.$$

Then there exists $\lambda_0 = \lambda_0(\varepsilon, \kappa, \tau, \rho, \sigma, \gamma, K, K_1, c_1, v)$ such that for any $\lambda > \lambda_0$ and any N, there exists $X_N \subset \mathbb{T}^2$ such that for any line segment $L \subset [0, 1)^2$,

(47)
$$\operatorname{Leb}(X_N \cap L) \le e^{-N^{(\sigma-1)(13/4-3\kappa)+(13/4-3\kappa)^2-\varepsilon}}$$

and for any $x \notin X_N$ and $Q_N \in \mathcal{E}_N^0$, we have

$$||G_{Q_N}(E,x)|| \le e^{N^{\sigma}}$$

and

$$|G_{Q_N}(E, x; n, n')| \le e^{-\frac{1}{2}c_1|n-n'|}$$
 for $|n-n'| \ge N/10$.

Theorem 3.21. Assume S is self-adjoint, v is nonconstant on any line segment contained in $[0,1)^2$, $\omega_1 \in DC(\kappa,\tau)$ and $\omega_2 \in DC(\kappa,\tau)$ with $1 \le \kappa < \frac{13}{12}$. Let H(x)be given by (46). Then for any ε , there exists $\lambda_0 = \lambda_0(\varepsilon,\kappa,\tau,\rho,\gamma,K,K_1,c_1,v)$ such that for any $\lambda > \lambda_0$,

$$|k(E_1) - k(E_2)| \le e^{-\left(\log \frac{1}{|E_1 - E_2|}\right)^{(13/4 - 3\kappa)^2 - \varepsilon}},$$

provided that $|E_1 - E_2|$ is sufficiently small.

Corollary 3.22. Assume S is self-adjoint and v is nonconstant on any line segments contained in $[0, 1)^2$. Let H(x) be given by (46). Then for almost every $\omega \in \mathbb{R}^2$ the following is true. For any $\varepsilon > 0$, there exists $\lambda_0 = \lambda_0(\varepsilon, \omega, \rho, \gamma, K, K_1, c_1, v)$ such that for any $\lambda > \lambda_0$,

$$|k(E_1) - k(E_2)| \le e^{-\left(\log \frac{1}{|E_1 - E_2|}\right)^{\frac{1}{16} - \varepsilon}},$$

provided that $|E_1 - E_2|$ is sufficiently small.

Remark 8. Theorems 3.20 and 3.21 follow the arguments in [22]. Our quantitative approaches developed in the paper allow us to obtain the explicit bounds.

3.6. Sub-exponentially decaying matrices with interactions. Our applications can be wider. Here are several examples. Instead of (26), assume

(48)
$$|S(x;n,n')| \le K e^{-c_1 |n-n'|^{\sigma}}, 0 < \tilde{\sigma} \le 1, c_1 > 0,$$

for any $n, n' \in \mathbb{Z}^d$.

Assume for any N > 1, $n, n' \in \mathbb{Z}^d$ with $|n| \leq N$ and $|n'| \leq N$, there exists a trigonometric polynomial $\tilde{S}(x; n, n')$ of degree less than e^{N^a} such that

(49)
$$\sup_{x \in \mathbb{T}^b} |S(x;n,n') - \tilde{S}(x;n,n')| \le Ke^{-N^2}$$

In this subsection, assume S satisfies (27), (48) and (49). Let \tilde{U} be a diagonal matrix on $\ell^2(\mathbb{Z}^d)$ satisfying

$$||U|| \le K.$$

Given $m \in \mathbb{Z}^d$, define the diagonal matrix \tilde{U}^m on $\ell^2(\mathbb{Z}^d)$ by

$$\tilde{U}^m(n) = \tilde{U}(m+n), n \in \mathbb{Z}^d$$

We say \tilde{U} has low complexity if there exists 0 < a < 1 such that for any N > 1,

(50)
$$#\{R_{Q_N}U^m(n)\delta_{nn'}R_{Q_N}: m \in \mathbb{Z}^d, Q_N \in \mathcal{E}_N^0\} \le Ke^{N^a}$$

Let

(51)
$$\tilde{H}(x) = H(x) + \lambda^{-1}U + \tilde{U} = \lambda^{-1}(S+U) + (\tilde{U}(n) + v(f(n,x)))\delta_{nn'}.$$

For any $m \in \mathbb{Z}^d$, let

(52)
$$\tilde{H}^m(x) = H(x) + \lambda^{-1}U^m + \tilde{U}^m = \lambda^{-1}(S + U^m) + (\tilde{U}^m + v(f(n,x)))\delta_{nn'}.$$

Denote by \tilde{G}^m the Green's function of \tilde{H}^m .

Theorem 3.23. Assume α is strong Diophantine, and U and \tilde{U} have low complexity in the sense of (19) and (50) repectively. Assume

$$1 - \frac{1}{b} < \sigma < \tilde{\sigma} \text{ and } a \le \frac{1}{4} \left\{ \frac{1}{K_1}, \frac{\sigma - 1}{b^2} + \frac{1}{b^3} \right\}$$

Let H(x) and $\tilde{H}^m(x)$ be given by (38) and (52) respectively. Then for any $\varepsilon > 0$, there exists

$$\lambda_0 = \lambda_0(\varepsilon, \alpha, \rho, c_1, \sigma, \tilde{\sigma}, \gamma, K, K_1, c_1, v)$$

such that for any $\lambda > \lambda_0$ and any N, there exists $X_N \subset \mathbb{T}^b$ such that

$$\operatorname{Leb}(X_N) \le e^{-N^{\frac{\sigma-1}{b^2} + \frac{1}{b^3} - \varepsilon}},$$

and for any $x \notin X_N$ and $m \in \mathbb{Z}$, we have

$$||\tilde{G}^m_{[-N,N]}(E,x)|| \le e^{N^{\sigma}},$$

and

$$|\tilde{G}^m_{[-N,N]}(E,x;n,n')| \le e^{-\frac{c}{2}|n-n'|}$$
 for $|n-n'| \ge N/10$,

where $c = \frac{5^{\tilde{\sigma}} - 1}{5^{\tilde{\sigma}}}$.

Theorem 3.24. Assume $\omega \in DC(\kappa, \tau)$, U and \tilde{U} have low complexity. Assume $0 < \sigma < \tilde{\sigma} \leq 1$ and $a \leq \frac{1}{4}\min\{\frac{1}{K_1}, \sigma\}$. Let H(x) and $\tilde{H}^m(x)$ be given by (39) and (52) respectively. Then for any $\varepsilon > 0$, there exists

$$\lambda_0 = \lambda_0(\varepsilon, \kappa, \tau, \sigma, \tilde{\sigma}, \rho, \gamma, K, K_1, c_1, v)$$

such that for any $\lambda > \lambda_0$ and any N, there exists $X_N \subset \mathbb{T}$ such that

(53) $\operatorname{Leb}(X_N) \le e^{-N^{\sigma-\varepsilon}},$

and for any $x \notin X_N$, any $m \in \mathbb{Z}^d$ and any $Q_N \in \mathcal{E}_N^0$, we have $||\tilde{G}_{Q_N}^m(E, x)|| \leq e^{N^{\sigma}},$

and

(54)
$$|\tilde{G}_{Q_N}^m(E,x;n,n')| \le e^{-\frac{c}{2}|n-n'|^{\tilde{\sigma}}} \text{ for } |n-n'| \ge N/10,$$

where $c = \frac{5^{\tilde{\sigma}} - 1}{5^{\tilde{\sigma}}}$.

Theorem 3.25. Assume S is self-adjoint, $\omega \in DC(\kappa, \tau)$ and $a \leq \frac{1}{4}\min\{\frac{1}{K_1}, \tilde{\sigma}\}$. Let H(x) be given by (39). Then for any $\varepsilon > 0$, there exists

$$\lambda_0 = \lambda_0(\varepsilon, \kappa, \tau, \tilde{\sigma}, \rho, \gamma, K, K_1, c_1, v)$$

such that for any $\lambda > \lambda_0$,

$$|k(E_1) - k(E_2)| \le e^{-\left(\log \frac{1}{|E_1 - E_2|}\right)^{1-\varepsilon}},$$

provided that $|E_1 - E_2|$ is sufficiently small.

Using Theorem 2.4 instead of Theorem 2.3, the proof of Theorems 3.23, 3.24 and 3.25 follows from that of Theorems 3.7, 3.11 and 3.12 respectively. In order to avoid repetitions, we skip the details.

4. Multi-scale analysis

4.1. Exhaustion construction for an elementary region. For $m \in \mathbb{Z}^d$ and $\Lambda \subset \mathbb{Z}^d$, define the distance by

$$\operatorname{dist}(m,\Lambda) = \inf_{n \in \Lambda} |m - n|.$$

Fix an elementary region $\Lambda \in \mathcal{E}_N$. Let $x \in \Lambda$. Given $M \leq N/10$, we will construct exhaustion at x with width M. Set

$$\begin{split} \tilde{S}_0(x) &= (x + [-2M, 2M]^d) \cap \Lambda \\ \tilde{S}_j(x) &= \bigcup_{y \in S_{j-1}(x)} (y + [-4M, 4M]^d) \cap \Lambda, 1 \le j \le \tilde{l} \end{split}$$

where \tilde{l} is the minimum such that $\tilde{S}_{\tilde{l}}(x) = \Lambda$. We set $S_{-1}(x) = \emptyset$ for convenience.

When $\tilde{S}_{j-1}(x)$ is very close to the boundary of Λ , $\tilde{A}_j(x) = \tilde{S}_j(x) \setminus \tilde{S}_{j-1}(x)$ and \tilde{S}_j may have width less than M. However, there are at most finitely many j with $0 \leq j \leq \tilde{l}$, saying C(d), such that $\tilde{A}_j(x) = \tilde{S}_j(x) \setminus \tilde{S}_{j-1}(x)$ has width less than M, where C(d) is a constant depending on d.

We will delete j if $A_j(x) = S_j(x) \setminus S_{j-1}(x)$ has small width and then rearrange exhaustion. Here are the details. Let $j_0 \in \{0, 1, \dots, \tilde{l} - 1\}$ be the possibly smallest number such that both $\tilde{S}_{j_0}(x)$ and $\tilde{S}_{\tilde{l}}(x) \setminus \tilde{S}_{j_0}(x)$ have width at least M. Otherwise, set $j_0 = \tilde{l}$. Let $S_0(x) = \tilde{S}_{j_0}(x)$. Let $j_1 \in \{j_0, j_0 + 1, \dots, \tilde{l} - 1\}$ be the possibly smallest number such that both $\tilde{S}_{j_1}(x) \setminus \tilde{S}_{j_0}(x)$ and $\tilde{S}_{\tilde{l}}(x) \setminus \tilde{S}_{j_1}(x)$ have width at least M. Otherwise, set $j_1 = \tilde{l}$. Let $S_1(x) = \tilde{S}_{j_1}(x)$. Suppose we have defined $j_0, j_1 \cdots, j_k$ and corresponding $S_1(x), S_2(x), \dots S_k(x)$. Let $j_{k+1} \in \{j_k, j_k + 1, \dots, \tilde{l} - 1\}$ be the possibly smallest number such that $\tilde{S}_{j_{k+1}}(x) \setminus \tilde{S}_{j_k}(x)$ and $\tilde{S}_{\tilde{l}}(x) \setminus \tilde{S}_{j_{k+1}}(x)$ have width at least M. Otherwise, set $j_{k+1} = \tilde{l}$. Let $S_{k+1}(x) = \tilde{S}_{j_{k+1}}(x)$. Let l be such that $S_l(x) = \Lambda$. By our constructions, $\tilde{l} - C(d) \leq l \leq \tilde{l}$.

Here is an example. Assume x locates exactly at the left upmost corner. In Fig.3, $\tilde{A}_k(x) = \tilde{S}_k(x) \setminus \tilde{S}_{k-1}(x)$ and $\tilde{S}_{\tilde{l}}(x) = \tilde{S}_{\tilde{l}}(x) \setminus \tilde{S}_{\tilde{l}-1}(x)$ are the only two annuli which have width less than M. Therefore,

• $l = \tilde{l} - 2$ • For $j = 0, 1, 2, \cdots, k - 2, S_j(x) = \tilde{S}_j(x)$. • For $j = k - 1, k - 2, \cdots, l - 3, S_j(x) = \tilde{S}_{j+1}(x)$. $S_{l-2}(x) = \tilde{S}_{\tilde{l}}(x)$.

For any elementary region Λ , $x \in \Lambda$ and M, we call $\{S_j(x)\}_{j=0}^l$ the exhaustion of Λ at x with width M. We call $A_j(x) = S_j(x) \setminus S_{j-1}(x)$ the *j*th annulus. For any $y \in S_j(x) \setminus S_{j-1}(x)$, $j = 1, 2, \dots, l$, one has

(55)
$$4(j-1)M \le |y-x| \le 4jM + C(d)M.$$

By our constructions, any $\{A_j(x)\}$ has width at least M. Namely, for any $n \in A_j(x)$ there exists $W(n) \in \mathcal{E}_M$ such that

$$n \in W(n) \subset A_j(x)$$

and

dist
$$(n, A_j(x) \setminus W(n)) \ge M/2.$$




Fig.3: Exhaustion Construction

4.2. Resolvent identities. For simplicity, assume K = 1, namely

(56)
$$|A(n,n')| \le e^{-c_1|n-n'|^{\sigma}}, 0 < \tilde{\sigma} \le 1, c_1 > 0,$$

for any $n, n' \in \mathbb{Z}^d$. For any $\Lambda \subset \mathbb{Z}^d$, denote by $A_{\Lambda} = R_{\Lambda}AR_{\Lambda}$, where R_{Λ} is the restriction on Λ , and the Green's function

$$G_{\Lambda} = (R_{\Lambda} A R_{\Lambda})^{-1},$$

provided $R_{\Lambda}AR_{\Lambda}$ is invertible. Denote by $G_{\Lambda}(n, n')$ its elements, $n, n' \in \Lambda \subset \mathbb{Z}^d$. Assume Λ_1 and Λ_2 are wo disjoint subsets of \mathbb{Z}^d . Namely, $\Lambda_1, \Lambda_2 \subset \mathbb{Z}^d$ and

Assume Λ_1 and Λ_2 are wo disjoint subsets of \mathbb{Z}^a . Namely, $\Lambda_1, \Lambda_2 \subset \mathbb{Z}^a$ and $\Lambda_1 \cap \Lambda_2 = \emptyset$. Let $\Lambda = \Lambda_1 \cup \Lambda_2$. Suppose that $R_{\Lambda}AR_{\Lambda}$ and $R_{\Lambda_i}AR_{\Lambda_i}$, i = 1, 2 are invertible. Then

$$G_{\Lambda} = G_{\Lambda_1} + G_{\Lambda_2} - (G_{\Lambda_1} + G_{\Lambda_2})(A_{\Lambda} - A_{\Lambda_1} - A_{\Lambda_2})G_{\Lambda}.$$

If $m \in \Lambda_1$ and $n \in \Lambda$, we have (57)

$$|G_{\Lambda}(m,n)| \le |G_{\Lambda_1}(m,n)|\chi_{\Lambda_1}(n) + \sum_{n' \in \Lambda_1, n'' \in \Lambda_2} e^{-c_1|n'-n''|^{\tilde{\sigma}}} |G_{\Lambda_1}(m,n')| |G_{\Lambda}(n'',n)|.$$

If $n \in \Lambda_2$ and $m \in \Lambda$, we have (58)

$$|G_{\Lambda}(m,n)| \le |G_{\Lambda_2}(m,n)|\chi_{\Lambda_2}(n) + \sum_{n' \in \Lambda_1, n'' \in \Lambda_2} e^{-c_1|n'-n''|^{\tilde{\sigma}}} |G_{\Lambda}(m,n')| |G_{\Lambda_2}(n'',n)|.$$

Lemma 4.1 (Schur test). Suppose $A = A_{ij}$ is a matrix. Then

$$||A|| \le \sqrt{\left(\sup_{i} \sum_{j} |A_{ij}|\right) \left(\sup_{j} \sum_{i} |A_{ij}|\right)}.$$

The following lemma is a generalization of Lemma 3.2 in [51].

Lemma 4.2. Let $c_2 \in [\tilde{c}_1, c_1]$, $\sigma < \tilde{\sigma}$ and $M_0 \leq M_1 \leq N$. Assume Λ is a subset of \mathbb{Z}^d with diam $(\Lambda) \leq 2N + 1$. Suppose that for any $n \in \Lambda$, there exists some $W = W(n) \in \mathcal{E}_M$ with $M_0 \leq M \leq M_1$ such that $n \in W \subset \Lambda$, dist $(n, \Lambda \setminus W) \geq \frac{M}{2}$ and

(59)
$$||G_{W(n)}|| \le 2e^{M^{\sigma}}$$

(60)
$$|G_{W(n)}(n,n')| \le 2e^{-c_2|n-n'|^{\tilde{\sigma}}} \text{ for } |n-n'| \ge \frac{M}{10}$$

,

We assume further that M_0 is large enough so that

(61)
$$\sup_{M_0 \le M \le M_1} \sup_{c_2 \in [\tilde{c}_1, c_1]} 2e^{M^{\sigma}} (2M+1)^d e^{\frac{c_2}{10^{\tilde{\sigma}}}M^{\tilde{\sigma}}} \sum_{j=0}^{\infty} (M+2j+1)^d e^{-c_2(j+M/2)^{\tilde{\sigma}}} \le \frac{1}{2}.$$

Then

$$||G_{\Lambda}|| \le 4(2M_1+1)^d e^{M_1^{\sigma}}.$$

Proof. Under the assumption of (61), it is easy to check that for any M with $M_0 \leq M \leq M_1$ and any $n \in \Lambda$,

(62)
$$2(2M+1)^{d}e^{M^{\sigma}+\frac{c_{2}}{10^{\tilde{\sigma}}}M^{\tilde{\sigma}}}\sum_{\substack{n_{2}\in\Lambda\\|n_{2}-n|\geq \frac{M}{2}}}e^{-c_{2}|n-n_{2}|^{\tilde{\sigma}}}\leq\frac{1}{2}.$$

By (59) and (60), one has

(63)
$$|G_{W(n)}(n,n')| \le 2e^{M^{\sigma} + \frac{c_2}{10^{\tilde{\sigma}}}M^{\tilde{\sigma}}}e^{-c_2|n-n'|^{\tilde{\sigma}}}$$

For each $n \in \Lambda$, applying (57) with $\Lambda_1 = W(n)$, one has

$$|G_{\Lambda}(n,n')| \le |G_{W(n)}(n,n')|\chi_{W(n)}(n') + \sum_{\substack{n_1 \in W(n)\\n_2 \in \Lambda \setminus W(n)}} e^{-c_1|n_1 - n_2|^{\tilde{\sigma}}} |G_{W(n)}(n,n_1)| |G_{\Lambda}(n_2,n')|$$

It is easy to see for $0 < \tilde{\sigma} \leq 1$,

(64)
$$|x+y|^{\tilde{\sigma}} \le |x|^{\tilde{\sigma}} + |y|^{\tilde{\sigma}}.$$

By (63) and the fact that $|W(n)| \leq (2M+1)^d$, one has

$$\begin{aligned} |G_{\Lambda}(n,n')| \leq &|G_{W(n)}(n,n')|\chi_{W(n)}(n') \\ &+ 2\sum_{\substack{n_{1}\in W(n)\\n_{2}\in\Lambda\setminus W(n)}} e^{M^{\sigma} + \frac{c_{2}}{10^{\sigma}}M^{\tilde{\sigma}}} e^{-c_{2}|n-n_{1}|^{\tilde{\sigma}}} e^{-c_{1}|n_{1}-n_{2}|^{\tilde{\sigma}}} |G_{\Lambda}(n_{2},n')| \\ \leq &|G_{W(n)}(n,n')|\chi_{W(n)}(n') \\ &+ 2(2M+1)^{d} e^{M^{\sigma} + \frac{c_{2}}{10^{\sigma}}M^{\tilde{\sigma}}} \sum_{\substack{n_{2}\in\Lambda\setminus W(n)\\n_{2}\in\Lambda\setminus W(n)}} e^{-c_{2}|n-n_{2}|^{\tilde{\sigma}}} |G_{\Lambda}(n_{2},n')| \\ \leq &|G_{W(n)}(n,n')|\chi_{W(n)}(n') \\ &+ 2(2M+1)^{d} e^{M^{\sigma} + \frac{c_{2}}{10^{\sigma}}M^{\tilde{\sigma}}} \sum_{\substack{n_{2}\in\Lambda\\|n_{2}-n|\geq \frac{M}{2}}} e^{-c_{2}|n-n_{2}|^{\tilde{\sigma}}} |G_{\Lambda}(n_{2},n')|. \end{aligned}$$
(65)

where the second inequality holds by (64) and the last inequality holds by the assumption dist $(n, \Lambda \setminus W(n)) \ge \frac{M}{2}$. Summing over $n' \in \Lambda$ in (65) and noticing (62) yields

(66)
$$\sup_{n \in \Lambda} \sum_{n' \in \Lambda} |G_{\Lambda}(n, n')| \leq 2(2M_1 + 1)^d e^{M_1^{\sigma}} + \frac{1}{2} \sup_{n_2 \in \Lambda} \sum_{n' \in \Lambda} |G_{\Lambda}(n_2, n')|.$$

Similarly, using (58) instead of (57), one has

(67)
$$\sup_{n \in \Lambda} \sum_{n' \in \Lambda} |G_{\Lambda}(n', n)| \le 2(2M_1 + 1)^d e^{M_1^{\sigma}} + \frac{1}{2} \sup_{n_2 \in \Lambda} \sum_{n' \in \Lambda} |G_{\Lambda}(n', n_2)|.$$

Now the lemma follows from (66), (67) and Lemma 4.1.

4.3. Proof of Theorem 2.1.

Proof. Choose a constant $\rho \in (1, 1 + \tilde{\sigma} - \sigma)$. Calculation shows $\rho \sigma < \tilde{\sigma}$.

Define inductively $M_{j+1} = \lfloor M_j^{\rho} \rfloor$, $M_0 = M$. Let $\gamma_0 = c_2$. Fix an elementary region $\tilde{\Lambda}_1 \in \mathcal{E}_{M_1}$ and $\tilde{\Lambda}_1 \subset \tilde{\Lambda}_0$. For any $x \in \tilde{\Lambda}_1$, consider the exhaustion $\{S_j(x)\}_{j=0}^l$ of $\tilde{\Lambda}_1$ at x with width M_0 . Denote by $\{A_k(x)\}$ the annuli.

We call the annulus $A_k(x)$ good, if for any $y \in A_k(x)$, there exists $W(y) \in \mathcal{E}_{M_0}$ such that

$$y \in W(y) \subset A_k(x), \operatorname{dist}(y, A_k(x) \setminus W(y)) \ge M_0/2,$$

and for $|n - n'| \ge \frac{M_0}{10}$,

(68)
$$|(R_{W(y)}AR_{W(y)})^{-1}(n,n')| \le e^{-\gamma_0|n-n'|^{\tilde{\sigma}}}$$

Otherwise, we call the annulus A_k bad.

Fix $\kappa > 0$, which will be determined later. An elementary region $\Lambda_1 \subset \Lambda_0$ is called bad if there exists $x \in \tilde{\Lambda}_1$ such that the number of bad annuli $\{A_k(x)\}$ exceeds

$$B_1 := \kappa \frac{M_1}{M_0}.$$

Otherwise, we call Λ_1 good. Let \mathcal{F}_1 be an arbitrary family of pairwise disjoint bad elementary regions in \mathcal{E}_{M_1} contained in $\tilde{\Lambda}_0$. Since every annulus in $\{A_k\}$ has width at least M_0 by our construction, one has that every bad annulus contains at least one elementary region in \mathcal{E}_{M_0} without satisfying (68) and hence

(69)
$$\#\mathcal{F}_1 \le \frac{N^{\varsigma}}{\kappa M_1}$$

Assume that $\Lambda_1 \subset \Lambda_0$ is a good elementary region in \mathcal{E}_{M_1} . We will first show that $\tilde{\Lambda}_1$ is in class G with slightly smaller γ_0 . Consider the exhaustion $\{S_j(x)\}$ of $\tilde{\Lambda}_1$ at x with width M_0 . By the assumption, there are no more than B_1 bad annuli in this exhaustion. Denote by $\{A_j(x)\}_{j=0}^l$ annuli. By putting adjacent good annuli or bad annuli together, we obtain a new exhaustion

(70)
$$\emptyset = J_{-1} \subset J_0 \subset J_1 \subset \cdots \subset J_g = \tilde{\Lambda}_1.$$

More precisely, $\{J_s(x)\}, s = 0, 1, 2, \cdots, g$, satisfies the following rules.

•
$$J_s(x) \setminus J_{s-1}(x) = \{A_j(x)\}_{j=t_s}^{j=t'_s} \text{ for some } t_s < t'_s.$$

• $x \in J_0(x).$

- The annuli $A_j(x)$, $j = t_s, t_s + 1, \dots, t'_s$ are either all good or all bad.
- Take $J_s(x)$ maximal with the above three properties.

We remind that $J_s(x) \setminus J_{s-1}(x)$ has width at least M_0 for any $s = 0, 1, 2, \dots, g$. By our construction, if all annuli in $J_s(x) \setminus J_{s-1}(x)$ are good (bad), then all annuli in $J_{s+1}(x) \setminus J_s(x)$ are bad (good).

For any $n \in \Lambda_1$, let k(n) be the number of good annuli between x and n. Namely, for any $n \in A_j(x)$,

$$k(n) = \#\{A_t(x) : A_t(x) \text{ is a good annulus}, 0 \le t \le j\}.$$

Before we start the estimates, let us give several facts first, which will be used constantly in the later proof. By our constructions, J_s is a generalized elementary region, $s = 0, 1, \dots, g$. By the assumption (4), one has for all $s = 0, 1, \dots, g$,

(71)
$$(R_{J_s}AR_{J_s})^{-1} \le e^{M_1^{\sigma}}$$

Assume

$$0 < c \leq (1 - 5^{-\tilde{\sigma}})c_1.$$

If
$$|n - n_2| \ge \frac{M}{2}$$
 and $|n - n_1| \le \frac{M}{10}$, one has
 $c_1 |n_1 - n_2|^{\tilde{\sigma}} \ge c_1 (|n - n_2|^{\tilde{\sigma}} - |n - n_1|^{\tilde{\sigma}})$
(72) $\ge c |n - n_2|^{\tilde{\sigma}}.$

(73)
$$4M_0k(n_1) + |n_1 - n_2| \ge 4M_0k(n_2) - 4M_0k(n_2)$$

Without loss of generality, assume all the annuli in J_0 are bad (the another case is similar).

For any $n \in \tilde{\Lambda}_1$, define

$$\Gamma_s(n) = \max\{4M_0k(n) - 10(s+1)M_0, 0\}$$

By (73), we have for any $n_1 \in \tilde{\Lambda}_1$ and $n_2 \in \tilde{\Lambda}_1$,

(74)
$$\Gamma_s(n_1) + |n_2 - n_1| \ge \max\{\Gamma_s(n_2) - 4M_0, 0\}.$$

We shall inductively obtain estimates of the form

(75)
$$|G_{J_s(x)}(x,z)| \le T_s e^{-\gamma_0 \Gamma_s^{\sigma}(z)},$$

where $z \in J_s$, $s = 0, 1, \cdots, g$.

First step: s = 0

Since all annuli in J_0 are bad, one has k(z) = 0 and hence

(76)
$$\Gamma_0(z) = 0.$$

By (71) and (76), one has for $z \in J_0(x)$,

$$\begin{array}{lcl} G_{J_s}(x,z)| &\leq & e^{M_1^{\sigma}} \\ &= & e^{M_1^{\sigma}} e^{-\gamma_0 \Gamma_0^{\tilde{\sigma}}(z)} \end{array}$$

It implies that (75) holds for

$$(77) T_0 = e^{M_1^{\sigma}}.$$

Assume (75) holds at sth step for a proper T_s .

Case 1: All annuli in $J_{s+1} \setminus J_s$ are bad.

Pick any $z \in J_{s+1}$. Let $\tilde{n}_1 \in J_s$ and $\tilde{n}_2 \in J_{s+1} \setminus J_s$ be such that

$$\Gamma_s(\tilde{n}_1) + |\tilde{n}_1 - \tilde{n}_2| = \inf_{\substack{n_1 \in J_s \\ n_2 \in J_{s+1} \setminus J_s}} (\Gamma_s(n_1) + |n_1 - n_2|).$$

Case 1₁: $z \in J_{s+1} \setminus J_s$. In this case, for any $n_2 \in J_{s+1} \setminus J_s$, one has

(78)
$$k(z) = k(n_2), \Gamma_s(z) = \Gamma_s(n_2),$$

since all annuli in $J_{s+1} \setminus J_s$ are bad.

Applying (57) $(\Lambda_1 = J_s \text{ and } \Lambda_2 = J_{s+1} \setminus J_s)$, one has

$$\begin{aligned} |G_{J_{s+1}}(x,z)| &\leq \sum_{\substack{n_1 \in J_s \\ n_2 \in J_{s+1} \setminus J_s}} |G_{J_s}(x,n_1)| e^{-c_1 |n_1 - n_2|^{\tilde{\sigma}}} |G_{J_{s+1}}(n_2,z)| \\ &\leq \sum_{\substack{n_1 \in J_s \\ n_2 \in J_{s+1} \setminus J_s}} T_s e^{-\gamma_0 \Gamma_s^{\tilde{\sigma}}(n_1)} e^{-\gamma_0 |n_1 - n_2|^{\tilde{\sigma}}} |G_{J_{s+1}}(n_2,z)| \\ &\leq e^{M_1^{\sigma}} T_s \sum_{\substack{n_1 \in J_s \\ n_2 \in J_{s+1} \setminus J_s}} e^{-\gamma_0 \Gamma_s^{\tilde{\sigma}}(n_1)} e^{-\gamma_0 |n_1 - n_2|^{\tilde{\sigma}}} \\ &\leq (2M_1 + 1)^{2d} e^{M_1^{\sigma}} T_s \sup_{\substack{n_1 \in J_s \\ n_2 \in J_{s+1} \setminus J_s}} e^{-\gamma_0 \Gamma_s^{\tilde{\sigma}}(n_1)} e^{-\gamma_0 |n_1 - n_2|^{\tilde{\sigma}}} \\ &\leq (2M_1 + 1)^{2d} e^{M_1^{\sigma}} T_s e^{-\gamma_0 (\Gamma_s(\tilde{n}_1) + |\tilde{n}_1 - \tilde{n}_2|)^{\tilde{\sigma}}} \\ &\leq (2M_1 + 1)^{2d} e^{M_1^{\sigma}} T_s e^{-\gamma_0 (\max\{\Gamma_s(\tilde{n}_2) - 4M_0, 0\})^{\tilde{\sigma}}}, \\ &\leq (2M_1 + 1)^{2d} e^{M_1^{\sigma}} T_s e^{-\gamma_0 (\max\{\Gamma_s(z) - 4M_0, 0\})^{\tilde{\sigma}}}, \end{aligned}$$

where the second inequality holds by the induction (75) and $\gamma_0 \leq c_1$, the third inequality holds by (71), the fifth inequality holds by (64), the sixth inequality holds (74), and the last inequality holds by (78).

Case 1_2 : $z \in J_s$.

In this case, we have for any $n_2 \in J_{s+1} \setminus J_s$,

(80)
$$k(n_2) \ge k(z).$$

Applying (57) ($\Lambda_1 = J_s$ and $\Lambda_2 = J_{s+1} \setminus J_s$), one has

$$\begin{aligned} |G_{J_{s+1}}(x,z)| &\leq |G_{J_s}(x,z)| + \sum_{\substack{n_1 \in J_s \\ n_2 \in J_{s+1} \setminus J_s}} |G_{J_s}(x,n_1)| e^{-c_1 |n_1 - n_2|^{\tilde{\sigma}}} |G_{J_{s+1}}(n_2,z)| \\ &\leq T_s e^{-\gamma_0 \Gamma_s^{\tilde{\sigma}}(z)} + \sum_{\substack{n_1 \in J_s \\ n_2 \in J_{s+1} \setminus J_s}} T_s e^{-\gamma_0 \Gamma_s^{\tilde{\sigma}}(n_1)} e^{-\gamma_0 |n_1 - n_2|^{\tilde{\sigma}}} |G_{J_{s+1}}(n_2,z)| \\ &\leq T_s e^{-\gamma_0 \Gamma_s^{\tilde{\sigma}}(z)} + (2M_1 + 1)^{2d} e^{M_1^{\sigma}} T_s \sup_{\substack{n_1 \in J_s \\ n_2 \in J_{s+1} \setminus J_s}} e^{-\gamma_0 (\Gamma_s(n_1) + |n_1 - n_2|)^{\tilde{\sigma}}} \\ &= T_s e^{-\gamma_0 \Gamma_s^{\tilde{\sigma}}(z)} + (2M_1 + 1)^{2d} e^{M_1^{\sigma}} T_s e^{-\gamma_0 (\Gamma_s(\tilde{n}_1) + |\tilde{n}_1 - \tilde{n}_2|)^{\tilde{\sigma}}} \\ &\leq T_s e^{-\gamma_0 \Gamma_s^{\tilde{\sigma}}(z)} + (2M_1 + 1)^{2d} e^{M_1^{\sigma}} T_s e^{-\gamma_0 (\max\{\Gamma_s(\tilde{n}_2) - 4M_0, 0\})^{\tilde{\sigma}}}, \\ &\leq T_s e^{-\gamma_0 \Gamma_s^{\tilde{\sigma}}(z)} + (2M_1 + 1)^{2d} e^{M_1^{\sigma}} T_s e^{-\gamma_0 (\max\{\Gamma_s(z) - 4M_0, 0\})^{\tilde{\sigma}}}, \\ &\leq 2(2M_1 + 1)^{2d} e^{M_1^{\sigma}} T_s e^{-\gamma_0 (\max\{\Gamma_s(z) - 4M_0, 0\})^{\tilde{\sigma}}}, \end{aligned}$$

$$(81)$$

where the second inequality holds by the induction (75), the third inequality holds by (71) and (64), the forth inequality holds by (74), and the fifth inequality holds by (80).

Case 2: All the annuli in $J_{s+1} \setminus J_s$ are good.

By our constructions, $J_{s+1} \setminus J_s$ has width at least M_0 . Therefore, for any $k \in J_{s+1} \setminus J_s$, there exists some $W = W(k) \in \mathcal{E}_{M_0}$ such that $k \in W \subset \Lambda$,

(82)
$$\operatorname{dist}(k, J_{s+1} \setminus J_s \setminus W) \ge \frac{M_0}{2},$$

and

(83)
$$||G_{W(k)}|| \le e^{M_0^{\sigma}},$$

(84)
$$|G_{W(k)}(n_1, n_2)| \le e^{-\gamma_0 |n_1 - n_2|^{\tilde{\sigma}}} \text{ for } |n_1 - n_2| \ge \frac{M_0}{10},$$

where (83) holds by the assumption (4).

Since M_0 is large enough, one has (61) is satisfied. Applying Lemma 4.2, we have

(85)
$$\|G_{J_{s+1}\setminus J_s}\| \le 4(2M_0+1)^d e^{M_0^{\sigma}}.$$

We remark that we can not use the assumption (4) to bound $G_{J_{s+1}\setminus J_s}$ since $J_{s+1}\setminus J_s$ is not necessary to be a generalized elementary region. It is worth to point out that $J_{s+1}\setminus J_s$ may not be connected.

We will first prove that for any $m, n \in J_{s+1} \setminus J_s$,

(86)
$$|G_{J_{s+1}\setminus J_s}(m,n)| \le M_1^{10dM_1^{\tilde{\sigma}}M_0^{\tilde{\sigma}-\tilde{\sigma}}} e^{-\gamma_0(\max\{|m-n|-2M_0,0\})^{\tilde{\sigma}}}.$$

Assume $|m - n| \le 2M_0$. (86) holds by (85).

Assume $|m - n| > 2M_0$. Applying (57) with $\Lambda_1 = W(m)$ and using that $|m - n| > 2M_0$, one has

$$(87) \quad |G_{J_{s+1}\setminus J_s}(m,n)| \leq \sum_{\substack{n_1 \in W(m)\\n_2 \in J_{s+1}\setminus J_s\setminus W(m)}} e^{-c_1|n_1-n_2|^{\tilde{\sigma}}} |G_{W(m)}(m,n_1)| |G_{\Lambda}(n_2,n)|.$$

Applying (82) with k = m and by (72), one has for any n_1 with $|n_1 - m| \leq \frac{M_0}{10}$ and $n_2 \in J_{s+1} \setminus J_s \setminus W(m)$, one has

(88)
$$c_1|n_1-n_2|^{\tilde{\sigma}} \ge c_2|m-n_2|^{\tilde{\sigma}}.$$

By (83), (84) and (87), we have

$$\begin{aligned} |G_{J_{s+1}\setminus J_{s}}(m,n)| &\leq \sum_{\substack{n_{1}\in W(m),|n_{1}-m|\leq \frac{M_{0}}{10}-1\\n_{2}\in J_{s+1}\setminus J_{s}\setminus W(m)}}} e^{-c_{1}|n_{1}-n_{2}|^{\tilde{\sigma}}}|G_{W(m)}(m,n_{1})||G_{J_{s+1}\setminus J_{s}}(n_{2},n)| \\ &+ \sum_{\substack{n_{1}\in W(m),|n_{1}-m|\leq \frac{M_{0}}{10}\\n_{2}\in J_{s+1}\setminus J_{s}\setminus W(m)}}} e^{-c_{1}|n_{1}-n_{2}|^{\tilde{\sigma}}}|G_{W(m)}(m,n_{1})||G_{J_{s+1}\setminus J_{s}}(n_{2},n)| \\ &\leq \sum_{\substack{n_{1}\in W(m),|n_{1}-m|\leq \frac{M_{0}}{10}\\n_{2}\in J_{s+1}\setminus J_{s}\setminus W(m)}} e^{A_{0}^{\sigma}}e^{-c_{1}|n_{1}-n_{2}|^{\tilde{\sigma}}}|G_{J_{s+1}\setminus J_{s}}(n_{2},n)| \\ &+ \sum_{\substack{n_{1}\in W(m),|n_{1}-m|\leq \frac{M_{0}}{10}\\n_{2}\in J_{s+1}\setminus J_{s}\setminus W(m)}} e^{-c_{1}|n_{1}-n_{2}|^{\tilde{\sigma}}}e^{-\gamma_{0}|m-n_{1}|^{\tilde{\sigma}}}|G_{J_{s+1}\setminus J_{s}}(n_{2},n)| \\ &\leq \sum_{\substack{n_{1}\in W(m),|n_{1}-m|\leq \frac{M_{0}}{10}}e^{-\gamma_{0}|m-n_{2}|^{\tilde{\sigma}}}|G_{J_{s+1}\setminus J_{s}}(n_{2},n)| \\ &+ \sum_{\substack{n_{1}\in W(m),|n_{1}-m|\leq \frac{M_{0}}{10}}e^{-\gamma_{0}|m-n_{2}|^{\tilde{\sigma}}}|G_{J_{s+1}\setminus J_{s}}(n_{2},n)| \\ &+ \sum_{\substack{n_{1}\in W(m),|n_{1}-m|\leq \frac{M_{0}}{10}}e^{-\gamma_{0}|m-n_{2}|^{\tilde{\sigma}}}|G_{J_{s+1}\setminus J_{s}}(n_{2},n)| \\ &+ \sum_{\substack{n_{1}\in W(m),|n_{1}-m|\geq \frac{M_{0}}{10}}e^{-\gamma_{0}|m-n_{2}|^{\tilde{\sigma}}}|G_{J_{s+1}\setminus J_{s}}(n_{2},n)| \\ &\leq (2M_{1}+1)^{2d}e^{M_{0}^{\sigma}}\sum_{n_{2}\in J_{s+1}\setminus J_{s}\setminus W(m)}e^{-\gamma_{0}|m-n_{2}|^{\tilde{\sigma}}}|G_{J_{s+1}\setminus J_{s}}(n_{2},n)|, \end{aligned}$$

where the third inequality holds because of (88). Recall that $|m - n_2| \ge \frac{M_0}{2}$. Iterating (89) until $|n_2 - n| \le 2M_0$ or at most $\lfloor \frac{2\tilde{\sigma}|m-n|^{\tilde{\sigma}}}{M_0^{\tilde{\sigma}}} \rfloor + 1$ times, we have

$$\begin{aligned} |G_{J_{s+1}\setminus J_s}(m,n)| &\leq e^{M_0^{\sigma} \left(\frac{2^{\tilde{\sigma}}|m-n|^{\tilde{\sigma}}}{M_0^{\tilde{\sigma}}}+1\right)} (2M_1+1)^{2d \left(\frac{2^{\tilde{\sigma}}|m-n|^{\tilde{\sigma}}}{M_0^{\tilde{\sigma}}}+1\right)} \\ &\times e^{-\gamma_0(|m-n|-2M_0)^{\tilde{\sigma}}} \|G_{J_{s+1}\setminus J_s}\| \\ &\leq M_1^{9dM_1^{\tilde{\sigma}}M_0^{\sigma-\tilde{\sigma}}} e^{-\gamma_0(|m-n|-2M_0)^{\tilde{\sigma}}} \|G_{J_{s+1}\setminus J_s}\| \\ &\leq M_1^{9dM_1^{\tilde{\sigma}}M_0^{\sigma-\tilde{\sigma}}} e^{-\gamma_0(|m-n|-2M_0)^{\tilde{\sigma}}} 4 (2M_0+1)^d e^{M_0^{\sigma}} \\ &\leq M_1^{10dM_1^{\tilde{\sigma}}M_0^{\sigma-\tilde{\sigma}}} e^{-\gamma_0(|m-n|-2M_0)^{\tilde{\sigma}}}, \end{aligned}$$

where the first inequality holds by $|m-n| \leq 2M_1$ and the third inequality holds by (85).

(90)

Case 2_1 : $z \in J_s$. For this case, following the proof of Case 1_2 (see (81)), one has

(91)
$$|G_{J_{s+1}}(x,z)| \leq 2(2M_1+1)^{2d} e^{M_1^{\sigma}} T_s e^{-\gamma_0 (\max\{\Gamma_s(z)-4M_0,0\})^{\tilde{\sigma}}}$$

Case 2₂: $z \in J_{s+1} \setminus J_s$. Applying (58) ($\Lambda_1 = J_s$ and $\Lambda_2 = J_{s+1} \setminus J_s$), one has

$$\begin{aligned} |G_{J_{s+1}}(x,z)| &\leq \sum_{\substack{n_1 \in J_s \\ n_2 \in J_{s+1} \setminus J_s}} |G_{J_{s+1} \setminus J_s}(n_2,z)| e^{-c_1 |n_1 - n_2|^{\tilde{\sigma}}} |G_{J_{s+1}}(x,n_1)| \\ &\leq 2(2M_1 + 1)^{2d} e^{M_1^{\sigma}} T_s \\ &\times \sum_{\substack{n_1 \in J_s \\ n_2 \in J_{s+1} \setminus J_s}} |G_{J_{s+1} \setminus J_s}(n_2,z)| e^{-c_1 |n_1 - n_2|^{\tilde{\sigma}}} e^{-\gamma_0 (\max\{\Gamma_s(n_1) - 4M_0, 0\})^{\tilde{\sigma}}} \\ &\leq 2(2M_1 + 1)^{4d} e^{M_1^{\sigma}} M_1^{10dM_1^{\tilde{\sigma}} M_0^{\sigma - \tilde{\sigma}}} T_s \\ &\times \sup_{\substack{n_1 \in J_s \\ n_2 \in J_{s+1} \setminus J_s}} e^{-\gamma_0 (\max\{|n_2 - z| - 2M_0, 0\})^{\tilde{\sigma}}} e^{-\gamma_0 (\max\{\Gamma_s(n_1) - 4M_0, 0\})^{\tilde{\sigma}}} \\ &\leq M_1^{11dM_1^{\tilde{\sigma}} M_0^{\sigma - \tilde{\sigma}}} T_s \sup_{n_1 \in J_s} e^{-\gamma_0 (\max\{|n_1 - z| - 2M_0, 0\})^{\tilde{\sigma}}} e^{-\gamma_0 (\max\{\Gamma_s(n_1) - 4M_0, 0\})^{\tilde{\sigma}}} \\ \end{aligned}$$

where the second inequality holds by (91), the third inequality holds by (86), the fourth inequality holds by (64) and the fifth inequality holds by (64) and (74).

Putting all cases together and by (79), (81), (91) and (92), one has if (75) holds at sth step, then

(93)
$$\begin{aligned} |G_{J_{s+1}}(x,z)| &\leq M_1^{12dM_1^{\tilde{\sigma}}M_0^{\sigma-\tilde{\sigma}}}T_s e^{-\gamma_0(\max\{\Gamma_s(z)-10M_0,0\})^{\tilde{\sigma}}} \\ &\leq M_1^{12dM_1^{\tilde{\sigma}}M_0^{\sigma-\tilde{\sigma}}}T_s e^{-\gamma_0\Gamma_{s+1}^{\tilde{\sigma}}(z)}. \end{aligned}$$

By (77) and (93), we obtain that (75) is true for

$$(94) T_0 = e^{M_1^\sigma},$$

(95)
$$T_{s+1} = M_1^{12dM_1^{\tilde{\sigma}}M_0^{\sigma-\tilde{\sigma}}} T_s.$$

By (75), (94) and (95), one has

(96)
$$|G_{J_g}(x,z)| \le M_1^{13gdM_1^{\tilde{\sigma}}M_0^{\sigma-\sigma}} e^{-\gamma_0 \Gamma_g^{\tilde{\sigma}}(z)}.$$

By the assumption that $\tilde{\Lambda}_1$ is good, one has

$$(97) g \le 2B_1 = 2\kappa \frac{M_1}{M_0},$$

and hence (by (55))

(98)
$$k(z) \ge \frac{|x-z|}{4M_0} - 2B_1 - C(d) \ge \frac{|x-z|}{4M_0} - 2\kappa \frac{M_1}{M_0} - C(d).$$

By (98) and the definition of Γ_s , we have for $|x - z| \ge \frac{M_1}{10}$,

(99)

$$\Gamma_{g}^{\tilde{\sigma}}(z) \geq (|x-z| - 8\kappa M_{1} - 10(g+1)M_{0})^{\tilde{\sigma}} \\
\geq (|x-z| - 30\kappa M_{1})^{\tilde{\sigma}} \\
\geq |x-z|^{\tilde{\sigma}}(1 - 160\kappa)^{\tilde{\sigma}} \\
\geq |x-z|^{\tilde{\sigma}}(1 - 200\tilde{\sigma}\kappa),$$

where κ will be chosen to be sufficiently small.

By (96), (97) and (99), we have for $|x - z| \ge \frac{M_1}{10}$,

(100)

$$\begin{aligned} |G_{\tilde{\Lambda}_{1}}(x,z)| &= |G_{J_{g}}(x,z)| \\ &\leq M_{1}^{13gdM_{1}^{\tilde{\sigma}}M_{0}^{\sigma-\tilde{\sigma}}}e^{-\gamma_{0}\Gamma_{g}^{\tilde{\sigma}}(z)} \\ &\leq e^{-\gamma_{0}(1-200\kappa\tilde{\sigma}-300d\kappa\rho\gamma_{0}^{-1}\frac{\log M_{0}}{M_{0}^{1-\rho+\tilde{\sigma}-\sigma}})|x-z|^{\tilde{\sigma}}} \end{aligned}$$

Inductions: Define

(101)
$$\gamma_m = \prod_{i=0}^{m-1} \gamma_0 (1 - C(d)\kappa\tilde{\sigma} - C(d)\kappa\rho\gamma_i^{-1} \frac{\log M_i}{M_i^{1-\rho+\tilde{\sigma}-\sigma}}).$$

We remind that $1 - \rho + \tilde{\sigma} - \sigma > 0$. Fix an elementary region $\tilde{\tilde{\Lambda}}_1 \in \mathcal{E}_{M_m}$ and $\tilde{\tilde{\Lambda}}_1 \subset \tilde{\Lambda}_0$. For any $x \in \tilde{\tilde{\Lambda}}_1$, consider the exhaustion $\{S_j(x)\}_{j=0}^l$ of $\tilde{\tilde{\Lambda}}_1$ at x with width M_{m-1} . We say the annulus $A_j(x)$ is good, if for any $y \in A_j(x)$, there exists $W(y) \in \mathcal{E}_{M_{m-1}}$ such that

$$y \in W(y) \subset A_j(x), \text{ dist } (y, A_j(x) \setminus W(y)) \ge M_{m-1}/2,$$

and for $|n - n'| \ge \frac{M_{m-1}}{10}$,

$$|(R_{W(y)}AR_{W(y)})^{-1}(n,n')| \le e^{-\gamma_m|n-n'|^{\tilde{\sigma}}}$$

Otherwise, we call the annulus bad. An elementary region $\tilde{\Lambda}_1 \subset \tilde{\Lambda}_0$ is called bad provided for some $x \in \tilde{\Lambda}_1$ the number of bad annuli $\{A_j(x)\}$ exceeds

$$B_m := \kappa \frac{M_m}{M_{m-1}}.$$

Otherwise, we call $\tilde{\Lambda}_1$ good. Let \mathcal{F}_m be an arbitrary family of pairwise disjoint bad elementary regions in \mathcal{E}_{M_m} contained in $\tilde{\Lambda}_0$. By induction, it is easy to see that

$$\#\mathcal{F}_m \le \frac{1}{\kappa^m} \frac{N^\varsigma}{M_m}.$$

Replace M_0 , M_1 , γ_0 , B_1 with M_{m-1} , M_m , γ_{m-1} , B_m . By induction and following the proof of (100), we have for good elementary regions $\tilde{\tilde{\Lambda}}_1 \subset \tilde{\Lambda}_0$ and $\tilde{\tilde{\Lambda}}_1 \in \mathcal{E}_{M_m}$, we have for $|x - z| \geq \frac{M_m}{10}$,

(102)
$$|G_{\tilde{\Lambda}_1}(x,z)| \leq e^{-\gamma_m |x-z|^{\tilde{\sigma}}}.$$

In order to reach $\tilde{\Lambda}_0$ after k step, we need

$$M_k = M^{\rho^k} = N,$$

hence

(103)
$$\rho^k \approx \frac{1}{\xi}.$$

We may modify ρ a little bit at the last few steps to ensure that k is a positive integer. However, this issue is rather small. Choose $\kappa = N^{-\delta}$ and

$$\delta = -\frac{1}{2}(1-\varsigma)\frac{\log\rho}{\log\xi^{-1}}$$

Direct computations show that

(104)
$$\#\mathcal{F}_k \le \frac{1}{\kappa^k} \frac{N^{\varsigma}}{M_k} < 1.$$

(104) implies that $M_k = N$ is good. Therefore, (6) holds for

$$c_3 = \gamma_k,$$

where k solves (103). Computations show that

$$c_3 = c_2 - N^{-\vartheta},$$

where $\vartheta = \vartheta(\sigma, \tilde{\sigma}, \xi, \varsigma) > 0$.

5. Proof of Theorem 2.2

The proof of Theorem 2.2 is based on matrix-valued Cartan-type estimates [15, 20, 36, 51]. For our purpose, a new version of Cartan's estimate, which works for non-self-adjoint matrices, is necessary. For convenience, we include a proof in the Appendix.

Lemma 5.1. Let T(x) be a $N \times N$ matrix function of a parameter $x \in [-\delta, \delta]^J$ $(J \in \mathbb{N})$ satisfying the following conditions:

(i) T(x) is real analytic in $x \in [-\delta, \delta]^J$ and has a holomorphic extension to

$$\mathcal{D}_{\delta,\delta_1} = \left\{ x = (x_i)_{1 \le i \le J} \in \mathbb{C}^J : \sup_{1 \le i \le J} |\Re x_i| \le \delta, \sup_{1 \le i \le J} |\Im x_i| \le \delta_1 \right\}$$

satisfying

(105)
$$\sup_{x \in \mathcal{D}_{\delta, \delta_1}} \|T(x)\| \le B_1, B_1 \ge 1.$$

(ii) For all
$$x \in [-\delta, \delta]^J$$
, there is subset $V \subset [1, N]$ with
 $|V| \le M$,

and

(106)
$$||(R_{[1,N]\setminus V}T(x)R_{[1,N]\setminus V})^{-1}|| \le B_2, B_2 \ge 1.$$

(iii)

(107)
$$\max\{x \in [-\delta, \delta]^J : \|T^{-1}(x)\| \ge B_3\} \le 10^{-3J} J^{-J} \delta_1^J (1+B_1)^{-J} (1+B_2)^{-J} J^{-J} \delta_2^J J$$

(108)
$$0 < \epsilon \le (1 + B_1 + B_2)^{-10M}$$

Then

(109)
$$\max\left\{x \in [-\delta/2, \delta/2]^J: \|T^{-1}(x)\| \ge \epsilon^{-1}\right\} \le C\delta^J e^{-c\left(\frac{\log \epsilon^{-1}}{M\log(B_1+B_2+B_3)}\right)^{1/J}},$$

where $C = C(J), c = c(J) > 0.$

Proof of Theorem 2.2. Without loss of generality, we assume i = 1. Fix $x_1 \in \mathbb{T}^{b_1}$ and $x_1^{\neg} \in \mathbb{T}^{b-b_1}$. Recall that $x = (x_1, x_1^{\neg}) \in \mathbb{T}^b$. Let $\Lambda = \mathcal{R} \subset [-N_3, N_3]^d$. By making $\mathcal{B}_{\mathcal{R}}(x)$ slightly larger, we have there exists $\overline{\Lambda} \subset \Lambda$ such that for any $j \in \Lambda \setminus \overline{\Lambda}$, there exists $W(j) \in \mathcal{E}_{N_1}$ such that $W(j) \subset \Lambda \backslash \overline{\Lambda},$

$$\operatorname{dist}(j, \Lambda \setminus \overline{\Lambda} \setminus W(j)) \ge N_1/2$$

and

(110)
$$||G_{W(j)}|| \le e^{N_1^{\sigma}},$$

(111)
$$|G_{W(j)}(n,n')| \le e^{-c_2|n-n'|^{\tilde{\sigma}}} \text{ for } |n-n'| \ge \frac{N_1}{10}$$

and

(112)
$$|\bar{\Lambda}| \le C(d)L^{1-\delta}N_1^{2d}.$$

Indeed, $\overline{\Lambda}$ can be chosen so that

(113)
$$\bar{\Lambda} \subset \{n \in \mathbb{Z}^d : \text{ dist } (n, B_{\mathcal{R}}(x)) \leq C(d)N_1.$$

Let $\eta = \frac{c_1}{\gamma}$. Let \mathcal{D} be the $e^{-\eta N_1}$ neighborhood of x_1 in the complex plane, i.e.,

$$\mathcal{D} = \{ z \in \mathbb{C}^{b_1} : |\Im z| \le e^{-\eta N_1}, |\Re z - x_1| \le e^{-\eta N_1} \}.$$

By the assumption that $N_3 \leq e^{N_1^{2\bar{K}_1}}$, one has for any $y \in \mathcal{D}$,

$$||x - y|| \le e^{-e^{(\log(2N_3 + 2))^{K_1}}}$$

and hence (by (8))

(114)
$$\begin{aligned} |A(x;n,n') - A(y;n,n')| &\leq K ||x-y||^{\gamma} \\ &\leq K e^{-c_1 N_1}, \end{aligned}$$

for $n, n' \in [-N_3, N_3]^d$ and large N_1 . By (110), (111), (114), and standard perturbation arguments, we have for any $y \in \mathcal{D}$, and $j \in \Lambda \setminus \overline{\Lambda}$,

(115)
$$||G_{W(j)}(x_1+y,x_1)|| \le 2e^{N_1^{\sigma}}$$

(116)
$$|G_{W(j)}(x_1 + y, x_1; n, n')| \le 2e^{-c_2|n-n'|^{\tilde{\sigma}}} \text{ for } |n-n'| \ge \frac{N_1}{10}$$

Substituting Λ with $\Lambda \setminus \overline{\Lambda}$ in Lemma 4.2, one has for any $y \in \mathcal{D}$,

(117)
$$\|G_{\Lambda\setminus\bar{\Lambda}}(x_1+y,x_1)\| \le e^{2N_1\sigma}.$$

We want to use Lemma 5.1. For this purpose, let

$$T(y) = R_{\Lambda}AR_{\Lambda}, J = b_1, \delta = \delta_1 = e^{-\eta N_1}.$$

Now we are in the position to check the assumptions of Lemma 5.1. By (114) and (7), one has $B_1 = O(1)$.

Let $V = \overline{\Lambda}$. By (112) and (117), one has

(118)
$$M = |\bar{\Lambda}| \le C(d) L^{1-\delta} N_1^{2d}, B_2 = e^{2N_1^{\sigma}}.$$

Applying Lemma 4.2 with $M_0 = M_1 = N_2$ and (12), one has

$$||T^{-1}(y)|| \le 4(2N_2+1)^d e^{N_2^{\sigma}} \le e^{2N_2^{\sigma}} =: B_3$$

except on a set of $y \in \mathbb{T}^{b_1}$ with measure less than $e^{-N_2^{\zeta}}$.

Since $N_2 \ge N_1^{\frac{2}{\zeta}}$, direct computation shows that

$$10^{-3b_1}b_1^{-b_1}\delta^{b_1}(1+B_1)^{-b_1}(1+B_2)^{-b_1} \ge e^{-N_2^{\zeta}}.$$

This verifies (iii) in Lemma 5.1.

Let $\epsilon = e^{-L^{\mu}}$. By (118) and the assumption that $L \ge N_2^{\frac{2d+b+2}{\mu-1+\delta}}$, one has $\epsilon < (1+B_1+B_2)^{-10M}$.

Let

$$Y = \{ y \in \mathcal{D} : ||T^{-1}(y)|| \ge e^{L^{\mu}} \}.$$

By (109) of Lemma 5.1,

(119)
$$\operatorname{mes}(Y) \le C e^{-c \left(\frac{L^{\mu-1+\delta}}{N_2^{\sigma} N_1^{2d+\sigma}}\right)^{1/b_1}}.$$

By covering \mathbb{T}^{b_1} with balls with radius $e^{-\eta N_1}$, we have

$$\operatorname{Leb}(\tilde{X}_{\mathcal{R}}(x_{1}^{\neg})) \leq e^{CN_{1}} e^{-c\left(\frac{L^{\mu-1+\delta}}{N_{2}^{\sigma}N_{1}^{2d+\sigma}}\right)^{1/b_{1}}}$$
$$\leq e^{-\left(\frac{L^{\mu-1+\delta}}{N_{2}^{2d+b+2}}\right)^{1/b_{1}}},$$

where the second inequality holds by the assumption $L \ge N_2^{\frac{2d+b+2}{\mu-1+\delta}}$. It implies (13).

6. Proof of Theorems 2.3 and 2.4

Theorem 6.1. Let $\sigma, \tilde{\sigma}, \kappa, s \in (0, 1)$ and $\tilde{\sigma} > \kappa$. Assume diam $(\Lambda) \leq 2N + 1$. Let $M_0 = (\log N)^{1/s}$. Assume

(120)
$$c_2 \in (0, (1-5^{-\tilde{\sigma}})c_1].$$

Suppose that for any $n \in \Lambda$, there exists some $W = W(n) \in \mathcal{E}_M$ with $M_0 \leq M \leq N^{\kappa}$ such that $n \in W$, $\operatorname{dist}(n, \Lambda \setminus W) \geq \frac{M}{2}$, $W \subset \Lambda$ and

$$||G_W|| \le 2e^{M^{\sigma}},$$

 $|G_W(n,n')| \le 2e^{-c_2|n-n'|^{\tilde{\sigma}}} \text{ for } |n-n'| \ge \frac{M}{10}.$

Then

(121)
$$||G_{\Lambda}|| \le 4(1+2N^{\kappa})^d e^{N^{\kappa\sigma}},$$

and

$$|G_{\Lambda}(n,n')| \le e^{-\overline{c}|n-n'|^{\tilde{\sigma}}} \text{ for } |n-n'| \ge N/10,$$

where

(122)
$$\bar{c} = c_2 - \frac{O(1)}{M_0^{\tilde{\sigma}-s}} - \frac{O(1)}{M_0^{\tilde{\sigma}-\sigma}} - \frac{O(1)}{N^{\tilde{\sigma}-\kappa}}.$$

Proof. (121) follows from Lemma 4.2 immediately.

Assume $|n-n'| \ge \frac{N}{10}$. Applying (57) with $\Lambda_1 = W = W(n)$, one has $n' \notin W(n)$ and

$$|G_{\Lambda}(n,n')| \leq \sum_{\substack{n_1 \in W \\ n_2 \in \Lambda \setminus W}} e^{-c_1 |n_1 - n_2|^{\tilde{\sigma}}} |G_W(n,n_1)| |G_{\Lambda}(n_2,n')|.$$

It implies

$$\begin{aligned} |G_{\Lambda}(n,n')| &\leq \sum_{\substack{n_{1} \in W, |n_{1}-n| \leq \frac{M}{10} = 1\\ n_{2} \in \Lambda \setminus W}} e^{-c_{1}|n_{1}-n_{2}|^{\tilde{\sigma}}} |G_{W}(n,n_{1})| |G_{\Lambda}(n_{2},n')|} \\ &+ \sum_{\substack{n_{1} \in W, |n_{1}-n| \geq \frac{M}{10}\\ n_{2} \in \Lambda \setminus W}} e^{-c_{1}|n_{1}-n_{2}|^{\tilde{\sigma}}} |G_{\Lambda}(n_{2},n')| \\ &\leq \sum_{\substack{n_{1} \in W, |n_{1}-n| \leq \frac{M}{10} = 1\\ n_{2} \in \Lambda \setminus W}} e^{-c_{1}|n_{1}-n_{2}|^{\tilde{\sigma}}} |G_{\Lambda}(n_{2},n')| \\ &+ \sum_{\substack{n_{1} \in W, |n_{1}-n| \leq \frac{M}{10} = 1\\ n_{2} \in \Lambda \setminus W}} e^{-c_{1}|n_{1}-n_{2}|^{\tilde{\sigma}}} |G_{\Lambda}(n_{2},n')| \\ &\leq \sum_{\substack{n_{1} \in W, |n_{1}-n| \leq \frac{M}{10} = 1\\ n_{2} \in \Lambda \setminus W}} e^{-c_{2}|n-n_{2}|^{\tilde{\sigma}}} |G_{\Lambda}(n_{2},n')| \\ &+ \sum_{\substack{n_{1} \in W, |n_{1}-n| \leq \frac{M}{10} = 1\\ n_{2} \in \Lambda \setminus W}} e^{-c_{2}|n-n_{2}|^{\tilde{\sigma}}} |G_{\Lambda}(n_{2},n')| \\ &\leq e^{M^{\sigma}} \sum_{\substack{n_{1} \in W, |n_{1}-n| \leq \frac{M}{10} = 1\\ n_{2} \in \Lambda \setminus W}} e^{-c_{2}|n-n_{2}|^{\tilde{\sigma}}} |G_{\Lambda}(n_{2},n')| \\ &+ \sum_{\substack{n_{1} \in W, |n_{1}-n| \leq \frac{M}{10} = 1\\ n_{2} \in \Lambda \setminus W}} e^{-c_{2}|n-n_{2}|^{\tilde{\sigma}}} |G_{\Lambda}(n_{2},n')| \\ &\leq e^{M^{\sigma}} (2N+1)^{2d} \sup_{\substack{n_{2} \in \Lambda \setminus W}} e^{-c_{2}|n-n_{2}|^{\tilde{\sigma}}} |G_{\Lambda}(n_{2},n')| \\ &\leq (2N+1)^{2d} \sup_{\substack{n_{2} \in \Lambda \setminus W}} e^{-c_{2}|n-n_{2}|^{\tilde{\sigma}}} |G_{\Lambda}(n_{2},n')| \end{aligned}$$
(123)

where the third inequality holds by (64) and the fourth inequality holds by (72). Iterating (123) until $|n_2 - n'| \leq 4N^{\kappa}$ (but at most $\lfloor \frac{2\tilde{\sigma}|n-n'|\tilde{\sigma}}{M_0^{\tilde{\sigma}}} \rfloor$ times) and applying (121), we have for $|n - n'| \geq \frac{N}{10}$,

$$\begin{aligned} |G_{\Lambda}(n,n')| &\leq (2N+1)^{\frac{2^{\tilde{\sigma}}|n-n'|^{\tilde{\sigma}}}{M_{0}^{\tilde{\sigma}}}} e^{-(c_{2}-\frac{O(1)}{M_{0}^{\tilde{\sigma}-\sigma}})(|n-n'|-4N^{\kappa})^{\tilde{\sigma}}} 4(1+2N^{\kappa})^{d} e^{N^{\kappa\sigma}} \\ &\leq e^{\frac{4|n-n'|^{\tilde{\sigma}}}{M_{0}^{\tilde{\sigma}}}\log N} e^{-(c_{2}-\frac{O(1)}{M_{0}^{\tilde{\sigma}-\sigma}})(|n-n'|-4N^{\kappa})^{\tilde{\sigma}}} 4(1+2N^{\kappa})^{d} e^{N^{\kappa\sigma}} \\ &\leq e^{\frac{4|n-n'|^{\tilde{\sigma}}}{M_{0}^{\tilde{\sigma}}}M_{0}^{s}} e^{-(c_{2}-\frac{O(1)}{M_{0}^{\tilde{\sigma}-\sigma}})(|n-n'|-4N^{\kappa})^{\tilde{\sigma}}} 4(1+2N^{\kappa})^{d} e^{N^{\kappa\sigma}} \\ &\leq e^{-\bar{c}|n-n'|}. \end{aligned}$$

It is easy to see that the number of generalized elementary regions in $[-N, N]^d$ with width larger or equal to N^{ξ} is bounded by $N^{C(d)}$, more precisely for any $\xi > 0$,

(124)
$$\#\{\Lambda \subset [-N,N]^d : \Lambda \subset \mathcal{R}_L^{N^{\xi}}\} \le N^{C(d)}.$$

Proof of Theorem 2.3. Since the Green's function satisfies properties P with parameters (μ, ζ, c_2) at size N_2 , we have there exists $\tilde{X}_{N_2} \subset \mathbb{T}^b$ with

(125)
$$\sup_{1 \le i \le k, x_i^{\gamma} \in \mathbb{T}^{b-b_i}} \operatorname{Leb}(\tilde{X}_{N_2}(x_i^{\gamma})) \le N_3^{C(d)} e^{-N_2^{\zeta}},$$

such that

$$||G_{m+Q_{N_2}}(x)|| \le e^{N_2^{\mu}},$$

and for $|n - n'| \ge N_2/10$,

$$|G_{m+Q_{N_2}}(x;n,n')| \le e^{-c_2|n-n'|^{\tilde{\sigma}}},$$

for any $Q_{N_2} \in \mathcal{E}_{N_2}^0$ and $|m| \leq N_3$. Indeed, we only need to set

$$\tilde{X}_{N_2} = \bigcup_{|m| \le N_3} X_{N_2}(f^m(x)).$$

By the assumption $N_3 \ge N_2^C$ and $N_2 \ge N_1^C$ with large C depending on ε , one has

(126)
$$N_2 \le N_3^{\varepsilon}, N_1 \le N_2^{\varepsilon}$$

Let $\xi = \delta - 5\varepsilon$. Applying (15) to Theorem 2.2, and by (124) and (126), there exists $X_{N_3} \subset [0, 1)^b$ such that

(127)
$$\sup_{x_i^{\neg} \in \mathbb{T}^{b-b_i}} \operatorname{Leb}(X_{N_3}(x_i^{\neg})) \leq N_3^{C(d)} e^{-N_3^{\xi(\frac{\sigma-1}{b_i} + \frac{\delta}{b_i}) - \varepsilon}} \leq e^{-N_3^{\frac{\sigma-1}{b_i}\delta + \frac{\delta^2}{b_i} - \varepsilon}},$$

and for any $x \notin X_{N_3}$, $\mathcal{R} \subset \mathcal{R}_L^{N_3^{\xi}}$ with $N_3^{\xi} \leq L \leq N_3$, (128) $||G_{\mathcal{R}}(x)|| \leq e^{L^{\sigma}}$.

Let $\tilde{\mathcal{F}}$ be any pairwise disjoint elementary regions in $[-N_3, N_3]^d$ with size $\lfloor N_3^{\xi} \rfloor$. By (15), it is easy to see that there are at most $N_1^{C(d)}N_3^{1-\delta} = N_3^{1-\delta+\varepsilon}$ in $\tilde{\mathcal{F}}$ will intersect elementary regions not in SG_{N_1} . By Theorem 6.1, any elementary region in $[-N_3, N_3]^d$ with size $\lfloor N_3^{\xi} \rfloor$, without intersecting any non- SG_{N_1} elementary regions, will satisfy (3). It implies (5) is true for $\varsigma = 1 - \varepsilon$. Applying Theorem 2.1 and (128), we obtain Theorem 2.3. Let us explain where the bound $c_2 - N_1^{-\vartheta_1} - N_3^{-\vartheta_2}$ in (18) is from. Since $N_3^{\xi} \leq e^{\xi N_1^c}$, one has $s = \frac{11}{10}c$ in Theorem

6.1. Applying $M_0 = N_1$, $N = N_3^{\xi}$, $\sigma = \mu$ to Theorem 6.1, we obtain the bound $c_2 - O(1)N_1^{-(\tilde{\sigma} - \frac{11}{10}c)} - O(1)N_1^{-(\tilde{\sigma} - \mu)} - N_3^{-\vartheta_2}$. Theorem 2.1 will only contribute $N_3^{-\vartheta_2}$.

Proof of Theorem 2.4. Fix any $m \in \mathbb{Z}^d$. Applying Theorem 2.3 with \tilde{A}^m , one has there exists a subset $X_{N_3}^m \subset \mathbb{T}^b$ such that

$$\sup_{1 \le i \le k, x_i^{\neg} \in \mathbb{T}^{b-b_i}} \operatorname{Leb}(X_{N_3}^m(x_i^{\neg})) \le e^{-N_3 \frac{\sigma-1}{b_i}\delta + \frac{\delta^2}{b_i} - \varepsilon},$$

and for any $x \notin X_{N_3}^m$ and $Q_{N_3} \in \mathcal{E}_{N_3}^0$,

$$||(R_{Q_{N_3}}\tilde{A}^m(x)R_{Q_{N_3}})^{-1}|| \leq e^{N_3^{\sigma}},$$

and for $|n - n'| \ge \frac{N_3}{10}$,

$$|(R_{Q_{N_3}}\tilde{A}^m R_{Q_{N_3}})^{-1}(x;n,n')| \leq e^{-(c_2 - N_1^{-\vartheta_1} - N_3^{-\vartheta_2})|n-n'|^{\tilde{\sigma}}}$$

Let

$$X_{N_3} = \bigcup_{m \in \mathbb{Z}^d} X_{N_3}^m$$

By (19) and (21), we have

$$\sup_{1 \le i \le k, x_i^{\neg} \in \mathbb{T}^{b-b_i}} \operatorname{Leb}(X_{N_3}(x_i^{\neg})) \le e^{N_3^a} e^{-N_3^{\frac{\sigma-1}{b_i}\delta + \frac{\delta^2}{b_i} - \varepsilon}} \le e^{-N_3^{\frac{\sigma-1}{b_i}\delta + \frac{\delta^2}{b_i} - \varepsilon}}.$$

7. Proof of Theorem 2.5

Proof. Once we have the LDT at hand, the modulus of continuity of the IDS is standard. The proof here follows from the corresponding part in [13, 65]. Let $N = |\log |E_1 - E_2||^{\frac{1}{\sigma}-\varepsilon}$. Without loss of generality, assume $E_1 < E_2$ and let E be the center of $[E_1, E_2]$. Therefore,

$$(129) |E_1 - E_2| \le e^{-N^{\sigma + \varepsilon}}$$

By the assumption, there exists a set $X_N \subset \mathbb{T}^b$ such that

$$\operatorname{Leb}(X_N) \le e^{-N^{\zeta}},$$

and for any $x \notin X_N$ and any $Q_N \in \mathcal{E}_N^0$,

$$||G_{Q_N}(E,x)|| \leq e^{N^{\sigma}} |G_{Q_N}(E,x;n,n')| \leq e^{-c|n-n'|^{\tilde{\sigma}}} \text{ for } |n-n'| \geq \frac{N}{10},$$

where c > 0. We should mention that X_N depends on E. By the assumption (23), for large N_1 , one has

$$#\{n \in \mathbb{Z}^d : |n| \le N_1, f^n(x) \in X_N\} \le 2(2N_1 + 1)^d e^{-N^{\zeta}}$$

Let $\Lambda = [-N_1, N_1]^d$. By making $\#\{n \in \mathbb{Z}^d : |n| \leq N_1, f^n(x) \in X_N\}$ slightly larger, we have there exists $\overline{\Lambda} \subset \Lambda$ such that for all $j \in \Lambda \setminus \overline{\Lambda}$, there exists $W(j) \in \mathcal{E}_N$ such that $W(j) \subset \Lambda \setminus \overline{\Lambda}$, $\operatorname{dist}(j, \Lambda \setminus \overline{\Lambda} \setminus W(j)) \geq N/2$ and

(130)
$$||G_{W(j)}|| \le e^{N^{\sigma}}$$

(131)
$$|G_{W(j)}(n,n')| \le e^{-c|n-n'|^{\tilde{\sigma}}} \text{ for } |n-n'| \ge \frac{N}{10}.$$

and

$$|\bar{\Lambda}| \le C(d) N^{2d} (2N_1 + 1)^d e^{-N^{\zeta}}.$$

Here, $\overline{\Lambda}$ is obtained in a similar way as (113).

Substituting Λ with $\Lambda \setminus \Lambda$ in Lemma 4.2, we have

$$\|G_{\Lambda\setminus\bar{\Lambda}}(E,x)\| \le 4(2N+1)^d e^{N^{\sigma}}.$$

By standard perturbation arguments, we have for any $\tilde{E} \in [E_1, E_2]$,

(132) $\|G_{\Lambda\setminus\bar{\Lambda}}(\tilde{E},x)\| \le 8(2N+1)^d e^{N^{\sigma}}.$

Denote by ξ_j , $j = 1, 2, \dots, M$, the normalized eigenfunctions of H_{Λ} with eigenvalues falling into the interval $[E_1, E_2]$. Let ξ be one of them with eigenvalue \tilde{E} . By definition,

(133)
$$R_{\Lambda\setminus\bar{\Lambda}}(H_{\Lambda}-E)R_{\Lambda\setminus\bar{\Lambda}}\xi + R_{\Lambda\setminus\bar{\Lambda}}(H_{\Lambda}-E)R_{\bar{\Lambda}}\xi = (\tilde{E}-E)R_{\Lambda\setminus\bar{\Lambda}}\xi.$$

Applying $G_{\Lambda\setminus\bar{\Lambda}}(E,x)$ to (133), one has

(134)
$$R_{\Lambda\setminus\bar{\Lambda}}\xi + G_{\Lambda\setminus\bar{\Lambda}}(E,x)R_{\Lambda\setminus\bar{\Lambda}}(H_{\Lambda}-E)R_{\bar{\Lambda}}\xi = (\tilde{E}-E)G_{\Lambda\setminus\bar{\Lambda}}(E,x)R_{\Lambda\setminus\bar{\Lambda}}\xi.$$

Denote by P the projection onto the range of $G_{\Lambda\setminus\bar{\Lambda}}(E, x)R_{\Lambda\setminus\bar{\Lambda}}(H_{\Lambda} - E)R_{\bar{\Lambda}}$. Clearly, the dimension of this range does not exceed $\bar{\Lambda}$. Thus rank $(P) \leq \bar{\Lambda}$. By (129) and (132), one has

(135)
$$||(\tilde{E} - E)G_{\Lambda \setminus \bar{\Lambda}}(E, x)R_{\Lambda \setminus \bar{\Lambda}}\xi|| \le \frac{1}{100}||\xi||.$$

Applying I - P to (134) and by (135), we have

(136)
$$||R_{\Lambda\setminus\bar{\Lambda}}\xi - PR_{\Lambda\setminus\bar{\Lambda}}\xi|| \le \frac{1}{100}||\xi||.$$

Applying (136) to each ξ_j , we have

$$M = \sum_{j=1}^{N} ||\xi_j||^2$$

$$\leq \frac{M}{2} + 4 \sum_{j=1}^{M} ||PR_{\Lambda \setminus \bar{\Lambda}}\xi_j||^2 + 2 \sum_{j=1}^{M} ||R_{\bar{\Lambda}}\xi_j||^2$$

$$\leq \frac{M}{2} + 4 \operatorname{Trace}(PR_{\Lambda \setminus \bar{\Lambda}}) + 2 \operatorname{Trace}(R_{\bar{\Lambda}})$$

$$\leq \frac{M}{2} + 6|\bar{\Lambda}|$$

$$\leq \frac{M}{2} + C(d)N^{2d}(2N_1 + 1)^d e^{-N^{\zeta}}.$$

Therefore,

$$M \le C(d) N^{2d} (2N_1 + 1)^d e^{-N^{\zeta}}.$$

It implies

$$k(x, E_1, E_2) \le C(d) N^{2d} e^{-N^{\zeta}} \le e^{-(\log \frac{1}{E_2 - E_1})^{\frac{\zeta}{\sigma} - \varepsilon}}.$$

8. The discrepancy and semi-algebraic sets

8.1. **Discrepancy.** Let $\vec{x}_1, ..., \vec{x}_N \in [0, 1)^b$ and $\mathcal{S} \subset [0, 1)^b$. Let $A(\mathcal{S}; \{\vec{x}_n\}_{n=1}^N)$ be the number of \vec{x}_n $(1 \leq n \leq N)$ such that $\vec{x}_n \in \mathcal{S}$. We define the discrepancy of the sequence $\{\vec{x}_n\}_{n=1}^N$ by

(137)
$$D_N(\{\vec{x}_n\}_{n=1}^N) = \sup_{\mathcal{S}\in\mathcal{C}} \left| \frac{A(\mathcal{S};\{\vec{x}_n\}_{n=1}^N)}{N} - \operatorname{Leb}(S) \right|$$

where \mathcal{C} is the family of all intervals in $[0,1)^b$, namely \mathcal{S} has the form of

$$\mathcal{S} = [\varrho_1, \beta_1] \times [\varrho_2, \beta_2] \times \cdots \times [\varrho_b, \beta_b]$$

with $0 \leq \varrho_n < \beta_n < 1$, $n = 1, 2, \dots, b$. Let $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_b) \in [0, 1)^b$. The *b*-dimensional sequence $\vec{x}_n = (n\alpha_1, n\alpha_2, \dots, n\alpha_b) \mod \mathbb{Z}^b$ ($n\alpha$ for short), $n = 1, 2, \dots$, is called the Kronecker sequence. We denote by the discrepancy of $\{n\alpha\}_{n=1}^N$, $D_N(\alpha)$. The following Lemmas are well known.

Lemma 8.1. [32] Assume $\alpha \in DC(\kappa, \tau)$. Then

$$D_N(\alpha) \le C(b,\kappa,\tau) N^{-\frac{1}{\kappa}} (\log N)^2$$

Lemma 8.2. [66] For almost every α , we have

$$D_N(\alpha) \le C(\alpha) N^{-1} (\log N)^{b+2}.$$

Let $f: \mathbb{T}^b \to \mathbb{T}^b$ be defined as follows

 $T(y_1, y_2, ..., y_b) = (y_1 + \alpha, y_2 + y_1, ..., y_b + y_{b-1}).$

Let T^n be the *n*th iteration of T and $\vec{Y}_n = T^n(y_1, ..., y_b)$.

Lemma 8.3. Assume $\alpha \in DC(\kappa, \tau)$. Then for any $\varepsilon > 0$,

$$D_N(\{\vec{Y}_n\}_{n=1}^N) \le C(b,\kappa,\tau,\varepsilon) N^{-\frac{1}{2^{b-1}\kappa}+\varepsilon}.$$

Remark 9. Lemma 8.3 follows from the Erdős-Turán inequality (see Corollary 1.1 in p.8 of [63]) and the Weyl's method (Theorem 2 in p.41 of [63]).

The Erdős-Turán inequality and Weyl's method also imply

Lemma 8.4. Assume $\alpha \in DC(\kappa, \tau)$. Let $Y_n = P_b(T^n(y_1, ..., y_b))$, where P_b is the bth coordinate projection. Then for any $\varepsilon > 0$,

$$D_N(\{Y_n\}_{n=1}^N) \le C(b,\kappa,\tau,\varepsilon) N^{-\frac{1}{2^{b-1_\kappa}}+\varepsilon}.$$

8.2. Semi-algebraic sets. A set $S \subset \mathbb{R}^n$ is called a semi-algebraic set if it is a finite union of sets defined by a finite number of polynomial equalities and inequalities. More precisely, let $\{P_1, \dots, P_s\} \subset \mathbb{R}[x_1, \dots, x_n]$ be a family of real polynomials whose degrees are bounded by d. A (closed) semi-algebraic set S is given by an expression

(138)
$$\mathcal{S} = \bigcup_{j} \bigcap_{\ell \in \mathcal{L}_{j}} \left\{ x \in \mathbb{R}^{n} : P_{\ell}(x)\varsigma_{j\ell} 0 \right\},$$

where $\mathcal{L}_j \subset \{1, \dots, s\}$ and $\varsigma_{j\ell} \in \{\geq, \leq, =\}$. Then we say that \mathcal{S} has degree at most sd. In fact, the degree of \mathcal{S} which is denoted by deg(\mathcal{S}), means the smallest sd over all representations as in (138).

The following lemma is a special case appearing [10]. It is restated in [15].

Lemma 8.5. [15, Theorem 9.3] [10, Theorem 1] Let $S \subset [0,1]^n$ be a semialgebraic set of degree B. Then the number of connected components of S does not exceed $(1+B)^{C(n)}$.

The following lemma follows from the Yomdin-Gromov triangulation theorem [39, 73], which has been stated in [15]. We refer readers to [11] and references therein for the complete proof of the Yomdin-Gromov triangulation theorem.

Lemma 8.6. [15, Corollary 9.6] Let $S \subset [0,1]^n$ be a semi-algebraic set of degree B. Let $\epsilon > 0$ be a small number and $\text{Leb}(S) \leq \epsilon^n$. Then S can be covered by a family of ϵ -balls with total number less than $\frac{(1+B)^{C(n)}}{\epsilon^{n-1}}$.

Theorem 8.7. Assume that the discrepancy of the sequence $\{\vec{x}_j\}_{j=1}^N$ satisfies

$$D_N(\{\vec{x}_j\}_{j=1}^N) \le N^{-\varsigma},$$

for some $\varsigma > 0$. Let $S \subset [0,1]^n$ be a semi-algebraic set with degree less than B. Suppose

$$\operatorname{Leb}(\mathcal{S}) \leq N^{-\varsigma}$$

Then

$$A(\mathcal{S}; \{\vec{x}_j\}_{j=1}^N) \le (1+B)^{C(n)} N^{1-\frac{\varsigma}{n}}$$

Proof. Let $\epsilon = N^{-\frac{\varsigma}{n}}$. By Lemma 8.6, S can be covered, at most $\frac{(1+B)^C}{\epsilon^{n-1}}$, ϵ -balls. Pick one ϵ -ball, say J. By the fact $D_N(\{\vec{x}_j\}_{j=1}^N) \leq N^{-\varsigma}$, one has

$$A(J; \{\vec{x}_j\}_{j=1}^N) \le CN\epsilon^n + N^{1-\varsigma} \le CN^{1-\varsigma},$$

where C depends on the dimension n. Since there are at most $\frac{(1+B)^C}{\epsilon^{n-1}}$ balls, we have

$$A(S; \{\vec{x}_j\}_{j=1}^N) \leq (1+B)^C \frac{1}{\epsilon^{n-1}} N^{1-\varsigma} = (1+B)^C N^{\frac{n-1}{n}\varsigma} N^{1-\varsigma} = (1+B)^C N^{1-\frac{\varsigma}{n}}.$$

- Remark 10. Theorem 8.7 says that there is a factor b loss (referred to as dimension loss) when passing discrepancy from intervals to semi-algebraic sets. The dimension loss is not surprising. For example, there is also a dimension loss passing the discrepancy to the isotropic discrepancy [56, Theorem 1.6].
 - The proof of Theorem 8.7 is taken from Bourgain [15], where no explicit bounds are given.

For a set $S \subset [0,1)^2$, denote by l(S) the length of the longest line segment contained in S.

Lemma 8.8. [22, Theorem 5.1] Assume $\alpha_1 \in DC(\kappa, \tau)$ and $\alpha_2 \in DC(\kappa, \tau)$. Let $S \subset [0,1)^2$ be a semi-algebraic set with degree less than B and

$$l(S) \le \frac{1}{2} \min_{1 \le |k| \le 2N} ||k\alpha||.$$

Then

$$#\{k = (k_1, k_2) \in \mathbb{Z}^2 : |k| \le N, (k_1\alpha_1, k_2\alpha_2) \in S \mod \mathbb{Z}^2\}$$

(139)
$$\leq (1+B)^{C(d)}C(\kappa,\tau)N^{3\kappa-\frac{9}{4}}.$$

9. Proof of all the results in Section 3

Applying Theorem 2.5 with $\sigma = 1 - \varepsilon$, Theorem 3.2 follows from Theorem 3.1, Theorem 3.8 follows from Theorem 3.7, Theorem 3.12 follows from Theorem 3.11, Theorem 3.15 follows from Theorem 3.14, Theorem 3.19 follows from Theorem 3.18 and Theorem 3.21 follows from Theorem 3.20.

Applying strong Diophantine frequencies to Theorems 3.15 and 3.21, we obtain Corollaries 3.16 and 3.22.

With large deviation theorems 3.7 and 3.11 at hand, the proof of Theorems 3.10 and 3.13 is rather standard. We refer the readers to [17, Section 3], [20, Section 6] and [15, Chapter XV] for details. We note that the only difference is that the degree of semi-algebraic sets is at most $e^{(\log N)^C}$ in our cases, not N^C .

By the discussion above, in order to prove all the results in Section 3, it suffices to prove Theorems 3.1, 3.3, 3.7, 3.9, 3.11, 3.14, 3.18, 3.20 and Corollary 3.17.

In this section, C(c) is always a large (small) constant. It may change even in the same formula.

Lemma 9.1. [15, Prop. 7.19] Let H(x) be given by (32) and the Lyapunov exponent is given by (35). Suppose L(E) > 0. Then there exist $0 < \sigma < 1$ and $\zeta > 0$ such that for large N, there exists $X_N \subset \mathbb{T}^b$ such that $\text{Leb}(X_N) \leq e^{-N^{\zeta}}$ and for $x \notin X_N$, one of the intervals

$$\Lambda = [1, N]; [1, N-1]; [2, N]; [2, N-1]$$

will satisfy

$$|G_{\Lambda}(n_1, n_2)| \le e^{-L(E)|n_1 - n_2| + N^{\sigma}}$$

Proof of Theorem 3.1. By Lemma 9.1, there exist $0 < \sigma_1 < 1$ and $\zeta_1 > 0$ such that for any large N_1 , there exists $X_{N_1} \subset \mathbb{T}^b$ such that $\text{Leb}(X_{N_1}) \leq e^{-N_1^{\zeta_1}}$ and for $x \notin X_{N_1}$, one of the intervals

(140)
$$\Lambda(N_1) = [1, N_1]; [1, N_1 - 1]; [2, N_1]; [2, N_1 - 1]$$

will satisfy

(141)
$$|G_{\Lambda(N_1)}(n_1, n_2)| \le e^{-L(E)|n_1 - n_2| + N_1^{\sigma_1}}.$$

By approximating the analytic function with trigonometric polynomials given by (28) and using Taylor expansions, we can further assume that X_{N_1} is a semialgebraic set with degree less than $e^{(\log N_1)^C}$. This argument is quite standard. We refer to [15] for details. By Lemma 8.1 and Theorem 8.7, for any $e^{(\log N_1)^C} \leq N_3 \leq e^{N_1^c}$,

$$A(X_{N_1}; \{n\omega\}_{n=1}^{N_3}) \le N_3^{1-\frac{1}{b\kappa}+\varepsilon}.$$

Let
$$N_2 = N_3^{\frac{1}{C}}$$
. Applying (141) to N_2 , one has
(142) $|G_{\Lambda(N_2)}(n_1, n_2)| \le e^{-L(E)|n_1 - n_2| + N_2^{\sigma_1}},$

except for a set of x with measure less than $e^{-N_2^{\sigma_1}}$. Now Theorem 3.1 follows from Theorem 2.3. We should mention that the elementary region is $[-N_1, N_1]$ in Theorem 2.3 which is slightly different from (140). However, the same statement is true.

Proof of Theorem 3.18. The proof of Theorem 3.18 is similar to that of Theorem 3.1. The difference is that instead of Lemma 9.1, we need to use the corresponding statements in p.3575 [70] for initial scales. We also need to use Lemma 8.4 instead of Lemma 8.3.

Proof of Theorem 3.11. Let $N_2 = e^{N_1^c}$. Assume the Green's function in Theorem 3.11 satisfies properties P with parameters (μ, ζ, c_2) at sizes N_1 and N_2 . Let $N_3 = N_2^C$. We can assume that X_{N_1} is a semi-algebraic set with degree less than $e^{(\log N_1)^C}$. By Lemma 8.5, X_{N_1} is consisted of at most $e^{(\log N_1)^C}$ intervals with measure less than $e^{-N_1^{\zeta}}$. Let I be one of the intervals. Since ω satisfies Diophantine condition, for any $x \in \mathbb{T}$, there is at most one $n \in \mathbb{Z}^d$ with $|n| \leq N_3$ such that $x + n\omega \mod \mathbb{Z} \in I$. Therefore,

(143)
$$A(X_{N_1}; \{n\omega\}_{n=1}^{N_3}) \le e^{(\log N_1)^C} \le N_3^{\varepsilon}.$$

By Theorem 2.3, we have the Green's function satisfies properties P with parameters $(\sigma, \sigma - \varepsilon, c_2 - N_3^{-\vartheta})$ at size N_3 . Standard Neumann series expansion ensures that for any large N_0 , there exists λ_0 such that for any $\lambda > \lambda_0$, the Green's functions have properties P with parameters $(\sigma, \sigma - \varepsilon, \frac{4}{5}c_1)$ at all sizes smaller than N_0 [51, Theorem 4.3]. Now Theorem 3.11 follows by standard induction. See pages 15 and 16 in [51] for details.

Proof of Theorem 3.7. Fix N_1 . Let $N_2 = e^{N_1^c}$ and $N_3 = N_2^C$. Assume the Green's function in Theorem 3.7 satisfies properties P with parameters (μ, ζ, c_2) at sizes N_1 and N_2 . We can again assume that X_{N_1} is a semi-algebraic set with degree less than $e^{(\log N_1)^C}$. By Lemma 8.1 and Theorem 8.7,

(144)
$$A(X_{N_1}; \{n\omega\}_{n=1}^{N_3}) \le N_3^{1-\frac{1}{b\kappa}+\varepsilon}$$

By Theorem 2.3, we have the Green's function satisfies properties P with parameters

$$\left(\sigma, \frac{\sigma-1}{b^2\kappa} + \frac{1}{b^3\kappa^2} - \varepsilon, c_2 - N_3^{-\vartheta}\right)$$

at size N_3 . As the arguments at the end of proof of Theorem 3.11, large λ will ensure the initial scales and hence Theorem 3.7 follows by induction.

Proof of Theorems 3.3 and 3.9. The proof of Theorems 3.3 and 3.9 closely follow that of Theorems 3.2 and 3.8. The difference is that we need to use Lemma 8.2 instead of Lemma 8.1. \Box

Proof of Theorem 3.14. Replacing Lemma 8.1 with Lemma 8.3, Theorem 3.14 follows Theorem 3.7. \Box

Proof of Corollary 3.17. By formula (3.53) in [19], one has for almost every α ,

(145)
$$A(X_{N_1}; \{n\omega\}_{n=1}^{N_3}) \le N_3^{1-\frac{1}{3}+\varepsilon}$$

Let $\delta = 1/3 - \varepsilon$. Applying $\tilde{\sigma} = 1$, $\sigma = 1 - \varepsilon$ and $b_i = 2$ in Theorem 2.3 and then Theorem 2.5, we obtain Corollary 3.17. Indeed, 1/18 comes from $(1/3)^2/b$.

Proof of Theorem 3.20. The proof of Theorem 3.20 is similar to that of Theorems 3.11 and 3.7. We only point out the modifications.

- The induction goes in the following way. The semi-algebraic set X_N intersecting with any line segments contained in $[0,1)^2$ has Lebesgue measure at most $e^{-N^{\zeta}}$. The assumption that v is not constant on any line segments ensure the initial scales.
- Replace (143) or (144) with (139).
- Since the induction is based on semi-algebraic sets only on line segments, the Cartan's estimate will not lead to dimension loss. In other words, when (16) is used to do the induction, $b_i = 1$.

Remark 11. (1) The calculation of the bound in Theorem 3.20 goes in the following way. By (139), the sublinear bound is

$$3\kappa - \frac{9}{4} = 1 - \delta$$
, where $\delta = \frac{13}{4} - 3\kappa$.

Therefore, the bound in (16) becomes $(b_i = 1)$

$$\frac{\sigma - 1}{b_i}\delta + \frac{\delta^2}{b_i} = (\sigma - 1)\delta + \delta^2$$
$$= (\sigma - 1)\left(\frac{13}{4} - 3\kappa\right) + \left(\frac{13}{4} - 3\kappa\right)^2.$$

(2) The induction of Theorem 3.20 follows the corresponding parts in [22]. Our quantitative approaches developed in the paper allow us to obtain the explicit bound.

APPENDIX A. CARTAN'S ESTIMATES FOR NON-SELF-ADJOINT MATRICES

In the following, we will prove the several variables matrix-valued Cartan estimate (Lemma 5.1). The proof is similar to that in [14, 15, 17, 20, 51]. The improvement is that we do not assume the matrix is self-adjoint.

Lemma A.1. Let T be the matrix

$$T = \left(\begin{array}{cc} T_1 & T_2 \\ T_3 & T_4 \end{array}\right),$$

where T_1 is an invertible $n \times n$ matrix, T_2 is an $n \times k$ matrix, T_3 is a $k \times n$ matrix, and T_4 is a $k \times k$ matrix. Let

$$S = T_4 - T_3 T_1^{-1} T_2.$$

Then T is invertible if and only if S is invertible, and (146) $||S^{-1}|| \leq ||T^{-1}|| \leq C(1+||T_2||)(1+||T_3||)(1+||T_1^{-1}||)^2(1+||S^{-1}||),$ where C is an absolute constant.

Proof. It is easy to check that

(147)
$$T = \begin{pmatrix} T_1 & T_2 \\ T_3 & T_4 \end{pmatrix} = \begin{pmatrix} I & 0 \\ T_3 T_1^{-1} & I \end{pmatrix} \begin{pmatrix} I & T_2 \\ 0 & S \end{pmatrix} \begin{pmatrix} T_1 & 0 \\ 0 & I \end{pmatrix}.$$

It implies T is invertible if and only if S is invertible. By (147), one has

(148)
$$T^{-1} = \begin{pmatrix} T_1 & 0 \\ 0 & I \end{pmatrix}^{-1} \begin{pmatrix} I & T_2 \\ 0 & S \end{pmatrix}^{-1} \begin{pmatrix} I & 0 \\ T_3 T_1^{-1} & I \end{pmatrix}^{-1} = \begin{pmatrix} T_1^{-1} & 0 \\ 0 & I \end{pmatrix} \begin{pmatrix} I & -T_2 S^{-1} \\ 0 & S^{-1} \end{pmatrix} \begin{pmatrix} I & 0 \\ -T_3 T_1^{-1} & I \end{pmatrix}$$

(149) $= \begin{pmatrix} \star & \star \\ \star & S^{-1} \end{pmatrix}.$

Now the second inequality of (146) follows from (148) and the first one follows from (149). $\hfill \Box$

Denote by $\mathcal{D}(z,r)$ the standard disk on \mathbb{C} of center z and radius r > 0.

Lemma A.2. [37, Lemma 2.15] Let $f(z_1, \dots, z_J)$ be an analytic function defined in a ploydisk $\mathcal{P} = \prod_{1 \leq i \leq J} \mathcal{D}(z_{i,0}, 1/2)$ and $\phi = \log |f|$. Let $\sup_{\underline{z} \in \mathcal{P}} \phi(\underline{z}) \leq M, m \leq \phi(\underline{z}_0), \underline{z}_0 = (z_{1,0}, \dots, z_{J,0})$. Given sufficiently large F, there exists a set $\mathcal{B} \subset \mathcal{P}$ such that

(150)
$$\phi(\underline{z}) > M - C(J)F(M - m), \text{ for any } \underline{z} \in \prod_{1 \le i \le J} \mathcal{D}(z_{i,0}, 1/4) \setminus \mathcal{B}_{\underline{z}}$$

and

(151)
$$\operatorname{mes}(\mathcal{B} \cap \mathbb{R}^J) \le C(J)e^{-F^{1/J}}.$$

Proof of Lemma 5.1. The proof is similar to that of Lemma 3.4 in [15]. In the following proof, C = C(J) and c = c(J).

Let

$$\mu = 10^{-2} J^{-1} \delta_1 (1 + B_1)^{-1} (1 + B_2)^{-1}.$$

Fix

$$x_0 \in [-\delta/2, \delta/2]^J$$

and consider T(z) with $|z - x_0| = \sup_{1 \le i \le J} |z_i - x_{0,i}| < \mu$. Thanks to Cauchy's estimate and (105), one obtains for $|z - x_0| < \mu$,

$$\|\partial_{z_i}T(z)\| \le \frac{4B_1}{\delta_1}, i = 1, 2, \cdots, J,$$

which implies

$$||T(z) - T(x_0)|| \le \frac{4JB_1\mu}{\delta_1} \le 25^{-1}(1+B_2)^{-1}$$

From the assumption (ii) of Lemma 5.1, we can find $V = V(x_0)$ so that $|V| = \tilde{M} \leq M$ and (106) is satisfied. Denote by $V^c = [1, N] \setminus V$. Thus using the standard Neumann series argument and (106), one has

(152) $||(R_{V^c}T(z)R_{V^c})^{-1}|| \le 2B_2 \text{ for } |z-x_0| < \mu.$

We define for $|z - x_0| < \mu$ the analytic function

(153)
$$S(z) = R_V T(z) R_V - R_V T(z) R_{V^c} (R_{V^c} T(z) R_{V^c})^{-1} R_{V^c} T(z) R_V.$$

Then by (152) and (153), we have

(154)
$$||S(z)|| \le 3B_1^2 B_2$$

Recalling Lemma A.1, if S(z) is invertible, so is T(z) and by (146),

(155)
$$||S^{-1}(z)|| \le C||T^{-1}(z)|| \le CB_1^2 B_2^2 (1 + ||S^{-1}(z)||)$$

For $x \in \mathbb{R}^J$, one has

(156)
$$||S(x)||^{\tilde{M}} \ge |\det S(x)|.$$

Let $\lambda = \min\{|\tilde{\lambda}| : \tilde{\lambda} \in \sigma(S(x))\}$. We have

(157)
$$|\det S(x)| \ge \lambda^M$$
$$\ge \|S^{-1}(x)\|^{-\tilde{M}}.$$

By Cramer's rule, one has every entry of $S^{-1}(x)$ is bounded by

$$\frac{||S(x)||^{\tilde{M}-1}}{|\det S(x)|}$$

and hence (by (154))

(158)
$$||S^{-1}(x)|| \le \frac{\tilde{M}(3B_1^2B_2)^{\tilde{M}}}{|\det S(x)|}.$$

Let

$$\phi(z) = \log |\det S(x_0 + \mu z)|, \ |z| < 1.$$

Then by (156) and (154),

(159)
$$\sup_{|z|<1} \phi(z) \le C\tilde{M} \log(B_1 + B_2).$$

By (107) and the definition of μ , there is some x_1 with $|x_0 - x_1| < \mu/10$ such that

(160)
$$||T^{-1}(x_1)|| \le B_3.$$

Hence by (155), $||S^{-1}(x_1)|| \le CB_3$, and from (157),

(161)
$$\phi(a) \ge -C\tilde{M}\log B_3,$$

where $a = \frac{x_1 - x_0}{\mu}$, so |a| < 1/10. Let

$$\mathcal{P} = \prod_{1 \le i \le J} \mathcal{D}(a_i, 1/2)$$

Therefore, one has

$$\sup_{z \in \mathcal{P}} \phi(z) \le C\tilde{M} \log(B_1 + B_2), \phi(a) \ge -C\tilde{M} \log B_3$$

Applying Lemma A.2 and recalling (150), (151), for any $F \gg 1$, there is some set $\mathcal{B} \subset \prod_{1 \leq i \leq J} \mathcal{D}(a_i, 1/4)$ with

(162)
$$\phi(z) \ge -CF\tilde{M}\log(B_1 + B_2 + B_3) \text{ for } z \in \prod_{1 \le i \le J} \mathcal{D}(a_i, 1/4) \setminus \mathcal{B},$$

and

(163)
$$\operatorname{mes}(\mathcal{B} \cap \mathbb{R}^J) \le C e^{-F^{1/J}}.$$

For $0 < \epsilon < 1$, let

$$F = \frac{-c\log\epsilon}{\tilde{M}\log(B_1 + B_2 + B_3)}$$

Then by (162) and (163),

$$\max \left\{ x \in \mathbb{R}^J : |x - x_1| < \mu/4 \text{ and } |\det(S(x))| \le \epsilon \right\}$$
$$= \mu^J \max \left\{ x \in \mathbb{R}^J : |x - a| < 1/4 \text{ and } \phi(x) \le \log \epsilon \right\}$$
$$\le C \mu^J e^{-F^{1/J}}.$$

Since $|x_0 - x_1| < \mu/10$, we have (164)

 $\max\left\{x \in \mathbb{R}^{J}: |x - x_{0}| < \mu/8 \text{ and } |\det(S(x))| \le \epsilon\right\} \le C\mu^{J} e^{-c\left(\frac{\log \epsilon^{-1}}{\tilde{M}\log(B_{1} + B_{2} + B_{3})}\right)^{1/J}}.$ $\text{Recalling (155), (158) \text{ and } (108), \text{ one has for } |x - x_{0}| < \mu/8 \text{ and } |\det S(x)| \ge \epsilon,$ $(165) \qquad ||T^{-1}(x)|| \le CB_{1}^{2}B_{2}^{2}\epsilon^{-1}\tilde{M}(3B_{1}^{2}B_{2})^{\tilde{M}} \le \epsilon^{-2}.$

Covering $\left[-\frac{\delta}{2}, \frac{\delta}{2}\right]^J$ by cubes of side $\mu/4$, and combining (164) and (165), one has

$$\max\left\{x \in \left[-\delta/2, \delta/2\right]^{J} : \|T^{-1}(x)\| \ge \epsilon^{-2}\right\} \le C\delta^{J} e^{-c\left(\frac{\log \epsilon^{-1}}{M\log(B_{1}+B_{2}+B_{3})}\right)^{1/J}} \\ \le C\delta^{J} e^{-c\left(\frac{\log \epsilon^{-1}}{M\log(B_{1}+B_{2}+B_{3})}\right)^{1/J}}.$$

Acknowledgments

I would like to thank Alan Haynes for telling me [63], which leads to Lemmas 8.3 and 8.4. I also wish to thank Svetlana Jitomirskaya for comments on earlier versions of the manuscript. This research was supported by NSF DMS-1700314/2015683, DMS-2000345 and the Southeastern Conference (SEC) Faculty Travel Grant 2020-2021.

References

- [1] A. Avila. The absolutely continuous spectrum of the almost mathieu operator. arXiv preprint arXiv:0810.2965, 2008.
- [2] A. Avila. Global theory of one-frequency Schrödinger operators. Acta Math., 215(1):1–54, 2015.
- [3] A. Avila and D. Damanik. Absolute continuity of the integrated density of states for the almost Mathieu operator with non-critical coupling. *Invent. Math.*, 172(2):439–453, 2008.
- [4] A. Avila, B. Fayad, and R. Krikorian. A KAM scheme for SL(2, ℝ) cocycles with Liouvillean frequencies. *Geom. Funct. Anal.*, 21(5):1001–1019, 2011.
- [5] A. Avila and S. Jitomirskaya. The Ten Martini Problem. Ann. of Math. (2), 170(1):303–342, 2009.
- [6] A. Avila and S. Jitomirskaya. Almost localization and almost reducibility. J. Eur. Math. Soc. (JEMS), 12(1):93–131, 2010.
- [7] A. Avila and S. Jitomirskaya. Hölder continuity of absolutely continuous spectral measures for one-frequency Schrödinger operators. *Comm. Math. Phys.*, 301(2):563–581, 2011.
- [8] A. Avila and R. Krikorian. Reducibility or nonuniform hyperbolicity for quasiperiodic Schrödinger cocycles. Ann. of Math. (2), 164(3):911–940, 2006.
- [9] A. Avila, J. You, and Q. Zhou. Sharp phase transitions for the almost Mathieu operator. Duke Math. J., 166(14):2697–2718, 2017.
- [10] S. Basu. On bounding the Betti numbers and computing the Euler characteristic of semi-algebraic sets. *Discrete Comput. Geom.*, 22(1):1–18, 1999.
- [11] G. Binyamini and D. Novikov. Complex cellular structures. Ann. of Math. (2), 190(1):145-248, 2019.

- [12] J. Bourgain. Quasi-periodic solutions of Hamiltonian perturbations of 2D linear Schrödinger equations. Ann. of Math. (2), 148(2):363–439, 1998.
- [13] J. Bourgain. Hölder regularity of integrated density of states for the almost Mathieu operator in a perturbative regime. *Lett. Math. Phys.*, 51(2):83–118, 2000.
- [14] J. Bourgain. Estimates on Green's functions, localization and the quantum kicked rotor model. Ann. of Math. (2), 156(1):249–294, 2002.
- [15] J. Bourgain. Green's function estimates for lattice Schrödinger operators and applications, volume 158 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 2005.
- [16] J. Bourgain. Positivity and continuity of the Lyapounov exponent for shifts on \mathbb{T}^d with arbitrary frequency vector and real analytic potential. J. Anal. Math., 96:313–355, 2005.
- [17] J. Bourgain. Anderson localization for quasi-periodic lattice Schrödinger operators on Z^d, d arbitrary. Geom. Funct. Anal., 17(3):682–706, 2007.
- [18] J. Bourgain and M. Goldstein. On nonperturbative localization with quasiperiodic potential. Ann. of Math. (2), 152(3):835–879, 2000.
- [19] J. Bourgain, M. Goldstein, and W. Schlag. Anderson localization for Schrödinger operators on Z with potentials given by the skew-shift. *Comm. Math. Phys.*, 220(3):583–621, 2001.
- [20] J. Bourgain, M. Goldstein, and W. Schlag. Anderson localization for Schrödinger operators on \mathbb{Z}^2 with quasi-periodic potential. *Acta Math.*, 188(1):41–86, 2002.
- [21] J. Bourgain and S. Jitomirskaya. Absolutely continuous spectrum for 1D quasiperiodic operators. *Invent. Math.*, 148(3):453–463, 2002.
- [22] J. Bourgain and I. Kachkovskiy. Anderson localization for two interacting quasiperiodic particles. *Geom. Funct. Anal.*, 29(1):3–43, 2019.
- [23] J. Bourgain and A. Klein. Bounds on the density of states for Schrödinger operators. *Invent. Math.*, 194(1):41–72, 2013.
- [24] J. Bourgain and W.-M. Wang. Anderson localization for time quasi-periodic random Schrödinger and wave equations. *Comm. Math. Phys.*, 248(3):429– 466, 2004.
- [25] J. Bourgain and W.-M. Wang. Quasi-periodic solutions of nonlinear random Schrödinger equations. J. Eur. Math. Soc. (JEMS), 10(1):1–45, 2008.
- [26] A. Cai, C. Chavaudret, J. You, and Q. Zhou. Sharp Hölder continuity of the Lyapunov exponent of finitely differentiable quasi-periodic cocycles. *Math.* Z., 291(3-4):931–958, 2019.
- [27] V. A. Chulaevsky and E. I. Dinaburg. Methods of KAM-theory for longrange quasi-periodic operators on \mathbf{Z}^{ν} . Pure point spectrum. *Comm. Math. Phys.*, 153(3):559–577, 1993.
- [28] V. A. Chulaevsky and Y. G. Sinaĭ. Anderson localization for the 1-D discrete Schrödinger operator with two-frequency potential. *Comm. Math. Phys.*,

125(1):91-112, 1989.

- [29] W. Craig and B. Simon. Log Hölder continuity of the integrated density of states for stochastic Jacobi matrices. *Comm. Math. Phys.*, 90(2):207–218, 1983.
- [30] D. Damanik, M. Goldstein, W. Schlag, and M. Voda. Homogeneity of the spectrum for quasi-periodic Schrödinger operators. J. Eur. Math. Soc. (JEMS), 20(12):3073–3111, 2018.
- [31] E. I. Dinaburg and J. G. Sinaĭ. The one-dimensional Schrödinger equation with quasiperiodic potential. *Funkcional. Anal. i Priložen.*, 9(4):8–21, 1975.
- [32] M. Drmota and R. F. Tichy. Sequences, discrepancies and applications, volume 1651 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 1997.
- [33] L. H. Eliasson. Floquet solutions for the 1-dimensional quasi-periodic Schrödinger equation. Comm. Math. Phys., 146(3):447–482, 1992.
- [34] J. Fröhlich, T. Spencer, and P. Wittwer. Localization for a class of onedimensional quasi-periodic Schrödinger operators. *Comm. Math. Phys.*, 132(1):5–25, 1990.
- [35] M. Goldstein and W. Schlag. Hölder continuity of the integrated density of states for quasi-periodic Schrödinger equations and averages of shifts of subharmonic functions. Ann. of Math. (2), 154(1):155–203, 2001.
- [36] M. Goldstein and W. Schlag. Fine properties of the integrated density of states and a quantitative separation property of the Dirichlet eigenvalues. *Geom. Funct. Anal.*, 18(3):755–869, 2008.
- [37] M. Goldstein and W. Schlag. Fine properties of the integrated density of states and a quantitative separation property of the Dirichlet eigenvalues. *Geom. Funct. Anal.*, 18(3):755–869, 2008.
- [38] M. Goldstein, W. Schlag, and M. Voda. On the spectrum of multi-frequency quasiperiodic Schrödinger operators with large coupling. *Invent. Math.*, 217(2):603–701, 2019.
- [39] M. Gromov. Entropy, homology and semialgebraic geometry. Number 145-146, pages 5, 225–240. 1987. Séminaire Bourbaki, Vol. 1985/86.
- [40] S. Hadj Amor. Hölder continuity of the rotation number for quasi-periodic co-cycles in SL(2, ℝ). Comm. Math. Phys., 287(2):565–588, 2009.
- [41] R. Han and S. Jitomirskaya. Full measure reducibility and localization for quasiperiodic Jacobi operators: a topological criterion. Adv. Math., 319:224– 250, 2017.
- [42] R. Han and S. Zhang. Large deviation estimates and Hölder regularity of the Lyapunov exponents for quasi-periodic Schrödinger cocycles. *International Mathematics Research Notices to appear.*
- [43] X. Hou and J. You. Almost reducibility and non-perturbative reducibility of quasi-periodic linear systems. *Invent. Math.*, 190(1):209–260, 2012.
- [44] S. Jitomirskaya. On point spectrum of critical almost Mathieu operators. Preprint.

- [45] S. Jitomirskaya and I. Kachkovskiy. L²-reducibility and localization for quasiperiodic operators. Math. Res. Lett., 23(2):431–444, 2016.
- [46] S. Jitomirskaya and I. Krasovsky. Critical almost mathieu operator: hidden singularity, gap continuity, and the Hausdorff dimension of the spectrum. *arXiv preprint arXiv:1909.04429*, 2019.
- [47] S. Jitomirskaya, H. Krüger, and W. Liu. Exact dynamical decay rate for the almost Mathieu operator. *Math. Res. Lett. to appear*.
- [48] S. Jitomirskaya and W. Liu. Arithmetic spectral transitions for the Maryland model. Comm. Pure Appl. Math., 70(6):1025–1051, 2017.
- [49] S. Jitomirskaya and W. Liu. Universal hierarchical structure of quasiperiodic eigenfunctions. Ann. of Math. (2), 187(3):721–776, 2018.
- [50] S. Jitomirskaya and W. Liu. Universal reflective-hierarchical structure of quasiperiodic eigenfunctions and sharp spectral transition in phase. arXiv preprint arXiv:1802.00781, 2018.
- [51] S. Jitomirskaya, W. Liu, and Y. Shi. Anderson localization for multifrequency quasi-periodic operators on \mathbb{Z}^d . *Geom. Funct. Anal.*, 30(2):457–481, 2020.
- [52] S. Jitomirskaya and C. A. Marx. Analytic quasi-perodic cocycles with singularities and the Lyapunov exponent of extended Harper's model. *Comm. Math. Phys.*, 316(1):237–267, 2012.
- [53] S. Jitomirskaya and S. Zhang. Quantitative continuity of singular continuous spectral measures and arithmetic criteria for quasiperiodic Schrödinger operators. *JEMS to appear*.
- [54] S. Y. Jitomirskaya. Anderson localization for the almost Mathieu equation: a nonperturbative proof. *Comm. Math. Phys.*, 165(1):49–57, 1994.
- [55] S. Y. Jitomirskaya. Metal-insulator transition for the almost Mathieu operator. Ann. of Math. (2), 150(3):1159–1175, 1999.
- [56] L. Kuipers and H. Niederreiter. Uniform distribution of sequences. Wiley-Interscience [John Wiley & Sons], New York-London-Sydney, 1974. Pure and Applied Mathematics.
- [57] W. Liu. Almost Mathieu operators with completely resonant phases. Ergodic Theory Dynam. Systems, 40(7):1875–1893, 2020.
- [58] W. Liu and Y. Shi. Upper bounds on the spectral gaps of quasiperiodic Schrödinger operators with Liouville frequencies. J. Spectr. Theory, 9(4):1223–1248, 2019.
- [59] W. Liu and X. Yuan. Anderson localization for the almost Mathieu operator in the exponential regime. J. Spectr. Theory, 5(1):89–112, 2015.
- [60] W. Liu and X. Yuan. Anderson localization for the completely resonant phases. J. Funct. Anal., 268(3):732–747, 2015.
- [61] W. Liu and X. Yuan. Hölder continuity of the spectral measures for onedimensional Schrödinger operator in exponential regime. J. Math. Phys., 56(1):012701, 21, 2015.

- [62] C. A. Marx and S. Jitomirskaya. Dynamics and spectral theory of quasiperiodic Schrödinger-type operators. *Ergodic Theory Dynam. Systems*, 37(8):2353–2393, 2017.
- [63] H. L. Montgomery. Ten lectures on the interface between analytic number theory and harmonic analysis, volume 84 of CBMS Regional Conference Series in Mathematics. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1994.
- [64] J. Moser and J. Pöschel. An extension of a result by Dinaburg and Sinaĭ on quasiperiodic potentials. *Comment. Math. Helv.*, 59(1):39–85, 1984.
- [65] W. Schlag. On the integrated density of states for Schrödinger operators on \mathbb{Z}^2 with quasi periodic potential. *Comm. Math. Phys.*, 223(1):47–65, 2001.
- [66] W. M. Schmidt. Metrical theorems on fractional parts of sequences. Trans. Amer. Math. Soc., 110:493–518, 1964.
- [67] J. Shi and X. Yuan. Anderson localization for Jacobi matrices associated with high-dimensional skew shifts. arXiv preprint arXiv:1908.00229, 2019.
- [68] B. Simon. Almost periodic Schrödinger operators. IV. The Maryland model. Ann. Physics, 159(1):157–183, 1985.
- [69] Y. G. Sinaĭ. Anderson localization for one-dimensional difference Schrödinger operator with quasiperiodic potential. J. Statist. Phys., 46(5-6):861–909, 1987.
- [70] K. Tao. Non-perturbative positive Lyapunov exponent of Schrödinger equations and its applications to skew-shift mapping. J. Differential Equations, 266(6):3559–3579, 2019.
- [71] K. Tao. Non-perturbative weak Hölder continuity of Lyapunov exponent of discrete analytic Jacobi operators with skew-shift mapping. *Electron. J. Differential Equations*, pages Paper No. 81, 17, 2019.
- [72] W.-M. Wang. Energy supercritical nonlinear Schrödinger equations: quasiperiodic solutions. *Duke Math. J.*, 165(6):1129–1192, 2016.
- [73] Y. Yomdin. C^k-resolution of semialgebraic mappings. Addendum to: "Volume growth and entropy". Israel J. Math., 57(3):301–317, 1987.
- [74] J. You. Quantitative almost reducibility and its applications. In Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. III. Invited lectures, pages 2113–2135. World Sci. Publ., Hackensack, NJ, 2018.
- [75] X. Zhao. Hölder continuity of absolutely continuous spectral measure for multi-frequency Schrödinger operators. arXiv preprint arXiv:2001.05121, 2020.

(W. Liu) Department of Mathematics, Texas A&M University, College Station, TX 77843-3368, USA

E-mail address: liuwencai1226@gmail.com