

**On the 2PN pericentre precession in the general theory of relativity and the recently discovered fast orbiting S-stars in Sgr A\***

Lorenzo Iorio<sup>1</sup>

Ministero dell'Istruzione, dell'Università e della Ricerca (M.I.U.R.)-Istruzione  
Permanent address for correspondence: Viale Unità di Italia 68, 70125, Bari (BA), Italy

lorenzo.iorio@libero.it

Received \_\_\_\_\_; accepted \_\_\_\_\_

## Abstract

Recently, the secular pericentre precession was analytically computed to the second post-Newtonian (2PN) order by the present author with the Gauss equations in terms of the osculating Keplerian orbital elements in order to obtain closer contact with the observations in astronomical and astrophysical scenarios of potential interest. A discrepancy with previous results by other authors was found. Moreover, some of such findings by the same authors were deemed as mutually inconsistent. In this paper, it is demonstrated that, in fact, two calculational errors plagued the most recent calculation by the present author. They are explicitly disclosed and corrected. As a result, all the examined approaches mutually agree yielding the same analytical expression for the total 2PN pericentre precession once the appropriate conversions from the adopted parameterizations are made. It is also shown that, in future, it may become measurable, at least in principle, for some of the recently discovered short-period S-stars in Sgr A\* like S62 and S4714.

keywords general relativity and gravitation; celestial mechanics

## 1. Introduction

The analytical calculation of the secular 2PN pericentre precession  $\dot{\omega}^{2\text{PN}}$  of a gravitationally bound two-body system made of two mass monopoles  $M_A$ ,  $M_B$  with the perturbative Gauss equations for the variation of the osculating Keplerian orbital elements (e.g. Kopeikin, Efroimsky & Kaplan 2011; Soffel & Han 2019) was the subject of Iorio (2020a). For the sake of simplicity, the test particle limit will be considered in most of the paper. In the following,  $c$  is the speed of light in vacuum,  $\mu \doteq GM$  is the gravitational parameter of the primary whose mass is  $M$ ,  $G$  is the Newtonian constant of gravitation,  $\mathbf{v}$ ,  $v_r$ ,  $v$  are the test particle’s velocity, radial velocity and speed, respectively,  $r$  is the test particle’s distance from the primary,  $\hat{\mathbf{r}}$  is the position unit vector of the test particle with respect to the primary,  $f_0$ ,  $a$ ,  $e$  are (Klioner & Kopeikin 1994) the osculating numerical values of the true anomaly, semimajor axis and eccentricity, respectively, at the same arbitrary moment of time  $t_0$ , and  $n_b = \sqrt{\mu/a^3}$  is the osculating mean motion.

The expression for the total 2PN pericentre precession derived by Iorio (2020a) consists of the sum of three contributions. The first one, dubbed as “direct”, is (Iorio 2020a, Equation (8))

$$\dot{\omega}_{\text{dir}}^{2\text{PN}} = \frac{n_b \mu^2 (28 - e^2)}{4 c^4 a^2 (1 - e^2)^2}, \quad (1)$$

arising straightforwardly from the 2PN acceleration

$$\mathbf{A}^{2\text{PN}} = \frac{\mu^2}{c^4 r^3} \left[ \left( 2 v_r^2 - \frac{9\mu}{r} \right) \hat{\mathbf{r}} - 2 v_r \mathbf{v} \right]. \quad (2)$$

There are also two further contributions, labeled as “mixed” or “indirect”. They account for the fact that, when the Gauss equation for the rate of change of the pericentre induced by the 1PN acceleration

$$\mathbf{A}^{1\text{PN}} = \frac{\mu}{c^2 r^2} \left[ \left( \frac{4\mu}{r} - v^2 \right) \hat{\mathbf{r}} + 4 v_r \mathbf{v} \right] \quad (3)$$

is averaged over one orbital period  $P_b$ , the latter one has to be considered as the time interval between two consecutive crossings of the moving pericentre. Moreover, also the instantaneous shifts of the other orbital elements due to Equation (3) itself are to be taken into account when the orbital average is performed. Both such effects contribute the total pericentre precession to the 2PN level. The first indirect effect yields (Iorio 2020a, Equation (14))

$$\dot{\omega}_{\text{indir}}^{2\text{PN(I)}} = \frac{n_b \mu^2 (9 + 37 e^2 + e^4)}{2 c^4 a^2 e^2 (1 - e^2)^2}, \quad (4)$$

while the second indirect contribution reads (Iorio 2020a, Equation (22))

$$\dot{\omega}_{\text{indir}}^{2\text{PN(II)}} = - \frac{n_b \mu^2 \{9 - 87 e^2 - 136 e^4 + 19 e^6 - 6 e^3 [(34 + 26 e^2) \cos f_0 + 15 e \cos 2f_0]\}}{2 c^4 e^2 a^2 (1 - e^2)^3}. \quad (5)$$

Thus, the sum of Equation (1), Equation (4), and Equation (5) gives the total 2PN pericentre precession

$$\dot{\omega}_{\text{tot}}^{2\text{PN}} = \frac{3 n_b \mu^2 [86 + 57 e^2 - 13 e^4 + 8 e (17 + 13 e^2) \cos f_0 + 60 e^2 \cos 2f_0]}{4 c^4 a^2 (1 - e^2)^3}. \quad (6)$$

Iorio (2020a) compared his results with those by Kopeikin & Potapov (1994), who used the perturbative approach relying upon the Gauss equations as well, and those by Damour & Schafer (1988), obtained with the Hamilton-Jacobi method. A discrepancy with such authors was found since Iorio (2020a) demonstrated that the total 2PN pericentre precession inferred by Kopeikin & Potapov (1994, Equation (5.2)) can be cast into the form

$$\dot{\omega}_{\text{tot}}^{2\text{PN}} = \frac{3 n_b \mu^2 (2 + e^2 - 32 e^2 \cos f_0)}{4 c^4 a^2 (1 - e^2)^2}. \quad (7)$$

Equation (7) was not explicitly shown by Kopeikin & Potapov (1994). Furthermore, Iorio (2020a) claimed that Damour & Schafer (1988, Equation (3.12)), which was shown to be coincident with Equation (7), and Damour & Schafer (1988, Equation (5.18)) would be mutually inconsistent.

Here, it will be proven that, actually, a mere calculational error occurred in the derivation of  $\dot{\omega}_{\text{indir}}^{2\text{PN(II)}}$  by Iorio (2020a) which prevented to obtain Equation (7) instead of the incorrect Equation (6). Once such an error is corrected, both the approaches by Iorio (2020a) and Kopeikin & Potapov (1994), which differ in how obtaining just  $\dot{\omega}_{\text{indir}}^{2\text{PN(II)}}$ , agree yielding the same

total 2PN pericentre precession of Equation (7). Moreover, it will be shown that also the alleged inconsistency of Equation (3.12) and Equation (5.18) by Damour & Schafer (1988) is, in fact, due to another error by Iorio (2020a), as correctly pointed out by Kopeikin (2020).

To the benefit of the reader, it is noted that Damour & Schafer (1988); Kopeikin & Potapov (1994) usually dealt with the fractional pericentre advance per orbit, i.e.,  $\Delta\omega/2\pi$ ; in order to obtain the corresponding precession, it is sufficient to multiply it by  $n_b$ .

The paper is organized as follows. In Section 2, the calculational error in working out  $\dot{\omega}_{\text{indir}}^{2\text{PN(II)}}$  is explicitly disclosed and corrected. Section 3 is devoted to showing, independently of Kopeikin (2020), that Equation (3.12) and Equation (5.18) of Damour & Schafer (1988) are, actually, mutually consistent yielding both the same total 2PN pericentre precession as Equation (7). Some aspects of the presence of  $f_0$  in Equation (7) are discussed in Section 4. Section 5 is devoted to calculating the 2PN pericentre precession for some of the recently discovered fast orbiting S-stars (Peißker et al. 2020) in the Galactic Center (GC) at Sgr A\*. Section 6 summarizes the present findings and offers concluding remarks.

## 2. Disclosing and correcting the error for $\dot{\omega}_{\text{indir}}^{2\text{PN(II)}}$

In Iorio (2020a), it turned out that Equation (1) and Equation (4) agree with the corresponding calculation by Kopeikin & Potapov (1994), despite such authors did neither recur to the schematization by Iorio (2020a) nor explicitly display their intermediate results.

Instead, Iorio (2020a) realized that the discrepancy among his results and those by Kopeikin & Potapov (1994) resides in  $\dot{\omega}_{\text{indir}}^{2\text{PN(II)}}$ , i.e., in that part of the indirect precession arising from the fact that the semimajor axis and the eccentricity do change instantaneously during an orbital revolution due to Equation (3) according to

$$\Delta a(f_0, f)^{\text{1PN}} = -\frac{2e\mu(\cos f - \cos f_0)[7 + 3e^2 + 5e(\cos f + \cos f_0)]}{c^2(1 - e^2)^2}, \quad (8)$$

$$\Delta e(f_0, f)^{\text{1PN}} = \frac{\mu(\cos f_0 - \cos f)[3 + 7e^2 + 5e(\cos f + \cos f_0)]}{c^2 a(1 - e^2)}. \quad (9)$$

The calculation of  $\dot{\omega}_{\text{indir}}^{2\text{PN(II)}}$  by Kopeikin & Potapov (1994) can be reproduced as follows (Iorio 2020a, pp. 13). Evaluate the Gauss equation of the pericentre for a perturbing in-plane acceleration

$$\frac{d\omega}{df} = \frac{r^2}{\mu e} \left[ -A_\rho \cos f + \left( 1 + \frac{r}{p} \right) \sin f A_\tau \right], \quad (10)$$

where  $p \doteq a(1 - e^2)$ , with the radial and transverse components of the 1PN acceleration of

Equation (3)

$$A_{\rho}^{1\text{PN}} = \frac{\mu^2 (1 + e \cos f)^2 (3 + e^2 + 2e \cos f - 2e^2 \cos 2f)}{c^2 a^3 (1 - e^2)^3}, \quad (11)$$

$$A_{\tau}^{1\text{PN}} = \frac{4e\mu^2 (1 + e \cos f)^3 \sin f}{c^2 a^3 (1 - e^2)^3}. \quad (12)$$

Then, make the replacement

$$a \rightarrow a + \Delta a(f_0, f)^{1\text{PN}}, \quad (13)$$

$$e \rightarrow e + \Delta e(f_0, f)^{1\text{PN}}, \quad (14)$$

by means of Equations (8)-(9), expand  $d\omega/df$  to the order of  $O(c^{-4})$ , and integrate the resulting expression

$$\begin{aligned} \left. \frac{d\omega}{df} \right|_{\text{indir}}^{2\text{PN(II)}} &= -\frac{\mu^2 (\cos f - \cos f_0)}{2c^4 a^2 e^2 (1 - e^2)^2} \left\{ e \left[ 15 - 43e^2 + 5(3 + 17e^2) \cos 2f \right] + \right. \\ &\quad \left. + 2 \cos f \left[ 9 + 48e^2 - e^4 + 5e(3 + e^2) \cos f_0 \right] \right\} \end{aligned} \quad (15)$$

from  $f_0$  to  $f_0 + 2\pi$ . By dividing the result by  $1/P_b = n_b/2\pi$ , the following formula is obtained

$$\dot{\omega}_{\text{indir}}^{2\text{PN(II)}} = \frac{n_b \mu^2 (-9 - 48e^2 + e^4 - 48e^3 \cos f_0)}{2c^4 a^2 e^2 (1 - e^2)^2}, \quad (16)$$

so that the sum of Equation (4) and Equation (16) gives the total 2PN indirect precession (Iorio 2020a, Equation (55))

$$\dot{\omega}_{\text{indir}}^{2\text{PN}} = \frac{n_b \mu^2 (-11 + 2e^2 - 48e \cos f_0)}{2c^4 a^2 (1 - e^2)^2}, \quad (17)$$

which is, actually, correct. It turns out that adding Equation (17) to Equation (1) yields just the total 2PN precession of Equation (7).

Iorio (2020a, pag. 5) followed another approach in calculating  $\dot{\omega}_{\text{indir}}^{2\text{PN(II)}}$ . In the specific case of the pericentre and of Equation (3), starting from Equation (10), calculated with Equations (11)-(12), the net 2PN pericentre shift per orbit due to the 1PN instantaneous variations of Equations (8)-(9) is worked out as

$$\Delta\omega_{\text{indir}}^{2\text{PN(II)}} = \int_{f_0}^{f_0+2\pi} \left[ \frac{\partial(d\omega/df)}{\partial a} \Delta a(f_0, f)^{1\text{PN}} + \frac{\partial(d\omega/df)}{\partial e} \Delta e(f_0, f)^{1\text{PN}} \right] df. \quad (18)$$

A calculational error<sup>1</sup> in the first addend of Equation (18) yielded the wrong result of Equation (5). After correcting it, it is possible to show that the function to be integrated in Equation (18) agrees with Equation (15). Thus, Equation (16) can be correctly obtained also with the method for calculating  $\dot{\omega}_{\text{indir}}^{2\text{PN}(\text{II})}$  used by Iorio (2020a).

By repeating the calculation by Iorio (2020a), corrected for the aforementioned error, one obtains, for the full two-body system,

$$\dot{\omega}_{\text{indir}}^{2\text{PN}} = \frac{n_b \mu^2 \left[ -44 + 8 \nu (-8 + 7 \nu) + e^2 (8 + 39 \nu + 48 \nu^2) + 96 e (-2 + \nu) \cos f_0 \right]}{8 c^4 a^2 (1 - e^2)^2} \quad (19)$$

which, added to Iorio (2020a, Equation (32)), returns

$$\dot{\omega}_{\text{tot}}^{2\text{PN}} = \frac{3 n_b \mu^2 \left[ 2 - 4 \nu + e^2 (1 + 10 \nu) + 16 e (-2 + \nu) \cos f_0 \right]}{4 c^4 a^2 (1 - e^2)^2}. \quad (20)$$

In Equations (19)-(20), it is

$$\nu \doteq \frac{M_A M_B}{(M_A + M_B)^2}, \quad (21)$$

$$\mu \doteq G (M_A + M_B). \quad (22)$$

As far as Mercury is concerned, for which it is (Iorio 2020a, Fig. 1)

$$\dot{\omega}_{\text{dir}}^{2\text{PN}} = 2.6 \mu\text{as cty}^{-1}, \quad (23)$$

where  $\mu\text{as cty}^{-1}$  stands for microarcseconds per century, Equation (17) yields

$$-4 \mu\text{as cty}^{-1} \leq \dot{\omega}_{\text{indir}}^{2\text{PN}} \leq -0.2 \mu\text{as cty}^{-1}, \quad (24)$$

for  $0^\circ \leq f_0 \leq 360^\circ$ ; Equation (24) corrects Iorio (2020a, Equation (24)).

For the double pulsar PSR J0737-3039A/B, for which it is (Iorio 2020a, Equation (33))

$$\dot{\omega}_{\text{dir}}^{2\text{PN}} = 0.00019^\circ \text{ yr}^{-1}, \quad (25)$$

from Equation (19), it turns out

$$-0.00022^\circ \text{ yr}^{-1} \leq \dot{\omega}_{\text{indir}}^{2\text{PN}} \leq -0.00013^\circ \text{ yr}^{-1} \quad (26)$$

---

<sup>1</sup>To be more specific,  $\mu^2$  entering Equations (11)-(12) was expressed in  $d\omega/df$  as  $n_b^4 a^6$ , thus altering the partial derivative of  $d\omega/df$  with respect to  $a$ .

for  $0^\circ \leq f_0 \leq 360^\circ$ . For the Hulse-Taylor binary pulsar PSR B1913+16, for which it is (Iorio 2020a, Equation (35))

$$\dot{\omega}_{\text{dir}}^{2\text{PN}} = 0.000038^\circ \text{ yr}^{-1}, \quad (27)$$

Equation (19) yields

$$-0.00009^\circ \text{ yr}^{-1} \leq \dot{\omega}_{\text{indir}}^{2\text{PN}} \leq 0.000034^\circ \text{ yr}^{-1} \quad (28)$$

for  $0^\circ \leq f_0 \leq 360^\circ$ . Equation (26) and Equation (28) correct Iorio (2020a, Equations (48)-(49)).

For the supermassive binary black hole in OJ 287, for which it is (Iorio 2020a, pag. 10)

$$\dot{\omega}_{\text{dir}}^{2\text{PN}} = 11.0^\circ \text{ cty}^{-1}, \quad (29)$$

Equation (19) returns an indirect 2PN perinigricon precession ranging within

$$-33.4^\circ \text{ cty}^{-1} \leq \dot{\omega}_{\text{indir}}^{2\text{PN}} \leq 17^\circ \text{ cty}^{-1} \quad (30)$$

for  $0^\circ \leq f_0 \leq 360^\circ$ . Equation (30) corrects the figures yielded in Iorio (2020a, pag. 12). In retrospect, they should have been a wake-up call concerning the validity of Iorio (2020a, Equation (47)) since the reported maximum value of  $516^\circ \text{ cty}^{-1}$  is even larger than than the 1PN precession itself amounting to (Iorio 2020a, Equation (37))  $\dot{\omega}^{1\text{PN}} = 206.8^\circ \text{ cty}^{-1}$ .

The discussion in Iorio (2020a) concerning the measurability of the 2PN pericentre precessions of Mercury and of the binary pulsars will not be repeated here.

### 3. Correcting the error for $e_T$

As correctly pointed out by Kopeikin (2020), Iorio (2020a) erroneously claimed that Equation (3.12) of Damour & Schafer (1988)

$$\frac{\Delta\omega_{\text{tot}}^{2\text{PN}}}{2\pi} = \frac{3}{c^2 h^2} \left[ 1 + \left( \frac{5}{2} - \nu \right) \frac{E}{c^2} + \left( \frac{35}{4} - \frac{5}{2}\nu \right) \frac{1}{c^2 h^2} \right], \quad (31)$$

and Equation (5.18) of Damour & Schafer (1988)

$$\begin{aligned} \frac{\Delta\omega_{\text{tot}}^{2\text{PN}}}{2\pi} = & \frac{3(\mu n)^{2/3}}{c^2 (1 - e_T^2)} \left[ 1 + \frac{(\mu n)^{2/3}}{c^2 (1 - e_T^2)} \left( \frac{39}{4} x_A^2 + \frac{27}{4} x_B^2 + 15 x_A x_B \right) - \right. \\ & \left. - \frac{(\mu n)^{2/3}}{c^2} \left( \frac{13}{4} x_A^2 + \frac{1}{4} x_B^2 + \frac{13}{3} x_A x_B \right) \right] \end{aligned} \quad (32)$$

would be mutually inconsistent after being expressed in terms of  $a$ ,  $e$ ,  $f_0$ . In Equations (31)-(32),  $h$  and  $E$  are the coordinate-invariant, reduced orbital angular momentum and energy, respectively,

$$x_A \doteq \frac{M_A}{M_A + M_B}, \quad (33)$$

$$x_B \doteq \frac{M_B}{M_A + M_B} = 1 - x_A, \quad (34)$$

$n$  is the PN mean motion (Damour & Deruelle 1985), and  $e_T$  is one of the several Damour-Deruelle (DD) parameters (Damour & Deruelle 1986). More precisely, in Iorio (2020a, pp. 14-15), it was correctly demonstrated that Equation (31) yields Equation (7). On the other hand, in Iorio (2020a, pag. 15), it was erroneously claimed that Equation (32) could not reduce to Equation (7). The error consists of the fact that Iorio (2020a) confused  $e_T$  (Damour & Deruelle 1986) entering Equation (32) with  $e_t$ , another member of the DD parameterization (Damour & Deruelle 1985).

Instead, it is (Damour & Deruelle 1986, pag. 272)

$$e_T = e_t(1 + \delta) + e_\theta - e_r. \quad (35)$$

The parameters entering Equation (35) are defined as (Damour & Deruelle 1985, Equation (3.8 b))

$$e_t = \frac{e_R}{1 + \frac{\mu}{c^2 a_R} \left(4 - \frac{3}{2} \nu\right)}, \quad (36)$$

(Damour & Deruelle 1986, Equation (20))

$$\delta = \frac{\mu}{c^2 a_R} (x_A x_B + 2x_B^2), \quad (37)$$

(Damour & Deruelle 1985, Equation (4.13))

$$e_\theta = e_R \left(1 + \frac{\mu \nu}{2 c^2 a_R}\right), \quad (38)$$

(Damour & Deruelle 1985, Equation (6.3 b))

$$e_r = e_R \left[1 - \frac{\mu}{2 c^2 a_R} (x_A^2 - \nu)\right]. \quad (39)$$

In Equations (36)-(39), the DD “semimajor axis” of the relative motion<sup>2</sup>  $a_R$  can be expressed as

---

<sup>2</sup>In Iorio (2020a), it is designed as  $a_r$ : in fact, such a choice may be confusing since, in Damour & Deruelle (1985, Equation (6.3 a)), such a quantity is meant as  $a_r = M_B a_R / (M_A + M_B)$ .



(Iorio 2020a, Equation (66))

$$\begin{aligned}
 4 (1 - e^2)^2 a_R &= 4 \left\{ a (1 - e^2)^2 - \frac{\mu}{c^2} [-3 + \nu + e^4 (1 + 2\nu) + e^2 (-13 + 7\nu)] \right\} + \\
 &+ e \frac{\mu}{c^2} \left\{ [56 + e^2 (24 - 31\nu) - 24\nu] \cos f_0 + \right. \\
 &\left. + e [4 (5 - 4\nu) \cos 2f_0 - e\nu \cos 3f_0] \right\}, \tag{40}
 \end{aligned}$$

and the DD “eccentricity”  $e_R$  is given by<sup>3</sup> Iorio (2020a, Equation (67))

$$\begin{aligned}
 8 a (-1 + e^2) e_R &= 4 e \left\{ 2 a (-1 + e^2) + \frac{\mu}{c^2} [-17 + 6\nu + e^2 (2 + 4\nu)] \right\} + \\
 &+ \frac{\mu}{c^2} \left\{ [8 (-3 + \nu) + e^2 (-56 + 47\nu)] \cos f_0 + \right. \\
 &\left. + e [4 (-5 + 4\nu) \cos 2f_0 + e\nu \cos 3f_0] \right\}. \tag{41}
 \end{aligned}$$

It turns out that, using Equations (35)-(41) and (Damour & Deruelle 1985, Equation (3.7))

$$n = \sqrt{\frac{\mu}{a_R^3}} \left[ 1 + \frac{\mu}{2 c^2 a_R} (-9 + \nu) \right] \tag{42}$$

in Equation (32) and expanding all to the order of  $\mathcal{O}(c^{-4})$  removes the previously mentioned alleged discrepancy. Indeed, now, the corresponding 2PN precession can be cast just into the form of Equation (7), inasmuch the same way as Equation (31) did.

#### 4. Some considerations about the occurrence of $f_0$ in the 2PN pericentre precession

The fact that  $f_0$  enters Equation (7) through Equation (16) seems to be likely a general feature of that part of the long-term orbital precessions arising from the inclusion of the instantaneous shifts of all the Keplerian orbital elements in performing the average of the right-hand-side of the Gauss equation of any of them over an orbital period. Indeed, such a feature explicitly occurs also with the interplay of, e.g., the Newtonian acceleration due to the primary’s quadrupole mass moment  $J_2$  and Equation (3) in calculating part of the indirect rates of change of the Keplerian

---

<sup>3</sup>In Iorio (2020a), it is denoted as  $e_r$ , but, in view of Equation (39), such a choice is misleading.

orbital elements of the order of  $J_2/c^2$  by means of the Gauss equations, as done by<sup>4</sup> Will (2014); Iorio (2015).

It should be remarked that Equation (7) does not pretend, by no means, to contain any new physics with respect to what was obtained earlier, with different computational approaches and different parameterizations, by Damour & Schafer (1988) and Kopeikin & Potapov (1994) in terms of first integrals of motion like  $E$  and  $h$ . Indeed, as it was shown here, Equation (5.2) by Kopeikin & Potapov (1994), and Equation (3.12) and Equation (5.18) by Damour & Schafer (1988) agree with Equation (7). The appearance of  $f_0$  in Equation (7) is of a gauge nature, in that it does not influence the values of the integrals of motion, for a fixed orbit. It should be remembered that, after all, in any practical data analysis of specific astronomical and astrophysical systems of interest, an underlying choice of a given parameterization and gauge is always unavoidably made. In most cases, it generally corresponds to that adopted in the present study, which is just targeted to practical people aiming to designing and/or performing tests using, say, Mercury, exoplanets close to their parent stars, and S-stars in Sgr A\*.

### 5. Looking at some of the short-period S-stars in Sgr A\*

Recently, Peißker et al. (2020) discovered some short-period S-stars orbiting the candidate supermassive black hole (SMBH) in the Galactic Center (GC) at Sgr A\* (Ghez et al. 2008; Genzel, Eisenhauer & Gillessen 2010); their relevant orbital parameters are reported in Table 1. Since some of them can move as fast as  $\approx 10$  per cent of the speed of light  $c$  at perinigricon,

Table 1: Key orbital parameters of some fast-revolving S-stars as per Table 2 of Peißker et al. (2020). Here, the semimajor axis  $a$  is in milliparsec (mpc), the orbital period  $P_b$  is in yr, while the eccentricity  $e$  is dimensionless.

star	$a$ (mpc)	$P_b$ (yr)	$e$
S4714	$4.079 \pm 0.012$	$12.0 \pm 0.3$	$0.985 \pm 0.011$
S62	$3.588 \pm 0.02$	$9.9 \pm 0.3$	$0.976 \pm 0.01$
S4711	$3.002 \pm 0.06$	$7.6 \pm 0.3$	$0.768 \pm 0.030$

it seems appropriate to look at their 2PN orbital effects. Lately, Fragione & Loeb (2020); Iorio

---

<sup>4</sup>A terminological confusion may arise if one does not properly note that Will (2014) used the adjective “mixed” to denote the PN-quadrupole acceleration of the order of  $J_2/c^2$  (Soffel et al. 1987; Soffel 1989; Brumberg 1991) entering the equations of motion. Moreover, Will (2014) dubbed “cross-term effects” the contributions to the overall orbital precessions labeled as “indirect”, or “mixed”, by Iorio (2015) and here.

(2020b); Peißker et al. (2020) investigated the 1PN gravitomagnetic Lense-Thirring orbital precessions, induced by the SMBH’s angular momentum  $\mathbf{J}_\bullet$ , in view of a possible detection in future.

Figure 1 displays the 2PN perinigricon precessions  $\dot{\omega}^{2\text{PN}}$  and shifts per orbit  $\Delta\omega^{2\text{PN}}$  of the S-stars S4711, S4714, S62 as functions of  $f_0$  according to Equation (7). It turns out that the largest




Fig. 1.— Upper panel: plot of the 2PN perinigricon precessions  $\dot{\omega}^{2\text{PN}}$  of the S-stars S4711, S4714, S62 (Peißker et al. 2020), in arcseconds per year (″ yr<sup>-1</sup>), as a function of the true anomaly  $f_0$  at epoch according to Equation (7). Lower panel: plot of the 2PN perinigricon shifts per orbit  $\Delta\omega^{2\text{PN}} = 2\pi/n_b \dot{\omega}^{2\text{PN}}$  of the S-stars S4711, S4714, S62 (Peißker et al. 2020), in arcseconds (″), as a function of the true anomaly  $f_0$  at epoch according to Equation (7).

precessions, of the order of  $|\dot{\omega}^{2\text{PN}}| \lesssim 4 - 6$  arcseconds per year (″ yr<sup>-1</sup>), occur for S62 and S4714, whose shifts per orbit can be as large as  $|\Delta\omega^{2\text{PN}}| \lesssim 40 - 90$ ″. Such figures are about of the same order of magnitude of the largest possible values of the 1PN Lense-Thirring precessions for  $\omega$ , as shown by Table 4 of Iorio (2020b). Although future improvements in the observational techniques

may possibly bring such post-Newtonian effects within the range of measurability (Peißker et al. 2020), it should be remarked that they may be biased by the currently much larger systematic bias due to the lingering uncertainties in the stellar orbital parameters and the SMBH’s mass entering the dominant 1PN component of the perinigricon precession

$$\dot{\omega}^{\text{1PN}} = \frac{3 n_b \mu}{c^2 a (1 - e^2)}; \tag{43}$$

see Table 5 of Iorio (2020b).

## 6. Summary and conclusions

After having disclosed and corrected the calculational errors affecting Iorio (2020a), it was demonstrated that the approaches by Damour & Schafer (1988); Kopeikin & Potapov (1994); Iorio (2020a) are, in fact, equivalent in analytically calculating the 2PN pericentre precession. Indeed, they yield the same result once the appropriate conversions from the adopted parameterizations are made. This demonstrates that the approach by Iorio (2020a) is correct.

The fact that the total 2PN pericentre rate of Equation (7) contains also  $f_0$  due to the indirect contributions to it arising from the cross-coupling of the 1PN acceleration with itself in the calculational procedure is an example of a general feature which seems to characterize also other mixed effects due to the interplay of different accelerations (Newtonian and PN) in the orbital averaging of the Gauss equations when the instantaneous shifts of all the Keplerian orbital elements are taken into account as well. The same feature occurs, e.g., also for the mixed precessions of the order of  $J_2/c^2$  due to the Newtonian quadrupole acceleration and the 1PN one, as explicitly calculated also by other authors in the literature. Nonetheless, this fact should not induce one to think about it as a sign of some new physics, being simply a matter of choice of a particular gauge. Choosing a given coordinate system and a gauge is a standard and unavoidable practice in actual data reductions and in designing possible future tests with astronomical and astrophysical systems of potential interest.

To this aim, Equation (7) was used to calculate the magnitude of the 2PN perinigricon precessions for some of the recently discovered fast-orbiting S-stars in Sgr A\* whose orbital configuration is given in the literature just in terms of the osculating Keplerian orbital elements adopted here. It turns out that the shifts per orbit of S62 and S4714 can be as large as  $\approx 40 - 90$  arcseconds, being, in principle, detectable in view of the expected progress in the observational techniques. Nonetheless, a major source of systematic bias lies in the lingering uncertainty in the much larger 1PN component of the stellar perinigricon rates.

### **Acknowledgements**

I am grateful to M. Efroimsky for useful advices.

## REFERENCES

- Brumberg V. A., 1991, *Essential Relativistic Celestial Mechanics*. Adam Hilger, Bristol
- Damour T., Deruelle N., 1985, *Ann. Inst. Henri Poincaré Phys. Théor.*, 43, 107
- Damour T., Deruelle N., 1986, *Ann. Inst. Henri Poincaré Phys. Théor.*, 44, 263
- Damour T., Schafer G., 1988, *Nuovo Cimento B*, 101, 127
- Fragione G., Loeb A., 2020, *ApJ*, 901, L32
- Genzel R., Eisenhauer F., Gillessen S., 2010, *Reviews of Modern Physics*, 82, 3121
- Ghez A. M. et al., 2008, *ApJ*, 689, 1044
- Iorio L., 2015, *Int. J. Mod. Phys. D*, 24, 1550067
- Iorio L., 2020a, *Universe*, 6, 53
- Iorio L., 2020b, *ApJ*, arXiv:2009.01158
- Klioner S. A., Kopeikin S. M., 1994, *ApJ*, 427, 951
- Kopeikin S., Efroimsky M., Kaplan G., 2011, *Relativistic Celestial Mechanics of the Solar System*. Weinheim: Wiley-VCH
- Kopeikin S. M., 2020, *Eur. J. Phys. Plus*, 135, 466
- Kopeikin S. M., Potapov V. A., 1994, *Astron. Rep.*, 38, 104
- Peißker F., Eckart A., Zajaček M., Ali B., Parsa M., 2020, *ApJ*, 899, 50
- Soffel M., Wirrer R., Schastok J., Ruder H., Schneider M., 1987, *Celest. Mech. Dyn. Astr.*, 42, 81
- Soffel M. H., 1989, *Relativity in Astrometry, Celestial Mechanics and Geodesy*. Springer, Heidelberg
- Soffel M. H., Han W.-B., 2019, *Applied General Relativity, Astronomy and Astrophysics Library*. Springer Nature Switzerland, Cham
- Will C. M., 2014, *Phys. Rev. D*, 89, 044043