

# Robust Inference for Mediated Effects in Partially Linear Models

Oliver Hines<sup>1</sup>, Stijn Vansteelandt<sup>1,2</sup>, and Karla Diaz-Ordaz<sup>1</sup>

<sup>1</sup>Department of Medical Statistics, London School of Hygiene and Tropical Medicine, London, U.K.

<sup>2</sup>Department of Applied Mathematics, Computer Science and Statistics, Ghent University, Ghent, Belgium

May 11, 2022

## Abstract

We consider mediated effects of an exposure,  $X$  on an outcome,  $Y$ , via a single mediator,  $M$ , under the assumption of no unmeasured confounding in the setting where models for the conditional expectation of the mediator and outcome are partially linear. In this setting the indirect effect is defined through a product of model coefficients, and the direct effect by a single model coefficient.

We propose G-estimators for the direct and indirect effect and demonstrate consistent asymptotically normality for indirect effects when models for  $M$ , or  $X$  and  $Y$  are correctly specified, and for direct effects, when models for  $Y$ , or  $X$  and  $M$  are correct. This marks an improvement over previous ‘triple’ robust methods, which do not assume partially linear mean models, but instead require correct specification of any pair of: the conditional expectation of  $Y$  and the conditional densities of  $M$  and  $X$ .

Testing of the no-mediation hypothesis is inherently problematic due to the composite nature of the test (either  $X$  has no effect on  $M$  or  $M$  no effect on  $Y$ ), leading to low power when both effect sizes are small. We use Generalized Methods of Moments (GMM) results to construct a new score testing framework, which includes as special cases the no-mediation and the no-direct-effect hypotheses. The proposed tests rely on a bias-reduction strategy for estimating parameters in nuisance confounder models.

Simulations show that the GMM based tests perform better in terms of power and small sample performance compared with traditional tests in the partially linear setting, with drastic improvement under model misspecification. New methods are illustrated in a mediation analysis of data from the COPERS trial, a randomized trial investigating the effect of a non-pharmacological intervention of patients suffering from chronic pain. An accompanying R package implementing these methods can be found at [github.com/ohines/plmed](https://github.com/ohines/plmed).

**Keywords:** G-estimation, Mediation, Robust Inference

## 1 Introduction

Testing and estimation of mediated effects is important in psychology, sociology, epidemiology, and econometrics, typically as a secondary analysis once a total effect has been established. These analyses infer how much of the effect is mediated by a given intermediate variable, drawing on  $n$  iid observations of an exposure ( $X$ ), outcome ( $Y$ ), mediating variable ( $M$ ), and a low dimensional collection of variables ( $Z$ ), that are sufficient to adjust for confounding of the association between  $X$  and  $M$  and between  $(X, M)$  and  $Y$ .

In this paper we consider continuous  $M$  and  $Y$  and assume the following partially linear mean models

$$E(M|X, Z) = \beta_1 X + f(Z, \gamma_m) \tag{1}$$

$$E(Y|M, X, Z) = \beta_2 M + \beta_3 X + g(Z, \gamma_y) \tag{2}$$

where  $f$  and  $g$  are known, possibly non-linear, differentiable functions parametrized by nuisance parameters  $\gamma_m$  and  $\gamma_y$ . Early work by Baron and Kenny (1986) defined indirect and direct effects as the product  $\beta_1\beta_2$  and the value  $\beta_3$  respectively, with the total effect given by the sum of the two effects:  $\beta_1\beta_2 + \beta_3$ . These cohere with later definitions of the natural indirect effect (NIDE), and the natural direct effect (NDE), which were developed for general non-linear models by Robins and Greenland (1992); Pearl (2001). When the total effect is nonzero, the ‘mediation proportion’ can be defined as

$$\alpha = \frac{\beta_1\beta_2}{\beta_1\beta_2 + \beta_3}.$$

Furthermore, if the NIDE and NDE are of the same sign (or one is zero), then  $\alpha \in [0, 1]$  may be interpreted as the proportion of the total effect which is mediated by  $M$ .

In this work we consider estimation of the NIDE and NDE, and testing of the null hypothesis  $H_\alpha : (\alpha - 1)\beta_1\beta_2 + \alpha\beta_3 = 0$ , that either the mediation proportion is equal to  $\alpha$ , or else (for  $\alpha \neq 0, 1$ ) the NIDE and NDE are both zero. The cases,  $\alpha = 0$  and  $\alpha = 1$  are of particular interest and correspond respectively to the no-mediation hypothesis ( $H_0 : \beta_1\beta_2 = 0$ ) and the no-direct-effect hypothesis ( $H_1 : \beta_3 = 0$ ).

Estimation of the NIDE and NDE in this linear setting is routinely based on Ordinary Least Squares (OLS), requiring that the models for  $E(M|X, Z)$  and  $E(Y|M, X, Z)$  are both correctly specified. In the more general, non-linear setting, ‘triple robust’ estimators of the NDE and NIDE have been constructed by Tchetgen Tchetgen and Shpitser (2012), so called due to their similarity to ‘double robust’ methods. Double robust methods of the average treatment effect, for example, typically produce Consistent Asymptotically Normal (CAN) point estimates provided that either a mean outcome model, or propensity score model is correctly specified (Kang and Schafer, 2007).

Triple robust estimators of the NDE and NIDE are proven to be CAN provided any pair of the following are correctly specified: (i) the conditional expectation,  $E(Y|M, X, Z)$ , (ii) the conditional density of  $M|X, Z$ , and (iii) the conditional density of  $X|Z$ . These methods are locally efficient provided all three models are specified correctly.

In this paper we construct G-estimators for the NIDE and NDE assuming that the mediator and outcome mean models are partially linear. We show that a G-estimator for the NIDE is CAN when either

- (i)  $E(M|X, Z)$  is correctly specified
- (ii) Both  $E(Y|M, X, Z)$  and  $E(X|Z)$  are correctly specified;

and that the G-estimator for the NDE is CAN provided either

- (i)  $E(Y|M, X, Z)$  is correctly specified
- (ii) Both  $E(M|X, Z)$  and  $E(X|Z)$  are correctly specified.

Hence, under the partially linear models in (1) and (2), G-estimators possess increased robustness compared with the estimator of Tchetgen Tchetgen and Shpitser (2012), as they rely only on the conditional mean models for  $M$  and  $X$ , whereas the triple robust estimator requires models for their full conditional densities, which may be difficult to specify in practice. This increased robustness is essential in applications involving continuous exposure and/or mediator.

Testing of the no-mediation hypothesis,  $H_0$ , is problematic since the function used to constrain the hypothesized parameter space ( $\psi_0(\beta) = \beta_1\beta_2 = 0$ ) has a Jacobian which is full rank almost everywhere, except for a singular point at  $\beta_1 = \beta_2 = 0$ . This generally gives rise to test statistics with different asymptotic behaviour at this singular point, and in finite samples, tests for  $H_0$  which are underpowered in its neighbourhood. We refer the interested reader to Dufour et al. (2013) for further details.

Classical tests of the no-mediation hypothesis are constructed from t-test statistics for the OLS parameter estimates of  $\beta_j$ , for  $j = 1, 2$ . Denoting by  $\hat{\beta}_j^{(OLS)}$  the OLS estimator, with estimated standard error,  $\hat{\sigma}_j^{(OLS)}$  and squared t-statistics,  $T_j^{(OLS)} = (\hat{\beta}_j^{(OLS)}/\hat{\sigma}_j^{(OLS)})^2$ , it can be demonstrated that  $T_1^{(OLS)}$  and  $T_2^{(OLS)}$  are independent under correct model specification.

Using these squared t-statistics, a Wald test for  $H_0$ , also known as the Sobel test (Sobel (1982)) can be constructed, based on the test statistic  $W^{(OLS)}$ . Alternatively, a joint test, based on the statistic  $LR^{(OLS)}$  (also known as a Likelihood Ratio (LR) test, (MacKinnon et al., 2002; Giersbergen, 2014)). These statistics are given by

$$W^{(OLS)} = \frac{T_1 T_2}{T_1 + T_2} = \frac{\hat{\beta}_1^2 \hat{\beta}_2^2}{\hat{\beta}_1^2 \hat{\sigma}_2^2 + \hat{\beta}_2^2 \hat{\sigma}_1^2} \quad (3)$$

$$LR^{(OLS)} = \min(T_1, T_2) \quad (4)$$

where for readability, the superscript (*OLS*) has been dropped from all terms on the right hand side. These statistics have received considerable attention, especially due to unexpected properties regarding the relative power of total, direct, and indirect effect tests under different true parameter values (Wang, 2018; Kenny and Judd, 2014; Fritz et al., 2012).

In this paper, we propose two alternative tests based on the moment conditions for the G-estimator: a Wald type approach, and an approach analogous to a classical score (or Lagrange Multiplier) test, but derived using a Generalized Methods of Moments (GMM) hypothesis testing framework. The relative merits of the new tests against the OLS based tests above are discussed in Section 4.3. From a robustness perspective, tests based on OLS estimating equations require that  $E(M|X, Z)$  and  $E(Y|M, X, Z)$  are correctly specified, whereas those based on the G-estimation equations inherit the same robustness to model misspecification as the G-estimator itself, provided that nuisance parameters are estimated orthogonally (in a sense discussed in Section 3). A simulation study is carried out in Section 5 to assess the behaviour of the new robust tests in finite samples, followed by an illustration on clinical data in Section 6. These methods are made available through the `p1med` package in R.

## 2 The G-estimator for mediation

Our objective is to derive direct and indirect effect estimators which are asymptotically linear and hence CAN in the sense that they asymptotically follow a normal distributions centred at the true value, with variances of order  $n^{-1}$ . Specifically, we shall say that an estimator  $\hat{\psi}$  of some true parameter  $\psi$ , based on observed data  $\{W_i\}_{i=1}^n$  is asymptotically linear if  $\hat{\psi} - \psi = E_n\{\varphi(W, \psi, \eta)\} + o_p(n^{-1/2})$ , where  $E_n(\cdot) = n^{-1} \sum_{i=1}^n (\cdot)_i$  denotes expectation with respect to the empirical distribution, and where  $\eta$  is a nuisance parameter vector and  $\varphi(W, \psi, \eta)$  is a unique ‘influence function’ with  $E\{\varphi(W, \psi, \eta)\} = 0$  and  $E\{\varphi(W, \psi, \eta)\varphi(W, \psi, \eta)^T\} < \infty$  in the positive definite sense. Here  $o_p$  denotes stochastic order notation so that  $A_n = o_p(r_n^{-1})$  means that  $A_n r_n \xrightarrow{p} 0$  where  $\xrightarrow{p}$  denotes convergence in probability. Later  $\xrightarrow{d}$  will be used to denote convergence in distribution.

We will consider the target parameter,  $\beta = (\beta_1, \beta_2, \beta_3)$ , and begin by introducing a model for the exposure given by  $E(X|Z) = h(Z, \gamma_x)$  where  $h$  is a known differentiable function parametrized by the nuisance parameter vector  $\gamma_x$ . The complete model nuisance parameter is therefore  $\gamma = (\gamma_x, \gamma_m, \gamma_y)$ . Throughout it is assumed that  $\gamma$  is estimated using a CAN

estimator  $\hat{\gamma} = \hat{\gamma}(\hat{\beta})$  which is consistent in the sense that the probability limit of each component of  $\gamma$  is equal to the true value under correct specification. This is defined more explicitly by assumptions A1 to A3 below.

The G-estimation moment conditions are based on the covariance of model residuals and are given by the vector  $U(\beta, \gamma)$  with components,

$$U_1(\beta, \gamma) = \{X - h(Z, \gamma_x)\}\{M - \beta_1 X - f(Z, \gamma_m)\} \quad (5)$$

$$U_2(\beta, \gamma) = \{M - \beta_1 X - f(Z, \gamma_m)\}\{Y - \beta_2 M - \beta_3 X - g(Z, \gamma_y)\} \quad (6)$$

$$U_3(\beta, \gamma) = \{X - h(Z, \gamma_x)\}\{Y - \beta_2 M - \beta_3 X - g(Z, \gamma_y)\} \quad (7)$$

When all models are correctly specified, these moment conditions have zero expectation evaluated at the true parameter values,  $E\{U(\beta^0, \gamma^0)\} = 0$ . The G-estimator for  $\beta$ , denoted by  $\hat{\beta}$  is the value which sets the sample average of moment conditions to zero, that is it solves the system of three equations given by

$$E_n[U\{\hat{\beta}, \hat{\gamma}(\hat{\beta})\}] = 0 \quad (8)$$

To examine the behaviour of this estimator under model misspecification, we introduce notation for the probability limit of this G-estimator,  $\beta^* = \text{plim } \hat{\beta}$ , with associated nuisance parameter estimator limit,  $\gamma^* = \text{plim } \hat{\gamma}(\beta^*)$ . The probability limit of the G-estimator is the solution to

$$E\{U(\beta^*, \gamma^*)\} = 0$$

We additionally assume that  $\beta^*$  is unique. This assumption can be partly justified by the linearity of the G-moment conditions, which implies that  $\beta^*$  is unique when  $\gamma^*$  is treated as a constant. It is sufficient, therefore, to assume that there is no pathological way by which the dependence of  $\gamma^*$  on  $\beta^*$  might introduce extra solutions.

In fact, we shall assume that each component of the nuisance parameter estimator is consistent when the associated part of the model is correctly specified. Denoting the correct target parameter value by superscript 0, this assumption may be written as

A1. If  $E(X|Z)$  is correctly specified then  $\text{plim } \hat{\gamma}_x(\beta) = \gamma_x^0$  for all  $\beta$

A2. If  $E(M|X, Z)$  is correctly specified then  $\text{plim } \hat{\gamma}_m(\beta) = \gamma_m^0$  for all  $\beta$  such that  $\beta_1 = \beta_1^0$

A3. If  $E(Y|M, X, Z)$  is correctly specified then  $\text{plim } \hat{\gamma}_y(\beta) = \gamma_y^0$  for all  $\beta$  such that  $(\beta_2, \beta_3) = (\beta_2^0, \beta_3^0)$

Under these assumptions, one can derive the conditions for which the G-estimator of  $(\beta_1, \beta_2)$  and  $(\beta_2, \beta_3)$  are consistent. These are given in Lemmas 1 and 2 respectively. For completeness, the union of these two cases, under which  $(\beta_1, \beta_2, \beta_3)$  is consistent, is given by Lemma 3

**Lemma 1 (Consistency of the G-estimator of  $(\beta_1, \beta_2)$ )** *Provided the models for  $M$  and  $Y$  are partially linear in  $X$  and  $M$  respectively, such that*

$$\begin{aligned} E(M - \beta_1^0 X|X, Z) &= E(M - \beta_1^0 X|Z) \\ E(Y - \beta_2^0 M|M, X, Z) &= E(Y - \beta_2^0 M|X, Z) \end{aligned}$$

and either

(i) *The model for  $E(M|X, Z)$  is correctly specified*

(ii) *The models for  $E(Y|M, X, Z)$  and  $E(X|Z)$  are both correctly specified*

then  $\beta^* = (\beta_1^0, \beta_2^0, \beta_3^*)$ , hence the G-estimator is consistent for  $(\beta_1, \beta_2)$ . See Supplementary material for proof.

**Lemma 2 (Consistency of the G-estimator of  $(\beta_2, \beta_3)$ )** *Provided the model for  $Y$  is partially linear in  $(M, X)$  such that*

$$E(Y - \beta_2^0 M - \beta_3^0 X | M, X, Z) = E(Y - \beta_2^0 M - \beta_3^0 X | Z)$$

and either

(i) *The model for  $E(Y|M, X, Z)$  is correctly specified*

(ii) *The models for  $E(X|Z)$  and  $E(M|X, Z)$  are both correctly specified*

then  $\beta^* = (\beta_1^*, \beta_2^0, \beta_3^0)$ , hence the G-estimator is consistent for  $(\beta_2, \beta_3)$ . See Supplementary material for proof.

**Lemma 3 (Robustness of G moment conditions)** *Provided that the models for  $M$  and  $Y$  are partially linear in  $X$  and  $(M, X)$  respectively, such that*

$$\begin{aligned} E(M - \beta_1^0 X | X, Z) &= E(M - \beta_1^0 X | Z) \\ E(Y - \beta_2^0 M - \beta_3^0 X | M, X, Z) &= E(Y - \beta_2^0 M - \beta_3^0 X | Z) \end{aligned}$$

and any pair of

(i) *The model for  $E(X|Z)$*

(ii) *The model for  $E(M|X, Z)$*

(iii) *The model for  $E(Y|M, X, Z)$*

are correctly specified, then  $\beta^* = (\beta_1^0, \beta_2^0, \beta_3^0)$ , hence the G-estimator is consistent for the full target parameter. For proof observe that the conditions in Lemmas 1 and 2 are satisfied.

Theorem 1 describes the conditions under which the G-estimator will be asymptotically linear.

**Theorem 1 (Asymptotically linearity of  $\hat{\beta}$ )** *Letting  $\beta^*$  denote the probability limit of the G-estimator, as set out in Lemmas 1 to 3, and assuming the nuisance parameter estimator,  $\hat{\gamma}(\beta)$  is CAN and obeys assumptions A1 to A3 such that*

$$\hat{\gamma} - \gamma^* = E_n\{\phi(\beta^*, \gamma^*)\} + o_p\left(n^{-1/2}\right) \quad (9)$$

Then, subject to regularity conditions, the estimator  $\hat{\beta}$  is CAN

$$\hat{\beta} - \beta^* = E_n\{\varphi(\beta^*, \gamma^*)\} + o_p\left(n^{-1/2}\right)$$

with influence function  $\varphi(\cdot)$  given by

$$\varphi(\beta^*, \gamma^*) = E\left\{-\frac{\partial U(\beta^*, \gamma^*)}{\partial \beta}\right\}^{-1} \left( U(\beta^*, \gamma^*) + E\left\{\frac{\partial U(\beta^*, \gamma^*)}{\partial \gamma}\right\} \phi(\beta^*, \gamma^*) \right) \quad (10)$$

See Supplementary material for proof.

The G-estimator of the NIDE is then given by the product  $\hat{\beta}_1 \hat{\beta}_2$  with influence function  $\omega(\cdot)$  given by

$$\omega(\beta^*, \gamma^*) = \beta_1^0 \varphi_2(\beta^*, \gamma^*) + \beta_2^0 \varphi_1(\beta^*, \gamma^*) \quad (11)$$

where  $\beta^* = (\beta_1^0, \beta_2^0, \beta_3^*)$  and for  $j = 1, 2, 3$ ,  $\varphi_j(\beta^*, \gamma^*)$  is the  $j$ th component of the influence function in 10. Derivation of this influence function can be found in the Supplementary material.

Similarly, the G-estimator of the NDE is  $\hat{\beta}_3$  with influence function  $\varphi_3(\beta^*, \gamma^*)$  where  $\beta^* = (\beta_1^*, \beta_2^0, \beta_3^0)$  and with consistency guaranteed under the conditions in Lemma 2.

In the case where  $(\beta_1^0, \beta_2^0) = (0, 0)$  then the influence function,  $\omega(\beta^*, \gamma^*)$  in (11) is exactly zero. In this case the NIDE estimator  $\hat{\beta}_1\hat{\beta}_2$  is asymptotically linear and CAN in the sense that  $n^{1/2}\hat{\beta}_1\hat{\beta}_2$  asymptotically follows a normal distribution with zero variance. Multiplying  $\hat{\beta}_1\hat{\beta}_2$  by higher powers of  $n$  yields more interesting behaviour. When all models are correctly specified,  $n\hat{\beta}_1\hat{\beta}_2$  asymptotically follows a ‘product normal’ distribution (the distribution of two mean zero normal variables with known variances) (Aroian, 1947).

### 3 Nuisance parameter estimation

Theoretical results show that the choice of nuisance parameter estimators does not impact the asymptotic variance of double robust estimators when both working models are correctly specified (Tsiatis, 2006). Similarly, in our case, it is straightforward to show that, when the models for  $X, M$  and  $Y$  are correctly specified then

$$E \left\{ \frac{\partial U(\beta^*, \gamma^*)}{\partial \gamma} \right\} = 0 \quad (12)$$

This property is sometimes referred to as orthogonality, with the intuition that the G-moment conditions are locally insensitive to nuisance parameters when all models are correctly specified. Orthogonal estimators are particularly useful for the construction of score tests since it follows that

$$\begin{aligned} n^{1/2} E_n \{ U(\beta^*, \hat{\gamma}(\beta^*)) \} &= n^{1/2} E_n \{ U(\beta^*, \gamma^*) \} + o_p(1) \\ E_n \{ U(\beta^*, \hat{\gamma}(\beta^*)) U(\beta^*, \hat{\gamma}(\beta^*))^T \} &= E_n \{ U(\beta^*, \gamma^*) U(\beta^*, \gamma^*)^T \} + o_p(1) \end{aligned}$$

However, when (12) is not satisfied then the influence function in (10) depends on the nuisance influence function  $\phi(\beta^*, \gamma^*)$ . Equation (10) therefore represents a class of estimators, indexed by the choice of nuisance parameter estimation method. Choosing the nuisance parameter estimator under misspecification is non-trivial as it may greatly affect the asymptotic variance of the estimator. Various proposals have been suggested for conventional double robust estimators which aim to minimize either the variance under misspecification (Rotnitzky and Vansteelandt, 2014) or the bias when both models are misspecified (Vermeulen and Vansteelandt, 2015).

An alternative method of constructing an estimator which is pseudo-orthogonal is by careful choice of  $\phi(\beta^*, \gamma^*)$  so that (12) is approximately satisfied. The bias-reduction strategy of Vermeulen and Vansteelandt (2015); Avagyan and Vansteelandt (2017) is an example of this.

To implement the bias-reduction strategy for our G-estimator, we must first augment the G-moment functions ((5)–(7)), such that each estimating equation has a unique nuisance parameter. In practice, this means, for example, that different estimators of  $\gamma_x$  may be used in (5) and (6), which are both consistent when  $E(X|Z)$  is correctly specified, with the same being true of  $\gamma_m$  and  $\gamma_y$ . The augmented moment functions are given by

$$U_1(\beta, \gamma) = \left\{ X - h \left( Z, \gamma_x^{(1)} \right) \right\} \left\{ M - \beta_1 X - f \left( Z, \gamma_m^{(1)} \right) \right\} \quad (13)$$

$$U_2(\beta, \gamma) = \left\{ M - \beta_1 X - f \left( Z, \gamma_m^{(2)} \right) \right\} \left\{ Y - \beta_2 M - \beta_3 X - g \left( Z, \gamma_y^{(1)} \right) \right\} \quad (14)$$

$$U_2(\beta, \gamma) = \left\{ X - h \left( Z, \gamma_x^{(2)} \right) \right\} \left\{ Y - \beta_2 M - \beta_3 X - g \left( Z, \gamma_y^{(2)} \right) \right\} \quad (15)$$

with full nuisance parameter,  $\gamma = (\gamma_x^{(1)}, \gamma_x^{(2)}, \gamma_m^{(1)}, \gamma_m^{(2)}, \gamma_y^{(1)}, \gamma_y^{(2)})$ . The bias-reduced nuisance parameter estimator is that which solves

$$E_n \left\{ \frac{\partial U(\hat{\beta}, \hat{\gamma})}{\partial \gamma} \right\} = 0$$

This estimator is asymptotically linear and obeys the consistency assumptions A1 to A3. For identifiability, we require models where  $\text{Dim}(\gamma_x) = \text{Dim}(\gamma_m) = \text{Dim}(\gamma_y)$ . Such restrictions are not uncommon e.g. (Rotnitzky et al., 2012) and may be satisfied by enlarging the working models.

## 4 Hypothesis testing

We now consider tests of the null hypothesis,  $H_\alpha : (\alpha - 1)\beta_1\beta_2 + \alpha\beta_3 = 0$ , with  $\alpha \in [0, 1]$  known. We begin by constructing a score test, for general  $\alpha$ , based on the G-moment conditions. Provided the nuisance parameters are orthogonally estimated, the score test is robust to certain model misspecification. Also, in the specific case where  $\alpha = 0$  and the true parameter takes the value  $(\beta_1^0, \beta_2^0) = (0, 0)$ , the score test is conservative, in the sense that the Type I error rate is below the nominal test size.

The score test is compared with a Wald test for the special cases of the no-mediation hypothesis ( $\alpha = 0$ ) and the no-direct-effect hypothesis ( $\alpha = 1$ ). These tests are again constructed using the G-moment conditions, and inherit the robustness properties of the G-estimator, without requiring orthogonal nuisance parameter estimation.

Sobel (1982) proposed a Wald test of the no-mediation hypothesis, based on the OLS moment conditions. Our Wald test for the no-mediation hypothesis is similar enough to Sobel's work that we shall refer to it as the Robust Sobel test (or Robust Wald test).

### 4.1 The Score Test

The score test is based on the observation that, provided  $E\{U(\beta^*, \gamma^*)\} = 0$ , the classical central limit theorem implies

$$\begin{aligned} n^{1/2} E_n\{U(\beta^*, \gamma^*)\} &\xrightarrow{d} \mathcal{N}(0, E\{U(\beta^*, \gamma^*)U(\beta^*, \gamma^*)^T\}) \\ n E_n\{U(\beta^*, \gamma^*)\}^T E\{U(\beta^*, \gamma^*)U(\beta^*, \gamma^*)^T\}^{-1} E_n\{U(\beta^*, \gamma^*)\} &\xrightarrow{d} \chi_3^2 \end{aligned} \quad (16)$$

The left hand side is similar in form to a GMM estimator, based on the objective function

$$D(\beta, \gamma, I) = E_n\{U(\beta, \gamma)\}^T I^{-1} E_n\{U(\beta, \gamma)\} \geq 0 \quad \forall(\beta, \gamma)$$

where  $I$  is a positive semi-definite 3 by 3 matrix. The GMM estimator of  $\beta$  is the minimizer  $\arg \min_\beta D(\beta, \hat{\gamma}(\beta), I)$ . In our case the GMM estimator is said to be exactly specified since  $\text{Dim}(\beta) = \text{Dim}(U(\beta, \gamma))$ . In this exactly specified setting, minimisation of the GMM objective function is equivalent to solving (8) and the estimator is independent of the choice of weighting matrix,  $I$ .

The minimisation of  $D(\beta, \hat{\gamma}(\beta), I)$  over a constrained parameter space, however, may be exploited for hypothesis testing, using results by Newey and West (1987) for the GMM Two-Step estimator, later extended by Dufour et al. (2017) to the GMM-Continuous Updating Estimator (CUE), both discussed below.

Work by Hansen (1982) in the over-specified setting, (i.e. when  $\text{Dim}(\beta) < \text{Dim}\{U(\beta, \gamma)\}$ ), showed that the optimal GMM estimator is constructed using weights proportional to the variance matrix,  $I \propto E\{U(\beta^*, \gamma^*)U(\beta^*, \gamma^*)^T\}$ . This is optimal in the sense that the asymptotic

covariance matrix of the resulting estimator is as small as possible (in the positive definite sense) among the class of GMM estimators.

This optimal choice also lends itself to hypothesis testing, as suggested by (16). In this work we consider minimisation of the objective function,  $D$ , under two proposals. The first (Two-Step) proposal first estimates nuisance parameters and the variance matrix  $E\{U(\beta^*, \gamma^*)U(\beta^*, \gamma^*)^T\}$ . Then, using these initial estimates, constrained estimates of  $\beta$  are obtained by a subsequent minimisation of the GMM estimator. The second, (CUE) proposal allows the estimates of nuisance parameters and the variance matrix  $I$  to be updated continuously. Writing,

$$\hat{I}(\beta, \gamma) = n^{-1}E_n\{U(\beta, \gamma)U(\beta, \gamma)^T\}$$

then the proposed Two-Step  $S_\alpha$  and CUE  $\tilde{S}_\alpha$  objective functions are respectively given by

$$S(\beta) = D\left(\beta, \hat{\gamma}(\hat{\beta}), \hat{I}(\hat{\beta}, \hat{\gamma}(\hat{\beta}))\right) \quad (17)$$

$$\tilde{S}(\beta) = D\left(\beta, \hat{\gamma}(\beta), \hat{I}(\beta, \hat{\gamma}(\beta))\right) \quad (18)$$

Where  $\hat{\beta}$  is the unconstrained G-estimate of  $\beta$ . Defining the null parameter space as  $B_\alpha = \{\beta | (\alpha - 1)\beta_1\beta_2 + \alpha\beta_3 = 0\}$ , the Two-step and CUE score type test statistics may be written as

$$S_\alpha = \min_{\beta \in B_\alpha} S(\beta)$$

$$\tilde{S}_\alpha = \min_{\beta \in B_\alpha} \tilde{S}(\beta)$$

In practice, computation of the Two-Step score statistic may be achieved by using the method of Lagrange Multipliers to construct estimating equations for the constrained minimisation problem. These may then be solved by a Newton-Raphson scheme. Similarly, computation of the CUE score statistic can be achieved using Lagrange Multipliers to construct estimating equations for  $\beta$ , however the Newton-Raphson procedure should additionally include the nuisance parameter estimating equations. When the bias-reduced nuisance estimation strategy is used, computation by Newton-Raphson requires that  $f(\cdot)$ ,  $g(\cdot)$  and  $h(\cdot)$  are twice continuously differentiable.

To derive the asymptotic distributions of  $S_\alpha$  and  $\tilde{S}_\alpha$ , we consider the general problem of minimising over some hypothesis set  $B_\psi = \{\beta | \psi(\beta) = 0\}$ , where  $\psi(\cdot)$  is a differentiable function. For example, when  $\psi(\beta) = \beta - \beta^*$  then  $B_\psi$  represents a single point, and the asymptotic distribution in (16) is recovered. Theorem 2 gives the general result, which follows from extending results by Newey and West (1987) and Dufour et al. (2017) to account for nuisance parameter estimation.

**Theorem 2 (Constrained GMM)** *Consider a null hypothesis  $H_0 : \psi(\beta^*) = 0$ , where  $\psi$  has dimension  $r \in \{1, 2, 3\}$  and is continuously differentiable in some non-empty, open neighbourhood,  $N$ , of the true limiting value  $\beta^*$ . Provided that for all  $\beta \in N$*

$$\text{Rank} \left( \frac{\partial \psi(\beta)}{\partial \beta} \right) = r \quad (19)$$

*and  $E\{U(\beta^*, \gamma^*)\} = 0$  with  $\hat{\gamma}$  estimated orthogonally, then*

$$\min_{\beta \in B_\psi} S(\beta) \xrightarrow{d} \chi_r^2$$

*where  $B_\psi = \{\beta | \psi(\beta) = 0\}$ . Furthermore, provided also that the nuisance parameter influence function  $\phi(\beta, \gamma)$  is twice continuously differentiable in  $\beta$  then*

$$\min_{\beta \in B_\psi} \tilde{S}(\beta) \xrightarrow{d} \chi_r^2$$

*Proof follows from Newey and West (1987) and Dufour et al. (2017).*



Applying Theorem 2 to the target hypothesis,  $H_\alpha$ , we see that the rank condition in (19) is not necessarily satisfied for the no-mediation hypothesis ( $\alpha = 0$ ). Letting  $\psi_\alpha(\beta) = (\alpha - 1)\beta_1\beta_2 + \alpha\beta_3$  then

$$\text{Rank} \left( \frac{\partial \psi_\alpha(\beta)}{\partial \beta} \right) = \text{Rank} \begin{pmatrix} (\alpha - 1)\beta_2 \\ (\alpha - 1)\beta_1 \\ \alpha \end{pmatrix} = \begin{cases} 0 & \text{for } \alpha = \beta_1 = \beta_2 = 0 \\ 1 & \text{otherwise} \end{cases}$$

Therefore for  $\alpha \neq 0$  one may apply the result in Theorem 2 directly to construct a test which rejects  $H_\alpha$  when  $S_\alpha > c$  for some critical value,  $c$ . This test size has size  $1 - F_{\chi_1^2}(c)$  where  $F_{\chi_1^2}(x)$  is the distribution function of a  $\chi_1^2$  variable.

One can show that this test is also a valid test of the no-mediation hypothesis,  $H_0$ . To do this, we define the null parameter space  $B_0 = \{\beta | \beta_1\beta_2 = 0\}$  and let  $B_j = \{\beta | \beta_j = 0\}$  for  $j = 1, 2$  so that  $B_0 = B_1 \cup B_2$ . Hence

$$S_0 = \min \left\{ \min_{\beta \in B_1} S(\beta), \min_{\beta \in B_2} S(\beta) \right\}$$

$$\tilde{S}_0 = \min \left\{ \min_{\beta \in B_1} \tilde{S}(\beta), \min_{\beta \in B_2} \tilde{S}(\beta) \right\}$$

Under  $H_0$  we know that either  $\beta_1 = 0$  or  $\beta_2 = 0$  and since the constraint function  $\psi_j(\beta) = \beta_j$  does satisfy the rank condition in (19), one can show, that under  $H_0$ ,

$$\sup_{\beta^* \in B_0} P_{\beta^*}(S_0 > x) \rightarrow 1 - F_{\chi_1^2}(x) \quad (20)$$

$$\sup_{\beta^* \in B_0} P_{\beta^*}(\tilde{S}_0 > x) \rightarrow 1 - F_{\chi_1^2}(x) \quad (21)$$

where  $P_{\beta^*}$  denotes the probability measure with a true limiting parameter value of  $\beta^*$  and  $\rightarrow$  denotes convergence as  $n$  tends to infinity. This result is justified in the Supplementary material.

## 4.2 Wald Tests

Using the asymptotic linearity of  $\hat{\beta}_1\hat{\beta}_2$  in (11) and under the conditions of Lemma 1, one obtains the CAN distribution of  $\hat{\beta}_1\hat{\beta}_2$ . Estimating the variance of  $n^{1/2}\hat{\beta}_1\hat{\beta}_2$  by  $E_n\{\omega^2(\hat{\beta}, \hat{\gamma}(\hat{\beta}))\}$  one arrives at the Wald test statistic for the no-mediation hypothesis:  $\beta_1\beta_2 = 0$

$$W = \frac{n^{1/2}(\hat{\beta}_1\hat{\beta}_2 - \beta_1\beta_2)}{E_n\{\omega^2(\hat{\beta}, \hat{\gamma}(\hat{\beta}))\}} \xrightarrow{d} \mathcal{N}(0, E\{\omega^2(\beta_0, \gamma^*)\})$$

$$= \frac{n\hat{\beta}_1^2\hat{\beta}_2^2}{E_n\{\omega^2(\hat{\beta}, \hat{\gamma}(\hat{\beta}))\}} = \frac{T_1T_2}{T_1 + T_2 + 2\rho\sqrt{T_1T_2}} = \frac{\hat{\beta}_1^2\hat{\beta}_2^2}{\hat{\beta}_1^2\hat{\sigma}_2^2 + \hat{\beta}_2^2\hat{\sigma}_1^2 + 2\hat{\beta}_1\hat{\beta}_2\Delta}$$

where for  $j = 1, 2, 3$ , the squared t-statistic is represented by  $T_j = \hat{\beta}_j^2/\hat{\sigma}_j^2$ , and  $\rho = \Delta/\hat{\sigma}_1\hat{\sigma}_2$  with  $\hat{\sigma}_j^2$  and  $\Delta$  given by  $n^{-1}E_n\{\varphi_j^2(\hat{\beta}, \hat{\gamma})\}$  and  $n^{-1}E_n\{\varphi_1(\hat{\beta}, \hat{\gamma})\varphi_2(\hat{\beta}, \hat{\gamma})\}$  respectively.

The distribution of  $W$  is problematic since at the true parameter value  $(\beta_1^0, \beta_2^0) = (0, 0)$ , we have  $\text{var}(n^{1/2}\hat{\beta}_1\hat{\beta}_2) \rightarrow 0$  and  $n \rightarrow \infty$ . A characterisation of Wald-type statistics for testing polynomial constraints with singular points is given by Dufour et al. (2013). For the constraint  $\beta_1\beta_2 = 0$ , Glonek (1993) demonstrated that

$$W \xrightarrow{d} \begin{cases} \frac{1}{4}\chi_1^2 & \text{for } \beta_1 = \beta_2 = 0 \\ \chi_1^2 & \text{otherwise} \end{cases}$$

This result suggests that one may reject the no-mediation hypothesis when the Wald statistic exceeds some critical value,  $c$ , chosen with reference to the  $\chi_1^2$  distribution. Such a test will have

size  $1 - F_{\chi_1^2}(c)$  when the null is satisfied but one of  $\beta_1$  or  $\beta_2$  is non-zero, and will be conservative when  $\beta_1 = \beta_2 = 0$ . The fact that  $W$  behaves differently at a singular point is known to greatly restrict the power of the Wald test to detect small indirect effects in finite samples.

Construction of a Wald based test for the NDE is fairly trivial. Under the conditions of Lemma 2, the squared t-statistic,  $T_3 \xrightarrow{d} \chi_1^2$  when  $\beta_3^0 = 0$ .

### 4.3 Comparison of methods

Revisiting the classical tests for the no-mediation hypothesis, as given in (3) and (4) we see that  $0 \leq W^{(OLS)} \leq LR^{(OLS)}$  with equality as  $T_j^{(OLS)}$  approaches infinity for either  $j = 1$  or  $j = 2$ , which occurs in the asymptotic limit when  $\beta_j \neq 0$ . Furthermore, a hypothesis test which rejects when  $LR^{(OLS)} > c$ , for some critical value  $c$ , has a test size given by:  $\lim_{n \rightarrow \infty} \sup_{\{\beta | \beta_1 \beta_2 = 0\}} P(LR^{(OLS)} > c)$ .

Since  $LR^{(OLS)}$  is monotonically increasing in  $T_j^{(OLS)}$ , the supremum above is achieved when  $T_j^{(OLS)}$  approaches infinity for either  $j = 1$  or  $j = 2$  (in the asymptotic limit when the true parameter value,  $\beta_j \neq 0$ ). In this limit, however,  $LR^{(OLS)} = W^{(OLS)}$ , and since  $W^{(OLS)}$  is also monotonically increasing in  $T_j$  we find that for all  $c$ ,

$$\lim_{n \rightarrow \infty} \sup_{\{\beta | \beta_1 \beta_2 = 0\}} P(LR^{(OLS)} > c) = \lim_{n \rightarrow \infty} \sup_{\{\beta | \beta_1 \beta_2 = 0\}} P(W^{(OLS)} > c)$$

Therefore a test which rejects when  $W^{(OLS)} > c$  has equal size to that which rejects when  $LR^{(OLS)} > c$  and since  $W^{(OLS)} \leq LR^{(OLS)}$  then the  $LR^{(OLS)}$  based test is uniformly more powerful than the  $W^{(OLS)}$  test. van Garderen and van Giersbergen (2019) showed that no test for  $H_0$  may be constructed from  $T_1^{(OLS)}$  and  $T_2^{(OLS)}$  for which the Type I error rate does not depend on  $(\beta_1^0, \beta_2^0)$ .

We make this comparison between the two classical tests because it gives some intuition as to why the G-estimation score test, which is analogous to  $LR^{(OLS)}$ , might be more powerful than the G-estimation Wald test, analogous to  $W^{(OLS)}$ . The fact that  $LR^{(OLS)}$  is analogous to our proposed score test can be seen by rewriting  $LR^{(OLS)}$  as a minimisation over an objective function.

$$S^{(OLS)}(\beta) = \sum_{j=1}^2 \left( \frac{\hat{\beta}_j^{(OLS)} - \beta_j}{\hat{\sigma}_j^{(OLS)}} \right)^2 \quad (22)$$

$$LR^{(OLS)} = \min_{\{\beta | \beta_1 \beta_2 = 0\}} S^{(OLS)}(\beta)$$

This objective function resembles a sum of OLS squared t-statistics, where, under the null, the minimisation step necessarily removes the contribution of a single term from the sum. To demonstrate the analogy between  $S^{(OLS)}(\beta)$  and our G-estimation score objective functions in (17) and (18), we consider the case where the three models for  $E(X|Z)$ ,  $E(M|X, Z)$ , and  $E(Y|M, X, Z)$  are all correctly specified.

In this setting, the G-estimating equations are always orthogonal to the nuisance parameter estimates. For illustration we shall additionally assume that  $\text{var}(Y|M, X, Z) = \text{var}(Y|X, Z)$ , so that the true covariance matrix,  $I = E\{U(\beta^0, \gamma^0)U(\beta^0, \gamma^0)^T\}$  is diagonal. Under these assumptions, the squared t-statistics for the null hypothesis  $\beta_j^0 = 0$ , are given by

$$T_j = \frac{\hat{\beta}_j^2}{\hat{\sigma}_j^2} = \frac{nE_n\{\varphi_j(\hat{\beta}_{-j}, \hat{\gamma})\}^2}{E_n\{\varphi_j^2(\hat{\beta}, \hat{\gamma})\}} \text{ for } j = 1, 2, 3$$

which reduces to

$$T_j = \frac{nE_n\{U_j(\hat{\beta}_{-j}, \hat{\gamma})\}^2}{E_n\{U_j^2(\hat{\beta}, \hat{\gamma})\}} \text{ for } j = 1, 2$$

$$T_3 = \frac{nE_n\{U_3(\hat{\beta}_{-3}, \hat{\gamma})\}^2}{E_n\{U_3^2(\hat{\beta}, \hat{\gamma})\} + \hat{\beta}_1^2 E_n\{\widehat{\text{var}}(X|Z)\} E_n\{\widehat{\text{var}}(M|X, Z)\}^{-1} E_n\{U_2^2(\hat{\beta}, \hat{\gamma})\}}$$

where  $\hat{\beta}_{-j}$  denotes the G-estimate of  $\beta$  with  $j$ th parameter set to zero and  $\widehat{\text{var}}(\cdot)$  denotes conditional variance estimated using the parameter G-estimates. Note that the denominator of  $T_3$  contains an additional term due to the non-zero value of  $E\{\partial U_3(\beta^0, \gamma^0)/\partial \beta_2\}$ , which happens to be the only non-zero off-diagonal term of the matrix  $E\{\partial U(\beta^0, \gamma^0)/\partial \beta\}$ .

Since the covariance matrix,  $I$  is diagonal in this setting, the two-step and CUE objective functions may be written as

$$S(\beta) = \sum_{j=1}^3 \frac{nE_n\{U_j(\beta, \hat{\gamma})\}^2}{E_n\{U_j^2(\hat{\beta}, \hat{\gamma})\}}$$

$$\tilde{S}(\beta) = \sum_{j=1}^3 \frac{nE_n\{U_j(\beta, \hat{\gamma}(\beta))\}^2}{E_n\{U_j^2(\beta, \hat{\gamma}(\beta))\}}$$

Similar to (22), therefore, these score test objective functions resemble sums of squared t-statistics. Theorem 2 may consequently be given the interpretation that the minimisation procedure under the null ‘minimises out’ independent  $\chi_1^2$  terms from this score test objective function, leaving a sum of independent  $\chi_1^2$  terms equal in number to the dimensions of the constraint.

## 5 Simulation Study

### 5.1 Simulation Study for estimation

A simulation study was carried out to examine the bias and variance of the G-estimators of the NIDE and NDE in finite samples and under misspecification. The performance of the score and Wald tests was also compared for the no-mediation hypothesis ( $H_0$ ) and the no-direct effect hypothesis ( $H_1$ ), with the bias-reduction strategy used for nuisance parameter estimation. Datasets of size  $n$  were generated for different  $(\beta_1, \beta_2, \beta_3)$  values using the hierarchical data generating process

$$Z \sim \mathcal{N}(0, 1)$$

$$X \sim \text{Bernoulli}(\text{expit}(Z + s_x Z^2))$$

$$M \sim \mathcal{N}(\beta_1 X + Z + s_m Z^2, 1)$$

$$Y \sim \mathcal{N}(\beta_2 M + \beta_3 X + Z + s_y Z^2, 1)$$

with  $s_x, s_m, s_y \in \{0, 1\}$  indicating model misspecification and where  $\text{expit}$  is the inverse-logit function. Analysis was conducted under the assumed model

$$E(X|Z) = \text{expit}(\gamma_{x1}Z + \gamma_{x2})$$

$$E(M|X, Z) = \beta_1 X + \gamma_{m1}Z + \gamma_{m2}$$

$$E(Y|M, X, Z) = \beta_2 M + \beta_3 X + \gamma_{y1}Z + \gamma_{y2}$$

Hence the assumed models are correctly specified when the corresponding misspecification indicator is equal to zero.

To investigate the bias and variance of the G-estimator, two parameter vectors were simulated,  $\beta = (0, 0, 0)$  and  $\beta = (1, 1, 1)$  with sample sizes  $n = 100, 500, 1000$  and under various levels of misspecification. 1000 dataset replicates were generated for each simulation, and for each dataset  $\hat{\beta}$  was estimated and  $\text{var}(\hat{\beta})$  was estimated based on the influence function, and by bootstrap with 1000 resampling iterations. Monte Carlo estimates for the expectation of each quantity were obtained across the 1000 dataset replicates.

Monte Carlo estimates of the bias of the G-estimators for  $\beta$  and the NIDE show that the bias remains close to zero even in small samples and under misspecification. Plots of these biases are given in the supplementary material. Figures 1 and 2 show plots of the variances of the NDE and NIDE G-estimates, respectively, scaled by the sample size,  $n$ . For all plots we see good agreement between the bootstrap and asymptotic estimates of the variance, which tend to overlap well with the Monte Carlo variance of  $\hat{\beta}$ , which we take as the empirical true value.

In Fig.1, correctly specifying  $E(M|X, Z)$  and  $E(X|Z)$  does little to change the variance of the  $\hat{\beta}_3$  provided that  $E(Y|M, X, Z)$  is correct. In Fig.2 the variance is reduced by correctly specifying all working models. There is also a noticeable difference in the NIDE variance under the true parameter  $\beta = (0, 0, 0)$ . This is to be expected from the form of (11). Additional plots showing the variances of  $\beta_1$  and  $\beta_2$  can be found in the supplementary material.

## 5.2 Simulation Study for hypothesis testing

To investigate the hypothesis testing methods a greater number of resampled datasets ( $10^4$ ) was used, since the computationally intensive bootstrap variance estimation procedure did not need to be carried out. The proposed tests based on G-estimation are compared with classical (non-robust) methods. Replicate datasets were generated for  $n$  in the range 50 to 500 with various true values of  $\beta$  under various levels of misspecification. Figures 3 and 4 respectively show the proportion of datasets for which the tests of  $H_0$  and  $H_1$  were rejected at the 5% level (indicated by a grey line). When the null is satisfied this rejection proportion corresponds to the Type I error rate, and otherwise corresponds to the statistical power (as in the right most column of both figures).

In Fig.3 all testing methods fail to achieve the nominal size when the true parameter takes the value  $(\beta_1, \beta_2) = (0, 0)$ , as expected by theory. All tests, however, do achieve nominal size when one of  $\beta_1$  or  $\beta_2$  differs from zero, given the requisite misspecification conditions. Under correct specification of all working models, the G-estimation score tests display similar power to the classical  $LR$  test, dominating both the robust and classical Sobel tests, which also have similar power to each other. This supports the heuristic argument in Section 4.3.

Although these results suggest that the Two-step procedure has greater power over the CUE score test, the Two-step method appears to have an inflated Type I error rate in small samples, which converges more slowly to the nominal level. This may explain the power discrepancy. This behaviour is reflected also in Fig.4, where the Robust Wald test suffers from a slightly inflated Type I error rate in small samples. In Fig.4 the robust tests perform better than classical test when  $E(Y|M, X, Z)$  is misspecified, as in the central row.

## 6 Motivating example: the COPERS trial

We now illustrate our G-estimation procedure and hypothesis testing methods, by analysing data from the COPERS (COping with persistent Pain, Effectiveness Research in Self-management) trial. COPERS was a multi-centre, pragmatic, randomized control trial examining the effectiveness of a novel non pharmacological intervention on the management of chronic musculoskeletal pain. Participants in the intervention arm ( $n = 384$ ) were offered to participate in group therapy sessions, while those in the control arm ( $n = 300$ ) received usual care. The group therapy introduced cognitive behavioural approaches to promote self-efficacy in managing chronic




Figure 1: Simulated results for estimates of  $n\text{var}(\hat{\beta}_3)$  from  $10^3$  dataset replicates estimated using: asymptotic results based on the influence function (circles), bootstrap estimation with  $10^3$  resampled datasets (triangles) and the Monte Carlo variance of  $\hat{\beta}$  over the  $10^3$  dataset replicates (squares). Each row represents a different true sample size and  $\beta$  parameter, whilst each column gives the models which are correctly specified (those for which the misspecification indicator is equal to zero)




Figure 2: Simulated results for estimates of  $nvar(\hat{\beta}_1, \hat{\beta}_2)$  from  $10^3$  dataset replicates estimated using: asymptotic results based on the influence function (circles), bootstrap estimation with  $10^3$  resampled datasets (triangles) and the Monte Carlo variance of  $\hat{\beta}$  over the  $10^3$  dataset replicates (squares). Each row represents a different true sample size and  $\beta$  parameter, whilst each column gives the models which are correctly specified (those for which the misspecification indicator is equal to zero)




Figure 3: Simulated results of the proportion of the  $10^4$  datasets for which the no-mediation hypothesis ( $H_0$ ) is rejected at the 5% level testing using the CUE score test (solid), Two-step score test (dotdashed), Robust Sobel test (dashed), Classical Sobel test (twodashed), and the Classical LR test (dotted). Each column represents a different true  $\beta$  parameter, whilst each row gives the models which are correctly specified (those for which the misspecification indicator is equal to zero)




Figure 4: Simulated results of the proportion of the  $10^4$  datasets for which the no-direct effect hypothesis ( $H_1$ ) is rejected at the 5% level testing using the CUE score test (solid), Two-step score test (dotdashed), Robust Wald test (dashed), and the Classical Wald test (twodashed). Each column represents a different true  $\beta$  parameter, whilst each row gives the models which are correctly specified (those for which the misspecification indicator is equal to zero)



pain. Attendance at the 24 group therapy sessions was observed to vary between participants in the intervention arm, with the original investigators denoting ‘compliers’ ( $X = 1, n = 260$ ) as participants who attended at least 12 sessions, and ‘non-compliers’ as those who did not ( $X = 0, n = 53$ ).

The primary outcome,  $Y$ , was pain-related disability at 12 months, measured on the Chronic Pain Grade (CPG) disability sub-scale. This is a continuous measure on a scale from 0 to 100, with higher scores indicating worse pain-related disability. The original analysis found no evidence that the COPERS intervention had an effect on improving pain-related disability at 12 months (The average treatment effect on the CPG scale was  $-1.0$ , with a 95% CI of  $-4.8$  to  $2.7$ ).

The COPERS researchers were interested in investigating whether the effect of therapy was mediated by feelings of self-coping with pain. To this effect, trial participants were also asked to fill out the Pain Self-Efficacy Questionnaire (PSEQ) at 12 weeks (shortly after receiving the intervention), which is intended to measure the participants confidence to live a normal life despite chronic pain. A mediation analysis was not performed due to the lack of overall treatment effect. In our analysis, we will treat the compliance indicator as the exposure of interest,  $X$ , and examine only the patients in the treatment arm, with the PSEQ score (a continuous measure) as the mediator of interest,  $M$ . Several patients ( $n = 71$ ) were excluded from analyses because of missing pre-treatment, mediation or outcome variables.

We use linear models for the conditional means of  $X$ ,  $M$ , and  $Y$ , given the same baseline covariates  $Z$  included in the original primary analysis models, namely: site of recruitment, age, gender, Hospital Anxiety and Depression Scale [HADS]), Health Education Impact Questionnaire for social integration subscale, and the corresponding outcome at baseline:

$$\begin{aligned} E(X|Z) &= \text{expit}(\gamma_x^T Z) \\ E(M|X, Z) &= \beta_1 X + \gamma_m^T Z \\ E(Y|M, X, Z) &= \beta_2 M + \beta_3 X + \gamma_y^T Z \end{aligned}$$

Table 1 gives mediated effect estimates obtained by G-estimation and OLS. Table 2 shows  $p$ -values for the no-mediation and no-indirect effect hypotheses, obtained using our Robust Sobel and score tests, along with the classical Sobel and LR methods.

Table 1: Estimated mediation effects for the COPERS trial using Ordinary Least Squares, and G-estimation

Parameter	OLS(sd)	G-estimate(sd)
$\beta_1$	6.80(1.46)	7.18(1.41)
$\beta_2$	-0.69(0.15)	-0.69(0.16)
NDE= $\beta_3$	3.79(3.82)	4.07(3.73)
NIDE= $\beta_1\beta_2$	-4.69(1.41)	-4.94(1.51)

This analysis suggests that attending at least half of the sessions had the effect of reducing pain-related disability, with almost all of the effect mediated by the participant’s confidence to live a normal life despite chronic pain. Indeed, on the basis of this analysis, there is not sufficient evidence to reject the hypothesis that there is no-direct effect at all, which is in agreement with the results of the original analysis. Given the possibility of strong mediated effects, researchers’ interest in cognitive behavioural therapy for chronic pain may want to design add-on interventions that can potentially also change self-coping perceptions.

## 7 Discussion

The main contribution of the current paper is a practical and robust method for carrying out inference of mediated effects in settings where partial linearity of mediator and outcome

Table 2: Hypothesis testing results on the COPERS dataset for the null hypotheses  $H_\alpha$  ( $H_0$  is the no-mediation hypothesis and  $H_1$  is the no-direct-effect hypothesis)

Null $\alpha$ value	Test	P-Value
0	Sobel	$9.20 \times 10^{-4}$
0	LR	$3.11 \times 10^{-6}$
0	Robust Sobel	$1.07 \times 10^{-3}$
0	Robust score (CUE)	$5.33 \times 10^{-5}$
0	Robust score (2-step)	$3.28 \times 10^{-7}$
1	Classical Wald	0.322
1	Robust Wald	0.275
1	Robust score (CUE)	0.275
1	Robust score (2-step)	0.277

conditional expectations can be assumed and confounder models are low dimensional. We recommend estimation of the NIDE and NDE by G-estimation, in settings where partial linearity may be assumed, but a mediator density function may be difficult to estimate, as required by the methods of Tchetgen Tchetgen and Shpitser (2012). Compared with OLS, the G-estimators are consistent for the NIDE and NDE, under misspecification of mediator and outcome models respectively. The variance of these estimators may be estimated by bootstrap or using asymptotic results with both giving similar results. We also make available the methods in the R package `plmed`, which calculates the NDE and NIDE by G-estimation with variances estimated using asymptotic results by default.

In terms of hypothesis testing we recommend the robust CUE score test over Two-step robust score methods, due to its faster convergence of the Type I error rate to the nominal size in small samples, and improved power to detect small NIDEs over Wald based testing methods, as demonstrated in simulation studies.

One limitation of these methods, aside from the partially linear model assumption is the requirement that the confounder models are low dimensional. G-estimators of the average treatment effect, however, have been shown to permit inference under high-dimensional confounding. In particular, by exploiting the orthogonality of the G-estimator when the exposure and outcome models are both correct, one can show that valid inference is still obtained when cross-validated data-adaptive methods (e.g. lasso or machine learning) are used to estimate the confounder models (such as  $f(\cdot)$ ,  $g(\cdot)$ , and  $h(\cdot)$  in the current paper), with the assumption that such methods will converge to the true model and an appropriate sample splitting scheme is applied (Chernozhukov et al., 2016).

Other work by Dukes and Vansteelandt (2019) obtains valid inference of the average treatment effect using G-estimators when confounder models are fitted using the bias-reduction strategy with a lasso  $l_1$  penalty on the nuisance parameter. This work does not require sample splitting nor does it require that both the exposure and outcome confounder models converge to the truth. Indeed their methods are valid even when the number of confounding variables is allowed to grow with sample size, provided certain sparsity assumptions on the nuisance parameter are satisfied. Similar results could be derived for the G-estimator of mediated effects in the current paper, with care taken to address the requirement of Theorem 2 that the nuisance parameter estimator be asymptotically linear.

## References

Aroian, L. A. (1947). The Probability Function of the Product of Two Normally Distributed Variables. *The Annals of Mathematical Statistics* 18(2), 265–271.

- Avagyan, V. and S. Vansteelandt (2017). Honest data-adaptive inference for the average treatment effect under model misspecification using penalised bias-reduced double-robust estimation. (2015), 1–33.
- Baron, R. M. and D. A. Kenny (1986). The moderator mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *Journal of Personality and Social Psychology* 51(6), 1173–1182.
- Chernozhukov, V., D. Chetverikov, M. Demirer, E. Duflo, C. Hansen, W. Newey, and J. Robins (2016). Double/Debiased Machine Learning for Treatment and Causal Parameters.
- Dufour, J.-M., E. Renault, and V. Zinde-Walsh (2013). Wald tests when restrictions are locally singular. (1).
- Dufour, J. M., A. Trognon, and P. Tuvaandorj (2017). Invariant tests based on M-estimators, estimating functions, and the generalized method of moments. *Econometric Reviews* 36(1-3), 182–204.
- Dukes, O. and S. Vansteelandt (2019). Uniformly valid confidence intervals for conditional treatment effects in misspecified high-dimensional models. pp. 1–40.
- Fritz, M. S., A. B. Taylor, and D. P. MacKinnon (2012). Explanation of Two Anomalous Results in Statistical Mediation Analysis. *Multivariate Behavioral Research* 47(1), 61–87.
- Giersbergen, N. P. A. V. (2014). Inference about the Indirect Effect : a Likelihood Approach.
- Glonek, G. F. V. (1993). On the Behaviour of Wald Statistics for the Disjunction of Two Regular Hypotheses. *Journal of the Royal Statistical Society. Series B (Methodological)* 55(3), 749–755.
- Hansen, L. P. (1982, jul). Large Sample Properties of Generalized Method of Moments Estimators. *Econometrica* 50(4), 1029.
- Kang, J. D. and J. L. Schafer (2007). Demystifying double robustness: A comparison of alternative strategies for estimating a population mean from incomplete data. *Statistical Science* 22(4), 523–539.
- Kenny, D. A. and C. M. Judd (2014). Power Anomalies in Testing Mediation. *Psychological Science* 25(2), 334–339.
- MacKinnon, D. P., C. M. Lockwood, J. M. Hoffman, S. G. West, and V. Sheets (2002). A comparison of methods to test mediation and other intervening variable effects. *Psychological Methods* 7(1), 83–104.
- Newey, W. K. and K. D. West (1987, oct). Hypothesis Testing with Efficient Method of Moments Estimation. *International Economic Review* 28(3), 777.
- Pearl, J. (2001). Direct and Indirect Effects. 411(20).
- Robins, J. M. and S. Greenland (1992). Identifiability and Exchangeability for Direct and Indirect Effects Linked references are available on JSTOR for this article :. *Epidemiology* 3(2), 143–155.
- Rotnitzky, A., Q. Lei, M. Sued, and J. M. Robins (2012). Improved double-robust estimation in missing data and causal inference models. *Biometrika* 99(2), 439–456.

- Rotnitzky, A. and S. Vansteelandt (2014). Double-robust methods. In G. Molenberghs, G. Fitzmaurice, M. Kenward, A. Tsiatis, and G. Verbeke (Eds.), *Handbook of missing data methodology*, Chapter 9, pp. 185–212. New York: CRC Press.
- Sobel, M. E. (1982). Asymptotic Confidence Intervals for Indirect Effects in Structural Equation Models. *Sociological Methodology* 13(1982), 290.
- Tchetgen Tchetgen, E. J. and I. Shpitser (2012). Semiparametric theory for causal mediation analysis: Efficiency bounds, multiple robustness and sensitivity analysis. *Annals of Statistics* 40(3), 1816–1845.
- Tsiatis, A. A. (2006). *Semiparametric Theory and Missing Data*. Springer Series in Statistics. New York, NY: Springer New York.
- van Garderen, K. J. and N. van Giersbergen (2019). Almost Similar Tests for Mediation Effects Hypotheses with Singularities Kees Jan van Garderen and Noud van Giersbergen.
- Vermeulen, K. and S. Vansteelandt (2015, jul). Bias-Reduced Doubly Robust Estimation. *Journal of the American Statistical Association* 110(511), 1024–1036.
- Wang, K. (2018). Understanding Power Anomalies in Mediation Analysis. *Psychometrika* 83(2), 387–406.

## A Proofs

### A.1 Proof of Lemma 1

In Step 1 the following expressions for each component of  $E\{U(\beta^*, \gamma^*)\}$  are derived. In Step 2 we consider the behaviour of these expressions in each of the two misspecification cases.

$$E\{U_1(\beta^*, \gamma^*)\} = E \left[ \{E(X|Z) - h(Z, \gamma_x^*)\} \{E(M - \beta_1^0 X|Z) - f(Z, \gamma_m^*)\} \right] \quad (23)$$

$$E\{U_2(\beta^*, \gamma^*)\} = E \left[ \{E(M - \beta_1^0 X|Z) - f(Z, \gamma_m^*)\} \{E(Y - \beta_2^0 M|X, Z) - \beta_3^* X - g(Z, \gamma_y^*)\} \right] \quad (24)$$

$$E\{U_3(\beta^*, \gamma^*)\} = E \left[ \{E(X|Z) - h(Z, \gamma_x^*)\} \{E(Y - \beta_2^0 M|X, Z) - \beta_3^* X - g(Z, \gamma_y^*)\} \right] \quad (25)$$

Step 1: By the partial linearity of  $M$  we obtain

$$\begin{aligned} E\{U_1(\beta^*, \gamma^*)|X, Z\} &= \left\{ X - h(Z, \gamma_x^*) \right\} \left\{ E(M - \beta_1^0 X|X, Z) - f(Z, \gamma_m^*) \right\} \\ &= \left\{ X - h(Z, \gamma_x^*) \right\} \left\{ E(M - \beta_1^0 X|Z) - f(Z, \gamma_m^*) \right\} \\ E\{U_1(\beta^*, \gamma^*)|Z\} &= \left\{ E(X|Z) - h(Z, \gamma_x^*) \right\} \left\{ E(M - \beta_1^0 X|Z) - f(Z, \gamma_m^*) \right\} \end{aligned}$$

For the second component, the partial linearity of  $M$  and  $Y$  gives

$$\begin{aligned} E\{U_2(\beta^*, \gamma^*)|M, X, Z\} &= \left\{ M - \beta_1^0 X - f(Z, \gamma_m^*) \right\} \left\{ E(Y - \beta_2^0 M|M, X, Z) - \beta_3^* X - g(Z, \gamma_y^*) \right\} \\ E\{U_2(\beta^*, \gamma^*)|X, Z\} &= \left\{ E(M - \beta_1^0 X|X, Z) - f(Z, \gamma_m^*) \right\} \left\{ E(Y - \beta_2^0 M|X, Z) - \beta_3^* X - g(Z, \gamma_y^*) \right\} \\ &= \left\{ E(M - \beta_1^0 X|Z) - f(Z, \gamma_m^*) \right\} \left\{ E(Y - \beta_2^0 M|X, Z) - \beta_3^* X - g(Z, \gamma_y^*) \right\} \end{aligned}$$

Likewise,

$$\begin{aligned} E\{U_3(\beta^*, \gamma^*)|M, X, Z\} &= \left\{ X - h(Z, \gamma_x^*) \right\} \left\{ E(Y - \beta_2^0 M|M, X, Z) - \beta_3^* X - g(Z, \gamma_y^*) \right\} \\ E\{U_3(\beta^*, \gamma^*)|X, Z\} &= \left\{ X - h(Z, \gamma_x^*) \right\} \left\{ E(Y - \beta_2^0 M|X, Z) - \beta_3^* X - g(Z, \gamma_y^*) \right\} \\ E\{U_3(\beta^*, \gamma^*)|Z\} &= \left\{ E(X|Z) - h(Z, \gamma_x^*) \right\} \left\{ E(Y - \beta_2^0 M|X, Z) - \beta_3^* X - g(Z, \gamma_y^*) \right\} \end{aligned}$$

Step 2: We shall consider the cases (i) and (ii) separately. In case (i) the conditions for assumption A2 are met, hence  $E(M - \beta_1^0 X|Z) = f(Z, \gamma_m^*)$  and so (23) and (24) are exactly zero. The proof for case (i) is completed by letting  $\beta_3^*$  be the value which solves (24) equal to zero.

For case (ii) the conditions of A1 are met, hence  $E(X|Z) = h(Z, \gamma_x^*)$  so (23) and (25) are exactly zero. Since  $E(Y|M, X, Z)$  is correctly specified, there exists  $\beta_3^0$  such that  $E(Y - \beta_2^0 M - \beta_3^0 X|M, X, Z) = E(Y - \beta_2^0 M - \beta_3^0 X|Z) = g(Z, \gamma_y^*)$ , where the final equality arises since the conditions of A3 are satisfied. Therefore (24) is exactly zero, which completes the proof for case (ii).

### A.2 Proof of Lemma 2

In Step 1 the following expressions for the second and third components of  $E\{U(\beta^*, \gamma^*)\}$  are derived. In Step 2 we consider the behaviour of these expressions in each of the two misspecification cases.

$$E\{U_2(\beta^*, \gamma^*)\} = E \left[ \left\{ E(M - \beta_1^* X|X, Z) - f(Z, \gamma_m^*) \right\} \left\{ E(Y - \beta_2^0 M - \beta_3^0 X|Z) - g(Z, \gamma_y^*) \right\} \right] \quad (26)$$

$$E\{U_3(\beta^*, \gamma^*)\} = E \left[ \left\{ E(X|Z) - h(Z, \gamma_x^*) \right\} \left\{ E(Y - \beta_2^0 M - \beta_3^0 X|Z) - g(Z, \gamma_y^*) \right\} \right] \quad (27)$$

Step 1. By the partial linearity of  $Y$ , we find

$$\begin{aligned} E\{U_2(\beta^*, \gamma^*)|M, X, Z\} &= \left\{M - \beta_1^* X - f(Z, \gamma_m^*)\right\} \left\{E(Y - \beta_2^0 M - \beta_3^0 X|M, X, Z) - g(Z, \gamma_y^*)\right\} \\ &= \left\{M - \beta_1^* X - f(Z, \gamma_m^*)\right\} \left\{E(Y - \beta_2^0 M - \beta_3^0 X|Z) - g(Z, \gamma_y^*)\right\} \\ E\{U_2(\beta^*, \gamma^*)|X, Z\} &= \left\{E(M - \beta_1^* X|X, Z) - f(Z, \gamma_m^*)\right\} \left\{E(Y - \beta_2^0 M - \beta_3^0 X|Z) - g(Z, \gamma_y^*)\right\} \end{aligned}$$

Likewise,

$$\begin{aligned} E\{U_3(\beta^*, \gamma^*)|M, X, Z\} &= \left\{X - h(Z, \gamma_x^*)\right\} \left\{E(Y - \beta_2^0 M - \beta_3^0 X|M, X, Z) - g(Z, \gamma_y^*)\right\} \\ &= \left\{X - h(Z, \gamma_x^*)\right\} \left\{E(Y - \beta_2^0 M - \beta_3^0 X|Z) - g(Z, \gamma_y^*)\right\} \\ E\{U_3(\beta^*, \gamma^*)|Z\} &= \left\{E(X|Z) - h(Z, \gamma_x^*)\right\} \left\{E(Y - \beta_2^0 M - \beta_3^0 X|Z) - g(Z, \gamma_y^*)\right\} \end{aligned}$$

Step 2. We shall consider the cases (i) and (ii) separately.

In case (i) the conditions for assumption A3 are met, hence  $E(Y - \beta_2^0 M - \beta_3^0 X|Z) = g(Z, \gamma_y^*)$  so the term inside 26 and 27 are exactly zero. Letting  $\beta_1^*$  be the solution to

$$E\{U_1(\beta^*, \gamma^*)\} = E \left[ \left\{X - h(Z, \gamma_x^*)\right\} \left\{M - \beta_1^* X - f(Z, \gamma_m^*)\right\} \right] = 0$$

completes the proof for case (i). For case (ii) the conditions of A1 are met, so  $E(X|Z) = h(Z, \gamma_x^*)$  and so the term inside (27) is zero. Since  $E(M|X, Z)$  is correctly specified there exists  $\beta_1^0$  such that  $E(M - \beta_1^0 X|X, Z) = E(M - \beta_1^0 X|Z)$ . Letting  $\beta_1^* = \beta_1^0$  then (26) becomes

$$E\{U_2(\beta^*, \gamma^*)\} = E \left[ \left\{E(M - \beta_1^0 X|Z) - f(Z, \gamma_m^*)\right\} \left\{E(Y - \beta_2^0 M - \beta_3^0 X|Z) - g(Z, \gamma_y^*)\right\} \right]$$

which is exactly zero since  $E(M - \beta_1^0 X|Z) = f(Z, \gamma_m^*)$  under A2. The same is true of the expression in (23), completing the proof for case (ii).

### A.3 Proof of Proposition 1

Consider the Taylor Expansion

$$E_n\{U(\hat{\beta}, \hat{\gamma})\} = E_n\{U(\beta^*, \gamma^*)\} + E_n \left\{ \frac{\partial U(\beta^*, \gamma^*)}{\partial \beta} \right\} (\hat{\beta} - \beta^*) + E_n \left\{ \frac{\partial U(\beta^*, \gamma^*)}{\partial \gamma} \right\} (\hat{\gamma} - \gamma^*) + o_p(n^{-1/2})$$

Since  $E_n\{U(\hat{\beta}, \hat{\gamma})\} = 0$  then

$$\hat{\beta} - \beta^* = E_n \left\{ -\frac{\partial U(\beta^*, \gamma^*)}{\partial \beta} \right\}^{-1} \left[ E_n\{U(\beta^*, \gamma^*)\} + E_n \left\{ \frac{\partial U(\beta^*, \gamma^*)}{\partial \gamma} \right\} (\hat{\gamma} - \gamma^*) \right] + o_p(n^{-1/2})$$

Using the estimator in (9) and rearranging gives

$$\hat{\beta} - \beta^* = E_n \left( E_n \left\{ -\frac{\partial U(\beta^*, \gamma^*)}{\partial \beta} \right\}^{-1} \left[ U(\beta^*, \gamma^*) + E_n \left\{ \frac{\partial U(\beta^*, \gamma^*)}{\partial \gamma} \right\} \phi(\beta^*, \gamma^*) \right] \right) + o_p(n^{-1/2})$$

Applying the weak law of large numbers to the partial derivative terms gives the form of the influence function  $\varphi(\cdot)$  in (10). We must further show that  $E\{\varphi(\beta^*, \gamma^*)\} = 0$

$$E\{\varphi(\beta^*, \gamma^*)\} = E \left\{ -\frac{\partial U(\beta^*, \gamma^*)}{\partial \beta} \right\}^{-1} \left[ E\{U(\beta^*, \gamma^*)\} + E \left\{ \frac{\partial U(\beta^*, \gamma^*)}{\partial \gamma} \right\} E\{\phi(\beta^*, \gamma^*)\} \right]$$

Since  $\phi(\cdot)$  is an influence function,  $E\{\phi(\beta^*, \gamma^*)\} = 0$ . Therefore provided  $E\{U(\beta^*, \gamma^*)\} = 0$  then  $E\{\varphi(\beta^*, \gamma^*)\} = 0$  as required.

#### A.4 Derivation of Equation (11)

From Proposition 1 we have

$$\begin{aligned}\hat{\beta}_1 &= \beta_1 + E_n\{\varphi_1(\beta^*, \gamma^*)\} + o_p\left(n^{-1/2}\right) \\ \hat{\beta}_2 &= \beta_2 + E_n\{\varphi_2(\beta^*, \gamma^*)\} + o_p\left(n^{-1/2}\right)\end{aligned}$$

Therefore, letting  $A = E_n\{\varphi_1(\beta^*, \gamma^*)\}$  and  $B = E_n\{\varphi_2(\beta^*, \gamma^*)\}$ , we obtain

$$\hat{\beta}_1\hat{\beta}_2 - \beta_1\beta_2 = E_n\{\omega(\beta^*, \gamma^*)\} + AB + o_p\left(n^{-1/2}\right)$$

The desired result follows provided that  $AB = o_p\left(n^{-1/2}\right)$ . Using Markov's inequality,

$$P(|n^{1/2}AB| \geq \epsilon) = P(n(AB)^2 \geq \epsilon^2) \leq \frac{nE\{(AB)^2\}}{\epsilon^2}$$

Examining the expectation term, we find a sum over four indices

$$E\{(AB)^2\} = n^{-4} \sum_{i=1}^n \sum_{j=1}^n \sum_{k=1}^n \sum_{l=1}^n E\{\varphi_1^{(i)}(\beta^*, \gamma^*)\varphi_1^{(j)}(\beta^*, \gamma^*)\varphi_2^{(k)}(\beta^*, \gamma^*)\varphi_2^{(l)}(\beta^*, \gamma^*)\}$$

where the superscript  $(i)$  denotes that the influence function is evaluated on the  $i$ th observation. Since the observations are iid and the influence function has mean zero, the terms of this quadruple sum are only non-zero when their indices are paired, i.e when  $(i = j \text{ and } k = l)$  or  $(i = k \text{ and } j = l)$  or  $(i = l \text{ and } j = k)$ . The number of non-zero terms in the sum is therefore of order  $n^2$ , and hence

$$P(|n^{1/2}AB| \geq \epsilon) \leq \mathcal{O}(n^{-1})$$

where  $\mathcal{O}$  denotes conventional big-O notation, i.e. for sufficiently large  $n$  there exists some constant  $k$  such that  $|\mathcal{O}(n^{-1})| \leq kn^{-1}$

#### A.5 Proof of Equations (20) and (21)

We will prove (20) with the result for (21) proceeding in a similar fashion.

Consider Theorem 2 under the null hypothesis  $\psi^{(0)}(\beta^*) = \beta_1\beta_2 = 0$ . With the null parameter space given by  $B_0 = \{\beta|\psi^{(0)}(\beta) = 0\}$ , with

$$\text{Rank}\left(\frac{\partial\psi^{(0)}(\beta)}{\partial\beta}\right) = \begin{cases} 0 & \text{for } \beta_1 = \beta_2 = 0 \\ 1 & \text{otherwise} \end{cases}$$

One may decompose the supremum in (21) as

$$\sup_{\beta^* \in B_0} P_{\beta^*}(S_0 > x) = \max\left\{\sup_{\beta^* \in B_0 \setminus A} P_{\beta^*}(S_0 > x), \sup_{\beta^* \in A} P_{\beta^*}(S_0 > x)\right\} \quad (28)$$

where  $A = \{\beta|\beta_1 = \beta_2 = 0\}$ . For the first term in the max bracket above, the rank condition of Theorem 2 holds, so for all  $\beta^* \in B_0 \setminus A$

$$P_{\beta^*}(S_0 > x) \rightarrow 1 - F_{\chi_1^2}(x)$$

Considering the second term, one may decompose the test statistic as

$$P_{\beta^*}(S_0 > x) = P_{\beta^*}(S_1 > x, S_2 > x) \leq P_{\beta^*}(S_1 > x)$$

where (for  $j = 1, 2$ )  $S_j = \min_{\beta \in B_j} S(\beta)$  and  $B_j = \{\beta|\beta_j = 0\}$ . By Theorem 2, for all  $\beta^*$  in  $A$ ,

$$P_{\beta^*}(S_1 > x) \rightarrow 1 - F_{\chi_1^2}(x)$$

Hence  $P_{\beta^*}(S_0 > x)$  is asymptotically bounded from above by  $1 - F_{\chi_1^2}(x)$  for all  $\beta^*$  in  $B_0$ , so (20) holds.




Figure 5: Simulated results to investigate the biases of  $\hat{\beta}_1$  (purple),  $\hat{\beta}_2$  (blue), the NDE= $\hat{\beta}_3$  (green) and the NIDE= $\hat{\beta}_1\hat{\beta}_2$  (red). The estimated bias from  $10^4$  dataset replicates is plotted on the x-axis with error bars giving a 95% confidence interval of the Monte Carlo estimate. Plots are arranged in a grid where each row represents a different sample size and true target parameter value, and the header of each row lists the correctly specified models (those for which the misspecification indicator is equal to zero). Where the requirements on model misspecification are not met, the bias of the estimator is not shown.

## B Additional Plots






Figure 6: Simulated results for estimates of  $n\text{var}(\hat{\beta}_1)$  from  $10^3$  dataset replicates estimated using: asymptotic results based on the influence function (circles), bootstrap estimation with  $10^3$  resampled datasets (triangles) and the Monte Carlo variance of  $\hat{\beta}$  over the  $10^3$  dataset replicates (squares). Each row represents a different true sample size and  $\beta$  parameter, whilst each column gives the models which are correctly specified (those for which the misspecification indicator is equal to zero)




Figure 7: Simulated results for estimates of  $n\text{var}(\hat{\beta}_2)$  from  $10^3$  dataset replicates estimated using: asymptotic results based on the influence function (circles), bootstrap estimation with  $10^3$  resampled datasets (triangles) and the Monte Carlo variance of  $\hat{\beta}$  over the  $10^3$  dataset replicates (squares). Each row represents a different true sample size and  $\beta$  parameter, whilst each column gives the models which are correctly specified (those for which the misspecification indicator is equal to zero)