

# Linear regression and its inference on noisy network-linked data\*

Can M. Le

Department of Statistics, University of California, Davis

Tianxi Li

Department of Statistics, University of Virginia

February 1, 2022

## Abstract

Linear regression on a set of observations linked by a network has been an essential tool in modeling the relationship between response and covariates with additional network data. Despite its wide range of applications in many areas, such as in social sciences and health-related research, the problem has not been well-studied in statistics so far. Previous methods either lack inference tools or rely on restrictive assumptions on social effects and usually assume that networks are observed without errors, which is unrealistic in many problems. This paper proposes a linear regression model with nonparametric network effects. The model does not assume that the relational data or network structure is exactly observed; thus, the method can be provably robust to a certain network perturbation level. A set of asymptotic inference results is established under a general requirement of the network observational errors, and the robustness of this method is studied in the specific setting when the errors come from random network models. We discover a phase-transition phenomenon of the inference validity concerning the network density when no prior knowledge of the network model is available while also showing a significant improvement achieved by knowing the network model. As a by-product of this analysis, we derive a rate-optimal concentration bound for random subspace projection that may be of independent interest. Extensive simulation studies are conducted to verify these theoretical results and demonstrate the advantage of the proposed method over existing work in terms of accuracy and computational efficiency under different data-generating models. The method is then applied to middle school students' network data to study the effectiveness of educational workshops in reducing school conflicts.

## 1 Introduction

Nowadays, networks appear frequently in many areas, including social sciences, transportation, and biology. In most cases, networks are used to represent relationships or interactions between units of a complex (social, physical, or biological) system, so analyzing network data may render

---

\*Both authors contributed equally to this work.

crucial insights into the dynamics and/or interaction mechanism of the system. One particular, yet commonly encountered, situation is when a group of units is observed connected by a network and a set of attributes for each of these units is available. Such data sets are sometimes called network-linked data [Li et al., 2019, 2020c] or multiview network data [Gao et al., 2019]. Network-linked data are widely available in almost all fields involving network analysis about social effects [Michell and West, 1996, Pearson and West, 2003], collaborations [Ji and Jin, 2016, Su et al., 2019], and causal experiments [Basse and Airoidi, 2018a,b]. In such a data set, rich information is available from the perspective of both the individual attributes and the network, and the challenge is to find proper statistical methods to incorporate both. Consider the situation where, for each node  $i$  of the network, we observe  $(x_i, y_i)$ , in which  $x_i \in \mathbb{R}^p$  is a vector of covariates while  $y_i \in \mathbb{R}$  is a scalar response. In particular, we aim for a regression model of  $y_i$  against  $x_i$  that also takes the network information into account. Such a model arises naturally in any problem when a prediction model or inference of a specific attribute is of interest.

Although the systematic study of regression on network-linked data has only recently begun to attract interest in statistics [Li et al., 2019, Zhu et al., 2017, Su et al., 2019], it has been studied in econometrics by many authors, who mostly focused on multiple networks [Manski, 1993, Lee, 2007, Bramoullé et al., 2009] or longitudinal data [Jackson and Rogers, 2007, Manresa, 2013]. Social effects are typically observed in the form that connected units share similar behaviors or properties. Such similarity or correlation may be due to either *homophily*, where social connections are established because of similarity, or *contagion*, where individuals become similar through the influence of their social ties. In general, one cannot distinguish homophily from contagion in a single snapshot of observational data [Shalizi and Thomas, 2011], as in our setting. Therefore, the two directions of causality will not be distinguished, and this type of generic similarity between connected nodes will be called “network cohesion”, as in Li et al. [2019, 2020c]. A significant class of models for such network regression problems is the class of autoregressive models, which is based on ideas in spatial statistics. The spatial autoregressive models have been widely used in econometrics, such as in Manski [1993], Lee [2007], Bramoullé et al. [2009], Hsieh and Lee [2016], Zhu et al. [2017], to name a few. A common form for such a model is

$$y_i = \gamma \left( \frac{1}{n_i} \sum_{i' \sim i} y_{i'} \right) + X\beta + \frac{1}{n_i} \left( \sum_{i' \sim i} x_{i'} \right)^T \eta + \epsilon,$$

where  $y_i$  is the response,  $x_i$  is the covariate vector and  $n_i$  is the number of neighbors of  $i$  (the notation  $i \sim i'$  indicates that  $i$  and  $i'$  are connected). In this model, the neighborhood averages of the response and covariates are used to model the social effect. The parameters  $\gamma$  and  $\eta$  are typically called “endogenous” and “exogenous” effects, respectively, while  $\beta$  is the standard regression slope. This class of models is also called the “linear in means” model or social interaction model (SIM). When multiple networks for different populations are available, the intercept can be treated as a group effect, called the “external effect” [Manski, 1993, Lee, 2007, Lee et al., 2010]. The SIM framework, though it has been a popular setup for regression on network-linked data, suffers from two crucial drawbacks. The first comes from its restrictive parametric form of the social effect. Assuming social effect in the form of autoregressive neighborhood average (or summation) is far from realistic, and such a stringent assumption significantly limits the usefulness of the framework. As shown in Li et al. [2019] and also in our empirical study, the restrictive assumption of social effect leads to poor prediction performance. The second limitation comes from the assumption that the network structure is precisely observed from the data. In practice, it is well known that most network data are subject to errors, due to missingness [Lakhina et al., 2003, Butts, 2003, Clauset and Moore, 2005], observational errors [Handcock and Gile, 2010, Rolland et al., 2014, Le and Levina, 2017, Khabbaziyan et al., 2017, Newman, 2018, Lunagómez et al., 2018], or data collection

method [Wu et al., 2018, Rohe et al., 2019]. If the imprecise network is used in SIM, the model is ill-defined and the inference would also be problematic, as shown by Chandrasekhar and Lewis [2011].

In this paper, a new regression model to fit the network-linked data is proposed that addresses both drawbacks mentioned above. The new model is based on a flexible network effect assumption and is robust to network observational errors. It also allows us to perform tests and construct confidence intervals for the model parameters. Our theoretical analysis provides the support for model estimation and inference and quantifies the magnitude of network observational errors under which the inference remains robust. Moreover, in the random network perturbation scenario, the tradeoff between the available information of the network structure and the level of robustness of the statistical inference is characterized. In its most difficult setting, when no prior knowledge about the network model is available, the result reveals a phase-transition phenomenon at the network average degree of  $\sqrt{n}$ , above which the inference is asymptotically correct, and below which the inference becomes invalid. The inference could remain valid for much sparser networks if more information about the network structure is available. For example, when the network model is known and an effective parametric estimation can be applied, the sparsity requirement can be relaxed to  $\log n$ . To the best of our knowledge, this paper is the first work that addresses the inference of the network-linked regression models and accounts for the network observational errors.

Related to the network-linked model proposed in this paper is the semi-parametric model called “regression with network cohesion” (RNC), introduced by Li et al. [2019]. In their model, the network effect is represented by individual parameters that are assumed to be “smooth” over the network, and a similar idea is used in a few other statistical estimation settings [Wang et al., 2016, Zhao and Shojaie, 2016, Fan and Guan, 2018]. Despite its flexibility of social effect assumption and excellent predictive performance, the RNC model lacks a valid inference framework and cannot be applied in many modern applications where statistical inference is needed [Ogburn, 2018, Su et al., 2019]. The result of Li et al. [2020c] indicates that the RNC estimator fails to guarantee valid inference under reasonable assumptions unless additional assumptions such sparsity are made [Zhao and Shojaie, 2016]. Moreover, little is known about the robustness of the RNC method to network observational errors, although preliminary results have been obtained in a particular scenario of network sparsification [Sadhanala et al., 2016, Li et al., 2019]. In contrast, our model does not assume the smoothness of network effects as in the RNC method. Instead, we use a general relational subspace to define the social effects. As can be seen later, the proposed model overcomes both of the two aforementioned limitations of RNC and is computationally more efficient.

Table 1 gives a high-level comparison between the proposed model and two popular benchmark regression models (discussed previously) in three aspects: the flexibility of modeling social effects, the availability of an inference method, and the provable robustness to the network perturbation. The model we introduced in this paper is the only one of the three which renders all of the desired properties.

Table 1: Comparison between the proposed method and two other popular methods for network-linked data.

| Models                | social effect flexibility | inference | network robustness |
|-----------------------|---------------------------|-----------|--------------------|
| SIM [Manski, 1993]    | ✗                         | ✓         | ✗                  |
| RNC [Li et al., 2019] | ✓                         | ✗         | ✗                  |
| Current method        | ✓                         | ✓         | ✓                  |

The rest of the paper is organized as follows. Section 2 introduces our model and the corresponding statistical inference algorithm. Section 3 and Section 4 are devoted to our main theoretical results; the generic theory for model estimation consistency and the asymptotic inference is given in Section 3; then, under the random network modeling framework, detailed discussions of the technical requirement are introduced in Section 4. Extensive simulation experiments are given in Section 5 to verify our theory and compare our method with a few benchmark methods mentioned above. Section 6 demonstrates the usefulness of the proposed method in analyzing the gender and racial effects on social activities among adolescents. Concluding remarks and future directions are discussed in Section 7. The proofs of all theoretical results and additional real data analysis are given in the supplementary material.

## 2 Network regression model

Throughout the paper,  $0_{k \times \ell}$  and  $I_k$  are used to denote the zero matrix of size  $k \times \ell$  and the identity matrix of size  $k \times k$ , respectively. The subscripts may be dropped when the dimensions are clear from the context. We use  $e_i$  to denote the vector whose  $i$ th coordinate is 1 and the remaining coordinates are zero; the dimension of  $e_i$  may vary according to the context. Given a matrix  $M \in \mathbb{R}^{k \times \ell}$  and  $1 \leq i \leq j \leq \ell$ ,  $M_{i:j} \in \mathbb{R}^{k \times (j-i+1)}$  is used to denote the matrix whose columns are those of  $M$  with indices  $i, i+1, \dots, j$ ; when  $i = j$ ,  $M_i$  is used instead of  $M_{i:i}$  for the simplicity of the notation. Throughout the paper,  $\|\cdot\|$  denotes the spectral norm, which is the largest singular value, for matrices and the Euclidean norm for vectors. The number of nodes in the network is denoted by  $n$ . We say that an event  $E$  occurs with high probability if  $\mathbb{P}(E) \geq 1 - n^{-c}$  for some constant  $c > 2$ . We use  $a_n = o(1)$  and  $b_n = O(1)$  to indicate that  $\lim_{n \rightarrow \infty} a_n = 0$  and  $b_n$  is a bounded sequence, respectively.

### 2.1 A semi-parametric regression model with network effects

The unobserved true network is captured by the relational matrix  $P \in \mathbb{R}^{n \times n}$ , where  $P_{ij}$  describes the strength of the relation between nodes  $i$  and  $j$ . For each node  $i$  of the network  $(x_i, y_i)$  is observed, where  $x_i \in \mathbb{R}^p$  is a vector of covariates while  $y_i \in \mathbb{R}$  is a scalar response. Denote by  $Y = (y_1, \dots, y_n)^T \in \mathbb{R}^n$  the vector of responses and by  $X = (x_1, \dots, x_n)^T \in \mathbb{R}^{n \times p}$  the design matrix, which sometimes has the first column as the all-one vector. Our goal is to construct a model that describes the dependence of  $Y$  on  $X$  and  $P$ .

The observed network is represented by an  $n \times n$  adjacency matrix  $A$ , where  $0 \leq A_{ij} \leq 1$  is the weight of the edge between  $i$  and  $j$ . In the special case of an unweighted network,  $A$  is a binary matrix and  $A_{ij} = 1$  if and only if node  $i$  and node  $j$  are connected. We view  $A$  as a perturbed version of  $P$  and focus on undirected networks, for which  $A$  and  $P$  are symmetric matrices. Moreover, we only consider the *fixed design* setting, to avoid unnecessary complication of jointly modeling covariates and relational data. That means  $X$  and  $P$  are always treated as fixed. The adjacency matrix  $A$  is also treated as fixed for the moment as the regression model and its generic inference framework are described. Later on, in only Section 4 where we study the robustness of our framework under deviation of  $A$  from  $P$ , we will assume  $A$  as a perturbation of  $P$  following random network models.

Intuitively, the structural assumption is that  $y_i$ 's tend to be similar for individuals having strong connections. This is called the ‘‘assortative mixing’’ or ‘‘network cohesion’’ property [Kolaczyk,

2009, Li et al., 2019]. To incorporate this intuition, consider the model

$$\mathbb{E}Y = X\zeta + \mu, \quad (1)$$

where  $\zeta \in \mathbb{R}^p$  is the vector of coefficients for the covariates  $X$  and  $\mu \in \mathbb{R}^n$  is the vector of individual effects reflecting the network cohesion property. This model was studied in Li et al. [2019]. Instead of modeling the network effect by using a specific auto-regressive dependence as in Manski [1993], (1) treats the network effect as a nonparametric component. As shown by Li et al. [2019] and the experiments later on in this paper, this approach is more flexible and gives significantly better predictive performance than the SIM framework of Manski [1993]. In Li et al. [2019],  $\mu$  is assumed to have small sum of squared differences  $\sum_{i,j:A_{ij}=1}(\mu_i - \mu_j)^2$ . In contrast, we rely on a different form of the network cohesion that proves to be more general and stable. Specifically, the cohesion requirement of  $\mu$  is formulated by assuming that

$$\mu \in S_K(P), \quad (2)$$

where  $S_K(P)$  is the subspace spanned by the  $K$  leading eigenvectors of  $P$ . The idea of using the eigenspace of  $P$  to encode the smoothness of a function over the network is related to the standard spectral embedding [Shi and Malik, 2000, Ng et al., 2001, Belkin and Niyogi, 2003, Tang et al., 2013], which maps the nodes of the network to a set of points in a Euclidean space so that the geometric relations between these points are similar to the topological relations of the nodes in the original network. Depending on specific problems it may be possible to replace  $P$  in (2) with other appropriate matrix-valued functions of  $P$  for which the inference framework remains valid. One such example is provided in Section 4.3. In summary, we assume the following mean structure:

**Definition 1.** The mean structure of the regression model with network effects is defined to be

$$\mathbb{E}Y \in \text{span}\{\text{col}(X), S_K(P)\}, \quad (3)$$

where  $\text{col}(X)$  is the column space of  $X$  and  $\text{span}$  denotes the linear subspace jointly spanned by  $\text{col}(X)$  and  $S_K(P)$ . This is equivalent to (1) with the cohesion property (2).

The advantage of our mean structure (1) lies in its generality. In particular, one distinction between our model and the more commonly assumed linear mean structure is that our model includes the situation when the two subspaces  $\text{col}(X)$  and  $S_K P$  have nontrivial intersection. For example, one of the covariates may be perfectly cohesive over the network, and this situation is allowed in our model. In this case, one could not uniquely determine  $X\zeta$  and  $\mu$ . However, such a tricky situation is unavoidable for the level of generality and robustness we want to achieve. As a matter of fact, this type of ambiguity is an instance of the general conclusion from Shalizi and Thomas [2011] that contagion, and different types of homophily effects from a single snapshot of observational data are not distinguishable. Nevertheless, the mean-structure, as well as an interpretable decomposition of the effects, can be uniquely identified with the following parameterization.

**Definition 2.** Let  $\mathcal{R} = \text{col}(X) \cap S_K(P)$  be the intersection of  $\text{col}(X)$  and  $S_K(P)$ . Model (3) can be reparametrized as

$$\mathbb{E}Y = X\beta + \xi + \alpha, \quad (4)$$

where

$$\xi \in \mathcal{R}, \quad X\beta \perp \mathcal{R}, \quad \alpha \in S_K(P), \quad \alpha \perp \mathcal{R}, \quad (5)$$

and  $\beta \in \mathbb{R}^p$ ,  $\xi, \alpha \in \mathbb{R}^n$ . We call it the *regression model with network effects*.

Although there are several ways to reparameterize the mean structure (3), the one in (5) is preferable because it combines the two sources of information (the covariates  $X$  and the relational information  $P$ ) in a natural way so that the interpretation of the corresponding parameters is straightforward. Specifically, we observe that

- When  $\beta = 0$ , we have  $\mathbb{E}Y \in S_K(P)$  so the model can be specified as a cohesive network effects without using the covariates  $X$ . Therefore,  $\beta$  represents the conditional covariate effects of  $X$  given  $P$ .
- When  $\alpha = 0$ , we have  $\mathbb{E}Y \in \text{col}(X)$  so the model can be specified by the covariates without using the relational information  $P$ . Therefore,  $\alpha$  represents the conditional network effect of  $P$  given  $X$ .

Thus,  $\beta$  and  $\alpha$  reflect the conditional effects similar to the covariate effects in the standard linear regression setting. In addition,  $\xi$  is the overlap of the two sources. The following result confirms the identifiability of our parameters.

**Proposition 1** (Parameter identifiability). The parameterization in Definition 2 is identifiable. That is, if there exist  $(\beta, \alpha, \xi)$  and  $(\beta', \alpha', \xi')$  satisfying (4) and (5) simultaneously then  $\beta = \beta'$ ,  $\alpha = \alpha'$ , and  $\xi = \xi'$ .

**Remark 1.** A seemingly easier way to combine the two sources of information is to treat the eigenvectors of  $P$  as another set of covariates and fit  $Y$  using both  $X$  and the eigenvectors. However, this approach has to assume  $\mathcal{R} = \{0\}$  for identifiability, which is a very strong assumption that we prefer to avoid. Note also that (5) does not require  $\alpha \perp X\beta$ , although this strong restriction would significantly simplify the fitting procedure and its analysis.

Finally, we assume that the noise in the regression model of Definition 2 is a multivariate Gaussian vector. Specifically,

$$Y = \mathbb{E}Y + \epsilon \text{ and } \epsilon \sim N(0, \sigma^2 I), \quad (6)$$

where  $I \in \mathbb{R}^{n \times n}$  is the identity matrix and  $\sigma^2$  is the variance of the noise. We leave the study of other noise distributions for future work.

**Remark 2.** As to be clear in our theory later, the reasonable interpretation of  $K$  in (2) is the index where the spectral space of  $P$  has a large eigen gap. The problem of estimating such a  $K$  has been extensively studied in network literature and many efficient methods are now available [Chatterjee, 2015, Chen and Lei, 2018, Le and Levina, 2015, Li et al., 2020b, Jin et al., 2020]. These methods all provide theoretical guarantees for the recovery of  $K$  with overwhelming probability and can be applied to our setting. Since the task of estimating  $K$  neither makes our problem conceptually more challenging nor brings more insight about it, for simplicity of presentation, we assume that  $K$  is known throughout the paper.

## 2.2 Statistical inference method

Our generic inference framework requires access to a certain approximation of  $P$ , denoted by  $\hat{P}$ . The specific  $\hat{P}$  is determined by the user according to the understanding of the data problem. For example, one may assume that the adjacency matrix  $A$  is a perturbed version of  $P$ . In this situation, without additional information, a natural option is to set  $\hat{P} = A$  and we refer to the corresponding estimation and inference procedure as the “model-free” version of our method, highlighting that no

specific model for the network structure is assumed. Section 4 discusses some other cases for which additional network model assumptions are considered and more accurate parametric estimates of  $P$  are available.

The identifiability condition (5) suggests a natural procedure for estimating parameters in model (4) via subspace projections. However, the fact that  $\alpha$  and  $X\beta$  need not be orthogonal complicates the estimation procedure and its analysis. To highlight the main idea, we first describe the population-level estimation, assuming that both  $P$  and  $\mathbb{E}Y$  are known.

Let  $Z \in \mathbb{R}^{n \times p}$  be a matrix whose columns form an orthonormal basis of the covariate subspace  $\text{col}(X)$ . Similarly, let  $W \in \mathbb{R}^{n \times K}$  be the matrix whose columns are eigenvectors of  $P$  that span the subspace  $S_K(P)$ . Define the singular value decomposition (SVD) of matrix  $Z^T W$  to be  $Z^T W = U \Sigma V^T$ . Here,  $U \in \mathbb{R}^{p \times p}$  and  $V \in \mathbb{R}^{K \times K}$  are orthonormal matrices of singular vectors while  $\Sigma \in \mathbb{R}^{p \times K}$  is the matrix with the following singular values on the main diagonal:

$$\sigma_1 = \sigma_2 = \dots = \sigma_r = 1 > \sigma_{r+1} \geq \dots \geq \sigma_{r+s} > 0 = \sigma_{r+s+1} = \dots = 0. \quad (7)$$

Thus,  $r$  is the dimension of  $\mathcal{R} = \text{col}(X) \cap S_K(P)$  and  $Z^T W$  has  $r + s$  non-zero singular values. Column vectors of the matrices

$$\tilde{Z} = ZU \quad \text{and} \quad \tilde{W} = WV \quad (8)$$

also form a basis of  $\text{col}(X)$  and  $S_K(P)$ , respectively. In particular, the first  $r$  column vectors of  $\tilde{Z}$  and  $\tilde{W}$  coincide and form a basis of  $\mathcal{R}$ . Moreover, the last  $p - r$  column vectors of  $\tilde{Z}$  form a basis of the subspace of  $\text{col}(X)$  that is perpendicular to  $\mathcal{R}$ ; similarly, the last  $K - r$  column vectors of  $\tilde{W}$  form a basis of the subspace of  $S_K(P)$  that is perpendicular to  $\mathcal{R}$ . The model assumption (5) then indicates

$$\xi \in \text{col}(\tilde{Z}_{1:r}) = \text{col}(\tilde{W}_{1:r}) = \mathcal{R}, \quad X\beta \in \text{col}(\tilde{Z}_{(r+1):p}) \perp \mathcal{R}, \quad \alpha \in \text{col}(\tilde{W}_{(r+1):K}) \perp \mathcal{R}. \quad (9)$$

Therefore,  $\xi$  can be recovered by  $\xi = \mathcal{P}_{\mathcal{R}} \mathbb{E}Y$ , where  $\mathcal{P}_{\mathcal{R}}$  is the orthogonal projection onto  $\mathcal{R}$  written as

$$\mathcal{P}_{\mathcal{R}} = \tilde{Z}_{1:r} \tilde{Z}_{1:r}^T = \tilde{W}_{1:r} \tilde{W}_{1:r}^T. \quad (10)$$

For the convenience of theoretical analysis later, we also introduce the projection coefficient  $\theta \in \mathbb{R}^p$

$$\theta = (X^T X)^{-1} X^T \xi.$$

Recall that  $\xi \in \mathcal{R} \subset \text{col}(X)$ , so we have  $\xi = X\theta$ , and we have the one-to-one correspondence between  $\theta$  and  $\xi$ . However, notice that  $\theta$  cannot be attributed to conditional effects of covariates, as explained in the previous section.

For estimating  $\alpha$  and  $\beta$ , we project  $\mathbb{E}Y$  on  $\text{col}(\tilde{Z}_{(r+1):p})$  and  $\text{col}(\tilde{W}_{(r+1):K})$ , respectively. Since these subspaces need not be orthogonal to each other (the principle angles between the two subspaces are the singular values  $\sigma_{r+1}, \dots, \sigma_{r+s}$ ), the corresponding projections are not necessarily orthogonal projections. Instead, they admit the following forms:

$$\mathcal{P}_{\mathcal{C}} = (\tilde{Z}_{(r+1):p}, 0_{n \times (K-r)})(M^T M)^{-1} M^T, \quad (11)$$

$$\mathcal{P}_{\mathcal{N}} = (0_{n \times (p-r)}, \tilde{W}_{(r+1):K})(M^T M)^{-1} M^T, \quad (12)$$

where  $M = (\tilde{Z}_{(r+1):p}, \tilde{W}_{(r+1):K})$ . Both  $\alpha$  and  $\beta$  can then be recovered by

$$\alpha = \mathcal{P}_{\mathcal{N}} \mathbb{E}Y \quad \text{and} \quad \beta = (X^T X)^{-1} X^T \mathcal{P}_{\mathcal{C}} \mathbb{E}Y. \quad (13)$$

Finally, to test the hypothesis  $H_0 : \alpha = 0$ , we can use  $\|\alpha\|^2$  as the statistic. We have  $\alpha = \sum_{i=1}^{K-r} \tilde{W}_{r+i} \gamma_i = \tilde{W}_{(r+1):K} \gamma$ , such that  $\|\alpha\| = \|\gamma\|$ . Although  $\gamma$  can not be uniquely determined because  $\tilde{W}_{(r+1):K}$  is only unique up to an orthogonal transformation, we can still uniquely identify its magnitude  $\|\gamma\|^2$  to perform a chi-squared test against the null hypothesis  $H_0 : \alpha = 0$ .

In practice, when  $P$  and  $\mathbb{E}Y$  are not observed, it is natural to replace them everywhere by the available approximations  $\hat{P}$  and  $Y$  in the procedure above. There is, however, one crucial issue that requires special attention: the plugin estimate  $\text{col}(X) \cap S_K(\hat{P})$  for  $\mathcal{R} = \text{col}(X) \cap S_K(P)$  is often a bad approximation of  $\mathcal{R}$ . This is because the intersection of subspaces is not robust with respect to small perturbations to the subspaces. For example, it is easy to see that in low dimensional settings, when  $\text{col}(X) \cap S_K(P)$  is nontrivial, a small perturbation of  $P$  can easily make  $\text{col}(X) \cap S_K(\hat{P})$  a null space. Therefore, rather than using  $\text{col}(X) \cap S_K(\hat{P})$  to approximate  $\mathcal{R}$ , the projections in (10), (11) and (12) are directly approximated using the eigenvectors of  $\hat{P}$  (this partially explains the detailed discussion of the population level estimation above). The whole estimation procedure is summarized in Algorithm 1. From here through Section 3, we will assume that the dimension  $r$  of  $\mathcal{R}$  is known for simplicity. This is because, so far,  $A$  has been treated as fixed while a discussion of selecting  $r$  naturally involves a detailed analysis of the perturbation mechanism from  $P$  to  $A$ . In Section 4 when the perturbation model for  $A$  is introduced, we provide a simple method to select  $r$  with a theoretical guarantee (see Corollary 3).

**Algorithm 1** (Spectral projection estimation). Given  $X, Y, \hat{P}, K$  and  $r$ .

1. Calculate an orthonormal basis of  $\text{col}(X)$  and forms  $Z \in \mathbb{R}^{n \times p}$ . Similarly, calculate  $K$  eigenvectors of  $\hat{P}$  and form  $\hat{W} \in \mathbb{R}^{n \times K}$ .

2. Calculate the singular value decomposition

$$Z^T \hat{W} = \hat{U} \hat{\Sigma} \hat{V}^T \quad (14)$$

and denote

$$\hat{Z} = Z \hat{U}, \quad \check{W} = \hat{W} \hat{V}. \quad (15)$$

3. Let  $\hat{M} = (\hat{Z}_{(r+1):p}, \check{W}_{(r+1):K})$ . Estimate  $\mathcal{P}_R, \mathcal{P}_C$  and  $\mathcal{P}_N$  by

$$\hat{\mathcal{P}}_R = \hat{Z}_{1:r} \hat{Z}_{1:r}^T, \quad (16)$$

$$\hat{\mathcal{P}}_C = (\hat{Z}_{(r+1):p}, 0_{n \times (K-r)}) (\hat{M}^T \hat{M})^{-1} \hat{M}^T, \quad (17)$$

$$\hat{\mathcal{P}}_N = (0_{n \times (p-r)}, \check{W}_{(r+1):K}) (\hat{M}^T \hat{M})^{-1} \hat{M}^T. \quad (18)$$

4. Estimate  $\xi, \beta$  and  $\alpha$  by

$$\hat{\xi} = \hat{\mathcal{P}}_R Y, \quad \hat{\beta} = (X^T X)^{-1} X^T \hat{\mathcal{P}}_C Y, \quad \hat{\alpha} = \hat{\mathcal{P}}_N Y. \quad (19)$$

And recover the projection coefficient of  $\xi$  to  $\text{col}(X)$  as  $\hat{\theta} = (X^T X)^{-1} X^T \hat{\xi}$ .

5. Let  $\hat{H} = \hat{\mathcal{P}}_R + \hat{\mathcal{P}}_C + \hat{\mathcal{P}}_N$ . Estimate the variance  $\sigma^2$  of  $\epsilon$  in (4) by

$$\hat{\sigma}^2 = \|Y - \hat{H}Y\|^2 / (n - p - K + r). \quad (20)$$

6. Estimate the covariance of  $\hat{\beta}$  by

$$\hat{\sigma}^2 (X^T X)^{-1} X^T \hat{\mathcal{P}}_C \hat{\mathcal{P}}_C^T X (X^T X)^{-1}. \quad (21)$$

The inference of  $\beta$  can be done according to Theorem 1.



7. To test the significance of  $\alpha$ , estimate  $\gamma$  by

$$\hat{\gamma} = \check{W}_{(r+1):K} \hat{\alpha} \quad (22)$$

and estimate the covariance matrix of  $\hat{\gamma}$  by  $\hat{\Sigma}_{\hat{\gamma}} = \hat{\sigma}^2 \check{W}_{(r+1):K} \hat{\mathcal{P}}_{\mathcal{N}} \hat{\mathcal{P}}_{\mathcal{N}}^T \check{W}_{(r+1):K}^T$ . Normalize  $\hat{\gamma}$  to obtain  $\hat{\gamma}_0 = \hat{\Sigma}_{\hat{\gamma}}^{-1/2} \hat{\gamma}$ . Use  $\|\hat{\gamma}_0\|^2$  for a chi-squared test (with  $K$  degrees of freedom) against the null hypothesis  $H_0 : \alpha = 0$  according to Theorem 2.

### 3 Generic inference theory of model parameters

For valid inference of parameters, we make the following standard assumptions about the data.

**Assumption A1** (Standardized scale). All columns of  $X$  satisfy  $\|X_i\| = \sqrt{n}$ . Moreover,

$$\|\mathbb{E}Y\| \leq C\sqrt{np}$$

for some constant  $C > 0$ .

**Assumption A2** (Weak dependence of  $X$ ). Denote  $\Theta = (X^T X/n)^{-1}$ . Assume that  $\Theta$  is well-conditioned, that is, there exists a constant  $\rho > 0$  such that

$$\rho \leq \lambda_{\min}(\Theta) \leq \lambda_{\max}(\Theta) \leq 1/\rho.$$

Due to the deviation of  $\hat{P}$  from the true signal  $P$ , the estimators would be biased. A common approach to ensure the valid inference is to control the bias levels of the parameter estimates [Van de Geer et al., 2014, Zhang and Zhang, 2014, Javanmard and Montanari, 2014]. In the current context, we need to guarantee that the biases are negligible compared to the standard deviations of the estimators. This requires  $\hat{P} - P$  to be controlled to some extent. In particular, the biases will depend on the magnitude of the perturbation of  $S_K(P)$  associated with  $\text{col}(X)$ , defined by

$$\tau_n = \|Z^T \hat{W} \hat{W}^T - Z^T W W^T\| = \|(\hat{W} \hat{W}^T - W W^T) Z\|. \quad (23)$$

As a warm-up, it is not difficult to get a bound on  $\|\hat{\Sigma} - \Sigma\|$  in terms of  $\tau_n$ . Indeed, from the previous discussion, the singular decompositions of  $Z^T W W^T$  and  $Z^T \hat{W} \hat{W}^T$  are

$$Z^T W W^T = U \Sigma \tilde{W}^T, \quad Z^T \hat{W} \hat{W}^T = \hat{U} \hat{\Sigma} \check{W}^T. \quad (24)$$

Therefore, by Weyl's inequality,

$$\|\hat{\Sigma} - \Sigma\| \leq \tau_n. \quad (25)$$

Since the estimates depend crucially on the singular value decomposition of  $Z^T W W^T$  and  $Z^T \hat{W} \hat{W}^T$ , (23), (24) and (25) play a central role in establishing the theoretical results. We now introduce our assumption on the error level that ensures the validity of the inference. It can be seen as a way to characterize the level of perturbation our framework could tolerate. This validity and theoretical insights about this assumption will be studied in more detail later (see Section 4). As we show there, the condition is mild in many commonly studied network settings.

**Assumption A3** (Small projection perturbation). Let  $Z$  and  $W$  be the matrices formed by the bases of  $\text{col}(X)$  and  $S_K(P)$ , respectively. Let  $Z^T W = U \Sigma V^T$  be the singular value decomposition with singular values specified in (7). Assume  $\hat{P}$  is a small projection perturbation of  $P$  with respect to

$X$ , by which we mean

$$\frac{\tau_n}{\min\{(1 - \sigma_{r+1})^3, \sigma_{r+s}^3\}} = o(1/\sqrt{np}).$$

The first property to be introduced is the estimation consistency of  $\hat{\sigma}^2$ , which serves as the critical building block for later inference.

**Proposition 2** (Consistency of variance estimation). Assume that **A1** holds and

$$40\tau_n \leq \min\{(1 - \sigma_{r+1})^2, \sigma_{r+s}^2\}. \quad (26)$$

Then with high probability, the estimate  $\hat{\sigma}^2$  defined by (20) satisfies

$$|\hat{\sigma}^2 - \sigma^2| \leq \frac{C\tau_n}{\min\{(1 - \sigma_{r+1})^3, \sigma_{r+s}^3\}} \cdot \frac{n(\sigma^2 + p)}{n - p - K + r}$$

for some constant  $C > 0$ . In particular, if **A3** holds and  $p + K = o(n)$  then  $\hat{\sigma}^2$  is consistent.

The covariate effects  $\beta$  will be the central target for inference. The first result is the following bound on the bias of  $\hat{\beta}$ , defined to be  $\|\mathbb{E}(\hat{\beta}) - \beta\|_\infty$ .

**Proposition 3** (The bias of  $\hat{\beta}$ ). If condition (26) holds then there exists a constant  $C > 0$  such that

$$\|\mathbb{E}(\hat{\beta}) - \beta\| \leq \frac{C\tau_n}{\min\{(1 - \sigma_{r+1})^3, \sigma_{r+s}^3\}} \cdot \|(X^T X)^{-1}\| \cdot \|X\| \cdot \|X\beta + X\theta + \alpha\|.$$

In particular, if **A1**, **A2** and **A3** hold then the bias of  $\hat{\beta}$  is of order  $o(1/\sqrt{n})$ .

In general, one may be interested in inferring the contrast  $\omega^T \beta$  for a given unit vector  $\omega$ , such as in comparing covariate effects or making predictions. Let  $\tilde{X} = X/\sqrt{n}$ , then from (19),

$$\text{Var}(\omega^T \hat{\beta}) = \frac{\sigma^2}{n} \omega^T \Theta \tilde{X}^T \hat{\mathcal{P}}_C \hat{\mathcal{P}}_C^T \tilde{X} \Theta \omega, \quad (27)$$

with  $\hat{\mathcal{P}}_C$  defined by (17). According to Proposition 3, a sufficient condition for valid inference of  $\omega^T \beta$  is  $\omega^T \Theta \tilde{X}^T \hat{\mathcal{P}}_C \hat{\mathcal{P}}_C^T \tilde{X} \Theta \omega \geq c$  for some constant  $c > 0$ . Note that this quantity is directly computable in practice. However, we will focus on the population condition with respect to the true matrix  $\mathcal{P}_C$  in the following result.

**Theorem 1** (Asymptotic distribution of  $\omega^T \hat{\beta}$ ). Assume that **A1**, **A2**, **A3** hold and

$$\sqrt{np} \cdot \min\{1 - \sigma_{r+1}, \sigma_{r+s}\} \geq 1. \quad (28)$$

For a given unit vector  $\omega \in \mathbb{R}^p$ , assume that

$$\|\tilde{Z}_{(r+1):p}^T \tilde{X} \Theta \omega\| \geq c \quad (29)$$

for some constant  $c > 0$  and sufficiently large  $n$ . Then as  $n \rightarrow \infty$ ,

$$\mathbb{P} \left( \frac{\omega^T \hat{\beta} - \omega^T \beta}{\sqrt{\frac{\hat{\sigma}^2}{n} \omega^T \Theta \tilde{X}^T \hat{\mathcal{P}}_C \hat{\mathcal{P}}_C^T \tilde{X} \Theta \omega}} \leq t \right) \rightarrow \Phi(t),$$

where  $\Phi$  is the cumulative distribution function of the standard normal distribution.

Next, we discuss the special case when  $\omega = e_j$ , which corresponds to making inference for the individual parameter  $\beta_j = e_j^T \beta$  for some  $1 \leq j \leq p$ . In this case, condition (29) has a simple interpretation.

**Corollary 1** (Valid inference of individual regression coefficient). Let  $\eta_j$  be the partial residual from regressing  $\tilde{X}_j$  against all other columns of  $\tilde{X}$  and  $R_j^2$  be the corresponding partial  $R^2$  for this regression. Assume that A1, A2, A3 hold and there exists a constant  $c > 0$  such that

$$\|\tilde{Z}_{(r+1):p}^T \eta_j\| / (1 - R_j^2) \geq c. \quad (30)$$

Then, as  $n \rightarrow \infty$ ,

$$\mathbb{P} \left( \frac{\hat{\beta}_j - \beta_j}{\sqrt{\frac{\hat{\sigma}^2}{n} e_j^T \Theta \tilde{X}^T \hat{P}_C \hat{P}_C^T \tilde{X} \Theta e_j}} \leq t \right) \rightarrow \Phi(t),$$

where  $\Phi$  is the cumulative distribution function of the standard normal distribution.

To understand condition (30), notice that  $\eta_j$  is the signal from  $X_j$ , after adjusting for the other covariates, while  $1 - R_j^2$  is the proportion of this additional information from  $\tilde{X}_j$ . Since  $\|\tilde{Z}_{(r+1):p}^T \eta_j\|$  is the magnitude of  $\eta_j$ 's projection on the subspace  $\text{col}(\tilde{Z}_{(r+1):p})$ , (30) essentially requires that the additional information contributed by  $X_j$  after conditioning on other covariates should nontrivially align with  $\text{col}(\tilde{Z}_{(r+1):p})$ , the subspace on which the effect of  $\beta_j$  lies. This type of condition is needed because if  $\tilde{Z}_{(r+1):p}^T \eta_j = 0$ , the parameter space of  $\beta_j$  degenerates to  $\{0\}$  and the inference is not meaningful. Notice that requirement (30) is completely different from assuming that  $|\beta_j|$  is large, and it does allow  $|\beta_j|$  to be zero or small.

The next result shows the consistency of estimating  $\alpha$ . It also provides bounds for the bias and variance of  $\hat{\gamma}$ , which are later used for testing the hypothesis  $H_0 : \alpha = 0$ .

**Proposition 4** (Consistency of  $\hat{\alpha}$  and  $\hat{\gamma}$ ). Assume that A1 and (26) hold. Then there exists a constant  $C > 0$  such that with high probability, the following hold.

1. Let  $\|W\|_{2 \rightarrow \infty}$  be the maximum of the Euclidean norms of the row vectors of  $W$ , then

$$\|\hat{\alpha} - \alpha\|_\infty \leq \frac{C}{\min\{(1 - \sigma_{r+1})^3, \sigma_{r+s}^3\}} \cdot \left( \|W\|_{2 \rightarrow \infty} \sqrt{K \sigma^2 \log n} + \tau_n \sqrt{n(p + \sigma^2)} \right).$$

2. There exists an orthogonal matrix  $O \in \mathbb{R}^{(K-r) \times (K-r)}$  such that

$$\|\mathbb{E} \hat{\gamma} - O \gamma\| \leq \frac{C \tau_n \sqrt{np}}{\min\{(1 - \sigma_{r+1})^3, \sigma_{r+s}^3\}}.$$

3. Let  $\Sigma_{\hat{\gamma}}$  be the covariance matrix of  $\hat{\gamma}$ , then

$$\left\| \Sigma_{\hat{\gamma}} - \sigma^2 \begin{pmatrix} (I_s - \Gamma^2)^{-1} & 0 \\ 0 & I_{K-r-s} \end{pmatrix} \right\| \leq \frac{C \tau_n}{\min\{(1 - \sigma_{r+1})^2, \sigma_{r+s}^2\}},$$

where  $\Gamma = \text{diag}(\sigma_{r+1}, \dots, \sigma_{r+s})$ .

Proposition 4 shows that  $\hat{\alpha}$  is an element-wise consistent estimate of  $\alpha$  if A3 holds,  $\min\{(1 - \sigma_{r+1})^3, \sigma_{r+s}^3\}$  is bounded away from zero, and  $\|W\|_{2 \rightarrow \infty} \sqrt{K \sigma^2 \log n} = o(1)$ . The last requirement holds when  $P$  is an incoherent matrix — the rows of  $W$  are in similar magnitudes — a commonly

observed property introduced by [Candès and Recht \[2009\]](#), [Candès and Tao \[2010\]](#). The other two inequalities of Proposition 4 show that the bounds for both bias and variance of  $\hat{\gamma}$  are of order  $o(1)$  if [A3](#) holds. These properties directly lead to the validity of the chi-squared test for network effects.

**Theorem 2** (Chi-squared test for  $\alpha$ ). Assume that [A1](#) and [A3](#) hold,  $K$  is a fixed integer, and  $p = o(n)$ . Then as  $n \rightarrow \infty$ , the statistic  $\|\hat{\gamma}_0\|^2$  constructed in Algorithm 1 satisfies

$$\|\hat{\gamma}_0\|^2 \xrightarrow{d} \chi_K^2,$$

where  $\chi_K^2$  is a random variable that follows the chi-squared distribution with  $K$  degrees of freedom and  $\xrightarrow{d}$  denotes the convergence in distribution.

The next theorem provides the theoretical result for  $\hat{\theta}$ .

**Theorem 3** (Inference of  $\theta$  and  $\xi$ ). The estimator  $\hat{\theta}$  in (19) follows a Gaussian distribution with covariance matrix  $\text{Cov}(\hat{\theta}) = \frac{\sigma^2}{n} \Theta \tilde{X}^T \hat{\mathcal{P}}_{\mathcal{R}} \tilde{X} \Theta$ . Furthermore, under [A1](#), [A2](#) and [A3](#),  $\|\mathbb{E}(\hat{\theta}) - \theta\| = o(1/\sqrt{n})$ . If in addition, there exists a constant  $c > 0$  such that

$$\|\tilde{Z}_{1:r}^T \eta_j\| / (1 - R_j^2) \geq c, \tag{31}$$

where  $\eta_j$  and  $R_j$  are defined in Corollary 1, then as  $n \rightarrow \infty$ ,

$$\mathbb{P} \left( \frac{\hat{\theta}_j - \theta_j}{\sqrt{\frac{\hat{\sigma}^2}{n} e_j^T \Theta \tilde{X}^T \hat{\mathcal{P}}_{\mathcal{R}} \tilde{X} \Theta e_j}} \leq t \right) \rightarrow \Phi(t).$$

The inference of each  $\xi_i, i \in [n]$  can be done by noticing  $\xi_i = X_i \cdot \theta$  where  $X_i$  is the  $i$ th row of  $X$ .

Condition (31) is in parallel of (30); see the discussion following Corollary 1 for its interpretation.

## 4 Small projection perturbation under random network models

We have introduced the small projection perturbation condition as a general way to characterize the structural perturbation level our framework could tolerate. To provide more insights about this characterization, in this section, we study the small projection perturbation assumption in a special setting when the perturbation of  $P$  comes from random network models. Specifically, we assume that the edge between each pair of nodes  $(i, j)$  is generated independently from a Bernoulli distribution with  $\mathbb{P}(A_{ij} = 1) = P_{ij}, 1 \leq i < j \leq n$ . In particular,  $\mathbb{E}A = P$ . This model is known as the inhomogeneous Erdős-Rényi model [[Bollobas et al., 2007](#)]. To clarify, the randomness discussed in this section comes from the random network model above, which is assumed to be independent of the randomness from the noise  $\epsilon$  in the regression model.

Section 4.1 investigates the most difficult setting when no prior information is available about the network model, while Section 4.2 presents the results in the arguably easiest setting when the true underlying network can be accurately estimated. Additionally, in Section 4.1, we show that our theoretical requirement for the network density in the completely non-informative situation is rate optimal. Section 4.3 further extends the results in another commonly seen setting when the Laplacian matrix represents the relational information.

## 4.1 The non-informative situation: small projection perturbation for adjacency matrices

In this section, we investigate the situation when no prior information about the network model is available. This can be seen as the most difficult, yet the most general, situation. Arguably, the only reasonable approximation of  $P$  is  $\hat{P} = A$ . We will study [A3](#) for this version of  $\hat{P}$ .

Recall that the main requirement of [A3](#) is  $\tau_n = o(1/\sqrt{n})$ , where  $\tau_n$  measures the level of perturbation of the network subspace  $S_K(P)$  associated with the covariate subspace  $\text{col}(X)$ , defined in [\(23\)](#). Existing results relevant for controlling  $\tau_n$ , such as [Abbe et al. \[2017\]](#), [Cape et al. \[2019\]](#), [Mao et al. \[2020\]](#), [Lei \[2019\]](#), do not give sufficiently tight bounds to support the inference, even for dense networks. The recent work of [Xia \[2019\]](#) contains useful tools for obtaining such error bound that allows sparsity when  $P$  is a low-rank matrix. However, since in general problems,  $P$  may be of full rank, a new tool is needed for theoretical analysis. The following assumption is made about  $P$ .

**Assumption A4** (Eigenvalue gap of the expected adjacency matrix). Let  $A$  be the adjacency matrix of a random network generated from the inhomogeneous Erdős-Rényi model with the edge probability matrix  $P = \mathbb{E}A$ . Assume that the  $K$  largest eigenvalues of  $P$  are well separated from the remaining eigenvalues and their range is not too large:

$$\min_{i \leq K, i' > K} |\lambda_i - \lambda_{i'}| \geq \rho' d, \quad \max_{i, i' \leq K} |\lambda_i - \lambda_{i'}| \leq d/\rho',$$

where  $\rho' > 0$  is a constant and  $d = n \cdot \max_{ij} P_{ij}$ .

The quantity  $d$  can be seen as an upper bound of the network node degree, which has been widely used to measure network density (e.g., [Lei and Rinaldo \[2014\]](#), [Le et al. \[2017\]](#)). [A4](#) is very general in the sense that it only assumes that  $P$  has a sufficiently large eigenvalue gap, but  $P$  need not be low-rank or even approximately low-rank. It appears that this is already sufficient to guarantee the small projection perturbation property of  $A$ , as stated in the next theorem. We believe the theorem itself can be used as a general tool for statistical analysis of independent interest.

**Theorem 4** (Concentration of perturbed projection for adjacency matrix). Let  $w_1, \dots, w_n$  and  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$  be eigenvectors and corresponding eigenvalues of  $P$  and similarly, let  $\hat{w}_1, \dots, \hat{w}_n$  and  $\hat{\lambda}_1 \geq \hat{\lambda}_2 \geq \dots \geq \hat{\lambda}_n$  be the eigenvectors and eigenvalues of  $A$ . Denote  $W = (w_1, \dots, w_K)$  and  $\hat{W} = (\hat{w}_1, \dots, \hat{w}_K)$ . Assume that [A4](#) holds and  $d \geq C \log n$  for a sufficiently large constant  $C > 0$ . Then for any fixed unit vector  $v$ , with high probability,

$$\|(\hat{W}\hat{W}^T - WW^T)v\| \leq \frac{2\sqrt{K \log n}}{d}. \quad (32)$$

Notice that the bound is for a given deterministic vector  $v$  instead of all unit vectors. The latter would be equivalent to a bound on  $\|\hat{W}\hat{W}^T - WW^T\|$  and is too large for our inference purpose. By restricting the scope of applicability, our result trades off for the tighter bound [\(32\)](#), which is crucial for our inference to work in relatively sparse network settings.

**Corollary 2** (Small projection perturbation, adjacency matrix case). Assume that  $\hat{P} = A$  is used in [Algorithm 1](#). If [A4](#) holds and  $d$  satisfies

$$\frac{d \cdot \min \{(1 - \sigma_{r+1})^3, \sigma_{r+s}^3\}}{(Knp \log n)^{1/2}} \rightarrow \infty$$

then [A3](#) holds with high probability for sufficiently large  $n$ .

Corollary 2 provides a sufficient condition for valid inference. If  $\min\{1 - \sigma_{r+1}, \sigma_{r+s}\}$  is bounded away from zero while  $K$  and  $p$  are fixed then  $d$  needs to grow faster than  $\sqrt{n \log n}$ . This is arguably a strong assumption, although it does allow for moderately sparse networks. A natural question is whether it can be relaxed. Unfortunately, the answer to this is negative. We now show that under the current assumptions, the bound (32) is rate optimal up to a logarithm factor and we cannot guarantee valid inference if  $d$  is of the order  $\sqrt{n}$ .

**Theorem 5** (Tightness of concentration and degree requirements). Assume  $C\sqrt{n/\log n} \leq d \leq n^{1-\xi}$  for some constant  $\xi \in (0, 1/2]$  and some sufficiently large constant  $C > 0$ . The following statements hold.

- (i) There exists a configuration of  $(K, v, P)$  satisfying the condition of Theorem 4 with  $K = O(1)$ , under which, for a sufficiently large  $n$ ,

$$\|(\hat{W}\hat{W}^T - WW^T)v\| \geq c/d$$

holds with high probability and a constant  $c > 0$ .

- (ii) There exists a configuration of  $(K, p, X, P, \beta, \theta, \alpha)$  satisfying A1, A2 and the conditions of Theorem 4 with  $K, p = O(1)$  and  $\min\{1 - \sigma_{r+1}, \sigma_{r+s}\} \geq 1/4$ , under which, for a sufficiently large  $n$ , we have

$$\|\mathbb{E}\hat{\beta} - \beta\|_\infty \geq c'/d \quad \text{and} \quad \max_j \text{Var}(\hat{\beta}_j) \leq C'/n$$

with high probability for some constant  $C', c' > 0$ .

Notice that the two statements of Theorem 5 are different and, in general, neither implies the other. The first statement is about the general concentration bound of Theorem 4. It indicates that for the given range of  $d$ , the bound of (32) is rate optimal (up to a logarithm order). The second statement is about the necessary condition of the inference problem, under the model-free setting. When  $d = \sqrt{n}$ , the bias is at least in the order of  $1/\sqrt{n}$ . In contrast, the standard deviation is at most of order  $1/\sqrt{n}$ . Therefore, we will not be able to give asymptotically correct testing or confidence intervals under these circumstances. Meanwhile, under the same condition, Corollary 2 shows that if  $d$  grows faster than  $\sqrt{n \log n}$ , valid inference could be achieved. In combination, we observe a phase transition at the network degree of  $\sqrt{n}$ .

To conclude this section, we address the problem of selecting the correct dimension  $r$ , which is used in Algorithm 1, as another application of Theorem 4. Specifically, denote  $\hat{d} = \frac{1}{n} \sum_{i,j=1}^n A_{ij}$  and  $\bar{d} = \frac{1}{n} \sum_{i,j=1}^n P_{ij}$ . The following rule is used to select  $r$ :

$$\hat{r} = \max \left\{ i : \hat{\sigma}_i \geq 1 - \frac{4\sqrt{pK \log n}}{\hat{d}} \right\}. \quad (33)$$

This estimate can be shown to give the correct dimension with high probability.

**Corollary 3** (Consistency of estimating the dimension of  $\mathcal{R}$ ). Assume that the conditions of Theorem 4 hold and

$$\bar{d} \geq \frac{16\sqrt{pK \log n}}{1 - \sigma_{r+1}}. \quad (34)$$

Then for a sufficiently large  $n$ , the dimension estimate  $\hat{r}$  defined in (33) satisfies  $\hat{r} = r$  with high probability.

## 4.2 The informative situation: small projection perturbation for parametric models

In the previous section, it is shown that if  $A$  is used as  $\hat{P}$  in Algorithm 1, the network degree needs to grow faster than  $\sqrt{n}$  for the small perturbation assumption to hold. In many cases, it may be reasonable to assume that  $P$  satisfies certain structural conditions. Leveraging such additional information can provide a more accurate estimate of  $P$  and, in turn, ensure the validity of inference for potentially much sparser networks. In this section, we investigate a few special cases for which the edge density requirement can be substantially relaxed, highlighting the benefit of knowing the correct model.

The discussion begins with the stochastic block model (SBM) as the true network generating model. Specifically, assume that there exists a vector  $g \in [K]^n$  specifying the community node labels and a symmetric matrix  $B \in [0, 1]^{K \times K}$  such that  $P_{ij} = B_{g_i g_j}$ . The SBM has been widely used to model community structures [Holland et al., 1983], and its theoretical properties have been well-understood thanks to intensive research in recent years. For more detail, refer to the review paper of Abbe [2018] and the references therein. Since the SBM is used in this section only to illustrate how the degree requirement can be relaxed, we will focus on the following special configuration.

**Assumption A5** (Stochastic block model). Let  $n_k$  be the number of nodes in the  $k$ th community. Assume that  $(1 - t)n/K \leq n_k \leq (1 + t)n/K$  for some constant  $t$  and all  $1 \leq k \leq K$ . Moreover, assume that  $B = \kappa_n B_0$ , where  $B_0 \in [0, 1]^{K \times K}$  is a fixed matrix and  $\kappa_n$  controls the dependence of network density on  $n$  and  $K = o(n^{1/4})$ .

Another example is the degree-corrected block model (DCBM) proposed by Karrer and Newman [2011], which generalizes the SBM by allowing the degree heterogeneity of the nodes. Specifically, in addition to the SBM parameters, the model has a degree parameter  $\nu \in \mathbb{R}^n$  such that  $P_{ij} = \nu_i \nu_j B_{g_i g_j}$ . Notice that  $\nu_i$  is only identifiable up to a scale so additional constraints are needed. The following simplified assumption is made about it.

**Assumption A6** (Identifiability of degree parameter). Assume  $\sum_{g_i=k} \nu_i = n_k$  and there exists a constant  $\zeta$  such that  $1/\zeta \leq \nu_i \leq \zeta$ .

Notice that under Assumptions A5 and A6, both the SBM and DCBM satisfy Assumption A4 with  $d = n\kappa_n$ . In particular, under the SBM, the subspace of the  $K$  leading eigenvectors of  $P$  has the same component for all nodes within the same community. Therefore, the model under the SBM reduces to a linear regression model with fixed group effects according to communities. The following assumption is further made for the two models.

**Assumption A7** (Exact recovery of community labels). Assume the community  $g$  is known with  $n\kappa_n / \log n \rightarrow \infty$ .

Although  $g$  is seldom known in practice, this assumption is based on the fact that in many cases  $g$  can be exactly recovered from the network. Indeed, there exist many polynomial-time algorithms that ensure the exact recovery of  $g$  such as Gao et al. [2017], Lei and Zhu [2017], Abbe et al. [2017], Li et al. [2020a], Lei et al. [2020] for the SBM and Lei and Zhu [2017], Chen et al. [2018], Gao et al. [2018] for the DCBM, to name a few. The corresponding regularity condition needed for such strong consistency of community detection is already reflected in the degree requirement in Assumption A7. Given the community labels, the commonly used estimator of  $P$  would be the MLE (see Section F of the supplementary material for details). When such parametric estimates are used as  $\hat{P}$  in Algorithm 1, the corresponding estimation and inference procedure are referred to as the *parametric version* of our method, in contrast to the model-free version when  $A$  is used as

$\hat{P}$ . The following theorem shows that the parametric version delivers valid inference under much weaker network density assumptions compared to its nonparametric counterpart.

**Theorem 6** (Small projection perturbation under block models). Let  $d = n\kappa_n$ . Let  $\hat{P}$  parametric estimator of  $P$  introduced in Section F. Under either of the following settings:

- (i)  $A$  is generated from  $P$  according to the SBM satisfying Assumption A5 and A7 with

$$\min \left\{ (1 - \sigma_{r+1})^6, \sigma_{r+s}^6 \right\} \cdot \frac{d}{p \log n} \rightarrow \infty;$$

- (ii)  $A$  is generated from  $P$  according to the DCBM satisfying Assumption A5, A6 and A7 with

$$\min \left\{ (1 - \sigma_{r+1})^6, \sigma_{r+s}^6 \right\} \cdot \frac{d}{K^2 p \log n} \rightarrow \infty;$$

the estimator  $\hat{P}$  satisfies Assumption A3 with high probability.

Again, if  $p$ ,  $K$  and  $\min \{(1 - \sigma_{r+1}), \sigma_{r+s}\}$  are fixed, then the degree requirement of the parametric version for the small projection perturbation to hold is  $d/\log n \rightarrow \infty$  for either of two block models. This is much weaker than the degree requirement for the nonparametric estimation, indicating the benefit of knowing the underlying model for  $P$ . Although the above result is only about two special classes of models for the network, it clearly shows the tradeoff between the model assumption and statistical efficiency: without any information about the true model, the model-free version of the proposed method requires the average degree to grow faster than  $\sqrt{n}$  to be effective; in contrast, if the class that the true model belongs to is known then the parametric version of our method performs well for much sparser networks.

### 4.3 Extension to Laplacian individual effects

So far, we have treated  $A$  as a perturbed version of  $P$ . A natural and popular alternative is to use the Laplacian matrix as the perturbed version of the relational information  $P$  [Smola and Kondor, 2003, Sadhanala et al., 2016, Li et al., 2019]. Specifically, we assume that  $\alpha$  lies in the subspace determined by  $K$  eigenvectors of  $P = \mathbb{E}L = \mathbb{E}D - \mathbb{E}A$  corresponding to the smallest eigenvalues, while the estimation procedure is based on the perturbed version  $\hat{P} = L = D - A$  of  $P$ , where  $D$  is the diagonal matrix with node degrees  $d_i$  on the diagonal. Similar to Assumption A4 about the spectrum of  $\mathbb{E}A$ , the following assumption about the spectrum of  $\mathbb{E}L$  is made.

**Assumption A8** (Eigenvalue gap of the expected Laplacian). Let  $L = D - A$  be the Laplacian of a random network generated from the inhomogeneous Erdős-Rényi model and  $P = \mathbb{E}L$ . Denote by  $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$  the eigenvalues of  $P$ . Assume that the  $K$  smallest eigenvalues of  $P$  are well separated from the remaining eigenvalues and their range is not too large:

$$\min_{i \leq K, i' > K} |\lambda_i - \lambda_{i'}| \geq \rho' d, \quad \max_{i, i' \leq K} |\lambda_i - \lambda_{i'}| \leq d/\rho',$$

where  $\rho' > 0$  is a constant and  $d = n \cdot \max_{ij} P_{ij}$ .

We show that, under the perturbation mechanism of Assumption A8, the eigenvectors of the Laplacian corresponding to the  $K$  smallest eigenvalues satisfy the small projection perturbation requirement, and therefore the inference framework based on Algorithm 1 with  $\hat{P} = L$  remains valid.



**Theorem 7** (Concentration of perturbed projection for the Laplacian). Let  $w_1, \dots, w_n$  and  $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$  be eigenvectors and corresponding eigenvalues of  $\mathbb{E}L = \mathbb{E}D - \mathbb{E}A$  and similarly, let  $\hat{w}_1, \dots, \hat{w}_n$  and  $\hat{\lambda}_1 \leq \hat{\lambda}_2 \leq \dots \leq \hat{\lambda}_n$  be the eigenvectors and eigenvalues of  $L = D - A$ . Denote  $W = (w_1, \dots, w_K)$  and  $\hat{W} = (\hat{w}_1, \dots, \hat{w}_K)$ . Assume that Assumption A8 holds and  $d \geq C \log n$  for a sufficiently large constant  $C$ . Then for any fixed unit vector  $v$ , with high probability we have

$$\|(\hat{W}\hat{W}^T - WW^T)v\| \leq \frac{C[K(1 + n\|W\|_\infty^2)]^{1/2} \log n}{d},$$

where  $\|W\|_\infty$  denotes the largest absolute value of entries in  $W$ .

Compared to the bound for the adjacency matrix in Theorem 4, the bound in Theorem 7 contains a new term  $n\|W\|_\infty^2$ . When  $\mathbb{E}L$  is incoherent,  $\|W\|_\infty^2$  is of order  $1/n$  (for example under stochastic block model) and therefore the upper bound is in the order of  $\sqrt{K \log^2 n}/d$ . Both the term  $n\|W\|_\infty^2$  and the extra  $\sqrt{\log n}$  factor are results of the deviation of node degrees from their expected values.

**Corollary 4** (Small projection perturbation, Laplacian case). Assume that  $L$  is used as  $\hat{P}$  in Algorithm 1 and  $P = \mathbb{E}L$ . If Assumption A8 holds and  $d$  is sufficiently large so that

$$\frac{d \cdot \min \{(1 - \sigma_{r+1})^3, \sigma_{r+s}^3\}}{[Kn p(1 + n\|W\|_\infty^2)]^{1/2} \log n} \rightarrow \infty,$$

then Assumption A3 holds with high probability. In particular, the proposed inference framework in Section 3 is valid with high probability.

## 5 Simulation

In this section, we present a simulation study to evaluate the proposed method and compare it with other benchmark methods for linear regression on networks. First, we evaluate the validity of the inference framework and demonstrate the predicted phase transition of the model-free method, as well as the advantage of the parametric version in the situation when the true network model is known. After that, we introduce comparisons with other benchmark methods under several network effect models.

### 5.1 Inference validation

We will evaluate our theoretical claims about the inference and phase transition in this subsection. For this purpose, we will again assume  $K$  to be known to exactly match our theoretical framework in this subsection. However, this does not lose generality because in all of our experiment settings, because the methods we mentioned before [Chen and Lei, 2018, Le and Levina, 2015, Li et al., 2020b] can accurately identify the  $K$  almost perfectly in our settings. We fix  $p = K = 4$  in all configurations. The network is generated by the SBM with  $K = 4$  communities. In Section H of the supplementary material, we provide additional results and also a study when the network is generated by the more general DCBM. Using the parameterization in Section 4.1, we set  $B = 0.21\mathbf{1}_4\mathbf{1}_4^T + 0.8I_4$ , where  $\mathbf{1}_k$  denotes the all-one vector of length  $k$ . Three levels of sample size  $n = 1000, 2000, 4000$ , and three levels of average expected degree  $\varphi_n = 2 \log n, \sqrt{n}$ , and  $n^{2/3}$  for each level of  $n$  are compared. Given the eigenvectors  $w_1, \dots, w_n$  from  $P$ , we generate  $X \in \mathbb{R}^{n \times p}$  in the following way: Set  $X_1/\sqrt{n} = w_1$ ; Set  $X_j/\sqrt{n} = \sqrt{\frac{1}{25}}w_j + \sqrt{\frac{24}{25}}w_{j+3}, j = 2, 3, 4$ . This

configuration gives a design with  $r = 1, s = 3$  and  $(\sigma_1, \sigma_2, \sigma_3, \sigma_4) = (1, 0.2, 0.2, 0.2)$ . In particular, the setup with  $\beta = (0, 1, 1, 1)^T$  and  $\theta = (1, 0, 0, 0)^T$  satisfies the proposed model and is fixed in all settings. Similarly, the  $\gamma$  can be any vector with the first coordinate being zero. We consider two cases:  $\gamma = (0, 1, 1, 1)^T$  and  $\gamma = (0, 0, 0, 0)^T$ . As discussed, the case of  $\gamma = (0, 0, 0, 0)^T$  indicates that there is no network effect. The two settings do not give any difference on inference of  $\beta$  while the latter is used to verify the validity of the  $\chi^2$  test. Also, notice that  $X_1$  violates assumption (30) because it is completely orthogonal to the space of  $\text{col}(\hat{Z}_{r+1:p})$ . Therefore, valid inference cannot be made for  $\beta_1$ . This setup highlights that having a degenerate parameter space on  $\beta_1$  does not impact the validity of our inference on the other parameters  $\beta_2, \beta_3, \beta_4$ , as shown by both the theoretical and numerical analyses. Both the model-free and parametric versions of the subspace projection (SP) methods are used to demonstrate the empirical difference, denoted by ‘‘SP’’ and ‘‘SP-SBM’’, respectively. All results are taken as the average of the 50 independent replications.

Table 2 shows the ratio between the absolute bias and standard deviation (SD) averaged across  $\hat{\beta}_2, \hat{\beta}_3$  and  $\hat{\beta}_4$  for different  $n$  and network densities, when the network is generated from the SBM. For average degree of order  $\sqrt{n}$  or below, the ratio does not vanish with increasing  $n$  for the SP, indicating that valid statistical inference for the parameters cannot be achieved. For denser networks, the bias becomes vanishing; thus, the asymptotic inference becomes valid. These observations coincide with the prediction of the phase transition at degree  $\sqrt{n}$ . In contrast, the SP-SBM results in much smaller bias-SD ratios and achieves vanishing ratio even at degree  $\sqrt{n}$ . At the borderline case  $2 \log n$ , such vanishing pattern is not clearly observed, even for the SP-SBM.

Table 2: The average bias-SD ratios for  $\beta_2, \beta_3$  and  $\beta_4$  when the network is generated from the SBM.

| Method | bias-SD ratio |                         |             |             | coverage probability    |            |           |
|--------|---------------|-------------------------|-------------|-------------|-------------------------|------------|-----------|
|        | $n$           | average expected degree |             |             | average expected degree |            |           |
|        |               | $2 \log n$              | $\sqrt{n}$  | $n^{2/3}$   | $2 \log n$              | $\sqrt{n}$ | $n^{2/3}$ |
| SP     | 1000          | 1.950                   | 0.414       | 0.326       | 0.7282                  | 0.9309     | 0.9401    |
|        | 2000          | 2.526                   | 0.344       | 0.237       | 0.5275                  | 0.9373     | 0.9453    |
|        | 4000          | 2.460                   | 0.408       | 0.175       | 0.5879                  | 0.9374     | 0.9462    |
| SP-SBM | 1000          | 0.922                   | 0.164       | $< 10^{-4}$ | 0.8763                  | 0.9501     | 0.9486    |
|        | 2000          | 0.931                   | 0.025       | $< 10^{-4}$ | 0.8502                  | 0.9505     | 0.9500    |
|        | 4000          | 0.847                   | $< 10^{-4}$ | $< 10^{-4}$ | 0.8423                  | 0.9500     | 0.9515    |

Correspondingly, Table 2 also shows the resulting average coverage probabilities  $p_{\text{cov}}$  of the 95% confidence intervals for  $\beta_2, \beta_3$  and  $\beta_4$ . Notice that, differently from Table 2, the coverage probability is calculated as the average of 500 Monte Carlo runs, thereby taking the errors of  $\hat{\sigma}$  into account. It can be seen that the confidence interval for the SP is not good enough for degree of  $\sqrt{n}$  or lower, but does give valid results for denser graphs. The parametric version is always more accurate than the model-free version, and in particular, remains valid for the order of  $\sqrt{n}$ .

## 5.2 Comparison with other benchmarks

In this subsection, we will evaluate the practical performance of our method in different settings of network effects and compare it with other benchmark methods. The first benchmark in the comparison is the ordinary least squares method (OLS). The other two are the SIM of Manski [1993] and the RNC of Li et al. [2019] introduced before. It has been shown that the RNC is a better modeling option than both OLS and the SIM [Li et al., 2019]. However, the RNC method cannot make inference of the parameters, and it is computationally more intensive than our method. In this experiment, the RNC is always tuned by 10-fold cross-validation. For our method, the number  $K$  is determined by

the method of [Le and Levina \[2015\]](#). The theory-driven tuning of  $r$  introduced in (33) is based on large-sample properties and may be conservative for small samples. We also propose a bootstrapping method in Section G of the supplementary material. This method is used to select  $r$  for the numerical and data examples from now on and works very well. Lastly, a systematic model fitting procedure is used for SP to take advantage of its available inference framework. Specifically, the SP methods would check the  $\chi^2$  test result of  $\gamma$ . If the p-value for the  $\chi^2$  test is more significant than 0.05 (any other reasonable level can be chosen), the method would instead return OLS fit, which is equivalent to constraining  $\gamma = 0$  in our estimation procedure.

As before, the network is generated by the SBM with  $K = 4$  and  $n = 1000$  with the same configurations. The results for the more general situation of DCBM can be found in Section H of the supplementary material. However, to avoid overly tailoring the setup for our methods,  $X$ 's are generated differently. Specifically,  $X_2, X_3, X_4$  are randomly generated from Gaussian distribution  $N(0, 1)$ , uniform distribution  $U(0, 1)$  and exponential distribution  $\text{Exp}(1)$  before standardization is applied.  $X_1$  is directly set to be  $\sqrt{n}w_1$  as before, where  $w_1$  is the first eigenvector of  $P$ . Meanwhile, three different schemes are used to generate the individual effect  $\alpha$ , so that the model misspecification situations can also be tested. In particular, the first scenario is exactly the same setting as in the last section, which corresponds to the assumed model. This individual effect setting is called "eigenspace". In the second scenario, we set  $\gamma = 0$ , thus the model becomes the standard linear regression model for which OLS is designed. In the last scenario, we set  $\alpha$  to be the average of the three eigenvectors corresponding to the three smallest nonzero eigenvalues of the observed Laplacian matrix  $L$ . This  $\alpha$  gives a small value of  $\sum_{i \sim j} (\alpha_i - \alpha_j)^2$ , and thus matches the assumption of the RNC framework [\[Li et al., 2019\]](#). As discussed in Section 4.3, our method should still work by using  $\hat{P} = L$  when  $\alpha$  is defined in this way. Therefore, for the evaluation in this section, the Laplacian version of the SP estimator is also included and labeled as "SP-L".

Since the data generating models have different parameterizations, we directly compare the relative mean estimation error (MSE) of the expected response  $\mathbb{E}Y$ . All of the results are averaged over 50 independent replications and are shown in Table 3.

Table 3: The mean squared error (MSE)  $\times 10^2$  of  $\mathbb{E}Y$  when the network is generated from the SBM with  $n = 1000, p = 4, K = 4$  with three types of individual effects.

| indiv. effects | avg. degree | SP    | SP-SBM | OLS   | SIM   | RNC   | SP-L  | SP-SBM-L |
|----------------|-------------|-------|--------|-------|-------|-------|-------|----------|
| eigenspace     | $2 \log n$  | 12.41 | 5.17   | 13.35 | 18.74 | 12.11 | 24.31 | 9.26     |
|                | $\sqrt{n}$  | 5.18  | 0.10   | 13.35 | 17.24 | 10.46 | 10.69 | 6.72     |
|                | $n^{2/3}$   | 1.37  | 0.10   | 13.35 | 15.83 | 7.38  | 7.51  | 6.72     |
| $\gamma = 0$   | $2 \log n$  | 0.13  | 0.13   | 0.13  | 27.07 | 0.30  | 0.13  | 0.13     |
|                | $\sqrt{n}$  | 0.13  | 0.14   | 0.13  | 25.88 | 0.62  | 0.13  | 0.14     |
|                | $n^{2/3}$   | 0.14  | 0.14   | 0.13  | 25.34 | 1.94  | 0.14  | 0.13     |
| smooth         | $2 \log n$  | 25.03 | 24.98  | 25.17 | 28.94 | 21.48 | 12.67 | 25.07    |
|                | $\sqrt{n}$  | 22.71 | 22.95  | 25.33 | 27.44 | 19.06 | 12.75 | 24.14    |
|                | $n^{2/3}$   | 1.14  | 1.19   | 9.51  | 26.10 | 11.75 | 4.84  | 5.35     |

Under the model with eigenspace individual effects (the proposed model), it can be seen that OLS never renders competitive performance while the SIM is only slightly better than OLS. The RNC gives much better results than OLS and the SIM because it effectively incorporates the network information. However, since the network is noisy, its performance is far from adequate. Both versions of our method (SP and SP-SBM) significantly outperform the other methods, with the parametric version performing better than the model-free version, as expected. When there are no individual effects ( $\gamma = 0$ ), OLS becomes the correct model and gives optimal results, as expected. The RNC and our methods all are correctly adaptive to this setting, but the SP methods are still

better than the RNC. The performance under the model with smooth individual effects (the RNC model) is noisier as it depends on the perturbation of the network. Overall, neither OLS nor the SIM works. However, under the SBM with average degree  $n^{2/3}$ , the network is too dense, such that the smooth individual effects become almost identical everywhere, and OLS becomes effective. In this setting, using the Laplacian matrix in our method still models the correct form of individual effects, so SP-L still works and is more effective than the RNC. Note that the parametric versions (SP-SBM-L) are no longer better than the SP in this situation because the individual effects depend on the observed network instead of the population signal; thus, the parametric methods are not using the correct model.

In conclusion, when there are network effects, our method outperforms the other benchmark methods in the experiments. When there are no network effects, it is adaptive to the simpler model and gives a similar result to OLS. Compared to the model-free SP, the parametric SP method is less robust to model misspecification.

## 6 School conflict reduction study

In [Paluck et al. \[2016\]](#), an experiment is conducted to study the impacts of educational workshops on reducing conflicts in schools. The experimenters randomly selected 26 middle schools in New Jersey in which they held educational workshops. In each school, the experimenters first determined a group of eligible students. A small proportion of them was selected (randomly after gender and race blocking) to participate in bi-monthly educational workshops about how to help reduce school conflicts. Students were also asked to name who they “spend time with” in the survey, and the answers were used to construct the social network structure. We ignore the directions of edges and treat two students as connected as long as either one identifies the other as a connection. For each school, we take the largest connected component of the social network as the final data. In total, there are 9,429 students in our analysis.

We follow a similar analysis as in [Paluck et al. \[2016\]](#). In particular, students provided how much their friends “influence their way of dealing with conflicts”, and how much their friends “talked about how to change the school climate”. These two variables are averaged as the response, as a measure of “friends’ support of reducing conflicts”. The covariates include whether the student participates the workshops, the proportion of the student’s friends involved in the workshops, and those selected based on the significance in an OLS model fitting, including gender, whether the student is Asian or not, and the number of conflicts the student experienced in the past year (summarized in nine categories of conflicts in the survey). The school effects are also included to accommodate the individual school differences.

Applying the model-free version of our SP method, the  $\chi^2$  test results in a very small p-value ( $< 10^{-5}$ ), indicating that the social effect is significant. The estimated  $r$  is zero, so there is no confounding between the covariates and the network, and  $\xi$  is trivially zero. The covariate effects are given in [Table 4](#), in comparison with the OLS results without modeling the social effect. The school effects are not presented due to space limit.

For the first three covariates, the estimated effects are similar between our method and the OLS estimates. The estimates reveal that female students and asian students tend to observe more supports from friends about reducing school conflicts. The more conflicts one experiences, the more supports they would see from friends. Meanwhile, the estimates of the educational program effects by OLS and our method are very different. Neither of the two covariates about the educational program are estimated to be significant in OLS, while both of them have reasonably small p-values under our

model. Because our method includes the OLS as a special case, and the  $\chi^2$  test indicates a strong network effect, we believe the inference of our method is more reasonable. Our conclusion can also be supported by the original analysis of experimenters [Paluck et al., 2016], who delivered the same conclusion. However, the analysis of Paluck et al. [2016] takes the experiment design and sampling probability into consideration with additional assumptions about the spill-over effects of the educational workshops and is based on complicated causal inference methods. Therefore, their conclusion is about the stronger causal relation. In contrast, our method does not take advantage of the design, nor do we know whether the network cohesion assumption has an explicit correspondence to the widely used spill-over effects assumptions. So our conclusion is not causal. We leave the development of causal inference under the current framework for future work.

Table 4: Regression coefficients of covariate effects in the school conflict example.

|                                | Our method |             | OLS    |             |
|--------------------------------|------------|-------------|--------|-------------|
|                                | coef.      | p-val.      | coef.  | p-val.      |
| Gender: Boy                    | -0.140     | $< 10^{-5}$ | -0.146 | $< 10^{-5}$ |
| Race: Asian                    | 0.175      | $< 10^{-5}$ | 0.150  | $< 10^{-5}$ |
| Conflicts in the past year     | 0.034      | $< 10^{-5}$ | 0.038  | $< 10^{-5}$ |
| Educated                       | 0.068      | 0.010       | 0.048  | 0.097       |
| Proportion of educated friends | 0.239      | 0.029       | 0.046  | 0.71        |

In addition to the current example, we also apply our method to analyze socially active levels of adolescents based on the AddHealth study of Harris [2009]. The details are included in Appendix I.

## 7 Conclusion

We have introduced a linear regression model on observations linked by a network. The model comes with a computationally efficient inference algorithm that can tolerate network observational errors. The study in this paper focuses on using Assumption A3 to control the estimation bias and deliver valid inference. It makes important progress toward the inference problem for the network-linked model.

The next step is to explore whether a certain bias correction can be applied so that the small perturbation assumption can be further relaxed. Developing such a technique will require an accurate estimate of the perturbation  $\hat{P} - P$  and, in turn, new tools for characterizing such random matrices. Meanwhile, the current focus is on the fixed design problem of  $X$  and  $P$ ; the framework remains valid if we condition on  $X$  and  $P$  while assuming that they are independent. Exploring the model with dependence between  $X$  and  $P$  is another promising research direction.

## Acknowledgement

C. M. Le is supported in part by the NSF grant DMS-2015134. T. Li is supported in part by the NSF grant DMS-2015298 and the Quantitative Collaborative Award from the College of Arts and Sciences at the University of Virginia.

## References

- E. Abbe. Community detection and stochastic block models: Recent developments. *Journal of Machine Learning Research*, 18(177):1–86, 2018.
- E. Abbe, J. Fan, K. Wang, and Y. Zhong. Entrywise eigenvector analysis of random matrices with low expected rank. *arXiv preprint arXiv:1709.09565*, 2017.
- G. W. Basse and E. M. Airoldi. Limitations of design-based causal inference and a/b testing under arbitrary and network interference. *Sociological Methodology*, 48(1):136–151, 2018a.
- G. W. Basse and E. M. Airoldi. Model-assisted design of experiments in the presence of network-correlated outcomes. *Biometrika*, 105(4):849–858, 2018b.
- M. Belkin and P. Niyogi. Laplacian eigenmaps for dimensionality reduction and data representation. *Neural Computation*, 15(6):1373–1396, 2003.
- Y. Benjamini and Y. Hochberg. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)*, 57(1):289–300, 1995.
- R. Bhatia. *Matrix Analysis*. Springer-Verlag New York, 1996.
- B. Bollobas, S. Janson, and O. Riordan. The phase transition in inhomogeneous random graphs. *Random Structures and Algorithms*, 31:3–122, 2007.
- Y. Bramoullé, H. Djebbari, and B. Fortin. Identification of peer effects through social networks. *Journal of Econometrics*, 150(1):41–55, 2009.
- V. V. Buldygin and K. K. Moskvichova. The sub-Gaussian norm of a binary random variable. *Theory of Probability and Mathematical Statistics*, (86):33–49, 2013.
- C. T. Butts. Network inference, error, and informant (in) accuracy: a bayesian approach. *social networks*, 25(2):103–140, 2003.
- E. J. Candès and B. Recht. Exact matrix completion via convex optimization. *Foundations of Computational mathematics*, 9(6):717–772, 2009.
- E. J. Candès and T. Tao. The power of convex relaxation: Near-optimal matrix completion. *IEEE Transactions on Information Theory*, 56(5):2053–2080, 2010.
- J. Cape, M. Tang, and C. E. Priebe. The two-to-infinity norm and singular subspace geometry with applications to high-dimensional statistics. *The Annals of Statistics*, 47(5):2405–2439, 2019.
- A. Chandrasekhar and R. Lewis. Econometrics of sampled networks. *Unpublished manuscript, MIT.[422]*, 2011.
- S. Chatterjee. Matrix estimation by universal singular value thresholding. *The Annals of Statistics*, 43(1):177–214, 2015.
- K. Chen and J. Lei. Network cross-validation for determining the number of communities in network data. *Journal of the American Statistical Association*, 113(521):241–251, 2018.
- Y. Chen, X. Li, J. Xu, et al. Convexified modularity maximization for degree-corrected stochastic block models. *The Annals of Statistics*, 46(4):1573–1602, 2018.

- A. Clauset and C. Moore. Accuracy and scaling phenomena in internet mapping. *Physical Review Letters*, 94(1):018701, 2005.
- Z. Fan and L. Guan. Approximate  $\ell_0$ -penalized estimation of piecewise-constant signals on graphs. *The Annals of Statistics*, 46(6B):3217–3245, 2018.
- C. Gao, Z. Ma, A. Y. Zhang, and H. H. Zhou. Achieving optimal misclassification proportion in stochastic block models. *The Journal of Machine Learning Research*, 18(1):1980–2024, 2017.
- C. Gao, Z. Ma, A. Y. Zhang, and H. H. Zhou. Community detection in degree-corrected block models. *The Annals of Statistics*, 46(5):2153–2185, 2018.
- L. L. Gao, D. Witten, and J. Bien. Testing for association in multi-view network data. *arXiv preprint arXiv:1909.11640*, 2019.
- M. S. Handcock and K. J. Gile. Modeling social networks from sampled data. *The Annals of Applied Statistics*, 4(1):5, 2010.
- K. M. Harris. *The National Longitudinal Study of Adolescent to Adult Health (Add Health), Waves I & II, 1994-1996; Wave III, 2001-2002; Wave IV, 2007-009 [machine-readable data file and documentation]*. Carolina Population Center, University of North Carolina at Chapel Hill, 2009.
- P. W. Holland, K. B. Laskey, and S. Leinhardt. Stochastic blockmodels: First steps. *Social Networks*, 5(2):109–137, 1983.
- C.-S. Hsieh and L. F. Lee. A social interactions model with endogenous friendship formation and selectivity. *Journal of Applied Econometrics*, 31(2):301–319, 2016.
- M. O. Jackson and B. W. Rogers. Relating network structure to diffusion properties through stochastic dominance. *The BE Journal of Theoretical Economics*, 7(1), 2007.
- A. Javanmard and A. Montanari. Confidence intervals and hypothesis testing for high-dimensional regression. *The Journal of Machine Learning Research*, 15(1):2869–2909, 2014.
- P. Ji and J. Jin. Coauthorship and citation networks for statisticians. *The Annals of Applied Statistics*, 10(4):1779–1812, 2016.
- J. Jin, Z. T. Ke, S. Luo, and M. Wang. Estimating the number of communities by stepwise goodness-of-fit. *arXiv preprint arXiv:2009.09177*, 2020.
- B. Karrer and M. E. Newman. Stochastic blockmodels and community structure in networks. *Physical Review E*, 83(1):016107, 2011.
- M. Khabbazian, B. Hanlon, Z. Russek, K. Rohe, et al. Novel sampling design for respondent-driven sampling. *Electronic Journal of Statistics*, 11(2):4769–4812, 2017.
- E. D. Kolaczyk. *Statistical Analysis of Network Data: Methods and Models*. Springer Publishing Company, Incorporated, 1st edition, 2009. ISBN 038788145X, 9780387881454.
- A. Lakhina, J. W. Byers, M. Crovella, and P. Xie. Sampling biases in ip topology measurements. In *IEEE INFOCOM 2003. Twenty-second Annual Joint Conference of the IEEE Computer and Communications Societies (IEEE Cat. No. 03CH37428)*, volume 1, pages 332–341. IEEE, 2003.
- C. M. Le and E. Levina. Estimating the number of communities in networks by spectral methods. *arXiv preprint arXiv:1507.00827*, 2015.

- C. M. Le and E. Levina. Estimating a network from multiple noisy realizations. *arXiv preprint arXiv:1710.04765*, 2017.
- C. M. Le, E. Levina, and R. Vershynin. Concentration and regularization of random graphs. *Random Structures & Algorithms*, 2017.
- L.-f. Lee. Identification and estimation of econometric models with group interactions, contextual factors and fixed effects. *Journal of Econometrics*, 140(2):333–374, 2007.
- L.-f. Lee, X. Liu, and X. Lin. Specification and estimation of social interaction models with network structures. *The Econometrics Journal*, 13(2):145–176, 2010.
- J. Lei and A. Rinaldo. Consistency of spectral clustering in stochastic block models. *The Annals of Statistics*, 43(1):215–237, 2014.
- J. Lei and L. Zhu. Generic sample splitting for refined community recovery in degree corrected stochastic block models. *Statistica Sinica*, pages 1639–1659, 2017.
- L. Lei. Unified  $\ell_{2 \rightarrow \infty}$  eigenspace perturbation theory for symmetric random matrices. *arXiv preprint arXiv:1909.04798*, 2019.
- L. Lei, X. Li, and X. Lou. Consistency of spectral clustering on hierarchical stochastic block models. *arXiv preprint arXiv:2004.14531*, 2020.
- T. Li, E. Levina, and J. Zhu. Prediction models for network-linked data. *The Annals of Applied Statistics*, 13(1):132–164, 2019.
- T. Li, L. Lei, S. Bhattacharyya, K. Van den Berge, P. Sarkar, P. J. Bickel, and E. Levina. Hierarchical community detection by recursive partitioning. *Journal of the American Statistical Association*, pages 1–18, 2020a.
- T. Li, E. Levina, and J. Zhu. Network cross-validation by edge sampling. *Biometrika*, 107(2): 257–276, 2020b.
- T. Li, C. Qian, E. Levina, and J. Zhu. High-dimensional gaussian graphical models on network-linked data. *Journal of Machine Learning Research*, 21(74):1–45, 2020c. URL <http://jmlr.org/papers/v21/19-563.html>.
- S. Lunagómez, M. Papamichalis, P. J. Wolfe, and E. M. Airolidi. Evaluating and optimizing network sampling designs: Decision theory and information theory perspectives. *arXiv preprint arXiv:1811.07829*, 2018.
- E. Manresa. Estimating the structure of social interactions using panel data. *Unpublished Manuscript. CEMFI, Madrid*, 2013.
- C. F. Manski. Identification of endogenous social effects: The reflection problem. *The Review of Economic Studies*, 60(3):531–542, 1993.
- X. Mao, P. Sarkar, and D. Chakrabarti. Estimating mixed memberships with sharp eigenvector deviations. *Journal of the American Statistical Association*, (just-accepted):1–24, 2020.
- L. Michell and P. West. Peer pressure to smoke: the meaning depends on the method. *Health Education Research*, 11(1):39–49, 1996.
- M. Newman. Estimating network structure from unreliable measurements. *Physical Review E*, 98(6):062321, 2018.



- A. Y. Ng, A. X. Zheng, and M. I. Jordan. Link analysis, eigenvectors and stability. In *International Joint Conference on Artificial Intelligence*, volume 17, pages 903–910. Lawrence Erlbaum Associates Ltd, 2001.
- E. L. Ogburn. Challenges to estimating contagion effects from observational data. In *Complex Spreading Phenomena in Social Systems*, pages 47–64. Springer, 2018.
- E. L. Paluck, H. Shepherd, and P. M. Aronow. Changing climates of conflict: A social network experiment in 56 schools. *Proceedings of the National Academy of Sciences*, 113(3):566–571, 2016.
- M. Pearson and P. West. Drifting smoke rings. *Connections*, 25(2):59–76, 2003.
- K. Rohe et al. A critical threshold for design effects in network sampling. *The Annals of Statistics*, 47(1):556–582, 2019.
- T. Rolland, M. Taşan, B. Charloteaux, S. J. Pevzner, Q. Zhong, N. Sahni, S. Yi, I. Lemmens, C. Fontanillo, and R. Mosca. A proteome-scale map of the human interactome network. *Cell*, 159(5):1212–1226, 2014.
- V. Sadhanala, Y.-X. Wang, and R. J. Tibshirani. Graph sparsification approaches for laplacian smoothing. In *Proceedings of the 19th International Conference on Artificial Intelligence and Statistics*, pages 1250–1259, 2016.
- C. R. Shalizi and A. C. Thomas. Homophily and contagion are generically confounded in observational social network studies. *Sociological Methods and Research*, 40(2):211–239, 2011.
- C. D. Sherbourne. The role of social support and life stress events in use of mental health services. *Social science & medicine*, 27(12):1393–1400, 1988.
- J. Shi and J. Malik. Normalized cuts and image segmentation. *Pattern Analysis and Machine Intelligence, IEEE Transactions on*, 22(8):888–905, 2000.
- A. J. Smola and R. Kondor. Kernels and regularization on graphs. In *Learning theory and kernel machines*, pages 144–158. Springer, 2003.
- L. Su, W. Lu, R. Song, and D. Huang. Testing and estimation of social network dependence with time to event data. *Journal of the American Statistical Association*, 0(0):1–28, 2019. doi: 10.1080/01621459.2019.1617153. URL <https://doi.org/10.1080/01621459.2019.1617153>.
- M. Tang, D. L. Sussman, and C. E. Priebe. Universally consistent vertex classification for latent positions graphs. *The Annals of Statistics*, 41(3):1406–1430, 2013.
- S. Van de Geer, P. Bühlmann, Y. Ritov, and R. Dezeure. On asymptotically optimal confidence regions and tests for high-dimensional models. *The Annals of Statistics*, 42(3):1166–1202, 2014.
- R. Vershynin. *High-dimensional probability: An introduction with applications in data science*, volume 47. Cambridge University Press, 2018.
- Y.-X. Wang, J. Sharpnack, A. Smola, and R. J. Tibshirani. Trend filtering on graphs. *Journal of Machine Learning Research*, 17(105):1–41, 2016.
- J. E. Wildes, A. D. Simons, and K. L. Harkness. Life events, number of social relationships, and twelve-month naturalistic course of major depression in a community sample of women. *Depression and Anxiety*, 16(3):104–113, 2002.

- Y.-J. Wu, E. Levina, and J. Zhu. Link prediction for egocentrically sampled networks. *arXiv preprint arXiv:1803.04084*, 2018.
- D. Xia. Data-dependent confidence regions of singular subspaces. *arXiv preprint arXiv:1901.00304*, 2019.
- Y. Yu, T. Wang, and R. J. Samworth. A useful variant of the davis–kahan theorem for statisticians. *Biometrika*, 102(2):315–323, 2014.
- C.-H. Zhang and S. S. Zhang. Confidence intervals for low dimensional parameters in high dimensional linear models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 76(1):217–242, 2014.
- S. Zhao and A. Shojaie. A significance test for graph-constrained estimation. *Biometrics*, 72(2):484–493, 2016.
- X. Zhu, R. Pan, G. Li, Y. Liu, and H. Wang. Network vector autoregression. *The Annals of Statistics*, 45(3):1096–1123, 2017.

## A Basic tools and properties for generic inference

We start with the following lemma, which among other things allows us to rewrite  $\mathcal{P}_C$  and  $\mathcal{P}_N$  in a more convenient way for theoretical analysis.

**Lemma 1** (Properties of population projections). Let  $\mathcal{P}_C$  and  $\mathcal{P}_N$  be the matrices defined by (11) and (12), respectively. Then the following statements hold.

- (i)  $\mathcal{P}_C + \mathcal{P}_N = M(M^T M)^{-1} M^T$  is the orthogonal projection onto the column space of  $M$ .
- (ii)  $\mathcal{P}_C$  and  $\mathcal{P}_N$  can be written as

$$\mathcal{P}_C = \sum_{i=r+1}^{r+s} \frac{1}{1-\sigma_i^2} \tilde{Z}_i \tilde{Z}_i^T - \sum_{i=r+1}^{r+s} \frac{\sigma_i}{1-\sigma_i^2} \tilde{Z}_i \tilde{W}_i^T + \sum_{i=r+s+1}^p \tilde{Z}_i \tilde{Z}_i^T, \quad (35)$$

$$\mathcal{P}_N = \sum_{i=r+1}^{r+s} \frac{1}{1-\sigma_i^2} \tilde{W}_i \tilde{W}_i^T - \sum_{i=r+1}^{r+s} \frac{\sigma_i}{1-\sigma_i^2} \tilde{W}_i \tilde{Z}_i^T + \sum_{i=r+s+1}^K \tilde{W}_i \tilde{W}_i^T. \quad (36)$$

- (iii) For any  $x \in \text{col}(\tilde{Z}_{(r+1):p})$  and  $y \in \text{col}(\tilde{W}_{(r+1):K})$ , we have

$$\mathcal{P}_C(x+y) = x, \quad \mathcal{P}_N(x+y) = y.$$

- (iv) For any  $x \in \mathbb{R}^n$ , we have  $\|\mathcal{P}_C x\| \geq \|\tilde{Z}_{(r+1):p}^T x\|$ .

A crucial step in proving our results is to bound the differences  $\hat{\mathcal{P}}_C - \mathcal{P}_C$  and  $\hat{\mathcal{P}}_N - \mathcal{P}_N$ ; see (17), (18), (11) and (12) for the definition of these matrices. Since they depend on  $M$  and  $\hat{M}$ , it is tempting to bound  $\hat{M} - M$  first and then use that bound to show that  $\hat{\mathcal{P}}_C - \mathcal{P}_C$  and  $\hat{\mathcal{P}}_N - \mathcal{P}_N$  are small in magnitude. Unfortunately,  $\hat{M} - M$  may not be small because in general singular vectors are unique only up to multiplying by a certain orthogonal matrix. Nevertheless, we will still be able to show that  $\hat{\mathcal{P}}_C - \mathcal{P}_C$  and  $\hat{\mathcal{P}}_N - \mathcal{P}_N$  are small by leveraging the projection-type properties of  $\hat{\mathcal{P}}_C$ ,  $\mathcal{P}_C$ ,  $\hat{\mathcal{P}}_N$  and  $\mathcal{P}_N$ , although the fact that they may not be orthogonal projections makes the analysis much more involved.

In light of Lemma 1, it is convenient to introduce the following notations:

$$\tilde{\mathcal{P}}_C = \sum_{i=r+1}^{r+s} \frac{1}{1-\hat{\sigma}_i^2} \hat{Z}_i \hat{Z}_i^T - \sum_{i=r+1}^{r+s} \frac{\hat{\sigma}_i}{1-\hat{\sigma}_i^2} \hat{Z}_i \tilde{W}_i^T + \sum_{i=r+s+1}^p \hat{Z}_i \hat{Z}_i^T, \quad (37)$$

$$\tilde{\mathcal{P}}_N = \sum_{i=r+1}^{r+s} \frac{1}{1-\hat{\sigma}_i^2} \tilde{W}_i \tilde{W}_i^T - \sum_{i=r+1}^{r+s} \frac{\hat{\sigma}_i}{1-\hat{\sigma}_i^2} \tilde{W}_i \hat{Z}_i^T + \sum_{i=r+s+1}^K \tilde{W}_i \tilde{W}_i^T. \quad (38)$$

The following lemma bound  $\hat{\mathcal{P}}_C - \tilde{\mathcal{P}}_C$  and  $\hat{\mathcal{P}}_N - \tilde{\mathcal{P}}_N$ .

**Lemma 2** (Perturbation of projections, first step). If condition (26) holds then

$$\max\{\|\hat{\mathcal{P}}_C - \tilde{\mathcal{P}}_C\|, \|\hat{\mathcal{P}}_N - \tilde{\mathcal{P}}_N\|\} \leq \frac{16\tau_n}{(1-\sigma_{r+1})^2}.$$

We will bound the differences  $\hat{\mathcal{P}}_C - \mathcal{P}_C$  and  $\hat{\mathcal{P}}_N - \mathcal{P}_N$  by bounding  $\tilde{\mathcal{P}}_C - \mathcal{P}_C$  and  $\tilde{\mathcal{P}}_N - \mathcal{P}_N$ , and

then combining them with Lemma 2. For that purpose, denote

$$\delta_1 = \sum_{i=r+1}^{r+s} \frac{1}{1-\hat{\sigma}_i^2} \hat{Z}_i \hat{Z}_i^T - \sum_{i=r+1}^{r+s} \frac{1}{1-\sigma_i^2} \tilde{Z}_i \tilde{Z}_i^T, \quad (39)$$

$$\delta_2 = \sum_{i=r+1}^{r+s} \frac{\hat{\sigma}_i}{1-\hat{\sigma}_i^2} \hat{Z}_i \check{W}_i^T - \sum_{i=r+1}^{r+s} \frac{\sigma_i}{1-\sigma_i^2} \tilde{Z}_i \check{W}_i^T, \quad (40)$$

$$\delta_3 = \sum_{i=r+s+1}^p \hat{Z}_i \hat{Z}_i^T - \sum_{i=r+s+1}^p \tilde{Z}_i \tilde{Z}_i^T, \quad (41)$$

$$\delta_4 = \sum_{i=r+1}^{r+s} \frac{1}{1-\hat{\sigma}_i^2} \check{W}_i \check{W}_i^T - \sum_{i=r+1}^{r+s} \frac{1}{1-\sigma_i^2} \tilde{W}_i \tilde{W}_i^T, \quad (42)$$

$$\delta_5 = \sum_{i=r+1}^{r+s} \frac{\hat{\sigma}_i}{1-\hat{\sigma}_i^2} \check{W}_i \hat{Z}_i^T - \sum_{i=r+1}^{r+s} \frac{\sigma_i}{1-\sigma_i^2} \tilde{W}_i \tilde{Z}_i^T, \quad (43)$$

$$\delta_6 = \sum_{i=r+s+1}^K \check{W}_i \check{W}_i^T - \sum_{i=r+s+1}^K \tilde{W}_i \tilde{W}_i^T. \quad (44)$$

By Lemma 1,  $\tilde{\mathcal{P}}_C - \mathcal{P}_C = \delta_1 - \delta_2 + \delta_3$  and  $\tilde{\mathcal{P}}_N - \mathcal{P}_N = \delta_4 - \delta_5 + \delta_6$ . The following lemma provides upper bounds for all  $\|\delta_i\|$ .

**Lemma 3** (Perturbation of projections, second step). If condition (26) holds then

$$\max_{1 \leq i \leq 6} \|\delta_i\| \leq \frac{60\tau_n}{\min\{(1-\sigma_{r+1})^3, \sigma_{r+s}^3\}}.$$

The proofs of Lemmas 1, 2 and 3 are postponed to later in this section. As a direct consequence of these results and the triangle inequality, we have the following bound.

**Corollary 5** (Perturbation of projections). If condition (26) holds then there exists a constant  $C > 0$  such that

$$\max\{\|\hat{\mathcal{P}}_C - \mathcal{P}_C\|, \|\hat{\mathcal{P}}_N - \mathcal{P}_N\|\} \leq \frac{C\tau_n}{\min\{(1-\sigma_{r+1})^3, \sigma_{r+s}^3\}}.$$

We are now ready to prove Proposition 3 about the bias of  $\hat{\beta}$ .

*Proof of Proposition 3.* Recall the estimate  $\hat{\beta}$  in (19), identity of  $\beta$  in (13), and the model  $Y = X\beta + X\theta + \alpha + \epsilon$  from (4). Taking the expectation with respect to the randomness of the noise  $\epsilon$ , we get

$$\mathbb{E}\hat{\beta} = (X^T X)^{-1} X^T \hat{\mathcal{P}}_C (X\beta + X\theta + \alpha). \quad (45)$$

By (13), (45) and Corollary 5, we have

$$\begin{aligned} \|\mathbb{E}\hat{\beta} - \beta\| &= \|(X^T X)^{-1} X^T (\hat{\mathcal{P}}_C - \mathcal{P}_C) (X\beta + X\theta + \alpha)\| \\ &\leq \|(X^T X)^{-1}\| \cdot \|X\| \cdot \|X\beta + X\theta + \alpha\| \cdot \|\hat{\mathcal{P}}_C - \mathcal{P}_C\| \\ &\leq \|(X^T X)^{-1}\| \cdot \|X\| \cdot \|X\beta + X\theta + \alpha\| \cdot \frac{C\tau_n}{(\min\{1-\sigma_{r+1}, \sigma_{r+s}\})^3}. \end{aligned}$$

The proof is complete.  $\square$

Next, we prove Proposition 4 about the consistency of  $\hat{\alpha}$  and the bias and variance of  $\hat{\gamma}$ .

*Proof of Proposition 4. Bounding  $\|\hat{\alpha} - \alpha\|$ .* From (13), (19) and (4) we have

$$\hat{\alpha} - \alpha = (\hat{\mathcal{P}}_N - \mathcal{P}_N)(X\beta + X\theta + \alpha + \epsilon) + \mathcal{P}_N\epsilon.$$

By Assumption A1 and the fact that  $\|\epsilon\| = O(\sqrt{n\sigma^2})$  with high probability,

$$\|X\beta + X\theta + \alpha + \epsilon\| \leq C\sqrt{n(p + \sigma^2)}.$$

Using the triangle inequality and Corollary 5, we get

$$\begin{aligned} \|\hat{\alpha} - \alpha\| &\leq \|\hat{\mathcal{P}}_N - \mathcal{P}_N\| \cdot \|X\beta + X\theta + \alpha + \epsilon\| + \|\mathcal{P}_N\epsilon\| \\ &\leq \frac{C\tau_n\sqrt{n(p + \sigma^2)}}{\min\{(1 - \sigma_{r+1})^3, \sigma_{r+s}^3\}} + \|\mathcal{P}_N\epsilon\|. \end{aligned} \quad (46)$$

To bound  $\|\mathcal{P}_N\epsilon\|$ , we first use (8) to rewrite  $\tilde{W}_i$  as  $\tilde{W}_i = WV_i$  for  $i \leq K$ . Then from (12), we can rewrite  $\mathcal{P}_N$  as follows:

$$\mathcal{P}_N = W \left( \sum_{i=r+1}^{r+s} \frac{1}{1 - \sigma_i^2} V_i \tilde{W}_i^T - \sum_{i=r+1}^{r+s} \frac{\sigma_i}{1 - \sigma_i^2} V_i \tilde{Z}_i^T + \sum_{i=r+s+1}^K V_i \tilde{W}_i^T \right) =: WQ,$$

where  $Q$  is a fixed matrix of rank at most  $2K$ . Moreover, for any unit vector  $x$ ,

$$\begin{aligned} \|Qx\| &\leq \left\| \sum_{i=r+1}^{r+s} \frac{1}{1 - \sigma_i^2} V_i \tilde{W}_i^T x + \sum_{i=r+s+1}^K V_i \tilde{W}_i^T x \right\| + \left\| \sum_{i=r+1}^{r+s} \frac{\sigma_i}{1 - \sigma_i^2} V_i \tilde{Z}_i^T x \right\| \\ &= \left( \sum_{i=r+1}^{r+s} \frac{(\tilde{W}_i^T x)^2}{(1 - \sigma_i^2)^2} + \sum_{i=r+s+1}^K (\tilde{W}_i^T x)^2 \right)^{1/2} + \left( \sum_{i=r+1}^{r+s} \frac{\sigma_i^2 (\tilde{Z}_i^T x)^2}{(1 - \sigma_i^2)^2} \right)^{1/2} \\ &\leq \frac{\|\tilde{W}^T x\|}{1 - \sigma_{r+1}^2} + \frac{\|\tilde{Z} x\|}{1 - \sigma_{r+1}^2} \\ &\leq \frac{2}{1 - \sigma_{r+1}^2}. \end{aligned}$$

Thus,  $\|Q\| \leq 2/(1 - \sigma_{r+1}^2)$  and  $Q\epsilon$  is a Gaussian vector of dimension at most  $2K$  and covariance matrix with spectral norm at most  $4\sigma^2/(1 - \sigma_{r+1}^2)^2$ . Therefore with high probability

$$\|Q\epsilon\| \leq \frac{C\sqrt{K\sigma^2} \cdot \log n}{1 - \sigma_{r+1}^2}.$$

In particular,

$$\|\mathcal{P}_N\epsilon\| \leq \|W\|_{2 \rightarrow \infty} \cdot \|Q\epsilon\| \leq \|W\|_{2 \rightarrow \infty} \cdot \frac{C\sqrt{K\sigma^2} \cdot \log n}{1 - \sigma_{r+1}^2}.$$

Together with (46), this implies the required bound for  $\|\hat{\alpha} - \alpha\|$ .

**Bounding the bias of  $\hat{\gamma}$ .** By (13),

$$\gamma = \tilde{W}_{(r+1):K}^T \alpha = \tilde{W}_{(r+1):K}^T \mathcal{P}_N (X\beta + X\theta + \alpha).$$

Similarly, by (22), (4) and taking the expectation with respect to the randomness of the noise  $\epsilon$ , we have

$$\mathbb{E}\hat{\gamma} = \check{W}_{(r+1):K} \hat{\mathcal{P}}_N \mathbb{E}Y = \check{W}_{(r+1):K} \hat{\mathcal{P}}_N (X\beta + X\theta + \alpha).$$

Let  $O = \check{W}_{(r+1):K}^T \tilde{W}_{(r+1):K}$ , then

$$\mathbb{E}\hat{\gamma} - O\gamma = \left( \check{W}_{(r+1):K}^T \hat{\mathcal{P}}_N - O \tilde{W}_{(r+1):K}^T \mathcal{P}_N \right) (X\beta + X\theta + \alpha). \quad (47)$$

To bound this bias, we first replace  $\hat{\mathcal{P}}_N$  with  $\mathcal{P}_N$  and then estimate the resulting expression. By Corollary 5, there exists a constant  $C > 0$  such that

$$\|\hat{\mathcal{P}}_N - \mathcal{P}_N\| \leq \frac{C\tau_n}{\min\{(1 - \sigma_{r+1})^3, \sigma_{r+s}^3\}}.$$

Together with Assumption A1, this implies

$$\|\check{W}_{(r+1):K}^T (\hat{\mathcal{P}}_N - \mathcal{P}_N) (X\beta + X\theta + \alpha)\| \leq \frac{C\tau_n \sqrt{n\bar{p}}}{\min\{(1 - \sigma_{r+1})^3, \sigma_{r+s}^3\}}.$$

Therefore from (47), (13) and the triangle inequality,

$$\begin{aligned} \|\mathbb{E}\hat{\gamma} - O\gamma\| &\leq \left\| \left( \check{W}_{(r+1):K}^T - O \tilde{W}_{(r+1):K}^T \right) \mathcal{P}_N (X\beta + X\theta + \alpha) \right\| + \frac{C\tau_n \sqrt{n\bar{p}}}{\min\{(1 - \sigma_{r+1})^3, \sigma_{r+s}^3\}} \\ &= \left\| \left( \check{W}_{(r+1):K}^T - O \tilde{W}_{(r+1):K}^T \right) \alpha \right\| + \frac{C\tau_n \sqrt{n\bar{p}}}{\min\{(1 - \sigma_{r+1})^3, \sigma_{r+s}^3\}} \\ &= \left\| \check{W}_{(r+1):K}^T \left( I - \tilde{W}_{(r+1):K} \tilde{W}_{(r+1):K}^T \right) \alpha \right\| + \frac{C\tau_n \sqrt{n\bar{p}}}{\min\{(1 - \sigma_{r+1})^3, \sigma_{r+s}^3\}} \\ &= \frac{C\tau_n \sqrt{n\bar{p}}}{\min\{(1 - \sigma_{r+1})^3, \sigma_{r+s}^3\}}. \end{aligned}$$

The last equality holds because  $\alpha \in \text{col}(\tilde{W}_{(r+1):K})$  by (9) and  $\tilde{W}_{(r+1):K} \tilde{W}_{(r+1):K}^T$  is the orthogonal projection onto  $\text{col}(\tilde{W}_{(r+1):K})$ .

**Bounding the covariance of  $\hat{\gamma}$ .** Recall  $\hat{\gamma} = \check{W}_{(r+1):K} \hat{\mathcal{P}}_N Y$  from (22). Since  $Y = X\beta + X\theta + \alpha + \epsilon$ , it follows that

$$\Sigma_{\hat{\gamma}} = \sigma^2 \check{W}_{(r+1):K}^T \hat{\mathcal{P}}_N \hat{\mathcal{P}}_N^T \check{W}_{(r+1):K}.$$

Define  $\Gamma = \text{diag}(\sigma_{r+1}, \dots, \sigma_{r+s})$  and  $\hat{\Gamma} = \text{diag}(\hat{\sigma}_{r+1}, \dots, \hat{\sigma}_{r+s})$ . A direct calculation shows that

$$\check{W}_{(r+1):K}^T \hat{\mathcal{P}}_N = \begin{pmatrix} (1 - \hat{\Gamma}^2)^{-1} \check{W}_{(r+1):(r+s)}^T & -\hat{\Gamma}(1 - \hat{\Gamma}^2)^{-1} \hat{Z}_{(r+1):(r+s)}^T \\ & \check{W}_{(r+s+1):K}^T \end{pmatrix}.$$

Therefore

$$\Sigma_{\hat{\gamma}} = \sigma^2 \cdot \begin{pmatrix} (1 - \hat{\Gamma}^2)^{-1} & & & \hat{\Gamma}(1 - \hat{\Gamma}^2)^{-1} \check{Z}_{(r+1):(r+s)}^T \check{W}_{(r+s+1):K} \\ \check{W}_{(r+s+1):K}^T \check{Z}_{(r+1):(r+s)} \hat{\Gamma} (1 - \hat{\Gamma}^2)^{-1} & & & \\ & & & I_{K-r-s} \end{pmatrix}.$$

Since  $Z^T \hat{W} = \hat{U} \hat{\Sigma} \hat{V}$  by (14), it follows from (15) that  $\hat{Z}^T \check{W} = \hat{\Sigma}$ , and therefore

$$\hat{Z}_{(r+1):(r+s)}^T \check{W}_{(r+s+1):K} = 0$$

because the left-hand side is an off-diagonal block of  $\hat{\Sigma}$ . Moreover, by (26) and inequality (56) of Lemma 4,

$$\|(1 - \hat{\Gamma}^2)^{-1} - (1 - \Gamma^2)^{-1}\| \leq \frac{4\tau_n}{\min\{(1 - \sigma_{r+1})^2, \sigma_{r+s}^2\}},$$

and the bound for  $\Sigma_{\hat{\gamma}}$  follows.  $\square$

*Proof of Lemma 1.* Since

$$M = (\tilde{Z}_{(r+1):p}, 0_{n \times (K-r)}) + (0_{n \times (p-r)}, \tilde{W}_{(r+1):K}),$$

it follows directly from (11) and (12) that  $\mathcal{P}_C + \mathcal{P}_N = M(M^T M)^{-1} M$ , and (i) is proved.

Next, we verify (35) and (36) in (ii). Recall from (7) that  $Z^T W = U \Sigma V^T$  and the nonzero singular values of  $Z^T W$  are

$$\sigma_1 = \sigma_2 = \dots = \sigma_r = 1 > \sigma_{r+1} \geq \dots \geq \sigma_{r+s} > 0.$$

Multiplying both sides of  $Z^T W = U \Sigma V^T$  with  $U^T$  on the left and  $V$  on the right, and then using (8), we get

$$\tilde{Z}^T \tilde{W} = \Sigma = \begin{pmatrix} I_r & 0_{r \times s} & 0_{r \times (K-r-s)} \\ 0_{s \times r} & \Gamma & 0_{s \times (K-r-s)} \\ 0_{(p-r-s) \times r} & 0_{(p-r-s) \times s} & 0_{(p-r-s) \times (K-r-s)} \end{pmatrix},$$

where

$$\Gamma = \text{diag}(\sigma_{r+1}, \dots, \sigma_{r+s}). \quad (48)$$

It follows that

$$\tilde{Z}_{(r+1):p}^T \tilde{W}_{(r+1):K} = \begin{pmatrix} \Gamma & 0_{s \times (K-r-s)} \\ 0_{(p-r-s) \times s} & 0_{(p-r-s) \times (K-r-s)} \end{pmatrix}. \quad (49)$$

Since  $M = (\tilde{Z}_{(r+1):p}, \tilde{W}_{(r+1):K})$ , the above identity implies

$$M^T M = \begin{pmatrix} I_s & 0 & \Gamma & 0 \\ 0 & I_{p-r-s} & 0 & 0 \\ \Gamma & 0 & I_s & 0 \\ 0 & 0 & 0 & I_{K-r-s} \end{pmatrix}, \quad (50)$$

$$(M^T M)^{-1} = \begin{pmatrix} (I - \Gamma^2)^{-1} & 0 & -(I - \Gamma^2)^{-1} \Gamma & 0 \\ 0 & I_{p-r-s} & 0 & 0 \\ -(I - \Gamma^2)^{-1} \Gamma & 0 & (I - \Gamma^2)^{-1} & 0 \\ 0 & 0 & 0 & I_{K-r-s} \end{pmatrix}. \quad (51)$$

Therefore

$$\begin{aligned}
(M^T M)^{-1} M^T &= \begin{pmatrix} (I - \Gamma^2)^{-1} & 0 & -(I - \Gamma^2)^{-1} \Gamma & 0 \\ 0 & I_{p-r-s} & 0 & 0 \\ -(I - \Gamma^2)^{-1} \Gamma & 0 & (I - \Gamma^2)^{-1} & 0 \\ 0 & 0 & 0 & I_{K-r-s} \end{pmatrix} \begin{pmatrix} \tilde{Z}_{(r+1):(r+s)}^T \\ \tilde{Z}_{(r+s+1):p}^T \\ \tilde{W}_{(r+1):(r+s)}^T \\ \tilde{W}_{(r+s+1):K}^T \end{pmatrix} \\
&= \begin{pmatrix} (I - \Gamma^2)^{-1} \tilde{Z}_{(r+1):(r+s)}^T - (I - \Gamma^2)^{-1} \Gamma \tilde{W}_{(r+1):(r+s)}^T \\ \tilde{Z}_{(r+s+1):p}^T \\ -(I - \Gamma^2)^{-1} \Gamma \tilde{Z}_{(r+1):(r+s)}^T + (I - \Gamma^2)^{-1} \tilde{W}_{(r+1):(r+s)}^T \\ \tilde{W}_{(r+s+1):K}^T \end{pmatrix}.
\end{aligned}$$

Multiplying this matrix on the left by either  $(\tilde{Z}_{(r+1):p}, 0_{n \times (K-r)})$  or  $(0_{n \times (p-r)}, \tilde{W}_{(r+1):K})$ , we see that (35) and (36) hold and (ii) is proved.

For (iii), we will show that  $\mathcal{P}_C(x + y) = x$ ; the identity  $\mathcal{P}_N(x + y) = y$  can be proved by using the same argument. Since  $x \in \text{col}(\tilde{Z}_{(r+1):p})$  and  $y \in \text{col}(\tilde{W}_{(r+1):K})$ , it is enough to show that the identity holds if  $x$  is any column vector of  $\tilde{Z}_{(r+1):p}$  and  $y$  is any column vector of  $\tilde{W}_{(r+1):K}$ . Therefore, assume that  $x = \tilde{Z}_{i'}$  and  $y = \tilde{W}_{j'}$  for some  $r + 1 \leq i' \leq p$  and  $r + 1 \leq j' \leq K$ . If  $r + 1 \leq i' \leq r + s$  then by (49),

$$\begin{aligned}
\mathcal{P}_C x &= \sum_{i=r+1}^{r+s} \frac{1}{1 - \sigma_i^2} \tilde{Z}_i \tilde{Z}_i^T \tilde{Z}_{i'} - \sum_{i=r+1}^{r+s} \frac{\sigma_i}{1 - \sigma_i^2} \tilde{Z}_i \tilde{W}_i^T \tilde{Z}_{i'} + \sum_{i=r+s+1}^p \tilde{Z}_i \tilde{Z}_i^T \tilde{Z}_{i'} \\
&= \frac{1}{1 - \sigma_{i'}^2} \tilde{Z}_{i'} - \frac{\sigma_{i'}^2}{1 - \sigma_{i'}^2} \tilde{Z}_{i'} \\
&= x.
\end{aligned}$$

If  $i' \geq r + s + 1$  then again by (49),

$$\mathcal{P}_C x = \sum_{i=r+1}^{r+s} \frac{1}{1 - \sigma_i^2} \tilde{Z}_i \tilde{Z}_i^T \tilde{Z}_{i'} - \sum_{i=r+1}^{r+s} \frac{\sigma_i}{1 - \sigma_i^2} \tilde{Z}_i \tilde{W}_i^T \tilde{Z}_{i'} + \sum_{i=r+s+1}^p \tilde{Z}_i \tilde{Z}_i^T \tilde{Z}_{i'} = \tilde{Z}_{i'} = x.$$

We have shown that  $\mathcal{P}_C x = x$  for all  $x \in \text{col}(\tilde{Z}_{(r+1):p})$ . Now, if  $y = \tilde{W}_{j'}$  and  $r + 1 \leq j' \leq r + s$  then by (49),

$$\begin{aligned}
\mathcal{P}_C y &= \sum_{i=r+1}^{r+s} \frac{1}{1 - \sigma_i^2} \tilde{Z}_i \tilde{Z}_i^T \tilde{W}_{j'} - \sum_{i=r+1}^{r+s} \frac{\sigma_i}{1 - \sigma_i^2} \tilde{Z}_i \tilde{W}_i^T \tilde{W}_{j'} + \sum_{i=r+s+1}^p \tilde{Z}_i \tilde{Z}_i^T \tilde{W}_{j'} \\
&= \frac{\sigma_{j'}}{1 - \sigma_{j'}^2} \tilde{Z}_{j'} - \frac{\sigma_{j'}}{1 - \sigma_{j'}^2} \tilde{Z}_{j'} \\
&= 0.
\end{aligned}$$

If  $j' \geq r + s + 1$  then also by (49),

$$\mathcal{P}_C y = \sum_{i=r+1}^{r+s} \frac{1}{1 - \sigma_i^2} \tilde{Z}_i \tilde{Z}_i^T \tilde{W}_{j'} - \sum_{i=r+1}^{r+s} \frac{\sigma_i}{1 - \sigma_i^2} \tilde{Z}_i \tilde{W}_i^T \tilde{W}_{j'} + \sum_{i=r+s+1}^p \tilde{Z}_i \tilde{Z}_i^T \tilde{W}_{j'} = 0.$$

Thus,  $\mathcal{P}_C y = 0$  for all  $y \in \text{col}(\tilde{W}_{(r+1):K})$  and (iii) is proved.



It remains to show (iv). By (35) we have

$$\begin{aligned}\mathcal{P}_C^T x &= \sum_{i=r+1}^{r+s} \frac{1}{1-\sigma_i^2} \tilde{Z}_i \tilde{Z}_i^T x - \sum_{i=r+1}^{r+s} \frac{\sigma_i}{1-\sigma_i^2} \tilde{W}_i \tilde{Z}_i^T x + \sum_{i=r+s+1}^p \tilde{Z}_i \tilde{Z}_i^T x \\ &= \sum_{i=r+1}^{r+s} \left( \frac{\tilde{Z}_i^T x}{1-\sigma_i^2} \cdot \tilde{Z}_i - \frac{\sigma_i \tilde{Z}_i^T x}{1-\sigma_i^2} \cdot \tilde{W}_i \right) + \sum_{i=r+s+1}^p \tilde{Z}_i^T x \cdot \tilde{Z}_i.\end{aligned}$$

By (8), we have  $\tilde{Z}^T \tilde{W} = \Sigma$ . This implies that  $\tilde{Z}_{r+s+1}, \dots, \tilde{Z}_p$  are orthogonal to  $\tilde{Z}_{r+1}, \dots, \tilde{Z}_{r+s}$  and  $\tilde{W}_{r+1}, \dots, \tilde{W}_{r+s}$ . Moreover,  $\tilde{Z}_i$  and  $\tilde{W}_j$  are orthogonal for any pair of  $i$  and  $j$  such that  $i \neq j$  and  $r+1 \leq i, j \leq r+s$ . Also,  $\tilde{Z}_i^T \tilde{W}_i = \sigma_i$  if  $r+1 \leq i \leq r+s$ . Therefore,

$$\begin{aligned}\|\mathcal{P}_C^T x\|^2 &= \left\| \sum_{i=r+1}^{r+s} \left( \frac{\tilde{Z}_i^T x}{1-\sigma_i^2} \cdot \tilde{Z}_i - \frac{\sigma_i \tilde{Z}_i^T x}{1-\sigma_i^2} \cdot \tilde{W}_i \right) \right\|^2 + \left\| \sum_{i=r+s+1}^p \tilde{Z}_i^T x \cdot \tilde{Z}_i \right\|^2 \\ &= \sum_{i=r+1}^{r+s} \left( \frac{(\tilde{Z}_i^T x)^2}{(1-\sigma_i^2)^2} + \frac{\sigma_i^2 (\tilde{Z}_i^T x)^2}{(1-\sigma_i^2)^2} - 2 \frac{\sigma_i^2 (\tilde{Z}_i^T x)^2}{(1-\sigma_i^2)^2} \right) + \sum_{i=r+s+1}^p (\tilde{Z}_i^T x)^2 \\ &= \sum_{i=r+1}^{r+s} \frac{(\tilde{Z}_i^T x)^2}{1-\sigma_i^2} + \sum_{i=r+s+1}^p (\tilde{Z}_i^T x)^2 \\ &\geq \|\tilde{Z}_{(r+1):p} x\|^2,\end{aligned}$$

and (iv) is proved.  $\square$

*Proof of Lemma 2.* According to (14) and (15), the singular value decomposition of  $\hat{Z}^T \check{W}$  admits the following form:

$$\hat{Z}^T \check{W} = \hat{\Sigma} = \begin{pmatrix} \hat{I} & 0 & 0 \\ 0 & \hat{\Gamma} & 0 \\ 0 & 0 & \hat{\Phi} \end{pmatrix} \in \mathbb{R}^{p \times K},$$

where  $\hat{I} \in \mathbb{R}^{r \times r}$  and  $\hat{\Gamma} \in \mathbb{R}^{s \times s}$  are square diagonal matrices, and  $\hat{\Phi} \in \mathbb{R}^{(p-r-s) \times (K-r-s)}$  is a rectangular diagonal matrix. It then follows from the definition of  $\hat{M}$  in Algorithm 1 that

$$\hat{M}^T \hat{M} = \begin{pmatrix} I_s & 0 & \hat{\Gamma} & 0 \\ 0 & I_{p-r-s} & 0 & \hat{\Phi} \\ \hat{\Gamma} & 0 & I_s & 0 \\ 0 & \hat{\Phi}^T & 0 & I_{K-r-s} \end{pmatrix}.$$

Let  $Q$  be the matrix obtained from  $\hat{M}^T \hat{M}$  by replacing  $\hat{\Phi}$  and  $\hat{\Phi}^T$  with zero matrices of the same dimensions. Using (50) and (25), we have

$$\max\{\|Q - \hat{M}^T \hat{M}\|, \|\hat{M}^T \hat{M} - M^T M\|, \|Q - M^T M\|\} \leq \|\hat{\Gamma} - \Gamma\| + \|\hat{\Phi}\| \leq 2\|\hat{\Sigma} - \Sigma\| \leq 2\tau_n.$$

Also, from (50) and a direct calculation we see that  $M^T M$  has  $p + K - 2r$  eigenvalues, of which  $p + K - 2r - 2s$  are identically one and the remaining  $2s$  eigenvalues are  $1 \pm \sigma_i$  where  $\sigma_i$  are diagonal elements of  $\Gamma$  and  $r+1 \leq i \leq r+s$ . Therefore by Weyl's inequality,

$$1 - \sigma_{r+1} - 2\tau_n \leq \|\hat{M}^T \hat{M}\|, \|Q\| \leq 1 + \sigma_{r+1} + 2\tau_n,$$

which implies that both  $(\hat{M}^T \hat{M})^{-1}$  and  $Q^{-1}$  exist by condition (26). A direct calculation shows

that

$$Q^{-1} = \begin{pmatrix} (I - \hat{\Gamma}^2)^{-1} & 0 & -(I - \hat{\Gamma}^2)^{-1}\hat{\Gamma} & 0 \\ 0 & I_{p-r-s} & 0 & 0 \\ -(I - \hat{\Gamma}^2)^{-1}\hat{\Gamma} & 0 & (I - \hat{\Gamma}^2)^{-1} & 0 \\ 0 & 0 & 0 & I_{K-r-s} \end{pmatrix}.$$

Moreover,  $\tilde{\mathcal{P}}_C$  and  $\tilde{\mathcal{P}}_N$  defined in (37) and (38), respectively, can be written as follows:

$$\tilde{\mathcal{P}}_C = (\hat{Z}_{(r+1):p}, 0_{n \times (K-r)})Q^{-1}\hat{M}^T, \quad (52)$$

$$\tilde{\mathcal{P}}_N = (0_{n \times (p-r)}, \check{W}_{(r+1):K})Q^{-1}\hat{M}^T. \quad (53)$$

Therefore by (26),

$$\begin{aligned} \|(\hat{M}^T\hat{M})^{-1} - Q^{-1}\| &\leq \|\hat{M}^T\hat{M} - Q\| \cdot \|(\hat{M}^T\hat{M})^{-1}\| \cdot \|Q^{-1}\| \\ &\leq \frac{2\tau_n}{(1 - \sigma_{r+1} - 2\tau_n)^2} \\ &\leq \frac{4\tau_n}{(1 - \sigma_{r+1})^2}. \end{aligned}$$

The claim of the lemma then follows from (17), (18), (52), (53), the inequality above and the fact that  $\|\hat{M}\| \leq 2$ .  $\square$

To prove Lemma 3, we need the following two lemmas, which will be proved at the end of this section.

**Lemma 4** (Perturbation of singular vectors and partial singular decomposition). Let  $X, \hat{X}$  be  $p \times k$  matrices with singular decompositions

$$X = U\Sigma V^T, \quad \hat{X} = \hat{U}\hat{\Sigma}\hat{V}^T.$$

Let  $\eta$  be a set of singular values of  $X$ ,  $\eta^c$  be the set of remaining singular values of  $X$  and  $\Delta = \min_{\sigma \in \eta, \sigma' \in \eta^c} |\sigma - \sigma'|$ . Denote by  $U_\eta$  the matrix whose columns are left singular vectors of  $X$  with corresponding singular values in  $\eta$ ; define  $V_\eta, \hat{U}_\eta$  and  $\hat{V}_\eta$  similarly. Also, let  $\Gamma$  and  $\hat{\Gamma}$  be diagonal matrices with  $\{\sigma_i, i \in \eta\}$  and  $\{\hat{\sigma}_i, i \in \eta\}$  on the diagonal, respectively. Assume that  $\|X\| \leq 1$ ,  $\|\hat{X}\| \leq 1$  and  $\Delta > 4\|\hat{X} - X\|$ . Then

$$\max \left\{ \|\hat{U}_\eta\hat{U}_\eta^T - U_\eta U_\eta^T\|, \|\hat{V}_\eta\hat{V}_\eta^T - V_\eta V_\eta^T\| \right\} \leq \frac{2\|\hat{X} - X\|}{\Delta - 4\|\hat{X} - X\|}, \quad (54)$$

$$\left\| \sum_{i \in \eta} \sigma_i U_{\cdot i} V_{\cdot i}^T - \sum_{i \in \eta} \hat{\sigma}_i \hat{U}_{\cdot i} \hat{V}_{\cdot i}^T \right\| \leq \frac{3\|\hat{X} - X\|}{\Delta - 4\|\hat{X} - X\|}. \quad (55)$$

Furthermore, if  $1 - \max_{i \in \eta} \sigma_i^2 > 2\|\hat{X} - X\|$ , we have

$$\|(1 - \hat{\Gamma}^2)^{-1} - (1 - \Gamma^2)^{-1}\| \leq \frac{2\|\hat{X} - X\|}{(1 - \max_{i \in \eta} \sigma_i^2 - 2\|\hat{X} - X\|)^2}, \quad (56)$$

and

$$\begin{aligned} & \max \left\{ \|\hat{U}_\eta(1 - \hat{\Gamma}^2)^{-1}\hat{U}_\eta^T - U_\eta(1 - \Gamma^2)^{-1}U_\eta^T\|, \|\hat{V}_\eta(1 - \hat{\Gamma}^2)^{-1}\hat{V}_\eta^T - V_\eta(1 - \Gamma^2)^{-1}V_\eta^T\| \right\} \\ & \leq \frac{6\|\hat{X} - X\|}{(1 - \max_{i \in \eta} \sigma_i^2 - 2\|\hat{X} - X\|)^2(\Delta - 4\|\hat{X} - X\|)}. \end{aligned} \quad (57)$$

**Lemma 5** (Unorthogonal projection bound). Let  $X, \hat{X}$  be two matrices of the same size with singular decompositions

$$X = U\Sigma V^T, \quad \hat{X} = \hat{U}\hat{\Sigma}\hat{V}^T.$$

Assume that  $\|X\| < 1$ ,  $\|\hat{X}\| < 1$  and  $\|X\|^2 + 2\|\hat{X} - X\| < 1$ . Then

$$\left\| U\Sigma(1 - \Sigma^2)^{-1}V^T - \hat{U}\hat{\Sigma}(1 - \hat{\Sigma}^2)^{-1}\hat{V}^T \right\| \leq \frac{3\|X - \hat{X}\|}{(1 - \|X\|^2 - 2\|\hat{X} - X\|)^2}.$$

*Proof of Lemma 3.* Recall that (23) provides a bound on the difference between two matrices  $Z^T\hat{W}\hat{W}^T$  and  $Z^TWW^T$ , the singular value decompositions of which are given by (24). Since  $\delta_1, \dots, \delta_6$  depend on singular values and singular vectors of these matrices, we will use Lemma 4 and Lemma 5 to bound them.

**Bounding  $\delta_1$ .** Since  $\tilde{Z} = ZU$  and  $\hat{Z} = Z\hat{U}$  by (8) and (15), respectively,

$$\begin{aligned} \delta_1 &= \sum_{i=r+1}^{r+s} \left( \frac{1}{1 - \hat{\sigma}_i^2} Z\hat{U}_i\hat{U}_i^T Z^T - \frac{1}{1 - \sigma_i^2} ZU_iU_i^T Z^T \right) \\ &= Z \left[ \hat{U}_{(r+1):(r+s)}(1 - \hat{\Gamma}^2)^{-1}\hat{U}_{(r+1):(r+s)}^T - U_{(r+1):(r+s)}(1 - \Gamma^2)^{-1}U_{(r+1):(r+s)}^T \right] Z^T. \end{aligned}$$

By (24), column vectors of  $U$  and  $\hat{U}$  are left singular vectors of  $Z^TWW^T$  and  $Z^T\hat{W}\hat{W}^T$ , respectively. Also,  $\|Z^TWW^T - Z^T\hat{W}\hat{W}^T\| \leq \tau_n$  by (23) and  $\min\{1 - \sigma_{r+1}, \sigma_{r+s}\} \geq \tau_n/40$  by (26). It then follows from inequality (57) of Lemma 4 that

$$\|\delta_1\| \leq \frac{12\tau_n}{(\min\{1 - \sigma_{r+1}, \sigma_{r+s}\})^3}.$$

**Bounding  $\delta_2$ .** The same argument for bounding  $\delta_1$  can be applied here to bound  $\delta_2$ . Since  $\tilde{Z} = ZU$  and  $\hat{Z} = Z\hat{U}$  by (8) and (15), we have

$$\delta_2 = Z \left[ \hat{U}_{(r+1):(r+s)}\hat{\Gamma}(1 - \hat{\Gamma}^2)^{-1}\check{W}_{(r+1):(r+s)}^T - U_{(r+1):(r+s)}\Gamma(1 - \Gamma^2)^{-1}\check{W}_{(r+1):(r+s)}^T \right].$$

By (26), (24), (23) and the bound (55) of Lemma 4,

$$\begin{aligned} \left\| \hat{U}_{(r+1):(r+s)}\hat{\Gamma}\check{W}_{(r+1):(r+s)}^T - U_{(r+1):(r+s)}\Gamma\check{W}_{(r+1):(r+s)}^T \right\| &\leq \frac{3\tau_n}{\min\{1 - \sigma_{r+1}, \sigma_{r+s}\} - 4\tau_n} \\ &\leq \frac{5\tau_n}{\min\{1 - \sigma_{r+1}, \sigma_{r+s}\}}. \end{aligned}$$

It remains to apply Lemma 5, for which the following condition must be satisfied:

$$\sigma_{r+1} + \frac{6\tau_n}{\min\{1 - \sigma_{r+1}, \sigma_{r+s}\} - 4\tau_n} < 1.$$

This follows from (26) because  $\min\{1 - \sigma_{r+1}, \sigma_{r+s}\} \geq \tau_n/c \geq 20\tau_n$  and

$$\frac{6\tau_n}{\min\{1 - \sigma_{r+1}, \sigma_{r+s}\} - 4\tau_n} \leq \frac{10\tau_n}{\min\{1 - \sigma_{r+1}, \sigma_{r+s}\}} \leq \min\{1 - \sigma_{r+1}, \sigma_{r+s}\} \leq 1 - \sigma_{r+1}.$$

Therefore by Lemma 5,

$$\begin{aligned} \|\delta_2\| &\leq \left\| \hat{U}_{(r+1):(r+s)} \hat{\Gamma}(1 - \hat{\Gamma}^2)^{-1} \check{W}_{(r+1):(r+s)}^T - U_{(r+1):(r+s)} \Gamma(1 - \Gamma^2)^{-1} \check{W}_{(r+1):(r+s)}^T \right\| \\ &\leq \frac{3 \cdot \frac{5\tau_n}{\min\{1 - \sigma_{r+1}, \sigma_{r+s}\}}}{\left(1 - \sigma_{r+1}^2 - 2 \cdot \frac{5\tau_n}{\min\{1 - \sigma_{r+1}, \sigma_{r+s}\}}\right)^2} \\ &\leq \frac{\frac{15\tau_n}{\min\{1 - \sigma_{r+1}, \sigma_{r+s}\}}}{\left(1 - \sigma_{r+1} - \frac{10\tau_n}{\min\{1 - \sigma_{r+1}, \sigma_{r+s}\}}\right)^2} \\ &\leq \frac{60\tau_n}{(\min\{1 - \sigma_{r+1}, \sigma_{r+s}\})^3}. \end{aligned}$$

**Bounding  $\delta_3$ .** Since  $\tilde{Z} = ZU$  and  $\hat{Z} = Z\hat{U}$  by (8) and (15),

$$\delta_3 = Z \left[ \hat{U}_{(r+s+1):p} \hat{U}_{(r+s+1):p}^T - U_{(r+s+1):p} U_{(r+s+1):p}^T \right] Z^T.$$

By (26), (24), (23) and the bound (54) of Lemma 4, we have

$$\|\delta_3\| \leq \left\| \hat{U}_{(r+s+1):p} \hat{U}_{(r+s+1):p}^T - U_{(r+s+1):p} U_{(r+s+1):p}^T \right\| \leq \frac{2\tau_n}{\sigma_{r+s} - 4\tau_n} \leq \frac{4\tau_n}{\min\{1 - \sigma_{r+1}, \sigma_{r+s}\}}.$$

**Bounding  $\delta_4$ .** We have

$$\delta_4 = \check{W}_{(r+s):(r+s)}(1 - \hat{\Gamma}^2)^{-1} \check{W}_{(r+s):(r+s)}^T - \check{W}_{(r+s):(r+s)}(1 - \Gamma^2)^{-1} \check{W}_{(r+s):(r+s)}^T.$$

By (26), (24), (23) and the bound (57) of Lemma 4, we have

$$\|\delta_4\| \leq \frac{12\tau_n}{(\min\{1 - \sigma_{r+1}, \sigma_{r+s}\})^3}.$$

**Bounding  $\delta_5$ .** Since  $\tilde{Z} = ZU$  and  $\hat{Z} = Z\hat{U}$  by (8) and (15),

$$\delta_5 = \left[ \check{W}_{(r+1):(r+s)} \hat{\Gamma}(1 - \hat{\Gamma}^2)^{-1} \hat{U}_{(r+1):(r+s)}^T - \check{W}_{(r+1):(r+s)} \Gamma(1 - \Gamma^2)^{-1} U_{(r+1):(r+s)}^T \right] Z^T = \delta_2^T,$$

and the bound for  $\delta_2$  applies.

**Bounding  $\delta_6$ .** We have

$$\delta_6 = \check{W}_{(r+s+1):K} \check{W}_{(r+s+1):K}^T - \check{W}_{(r+s+1):K} \check{W}_{(r+s+1):K}^T.$$

By (26), (24), (23) and the bound (54) of Lemma 4, we have

$$\|\delta_6\| \leq \frac{2\tau_n}{\sigma_{r+s} - 4\tau_n} \leq \frac{4\tau_n}{\min\{1 - \sigma_{r+1}, \sigma_{r+s}\}}.$$

The proof of Lemma 3 is complete by combining the bounds for  $\delta_1$  to  $\delta_6$ .  $\square$

*Proof of Lemma 4.* Note that  $X^T X$  and  $\hat{X}^T \hat{X}$  are symmetric matrices with eigenvectors  $V_i, \hat{V}_i$  and eigenvalues being squares of singular values of  $X$  and  $\hat{X}$ , respectively. Also, by the triangle inequality,

$$\|\hat{X}^T \hat{X} - X^T X\| = \|(\hat{X} - X)^T \hat{X} - X^T (\hat{X} - X)\| \leq 2\|\hat{X} - X\|.$$

Therefore by Theorem 7.3.1 of [Bhatia \[1996\]](#),

$$\|\hat{V}_\eta \hat{V}_\eta^T - V_\eta V_\eta^T\| \leq \frac{\|\hat{X}^T \hat{X} - X^T X\|}{\Delta - 2\|\hat{X}^T \hat{X} - X^T X\|} \leq \frac{2\|\hat{X} - X\|}{\Delta - 4\|\hat{X} - X\|}.$$

A similar bound holds for  $\|\hat{U}_\eta \hat{U}_\eta^T - U_\eta U_\eta^T\|$  if we consider  $XX^T$  instead of  $X^T X$ , and (54) is proved.

To show (56), we use Weyl's inequality

$$\begin{aligned} \|(1 - \hat{\Gamma}^2)^{-1} - (1 - \Gamma^2)^{-1}\| &= \max_{i \in \eta} \frac{|\hat{\sigma}_i^2 - \sigma_i^2|}{(1 - \hat{\sigma}_i^2)(1 - \sigma_i^2)} \\ &\leq \frac{2\|\hat{X} - X\|}{(1 - \max_{i \in \eta} \sigma_i^2)(1 - \max_{i \in \eta} \sigma_i^2 - 2\|\hat{X} - X\|)} \\ &\leq \frac{2\|\hat{X} - X\|}{(1 - \max_{i \in \eta} \sigma_i^2 - 2\|\hat{X} - X\|)^2}. \end{aligned}$$

To show (55), let  $S$  and  $\hat{S}$  be diagonal matrices with  $\sigma_i, i \in \eta$  and  $\hat{\sigma}_i, i \in \eta$  on the diagonal, respectively. For every unit vector  $x$ , we need to bound  $\|U_\eta S V_\eta^T x - \hat{U}_\eta \hat{S} \hat{V}_\eta^T x\|$ . Let  $H = V_\eta V_\eta^T, H^\perp = I - H, \hat{H} = \hat{V}_\eta \hat{V}_\eta^T$  and  $\hat{H}^\perp = I - \hat{H}$ . Since

$$x = Hx + H^\perp x = \hat{H}x + \hat{H}^\perp x,$$

it follows that

$$U_\eta S V_\eta^T x - \hat{U}_\eta \hat{S} \hat{V}_\eta^T x = U_\eta S V_\eta^T Hx - \hat{U}_\eta \hat{S} \hat{V}_\eta^T \hat{H}x = XHx - \hat{X}\hat{H}x.$$

By (54), we have

$$\|\hat{X}\hat{H}x - \hat{X}Hx\| \leq \|(\hat{H} - H)x\| \leq \frac{2\|\hat{X} - X\|}{\Delta - 4\|\hat{X} - X\|}.$$

Therefore

$$\begin{aligned} \|U_\eta S V_\eta^T x - \hat{U}_\eta \hat{S} \hat{V}_\eta^T x\| &\leq \|XHx - \hat{X}Hx\| + \|\hat{X}Hx - \hat{X}\hat{H}x\| \\ &\leq \|X - \hat{X}\| + \frac{2\|\hat{X} - X\|}{\Delta - 4\|\hat{X} - X\|} \\ &\leq \frac{3\|\hat{X} - X\|}{\Delta - 4\|\hat{X} - X\|}. \end{aligned}$$

Since  $x$  is an arbitrary unit vector, (55) is proved.

Finally, to show (57), define  $\hat{Z} = \hat{U}_\eta \hat{\Gamma} \hat{U}_\eta^T$  and  $Z = U_\eta \Gamma U_\eta^T$ . We have

$$\begin{aligned} \|\hat{U}_\eta(I - \hat{\Gamma}^2)^{-1}\hat{U}_\eta^T - U_\eta(I - \Gamma^2)^{-1}U_\eta^T\| &= \|(I - \hat{Z}^2)^{-1} - (I - Z^2)^{-1}\| \\ &= \|(I - \hat{Z}^2)^{-1}(\hat{Z}^2 - Z^2)(I - Z^2)^{-1}\| \\ &\leq \|\hat{Z}^2 - Z^2\| \|(I - \hat{Z}^2)^{-1}\| \|(I - Z^2)^{-1}\|. \end{aligned}$$

It is easy to see that

$$\|(I - Z^2)^{-1}\| = \frac{1}{1 - \max_{i \in \eta} \sigma_i^2}$$

and

$$\|(I - \hat{Z}^2)^{-1}\| = \frac{1}{1 - \max_{i \in \eta} \hat{\sigma}_i^2} \leq \frac{1}{1 - \max_{i \in \eta} \sigma_i^2 - 2\|\hat{X} - X\|}.$$

Furthermore, by (55),

$$\begin{aligned} \|\hat{Z}^2 - Z^2\| &= \|\hat{U}_\eta \hat{\Gamma} \hat{V}_\eta^T \hat{V}_\eta \hat{\Gamma} \hat{U}_\eta^T - U_\eta \Gamma V_\eta^T V_\eta \Gamma U_\eta^T\| \\ &\leq \|\hat{U}_\eta \hat{\Gamma} \hat{V}_\eta^T (\hat{V}_\eta \hat{\Gamma} \hat{U}_\eta^T - V_\eta \Gamma U_\eta^T)\| + \|(\hat{U}_\eta \hat{\Gamma} \hat{V}_\eta^T - U_\eta \Gamma V_\eta^T) V_\eta \Gamma U_\eta^T\| \\ &\leq 2\|\hat{U}_\eta \hat{\Gamma} \hat{V}_\eta^T - U_\eta \Gamma V_\eta^T\| \\ &\leq \frac{6\|\hat{X} - X\|}{\Delta - 4\|\hat{X} - X\|}. \end{aligned}$$

Combining the above inequalities gives the stated result. The counterpart for  $V$  is proved in the same way.  $\square$

*Proof of Lemma 5.* Since  $\|\Sigma\| = \|X\| < 1$  and  $\|\hat{\Sigma}\| = \|\hat{X}\| < 1$ , we have

$$\Sigma(1 - \Sigma^2)^{-1} = \sum_{i=0}^{\infty} \Sigma^{2i+1}, \quad \hat{\Sigma}(1 - \hat{\Sigma}^2)^{-1} = \sum_{i=0}^{\infty} \hat{\Sigma}^{2i+1}.$$

Therefore

$$\begin{aligned} U\Sigma(1 - \Sigma^2)^{-1}V^T - \hat{U}\hat{\Sigma}(1 - \hat{\Sigma}^2)^{-1}\hat{V}^T &= \sum_{i=0}^{\infty} (U\Sigma^{2i+1}V^T - \hat{U}\hat{\Sigma}^{2i+1}\hat{V}^T) \\ &= \sum_{i=0}^{\infty} (U\Sigma V^T V\Sigma^{2i}V^T - \hat{U}\hat{\Sigma}\hat{V}^T \hat{V}\hat{\Sigma}^{2i}\hat{V}^T) \\ &= \sum_{i=0}^{\infty} (XV\Sigma^{2i}V^T - \hat{X}\hat{V}\hat{\Sigma}^{2i}\hat{V}^T) \\ &= \sum_{i=0}^{\infty} (XY^{2i} - \hat{X}\hat{Y}^{2i}), \end{aligned}$$

where  $Y = V\Sigma V^T$  and  $\hat{Y} = \hat{V}\hat{\Sigma}\hat{V}^T$ . Since  $X^T X = Y^2$  and  $\hat{X}^T \hat{X} = \hat{Y}^2$ , it follows that  $\|Y^2\| < 1$  and

$$\|\hat{Y}^2 - Y^2\| \leq \|\hat{X}^T(\hat{X} - X)\| + \|(\hat{X} - X)X\| \leq 2\|\hat{X} - X\|.$$

By the triangle inequality and the assumption of the lemma,

$$\|\hat{Y}^2\| \leq \|Y^2\| + 2\|\hat{X} - X\| = \|X\|^2 + 2\|\hat{X} - X\| < 1.$$

Therefore  $\sum_{i=0}^{\infty} Y^{2i}$  and  $\sum_{i=0}^{\infty} \hat{Y}^{2i}$  are convergent series. By the triangle inequality, we have

$$\begin{aligned} \left\| \sum_{i=0}^{\infty} (XY^{2i} - \hat{X}\hat{Y}^{2i}) \right\| &= \left\| \sum_{i=0}^n (X - \hat{X})Y^{2i} + \sum_{i=0}^{\infty} \hat{X}(Y^{2i} - \hat{Y}^{2i}) \right\| \\ &\leq \|X - \hat{X}\| \cdot \left\| \sum_{i=0}^{\infty} Y^{2i} \right\| + \left\| \sum_{i=0}^{\infty} Y^{2i} - \sum_{i=0}^{\infty} \hat{Y}^{2i} \right\| \\ &= \|X - \hat{X}\| \cdot \|(1 - Y^2)^{-1}\| + \left\| (1 - \hat{Y}^2)^{-1} - (1 - Y^2)^{-1} \right\| \\ &= \|X - \hat{X}\| \cdot \|(1 - Y^2)^{-1}\| + \left\| (1 - \hat{Y}^2)^{-1}(\hat{Y}^2 - Y^2)(1 - Y^2)^{-1} \right\| \\ &\leq \|X - \hat{X}\| \cdot \|(1 - Y^2)^{-1}\| + \|\hat{Y}^2 - Y^2\| \cdot \|(1 - \hat{Y}^2)^{-1}\| \cdot \|(1 - Y^2)^{-1}\| \\ &\leq \|X - \hat{X}\| \cdot \|(1 - Y^2)^{-1}\| + 2\|\hat{X} - X\| \cdot \|(1 - \hat{Y}^2)^{-1}\| \cdot \|(1 - Y^2)^{-1}\|. \end{aligned}$$

Note that  $\|(1 - Y^2)^{-1}\| = (1 - \|Y^2\|)^{-1} = (1 - \|X\|^2)^{-1}$  and

$$\|(1 - \hat{Y}^2)^{-1}\| = \frac{1}{1 - \|\hat{Y}^2\|} \leq \frac{1}{1 - \|Y^2\| - 2\|\hat{X} - X\|} = \frac{1}{1 - \|X\|^2 - 2\|\hat{X} - X\|}.$$

Putting these inequalities together, we get

$$\left\| \sum_{i=0}^{\infty} (XY^{2i} - \hat{X}\hat{Y}^{2i}) \right\| \leq \frac{3\|X - \hat{X}\|}{(1 - \|X\|^2 - 2\|\hat{X} - X\|)^2}$$

and the proof is complete.  $\square$

## B Consistency of variance estimation

Denote by  $H$  and  $H^\perp$  the orthogonal projections onto  $\text{span}(\text{col}(X), S_K(P))$  and its orthogonal complement, respectively. Then  $H^\perp Y = H^\perp \epsilon$  and therefore

$$\mathbb{E}\|H^\perp Y\|^2 = \mathbb{E}\text{Trace}(H^\perp \epsilon \epsilon^T) = (n - p - K + r)\sigma^2, \quad (58)$$

where  $p + K - r$  is the dimension of  $\text{span}(\text{col}(X), S_K(P))$ . Since  $\mathcal{P}_{\mathcal{R}}$  is the orthogonal projection onto  $\mathcal{R} = \text{col}(X) \cap S_K(P)$  by (16) and  $\mathcal{P}_{\mathcal{C}} + \mathcal{P}_{\mathcal{N}}$  is the orthogonal projection onto the subspace of  $\text{span}(\text{col}(X), S_K(P))$  that is orthogonal to  $\mathcal{R}$  by Lemma 1, we have

$$H = \mathcal{P}_{\mathcal{R}} + \mathcal{P}_{\mathcal{C}} + \mathcal{P}_{\mathcal{N}}.$$

Recall  $\hat{H} = \hat{\mathcal{P}}_R + \hat{\mathcal{P}}_C + \hat{\mathcal{P}}_N$  from Algorithm 1. Denote  $\hat{H}^\perp = I - \hat{H}$  and  $H^\perp = I - H$ .

*Proof of Proposition 2.* We first bound the difference

$$\hat{H} - H = (\hat{\mathcal{P}}_R - \mathcal{P}_{\mathcal{R}}) + (\hat{\mathcal{P}}_C - \mathcal{P}_{\mathcal{C}}) + (\hat{\mathcal{P}}_N - \mathcal{P}_{\mathcal{N}}).$$

Recall that  $\hat{\mathcal{P}}_R = \hat{Z}_{1:r} \hat{Z}_{1:r}^T$  by (16) and  $\mathcal{P}_{\mathcal{R}} = \tilde{Z}_{1:r} \tilde{Z}_{1:r}^T$  by (10). Also,  $\tilde{Z}_{1:r} = ZU_{1:r}$  and  $\hat{Z}_{1:r} = Z\hat{U}_{1:r}$  by (8) and Algorithm 1, respectively. By (24), column vectors of  $U$  and  $\hat{U}$  are left singular vectors of  $Z^T W W^T$  and  $Z^T \hat{W} \hat{W}^T$ , respectively. Also,  $\|Z^T W W^T - Z^T \hat{W} \hat{W}^T\| \leq \tau_n$  by (23) and  $1 - \sigma_{r+1} \geq 40\tau_n$  by (26). It then follows from the bound (54) of Lemma 4 that

$$\|\hat{\mathcal{P}}_R - \mathcal{P}_{\mathcal{R}}\| = \|Z (\hat{U}_{1:r} \hat{U}_{1:r}^T - U_{1:r} U_{1:r}^T) Z^T\| \leq \frac{2\tau_n}{1 - \sigma_{r+1} - 4\tau_n} \leq \frac{4\tau_n}{1 - \sigma_{r+1}}. \quad (59)$$

Together with the bound of Corollary 5 and the triangle inequality, we see that there exists a constant  $C > 0$  such that with high probability

$$\begin{aligned} \|\hat{H} - H\| &\leq \|\hat{\mathcal{P}}_R - \mathcal{P}_{\mathcal{R}}\| + \|\hat{\mathcal{P}}_C - \mathcal{P}_{\mathcal{C}}\| + \|\hat{\mathcal{P}}_N - \mathcal{P}_{\mathcal{N}}\| \\ &\leq \frac{C\tau_n}{\min\{(1 - \sigma_{r+1})^3, \sigma_{r+s}^3\}}. \end{aligned}$$

Since  $\hat{H}^\perp - H^\perp = \hat{H} - H$ , we obtain

$$\begin{aligned} |\hat{\sigma}^2 - \sigma^2| &= \left| \frac{\|\hat{H}^\perp Y\|^2}{n - p - K + r} - \frac{\|H^\perp Y\|^2}{n - p - K + r} \right| \\ &\leq \|\hat{H}^\perp - H^\perp\| \cdot \frac{\|Y\|^2}{n - p - K + r} \\ &\leq \frac{C\tau_n}{\min\{(1 - \sigma_{r+1})^3, \sigma_{r+s}^3\}} \cdot \frac{\|Y\|^2}{n - p - K + r}. \end{aligned}$$

The claim of the lemma then follows because  $\|Y\| \leq \|X\beta + X\theta + \alpha\| + \|\epsilon\| = O(\sqrt{n(\sigma^2 + p)})$  with high probability because of Assumption A1 and the fact that  $\|\epsilon\|^2 = O(\sigma^2 n)$  with high probability.  $\square$



## C Inference of $\alpha$ and $\beta$

*Proof of Theorem 1.* Recall the variance of  $\omega^T \hat{\beta}$  in (27). By the triangle inequality,

$$\omega^T \Theta \tilde{X}^T \hat{\mathcal{P}}_C \hat{\mathcal{P}}_C^T \tilde{X} \Theta \omega \geq \omega^T \Theta \tilde{X}^T \mathcal{P}_C \mathcal{P}_C^T \tilde{X} \Theta \omega - |\omega^T \Theta \tilde{X}^T (\hat{\mathcal{P}}_C \hat{\mathcal{P}}_C^T - \mathcal{P}_C \mathcal{P}_C^T) \tilde{X} \Theta \omega|$$

From part (iv) of Lemma 1 we have

$$\omega^T \Theta \tilde{X}^T \mathcal{P}_C \mathcal{P}_C^T \tilde{X} \Theta \omega = \|\mathcal{P}_C^T \tilde{X} \Theta \omega\|^2 \geq \|\tilde{Z}_{(r+1):p}^T \tilde{X} \Theta \omega\|^2.$$

It follows from (35) that  $\|\mathcal{P}_C\| \leq 2/(1 - \sigma_{r+1})$ . By the triangle inequality, Corollary 5 and Assumption A3,

$$\|\hat{\mathcal{P}}_C\| \leq \|\hat{\mathcal{P}}_C - \mathcal{P}_C\| + \|\mathcal{P}_C\| \leq \frac{C}{1 - \sigma_{r+1}}.$$

Using Corollary 5 and above inequalities, we get

$$\begin{aligned} \|\hat{\mathcal{P}}_C \hat{\mathcal{P}}_C^T - \mathcal{P}_C \mathcal{P}_C^T\| &\leq \|\hat{\mathcal{P}}_C (\hat{\mathcal{P}}_C^T - \mathcal{P}_C)\| + \|(\hat{\mathcal{P}}_C - \mathcal{P}_C) \mathcal{P}_C^T\| \\ &\leq \frac{C \tau_n}{\min\{(1 - \sigma_{r+1})^4, \sigma_{r+s}^4\}}. \end{aligned}$$

Therefore by (29), Assumption A3 and (28)

$$\begin{aligned} \omega^T \Theta \tilde{X}^T \hat{\mathcal{P}}_C \hat{\mathcal{P}}_C^T \tilde{X} \Theta \omega &\geq \|\tilde{Z}_{(r+1):p}^T \tilde{X} \Theta \omega\|^2 - \frac{C \tau_n \|\tilde{X} \Theta \omega\|^2}{\min\{(1 - \sigma_{r+1})^4, \sigma_{r+s}^4\}} \\ &\geq c^2 - \frac{C \tau_n \|\Theta \omega\|^2}{\min\{(1 - \sigma_{r+1})^4, \sigma_{r+s}^4\}} \\ &\geq c^2/2. \end{aligned}$$

The last inequality implies that the variance of  $\omega^T \hat{\beta}$  satisfies  $\text{Var}(\omega^T \hat{\beta}) \geq c^2 \sigma^2 / (2n)$ . Since the bias of  $\omega^T \hat{\beta}$  is of order  $o(1/\sqrt{n})$  by Proposition 3, it follows that the term

$$\frac{\omega^T \hat{\beta} - \omega^T \beta}{\sqrt{\frac{\sigma^2}{n} \omega^T \Theta \tilde{X}^T \hat{\mathcal{P}}_C \hat{\mathcal{P}}_C^T \tilde{X} \Theta \omega}}$$

is asymptotically standard normal. The proof is complete because  $\hat{\sigma}^2$  is consistent by Proposition 2.  $\square$

*Proof of Corollary 1.* Let  $\eta_j$  be the partial residual from regressing  $\tilde{X}_j$  on all other columns in  $\tilde{X}$ . Therefore, we have  $\tilde{X}_j^T \eta_j = 0$  if  $j' \neq j$ . When making inference on  $\beta_j$ , we have  $\omega = e_j$  and in particular, we can write it as

$$\omega = (\tilde{X}_j^T \eta_j)^{-1} \tilde{X}^T \eta_j.$$

Therefore,

$$\tilde{Z}_{(r+1):p}^T \tilde{X} \Theta \omega = (\tilde{X}_j^T \eta_j)^{-1} \tilde{Z}_{(r+1):p}^T \tilde{X} (\tilde{X}^T \tilde{X})^{-1} \tilde{X}^T \eta_j.$$

Since  $\tilde{X} (\tilde{X}^T \tilde{X})^{-1} \tilde{X}^T$  is the orthogonal projection onto the column space of  $\tilde{X}$  and the column

vectors of  $\tilde{Z}$  form an orthonormal basis of that space, it follows that

$$\tilde{Z}_{(r+1):p}^T \tilde{X} \Theta \omega = (\tilde{X}_j^T \eta_j)^{-1} \tilde{Z}_{(r+1):p}^T \left( \tilde{Z}_{1:r} \tilde{Z}_{1:r}^T + \tilde{Z}_{(r+1):p} \tilde{Z}_{(r+1):p}^T \right) \eta_i = (\tilde{X}_j^T \eta_j)^{-1} \tilde{Z}_{(r+1):p}^T \eta_j.$$

Therefore, we need  $\|\tilde{Z}_{(r+1):p}^T \eta_j\| / |\tilde{X}_j^T \eta_j| \geq c$  to use (29). To make it more interpretable, notice that

$$|\tilde{X}_j^T \eta_j| = \|\eta_j\|^2 = \|\eta_j\|^2 / \|\tilde{X}_j\|^2 = 1 - \|\tilde{X}_j - \eta_j\|^2 / \|\tilde{X}_j\|^2 = 1 - R_j^2,$$

where  $R_j^2$  is the partial  $R^2$  of regressing  $\tilde{X}_j$  to the other columns in  $\tilde{X}$ . Thus, (30) implies (29) and Corollary 1 is a consequence of Theorem 1.  $\square$

*Proof of Theorem 2.* Let  $\text{Bias}(\hat{\gamma}) = \mathbb{E}\hat{\gamma} - O\gamma$  be the bias of  $\hat{\gamma}$  in estimating  $O\gamma$ , where  $O$  is the orthogonal matrix in Proposition 4. Then using the notation from Algorithm 1,

$$\hat{\gamma} \sim N(O\gamma + \text{Bias}(\hat{\gamma}), \Sigma_{\hat{\gamma}}).$$

Therefore, under the null hypothesis  $H_0 : \alpha = 0$ , we have  $\gamma = 0$  and

$$\Sigma_{\hat{\gamma}}^{-1/2} \hat{\gamma} \sim N\left(\Sigma_{\hat{\gamma}}^{-1/2} \text{Bias}(\hat{\gamma}), I\right).$$

From Assumption A3, which implies (26), and Proposition 4,

$$\begin{aligned} \|\Sigma_{\hat{\gamma}}^{-1/2} \text{Bias}(\hat{\gamma})\| &\leq \|\Sigma_{\hat{\gamma}}\|^{-1/2} \cdot \|\text{Bias}(\gamma)\| \\ &\leq \left( \sigma^2 - \frac{C\tau_n}{\min\{(1 - \sigma_{r+1})^2, \sigma_{r+s}^2\}} \right)^{-1/2} \cdot \frac{C\tau_n \sqrt{np}}{\min\{(1 - \sigma_{r+1})^3, \sigma_{r+s}^3\}} \\ &= o(1). \end{aligned}$$

Therefore, as  $n \rightarrow \infty$  we have

$$\Sigma_{\hat{\gamma}}^{-1/2} \hat{\gamma} \xrightarrow{d} N(0, I).$$

Note that  $\|\hat{\Sigma}_{\hat{\gamma}} - \Sigma_{\hat{\gamma}}\| \rightarrow 0$  because  $\hat{\sigma}^2$  is consistent by Proposition 2. Using Slutsky's theorem and the continuously mapping theorem, we obtain

$$\|\hat{\gamma}_0\|^2 = \|\hat{\Sigma}_{\hat{\gamma}}^{-1/2} \hat{\gamma}\|^2 \xrightarrow{d} \chi_K^2.$$

The proof is complete.  $\square$

*Proof of Theorem 3.* Since  $Y = X\beta + X\theta + \alpha + \epsilon$ , it follows from (19) that

$$\hat{\theta} = (X^T X)^{-1} X^T \hat{\mathcal{P}}_{\mathcal{R}} Y = (X^T X)^{-1} X^T \hat{\mathcal{P}}_{\mathcal{R}} (X\beta + X\theta + \alpha) + (X^T X)^{-1} X^T \hat{\mathcal{P}}_{\mathcal{R}} \epsilon.$$

Recall that  $\epsilon \sim N(0, \sigma^2 I_p)$  and  $\tilde{X} = X/\sqrt{n}$ . Therefore,  $\hat{\theta}$  is a multivariate Gaussian vector with variance

$$\text{Var}(\hat{\theta}) = \frac{\sigma^2}{n} \Theta \tilde{X}^T \hat{\mathcal{P}}_{\mathcal{R}} \hat{\mathcal{P}}_{\mathcal{R}}^T \tilde{X} \Theta$$

and expectation

$$\begin{aligned} \mathbb{E}\hat{\theta} &= (X^T X)^{-1} X^T \hat{\mathcal{P}}_{\mathcal{R}} (X\beta + X\theta + \alpha) \\ &= \theta + (X^T X)^{-1} X^T (\hat{\mathcal{P}}_{\mathcal{R}} - \mathcal{P}_{\mathcal{R}}) (X\beta + X\theta + \alpha). \end{aligned}$$

By [A1](#), [A2](#), [A3](#), and [\(59\)](#) we have

$$\|\mathbb{E}\hat{\theta} - \theta\| \leq \|(X^T X)^{-1}\| \cdot \|X\| \cdot \|X\beta + X\theta + \alpha\| \cdot \|\hat{\mathcal{P}}_{\mathcal{R}} - \mathcal{P}_{\mathcal{R}}\| = o(n^{-1/2}).$$

The claim about limiting distributions of  $\hat{\theta}_i$  follows from [\(31\)](#) and the same argument used in the proof of [Corollary 1](#). □

## D Bounding projection perturbation for adjacency matrix

In this section we prove Theorem 4, Corollary 2 and Theorem 5. For the proof of Theorem 4, we need the following lemma.

**Lemma 6** (Lei and Rinaldo [2014]). Let  $A$  be the adjacency matrix of a random graph on  $n$  nodes generated from the inhomogeneous Erdős-Rényi model and denote  $d = n \cdot \max_{ij} \mathbb{E}A_{ij}$ . Then there exist some constants  $C, c_0 > 0$  such that if  $d \geq C \log n$  then  $\|A - \mathbb{E}A\| \leq C\sqrt{d}$  with probability at least  $1 - n^{-c_0}$ .

*Proof of Theorem 4.* Recall the eigen-projection representation (see for example Xia [2019])

$$\hat{W}\hat{W}^T - WW^T = \sum_{k=1}^{\infty} S_k, \quad S_k = \frac{1}{2\pi i} \int_{\Omega} [R(z)E]^k R(z) dz. \quad (60)$$

Here  $E = A - \mathbb{E}A$ ,  $R(z) = (zI - \mathbb{E}A)^{-1}$  is the resolvent and  $\Gamma$  is a contour separating  $\{\lambda_1, \dots, \lambda_K\}$  from the remaining eigenvalues such that  $\Omega$  is sufficiently far away from all eigenvalues of  $\mathbb{E}A$ , that is, for some constant  $c_1 > 0$ ,

$$\min_{j \in [n], z \in \Gamma} |z - \lambda_j| \geq \frac{d}{2c_1}.$$

Such a contour exists because of the eigenvalue gap assumption on  $\mathbb{E}A$ . This property of  $\Omega$  implies that for any  $z \in \Omega$ , we have

$$\|R(z)\| \leq \frac{2c_1}{d}.$$

Note that the range of  $\{\lambda_1, \dots, \lambda_K\}$  is of order  $O(d)$ , therefore the length of  $\Omega$  is of order  $O(d)$  as well. According to Lemma 6, there exists constant  $C > 0$  such that the event

$$T_0 = \left\{ \|E\| \leq C\sqrt{d} \right\}$$

occurs with probability at least  $1 - n^{-c_0}$ . Under  $T_0$ , the bound on the resolvent yields an upper bound for  $S_k$ :

$$\|S_k\| \leq \frac{1}{2\pi} \left( \frac{2c_1}{d} \right)^k \cdot (C\sqrt{d})^{k/2} \cdot \frac{2c_1}{d} \cdot \text{length}(\Omega) \leq \frac{c}{\pi} \left( \frac{4c_1^2 C}{d} \right)^{k/2}. \quad (61)$$

We will use this inequality to upper bound  $S_k$  for all  $k \geq 2$ . It remains to show that  $S_1$  is small. Since

$$R(z) = (zI - \mathbb{E}A)^{-1} = \sum_{j=1}^n \frac{1}{z - \lambda_j} w_j w_j^T,$$

it follows that

$$S_1 = \frac{1}{2\pi i} \int_{\Omega} R(z) E R(z) dz = \sum_{j, j'=1}^n \frac{1}{2\pi i} \int_{\Omega} \frac{w_j^T E w_{j'}}{(z - \lambda_j)(z - \lambda_{j'})} dz \cdot w_j w_{j'}^T.$$

If  $j$  and  $j'$  are both either at most  $K$  or greater than  $K$  then the integral is zero. If one of them is at most  $K$  and the other does not, say  $j \leq K$  and  $j' > K$ , then the integral may not be zero; the

contribution of all such terms is

$$M := 2 \sum_{j \leq K, j' > K} \frac{1}{2\pi i} \int_{\Omega} \frac{w_j^T E w_{j'}}{(z - \lambda_j)(z - \lambda_{j'})} dz \cdot w_j w_{j'}^T = \sum_{j \leq K} \frac{1}{2\pi i} \sum_{j' > K} \frac{w_j^T E w_{j'}}{\lambda_j - \lambda_{j'}} \cdot w_j w_{j'}^T.$$

Let  $v$  be a fixed unit vector. Then

$$\begin{aligned} \|Mv\|^2 &= \left\| 2 \sum_{j \leq K} \frac{1}{2\pi i} w_j \sum_{j' > K} \frac{w_j^T E w_{j'}}{\lambda_j - \lambda_{j'}} \cdot w_{j'}^T v \right\|^2 \\ &= \frac{1}{\pi^2} \sum_{j \leq K} \left| \sum_{j' > K} \frac{w_j^T E w_{j'}}{\lambda_j - \lambda_{j'}} \cdot w_{j'}^T v \right|^2 \\ &= \frac{1}{\pi^2} \sum_{j \leq K} |w_j^T E Q_j v|^2, \end{aligned} \tag{62}$$

in which we denote

$$Q_j = \sum_{j' > K} \frac{1}{\lambda_j - \lambda_{j'}} w_{j'} w_{j'}^T.$$

Note that  $Q_j$  depends only on  $P$  and since  $|\lambda_j - \lambda_{j'}| \geq cd$  for  $j \leq K, j' > K$ , we have  $\|Q_j\| \leq 1/(cd)$  for all  $j \leq K$ , which further implies  $\|Q_j v\| \leq 1/(c_1 d)$ . For any fixed vectors  $x, y \in \mathbb{R}^n$ ,

$$x^T E y = \sum_{i < j} E_{ij} (x_i y_j + x_j y_i) + \sum_i E_{ii} x_i y_i,$$

where  $E_{ij}, i \leq j$  are independent Bernoulli errors. By Hoeffding's inequality (see for example Theorem 2.6.3 in [Vershynin \[2018\]](#)), for some absolute constant  $c'$  we have

$$\begin{aligned} \mathbb{P}(|x^T E y| > t) &\leq 2 \exp \left( - \frac{c' t^2}{\sum_{i < j} (x_i y_j + x_j y_i)^2 + \sum_i x_i^2 y_i^2} \right) \\ &\leq 2 \exp \left( - \frac{c' t^2}{2 \|x\|^2 \|y\|^2} \right). \end{aligned}$$

Applying the above inequality to each  $1 \leq j \leq K$  and using the bound  $\|Q_j v\| \leq 1/(c_1 d)$ , we have

$$\begin{aligned} \mathbb{P} \left( |w_j^T E Q_j v| > \frac{\sqrt{\log n}}{d} \right) &\leq 2 \exp \left( - \frac{c' \log n / d^2}{2 \|w_j\|^2 \|Q_j v\|^2} \right) \\ &\leq 2 \exp \left( - \frac{c' \log n / d^2}{2 / (c_1 d)^2} \right) \\ &= 2n^{-c' / 2c_1^2}. \end{aligned}$$

Denote  $T_j = \{ |w_j^T E Q_j v| > \sqrt{\log n} / d \}$ . Then under  $\bigcap_{j=1}^K T_j$ , by (62) we have

$$\|Mv\|^2 \leq \frac{K \log n}{\pi^2 d^2}.$$

Finally, under  $\bigcap_{j=0}^K T_j$ , using (60) and (61), we obtain

$$\begin{aligned}
\|(\hat{W}\hat{W}^T - WW^T)v\| &\leq \|S_1v\| + \sum_{k \geq 2} \|S_kv\| \\
&\leq \frac{\sqrt{K \log n}}{d} + \sum_{k \geq 2} \left(\frac{4c^2C}{d}\right)^{k/2} \\
&\leq \frac{\sqrt{K \log n}}{d} + \frac{2}{d} \\
&\leq \frac{2\sqrt{K \log n}}{d}
\end{aligned}$$

with probability at least  $1 - (1 + 2K)n^{-c}$  for  $c = \min(c_0, c'/2c_1^2) > 0$ , in which the summation of the sequence is valid as long as  $4c_1^2C/d < 1/2$ . Notice that  $c$  does not depend on  $v$ .  $\square$

*Proof of Corollary 2.* Notice that  $Z$  is fixed matrix. Directly applying Theorem 4 for each column vector of  $Z$  yields

$$\|(\hat{W}\hat{W} - WW^T)Z_j\| \leq \frac{2\sqrt{Kn \log n}}{d}$$

for every  $1 \leq j \leq p$ . This implies

$$\|(\hat{W}\hat{W} - WW^T)Z\| \leq \frac{2\sqrt{pKn \log n}}{d}$$

with probability at least  $1 - pKn^{-c}$ . The proof is complete.  $\square$

*Proof of Corollary 3.* For sufficiently large  $n$ , it is easy to see that by Bernstein inequality the following event

$$T_0 = \left\{ \bar{d}/2 \leq \hat{d} \leq 2\bar{d} \right\}$$

happens with probability at least  $1 - n^{-c}$  for some constant  $c > 0$ . By Theorem 4, the following event

$$T_1 = \left\{ \|(\hat{W}\hat{W}^T - WW^T)Z\| \leq \frac{2\sqrt{pKn \log n}}{d} \right\}$$

happens with probability at least  $1 - pKn^{-c}$ . Under  $T_1$ , according to (23), (24) and (25), we have for every  $1 \leq i \leq \min\{p, K\}$ :

$$|\sigma_i - \hat{\sigma}_i| \leq \|\hat{\Sigma} - \Sigma\| \leq \|(\hat{W}\hat{W}^T - WW^T)Z\| \leq \frac{2\sqrt{pKn \log n}}{d}.$$

Since  $\sigma_r = 1$ , this implies that under  $T_0 \cap T_1$  we have

$$\hat{\sigma}_r \geq 1 - \frac{2\sqrt{pKn \log n}}{d} \geq 1 - \frac{2\sqrt{pKn \log n}}{\bar{d}} \geq 1 - \frac{4\sqrt{pKn \log n}}{\hat{d}}.$$

Similarly, under  $T_0 \cap T_1$  we have

$$\hat{\sigma}_{r+1} \leq \sigma_{r+1} + \frac{2\sqrt{pKn \log n}}{d} \leq \sigma_{r+1} + \frac{4\sqrt{pKn \log n}}{\hat{d}}.$$

The right-hand side of this inequality is strictly smaller than the threshold  $1 - \frac{4\sqrt{pK \log n}}{\hat{d}}$  if

$$\hat{d} > \frac{8\sqrt{pK \log n}}{1 - \sigma_{r+1}},$$

which holds if  $T_0$  occurs and condition (34) is satisfied. Therefore,

$$\mathbb{P}(\hat{r} = r) \geq \mathbb{P}(T_0 \cap T_1) \geq 1 - 2pKn^{-c}$$

and the proof is complete.  $\square$

To prove Theorem 5, we need the following two lemmas to deal with the special case when  $P$  is a matrix of a fixed rank.

**Lemma 7** (Theorem 1 of Xia [2019]). Let  $A$  be the adjacency matrix of a random graph generated from the inhomogeneous Erdős-Rényi model and  $E = A - \mathbb{E}A$ . Assume that  $\mathbb{E}A$  is of a fixed rank  $K$  and  $\|E\| < \lambda_K/2$ , where  $\lambda_K$  is the smallest nonzero eigenvalue of  $\mathbb{E}A$ . Denote by  $W \in \mathbb{R}^{n \times K}$  the matrix whose columns are eigenvectors of  $\mathbb{E}A$  corresponding to the  $K$  largest eigenvalues of  $\mathbb{E}A$ ; define  $\hat{W} \in \mathbb{R}^{n \times K}$  similarly for  $A$ . Then

$$\hat{W}\hat{W}^T - WW^T = \sum_{k \geq 1} S_k,$$

where

$$S_k = \sum_s (-1)^{1+\tau(s)} \tilde{P}^{-s_1} E \tilde{P}^{-s_2} \dots \tilde{P}^{-s_k} E \tilde{P}^{-s_{k+1}},$$

the sum is over all vectors of non-negative integers  $s = (s_1, \dots, s_{k+1})$  such that  $\sum_{i=1}^{k+1} s_i = k$ , and  $\tau(s)$  is the number of positive entries of  $s$ . Moreover,

$$\tilde{P}^0 = W_\perp W_\perp^T, \quad \tilde{P}^{-k} = W \Lambda^{-k} W^T \text{ for } k \geq 1,$$

where  $\Lambda \in \mathbb{R}^{K \times K}$  is the diagonal matrix of  $K$  non-zero eigenvalues of  $\mathbb{E}A$ , and  $W_\perp \in \mathbb{R}^{n \times (n-K)}$  is the matrix whose columns are eigenvectors of  $\mathbb{E}A$  associated with the eigenvalue zero.

**Lemma 8** (Theorem 6.3.2 of Vershynin [2018]). Let  $R \in \mathbb{R}^{m \times n}$  be a non-random matrix and  $X = (X_1, \dots, X_n)^T \in \mathbb{R}^n$  be a random vector with independent, mean zero, unit variance, sub-gaussian coordinates. Then

$$\left| \|RX\| - \|R\|_F \right|_{\psi_2} \leq C \max_i \|X_i\|_{\psi_2}^2 \|R\|,$$

in which  $\|\cdot\|_{\psi_2}$  is the sub-gaussian Orlicz norm.

*Proof of Theorem 5. Part (i).* It is sufficient to find one configuration satisfying the conditions of Theorem 4, such that the inequality holds. In particular, consider an Erdős-Rényi model with  $K = 1$  and  $P = \rho_n \mathbf{1}_n \mathbf{1}_n^T$ , where  $\mathbf{1}_n$  denotes the all-one vector. In this case,  $d = n\rho_n$ . Denote  $v = \frac{1}{\sqrt{n}} \mathbf{1}_n$ . Since

$$|v^T(\hat{W}\hat{W}^T - WW^T)v| \leq \|(\hat{W}\hat{W}^T - WW^T)v\|,$$

it is sufficient to show that

$$|v^T(\hat{W}\hat{W}^T - WW^T)v| \geq c/d. \tag{63}$$

By Lemma 7, we have

$$v^T(\hat{W}\hat{W}^T - WW^T)v = v^T(S_1(E) + S_2(E))v + v^T\sum_{k \geq 3} S_k(E)v, \quad (64)$$

where  $S_k$ 's are defined accordingly in Lemma 7. In particular, if we denote  $\tilde{P}^\perp = \tilde{P}^0$  then

$$\begin{aligned} S_1(E) &= \tilde{P}^{-1}E\tilde{P}^\perp + \tilde{P}^\perp E\tilde{P}^{-1}, \\ S_2(E) &= (\tilde{P}^{-2}E\tilde{P}^\perp E\tilde{P}^\perp + \tilde{P}^\perp E\tilde{P}^{-2}E\tilde{P}^\perp + \tilde{P}^\perp E\tilde{P}^\perp E\tilde{P}^{-2}) \\ &\quad - (\tilde{P}^\perp E\tilde{P}^{-1}E\tilde{P}^{-1} + \tilde{P}^{-1}E\tilde{P}^\perp E\tilde{P}^{-1} + \tilde{P}^{-1}E\tilde{P}^{-1}E\tilde{P}^\perp), \end{aligned}$$

where  $\tilde{P}^{-1} = \frac{1}{n\rho_n}vv^T$ . Since  $\tilde{P}^\perp v = 0$ , we have

$$v^T(S_1(E) + S_2(E))v = -v^T\tilde{P}^{-1}E\tilde{P}^\perp E\tilde{P}^{-1}v = -\frac{1}{d^2n}\|\tilde{P}^\perp E\mathbf{1}_n\|^2.$$

Note that  $\tilde{P}^\perp = I - (1/n)\mathbf{1}_n\mathbf{1}_n^T$  is the centralization operator, therefore

$$\|\tilde{P}^\perp E\mathbf{1}_n\| \geq \|E\mathbf{1}_n\| - \frac{1}{n}\|\mathbf{1}_n\mathbf{1}_n^T E\mathbf{1}_n\| = \|E\mathbf{1}_n\| - \frac{1}{\sqrt{n}}\left|\sum_{i,j} E_{ij}\right|. \quad (65)$$

We will use Lemma 8 to estimate the first term. In particular, let  $\tilde{E} = E/\sqrt{\rho_n(1-\rho_n)}$  and let  $\Psi \in \mathbb{R}^{(n(n+1)/2)}$  be the vector of on and above-diagonal entries of  $\tilde{E}$ , indexed by

$$\Psi = (\tilde{E}_{11}, \tilde{E}_{12}, \dots, \tilde{E}_{1n}, \tilde{E}_{22}, \tilde{E}_{23}, \dots, \tilde{E}_{2n}, \tilde{E}_{33}, \dots, \tilde{E}_{nn})^T.$$

Notice that coordinates of  $\Psi$  are independent random variables with mean zero and unit variance. Then  $E\mathbf{1}_n$  can be written as  $\sqrt{\rho_n(1-\rho_n)}R\Psi$  for the matrix  $R \in \mathbb{R}^{n \times n(n+1)/2}$  defined by

$$R = \begin{pmatrix} \mathbf{1}_n^T & 0 & 0 & \cdots & 0 \\ e_2^{(n)T} & \mathbf{1}_{n-1}^T & 0 & \cdots & 0 \\ e_3^{(n)T} & e_2^{(n-1)T} & \mathbf{1}_{n-2}^T & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ e_n^{(n)T} & e_{n-1}^{(n-1)T} & e_{n-2}^{(n-2)T} & \cdots & 1 \end{pmatrix},$$

in which  $e_j^{(m)} \in \mathbb{R}^m$  is the vector with only its  $j$ th coordinate being 1 while the rest being zeros. Moreover, it is easy to check that (since there are exactly  $n^2$  entries in  $E$ )

$$\|R\|_F^2 = n^2.$$

To use Lemma 8, we will find the spectral norm of  $R$  first. This can be achieved by checking for what unit vector  $u \in \mathbb{R}^{n(n+1)/2}$  the norm  $\|Ru\|^2$  achieves its maximum. If we index  $u$  in the same way as  $\tilde{E}$  (in matrix form), we see that

$$\|Ru\|^2 = \sum_{i=1}^n \left( \sum_{j=1}^n u_{ij} \right)^2,$$

where  $u_{ij} = u_{ji}$  for  $i > j$ . It is easy to see that the maximum is achieved when the mass 1 is equally assigned to all the upper triangular coordinates, as well as the diagonal coordinates but the ratio is



two to one such that

$$u_{ij} = \frac{2}{\sqrt{(2n-1)n}}, \quad i < j \quad \text{and} \quad u_{ii} = \frac{1}{\sqrt{(2n-1)n}}.$$

In this case,

$$\|Ru\|^2 = 2n - 1.$$

So the spectral norm of  $R$  is  $\|R\| = \sqrt{2n-1}$ . Now by using Lemma 8, we have

$$\begin{aligned} \left\| \|E\mathbf{1}_n\| - \sqrt{\rho_n(1-\rho_n)n} \right\|_{\psi_2} &= \left\| \sqrt{\rho_n(1-\rho_n)} \|R\Psi\| - \sqrt{\rho_n(1-\rho_n)n} \right\|_{\psi_2} \\ &\leq C \sqrt{n\rho_n(1-\rho_n)} \|\tilde{E}_{12}\|_{\psi_2}^2 \\ &= C \left( \frac{n}{\rho_n(1-\rho_n)} \right)^{1/2} \|E_{12}\|_{\psi_2}^2. \end{aligned}$$

Since  $E_{12}$  is a centered Bernoulli random variable with success probability  $\rho_n$ , according to [Buldugin and Moskvichova \[2013\]](#), the squared Gaussian norm of  $E_{12}$  for  $\rho_n = o(1)$  is

$$\|E_{12}\|_{\psi_2}^2 = \frac{1 - 2\rho_n}{2(\log(1-\rho_n) - \log \rho_n)} \approx \frac{1}{\log 1/\rho_n}.$$

That means, for some constant  $c$ ,

$$\mathbb{P} \left( \left| \|E\mathbf{1}_n\| - \sqrt{\rho_n(1-\rho_n)n} \right| > t \right) \leq 2 \exp \left( - \frac{ct^2 \rho_n \log^2 1/\rho_n}{n} \right).$$

In particular, for  $\sqrt{n/\log n} \ll d \leq n^{1-\varepsilon}$ , with high probability we have

$$\|E\mathbf{1}_n\| > \sqrt{dn}/2.$$

We turn to upper bound the second term of (65). By the Bernstein inequality,

$$\frac{1}{\sqrt{n}} \left| \sum_{i,j} E_{ij} \right| \leq \sqrt{2d \log n}$$

with high probability. Therefore for sufficiently large  $n$ , the following happens with high probability

$$|v^T (S_1(E) + S_2(E)) v| = \frac{\|\tilde{P}^\perp E\mathbf{1}_n\|^2}{d^2 n} \geq \frac{1}{d^2 n} \left( \sqrt{dn}/2 - \sqrt{2d \log n} \right)^2 \geq \frac{1}{16d}. \quad (66)$$

Next, we look at the term  $v^T \sum_{k \geq 3} S_k(E) v$ . By the definition of  $S_k$ , we have

$$\left| v^T \sum_{k \geq 3} S_k(E) v \right| \leq \sum_{k \geq 3} \|S_k(E)\| \leq \sum_{k \geq 3} \left( \frac{4\|E\|}{d} \right)^k \leq \frac{C}{d^{3/2}} \quad (67)$$

with high probability according to Lemma 6. Combining (66), (67) and (64) yields part (i).

**Part (ii).** Notice that part (i) does not directly indicate the correctness of part (ii). But it motivates the configuration we will use. Specifically, let  $p = K = 1$  and  $P = \rho_n \mathbf{1}_n \mathbf{1}_n^T$ . Notice that in this case,  $W = w = \frac{1}{\sqrt{n}} \mathbf{1}_n$ . Also, let  $w_\perp \in \mathbb{R}^n$  be a unit vector that is orthogonal to  $w$ , the exact value

of which will be chosen later. We choose

$$X = x = \sqrt{n} \left( \frac{1}{\sqrt{2}} w + \frac{1}{\sqrt{2}} w_{\perp} \right).$$

So  $Z = x/\sqrt{n}$ . Moreover, set  $\alpha = \sqrt{n}w$ . Denote by  $\hat{w}$  the leading eigenvector of  $A$ .

It is easy to see that  $r = 0$  and  $s = 1$ . Since  $p = K = 1$ , we have  $\hat{Z} = \tilde{Z} = Z =: z$ ,  $\tilde{W} = W = w$  and  $\tilde{W} = \hat{W} = \hat{w}$ . In particular, a simple calculation shows that  $\hat{\mathcal{P}}_{\mathcal{C}}$  in (17) takes the form

$$\hat{\mathcal{P}}_{\mathcal{C}} = \frac{1}{1 - \hat{\sigma}_1^2} z z^T - \frac{\hat{\sigma}_1}{1 - \hat{\sigma}_1^2} z \hat{w}^T, \quad (68)$$

where  $\hat{\sigma}_1 = \hat{w}^T z$ . It then follows from (19) that

$$\begin{aligned} \mathbb{E} \hat{\beta} &= (x^T x)^{-1} x^T \hat{\mathcal{P}}_{\mathcal{C}} (x\beta + \alpha) \\ &= z^T \left( \frac{1}{1 - \hat{\sigma}_1^2} z z^T - \frac{\hat{\sigma}_1}{1 - \hat{\sigma}_1^2} z \hat{w}^T \right) (z\beta + w) \\ &= \left( \frac{1}{1 - \hat{\sigma}_1^2} z^T z - \frac{\hat{\sigma}_1}{1 - \hat{\sigma}_1^2} \hat{w}^T z \right) (z\beta + w) \\ &= \left( \frac{1}{1 - \hat{\sigma}_1^2} - \frac{\hat{\sigma}_1}{1 - \hat{\sigma}_1^2} \hat{w}^T z \right) \beta + \frac{1}{\sqrt{2}(1 - \hat{\sigma}_1^2)} - \frac{\hat{\sigma}_1}{1 - \hat{\sigma}_1^2} \hat{w}^T w \\ &= \beta + \frac{1}{\sqrt{2}(1 - \hat{\sigma}_1^2)} - \frac{\hat{\sigma}_1}{1 - \hat{\sigma}_1^2} \hat{w}^T w. \end{aligned}$$

This indicates that the bias of  $\hat{\beta}$  satisfies

$$|B(\hat{\beta})| = \frac{1}{1 - \hat{\sigma}_1^2} \left| \hat{\sigma}_1 \hat{w}^T w - \frac{1}{\sqrt{2}} \right| \geq \left| \hat{\sigma}_1 \hat{w}^T w - \frac{1}{\sqrt{2}} \right|.$$

To show that this is further lower bounded by  $c/d$ , notice that

$$\begin{aligned} \left| \hat{\sigma}_1 \hat{w}^T w - \frac{1}{\sqrt{2}} \right| &= \left| z^T \hat{w} \hat{w}^T w - \frac{1}{\sqrt{2}} \right| \\ &= \left| \frac{1}{\sqrt{2}} w^T \hat{w} \hat{w}^T w - \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}} w_{\perp}^T \hat{w} \hat{w}^T w \right| \\ &= \left| \frac{1}{\sqrt{2}} w^T \hat{w} \hat{w}^T w - \frac{1}{\sqrt{2}} w^T w w^T w + \frac{1}{\sqrt{2}} w_{\perp}^T \hat{w} \hat{w}^T w \right| \\ &\geq \frac{1}{\sqrt{2}} \left| w^T (\hat{w} \hat{w}^T - w w^T) w \right| - \frac{1}{\sqrt{2}} \left| w_{\perp}^T \hat{w} \hat{w}^T w \right| \\ &\geq \frac{1}{\sqrt{2}} \left| w^T (\hat{w} \hat{w}^T - w w^T) w \right| - \frac{1}{\sqrt{2}} \left| w_{\perp}^T \hat{w} \right|. \end{aligned}$$

By part (i), with high probability, we already have

$$\frac{1}{\sqrt{2}} \left| w^T (\hat{w} \hat{w}^T - w w^T) w \right| \geq \frac{c}{\sqrt{2}d}.$$

Now we show that the second term of the lower bound above is of order  $o(1/d)$ . For that purpose, we choose a special configuration  $w_{\perp} = e_1/\sqrt{2} - e_2/\sqrt{2}$ , where  $e_1$  and  $e_2$  are unit vectors such that the first and second coordinates are one, respectively, and all other coordinates are zero. In this

case, we have

$$|w_{\perp}^T \hat{w}| = \frac{1}{\sqrt{2}} |\hat{w}_1 - \hat{w}_2| \leq \frac{1}{\sqrt{2}} (|\hat{w}_1 - w_1| + |\hat{w}_2 - w_2|) \leq \sqrt{2} \|\hat{w} - w\|_{\infty}.$$

Notice that  $w = \mathbf{1}/\sqrt{n}$  so it is perfectly incoherent. Therefore, by using Corollary 3.6 of [Lei \[2019\]](#), or the similarly result of [Abbe et al. \[2017\]](#), we have

$$|w_{\perp}^T \hat{w}| \leq \sqrt{2} \|\hat{w} - w\|_{\infty} \leq \sqrt{\frac{\log n}{n^2 \rho_n}} = \sqrt{\frac{\log n}{dn}}$$

with high probability. Consequently, when  $d \ll n/\log n$ , we have  $|w_{\perp}^T \hat{w}| = o(1/d)$  and the lower bound on the bias of  $\hat{\beta}$  follows.

Finally, assume that the variance of the noise in the regression model (4) is  $\sigma^2 = 1$ . From (19) and (68) we have

$$\begin{aligned} \text{Var}(\hat{\beta}) &= (x^T x)^{-1} x^T \hat{\mathcal{P}}_C \hat{\mathcal{P}}_C^T x (x^T x)^{-1} \\ &= \frac{1}{n} z^T \left( \frac{1}{1 - \hat{\sigma}_1^2} z z^T - \frac{\hat{\sigma}_1}{1 - \hat{\sigma}_1^2} z \hat{w}^T \right) \left( \frac{1}{1 - \hat{\sigma}_1^2} z z^T - \frac{\hat{\sigma}_1}{1 - \hat{\sigma}_1^2} \hat{w} z^T \right) z \\ &= \frac{1}{n} \left( \frac{1}{1 - \hat{\sigma}_1^2} z^T - \frac{\hat{\sigma}_1}{1 - \hat{\sigma}_1^2} \hat{w}^T \right) \left( \frac{1}{1 - \hat{\sigma}_1^2} z - \frac{\hat{\sigma}_1}{1 - \hat{\sigma}_1^2} \hat{w} \right) \\ &= \frac{1}{n(1 - \hat{\sigma}_1^2)^2} \|z - \hat{\sigma}_1 \hat{w}\|^2 \\ &\leq \frac{4}{n(1 - \hat{\sigma}_1^2)^2} \\ &\leq \frac{16}{n} \end{aligned}$$

whenever  $\hat{\sigma}_1 < 1/\sqrt{2}$ . This completes the proof.  $\square$

## E Bounding projection perturbation for the Laplacian

*Proof of Theorem 7.* The proof of Theorem 7 is very similar to the proof of Theorem 4 in Section D. We highlight here the main differences. Denote

$$E' = L - \mathbb{E}L = (D - \mathbb{E}D) - (A - \mathbb{E}A) = (D - \mathbb{E}D) - E.$$

Then by Bernstein's inequality, there exists a constant  $C > 0$  such that

$$\|D - \mathbb{E}D\| \leq C\sqrt{d \log n}$$

with high probability. Together with Lemma 6, this implies  $\|E'\| \leq C\sqrt{d \log n}$ . Using Assumption A8 about eigenvalue gap of  $P = \mathbb{E}L$ , eigen-projection representation (60) (where  $E$  is replaced with  $E'$ ) and arguing as in the proof of Theorem 4, we see that similar to (61),

$$\|S_k\| \leq \frac{c}{\pi} \left( \frac{4c^2 C \log n}{d} \right)^{k/2}, \quad k \geq 2.$$

It remains to bound  $S_1$ . According to (62),

$$\|S_1 v\|^2 = \frac{2}{\pi^2} \sum_{j \leq K} |w_j^T E' Q_j v|^2 \leq \frac{4}{\pi^2} \sum_{j \leq K} |w_j^T E Q_j v|^2 + \frac{4}{\pi^2} \sum_{j \leq K} |w_j^T (D - \mathbb{E}D) Q_j v|^2,$$

where  $Q_j$  are fixed matrices with norms of order  $1/d$ . The first term on the right-hand side is of order  $K \log n/d^2$ , as in the proof of Theorem 4. Regarding the second term, for  $x = w_j$  and  $y = Q_j x$  with  $\|x\| = 1$  and  $\|y\| = O(1/d)$ , we have

$$x^T (D - \mathbb{E}D) y = \sum_{i, i'=1}^n x_i y_i (A_{ij} - \mathbb{E}A_{ii'})$$

Again, by Hoeffding's inequality [Vershynin, 2018, Theorem 2.6.3], we have for some absolute constant  $c' > 0$ :

$$\mathbb{P}(|x^T (D - \mathbb{E}D) y| > t) \leq 2 \exp\left(-\frac{c't^2}{\sum_{i < i'} x_i^2 y_i^2}\right).$$

Using  $x = w_j$  with  $|x_i| \leq \|W\|_\infty$  and  $\|y\| = O(1/d)$ , we have

$$\sum_{i < i'} x_i^2 y_i^2 \leq n \sum_{i=1}^n x_i^2 y_i^2 \leq n \|W\|_\infty^2 \cdot \|y\|^2 = O(n \|W\|_\infty^2 / d^2).$$

Therefore with high probability,  $|x^T (D - \mathbb{E}D) y|^2 = O(\|W\|_\infty^2 n \log n / d^2)$ . Putting these inequalities together, we obtain

$$\|S_1 v\|^2 = O\left(\frac{(K + \|W\|_\infty^2 n) \log n}{d}\right).$$

The proof is complete by using the triangle inequality the bound for  $\|S_k\|$  with  $k \geq 2$ .  $\square$

*Proof of Corollary 4.* The proof of Corollary 4 is very similar to the proof of Corollary 2 in Section D, but with the help of Theorem 7 instead of Theorem 4. We omit the detail.  $\square$

## F Proofs for small projection perturbation for parametric estimation under block models

First, we introduce the parametric estimators for  $P$  under block models. Given the community label, the maximum likelihood estimator (MLE) of  $P$  is used. Under the SBM, the MLE of  $B$  can be obtained by

$$\hat{B}_{k\ell} = \frac{\sum_{g_i=k, g_j=\ell} A_{ij}}{n_k n_\ell}. \quad (69)$$

with the follow-up estimator  $\hat{P}_{ij} = \hat{B}_{g_i g_j}$ . Under the DCBM, let  $d_i$  be the degree of node  $i$ . The  $\nu_i$ 's can be estimated by

$$\hat{\nu}_i = \frac{n_k d_i}{\sum_{g_j=k} d_j} \quad (70)$$

while the  $B$  matrix can be estimated by

$$\hat{B}_{k\ell} = \frac{1}{n_k n_\ell} \sum_{g_i=k, g_j=\ell} \frac{A_{ij}}{\hat{\nu}_i \hat{\nu}_j}. \quad (71)$$

The resulting  $\hat{P}_{ij}$  is given by  $\hat{\nu}_i \hat{\nu}_j \hat{B}_{g_i g_j}$ . These will be the estimators we use for the parametric version of our method.

*Proof of Theorem 6. Part (i).* Note that the estimator  $\hat{B}_{k\ell}$  is the sample average of independent Bernoulli random variables. Using Bernstein's inequality and  $K = o(\sqrt{nd/\log n})$  from A5, with high probability we have

$$|\hat{B}_{k\ell} - B_{k\ell}| = O\left(\frac{K\sqrt{\kappa_n \log n}}{n}\right)$$

holds for all  $1 \leq k, \ell \leq K$ . Therefore, we have

$$\|\hat{P} - P\|_\infty = O\left(\frac{K\sqrt{\kappa_n \log n}}{n}\right).$$

Notice that under the current assumption, the eigen-gap of  $P$  for the  $K$  leading eigenvectors is in the order of  $d/K$  [Lei and Rinaldo, 2014]. Using David-Kahan theorem [Yu et al., 2014] together with A5 and A7, we have

$$\|\hat{W}\hat{W}^T - WW^T\| = O\left(\frac{\|\hat{P} - P\|}{d/K}\right) = O\left(\frac{\sqrt{n}\|\hat{P} - P\|_\infty}{d/K}\right) = O\left(\frac{K^2\sqrt{\log n}}{n\sqrt{d}}\right).$$

The condition in part (i) then implies that Assumption A3 holds.

**Part (ii).** Since  $d \gg \log n$ , for sufficiently large  $n$ , by Bernstein inequality we have

$$\max_i |d_i - \mathbb{E}d_i| \leq C\sqrt{d \log n}$$

with high probability. Denote  $\bar{d}_k = \sum_{g_j=k} d_j/n_k$ , then  $\hat{\nu}_i = d_i/\bar{d}_{g_i}$  and  $\nu_i = \mathbb{E}d_i/\mathbb{E}\bar{d}_{g_i}$ . Therefore by the above inequality, Assumption A6 and the triangle inequality,

$$|\hat{\nu}_i - \nu_i| = \left| \frac{d_i}{\bar{d}_{g_i}} - \frac{\mathbb{E}d_i}{\mathbb{E}\bar{d}_{g_i}} \right| \leq \frac{|d_i - \mathbb{E}d_i|}{\mathbb{E}\bar{d}_{g_i}} + \frac{d_i|\bar{d}_{g_i} - \mathbb{E}\bar{d}_{g_i}|}{\bar{d}_{g_i}\mathbb{E}\bar{d}_{g_i}} = O\left(\sqrt{\frac{\log n}{d}}\right). \quad (72)$$

Now we bound the error  $\hat{B}_{k\ell} - B_{k\ell}$ . Recall that

$$\hat{B}_{k\ell} = \frac{1}{n_k n_\ell} \sum_{g_i=k, g_j=\ell} \frac{A_{ij}}{\hat{\nu}_i \hat{\nu}_j}.$$

Define

$$\hat{B}_{k\ell}^* = \frac{1}{n_k n_\ell} \sum_{g_i=k, g_j=\ell} \frac{A_{ij}}{\nu_i \nu_j}.$$

We have  $|\hat{B}_{k\ell} - B_{k\ell}| \leq |\hat{B}_{k\ell} - \hat{B}_{k\ell}^*| + |\hat{B}_{k\ell}^* - B_{k\ell}|$ . The second term  $|\hat{B}_{k\ell}^* - B_{k\ell}|$  is simply a sum of centered Bernoulli random variables and can be bounded by  $O_P\left(\frac{K\sqrt{\kappa_n \log n}}{n}\right)$  as before (we use Assumption A6 to bound  $\nu_i$ ). For the first term, notice that

$$\begin{aligned} |\hat{B}_{k\ell} - \hat{B}_{k\ell}^*| &\leq \frac{1}{n_1 n_2} \sum_{g_i=k, g_j=\ell} A_{ij} \left| \frac{1}{\hat{\nu}_i \hat{\nu}_j} - \frac{1}{\nu_i \nu_j} \right| \\ &\leq \max_{ij} \left| \frac{1}{\hat{\nu}_i \hat{\nu}_j} - \frac{1}{\nu_i \nu_j} \right| \cdot \frac{1}{n_k n_\ell} \sum_{g_i=k, g_j=\ell} A_{ij}. \end{aligned}$$

By Assumption A6 and (72), we have

$$\left| \frac{1}{\hat{\nu}_i \hat{\nu}_j} - \frac{1}{\nu_i \nu_j} \right| \leq \frac{|\nu_j - \hat{\nu}_j|}{\hat{\nu}_i \hat{\nu}_j \nu_j} + \frac{|\nu_i - \hat{\nu}_i|}{\hat{\nu}_i \hat{\nu}_j \nu_i} = O\left(\max_i |\hat{\nu}_i - \nu_i|\right) = O\left(\sqrt{\frac{\log n}{d}}\right).$$

Furthermore, using the assumption  $K = o(\sqrt{nd/\log n})$  and Bernstein's inequality as before, we see that with high probability we have

$$\max_{k,\ell} \frac{1}{n_k n_\ell} \sum_{g_i=k_1, g_j=k_2} A_{ij} = O(\kappa_n).$$

Therefore,  $|\hat{B}_{k\ell} - \hat{B}_{k\ell}^*| = O_P(\sqrt{\kappa_n \log n/n})$ . With the bound on  $|\hat{B}_{k\ell}^* - B_{k\ell}|$ , we have

$$|\hat{B}_{k\ell} - B_{k\ell}| = O\left(\sqrt{\frac{\kappa_n \log n}{n}} + \frac{K\sqrt{\kappa_n \log n}}{n}\right) = O\left(\sqrt{\frac{\kappa_n \log n}{n}}\right).$$

Finally, for any  $i$  and  $j$ , we have

$$|\hat{B}_{g_i g_j} \hat{\nu}_i \hat{\nu}_j - B_{g_i g_j} \nu_i \nu_j| \leq \hat{\nu}_i \hat{\nu}_j |\hat{B}_{g_i g_j} - B_{g_i g_j}| + B_{g_i g_j} |\hat{\nu}_i \hat{\nu}_j - \nu_i \nu_j| = O\left(\sqrt{\frac{\kappa_n \log n}{n}}\right).$$

Similar to the SBM case, we use the Davis-Kahan theorem and the bound on eigen-gap in [Lei and Rinaldo \[2014\]](#) to achieve the bound

$$\|\hat{W}\hat{W}^T - WW^T\| = O\left(\frac{\|\hat{P} - P\|}{d/K}\right) = O\left(\frac{\sqrt{n}\|\hat{P} - P\|_\infty}{d/K}\right) = O\left(\frac{K\sqrt{\log n}}{\sqrt{nd}}\right).$$

The condition in part (ii) then implies that Assumption A3 holds.  $\square$

## G A bootstrapping method for selecting $r$

The theory-driven tuning of  $r$  introduced in (33) is based on large-sample properties and may be conservative when  $n$  is not sufficiently large. Here, we propose an alternative way of tuning  $r$  that provides better performance for small networks, according to our empirical evaluation. The idea is to approximate the actual deviation of principal angles using a bootstrapping procedure. Specifically, we construct a singular value thresholding estimator  $P^*$  as in Chatterjee [2015], but with use of the given rank  $K$  instead. Then  $P^*$  is rescaled so that the average degree of  $P^*$  matches the average degree  $\hat{d}$  of the original network. Note that the informative eigenvectors of  $P^*$  are still  $\hat{W}$ . After that, we randomly generate 50 networks  $\hat{A}^{(b)}$  with  $b = 1, \dots, 50$  from  $P^*$  and denote their  $K$  leading eigenvectors by  $\hat{W}^{(b)}$ . Furthermore, the singular values of  $Z^T \hat{W}^{(b)}$  are computed, denoted by  $(\hat{\sigma}_1^{(b)}, \dots, \hat{\sigma}_K^{(b)})$  with  $b = 1, \dots, 50$ . Finally, the maximum deviance  $\delta = \max_{k,b} |\hat{\sigma}_k^{(b)} - \hat{\sigma}_k|$  is taken to construct

$$\hat{r} = \max_k \{k : \hat{\sigma}_k > 1 - \delta\}.$$

Intuitively, the sampling step of  $\hat{A}^{(b)}$  mimics the generating procedure of  $A$ . Thus,  $\delta$  should give an accurate estimate of the possible perturbation on the principal angles.

## H Additional results of the simulation examples

Table 5 summarizes the type-I error probability of the  $\chi^2$  test from the setting  $\gamma = (0, 0, 0, 0)^T$  in the configuration of Section 5. It shows the desired type-I error control to test  $H_0 : \alpha = 0$

Table 5: The type-I error probability of the  $\chi^2$  test for  $\gamma$  when the network is generated from the SBM and  $\gamma = 0$ .

| Method | $n$  | average expected degree |            |           |
|--------|------|-------------------------|------------|-----------|
|        |      | $2 \log n$              | $\sqrt{n}$ | $n^{2/3}$ |
| SP     | 1000 | 0.0490                  | 0.0497     | 0.0510    |
|        | 2000 | 0.0504                  | 0.0497     | 0.0506    |
|        | 4000 | 0.0496                  | 0.0490     | 0.0501    |
| SP-SBM | 1000 | 0.0496                  | 0.0500     | 0.0511    |
|        | 2000 | 0.0504                  | 0.0495     | 0.0503    |
|        | 4000 | 0.0499                  | 0.0493     | 0.0500    |

As a counterpart of the study in Section 5, we generate networks from the DCBM. For the DCBM, the degree parameters are generated from a uniform distribution between 0.2 and 1. All of the other configurations remain the same, as in the previous setting. Table 6 shows the average bias-SD ratio for  $\hat{\beta}_2, \hat{\beta}_3$  and  $\hat{\beta}_4$  under the DCBM, while Table 7 shows the empirical coverage of the 95% confidence intervals in this setting. The high-level message remains similar to the SBM setting and matches the theory. It can be seen that both versions of SP are valid for large enough  $n$  when  $\varphi_n = n^{2/3}$ , while the parametric version also works for  $\varphi_n = 2\sqrt{n}$ . However, the advantage of the parametric version is smaller compared with the SBM setting because the DCBM estimation is more complicated. Lastly, Table 8 shows the type I error probability of the  $\chi^2$  test, in which the validity is evident.

Table 6: The average theoretical bias-SD ratio for  $\beta_2, \beta_3$  and  $\beta_4$  when the network is generated from the DCBM.

| Method  | $n$  | average expected degree |            |           |
|---------|------|-------------------------|------------|-----------|
|         |      | $2 \log n$              | $\sqrt{n}$ | $n^{2/3}$ |
| SP      | 1000 | 1.509                   | 0.675      | 0.268     |
|         | 2000 | 2.084                   | 0.681      | 0.204     |
|         | 4000 | 3.105                   | 0.636      | 0.167     |
| SP-DCBM | 1000 | 0.950                   | 0.286      | 0.096     |
|         | 2000 | 1.040                   | 0.238      | 0.089     |
|         | 4000 | 1.470                   | 0.180      | 0.081     |



Table 7: The average coverage probability of 95% confidence intervals of  $\beta_2, \beta_3, \beta_4$  when the network is generated from the DCBM.

| Method  | $n$  | average expected degree |            |           |
|---------|------|-------------------------|------------|-----------|
|         |      | $2 \log n$              | $\sqrt{n}$ | $n^{2/3}$ |
| SP      | 1000 | 0.775                   | 0.912      | 0.944     |
|         | 2000 | 0.605                   | 0.901      | 0.946     |
|         | 4000 | 0.329                   | 0.892      | 0.947     |
| SP-DCBM | 1000 | 0.881                   | 0.943      | 0.950     |
|         | 2000 | 0.848                   | 0.945      | 0.948     |
|         | 4000 | 0.754                   | 0.946      | 0.951     |

Table 8: The type-I error probability of the  $\chi^2$  test for  $\gamma$  when the network is generated from the DCBM and  $\gamma = 0$ .

| Method  | $n$  | average expected degree |            |           |
|---------|------|-------------------------|------------|-----------|
|         |      | $2 \log n$              | $\sqrt{n}$ | $n^{2/3}$ |
| SP      | 1000 | 0.0518                  | 0.0514     | 0.0506    |
|         | 2000 | 0.0515                  | 0.0517     | 0.0497    |
|         | 4000 | 0.0508                  | 0.0504     | 0.0501    |
| SP-DCBM | 1000 | 0.0512                  | 0.0515     | 0.0502    |
|         | 2000 | 0.0518                  | 0.0510     | 0.0492    |
|         | 4000 | 0.0498                  | 0.0489     | 0.0498    |

Table 9 shows the comparison of our methods with the benchmark methods in the DCBM setting.

Table 9: The mean squared error (MSE)  $\times 10^2$  of  $\mathbb{E}Y$  when the network is generated from the DCBM with  $n = 1000, p = 4, K = 4$  with three types of individual effects.

| indiv. effects | avg. degree | SP    | SP-DCBM | OLS   | SIM   | RNC   | SP-L  | SP-DCBM-L |
|----------------|-------------|-------|---------|-------|-------|-------|-------|-----------|
| eigenspace     | $2 \log n$  | 11.07 | 6.70    | 41.51 | 29.63 | 12.28 | 41.51 | 41.51     |
|                | $\sqrt{n}$  | 4.47  | 1.26    | 41.51 | 27.89 | 10.95 | 41.51 | 41.51     |
|                | $n^{2/3}$   | 1.23  | 0.38    | 41.51 | 24.22 | 9.47  | 41.51 | 41.51     |
| $\gamma = 0$   | $2 \log n$  | 0.14  | 0.14    | 0.12  | 27.48 | 0.26  | 0.13  | 0.12      |
|                | $\sqrt{n}$  | 0.14  | 0.13    | 0.12  | 26.00 | 0.64  | 0.13  | 0.13      |
|                | $n^{2/3}$   | 0.13  | 0.13    | 0.12  | 25.34 | 1.57  | 0.13  | 0.12      |
| smooth         | $2 \log n$  | 25.20 | 25.20   | 25.20 | 28.34 | 16.03 | 12.69 | 17.92     |
|                | $2\sqrt{n}$ | 23.83 | 23.83   | 23.81 | 25.24 | 14.79 | 12.00 | 12.26     |
|                | $n^{2/3}$   | 26.21 | 26.22   | 26.20 | 26.70 | 12.80 | 13.20 | 15.00     |

## I Adolescents’ social activity study

The AddHealth study [Harris, 2009] is a longitudinal study about various health-related behaviors of students in grade 7–12 in the U.S.. In this section, we use a subset of the AddHealth data to study the gender and racial effects on social activity diversity. The data set contains 82 local communities, and each local community contains one public high school. The students listed up to 10 friends and how many different types of social activities they had with each friend. The average number of social activities per friend is calculated as a measure of social activity diversity as an important factor in mental health [Sherbourne, 1988, Wildes et al., 2002]. Besides this, the gender, grade, and race information are also available for each student. In some of the local communities, a feeder middle school of the high school is also included. In this case, the data set also contains the school labels of the students. We want to learn whether there are significant gender and racial differences in the social activity diversity level of the students. This type of analysis may bring insights about health-related behaviors in adolescents’ life. In the context, it is evident that social network information should be incorporated in modeling because each social activity may involve multiple students, and ignoring this factor may lead to misleading results.

The SP method is applied to all 82 data sets (local communities) and we always report the summary results with the 0.05 FDR control [Benjamini and Hochberg, 1995]. Out of the 82 data sets, 72 data sets present significant gender effect, and 14 data sets reveal significant racial effect. The network effect is significant in 37 communities. Here, we pick the largest network with significant network effects in the data set ( $n = 2152$  nodes and two schools) to show a detailed analysis. The network is shown in Figure 1, in which the node sizes represent the estimated  $\alpha$  in the SP. Large size indicates that the individual is believed to have higher social diversity.




Figure 1: The friendship network of 2152 students with school information.

Cross-validation of the network [Li et al., 2020b] indicates that neither of the block models is proper. Therefore, focus is instead placed on the model-free version of the SP on this data set. The p-value

for the  $\chi^2$  test of  $\gamma$  is close to 0, indicating a significant network effect. Multiple insignificantly different levels of the race variable are then merged, resulting in a binary variable of “white” and “non-white” (reference level), and the models are fit again. In both cases,  $r$  is estimated to be zero indicating no strong confounding between the covariates and the network structure. And the estimate  $\beta$  is the sole characterization of the covariate effects. Both two versions of model fitting results are shown in Table 10.

Table 10: Regression coefficients  $\hat{\beta}$  and the confidence intervals from the largest local community in the data set.

|                | original variables |             |        |             | merged variables |             |        |             |
|----------------|--------------------|-------------|--------|-------------|------------------|-------------|--------|-------------|
|                | SP                 |             | OLS    |             | SP               |             | OLS    |             |
|                | coef.              | p-val.      | coef.  | p-val.      | coef.            | p-val.      | coef.  | p-val.      |
| sex            | 0.180              | $< 10^{-9}$ | 0.173  | $< 10^{-7}$ | 0.180            | $< 10^{-8}$ | 0.172  | $< 10^{-7}$ |
| grade          | 0.246              | $< 10^{-3}$ | 0.206  | $< 10^{-5}$ | 0.245            | $< 10^{-7}$ | 0.203  | $< 10^{-5}$ |
| school         | 0.003              | 0.476       | -0.039 | 0.380       | 0.006            | 0.448       | -0.035 | 0.427       |
| race: white    | 0.484              | 0.075       | 0.746  | 0.030       | 0.380            | $< 10^{-7}$ | 0.598  | $< 10^{-9}$ |
| race: black    | 0.038              | 0.456       | 0.044  | 0.901       | –                | –           | –      | –           |
| race: hispanic | 0.106              | 0.376       | 0.163  | 0.633       | –                | –           | –      | –           |
| race: other    | 0.167              | 0.315       | 0.189  | 0.593       | –                | –           | –      | –           |

It can be seen that sex has a significant effect, indicating that girls tend to involve in 0.18 more social activities on average. The grade is also statistically significant, indicating increasingly diverse social activities among students as they grow up. In these refined results, The race effect estimated by OLS is about 1.5 times the effect size estimated by the SP. OLS also estimates an obviously weaker effect on grade. These differences can be explained by network interference, demonstrated in Figure 2. It appears that friendship ties between students exhibit evident race preference — most white (non-white) students tend to make friends with other white (nonwhite) students. A similar pattern is also observed for the grade, as expected because students tend to make friends with others in the same grade. These correlations between the network structure and the variables explains why our model gives very different estimates of racial and grade effects. Notice that the proposed model includes the standard linear model as a special case, and the  $\chi^2$  test indicates a strong social effect. We believe that the proposed model may be preferable as it is reasonable to believe that social effects exist in this situation and the inference framework can test the significance of the network effects.




Figure 2: The friendship network of 2152 students with race and grade information.