

Unlinked monotone regression

Fadoua Balabdaoui

*Seminar für Statistik
ETH, Zurich
Rämistrasse 101
8092, Zurich, Switzerland*

FADOUA.BALABDAOUI@STAT.MATH.ETHZ.CH

Charles R. Doss

*School of Statistics
University of Minnesota
Minneapolis, MN 55455, USA*

CDOSS@UMN.EDU

Cécile Durot

*Modal'x
Université Paris Nanterre
F92000, Nanterre, France*

CECILE.DUROT@GMAIL.COM

Editor: Editor name

Abstract

We consider so-called univariate unlinked (sometimes “decoupled,” or “shuffled”) regression when the unknown regression curve is monotone. In standard monotone regression, one observes a pair (X, Y) where a response Y is linked to a covariate X through the model $Y = m_0(X) + \epsilon$, with m_0 the (unknown) monotone regression function and ϵ the unobserved error (assumed to be independent of X). In the unlinked regression setting one gets only to observe a vector of realizations from both the response Y and from the covariate X where now $Y \stackrel{d}{=} m_0(X) + \epsilon$. There is no (observed) pairing of X and Y . Despite this, it is actually still possible to derive a consistent non-parametric estimator of m_0 under the assumption of monotonicity of m_0 and knowledge of the distribution of the noise ϵ . In this paper, we establish an upper bound on the rate of convergence of such an estimator under minimal assumption on the distribution of the covariate X . We discuss extensions to the case in which the distribution of the noise is unknown. We develop a gradient-descent-based algorithm for its computation, and we demonstrate its use on synthetic data. Finally, we apply our method (in a fully data driven way, without knowledge of the error distribution) on longitudinal data from the US Consumer Expenditure Survey.

Keywords: deconvolution, quantile, minimax, monotone regression, rates

Contents

1	Introduction	2
2	The minimum contrast estimator	6
2.1	Setup, terminology, and notation	6
2.2	Existence	7
2.3	Convergence rate	11
2.4	Convergence rate in the case of different sample sizes	14
3	Fenchel optimality conditions	14

©2000 Marina Meilă and Michael I. Jordan.

License: CC-BY 4.0, see <https://creativecommons.org/licenses/by/4.0/>. Attribution requirements are provided at <http://jmlr.org/papers/v1/meila00a.html>.

4	Computation	15
5	Extension to the case of unknown noise distribution	15
5.1	Estimation of the noise distribution in the semi-supervised setting . . .	15
5.2	Estimation of the noise distribution with longitudinal responses	17
6	Demonstrations on synthetic and real data	18
6.1	Computations on synthetic data	18
6.2	Computations on CEX data	19
7	Conclusions and directions for future research	20
A	Bounding the integrals in (36) and (37).	24
B	Basic empirical process theory definitions	25
C	Gradient computation	25
C.1	Proof of (26)	26
C.2	Computations for Laplace distribution	26
C.3	Computations for Gaussian errors	26
C.4	On implementation of Algorithm 1	27
D	Proofs	27
D.1	Proofs for Section 2.3	27
D.2	Proof of Theorem 2.2	34
E	Proof of Proposition 3.1	37

1. Introduction

An important part of data science is the construction of a dataset; nowadays, because there are so many different entities collecting increasing amounts of data, datasets are often constructed by combining separate sub-datasets or datastreams. Also, datasets sometimes undergo some form of anonymization: this can be due to the increasing prevalence of privacy concerns, or in some cases due to concerns about having limited data-transmission bandwidth when many separate sensors are streaming data to a central server (Pananjady et al., 2016). Thus, it is increasingly common for data scientists/analysts to want to relate variables in one dataset to variables in another dataset when the two datasets are *unlinked*. In this paper, we consider the problem of unlinked regression, specifically when the regression function is assumed to satisfy a monotonicity constraint.

In the standard regression setting, we have

$$Y_i = m_0(X_i) + \epsilon_i, \quad E(\epsilon_i) = 0, \quad i = 1, \dots, n, \quad (1)$$

for a random noise variable ϵ_i that is independent of X_i . The most basic assumption of this model is that for each index $i = 1, \dots, n$, the *pair* (X_i, Y_i) is observed. For now, we assume that the covariates X_i , $i = 1, \dots, n$ are univariate random variables. A more general model than the above standard regression model is the *shuffled* regression, in which we do not get to see the pairs (X_i, Y_i) , $i = 1, \dots, n$; rather, we only observe

(X_1, \dots, X_n) and (Y_1, \dots, Y_n) , without knowing which X_i is paired with which Y_i . Thus, we have the same model as (1) except that the first equality is now only an equality in distribution and there is an unknown permutation π on $\{1, \dots, n\}$ such that $Y_i = m_0(X_{\pi(i)}) + \epsilon_i$ for all i . This happens for instance in case of anonymized data. An even more general model is the *unlinked* regression model that we consider in this paper, where again, the first equality only holds in distribution but where in addition, the X_i 's could be observed on different individuals from the Y_i 's so that the two samples are not necessarily connected through a permutation π . This happens for instance if the two samples have been collected independently (by separate entities). The number of observed X_i 's may even differ from the number of observed Y_i 's so we observe independent and identically distributed (i.i.d.) variables Y_1, \dots, Y_{n_y} and i.i.d. variables X_1, \dots, X_{n_x} such that

$$Y \stackrel{d}{=} m_0(X) + \epsilon \quad (2)$$

where $Y \sim Y_i$, $X \sim X_i$, and ϵ is independent of X with distribution function Φ_ϵ . In shuffled or unlinked regression models, it may seem hopeless to even try to learn the regression function m_0 ; however, it turns out that if m_0 is assumed to be monotonically increasing, as we will do in this paper, and if the error distribution is known a priori, then in fact one can construct consistent estimators of m_0 . (We will return in Section 5 to discuss the case where the error distribution is unknown.)

Unlinked estimation can be considered in a variety of settings (e.g., DeGroot and Goel (1980)). It appears that unlinked monotone regression was recently introduced by Carpentier and Schlüter (2016). One motivating example discussed in Carpentier and Schlüter (2016) is about expenditure on goods or services, such as housing: the price an individual is willing to pay for housing is expected to be monotonically increasing (at least on average, if not individually) as a function of the individual's salary. However, estimating the monotonic relationship is hindered by the simple fact that the data on wages and housing transactions are often gathered by different agents. There are many other motivating examples for unlinked or shuffled regression, besides the ones already discussed. (In some examples, there may be information allowing partial matching; see our discussion point 6 in Section 7.) In flow cytometry, cells suspended in a fluid flow past a laser, and the response (scattering of light) reveals information about the cell, which may be explained by its features (e.g., gene expression). However, the order of the cells as they pass the laser is unknown, so we are in a shuffled regression setting (Abid et al., 2017). In "image stitching" (related to the so-called pose-correspondence problem) one wants to find the unknown correspondence between point clouds constructed from multiple camera angles of the same image (Pananjady et al., 2018). Several other motivating examples are discussed by Pananjady et al. (2018) (in the context of permuted/shuffled linear regression).

The method of Carpentier and Schlüter (2016) is based on the fact that monotone unlinked regression can be rephrased as the so-called deconvolution problem, as is apparent from (2). Below, we will discuss links to this problem in more detail. Note that, as is also true in the deconvolution setting, m_0 is not identifiable if we do not know the distribution of the noise ϵ (or have at least an estimate thereof). The following simple example explains why. Suppose that $m_0(x) = 2x$ and $X \sim \mathcal{N}(0, 1)$ is independent of $\epsilon \sim \mathcal{N}(0, 1)$. Then, this model is the same as $m_0(x) = x$ and $X \sim \mathcal{N}(0, 1)$ independent of $\epsilon \sim \mathcal{N}(0, 4)$. Let F_0 be the cumulative distribution function (CDF) of X and L_0 the

CDF of $m_0(X)$. If both m_0 and F_0 are assumed to be one-to-one, then L_0 satisfies

$$L_0(w) = P(m_0(X) \leq w) = F_0 \circ m_0^{-1}(w) \quad (3)$$

for all w in the range of m_0 , and we have

$$m_0 = L_0^{-1} \circ F_0. \quad (4)$$

Thus, the estimator constructed by Carpentier and Schlüter (2016) takes the form

$$\tilde{m}(x) = \tilde{L}_{n_y}^{-1} \circ \mathbb{F}_{n_x}(x) \quad (5)$$

where \tilde{L}_{n_y} is an estimator of L_0 obtained by deconvolution methods and is based on the sample (Y_1, \dots, Y_{n_y}) and knowledge of the distribution of ϵ , and where \mathbb{F}_{n_x} can be taken for example to be the empirical distribution function of F_0 based on the sample X_1, \dots, X_{n_x} . Thus, the estimator in (5) at a point x is equal to the deconvolution estimator of the quantile of $m_0(X)$ corresponding to the random level $\mathbb{F}_{n_x}(x)$. In the presence of “contextual variables” (i.e., covariates that are paired with both X and Y), Carpentier and Schlüter (2016) gave some consistency and rate of convergence results in their Theorem 3.2 under the assumption that m_0 and F_0^{-1} belong to Hölder classes. Carpentier and Schlüter (2016) do not discuss how to choose the optimal bandwidth, as the main focus in that paper is to show how their estimation approach can be easily implemented for real datasets.

Beside Carpentier and Schlüter (2016), the only other work of which we are aware on a similar problem is the very recent article of Rigollet and Weed (2019). In their setting, the authors consider a shuffled monotone regression model in a fixed design setting but use the term “uncoupled” to describe the problem. In fact, the authors do not make any attempt to recover the unknown permutation, and focus entirely, as we do in this paper, on estimating the unknown regression function. Rigollet and Weed (2019) assume that the known distribution of the noise is sub-exponential and the true monotone function is bounded by some known constant, but they do not make any smoothness assumption on that function. Using the Wasserstein’s distance and arguments from optimal transport, they showed that their estimator converges to the truth at a rate no slower than $\log \log n / \log n$ and that this rate is minimax when the distribution of the noise is Gaussian. Although Rigollet and Weed (2019) describe in their Section 2.2 an algorithm for computing their monotone estimator, the authors do not present simulation results. It comes as a surprise that the rate of convergence they obtain does not depend on the smoothness of the distribution of the noise, whereas optimal rates of convergence in the deconvolution problem are closely connected to this smoothness; see e.g. Fan (1991). For instance, Rigollet and Weed (2019)’s rate applies in both cases of a double exponential and a Gaussian error which are of different smoothness: the former is ordinary smooth of order 1 while the latter is supersmooth of order 2 in Fan’s terminology; see Section 1 in Fan (1991).

It is worth mentioning that more research seems to have been done in the shuffled (permuted) linear regression model than in the monotone regression model. We refer here to the work of Abid et al. (2017), Pananjady et al. (2017), Pananjady et al. (2018), and Unnikrishnan et al. (2018). The main focus in the former three papers is to find conditions on the signal-to-noise ratio that guarantee recovery of the unknown permutation. Abid et al. (2017) show that the least-squares estimator in this model is inconsistent in general, but they construct a method-of-moments type estimator and

prove that this estimator is consistent assuming that $E(X_i) \neq 0$. A common feature of these works is to restrict attention to the case of Gaussian noise. As we show in the present paper, the noise distribution may be of fundamental importance in these problems.

As announced above, we give now some more detail to the existing link between unlinked monotone regression and estimation in deconvolution problems. As noted earlier, (4) shows clearly that there is a tight connection to quantile estimation in a deconvolution setting. In fact, consider the deconvolution setting in which one observes n i.i.d. copies of Y where $Y = X + \epsilon$ for independent random variables X and ϵ . In this problem, the goal is learn the distribution of X under the assumption that the distribution of ϵ is known (such an assumption can be relaxed if this distribution can be estimated). Estimation of the distribution and quantile functions of X has been considered in Hall and Lahiri (2008), and revisited in Dattner et al. (2011, 2016) under slightly different assumptions. In particular, contrary to Hall and Lahiri (2008), no moment assumptions are made about the covariate X or ϵ in Dattner et al. (2011, 2016). There, the smoothness of the density of X is measured in terms of belonging to Sobolev or Hölder balls. In the case where the error is ordinary smooth of order larger than $1/2$, Dattner et al. (2011) recover the rates of convergence that Hall and Lahiri (2008) obtained for the integrated risk when estimating the distribution function, and moreover provide new rates of convergence for the case of smoother error distributions; the square-root rate is shown to be achieved for smooth enough distribution of X . The convergence results obtained in these previous papers do not apply directly in this present paper, as we do not assume that the covariate X (from (2)) admits even a Lebesgue density, which also means that $m_0(X)$ is not assumed to have a density.

While Carpentier and Schlüter (2016) and Rigollet and Weed (2019) are the only other articles of which we are aware on unlinked monotone regression besides the present paper, the classical isotonic regression model given in (1) is a very well-known estimation problem with a vast literature. The most known estimator in this problem is certainly the Grenander-type estimator, obtained by taking the right derivative of the greatest convex majorant of the cumulative sum diagram associated with the data $(X_i, Y_i), i = 1, \dots, n$ (Barlow et al., 1972; Robertson et al., 1988; Groeneboom and Jongbloed, 2014). The pointwise non-standard rate of convergence of the Grenander estimator is $n^{-1/3}$ if m_0 is continuously differentiable with a non-vanishing derivative and \sqrt{n} in case m_0 is locally flat (Groeneboom, 1983; Carolan and Dykstra, 1999; Zhang, 2002; Cator, 2011; Chatterjee et al., 2015). Asymptotic properties, including the pointwise limit distribution and convergence in the L_p -norms for $p \in [1, 5/2) \cup \{\infty\}$ have been fully described in Brunk (1970), Durot (2002), Durot (2007) and Durot et al. (2012); see also Groeneboom (1985, 1989). One can also combine kernel estimation with the monotonicity constraint to improve rates of convergence if m_0 has higher orders of smoothness (Mammen, 1991; Durot and Lopuhaä, 2014).

In this paper, we investigate the unlinked monotone regression following the method introduced by Edelman (1988) for estimating the mixing distribution in a mixture problem with Normal noise. Let $(X_1, \theta_1), \dots, (X_n, \theta_n)$ be independent random vectors such that $X_i | \theta_i \sim \mathcal{N}(\theta_i, 1)$, that is, conditionally on $\theta_1, \dots, \theta_n$ the random variables X_1, X_2, \dots, X_n are generated from the unknown distribution $\Phi(\cdot - \theta_1), \Phi(\cdot - \theta_2), \dots, \Phi(\cdot - \theta_n)$, where Φ denotes the cumulative distribution function of $\mathcal{N}(0, 1)$. The approach of Edelman (1988) consists of finding the vector $(\tilde{\theta}_1, \dots, \tilde{\theta}_n)$ which minimizes the integrated difference between $n^{-1} \sum_{i=1}^n \Phi(\cdot - \theta_i)$ and the empirical distribution \mathbb{F}_n

based on the observations $X_i, i = 1, \dots, n$ among all vectors $(\theta_1, \dots, \theta_n)$. As Edelman (1988) already noted, the normal distribution can be replaced by other distributions, which is exactly what we do here. The merits of this approach are the facts that it does not depend on some bandwidth, and that it is easily implementable. The link between our problem and the work Edelman (1988) is made clear in Section 2. There, we introduce our estimator, which we call the minimum contrast estimator, and we establish its existence and rate of convergence under some fairly general conditions on the distribution function of the covariates. In that section, we assume that the noise distribution is known and is also ordinary smooth with a given smoothness parameter $\beta \in (0, \infty)$. The rate of convergence of our estimator is obtained after converting a consistency result about the generalized inverse of this estimator, an interesting result in its own right. We first focus on the case of equal sample sizes $n_x = n_y$ and then extend the construction and rate to the case of different sample sizes. Although our estimator cannot be shown to be unique, we show in Section 3 that any solution has to satisfy a necessary optimizing condition. This condition is used to develop a gradient-descent algorithm to compute the estimator, see Section 4. In Section 5, we discuss how our method can be extended to the more realistic situation where the noise distribution is unknown. In Section 6, we illustrate our approach through synthetic and real data. We finish this manuscript with some concluding remarks and future research directions; see Section 7. Technical proofs are deferred to an Appendix.

2. The minimum contrast estimator

2.1 Setup, terminology, and notation

Let m_0 be the monotone function appearing in the model in (2), and in which we are interested. In the model that we consider, the response Y has the same distribution as the convolution of $m_0(X)$ and the noise ϵ with m_0 monotone non-decreasing. Note for the case where m_0 is non-increasing it is enough to consider $-Y$ instead of Y and all our results will still apply. Denote by \mathcal{M} the set of all bounded non-decreasing and right continuous functions defined on $[0, 1]$. This class accomodates for the assumption made in the sequel that the covariate $X \in [0, 1]$ almost surely.

Here, we describe the working framework of our estimation approach. We observe two independent samples (X_1, \dots, X_n) and (Y_1, \dots, Y_n) such that the following holds.

Assumption A0. $X_1, \dots, X_n \stackrel{iid}{\sim} F_0$, $0 \leq X_i \leq 1$ almost surely, and $Y_1, \dots, Y_n \stackrel{iid}{\sim} H_0$. The unobserved error ϵ is independent of X , satisfies $E(\epsilon) = 0$, and has CDF Φ_ϵ .

Thus F_0 , H_0 , and Φ_ϵ are the true CDF's of X_i , Y_i , and ϵ_i , respectively. Let $\mathbb{F}_n(x) = n^{-1} \sum_{i=1}^n \mathbb{1}_{[X_i, \infty)}(x)$ and $\mathbb{H}_n(x) = n^{-1} \sum_{i=1}^n \mathbb{1}_{[Y_i, \infty)}(x)$ where $x \mapsto \mathbb{1}_A(x)$ is the indicator function for the set A . Also, let $\|\cdot\|_\infty$ be the supremum norm, i.e. $\|m\|_\infty = \sup_{t \in [0, 1]} |m(t)|$. For $K > 0$, let \mathcal{M}_K be the set of functions $m \in \mathcal{M}$ such that $\|m\|_\infty \leq K$. For $m \in \mathcal{M}$, m^{-1} denotes the *generalized inverse* of m , i.e.

$$m^{-1}(y) := \inf \{x \in [0, 1] : m(x) \geq y\} \quad (6)$$

where the infimum of an empty set is defined to be 1. Hence, we have $m^{-1}(y) = 1$ for all $y > m(1)$. Here as is customary, $X_{(1)} \leq \dots \leq X_{(n)}$ denote the order statistics corresponding to X_1, \dots, X_n .

In this section, we assume that Φ_ϵ is known. This assumption will be relaxed in Section 5. Also, although we take the respective sizes of the samples of covariates

and responses to be equal, our method can be easily adapted to the case where these sizes are different. In that case, the convergence rate of our estimator is driven by the minimum of the sample sizes; see Subsection 2.4.

Let \mathcal{F} be the set of all distribution functions on \mathbb{R} . A contrast function \mathcal{C} defined on the Cartesian product $\mathcal{F} \times \mathcal{F}$ is any non-negative function such that $\mathcal{C}(F_1, F_2) = 0$ if and only if $F_1 = F_2$. Consider the estimator

$$\hat{m}_n = \operatorname{argmin}_{m \in \mathcal{M}} \mathcal{C} \left(\mathbb{H}_n, n^{-1} \sum_{i=1}^n \Phi_\epsilon(\cdot - m(X_i)) \right) \quad (7)$$

provided that a minimizer exists. Since the criterion depends on m only through its values at the observations $X_i, i = 1, \dots, n$, it follows that any candidate for the minimization problem in (7), m in \mathcal{M} , can be well replaced by the non-decreasing function \tilde{m} such that

$$\tilde{m}(t) = \begin{cases} m(X_{(1)}), & \text{for } t \in [0, X_{(1)}] \\ m(X_{(n)}), & \text{for } t \in [X_{(n)}, 1] \end{cases}$$

and \tilde{m} is constant between the remaining order statistics such that \tilde{m} is right continuous (by definition of \mathcal{M}).

In addition, note that \hat{m}_n does not have to be unique at the data points and thus \hat{m}_n denotes any solution of the minimization problem.

2.2 Existence

In this sequel, we consider the following contrast function

$$\mathcal{C}(F_1, F_2) = \int_{\mathbb{R}} \left\{ F_1(y) - F_2(y) \right\}^2 dy$$

whenever this integral is finite, which is the case when F_1 and F_2 are distribution functions of random variables admitting finite expectations. The choice of such contrast function is mainly motivated by application of the Parseval-Plancherel's Theorem. The estimator we consider here is reminiscent of the one studied in Edelman (1988) for deconvoluting a distribution function from a Gaussian noise. However, our goal here is different since the main target in our problem is the monotone transformation m_0 .

Before starting the analysis of the estimator, we establish first its existence. Denote

$$\mathbb{M}_n(m) = \int_{\mathbb{R}} \left\{ \mathbb{H}_n(y) - n^{-1} \sum_{i=1}^n \Phi_\epsilon(y - m(X_i)) \right\}^2 dy$$

and let

$$\mathbb{M}(m) = \int_{\mathbb{R}} \left\{ H_0(y) - \int_{\mathbb{R}} \Phi_\epsilon(y - m(x)) dF_0(x) \right\}^2 dy$$

be its deterministic counterpart. Then the minimizer \hat{m}_n in (7) (if it exists) can also be written as

$$\hat{m}_n := \operatorname{argmin}_{m \in \mathcal{M}} \mathbb{M}_n(m). \quad (8)$$

The following assumptions will be needed.

Assumption A1. For some $K_0 \in [0, \infty)$, we have $\|m_0\|_\infty = K_0$.

Assumption A2. The distribution function Φ_ϵ is continuous on \mathbb{R} .

Proposition 2.1. Let Assumption A0 hold. Then,

1. $\mathbb{M}_n(m)$ is finite for any $m \in \mathcal{M}$ for all $n \geq 1$;
2. If Assumption A1 also holds then $\mathbb{M}(m)$ is finite for any $m \in \mathcal{M}$;
3. If Assumptions A1 and A2 also hold, then there exists at least a solution to (8) that is piecewise constant and right-continuous, for all $n \geq 1$.

Note that part 3 of the proposition gives existence but not necessarily uniqueness of \hat{m}_n . In fact, it can be seen from the proof of Proposition 2.1 that if \hat{m}_n is a solution to (8), then any monotone function that coincides with \hat{m}_n at the observed covariates X_1, \dots, X_n gives another solution to (8). In what follows, we will consider a solution \hat{m}_n that takes constant values between successive covariates and that is right continuous. This choice is consistent with the way the Grenander-type estimator is defined, that is, the estimator in the classical monotone regression estimation problem.

Proof Recall that any element $m \in \mathcal{M}$ is bounded, and hence there exists $K > 0$ such that $\|m\|_\infty \leq K$. Denote by $Y_{(1)} \leq \dots \leq Y_{(n)}$ the order statistics corresponding to Y_1, \dots, Y_n . We have for all $y < Y_{(1)}$ that $\mathbb{H}_n(y) = 0$. Moreover, it follows from monotonicity of Φ_ϵ and m that $0 \leq \Phi_\epsilon(y - m(X_i)) \leq \Phi_\epsilon(y - m(X_{(1)})) \leq 1$ for all $i \in \{1, \dots, n\}$ and therefore,

$$\begin{aligned} \left\{ \mathbb{H}_n(y) - n^{-1} \sum_{i=1}^n \Phi_\epsilon(y - m(X_i)) \right\}^2 &= n^{-2} \left(\sum_{i=1}^n \Phi_\epsilon(y - m(X_i)) \right)^2 \\ &\leq \Phi_\epsilon(y - m(X_{(1)}))^2 \\ &\leq \Phi_\epsilon(y - m(X_{(1)})). \end{aligned}$$

Now, existence of expectation of ϵ implies that

$$\int_{-\infty}^c \Phi_\epsilon(t) dt < \infty, \quad \text{and} \quad \int_c^\infty (1 - \Phi_\epsilon(t)) dt < \infty \quad (9)$$

for arbitrary $c \in \mathbb{R}$ and therefore,

$$\begin{aligned} \int_{-\infty}^{Y_{(1)}} \left\{ \mathbb{H}_n(y) - n^{-1} \sum_{i=1}^n \Phi_\epsilon(y - m(X_i)) \right\}^2 dy &\leq \int_{-\infty}^{Y_{(1)}} \Phi_\epsilon(y - m(X_{(1)})) dy \\ &= \int_{-\infty}^{Y_{(1)} + m(X_{(1)})} \Phi_\epsilon(y) dy \\ &< \infty. \end{aligned} \quad (10)$$

Similarly, $\mathbb{H}_n(y) = 1$ for $y > Y_{(n)}$ and hence

$$\begin{aligned} \left\{ \mathbb{H}_n(y) - n^{-1} \sum_{i=1}^n \Phi_\epsilon(y - m(X_i)) \right\}^2 &= \left(1 - n^{-1} \sum_{i=1}^n \Phi_\epsilon(y - m(X_i)) \right)^2 \\ &\leq \left(1 - \Phi_\epsilon(y - m(X_{(n)})) \right)^2 \\ &\leq 1 - \Phi_\epsilon(y - m(X_{(n)})). \end{aligned}$$

Combined with (9), this proves that

$$\int_{Y_{(n)}}^{\infty} \left\{ \mathbb{H}_n(y) - n^{-1} \sum_{i=1}^n \Phi_{\epsilon}(y - m(X_i)) \right\}^2 dy < \infty. \quad (11)$$

Since the integrand is bounded, (10) and (11) yield

$$\int_{\mathbb{R}} \left\{ \mathbb{H}_n(y) - n^{-1} \sum_{i=1}^n \Phi_{\epsilon}(y - m(X_i)) \right\}^2 dy < \infty,$$

which proves that \mathbb{M}_n is well defined on \mathcal{M} .

We turn now to \mathbb{M} . Assume that Assumption A1 holds and consider $m \in \mathcal{M}$. Then, we have that

$$\begin{aligned} \mathbb{M}(m) &= \int_{\mathbb{R}} \left\{ \int_{\mathbb{R}} \left(\Phi_{\epsilon}(y - m_0(x)) - \Phi_{\epsilon}(y - m(x)) \right) dF_0(x) \right\}^2 dy \\ &= \int_{[0, \infty)} \left\{ \int_{\mathbb{R}} \left(1 - \Phi_{\epsilon}(y - m(x)) - (1 - \Phi_{\epsilon}(y - m_0(x))) \right) dF_0(x) \right\}^2 dy \end{aligned} \quad (12)$$

$$+ \int_{(-\infty, 0]} \left\{ \int_{\mathbb{R}} \left(\Phi_{\epsilon}(y - m_0(x)) - \Phi_{\epsilon}(y - m(x)) \right) dF_0(x) \right\}^2 dy. \quad (13)$$

We further bound above the integral in (12) by

$$\begin{aligned} &\leq 2 \int_{[0, \infty)} \left\{ \int_{\mathbb{R}} (1 - \Phi_{\epsilon}(y - m(x))) dF_0(x) \right\}^2 dy \\ &+ 2 \int_{[0, \infty)} \left\{ \int_{\mathbb{R}} (1 - \Phi_{\epsilon}(y - m_0(x))) dF_0(x) \right\}^2 dy \\ &\leq 4 \int_{[0, \infty)} \left(1 - \Phi_{\epsilon}(y - \max(K_0, K)) \right)^2 dy \end{aligned} \quad (14)$$

where we recall that $K \geq \|m\|_{\infty}$. The latter integral is finite using again (9). Similarly, we argue that the integral in (13) can be also bounded above by

$$4 \int_{(-\infty, 0]} \left(\Phi_{\epsilon}(y + \max(K_0, K)) \right)^2 dy < \infty.$$

This completes the proof that $\mathbb{M}(m)$ is finite.

Now, we show existence of a minimizer, as defined in (8). Using again that $\mathbb{H}_n(y) = 0$ for all $y < Y_{(1)}$, together with monotonicity of Φ_{ϵ} and m , we have that

$$\begin{aligned} \mathbb{M}_n(m) &\geq \int_{-\infty}^{Y_{(1)}} n^{-2} \left(\sum_{i=1}^n \Phi_{\epsilon}(y - m(X_i)) \right)^2 dy \\ &\geq n^{-2} \int_{-\infty}^{Y_{(1)}} \Phi_{\epsilon}(y - m(X_{(1)}))^2 dy \\ &= n^{-2} \int_{-\infty}^{Y_{(1)} - m(X_{(1)})} \Phi_{\epsilon}(t)^2 dt \\ &\rightarrow \infty, \text{ if } m(X_{(1)}) \rightarrow -\infty. \end{aligned}$$

Similarly, for $y \geq Y_{(n)}$, it holds that

$$\begin{aligned} \mathbb{M}_n(m) &\geq n^{-2} \int_{Y_{(n)}}^{\infty} (1 - \Phi_{\epsilon}(y - m(X_{(n)})))^2 dy \\ &\geq n^{-2} \int_{Y_{(n)} - m(X_{(n)})}^{\infty} (1 - \Phi_{\epsilon}(t))^2 dt \\ &\rightarrow \infty, \text{ if } m(X_{(n)}) \rightarrow \infty. \end{aligned}$$

Hence, there exists some $K > 0$ (which may depend on n) such that any candidate $m \in \mathcal{M}$ for the minimization problem in (8) should satisfy $-K \leq m(X_{(1)}) \leq \dots \leq m(X_{(n)}) \leq K$. By identifying an element $m \in \mathcal{M}_K$ by the corresponding vector $\theta = (m(X_{(1)}), \dots, m(X_{(n)}))^T$, it is easy to see that the original minimization problem is equivalent to minimizing

$$\tilde{\mathbb{M}}_n(\theta) =: \int_{\mathbb{R}} \left\{ \mathbb{H}_n(y) - n^{-1} \sum_{i=1}^n \Phi_{\epsilon}(y - \theta_i) \right\}^2 dy$$

on the compact finite dimensional subset

$$\mathcal{S}_K =: \{(\theta_1, \dots, \theta_n)^T \in \mathbb{R}^n : -K \leq \theta_1 \leq \dots \leq \theta_n \leq K\}.$$

Now, the function $\tilde{\mathbb{M}}_n$ is continuous on \mathcal{S}_K since for any sequence $(\theta_p)_{p \geq 0}$ in $\mathcal{S}_K^{\mathbb{N}}$ converging (in any distance) to $\theta \in \mathcal{S}_K$, the sequence of functions

$$y \mapsto \left(\mathbb{H}_n(y) - n^{-1} \sum_{i=1}^n \Phi_{\epsilon}(y - \theta_{p,i}) \right)^2$$

converges pointwise by continuity of Φ_{ϵ} (see Assumption A2) to the limit

$$y \mapsto \left(\mathbb{H}_n(y) - n^{-1} \sum_{i=1}^n \Phi_{\epsilon}(y - \theta_i) \right)^2. \quad (15)$$

Also, for $y \in \mathbb{R}$, we have that (15) is no larger than

$$\begin{cases} (n^{-1} \sum_{i=1}^n \Phi_{\epsilon}(y + K))^2, & \text{for } y < Y_{(1)} \\ 4, & \text{for } Y_{(1)} \leq y \leq Y_{(n)} \\ \left(1 - n^{-1} \sum_{i=1}^n \Phi_{\epsilon}(y - K) \right)^2, & \text{for } y > Y_{(n)} \end{cases}$$

where the function on the right side can be shown to be integrable using similar arguments as above. By the Lebesgue dominated convergence theorem, it follows that

$$\lim_{p \rightarrow \infty} \tilde{\mathbb{M}}_n(\theta_p) = \tilde{\mathbb{M}}_n(\theta).$$

Thus, $\tilde{\mathbb{M}}_n$ admits at least a minimizer in \mathcal{S}_K , $\hat{\theta}_n$ say. We conclude that \mathbb{M}_n admits at least a minimizer \hat{m}_n which is bounded by K , and such that $(\hat{m}_n(X_1), \dots, \hat{m}_n(X_n))^T = \hat{\theta}_n$. The values of the minimizer being given by $\hat{\theta}_n$ at the observed covariates X_1, \dots, X_n , any monotone interpolation of these values gives a solution to (8). In particular, there exists a solution \hat{m}_n that takes constant values between successive covariates and that is right continuous. This completes the proof. \blacksquare

2.3 Convergence rate

In the following, we use the notation

$$\psi_F(x) = \int_{\mathbb{R}} e^{itx} dF(t) \quad (16)$$

for any distribution function F on \mathbb{R} , and

$$\phi_g(x) = \int_{\mathbb{R}} e^{itx} g(t) dt \quad (17)$$

the Fourier transform of g , whenever g is integrable. Note that when F is absolutely continuous with density f , then it follows immediately that $\psi_F = \phi_f$.

In this section, we assume that the Assumptions A0, A1 and A2 hold. By Proposition 2.1, this guarantees that the minimization problem in (8) admits a piecewise constant and right-continuous solution. From now on, we define \hat{m}_n to be any such solution. In this section, we derive the convergence rate of \hat{m}_n , and for this task we need some smoothness assumptions on the distributions of the design points and the error terms.

Assumption A3. *The common distribution function F_0 of the covariates X_1, \dots, X_n is continuous.*

Assumption A4. *The distribution function Φ_ϵ is absolutely continuous with even (zero-symmetric) ordinary smooth density f_ϵ in the sense that*

$$\frac{d_0}{|t|^\beta} \leq |\phi_{f_\epsilon}(t)| \leq \frac{d_1}{|t|^\beta} \quad (18)$$

as $|t| \rightarrow \infty$, for some $\beta > 0$ and constants $d_0 > 0, d_1 > 0$.

Assumption A4 is common in deconvolution problems, see for instance Dattner et al. (2011). The symmetry assumption ensures that the function ϕ_{f_ϵ} is real valued. The positive real number β is usually referred to as the order of smoothness. Known examples include the Laplace distribution (with $\beta = 1$) and more generally symmetric Gamma distributions. See for example the examples given in Fan (1991) after (1.4).

We consider below the L_1 -error over a given interval $[a, b] \subset (0, 1)$ with respect to the distribution of the X_i 's: if X denotes a random variable with the same distribution as the X_i 's that is independent of the X_i , and E^X denotes the expectation with respect to X , then the estimation error that we consider is

$$\int_a^b |\hat{m}_n(x) - m_0(x)| dF_0(x) = E^X [|\hat{m}_n(X) - m_0(X)| \mathbb{1}_{\{X \in [a, b]\}}].$$

Our main theorem below provides the rate of convergence of the above L_1 -error.

Assumption A5. *Assume that there exists T^* such that $|\phi_{f_\epsilon}(t)| \geq |\phi_{f_\epsilon}(T)| > 0$ for all $T > T^*$ and $|t| \leq T$.*

Theorem 2.1. *Suppose that the Assumptions A0 to A5 hold. Let $[a, b] \subset (0, 1)$ be a fixed interval such that $0 < F_0(a) \leq F_0(b) < 1$. Then it holds that*

$$\widehat{m}_n(a) = O_p(1) \text{ and } \widehat{m}_n(b) = O_p(1). \quad (19)$$

Moreover,

$$\int_a^b |\widehat{m}_n(x) - m_0(x)| dF_0(x) = O_p(n^{-1/(2(2\beta+1))}).$$

We give below the main steps of the proof. Details are postponed to Section D.1. Let us define

$$\widehat{\mathbb{L}}_n(w) := \frac{1}{n} \sum_{i=1}^n \mathbb{1}_{\{\widehat{m}_n(X_i) \leq w\}} \text{ and } \mathbb{L}_n(w) := \frac{1}{n} \sum_{i=1}^n \mathbb{1}_{\{m_0(X_i) \leq w\}} \quad (20)$$

for all $w \in \mathbb{R}$. Using deconvolution arguments we show closeness of the two processes in the following proposition.

Proposition 2.2. *Assume that Assumptions A4 and A5 hold. Then,*

$$E \left[\int_{\mathbb{R}} \left(\widehat{\mathbb{L}}_n(w) - \mathbb{L}_n(w) \right)^2 dw \right] = O(n^{-1/(2\beta+1)}).$$

Next, using arguments from empirical process theory, and in particular entropy arguments, we show in the following proposition that on possibly random intervals $[A, B]$, the empirical processes $\widehat{\mathbb{L}}_n$ and \mathbb{L}_n are close to their population counterparts \widehat{L}_n^0 and L_0 respectively, where for all $w \in \mathbb{R}$,

$$\widehat{L}_n^0(w) = \int \mathbb{1}_{\{\widehat{m}_n(x) \leq w\}} dF_0(x) \text{ and } L_0(w) = \int \mathbb{1}_{\{m_0(x) \leq w\}} dF_0(x), \quad (21)$$

and we derive the rate of convergence of the squared distance between the population counterparts on intervals $[A, B]$.

Proposition 2.3. *Under Assumptions A3 and A4, for all random variables $A < B$ (that may depend on n) it holds that*

$$\int_A^B \left(\widehat{\mathbb{L}}_n(w) - \widehat{L}_n^0(w) \right)^2 dw \leq (B - A) O_p(1/n),$$

$$\int_A^B \left(\mathbb{L}_n(w) - L_0(w) \right)^2 dw \leq (B - A) O_p(1/n),$$

where $O_p(1/n)$ is uniform in A and B . Moreover, if $B - A = O_p(n^{2\beta/(2\beta+1)})$ and Assumption A5 holds, then

$$\int_A^B \left(\widehat{L}_n^0(w) - L_0(w) \right)^2 dw = O_p(n^{-1/(2\beta+1)}). \quad (22)$$

The following proposition makes the connection between the above error and a squared distance between the inverse functions of \widehat{m}_n and m_0 composed with the distribution function F_0 of the X_i 's; a rate of convergence of that squared distance is derived. We recall that for $m \in \mathcal{M}$, the inverse of m is defined by (6), where the infimum of an empty set is defined to be 1.

Proposition 2.4. *Under Assumptions A3, A4, and A5 for all random variables $A < B$ such that $B - A = O_p(1)$ it holds that*

$$\begin{aligned} \int_A^B (F_0 \circ \widehat{m}_n^{-1}(w) - F_0 \circ m_0^{-1}(w))^2 dw &= \int_A^B (\widehat{L}_n^0(w) - L_0(w))^2 dw \\ &= O_p(n^{-1/(2\beta+1)}). \end{aligned}$$

The last step is to make the connection between the above squared distance and the L_1 -error of \widehat{m}_n .

Proposition 2.5. *Suppose that the Assumptions A3 and A4 hold. Let $[a, b] \subset (0, 1)$ be a fixed interval such that $0 < F_0(a) \leq F_0(b) < 1$. Then it holds that*

$$\int_a^b |\widehat{m}_n(x) - m_0(x)| dF_0(x) \leq \left((B_n - A_n) \int_{A_n}^{B_n} (F_0 \circ \widehat{m}_n^{-1}(x) - F_0 \circ m_0^{-1}(x))^2 dx \right)^{1/2}$$

where $A_n = m_0(a) \wedge \widehat{m}_n(a)$ and $B_n = m_0(b) \vee \widehat{m}_n(b)$.

Finally, we note that the same proofs also apply to the case of a supersmooth noise distribution. In this case, Assumption A4 should be replaced by the following.

Assumption A4'. *The distribution function Φ_ϵ is absolutely continuous with even (zero-symmetric) ordinary supersmooth density f_ϵ in the sense that*

$$d_0 |t|^\alpha \exp(-|t|^\beta/\gamma) \leq |\phi_{f_\epsilon}(t)| \leq d_1 |t|^\alpha \exp(-|t|^\beta/\gamma)$$

as $|t| \rightarrow \infty$, for some $\alpha > 0, \beta > 0$ and constants $d_0 > 0, d_1 > 0$.

In the above definition we provide for supersmoothness we deviate from the one given by Fan (1991) in (1.3) by taking the same exponent α in the lower and upper bound, for simplicity.

Remark 1. *Under the same Assumptions A0–A3, A4', and A5, we have for any $[a, b] \subset (0, 1)$ such that $0 < F_0(a) \leq F_0(b) < 1$ that*

$$\int_a^b |\widehat{m}_n(x) - m_0(x)| dF_0(x) = O_p((\log n)^{-1/\beta}).$$

For instance, in the case of a Gaussian noise, in which case $\beta = 2$, the rate is $1/\sqrt{\log n}$, which matches the conclusion of Edelman (1988) right after the proof of his Theorem 1. On the other hand, this rate is slower than the minimax rate, $\log \log n / \log n$, obtained by Rigollet and Weed (2019) (also under the assumption that the true monotone regression function is bounded). One surprising feature is that the rate $\log \log n / \log n$ found by the authors does not depend on the smoothness of the noise distribution. Such a dependence is seen in our results, whether the noise is ordinary smooth or supersmooth. For a proof of the rate given in Remark 1 we refer to Appendix D.

2.4 Convergence rate in the case of different sample sizes

In this section we briefly consider the case where one observes Y_1, \dots, Y_{n_y} and X_1, \dots, X_{n_x} with possibly different sample sizes $n_x \neq n_y$. In that case, the estimator is defined as

$$\hat{m}_{n_x, n_y} = \operatorname{argmin}_{m \in \mathcal{M}} \int_{\mathbb{R}} \left\{ \mathbb{H}_{n_y}(y) - n_x^{-1} \sum_{i=1}^{n_x} \Phi_{\epsilon}(y - m(X_i)) \right\}^2 dy$$

where \mathbb{H}_{n_y} denotes the empirical distribution function corresponding to the sample Y_1, \dots, Y_{n_y} . The asymptotic here has to be understood in the sense that both sample sizes n_x and n_y go to infinity. This means that $n_x \wedge n_y \rightarrow \infty$, where $n_x \wedge n_y$ denotes the infimum between n_x and n_y .

Theorem 2.2. *Suppose that the Assumptions A0, A1, A2, A3, A4 and A5 hold. Let $[a, b] \subset (0, 1)$ be a fixed interval such that $0 < F_0(a) \leq F_0(b) < 1$. Then it holds that as $n_x \wedge n_y \rightarrow \infty$,*

$$\hat{m}_{n_x, n_y}(a) = O_p(1) \text{ and } \hat{m}_{n_x, n_y}(b) = O_p(1).$$

Moreover,

$$\int_a^b |\hat{m}_{n_x, n_y}(x) - m_0(x)| dF_0(x) = O_p((n_x \wedge n_y)^{-1/(2(2\beta+1))}).$$

3. Fenchel optimality conditions

In view of the computational section below, we derive in this section the optimality conditions related to the optimization problem defining the estimator \hat{m}_n . Recall that by Assumption A0, the covariates X_1, \dots, X_n are assumed to belong to $[0, 1]$. In the following, we denote by \hat{m}_n a piecewise constant and right-continuous solution to (8), see Proposition 2.1. For $1 \leq p \leq n$, we write $\hat{m}_1 < \dots < \hat{m}_p$ for the distinct values taken by \hat{m}_n on $[0, 1]$.

We will use the following assumption on the density f_{ϵ} .

Assumption A6. *The density f_{ϵ} is continuously differentiable such that*

$$\sup_{t \in \mathbb{R}} |f'_{\epsilon}(t)| \leq D,$$

for some constant $D > 0$, and $\int_{\mathbb{R}} |f'_{\epsilon}(t)| dt < \infty$.

Proposition 3.1. *Let $\hat{m}_1 < \dots < \hat{m}_p$ be the distinct values of the estimator \hat{m}_n . Let Assumption A6 hold. Then, for any $k \in \{1, \dots, p\}$*

$$\int_{\mathbb{R}} \left(\mathbb{H}_n(y) - n^{-1} \sum_{i=1}^n \Phi_{\epsilon}(y - \hat{m}_n(X_i)) \right) f_{\epsilon}(y - \hat{m}_k) dy = 0. \quad (23)$$

Furthermore, this condition can be equivalently re-written as

$$1 - \frac{1}{n} \sum_{i=1}^n \Phi_{\epsilon}(Y_i - \hat{m}_k) = \frac{1}{n} \sum_{i=1}^n \int_{\mathbb{R}} \Phi_{\epsilon}(y - \hat{m}_n(X_i)) f_{\epsilon}(y - \hat{m}_k) dy \quad (24)$$

for $k = 1, \dots, p$.

Remark 2. The alternative form in (24) gives a useful way of verifying numerically the second equality condition via numerical integration. Indeed, with $m = \hat{m}_n(X_i)$ and $m' = \hat{m}_k$, the integral on the right side of (24) takes the form

$$\int_{\mathbb{R}} \Phi_{\epsilon}(y - m) f_{\epsilon}(y - m') dy = B(m' - m),$$

where for all $m \in \mathbb{R}$, $B(m) = E\Phi_{\epsilon}(\epsilon + m) = \int \Phi_{\epsilon}(y) f_{\epsilon}(y - m) dy$. Explicit formulas of $B(m)$, for $m \in \mathbb{R}$, can be even found for some distributions such as Laplace or Gaussian; see Subsections C.2 and C.3.

Remark 3. Consider the case where \hat{m}_n takes one unique value, denoted \hat{m}_1 . Then the right side of (24) equals $\int_{\mathbb{R}} \Phi_{\epsilon}(y - \hat{m}_1) f_{\epsilon}(y - \hat{m}_1) dy = \int_{\mathbb{R}} \Phi_{\epsilon}(y) f_{\epsilon}(y) dy$ which equals $E(\Phi_{\epsilon}(\epsilon)) = E(U) = 1/2$ where U is $\text{Uniform}(0, 1)$.

4. Computation

Recall that our goal is to minimize \mathbb{M}_n . Write $\mathbf{m} := (m_1, \dots, m_n)$, so that

$$\mathbb{M}_n(\mathbf{m}) := \int_{\mathbb{R}} \left\{ \mathbb{H}_n - n^{-1} \sum_{i=1}^n \Phi_{\epsilon}(\cdot - m_i) \right\}^2 \quad (25)$$

(by a slight abuse of notation). To minimize \mathbb{M}_n , we can compute an unconstrained minimizer $\tilde{\mathbf{m}}$ of \mathbb{M}_n (i.e., we do not force $\tilde{m}_i \leq \tilde{m}_{i+1}$ for $i = 1, \dots, n-1$), and then the overall solution would be given by reordering the entries of $\tilde{\mathbf{m}}$ so that it is nondecreasing; i.e., $\hat{m}_n(X_{(i)}) := \tilde{m}_{(i)}$ (where $\tilde{m}_{(1)} \leq \dots \leq \tilde{m}_{(n)}$). The gradients $\frac{\partial}{\partial m_i} \mathbb{M}_n(\mathbf{m})$ can be computed using that

$$\frac{\partial}{\partial m_i} \mathbb{M}_n(\mathbf{m}) = 2n^{-1} - 2n^{-2} \sum_{\alpha=1}^n \{ \Phi_{\epsilon}(Y_{\alpha} - m_i) - B(m_i - m_{\alpha}) \} \quad (26)$$

where for all $m \in \mathbb{R}$, $B(m) = E\Phi_{\epsilon}(\epsilon + m)$, see Appendix C.1.

In Appendix C we show how to derive the gradient of \mathbb{M}_n when Φ_{ϵ} is either a Gaussian or a Laplace distribution. Thus, we can consider using a (first order) gradient descent algorithm for computation.

Because of the structure of the solution, the above naive gradient descent algorithm can be modified to improve its speed. In practice, we have found that the solution generally is piecewise constant in the sense that the entries of \mathbf{m} only contain a small (relative to n) number of unique values (although we don't have any quantitative theory for this observation yet). In Algorithm 1 we give a modified gradient descent algorithm, which makes use of this observation and of the fact that $\frac{\partial}{\partial m_i} \mathbb{M}_n(\mathbf{m})$ and $\frac{\partial}{\partial m_j} \mathbb{M}_n(\mathbf{m})$ are equal if $m_i = m_j$. In Algorithm 1 we use the notation $\mathbf{m}_{i:j}$ to denote the subvector of \mathbf{m} given by the indices $\{i, \dots, j\}$. In Appendix C.4 we provide a few comments about practical implementation of the algorithm.

5. Extension to the case of unknown noise distribution

5.1 Estimation of the noise distribution in the semi-supervised setting

In general, full knowledge of the distribution of ϵ might not be available which means that one needs to estimate it. In this case, it may be possible to collect a sample of ϵ 's,

Algorithm 1: Modified gradient descent (for piecewise constant functions)

input : $\mathbf{m}^{(0)} := (m_1^{(0)}, \dots, m_n^{(0)})$ (non-decreasing), Tolerance parameter ϵ ,
 Number iterations K , Stepsize η
output: $\hat{m}_n(X_{(1)}), \dots, \hat{m}_n(X_{(n)})$
 $n_1 \leftarrow n$;
 $\text{counts}^{(0)} \leftarrow (1, \dots, 1)$ (length n);
for $i \leftarrow 1$ **to** K **do**
 /* Group (approximately) non-unique entries */
 newidx $\leftarrow 1$, begidx $\leftarrow 1$;
 for $j \leftarrow 2$ **to** $n_i + 1$ **do**
 if $(j == n_i + 1)$ *OR* $(m_j^{(i-1)} - m_{\text{begidx}}^{(i-1)} > \epsilon)$ **then**
 $m_{\text{newidx}}^{(i)} \leftarrow \text{mean}(m_{\text{begidx}:(j-1)}^{(i-1)})$;
 $\text{counts}_{\text{newidx}}^{(i)} \leftarrow \text{sum of counts}_{\text{begidx}:(j-1)}^{(i-1)}$;
 begidx $\leftarrow j$;
 newidx $\leftarrow \text{newidx} + 1$;
 end
 $n_i \leftarrow \text{length of } \mathbf{m}^{(i)}$;
 end
 /* Compute gradient, take step (Appendix C has gradient computations) */
 Let $\nabla \mathbb{M}_n(\mathbf{m}^{(i)})$ be the vector containing $\frac{\partial}{\partial m_j^{(i)}} \mathbb{M}_n(\mathbf{m}^{(i)})$, $j = 1, \dots, n_i$;
 $\mathbf{m}^{(i)} \leftarrow \mathbf{m}^{(i)} - \eta \nabla \mathbb{M}_n(\mathbf{m}^{(i)})$;
end
 /* Reconstruct full length solution */
 The solution vector is given by the (unique) elements $m_i^{(K)}$, $i = 1, \dots, n_K$, each
 repeated $\text{counts}_i^{(K)}$ times, respectively.

$\epsilon_1^*, \dots, \epsilon_N^*$, from a separate data source. These can be used to construct an estimate of Φ_ϵ which can be then plugged into the objective function. The sample of ϵ 's does not necessarily need to be independent of the Y or X samples (note, for instance, Dattner et al. 2016). There are a variety of ways one may arrive at the sample of ϵ 's. In some cases, the main dataset may consist of unlinked covariates and responses, but there may be a smaller (or sub) dataset of linked/paired covariates and responses, $(X_1^*, Y_1^*), \dots, (X_N^*, Y_N^*)$. In this case, one may run the traditional monotone regression on this subset to obtain a monotone estimator \hat{m}_N^* from which one can compute the estimated residuals by $\hat{\epsilon}_i^* := Y_i^* - \hat{m}_N^*(X_i^*)$, $i = 1, \dots, N$. In general, the previously-described setting might be considered to be one of *semi-supervised learning*, where only a part of the data is unlinked. It would be useful with such data to learn from all of it simultaneously. This may be possible using the M-estimation framework we have proposed in this paper, but we leave an investigation of that question for future research.

5.2 Estimation of the noise distribution with longitudinal responses

Another common framework in which we may want to estimate Φ_ϵ from data is the one where we have repeated (or longitudinal) observations on the response Y . Assume we observe X_1, \dots, X_{n_x} as before and also $Y_{1,j}, \dots, Y_{n_y,j}$, where we take $j \in \{1, 2\}$ (for simplicity). We will impose the assumption that the distribution of ϵ is symmetric around 0. We also assume that $Y_{i,j} = m_0(\tilde{X}_i) + \epsilon_{i,j}$ (for some \tilde{X}_i which does not belong to our dataset and need not be observed), where $\epsilon_{i,1}$ is independent of $\epsilon_{i,2}$, and both are independent of all other error terms and all X variables. Then, as in Carroll et al. (2006) and Dattner et al. (2016), we can let $Y_i^* := (Y_{i,1} + Y_{i,2})/2 = m_0(\tilde{X}_i) + \epsilon'_i$ where $\epsilon'_i := (\epsilon_{i,1} + \epsilon_{i,2})/2$. If we let $\epsilon_i^* := (Y_{i,1} - Y_{i,2})/2 = (\epsilon_{i,1} - \epsilon_{i,2})/2$, then it follows from the assumption of 0-symmetry that $\epsilon_i^* \sim \epsilon'_i$. Then, we can use X_1, \dots, X_{n_x} and $Y_1^*, \dots, Y_{n_y}^*$ as our unlinked data and $\epsilon_1^*, \dots, \epsilon_{n_y}^*$ to estimate $\Phi_{\epsilon'}$.

Note that computing the estimator of m in practice as described in Algorithm 1 requires computation of the gradients $\frac{\partial}{\partial m_i} \mathbb{M}_n(\mathbf{m})$, where we use the same slight abuse of notation as in (25). The gradient depends on Φ_ϵ and is given by (26). Hence, in the case where Φ_ϵ is unknown, the gradient cannot be computed directly and has to be replaced by an appropriate estimator. In the setting of longitudinal responses, we have

$$\frac{\partial}{\partial m_i} \mathbb{M}_n(\mathbf{m}) = 2n^{-1} - 2n^{-2} \sum_{\alpha=1}^n \{\Phi_{\epsilon'}(Y_\alpha^* - m_i) - B(m_i - m_\alpha)\}$$

where $\Phi_{\epsilon'}$ denotes the common distribution function of $\epsilon'_1, \dots, \epsilon'_n$ and where for all $m \in \mathbb{R}$, $B(m) = E(\Phi_{\epsilon'}(\epsilon' + m))$. With $\hat{\Phi}$ the empirical distribution function based on the sample $\epsilon_1^*, \dots, \epsilon_{n_y}^*$, the gradient can be estimated by

$$2n^{-1} - 2n^{-2} \sum_{\alpha=1}^n \left\{ \hat{\Phi}(Y_\alpha^* - m_i) - \hat{B}(m_i - m_\alpha) \right\}$$

where $\hat{B}(m) = n_y^{-1} \sum_{i=1}^{n_y} \hat{\Phi}(\epsilon_i^* + m)$. The advantage of this estimator is that it does not require any choice of tuning parameters.

6. Demonstrations on synthetic and real data

6.1 Computations on synthetic data

In this subsection we present simulation studies for our method and compare our minimum contrast estimator to the deconvolution method of Carpentier and Schlüter (2016). We also compare to classical/linked (oracle) isotonic regression (which uses matching information that the other estimators do not use). We will use mean-squared errors (MSE's) for comparison: for an estimator \hat{m} , we report $n^{-1} \sum_{i=1}^n (\hat{m}(X_{(i)}) - m_0(X_{(i)}))^2$. There are 2 output tables, Tables 2 and 3, containing Monte Carlo estimates of MSE's. The sample sizes are taken to be $n = 100$ and $n = 1000$ for the first and second tables respectively. In both tables, we used 1000 Monte Carlo replications, and used $20 \times n$ gradient iterations for implementing our method using Algorithm 1. We used 5 different true mean functions m_0 . They are (up to translation and additive constants), together with the abbreviations that depict them, gathered in Table 1. In

m_0	Abbreviation
x	“lin”
0	“const”
$2\mathbb{1}_{[0,5)}(x) + 8\mathbb{1}_{[5,10]}(x)$	“step2”
$5\mathbb{1}_{[10/3,20/3)}(x) + 10\mathbb{1}_{[20/3,10]}(x)$	“step3”
$(x^4\mathbb{1}_{(0,5]}(x) - x^4\mathbb{1}_{[-5,0)}(x))/120$	“power”

Table 1: The true monotone regression function used in the simulations and their abbreviations.

the tables, our estimator is “UL BDD” (where “UL” stands for “unlinked”), Carpentier and Schlüter (2016)’s is “UL CS”, and isotonic regression is “L mono” (for linked regression, that is, based on the classical case where all covariates and responses are perfectly linked). Our simulations were performed taking both Laplace and Gaussian errors with standard deviation 1 (and both unlinked methods are well-specified). In Figure 1 we present the output from a single Monte Carlo run, so that the true functions along with sample data and estimates can be visualized. The 10 plots in the figure are all based on $n = 100$ samples for each of 10 settings: the 5 monotone regression functions m_0 of Table 1 with Laplace distributed errors (left column) and Gaussian errors (right column).

To implement the deconvolution method of Carpentier and Schlüter (2016), we needed to monotoneize the estimated CDF so that we could compute its generalized inverse. Carpentier and Schlüter (2016) do not mention a specific method for doing this; we chose to replace $\hat{F}(x)$ with $\max(\hat{F}(y) : y \leq x)$. The deconvolution estimator at $X_{(1)}$ or $X_{(n)}$ was occasionally unstable because of the steepness of the deconvolution-estimated quantile function. In that case, we have dropped those values out without a noticeable effect. Furthermore, the bandwidth for the deconvolution estimator of the CDF was chosen using the bootstrap method of Delaigle and Gijbels (2004).

From the output one can see that it is not always the case that the Gaussian noise is harder for our method than Laplace is. This is an interesting finding because it is known that deconvolution is harder with Gaussian noise than it is with Laplace; see e.g. Fan (1991). This suggests that, although unlinked regression is tightly connected

to deconvolution, considering the problem from this point of view may not be the most efficient way. The output also shows that the CS estimator performs poorly especially when m_0 has discontinuities. In general our estimator is competitive with the CS deconvolution estimator and in some cases is significantly better.

6.2 Computations on CEX data

Figure 2 shows plots based on the United States’ Consumer Expenditure Survey (CEX). The CEX survey has detailed data on both income and the expenditure patterns of so-called “U.S. consumer units” (roughly, households), see Ruggles et al. (2020). The CEX consists of two surveys, the “Interview” and the “Diary”; the data we use here come from the former. Since the CEX survey has data on both income and expenditures, we can use regular (matched / standard) regression techniques. We compare a U.S. consumer unit’s food expenditure to its income by regressing the former on the latter. Moreover, by simply ignoring the X - Y pairing information we can also use our approach detailed above for the unlinked setting, and then compare the results obtained with both approaches. Note that we prefer here to use data that are matched, for the sake of being able to “validate” our results, but there are many settings where matching naturally lacks; for instance a firm may be able to gather information on an individual consumer’s expenditures on the firm’s products, but the firm would not be able to know the individual’s income information. They would be able to access that expenditure information (at least nationally) through the CEX, which creates a dataset with unlinked covariates and responses.

We consider the interview data only from the second quarter of 2018, for which there are approximately 6000 respondents. We narrow this down to 2164 respondents who provided the relevant information, had income no larger than \$250,000, and reported a positive response for both income and food expenditure. The survey actually follows each individual for four consecutive quarters, but we only included those who were surveyed in both quarter 2 and quarter 3. The “residuals” were computed as described in the previous section: for each individual i , we computed $\tilde{\epsilon}_i := (Y_{i,1} - Y_{i,2})/2$ where $Y_{i,1}$ is the quarter 2 response and $Y_{i,2}$ is the quarter 3 response. The error distribution is assumed to be Laplace distributed with $\lambda = \sqrt{\hat{\sigma}^2/2}$ where $\hat{\sigma}^2$ is the empirical variance of $\tilde{\epsilon}_1, \dots, \tilde{\epsilon}_{2164}$. We chose to assume that the errors follow the Laplace distribution rather than to use the full method detailed in Subsection 5.2 because the former

	UL BDD	UL CS	L mono
lin, Laplace	0.31	0.27	0.16
const, Laplace	0.18	0.25	0.05
step2, Laplace	0.33	2.84	0.09
step3, Laplace	0.43	2.74	0.12
power, Laplace	0.29	0.47	0.13
lin, Gauss	0.48	0.78	0.16
const, Gauss	0.10	1.23	0.05
step2, Gauss	0.19	2.48	0.09
step3, Gauss	0.32	2.59	0.12
power, Gauss	0.43	0.59	0.14

Table 2: Monte Carlo’d MSE’s, $n = 100$.

is much faster to run. Choosing the Laplace distribution was based on observing that it fits much better the distribution of the residuals than the Gaussian one. In Figure 2, the “UL BDD” line is the unlinked monotone minimum contrast estimator proposed in this paper. This estimate is *fully* data driven, using no oracle (matching) information. The “L mono” line corresponds to monotone regression estimator based on the matched data; i.e., the Grenander-type estimator. Similarly the “L linear” line is a linear regression estimator based on the matched data. The “UL CS” line corresponds to the deconvolution estimator of Carpentier and Schlüter (2016) (using the same choice of λ we used in our method). Finally, the “UL quantile” line is based on matching the empirical quantiles of the Y and X samples: it is simply given by (connecting linearly) the points $(X_{(1)}, Y_{(1)}), \dots, (X_{(n)}, Y_{(n)})$.

Our estimator is somewhat accurate although it does differ noticeably from the oracle isotonic regression. The estimator of Carpentier and Schlüter (2016) does very poorly on this dataset. We suspect that inaccuracy in choice of the error distribution causes difficulty for both of the unlinked estimators, which is of course expected. An important avenue for future research is to learn how to use the residuals $\tilde{\epsilon}_i$ more efficiently. This could be done for example by studying faster algorithms so that it is computationally feasible to use an estimate of the gradient at each iteration.

7. Conclusions and directions for future research

In this paper, we have presented a general method for unlinked regression with a monotonic regression function, and developed basic theory for the resulting estimator. We believe our approach will generalize to other (identifiable) unlinked regression settings. We have introduced a variant of a gradient descent algorithm for computing the estimator, and demonstrated its use on a real data example in a fully data driven way in which we estimated the unknown error distribution. There are many remaining questions about this problem and about our method that future work could answer.

1. For our method to be useful on datasets with n larger than 5,000-10,000, a faster algorithm will be needed.
2. A further difficulty related to computation is the practical setting in which the error distribution is not known a priori. We presented a method for estimating the error distribution based on data. However, unless n is only a few hundred, this method

MSE's	UL BDD	UL CS	L mono
lin, Laplace	0.10	0.10	0.03
const, Laplace	0.06	0.13	0.01
step2, Laplace	0.12	3.58	0.01
step3, Laplace	0.15	3.34	0.02
power, Laplace	0.09	0.36	0.03
lin, Gauss	0.29	0.18	0.03
const, Gauss	0.03	0.70	0.01
step2, Gauss	0.07	1.04	0.01
step3, Gauss	0.13	1.40	0.02
power, Gauss	0.25	0.25	0.03

Table 3: Monte Carlo'd MSE's, $n = 1000$.

is not very computationally feasible (and so we used a modified version of it in our analysis of a real data set). Methods and algorithms that are more practical for the case of unknown error distribution are necessary for this model to achieve widespread use. One avenue for future work would be to consider studying so-called log- or s -concave density estimates of the error distribution (see, e.g., Koenker and Mizera (2010); Doss and Wellner (2016); Samworth (2018)); one reason these estimates may be useful is because they are flexible (nonparametric) but also do not require the user to select any tuning parameters and they can be stored very compactly and so using them for every iteration of a gradient-based method could be feasible. Possibly some kernel density estimators could be implemented such that they could be used for every iteration of a gradient-based algorithm.

3. Our current study is restricted to the case of a univariate predictor. Studying both theory and practice when dimension is larger than 1 will be an important avenue for future work. In the case of linear regression, several works (Abid et al. (2017); Pananjady et al. (2017, 2018); Unnikrishnan et al. (2018)) have already begun this study, although those works focus mostly on the case of Gaussian noise.
4. Finding (minimax) lower bounds for the rate of convergence seems to be hard to obtain in our setting. Such bounds are needed for a more complete theoretical understanding of the problem setting. Rigollet and Weed (2019) provide a minimax lower bound in the case of one fixed noise distribution (the Gaussian).
5. One of the major differences in unlinked regression from linked regression is that in the former the specification of the error distribution is crucially important. As shown in the introduction, if the error distribution is unknown then the model is not even identifiable. It would be helpful to understand the general properties of unlinked regression models and of our method in particular when one has partial but incomplete knowledge of the error distribution (e.g., some moment parameters can be estimated well but the full distribution is not known precisely).
6. Carpentier and Schlüter (2016) allowed for so-called “contextual variables”; for instance, if the unit i of observation is an individual, both Y_i and X_i may be paired with a contextual variable $Z_{i,Y}$ and $Z_{i,X}$ such as the individual’s age. One may “match” Y ’s and X ’s with equal (or similar) ages, and then one may consider unmatched regression on these partially matched datasets. This is what Carpentier and Schlüter (2016) proposed in the case of discrete, perfectly (noiselessly) observed contextual variables. More broadly, one may use so-called linkage methods (Herzog et al., 2007) to partially link Y and X (effectively reducing the noise level) when the contextual variables are not as idealized, and then perform linked regression. This methodology could be broadly useful in the linkage literature and warrants further study.

Acknowledgements

The second author is supported in part by NSF Grant DMS-1712664. The third author is supported in part by MME-DII (ANR11-LBX-0023-01) and by the FP2M federation (CNRS FR 2036).

References

- Abubakar Abid, Ada Poon, and James Zou. Linear regression with shuffled labels. *arXiv*, May 2017.
- R. E. Barlow, D. J. Bartholomew, J. M. Bremner, and H. D. Brunk. *Statistical Inference under Order Restrictions. The Theory and Application of Isotonic Regression*. John Wiley, London-New York-Sydney, 1972.
- H. D. Brunk. Estimation of isotonic regression. In *Nonparametric Techniques in Statistical Inference (Proc. Sympos., Indiana Univ., Bloomington, Ind., 1969)*, pages 177–197. Cambridge Univ. Press, London, 1970.
- Chris Carolan and Richard Dykstra. Asymptotic behavior of the Grenander estimator at density flat regions. *The Canadian Journal of Statistics*, 27(3):557–566, 1999. ISSN 0319-5724.
- Alexandra Carpentier and Teresa Schlüter. Learning relationships between data obtained independently. In *Proceedings of the 19th International Conference on Artificial Intelligence and Statistics*, pages 658–666, 2016.
- Raymond J. Carroll, David Ruppert, Leonard A. Stefanski, and Ciprian M. Crainiceanu. *Measurement error in nonlinear models*, volume 105 of *Monographs on Statistics and Applied Probability*. Chapman & Hall/CRC, Boca Raton, FL, second edition, 2006. ISBN 978-1-58488-633-4; 1-58488-633-1. doi: 10.1201/9781420010138. URL <https://doi.org/10.1201/9781420010138>. A modern perspective.
- Eric Cator. Adaptivity and optimality of the monotone least-squares estimator. *Bernoulli*, 17(2):714–735, 2011. ISSN 1350-7265. doi: 10.3150/10-BEJ289. URL <https://doi.org/10.3150/10-BEJ289>.
- Sabyasachi Chatterjee, Adityanand Guntuboyina, and Bodhisattva Sen. On risk bounds in isotonic and other shape restricted regression problems. *The Annals of Statistics*, 43(4):1774–1800, 2015.
- Itai Dattner, Alexander Goldenshluger, and Anatoli Juditsky. On deconvolution of distribution functions. *The Annals of Statistics*, 39(5):2477–2501, 2011.
- Itai Dattner, Markus Reiß, Mathias Trabs, et al. Adaptive quantile estimation in deconvolution with unknown error distribution. *Bernoulli*, 22(1):143–192, 2016.
- Morris H DeGroot and Prem K Goel. Estimation of the correlation coefficient from a broken random sample. *The Annals of Statistics*, 8(2):264–278, March 1980.
- A Delaigle and I Gijbels. Bootstrap bandwidth selection in kernel density estimation from a contaminated sample. *Ann. Inst. Statist. Math.*, 56(1):19–47, 2004.
- Charles R Doss and Jon A Wellner. Global rates of convergence of the mles of log-concave and s -concave densities. *The Annals of Statistics*, 44(3):954–981, 2016.
- Cécile Durot. Sharp asymptotics for isotonic regression. *Probab. Theory Related Fields*, 122(2):222–240, 2002. ISSN 0178-8051. doi: 10.1007/s004400100171. URL <https://doi.org/10.1007/s004400100171>.

- Cécile Durot. On the \mathbb{L}_p -error of monotonicity constrained estimators. *Ann. Statist.*, 35(3):1080–1104, 2007. ISSN 0090-5364. doi: 10.1214/009053606000001497. URL <https://doi.org/10.1214/009053606000001497>.
- Cécile Durot and Hendrik P. Lopuhaä. A Kiefer-Wolfowitz type of result in a general setting, with an application to smooth monotone estimation. *Electron. J. Stat.*, 8(2):2479–2513, 2014. ISSN 1935-7524. doi: 10.1214/14-EJS958. URL <https://doi.org/10.1214/14-EJS958>.
- Cécile Durot, Vladimir N. Kulikov, and Hendrik P. Lopuhaä. The limit distribution of the L_∞ -error of Grenander-type estimators. *Ann. Statist.*, 40(3):1578–1608, 2012. ISSN 0090-5364. doi: 10.1214/12-AOS1015. URL <https://doi.org/10.1214/12-AOS1015>.
- David Edelman. Estimation of the mixing distribution for a normal mean with applications to the compound decision problem. *Ann. Statist.*, 16(4):1609–1622, 1988. ISSN 0090-5364. doi: 10.1214/aos/1176351056. URL <https://doi.org/10.1214/aos/1176351056>.
- Bob E Ellison. Two theorems for inferences about the normal distribution with applications in acceptance sampling. *Journal of the American Statistical Association*, 59(305):89–95, 1964. doi: 10.2307/2282860.
- Jianqing Fan. On the optimal rates of convergence for nonparametric deconvolution problems. *Ann. Statist.*, 19(3):1257–1272, 1991. ISSN 0090-5364. doi: 10.1214/aos/1176348248. URL <https://doi.org/10.1214/aos/1176348248>.
- P. Groeneboom. Estimating a monotone density. In *Proceedings of the Berkeley conference in honor of Jerzy Neyman and Jack Kiefer, Vol. II (Berkeley, Calif., 1983)*, Wadsworth Statist./Probab. Ser., pages 539–555. Wadsworth, Belmont, CA, 1985.
- Piet Groeneboom. The concave majorant of brownian motion. *Ann. Probab.*, 11(4):1016–1027, 1983.
- Piet Groeneboom. Brownian motion with a parabolic drift and Airy functions. *Probab. Theory Related Fields*, 81(1):79–109, 1989. ISSN 0178-8051. doi: 10.1007/BF00343738. URL <https://doi.org/10.1007/BF00343738>.
- Piet Groeneboom and Geurt Jongbloed. *Nonparametric estimation under shape constraints*, volume 38 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, New York, Cambridge, 2014.
- Peter Hall and Soumendra N Lahiri. Estimation of distributions, moments and quantiles in deconvolution problems. *The Annals of Statistics*, 36(5):2110–2134, 2008.
- Thomas N Herzog, Fritz J Scheuren, and William E Winkler. *Data quality and record linkage techniques*. Springer Science & Business Media, 2007.
- Roger Koenker and Ivan Mizera. Quasi-concave density estimation. *Ann. Statist.*, 38(5):2998–3027, 2010.
- Enno Mammen. Estimating a smooth monotone regression function. *Ann. Statist.*, 19(2):724–740, 1991. ISSN 0090-5364. doi: 10.1214/aos/1176348117. URL <https://doi.org/10.1214/aos/1176348117>.

- Ashwin Pananjady, Martin J Wainwright, and Thomas A Courtade. Linear regression with an unknown permutation: Statistical and computational limits. In *Proceedings of the 2016 54th Annual Allerton Conference on Communication, Control, and Computing*, pages 417–424. IEEE, 2016.
- Ashwin Pananjady, Martin J Wainwright, and Thomas A Courtade. Denoising linear models with permuted data. In *2017 IEEE International Symposium on Information Theory (ISIT)*, pages 446–450. IEEE, 2017.
- Ashwin Pananjady, Martin J Wainwright, and Thomas A Courtade. Linear regression with shuffled data: statistical and computational limits of permutation recovery. *IEEE Trans. Inform. Theory*, 64(5):3286–3300, 2018.
- Philippe Rigollet and Jonathan Weed. Uncoupled isotonic regression via minimum wasserstein deconvolution. *Information and Inference (to appear)*, 2019.
- Tim Robertson, F T Wright, and R L Dykstra. *Order restricted statistical inference*. Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons, Ltd., Chichester, 1988.
- Steven Ruggles, Sarah Flood, Ronald Goeken, Josiah Grover, Erin Meyer, Jose Pacas, and Matthew Sobek. *IPUMS, USA: Version 10.0 [dataset]*. Minneapolis, MN: IPUMS, 2020.
- Richard J Samworth. Recent progress in log-concave density estimation. *Statist. Sci.*, 33(4):493–509, 2018.
- Jayakrishnan Unnikrishnan, Saeid Haghighatshoar, and Martin Vetterli. Unlabeled sensing with random linear measurements. *IEEE Trans. Inform. Theory*, 64(5):3237–3253, 2018.
- Aad W. van der Vaart and Jon A. Wellner. *Weak convergence and empirical processes*. Springer Series in Statistics. Springer-Verlag, 1996.
- Cun-Hui Zhang. Risk bounds in isotonic regression. *The Annals of Statistics*, 30(2):528–555, 2002.

Appendix A. Bounding the integrals in (36) and (37).

Using Assumptions A0 and A1 we can write that

$$\begin{aligned} \int_{-\infty}^0 H_0(y)dy &= \int_{-\infty}^0 \int_{\mathbb{R}} \Phi_{\epsilon}(y - m_0(x)) dF_0(x) dy \\ &\leq \int_{-\infty}^0 \Phi_{\epsilon}(y + K_0) dy < \infty, \end{aligned}$$

(note that this also follows from $E(|Y|) < \infty$ and integration by parts.) Similarly it can be shown that $\int_0^{\infty} (1 - H_0(y))dy < \infty$. Hence, $I_1 \leq \int_0^{\infty} (1 - H_0(y))dy + \int_{-\infty}^0 H_0(y)dy < \infty$.

∞ . Also,

$$\begin{aligned}
 \int_{\mathbb{R}} (\Phi_{\epsilon}(y - m_0(x)) - H_0(y))^2 dy &\leq 2 \int_{-\infty}^0 (\Phi_{\epsilon}(y - m_0(x))^2 + H_0(y)^2) dy \\
 &\quad + \int_0^{\infty} (1 - H_0(y))^2 dy \\
 &\leq 2 \int_{-\infty}^0 \Phi_{\epsilon}(y + K_0) dy + 2 \int_{-\infty}^0 H_0(y) dy \\
 &\quad + \int_0^{\infty} (1 - H_0(y)) dy \\
 &< \infty,
 \end{aligned}$$

as shown above; this implies, by Fubini's Theorem, that $I_2 < \infty$. \square

Appendix B. Basic empirical process theory definitions

For a (possibly random) signed measure Q on a (measurable) space \mathcal{X} and a measurable function f on \mathcal{X} , we denote $Qf := \int_{\mathcal{X}} f dQ$. For some class of functions \mathcal{G} , we can define

- its ϵ -covering number $N(\epsilon, \mathcal{G}, \|\cdot\|)$ with respect to some norm $\|\cdot\|$ is defined as the smallest integer $N > 0$ such that there exists g_1, \dots, g_N satisfying that for any $g \in \mathcal{G}$, there exists $i \in \{1, \dots, N\}$ such that

$$\|g - g_i\| < \epsilon,$$

- its ϵ -bracketing number $N_B(\epsilon, \mathcal{G}, \|\cdot\|)$ with respect to some norm $\|\cdot\|$ is defined as the smallest integer $N > 0$ such that there exist pairs $(h_1, k_1), \dots, (h_N, k_N)$ satisfying that for any $g \in \mathcal{G}$, there exist $i \in \{1, \dots, N\}$ such that $h_i \leq g \leq k_i$ and

$$\|k_i - h_i\| < \epsilon.$$

From the definition of the covering and bracketing numbers it can be easily shown (van der Vaart and Wellner, 1996, pp. 83–84) that if $\|\cdot\|$ is an L_p norm, for some $1 \leq p \leq \infty$, then for any $\delta > 0$,

$$N(\delta, \mathcal{G}, \|\cdot\|) \leq N_B(2\delta, \mathcal{G}, \|\cdot\|). \quad (27)$$

Also, if the class \mathcal{G} admits an envelope F , then define for $\eta > 0$ the number

$$J(\eta, \mathcal{G}) = \sup_Q \int_0^{\eta} \sqrt{1 + \log N(\delta \|F\|_{Q,2}, \mathcal{G}, L_2(Q))} d\delta \quad (28)$$

where the supremum is taken over all discrete probability measures Q such that $\|F\|_{Q,2} := \left(\int |F(x)|^2 dQ(x) \right)^{1/2} < \infty$.

Appendix C. Gradient computation

In this section, we prove (26) and we give an explicit formula of $B(m)$ in the case of Laplace and Gaussian distributions.

C.1 Proof of (26)

We have

$$\begin{aligned}
\frac{\partial}{\partial m_i} \mathbb{M}_n(\mathbf{m}) &= \int_{\mathbb{R}} 2 \left(\mathbb{H}_n(y) - n^{-1} \sum_{\alpha=1}^n \Phi_{\epsilon}(y - m_{\alpha}) \right) n^{-1} f_{\epsilon}(y - m_i) dy \\
&= 2n^{-2} \sum_{\alpha=1}^n \int_{\mathbb{R}} \left\{ \mathbb{1}_{[Y_{\alpha}, \infty)}(y) f_{\epsilon}(y - m_i) - \Phi_{\epsilon}(y - m_{\alpha}) f_{\epsilon}(y - m_i) \right\} dy \\
&= 2n^{-2} \sum_{\alpha=1}^n \left\{ \int_{Y_{\alpha} - m_i}^{\infty} f_{\epsilon}(y) dy - \int_{\mathbb{R}} \Phi_{\epsilon}(y - m_{\alpha} + m_i) f_{\epsilon}(y) dy \right\},
\end{aligned} \tag{29}$$

and (26) follows.

C.2 Computations for Laplace distribution

Let $\lambda > 0$ and assume that

$$\Phi_{\epsilon}(z) := \begin{cases} 2^{-1} e^{-|z|/\lambda} & \text{if } z \leq 0 \\ 1 - 2^{-1} e^{-z/\lambda} & \text{if } z > 0 \end{cases}$$

and let $f_{\epsilon}(y) := \Phi'_{\epsilon}(y) = e^{-|z|/\lambda}/2\lambda$ for $z \in \mathbb{R}$. We compute $B(m)$ for $m \in \mathbb{R}$. We have

$$B(m) = \int_{-\infty}^0 (4\lambda)^{-1} e^{y/\lambda} e^{-|y-m|/\lambda} dy + \int_0^{\infty} (1 - 2^{-1} e^{-y/\lambda}) e^{-|y-m|/\lambda} (2\lambda)^{-1} dy$$

which equals

$$\begin{aligned}
&\int_{-\infty}^{m \wedge 0} (4\lambda)^{-1} e^{(y-m/2)2/\lambda} dy + \int_{m \wedge 0}^0 (4\lambda)^{-1} e^{m/\lambda} dy \\
&+ \int_0^{m \vee 0} (2\lambda)^{-1} (e^{(y-m)/\lambda} - 2^{-1} e^{-m/\lambda}) dy \\
&+ \int_{m \vee 0}^{\infty} \frac{1}{2\lambda} e^{-(y-m)/\lambda} - \frac{1}{4\lambda} e^{-(y-m/2)2/\lambda} dy.
\end{aligned}$$

If $m \leq 0$, then

$$B(m) = \frac{1}{8} e^{m/\lambda} + \frac{|m|}{4\lambda} e^{m/\lambda} + 0 + \frac{3}{8} e^{m/\lambda}. \tag{30}$$

If $m \geq 0$, then

$$B(m) = \frac{1}{8} e^{-m/\lambda} + 0 + \left(\frac{1}{2} - \frac{1}{4\lambda} e^{-m/\lambda} (2\lambda + m) \right) + \left(\frac{1}{2} - \frac{1}{8} e^{-m/\lambda} \right). \tag{31}$$

This gives an explicit formula for $B(m)$.

C.3 Computations for Gaussian errors

Now we consider the case where, for some $\sigma > 0$, $\Phi_{\epsilon} = \Phi(\cdot/\sigma)$ is the cumulative distribution function of a $N(0, \sigma^2)$ random variable. It turns out we can write $B(\cdot)$ in terms of Φ : By Theorem 2 of Ellison (1964), $B(m) = \Phi(m/\sigma\sqrt{2})$.

C.4 On implementation of Algorithm 1

A few notes about the implementation of Algorithm 1 are as follows. Rather than iterating for a fixed number K of steps, one can iterate until a stopping rule (e.g., the objective function decrease is smaller than a fixed tolerance level) is satisfied. A heuristic choice for the parameter eps is $\text{eps} = (Y_{(n)} - Y_{(1)})/(n^{1/3}\sigma)$ where σ^2 is the variance of ϵ , and $n^{1/3}$ is motivated by properties of classical isotonic regression. One can also take multiple gradient steps after each “for j ” loop, which may be useful in practice. The usual initialization vector is the sorted response vector, $(Y_{(1)}, \dots, Y_{(n_y)})$.

Appendix D. Proofs

D.1 Proofs for Section 2.3

We first prove the propositions in Section 2.3 and end up with the proof of the main Theorem 2.1.

Proof [Proof of Proposition 2.2.] Let $\widehat{\mathbb{H}}_n$ and \mathbb{H}_n^0 be the distribution functions defined as

$$\widehat{\mathbb{H}}_n(y) = \frac{1}{n} \sum_{i=1}^n \Phi_\epsilon(y - \widehat{m}_n(X_i)), \text{ and } \mathbb{H}_n^0(y) = \frac{1}{n} \sum_{i=1}^n \Phi_\epsilon(y - m_0(X_i)), \quad (32)$$

for $y \in \mathbb{R}$. Recall the Plancherel’s identity for Fourier transforms: for a function $g \in L_1(\mathbb{R}) \cap L_2(\mathbb{R})$, where $L_1(\mathbb{R})$ and $L_2(\mathbb{R})$ denote respectively the set of integrable, and the set of square integrable functions from \mathbb{R} to \mathbb{R} with respect to the Lebesgue measure it holds that

$$\int_{\mathbb{R}} g(x)^2 dx = \frac{1}{2\pi} \int_{\mathbb{R}} |\phi_g(x)|^2 dx$$

where ϕ_g is defined in (17). If F_1 and F_2 are two distribution functions with finite expectations, it follows using integration by parts that

$$\psi_{F_2}(x) - \psi_{F_1}(x) = -ix \int_{\mathbb{R}} (F_2(t) - F_1(t)) e^{itx} dt$$

implying that

$$\phi_{F_2-F_1}(x) = i \frac{\psi_{F_2}(x) - \psi_{F_1}(x)}{x} \quad (33)$$

for $x \neq 0$. Moreover, if F_1 and F_2 have finite expectations then

$$\int_{-\infty}^0 F_j(x) dx < \infty \text{ and } \int_0^\infty (1 - F_j(x)) dx < \infty,$$

for $j \in \{1, 2\}$, implying that $F_1 - F_2 \in L_1(\mathbb{R}) \cap L_2(\mathbb{R})$. Therefore, the Plancherel identity implies that

$$\int_{\mathbb{R}} (F_2(x) - F_1(x))^2 dx = \frac{1}{2\pi} \int_{\mathbb{R}} \frac{1}{x^2} |\psi_{F_2}(x) - \psi_{F_1}(x)|^2 dx.$$

We apply below this identity with F_1 and F_2 replaced respectively by $\widehat{\mathbb{L}}_n$ and \mathbb{L}_n , defined in (20). Note that the two corresponding distributions have finite expectations

since they are supported on a finite set. Hence,

$$\int_{\mathbb{R}} \left(\widehat{\mathbb{L}}_n(w) - \mathbb{L}_n(w) \right)^2 dw = \frac{1}{2\pi} \int_{\mathbb{R}} \frac{1}{t^2} \left| \psi_{\widehat{\mathbb{L}}_n}(t) - \psi_{\mathbb{L}_n}(t) \right|^2 dt.$$

By Assumption A5, we can find $T^* > 0$ such that $|\phi_{f_\epsilon}(t)| \geq |\phi_{f_\epsilon}(T)| > 0$ for all $T > T^*$ and $|t| \leq T$. Using that $|\psi_F| \leq 1$ for any distribution function F , it follows from the previous display that for all $T > T^*$ we have

$$\begin{aligned} \int_{\mathbb{R}} \left(\widehat{\mathbb{L}}_n(w) - \mathbb{L}_n(w) \right)^2 dw &\leq \frac{1}{2\pi |\phi_{f_\epsilon}(T)|^2} \int_{-T}^T \frac{|\phi_{f_\epsilon}(t)|^2}{t^2} \left| \psi_{\widehat{\mathbb{L}}_n}(t) - \psi_{\mathbb{L}_n}(t) \right|^2 dt + \frac{4}{\pi T} \\ &\leq \frac{1}{2\pi |\phi_{f_\epsilon}(T)|^2} \int_{\mathbb{R}} \frac{|\phi_{f_\epsilon}(t)|^2}{t^2} \left| \psi_{\widehat{\mathbb{L}}_n}(t) - \psi_{\mathbb{L}_n}(t) \right|^2 dt + \frac{4}{\pi T}. \end{aligned}$$

Now, using again Plancherel's identity we have

$$\begin{aligned} \int_{\mathbb{R}} \frac{|\phi_{f_\epsilon}(t)|^2}{t^2} \left| \psi_{\widehat{\mathbb{L}}_n}(t) - \psi_{\mathbb{L}_n}(t) \right|^2 dt &= \int_{\mathbb{R}} |\phi_{f_\epsilon}(t)|^2 |\widehat{\mathbb{L}}_n - \mathbb{L}_n(t)|^2 dt \\ &= \int_{\mathbb{R}} |\phi_{f_\epsilon \star (\widehat{\mathbb{L}}_n - \mathbb{L}_n)}(t)|^2 dt \\ &= 2\pi \int_{\mathbb{R}} \left(\widehat{\mathbb{H}}_n(y) - \mathbb{H}_n^0(y) \right)^2 dy \end{aligned}$$

since $\widehat{\mathbb{H}}_n = f_\epsilon \star \widehat{\mathbb{L}}_n$ and $\mathbb{H}_n^0 = f_\epsilon \star \mathbb{L}_n$. Here, $(f \star g)(y) := \int_{\mathbb{R}} f(z)g(y-z)dz$. Hence, it follows from Assumption A4 that for sufficiently large T ,

$$\int_{\mathbb{R}} \left(\widehat{\mathbb{L}}_n(w) - \mathbb{L}_n(w) \right)^2 dw \leq \frac{T^{2\beta}}{d_0^2} \int_{\mathbb{R}} \left(\widehat{\mathbb{H}}_n(y) - \mathbb{H}_n^0(y) \right)^2 dy + \frac{4}{\pi T}. \quad (34)$$

Assuming that we have

$$\int_{\mathbb{R}} E \left(\widehat{\mathbb{H}}_n(y) - \mathbb{H}_n^0(y) \right)^2 dy = O(n^{-1}), \quad (35)$$

it will follow that for all sufficiently large T ,

$$\int_{\mathbb{R}} E \left(\widehat{\mathbb{L}}_n(w) - \mathbb{L}_n(w) \right)^2 dw \leq O(T^{2\beta} n^{-1}) + \frac{4}{\pi T}.$$

For $T = T_n \sim n^{1/(2\beta+1)}$ we get

$$\int_{\mathbb{R}} E \left(\widehat{\mathbb{L}}_n(w) - \mathbb{L}_n(w) \right)^2 dw \leq O \left(\frac{1}{n^{1/(2\beta+1)}} \right),$$

which proves Proposition 2.2.

Now, we will show (35). From the inequality $(a+b)^2 \leq 2(a^2 + b^2)$, which holds for any a and b in \mathbb{R} , and the definition of \widehat{m}_n it follows that

$$\begin{aligned} \int_{\mathbb{R}} \left(\widehat{\mathbb{H}}_n(y) - \mathbb{H}_n^0(y) \right)^2 dy &\leq 2 \int_{\mathbb{R}} \left(\widehat{\mathbb{H}}_n(y) - \mathbb{H}_n(y) \right)^2 dy + 2 \int_{\mathbb{R}} \left(\mathbb{H}_n^0(y) - \mathbb{H}_n(y) \right)^2 dy \\ &\leq 4 \int_{\mathbb{R}} \left(\mathbb{H}_n^0(y) - \mathbb{H}_n(y) \right)^2 dy \\ &\leq 8 \int_{\mathbb{R}} \left(\mathbb{H}_n^0(y) - H_0(y) \right)^2 dy + 8 \int_{\mathbb{R}} \left(\mathbb{H}_n(y) - H_0(y) \right)^2 dy \end{aligned}$$

where

$$E[(\mathbb{H}_n(y) - H_0(y))^2] = n^{-1}H_0(y)(1 - H_0(y)),$$

and

$$\begin{aligned} E[(\mathbb{H}_n^0(y) - H_0(y))^2] &= \frac{1}{n} \text{Var } \Phi_\epsilon(y - m_0(X)) \\ &= \frac{1}{n} \int_{\mathbb{R}} (\Phi_\epsilon(y - m_0(x)) - H_0(y))^2 dF_0(x). \end{aligned}$$

Both the integrals

$$I_1 = \int_{\mathbb{R}} H_0(y)(1 - H_0(y))dy \quad (36)$$

and

$$I_2 = \int_{\mathbb{R}} \int_{\mathbb{R}} (\Phi_\epsilon(y - m_0(x)) - H_0(y))^2 dF_0(x)dy \quad (37)$$

are finite, since it can be shown that $\int_{-\infty}^0 H_0(y)dy < \infty$ and $\int_0^\infty (1 - H_0(y))dy < \infty$; see Appendix A. This yields the result. \blacksquare

Proof [Proof of Proposition 2.3.] In the sequel we denote by \mathbb{P}_n^X and P^X the empirical probability measure associated with X_1, \dots, X_n and the true corresponding probability measure. Then, the two integrals in Proposition 2.3 are the integrated square of the empirical processes

$$\widehat{\mathbb{L}}_n(w) - \widehat{L}_n^0(w) = (\mathbb{P}_n^X - P^X) \mathbb{1}_{\{\widehat{m}_n(\cdot) \leq w\}}$$

and

$$\mathbb{L}_n(w) - L_0(w) = (\mathbb{P}_n^X - P^X) \mathbb{1}_{\{m_0(\cdot) \leq w\}}.$$

In Appendix B, we recall some of the basic tools of empirical processes that we need in this proof. In what follows the notation \lesssim means smaller or equal modulo a universal positive multiplicative constant. Consider the set of functions

$$\mathcal{I} := \left\{ k : k(x) = k_w(x) = \mathbb{1}_{m(x) \leq w}, \text{ with } m \in \mathcal{M} \text{ and } (x, w) \in [0, 1] \times [A, B] \right\}.$$

Using the same notation as in van der Vaart and Wellner (1996) (for completeness, we provide definitions in Appendix B), let us write $\mathbb{G}_n k = \sqrt{n}(\mathbb{P}_n^X - P^X)k$ for $k \in \mathcal{I}$. Since \mathcal{I} is a subset of the class of monotone non-increasing functions $f : \mathbb{R} \mapsto [0, 1]$, it follows from van der Vaart and Wellner (1996, Theorem 2.7.5) that there exists a universal constant $C > 0$, such that for any $\delta > 0$ and any probability measure Q

$$\log N_B(\delta, \mathcal{I}, L_2(Q)) \leq \frac{C}{\delta}$$

(where $N_B(\cdot, \cdot, \cdot)$ is defined in Appendix B). Since \mathcal{I} admits $F(t) = 1$ as an envelope, this and the inequality in (27) imply that

$$\begin{aligned} J(1, \mathcal{I}) &\leq \sup_Q \int_0^1 \sqrt{1 + \log N_B(2\delta, \mathcal{I}, L_2(Q))} d\delta \\ &\leq \int_0^1 \sqrt{1 + \frac{C}{2\delta}} d\delta \leq 1 + \sqrt{2C} < \infty, \end{aligned}$$

where $J(\delta, \mathcal{F})$ is defined in (28). It follows now from van der Vaart and Wellner (1996, Theorem 2.14.1) that

$$(E [\|\mathbb{G}_n\|_{\mathcal{I}}^2])^{1/2} \lesssim J(1, \mathcal{I}). \quad (38)$$

Let us denote

$$M_n = \max \left(\int_A^B \left(\widehat{\mathbb{L}}_n(w) - \widehat{L}_n^0(w) \right)^2 dw, \int_A^B \left(\mathbb{L}_n(w) - L_0(w) \right)^2 dw \right).$$

Then,

$$0 \leq M_n \leq \frac{B-A}{n} \|\mathbb{G}_n\|_{\mathcal{I}}^2.$$

The first two claims in the proposition now follow from (38) combined to the Markov's inequality.

Now, using the inequality $(a+b+c)^2 \leq 3(a^2+b^2+c^2)$ for any real numbers a, b and c , we have

$$\begin{aligned} \int_A^B \left(\widehat{L}_n^0(w) - L_0(w) \right)^2 dw &\leq 3 \int_A^B \left(\widehat{\mathbb{L}}_n(w) - \widehat{L}_n^0(w) \right)^2 dw \\ &\quad + 3 \int_A^B \left(\widehat{\mathbb{L}}_n(w) - \mathbb{L}_n(w) \right)^2 dw \\ &\quad + 3 \int_A^B \left(\mathbb{L}_n(w) - L_0(w) \right)^2 dw. \end{aligned}$$

It thus follows from Proposition 2.2 that (22) holds provided that $B-A = O_p(n^{2\beta/(2\beta+1)})$. ■

Proof [Proof of Proposition 2.4] The first equality in Proposition 2.4 follows from Lemma 4 below combined with the definition of \widehat{L}_n^0 and L_n^0 , while the second equality follows from Proposition 2.3. ■

Lemma 4. *Let $m \in \mathcal{M}$. If F_0 is continuous, then*

$$\int_{\mathbb{R}} \left(\int \mathbb{1}_{\{m(x) \leq w\}} dF_0(x) - F_0 \circ m^{-1}(w) \right)^2 dw = 0. \quad (39)$$

Proof [Proof of Lemma 4] Recall that m^{-1} is defined by (6) where the infimum of an empty set is defined to be 1. If the set in (6) is non-empty, then the infimum is achieved by right-continuity of m . Hence, we have $m \circ m^{-1}(y) \geq y$ for all $y \leq m(1)$. Now, consider $x \in [0, 1]$ and $y \leq m(1)$ such that $m(x) \geq y$. Since the infimum in (6) is achieved this implies that $x \geq m^{-1}(y)$. Conversely, if we have $x \geq m^{-1}(y)$ then monotonicity of m implies that $m(x) \geq m \circ m^{-1}(y)$ where as mentioned above, $m \circ m^{-1}(y) \geq y$. It follows that for all $x \in [0, 1]$ and $y \leq m(1)$ we have the equivalence

$$m(x) \geq y \Leftrightarrow x \geq m^{-1}(y). \quad (40)$$

Now, consider $y > m(1)$. The set in (6) is empty and therefore, $m^{-1}(y) = 1$ by definition. The left-hand inequality in (40) does not hold if $x \in [0, 1]$, and the right-hand inequality does not hold neither if $x < 1$ since $m^{-1}(y) = 1$. This mean that the above equivalence holds for all $x \in [0, 1)$ and $y \in \mathbb{R}$. Let X be a random variable with distribution function F_0 . Since $P(X = 1) = 0$ by assumption, it follows that for all $w \in \mathbb{R}$

$$P(m(X) < w) = P(X < m^{-1}(w))$$

and therefore,

$$P(m(X) \leq w) - P(X \leq m^{-1}(w)) = P(m(X) = w) - P(X = m^{-1}(w))$$

where the second probability on the right hand side equals zero since X has a continuous distribution function. It follows that

$$\begin{aligned} \int_{\mathbb{R}} (P(m(X) \leq w) - F_0 \circ m^{-1}(w))^2 dw &= \int_{\mathbb{R}} P(m(X) = w)^2 dw \\ &= 0. \end{aligned} \quad (41)$$

To see why the preceding equality holds true, note that since the distribution function of X , F_0 , is assumed to be continuous, then it follows that

$$\int_{\mathbb{R}} P(m(X) = w)^2 dw = \int_{\mathcal{W}} P(X \in [a(w), b(w)))^2 dw$$

where \mathcal{W} is the set of point $w \in \mathbb{R}$ such that there exist $x \neq x'$ that satisfy $m(x) = m(x') = w$, and for $w \in \mathcal{W}$, $a(w) < b(w)$ are such that m takes the constant value w on $[a(w), b(w))$, and $a(w) = m^{-1}(w)$. Using the well-known fact that a monotone function admits at most countably many constant parts, the set \mathcal{W} is at most countable and therefore,

$$\int_{\mathbb{R}} P(m(X) = w)^2 dw \leq \int_{\mathcal{W}} dw = \lambda(\mathcal{W}) = 0$$

where λ denotes the Lebesgue measure on \mathbb{R} . Lemma 4 follows from (41) since

$$P(m(X) \leq w) = \int \mathbb{1}_{\{m(x) \leq w\}} dF_0(x).$$

■

Proof [Proof of Proposition 2.5] It follows from Lemma 5 below that

$$\int_a^b |\hat{m}_n(x) - m_0(x)| dF_0(x) \leq \int_{A_n}^{B_n} |F_0 \circ \hat{m}_n^{-1}(x) - F_0 \circ m_0^{-1}(x)| dx.$$

The proposition then follows from applying the Cauchy-Schwarz inequality. ■

Lemma 5. *Let $f : [0, 1] \rightarrow \mathbb{R}$ and $g : [0, 1] \rightarrow \mathbb{R}$ be right-continuous non-decreasing functions. Let f^{-1} and g^{-1} be the corresponding generalized inverses, see (6) where the infimum of an empty set is defined to be one. Let $H : [0, 1] \rightarrow [0, 1]$ be a continuous non-decreasing function. Then, for all $a < b$ in $[0, 1]$ we have*

$$\int_a^b |f(t) - g(t)| dH(t) \leq \int_{I(a)}^{S(b)} |H \circ g^{-1}(x) - H \circ f^{-1}(x)| dx$$

where

$$I(a) = f(a) \wedge g(a) ; S(b) = f(b) \vee g(b).$$

Proof [Proof of Lemma 5.]

For all real numbers u , let $u_+ = \max(u, 0)$. We then have

$$\int_a^b |f(t) - g(t)| dH(t) = I_1 + I_2 \tag{42}$$

where

$$I_1 = \int_a^b (f(t) - g(t))_+ dH(t) \text{ and } I_2 = \int_a^b (g(t) - f(t))_+ dH(t).$$

Let us deal first with I_1 . We have

$$\begin{aligned} I_1 &= \int_a^b \int_0^\infty \mathbb{I}_{\{x \leq f(t) - g(t)\}} dx dH(t) = \int_a^b \int_{g(t)}^\infty \mathbb{I}_{\{x \leq f(t)\}} dx dH(t) \\ &= \int_a^b \int_{g(t)}^{f(b)} \mathbb{I}_{\{x \leq f(t)\}} dx dH(t), \end{aligned}$$

where we use a change of variable for the second equality and the monotonicity of f for the third one. Similar to (40), the equivalence

$$t \geq f^{-1}(x) \iff f(t) \geq x$$

holds for all $t \in [0, 1]$ and $x \in \mathbb{R}$. Combining this with the Fubini theorem, we arrive at

$$\begin{aligned} I_1 &= \int_a^b \int_{g(t)}^{f(b)} \mathbb{I}_{\{t \geq f^{-1}(x)\}} dx dH(t) \\ &= \int_{g(a)}^{f(b)} \int_a^b \mathbb{I}_{\{t \geq f^{-1}(x)\}} \mathbb{I}_{\{t < g^{-1}(x)\}} dH(t) dx. \end{aligned}$$

Hence, it follows from the continuity of H that

$$\begin{aligned} I_1 &= \int_{g(a)}^{f(b)} (H(g^{-1}(x) \wedge b) - H(f^{-1}(x) \vee a))_+ dx \\ &\leq \int_{g(a)}^{f(b)} (H(g^{-1}(x)) - H(f^{-1}(x)))_+ dx, \end{aligned}$$

since H is non-decreasing. Since $I(a) \leq g(a)$ and $S(b) \geq f(b)$, this implies that

$$I_1 \leq \int_{I(a)}^{S(b)} (H \circ g^{-1}(x) - H \circ f^{-1}(x))_+ dx.$$

Interchanging the roles of f and g , we obtain

$$I_2 \leq \int_{I(a)}^{S(b)} (H \circ f^{-1}(x) - H \circ g^{-1}(x))_+ dx$$

and therefore,

$$I_1 + I_2 \leq \int_{I(a)}^{S(b)} |H \circ f^{-1}(x) - H \circ g^{-1}(x)| dx.$$

Lemma 5 then follows from (42). ■

Proof [Proof of Theorem 2.1] If we have (19), then A_n and B_n from Proposition 2.5 are both of the order $O_p(1)$. In that case, the second assertion in Theorem 2.1 is an immediate consequence of Proposition 2.5 combined with Proposition 2.4.

It remains to show that (19) holds. It follows from the definition of $\widehat{\mathbb{L}}_n$ and \mathbb{L}_n together with the Hölder inequality and Proposition 2.2 that

$$\begin{aligned} \int_{\|m_0\|_\infty}^{2\|m_0\|_\infty} \left(1 - \frac{1}{n} \sum_{i=1}^n \mathbb{1}_{\{\widehat{m}_n(X_i) \leq w\}}\right) dw &= \int_{\|m_0\|_\infty}^{2\|m_0\|_\infty} |\widehat{\mathbb{L}}_n(w) - \mathbb{L}_n(w)| dw \\ &\leq \left(\|m_0\|_\infty \int_{\mathbb{R}} (\widehat{\mathbb{L}}_n(w) - \mathbb{L}_n(w))^2 dw\right)^{1/2} \\ &= o_p(1). \end{aligned}$$

On the other hand,

$$\int_{\|m_0\|_\infty}^{2\|m_0\|_\infty} \left(1 - \frac{1}{n} \sum_{i=1}^n \mathbb{1}_{\{\widehat{m}_n(X_i) \leq w\}}\right) dw \geq \|m_0\|_\infty n^{-1} \sum_{i=1}^n \mathbb{1}_{\{\widehat{m}_n(X_i) > 2\|m_0\|_\infty\}}$$

and therefore,

$$\sum_{i=1}^n \mathbb{1}_{\{\widehat{m}_n(X_i) > 2\|m_0\|_\infty\}} = o_p(n).$$

By monotonicity of \widehat{m}_n this implies that

$$\mathbb{1}_{\{\widehat{m}_n(b) > 2\|m_0\|_\infty\}} \sum_{i=1}^n \mathbb{1}_{\{X_i > b\}} \leq \sum_{i=1}^n \mathbb{1}_{\{\widehat{m}_n(X_i) > 2\|m_0\|_\infty\}} = o_p(n).$$

By the law of large numbers, $n^{-1} \sum_{i=1}^n \mathbb{1}_{\{X_i > b\}}$ converges in probability to $1 - F_0(b) > 0$ and therefore, it follows from the previous display that

$$\mathbb{1}_{\{\widehat{m}_n(b) > 2\|m_0\|_\infty\}} = o_p(1).$$

This implies that

$$\lim_{n \rightarrow \infty} P(\widehat{m}_n(b) > 2\|m_0\|_\infty) = 0.$$

One can prove similarly that

$$\lim_{n \rightarrow \infty} P(\widehat{m}_n(a) < -2\|m_0\|_\infty) = 0.$$

This implies (19) by monotonicity of \widehat{m}_n , which completes the proof of Theorem 2.1. ■

Proof [Proof of Remark 1] This follows similar logic as leads to Theorem 2.1. Under Assumptions A0–A3, Assumption A4') and Assumption A5, we can show that the rate of convergence will be driven by

$$O\left(\exp\left(\frac{2T^\beta}{\gamma}\right)T^{-2\alpha}n^{-1}\right) + \frac{4}{\pi T}$$

for $T = T_n \rightarrow \infty$ as $n \rightarrow \infty$ which should be determined so that the above expression is smallest. This means that the first term should converge to 0 or equivalently that there exists a sequence $(K_n)_n$ such that $\log(T_n) = K_n \rightarrow \infty$ and $K_n \leq \log n$ such that

$$\frac{2T_n^\beta}{\gamma} = \log n + (2\alpha - 1)K_n$$

or equivalently

$$T_n = c\left(\log n + (2\alpha - 1)K_n\right)^{1/\beta}, \quad \text{for } c = (\gamma/2)^{1/\beta}.$$

It is not difficult to see that the optimal choice of the sequence $(K_n)_n$ is $K_n = (1 - a)\log n$ for some $a \in (0, 1)$ (the case $a = 0$ is impossible because otherwise we would have $T_n = c(2\alpha)^{1/\beta}(\log(T_n))^{1/\beta}$). This in turn yields

$$T_n = c\left(a\log n + 2\alpha\log(T_n)\right)^{1/\beta},$$

implying that $T_n \sim ca^{1/\beta}(\log n)^{1/\beta}$ and that rate of convergence is $(\log n)^{-1/\beta}$. ■

D.2 Proof of Theorem 2.2

The proof of Theorem 2.2 follows the lines of the proof of Theorem 2.1 so details are omitted and we only point out the main differences between the two proofs.

Similar to (20) and (32), we define

$$\widehat{\mathbb{L}}_{n_x, n_y}(w) := \frac{1}{n_x} \sum_{i=1}^{n_x} \mathbb{1}_{\{\widehat{m}_{n_x, n_y}(X_i) \leq w\}} \quad \text{and} \quad \mathbb{L}_{n_x}(w) := \frac{1}{n_x} \sum_{i=1}^{n_x} \mathbb{1}_{\{m_0(X_i) \leq w\}}$$

for all $w \in \mathbb{R}$, and

$$\widehat{\mathbb{H}}_{n_x, n_y}(y) = \frac{1}{n_x} \sum_{i=1}^{n_x} \Phi_\epsilon(y - \widehat{m}_{n_x, n_y}(X_i)), \quad \text{and} \quad \mathbb{H}_{n_x}^0(y) = \frac{1}{n_x} \sum_{i=1}^{n_x} \Phi_\epsilon(y - m_0(X_i))$$

for all $y \in \mathbb{R}$. With similar arguments as for the proof of (34) we obtain that for all sufficiently large T ,

$$\int_{\mathbb{R}} \left(\widehat{\mathbb{L}}_{n_x, n_y}(w) - \mathbb{L}_{n_x}(w)\right)^2 dw \leq \frac{T^{2\beta}}{d_0^2} \int_{\mathbb{R}} \left(\widehat{\mathbb{H}}_{n_x, n_y}(y) - \mathbb{H}_{n_x}^0(y)\right)^2 dy + \frac{4}{\pi T}. \quad (43)$$

Moreover, it follows from the definition of \widehat{m}_{n_x, n_y} that

$$\begin{aligned} \int_{\mathbb{R}} \left(\widehat{\mathbb{H}}_{n_x, n_y}(y) - \mathbb{H}_{n_x}^0(y) \right)^2 dy &\leq 2 \int_{\mathbb{R}} \left(\widehat{\mathbb{H}}_{n_x, n_y}(y) - \mathbb{H}_{n_y}(y) \right)^2 dy + 2 \int_{\mathbb{R}} \left(\mathbb{H}_{n_x}^0(y) - \mathbb{H}_{n_y}(y) \right)^2 dy \\ &\leq 4 \int_{\mathbb{R}} \left(\mathbb{H}_{n_x}^0(y) - \mathbb{H}_{n_y}(y) \right)^2 dy \\ &\leq 8 \int_{\mathbb{R}} \left(\mathbb{H}_{n_x}^0(y) - H_0(y) \right)^2 dy + 8 \int_{\mathbb{R}} \left(\mathbb{H}_{n_y}(y) - H_0(y) \right)^2 dy \end{aligned}$$

where

$$E[(\mathbb{H}_{n_y}(y) - H_0(y))^2] = n_y^{-1} H_0(y)(1 - H_0(y)),$$

and

$$\begin{aligned} E[(\mathbb{H}_{n_x}^0(y) - H_0(y))^2] &= \frac{1}{n_x} \text{Var} \Phi_\epsilon(y - m_0(X)) \\ &= \frac{1}{n_x} \int_{\mathbb{R}} (\Phi_\epsilon(y - m_0(x)) - H_0(y))^2 dF_0(x). \end{aligned}$$

The integrals I_1 and I_2 defined in (36) and (37) are finite, since it can be shown that $\int_{-\infty}^0 H_0(y) dy < \infty$ and $\int_0^\infty (1 - H_0(y)) dy < \infty$; see Appendix A. Hence, we obtain that

$$\int_{\mathbb{R}} E \left(\widehat{\mathbb{H}}_{n_x, n_y}(y) - \mathbb{H}_{n_x}^0(y) \right)^2 dy = O((n_x \wedge n_y)^{-1}).$$

Combining this with (43) proves that for all sufficiently large T ,

$$\int_{\mathbb{R}} E \left(\widehat{\mathbb{L}}_{n_x, n_y}(w) - \mathbb{L}_{n_x}(w) \right)^2 dw \leq T^{2\beta} O((n_x \wedge n_y)^{-1}) + \frac{4}{\pi T}.$$

For $T \sim (n_x \wedge n_y)^{1/(2\beta+1)}$ we get

$$\int_{\mathbb{R}} E \left(\widehat{\mathbb{L}}_{n_x, n_y}(w) - \mathbb{L}_{n_x}(w) \right)^2 dw \leq O \left(\frac{1}{(n_x \wedge n_y)^{1/(2\beta+1)}} \right), \quad (44)$$

which proves an analogue of Proposition 2.2 in the case of possibly unequal sample sizes.

Next, we consider an analogue of Proposition 2.3. For this task, we denote by $\mathbb{P}_{n_x}^X$ and P^X the empirical probability measure associated with X_1, \dots, X_{n_x} and the true corresponding probability measure. We consider the empirical processes

$$\widehat{\mathbb{L}}_{n_x, n_y}(w) - \widehat{L}_{n_x, n_y}^0(w) = (\mathbb{P}_{n_x}^X - P^X) \mathbb{1}_{\{\widehat{m}_{n_x, n_y}(\cdot) \leq w\}}$$

and

$$\mathbb{L}_{n_x}(w) - L_0(w) = (\mathbb{P}_{n_x}^X - P^X) \mathbb{1}_{\{m_0(\cdot) \leq w\}};$$

where

$$\widehat{L}_{n_x, n_y}^0(w) = \int \mathbb{1}_{\{\widehat{m}_{n_x, n_y}(x) \leq w\}} dF_0(x)$$

and L_0 is defined in (21). Then, with similar arguments as in the proof of Proposition 2.3, we obtain that for all random variables $A < B$ (that may depend on n) it holds that

$$\int_A^B \left(\widehat{\mathbb{L}}_{n_x, n_y}(w) - \widehat{L}_{n_x, n_y}^0(w) \right)^2 dw \leq (B - A) O_p(1/n_x) \leq (B - A) O_p(1/(n_x \wedge n_y)),$$

$$\int_A^B \left(\mathbb{L}_{n_x}(w) - L_0(w) \right)^2 dw \leq (B - A) O_p(1/n_x) \leq (B - A) O_p(1/(n_x \wedge n_y)),$$

where $O_p(1/n_x)$ is uniform in A and B . Moreover, if $B - A = O_p((n_x \wedge n_y)^{2\beta/(2\beta+1)})$, then

$$\int_A^B \left(\widehat{L}_{n_x, n_y}^0(w) - L_0(w) \right)^2 dw = O_p((n_x \wedge n_y)^{-1/(2\beta+1)}).$$

Next, similar to Proposition 2.4 we obtain that for all random variables $A < B$ such that $B - A = O_p(1)$ it holds that

$$\begin{aligned} \int_A^B \left(F_0 \circ \widehat{m}_{n_x, n_y}^{-1}(w) - F_0 \circ m_0^{-1}(w) \right)^2 dw &= \int_A^B \left(\widehat{L}_n^0(w) - L_0(w) \right)^2 dw \\ &= O_p((n_x \wedge n_y)^{-1/(2\beta+1)}). \end{aligned}$$

Proposition 2.5 still holds in the case of possibly different sample sizes with \widehat{m}_n replaced by \widehat{m}_{n_x, n_y} . If (19) also holds, then A_n and B_n from Proposition 2.5 are both of the order $O_p(1)$. In that case, the second assertion in Theorem 2.2 is an immediate consequence of Proposition 2.5 combined with the preceding display. Hence, it remains to prove that (19) holds. It follows from the definition of $\widehat{\mathbb{L}}_{n_x, n_y}$ and \mathbb{L}_{n_x} together with the Cauchy-Schwarz inequality and (44) that

$$\begin{aligned} \int_{\|m_0\|_\infty}^{2\|m_0\|_\infty} \left(1 - \frac{1}{n_x} \sum_{i=1}^{n_x} \mathbb{1}_{\{\widehat{m}_{n_x, n_y}(X_i) \leq w\}} \right) dw &= \int_{\|m_0\|_\infty}^{2\|m_0\|_\infty} \left| \widehat{\mathbb{L}}_{n_x, n_y}(w) - \mathbb{L}_{n_x}(w) \right| dw \\ &\leq \left(\|m_0\|_\infty \int_{\mathbb{R}} \left(\widehat{\mathbb{L}}_{n_x, n_y}(w) - \mathbb{L}_{n_x}(w) \right)^2 dw \right)^{1/2} \\ &= o_p(1). \end{aligned}$$

On the other hand,

$$\int_{\|m_0\|_\infty}^{2\|m_0\|_\infty} \left(1 - \frac{1}{n_x} \sum_{i=1}^{n_x} \mathbb{1}_{\{\widehat{m}_{n_x, n_y}(X_i) \leq w\}} \right) dw \geq \|m_0\|_\infty n_x^{-1} \sum_{i=1}^{n_x} \mathbb{1}_{\{\widehat{m}_{n_x, n_y}(X_i) > 2\|m_0\|_\infty\}}$$

and therefore,

$$\sum_{i=1}^{n_x} \mathbb{1}_{\{\widehat{m}_{n_x, n_y}(X_i) > 2\|m_0\|_\infty\}} = o_p(n_x).$$

By monotonicity of \widehat{m}_{n_x, n_y} this implies that

$$\mathbb{1}_{\{\widehat{m}_{n_x, n_y}(b) > 2\|m_0\|_\infty\}} \sum_{i=1}^{n_x} \mathbb{1}_{\{X_i > b\}} \leq \sum_{i=1}^{n_x} \mathbb{1}_{\{\widehat{m}_{n_x, n_y}(X_i) > 2\|m_0\|_\infty\}} = o_p(n_x).$$

By the law of large numbers, $n_x^{-1} \sum_{i=1}^{n_x} \mathbb{1}_{\{X_i > b\}}$ converges in probability to $1 - F_0(b) > 0$ and therefore, it follows from the previous display that

$$\mathbb{1}_{\{\hat{m}_{n_x, n_y}(b) > 2\|m_0\|_\infty\}} = o_p(1).$$

This implies that

$$\lim_{n \rightarrow \infty} P(\hat{m}_{n_x, n_y}(b) > 2\|m_0\|_\infty) = 0.$$

One can prove similarly that

$$\lim_{n \rightarrow \infty} P(\hat{m}_{n_x, n_y}(a) < -2\|m_0\|_\infty) = 0.$$

This implies (19) by monotonicity of \hat{m}_n , which completes the proof of Theorem 2.2. \square

Appendix E. Proof of Proposition 3.1

To make the notation less cumbersome, we write in the following Z_i for the i -th order statistic $X_{(i)}$. Suppose that \hat{m}_n takes at least two distinct values and let $1 \leq j < j' \leq n$ be such that \hat{m}_n is constant on $[Z_j, Z_{j'})$, where $Z_j < Z_{j'}$ are two successive jump points of \hat{m}_n . Consider the function m_δ which is right-continuous, constant between the order statistics Z_1, \dots, Z_n , and

$$m_\delta(Z_i) = \begin{cases} \hat{m}_n(Z_i) + \delta, & i \in \{j, \dots, j' - 1\} \\ \hat{m}_n(Z_i), & \text{otherwise.} \end{cases}$$

Then, the function m_δ as defined above belongs to \mathcal{M} , provided that $|\delta|$ is small enough. It follows from the definition (8) that $\mathbb{M}_n(m_\delta) \geq \mathbb{M}_n(\hat{m}_n)$. Using Taylor expansion of Φ_ϵ with the integral remainder term we can write that for $i \in \{j, \dots, j' - 1\}$

$$\Phi_\epsilon(y - \hat{m}_n(Z_i) - \delta) = \Phi_\epsilon(y - \hat{m}_n(Z_i)) - \delta f_\epsilon(y - \hat{m}_n(Z_i)) + R_{\delta, i}(y)$$

where the remainder term $R_{\delta, i}$ is given below. Hence,

$$\begin{aligned} 0 &\leq \mathbb{M}_n(m_\delta) - \mathbb{M}_n(\hat{m}_n) \\ &= \int_{\mathbb{R}} \left\{ \mathbb{H}_n(y) - \frac{1}{n} \sum_{i \notin \{j, \dots, j' - 1\}} \Phi_\epsilon(y - \hat{m}_n(Z_i)) - \frac{1}{n} \sum_{i=j}^{j'-1} \Phi_\epsilon(y - \hat{m}_n(Z_i) - \delta) \right\}^2 dy \\ &\quad - \int_{\mathbb{R}} \left\{ \mathbb{H}_n(y) - \frac{1}{n} \sum_{i=1}^n \Phi_\epsilon(y - \hat{m}_n(X_i)) \right\}^2 dy \\ &= \int_{\mathbb{R}} \left\{ \mathbb{H}_n(y) - \frac{1}{n} \sum_{i \notin \{j, \dots, j' - 1\}} \Phi_\epsilon(y - \hat{m}_n(Z_i)) \right. \\ &\quad \left. - \frac{1}{n} \sum_{i=j}^{j'-1} \Phi_\epsilon(y - \hat{m}_n(Z_i)) + \delta \frac{1}{n} \sum_{i=j}^{j'-1} f_\epsilon(y - \hat{m}_n(Z_i)) - \frac{1}{n} \sum_{i=j}^{j'-1} R_{\delta, i}(y) \right\}^2 dy \\ &\quad - \int_{\mathbb{R}} \left\{ \mathbb{H}_n(y) - \frac{1}{n} \sum_{i=1}^n \Phi_\epsilon(y - \hat{m}_n(X_i)) \right\}^2 dy \end{aligned} \tag{45}$$

which equals

$$\begin{aligned} & \int_{\mathbb{R}} \left\{ \mathbb{H}_n(y) - \frac{1}{n} \sum_{i=1}^n \Phi_{\epsilon}(y - \hat{m}_n(X_i)) + \delta \frac{1}{n} \sum_{i=j}^{j'-1} f_{\epsilon}(y - \hat{m}_n(Z_i)) - \frac{1}{n} \sum_{i=j}^{j'-1} R_{\delta,i}(y) \right\}^2 dy \\ & - \int_{\mathbb{R}} \left\{ \mathbb{H}_n(y) - \frac{1}{n} \sum_{i=1}^n \Phi_{\epsilon}(y - \hat{m}_n(X_i)) \right\}^2 dy \end{aligned}$$

which equals

$$\begin{aligned} & \frac{2}{n} \int_{\mathbb{R}} \left(\mathbb{H}_n(y) - \frac{1}{n} \sum_{i=1}^n \Phi_{\epsilon}(y - \hat{m}_n(X_i)) \right) \left(\delta \sum_{i=j}^{j'-1} f_{\epsilon}(y - \hat{m}_n(Z_i)) - \sum_{i=j}^{j'-1} R_{\delta,i}(y) \right) dy \\ & + \frac{1}{n^2} \int_{\mathbb{R}} \left(\delta \sum_{i=j}^{j'-1} f_{\epsilon}(y - \hat{m}_n(Z_i)) - \sum_{i=j}^{j'-1} R_{\delta,i}(y) \right)^2 dy \end{aligned}$$

which equals

$$\begin{aligned} & \frac{2}{n} \int_{\mathbb{R}} \left(\mathbb{H}_n(y) - \frac{1}{n} \sum_{i=1}^n \Phi_{\epsilon}(y - \hat{m}_n(X_i)) \right) \left(\delta(j' - j) f_{\epsilon}(y - \hat{m}_n(Z_j)) - \sum_{i=j}^{j'-1} R_{\delta,i}(y) \right) dy \\ & + \frac{1}{n^2} \int_{\mathbb{R}} \left(\delta(j' - j) f_{\epsilon}(y - \hat{m}_n(Z_j)) - \sum_{i=j}^{j'-1} R_{\delta,i}(y) \right)^2 dy \end{aligned}$$

where for $i = j, \dots, j' - 1$

$$\begin{aligned} R_{\delta,i}(y) &= \int_{y - \hat{m}_n(Z_i)}^{y - \hat{m}_n(Z_i) - \delta} f'_{\epsilon}(t) \cdot (y - \hat{m}_n(Z_i) - \delta - t) dt \\ &= - \int_0^{\delta} f'_{\epsilon}(y - \hat{m}_n(Z_i) - u) \cdot (u - \delta) du, \text{ using the change of variable } u = y - \hat{m}_n(Z_i) - t \\ &= -\delta^2 \int_0^1 f'_{\epsilon}(y - \hat{m}_n(Z_i) - \delta v) (v - 1) dv, \text{ using the change of variable } v = u/\delta. \end{aligned}$$

Thus,

$$\begin{aligned} & \frac{\mathbb{M}_n(m_{\delta}) - \mathbb{M}_n(\hat{m}_n)}{\delta} \\ &= \frac{2}{n} (j' - j) \int_{\mathbb{R}} \left(\mathbb{H}_n(y) - \frac{1}{n} \sum_{i=1}^n \Phi_{\epsilon}(y - \hat{m}_n(X_i)) \right) f_{\epsilon}(y - \hat{m}_n(Z_j)) dy \\ & - \frac{2}{n} \int_{\mathbb{R}} \left(\mathbb{H}_n(y) - \frac{1}{n} \sum_{i=1}^n \Phi_{\epsilon}(y - \hat{m}_n(X_i)) \right) \frac{1}{\delta} \sum_{i=j}^{j'-1} R_{\delta,i}(y) dy \\ & + \frac{1}{n^2} \int_{\mathbb{R}} \left\{ \delta(j' - j)^2 f_{\epsilon}(y - \hat{m}_n(Z_j))^2 - 2(j' - j) f_{\epsilon}(y - \hat{m}_n(Z_j)) \sum_{i=j}^{j'-1} R_{\delta,i}(y) + \frac{\left(\sum_{i=j}^{j'-1} R_{\delta,i}(y) \right)^2}{\delta} \right\} dy \end{aligned}$$

which equals

$$\begin{aligned}
 & \frac{2}{n}(j' - j) \int_{\mathbb{R}} \left(\mathbb{H}_n(y) - \frac{1}{n} \sum_{i=1}^n \Phi_{\epsilon}(y - \hat{m}_n(X_i)) \right) f_{\epsilon}(y - \hat{m}_n(Z_j)) dy \\
 & + \frac{2\delta}{n} \int_{\mathbb{R}} \left(\mathbb{H}_n(y) - \frac{1}{n} \sum_{i=1}^n \Phi_{\epsilon}(y - \hat{m}_n(X_i)) \right) \cdot \left(\sum_{i=j}^{j'-1} \int_0^1 f'_{\epsilon}(y - \hat{m}_n(Z_i) - \delta v) (v - 1) dv \right) dy \\
 & + \frac{\delta(j' - j)^2}{n^2} \int_{\mathbb{R}} f_{\epsilon}(y - \hat{m}_n(Z_j))^2 dy \\
 & + \frac{2\delta(j' - j)}{n^2} \int_{\mathbb{R}} f_{\epsilon}(y - \hat{m}_n(Z_j)) \sum_{i=j}^{j'-1} \int_0^1 f'_{\epsilon}(y - \hat{m}_n(Z_i) - \delta v) (v - 1) dv dy \\
 & + \frac{\delta^3}{n^2} \int_{\mathbb{R}} \left(\int_0^1 \sum_{i=j}^{j'-1} f'_{\epsilon}(y - \hat{m}_n(Z_i) - \delta v) (v - 1) dv \right)^2 dy.
 \end{aligned}$$

We show below that each term on the right hand side that depends on δ takes the form of $\delta^i, i = 1, 2, 3$ times a finite integral, so that it tends to zero as $\delta \rightarrow 0$. From Assumption A6, it follows that

$$\begin{aligned}
 & \left| \left(\mathbb{H}_n(y) - \frac{1}{n} \sum_{i=1}^n \Phi_{\epsilon}(y - \hat{m}_n(X_i)) \right) \cdot \left(\sum_{i=j}^{j'-1} \int_0^1 f'_{\epsilon}(y - \hat{m}_n(Z_i) - \delta v) (v - 1) dv \right) \right| \\
 & \leq D(j' - j) \left| \mathbb{H}_n(y) - \frac{1}{n} \sum_{i=1}^n \Phi_{\epsilon}(y - \hat{m}_n(X_i)) \right|
 \end{aligned}$$

which can be shown to be integrable on \mathbb{R} using the property of Φ_{ϵ} in (9). Also, Assumption A6 implies that there exists $D' > 0$ such that $\sup_{t \in \mathbb{R}} f_{\epsilon}(t) \leq D'$. Then,

$$\int_{\mathbb{R}} f_{\epsilon}(y - \hat{m}_n(Z_j))^2 dy \leq D' \int_{\mathbb{R}} f_{\epsilon}(y - \hat{m}_n(Z_j)) dy = D',$$

and by Fubini's Theorem

$$\begin{aligned}
 & \int_{\mathbb{R}} \sum_{i=j}^{j'-1} \int_0^1 f_{\epsilon}(y - \hat{m}_n(Z_j)) |f'_{\epsilon}(y - \hat{m}_n(Z_i) - \delta v)| (v - 1) dv dy \\
 & = \sum_{i=j}^{j'-1} \int_0^1 \left(\int_{\mathbb{R}} f_{\epsilon}(y - \hat{m}_n(Z_j)) |f'_{\epsilon}(y - \hat{m}_n(Z_i) - \delta v)| dy \right) (v - 1) dv \\
 & \leq D \sum_{i=j}^{j'-1} \int_0^1 (v - 1) dv = \frac{D(j' - j)}{2}
 \end{aligned}$$

using Assumption A6 and the fact that f_{ϵ} is a density. Finally, using again Assumption A6 and Fubini's Theorem we have

$$\begin{aligned}
 & \int_{\mathbb{R}} \left(\int_0^1 \sum_{i=j}^{j'-1} f'_{\epsilon}(y - \hat{m}_n(Z_i) - \delta v) (v - 1) dv \right)^2 dy \\
 & \leq \frac{D(j' - j)}{2} \int_{\mathbb{R}} \int_0^1 \sum_{i=j}^{j'-1} |f'_{\epsilon}(y - \hat{m}_n(Z_i) - \delta v)| (1 - v) dv dy
 \end{aligned}$$

which equals

$$\frac{D(j' - j)}{2} \int_0^1 \left(\sum_{i=j}^{j'-1} \int_{\mathbb{R}} |f'_\epsilon(y - \hat{m}_n(Z_i) - \delta v)| dy \right) (1 - v) dv = \frac{D(j' - j)^2}{4} \int_{\mathbb{R}} |f'_\epsilon(t)| dt < \infty,$$

by Assumption A6. By using (45) and distinguishing between the cases of positive and negative values of δ it follows that

$$\begin{aligned} 0 &= \lim_{\delta \rightarrow 0} \frac{\mathbb{M}_n(m_\delta) - \mathbb{M}_n(\hat{m}_n)}{\delta} \\ &= \frac{2}{n} (j' - j) \int_{\mathbb{R}} \left(\mathbb{H}_n(y) - \frac{1}{n} \sum_{i=1}^n \Phi_\epsilon(y - \hat{m}_n(X_i)) \right) f_\epsilon(y - \hat{m}_n(Z_j)) dy \end{aligned}$$

and therefore,

$$0 = \int_{\mathbb{R}} \left(\mathbb{H}_n(y) - n^{-1} \sum_{i=1}^n \Phi_\epsilon(y - \hat{m}_n(X_i)) \right) f_\epsilon(y - \hat{m}_k) dy$$

where $\hat{m}_k = \hat{m}_n(Z_j) = \dots = \hat{m}_n(Z_{j'-1})$. This is precisely the condition given in (23).

In the case \hat{m}_n takes a unique value, a similar reasoning give the same result, characterizing \hat{m}_k for $k = 1$.

Now, the alternative expression in (24) follows from the fact that for any $a \in \mathbb{R}$

$$\int_{\mathbb{R}} \mathbb{H}_n(y) f_\epsilon(y - a) dy = \frac{1}{n} \sum_{i=1}^n \int_{Y_i}^{\infty} f_\epsilon(y - a) dy = 1 - \frac{1}{n} \sum_{i=1}^n \Phi_\epsilon(Y_i - a)$$

which completes the proof. □




Figure 1: Output from a single Monte Carlo simulation, with $n = 100$, and X_i i.i.d. uniform on $[0, 10]$. The left column has Laplace errors and the right has Gaussian errors, both with standard deviation 1. The dotted gray line is the true m_0 , the red line is our minimum contrast estimator, the blue line is the deconvolution estimator of Carpentier and Schlüter (2016), and the green line is a classical/linked isotonic regression.




Figure 2: CEX Interview Survey: Family income vs. Food expenditure