## The KSVZ Axion and Pseudo-Nambu-Goldstone Boson Models for the XENON1T Excess

Tianjun Li

CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China and

School of Physical Sciences, University of Chinese Academy of Sciences, Beijing, 100049, China

The XENON1T excess can be explained by the Axion Like Particle (ALP) dark matter with mass around 2.5 keV. However, there are three problems needed to be solved: suppressing the coupling  $g_{a\gamma}$  between the ALP and photon, and generating the proper coupling  $g_{ae}$  between the ALP and electron as well as the correct ALP mass. We propose three models to solve these problems. In our models, the  $g_{ae}$  couplings are produced by integrating out the vector-like leptons, and the correct ALP masses arise from high-dimensional operators. In the KSVZ axion model, the coupling  $g_{a\gamma}$ can be suppressed by choosing proper sets of vector-like fermions, but we need some fine-tunings to obtain the ALP mass. Similarly, one can study the DFSZ axion model. In the  $Z_8$  and  $U(1)_X$  models with approximate Pseudo-Nambu-Goldstone Bosons (PNGBs), the coupling  $g_{a\gamma}$  is suppressed due to  $SU(3)_C \times U(1)_{\rm EM}$  anomaly free. In the  $Z_8$  model, the PNGB mass can be generated naturally at the keV scale via the dimension-8 operator. To solve the PNGB quality problem in the  $Z_8$  model, we embed it into the model with  $U(1)_X$  gauge symmetry.

Introduction. – Using the low-energy electronic recoil data with an exposure of 0.65 ton-years, the XENON Collaboration recently reported the results for new physics search [1]. They have observed 285 events over an expected background of  $232\pm15$  events, and found an excess for the electron recoil energies below 7 keV, rising towards lower energies and prominent between 2 and 3 keV. Also, they showed that the solar axion and the solar neutrino with magnetic moment can provide  $3.5\sigma$  and  $3.2\sigma$  significance fits to the excess, respectively. Unfortunately, the corresponding parameter spaces are in tension with stellar cooling bounds [2-4]. With an unconstrained tritium component in the fitting, both the solar axion and the solar neutrino magnetic moment hypotheses no longer have the substantial statistical significance, and their significance levels are respectively reduced to  $2.1\sigma$ and  $0.9\sigma$ . This excess has been studied extensively via solar axion, Axion Like Particles (ALPs), the non-standard neutrino-electron interactions with light mediators, and dark photon, etc [5-50].

It is well-known that the Peccei-Quinn mechanism [51, 52 provides a natural solution to the strong CP problem in the Quantum Chromodynamics (QCD), and predicts a light Pseudo-Nambu-Goldstone Boson (PNGB), dubbed as axion a from QCD anomalous  $U(1)_{PQ}$  global symmetry breaking. The electroceak axion [51-54]was ruled out by the  $K \rightarrow \pi a$  and  $J/\Psi \rightarrow a\gamma$  experiments. And there are two viable invisible axion models: the Dine-Fischler-Srednicki-Zhitnitsky (DFSZ) model [55, 56] and Kim-Shifman-Vainshtein-Zakharov (KSVZ) model [57, 58] with  $U(1)_{PQ}$  symmetry breaking scale from about  $10^{10}$  GeV to  $10^{12}$  GeV. Interestingly, the ALPs, which are the generalizations of axion, may be intrinsic the structure of string theory. The ALP dark matter can explain the XENON1T excess via the electron absorption [5, 18], and let us study its properties before our model building. The Lagrangian between axion and photon/fermions is

$$\mathcal{L}_{a}^{\text{int}} \supset \frac{\alpha_{\text{EM}}}{8\pi} \frac{C_{a\gamma}}{f_{a}} aF\tilde{F} + C_{af} \frac{\partial_{\mu}a}{2f_{a}} \bar{f}\gamma^{\mu}\gamma_{5}f, \qquad (1)$$

where  $\alpha_{\rm EM}$  is structure constant,  $f_a$  is the axion decay constant, and  $C_{a\gamma}$  and  $C_{af}$  are the couplings. The above Lagragian can be rewritten as

$$\mathcal{L}_a^{\text{int}} \supset \frac{1}{4} g_{a\gamma} a F \tilde{F} - i g_{af} a \bar{f} \gamma_5 f , \qquad (2)$$

where

$$g_{a\gamma} = \frac{\alpha_{\rm EM}}{2\pi} \frac{C_{a\gamma}}{f_a}, \qquad g_{af} = C_{af} \frac{m_f}{f_a}$$

The best fit for the XENON1T excess gives [18]

$$m_a = 2.5 \text{ keV}$$
,  $g_{ae} = 2.5 \times 10^{-14}$ . (3)

In particular, the cooling constraint  $g_{ae} < 2.5 \times 10^{-13}$  can be satisfied [3, 4]. The stronger constraint on the decay width for the axion decay into diphoton arises from the observation of the cosmic X-ray backgroud (CXB) gives [59]

$$\frac{C_{a\gamma}}{C_{ae}} \lesssim 2.9 \times 10^{-3} \left(\frac{2.5 \text{ keV}}{m_a}\right)^{3/2} \left(\frac{2.5 \times 10^{-14}}{g_{ae}}\right) . (4)$$

And then we obtain

$$C_{a\gamma} \lesssim 2.9 \times 10^{-3} \left( \frac{f_a}{2 \times 10^{10} \text{ GeV}} \right)$$
 (5)

For the QCD axion models, we have

$$C_{a\gamma} = \frac{E}{N} - 1.92(4).$$
 (6)

where E and N are respectively the electromagnetic and QCD anomaly factors, and 1.92(4) is generated by the

mixing of the axion with the QCD mesons below the confinement scale.

Next, let us discuss the properites of the ALP dark matter particle, which can explain the XENON1T excess. First, we shall show  $f_a \simeq 2 \times 10^{10}$  GeV later, and then the traditional QCD axion will have a mass around  $2.85 \times 10^{-4}$  eV. Thus, the ALP dark matter particle cannot be the traditional QCD axion. Second, from Eq. (5), we obtain  $C_{a\gamma} \lesssim 2.9 \times 10^{-3}$ . In general, there exists about 0.1% fine-tuning for Eq. (6), and the natural solution to it is that both the first term and the second term on the right-handed side vanish: the first condition implies that we do not have  $[U(1)_{\rm EM}]^2 U(1)_{PQ}$ anomaly, while the second condition means no mixing between axion and QCD mesons and thus we do not have  $[SU(3)]^2 U(1)_{PQ}$  anomaly. Therefore, the ALP dark matter particle, which can explain the XENON1T excess, might arise from breaking of a  $SU(3)_C \times U(1)_{\rm EM}$  anomaly free  $U(1)_X$  symmetry (or its discrete subgroup) and is a PNGB.

In short, to explain the XENON1T excess via a PNGB dark matter, we need to address three problems: how to suppress the coupling  $g_{a\gamma}$ , and how to generate the coupling  $g_{ae}$  as well as the correct ALP mass. We shall propose three models to solve these problems: the KSVZ axion model with  $U(1)_{PQ}$  symmetry, the model with  $Z_8$  discrete symmetry, and the model with  $U(1)_X$ gauge symmetry. In our models, assuming that the righthanded electron is charged under  $U(1)_{PQ}$ ,  $Z_8$ , and  $U(1)_X$ symmetries, we can produce the  $g_{ae}$  couplings by integrating out the vector-like leptons. In the KSVZ axion model, the coupling  $g_{a\gamma}$  can be suppressed by choosing proper sets of vector-like fermions. And with some finetuning, we can obtain the correct axion mass from highdimensional operators via quantum gravity effects. Similarly, one can study the DFSZ model, where the coupling  $g_{ae}$  is present and thus we do not need to generate it. In the  $Z_8$  and  $U(1)_X$  models, we do not have  $SU(3)_C \times U(1)_{\rm EM}$  anomaly, so the coupling  $g_{a\gamma}$  is suppressed. In the  $SU(3)_C \times SU(2)_L \times U(1)_Y \times Z_8$  model, we obtain the decay constant around  $2\times 10^{10}~{\rm GeV}$  for the best fit. The correct PNGB a mass around 2.5 keV can be generated from dimension-8 operator naturally. We also show that a has a lifetime long enough to be a dark matter candidate. Moreover, the PNGB dark matter density around the observed value can be generated via the misaligment mechanism, while its thermal density is negligible. Furthermore, to solve the PNGB quality problem via quantum gravity effects in the  $Z_8$  model, we embed it into a  $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_X$ model. The  $U(1)_X$  gauge symmetry is broken down to a  $Z_8$  discrete symmetry around the string scale  $10^{17}$  GeV, and then the  $Z_8$  model can be realized.

The KSVZ Axion Model.– First, we construct the KSVZ axion model which can explain the XENON1T excess. We introduce the vector-like fermions  $(XQ_i^c, XQ_i^c), (XU_i^c, XU_i^c), (XD_i^c, XD_i^c), (XL_i^c, XL_i),$ and  $(XE_i^c, XE_i^c)$ , as well as a SM singlet axion field

$XQ_i$	$({f 3},{f 2},{f 1/6},{f 1})$	$XQ_i^c$	$({f 3},{f 2},-{f 1}/{f 6},{f 1})$
$XU_i$	$({f 3},{f 1},{f 2}/{f 3},{f 1})$	$XU_i^c$	$(\overline{f 3}, {f 1}, -{f 2}/{f 3}, {f 1})$
$XD_i$	$({f 3},{f 1},-{f 1}/{f 3},{f 1})$	$XD_i^c$	$(\overline{f 3}, f 1, f 1/f 3, f 1)$
$XL_i$	$({f 1},{f 2},-{f 1}/{f 2},{f 1})$	$XL_i^c$	$({f 1},{f 2},{f 1}/{f 2},{f 1})$
$XE_i$	$({f 1},{f 1},-{f 1},{f 1})$	$XE_i^c$	$({f 1},{f 1},{f 1},{f 1},{f 1})$
S	$({f 1},{f 1},{f 0},-{f 2})$		

TABLE I. The particles and their quantum numbers under the  $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{PQ}$  gauge and global symmetries.

S. For simplicity, we assme the vector-like fermions have  $U(1)_{PQ}$  charge +1, while S has  $U(1)_{PQ}$  charge -2 These particles and their quantum numbers under the  $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{PQ}$  gauge and global symmetries are summarized in Table I.

The Lagrangian is given by

$$-\mathcal{L} = -m_S^2 |S|^2 + \lambda_S |S|^4 + \left( y_{ij}^{XQ} SXQ_i XQ_j^c + y_{ij}^{XU} SXU_i XU_j^c + y_{ij}^{XD} SXD_i XD_j^c + y_{ij}^{XL} SXL_i XL_j^c + y_{ij}^{XU} SXE_i XE_j^c + \text{H.C.} \right) .$$
(7)

To have small  $C_{a\gamma}$ , we need to find the sets of vectorlike fermions which gives E/N close to 1.92(4). Because the contribution to the electromagnetic anomaly factor from  $(XL_i^c, XL_i)$  is the same as the  $(XE_i^c, XE_i^c)$ , we do not consider  $(XE_i^c, XE_i)$  for simplicity. Of course, any  $(XL_i^c, XL_i^c)$  can be replaced by a  $(XE_i^c, XE_i)$  in the following discussions. For *n* pairs of  $(XQ_i^c, XQ_i^c)$ , *m* pairs of  $(XU_i^c, XU_i^c)$ , *k* pairs of  $(XD_i^c, XD_i^c)$ , and *l* pairs of  $(XL_i^c, XL_i^c)$ , we obtain the condition  $C_{a\gamma} \simeq 0$ 

$$\frac{10n + 8m + 2k + 6l}{6n + 3m + 3k} \simeq 1.92(4) . \tag{8}$$

It is not difficult to find the approximate solution to the above equation, for example  $\frac{10n+8m+2k+6l}{6n+3m+3k} = 2$  for n = m = 0, k = 6, and l = 4. In addition, assuming that the right-handed electron and muon are charged under  $U(1)_{PQ}$  symmetry and introducing the vector-like fermions  $(XL_1, XL_1^c)$  and  $(XL_2, XL_2^c)$ , we can generate the coupling  $g_{aee}$  as we discuss in the following  $Z_8$  and  $U(1)_X$  models. If the QCD axion only obtains mass via instanton effect, its mass will be too small since the decay constant is around  $10^{10}$  GeV as in the following discussions. Therefore, the key question is how to generate the correct axion mass around 2.5 keV. As we know, the global  $U(1)_{PQ}$  symmetry can be broken by the quantum gravity effects. To be concrete, we consider the following effective operator with dimension d = 2m + n that violates the PQ symmetry by n units [60]

$$V \supset \frac{\lambda_n^m |S|^{2m} \left( e^{-i\delta_n^m} S^n + e^{i\delta_n^m} S^{\dagger^n} \right)}{M_{\rm Pl}^{d-4}} \approx m_*^2 f_a^2 \left( \frac{\theta^2}{2} - \frac{\theta}{n} \tan \delta_n^m \right) \,,$$

where we have expanded for  $\theta = \frac{a}{f_a} \ll 1$  by neglecting an irrelevant constant. Here,  $M_{\rm Pl}$  is the reduced Planck scale,  $\lambda_n^m$  is real and  $\delta_n^m$  the phase of the coupling,  $S = \frac{1}{\sqrt{2}}(f_a + s)e^{ia/f_a}$ , and  $m_*^2 = \frac{\lambda_n^m f_a^2}{2} \left(f_a/(\sqrt{2}M_{\rm Pl})\right)^{d-4} \cos \delta_n^m$ . In particular, the linear term or tadpole term will shift the QCD vacuum from  $\langle \theta \rangle = 0$ . Therefore, if we have multiple high-dimensional operators, we can find the fine-tuned solution where the sum of the linear terms is zero or so small that the solution to the strong CP problem can be preserved. And the condition is

$$\sum_{m,n} \frac{\tan \delta_n^m}{n} \simeq 0 . \tag{9}$$

Also, the axion mass is given by

$$m_a = \sqrt{\sum_{m,n} \frac{\lambda_n^m f_a^2}{2} \left( f_a / (\sqrt{2} \mathrm{M}_{\mathrm{Pl}}) \right)^{d-4} \cos \delta_n^m} \,. \tag{10}$$

Therefore, with some fine-tuning, we have shown that the KSVZ axion model can explain the XENON1T excess. Similarly, one can study the DFSZ model, where the coupling  $g_{ae}$  is present, and then we do not need to generate it.

The  $Z_8$  Model.– We shall propose a  $SU(3)_C \times$  $SU(2)_L \times U(1)_Y \times Z_8$  model where  $Z_8$  is a global discrete symmetry. First, let us explain our convention, which is the same as the supersymmetric Standard Model (SM). The SM quark doublets, right-handed up-type quarks, right-handed down-type quarks, lepton doublets, righthanded charged leptons, right-handed neutrinos, and the SM Higgs doublet are denoted as  $Q_i, U_i^c, D_i^c, L_i, E_i^c, N_i^c$ and H, respectively. We shall construct the models where the masses and mixings for the SM quarks and neutrinos are generated in a traditional way. Thus,  $Q_i$ ,  $U_i^c$ ,  $D_i^c$ ,  $L_i, N_i^c$ , and H are not charged under  $Z_8$  discrete symmetry. Also, we assume that the  $Z_8$  quantum numbers for right-handed electron  $E_1^c$ , muon  $E_2^c$ , and tau  $E_3^c$  are +1, -1,and 0, respectively. To break the  $Z_8$  gauge symmetry and have a approximate PNGB, we introduce a SM singlet scalar S with charge -1 under  $Z_8$ . Moreover, to generate the electron and mun Yukawa couplings, we introduce two pairs of vector-like fermions  $(XL_1, XL_1^c)$ and  $(XL_2, XL_2^c)$ . These particles and their quantum numbers under the  $SU(3)_C \times SU(2)_L \times U(1)_Y \times Z_8$  gauge and discrete symmetries are summarized in Table II.

The scalar potential in our model is given by

$$V = -m_S^2 |S|^2 - m_H^2 |H|^2 + \lambda_S |S|^4 + \lambda_{SH} |S|^2 |H|^2 + \lambda_H |H|^4 + \frac{y}{M_{\rm Pl}^4} |S|^8 + \frac{1}{M_{\rm Pl}^4} \left( y' S^8 + \text{H.C.} \right) .(11)$$

For simplicity, we assume y > |y'| so that the potential is stabilized. From the the dimension-8 operator  $y'S^8/M_{\rm Pl}^4$ , we obtain the mass of the PNGB *a* is at the order of  $|\langle S \rangle|^6/M_{\rm Pl}^4$ .

$Q_i$	$({f 3},{f 2},{f 1/6},{f 0})$	$U_i^c$	$({f \overline 3},{f 1},-{f 2}/{f 3},{f 0})$
$D_i^c$	$(\overline{f 3}, {f 1}, {f 1}/{f 3}, {f 0})$	$L_i$	$({f 1},{f 2},-{f 1}/{f 2},{f 0})$
$E_1^c$	$({f 1},{f 1},{f 1},{f 1},{f 1})$	$E_2^c$	$({f 1},{f 1},{f 1},-{f 1})$
$E_3^c$	$({f 1},{f 1},{f 1},{f 0})$	$N_i^c$	$({f 1},{f 1},{f 0},{f 0})$
$XL_1$	$({f 1},{f 2},-{f 1}/{f 2},-{f 1})$	$XL_1^c$	$({f 1},{f 2},{f 1/2},{f 1})$
$XL_2$	$({f 1},{f 2},-{f 1}/{f 2},{f 1})$	$XL_2^c$	$({f 1},{f 2},{f 1/2},-{f 1})$
Н	$({f 1},{f 2},-{f 1}/{f 2},{f 1})$	S	$({f 1},{f 1},{f 0},-{f 1})$

TABLE II. The particles and their quantum numbers under the  $SU(3)_C \times SU(2)_L \times U(1)_Y \times Z_8$  gauge and discrete symmetries.

The Lagrangian for the Yukawa couplings and vector-like fermion masses is

$$-\mathcal{L} = y_{ij}^{U} Q_i U_j^c \overline{H} + y_{ij}^{D} Q_i D_j^c H + y_{i3}^{E} L_i E_3^c H + y_{ij}^{\nu} L_i N_j^c \overline{H} + y_1^{XL} X L_1 E_1^c H + y_2^{XL} X L_2 E_2^c H + y_i^S S L_i X L_1^c + y_i^{\overline{S}} \overline{S} L_i X L_2^c + M_{ij}^N N_i^c N_j^c + M_1^{XL} X L_1 X L_1^c + M_2^{XL} X L_2 X L_2^c + \text{H.C.}, \qquad (12)$$

Using  $y_{ij}^{\nu}L_i N_j^c \overline{H}$  and  $M_{ij}^N N_i^c N_j^c$  terms, we can generate the neutrino masses and mixings via Type I seesaw mechanism. For simplicity, we choose  $y_1^S \neq 0$  and  $y_2^{\overline{S}} = 0$ , while  $y_2^S = y_3^S = y_1^{\overline{S}} = y_3^{\overline{S}} = 0$ . After integrating out the vector-like particles  $(XL_1, XL_1^c)$  and  $(XL_2, XL_2^c)$ , we obtain

$$-\mathcal{L} \supset -\frac{1}{M_1^{XL}} y_1^{XL} y_1^S S L_1 E_1^c H$$
$$-\frac{1}{M_2^{XL}} y_2^{XL} y_2^S \overline{S} L_2 E_2^c H + \text{H.C.} . \qquad (13)$$

Thus, we obtain

$$f_a \equiv \langle S \rangle = \frac{m_e}{g_{ae}} = 2 \times 10^{10} \text{ GeV} \times \left(\frac{2.5 \times 10^{-14}}{g_{ae}}\right) .$$

Therefore, for the best fit, we have  $f_a = 2 \times 10^{10}$  GeV. And then the mass of the PNGB *a* is around keV scale from the dimension-8 operator  $y'S^8/M_{\rm Pl}^4$  in Eq.(11), and we can indeed take it as 2.5 keV.

After integrating out the electron and muon, we obtain the effective Lagrangian between the PNGB a and photon [61]

$$\mathcal{L}_{\text{eff}} = \frac{\alpha_{\text{em}} m_a^2}{48\pi f_a} \left( \frac{1}{m_e^2} - \frac{1}{m_\mu^2} \right) a F_{\mu\nu} \tilde{F}^{\mu\nu} \,. \tag{14}$$

And then we get

$$C_{a\gamma} = \frac{1}{6} \left( \frac{m_a^2}{m_e^2} - \frac{m_a^2}{m_\mu^2} \right) \simeq 4.17 \times 10^{-6} , \qquad (15)$$

which is much smaller than  $2.9 \times 10^{-3}$  and is negligible. The PNGB *a* can decay into two photons via the above effective interaction, and the decay rate is [61]

$$\Gamma_{a \to \gamma \gamma} \simeq \frac{\alpha_{\rm em}^2 q_e^2}{9216\pi^3} \frac{m_a^7}{m_e^4 f_a^2}$$
  

$$\simeq 4.17 \times 10^{-57} \,\text{GeV} \, \left(\frac{m_a}{2 \text{ keV}}\right)^7 \left(\frac{2 \times 10^{10} \text{ GeV}}{f_a}\right)^2.$$

So the constraint on the flux of the X-ray photons produced by the PNGB decay can be satisfied [5, 62].

In our model, the relativistic PNGBs can be produced from the scatterings between electron/muon and the Higgs bosons in the thermal bath. The resulting abundance is [61]

$$\Omega_a^{(\mathrm{th})} h^2 \sim 3.28 \times 10^{-4} \left( \frac{T_R}{3 \times 10^5 \text{ GeV}} \right) \left( \frac{m_a}{2.5 \text{ keV}} \right) \\ \times \left( \frac{2 \times 10^{10} \text{ GeV}}{f_a} \right)^2 , \qquad (16)$$

where  $T_R$  is the reheating temperature. Thus, the thermal relic density of a is negligible.

The PNGB *a* can be produced by the misalignment mechanism as well. When the Hubble parameter is smaller than the mass of *a*, it begins to oscillate around its potential minimum. The temperature  $T_{\rm osc}$  at the onset of the PNGB oscillation is [5, 61]

$$T_{\rm osc} \sim 1.12 \times 10^6 \,\,{\rm GeV} \left(\frac{m_a}{2.5 \,\,{\rm keV}}\right)^{1/2}$$
 (17)

For the temperature higher than  $T_{\rm osc}$ , the PNGB field a has a field value which is not the potential minimum in general. We define the initial oscillation amplitude as  $a_{\rm Initial} \equiv \theta_{\rm mis} f_a$  with  $\theta_{\rm mis}$  the misalignment angle, and obtain the oscillation energy of the PNGB a [5, 61]

$$\Omega_a^{(\text{mis})} h^2 \sim 0.1 \left(\frac{\theta_*}{4}\right)^2 \left(\frac{f_a}{2 \times 10^{10} \text{ GeV}}\right)^2 \\ \times \begin{cases} \left(\frac{T_R}{10^6 \text{ GeV}}\right) & \text{for } T_R \lesssim T_{\text{osc}} \\ \left(\frac{m_a}{2.5 \text{ keV}}\right)^{1/2} & \text{for } T_R \gtrsim T_{\text{osc}} \end{cases} .$$
(18)

Thus, to realize the observed dark matter relic density, we need large initial misalignment angle. We consider the reheating temperature is higher than the oscillation temperature, and the  $Z_8$  symmetry breaking is after inflation. Thus, the decays of the topological defects such as cosmic string and domain wall might contribute to the relic density of the PNGB *a* as well.

The  $U(1)_X$  Model. In the above model,  $Z_8$  is a discrete symmetry, and can be broken via the quantum gravity effects. Thus, the above discussions might not be valid in general if we consider quantum gravity corrections, which is called the PNGB quality problem. Because we do not solve the strong CP problem, in principle we are fine with quantum gravity corrections since we can fine-tune some parameters in our models. To solve the PNGB quality problem, we propose the  $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_X$  model where the  $U(1)_X$  gauge symmetry is broken down to the  $Z_8$  discrete symmetry around the string scale  $10^{17}$  GeV. In addition to the particles in the  $Z_8$  model, we shall introduce two pairs of vector-like particles  $(XE_1, XE_1^c)$  and  $(XE_2, XE_2^c)$ , as well as a SM singlet Higgs scalar field T with  $U(1)_X$  charge 8. The particles and their quantum numbers under the  $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_X$ 

$Q_i$	$({f 3},{f 2},{f 1/6},{f 0})$	$U_i^c$	$({f \overline 3}, {f 1}, -{f 2}/{f 3}, {f 0})$
$D_i^c$	$({f \overline 3},{f 1},{f 1}/{f 3},{f 0})$	$L_i$	$({f 1},{f 2},-{f 1}/{f 2},{f 0})$
$E_1^c$	$({f 1},{f 1},{f 1},{f 1},{f 1})$	$E_2^c$	$({f 1},{f 1},{f 1},-{f 1})$
$E_3^c$	$({f 1},{f 1},{f 1},{f 0})$	$N_i^c$	$({f 1},{f 1},{f 0},{f 0})$
$XL_1$	$({f 1},{f 2},-{f 1}/{f 2},-{f 1})$	$XL_1^c$	$({f 1},{f 2},{f 1/2},{f 1})$
$XL_2$	$({f 1},{f 2},-{f 1}/{f 2},{f 1})$	$XL_2^c$	$({f 1},{f 2},{f 1}/{f 2},-{f 1})$
$XE_1$	$({f 1},{f 1},-{f 1},-{f 1})$	$XE_1^c$	$({f 1},{f 1},{f 1},{f 0})$
$XE_2$	$({f 1},{f 1},-{f 1},{f 1})$	$XE_2^c$	$({f 1},{f 1},{f 1},{f 0})$
Н	$({f 1},{f 2},-{f 1}/{f 2},{f 1})$	S	$({f 1},{f 1},{f 0},-{f 1})$
T	(1, 1, 0, 8)		

TABLE III. The particles and their quantum numbers under the  $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_X$  gauge symmetry.

gauge symmetry are given in Table III. And one can easily show that our model is anomaly free.

The scalar potential is given by

$$V = -m_S^2 |S|^2 - m_T^2 |T|^2 - m_H^2 |H|^2 + \lambda_S |S|^4 + \lambda_T |T|^4 + \lambda_H |H|^4 + \lambda_{ST} |S|^2 |T|^2 + \lambda_{SH} |S|^2 |H|^2 + \lambda_{TH} |T|^2 |H|^2 + \frac{y}{M_{\rm Pl}^6} |T^2| |S|^8 + \frac{1}{M_{\rm Pl}^5} \left( y'TS^8 + \text{H.C.} \right) .$$
(19)

To stabilize the potential after  $U(1)_X$  gauge symmetry breaking, we require

$$\frac{y}{M_{\rm Pl}^6} |\langle T \rangle|^2 > \frac{1}{M_{\rm Pl}^5} |y' \langle T \rangle| . \tag{20}$$

The Lagrangian for the Yukawa couplings and vectorlike fermion masses is

$$\begin{aligned} -\mathcal{L} &= y_{ij}^{U} Q_{i} U_{j}^{c} \overline{H} + y_{ij}^{D} Q_{i} D_{j}^{c} H + y_{i3}^{E} L_{i} E_{3}^{c} H + y_{ij}^{\nu} L_{i} N_{j}^{c} \overline{H} \\ &+ y_{1}^{XL} X L_{1} E_{1}^{c} H + y_{2}^{XL} X L_{2} E_{2}^{c} H + y_{ik}^{XE} L_{i} X E_{k}^{c} H \\ &+ y_{i}^{S} S L_{i} X L_{1}^{c} + y_{i}^{\overline{S}} \overline{S} L_{i} X L_{2}^{c} + y_{k}^{'\overline{S}} \overline{S} X E_{1} X E_{k}^{c} \\ &+ y_{i}^{'S} S X E_{2} X E_{k}^{c} + M_{ij}^{N} N_{i}^{c} N_{j}^{c} + M_{1}^{XL} X L_{1} X L_{1}^{c} \\ &+ M_{2}^{XL} X L_{2} X L_{2}^{c} + \text{H.C.} , \end{aligned}$$

where i, j = 1, 2, 3, and k = 1, 2.

We assume that T acquires a Vacuum Expectation Value (VEV) around string scale  $10^{17}$  GeV, and then the  $U(1)_X$  gauge symmetry is broken down to a discrete  $Z_8$  symmetry. To realize the Lagrangian in Eq. (12), we require the Yukawa couplings  $y_{ik}^{XE}$  to be zero or very small. This can be done in two ways. First, we introduce a  $Z_2$  symmetry under which  $(XE_1, XE_1^c)$ and  $(XE_2, XE_2^c)$  are odd while all the other particles are even. So, the  $y_{ik}^{XE}L_iXE_k^cH$  terms will be forbidden. Because  $(XE_1, XE_1^c)$  and  $(XE_2, XE_2^c)$  cannot decay into the SM particles completely and they are charged particles, we need to require that the reheating temperature is smaller than their masses, for example, around  $10^{10}$  GeV. Second, we consider the five-dimensional space-time on  $S^1/Z_2$ , and assume that the  $SU(3)_C \times SU(2)_L \times U(1)_Y \times$  $U(1)_X$  gauge bosons,  $S, T, (XE_1, XE_1^c)$  and  $(XE_2, XE_2^c)$ are in the bulk, while all the rest particles are on the 3brane at y = 0. In addition, we assume that the wave functions for  $(XE_1, XE_1^c)$  and  $(XE_2, XE_2^c)$  are highly suppressed on the 3-brane at y = 0, and then the Yukawa couplings  $y_{ik}^{XE}$  will be very small. The rest discussions are similar to the above Section, so we shall not repeat it here. In short, we can sovel the PNGB quality problem in the  $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_X$  model.

**Conclusion.**– We proposed three models to explain the XENON1T excess. In our models, the  $g_{ae}$  couplings are generated by integrating out the vector-like leptons, and the correct PNGB mass arises from high-dimensional operators. In the KSVZ axion model, the coupling  $g_{a\gamma}$ can be suppressed by choosing proper sets of vector-like

 $\begin{array}{cccc} [1] & \text{E. Aprile} & \underline{\text{et al.}} & (\text{XENON}), & (2020), \\ & & \text{arXiv:} 2006.09721 & \hline [\text{hep-ex}]. & \end{array}$ 

- [2] A. H. Córsico, L. G. Althaus, M. M. Miller Bertolami, S. Kepler, and E. García-Berro, JCAP 08, 054 (2014), arXiv:1406.6034 [astro-ph.SR].
- [3] M. Giannotti, I. G. Irastorza, J. Redondo, A. Ringwald, and K. Saikawa, JCAP 10, 010 (2017), arXiv:1708.02111 [hep-ph].
- [4] S. A. Díaz, K.-P. Schröder, K. Zuber, D. Jack, and E. E. B. Barrios, (2019), arXiv:1910.10568 [astro-ph.SR].
- [5] F. Takahashi, M. Yamada, and W. Yin, (2020), arXiv:2006.10035 [hep-ph].
- [6] K. Kannike, M. Raidal, H. Veermäe, A. Strumia, and D. Teresi, (2020), arXiv:2006.10735 [hep-ph].
- [7] d. Amaral, Dorian Warren Praia, D. G. Cerdeno, P. Foldenauer, and E. Reid, (2020), arXiv:2006.11225 [hep-ph].
- [8] C. Boehm, D. G. Cerdeno, M. Fairbairn, P. A. Machado, and A. C. Vincent, (2020), arXiv:2006.11250 [hep-ph].
- [9] A. Bally, S. Jana, and A. Trautner, (2020), arXiv:2006.11919 [hep-ph].
- [10] D. Aristizabal Sierra, V. De Romeri, L. Flores, and D. Papoulias, (2020), arXiv:2006.12457 [hep-ph].
- [11] A. N. Khan, (2020), arXiv:2006.12887 [hep-ph].
- [12] H. An, M. Pospelov, J. Pradler, and A. Ritz, (2020), arXiv:2006.13929 [hep-ph].
- [13] M. Lindner, Y. Mambrini, T. B. de Melo, and F. S. Queiroz, (2020), arXiv:2006.14590 [hep-ph].
- [14] L. Di Luzio, M. Fedele, M. Giannotti, F. Mescia, and E. Nardi, (2020), arXiv:2006.12487 [hep-ph].
- [15] C. Gao, J. Liu, L.-T. Wang, X.-P. Wang, W. Xue, and Y.-M. Zhong, (2020), arXiv:2006.14598 [hep-ph].
- [16] G. Alonso-Álvarez, F. Ertas, J. Jaeckel, F. Kahlhoefer, and L. Thormaehlen, (2020), arXiv:2006.11243 [hep-ph].
- [17] K. Nakayama and Y. Tang, (2020), arXiv:2006.13159 [hep-ph].
- [18] I. M. Bloch, A. Caputo, R. Essig, D. Redigolo, M. Sholapurkar, and T. Volansky, (2020), arXiv:2006.14521 [hep-ph].
- [19] B. Fornal, P. Sandick, J. Shu, M. Su, and Y. Zhao, (2020), arXiv:2006.11264 [hep-ph].
- [20] Y. Chen, J. Shu, X. Xue, G. Yuan, and Q. Yuan, (2020), arXiv:2006.12447 [hep-ph].
- [21] Q.-H. Cao, R. Ding, and Q.-F. Xiang, (2020), arXiv:2006.12767 [hep-ph].
- [22] Y. Jho, J.-C. Park, S. C. Park, and P.-Y. Tseng, (2020), arXiv:2006.13910 [hep-ph].

fermions, but we need some fine-tuning to obtain the

Acknowledgments.— This research is supported in part by the Projects 11875062 and 11947302 supported by the National Natural Science Foundation of China, and by the Key Research Program of Frontier Science, CAS.

- [23] K. Harigaya, Y. Nakai, and M. Suzuki, (2020), arXiv:2006.11938 [hep-ph].
- [24] L. Su, W. Wang, L. Wu, J. M. Yang, and B. Zhu, (2020), arXiv:2006.11837 [hep-ph].
- [25] H. M. Lee, (2020), arXiv:2006.13183 [hep-ph].
- [26] J. Bramante and N. Song, (2020), arXiv:2006.14089 [hep-ph].
- [27] M. Baryakhtar, A. Berlin, H. Liu, and N. Weiner, (2020), arXiv:2006.13918 [hep-ph].
- [28] M. Du, J. Liang, Z. Liu, V. Q. Tran, and Y. Xue, (2020), arXiv:2006.11949 [hep-ph].
- [29] G. Choi, M. Suzuki, and T. T. Yanagida, (2020), arXiv:2006.12348 [hep-ph].
- [30] J. Buch, M. A. Buen-Abad, J. Fan, and J. S. C. Leung, (2020), arXiv:2006.12488 [hep-ph].
- [31] U. K. Dey, T. N. Maity, and T. S. Ray, (2020), arXiv:2006.12529 [hep-ph].
- [32] N. F. Bell, J. B. Dent, B. Dutta, S. Ghosh, J. Kumar, and J. L. Newstead, (2020), arXiv:2006.12461 [hep-ph].
- [33] G. Paz, A. A. Petrov, M. Tammaro, and J. Zupan, (2020), arXiv:2006.12462 [hep-ph].
- [34] J. B. Dent, B. Dutta, J. L. Newstead, and A. Thompson, (2020), arXiv:2006.15118 [hep-ph].
- [35] D. McKeen, M. Pospelov, and N. Raj, (2020), arXiv:2006.15140 [hep-ph].
- [36] A. E. Robinson, (2020), arXiv:2006.13278 [hep-ex].
- [37] R. Primulando, J. Julio, and P. Uttayarat, (2020), arXiv:2006.13161 [hep-ph].
- [38] W. DeRocco, P. W. Graham, and S. Rajendran, (2020), arXiv:2006.15112 [hep-ph].
- [39] H. An and D. Yang, (2020), arXiv:2006.15672 [hep-ph].
- [40] P. Ko and Y. Tang, (2020), arXiv:2006.15822 [hep-ph].
- [41] L. Delle Rose, G. Hütsi, C. Marzo, and L. Marzola, (2020), arXiv:2006.16078 [hep-ph].
- [42] W. Chao, Y. Gao, and M. j. Jin, (2020), arXiv:2006.16145 [hep-ph].
- [43] B. Bhattacherjee and R. Sengupta, (2020), arXiv:2006.16172 [hep-ph].
- [44] Y. Gao and T. Li, (2020), arXiv:2006.16192 [hep-ph].
- [45] C. Dessert, J. W. Foster, Y. Kahn, and B. R. Safdi, (2020), arXiv:2006.16220 [hep-ph].
- [46] H. Alhazmi, D. Kim, K. Kong, G. Mohlabeng, J.-C. Park, and S. Shin, (2020), arXiv:2006.16252 [hep-ph].
- [47] G. Cacciapaglia, C. Cai, M. T. Frandsen, M. Rosenlyst, and H. H. Zhang, (2020), arXiv:2006.16267 [hep-ph].
- [48] S. Baek, J. Kim, and P. Ko, (2020),

arXiv:2006.16876 [hep-ph].

- [49] K. Zioutas, G. Cantatore, M. Karuza, A. Kryemadhi, M. Maroudas, and Y. Semertzidis, (2020), arXiv:2006.16907 [hep-ph].
- [50] J. Sun and X.-G. He, (2020), arXiv:2006.16931 [hep-ph].
- [51] R. Peccei and H. R. Quinn, Phys. Rev. Lett. 38, 1440 (1977).
- [52] R. Peccei and H. R. Quinn, Phys. Rev. D 16, 1791 (1977).
- [53] S. Weinberg, Phys. Rev. Lett. 40, 223 (1978).
- [54] F. Wilczek, Phys. Rev. Lett. 40, 279 (1978).
- [55] M. Dine, W. Fischler, and M. Srednicki, Phys. Lett. B 104, 199 (1981).
- [56] A. Zhitnitsky, Sov. J. Nucl. Phys. **31**, 260 (1980).

- [57] J. E. Kim, Phys. Rev. Lett. 43, 103 (1979).
- [58] M. A. Shifman, A. Vainshtein, and V. I. Zakharov, Nucl. Phys. B 166, 493 (1980).
- [59] R. Hill, K. W. Masui, and D. Scott, Appl. Spectrosc. 72, 663 (2018), arXiv:1802.03694 [astro-ph.CO].
- [60] I. G. Irastorza and J. Redondo, Prog. Part. Nucl. Phys. **102**, 89 (2018), arXiv:1801.08127 [hep-ph].
- [61] K. Nakayama, F. Takahashi, and T. T. Yanagida, Phys. Lett. B 734, 178 (2014), arXiv:1403.7390 [hep-ph].
- [62] A. Caputo, M. Regis, and M. Taoso, JCAP 03, 001 (2020), arXiv:1911.09120 [astro-ph.CO].