

Single-molecule-resolution ultrafast near-field optical microscopy via plasmon lifetime extension

Rasim Volga Ovali,¹ Ramazan Sahin,² Alpan Bek,³ and Mehmet Emre Tasgin^{4,*}

¹*Department of Physics, Recep Tayyip Erdogan University, 53100 Rize, Turkey*

²*Faculty of Science, Department of Physics, Akdeniz University, 07058 Antalya, Turkey*

³*Department of Physics, Middle East Technical University, 06800 Ankara, Turkey*

⁴*Institute of Nuclear Sciences, Hacettepe University, 06800 Ankara, Turkey*

(Dated: November 24, 2021)

A recent study shows that: when a long lifetime particle is positioned near a plasmonic metal nanoparticle, lifetime of plasmon oscillations extends, but, *only* near that long-life particle [PRB 101, 035416 (2020)]. Here, we show that this phenomenon can be utilized for ultrahigh (single-molecule) resolution ultrafast apertureless (scattering) SNOM applications. We use the exact solutions of 3D Maxwell equations. We illuminate a metal-coated silicon tip, a quantum emitter (QE) placed on the tip apex, with a femtosecond laser. The induced near-field in the apex decays rapidly except in the vicinity of the sub-nm-sized QE. Thus, the resolution becomes solely limited by the size of the QE. As positioning of a QE on the tip apex is challenging, we propose the use of a newly-discovered phenomenon; stress-induced defect formation in 2D materials. When a monolayer, e.g., transition metal dichalcogenide (TMD) is transferred to the AFM tip, the tip indentation of 2D TMD originates a defect-center located right at the sharpest point of the tip; that is exactly at its apex. Moreover, the resonance of the defect is tunable via a voltage applied to the tip. Our method can equally be used for background-noise-free nonlinear imaging and for facilitating single-molecule-size chemical manipulation.

Metal nanoparticles (MNPs) interact very strongly with incident optical light. Surface plasmons, oscillations of free electrons, localize the incident light into nm-sized hotspots. Hotspot intensity can be 5 to 7 orders of magnitude larger than one of the incident light [1]. Hotspots can enhance both linear [2, 3] and nonlinear optical response [4]. Enhanced hotspot intensity allows detection of even a single molecule [5]. In a nonlinear process both incident and converted fields are enhanced. Thus, a nonlinear process (for instance Raman scattering) can be enhanced quadratically [4].

Localization of light not only strengthens the near-fields, but also enables a hotspot-size resolution at optical imaging [6] and surface manipulation [7]. Hotspot of a metal-coated atomic force microscope (AFM) tip can provide surface images with resolution down to ~ 10 nm [6, 8], limited by the tip apex size, at optical and near-infrared wavelengths. This method, scanning near-field optical microscopy (SNOM), is essential in particular for investigations of light-matter interaction in strongly-correlated condensed phase and 2D materials [9]. This is because, amplitude/phase of the light scattered by the probe (tip) contains information about the local dielectric function of the scanned material [10]. For instance; structures of different phases in VO_2 [11, 12] and organic thin films (for energy harvesting) [13], plasmon mode profiles [14–16] and exciton-polariton transport in 2D materials [17, 18] can be imaged via apertureless SNOM (a-SNOM) (also referred as scattering-type SNOM or s-SNOM) [19]. SNOM has numerous ap-

plications in physics, biology, chemistry and engineering [20, 21].

Although former experiments with SNOM confined to continuous-wave (cw) sources, recent studies are focused on femtosecond (fs) optical/infrared sources [22] which resolve the ultrafast transient spatiotemporal dynamics, for instance, of exciton-polariton dynamics in 2D materials (~ 10 fs, ~ 10 nm) [23–27] and solid state phase transition [28] far from equilibrium. Ultrafast SNOM enabled the observation of intriguing physics, for instance, negative “phase velocity” for phonon-polaritons in 2D hexagonal boron nitride [29]. Such a fs time-resolved imaging of electric fields is important for tracing ultrashort processes taking place in biology and chemistry [30]. Limitations on pulse repetition rate and tip oscillation frequencies, faced in initial experiments, are circumvented by correlating the tip oscillations with the tip-scattered field [31].

Spatial resolution of SNOM, however, is still limited with the tip apex size (~ 10 nm) [23–27]. Furthermore, manufacturing of metal-coated AFM tip of apex size ~ 10 nm is quite challenging. Fortunately, here we show that findings of a recent study [32] can overcome the challenges faced in the spatial resolution, intriguingly, via utilizing a phenomenon in its temporal dynamics. The method, described below, relies on the enhanced lifetime of charge oscillations taking place “only” near a quantum object (QO) [33].

Dark-hot resonances [34], also referred as Fano resonances [35, 36], appear when a bright-mode is coupled to a longer-life dark plasmon mode [37]. The local field can be further enhanced via extended lifetime of bright plasmons [32], on top of the enhancement due to localization. A nonlinear process, e.g., Raman scattering, can be further enhanced (again quadratically) by align-

* metasgin@hacettepe.edu.tr

ing both incident and converted fields with two dark-hot (Fano) resonances [38, 39]. A similar lifetime enhancement enable the operation of spasers —metal nanoparticles (MNPs) coated with molecules [40]. A Fano resonance, demonstrating a dip in its steady-state excitation spectrum [36, 41], ironically, enhances the plasmon energy accumulation in its temporal dynamics [32, 34].

Ref. [32] demonstrates that surface charge oscillations of a MNP near a longlife nanomaterial last much longer than the lifetime of the bright plasmon. As the lifetime extends only near the long lifetime nanomaterial, it shows itself as a weak narrowing in the absorption spectrum [32]. This is unlike a spaser whose surface is covered completely with a vast number of molecules [40].

In this paper, we utilize this phenomenon for achieving a quantum object (QO)-sized resolution ultrafast SNOM. We illuminate a gold-coated tip with a 5 fs-long laser pulse, Fig. 1. First, (i) we show that electric field of a gold-coated AFM tip lasts ~ 17 times longer compared to a plasmon lifetime, but, *only near a* (auxiliary) *quantum object* attached to the tip apex, see Figs. 1 and 2. We perform FDTD simulations of 3D Maxwell equations and use a Lorentzian dielectric for the QO [42–44]. Field below the 1 nm-sized auxiliary QO, which scans two 2 nm-sized nanoparticles (for instance can be proteins), also lasts that much longer, see Fig. 3. Second, (ii) we record “total intensities” (or fluorescence) scattered by the two 2 nm-sized nanoparticles. We show that lifetime-enhanced near-field below the auxiliary quantum object (Figs. 3 and 4) can resolve the 2 nm-sized particle successfully (see Fig. 5). In our simulations we use a 30 nm-thick tip apex.

Positioning a single quantum emitter at the gold-coated tip apex, although possible within current nanotechnological techniques [45], can be challenging. Thus, here we propose to employ a “strain-induced” quantum defect center of a bended 2D materials, e.g., transition metal dichalcogenide (TMD) [46, 47]. When such a 2D material is transferred to an AFM tip, the defect sits at the sharpest (bended) edge of the tip [47], that is at its apex, see Fig. 1. In our approach, the QO and hence the plasmon lifetime extension is generated exactly where it is needed, at the tip apex, which is the part of the tip that actually is responsible for image formation.

Strain-induced defects in TMDs create spatially and spectrally isolated (quantum emitter) centers for localized (0-dimensional) excitons [46–48]. Excitonic nature [49–53] of the centers enables strong near-field coupling [45, 54–56] and highly-enhanced narrow bandwidth emission [46, 48, 57–62] with single-photon characteristics [46, 47]; which makes them perfect candidates for imaging applications [46, 47]. Such QEs are already shown to demonstrate the spontaneous emission enhancement near metallic surfaces [54], reverse of the lifetime enhancement [32], which were shown for molecules or quantum dots [63] previously. Moreover, defect resonance can be controlled with an external voltage [64, 65] which can change the operation wavelength of the locally




FIG. 1. A 10nm-thick silicon AFM tip is coated with 10nm gold. Gold surface is also coated with a 1.5nm insulator (SiO₂) layer to avoid charge transfer between the quantum object and gold surface. A 2D material is grown on the tip, with maximum bending at the tip apex, creates a stress-induced excitonic defect-center which scans two 2-nm-sized nanoparticles. Tip is excited by a z-polarized 5 fs ultrashort pulse propagating in the x-direction. While the near-field excitations decay in 13.7 fs on the gold surface, near-field on the QO lasts much longer, i.e. 240 fs, see Fig. 3. This allows the utilization of the setup as a single-QO-sized resolution SNOM and chemical manipulation.

lifetime-enhanced device slightly. That is, such a setup is a perfect business for high-resolution ultrafast SNOM imaging we introduce here. The new method has the potential to trace electric field evolution of, for instance, protein bioactivities [30] at \sim fs and 1 nm spatiotemporal resolution at different frequencies. For completeness, we also perform similar simulations for a $f = 0.1$ QO [43], a typical value for molecules/proteins [43, 44], placed on tip apex.

The setup we study here can also be utilized for nm-size chemical manipulation [7, 66] and in Fano-enhanced nonlinear processes [39, 67, 68].

Setup.— We calculate the exact solutions of 3D Maxwell equations for a sample setup depicted in Fig. 1. We consider a 10 nm-thick silicon AFM tip coated with 10 nm gold for plasmonic operations [23–27]. Gold surface is also coated with a 1.5 nm insulator (SiO₂) layer to avoid charge-transfer between the gold surface and the auxiliary quantum object. A 2D material, displaying semiconductor feature along its body, is grown on the tip [47] as demonstrated in Fig. 1. The bending of the 2D material creates a stress-induced defect center [46, 47]. The stress-induced defect center appears at the most-curved position (i.e. apex) of the tip [47]. The defect, interacting strongly with the near-field, possesses a strong dipole-moment with a typical oscillator strength of $f \sim 1$. The AFM tip scans the two nanoparticles, of diameter 2 nm, sitting on a silicon substrate. We carry out




FIG. 2. Lifetime of electric field oscillations “between” the tip apex and the auxiliary QO when the system is illuminated with a 5 fs pulse. Oscillator strength of the auxiliary QO is chosen as $f = 1$, corresponding to a stress-induced excitonic defect center in a 2D material, and $f = 0.1$ for a molecule [43, 44].

3D FDTD simulations of the setup using the experimental dielectric functions of the materials, i.e., for silicon, SiO_2 , and gold [69]. We use a Lorentzian dielectric function, $\epsilon(\omega) = 1 - f\omega_0^2/(\omega_0^2 + i2\gamma\omega - \omega^2)$, for the auxiliary quantum object [42–44]. ω_0 and γ are the resonance and decay rate of the auxiliary quantum object.

Lifetime of such defect centers, in the order of $\sim \text{ns}$ [47], are quite long compared to the plasmonic oscillations (~ 10 fs in gold). So, it is possible to extend the lifetime of the near-field (plasmon) oscillations between the defect (QO) and the gold surface as in Ref. [32]. Such a lifetime extension, we expect and observe here, in Fig. 2, is the analog of the phenomenon observed in Ref. [32]. Ref. [32] employs longer lifetime of dark plasmon modes which can be ~ 10 -100 times longer compared to bright plasmon modes. Here, we employ the quantum object as the longlife particle whose lifetime is $\sim \text{ns}$, which is 5-orders of magnitude larger than the one for plasmon oscillations on the gold surface (~ 10 fs).

Lifetime extension.— In Fig. 2, we show that lifetime of plasmon oscillations “between” the auxiliary QO and the gold surface (~ 240 fs) increases ~ 17 times compared to the lifetime of localized surface plasmons (LSPs) in the bare gold surface ($\simeq 13.7$ fs). The plasmonic near-field oscillations out of the QO, in contrast, decay within 13.7 fs.

Lifetime extension phenomenon, employing an auxiliary quantum object, displays a common behavior with the one employing longer-life dark plasmon modes [32]. Enhancement increases for closer positioning (stronger coupling) of the auxiliary QO to the gold surface. But after a critical (strong) coupling, lifetime extension decreases, a feature also observed in fluorescence enhancement of molecules near plasmonic surfaces [63].

Utilization.— Therefore, such a setup (Fig. 1) is possible to be utilized as a single-QO-size resolution SNOM. For this utilization, however, a similar lifetime extension is needed to be observed also *below* the auxiliary QO, the location which scans the two 2 nm-sized nanoparticles sitting on the substrate.

In Fig. 3, we plot the temporal evaluation of the electric field, just (0.25 nm) below the QO. We simulate




FIG. 3. (black) Time evolution of the electric field on the gold surface, at the tip apex, when there is no QO. Oscillations decay quickly after ~ 25 fs. (red) An auxiliary QO is utilized to extend the lifetime of plasmonic oscillations. Electric field “0.25 nm-below” the auxiliary QO lasts much longer.

the auxiliary QO as a 1 nm-sized sphere of Lorentzian width $\gamma = 10^9 \text{ Hz}$, resonance $f_0 = 491 \text{ THz}$ and oscillator strength $f = 1$. We illuminate the setup (Fig. 1) with a z-polarized, 5 fs-width, ultrashort pulse propagating in the x-direction. We set the carrier frequency of the ultrashort pulse to $f_c = 491 \text{ THz}$ ($\lambda_c = 610 \text{ nm}$), the resonance of the auxiliary QO.

We kindly remark that in this *prominent* work we aim to demonstrate how the local lifetime enhancement method works for imaging implementations. Other spectral regions, e.g., 610-680 nm and 740-820 nm [46, 47], can also be used for nm-size-resolution imaging.

Fig. 3 shows that electric field oscillations, also below the QO, last much longer compared to the surface of a gold tip. So, near-field below the QO scan the two nanoparticles much longer times compared to another position on the gold surface (even of the tip apex would be a flat one). While the near-field of gold surface decays, after ~ 13.7 fs, the 1 nm-thick QO continues to provide scattering signal from the two nanoparticles. Therefore, one can achieve ~ 1 nm-size resolution SNOM either (i) by recording the “total scattered intensity”, i.e., $J_{\text{sca}} = \int dt I_{\text{sca}}(t)$, or, perfectly (ii) by simply omitting the signal before, e.g., $t = 25$ fs in Fig. 3. In this work, we study the former approach: we record the total scattered field.

Fig 4 demonstrates a clear picture how the “total near-field intensity”, $J_{\text{near}}(\mathbf{r}) = \int dt I_{\text{near}}(\mathbf{r}, t)$, changes for different locations on the tip apex. As evident from Fig. 4a, total near-field intensity is 1-order of magnitude larger near and below the QO compared to other locations on tip surface. Thus, it provides a ~ 1 nm-size resolution. Fig. 4b plots the total near-field intensity for different positions below the auxiliary QO for two different values of the oscillator strength $f = 1$ and $f = 0.1$ [70].

In Fig. 5, we present the “total scattered intensity”, J_{sca} , when the tip scans the nanoparticles, that is, for different positions of the tip-center. We calculate the J_{sca} in our 3D FDTD simulations as follows. We place




FIG. 4. (a) Total near-field intensity, $J_{\text{near}}(\mathbf{r}) = \int dt I_{\text{near}}(\mathbf{r}, t)$, scanning the two 2-nm sized nanoparticles in Fig. 1, is confined around the QO. The fringe-like behavior appears because the 5 fs pulse contains a broadband frequency. Hence, both dipole and quadrupole modes are excited [71]. (c) Total near-field intensity, just below the QO, for different distances between the QO and the gold surface.

3D intensity monitors encapsulating only on the scanned 2-nm sized nanoparticles. Our logic is simple: the field scattered by the scanned nanoparticles is proportional to the induced near-field intensity on (and in) the nanoparticles [72, 73]. In Figs. 5a and 5b, we scan a single 2 nm-sized nanoparticle; using a gold tip alone and by employing the auxiliary QO, respectively. While change in the resolution between Figs. 5a and 5b is obvious, Fig. 5a presents a “deceptive” high resolution for a 30 nm-thick gold apex. This appears because, we use a perfect (and fine-meshed) hemisphere shape for the AFM tip in the simulations while in actual experiments such a fault-free tip is almost impossible to manufacture. In an actual experiment, the stress-induced defect center, however, appears at the apex where bending is maximum [47].

Scanning two 2 nm-sized nanoparticles, separated 2 nm from each other, demonstrates the contrast between the two resolutions more explicitly. Gold-coated tip, without a QO, fails to resolve the two particles (Fig. 5c) this time. Only a small wiggle between the two nanoparticles, $x = 0$, appears, again, due to perfect hemisphere tip shape. A tip employing the auxiliary QO, resolves the separation between the two nanoparticles clearly, Fig. 5.

Complications due to oscillating tip.— In a SNOM experiment, the AFM tip oscillates with the natural frequency of the cantilever, typically $\Omega_{\text{tip}} \sim 10^5$ Hz or less. That is, the tip does not scan a surface as if it moves along the x-direction with a constant altitude. Hence, a high repetition-rate source is required for non-sophisticated-tracking of the scattered signal with respect to the tip altitude (or tip-sample distance). This constrained the




FIG. 5. AFM tip scans (a,b) a single 2 nm-sized nanoparticle and (c,d) two 2 nm-sized nanoparticles separated 2 nm from each other. Total scattered intensity in the (upper) absence and (lower) presence of the auxiliary QO. The tip apex is 30 nm. The *deceptive* high-resolution of nanoparticles in (a,c) is due to the perfect hemispherical body of the tip. (d) The auxiliary QO is very successful in resolving the two nanoparticles, while the gold tip without the QO fails despite a perfectly-shaped tip.

initial experiments, on ultrafast SNOM, to sources with repetition rates much larger than the Ω_{tip} . A recent experimental study [31], however, demonstrates a method for conducting ultrafast SNOM experiments with low-repetition rate sources. Ref. [31] obtains the dependence of the scattered signal (S) on the instantaneous oscillation phase of the tip (tip-sample distance) Φ , $S(\Phi)$ [74]. The time interval we use to calculate the total scattered intensity is only 500 fs, 7-orders shorter than the oscillation frequency of the tip. Hence, the “total scattering” data we obtain is like a point in the oscillation frame of the tip. Thus, a similar technique [31] can be used also in our setup in order to elect the scattering signal for an appropriate altitude of the tip.

In summary, we demonstrate a novel method for obtaining sub-nm-size ultrafast SNOM imaging, which heals the current limit (~ 10 nm) by 1-order of magnitude. We also propose a novel technique for manufacturing an AFM (SNOM) tip in which a large oscillator strength quantum object (a defect-center) is located necessarily at the lowest-altitude of tip apex [46, 47]. The method works both with low and high repetition rate sources.

Though we concentrate on on SNOM applications, our setup (Fig. 1) can equally be utilized for nonlinear microscopy techniques [39, 67, 68, 75] where small adjustments of the QO resonance is vital for Fano-enhanced phenomena to take place. Higher localization, compared to the ones using Fano resonances with dark-modes [39, 68], can be achieved since a QO is much

smaller than dark-mode profiles. The new method also enables ultrahigh resolution chemical manipulation on surfaces [7, 66].

ACKNOWLEDGMENT

We gratefully thank Ceyhun Bulutay for illuminating discussions. The full-credit for the utilization of a stress-induced defect-center for the new method belongs to AB. MET acknowledges support from TUBA GEBIP 2017 and TUBITAK 1001 No:117F118. RS, AB and MET acknowledge support from TUBITAK 1001 No: 119F101.

-
- [1] Christiane Höppener, Zachary J Lapin, Palash Bharadwaj, and Lukas Novotny, “Self-similar gold-nanoparticle antennas for a cascaded enhancement of the optical field,” *Physical Review Letters* **109**, 017402 (2012).
 - [2] Ali Panahpour, Abolfazl Mahmoodpoor, and Andrei V Lavrinenko, “Refraction enhancement in plasmonics by coherent control of plasmon resonances,” *Physical Review B* **100**, 075427 (2019).
 - [3] Mehmet Günay, You-Lin Chuang, and Mehmet Emre Tasgin, “Tunable Cherenkov radiation-based detectors via plasmon index enhancement,” Accepted in *Nanophotonics*; arXiv preprint arXiv:1911.08159 (2019).
 - [4] Martti Kauranen and Anatoly V Zayats, “Nonlinear plasmonics,” *Nature Photonics* **6**, 737 (2012).
 - [5] Eric Betzig and Robert J Chichester, “Single molecules observed by near-field scanning optical microscopy,” *Science* **262**, 1422–1425 (1993).
 - [6] Paul Bazylewski, Sabastine Ezugwu, and Giovanni Fanchini, “A review of three-dimensional scanning near-field optical microscopy (3d-snom) and its applications in nanoscale light management,” *Applied Sciences* **7**, 973 (2017).
 - [7] Chao Zhan, Xue-Jiao Chen, Jun Yi, Jian-Feng Li, De-Yin Wu, and Zhong-Qun Tian, “From plasmon-enhanced molecular spectroscopy to plasmon-mediated chemical reactions,” *Nature Reviews Chemistry* **2**, 216–230 (2018).
 - [8] Alpan Bek, Ralf Vogelgesang, and Klaus Kern, “Apertureless scanning near field optical microscope with sub-10 nm resolution,” *Review of Scientific Instruments* **77**, 043703 (2006).
 - [9] Mengkun Liu, Aaron J Sternbach, and DN Basov, “Nanoscale electrodynamics of strongly correlated quantum materials,” *Reports on Progress in Physics* **80**, 014501 (2016).
 - [10] Rainer Hillenbrand and Fritz Keilmann, “Complex optical constants on a subwavelength scale,” *Physical Review Letters* **85**, 3029 (2000).
 - [11] Mumtaz M Qazilbash, Markus Brehm, Byung-Gyu Chae, P-C Ho, Gregory O Andreev, Bong-Jun Kim, Sun Jin Yun, AV Balatsky, MB Maple, Fritz Keilmann, *et al.*, “Mott transition in VO_2 revealed by infrared spectroscopy and nano-imaging,” *Science* **318**, 1750–1753 (2007).
 - [12] MK Liu, M Wagner, E Abreu, S Kittiwatanakul, A McLeod, Z Fei, M Goldflam, S Dai, MM Fogler, J Lu, *et al.*, “Anisotropic electronic state via spontaneous phase separation in strained vanadium dioxide films,” *Physical Review Letters* **111**, 096602 (2013).
 - [13] Christian Westermeier, Adrian Cernescu, Sergiu Amarie, Clemens Liewald, Fritz Keilmann, and Bert Nickel, “Sub-micron phase coexistence in small-molecule organic thin films revealed by infrared nano-imaging,” *Nature Communications* **5**, 1–6 (2014).
 - [14] Jianing Chen, Michela Badioli, Pablo Alonso-González, Sukosin Thongrattanasiri, Florian Huth, Johann Osmond, Marko Spasenović, Alba Centeno, Amaia Pesquera, Philippe Godignon, *et al.*, “Optical nano-imaging of gate-tunable graphene plasmons,” *Nature* **487**, 77–81 (2012).
 - [15] Zhe Fei, AS Rodin, Gregory O Andreev, W Bao, AS McLeod, M Wagner, LM Zhang, Z Zhao, M Thiemens, G Dominguez, *et al.*, “Gate-tuning of graphene plasmons revealed by infrared nano-imaging,” *Nature* **487**, 82–85 (2012).
 - [16] Benedikt Hauer, Tobias Saltzmann, Ulrich Simon, and Thomas Taubner, “Solvothermally synthesized sb2te3 platelets show unexpected optical contrasts in mid-infrared near-field scanning microscopy,” *Nano Letters* **15**, 2787–2793 (2015).
 - [17] F Hu, Y Luan, ME Scott, J Yan, DG Mandrus, X Xu, and Z Fei, “Imaging exciton–polariton transport in mose 2 waveguides,” *Nature Photonics* **11**, 356–360 (2017).
 - [18] Z Fei, ME Scott, DJ Gosztola, JJ Foley IV, J Yan, DG Mandrus, H Wen, P Zhou, DW Zhang, Y Sun, *et al.*, “Nano-optical imaging of wse_2 waveguide modes revealing light-exciton interactions,” *Physical Review B* **94**, 081402 (2016).
 - [19] Xinzhong Chen, Debo Hu, Ryan Mescall, Guanjuan You, DN Basov, Qing Dai, and Mengkun Liu, “Modern scattering-type scanning near-field optical microscopy for advanced material research,” *Advanced Materials* **31**, 1804774 (2019).
 - [20] Iban Amenabar, Simon Poly, Wiwat Nuansing, Elmar H Hubrich, Alexander A Govyadinov, Florian Huth, Roman Krutokhvostov, Lianbing Zhang, Mato Knez, Joachim Heberle, *et al.*, “Structural analysis and mapping of individual protein complexes by infrared nanospectroscopy,” *Nature Communications* **4**, 1–9 (2013).
 - [21] Robert C Dunn, “Near-field scanning optical microscopy,” *Chemical Reviews* **99**, 2891–2928 (1999).
 - [22] Ziheng Yao, Suheng Xu, Debo Hu, Xinzhong Chen, Qing Dai, and Mengkun Liu, “Nanoimaging and nanospectroscopy of polaritons with time resolved s-snom,” *Advanced Optical Materials* **8**, 1901042 (2020).
 - [23] M Mrejen, L Yadgarov, A Levanyon, and H Suchowski, “Transient exciton-polariton dynamics in wse_2 by ultrafast near-field imaging,” *Science Advances* **5**, eaat9618 (2019).
 - [24] Philipp Steinleitner, Philipp Merkl, Philipp Nagler, Joshua Mornhinweg, Christian Schuller, Tobias Korn, Alexey Chernikov, and Rupert Huber, “Direct observation of ultrafast exciton formation in a monolayer of

- wse2,” *Nano Letters* **17**, 1455–1460 (2017).
- [25] GX Ni, L Wang, MD Goldflam, M Wagner, Z Fei, AS McLeod, MK Liu, F Keilmann, B Özyilmaz, AH Castro Neto, *et al.*, “Ultrafast optical switching of infrared plasmon polaritons in high-mobility graphene,” *Nature Photonics* **10**, 244–247 (2016).
 - [26] Martin Wagner, Zhe Fei, Alexander S McLeod, Aleksandr S Rodin, Wenzhong Bao, Eric G Iwinski, Zeng Zhao, Michael Goldflam, Mengkun Liu, Gerardo Dominguez, *et al.*, “Ultrafast and nanoscale plasmonic phenomena in exfoliated graphene revealed by infrared pump-probe nanoscopy,” *Nano Letters* **14**, 894–900 (2014).
 - [27] Vasily Kravtsov, Ronald Ulbricht, Joanna M Atkin, and Markus B Raschke, “Plasmonic nanofocused four-wave mixing for femtosecond near-field imaging,” *Nature Nanotechnology* **11**, 459 (2016).
 - [28] Sven A Donges, Omar Khatib, Brian T OCallahan, Joanna M Atkin, Jae Hyung Park, David Cobden, and Markus B Raschke, “Ultrafast nanoimaging of the photoinduced phase transition dynamics in vo_2 ,” *Nano Letters* **16**, 3029–3035 (2016).
 - [29] Edward Yoxall, Martin Schnell, Alexey Y Nikitin, Oihana Txoperena, Achim Woessner, Mark B Lundberg, Félix Casanova, Luis E Hueso, Frank HL Koppens, and Rainer Hillenbrand, “Direct observation of ultraslow hyperbolic polariton propagation with negative phase velocity,” *Nature Photonics* **9**, 674 (2015).
 - [30] Monique M Martin and James T Hynes, *Femtochemistry and femtobiology: ultrafast events in molecular science* (Elsevier, 2004).
 - [31] Haomin Wang, Le Wang, and Xiaoji G Xu, “Scattering-type scanning near-field optical microscopy with low-repetition-rate pulsed light source through phase-domain sampling,” *Nature Communications* **7**, 1–8 (2016).
 - [32] Bilge Can Yildiz, Alpan Bek, and Mehmet Emre Tasgin, “Plasmon lifetime enhancement in a bright-dark mode coupled system,” *Physical Review B* **101**, 035416 (2020).
 - [33] Here, we use the word QO intentionally, since for such a utilization the quantized particle does not need to emit (fluoresce).
 - [34] Mark I Stockman, “Dark-hot resonances,” *Nature* **467**, 541–542 (2010).
 - [35] Boris Luk’yanchuk, Nikolay I Zheludev, Stefan A Maier, Naomi J Halas, Peter Nordlander, Harald Giessen, and Chong Tow Chong, “The Fano resonance in plasmonic nanostructures and metamaterials,” *Nature Materials* **9**, 707 (2010).
 - [36] Mikhail F Limonov, Mikhail V Rybin, Alexander N Podubny, and Yuri S Kivshar, “Fano resonances in photonics,” *Nature Photonics* **11**, 543–554 (2017).
 - [37] Philippe Tassin, Lei Zhang, Rongkuo Zhao, Aditya Jain, Thomas Koschny, and Costas M Soukoulis, “Electromagnetically induced transparency and absorption in metamaterials: the radiating two-oscillator model and its experimental confirmation,” *Physical Review Letters* **109**, 187401 (2012).
 - [38] Jinna He, Chunzhen Fan, Pei Ding, Shuangmei Zhu, and Erjun Liang, “Near-field engineering of fano resonances in a plasmonic assembly for maximizing cars enhancements,” *Scientific Reports* **6**, 20777 (2016).
 - [39] Jian Ye, Fangfang Wen, Heidar Sobhani, J Britt Lassiter, Pol Van Dorpe, Peter Nordlander, and Naomi J Halas, “Plasmonic nanoclusters: near field properties of the fano resonance interrogated with sers,” *Nano Letters* **12**, 1660–1667 (2012).
 - [40] MA Noginov, G Zhu, AM Belgrave, Reuben Bakker, VM Shalaev, EE Narimanov, S Stout, E Herz, T Suteewong, and U Wiesner, “Demonstration of a spaser-based nanolaser,” *Nature* **460**, 1110–1112 (2009).
 - [41] Mehmet Emre Tasgin, Alpan Bek, and Selen Postaci, “Fano resonances in the linear and nonlinear plasmonic response,” in *Fano Resonances in Optics and Microwaves* (Springer, 2018) pp. 1–31.
 - [42] MA Khan and Michael N Leuenberger, “Optoelectronics with single layer group-vib transition metal dichalcogenides,” *Nanophotonics* **7**, 1589–1600 (2018).
 - [43] Xiaohua Wu, Stephen K Gray, and Matthew Pelton, “Quantum-dot-induced transparency in a nanoscale plasmonic resonator,” *Optics Express* **18**, 23633–23645 (2010).
 - [44] Malin Premaratne and Mark I Stockman, “Theory and technology of spasers,” *Advances in Optics and Photonics* **9**, 79–128 (2017).
 - [45] Andreas W Schell, Philip Engel, Julia FM Werra, Christian Wolff, Kurt Busch, and Oliver Benson, “Scanning single quantum emitter fluorescence lifetime imaging: quantitative analysis of the local density of photonic states,” *Nano Letters* **14**, 2623–2627 (2014).
 - [46] Carmen Palacios-Berraquero, Dhiren M Kara, Alejandro R-P Montblanch, Matteo Barbone, Pawel Latawiec, Duhee Yoon, Anna K Ott, Marko Loncar, Andrea C Ferrari, and Mete Atatüre, “Large-scale quantum-emitter arrays in atomically thin semiconductors,” *Nature Communications* **8**, 1–6 (2017).
 - [47] Wei Wu, Chandriker K Dass, Joshua R Hendrickson, Raul D Montañó, Robert E Fischer, Xiaotian Zhang, Tanushree H Choudhury, Joan M Redwing, Yongqiang Wang, and Michael T Pettes, “Locally defined quantum emission from epitaxial few-layer tungsten diselenide,” *Applied Physics Letters* **114**, 213102 (2019).
 - [48] Artur Branny, Santosh Kumar, Raphaël Proux, and Brian D. Gerardot, “Deterministic strain-induced arrays of quantum emitters in a two-dimensional semiconductor,” *Nature Communications* **8**, 1–7 (2017), arXiv:1610.01406.
 - [49] Galan Moody, John Schaibley, and Xiaodong Xu, “Exciton dynamics in monolayer transition metal dichalcogenides *invited*,” *J. Opt. Soc. Am. B* **33**, C39–C49 (2016).
 - [50] Ashish Arora, Maciej Koperski, Karol Nogajewski, Jacques Marcus, Clément Faugeras, and Marek Potemski, “Excitonic resonances in thin films of WSe_2 : From monolayer to bulk material,” *Nanoscale* **7**, 10421–10429 (2015), arXiv:1503.01682.
 - [51] Iann C. Gerber, Emmanuel Courtade, Shivangi Shree, Cedric Robert, Takashi Taniguchi, Kenji Watanabe, Andrea Balocchi, Pierre Renucci, Delphine Lagarde, Xavier Marie, and Bernhard Urbaszek, “Interlayer excitons in bilayer MoS_2 with strong oscillator strength up to room temperature,” *Physical Review B* **99**, 1–8 (2019), 1811.06469.
 - [52] Christian Schneider, Mikhail M. Glazov, Tobias Korn, Sven Höfling, and Bernhard Urbaszek, “Two-dimensional semiconductors in the regime of strong light-matter coupling,” *Nature Communications* **9** (2018), 10.1038/s41467-018-04866-6, 1804.06771.

- [53] Haining Wang, Changjian Zhang, Weimin Chan, Christina Manolatos, Sandip Tiwari, and Farhan Rana, “Radiative lifetimes of excitons and trions in monolayers of the metal dichalcogenide MoS₂,” *Physical Review B* **93**, 1–11 (2016).
- [54] Laxmi Narayan Tripathi, Oliver Iff, Simon Betzold, Lukasz Dusanowski, Monika Emmerling, Kihwan Moon, Young Jin Lee, Soon-Hong Kwon, Sven Hofling, and Christian Schneider, “Spontaneous emission enhancement in strain-induced wse₂ monolayer-based quantum light sources on metallic surfaces,” *ACS Photonics* **5**, 1919–1926 (2018).
- [55] Yongjun Lee, Seok Joon Yun, Youngbum Kim, Min Su Kim, Gang Hee Han, AK Sood, and Jeongyong Kim, “Near-field spectral mapping of individual exciton complexes of monolayer ws₂ correlated with local defects and charge population,” *Nanoscale* **9**, 2272–2278 (2017).
- [56] Wei Bao, Nicholas J Borys, Changhyun Ko, Joonki Suh, Wen Fan, Andrew Thron, Yingjie Zhang, Alexander Buyanin, Jie Zhang, Stefano Cabrini, *et al.*, “Visualizing nanoscale excitonic relaxation properties of disordered edges and grain boundaries in monolayer molybdenum disulfide,” *Nature Communications* **6**, 1–7 (2015).
- [57] Michele Cotrufo, Liuyang Sun, Junho Choi, Andrea Alù, and Xiaoqin Li, “Enhancing functionalities of atomically thin semiconductors with plasmonic nanostructures,” *Nanophotonics* **8**, 577–598 (2019).
- [58] Shikai Deng, Anirudha V. Sumant, and Vikas Berry, “Strain engineering in two-dimensional nanomaterials beyond graphene,” *Nano Today* **22**, 14–35 (2018).
- [59] Ji Feng, Xiaofeng Qian, Cheng Wei Huang, and Ju Li, “Strain-engineered artificial atom as a broad-spectrum solar energy funnel,” *Nature Photonics* **6**, 866–872 (2012).
- [60] Yu Ming He, Genevieve Clark, John R. Schaibley, Yu He, Ming Cheng Chen, Yu Jia Wei, Xing Ding, Qiang Zhang, Wang Yao, Xiaodong Xu, Chao Yang Lu, and Jian Wei Pan, “Single quantum emitters in monolayer semiconductors,” *Nature Nanotechnology* **10**, 497–502 (2015).
- [61] Jianchen Dang, Sibai Sun, Xin Xie, Yang Yu, Kai Peng, Chenjiang Qian, Shiyao Wu, Feilong Song, Jingnan Yang, Shan Xiao, Longlong Yang, Yunuan Wang, M. A. Rafiq, Can Wang, and Xiulai Xu, “Identifying defect-related quantum emitters in monolayer WSe₂,” *npj 2D Materials and Applications* **4**, 1–7 (2020).
- [62] Krishna P. Dhakal, Shrawan Roy, Houk Jang, Xiang Chen, Won Seok Yun, Hyunmin Kim, Jaedong Lee, Jeongyong Kim, and Jong Hyun Ahn, “Local Strain Induced Band Gap Modulation and Photoluminescence Enhancement of Multilayer Transition Metal Dichalcogenides,” *Chemistry of Materials* **29**, 5124–5133 (2017).
- [63] Pascal Anger, Palash Bharadwaj, and Lukas Novotny, “Enhancement and quenching of single-molecule fluorescence,” *Physical Review Letters* **96**, 113002 (2006).
- [64] Chitrleema Chakraborty, Laura Kinnischtzke, Kenneth M Goodfellow, Ryan Beams, and A Nick Vamvakas, “Voltage-controlled quantum light from an atomically thin semiconductor,” *Nature Nanotechnology* **10**, 507 (2015).
- [65] S Schwarz, A Kozikov, F Withers, JK Maguire, AP Foster, S Dufferwiel, L Hague, MN Makhonin, LR Wilson, AK Geim, *et al.*, “Electrically pumped single-defect light emitters in wse₂,” *2D Materials* **3**, 025038 (2016).
- [66] Emiko Kazuma, Minhui Lee, Jaehoon Jung, Michael Trenary, and Yousoo Kim, “Single-molecule study of a plasmon-induced reaction for a strongly chemisorbed molecule,” *Angewandte Chemie International Edition* **59**, 7960–7966 (2020).
- [67] Selen Postaci, Bilge Can Yildiz, Alpan Bek, and Mehmet Emre Tasgin, “Silent enhancement of sers signal without increasing hot spot intensities,” *Nanophotonics* **7**, 1687–1695 (2018).
- [68] Yu Zhang, Fangfang Wen, Yu-Rong Zhen, Peter Nordlander, and Naomi J Halas, “Coherent fano resonances in a plasmonic nanocluster enhance optical four-wave mixing,” *Proceedings of the National Academy of Sciences* **110**, 9215–9219 (2013).
- [69] P. B. Johnson and R. W. Christy, “Optical constants of the noble metals,” *Phys. Rev. B* **6**, 4370–4379 (1972).
- [70] For the $f = 0.1$ auxiliary QO, we use a 0.5 nm thick layer of SiO₂.
- [71] Here such different modes appear due to the perfect shape of the spherical auxiliary QO. When we increase the mesh-size, fringes disappear. Real-world quantum emitters are not spheres.
- [72] Vladimir M Shalaev and Satoshi Kawata, *Nanophotonics with surface plasmons* (Elsevier, 2006).
- [73] Naturally, there will be complications in scattering process. In this prominent work, however, we aim a basic demonstration of the novel phenomenon, rather than exactly simulating a phenomenon conducted experimentally.
- [74] $S(\Phi)$ relation can also be obtained for high-repetition rate sources.
- [75] Renhe Zhang, Yao Zhang, ZC Dong, S Jiang, C Zhang, LG Chen, L Zhang, Y Liao, J Aizpurua, Y ea Luo, *et al.*, “Chemical mapping of a single molecule by plasmon-enhanced raman scattering,” *Nature* **498**, 82–86 (2013).