

GLOBAL WELL-POSEDNESS AND SCATTERING FOR THE DYSTHE EQUATION IN $L^2(\mathbb{R}^2)$

RAZVAN MOSINCAT, DIDIER PILOD, AND JEAN-CLAUDE SAUT

Dedicated to Kristian B. Dysthe, with friendship and admiration.

ABSTRACT. This paper focuses on the Dysthe equation which is a higher order approximation of the water waves system in the modulation (Schrödinger) regime and in the infinite depth case. We first review the derivation of the Dysthe and related equations. Then we study the initial-value problem. We prove a small data global well-posedness and scattering result in the critical space $L^2(\mathbb{R}^2)$. This result is sharp in view of the fact that the flow map cannot be C^3 continuous below $L^2(\mathbb{R}^2)$. Our analysis relies on linear and bilinear Strichartz estimates in the context of the Fourier restriction norm method. Moreover, since we are at a critical level, we need to work in the framework of the atomic space U_S^2 and its dual V_S^2 of square bounded variation functions. We also prove that the initial-value problem is locally well-posed in $H^s(\mathbb{R}^2)$, $s > 0$. Our results extend to the finite depth version of the Dysthe equation.

Keywords : Dysthe equation; Cauchy problem; scattering.

1. INTRODUCTION

1.1. Introduction of the model and physical motivation. It is well-known that the full water waves system is too complex to be used to describe rigorously long time dynamics of waves except in physically trivial situations. A well-known procedure (going back to Lagrange!) is instead to derive simpler asymptotic models in various relevant regimes. We refer to the book [31] for a systematic, up to date and rigorous approach. The first step is of course to find suitable non-dimensional small parameters. This will allow to derive approximations of the water waves system by expansions with respect to the parameters. For instance, in order to derive rigorous error estimates, one expands the nonlocal Dirichlet to Neumann operator which appears in the Zakharov-Craig-Sulem formulation of the water waves problem, see [31].

In this introductory section we motivate and recall the derivation of higher order nonlinear Schrödinger-type equations in the context of surface water waves.

We are interested here in the so-called modulation regime which aims to describe the weakly nonlinear modulation of a train of surface gravity (or gravity-capillary) waves. In other words, one investigates the behavior of wave packets, which are slowly modulated oscillating waves. The modulation equations describe the time evolution of the amplitude of these oscillations. We refer to [31], Chapter 8 for an extensive description of this regime and of the many related asymptotic models it leads to.

To be more precise, one introduces h a typical depth of the fluid layer, a a typical amplitude of the wave and λ a typical wavelength in the horizontal directions (assumed to be isotropic). Then we set $\epsilon = a/h$ and $\mu = h^2/\lambda^2$. We say that we are in deep water when the shallowness assumption $\mu \ll 1$ (which is made for instance in the Boussinesq-KdV regime) is not satisfied. In this situation it does not make sense to derive asymptotic models to the water waves system in terms of the parameter μ .

Date: March 18, 2022.

2010 Mathematics Subject Classification. Primary: 35A01, 35Q53 ; Secondary: 35Q60 .

Key words and phrases. Dysthe equation, Initial-value problem, Well-posedness.

In the *modulation regime*, the relevant small parameter is the *wave steepness*, $\varepsilon = a/\lambda$ assumed to be small. Following [47] it will be convenient to define finite depth, deep water and infinite depth as $\frac{\lambda}{h}$ being $O(1)$, $O(\varepsilon)$ and 0 respectively.

One first fixes a characteristic carrying wave vector $\mathbf{k}_0 \in \mathbb{R}^d$, $d = 1, 2$ (frequently taken as the unit vector \mathbf{e}_x in the x-direction) and considers perturbations $\mathbf{k} = \mathbf{k}_0 + \Delta\mathbf{k}$. We will denote $k = |\mathbf{k}|$ and $k_0 = |\mathbf{k}_0|$ the corresponding wave numbers. In the physical literature ε is frequently expressed as ak .

In order to derive the NLS equation or higher order equations such as the Dysthe equation, one assumes a narrow bandwidth requirement, that is

$$\delta := \frac{|\Delta\mathbf{k}|}{k_0} \ll 1,$$

where $\Delta\mathbf{k}$ is the modulated wave vector. The parameter δ measures the dispersive effects. More precisely, one takes $\delta = O(\varepsilon)$, so that the dispersive effects and the wave steepness are of same order and actually one can take $\delta = \varepsilon$.

The dispersion relation will be in particular Taylor expanded around the wave number k_0 . At first order in ε (that is neglecting the $O(\varepsilon^2)$ terms in the expansion), one obtains nonlinear Schrödinger (NLS) type equations or systems (Benney-Roskes, Davey-Stewartson); we refer to [31] Chapter 8 for a rigorous derivation.

However, as it is recalled in [31], when the wave steepness ε is not very small, the NLS approximation does not match very well with some exact computations, see for instance Longuet-Higgins [36, 37]. As recalled in [8], the NLS equation exhibits an unbounded region of Benjamin-Feir instability in the case of two-dimensional sideband perturbations, which extend outside the regime of a narrow-band spectrum. As a result, energy initially contained at low wave numbers can leak out to higher modes, as shown in the numerical simulations of Martin and Yuen [39].

To overcome those shortcomings Dysthe [12], using the method of multiscales, (see also [21]) proposed in the infinite depth case ($h = +\infty$) to include the $O(\varepsilon^2)$ terms neglected in the derivation of the NLS equation, still under the bandwidth assumption $\delta = O(\varepsilon)$,

A consequence of the higher order expansion is the appearance of an induced mean field ϕ which solves an elliptic boundary-value problem and which can be expressed as a function of the amplitude ψ via a Riesz transform.

In the infinite depth case (then the dispersion relation of the water waves system is $\omega(\mathbf{k}) = |\mathbf{k}|^{1/2}$), and taking (without loss of generality) the carrier wave number $\mathbf{k}_0 = \mathbf{e}_x$, one obtains the following equation in dimensionless form for the slow envelope of the wave (see [12, 21] for a formal derivation and [31] for a rigorous one):¹

$$\begin{aligned} (1.1) \quad & 2i(\partial_t\psi + \frac{1}{2}\partial_x\psi) + \varepsilon(-\frac{1}{4}\partial_x^2 + \frac{1}{2}\partial_y^2)\psi - 4\varepsilon|\psi|^2\psi \\ & = \varepsilon^2\frac{i}{8}(\partial_x^3 - 6\partial_x\partial_y^2)\psi + 2i\varepsilon^2\psi(\psi\partial_x\bar{\psi} - \bar{\psi}\partial_x\psi) - 10\varepsilon^2i|\psi|^2\partial_x\psi - 4\varepsilon^2\psi\partial_x\mathcal{R}_x(|\psi|^2), \end{aligned}$$

where \mathcal{R}_x is the Riesz transform defined by $\widehat{\mathcal{R}_x f}(\xi, \mu) = -i\frac{\xi}{|(\xi, \mu)|}\hat{f}(\xi, \mu)$. If τ and (X, Y) are the physical variables, the derivatives $\partial_t, \partial_x, \partial_y$ are taken with respect to the slow variables $t = \varepsilon\tau, x = \varepsilon X, y = \varepsilon Y$. Note that the relevant time scale for (1.1) is $t = O(\frac{1}{\varepsilon})$, (and $\tau = O(\frac{1}{\varepsilon^2})$ for the physical time).

One can obtain similar equations when surface tension is taken into account (gravity-capillary waves), we refer for instance to [20] and to the survey article [10]. They write

$$\begin{aligned} (1.2) \quad & 2i(\partial_t\psi + c_g\partial_x\psi) + \varepsilon(p\partial_x^2 + q\partial_y^2) - \varepsilon\gamma|\psi|^2\psi \\ & = \varepsilon^2(-is\partial_x\partial_y^2\psi - ir\partial_x^3\psi - iu\psi^2\partial_x\bar{\psi} + iv|\psi|^2\partial_x\psi + \psi\partial_x\mathcal{R}_x(|\psi|^2)), \end{aligned}$$

¹The order one expansion in ε leads to the nonelliptic nonlinear Schrödinger equation, see [49].

where $c_g = \frac{\omega_0}{2k_0} \frac{1+3\kappa}{(1+\kappa)}$ is the group velocity and p, q, s, r, u and v are real parameters depending on the surface tension parameter $\kappa \geq 0$. More precisely

$$\begin{aligned} p &= \frac{3\kappa^2 + 6\kappa - 1}{4(1+\kappa)^2}, \quad q = \frac{1+3\kappa}{2(1+\kappa)}, \\ r &= -\frac{(1-\kappa)(1+6\kappa+\kappa^2)}{8(1+\kappa)^3}, \quad s = \frac{3+2\kappa+3\kappa^2}{4(1+\kappa)^2}, \\ \gamma &= \frac{8+\kappa+2\kappa^2}{8(1-2\kappa)(1+\kappa)}, \quad u = \frac{(1-\kappa)(8+\kappa+2\kappa^2)}{16(1-2\kappa)(1+\kappa)^2}, \\ v &= \frac{3(4\kappa^4 + 4\kappa^3 - 9\kappa^2 + \kappa - 8)}{8(1+\kappa)^2(1-2\kappa)^2}. \end{aligned}$$

When $\kappa = 0$, one obtains a variant of the Dysthe equation. Note that q and s are strictly positive, while p can achieve both signs. In particular it is negative for pure gravity waves as in the original Dysthe equation and positive for pure capillary waves. In the latter case (κ infinite) (1.2) becomes a variant of the modified Zakharov-Kuznetsov equation, i.e.

$$\begin{aligned} (1.3) \quad & 2i(\partial_t \psi + \frac{3}{2}\partial_x \psi) + \frac{3}{4}\partial_x^2 \psi + \frac{3}{2}\partial_y^2 \psi + \frac{1}{8}|\psi|^2 \psi \\ & = -\frac{3}{4}i\partial_{xyy}^3 \psi - \frac{i}{8}\partial_x^3 \psi - \frac{1}{16}\psi^2 \overline{\partial_x \psi} + \frac{3}{8}i|\psi|^2 \partial_x \psi + \psi \partial_x \mathcal{R}_x(|\psi|^2). \end{aligned}$$

The solutions of the Dysthe equation have been compared to those of the NLS equation (see [34, 35, 19]) showing differences with solutions of the NLS equation, and in particular an increase of the group velocity and asymmetry of the envelope of the surface elevation with respect to the peak of the wave profile. We also refer to [13, 22] for comparisons with experimental data.

Remark 1.1. Stiassnie [43] and Stiassnie-Shemer [44] have derived the Dysthe equation as a particular limit of the integro-differential equation which describes nonlinear four-wave interactions within a spectrum of surface waves introduced by Zakharov in [49]. We refer to the survey article [10] for a pedagogical description of the derivation by both methods (the multiscales and the Zakharov equation).

In the finite depth situation, that is when the domain of the fluid is the strip $[-h < z < 0]$, the complex envelope ψ of the wave is coupled to the potential ϕ of the induced current (see [34]). One gets (dropping the dependence in ε):

$$(1.4) \quad \begin{cases} \partial_t \psi + \frac{\omega}{2k_0} \partial_x \psi + i \frac{\omega}{2k_0^3} \left(\frac{1}{4} \partial_x^2 \psi - \frac{1}{2} \partial_y^2 \psi \right) + \frac{i}{2} \omega k^2 |\psi|^2 \psi \\ \quad - \frac{1}{16} \frac{\omega}{k^3} (\partial_x^3 \psi - 6 \partial_{xyy}^3 \psi) - \frac{\omega k_0}{4} \psi^2 \partial_x \bar{\psi} + \frac{3}{2} \omega k_0 |\psi|^2 \partial_x \psi + i k_0 \psi \partial_x \phi|_{z=0} = 0, \\ \partial_x^2 \phi + \partial_y^2 \phi + \partial_z^2 \phi = 0 \quad (-h < z < 0), \\ \partial_z \phi = \frac{\omega}{2} \partial_x |\psi|^2 \quad (z = 0), \\ \partial_z \phi = 0 \quad (z = -h). \end{cases}$$

Although this paper is devoted to the two-dimensional case we comment briefly on the interesting equation arising in the one-dimensional case obtained from (1.4) by dropping the terms with a partial derivative in y . The system for ϕ is easily solved by taking the Fourier transform in x and we find that

$$\partial_x \phi|_{z=0} = \frac{\omega}{2} \partial_x \mathcal{L}_h(|\psi|^2)$$

where the nonlocal operator \mathcal{L}_h is defined in Fourier variables by

$$\widehat{\mathcal{L}_h f}(\xi) = i \coth(h\xi) \hat{f}(\xi).$$

Note that in the infinite depth case $h = +\infty$, $\mathcal{L}_{+\infty}$ is given by $\widehat{\mathcal{L}_{+\infty} f}(\xi) = i \operatorname{sign}(\xi) \hat{f}(\xi)$ that is $\mathcal{L}_{+\infty} = -\mathcal{H}$, where \mathcal{H} is the Hilbert transform.

We thus can write (1.4) in the one-dimensional case as a single equation:

$$(1.5) \quad \begin{aligned} & \partial_t \psi + \frac{\omega}{2k_0} \partial_x \psi + i \frac{\omega}{8k_0^2} \partial_x^2 \psi + \frac{i}{2} \omega k_0^2 |\psi|^2 \psi \\ & - \frac{1}{16} \frac{\omega}{k_0^3} \partial_x^3 \psi - \frac{\omega k_0}{4} \psi^2 \partial_x \bar{\psi} + \frac{3}{2} \omega k_0 |\psi|^2 \partial_x \psi + i \frac{\omega k_0}{2} \psi \partial_x \mathcal{L}_h(|\psi|^2) = 0. \end{aligned}$$

This equation is reminiscent of the complex modified KdV equation

$$(1.6) \quad \partial_t u + u^2 \partial_x u + \partial_x^3 u = 0.$$

In fact by eliminating the transport term in (1.5) by the change of variable $X = x - \frac{\omega}{2k_0} t$ (we will keep the notation x for the spatial variable), then writing, with $\alpha = -\frac{1}{16} \frac{\omega}{k^3}$ and $\beta = \frac{\omega}{8k^2}$

$$\left(\xi + \frac{\beta}{3\alpha} \right)^3 = \xi^3 + \frac{\beta}{\alpha} \xi^2 + \frac{\beta^2 \xi}{3\alpha^2} + \frac{\beta^3}{27\alpha^3},$$

an easy computation shows that the fundamental solution of the linearization of (1.5) can be expressed as

$$\mathfrak{A}(x, t) = \frac{1}{(t\alpha)^{1/3}} \exp\left(\frac{2it\beta^3}{27\alpha^2}\right) \exp\left(\frac{-i\beta x}{2\alpha^2}\right) \text{Ai}\left(\frac{1}{t^{1/3}\alpha^{1/3}}\left(x - \frac{\beta^2}{3\alpha}t\right)\right),$$

where we have used here the Airy function

$$\text{Ai}(z) = \int_{-\infty}^{\infty} e^{i(\xi^3 + iz\xi)} d\xi.$$

It follows that the dispersive estimates for \mathfrak{A} are essentially the same as those of the linearized KdV equation and one obtains for (1.5) the same results as for the complex modified KdV equation (1.6) (see [25]), that is the local well-posedness of the initial value problem in $H^s(\mathbb{R})$, $s \geq \frac{1}{4}$.

In the two-dimensional case, the mean flow potential is solution of the system (see [44])

$$(1.7) \quad \begin{cases} \partial_x^2 \phi + \partial_y^2 \phi + \partial_z^2 \phi = 0, & -h < z < 0, \\ \partial_z \phi = \frac{\omega}{2} \partial_x |\psi|^2, & z = 0, \\ \partial_z \phi = 0, & z = -h. \end{cases}$$

This system is easily solved after Fourier transform in the (x, y) variables and one finds

$$(1.8) \quad \partial_x \phi|_{z=0} = \frac{\omega}{2} \partial_x \mathcal{L}_h(|\psi|^2), \quad \text{where} \quad \widehat{\mathcal{L}_h f}(\xi, \mu) = i \frac{\xi}{\sqrt{\xi^2 + \mu^2}} \coth(h\sqrt{\xi^2 + \mu^2}) \hat{f}(\xi, \mu),$$

leading us to the following Dysthe equation in finite depth, having the same structure as the previous ones,

$$(1.9) \quad \begin{aligned} & \partial_t \psi + \frac{\omega}{2k_0} \partial_x \psi + i \frac{\omega}{2k_0^2} \left(\frac{1}{4} \partial_x^2 \psi - \frac{1}{2} \partial_y^2 \psi \right) + \frac{i}{2} \omega k_0^2 |\psi|^2 \psi \\ & - \frac{1}{16} \frac{\omega}{k_0^3} (\partial_x^3 \psi - 6 \partial_{xyy}^3 \psi) - \frac{\omega k_0}{4} \psi^2 \partial_x \bar{\psi} + \frac{3}{2} \omega k_0 |\psi|^2 \partial_x \psi + i \frac{\omega k_0}{2} \psi \partial_x \mathcal{L}_h(|\psi|^2) = 0. \end{aligned}$$

Observe that formally $\mathcal{L}_h \rightarrow -\mathcal{R}_x$ as $h \rightarrow +\infty$. Moreover, the symbol of the operator $\partial_x \mathcal{L}_h$ is bounded close to the origin since

$$(1.10) \quad i \frac{\xi^2}{\sqrt{\xi^2 + \mu^2}} \coth[(\xi^2 + \mu^2)^{1/2} h] \sim \frac{i}{h} \frac{\xi^2}{\xi^2 + \mu^2} \lesssim 1 \quad \text{whenever} \quad |(\xi, \mu)| \lesssim 1.$$

Remark 1.2. Trulsen and Dysthe [47, 46] have extended the Dysthe equation by relaxing the narrow bandwidth condition to

$$\delta = \frac{|\Delta \mathbf{k}|}{k_0} = O(\varepsilon^{1/2}).$$

More precisely, in the case $\frac{\lambda}{h} = O(\varepsilon^{1/2})$, Trulsen and Dysthe obtained the system posed on the strip $\mathbb{R}_{xy}^2 \times (-h, 0)_z$ (dropping the dependence on ε), where the complex envelope ψ of the wave is coupled to the potential ϕ of the induced current:

$$(1.11) \quad \begin{cases} \partial_t \psi + \frac{1}{2} \partial_x \psi + \frac{i}{8} \partial_x^2 \psi - \frac{i}{4} \partial_y^2 \psi - \frac{1}{16} \partial_x^3 \psi + \frac{3}{8} \partial_{xyy}^3 \psi \\ \quad - \frac{5i}{128} \partial_x^4 \psi + \frac{15i}{32} \partial_{xxyy}^4 \psi - \frac{3i}{32} \partial_y^4 \psi + \frac{i}{2} |\psi|^2 \psi + \frac{7}{256} \partial_x^5 \psi - \frac{35}{64} \partial_{3x2y}^5 \psi \\ \quad + \frac{21}{64} \partial_{x4y}^5 \psi + \frac{3}{2} |\psi|^2 \partial_x \psi - \frac{1}{4} \psi^2 \partial_x \bar{\psi} + i \psi \partial_x \phi = 0 \quad \text{at } z = 0, \\ \partial_x^2 \phi + \partial_y^2 \phi + \partial_z^2 \phi = 0 \quad (-h < z < 0), \\ \partial_z \phi = \frac{1}{2} \partial_x |\psi|^2 \quad (z = 0), \\ \partial_z \phi = 0 \quad (z = -h). \end{cases}$$

This has the effect of adding to the Dysthe equation fourth- and fifth-order linear dispersive terms, while keeping the same nonlinear terms. This makes the Cauchy theory easier and this will be developed in a subsequent paper.

As noticed in [47], this equation fits better than the Dysthe type equations with the exact results in [38].

In the rest of this paper we normalize out the wave steepness parameter ε in (1.1) by the change of variables $(t, x, y, \psi) \mapsto \left(\frac{t}{\varepsilon}, \frac{x}{\varepsilon}, \frac{y}{\varepsilon}, \frac{\psi}{\varepsilon}\right)$ and we study the initial-value problem for the Dysthe equation written in the form

$$(1.12) \quad \begin{aligned} 2i(\partial_t v + \frac{1}{2} \partial_x v) + (-\frac{1}{4} \partial_x^2 v + \frac{1}{2} \partial_y^2 v) - \frac{i}{8} (\partial_x^3 v - 6 \partial_{xyy}^3 v) \\ = 4 |v|^2 v + 2iv(v \partial_x \bar{v} - \bar{v} \partial_x v) - 10i |v|^2 \partial_x v - 4v \partial_x \mathcal{R}_x(|v|^2), \end{aligned}$$

where $v = v(t, x, y)$ is a complex-valued function, $t \in \mathbb{R}$, $(x, y) \in \mathbb{R}^2$ and \mathcal{R}_x denotes the Riesz transform in the x variable.

Concerning the conservation laws of Dysthe type systems, one checks readily that they conserve formally the L^2 -norm, namely that

$$(1.13) \quad M[v](t) = \int |v(t, x, y)|^2 dx dy = M[v](0).$$

On the other hand and contrary to the nonlinear Schrödinger equation, they do not seem to possess a conserved energy. Note however that *Hamiltonian versions* of the Dysthe equation were derived in [8, 7, 9] by expanding the Hamiltonian in the Zakharov-Craig-Sulem formulation of the water waves system and in [15] starting from the Zakharov integro-differential equation ([49]) enhanced with the new kernel of Krasitskii [30] which yields an Hamiltonian version of the Zakharov equation.²

As explained in the Introduction of [15] one reason of the non Hamiltonian form of the “classical” Dysthe equation is the choice of variable. Usually, the higher order NLS type equations are expressed in either the first-harmonic complex amplitude of the surface elevation or in the first complex amplitude of the velocity potential evaluated at the water surface at rest. While these variables are natural from a physical point of view, they are not optimal from a mathematical point of view since they lead to a non-Hamiltonian equation. We will study the mathematical properties of the Hamiltonian higher order NLS equations in a subsequent paper.

1.2. Statement of main results and comments. We now review known results on the initial-value problem (IVP) for Dysthe type systems. Local well-posedness in analytic classes for a family of systems comprising the Dysthe ones has been established by A. de Bouard [3]. The local well-posedness for initial data in $H^3(\mathbb{R}^2)$ has been proved by Chihara [6], who applied techniques developed for derivative nonlinear Schrödinger equations to these third-order equations. Koch and Saut [27] obtained the local well-posedness for data in $H^{\frac{3}{2}}(\mathbb{R}^2)$ by using solely the local smoothing effects of the underlying group.

²A Hamiltonian form of the one-dimensional Dysthe equation is obtained in [14] via a gauge transform.

The result in [27] was improved very recently to $s > 1$ by Grande, Kurianski and Staffilani [16]. The proof is based on a nontrivial adaptation of the Kenig, Ponce and Vega techniques [25] to this two-dimensional problem. In particular a new maximal function estimate associated to the linear part of Dysthe is proved. The authors also proved that the IVP associated to (1.12) is ill-posed in $H^s(\mathbb{R}^2)$ for $s < 0$ in the sense that the flow map data-solution cannot be C^3 in $H^s(\mathbb{R}^2)$, $s < 0$.

Finally we recall that other papers have been devoted to the initial-value problem for relevant third-order nonlinear Schrödinger equations involving spatial derivatives in the nonlinearities, see for instance [31].

The aim of this article is to improve these results by using the Fourier restriction norm method. Our first theorem proves global well-posedness and scattering for small initial data down to the critical space $L^2(\mathbb{R}^2)$.

Theorem 1.1. *There exists $\delta > 0$ such that for any $v_0 \in L^2(\mathbb{R}^2)$ with $\|v_0\|_{L^2} < \delta$, there exists a unique $v \in \mathcal{Y} \subset C(\mathbb{R} : L^2(\mathbb{R}^2))$ solution to (1.12) with $v|_{t=0} = v_0$, the flow map $v_0 \mapsto v$ from $\{v_0 \in L^2(\mathbb{R}^2) : \|v_0\|_{L^2} < \delta\}$ to \mathcal{Y} is smooth, and there exist $v_0^\pm \in L^2(\mathbb{R}^2)$ such that*

$$v(t) - v_{\text{lin}}^\pm(t) \rightarrow 0 \text{ in } L^2(\mathbb{R}^2), \text{ as } t \rightarrow \pm\infty,$$

where v_{lin}^\pm is the solution to the linear Dysthe equation starting from v_0^\pm .

Remark 1.3. The resolution space \mathcal{Y} is defined in Section 4 – see (4.11) and (2.11).

Remark 1.4. This result is sharp due to the ill-posedness result proved in [16].

Remark 1.5. Theorem 1.1 can be seen as an analogous result to the one by Cazenave and Weissler in [5] for the L^2 -critical two-dimensional cubic NLS, which is the first-order approximation to the water wave system in the modulation regime and narrow bandwidth, while the Dysthe equation arises as a second-order approximation.

We also obtain a local well-posedness result in $H^s(\mathbb{R}^2)$, $s > 0$.

Theorem 1.2. *Let $s > 0$. For any $v_0 \in H^s(\mathbb{R}^2)$ there exist $T = T(\|v_0\|_{H^s}) > 0$ and a unique solution $v \in \mathcal{X}_T^{s, \frac{1}{2}+} \subset C([0, T] : H^s(\mathbb{R}^2))$ to (1.12) with $v|_{t=0} = v_0$. Moreover, for any $T' \in (0, T)$, there exists a neighborhood \mathcal{U} of $v_0 \in H^s(\mathbb{R}^2)$ such that the flow map data-solution $w_0 \in \mathcal{U} \mapsto w \in C([0, T'] : H^s(\mathbb{R}^2))$ is smooth.*

Remark 1.6. The well-posedness results obtained in Theorems 1.1 and 1.2 for the Dysthe equation in dimension $d = 2$ hold in lower regularity than the ones for the corresponding one-dimensional equation (1.5), for which well-posedness only holds in $H^s(\mathbb{R})$, $s \geq \frac{1}{4}$ by using the Kenig, Ponce and Vega techniques of [25]. This is due to the fact that the two-dimensional equation is more dispersive than its one-dimensional counterpart which allows to recover $1/4$ of a derivative instead of only $1/8$ of a derivative in the L^4 -Strichartz estimates. As explained below, these estimates are crucial to handling the resonant high \times high \times high frequency interactions region in the nonlinear terms.

Remark 1.7. It is clear from the proofs in Section 4 and the bound (1.10) that the results of Theorems 1.1 and 1.2 also hold for the finite depth Dysthe equation (1.9).

We now discuss the main ingredients for the proofs of Theorems 1.1 and 1.2. First observe that by a change of variables of the form

$$(1.14) \quad (t, x, y, v) \mapsto (t, x + a_1 t, y, e^{ia_2 x} e^{ia_3 t} v)$$

where a_1, a_2, a_3 are real numbers (see Lemma 2.1 below) we can remove the first- and second-order derivative linear terms on the left-hand side of (1.12). Thus, for the sake of simplicity, we work on the following simplified version³ of the Dysthe equation

$$(1.15) \quad \partial_t u + \partial_x (\partial_x^2 - 3\partial_y^2) u = \sum_{j=1}^4 c_j \mathcal{N}_j(u, u, u),$$

³We prefer not to normalize both constants in $\partial_x(\partial_x^2 - 3\partial_y^2)$ and thus work with a simpler Hessian determinant expression for its Fourier symbol (see (3.7)).

where $c_1, c_2, c_3, c_4 \in \mathbb{C}$ are constants and the nonlinearities $\mathcal{N}_j(u, u, u)$ are given by

$$\begin{aligned}
 \mathcal{N}_1(u, u, u) &= |u|^2 u, \\
 \mathcal{N}_2(u, u, u) &= |u|^2 \partial_x u, \\
 \mathcal{N}_3(u, u, u) &= u^2 \partial_x \bar{u}, \\
 \mathcal{N}_4(u, u, u) &= u \partial_x \mathcal{R}_x(|u|^2).
 \end{aligned}
 \tag{1.16}$$

We note that for $c_1 = 0$, the equation is invariant under the scaling transformation

$$u(t, x, y) \mapsto u_\lambda(t, x, y) := \lambda u(\lambda^3 t, \lambda x, \lambda y)
 \tag{1.17}$$

and since $\|u_\lambda(0)\|_{L^2(\mathbb{R}^2)} = \|u(0)\|_{L^2(\mathbb{R}^2)}$, we regard $L^2(\mathbb{R}^2)$ as the scaling critical Sobolev space of (1.15) (and of (1.12) due to the invariance of the Sobolev norms under (1.14)). We will prove the equivalent results of Theorems 1.1 and 1.2 for the IVP associated to (1.15) which has the advantage of having a homogeneous linear part. Theorems 1.1 and 1.2 will then be concluded by inverting the change of variables (1.14).

The proof of the small data global well-posedness and scattering result in $L^2(\mathbb{R}^2)$ for (1.15) relies on linear and bilinear Strichartz estimates in the context of the Fourier restriction norm method. We refer for instance to the work of Hadac, Herr and Koch [18] and Molinet, Saut and Tzvetkov [41] for the KP-II equation, Grünrock and Herr [17] and Molinet, Pilod [40] for the Zakharov-Kuznetsov equation, Kinoshita [26] for the modified Zakharov-Kuznetsov equation and Angelopoulos [1] and Kazeykina, Muñoz [23] for the Novikov-Veselov equation. The bilinear Strichartz estimates are used to deal with the low \times low \times high and low \times high \times high frequency interactions in the nonlinearity. To handle the resonant case corresponding to the high \times high \times high frequency interactions, we use a sharp linear Strichartz estimate proved by Kenig, Ponce and Vega in [24]. This estimate could also be obtained from the dispersion estimates of Ben-Artzi, Koch and Saut in [2] or as a corollary of a result of Carbery, Kenig, and Ziesler [11] for homogeneous dispersive operators. Finally, since we are at a critical level, we need to work in the framework of the atomic space U_S^2 and its dual V_S^2 of square bounded variation functions introduced by Koch and Tataru in [28, 29] (see also [18]) in order to derive the main trilinear estimate.

The proof of the arbitrarily large data local well-posedness in $H^s(\mathbb{R}^2)$, $s > 0$, for (1.15) follows the same strategy as above. Note however that the use of $U^2 - V^2$ space is not needed anymore since the problem is subcritical. It is thus enough to work with the usual $X^{s,b}$ spaces.

It is worth noting that this method of proof only relies on the fact that the nonlinearities $\mathcal{N}_j(u, u, u)$, $j = 1, \dots, 4$ in (1.15) are cubic with at most one derivative, but not on their specific structures. Indeed, it is clear from Plancherel's identity that the linear Strichartz estimate (3.5) and the bilinear Strichartz estimates (3.8)-(3.9) also hold if one replace φ , u_1 , u_2 by $\mathcal{F}^{-1}(|\widehat{\varphi}|)$, $\mathcal{F}^{-1}(|\widehat{u}_1|)$ and $\mathcal{F}^{-1}(|\widehat{u}_2|)$.

We observe that contrary to the linear Zakharov-Kuznetsov symbol, the Hessian of the dispersion relation of the Dysthe equation (1.15) is sign-definite, which allows us to recover $1/4$ of a derivative in the L^4 -Strichartz estimate (see (3.6)). A similar estimate only holds outside of cones centered at the origin for the linear Zakharov-Kuznetsov equation or with a weaker gain of $1/8$ derivatives (see the discussion in Remark 3.1 of [40]). This is the reason why we can reach $L^2(\mathbb{R}^2)$ in Theorem 1.1, while a similar result only holds for $H^s(\mathbb{R}^2)$, $s \geq \frac{1}{4}$ for the modified Zakharov-Kuznetsov equation (see [32, 42, 26]). In particular, Kinoshita constructed in [26] a counterexample localized in the resonant high \times high \times high frequency interaction region proving the sharpness of this result.

The argument of Kinoshita [26] could be applied to the limit version of the Dysthe equation with surface tension (1.3) providing a local well-posedness result in $H^{\frac{1}{4}}(\mathbb{R}^2)$ for the associated Cauchy problem. Note that a change of variable similar to the one in Lemma 2.1 could also be applied in this case to obtain a homogeneous linear part, but this time of the Zakharov-Kuznetsov type.

The rest of the paper is organized as follows: in the next section we introduce the notations, the crucial change of variable which allows to work with a homogenous linear part, define the function

spaces and recall some of their important properties. In Section 3, we recall the linear Strichartz estimates and derive the bilinear estimates. Those estimates are used in Section 4 to prove the trilinear estimates in \mathbb{R}^2 . Finally, we conclude the paper in Section 5 with some interesting open questions.

2. PRELIMINARIES AND LINEAR ESTIMATES

2.1. Notation. For any positive numbers a and b , the notation $a \lesssim b$ means that there exists a positive constant c such that $a \leq cb$, we use $a \lesssim_\theta b$ when we find it necessary to make it explicit that the constant c depends on the parameter θ , and $a \ll b$ when c is small (e.g. $c \leq 10^{-2}$). We also write $a \sim b$ when $a \lesssim b$ and $b \lesssim a$. We denote by $|S|$ the Lebesgue measure of a measurable set S of \mathbb{R}^d , whereas $\#F$ denotes the number of elements of a finite set F . The indicator function of a set A is denoted by $\mathbb{1}_A$. For $u = u(t, x, y) \in \mathcal{S}(\mathbb{R}^3)$, we use the notation $\mathcal{F}(u)$, or \widehat{u} to denote the space-time Fourier transform of u , whereas $\mathcal{F}_{xy}(u)$, or $(u)^{\wedge_{xy}}$, respectively $\mathcal{F}_t(u) = (u)^{\wedge_t}$, will denote its Fourier transform in space, respectively in time.

For $s \in \mathbb{R}$, we define J^s and D^s , the Bessel and Riesz potentials of order $-s$, by

$$J^s u = \mathcal{F}_{xy}^{-1}((1 + |(\xi, \mu)|^2)^{\frac{s}{2}} \mathcal{F}_{xy}(u)) \quad \text{and} \quad D^s u = \mathcal{F}_{xy}^{-1}(|(\xi, \mu)|^s \mathcal{F}_{xy}(u)).$$

We also recall here that the Riesz transform \mathcal{R}_x that appears in the nonlinearity of the Dysthe equation is given by

$$\mathcal{R}_x u = -i \mathcal{F}_{xy}^{-1} \left(\frac{\xi}{\sqrt{\xi^2 + \mu^2}} \mathcal{F}_{xy}(u)(\xi, \mu) \right).$$

Let $S(t) := e^{-t\partial_x(\partial_x^2 - 3\partial_y^2)}$ denote the unitary group associated with the linear part of equation (1.15), which is to say,

$$(2.1) \quad \mathcal{F}_{xy}(S(t)\varphi)(\xi, \mu) = e^{itw(\xi, \mu)} \mathcal{F}_{xy}(\varphi)(\xi, \mu),$$

where

$$w(\xi, \mu) = \xi^3 - 3\xi\mu^2.$$

We also define the resonance function R by

$$(2.2) \quad \begin{aligned} R(\xi_1, \mu_1, \xi_2, \mu_2) &:= w(\xi_1 + \xi_2, \mu_1 + \mu_2) - w(\xi_1, \mu_1) - w(\xi_2, \mu_2) \\ &= 3\xi_1\xi_2(\xi_1 + \xi_2) - 3\xi_2\mu_1^2 - 3\xi_1\mu_2^2 - 6(\xi_1 + \xi_2)\mu_1\mu_2. \end{aligned}$$

Throughout the paper, we fix a smooth cutoff function η such that

$$\eta \in C_0^\infty(\mathbb{R}), \quad 0 \leq \eta \leq 1, \quad \eta|_{[-5/4, 5/4]} = 1 \quad \text{and} \quad \text{supp}(\eta) \subset [-8/5, 8/5].$$

For $k \in \mathbb{N}^* = \mathbb{Z} \cap [1, +\infty)$, we define

$$\phi(\xi) = \eta(\xi) - \eta(2\xi), \quad \phi_{2^k}(\xi, \mu) := \phi(2^{-k}|(\xi, \mu)|).$$

and

$$\psi_{2^k}(\xi, \mu, \tau) = \phi(2^{-k}(\tau - w(\xi, \mu))).$$

By convention, we also denote

$$\phi_1(\xi, \mu) = \eta(|(\xi, \mu)|), \quad \text{and} \quad \psi_1(\xi, \mu, \tau) = \eta(\tau - w(\xi, \mu)).$$

Any summations over capitalized variables such as N , L , K or M are presumed to be dyadic with N , L , K or $M \geq 1$, i.e. these variables range over numbers of the form $\{2^k : k \in \mathbb{N}\}$. Then, we have that

$$(2.3) \quad \begin{aligned} \text{supp}(\phi_N) &\subset \left\{ \frac{5}{8}N \leq |(\xi, \mu)| \leq \frac{8}{5}N \right\} =: I_N, \quad N \geq 2, \\ \text{supp}(\phi_1) &\subset \left\{ |(\xi, \mu)| \leq \frac{8}{5} \right\} =: I_1, \end{aligned}$$

and

$$\sum_N \phi_N(\xi, \mu) = 1.$$

Let us define the Littlewood-Paley multipliers by

$$P_N u = \mathcal{F}_{xy}^{-1}(\phi_N \mathcal{F}_{xy}(u)), \quad Q_L u = \mathcal{F}^{-1}(\psi_L \mathcal{F}(u)).$$

2.2. Linear transformation.

Lemma 2.1. *The function $v = v(t, x, y)$ is a solution to*

$$(2.4) \quad \partial_t v + \alpha_1 \partial_x (\partial_x^2 - 3\partial_y^2) v + \alpha_2 i (\partial_x^2 - \partial_y^2) v + \alpha_3 \partial_x v = \sum_{j=1}^4 c_j \mathcal{N}_j(v, v, v),$$

where $\alpha_1 \in \mathbb{R} \setminus \{0\}$, $\alpha_2, \alpha_3 \in \mathbb{R}$, and $\mathcal{N}_j(v, v, v)$, $j = 1, \dots, 4$ are the nonlinearities defined in (1.16), if and only if $u = \mathcal{T}(v)$ is a solution to

$$(2.5) \quad \partial_t u + \partial_x (\partial_x^2 - 3\partial_y^2) u = \sum_{j=1}^4 \tilde{c}_j \mathcal{N}_j(u, u, u),$$

where

$$(2.6) \quad \mathcal{T}(v)(t, x, y) := e^{ia_2 x} e^{-ia_3 t} v(t, x + a_1 t, y)$$

and

$$a_1 = \frac{\alpha_2^2}{3\alpha_1^2} + \frac{\alpha_3}{\alpha_1}, \quad a_2 = \frac{\alpha_2}{3\alpha_1}, \quad a_3 = \frac{\alpha_2 \alpha_3}{3\alpha_1^2} + \frac{2}{27} \frac{\alpha_2^3}{\alpha_1^3}.$$

Remark 2.1. Before applying the transformation in Lemma 2.1 to equation (1.12) with $\epsilon = 1$, we first need to rescale the y variable by a factor $\sqrt{2}$ to be exactly in the configuration (2.4).

Proof. First, we observe that the linear symbol $\omega(\xi, \mu)$ associated to (2.4) can be factorized as

$$(2.7) \quad \begin{aligned} \omega(\xi, \mu) &= \alpha_1 (\xi^3 - 3\xi\mu^2) + \alpha_2 (\xi^2 - \mu^2) - \alpha_3 \xi \\ &= \alpha_1 \left(\xi + \frac{\alpha_2}{3\alpha_1} \right)^3 - 3\alpha_1 \left(\xi + \frac{\alpha_2}{3\alpha_1} \right) \mu^2 - \left(\frac{\alpha_2^2}{3\alpha_1} + \alpha_3 \right) \xi - \frac{\alpha_2^3}{27\alpha_1^3}. \end{aligned}$$

Next we proceed arguing as in Section 2 in [33]. If v is a solution of the linear part of (2.4), we first define $f(x, y, t) = e^{i\frac{\alpha_2}{3\alpha_1}x} v(x, y, t)$ and verify that

$$\partial_t f + \alpha_1 \partial_x (\partial_x^2 f - 3\partial_y^2 f) + \left(\alpha_3 + \frac{\alpha_2^2}{3\alpha_1} \right) \partial_x f - i \left(\frac{\alpha_2 \alpha_3}{3\alpha_1} + \frac{2}{27} \frac{\alpha_2^3}{\alpha_1^3} \right) f = 0.$$

Now define $g(x, y, t) = f(x + (\alpha_3 + \frac{\alpha_2^2}{3\alpha_1})t, y, t)$, so that

$$\partial_t g + \alpha_1 \partial_x (\partial_x^2 g - 3\partial_y^2 g) - i \left(\frac{\alpha_2 \alpha_3}{3\alpha_1} + \frac{2}{27} \frac{\alpha_2^3}{\alpha_1^3} \right) g = 0.$$

Finally, by defining $u(t, x, y) = e^{i\left(\frac{\alpha_2 \alpha_3}{3\alpha_1} + \frac{2}{27} \frac{\alpha_2^3}{\alpha_1^3}\right) \frac{t}{\alpha_1}} g(x, y, \frac{t}{\alpha_1}) = \mathcal{T}(v)(t, x, y)$, we conclude that

$$\partial_t u + \partial_x (\partial_x^2 u - 3\partial_y^2 u) = 0.$$

Under this change of variables, one easily checks that the nonlinear terms of (1.12) transform as follows:

$$\begin{aligned} |v|^2 v &\mapsto e^{-ia_2 x} e^{ia_3 t} |u|^2 u, \\ |v|^2 \partial_x v &\mapsto e^{-ia_2 x} e^{ia_3 t} \left(-ia_2 |u|^2 u + |u|^2 \partial_x u \right), \\ v^2 \partial_x \bar{v} &\mapsto e^{-ia_2 x} e^{ia_3 t} \left(ia_2 |u|^2 u + u^2 \partial_x \bar{u} \right), \\ v \partial_x \mathcal{R}_x(|v|^2) &\mapsto e^{-ia_2 x} e^{ia_3 t} u \partial_x \mathcal{R}_x(|u|^2), \end{aligned}$$

which concludes the proof of the lemma. \square

2.3. Function spaces. For $1 \leq p \leq \infty$, $L^p(\mathbb{R}^d)$ is the usual Lebesgue space with the usual norm $\|\cdot\|_{L^p}$, and for $s \in \mathbb{R}$, the Sobolev space $H^s(\mathbb{R}^d)$ denotes the space of all complex-valued functions with the norm $\|u\|_{H^s} = \|J^s u\|_{L^2}$. If $u = u(x, y, t)$ is a function defined for $(x, y) \in \mathbb{R}^2$ and $t \in [0, T]$ with $T > 0$, and if B is one of the spaces defined above, we define the mixed space-time spaces $L_T^p B_{xy}$, $L_t^p B_{xy}$, $L_{xy}^q L_T^p$ by the norms

$$\|u\|_{L_T^p B_{xy}} = \left(\int_0^T \|u(\cdot, \cdot, t)\|_B^p dt \right)^{\frac{1}{p}}, \quad \|u\|_{L_t^p B_{xy}} = \left(\int_{\mathbb{R}} \|u(\cdot, \cdot, t)\|_B^p dt \right)^{\frac{1}{p}},$$

and

$$\|u\|_{L_{xy}^q L_T^p} = \left(\int_{\mathbb{R}^2} \left(\int_0^T |u(x, y, t)|^p dt \right)^{\frac{q}{p}} dx \right)^{\frac{1}{q}},$$

for $1 \leq p, q < \infty$ with the obvious modifications in the case $p = +\infty$ or $q = +\infty$.

Next, we introduce the functional framework introduced by Koch and Tataru in [28] that allows us to reach the critical regularity for the Dysthe equation. We mainly work with the function spaces U^2 and V^2 , however the machinery also requires the use of U^p and V^p spaces with $p \neq 2$ (for example in the proof of (3.16) in Corollary 3.6 below).

Definition 2.2. Let $1 \leq p < \infty$ and let \mathcal{Z} be the set of finite partitions $-\infty = t_0 < t_1 < \dots < t_K = +\infty$. For $\{t_k\}_{k=0}^K \in \mathcal{Z}$ and $\{\phi_k\}_{k=0}^{K-1} \subset L^2(\mathbb{R}^2)$ with $\sum_{k=0}^{K-1} \|\phi_k\|_{L^2}^p = 1$ and $\phi_0 = 0$ we call the function $a : \mathbb{R} \rightarrow L^2(\mathbb{R}^2)$ given by

$$a = \sum_{k=1}^K \mathbb{1}_{[t_{k-1}, t_k)} \phi_{k-1}$$

a U^p -atom and we define the atomic space

$$U^p := \left\{ u = \sum_{j=1}^{\infty} \lambda_j a_j : a_j \text{ } U^p\text{-atom and } \lambda_j \in \mathbb{R} \text{ with } \sum_{j=1}^{\infty} |\lambda_j| < \infty \right\}$$

with norm

$$(2.8) \quad \|u\|_{U^p} := \inf \left\{ \sum_{j=1}^{\infty} |\lambda_j| : u = \sum_{j=1}^{\infty} \lambda_j a_j \text{ with } \lambda_j \in \mathbb{R} \text{ and } a_j \text{ } U^p\text{-atom} \right\}.$$

The function space V^p is defined as the normed space of all functions $v : \mathbb{R} \rightarrow L^2(\mathbb{R}^2)$ such that the limits $\lim_{t \rightarrow -\infty} v(t)$, $\lim_{t \rightarrow +\infty} v(t)$ exist, and for which the norm

$$(2.9) \quad \|v\|_{V^p} := \sup_{\{t_k\}_{k=0}^K \in \mathcal{Z}} \left(\sum_{k=1}^K \|v(t_k) - v(t_{k-1})\|_{L^2}^p \right)^{1/p}$$

is finite, where we use the convention that $v(-\infty) := \lim_{t \rightarrow -\infty} v(t)$ and $v(+\infty) := 0$.

For basic properties of these spaces we refer the reader to [18, 28]. Here we recall without proof the following results.

Lemma 2.3. *Let $1 \leq p < \infty$ and let V_-^p denote the space of all functions $v : \mathbb{R} \rightarrow L^2(\mathbb{R}^2)$ such that $\lim_{t \rightarrow -\infty} v(t) = 0$ and $\lim_{t \rightarrow +\infty} v(t)$ exists, endowed with the norm $\|\cdot\|_{V^p}$ given by (2.9). Also, let $V_{-,rc}^p$ denote the closed subspace of all right-continuous V_-^p -functions. Then, the following embeddings are continuous:*

$$U^p \subset V_{-,rc}^p \quad \text{and} \quad V_{-,rc}^p \subset U^q, \quad p < q.$$

Also, $U^p \subset U^q \subset L^\infty(\mathbb{R}; L^2(\mathbb{R}^2))$, $V^p \subset V^q$, $V_-^p \subset V_-^q$, and $V_{-,rc}^p \subset V_{-,rc}^q$, provided $1 \leq p < q < \infty$.

Lemma 2.4 (Duality lemma). *Let $1 < p < \infty$ and p' denote the Hölder conjugate, i.e. $\frac{1}{p} + \frac{1}{p'} = 1$. There exists a unique bilinear form $B : U^p \times V^{p'} \mapsto \mathbb{C}$ such that $V^{p'} \ni v \mapsto B(\cdot, v) \in (U^p)^*$ is an*

isometric isomorphism. In particular,

$$\begin{aligned}\|u\|_{U^p} &= \sup_{v \in V^{p'}, \|v\|_{V^{p'}}=1} |B(u, v)|, \\ \|v\|_{V^p} &= \sup_{u \in U^{p'} \text{ atom}} |B(u, v)|.\end{aligned}$$

Moreover, if $u \in V_-^1$ is absolutely continuous on compact intervals, then

$$B(u, v) = - \int_{\mathbb{R}} \langle u'(t), v(t) \rangle_{L^2(\mathbb{R}^2)} dt.$$

Next, by recalling the notation $S(t) := e^{-t\partial_x(\partial_x^2 - 3\partial_y^2)}$ of the linear group associated to the linear part of the Dysthe equation (1.15), we define the spaces

$$U_S^p = S(\cdot)U^p \text{ with norm } \|u\|_{U_S^p} = \|S(-\cdot)u\|_{U^p}$$

$$\text{and } V_S^p = S(\cdot)V^p \text{ with norm } \|u\|_{V_S^p} = \|S(-\cdot)u\|_{V^p},$$

and similarly $V_{-,S}^p$ and $V_{-,rc,S}^p$. In particular, by Lemma 2.3, any $u \in U_S^2$ also lies in $V_{-,rc,S}^2$ and we have

$$(2.10) \quad \|u\|_{V_S^2} \lesssim \|u\|_{U_S^2}.$$

The above space behaves well with respect to sharp cut-off functions since $\|\mathbb{1}_{[a,b]}f\|_{U^2} \leq \|f\|_{U^2}$ and $\|f\|_{V^2} \leq \|\mathbb{1}_{[a,b]}f\|_{V^2} \leq 2\|f\|_{V^2}$, for any $-\infty < a < b \leq +\infty$.

A simple application of the duality Lemma 2.4 provides the following linear non-homogeneous estimate.

Lemma 2.5. *Let $1 < p, p' < \infty$ with $\frac{1}{p} + \frac{1}{p'} = 1$, and $-\infty < a < b \leq +\infty$. If $f \in L^1([a, b]; L^2(\mathbb{R}^2))$, then*

$$\left\| \mathbb{1}_{[a,b]}(t) \int_0^t S(t-t')f(t') dt' \right\|_{U_S^p} \lesssim \sup_{\|g\|_{V_S^{p'}}=1} \left| \int_a^b \int_{\mathbb{R}^2} fg \right|.$$

We are now ready to define our resolution space for constructing solutions to (1.15). We denote by Y^s the closure of all functions $u \in C(\mathbb{R} : H^s(\mathbb{R}^2))$ with respect to the norm

$$(2.11) \quad \|u\|_{Y^s} := \left(\sum_N N^{2s} \|P_N u\|_{U_S^2}^2 \right)^{1/2} < \infty.$$

The following proposition is a transference principle in the U_S^p setting.

Proposition 2.6 ([18, Proposition 2.19.(i)]). *Let $\mathcal{L}_0 : L^2(\mathbb{R}^3) \times \cdots \times L^2(\mathbb{R}^3) \rightarrow L_{\text{loc}}^1(\mathbb{R}^3)$ be a n -linear operator. Assume that for some $1 \leq p, q \leq \infty$, we have*

$$\|\mathcal{L}_0(S(\cdot)\phi_1, \dots, S(\cdot)\phi_n)\|_{L_t^p(\mathbb{R} : L_{x,y}^q(\mathbb{R}^2))} \lesssim \prod_{i=1}^n \|\phi_i\|_{L^2}.$$

Then, there exists $\mathcal{L} : U_S^p \times \cdots \times U_S^p \rightarrow L_t^p(\mathbb{R} : L_{x,y}^q(\mathbb{R}^2))$ satisfying

$$\|\mathcal{L}(u_1, \dots, u_n)\|_{L_t^p(\mathbb{R} : L_{x,y}^q(\mathbb{R}^2))} \lesssim \prod_{i=1}^n \|u_i\|_{U_S^p}$$

such that $\mathcal{L}(u_1, \dots, u_n)(t)(x, y) = \mathcal{L}_0(u_1(t), \dots, u_n(t))(x, y)$ almost everywhere.

For the proof of Theorem 1.2, we use Bourgain spaces related to the linear part of (1.15). Thus, for any $s, b \in \mathbb{R}$, let $X^{s,b}$ denote the completion of the Schwartz space $\mathcal{S}(\mathbb{R}^3)$ under the norm

$$(2.12) \quad \|u\|_{X^{s,b}} = \left(\sum_{N,L} N^{2s} L^{2b} \|P_N Q_L u\|_{L^2(\mathbb{R}^3)}^2 \right)^{\frac{1}{2}},$$

and we define the localized-in-time version of these spaces by the norm

$$(2.13) \quad \|u\|_{X_T^{s,b}} = \inf \left\{ \|\tilde{u}\|_{X^{s,b}} : \tilde{u} \in \mathcal{S}(\mathbb{R}^3), \tilde{u}|_{[0,T]} = u \right\}.$$

The basic properties of these spaces that we need here are stated in the following proposition (see e.g. [45]).

Proposition 2.7. *Let $s \in \mathbb{R}$ and $\chi \in C_0^\infty(\mathbb{R})$ be a time cutoff function.*

(i) *For any $b > \frac{1}{2}$,*

$$(2.14) \quad \|\chi(t)S(t)\varphi\|_{X^{s,b}} \lesssim \|\varphi\|_{H^s(\mathbb{R}^2)}.$$

(ii) *For any $0 < \delta < \frac{1}{2}$,*

$$(2.15) \quad \left\| \chi(t) \int_0^t S(t-t')f(t') \right\|_{X^{s, -\frac{1}{2}+\delta}} \lesssim \|f\|_{X^{s, \frac{1}{2}+\delta}}.$$

(iii) *For any $T > 0$ and $-\frac{1}{2} < b' \leq b < \frac{1}{2}$,*

$$(2.16) \quad \|\chi(t/T)u\|_{X^{s,b'}} \lesssim T^{b-b'} \|f\|_{X^{s,b}}.$$

(iv) *For any $T > 0$ and $b > \frac{1}{2}$, we have $X_T^{s,b} \subset C([0, T] : H^s(\mathbb{R}^2))$.*

3. LINEAR AND BILINEAR STRICHARTZ ESTIMATES

3.1. Linear Strichartz estimate. Let Ω be a homogeneous polynomial in \mathbb{R}^2 . Let H_Ω be the Hessian determinant of Ω , i.e.

$$(3.1) \quad H_\Omega(\xi, \mu) = \det \text{Hess } \Omega(\xi, \mu).$$

We denote by $|H_\Omega(D_x, D_y)|^{\frac{1}{8}}$ and $e^{it\Omega(D_x, D_y)}$ the Fourier multipliers associated to $|H_\Omega(\xi, \mu)|^{\frac{1}{8}}$ and $e^{it\Omega(\xi, \mu)}$, i.e.

$$(3.2) \quad \mathcal{F}_{xy} \left(|H_\Omega(D_x, D_y)|^{\frac{1}{8}} \varphi \right) (\xi, \mu) = |H_\Omega(\xi, \mu)|^{\frac{1}{8}} \mathcal{F}_{xy}(\varphi)(\xi, \mu)$$

and

$$(3.3) \quad \mathcal{F}_{xy} \left(e^{it\Omega(D_x, D_y)} \varphi \right) (\xi, \mu) = e^{it\Omega(\xi, \mu)} \mathcal{F}_{xy}(\varphi)(\xi, \mu).$$

As a consequence of Carbery, Kenig and Ziesler in [11], we have the following sharp L^4 -Strichartz estimate for the unitary group $e^{it\Omega(D_x, D_y)}$ (see for example the proof of Corollary 3.4 in [40]).

Theorem 3.1. *For all $\varphi \in L^2(\mathbb{R}^2)$, we have*

$$(3.4) \quad \left\| |H_\Omega(D_x, D_y)|^{\frac{1}{8}} e^{it\Omega(D)} \varphi \right\|_{L_{xyt}^4} \lesssim \|\varphi\|_{L^2},$$

Now, we apply Theorem 3.1 in the case of the unitary group $e^{-t\partial_x(\partial_x^2 - 3\partial_y^2)}$.

Proposition 3.2. (i) *For all $\varphi \in L^2(\mathbb{R}^2)$, we have*

$$(3.5) \quad \left\| D^{\frac{1}{4}} e^{-t\partial_x(\partial_x^2 - 3\partial_y^2)} \varphi \right\|_{L_{x,y,t}^4(\mathbb{R}^3)} \lesssim \|\varphi\|_{L^2(\mathbb{R}^2)}.$$

(ii) *Let N be a dyadic number in $\{2^k : k \in \mathbb{N}^*\}$. For all $u \in V_{-,S}^2$ we have*

$$(3.6) \quad \|P_N u\|_{L^4(\mathbb{R}^3)} \lesssim N^{-\frac{1}{4}} \|P_N u\|_{V_S^2}.$$

Proof. The symbol associated to $e^{-t\partial_x(\partial_x^2 - 3\partial_y^2)}$ is given by $w(\xi, \mu) = \xi^3 - 3\xi\mu^2$. An easy computation shows that

$$(3.7) \quad H_w(\xi, \mu) := \det \text{Hess } w(\xi, \mu) = -36(\xi^2 + \mu^2).$$

Estimate (3.5) follows then as a direct application of Corollary 3.1 and estimate (3.6) follows from Proposition 2.6. \square

Remark 3.1. In the case of the linear Dysthe group $e^{-t\partial_x(\partial_x^2-3\partial_y^2)}$, Theorem 3.1 follows actually from the dispersive estimates proved by Kenig, Ponce and Vega in [24]. Indeed the symbol $w(\xi, \mu) = \xi^3 - 3\xi\mu^2$ clearly satisfies the conditions of Lemma 3.4 in [24] on \mathbb{R}^2 , which provides the sharp decay estimate (see also Ben-Artzi, Koch and Saut in [2])

$$\left| \int_{\mathbb{R}^2} e^{i(tw(\xi, \mu) + x\xi + y\mu)} H_w(\xi, \mu) d\xi d\mu \right| \lesssim |t|^{-1}, \quad \forall (x, y) \in \mathbb{R}^2, \quad \forall t \in \mathbb{R}.$$

Moreover, since $w(\xi, \mu)$ is homogeneous, estimate (3.5) is then deduced from (3.7) arguing as in Theorem 3.1 and Remark (a) in [24].

3.2. Bilinear Strichartz estimates. In this subsection, we state crucial bilinear estimates related to the Dysthe dispersion relation for functions defined on \mathbb{R}^3 .

Proposition 3.3. *Let N_1, N_2, L_1, L_2 be dyadic numbers in $\{2^k : k \in \mathbb{N}\}$ and assume that u_1, u_2 are two functions in $L^2(\mathbb{R}^3)$.*

(i) *We have the basic estimate*

$$(3.8) \quad \begin{aligned} & \| (P_{N_1} Q_{L_1} u_1) (P_{N_2} Q_{L_2} u_2) \|_{L^2(\mathbb{R}^3)} \\ & \lesssim \min\{N_1, N_2\} \min\{L_1, L_2\}^{\frac{1}{2}} \|P_{N_1} Q_{L_1} u_1\|_{L^2(\mathbb{R}^3)} \|P_{N_2} Q_{L_2} u_2\|_{L^2(\mathbb{R}^3)}. \end{aligned}$$

(ii) *If $N_1 \geq 4N_2$, then*

$$(3.9) \quad \| (P_{N_1} Q_{L_1} u_1) (P_{N_2} Q_{L_2} u_2) \|_{L^2(\mathbb{R}^3)} \lesssim N_2^{\frac{1}{2}} N_1^{-1} L_1^{\frac{1}{2}} L_2^{\frac{1}{2}} \|P_{N_1} Q_{L_1} u_1\|_{L^2(\mathbb{R}^3)} \|P_{N_2} Q_{L_2} u_2\|_{L^2(\mathbb{R}^3)}.$$

The proof of Proposition 3.3 is given in Lemma 9 in [1] (see also Proposition 3.6 in [40]). For the sake of completeness we give the argument here. The proof relies on some basic Lemmas stated for example in [41].

Lemma 3.4. *Consider a set $\Lambda \subset \mathbb{R} \times \mathbb{R}$. Let the projection on the μ axis be contained in a set $I \subset \mathbb{R}$. Assume in addition that there exists $C > 0$ such that for any fixed $\mu_0 \in I$, $|\{\xi \in \mathbb{R} : (\xi, \mu_0) \in \Lambda\}| \leq C$. Then, we get that $|\Lambda| \leq C|I|$.*

The second one is a direct consequence of the mean value theorem.

Lemma 3.5. *Let I and J be two intervals on the real line and $f : J \rightarrow \mathbb{R}$ be a smooth function. Then,*

$$(3.10) \quad |\{x \in J : f(x) \in I\}| \leq \frac{|I|}{\inf_{\xi \in J} |f'(\xi)|}.$$

Proof of Proposition 3.3. The Cauchy-Schwarz inequality and Plancherel's identity yield

$$(3.11) \quad \begin{aligned} & \| (P_{N_1} Q_{L_1} u_1) (P_{N_2} Q_{L_2} u_2) \|_{L^2(\mathbb{R}^3)} \\ & = \| (P_{N_1} Q_{L_1} u_1)^\wedge \star (P_{N_2} Q_{L_2} u_2)^\wedge \|_{L^2(\mathbb{R}^3)} \\ & \lesssim \sup_{(\xi, \mu, \tau) \in \mathbb{R}^3} |A_{\xi, \mu, \tau}|^{\frac{1}{2}} \|P_{N_1} Q_{L_1} u_1\|_{L^2(\mathbb{R}^3)} \|P_{N_2} Q_{L_2} u_2\|_{L^2(\mathbb{R}^3)}, \end{aligned}$$

where

$$\begin{aligned} A_{\xi, \mu, \tau} := & \left\{ (\xi_1, \mu_1, \tau_1) \in \mathbb{R}^3 : |(\xi_1, \mu_1)| \in I_{N_1}, |(\xi - \xi_1, \mu - \mu_1)| \in I_{N_2} \right. \\ & \left. |\tau_1 - w(\xi_1, \mu_1)| \in I_{L_1}, |\tau - \tau_1 - w(\xi - \xi_1, \mu - \mu_1)| \in I_{L_2} \right\}. \end{aligned}$$

It remains then to estimate the measure of the set $A_{\xi, \mu, \tau}$ uniformly in $(\xi, \mu, \tau) \in \mathbb{R}^3$.

To obtain (3.8), we use the trivial estimate

$$|A_{\xi, \mu, \tau}| \lesssim \min\{L_1, L_2\} \min\{N_1, N_2\}^2,$$

for all $(\xi, \mu, \tau) \in \mathbb{R}^3$.

Now we turn to the proof of estimate (3.9). First, we get easily from the triangle inequality that

$$(3.12) \quad |A_{\xi, \mu, \tau}| \lesssim \min\{L_1, L_2\} |B_{\xi, \mu, \tau}|,$$

where

$$(3.13) \quad B_{\xi, \mu, \tau} := \left\{ (\xi_1, \mu_1) \in \mathbb{R}^2 : |(\xi_1, \mu_1)| \in I_{N_1}, |(\xi - \xi_1, \mu - \mu_1)| \in I_{N_2} \right. \\ \left. |\tau - w(\xi, \mu) - R(\xi_1, \mu_1, \xi - \xi_1, \mu - \mu_1)| \lesssim \max\{L_1, L_2\} \right\}$$

and where $R(\xi_1, \mu_1, \xi_2, \mu_2)$ is the resonance function defined in (2.2).

In the case where $|\xi_1| \gg |\mu_1|$ or $|\mu_1| \gg |\xi_1|$, we observe from the hypothesis $N_1 \geq 4N_2$ that

$$\left| \frac{\partial R}{\partial \xi_1}(\xi_1, \xi - \xi_1, \mu_1, \mu - \mu_1) \right| = |3(\xi_1^2 - \mu_1^2) - 3((\xi - \xi_1)^2 - (\mu - \mu_1)^2)| \gtrsim N_1^2.$$

Then, if we define $B_{\xi, \mu, \tau}(\mu_1) = \{\xi_1 \in \mathbb{R} : (\xi_1, \mu_1) \in B_{\xi, \mu, \tau}\}$, we deduce applying estimate (3.10) that

$$|B_{\xi, \mu, \tau}(\mu_1)| \lesssim \frac{\max\{L_1, L_2\}}{N_1^2},$$

for all $\mu_1 \in \mathbb{R}$. Thus, it follows from Lemma 3.4 that

$$(3.14) \quad |B_{\xi, \mu, \tau}| \lesssim \frac{N_2}{N_1^2} \max\{L_1, L_2\}.$$

In the case where $|\xi_1| \sim |\mu_1|$, then we use that

$$\left| \frac{\partial R}{\partial \mu_1}(\xi_1, \xi - \xi_1, \mu_1, \mu - \mu_1) \right| = |6\xi_1\mu_1 - 6(\xi - \xi_1)(\mu - \mu_1)| \gtrsim N_1^2.$$

Then, if we define $B_{\xi, \mu, \tau}(\xi_1) = \{\mu_1 \in \mathbb{R} : (\xi_1, \mu_1) \in B_{\xi, \mu, \tau}\}$, we deduce applying estimate (3.10) that

$$|B_{\xi, \mu, \tau}(\xi_1)| \lesssim \frac{\max\{L_1, L_2\}}{N_1^2},$$

for all $\xi_1 \in \mathbb{R}$, so that estimate (3.14) also follows in this case.

Finally, we conclude the proof of the estimate (3.9) by gathering estimates (3.11)–(3.14). \square

Corollary 3.6. *Let N_1, N_2 be dyadic numbers in $\{2^k : k \in \mathbb{N}\}$.*

(i) *If $N_1 \geq 4N_2$, then*

$$(3.15) \quad \|P_{N_1} u_1 P_{N_2} u_2\|_{L^2(\mathbb{R}^3)} \lesssim N_2^{\frac{1}{2}} N_1^{-1} \|P_{N_1} u_1\|_{U_S^2} \|P_{N_2} u_2\|_{U_S^2}, \text{ for any } u_1, u_2 \in U_S^2, \text{ and}$$

$$(3.16) \quad \|P_{N_1} u_1 P_{N_2} u_2\|_{L^2(\mathbb{R}^3)} \lesssim N_2^{\frac{1}{2}} N_1^{-1} \ln^2\left(\frac{N_1}{N_2}\right) \|P_{N_1} u_1\|_{V_S^2} \|P_{N_2} u_2\|_{V_S^2}, \text{ for any } u_1, u_2 \in V_{-,S}^2.$$

(ii) *If $N_1, N_2 \lesssim 1$, we have*

$$(3.17) \quad \|P_{N_1} u_1 P_{N_2} u_2\|_{L^2(\mathbb{R}^3)} \lesssim \|P_{N_1} u_1\|_{V_S^2} \|P_{N_2} u_2\|_{V_S^2}, \text{ for any } u_1, u_2 \in V_{-,S}^2.$$

Proof. Let $\chi \in C_c^\infty(\mathbb{R})$ be a cut-off function satisfying $0 \leq \chi \leq 1$, $\chi|_{[-1,1]} = 1$ and $\text{supp } \chi \in [-2, 2]$. We deduce by using Plancherel's identity that for $\varphi \in L^2(\mathbb{R}^2)$ and any dyadic number $L \geq 1$

$$\mathcal{F}(Q_L \chi S(\cdot) \varphi)(\xi, \mu, \tau) = \phi_L(\tau - w(\xi, \mu)) \widehat{\chi}(\tau - w(\xi, \mu)) \mathcal{F}_{xy}(\varphi)(\xi, \mu).$$

Then by using again Plancherel's identity and taking into account that $\widehat{\chi}$ is rapidly decreasing, it holds that

$$\|Q_L \chi S(\cdot) \varphi\|_{L^2_{x,y,t}} \leq \|\phi_L \widehat{\chi}\|_{L^2(\mathbb{R})} \|\varphi\|_{L^2(\mathbb{R}^2)} \lesssim L^{-4} \|\varphi\|_{L^2(\mathbb{R}^2)}.$$

This ensures by using (3.9) that

$$(3.18) \quad \|(P_{N_1} S(\cdot) \varphi_1)(P_{N_2} S(\cdot) \varphi_2)\|_{L^2([-1,1] \times \mathbb{R}^2)} \\ \leq \|(\chi^{\frac{1}{2}} P_{N_1} S(\cdot) \varphi_1)(\chi^{\frac{1}{2}} P_{N_2} S(\cdot) \varphi_2)\|_{L^2(\mathbb{R}^3)} \lesssim N_2^{\frac{1}{2}} N_1^{-1} \|P_{N_1} \varphi_1\|_{L^2(\mathbb{R}^2)} \|P_{N_2} \varphi_2\|_{L^2(\mathbb{R}^2)}.$$

We can remove the time cut off χ by using the scaling invariance (1.17). Indeed let $u_1 = S(t) \varphi_1$ and $u_2 = S(t) \varphi_2$. Then it follows by using the notation in (1.17) that for any $\lambda > 0$

$$\|P_{N_1} u_1 P_{N_2} u_2\|_{L^2([- \lambda^{-3}, \lambda^3] \times \mathbb{R}^2)} = \lambda^{\frac{1}{2}} \|(P_{N_1} u_1)_\lambda (P_{N_2} u_2)_\lambda\|_{L^2([-1,1] \times \mathbb{R}^2)}.$$

Since $(P_N u)_\lambda = P_{\lambda N}(u)_\lambda$, we conclude from (3.18) that

$$\|P_{N_1} S(\cdot) \varphi_1\|_{L^2([- \lambda^3, \lambda^3] \times \mathbb{R}^2)} \|P_{N_2} S(\cdot) \varphi_2\|_{L^2(\mathbb{R}^2)} \lesssim N_2^{\frac{1}{2}} N_1^{-1} \|P_{N_1} \varphi_1\|_{L^2(\mathbb{R}^2)} \|P_{N_2} \varphi_2\|_{L^2(\mathbb{R}^2)},$$

which implies by letting λ to $+\infty$,

$$(3.19) \quad \|P_{N_1} S(\cdot) \varphi_1\|_{L^2(\mathbb{R}^3)} \|P_{N_2} S(\cdot) \varphi_2\|_{L^2(\mathbb{R}^3)} \lesssim N_2^{\frac{1}{2}} N_1^{-1} \|P_{N_1} \varphi_1\|_{L^2(\mathbb{R}^2)} \|P_{N_2} \varphi_2\|_{L^2(\mathbb{R}^2)}.$$

Estimate (3.15) follows then by applying Proposition 2.6, while (3.16) follows as in [18, Corollary 2.21]. We obtain (3.17) analogously. \square

Corollary 3.7. *For $0 < \theta \leq \frac{1}{4}$, we have*

$$(3.20) \quad \|P_N u\|_{L^4(\mathbb{R}^3)} \lesssim_\theta N^{-\frac{1}{4} + \frac{3\theta}{1+2\theta}} \|P_N u\|_{X^{0, \frac{1}{2}-\theta}}.$$

Proof. By using the transference principle for $X^{s,b}$ -spaces (see e.g. [45, Lemma 2.9]), from (3.5) we have

$$(3.21) \quad \|P_N u\|_{L^4(\mathbb{R}^3)} \lesssim_\theta N^{-\frac{1}{4}} \|P_N u\|_{X^{0, \frac{1}{2}+\theta}}.$$

The estimate (3.20) follows by interpolating (3.21) with the trivial L^4 -estimate that follows from (3.8), namely with

$$\|P_N u\|_{L^4(\mathbb{R}^3)} \lesssim N^{\frac{1}{2}} \|P_N u\|_{X^{0, \frac{1}{4}}}.$$

\square

4. PROOF OF THE MAIN RESULTS

For $T \in (0, +\infty]$, we consider

$$(4.1) \quad I_{T,j}(u_1, u_2, u_3)(t) := \mathbb{1}_{[0,T)}(t) \int_0^t S(t-t') \mathcal{N}_j(u_1, u_2, u_3)(t') dt'.$$

Proposition 4.1. *Let $s \geq 0$ and $u_1, u_2, u_3 \in Y^s$.*

(i) *There exists $C > 0$ such that for all $0 < T < +\infty$ and $1 \leq j \leq 4$, we have*

$$(4.2) \quad \|I_{T,j}(u_1, u_2, u_3)\|_{Y^s} \leq C \|u_1\|_{Y^s} \|u_2\|_{Y^s} \|u_3\|_{Y^s}.$$

(ii) *The estimate (4.2) also holds for $T = +\infty$ and for any $1 \leq j \leq 4$, we have*

$$(4.3) \quad \|I_{T,j}(u_1, u_2, u_3) - I_{+\infty,j}(u_1, u_2, u_3)\|_{Y^s} \rightarrow 0 \text{ as } T \rightarrow +\infty$$

and the limit

$$(4.4) \quad \lim_{t \rightarrow +\infty} S(-t) I_{+\infty,j}(u_1, u_2, u_3)(t)$$

exists in $L^2(\mathbb{R}^2)$.

Proof. (i) Note that it suffices to prove the estimates for $s = 0$ and by Littlewood-Paley decomposition it suffices to prove the following

$$(4.5) \quad \left(\sum_{N_4} \left\| \sum_{N_1, N_2, N_3} P_{N_4} \mathcal{D}_j(P_{N_1} u_1, P_{N_2} u_2, P_{N_3} u_3) \right\|_{U_S^2}^2 \right)^{\frac{1}{2}} \lesssim \|u_1\|_{Y^0} \|u_2\|_{Y^0} \|u_3\|_{Y^0},$$

where

$$(4.6) \quad \mathcal{D}_j(P_{N_1} u_1, P_{N_2} u_2, P_{N_3} u_3) := \int_0^t S(t-t') \mathcal{N}_j(P_{N_1} u_1, P_{N_2} u_2, P_{N_3} u_3)(t') dt',$$

with $u_i := \mathbb{1}_{[0,T)} u_i$, $1 \leq i \leq 4$. We treat the contribution of the worst parts of the nonlinearities \mathcal{N}_j 's, namely when there is a derivative and it falls on the largest frequency factor; the nonlinearity involving the Riesz transform \mathcal{R}_x (which is bounded on $L^2(\mathbb{R}^2)$) does not pose a further difficulty.

Thus, in what follows we focus on the nonlinear term

$$(4.7) \quad \mathcal{N}(P_{N_1} u_1, P_{N_2} u_2, P_{N_3} u_3) := (P_{N_1} u_1)(P_{N_2} u_2) \partial_x (P_{N_3} u_3)$$

with $N_1 \leq N_2 \leq N_3$. Thus, by Lemma 2.5,⁴ it remains to estimate the following term

$$(4.8) \quad \left(\sum_{N_4} \sup_{\|u_4\|_{V_S^2}=1} |I_{N_1, \dots, N_4}|^2 \right)^{\frac{1}{2}},$$

where

$$I_{N_1, \dots, N_4} := \sum_{N_1 \leq N_2 \leq N_3} N_3 \int_{\mathbb{R}^3} (P_{N_1} u_1)(P_{N_2} u_2)(P_{N_3} u_3)(P_{N_4} u_4) dx dy dt,$$

and where we assume without loss of generality that the Fourier transforms $\mathcal{F}_{xy}(u_i)$, $1 \leq i \leq 4$, are non-negative.

We recall that due to the frequency hyperplane, the largest two dyadic numbers are comparable and thus we distinguish the following cases.

Case 1: $N_1 \leq N_2 \ll N_3 \sim N_4$. We apply the bilinear estimates (3.15) and (3.16) together with (2.10). For fixed N_4 , we have

$$\begin{aligned} |I_{N_1, \dots, N_4}| &\lesssim \sum_{N_1 \leq N_2 \ll N_3 \sim N_4} N_3 \|P_{N_1} u_1 P_{N_3} u_3\|_{L_{x,y,t}^2} \|P_{N_2} u_2 P_{N_4} u_4\|_{L_{x,y,t}^2} \\ &\lesssim \|u_1\|_{Y^0} \|u_2\|_{Y^0} \sum_{N_1 \leq N_2 \ll N_3 \sim N_4} \frac{N_1^{\frac{1}{2}} N_2^{\frac{1}{2}-\varepsilon}}{N_4^{1-\varepsilon}} \|P_{N_3} u_3\|_{U_S^2} \|P_{N_4} u_4\|_{V_S^2} \\ &\lesssim \|u_1\|_{Y^0} \|u_2\|_{Y^0} \sum_{N_3 \sim N_4} \|P_{N_3} u_3\|_{U_S^2}, \end{aligned}$$

where in the last step we used $\|P_{N_4} u_4\|_{V_S^2} \leq 1$. Then, by the Cauchy-Schwarz inequality we have

$$(4.8) \lesssim \|u_1\|_{Y^0} \|u_2\|_{Y^0} \left(\sum_{N_4} \left(\sum_{N_3 \sim N_4} \|P_{N_3} u_3\|_{U_S^2} \right)^2 \right)^{\frac{1}{2}} \lesssim \|u_1\|_{Y^0} \|u_2\|_{Y^0} \|u_3\|_{Y^0}.$$

Case 2: $N_1, N_4 \ll N_2 \sim N_3$. By arguing similarly to Case 1, for fixed N_4 we now have

$$\begin{aligned} |I_{N_1, \dots, N_4}| &\lesssim \|u_1\|_{Y^0} \sum_{N_1 \ll N_2 \sim N_3} \frac{N_1^{\frac{1}{2}} N_4^{\frac{1}{2}-\varepsilon}}{N_2^{1-\varepsilon}} \|P_{N_2} u_2\|_{U_S^2} \|P_{N_3} u_3\|_{U_S^2} \|P_{N_4} u_4\|_{V_S^2} \\ &\lesssim \|u_1\|_{Y^0} \sum_{N_2 \sim N_3} \left(\frac{N_4}{N_2} \right)^{\frac{1}{2}-\varepsilon} \|P_{N_2} u_2\|_{U_S^2} \|P_{N_3} u_3\|_{U_S^2}. \end{aligned}$$

By Minkowski and Cauchy-Schwarz inequalities we then get

$$(4.8) \lesssim \|u_1\|_{Y^0} \sum_{N_2 \sim N_3} \left(\sum_{N_4 \ll N_2} \left(\frac{N_4}{N_2} \right)^{1-2\varepsilon} \right)^{\frac{1}{2}} \|P_{N_2} u_2\|_{U_S^2} \|P_{N_3} u_3\|_{U_S^2} \lesssim \|u_1\|_{Y^0} \|u_2\|_{Y^0} \|u_3\|_{Y^0}.$$

Case 3: $N_1 \ll N_2 \sim N_3 \sim N_4$. We apply the bilinear estimate of Corollary 3.6 and the L^4 -Strichartz estimate (3.6) twice. Thus, arguing similarly to Case 1, we obtain

$$\begin{aligned} (4.8) &\lesssim \left(\sum_{N_4} \left(\sum_{N_1 \ll N_2 \sim N_3 \sim N_4} N_3 \|P_{N_1} u_1 P_{N_4} u_4\|_{L_{x,y,t}^2} \|P_{N_2} u_2\|_{L_{x,y,t}^4} \|P_{N_3} u_3\|_{L_{x,y,t}^4} \right)^2 \right)^{\frac{1}{2}} \\ &\lesssim \|u_1\|_{Y^0} \|u_2\|_{Y^0} \|u_3\|_{Y^0}. \end{aligned}$$

Case 4: $N_4 \ll N_1 \sim N_2 \sim N_3$. In this case the desired estimate follows as in Case 3 via an application of Minkowski's inequality (similarly to the last step of Case 2).

⁴The estimates for $T = +\infty$ need to be derived separately as in this case we cannot apply Lemma 2.5.

Case 5: $1 \ll N_1 \sim N_2 \sim N_3 \sim N_4$. In this case we use the L^4 -Strichartz estimate (3.6) together with (2.10) in order to get

$$(4.8) \lesssim \left(\sum_{N_4} \left(\sum_{N_1 \ll N_2 \sim N_3 \sim N_4} N_3 \|P_{N_1} u_1\|_{L^4_{x,y,t}} \|P_{N_2} u_2\|_{L^4_{x,y,t}} \|P_{N_3} u_3\|_{L^4_{x,y,t}} \|P_{N_4} u_4\|_{L^4_{x,y,t}} \right)^2 \right)^{\frac{1}{2}} \\ \lesssim \|u_1\|_{Y^0} \|u_2\|_{Y^0} \|u_3\|_{Y^0}.$$

Case 6: $N_1 \sim N_2 \sim N_3 \sim N_4 \lesssim 1$. We simply use the bilinear estimate (3.17) twice together with (2.10) and we get (4.5).

Part (ii) follows analogously to the proof of [18, Corollary 3.4]. \square

Proof of Theorem 1.1. It uses a standard argument via contraction mapping principle applied to the simplified equation (1.15) after which we undo the transformation (2.6).

Let $v_0 \in L^2(\mathbb{R}^2)$ such that $\|v_0\|_{L^2(\mathbb{R}^2)} < \delta$, with δ to be chosen later. First, we construct a solution $u(t)$ to (1.15) for $t \in (0, \infty)$ with initial data $u_0(x, y) = e^{ia_2 x} v_0(x, \sqrt{2}y)$, for some $a_2 \in \mathbb{R}$ (see Lemma 2.1 and Remark 2.1). Thus, we consider $\Phi(u)$ given by

$$(4.9) \quad \Phi(u)(t) = \mathbb{1}_{[0,\infty)}(t) S(t) u_0 + \sum_{j=1}^4 c_j \mathbb{1}_{[0,\infty)}(t) \int_0^t S(t-t') \mathcal{N}_j(u, u, u)(t') dt'.$$

Since $\mathbb{1}_{[0,\infty)}(t) u_0$ is a U^2 -atom, one easily checks that $\|\mathbb{1}_{[0,\infty)}(t) S(t) u_0\|_{Y^0} \sim \|u_0\|_{L^2}$ and thus by Proposition 4.1, we get

$$\|\Phi(u)\|_{Y^0} \leq C\delta + C\|u\|_{Y^0}^3,$$

for some $C > 0$. By setting $r := 2C\delta$, we have that Φ maps $B_r := \{u \in Y^0 : \|u\|_{Y^0} \leq r\}$ into itself, provided that $8C^3\delta^2 \leq 1$. Similarly, by using telescoping sums, there exists $\tilde{C} > 0$ such that

$$\|\Phi(u_1) - \Phi(u_2)\|_{Y^0} \leq \tilde{C} \|u_1 - u_2\|_{Y^0} (\|u_1\|_{Y^0}^2 + \|u_2\|_{Y^0}^2).$$

By ensuring that $8C^2\tilde{C}\delta^2 < 1$, we have that Φ is a contraction on B_r , and thus it has a unique fixed point in B_r which solves (1.15) for $t \in [0, +\infty)$.

For the scattering claim, we use Proposition 4.1 (ii) and take

$$u_0^+ := u_0 + \sum_{j=1}^4 c_j \lim_{t \rightarrow +\infty} S(-t) I_{+\infty,j}(u, u, u)(t) \in L^2(\mathbb{R}^2).$$

Since the embedding $Y^0 \subset L^\infty(\mathbb{R} : L^2(\mathbb{R}^2))$ is continuous, we get that for any $T > t$,

$$\|u(t) - S(t) u_0^+\|_{L^2(\mathbb{R}^2)} \leq \sum_{j=1}^4 |c_j| \left(\|I_{T,j}(u, u, u) - I_{+\infty,j}(u, u, u)\|_{Y^0} \right. \\ \left. + \|S(-t) I_{+\infty,j}(u, u, u)(t) - \lim_{\tau \rightarrow +\infty} S(-\tau) I_{+\infty,j}(u, u, u)(\tau)\|_{L^2(\mathbb{R}^2)} \right)$$

and by (4.3) and (4.4), we then get $\|u(t) - S(t) u_0^+\|_{L^2(\mathbb{R}^2)} \rightarrow 0$ as $t \rightarrow +\infty$.

Second, we construct the solution $u_{\text{neg}}(t)$ to (1.15) for $t \in (-\infty, 0)$ and the scattering data u_0^- , by running the analogous argument for the equation obtained from (1.15) after the change of variable

$$(4.10) \quad u(t, x, y) \mapsto \overline{u(-t, -x, -y)} =: \mathcal{I}(u),$$

with initial data $\overline{u_0(-x, -y)}$, and then undoing the transformation (4.10). Since the transformations (2.6) and (4.10) commute, the solution v to the original Dysthe equation (1.12) is given by

$$(4.11) \quad v = \mathcal{T}^{-1}(u + u_{\text{neg}}) \in \mathcal{Y} := \{\mathbb{1}_{[0,\infty)} v_1 + \mathbb{1}_{(-\infty,0)} v_2 : \mathcal{T}(v_1) \in Y^0 \text{ and } \mathcal{T}(\mathcal{I}(v_2)) \in Y^0\},$$

and the corresponding scattering data are $v_0^\pm(x, y) := e^{-ia_2 x} u_0^\pm(x, \frac{y}{\sqrt{2}})$. \square

In order to obtain the local well-posedness result for arbitrarily large initial data (i.e. Theorem 1.2), we prove the trilinear estimates in $X^{s,b}$ spaces.

Proposition 4.2. *Let $s > 0$ and $\nu > 0$ sufficiently small. For all $1 \leq j \leq 4$, we have*

$$(4.12) \quad \|\mathcal{N}_j(u_1, u_2, u_3)\|_{X^{s, -\frac{1}{2}+2\nu}} \lesssim \|u_1\|_{X^{s, \frac{1}{2}+\nu}} \|u_2\|_{X^{s, \frac{1}{2}+\nu}} \|u_3\|_{X^{s, \frac{1}{2}+\nu}}.$$

Proof. As in the proof of Proposition 4.1, we focus on the contribution of the worst parts of the nonlinearities and thus we prove the estimate for the nonlinear term

$$\mathcal{N}(P_{N_1}u_1, P_{N_2}u_2, P_{N_3}u_3) := (P_{N_1}u_1)(P_{N_2}u_2)\partial_x(P_{N_3}u_3)$$

with $N_1 \leq N_2 \leq N_3$. By duality and Littlewood-Paley decomposition it suffices to prove the following

$$(4.13) \quad \left(\sum_{\substack{N_1 \leq N_2 \leq N_3 \\ N_4 \\ L_1, L_2, L_3, L_4}} \Gamma_{N_1, \dots, N_4}^2 \Lambda_{L_1, \dots, L_4}^2 |I_{N_1, \dots, N_4}^{L_1, \dots, L_4}(v_1, v_2, v_3, v_4)|^2 \right)^{\frac{1}{2}} \lesssim \prod_{i=1}^4 \|v_i\|_{L^2(\mathbb{R}^3)},$$

where

$$(4.14) \quad \Gamma_{N_1, \dots, N_4} := N_1^{-s} N_2^{-s} N_3^{1-s} N_4^s,$$

$$(4.15) \quad \Lambda_{L_1, \dots, L_4} := L_1^{-\frac{1}{2}-\nu} L_2^{-\frac{1}{2}-\nu} L_3^{-\frac{1}{2}-\nu} L_4^{-\frac{1}{2}+2\nu},$$

$$(4.16) \quad I_{N_1, \dots, N_4}^{L_1, \dots, L_4}(v_1, v_2, v_3, v_4) := \int_{\mathbb{R}^3} (P_{N_1}Q_{L_1}v_1)(P_{N_2}Q_{L_2}v_2)(P_{N_3}Q_{L_3}v_3)(P_{N_4}Q_{L_4}v_4) dx dy dt,$$

and where we assume without loss of generality that the Fourier transforms $\mathcal{F}_{xy}(v_i)$, $1 \leq i \leq 4$, are non-negative.

We discuss the same cases as in the proof of Proposition 4.1.

Case 1: $N_1 \leq N_2 \ll N_3 \sim N_4$. As in the proof of [40, Proposition 4.1], we interpolate between the two bilinear Strichartz estimates (3.8) and (3.9), and so we have

$$(4.17) \quad \begin{aligned} & \|(P_{N_2}Q_{L_2}u_2)(P_{N_4}Q_{L_4}u_4)\|_{L^2(\mathbb{R}^3)} \\ & \lesssim \frac{N_2^{\frac{1+\theta}{2}}}{N_4^{1-\theta}} \min\{L_2, L_4\}^{\frac{1}{2}} \max\{L_2, L_4\}^{\frac{1-\theta}{2}} \|P_{N_2}Q_{L_2}u_2\|_{L^2(\mathbb{R}^3)} \|P_{N_4}Q_{L_4}u_4\|_{L^2(\mathbb{R}^3)}. \end{aligned}$$

We now assume that $L_2 \leq L_4$ as the case $L_4 < L_2$ is easier. Then, by the Cauchy-Schwarz inequality, (4.17), and (3.9) for $(P_{N_1}Q_{L_1}u_1)(P_{N_3}Q_{L_3}u_3)$, we get

$$(4.18) \quad \begin{aligned} & \Gamma_{N_1, \dots, N_4} \Lambda_{L_1, \dots, L_4} |I_{N_1, \dots, N_4}^{L_1, \dots, L_4}(v_1, v_2, v_3, v_4)| \\ & \lesssim \frac{N_1^{\frac{1}{2}-s} N_2^{\frac{1+\theta}{2}-s}}{N_4^{1-\theta}} L_1^{-\nu} L_2^{-\nu} L_3^{-\nu} L_4^{2\nu-\frac{\theta}{2}} \prod_{i=1}^4 \|v_i\|_{L^2(\mathbb{R}^3)}. \end{aligned}$$

By choosing θ and ν such that $\frac{1+\theta}{2} - s + \frac{1}{2} - s < 1 - \theta$ and $2\nu - \frac{\theta}{2} < 0$, the estimate (4.13) follows immediately.

Case 2: $N_1, N_4 \ll N_2 \sim N_3$. The estimate (4.13) follows as in Case 1 above (by interchanging N_2 and N_4 in (4.18)) via Minkowski's inequality for the summation in N_4 (see also Case 2 in the proof of Proposition 4.1).

For the remaining cases, we follow essentially the same arguments as in the proof of Proposition 4.1.

Case 3: $N_1 \ll N_2 \sim N_3 \sim N_4$. We apply the bilinear Strichartz estimate (3.9) and the L^4 -Strichartz estimate (3.20) twice (with $\theta = \frac{\nu}{2}$ and $\theta = 3\nu$) and we get

$$\Gamma_{N_1, \dots, N_4} \Lambda_{L_1, \dots, L_4} |I_{N_1, \dots, N_4}^{L_1, \dots, L_4}(v_1, v_2, v_3, v_4)| \lesssim N_1^{-s} N_2^{-s+\frac{3\nu}{2(1+\nu)}+\frac{9\nu}{1+6\nu}} \prod_{i=1}^4 L_i^{-\nu} \|v_i\|_{L^2(\mathbb{R}^3)}.$$

By choosing $\nu > 0$ such that $\frac{3\nu}{2(1+\nu)} + \frac{9\nu}{1+6\nu} < s$ we ensure summability over $N_2 \sim N_3 \sim N_4$ and thus we get (4.13).

Case 4: $N_4 \ll N_1 \sim N_2 \sim N_3$. We use the interpolated bilinear Strichartz estimate as in Case 1 (namely (4.17) with $N_2^{\frac{1+\theta}{2}}/N_4^{1-\theta}$ replaced by $N_4^{\frac{1+\theta}{2}}/N_2^{1-\theta}$) and the L^4 -Strichartz estimate (3.20). We have

$$\Gamma_{N_1, \dots, N_4} \Lambda_{L_1, \dots, L_4} |I_{N_1, \dots, N_4}^{L_1, \dots, L_4}(v_1, v_2, v_3, v_4)| \lesssim N_1^{\theta-3s} N_4^{\frac{\theta}{2}} L_1^{-\nu} L_2^{-\nu} L_3^{-\nu} L_4^{2\nu-\frac{\theta}{2}} \prod_{i=1}^4 \|v_i\|_{L^2(\mathbb{R}^3)}$$

and we choose θ and ν such that $0 < 4\nu < \theta < 2s$.

Case 5: $1 \ll N_1 \sim N_2 \sim N_3 \sim N_4$. We use the L^4 -Strichartz estimate (3.20) to obtain

$$\Gamma_{N_1, \dots, N_4} \Lambda_{L_1, \dots, L_4} |I_{N_1, \dots, N_4}^{L_1, \dots, L_4}(v_1, v_2, v_3, v_4)| \lesssim N_1^{-2s+\frac{9\nu}{2(1+\nu)}+\frac{9\nu}{1+6\nu}} \prod_{i=1}^4 L_i^{-\nu} \|v_i\|_{L^2(\mathbb{R}^3)}$$

and we choose $\nu > 0$ such that $\frac{9\nu}{2(1+\nu)} + \frac{9\nu}{1+6\nu} < 2s$.

Case 6: $N_1 \sim N_2 \sim N_3 \sim N_4 \lesssim 1$. We simply use the bilinear estimate (3.8) twice to obtain

$$\Gamma_{N_1, \dots, N_4} \Lambda_{L_1, \dots, L_4} |I_{N_1, \dots, N_4}^{L_1, \dots, L_4}(v_1, v_2, v_3, v_4)| \lesssim L_1^{-\nu} L_2^{-\nu} L_3^{-\nu} L_4^{-\frac{1}{2}+2\nu} \prod_{i=1}^4 \|v_i\|_{L^2(\mathbb{R}^3)}$$

and we choose $0 < \nu < \frac{1}{4}$. □

Proof of Theorem 1.2. It uses a standard argument via contraction mapping principle and Proposition 2.7 applied to

$$(4.19) \quad \Phi(u)(t) = \chi(t)S(t)u_0 + \sum_{j=1}^4 c_j \chi(t) \int_0^t \chi(t'/T) S(t-t') \mathcal{N}_j(u, u, u)(t') dt'$$

in a ball of $X_T^{s, \frac{1}{2}+\nu}$, by choosing $T = C \|u_0\|_{H^s(\mathbb{R}^2)}^{-\frac{2}{\nu}}$, $\nu > 0$ sufficiently small, some constant $C > 0$, and some $\chi \in C_0^\infty(\mathbb{R})$ with $\chi \equiv 1$ on $[0, 1]$. Lastly, we note that the solution v corresponding to the original equation lies in the space $\mathcal{X}_T^{s, b}$ defined analogous to $X_T^{s, b}$, using the non-homogeneous symbol ω (defined by (2.7)) instead of the homogeneous version w . □

5. FINAL COMMENTS

Some interesting issues remain open for the Cauchy problem of Dysthe type equations, for instance the possible finite time blow-up of large local solutions.

Other questions concern the Dysthe equations as a water waves model and are summarized in [31] Section 8.5.4, for instance the comparison of the Dysthe equation with other models with improved dispersion.

More precisely, as recalled in Lannes [31], Section 8.5.4, the Dysthe equation contains both higher nonlinear and dispersive terms neglected in the standard cubic NLS equation, while the full dispersion equation derived in [31] Section 8.5.3 contains only higher order dispersive terms (at infinite order actually). A comparison between these models (and between a possible full-dispersion Dysthe equation) would bring some insight to the relative importance of dispersive and nonlinear effects in situations where the standard NLS approximation compares poorly with the experiments (see for instance [39]). We plan to come back to those issues in a subsequent paper.

Acknowledgements. D.P. and R.M. were supported by a Trond Mohn Foundation grant. J.-C. S. was partially supported by the ANR project ANuI (ANR-17-CE40-0035-02). The authors would like to thank Karsten Trulsen for several interesting comments which clarified for us the derivation of the models and for relevant additional references and Felipe Linares for correcting several typos and pointing out that the linear Strichartz estimate associated to the homogeneous Dysthe symbol was already proved in [24].

REFERENCES

- [1] Y. ANGELOPOULOS, *Well-posedness and ill-posedness results for the Novikov-Veselov equation*, Comm. Pure Appl. Anal., **15** (2016), 727–760.
- [2] M. BEN-ARTZI, H. KOCH AND J.-C. SAUT, *Dispersion estimates for third-order equations in two dimensions*, Comm. Part. Diff. Eq., **28** (2003), 1943–1974.
- [3] A. DE BOUARD, *Analytic solutions to nonelliptic nonlinear Schrödinger equations*, J. Diff. Equ, **104** (1) (1993), 196–213.
- [4] U. BRINCH-NIELSEN AND I.G. JONSSON, *Fourth order evolution equations and stability analysis for Stokes waves on arbitrary water depth*, Wave Motion, **8** (1986), 455–472.
- [5] T. CAZENAVE, F. B. WEISSLER, *The Cauchy problem for the critical nonlinear Schrödinger equation in H^s* , Nonlinear Anal. Theory Methods Appl., **14** (10) (1990), 807–36.
- [6] H. CHIHARA, *Third order dispersive equations related to deep water waves*, preprint (2004), arXiv:math/0404005.
- [7] W.CRAIG, P. GUYENNE, D. NICHOLS AND C. SULEM, *A Hamiltonian approach to nonlinear modulation of surface water waves*, Wave Motion, **47** (2010), 552–563.
- [8] W. CRAIG, P. GUYENNE AND C. SULEM, *Hamiltonian higher-order nonlinear Schrödinger equations for broader-banded waves on deep water*, European J. Mechanics B/Fluids, **32** (2010), 22–31.
- [9] W. CRAIG, P. GUYENNE AND C. SULEM, *Normal form transformations and Dysthe’s equation for the nonlinear modulation of deep water waves*, Water Waves (2020), in press.
- [10] F. DIAS AND C. KHARIF, *Nonlinear gravity and capillary -gravity waves*, Annu. Rev. Fluid Mech., **31** (1999), 301–346.
- [11] A. CARBERY, C. E. KENIG AND S. ZIESLER, *Restriction for homogeneous polynomial surfaces in \mathbb{R}^3* , Trans. Amer. Math. Soc., **365** (2013), 2367–2407.
- [12] K. B. DYSTHE, *Note on a modification to the nonlinear Schrödinger equation for application to deep water waves*, Proc. R. Soc. Lond. A., **369** (1979), 105–114.
- [13] K. B. DYSTHE AND K. TRULSEN, *The evolution of an evolution equation*, Frontiers of Nonlinear Physics, volume **II**, 125–138.
- [14] F. FEDELE AND D. DUTYKH, *Hamiltonian form and solitary waves of the spatial Dysthe equation*, JETP Letters, **94** (2011), 840–844.
- [15] O. GRAMSTAD AND K. TRULSEN, *Hamiltonian form of the modified nonlinear Schrödinger equation for gravity waves on arbitrary depth*, J. Fluid Mech. **670** (2011), 404–426.
- [16] R. GRANDE, K. KURIANSKI AND G. STAFFILANI, *On the nonlinear Dysthe equation*, preprint (2020), arXiv:2006.13392.
- [17] A. GRÜNROCK AND S. HERR, *The Fourier restriction norm method for the Zakharov-Kuznetsov equation*, Disc. Contin. Dyn. Syst. Ser. A, **34** (2014), 2061–2068.
- [18] M. HADAC, S. HERR AND H. KOCH, *Well-posedness and scattering for the KP-II equation in a critical space*, Ann. Inst. H. Poincaré Anal. Non Lin., **26** (2009), 917–941. ERRATUM Ann. Inst. H. Poincaré Anal. Non Lin., **27** (2010), 971–972.
- [19] D. HENDERSON, D.H. PEREGRINE AND J.W. DOLD, *Unsteady water wave modulations: Fully nonlinear solutions and comparison with the nonlinear Schrödinger equation*, Wave Motion, **29** (1999), 341–361.
- [20] S. J. HOGAN, *The fourth-order evolution equation for deep-water gravity-capillary waves.*, Proc. R. Soc. Lond. A, **402** (1985), 359–372.
- [21] P. A. E. M. JANSSEN, *Fourth-order envelope equation for deep-water waves*, J. Fluid Mech. **126** (1983), 1–11.
- [22] H.-H. HWUNG, W.-S. CHIANG AND S.-C. HSIAO, *Observations on the evolution of wave modulation*, Proc. Royal Society A, **463** (2007), 85–112.
- [23] A. KAZEYKINA AND C. MUÑOZ, *Dispersive estimates for rational symbols and local well-posedness of the nonzero energy NV equation*, J. Funct. Anal., **270** (2016), no. 5, 1744–1791.
- [24] C. E. KENIG, G. PONCE AND L. VEGA, *Oscillatory integrals and regularity of dispersive equations*, Indiana Math. Univ. J., **40** (1991), 33–69.
- [25] C. E. KENIG, G. PONCE AND L. VEGA, *Well-posedness and scattering results for the generalized Korteweg- de Vries equation via the contraction principle*, Comm. Pure Appl. Math., **46** (1993), 527–620.
- [26] S. KINOSHITA, *Well-posedness for the Cauchy problem of the modified Zakharov-Kuznetsov equation*, preprint (2019), arXiv:1911.13265.
- [27] H. KOCH AND J.-C. SAUT, *Local smoothing and local solvability for third order dispersive equations*, SIAM J. Math. Anal., **38** (2007), 1528–1541.
- [28] H. KOCH AND D. TATARU, *Dispersive estimates for principally normal pseudodifferential operators*, Comm. Pure Appl. Math., **58** (2005), 217–284.
- [29] H. KOCH AND D. TATARU, *A priori bounds for the 1D cubic NLS in negative Sobolev spaces*, Int. Math. Res. Not., (2007), article ID mm053, 36 pages.
- [30] V.P. KRASITSKII, *On reduced equations in the Hamiltonian theory of weakly nonlinear surface waves*, J. Fluid Mech. **272** (1994), 1–20.
- [31] D. LANNES, *Water waves : mathematical theory and asymptotics*, Mathematical Surveys and Monographs, vol 188 (2013), AMS, Providence.

- [32] F. LINARES AND A. PASTOR, *Well-posedness for the two-dimensional modified Zakharov-Kuznetsov equation*, SIAM J. Math. Anal., **41** (2009), 1323–1339.
- [33] F. LINARES, D. PILOD AND G. PONCE, *Well-posedness for a higher-order Benjamin-Ono equation*, J. Diff. Eq., **250** (2011) 450–475.
- [34] E. LO AND CHIANG C. MEI, *A numerical study of water-wave modulation based on a higher-order nonlinear Schrödinger equation*, J. Fluid Mech., **150** (1985), 395–416.
- [35] E. LO AND CHIANG C. MEI, *Slow evolution of nonlinear deep water waves in two horizontal dimensions: A numerical approach*, Wave Motion, **9** (1987), 245–259.
- [36] M.S. LONGUET-HIGGINS, *The instabilities of gravity waves of finite amplitude in deep water. I. Superharmonics*, Proc. Royal Society A, **360** (1978), 47–488.
- [37] M.S. LONGUET-HIGGINS, *The instabilities of gravity waves of finite amplitude in deep water. II. Subharmonics*, Proc. Royal Society A, **360** (1978), 489–505.
- [38] J.W. MCLEAN, Y.C. MA, D.U. MARTIN, P.G. SAFFMAN AND H.C. YUEN, *Three-dimensional instability of finite-amplitude water waves*, Phys. Rev. Lett. **46** (1981), 817–820.
- [39] D.U. MARTIN AND H.C. YUEN, *Quasi-recurring energy leakage in the two-space dimensional nonlinear Schrödinger equation*, Phys. Fluids, **13**(1980), 881–883.
- [40] L. MOLINET AND D. PILOD, *Bilinear Strichartz estimates for the Zakharov-Kuznetsov equation and applications*, Ann. Inst. H. Poincaré Anal. Non Lin., **32** (2015), 347–371.
- [41] L. MOLINET, J.-C. SAUT AND N. TZVETKOV, *Global well-posedness for the KP-II equation on the background of non localized solution*, Ann. Inst. H. Poincaré Anal. Non Lin., **28** (2011), no. 5, 653–676.
- [42] F. RIBAUD AND S. VENTO, *A note on the Cauchy problem for the 2D generalized Zakharov-Kuznetsov equations*, C. R. Math. Acad. Sci. Paris, **350** (2012), 499–503.
- [43] M. STIASSNIE, *Note on the modified nonlinear equation for deep water waves*, Wave Motion, **6** (1984), 43–433.
- [44] M. STIASSNIE AND L. SHEMER, *On modifications of the Zakharov equation for surface gravity waves*, J. Fluid Mech., **143** (1984), 47–67.
- [45] T. TAO, *Nonlinear Dispersive Equations: Local and Global Analysis*, no. 106 in CBMS, Amer. Math. Soc., 2007.
- [46] K. TRULSEN AND K. B. DYSTHE, *Frequency downshift in three-dimensional wave trains in a deep basin*, J. Fluid Mech. **352** (1997), 359–373.
- [47] K. TRULSEN AND K. B. DYSTHE, *A modified nonlinear Schrödinger equation for broader bandwidth gravity waves on deep water*, Wave Motion, **24** (1996), 281–289.
- [48] K. TRULSEN, I. KLIAKHANDLER, K. B. DYSTHE AND M. G. VELARDE, *On weakly nonlinear modulation of waves on deep water*, Phys. Fluids **12** (2000), 2432–2437.
- [49] V.E. ZAKHAROV, *Stability of periodic waves of finite amplitude on the surface of a deep fluid*, J. Appl. Mech. Tech. Phys., **9** (1968), 190–194.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BERGEN, POSTBOX 7800, 5020 BERGEN, NORWAY
E-mail address: Razvan.Mosincat@uib.no

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BERGEN, POSTBOX 7800, 5020 BERGEN, NORWAY
E-mail address: Didier.Pilod@uib.no

LABORATOIRE DE MATHÉMATIQUES, UMR 8628, UNIVERSITÉ PARIS-SACLAY AND CNRS, 91405 ORSAY, FRANCE
E-mail address: jean-claude.saut@universite-paris-saclay.fr