

On Second order correctness of Bootstrap in Logistic Regression

Debraj Das¹ and Priyam Das²

¹ Indian Institute of Technology, Kanpur, India
rajdas@iitk.ac.in

² Department of Biomedical Informatics, Harvard Medical School, Boston, USA
priyam.das@hms.harvard.edu

Abstract. In the fields of clinical trials, biomedical surveys, marketing, banking, with dichotomous response variable, the logistic regression is considered as an alternative convenient approach to linear regression. In this paper, we develop a novel bootstrap technique based on perturbation resampling method for approximating the distribution of the maximum likelihood estimator (MLE) of the regression parameter vector. We establish second order correctness of the proposed bootstrap method after proper studentization and smoothing. It is shown that inferences drawn based on the proposed bootstrap method are more accurate compared to that based on asymptotic normality. The main challenge in establishing second order correctness remains in the fact that the response variable being binary, the resulting MLE has a lattice structure. We show the direct bootstrapping approach fails even after studentization. We adopt smoothing technique developed in Lahiri (1993) to ensure that the smoothed studentized version of the MLE has a density. Similar smoothing strategy is employed to the bootstrap version also to achieve second order correct approximation.

Keywords: Logistic Regression, MLE, SOC, Lattice, Smoothing, Perturbation Bootstrap

1 Introduction

Logistic regression is one of the most widely used regression techniques when the response variable is binary. Use of the logistic or the ‘logit’ function as a statistical tool dates back to Berkson (1944) and later it was popularized in the field of regression by D. R. Cox in his seminal paper Cox (1958). Suppose y denotes binary response variable which can take values 0 or 1 and the value of y depends on the p independent variables $\mathbf{x} = (x_1, \dots, x_p)'$. Instead of capturing this dependence by modelling y directly, in logistic regression, log-odds corresponding to the success of y , denoted by $p(\mathbf{x}) = P(y = 1)$, is modeled as a linear function of x_1, \dots, x_p . The odds ratio for the event $\{Y = 1\}$ is given by $odd(\mathbf{x}) = \frac{p(\mathbf{x})}{1 - p(\mathbf{x})}$. The logistic regression model is given by

$$\text{logit}(p(\mathbf{x})) = \log \left[\frac{p(\mathbf{x})}{1 - p(\mathbf{x})} \right] = \mathbf{x}'\boldsymbol{\beta}, \quad (1.1)$$

where $\boldsymbol{\beta} = (\beta_1, \dots, \beta_p)$ is the p -dimensional vector of regression parameters. It is routine to consider the maximum likelihood estimator (MLE) of $\boldsymbol{\beta}$ for the purpose of inference. In this paper we also stick to the MLE of $\boldsymbol{\beta}$. For a sample $\{(\mathbf{x}_i, y_i)\}_{i=1}^n$, the likelihood is given by

$$L(\boldsymbol{\beta}|y_1, \dots, y_n, \mathbf{x}_1, \dots, \mathbf{x}_n) = \prod_{i=1}^n p(\mathbf{x}_i)^{y_i} (1 - p(\mathbf{x}_i))^{1-y_i},$$

where $p(\boldsymbol{\beta}|\mathbf{x}_i) = \frac{e^{\mathbf{x}_i'\boldsymbol{\beta}}}{1 + e^{\mathbf{x}_i'\boldsymbol{\beta}}}$, as implied from the model (1.1). The MLE $\hat{\boldsymbol{\beta}}_n$ of $\boldsymbol{\beta}$ is defined as the maximizer of $L(\boldsymbol{\beta}|y_1, \dots, y_n, \mathbf{x}_1, \dots, \mathbf{x}_n)$, which is obtained by solving

$$\sum_{i=1}^n (y_i - p(\boldsymbol{\beta}|\mathbf{x}_i))\mathbf{x}_i = 0, \quad (1.2)$$

since the second derivative of the left hand side (LHS) of (1.2) with respect to $\boldsymbol{\beta}$ is negative definite. We assume that the design vectors $\mathbf{x}_1, \dots, \mathbf{x}_n$ are fixed. In order to perform statistical inference, it is of interest to approximate the distribution of $\hat{\boldsymbol{\beta}}_n$. Since $\hat{\boldsymbol{\beta}}_n$ is the MLE, under certain regularity conditions it follows normal distribution asymptotically. This asymptotic distribution can be used for inferences regarding the regression parameter $\boldsymbol{\beta}$. In literature, asymptotic normality and other large sample properties of $\hat{\boldsymbol{\beta}}_n$ have been studied extensively (cf. Haberman (1974), McFadden (1974), Amemiya (1976), Gourieroux and Monfort (1981), Fahrmeir and Kaufmann (1985)). For the sake of completeness, we state the rate of convergence of $\hat{\boldsymbol{\beta}}_n$ to normality. See section 3 for details.

Bootstrap inference is an alternative to the asymptotic inference for underlying parameter of interest. Sometime the bootstrap inference is more accurate as is the case in linear regression. Hence it is also important to explore different bootstrap methods in logistic regression for the purpose of inference. Freedman (1981) proposed the residual bootstrap or the paired bootstrap depending on whether the covariates are non-random or random in linear regression. Some other bootstrap methods in linear regression are the wild bootstrap, the generalized bootstrap and the perturbation bootstrap. For the understanding of these bootstrap methods and their higher order properties see for example Liu (1988), Lahiri (1992), Mammen (1993), Chatterjee and Bose (2005), Das and Lahiri (2019) and references therein. Using similar mechanism of the residual and the paired bootstrap, Moulton and Zeger (1989, 1991) developed the standardized Pearson residual resampling and the observation vector resampling in generalized linear models (GLM). Lee (1990) considered the logistic regression model and showed that the conditional distribution of these resample based bootstrap estimators for the given data are close to the distribution of the original estimator in almost sure sense. Claeskens et al. (2003) developed two bootstrap methods for logistic regression in univariate case. The first method is ‘linear one-step bootstrap’ which was developed following the linearization principle proposed in Davison et al. (1986). The second approach known as ‘quadratic one-step bootstrap’ was constructed based on the quadratic approximation of the estimators discussed in Ghosh (1994). Claeskens et al. (2003) established the validity of these two bootstrap methods for approximating the underlying distribution in almost sure sense. Claeskens et al. (2003) also developed a finite sample bias correction of logistic regression estimator using their quadratic one-step bootstrap method and validated its effectiveness using simulations.

Despite the development of the above-mentioned bootstrap methods, to the best of our knowledge, the error rate of the approximations has not been studied. However, it is essential to study the error rate of the bootstrap methods to have an understanding about the sample size requirement for practical implementations. In case of linear regression, the bootstrap methods generally achieve second order correctness (SOC), i.e. having the error rate $o(n^{-1/2})$ uniformly over a class of sets. SOC is important to draw more accurate inference results compared to that based on asymptotic normal distribution. For elaborate description on the results on SOC of residual, generalized and perturbation bootstrap methods in linear regression one can look into Lahiri (1992), Barbe and Bertail (2012) and Das and Lahiri (2019) and references their in. In the same spirit, it is expected for any bootstrap method to achieve SOC in logistic regression as well. However SOC is explored for none of the bootstrap methods that are available in the literature for logistic regression. In this paper, we develop a perturbation bootstrap method in logistic regression. We show that the perturbation bootstrap attains SOC in approximating the distribution of $\hat{\boldsymbol{\beta}}_n$. SOC implies that the inference based on our proposed bootstrap method is much more accurate than that based on

asymptotic normality. Here the main ingredients are the appropriately chosen perturbation factors. Then we need to change (1.2) based on these perturbation factors and define the bootstrap estimator as the solution of the changed equation. See section 2 for details on the construction of the bootstrap version of the logistic regression estimator.

To establish SOC, we start with a studentization of $\sqrt{n}(\hat{\beta}_n - \beta)$ and its perturbation bootstrap version. We show that SOC cannot in general be achieved only by studentization, unlike in the case of multiple linear regression. This is due to the lattice nature of the distribution of the logistic regression estimator $\hat{\beta}_n$. The lattice nature is the result of the binary nature of the responses y_1, \dots, y_n . SOC is generally established by comparing the Edgeworth expansions in original and bootstrap case. However the usual Edgeworth expansion does not exist when the underlying setup is lattice. Correction terms are required to take care the lattice nature. For example one can look into Theorem 20.8 and corollary 23.2 in BR(86) and compare them. These correction terms in general can not be approximated with an error of $o(n^{-1/2})$, which makes SOC unachievable even with studentization. As a remedy we adopt the novel smoothing technique developed in Lahiri (1993). First, this smoothing technique is applied to transform the lattice nature of the distribution of the studentized version into absolute continuity. Then we show that the smoothed studentized version has edgeworth expansion without any correction term. Further we use the same smoothing technique for the bootstrap version also and we establish SOC by comparing the edgeworth expansions in the original and the bootstrap case. In this regard, one of the important results that we establish is the edgeworth expansion of a smoothed version of a sequence of sample means of independent (not necessarily identically distributed) random vectors (cf. Lemma 3). Lemma 3 may be of some independent interest. Moreover, an interesting property of the smoothing is that it has negligible effect on the asymptotic variance of $\hat{\beta}_n$ and hence it is not required to incorporate the effect of the smoothing in the form of the studentization. To compare the error in bootstrap approximation with that of normal approximation, we also establish the rate of convergence of the studentized $\hat{\beta}_n$ to normality.

The rest of the paper is organized as follows. The perturbation bootstrap version of the logistic regression estimator is described in Section 2. Main results are stated in Section 3. Proofs are presented in Section 4.

2 Description of the Perturbation Bootstrap

Let G_1^*, \dots, G_n^* be n independent copies of a non-negative and non-degenerate random variable G^* having mean μ_{G^*} , $Var(G^*) = \mu_{G^*}^2$ and $\mathbf{E}(G^* - \mu_{G^*})^3 = \mu_{G^*}^3$. These quantities serve as perturbation quantities in the construction of the perturbation bootstrap version of the logistic regression estimator. We define the bootstrap version as the minimizer of a carefully constructed objective function which involves the observed values y_1, \dots, y_n as well as the estimated probability of successes $\hat{p}(\mathbf{x}_i) = \frac{e^{\mathbf{x}_i' \hat{\beta}_n}}{1 + e^{\mathbf{x}_i' \hat{\beta}_n}}$, $i = 1, \dots, n$. Formally, the perturbation bootstrapped logistic regression estimator $\hat{\beta}_n^*$ is defined as

$$\hat{\beta}_n^* = \arg \max_{\mathbf{t}} \left[\sum_{i=1}^n \left\{ (y_i - \hat{p}(\mathbf{x}_i)) \mathbf{x}_i' \mathbf{t} \right\} (G_i^* - \mu_{G^*}) + \mu_{G^*} \sum_{i=1}^n \left\{ \hat{p}(\mathbf{x}_i) (\mathbf{x}_i' \mathbf{t}) - \log(1 + e^{\mathbf{x}_i' \mathbf{t}}) \right\} \right].$$

In other words, $\hat{\beta}_n^*$ is the solution of the equation

$$\sum_{i=1}^n (y_i - \hat{p}(\mathbf{x}_i)) \mathbf{x}_i (G_i^* - \mu_{G^*}) \mu_{G^*}^{-1} + \sum_{i=1}^n (\hat{p}(\mathbf{x}_i) - p(\mathbf{t}|\mathbf{x}_i)) \mathbf{x}_i = 0, \quad (2.1)$$

since the derivative of the LHS of (2.1) with respect to \mathbf{t} is negative definite. If bootstrap equation (2.1) is compared to the original equation (1.2), it is easy to note that the second part of the LHS of (2.1) is the estimated version of the LHS of (1.2). Obviously the bootstrap randomness is coming from the first part. The first part is also the main contributing term in the asymptotic expansion of the studentized version of $\hat{\beta}_n^*$. One immediate choice of the distribution of G^* is Beta(1/2, 3/2). Other choices can be found in Liu (1988), Mammen (1993) and Das et al. (2019). The moment characteristics of G^* are assumed to be true for the rest of this paper. Any additional assumption on G^* will be stated in respective theorems.

3 Main Results

This section is divided into two sub-sections. We begin with a Berry-Esseen type theorem for a studentized version of the logistic regression estimator $\hat{\beta}_n$. Theorem 1 is pointing out the error rate of the normal approximation which is generally used for inference in logistic regression. In the next sub-section, we start with exploring the effectiveness of bootstrap in approximating the distribution of the studentized version. Theorem 2 shows that we can not in general achieve SOC solely by studentization even when $p = 1$. As a remedy, we introduce a smoothing in the studentization and show that bootstrap achieves SOC.

For any natural number m , the class of sets \mathcal{A}_m is the collection of Borel subsets of \mathcal{R}^m satisfying

$$\sup_{B \in \mathcal{A}_m} \Phi((\delta B)^\epsilon) = O(\epsilon) \quad \text{as } \epsilon \downarrow 0.$$

Here Φ denotes the normal distribution with mean $\mathbf{0}$ and dispersion matrix being the identity matrix. We are going to use the class \mathcal{A}_p for the uniform asymptotic results on normal and bootstrap approximations. \mathbf{P}_* denotes the conditional bootstrap probability of G^* given data $\{y_1, \dots, y_n\}$.

3.1 Rate of Normal Approximation

In this sub-section we are going to explore the rate of normal approximation of suitable studentized version of the logistic regression estimator $\hat{\beta}_n$, uniformly over the class of sets \mathcal{C}_p . From the definition (1.2) of $\hat{\beta}_n$, we have that $\sum_{i=1}^n (y_i - \hat{p}(\mathbf{x}_i)) \mathbf{x}_i = 0$. Now using Taylor's expansion of $\sqrt{n}(\hat{\beta}_n - \beta)$, it is easy to see that the asymptotic variance of $\sqrt{n}(\hat{\beta}_n - \beta)$ is \mathbf{L}_n^{-1} where $\mathbf{L}_n = n^{-1} \sum_{i=1}^n \mathbf{x}_i \mathbf{x}_i' e^{\mathbf{x}_i' \beta} (1 + e^{\mathbf{x}_i' \beta})^{-2}$. An estimator of \mathbf{L}_n can be obtained by replacing β by $\hat{\beta}_n$ in the form of $\hat{\mathbf{L}}_n$. Hence we can define the studentized version of $\hat{\beta}_n$ as

$$\tilde{\mathbf{H}}_n = \sqrt{n} \hat{\mathbf{L}}_n^{1/2} (\hat{\beta}_n - \beta),$$

where $\hat{\mathbf{L}}_n = n^{-1} \sum_{i=1}^n \mathbf{x}_i \mathbf{x}_i' e^{\mathbf{x}_i' \hat{\beta}_n} (1 + e^{\mathbf{x}_i' \hat{\beta}_n})^{-2}$. Other studentized versions can be constructed by considering other estimators of \mathbf{L}_n . For details of the construction of different studentized versions, one can look into Lahiri (1994). The result on normal approximation will hold for other studentized versions also as long as it involves the estimator of \mathbf{L}_n which is \sqrt{n} -consistent.

Berry-Esseen theorem states that the error in normal approximation for the distribution of the mean of a sequence of independent random variables is $O(n^{-1/2})$, provided the average third absolute moment is bounded [cf. Theorem 12.4 in BR(86)]. Here we will see that there is an extra "log n " term in the error rate of the normal approximation. This is essentially due to the error incurred in Taylor's approximation of $\sqrt{n}(\hat{\beta}_n - \beta)$. This error can not in general be corrected by higher order approximations, like Edgeworth expansions, since the underlying setup in logistic regression has lattice nature. One important tool in deriving the error rate in normal approximation and later for deriving the higher order result for the bootstrap is to find the rate of convergence of $\hat{\beta}_n$ to β . To this end, we are now ready to state our first theorem.

Theorem 1 Suppose $n^{-1} \sum_{i=1}^n \|\mathbf{x}_i\|^3 = O(1)$ and $\mathbf{L}_n \rightarrow \mathbf{L}$ as $n \rightarrow \infty$ where \mathbf{L} is a pd matrix. Then

(a) there exists a positive constant C_0 such that when $n > C_0$ we have

$$\mathbf{P}\left(\hat{\boldsymbol{\beta}}_n \text{ solves (1.2) and } \|\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta}\| \leq C_0 n^{-1/2} (\log n)^{1/2}\right) = 1 - o(n^{-1/2}).$$

(b) we have

$$\sup_{\mathbf{B} \in \mathcal{A}_p} |\mathbf{P}(\tilde{\mathbf{H}}_n \in \mathbf{B}) - \Phi(\mathbf{B})| = O(n^{-1/2} \log n).$$

3.2 Rate of Bootstrap Approximation

We need to define the suitable studentized versions in both original and bootstrap setting before exploring the rate of convergence of bootstrap. Similar to original case, we have to find the asymptotic variance of the bootstrapped logistic regression estimator $\hat{\boldsymbol{\beta}}_n^*$ to define the studentized version in the bootstrap setting. Using Taylor's expansion, from (2.1) it is easy to see that the asymptotic variance of $\sqrt{n}(\hat{\boldsymbol{\beta}}_n^* - \hat{\boldsymbol{\beta}}_n)$ is $\hat{\mathbf{L}}_n^{-1} \hat{\mathbf{M}}_n \hat{\mathbf{L}}_n^{-1}$ where $\hat{\mathbf{L}}_n = n^{-1} \sum_{i=1}^n \mathbf{x}_i \mathbf{x}_i' e^{\mathbf{x}_i' \hat{\boldsymbol{\beta}}_n} (1 + e^{\mathbf{x}_i' \hat{\boldsymbol{\beta}}_n})^{-2}$ and $\hat{\mathbf{M}}_n = n^{-1} \sum_{i=1}^n (y_i - \hat{p}(\mathbf{x}_i))^2 \mathbf{x}_i \mathbf{x}_i'$. Therefore the studentized version in bootstrap setting can be defined as

$$\mathbf{H}_n^* = \sqrt{n} \hat{\mathbf{M}}_n^{*-1/2} \mathbf{L}_n^* (\hat{\boldsymbol{\beta}}_n^* - \hat{\boldsymbol{\beta}}_n),$$

where $\mathbf{L}_n^* = n^{-1} \sum_{i=1}^n \mathbf{x}_i \mathbf{x}_i' e^{\mathbf{x}_i' \hat{\boldsymbol{\beta}}_n^*} (1 + e^{\mathbf{x}_i' \hat{\boldsymbol{\beta}}_n^*})^{-2}$ and $\hat{\mathbf{M}}_n^* = n^{-1} \sum_{i=1}^n (y_i - \hat{p}(\mathbf{x}_i))^2 \mathbf{x}_i \mathbf{x}_i' \mu_{G^*}^{-2} (G_i^* - \mu_{G^*})^2$. Analogously, we define the original studentized version as

$$\mathbf{H}_n = \sqrt{n} \hat{\mathbf{M}}_n^{-1/2} \hat{\mathbf{L}}_n (\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta}),$$

which will be used for investigating SOC of bootstrap for rest of this section. In the next theorem we show that \mathbf{H}_n^* fails to be SOC in approximating the distribution of \mathbf{H}_n even when $p = 1$.

Theorem 2 Suppose $p = 1$ and denote the only covariate by x in the model (1.1). Let x_1, \dots, x_n be the observed values of x and β be the true value of the regression parameter. Define, $\mu_n = n^{-1} \sum_{i=1}^n x_i p(\beta | x_i)$. Assume the following conditions hold:

- (C.1) x_1, \dots, x_n are non random and are all integers.
- (C.2) $x_{i_1}, \dots, x_{i_m} = 1$ where $\{i_1, \dots, i_m\} \subseteq \{1, \dots, n\}$ with $m \geq (\log n)^2$.
- (C.3) $\max\{|x_i| : i = 1, \dots, n\} = O(1)$ and $\liminf_{n \rightarrow \infty} [n^{-1} \sum_{i=1}^n |x_i|^6] > 0$.
- (C.4) $\sqrt{n} |\mu_n| < M_1$ for $n \geq M_1$ where M_1 is a positive constant.
- (C.5) The distribution of G_1^* has an absolutely continuous component with respect to Lebesgue measure.
- (C.6) $\mathbf{E}G^{*4} < \infty$.

Then there exist an interval \mathbf{B}_n and a positive constants M_2 (does not depend on n) such that

$$\limsup_{n \rightarrow \infty} \mathbf{P}\left(\sqrt{n} |\mathbf{P}^*(\mathbf{H}_n^* \in \mathbf{B}_n) - \mathbf{P}(\mathbf{H}_n \in \mathbf{B}_n)| \geq M_2\right) = 1$$

Theorem 2 shows that the bootstrap can not in general achieve SOC even with studentization, unlike the scenario in multiple linear regression. Now let us discuss the form of the set \mathbf{B}_n . \mathbf{B}_n is of the form $f_n(\mathbf{E}_n \times \mathcal{R})$ with $\mathbf{E}_n = (-\infty, z_n]$ and $z_n = \left(\frac{3}{4n} - \mu_n\right)$. $f_n(\cdot)$ is a continuous function which is obtained from the Taylor expansion of \mathbf{H}_n . Since $\mathbf{E}_n \times \mathcal{R}$ is a convex subset of \mathcal{R}^2 , it is

also a connected set. Since $f_n(\cdot)$ is a continuous function, \mathbf{B}_n is a connected subset of \mathcal{R} and hence is an interval.

We are now going to define the smoothed versions of \mathbf{H}_n and \mathbf{H}_n^* which will result in achieving SOC by the bootstrap for general p . Note that the primary reason behind bootstrap's failure is the lattice nature of the distribution of $\sqrt{n}(\hat{\beta}_n - \beta)$. Hence if we can somehow smooth the distribution $\sqrt{n}(\hat{\beta}_n - \beta)$, or more generally the distribution of \mathbf{H}_n , so that the smoothed version has density with respect to Lebesgue measure then bootstrap may be shown to achieve SOC by employing theory of Edgeworth expansions. To that end, suppose Z is a p -dimensional standard normal random vector, independent of y_1, \dots, y_n . Define the smoothed version of \mathbf{H}_n as

$$\check{\mathbf{H}}_n = \mathbf{H}_n + \hat{M}_n^{-1/2} b_n Z, \quad (3.1)$$

where $\{b_n\}_{n \geq 1}$ is a suitable sequence such that it has negligible effect on the variance of $\sqrt{n}(\hat{\beta}_n - \beta)$ and hence on the studentization factor. See Theorem 3 for the conditions on $\{b_n\}_{n \geq 1}$. To define the smoothed studentized version in bootstrap setting, consider another p -dimensional standard normal vector by Z^* which is independent of $y_1, \dots, y_n, G_1^*, \dots, G_n^*$ and Z . Define the smoothed version of \mathbf{H}_n^* as

$$\check{\mathbf{H}}_n^* = \mathbf{H}_n^* + \hat{M}_n^{*-1/2} b_n Z^*. \quad (3.2)$$

The next theorem is the main theorem of this section. This shows that the smoothing does the trick for bootstrap to achieve SOC. Thus the inference on β based on the bootstrap after smoothing is much more accurate than the normal approximation. To state the main theorem, define $W_i = (Y_i \mathbf{x}_i', [Y_i^2 - \mathbf{E}Y_i^2] \mathbf{z}_i')'$ where $Y_i = (y_i - p(\beta | \mathbf{x}_i))$ and $\mathbf{z}_i = (x_{i1}^2, x_{i1}x_{i2}, \dots, x_{i1}x_{ip}, x_{i2}^2, x_{i2}x_{i3}, \dots, x_{i2}x_{ip}, \dots, x_{ip}^2)'$ with $\mathbf{x}_i = (x_{i1}, \dots, x_{ip})'$, $i \in \{1, \dots, n\}$.

Theorem 3 *Suppose $n^{-1} \sum_{i=1}^n \|\mathbf{x}_i\|^6 = O(1)$ and the matrix $n^{-1} \sum_{i=1}^n \text{Var}(W_i)$ converges to some positive definite matrix as $n \rightarrow \infty$. Also choose the sequence $\{b_n\}_{n \geq 1}$ such that $b_n = O(n^{-d})$ and $n^{-1/p_1} \log n = o(b_n^2)$ where $d > 0$ is a constant and $p_1 = \max\{p+1, 4\}$. Then*

(a) *there exists two positive constant C_2 such that when $n > C_2$ we have*

$$\mathbf{P}_* \left(\hat{\beta}_n^* \text{ solves (2.1) and } \|\hat{\beta}_n^* - \hat{\beta}_n\| \leq C_2 \cdot n^{-1/2} \cdot (\log n)^{1/2} \right) = 1 - o_p(n^{-1/2}).$$

(b) *we have*

$$\sup_{\mathbf{B} \in \mathcal{A}_p} |\mathbf{P}_*(\check{\mathbf{H}}_n^* \in \mathbf{B}) - \mathbf{P}(\check{\mathbf{H}}_n \in \mathbf{B})| = o_p(n^{-1/2}).$$

Remark 1. The class of sets \mathcal{A}_p used to state the uniform asymptotic results is somewhat abstract. There are two main reasons behind considering this class. One is to obtain asymptotic normality or to obtain valid Edgeworth expansions for the normalized part of the underlying pivot and the other one is to bound the remainder term by required small magnitude with sufficiently large probability (or bootstrap probability). A natural choice for \mathcal{A} is the collection of all Borel measurable convex subsets of \mathcal{R}^p , due to Theorem 3.1 in BR(86).

Remark 2. The results on bootstrap approximation presented in Theorem 3, may be established in almost sure sense also. In that case the only requirement is to have $n^{-1} \sum_{i=1}^n \|\mathbf{x}_i\|^{12} = O(1)$, since y_1, \dots, y_n can take either 0 or 1. Actually an almost sure version of part (a) of Theorem 3 is necessary to establish Theorem 2. Note that the requirement for almost sure version is met under the assumptions of Theorem 2.

4 Proof of the Results

Before going to the proofs we are going to define few notations. Suppose, $\Phi_{\mathbf{V}}$ and $\phi_{\mathbf{V}}$ respectively denote the normal distribution and its density with mean $\mathbf{0}$ and covariance matrix \mathbf{V} . We will write $\Phi_{\mathbf{V}} = \Phi$ and $\phi_{\mathbf{V}} = \phi$ when the dispersion matrix \mathbf{V} is the identity matrix. C, C_1, C_2, \dots denote generic constants that do not depend on the variables like n, x , and so on. ν_1, ν_2 denote vectors in \mathcal{R}^p , sometimes with some specific structures (as mentioned in the proofs). $(\mathbf{e}_1, \dots, \mathbf{e}_p)'$ denote the standard basis of \mathcal{R}^p . For a non-negative integral vector $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_l)'$ and a function $f = (f_1, f_2, \dots, f_l) : \mathcal{R}^l \rightarrow \mathcal{R}^l, l \geq 1$, let $|\alpha| = \alpha_1 + \dots + \alpha_l, \alpha! = \alpha_1! \dots \alpha_l!, f^\alpha = (f_1^{\alpha_1}) \dots (f_l^{\alpha_l}), D^\alpha f_1 = D_1^{\alpha_1} \dots D_l^{\alpha_l} f_1$, where $D_j f_1$ denotes the partial derivative of f_1 with respect to the j th component of $\alpha, 1 \leq j \leq l$. We will write $D^\alpha = D$ if α has all the component equal to 1. For $\mathbf{t} = (t_1, t_2, \dots, t_l)' \in \mathcal{R}^l$ and α as above, define $\mathbf{t}^\alpha = t_1^{\alpha_1} \dots t_l^{\alpha_l}$. For any two vectors $\alpha, \beta \in \mathcal{R}^k, \alpha \leq \beta$ means that each of the component of α is smaller than that of β . For a set \mathbf{A} and real constants $a_1, a_2, a_1 \mathbf{A} + a_2 = \{a_1 y + a_2 : y \in \mathbf{A}\}, \partial A$ is the boundary of A and A^ϵ denotes the ϵ -neighbourhood of A for any $\epsilon > 0$. \mathcal{N} is the set of natural numbers. $C(\cdot), C_1(\cdot), \dots$ denote generic constants which depend on only their arguments. Given two probability measures P_1 and P_2 defined on the same space $(\Omega, \mathcal{F}), P_1 * P_2$ defines the measure on (Ω, \mathcal{F}) by convolution of P_1 & P_2 and $\|P_1 - P_2\| = |P_1 - P_2|(\Omega), |P_1 - P_2|$ being the total variation of $(P_1 - P_2)$. For a function $g : \mathcal{R}^k \rightarrow \mathcal{R}^m$ with $g = (g_1, \dots, g_m)'$,

$$\text{Grad}[g(\mathbf{x})] = \left(\left(\frac{\partial g_i(\mathbf{x})}{\partial x_j} \right) \right)_{m \times k}$$

Before moving to the proofs of the theorems, we are going to state few essential Lemmas. Also we are going to present the proof of Theorem 2 at last, since some proof steps of Theorem 3 will be essential in proving Theorem 2. For Lemma 2 below, define $\xi_{1,n,s}(\mathbf{t}) = \left(1 + \sum_{i=1}^{s-2} n^{-r/2} \tilde{P}_r(it : \{\bar{\chi}_{\nu,n}\}) \right) \exp \left\{ -\mathbf{t}' \mathbf{E}_n \mathbf{t} / 2 \right\}$ where $\mathbf{E}_n = n^{-1} \sum_{i=1}^n \text{Var}(Y_i)$ and $\bar{\chi}_{\nu,n}$ is the average ν th cumulant of Y_1, \dots, Y_n . Define $\bar{\rho}_l = n^{-1} \sum_{i=1}^n \mathbf{E} \|Y_i\|^l$, the average l th absolute moment of $\{Y_1, \dots, Y_n\}$. The polynomials $\tilde{P}_r(z : \{\bar{\chi}_{\nu,n}\})$ are defined on the pages of 51 – 53 of BR(86). Define the identity

$$\xi_{1,n,s}(\mathbf{t}) \left(\sum_{j=0}^{\infty} (-\|\mathbf{t}\|^2 b_n^2)^j / j! \right) = \xi_{n,s}(\mathbf{t}) + o(n^{-(s-2)/2}),$$

uniformly in $\|\mathbf{t}\| < 1$, where c_n is defined in Lemma 2. $\psi_{n,s}(\cdot)$ is the Fourier inverse of $\xi_{n,s}(\cdot)$.

Lemma 1 Suppose Y_1, \dots, Y_n are zero mean independent r.v.s with $\mathbf{E}(|Y_i|^t) < \infty$ for $i = 1, \dots, n$ and $S_n = \sum_{i=1}^n Y_i$. Let $\sum_{i=1}^n \mathbf{E}(|Y_i|^t) = \sigma_t, c_t^{(1)} = (1 + \frac{2}{t})^t$ and $c_t^{(2)} = 2(2+t)^{-1} e^{-t}$. Then, for any $t \geq 2$ and $x > 0$,

$$P[|S_n| > x] \leq c_t^{(1)} \sigma_t x^{-t} + \exp(-c_t^{(2)} x^2 / \sigma_2)$$

Proof of Lemma 1. This inequality was proved in Fuk and Nagaev (1971).

Lemma 2 For any $t > 0, \frac{1 - N(t)}{n(t)} \leq \frac{1}{t}$ wher $N(\cdot)$ and $n(\cdot)$ respectively denote the cdf and pdf of real valued standard normal rv.

Proof of Lemma 3: This inequality is proved in Birnbaum (1942).

Lemma 3 Suppose Y_1, \dots, Y_n are mean zero independent random vectors in \mathcal{R}^k with $\mathbf{E}_n = n^{-1} \sum_{i=1}^n \text{Var}(Y_i)$ converging to some positive definite matrix V . Let $s \geq 3$ be an integer and $\bar{\rho}_{s+\delta} = O(1)$ for some $\delta > 0$. Additionally assume Z to be a $N(\mathbf{0}, \mathbf{I}_k)$ random vector which is independent of Y_1, \dots, Y_n and the sequence $\{c_n\}_{n \geq 1}$ to be such that $c_n = O(n^{-d})$ & $n^{-(s-2)/\tilde{k}} \log n = o(c_n^2)$ where $\tilde{k} = \max\{k+1, s+1\}$ & $d > 0$ is a constant. Then for any Borel set B of \mathcal{R}^k ,

$$\left| \mathbf{P}(\sqrt{n}\bar{Y} + c_n Z \in B) - \int_B \psi_{n,s}(x) dx \right| = o\left(n^{-(s-2)/2}\right), \quad (4.1)$$

where $\psi_{n,s}(\cdot)$ is defined above.

Proof of Lemma 3. Define $V_i = Y_i I(\|Y_i\| \leq \sqrt{n})$ and $W_i = V_i - EV_i$. Suppose $\bar{\chi}_{\nu,n}$ is the average cumulant of W_1, \dots, W_n and $D_n = n^{-1} \sum_{i=1}^n \text{Var}(W_i)$. Let $\tilde{\xi}_{1,n,s}, \tilde{\xi}_{n,s}$ and $\tilde{\psi}_{n,s}$ are respectively obtained from $\xi_{1,n,s}, \xi_{n,s}$ and $\psi_{n,s}$ with $\bar{\chi}_{\nu,n}$ replaced by $\bar{\chi}_{\nu,n}$ and E_n replaced by D_n . For any Borel set $B \in \mathcal{R}^k$, define $B_n = B - n^{-1/2} \sum_{i=1}^n EV_i$. Then we have

$$\begin{aligned} & \left| \mathbf{P}(\sqrt{n}\bar{Y}_n + c_n Z \in B) - \int_B \psi_{n,s}(x) dx \right| \\ & \leq \left| \mathbf{P}(\sqrt{n}\bar{Y}_n + c_n Z \in B) - \mathbf{P}(\sqrt{n}\bar{V}_n + c_n Z \in B) \right| \\ & \quad + \left| \mathbf{P}(\sqrt{n}\bar{W}_n + c_n Z \in B_n) - \int_{B_n} \tilde{\psi}_{n,s}(x) dx \right| + \left| \int_{B_n} \tilde{\psi}_{n,s}(x) dx - \int_B \psi_{n,s}(x) dx \right| \\ & = I_1 + I_2 + I_3 \quad (\text{say}). \end{aligned} \quad (4.2)$$

First we are going to show that $I_1 = o\left(n^{-(s-2)/2}\right)$. Now writing G_j and G'_j to be distributions of $n^{-1/2}Y_j$ and $n^{-1/2}V_j$, $j \in \{1, \dots, n\}$, we have

$$\begin{aligned} I_1 & \leq \sum_{j=1}^n \|G_j - G'_j\| \\ & = 2 \sum_{j=1}^n P\left(\|Y_j\| > n^{1/2}\right) \\ & = o\left(n^{-(s-2)/2}\right), \end{aligned} \quad (4.3)$$

due to the fact that $n^{-1} \sum_{j=1}^n E\|Y_j\|^{s+\delta} = O(1)$. Next we are going to show $I_3 = o\left(n^{-(s-2)/2}\right)$.

Define $m_1 = \inf\{j : b_n^{2j} = o(n^{-(s-2)/2})\}$. Again note that the eigen values of D_n are bounded away from 0, due to (14.18) in corollary 14.2 of BR(86) and the fact that E_n converges to some positive definite matrix. Therefore we have

$$I_3 = \left| \int_{B_n} \tilde{\psi}_{n,s}^{m_1}(x) dx - \int_B \psi_{n,s}^{m_1}(x) dx \right| + o\left(n^{-(s-2)/2}\right) = I_{31} + o\left(n^{-(s-2)/2}\right) \quad (\text{say}), \quad (4.4)$$

uniformly for any Borel set B of \mathcal{R}^k , where

$$\begin{aligned} \psi_{n,s}^{m_1}(x) & = \left\{ \left[\sum_{r=0}^{s-2} n^{-r/2} \tilde{P}_r(-D : \{\bar{\chi}_{\nu,n}\}) \right] \left[\sum_{j=0}^{m_1-1} 2^{-j} (j!)^{-1} c_n^{2j} (D'D)^j \right] \right\} \phi_{E_n}(x) \quad \text{and} \\ \tilde{\psi}_{n,s}^{m_1}(x) & = \left\{ \left[\sum_{r=0}^{s-2} n^{-r/2} \tilde{P}_r(-D : \{\bar{\chi}_{\nu,n}\}) \right] \left[\sum_{j=0}^{m_1-1} 2^{-j} (j!)^{-1} c_n^{2j} (D'D)^j \right] \right\} \phi_{D_n}(x). \end{aligned}$$

Now writing $l(\mathbf{u}) = \|\mathbf{u}\|/2$, $\mathbf{u} \in \mathcal{R}^k$, and $a_n = n^{-1/2} \sum_{i=1}^n EV_i$, from (4.3) we have

$$\begin{aligned}
I_{31} &\leq \sum_{r=0}^{s-2} \sum_{j=0}^{m_1-1} n^{-r/2} b_n^{2j} \left[\int_{B_n} \left| \left\{ \tilde{P}_r(-D : \{\bar{\chi}_{\nu,n}\}) \frac{l(-D)}{j!} \right\} \phi_{E_n}(x) - \left\{ \tilde{P}_r(-D : \{\bar{\chi}_{\nu,n}\}) \frac{l(-D)}{j!} \right\} \phi_{D_n}(x) \right| dx \right. \\
&\quad \left. + \int_B \left| \left\{ \tilde{P}_r(-D : \{\bar{\chi}_{\nu,n}\}) \frac{l(-D)}{j!} \right\} \phi_{E_n}(x) - \left\{ \tilde{P}_r(-D : \{\bar{\chi}_{\nu,n}\}) \frac{l(-D)}{j!} \right\} \phi_{E_n}(x - a_n) \right| dx \right] \\
&\quad + o\left(n^{-(s-2)/2}\right) \\
&= I_{311} + I_{312} + o\left(n^{-(s-2)/2}\right) \quad (\text{say}). \tag{4.5}
\end{aligned}$$

Now assume $E_n = I_k$, the $k \times k$ identity matrix. Then following the proof of Lemma 14.6 of BR(86), it can be shown that $I_{311} + I_{312} = o\left(n^{-(s-2)/2}\right)$. Main ingredients of the proof are (14.74), (14.78), (14.79) and bounds similar to (14.80) and (14.86) in BR(86). The general case when E_n converges to a positive definite matrix, will follow essentially through the same line. Hence from (4.4) and (4.5), we have $I_3 = o\left(n^{-(s-2)/2}\right)$. The last step is to show $I_2 = o\left(n^{(s-2)/2}\right)$. Now let us write $\Gamma_n = \sqrt{n}\bar{W}_n + c_n Z$. Then recall that

$$I_2 = \left| \mathbf{P}(\Gamma_n \in B_n) - \int_{B_n} \tilde{\psi}_{n,s}(x) dx \right|.$$

By Theorem 4 of chapter 5 of Feller(2014), we can say that Γ_n has density with respect to the Lebesgue measure. Let us call that density by $q_n(\cdot)$. Then we have

$$I_2 \leq \int |q_n(x) - \tilde{\psi}_{n,s}(x)| dx \leq \int |q_n(x) - \tilde{\psi}_{n,(\bar{k}-1)}(x)| dx + \int |\tilde{\psi}_{n,s}(x) - \tilde{\psi}_{n,(\bar{k}-1)}(x)| dx, \tag{4.6}$$

where $\bar{k} = \max\{k+1, s+1\}$. Note that $\int \|x\|^j |q_n(x) - \tilde{\psi}_{n,(\bar{k}-1)}(x)| dx < \infty$ for any $j \in \mathcal{N}$, since $\tilde{\psi}_{n,(\bar{k}-1)}(x)$ has negative exponential term and \bar{W}_n is bounded. Therefore by Lemma 11.6 of BR(86) we have

$$\begin{aligned}
I_2 &\leq C(k) \left[\max_{|\beta| \in \{0, \dots, (k+1)\}} \int \left| D^\beta (\hat{q}(t) - \check{\xi}_{n,(\bar{k}-1)}(t)) \right| dt \right] + \int |\tilde{\psi}_{n,s}(x) - \tilde{\psi}_{n,(\bar{k}-1)}(x)| dx \\
&= I_{21} + I_{22} \quad (\text{say}). \tag{4.7}
\end{aligned}$$

Here $\hat{q}_n(\cdot)$ is the Fourier transform of the density $q(\cdot)$. Clearly $I_{22} = o\left(n^{-(s-2)/2}\right)$ by looking into the definition of $\tilde{\psi}_{n,s}(\cdot)$. Now define

$$\check{\xi}_{n,(\bar{k}-1)}(t) = \left[\sum_{r=0}^{\bar{k}-3} n^{-r/2} \tilde{P}_r(it : \{\bar{\chi}_{\nu,n}\}) \right] \exp\left(\frac{-t' D_n t - c_n^2 \|t\|^2}{2}\right).$$

Then we have

$$\begin{aligned}
I_{21} &\leq C(k) \max_{|\beta| \in \{0, \dots, (k+1)\}} \left[\int \left| D^\beta (\hat{q}_n(t) - \check{\xi}_{n,(\bar{k}-1)}(t)) \right| dt + \int \left| D^\beta (\check{\xi}_{n,(\bar{k}-1)}(t) - \check{\xi}_{n,(\bar{k}-1)}(t)) \right| dt \right] \\
&= I_{211} + I_{212} \quad (\text{say}) \tag{4.8}
\end{aligned}$$

First, we are going to show that $I_{212} = o\left(n^{-(s-2)/2}\right)$. Note that

$$\check{\xi}_{n,(\bar{k}-1)}(t) - \check{\xi}_{n,(\bar{k}-1)}(t) = \left[\sum_{r=0}^{\bar{k}-3} n^{-r/2} \tilde{P}_r(it : \{\bar{\chi}_{\nu,n}\}) \right] \exp\left(\frac{-t' D_n t}{2}\right) \sum_{j=m_2}^{\infty} \frac{c_n^{2j} \|t\|^{2j} (-1)^j}{2^j j!},$$

where $m_2 = m_2(r) = (s-2)^{-1}m_1(\tilde{k}-3-r)$. Therefore for any $\beta \in \mathcal{N}^k$ with $|\beta| \in \{0, \dots, k+1\}$ we have

$$\begin{aligned} & D^\beta \left(\check{\xi}_{n,(\tilde{k}-1)}(t) - \check{\xi}_{n,(\tilde{k}-1)}(t) \right) \\ &= \sum_{r=0}^* \sum_{j=m_2}^{\tilde{k}-3} \sum_{\infty} C_1(\alpha, \beta, \gamma) \frac{n^{-r/2}(-1)^j c_n^{2j}}{2^j j!} \left[D^\alpha \left(\tilde{P}_r(it : \{\check{\chi}_{\nu,n}\}) \right) \right] \left[D^\gamma \left(\exp \left(\frac{-t' \mathbf{D}_n t}{2} \right) \right) \right] D^{\beta-\alpha-\gamma} (\|t\|^{2j}), \end{aligned} \quad (4.9)$$

where \sum^* is over $\alpha, \gamma \in \mathcal{N}^k$ such that $0 \leq \alpha, \gamma \leq \beta$. Since the degree of the polynomial $\tilde{P}_r(it : \{\check{\chi}_{\nu,n}\})$ is $3r$, $D^\alpha \left(\tilde{P}_r(it : \{\check{\chi}_{\nu,n}\}) \right) = 0$ if $|\alpha| > 3r$. When $|\alpha| \leq 3r$, then recalling that $n^{-1} \sum_{i=1}^n E \|Y_i\|^s = O(1)$ and by Lemma 9.5 & Lemma 14.1(v) of BR(86) we have

$$\left| D^\alpha \left(\tilde{P}_r(it : \{\check{\chi}_{\nu,n}\}) \right) \right| \leq \begin{cases} C_2(\alpha, r) (\bar{\rho}_s)^{r/(s-2)} \left(1 + (\bar{\rho}_2)^{r(s-3)/(s-2)} \right) (1 + \|t\|^{3r-|\alpha|}), & \text{if } 0 \leq r \leq (s-2) \\ C_3(\alpha, r) n^{(r+2-s)/2} \bar{\rho}_s \left(1 + (\bar{\rho}_2)^{r-1} \right) (1 + \|t\|^{3r-|\alpha|}), & \text{if } r > (s-2). \end{cases} \quad (4.10)$$

Again note that

$$\left| D^\gamma \left(\exp \left(\frac{-t' \mathbf{D}_n t}{2} \right) \right) \right| \leq C_4(\gamma) (1 + \|t\|)^{|\gamma|} \|\mathbf{D}_n\|^{|\gamma|} \left(\exp \left(\frac{-t' \mathbf{D}_n t}{2} \right) \right) \quad (4.11)$$

$$\text{and } \sum_{j=m_2}^{\infty} \left| \frac{c_n^{2j} D^{\beta-\alpha-\gamma} (\|t\|^{2j})}{2^j j!} \right| \leq C_5(\alpha, \beta, \gamma) c_n^{2m_3} \left[e^{c_n^2/2} + \|t\|^{m_3} \exp(c_n^2 \|t\|^2/2) \right], \quad (4.12)$$

where $m_3 = m_3(\alpha, \beta, \gamma, r) = \max\{m_2, |\beta - \alpha - \gamma|/2\}$. Now combining (4.10)-(4.12), from (4.9) we have $I_{212} = o(n^{-(s-2)/2})$. Last step is to show $I_{211} = o(n^{-(s-2)/2})$. Recall that

$$\begin{aligned} I_{211} &= C(k) \max_{|\beta| \in \{0, \dots, (k+1)\}} \left[\int \left| D^\beta \left(\hat{q}_n(t) - \check{\xi}_{n,(\tilde{k}-1)}(t) \right) \right| dt \right] \\ &\leq C(k) \max_{|\beta| \in \{0, \dots, (k+1)\}} \left[\int_{A_n} \left| D^\beta \left(\hat{q}_n(t) - \check{\xi}_{n,(\tilde{k}-1)}(t) \right) \right| dt + \int_{A_n^c} \left| D^\beta \left(\hat{q}_n(t) - \check{\xi}_{n,(\tilde{k}-1)}(t) \right) \right| dt \right] \\ &= I_{2111} + I_{2112} \quad (\text{say}), \end{aligned} \quad (4.13)$$

where

$$A_n = \left\{ t \in \mathcal{R}^k : \|t\| \leq C_6(k) \lambda_n^{-1/2} \left(\frac{n^{1/2}}{\eta_{\tilde{k}}^{1/(\tilde{k}-2)}} \right)^{(\tilde{k}-2)/\tilde{k}} \right\},$$

with $C_6(k)$ being some fixed positive constant, λ_n being the largest eigen value of \mathbf{D}_n , $\eta_{\tilde{k}} = n^{-1} \sum_{i=1}^n E \|\mathbf{B}_n W_i\|^{\tilde{k}}$ and $\mathbf{B}_n^2 = \mathbf{D}_n^{-1}$. Note that

$$\begin{aligned} & D^\beta \left(\hat{q}_n(t) - \check{\xi}_{n,(\tilde{k}-1)}(t) \right) \\ &= \sum_{0 \leq \alpha \leq \beta} C_7(\alpha, \beta) D^\alpha \left(E \left(e^{i\sqrt{n}t'} \bar{W}_n \right) - \exp \left(\frac{-t' \mathbf{D}_n t}{2} \right) \sum_{r=0}^{\tilde{k}-3} n^{-r/2} \tilde{P}_r(it : \{\check{\chi}_{\nu,n}\}) \right) D^{\beta-\alpha} \left(\exp \left(\frac{-c_n^2 \|t\|^2}{2} \right) \right), \end{aligned} \quad (4.14)$$

where

$$\left| D^{\beta-\alpha} \left(\exp \left(\frac{-c_n^2 \|\mathbf{t}\|^2}{2} \right) \right) \right| \leq C_8(\boldsymbol{\alpha}, \boldsymbol{\beta}) c_n^{2|\beta-\alpha|} \|\mathbf{t}\|^{|\beta-\alpha|} \exp \left(\frac{-c_n^2 \|\mathbf{t}\|^2}{2} \right) \quad \text{and}$$

by Theorem 9.11 and the following remark of BR(86) we have

$$\begin{aligned} & \left| D^\alpha \left(E \left(e^{i\sqrt{n}\mathbf{t}'\tilde{W}_n} \right) - \exp \left(\frac{-\mathbf{t}'\mathbf{D}_n\mathbf{t}}{2} \right) \sum_{r=0}^{\tilde{k}-3} n^{-r/2} \tilde{P}_r \left(i\mathbf{t} : \left\{ \bar{\chi}_{\nu,n} \right\} \right) \right) \right| \\ & \leq C_9(k) \lambda_n^{|\alpha|/2} \eta_{\tilde{k}} n^{-(\tilde{k}-2)/2} \left[(\mathbf{t}'\mathbf{D}_n\mathbf{t})^{(\tilde{k}-|\alpha|/2)} + (\mathbf{t}'\mathbf{D}_n\mathbf{t})^{(3(\tilde{k}-2)+|\alpha|/2)} \right] \exp \left(\frac{-\mathbf{t}'\mathbf{D}_n\mathbf{t}}{4} \right). \end{aligned} \quad (4.15)$$

Now note that $\bar{\rho}_{s+\delta} = O(1)$ and E_n converges to a positive definite matrix \mathbf{E} . Hence applying Lemma 14.1(v) (with $s' = \tilde{k}$) and corollary 14.2 of BR(86), from (4.14) we have $I_{2111} = o(n^{-(s-2)/2})$. Again applying Lemma 14.1(v) and corollary 14.2 of BR(86) we have $\eta_{\tilde{k}} \leq C_{10}(\tilde{k}, s) n^{(\tilde{k}-s)/2} \bar{\rho}_s$ for large enough n and λ_n being converged to some positive number. Therefore we have for large enough n ,

$$A_n^c \subseteq B_n \quad \text{where} \quad B_n = \left\{ \mathbf{t} \in \mathcal{R}^k : \|\mathbf{t}\| > C_{11}(k, \mathbf{E}) n^{(s-2)/2\tilde{k}} \right\},$$

implying

$$\begin{aligned} I_{2112} & \leq C(k) \max_{|\beta| \in \{0, \dots, (k+1)\}} \int_{B_n} \left| D^\beta \left(\hat{q}_n(t) - \check{\xi}_{n,(\tilde{k}-1)}(t) \right) \right| dt \\ & \leq C(k) \max_{|\beta| \in \{0, \dots, (k+1)\}} \left[\int_{B_n} \left| D^\beta \left(\hat{q}_n(t) \right) \right| dt + \int_{B_n} \left| D^\beta \left(\check{\xi}_{n,(\tilde{k}-1)}(t) \right) \right| dt \right] \\ & = I_{21121} + I_{21122} \quad (\text{say}), \end{aligned} \quad (4.16)$$

for large enough n . To establish $I_{2112} = o(n^{-(s-2)/2})$, first we are going to show $I_{21122} = o(n^{-(s-2)/2})$. Note that

$$D^\beta \left(\check{\xi}_{n,(\tilde{k}-1)}(t) \right) = \sum_{0 \leq \alpha \leq \beta} C_{12}(\boldsymbol{\alpha}, \boldsymbol{\beta}) D^\alpha \left(\sum_{r=0}^{\tilde{k}-3} n^{-r/2} \tilde{P}_r \left(i\mathbf{t} : \left\{ \bar{\chi}_{\nu,n} \right\} \right) \right) D^{\beta-\alpha} \left(\exp \left(\frac{-\mathbf{t}'\tilde{\mathbf{D}}_n\mathbf{t}}{2} \right) \right),$$

where $\tilde{\mathbf{D}}_n = \mathbf{D}_n + c_n^2 \mathbf{I}_k$. We are going to use bounds (4.10) and (4.11) with \mathbf{D}_n being replaced by $\tilde{\mathbf{D}}_n$. Note that by Corollary 14.2 of BR(86) and the fact that $c_n = O(n^{-d})$, $\tilde{\mathbf{D}}_n$ converges to the positive definite matrix \mathbf{E} , which is the limit of \mathbf{E}_n . Hence those bounds will imply that for large enough n ,

$$\begin{aligned} I_{21122} & = C(k) \max_{|\beta| \in \{0, \dots, (k+1)\}} \int_{B_n} \left| D^\beta \left(\check{\xi}_{n,(\tilde{k}-1)}(t) \right) \right| dt \\ & \leq C_{13}(k, \mathbf{E}) n^{(\tilde{k}+1-s)/2} \int_{B_n} \left(1 + \|\mathbf{t}\|^{3(\tilde{k}-1)} \right) \exp \left(-C_{14}(\mathbf{E}) \|\mathbf{t}\|^2/2 \right) dt \\ & \leq C_{15}(k, \mathbf{E}) n^{(\tilde{k}+1-s)/2} \int_{B_n} \exp \left(-C_{14}(\mathbf{E}) \|\mathbf{t}\|^2/4 \right) dt. \end{aligned} \quad (4.17)$$

Now apply Lemma 2 to conclude that $I_{21122} = o(n^{-(s-2)/2})$. Only remaining thing to show is $I_{21121} = o(n^{-(s-2)/2})$. Note that

$$D^\beta \left(\hat{q}_n(t) \right) = \sum_{0 \leq \alpha \leq \beta} C_{16}(\boldsymbol{\alpha}, \boldsymbol{\beta}) D^\alpha \left(E \left(e^{i\sqrt{n}\mathbf{t}'\tilde{W}_n} \right) \right) D^{\beta-\alpha} \left(\exp \left(\frac{-c_n^2 \|\mathbf{t}\|^2}{2} \right) \right), \quad (4.18)$$

where

$$\begin{aligned} & \left| D^\alpha \left(E \left(e^{i\sqrt{n}\mathbf{t}'\bar{W}_n} \right) \right) \right| \leq \left| D^\alpha \left(\prod_{i=1}^n E \left(e^{i\mathbf{t}'W_i/\sqrt{n}} \right) \right) \right| \\ \text{and } & \left| D^{\beta-\alpha} \left(\exp \left(\frac{-c_n^2 \|\mathbf{t}\|^2}{2} \right) \right) \right| \leq C_{17}(\boldsymbol{\alpha}, \boldsymbol{\beta}) \left(1 + \|\mathbf{t}\|^{|\beta-\alpha|} \right) \exp \left(\frac{-c_n^2 \|\mathbf{t}\|^2}{2} \right). \end{aligned}$$

Now by Leibniz's rule of differentiation, $D^\alpha \left(E \left(e^{i\sqrt{n}\mathbf{t}'\bar{W}_n} \right) \right)$ is the sum of $n^{|\alpha|}$ terms. A typical term is of the form

$$\prod_{i \notin C_r} E \left(e^{i\mathbf{t}'W_i/\sqrt{n}} \right) \prod_{l=1}^r D^{\beta_l} \left(E \left(e^{i\mathbf{t}'W_{i_l}/\sqrt{n}} \right) \right),$$

where $C_r = \{i_1, \dots, i_r\} \subset \{1, \dots, n\}$, $1 \leq r \leq |\alpha|$. β_1, \dots, β_r are non-negative integral vectors satisfying $|\beta_j| \geq 1$ for all $j \in \{1, \dots, r\}$ and $\sum_{j=1}^r \beta_j = \alpha$. Note that $\left| D^{\beta_l} \left(E \left(e^{i\mathbf{t}'W_{i_l}/\sqrt{n}} \right) \right) \right| \leq n^{-|\beta_l|/2} E \|W_{i_l}\|^{|\beta_l|}$ and $W_{j_l} \leq 2\sqrt{n}$, which imply that

$$\begin{aligned} & \left| \prod_{i \notin C_r} E \left(e^{i\mathbf{t}'W_i/\sqrt{n}} \right) \prod_{l=1}^r D^{\beta_l} \left(E \left(e^{i\mathbf{t}'W_{i_l}/\sqrt{n}} \right) \right) \right| \leq 2^{\sum_{l=1}^r |\beta_l|} = 2^{|\alpha|} \\ \Rightarrow & \left| D^\alpha \left(E \left(e^{i\sqrt{n}\mathbf{t}'\bar{W}_n} \right) \right) \right| \leq (2n)^{|\alpha|}. \end{aligned}$$

Let $K_n = C_{11}(k, \mathbf{E})n^{(s-2)/2\bar{k}}$. Therefore from (4.18), for large enough n we have

$$\begin{aligned} I_{21121} & \leq \left[\max_{|\beta| \in \{0, \dots, (k+1)\}} \sum_{\mathbf{0} \leq \alpha \leq \beta} C_{16}(\boldsymbol{\alpha}, \boldsymbol{\beta}) \right] (2n)^{k+1} \left[\int_{B_n} \left(1 + \|\mathbf{t}\|^{k+1} \right) \exp \left(\frac{-c_n^2 \|\mathbf{t}\|^2}{2} \right) \right] \\ & \leq C_{18}(k) (2n)^{k+1} \int_{r \geq K_n} r^{k-1} (1 + r^{k+1}) e^{-c_n^2 r^2/2} dr \\ & \leq C_{19}(k) (2n)^{k+1} c_n^{-1} \int_{r \geq K_n} \frac{1}{2\sqrt{\pi}c_n^{-1}} e^{-c_n^2 r^2/4} dr \\ & \leq C_{20}(k) n^{k+d+1} \int_{c_n K_n/\sqrt{2}}^{\infty} \frac{1}{\sqrt{2\pi}} e^{-z^2/2} dz \\ & = o\left(n^{-(s-2)/2}\right). \end{aligned} \tag{4.19}$$

The second inequality follows by considering polar transformation. Third inequality follows due to the assumptions that $n^{-(s-2)/\bar{k}}(\log n) = o(c_n^2)$ and $c_n = O(n^{-d})$. The last equality is the implication of Lemma 2. Therefore the proof of Lemma 3 is now complete.

Lemma 4 *Suppose all the assumptions of Lemma 2 are true. Define $d_n = n^{-1/2}c_n$ and $A_\delta = \{x \in \mathcal{R}^k : \|\mathbf{x}\| < \delta\}$ for some $\delta > 0$. Let $H : \mathcal{R}^k \rightarrow \mathcal{R}^m$ ($k \geq m$) has continuous partial derivatives of all orders on A_δ and $\text{Grad}[H(\mathbf{0})]$ is of full row rank. Then for any Borel set B of \mathcal{R}^m we have*

$$\left| \mathbf{P} \left(\sqrt{n} (H(\bar{Y}_n + d_n Z) - H(\mathbf{0})) \in B \right) - \int_B \check{\psi}_{n,s}(x) dx \right| = o\left(n^{-(s-2)/2}\right), \tag{4.20}$$

where $\check{\psi}_{n,s}(\mathbf{x}) = \left[1 + \sum_{r=1}^{s-2} n^{-r/2} a_{1,r}(Q_n, \mathbf{x}) \phi_{\bar{M}_n}(\mathbf{x}) \right] \left[\sum_{j=1}^{m_1-1} c_n^{2j} a_{2,j}(\mathbf{x}) \right]$ with $m_1 = \inf \{j : c_n^{2j} = o(n^{-(s-2)/2})\}$ and Q_n being the distribution of $\sqrt{n}\bar{Y}_n$. $a_{1,r}(Q_n, \cdot)$, $r \in \{1, \dots, (s-2)\}$, are polynomials whose coefficients are continuous functions of first s average cumulants of $\{Y_1, \dots, Y_n\}$. $a_{2,j}(\cdot)$, $j \in \{1, \dots, (m-1)\}$, are polynomials whose coefficients are continuous functions of partial derivatives of H of order $(s-1)$ or less. $\bar{M}_n = \bar{\mathbf{B}}\mathbf{E}_n\bar{\mathbf{B}}'$ with $\bar{\mathbf{B}} = \text{Grad}[H(\mathbf{0})]$ and $\mathbf{E}_n = n^{-1} \sum_{i=1}^n \text{Var}(Y_i)$.

Proof of Lemma 4. This follows exactly through the same line of the proof of Lemma 3.2 in Lahiri (1989).

Lemma 5 Let Y_1, \dots, Y_n be mean zero independent random vectors in \mathcal{R}^k with $n^{-1} \sum_{i=1}^n E\|Y_i\|^3 = O(1)$. Suppose $T_n^2 = E_n^{-1}$ where $E_n = n^{-1} \sum_{i=1}^n \text{Var}(Y_i)$ is the average positive definite covariance matrix and E_n converges to some positive definite matrix as $n \rightarrow \infty$. Then for any Borel subset B of \mathcal{R}^k we have

$$\left| \mathbf{P}\left(n^{-1/2} T_n \sum_{i=1}^n Y_i \in B\right) - \Phi(B) \right| \leq C_{22}(k) n^{-1/2} \rho_3 + 2 \Phi\left((\partial B)^{C_{22}(k) \rho_3 n^{-1/2}}\right),$$

where $\rho_3 = n^{-1} \sum_{i=1}^n E\|T_n Y_i\|^3$.

Proof of Lemma 5. This is a direct consequence of part (a) of corollary 24.3 in BR(86).

Lemma 6 Suppose A, B are matrices such that $(A - aI)$ and $(B - aI)$ are positive semi-definite matrices of same order, for some $a > 0$. For some $r > 0$, A^r, B^r are defined in the usual way. Then for any $0 < r < 1$, we have

$$\|A^r - B^r\| \leq r a^{r-1} \|A - B\|.$$

Proof of Lemma 6. More general version of this lemma is stated as corollary (X.46) in Bhatia (1996).

Lemma 7 Suppose all the assumptions of Lemma 4 are true and $\check{M}_n = I_m$, the $m \times m$ identity matrix. Define $\hat{H}_n = \left[\sqrt{n}(H(\bar{Y}_n + d_n Z) - H(\mathbf{0})) \right] + R_n$ where $\mathbf{P}\left(\|R_n\| = o(n^{-(s-2)/2})\right) = 1 - o(n^{-(s-2)/2})$ and s is as defined in Lemma 3. Then we have

$$\sup_{B \in \mathcal{A}_m} \left| \mathbf{P}\left(\hat{H}_n \in B\right) - \int_B \check{\psi}_{n,s}(x) dx \right| = o\left(n^{-(s-2)/2}\right), \quad (4.21)$$

where the class of sets \mathcal{A}_m is as defined in section 3.

Proof of Lemma 7. Recall the definition of $(\partial B)^\epsilon$ which is given in section 3. For some $B \subseteq \mathcal{R}^m$ and $\delta > 0$, define $B^{n,s,\delta} = (\partial B)^{\delta n^{-(s-2)/2}}$. Hence using Lemma 4, for any $B \in \mathcal{A}_m$ we have

$$\begin{aligned} & \left| \mathbf{P}\left(\hat{H}_n \in B\right) - \int_B \check{\psi}_{n,s}(x) dx \right| = o\left(n^{-(s-2)/2}\right) \\ & \leq \left| \mathbf{P}\left(\hat{H}_n \in B\right) - \mathbf{P}\left(\sqrt{n}(H(\bar{Y}_n + d_n Z) - H(\mathbf{0})) \in B\right) \right| + o\left(n^{-(s-2)/2}\right) \\ & \leq \mathbf{P}\left(\|R_n\| \neq o\left(n^{-(s-2)/2}\right)\right) + 2\mathbf{P}\left(\sqrt{n}(H(\bar{Y}_n + d_n Z) - H(\mathbf{0})) \in B^{n,s,\delta}\right) + o\left(n^{-(s-2)/2}\right) \\ & = 2\mathbf{P}\left(\sqrt{n}(H(\bar{Y}_n + d_n Z) - H(\mathbf{0})) \in B^{n,s,\delta}\right) + o\left(n^{-(s-2)/2}\right) \\ & = 2 \int_{B^{n,s,\delta}} \check{\psi}_{n,s}(x) dx + o\left(n^{-(s-2)/2}\right) \end{aligned} \quad (4.22)$$

for any $\delta > 0$. Now calculations at page 213 of BR(86) and arguments at page 58 of Lahiri(1989) imply that for any $B \in \mathcal{A}_m$,

$$\int_{B^{n,s,\delta}} \check{\psi}_{n,s}(x) dx \leq C_{21}(s) \sup_{B \in \mathcal{A}_m} \Phi(B^{n,s,\delta}) + o\left(n^{-(s-2)/2}\right) = o\left(n^{-(s-2)/2}\right),$$

since $\delta > 0$ is arbitrary. Therefore (4.21) follows from (4.22).

Lemma 8 Let \mathbf{A} and \mathbf{B} be positive definite matrices of same order. Then for some given matrix \mathbf{C} , the solution of the equation $\mathbf{A}\mathbf{X} + \mathbf{X}\mathbf{B} = \mathbf{C}$ can be expressed as

$$\mathbf{X} = \int_0^\infty e^{-t\mathbf{A}}\mathbf{C}e^{-t\mathbf{B}}dt,$$

where $e^{-t\mathbf{A}}$ and $e^{-t\mathbf{B}}$ are defined in the usual way.

Proof of Lemma 8. This lemma is an easy consequence of Theorem VII.2.3 in Bhatia (1996).

Lemma 9 Let W_1, \dots, W_n be n independent mean 0 random variables with average variance $s_n^2 = n^{-1} \sum_{i=1}^n \mathbf{E}W_i^2$ and $\mathbf{P}(\max\{|W_j| : j \in \{1, \dots, n\}\} \leq C_{30}) = 1$ for some positive constant C_{30} and integer $s \geq 3$. $\bar{\chi}_{\nu, n}$ is the average ν th cumulant. Recall the polynomial \tilde{P}_r for any non-negative integer r , as defined in the beginning of this section. Then there exists two constants $0 < C_{31}(s) < 1$ and $C_{32}(s) > 0$ such that whenever $|t| \leq C_{31}(s)\sqrt{n} \min\{C_{30}^{-2}s_n, C_{30}^{-s/(s-2)}s_n^{s/(s-2)}\}$, we have

$$\left| \prod_{j=1}^n \mathbf{E}\left(e^{in^{-1/2}tW_j}\right) - \sum_{r=0}^{s-2} n^{-r/2} \tilde{P}_r(it : \{\bar{\chi}_{\nu, n}\}) e^{-(s_n^2 t^2)/2} \right| \leq C_{32}(s) C_{30}^s s_n^{-s} n^{-(s-2)/2} \left[(s_n t)^s + (s_n t)^{3(s-2)} \right] e^{-(s_n^2 t^2)/4}$$

Proof of Lemma 9. In view of Theorem 9.9 of BR(86), enough to show that for any $j \in \{1, \dots, n\}$, $\left| \mathbf{E}(e^{its^{-1}n^{-1/2}W_j}) - 1 \right| \leq 1/2$ whenever $|t| \leq C_{31}(s)\sqrt{n} \min\{C_{30}^{-2}s_n, C_{30}^{-s/(s-2)}s_n^{s/(s-2)}\}$. This is indeed the case due to the fact that

$$\left| \mathbf{E}(e^{itn^{-1/2}W_j}) - 1 \right| \leq \frac{t^2 \mathbf{E}W_j^2}{2ns_n^2}.$$

Lemma 10 Assume the setup of Theorem 2 and let $X_i = y_i x_i$, $i \in \{1, \dots, n\}$. Define $\sigma_n^2 = n^{-1} \sum_{i=1}^n \text{Var}(X_i)$ and $\bar{\chi}_{\nu, n}$ as the ν th average cumulant of $\{(X_1 - E(X_1)), \dots, (X_n - E(X_n))\}$. $P_r(-\Phi_{\sigma_n^2} : \{\bar{\chi}_{\nu, n}\})$ is the finite signed measure on \mathcal{R} whose density is $\tilde{P}_r(-D : \{\bar{\chi}_{\nu, n}\})\phi_{\sigma_n^2}(x)$. Let $S_0(x) = 1$ and $S_1(x) = x - 1/2$. Suppose σ_n^2 is bounded away from both 0 & ∞ and assumptions (C.1)-(C.3) of Theorem 2 hold. Then we have

$$\sup_{x \in \mathcal{R}} \left| \mathbf{P}\left(n^{-1/2} \sum_{i=1}^n (X_i - E(X_i)) \leq x\right) - \sum_{r=0}^1 n^{-r/2} (-1)^r S_r(n\mu_n + n^{1/2}x) \frac{d^r}{dx^r} \Phi_{\sigma_n^2}(x) - n^{-1/2} P_1(-\Phi_{\sigma_n^2} : \{\bar{\chi}_{\nu, n}\})(x) \right| = o(n^{-1/2}), \quad (4.23)$$

where $P_r(-\Phi_{\sigma_n^2} : \{\bar{\chi}_{\nu, n}\})(x)$ is the $P_r(-\Phi_{\sigma_n^2} : \{\bar{\chi}_{\nu, n}\})$ -measure of the set $(-\infty, x]$.

Proof of Lemma 10. For any integer α , define $p_n(x) = \mathbf{P}(\sum_{i=1}^n X_i = \alpha)$ and $x_{\alpha, n} = n^{-1/2}(\alpha - n\mu_n)$. Also define $\tilde{X}_n = n^{-1/2} \sum_{i=1}^n (X_i - E(X_i))$ and $q_{n,3}(x) = n^{-1/2} \sum_{r=0}^1 n^{-r/2} \tilde{P}_r(-D : \{\bar{\chi}_{\nu, n}\})\phi_{\sigma_n^2}(x)$. Note that

$$\begin{aligned} & \sup_{x \in \mathcal{R}} \left| \mathbf{P}\left(n^{-1/2} \sum_{i=1}^n (X_i - E(X_i)) \leq x\right) - \sum_{r=0}^1 n^{-r/2} (-1)^r S_r(n\mu_n + n^{1/2}x) \frac{d^r}{dx^r} \Phi_{\sigma_n^2}(x) - n^{-1/2} P_1(-\Phi_{\sigma_n^2} : \{\bar{\chi}_{\nu, n}\})(x) \right| \\ & \leq \sup_{x \in \mathcal{R}} \left| \mathbf{P}(\tilde{X}_n \leq x) - Q_{n,3}(x) \right| + \sup_{x \in \mathcal{R}} \left| Q_{n,3}(x) - \sum_{r=0}^1 n^{-r/2} (-1)^r S_r(n\mu_n + n^{1/2}x) \frac{d^r}{dx^r} \Phi_{\sigma_n^2}(x) - n^{-1/2} P_1(-\Phi_{\sigma_n^2} : \{\bar{\chi}_{\nu, n}\})(x) \right| \\ & = J_1 + J_2 \quad (\text{say}), \end{aligned} \quad (4.24)$$

where $Q_{n,3}(x) = \sum_{\{\alpha: x_{\alpha,n} \leq x\}} q_{n,3}(x_{\alpha,n})$. Now the fact that $J_2 = o(n^{-1/2})$ follows from Theorem A.4.3 of BR(86) and dropping terms of order n^{-1} . Now we are going to show $J_1 = O(n^{-1})$. Note that

$$J_1 \leq \sum_{\alpha \in \Theta} |p_n(x_{\alpha,n}) - q_{n,3}(x_{\alpha,n})| = J_3 \quad (\text{say}),$$

where Θ has cardinality $\leq C_{33}n$, since $\mathbf{P}(|n^{-1} \sum_{i=1}^n X_i| \leq C_{33}) = 1$ for some constant $C_{33} > 0$, due to the assumption that $\max\{|x_j|^5 : j \in \{1, \dots, n\}\} = O(1)$. Hence $n^{-1}J_3 \leq C_{33} \sup_{\alpha \in \Theta} |p_n(x_{\alpha,n}) - q_{n,3}(x_{\alpha,n})| = C_{33} \sup_{\alpha \in \Theta} J_4(\alpha)$ (say). Hence enough to show $\sup_{\alpha \in \Theta} J_4(\alpha) = O(n^{-2})$. Now define $g_j(t) = \mathbf{E}(e^{itX_j})$ and $f_n(t) = \mathbf{E}(it\tilde{X}_n)$. Then we have

$$f_n(\sqrt{nt}) = \sum_{\alpha \in \Theta} p_n(x_{\alpha,n}) e^{i\sqrt{nt}x_{\alpha,n}}.$$

Hence by Fourier inversion formula for lattice random variables (cf. page 230 of BR(86)), we have

$$\begin{aligned} p_n(x_{\alpha,n}) &= (2\pi)^{-1} \int_{\mathcal{F}^*} e^{-i\sqrt{nt}x_{\alpha,n}} f_n(\sqrt{nt}) dt \\ &= (2\pi)^{-1} n^{-1/2} \int_{\sqrt{n}\mathcal{F}^*} e^{-itx_{\alpha,n}} f_n(t) dt, \end{aligned} \quad (4.25)$$

where $\mathcal{F}^* = (-\pi, \pi)$, the fundamental domain corresponding to the lattice distribution of $\sum_{i=1}^n X_i$. Again note that

$$q_{n,3}(x_{\alpha,n}) = (2\pi)^{-1} n^{-1/2} \int_{\mathcal{R}} e^{-itx_{\alpha,n}} \sum_{r=0}^1 n^{-r/2} \tilde{P}_r(it : \{\bar{\chi}_{\nu,n}\}) e^{-\sigma_n^2 t^2 / 2} dt. \quad (4.26)$$

Now defining the set $E = \{t \in \mathcal{R} : |t| \leq C_{31}(s)\sqrt{n} \min\{C_{33}^{-2}\sigma_n, C_{33}^{-5/3}\sigma_n^{5/3}\}\}$, from (4.25) & (4.26) we have

$$\begin{aligned} \sup_{\alpha \in \Theta} J_4(\alpha) &\leq (2\pi)^{-1} n^{-1/2} \left[\int_E |f_n(t) - \sum_{r=0}^1 n^{-r/2} \tilde{P}_r(it : \{\bar{\chi}_{\nu,n}\}) e^{-\sigma_n^2 t^2 / 2}| dt \right. \\ &\quad \left. + \int_{\sqrt{n}\mathcal{F}^* \cap E^c} |f_n(t)| dt + \int_{\mathcal{R} \cap (\sqrt{n}\mathcal{F}^*)^c} \left| \sum_{r=0}^1 n^{-r/2} \tilde{P}_r(it : \{\bar{\chi}_{\nu,n}\}) e^{-\sigma_n^2 t^2 / 2} \right| dt \right] \\ &= (2\pi)^{-1} n^{-1/2} (J_{41} + J_{42} + J_{43}) \quad (\text{say}). \end{aligned} \quad (4.27)$$

Note that $J_{41} = O(n^{-3/2})$ by applying Lemma 9 with $s = 5$. $J_{43} = O(n^{-3/2})$ due to the presence of the exponential term in the integrand and the form of the set E . Moreover noting the form of the set \mathcal{F}^* , we can say that there exists constants $C_{34} > 0$, $0 < C_{35}, C_{36} < \pi$ such that

$$J_{42} \leq C_{34} \sup_{t \in \sqrt{n}\mathcal{F}^* \cap E^c} \prod_{i=1}^n |g_j(n^{-1/2}t)| \leq C_{34} \sup_{C_{35} \leq |t| \leq C_{36}} |\mathbf{E}(e^{ity_{i_1}})|^m \leq C_{34} \delta^m, \quad (4.28)$$

for some $0 < \delta < 1$. Recall that $x_{i_j} = 1$ for all $j \in \{1, \dots, m\}$. The last inequality is due to the fact that there is no period of $\mathbf{E}(e^{ity_{i_1}})$ in the interval $[C_{35}, C_{36}] \cup [-C_{36}, -C_{35}]$. Now $J_{42} = O(n^{-3/2})$ follows from (4.28) since $m \geq (\log n)^2$. Therefore the proof is complete.

Lemma 11 Let $\check{W}_1, \dots, \check{W}_n$ be iid mean $\mathbf{0}$ non-degenerate random vectors in \mathcal{R}^l for some natural number l , with finite fourth absolute moment and $\limsup_{\|\mathbf{t}\| \rightarrow \infty} |\mathbf{E} e^{i\mathbf{t}'\check{W}_1}| < 1$ (i.e. Cramer's condition holds). Suppose $\tilde{W}_i = (\check{W}'_{i1}, \dots, \check{W}'_{im})'$ where \check{W}_{ij} is a random vector in \mathcal{R}^{l_j} and $\sum_{j=1}^m l_j = l$, m being a fixed natural number. Consider the sequence of random variables $\tilde{W}_1, \dots, \tilde{W}_n$ where $\tilde{W}_i = (c_{i1}\check{W}'_{i1}, \dots, c_{im}\check{W}'_{im})'$. $\{c_{ij} : i \in \{1, \dots, n\}, j \in \{1, \dots, m\}\}$ is a collection of real numbers such that for any $j \in \{1, \dots, m\}$, $\left\{n^{-1} \sum_{i=1}^n |c_{ij}|^4\right\} = O(1)$ and $\liminf_{n \rightarrow \infty} n^{-1} \sum_{i=1}^n c_{ij}^2 > 0$. Also assume that $\tilde{\mathbf{V}}_n = \text{Var}(\tilde{W}_i)$ converges to some positive definite matrix and $\bar{\chi}_{\nu, n}$ denotes the average ν th cumulant of $\tilde{W}_1, \dots, \tilde{W}_n$. Then we have

$$\sup_{\mathbf{B} \in \mathcal{A}_l} \left| \mathbf{P}\left(n^{-1/2} \sum_{i=1}^n \tilde{W}_i \in \mathbf{B}\right) - \int_{\mathbf{B}} \left[1 + n^{-1/2} \tilde{P}_r(-D : \{\bar{\chi}_{\nu, n}\})\right] \phi_{\tilde{\mathbf{V}}_n}(\mathbf{t}) d\mathbf{t} \right| = o(n^{-1/2}), \quad (4.29)$$

where the collection of sets \mathcal{A}_l is as defined in section 3.

Proof of Lemma 11. First note that $\tilde{W}_1, \dots, \tilde{W}_n$ is a sequence of independent random variables. Hence (4.29) follows by Theorem 20.6 of BR(86), provided there exists $\delta_4 \in (0, 1)$, independent of n , such that for all $\nu \leq \delta_4$,

$$n^{-1} \sum_{i=1}^n \mathbf{E} \|\tilde{W}_i\|^3 \mathbf{1}\left(\|\tilde{W}_i\| > \nu\sqrt{n}\right) = o(1) \quad (4.30)$$

and

$$\max_{|\alpha| \leq l+2} \int_{\|\mathbf{t}\| \geq \nu\sqrt{n}} \left| D^\alpha \mathbf{E} \exp(i\mathbf{t}' \mathbf{R}_{1n}^\dagger) \right| dt = o(n^{-1/2}) \quad (4.31)$$

where $\mathbf{R}_{1n}^\dagger = n^{-1/2} \sum_{i=1}^n (\mathbf{Z}_i - \mathbf{E}\mathbf{Z}_i)$ with

$$\mathbf{Z}_i = \tilde{W}_i \mathbf{1}\left(\|\tilde{W}_i\| \leq \nu\sqrt{n}\right).$$

First consider (4.30). Note that $\max\{|c_{ij}| : i \in \{1, \dots, n\}, j \in \{1, \dots, m\}\} = O(n^{1/4})$. Therefore, we have for any $\nu > 0$,

$$\begin{aligned} & n^{-1} \sum_{i=1}^n \mathbf{E} \|\tilde{W}_i\|^3 \mathbf{1}\left(\|\tilde{W}_i\| > \nu\sqrt{n}\right) \\ & \leq n^{-1} \sum_{i=1}^n \mathbf{E} \left(\sum_{j=1}^m c_{ij}^2 \|\check{W}_{ij}\|^2 \right)^{3/2} \mathbf{1}\left(\sum_{j=1}^m c_{ij}^2 \|\check{W}_{ij}\|^2 > \nu^2 n \right) \\ & \leq n^{-1} \sum_{i=1}^n \left(1 + \sum_{j=1}^m c_{ij}^2 \right)^2 \mathbf{E} \left[\|\check{W}_1\|^3 \mathbf{1}\left(\|\check{W}_1\|^2 > C_{37} \nu^2 n^{1/2}\right) \right] \\ & = o(1). \end{aligned}$$

Now consider (4.31). Note that for any $|\alpha| \leq l+2$, $|D^\alpha \mathbf{E} \exp(i\mathbf{t}' \mathbf{R}_{1n}^\dagger)|$ is bounded above by a sum of $n^{|\alpha|}$ -terms, each of which is bounded above by

$$C_{38}(\alpha) \cdot n^{-|\alpha|/2} \max\{\mathbf{E} \|\mathbf{Z}_i - \mathbf{E}\mathbf{Z}_i\|^{|\alpha|} : k \in \mathbf{I}_n\} \cdot \prod_{i \in \mathbf{I}_n^c} |\mathbf{E} \exp(i\mathbf{t}' \mathbf{Z}_i / \sqrt{n})| \quad (4.32)$$

where $\mathbf{I}_n \subset \{1, \dots, n\}$ is of size $|\alpha|$ and $\mathbf{I}_n^c = \{1, \dots, n\} \setminus \mathbf{I}_n$. Now for any $\omega > 0$ and $\mathbf{t} \in \mathcal{R}^{l_j}$, define the set

$$\mathbf{B}_n^{(j)}(\mathbf{t}, \omega) = \left\{ i : 1 \leq i \leq n \text{ and } |c_{ij}| \|\mathbf{t}\| > \omega \right\}.$$

Hence for any $\mathbf{t} \in \mathcal{R}^l$ writing $\mathbf{t} = (\mathbf{t}'_1, \dots, \mathbf{t}'_m)'$, \mathbf{t}_j is of length l_j , we have

$$\begin{aligned} & \sup \left\{ \prod_{i \in \mathbf{I}_n^c} |\mathbf{E} \exp(it' \mathbf{Z}_k / \sqrt{n})| : \|\mathbf{t}\| \geq v\sqrt{n} \right\} \\ &= \sup \left\{ \prod_{i \in \mathbf{I}_n^c} |\mathbf{E} \exp(it' \mathbf{Z}_k)| : \|\mathbf{t}\|^2 \geq v^2 \right\} \\ &\leq \max \left\{ \sup \left\{ \prod_{i \in \mathbf{I}_n^c \cap \mathbf{B}_n^{(j)}\left(\frac{\mathbf{t}_j}{\|\mathbf{t}_j\|}, v/\sqrt{2}\right)} \left[|\mathbf{E} \exp\left(ic_{ij} \mathbf{t}'_j \check{W}_{1j}\right)| + \mathbf{P}\left(\|\check{W}_1\| > C_{37} v^2 n^{1/2}\right) \right] \right. \right. \\ &\quad \left. \left. : \|\mathbf{t}_j\| \geq v/\sqrt{2} \right\} : j \in \{1, \dots, m\} \right\} \end{aligned}$$

Now since $|\mathbf{I}_n^c| \geq \left| \mathbf{I}_n^c \cap \mathbf{B}_n^{(j)}\left(\frac{\mathbf{t}_j}{\|\mathbf{t}_j\|}, v/\sqrt{2}\right) \right| \geq |\mathbf{B}_n^{(j)}\left(\frac{\mathbf{t}_j}{\|\mathbf{t}_j\|}, v/\sqrt{2}\right)| - |\alpha|$, due to Cramer's condition we have

$$\sup \left\{ \prod_{i \in \mathbf{I}_n^c \cap \mathbf{B}_n^{(j)}\left(\frac{\mathbf{t}_j}{\|\mathbf{t}_j\|}, v/\sqrt{2}\right)} \left[|\mathbf{E} \exp\left(ic_{ij} \mathbf{t}'_j \check{W}_{1j}\right)| + \mathbf{P}\left(\|\check{W}_1\| > C_{37} v^2 n^{1/2}\right) \right] : \|\mathbf{t}_j\| \geq v/\sqrt{2} \right\} \leq \theta \left| \mathbf{B}_n^{(j)}\left(\frac{\mathbf{t}_j}{\|\mathbf{t}_j\|}, v/\sqrt{2}\right) \right| - |\alpha| \quad (4.33)$$

Next note that $\liminf_{n \rightarrow \infty} n^{-1} \sum_{i=1}^n c_{ij}^2 > 0$ for all $j \in \{1, \dots, m\}$. Therefore for any $j \in \{1, \dots, m\}$, $\mathbf{u} \in \mathcal{R}^{l_j}$ with $|\mathbf{u}| = 1$, there exists $0 < \delta_5 < 1$ such that for sufficiently large n we have

$$\begin{aligned} \frac{n\delta_5}{2} &\leq \sum_{i=1}^n |uc_{ij}|^2 \\ &\leq \max \left\{ |c_{ij}|^2 : 1 \leq i \leq n \right\} \cdot |\mathbf{B}_n^{(j)}(\mathbf{u}, \omega)| + (n - |\mathbf{B}_n^{(j)}(\mathbf{u}, \omega)|) \cdot \omega^2 \\ &\leq C_{38} \cdot n^{1/2} \cdot |\mathbf{B}_n^{(j)}(\mathbf{u}, \omega)| + n\omega^2 \end{aligned}$$

which implies $|\mathbf{B}_n^{(j)}(\mathbf{u}, \omega)| \geq C_{39} \cdot n^{1/2}$ whenever $\omega < \sqrt{\delta_5/2}$. Therefore taking $\delta_4 = \sqrt{\delta_5/3}$, (4.31) follows from (4.32) and (4.33).

Proof of Theorem 1. Recall that the studentized pivot is

$$\tilde{\mathbf{H}}_n = \sqrt{n} \hat{\mathbf{L}}_n^{1/2} (\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta}),$$

where $\hat{\mathbf{L}}_n = n^{-1} \sum_{i=1}^n \mathbf{x}_i \mathbf{x}'_i e^{\mathbf{x}'_i \hat{\boldsymbol{\beta}}_n} (1 + e^{\mathbf{x}'_i \hat{\boldsymbol{\beta}}_n})^{-2}$. $\hat{\boldsymbol{\beta}}_n$ is the solution of (1.2). By Taylor's theorem, from (1.2) we have

$$\mathbf{L}_n(\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta}) = n^{-1} \sum_{i=1}^n (y_i - p(\boldsymbol{\beta}|\mathbf{x}_i)) \mathbf{x}_i - (2n)^{-1} \sum_{i=1}^n \mathbf{x}_i e^{z_i} (1 - e^{z_i})(1 + e^{z_i})^{-3} [\mathbf{x}'_i (\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta})]^2, \quad (4.34)$$

where $|z_i - \mathbf{x}'_i \boldsymbol{\beta}| \leq |\mathbf{x}'_i(\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta})|$ for all $i \in \{1, \dots, n\}$. Now due to the assumption $n^{-1} \sum_{i=1}^n \|\mathbf{x}_i\|^3 = O(1)$, by Lemma 1 (with $t = 3$) we have

$$\mathbf{P}\left(\left|n^{-1} \sum_{i=1}^n (y - p(\boldsymbol{\beta}|\mathbf{x}_i))x_{ij}\right| \leq C_{40}(p)n^{-1/2}(\log n)^{1/2}\right) = o(n^{-1/2}), \quad (4.35)$$

for any $j \in \{1, \dots, p\}$. Again by assumption \mathbf{L}_n converges to some positive definite matrix \mathbf{L} . Moreover,

$$\left\| (2n)^{-1} \sum_{i=1}^n \mathbf{x}_i e^{z_i} (1 - e^{z_i})(1 + e^{z_i})^{-3} [\mathbf{x}'_i(\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta})]^2 \right\| \leq \left(n^{-1} \sum_{i=1}^n \|\mathbf{x}_i\|^3 \right) \|\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta}\|^2.$$

Hence (4.34) can be rewritten as

$$(\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta}) = f_n(\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta}),$$

where f_n is a continuous function from \mathcal{R}^p to \mathcal{R}^p satisfying $\mathbf{P}\left(\|f_n(\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta})\| \leq C_{40}n^{-1/2}(\log n)^{1/2}\right) = 1 - o(n^{-1/2})$ whenever $\|(\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta})\| \leq C_{40}n^{-1/2}(\log n)^{1/2}$. Therefore, part (a) of Theorem 1 follows by Brouwer's fixed point theorem. Now we are going to prove part (b). Note that from (4.34) and the fact that \mathbf{L}_n converges to some positive definite matrix \mathbf{L} , we have for large enough n ,

$$\tilde{\mathbf{H}}_n = \hat{\mathbf{L}}_n^{1/2} [\mathbf{L}_n^{-1} A_n + R_{1n}]. \quad (4.36)$$

Here $A_n = n^{-1/2} \sum_{i=1}^n (y - p(\boldsymbol{\beta}|\mathbf{x}_i)) \mathbf{x}_i$ and $R_{1n} = -\mathbf{L}_n^{-1} \frac{1}{2\sqrt{n}} \sum_{i=1}^n \mathbf{x}_i e^{z_i} (1 - e^{z_i})(1 + e^{z_i})^{-3} [\mathbf{x}'_i(\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta})]^2$ with $|z_i - \mathbf{x}'_i \boldsymbol{\beta}| \leq |\mathbf{x}'_i(\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta})|$ for all $i \in \{1, \dots, n\}$. \mathbf{L}_n and $\hat{\mathbf{L}}_n$ are as defined earlier. Now applying part (a) we have $\mathbf{P}\left(\|R_{1n}\| = O\left(n^{-1/2}(\log n)\right)\right) = 1 - o\left(n^{-1/2}\right)$. Again by Taylor's theorem we have

$$\hat{\mathbf{L}}_n - \mathbf{L}_n = n^{-1} \sum_{i=1}^n \mathbf{x}_i \mathbf{x}'_i e^{\mathbf{x}'_i \boldsymbol{\beta}} (1 - e^{\mathbf{x}'_i \boldsymbol{\beta}}) (1 + e^{\mathbf{x}'_i \boldsymbol{\beta}})^{-3} [\mathbf{x}'_i(\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta})] + \mathbf{L}_{1n}, \quad (4.37)$$

where by part (a), we have $\mathbf{P}\left(\|\mathbf{L}_{1n}\| = O\left(n^{-1}(\log n)\right)\right) = 1 - o\left(n^{-1/2}\right)$. Hence using Lemma 6, part (a) and Taylor's theorem, one can show that $\mathbf{P}\left(\|\hat{\mathbf{L}}_n^{1/2} - \mathbf{L}_n^{1/2}\| = O\left(n^{-1/2}(\log n)^{1/2}\right)\right) = 1 - o\left(n^{-1/2}\right)$. Therefore (4.35) and (4.37) will imply that

$$\tilde{\mathbf{H}}_n = \mathbf{L}_n^{-1/2} A_n + R_{2n},$$

where $\mathbf{P}\left(\|R_{2n}\| = O\left(n^{-1/2}(\log n)\right)\right) = 1 - o\left(n^{-1/2}\right)$. Hence for any set $B \in \mathcal{A}_p$, there exists a constant $C_{41}(p) > 0$ such that

$$\begin{aligned} & \left| \mathbf{P}\left(\tilde{\mathbf{H}}_n \in B\right) - \Phi(B) \right| \\ & \leq \left| \mathbf{P}\left(\tilde{\mathbf{H}}_n \in B\right) - \mathbf{P}\left(\mathbf{L}_n^{-1/2} A_n \in B\right) \right| + \left| \mathbf{P}\left(\mathbf{L}_n^{-1/2} A_n \in B\right) - \Phi(B) \right| \\ & \leq \mathbf{P}\left(\|R_{2n}\| > C_{41}(p)n^{-1/2}(\log n)\right) + 2\mathbf{P}\left(\mathbf{L}_n^{-1/2} A_n \in (\partial B)^{C_{41}(p)n^{-1/2}(\log n)}\right) + \left| \mathbf{P}\left(\mathbf{L}_n^{-1/2} A_n \in B\right) - \Phi(B) \right| \\ & = O\left(n^{-1/2}(\log n)\right). \end{aligned}$$

The last equality is a consequence of Lemma 5 and the bound on $\|R_{2n}\|$. Therefore part (b) is proved.

Proof of Theorem 3. By applying Taylor's theorem, it follows from (2.1) that

$$\hat{\mathbf{L}}_n(\hat{\boldsymbol{\beta}}_n^* - \hat{\boldsymbol{\beta}}_n) = n^{-1} \sum_{i=1}^n (y - \hat{p}(\mathbf{x}_i)) \mathbf{x}_i - (2n)^{-1} \sum_{i=1}^n \mathbf{x}_i e^{z_i^*} (1 - e^{z_i^*}) (1 + e^{z_i^*})^{-3} [\mathbf{x}_i' (\hat{\boldsymbol{\beta}}_n^* - \hat{\boldsymbol{\beta}}_n)]^2, \quad (4.38)$$

where $|z_i^* - \mathbf{x}_i' \boldsymbol{\beta}| \leq |\mathbf{x}_i' (\hat{\boldsymbol{\beta}}_n^* - \hat{\boldsymbol{\beta}}_n)|$ for all $i \in \{1, \dots, n\}$. Now rest of part (a) of Theorem 3 follows exactly in the same line as the proof of part (a) of Theorem 1. To establish part (b), assume that $W_i = \left(Y_i \mathbf{x}_i', [Y_i^2 - \mathbf{E}Y_i^2] \mathbf{z}_i' \right)'$ and $W_i^* = \left(\hat{Y}_i [(G_i^* - \mu_{G^*}) \mu_{G^*}^{-1}] \mathbf{x}_i', \hat{Y}_i^2 [\mu_{G^*}^{-2} (G_i^* - \mu_{G^*})^2 - 1] \mathbf{z}_i' \right)'$. Here $Y_i = (y_i - p(\boldsymbol{\beta} | \mathbf{x}_i))$ and $\hat{Y}_i = (y_i - \hat{p}(\mathbf{x}_i))$. First we are going to show that

$$\check{\mathbf{H}}_n = \sqrt{n} \left(H(\bar{W}_n + n^{-1/2} b_n Z) \right) + R_n \quad \text{and} \quad \check{\mathbf{H}}_n^* = \sqrt{n} \left(\hat{H}(\bar{W}_n^* + n^{-1/2} b_n Z) \right) + R_n^*,$$

for some functions $H, \hat{H} : \mathcal{R}^k \rightarrow \mathcal{R}^p$ where $k = p + q$ with $q = \frac{p(p+1)}{2}$. $H(\cdot), \hat{H}(\cdot)$ have continuous partial derivatives of all orders, $H(\mathbf{0}) = \hat{H}(\mathbf{0}) = \mathbf{0}$ and $\mathbf{P}(\|R_n\| = o(n^{-1/2})) = 1 - o(n^{-1/2})$ & $\mathbf{P}_*(\|R_n^*\| = o(n^{-1/2})) = 1 - o_p(n^{-1/2})$. Next step is to apply Lemma 3, Lemma 4 and Lemma 7 to claim that suitable Edgeworth expansions exist for both $\check{\mathbf{H}}_n$ and $\check{\mathbf{H}}_n^*$. The last step is to conclude SOC of bootstrap by comparing the Edgeworth expansions. Now (4.34) and part (a) of Theorem 1 imply that

$$\sqrt{n}(\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta}) = \mathbf{L}_n^{-1} [A_n - \xi_n/2] + R_{3n}, \quad (4.39)$$

where $\mathbf{P}(\|R_{3n}\| \leq C_{42}(p)n^{-1}(\log n)^{3/2}) = 1 - o(n^{-1/2})$. Here $A_n = n^{-1/2} \sum_{i=1}^n Y_i \mathbf{x}_i$ and $\xi_n = n^{-3/2} \sum_{i=1}^n \mathbf{x}_i e^{\mathbf{x}_i' \boldsymbol{\beta}} (1 - e^{\mathbf{x}_i' \boldsymbol{\beta}}) (1 + e^{\mathbf{x}_i' \boldsymbol{\beta}})^{-3} [\mathbf{x}_i' (\mathbf{L}_n^{-1} A_n)]^2$. Clearly, $\mathbf{P}(\|\xi_n\| \leq C_{43}(p)n^{-1/2}(\log n)) = 1 - o(n^{-1/2})$. Therefore, by Taylor's theorem we have

$$\sqrt{n}(\hat{\mathbf{L}}_n - \mathbf{L}_n)(\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta}) = \xi_n + R_{4n}, \quad (4.40)$$

where $\mathbf{P}(\|R_{4n}\| \leq C_{44}(p)n^{-1}(\log n)^2) = 1 - o(n^{-1/2})$. Again noting (4.40), by equation (5) at page 52 of Turnbull (1929) we have

$$\hat{\mathbf{M}}_n^{-1/2} - \mathbf{L}_n^{-1/2} = -\mathbf{L}_n^{-1/2} \mathbf{Z}_{1n} \mathbf{L}_n^{-1/2} + \mathbf{Z}_{2n}, \quad (4.41)$$

where $(\hat{\mathbf{M}}_n - \mathbf{L}_n) = \mathbf{L}_n^{1/2} \mathbf{Z}_{1n} + \mathbf{Z}_{1n} \mathbf{L}_n^{1/2}$. Also easy to show that

$$\mathbf{P}(\|\hat{\mathbf{M}}_n - \mathbf{M}_n\| \leq C_{45}(p)n^{-1}(\log n)) = 1 - o(n^{-1/2}),$$

where $\mathbf{M}_n = n^{-1} \sum_{i=1}^n Y_i^2 \mathbf{x}_i \mathbf{x}_i'$. Hence using Lemma 6 we have $\mathbf{P}(\|\mathbf{Z}_{2n}\| \leq C_{46}(p)n^{-1}(\log n)^2) = 1 - o(n^{-1/2})$. Therefore from (4.39)-(4.41), Lemma 8 and the fact that $b_n = O(n^{-d})$ (for some $d > 0$) will imply that

$$\check{\mathbf{H}}_n = \mathbf{L}_n^{-1/2} [A_n + b_n Z + \xi_n/2] - \mathbf{L}_n^{-1/2} \left[\int_0^\infty e^{-t\mathbf{L}_n^{1/2}} (\mathbf{M}_n - \mathbf{L}_n) e^{-t\mathbf{L}_n^{1/2}} dt \right] \mathbf{L}_n^{-1/2} A_n + R_{5n}, \quad (4.42)$$

where $\mathbf{P}\left(\|R_{5n}\| \leq C_{47}(p)n^{-1/2}(\log n)^{-1}\right) = 1 - o(n^{-1/2})$. Now writing $W_i = (W'_{i1}, W'_{i2})'$ and $\bar{W}_n = n^{-1} \sum_{i=1}^n W_i = (\bar{W}'_{n1}, \bar{W}'_{n2})'$ with W_{i1} has first p components of W_i for all $i \in \{1, \dots, n\}$, we have

$$\begin{aligned} \Lambda_n + b_n Z &= \sqrt{n}(\bar{W}_{n1} + n^{-1/2}b_n Z) \\ \xi_n &= n^{-1/2} \sum_{i=1}^n \mathbf{x}_i e^{\mathbf{x}'_i \beta} (1 - e^{\mathbf{x}'_i \beta}) (1 + e^{\mathbf{x}'_i \beta})^{-3} \left[\bar{W}'_{n1} \mathbf{L}_n^{-1} \mathbf{x}_i \mathbf{x}'_i \mathbf{L}_n^{-1} \bar{W}_{n1} \right]^2 \\ &= \sqrt{n} \left(\bar{W}'_{n1} \check{\mathbf{M}}_1 \bar{W}_{n1}, \dots, \bar{W}'_{n1} \check{\mathbf{M}}_p \bar{W}_{n1} \right)', \end{aligned}$$

where $\check{\mathbf{M}}_k = n^{-1} \sum_{i=1}^n x_{ik} e^{\mathbf{x}'_i \beta} (1 - e^{\mathbf{x}'_i \beta}) (1 + e^{\mathbf{x}'_i \beta})^{-3} \left(\mathbf{L}_n^{-1} \mathbf{x}_i \mathbf{x}'_i \mathbf{L}_n^{-1} \right)$ for $k \in \{1, \dots, p\}$. Hence writing $\check{W}_{n1} = \bar{W}_{n1} + n^{-1/2}b_n Z$ we have

$$\mathbf{L}_n^{-1/2} \left[\Lambda_n + b_n Z + \xi_n / 2 \right] = \sqrt{n} \left[\mathbf{L}_n^{-1/2} \check{W}_n + \left(\check{W}'_{n1} \check{\mathbf{M}}_1 \check{W}_{n1}, \dots, \check{W}'_{n1} \check{\mathbf{M}}_p \check{W}_{n1} \right)' \right], \quad (4.43)$$

since $b_n = O(n^{-d})$ and $\|\check{\mathbf{M}}_k\| = O(1)$ for any $k \in \{1, \dots, p\}$. Here $\check{\mathbf{M}}_k = \sum_{j=1}^p L_{kjn}^{-1/2} \check{\mathbf{M}}_k$, $k \in \{1, \dots, p\}$, with $L_{kjn}^{-1/2}$ being the (k, j) th element of $\mathbf{L}_n^{-1/2}$. Again the j th row of $(\mathbf{M}_n - \mathbf{L}_n)$ is $\bar{W}'_{n2} \mathbf{E}_{jn}$ where \mathbf{E}_{jn} is a matrix of order $q \times p$ with $\|\mathbf{E}_{jn}\| \leq q$, $j \in \{1, \dots, p\}$. Therefore from (4.34) and (4.37) we have

$$\mathbf{L}_n^{-1/2} \left[\int_0^\infty e^{-t\mathbf{L}_n^{1/2}} (\mathbf{M}_n - \mathbf{L}_n) e^{-t\mathbf{L}_n^{1/2}} dt \right] \mathbf{L}_n^{-1/2} \Lambda_n = \sqrt{n} \left(\check{W}'_{n2} \check{\mathbf{M}}_1 \check{W}_{n2}, \dots, \check{W}'_{n2} \check{\mathbf{M}}_p \check{W}_{n2} \right)' + R_{6n}, \quad (4.44)$$

where $\check{W}_{n2} = \bar{W}_{n2} + n^{-1/2}b_n Z_1$ with $Z_1 \sim N_q(\mathbf{0}, \mathbf{I}_q)$, independent of Z & $\{y_1, \dots, y_n\}$. $\bar{\mathbf{M}}_k = \int_0^\infty \left[\sum_{j=1}^p m_{kjn}(t) \check{\mathbf{M}}_j(t) \right] dt$ where $m_{kjn}(t)$ is the (k, j) th element of the matrix $\mathbf{L}_n^{-1/2} e^{t\mathbf{L}_n^{1/2}}$ and $\check{\mathbf{M}}_j(t) = \mathbf{E}_{jn} e^{t\mathbf{L}_n^{1/2}} \mathbf{L}_n^{-1/2}$, $k, j \in \{1, \dots, p\}$. Moreover, $\mathbf{P}\left(\|R_{6n}\| \leq C_{48}n^{-1/2}(\log n)^{-1}\right) = 1 - o(n^{-1/2})$. Now define the $(p+q) \times (p+q)$ matrices $\{\mathbf{M}_1^\dagger, \dots, \mathbf{M}_p^\dagger\}$ where $\mathbf{M}_k^\dagger = \begin{bmatrix} \check{\mathbf{M}}_k & \mathbf{0} \\ \bar{\mathbf{M}}_k & \mathbf{0} \end{bmatrix}$.

Therefore from (4.42)-(4.44) we have

$$\begin{aligned} \check{\mathbf{H}}_n &= \sqrt{n} \left[\left(\mathbf{L}_n^{-1/2} \ \mathbf{0} \right) \check{W}_n + \left(\check{W}'_{n1} \mathbf{M}_1^\dagger \check{W}_{n1}, \dots, \check{W}'_{n1} \mathbf{M}_p^\dagger \check{W}_{n1} \right)' \right] + R_n \\ &= \sqrt{n} H(\check{W}_n) + R_n, \end{aligned} \quad (4.45)$$

where the function $H(\cdot)$ has continuous partial derivatives of all orders, $\check{W}_n = (\check{W}'_{n1}, \check{W}'_{n2})'$ and $R_n = R_{5n} + R_{6n}$.

Through the same line of arguments, writing $\bar{W}_{n1}^* = n^{-1} \sum_{i=1}^n W_{i1}^* = n^{-1} \sum_{i=1}^n \hat{Y}_i \mu_{G^*}^{-1} (G_i^* - \mu_{G^*}) \mathbf{x}_i$ and $\bar{W}_{n2}^* = n^{-1} \sum_{i=1}^n W_{i2}^* = n^{-1} \sum_{i=1}^n \hat{Y}_i^2 [\mu_{G^*}^{-2} (G_i^* - \mu_{G^*})^2 - 1] \mathbf{z}_i$, it can be shown that

$$\begin{aligned} \check{\mathbf{H}}_n^* &= \sqrt{n} \left[\left(\hat{\mathbf{M}}_n^{-1/2} \ \mathbf{0} \right) \check{W}_n^* + \left(\check{W}_n^{*'} \mathbf{M}_1^{*\dagger} \check{W}_n^*, \dots, \check{W}_n^{*'} \mathbf{M}_p^{*\dagger} \check{W}_n^* \right)' \right] + R_n^* \\ &= \sqrt{n} \hat{H}(\check{W}_n^*) + R_n^*, \end{aligned} \quad (4.46)$$

where $\check{W}_n^* = (\check{W}_n^{*'}, \check{W}_n^{*'})'$ with $\check{W}_{n1}^* = \bar{W}_{n1}^* + n^{-1/2}b_n Z^*$ and $\check{W}_{n2}^* = \bar{W}_{n2}^* + n^{-1/2}b_n Z_1^*$, Z_1^* being a $N_q(\mathbf{0}, \mathbf{I}_q)$ distributed random vector independent of $\{G_1^*, \dots, G_n^*\}$ and Z^* . $\mathbf{M}_j^{*\dagger} = \begin{bmatrix} \check{\mathbf{M}}_j^* & \mathbf{0} \\ \bar{\mathbf{M}}_j^* & \mathbf{0} \end{bmatrix}$

where $\check{\mathbf{M}}_j^* = \sum_{j=1}^p \hat{M}_{kjn}^{-1/2} \check{\mathbf{M}}_j^*$ with $\hat{M}_{kjn}^{-1/2}$ being the (k, j) th element of $\hat{\mathbf{M}}_n^{-1/2}$, $\check{\mathbf{M}}_j^*$ being same as $\check{\mathbf{M}}_j$ after replacing $\boldsymbol{\beta}$ by $\hat{\boldsymbol{\beta}}_n$. $\check{\mathbf{M}}_j^* = \int_0^\infty [\sum_{j=1}^p m_{kjn}^* \check{\mathbf{M}}_j(t)] dt$ where $m_{kjn}^*(t)$ is the (k, j) th element of the matrix $\hat{\mathbf{M}}_n^{-1/2} e^{-t\hat{\mathbf{M}}_n^{1/2}}$ and $\check{\mathbf{M}}_j^*(t) = \mathbf{E}_{jn} e^{-t\hat{\mathbf{M}}_n^{1/2}} \hat{\mathbf{M}}_n^{-1/2}$. Also $\mathbf{P}_*(\|\mathbf{R}_n^*\| \leq C_{49} n^{-1/2} (\log n)^{-1}) = 1 - o_p(n^{-1/2})$. Now by applying Lemma 3, Lemma 4 and Lemma 7 with $s = 3$, Edgeworth expansions of the densities of $\check{\mathbf{H}}_n$ and $\check{\mathbf{H}}_n^*$ can be found uniformly over the class \mathcal{A}_p upto an error $o(n^{-1/2})$ and $o_p(n^{-1/2})$ respectively. Call those Edgeworth expansions $\check{\psi}_{n,3}(\cdot)$ and $\check{\psi}_{n,3}^*(\cdot)$ respectively. Now if $\check{\psi}_{n,3}(\cdot)$ is compared with $\check{\psi}_{n,3}(\cdot)$ of Lemma 4, then $\check{\mathbf{M}}_n = I_p$. Similarly for $\check{\psi}_{n,3}^*(\cdot)$ also $\check{\mathbf{M}}_n = I_p$. Therefore, $\check{\psi}_{n,3}(\cdot)$ and $\check{\psi}_{n,3}^*(\cdot)$ have the forms

$$\begin{aligned} \check{\psi}_{n,3}(\mathbf{x}) &= \left[1 + n^{-1/2} q_1(\boldsymbol{\beta}, \mu_W, \mathbf{x}) + \sum_{j=1}^{m_2-1} b_n^{2j} q_{2j}(\boldsymbol{\beta}, \mathbf{L}_n, \mathbf{x}) \right] \phi(\mathbf{x}) \\ \check{\psi}_{n,3}^*(\mathbf{x}) &= \left[1 + n^{-1/2} q_1(\hat{\boldsymbol{\beta}}_n, \hat{\mu}_W, \mathbf{x}) + \sum_{j=1}^{m_2-1} b_n^{2j} q_{2j}(\hat{\boldsymbol{\beta}}_n, \hat{\mathbf{M}}_n, \mathbf{x}) \right] \phi(\mathbf{x}), \end{aligned}$$

where $m_2 = \inf\{j : b_n^{2j} = o(n^{-1/2})\}$, μ_W is the vector of $\{n^{-1} \sum_{i=1}^n \mathbf{E}(y_i - p(\boldsymbol{\beta}|\mathbf{x}_i))^2 x_{ij}^{l_1} x_{ij'}^{l_2} : j, j' \in \{1, \dots, p\}, l_1, l_2 \in \{0, 1, 2\}, l_1 + l_2 = 2\}$ and $\{n^{-1} \sum_{i=1}^n \mathbf{E}(y_i - p(\boldsymbol{\beta}|\mathbf{x}_i))^3 x_{ij}^{l_1} x_{ij'}^{l_2} x_{ij''}^{l_3} : j, j', j'' \in \{1, \dots, p\}, l_1, l_2, l_3 \in \{0, 1, 2, 3\}, l_1 + l_2 + l_3 = 3\}$. $\hat{\mu}_W$ is the vector of $\{n^{-1} \sum_{i=1}^n (y_i - \hat{p}(\mathbf{x}_i))^2 x_{ij}^{l_1} x_{ij'}^{l_2} : j, j' \in \{1, \dots, p\}, l_1, l_2 \in \{0, 1, 2\}, l_1 + l_2 = 2\}$ and $\{n^{-1} \sum_{i=1}^n (y_i - \hat{p}(\mathbf{x}_i))^3 x_{ij}^{l_1} x_{ij'}^{l_2} x_{ij''}^{l_3} : j, j', j'' \in \{1, \dots, p\}, l_1, l_2 \in \{0, 1, 2, 3\}, l_1 + l_2 + l_3 = 3\}$. $q_1(\mathbf{a}, \mathbf{b}, \mathbf{c})$ is a polynomial in \mathbf{c} whose coefficients are continuous functions of $(\mathbf{a}, \mathbf{b})'$. $q_{2j}(\mathbf{a}, \mathbf{b}, \mathbf{c})$ are polynomials in \mathbf{c} whose coefficients are continuous functions of \mathbf{a} and \mathbf{b} . Now Theorem 3 follows by comparing $\check{\psi}_{n,3}(\cdot)$ and $\check{\psi}_{n,3}^*(\cdot)$ and due to part (a) of Theorem 1.

Proof of Theorem 2. Recall that here $p = 1$ and hence $q = 1$. Define, $\mathbf{B}_n = \sqrt{n}H(\mathbf{E}_n \times \mathcal{R})$ with $\mathbf{E}_n = (-\infty, z_n]$ and $z_n = \left(\frac{3}{4n} - \mu_n\right)$. Here $\mu_n = n^{-1} \sum_{i=1}^n x_i p(\beta|x_i)$. Note that \mathbf{B}_n is an interval, as argued in section 3 just after the description of Theorem 2. The function $H(\cdot)$ is defined in (4.45). We are going to show that there exist a positive constant M_2 such that

$$\limsup_{n \rightarrow \infty} \mathbf{P}\left(\sqrt{n} \left| \mathbf{P}_*(\mathbf{H}_n^* \in \mathbf{B}_n) - \mathbf{P}(\mathbf{H}_n \in \mathbf{B}_n) \right| \geq M_3\right) = 1.$$

Define the set $\mathbf{Q}_n = \left\{ |\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta}| = o(n^{-1/2}(\log n)) \right\} \cap \left\{ \left| n^{-1} \sum_{i=1}^n [(y_i - p(\beta|x_i))^2 - \mathbf{E}(y_i - p(\beta|x_i))^2] x_i^2 \right| = o(n^{-1/2}(\log n)) \right\} \cap \left\{ \left| n^{-1} \sum_{i=1}^n [(y_i - p(\beta|x_i))^3 - (y_i - p(\beta|x_i))^3] x_i^3 \right| = o(1) \right\}$. Now due to a stronger version of (4.35), it is easy to see that $\mathbf{P}\left(|\hat{\boldsymbol{\beta}}_n - \boldsymbol{\beta}| = o(n^{-1/2}(\log n))\right) = 1$ for all but finitely many n , upon application of Borel-cantelli lemma and noting that $\max\{|x_i| : i \in \{1, \dots, n\}\} = O(1)$. Again by applying Lemma 1, it is easy to show that $\mathbf{P}\left(\left\{ \left| n^{-1} \sum_{i=1}^n [(y_i - p(\beta|x_i))^2 - (y_i - p(\beta|x_i))^2] \right| = o(n^{-1/2}(\log n)) \right\} \cap \left\{ \left| n^{-1} \sum_{i=1}^n [(y_i - p(\beta|x_i))^3 - (y_i - p(\beta|x_i))^3] \right| = o(1) \right\}\right) = 1$ for large enough n . Hence $\mathbf{P}(\mathbf{Q}_n) = 1$ for large enough n . Similarly define the bootstrap version of \mathbf{Q}_n as $\mathbf{Q}_n^* = \left\{ |\hat{\boldsymbol{\beta}}_n^* - \hat{\boldsymbol{\beta}}_n| = o(n^{-1/2}(\log n)) \right\} \cap \left\{ \left| n^{-1} \sum_{i=1}^n [(y_i - \hat{p}(x_i))^2 (\mu_{G^*}^{-2}(G_i^* - \mu_{G^*})^2 - 1)] x_i^2 \right| = o(n^{-1/2}(\log n)) \right\} \cap \left\{ \left| n^{-1} \sum_{i=1}^n [(y_i - \hat{p}(x_i))^3 (\mu_{G^*}^{-3}(G_i^* - \mu_{G^*})^3 - 1)] x_i^3 \right| = o(1) \right\}$. Through the same line, it is easy to establish that $\mathbf{P}(\mathbf{P}^*(\mathbf{Q}_n^*) = 1) = 1$ for large enough n . Hence

enough to show

$$\limsup_{n \rightarrow \infty} \mathbf{P} \left(\left\{ \sqrt{n} \left| \mathbf{P}_* \left(\{\mathbf{H}_n^* \in \mathbf{B}_n\} \cap \mathbf{Q}_n^* \right) - \mathbf{P} \left(\{\mathbf{H}_n \in \mathbf{B}_n\} \cap \mathbf{Q}_n \right) \right| \geq M_2 \right\} \cap \mathbf{Q}_n \right) = 1. \quad (4.47)$$

Recall the definitions of \bar{W}_n and \bar{W}_n^* from the proof of Theorem 3. Similar to (4.45) and (4.46), it is easy to observe that

$$\mathbf{H}_n = \sqrt{n}H(\bar{W}_n) + R_n \quad \text{and} \quad \mathbf{H}_n^* = \sqrt{n}\hat{H}(\bar{W}_n^*) + R_n^*, \quad (4.48)$$

where $\{|R_n| = O(n^{-1/2}(\log n)^{-1})\} \subseteq \mathbf{Q}_n$ and $\{|R_n^*| = O(n^{-1/2}(\log n)^{-1})\} \subseteq \mathbf{Q}_n^*$. To prove (4.47), first we are going to show for large enough n ,

$$\begin{aligned} & \left\{ \left\{ \sqrt{n} \left| \mathbf{P}_* \left(\{\mathbf{H}_n^* \in \mathbf{B}_n\} \cap \mathbf{Q}_n^* \right) - \mathbf{P} \left(\{\mathbf{H}_n \in \mathbf{B}_n\} \cap \mathbf{Q}_n \right) \right| \geq M_2 \right\} \cap \mathbf{Q}_n \right\} \\ & \supseteq \left\{ \left\{ \sqrt{n} \left| \mathbf{P}_* \left(\{\sqrt{n}\hat{H}(\bar{W}_n^*) \in \mathbf{B}_n\} \cap \mathbf{Q}_n^* \right) - \mathbf{P} \left(\{\sqrt{n}H(\bar{W}_n) \in \mathbf{B}_n\} \cap \mathbf{Q}_n \right) \right| \geq 2M_2 \right\} \cap \mathbf{Q}_n \right\}. \end{aligned} \quad (4.49)$$

Now due to (4.48), we have

$$\begin{aligned} & \left| \mathbf{P} \left(\mathbf{H}_n \in \mathbf{B}_n \right) - \mathbf{P} \left(\sqrt{n}H(\bar{W}_n) \in \mathbf{B}_n \right) \right| \leq \mathbf{P} \left(\sqrt{n}H(\bar{W}_n) \in (\partial \mathbf{B}_n)^{(n \log n)^{-1/2}} \right) \\ & \quad + \mathbf{P} \left(|R_n| \neq o(n^{-1/2}(\log n)^{-1}) \right) \\ & \left| \mathbf{P}_* \left(\mathbf{H}_n^* \in \mathbf{B}_n \right) - \mathbf{P}_* \left(\sqrt{n}\hat{H}(\bar{W}_n^*) \in \mathbf{B}_n \right) \right| \leq \mathbf{P}_* \left(\sqrt{n}\hat{H}(\bar{W}_n^*) \in (\partial \mathbf{B}_n)^{(n \log n)^{-1/2}} \right) \\ & \quad + \mathbf{P}_* \left(|R_n^*| \neq o(n^{-1/2}(\log n)^{-1}) \right) \end{aligned}$$

To establish (4.49), enough to show $\mathbf{P} \left(\sqrt{n}\hat{H}(\bar{W}_n) \in (\partial \mathbf{B}_n)^{(n \log n)^{-1/2}} \right) = o(n^{-1/2})$ and $\mathbf{P} \left(\left\{ \mathbf{P}_* \left(\sqrt{n}\hat{H}(\bar{W}_n^*) \in (\partial \mathbf{B}_n)^{(n \log n)^{-1/2}} \right) = o(n^{-1/2}) \right\} \cap \mathbf{Q}_n \right) = 1$ for large enough n . An Edgeworth expansion of $\sqrt{n}\bar{W}_n^*$ with an error $o(n^{-1/2})$ (in almost sure sense) can be established using Lemma 11. Then we can use transformation technique of Bhattacharya and Ghosh (1978) to find an Edgeworth expansion $\hat{\eta}_n(\cdot)$ of the density of $\sqrt{n}\hat{H}(\bar{W}_n^*)$ with an error $o(n^{-1/2})$ (in almost sure sense). Now the calculations similar to page 213 of BR(86) will imply that $\mathbf{P} \left(\left\{ \mathbf{P}_* \left(\sqrt{n}\hat{H}(\bar{W}_n^*) \in (\partial \mathbf{B}_n)^{(n \log n)^{-1/2}} \right) = o(n^{-1/2}) \right\} \cap \mathbf{Q}_n \right) = 1$, since \mathbf{B}_n is an interval. Next we are going to show that $\mathbf{P} \left(\sqrt{n}\hat{H}(\bar{W}_n) \in (\partial \mathbf{B}_n)^{(n \log n)^{-1/2}} \right) = 0$ for large enough n and to show that we need to utilize the form of \mathbf{B}_n , as Edgeworth expansion of $\sqrt{n}H(\bar{W}_n)$ similar to $\sqrt{n}\hat{H}(\bar{W}_n^*)$ does not exist due to the lattice nature of W_1^*, \dots, W_n^* . To this end define $k_n(\mathbf{x}) = (\sqrt{n}H(\mathbf{x}/\sqrt{n}), x_2)'$ where $\mathbf{x} = (x_1, x_2)'$. Note that $k_n(\cdot)$ is a diffeomorphism (cf. proof of lemma 3.2 in Lahiri (1989)). Hence $k_n(\cdot)$ is bijection and $k_n(\cdot)$ & $k_n^{-1}(\cdot)$ have derivatives of all orders. Therefore, arguments given between (2.15) and (2.18) at page 444 of Bhattacharya and Ghosh (1978) with g_n replaced by $k_n^{-1}(\cdot)$ will imply that

$$\begin{aligned} & \left| \mathbf{P} \left(\mathbf{H}_n \in \mathbf{B}_n \right) - \mathbf{P} \left(\sqrt{n}H(\bar{W}_n) \in \mathbf{B}_n \right) \right| \leq \mathbf{P} \left((\sqrt{n}\bar{W}_n \in (\partial k_n^{-1}(\mathbf{B}_n \times \mathcal{R}))^{d_n(n \log n)^{-1/2}}) \right) + o(n^{-1/2}) \\ & \quad = \mathbf{P} \left(\sqrt{n}\bar{W}_{n1} \in (\partial \mathbf{E}_n)^{d_n(n \log n)^{-1/2}} \right) + o(n^{-1/2}), \end{aligned}$$

where $d_n \leq \max \{ |\det(\text{Grad}[k_n(x)])|^{-1} : |x| = O(\sqrt{\log n}) \}$. Now by looking into the form of $H(\cdot)$ in (4.35), it is easy to see that $d_n = O(1)$, say $d_n \leq C_{44}$ for some positive constant C_{44} . Now note that

$$\begin{aligned} & \mathbf{P}\left(\sqrt{n}\bar{W}_{n1} \in (\partial\mathbf{E}_n)^{C_{44}(n \log n)^{-1/2}}\right) \\ &= \mathbf{P}\left(\left[n^{-1/2} \sum_{i=1}^n y_i x_i - \sqrt{n}\mu_n\right] \in (z_n - C_{44}(n \log n)^{-1/2}, z_n + C_{44}(n \log n)^{-1/2})\right) \\ &= \mathbf{P}\left(\sum_{i=1}^n y_i x_i \in (3/4 - C_{44}(\log n)^{-1/2}, 3/4 + C_{44}(\log n)^{-1/2})\right) \\ &= 0, \end{aligned}$$

for large enough n , since $\sum_{i=1}^n y_i x_i$ can take only integer values. Therefore (4.49) is established. Now recalling that $\hat{\eta}_n(\cdot)$ is the Egeworth expansion of the density of $\sqrt{n}\hat{H}(\bar{W}_n^*)$ with an almost sure error $o(n^{-1/2})$, we have for large enough n ,

$$\mathbf{P}\left(\sqrt{n}\left|\mathbf{P}_*\left(\sqrt{n}\hat{H}(\bar{W}_n^*) \in \mathbf{B}_n\right) - \int_{\mathbf{B}_n} \hat{\eta}_n(x)dx\right| = o(1)\right) = 1. \quad (4.50)$$

Now define $U_i = \left((y_i - p(\beta|x_i))x_i V_i, (y_i - p(\beta|x_i))^2 x_i^2 [V_i^2 - 1]\right)'$, $i \in \{1, \dots, n\}$, where V_1, \dots, V_n are iid continuous random variables which are independent of $\{y_1, \dots, y_n\}$. Also $\mathbf{E}(V_1) = 0$, $\mathbf{E}(V_1^2) = \mathbf{E}(V_1^3) = 1$ and $\mathbf{E}V_1^8 < \infty$. An immediate choice of the distribution of V_1 is that of $(G_1^* - \mu_{G^*})\mu_{G^*}^{-1}$. Other choices of $\{V_1, \dots, V_n\}$ can be found in Liu(1988), Mammen (1993) and Das et al. (2019). Now since $\max\{|x_i| : i \in \{1, \dots, n\}\} = O(1)$, there exists a natural number n_0 and constants $0 < \delta_2 \leq \delta_1 < 1$ such that $\sup_{n \geq n_0} p(\beta|x_n) \leq \delta_1$ and $\inf_{n \geq n_0} p(\beta|x_n) \geq \delta_2$. Again V_1, \dots, V_n are iid continuous random variables. Hence writing $p_n = p(\beta|x_n)$, for any $b > 0$ we have

$$\begin{aligned} & \sup_{n \geq n_0} \sup_{\|\mathbf{t}\| > b} \left| \mathbf{E}e^{i\mathbf{t}'U_n} \right| \\ & \leq \sup_{n \geq n_0} \left[p_n \sup_{\|\mathbf{t}\| > b(1-\delta_1)^2} \left| \mathbf{E}e^{it_1(1-p_n)V_1 + it_2(-p_n)^2[V_1^2-1]} \right| + (1-p_n) \sup_{\|\mathbf{t}\| > b\delta_2^2} \left| \mathbf{E}e^{it_1(-p_n)V_1 + it_2(-p_n)^2[V_1^2-1]} \right| \right] \\ & < 1, \end{aligned}$$

i.e. uniform Cramer's condition holds. Also the minimum eigen value condition of Theorem 20.6 of BR(86) holds due to $\max\{|x_i| : i \in \{1, \dots, n\}\} = O(1)$ and $\liminf_{n \rightarrow \infty} n^{-1} \sum_{i=1}^n x_i^6 > 0$. Hence applying Theorem 20.6 of BR(86) and then applying transformation technique of Bhattacharya and Ghosh (1978) we have

$$\left| \mathbf{P}\left(\sqrt{n}H(\bar{U}_n) \in \mathbf{B}_n\right) - \int_{\mathbf{B}_n} \eta_n(x)dx \right| = o(n^{-1/2}), \quad (4.51)$$

where $\bar{U}_n = n^{-1} \sum_{i=1}^n U_i$. Note that in both the expansions $\eta_n(\cdot)$ and $\hat{\eta}_n(\cdot)$ the variances corresponding to normal terms are 1. Also $\hat{H}(\cdot)$ can be obtained from $H(\cdot)$ first replacing \mathbf{L}_n by \hat{M}_n and then β by $\hat{\beta}_n$ (cf. (4.45) and (4.46)). Hence we can conclude that for any Borel set \mathbf{C} ,

$$\mathbf{P}\left(\left\{\sqrt{n}\left|\int_{\mathbf{C}} \eta_n(x)dx - \int_{\mathbf{C}} \hat{\eta}_n(x)dx\right| = o(1)\right\} \cap \mathbf{Q}_n\right) = 1$$

Hence from (4.50) and (4.51), we have

$$\mathbf{P}\left(\left\{\sqrt{n}\left|\mathbf{P}_*\left(\sqrt{n}\hat{H}(\bar{W}_n^*) \in \mathbf{B}_n\right) - \mathbf{P}\left(\sqrt{n}H(\bar{U}_n) \in \mathbf{B}_n\right)\right| = o(1)\right\} \cap \mathbf{Q}_n\right) = 1, \quad (4.52)$$

for large enough n . To establish (4.47), in view of (4.49) and (4.52) it is enough to find a positive constant M_3 such that

$$\sqrt{n} \left| \mathbf{P} \left(\sqrt{n} H(\bar{W}_n) \in \mathbf{B}_n \right) - \mathbf{P} \left(\sqrt{n} H(\bar{U}_n) \in \mathbf{B}_n \right) \right| = \sqrt{n} \left| \mathbf{P} \left(\sqrt{n} \bar{W}_{n1} \in \mathbf{E}_n \right) - \mathbf{P} \left(\sqrt{n} \bar{U}_{n1} \in \mathbf{E}_n \right) \right| \geq 4M_3.$$

Note that since $\mathbf{E}V_1^2 = \mathbf{E}V_i^3 = 1$, the first three average moments of $\{W_{11}, \dots, W_{n1}\}$ are same as that of $\{U_{11}, \dots, U_{n1}\}$. However $\{W_{11}, \dots, W_{n1}\}$ are independent lattice random variables whereas $\{U_{11}, \dots, U_{n1}\}$ are independent random variables for which uniform Cramer's condition holds. Therefore by Lemma 10 and Theorem 20.6 of BR(86) we have

$$\begin{aligned} & \sup_{x \in \mathcal{R}} \left| \mathbf{P} \left(\sqrt{n} \bar{W}_{n1} \leq x \right) - \Phi_{\sigma_n^2}(x) - n^{-1/2} P_1 \left(-\Phi_{\sigma_n^2} : \{\bar{\chi}_{\nu, n}\} \right)(x) \right. \\ & \quad \left. + n^{-1/2} (n\mu_n + \sqrt{n}x - 1/2) \frac{d}{dx} \Phi_{\sigma_n^2}(x) \right| = o(n^{-1/2}) \\ \text{and } & \sup_{x \in \mathcal{R}} \left| \mathbf{P} \left(\sqrt{n} \bar{U}_{n1} \leq x \right) - \Phi_{\sigma_n^2}(x) - n^{-1/2} P_1 \left(-\Phi_{\sigma_n^2} : \{\bar{\chi}_{\nu, n}\} \right)(x) \right| = o(n^{-1/2}), \end{aligned} \quad (4.53)$$

where $P_1 \left(-\Phi_{\sigma_n^2} : \{\bar{\chi}_{\nu, n}\} \right)(x)$ is as defined in Lemma 10. Recall that $\mathbf{E}_n = (-\infty, z_n]$ where $z_n = \left(\frac{3}{4n} - \mu_n \right)$. Therefore for some positive constants C_{46}, C_{47}, C_{48} we have

$$\begin{aligned} \sqrt{n} \left| \mathbf{P} \left(\sqrt{n} \bar{W}_{n1} \in \mathbf{E}_n \right) - \mathbf{P} \left(\sqrt{n} \bar{U}_{n1} \in \mathbf{E}_n \right) \right| &= \sqrt{n} \left| \mathbf{P} \left(\sqrt{n} \bar{W}_{n1} \leq \sqrt{n} z_n \right) - \mathbf{P} \left(\sqrt{n} \bar{U}_{n1} \leq \sqrt{n} z_n \right) \right| \\ &\geq (n\mu_n + nz_n - 1/2) (\sqrt{2\pi}\sigma_n)^{-1} e^{-(nz_n^2)/(2\sigma_n^2)} - o(1) \\ &= (4\sqrt{2\pi}\sigma_n)^{-1} e^{-(nz_n^2)/(2\sigma_n^2)} - o(1) \\ &\geq C_{46} \exp \left\{ -C_{47} n^{-1} \left(\frac{9}{16} + n^2 \mu_n^2 - \frac{3n\mu_n}{2} \right) \right\} - o(1) \\ &\geq C_{48} \exp \left\{ -C_{47} M_1^2 \right\}. \end{aligned}$$

The first inequality follows due to (4.53). Second one is due to $\max\{|x_i| : i \in \{1, \dots, n\}\} = O(1)$ and the last one is due to the assumption $\sqrt{n}|\mu_n| < M_1$. Taking $4M_2 = C_{48} \exp \left\{ -C_{47} M_1^2 \right\}$, the proof of Theorem 2 is now complete.

References

1. BARBE, P. and BERTAIL, P. (2012). *The weighted bootstrap*. Lecture Notes in Statistics, 98.
2. BERKSON, J. (1944). Application of the Logistic Function to Bio-Assay. *Journal of the American Statistical Association*. **39** 357-365.
3. BHATIA, R. (1996). *Matrix Analysis*. Springer.
4. BHATTACHARYA, R. N. and GHOSH, J. K. (1978). On the validity of the formal Edgeworth expansion. *Ann. Statist.* **6** 434-451.
5. BHATTACHARYA, R. N. and RANGA RAO, R. (1986). *Normal approximation and asymptotic expansions*. John Wiley & Sons.
6. BIRNBAUM, Z. W. (1942). An Inequality for Mill's Ratio. *Ann. Math. Statist.* **13**. 245-246.
7. CHATTERJEE, S. and BOSE, A. (2005). Generalized bootstrap for estimating equations. *Ann. Statist.* **33** 414-436.
8. CLAESKENS, G., AERTS, M. and MOLENBERGHS, G. (2003). A quadratic bootstrap method and improved estimation in logistic regression. *Statistics and Probability Letters*. **61** 383-394.
9. COX, D. R. (1944). Some Problems Connected with Statistical Inference. *Annals of Mathematical Statistics*. **29** 357-372.

10. DAVISON, A.C., HINKLEY, D.V. and SCHECHTMAN, E. (1986). Efficient bootstrap simulations. *Biometrika*. **73** 555—566.
11. DAS, D. and LAHIRI S. N. (2019). Second Order Correctness of Perturbation Bootstrap M-Estimator of Multiple Linear Regression Parameter. *Bernoulli*. **25** 654-682.
12. FAHRMEIR, L. and KAUFMANN, H. (1985). Consistency and Asymptotic Normality of the Maximum Likelihood Estimator in Generalized Linear Models. *Annals of Statistics*. **13**. 342–368.
13. FREDMAN, D. A. (1981). Bootstrapping Regression Models. *Ann. Statist.* **9** 1218–1228.
14. FUK, D. H. and NAGAEV, S. V. (1971). Probabilistic inequalities for sums of independent random variables. *Teor. Veroyatnost. i Primenen.* **16** 660-675.
15. GHOSH, J. K. (1994). *Higher Order Asymptotics*. NSF-CBMS Regional Conference Series in Probability and Statistics Vol. 4.
16. GOURIEROUX, C. and MONFORT, A. (1981). Asymptotic properties of the maximum likelihood estimator in dichotomous logit models. *Journal of Econometrics*. **17** 83–97.
17. HABERMAN, S. J. (1974). Log-Linear Models for Frequency Tables Derived by Indirect Observation: Maximum Likelihood Equations. *Annals of Statistics*. **2** 911–924.
18. LAHIRI, S. N. (1989) *Bootstrap approximations to the distributions of m-estimators*. Thesis.
19. LAHIRI, S. N. (1992). Bootstrapping M-estimators of a multiple linear regression parameter. *Ann. Statist.* **20** 1548-1570.
20. LAHIRI S. N. (1993). Bootstrapping the Studentized Sample Mean of Lattice Variables. *Journal of Multivariate Analysis*. **45** 247–256.
21. LAHIRI, S. N. (1994). On two-term Edgeworth expansions and bootstrap approximations for Studentized multivariate M-estimators. *Sankhya A*. **56** 201-226.
22. LEE, K. W. (1990). Bootstrapping logistic regression models with random regressors, *Comm. Statist.* **19** 2527—2539.
23. LIU, R. Y. (1988). Bootstrap Procedures under some Non-IID Models. . *Ann. Statist.* **16** 1696–1708.
24. MAMMEN E. (1993). Bootstrap and Wild Bootstrap for High Dimensional Linear Models. *Ann. Statist.* **21** 255–285.
25. MCFADDEN, D. (1974). Conditional logit analysis of qualitative choice behavior. *Frontiers in Econometrics*. *Academic Press*. 105–142.
26. MOULTON, L. H., and ZEGER, S. L. (1989). Analyzing repeated measures on generalized linear models via the bootstrap. *Biometrics* **45**, 381–394.
27. MOULTON, L. H., and ZEGER, S. L. (1991). Bootstrapping generalized linear models. *Computational Statistics and Data Analysis*. **11** 53–63.
28. TURNBULL, H. W. (1930). A Matrix Form of Taylor’s Theorem. *Proc. Edinburgh Math. Soc.* **33** 33-54.