

Analytic solution of system of singular nonlinear differential equations with Neumann-Robin boundary conditions arising in astrophysics

Randhir Singh*

* Department of Mathematics, Birla Institute of Technology Mesra, Ranchi-835215, India.

Abstract

In this paper, we propose a new approach for the approximate analytic solution of system of Lane-Emden-Fowler type equations with Neumann-Robin boundary conditions. The algorithm is based on Green's function and the homotopy analysis method. This approach depends on constructing Green's function before establishing the recursive scheme for the approximate analytic solution of the equivalent system of integral equations. Unlike Adomian decomposition method (ADM) [1], the present method contains adjustable parameters to control the convergence of the approximate series solution. Convergence and error estimation of the present is provided under quite general conditions. Several examples are considered to demonstrate the accuracy of the current algorithm. Computational results reveal that the proposed approach produces better results as compared to some existing iterative methods.

Keyword: System of Lane-Emden equations; Catalytic diffusion reactions; Green's function; Adomian decomposition method; homotopy analysis method.

1 Introduction

We consider the following system of singular boundary value problems (SBVPs)

$$\begin{cases} (p_1(x)y_1'(x))' = p_1(x)f_1(x, y_1(x), y_2(x)), & x \in (0, 1), \\ (p_2(x)y_2'(x))' = p_2(x)f_2(x, y_1(x), y_2(x)), \\ y_1'(0) = 0, \quad a_1y_1(1) + b_1y_1'(1) = c_1, \quad y_2'(0) = 0, \quad a_2y_2(1) + b_2y_2'(1) = c_2, \end{cases} \quad (1.1)$$

where $a_1, a_2, b_1, b_2, c_1, c_2$ are real constants. Here, $p_1(x) = x^{k_1}g_1(x)$, $p_2(x) = x^{k_2}g_2(x)$, $g_1(0) \neq 0$ and $g_2(0) \neq 0$ with $p_1(0) = p_2(0) = 0$.

*Corresponding author. E-mail: randhir.math@gmail.com,

We next consider the following system of Lane-Emden equations [1–7] a particular case of (1.1) with $p_1(x) = x^{k_1}$ and $p_2(x) = x^{k_2}$ as

$$\begin{cases} (x^{k_1} y_1'(x))' = x^{k_1} f_1(x, y_1(x), y_2(x)), & x \in (0, 1), \\ (x^{k_2} y_2'(x))' = x^{k_2} f_2(x, y_1(x), y_2(x)), \\ y_1'(0) = 0, \quad a_1 y_1(1) + b_1 y_1'(1) = c_1, \quad y_2'(0) = 0, \quad a_2 y_2(1) + b_2 y_2'(1) = c_2. \end{cases} \quad (1.2)$$

In recent years, the SBVPs have been studied extensively in [8–30] and references therein. But as far as we know, we find only the following results on system of Lane-Emden equations. Recently, in [5, 31, 32] authors studied (1.2) with boundary conditions $y_1'(0) = y_2'(0) = 0$, $y_1(1) = y_2(1) = 1$ and shape factors $k_1 = k_2 = 2$ and $f_1 = -b + a g_1(y_1, y_2) + c g_2(y_1, y_2)$, $f_2 = d g_1(y_1, y_2) + e g_2(y_1, y_2)$, that relates the concentration of the carbon substrate and the concentration of oxygen, the Michaelis-Menten functions

$$g_i(y_1, y_2) = M_1 \times M_2, \quad M_1 = \frac{y_1}{l_i + y_1}, \quad M_2 = \frac{y_2}{m_i + y_2}, \quad i = 1, 2,$$

where M_1 and M_2 are the respective Michaelis-Menten nonlinear operators.

In [3, 4, 33], authors considered the system of Lane-Emden equations (1.2) with boundary conditions $y_1'(0) = y_2'(0) = 0$, $y_1(1) = 1$, $y_2(1) = 2$ and shape factors $k_1 = k_2 = 2$ and $f_1 = a y_1^2 + b y_1 y_2$, $f_2 = c y_1^2 - d y_1 y_2$, occurs in catalytic diffusion reactions with the parameters a, b, c, d are actual chemical reactions.

In [4, 32], the ADM was applied to obtain a convergent analytic approximate solution of (1.2) with $k_1 = k_2 = 2$. Later, in [34], the variational iteration method (VIM) was applied to obtain approximations to solutions of (1.2) for shape factors $k_i = 1, 2, 3$, $i = 1, 2$. Most recently, a numerical procedure based on sinc-collocation method was developed in [35] to obtain the solution of (1.2). In [36] authors used the reproducing kernel Hilbert space method for solving to obtain the solution of (1.2). In [1], the ADM with Green's function used to find numerical approximation of the solutions of (1.2).

The homotopy analysis method was developed and improved by S. Liao [37–40] for solving a broad class of functional equations. Various modifications of the homotopy analysis method have also been elaborated, for example, the optimal homotopy asymptotic method was proposed by Marinca and Herisanu [41, 42], the optimal homotopy analysis method was introduced in [43, 44]. Recently, the homotopy analysis method with Green's function was applied to solve singular boundary value problems in [45–49].

In this work, we present the homotopy analysis method combined with the Green's function strategy for obtaining approximate solutions of coupled singular boundary value problems (1.1) and (1.2). Unlike standard HAM, our approach avoids solving the transcendental equations for the undetermined coefficients. Unlike the ADM and VIM, our proposed technique contains the convergence-control parameters which gives fast convergence of the series solution. Numerical results reveal that the present process provides better results as compared to some existing iterative methods [1, 4, 5].

2 Integral form of system of Lane-Emden-Fowler types equations

By following [20], we transform the following system of Lane-Emden-Fowler types equations with Neumann-Robin boundary conditions

$$\begin{cases} (p_1(x)y_1'(x))' = p_1(x)f_1(x, y_1(x), y_2(x)), & x \in (0, 1), \\ (p_2(x)y_2'(x))' = p_2(x)f_2(x, y_1(x), y_2(x)), \\ y_1'(0) = 0, \quad a_1y_1(1) + b_1y_1'(1) = c_1, \quad y_2'(0) = 0, \quad a_2y_2(1) + b_2y_2'(1) = c_2, \end{cases} \quad (2.1)$$

into the equivalent system of integral equations

$$\begin{cases} y_1(x) = \frac{c_1}{a_1} + \int_0^1 G_1(x, s) p_1(s) f_1(s, y_1(s), y_2(s)) ds, \\ y_2(x) = \frac{c_2}{a_2} + \int_0^1 G_2(x, s) p_2(s) f_2(s, y_1(s), y_2(s)) ds, \end{cases} \quad (2.2)$$

where $G_1(x, s)$ and $G_2(x, s)$ are given by

$$G_1(x, s) = \begin{cases} h_1(1) - h_1(x) + \frac{b_1}{a_1}h_1'(1), & s \leq x, \\ h_1(1) - h_1(s) + \frac{b_1}{a_1}h_1'(1), & x \leq s, \end{cases} \quad (2.3)$$

with $h_1(x) = \int_0^x \frac{dt}{p_1(t)}$, $h_1'(1) = \frac{1}{p_1(1)}$, $h_1(1) - h_1(s) = \int_s^1 \frac{dt}{p_1(t)}$, $h_1(1) - h_1(x) = \int_s^1 \frac{dt}{p_1(t)} - \int_s^x \frac{dt}{p_1(t)}$,

$$G_2(x, s) = \begin{cases} h_2(1) - h_2(x) + \frac{b_2}{a_2}h_2'(1), & s \leq x, \\ h_2(1) - h_2(s) + \frac{b_2}{a_2}h_2'(1), & x \leq s, \end{cases} \quad (2.4)$$

with $h_2(x) = \int_0^x \frac{dt}{p_2(t)}$, $h_2'(1) = \frac{1}{p_2(1)}$, $h_2(1) - h_2(s) = \int_s^1 \frac{dt}{p_2(t)}$, $h_2(1) - h_2(x) = \int_s^1 \frac{dt}{p_2(t)} - \int_s^x \frac{dt}{p_2(t)}$.

Similarly, by following [1, 17], we transform the following system of Lane-Emden types equations with Neumann-Robin boundary conditions

$$\begin{cases} (x^{k_1}y_1'(x))' = x^{k_1}f_1(x, y_1(x), y_2(x)), & x \in (0, 1), \\ (x^{k_2}y_2'(x))' = x^{k_2}f_2(x, y_1(x), y_2(x)), \\ y_1'(0) = 0, \quad a_1 y_1(1) + b_1 y_1'(1) = c_1, \quad y_2'(0) = 0, \quad a_2 y_2(1) + b_2 y_2'(1) = c_2, \end{cases} \quad (2.5)$$

into the equivalent system of integral equations

$$\begin{cases} y_1(t) = \frac{c_1}{a_1} + \int_0^1 G_1(x, s) s^{k_1} f_1(s, y_1(s), y_2(s)) ds, \\ y_2(x) = \frac{c_2}{a_2} + \int_0^1 G_2(x, s) s^{k_2} f_2(s, y_1(s), y_2(s)) ds, \end{cases} \quad (2.6)$$

where $G_1(x, s)$ and $G_2(x, s)$ for $k_1 = k_2 = 1$ are

$$G_1(x, s) = \begin{cases} \ln x - \frac{b_1}{a_1}, & s \leq x, \\ \ln s - \frac{b_1}{a_1}, & x \leq s, \end{cases} \quad (2.7)$$

$$G_2(x, s) = \begin{cases} \ln x - \frac{b_2}{a_2}, & s \leq x, \\ \ln s - \frac{b_2}{a_2}, & x \leq s \end{cases} \quad (2.8)$$

and $G_1(x, s)$ and $G_2(x, s)$ for $k_1 \neq 1, k_2 \neq 1$ are

$$G_1(x, s) = \begin{cases} \frac{x^{1-k_1} - 1}{1 - k_1} - \frac{b_1}{a_1}, & s \leq x \\ \frac{s^{1-k_1} - 1}{1 - k_1} - \frac{b_1}{a_1}, & x \leq s, \end{cases} \quad (2.9)$$

$$G_2(x, s) = \begin{cases} \frac{x^{1-k_2} - 1}{1 - k_2} - \frac{b_2}{a_2}, & s \leq x, \\ \frac{s^{1-k_2} - 1}{1 - k_2} - \frac{b_2}{a_2}, & x \leq s. \end{cases} \quad (2.10)$$

3 Solution method

We consider the system of integral operator equation (2.2) as

$$\begin{cases} N_1[y_1] = y_1(x) - \frac{c_1}{a_1} - \int_0^1 G_1(x, s) p_1(s) f_1(s, y_1(s), y_2(s)) ds = 0, \\ N_2[y_2] = y_2(x) - \frac{c_2}{a_2} - \int_0^1 G_2(x, s) p_2(s) f_2(s, y_1(s), y_2(s)) ds = 0. \end{cases} \quad (3.1)$$

Note that the above system of integral equations reduce to (2.6) with $p_1(x) = x^{k_1}$ and $p_2(x) = x^{k_2}$.

In order to apply the HAM with Green's function [45–49], we construct the zeroth-order deformation equations for (3.1) as

$$\begin{cases} (1-q)[\varphi_1(x, q) - y_{10}(x)] = q \, c_{10} \, N_1[\varphi_1(x, q)], \\ (1-q)[\varphi_2(x, q) - y_{20}(x)] = q \, c_{20} \, N_2[\varphi_2(x, q)], \quad q \in [0, 1] \end{cases} \quad (3.2)$$

where q is an embedding parameter, $(y_{10}(x), y_{20}(x))$ are initial approximations, $(c_{10} \neq 0, c_{20} \neq 0)$ are convergence control parameters and $(\varphi_1(x, q), \varphi_2(x, q))$ are unknown functions.

If $q = 0$ and $q = 1$ the homotopy equations (3.2) vary from $(\varphi_1(x, 0), \varphi_2(x, 0)) = (y_{10}(x), y_{20}(x)) \rightarrow (\varphi_1(x, 1), \varphi_2(x, 1)) = (y_1(x), y_2(x))$. Now as q varies from 0 to 1, the solutions of (3.1) will vary from the initial guesses $(y_{10}(x), y_{20}(x))$ to the exact solutions $(y_1(x), y_2(x))$ of equations (3.1).

Expanding $\varphi_1(x, q)$ and $\varphi_2(x, q)$ as a Taylor series with respect to q yields

$$\begin{cases} \varphi_1(x, q) = \sum_{k=0}^{\infty} y_{1k}(x) q^k, \\ \varphi_2(x, q) = \sum_{k=0}^{\infty} y_{2k}(x) q^k, \end{cases} \quad (3.3)$$

where

$$y_{1k}(x) = \frac{1}{k!} \frac{\partial^k \varphi_1(x, q)}{\partial q^k} \Big|_{q=0}, \quad y_{2k}(x) = \frac{1}{k!} \frac{\partial^k \varphi_2(x, q)}{\partial q^k} \Big|_{q=0}. \quad (3.4)$$

The series (3.3) converge for $q = 1$ if $c_{10} \neq 0$, $c_{20} \neq 0$ are chosen properly, and they take the form

$$\begin{cases} \varphi_1(x, 1) \equiv y_1(x) = \sum_{k=0}^{\infty} y_{1k}(x), \\ \varphi_2(x, 1) \equiv y_2(x) = \sum_{k=0}^{\infty} y_{2k}(x), \end{cases} \quad (3.5)$$

which will be the solutions of (3.1).

For further analysis, we define the vectors as

$$\begin{cases} \vec{y}_{1k} = \{y_{10}(x), y_{11}(x), \dots, y_{1k}(x)\}, \\ \vec{y}_{2k} = \{y_{20}(x), y_{21}(x), \dots, y_{2k}(x)\}. \end{cases} \quad (3.6)$$

Differentiating (3.2) k -times with respect to q , dividing them by $k!$, setting $q = 0$, we obtain the k th-order deformation equations as

$$\begin{cases} y_{1k}(x) - \chi_k \, y_{1(k-1)}(x) = c_{10} \, R_{1k}[\vec{y}_{1(k-1)}], \\ y_{2k}(x) - \chi_k \, y_{2(k-1)}(x) = c_{20} \, R_{2k}[\vec{y}_{2(k-1)}], \end{cases} \quad (3.7)$$

where $\chi_k = \begin{cases} 0, & k \leq 1, \\ 1, & k > 1, \end{cases}$ and

$$\begin{cases} R_{1k}[\vec{y}_{1(k-1)}] = \frac{1}{(k-1)!} \frac{\partial^{k-1}}{\partial q^{k-1}} \left[\varphi_1(x, q) - \frac{c_1}{a_1} - \int_0^1 G_1(x, s) p_1(s) f_1\left(s, \varphi_1(s, q), \varphi_2(s, q)\right) ds \right] \Big|_{q=0} \\ R_{2k}[\vec{x}_{2(k-1)}] = \frac{1}{(k-1)!} \frac{\partial^{k-1}}{\partial q^{k-1}} \left[\varphi_2(x, q) - \frac{c_2}{a_2} - \int_0^1 G_2(x, s) p_2(s) f_2\left(s, \varphi_1(s, q), \varphi_2(s, q)\right) ds \right] \Big|_{q=0} \end{cases}$$

$$\begin{cases} = \frac{1}{(k-1)!} \frac{\partial^{k-1}}{\partial q^{k-1}} \left[\sum_{i=0}^{\infty} y_{1i} q^i - \frac{c_1}{a_1} - \int_0^1 G_1(x, s) p_1(s) f_1\left(s, \sum_{i=0}^{\infty} y_{1i} q^i, \sum_{i=0}^{\infty} y_{2i} q^i\right) ds \right] \Big|_{q=0}, \\ = \frac{1}{(k-1)!} \frac{\partial^{k-1}}{\partial q^{k-1}} \left[\sum_{i=0}^{\infty} y_{2i} q^i - \frac{c_2}{a_2} - \int_0^1 G_2(x, s) p_2(s) f_2\left(s, \sum_{i=0}^{\infty} y_{1i} q^i, \sum_{i=0}^{\infty} y_{2i} q^i\right) ds \right] \Big|_{q=0}. \end{cases}$$

On simplification we get

$$\begin{cases} R_{1k}[\vec{y}_{1(k-1)}] = y_{1(k-1)} - \left(1 - \chi_k\right) \frac{c_1}{a_1} - \int_0^1 G_1(x, s) p_1(s) H_{1(k-1)} ds, \\ R_{2k}[\vec{y}_{2(k-1)}] = y_{2(k-1)} - \left(1 - \chi_k\right) \frac{c_2}{a_2} - \int_0^1 G_2(x, s) p_2(s) H_{2(k-1)} ds, \end{cases} \quad (3.8)$$

where H_{1k} and H_{2k} are given by

$$\begin{cases} H_{1k} = \frac{1}{k!} \frac{\partial^k}{\partial q^k} f_1\left(s, \sum_{i=0}^{\infty} y_{1i} q^i, \sum_{i=0}^{\infty} y_{2i} q^i\right) \Big|_{q=0}, \\ H_{2k} = \frac{1}{k!} \frac{\partial^k}{\partial q^k} f_2\left(s, \sum_{i=0}^{\infty} y_{1i} q^i, \sum_{i=0}^{\infty} y_{2i} q^i\right) \Big|_{q=0}. \end{cases} \quad (3.9)$$

Making use of (3.8), the k th-order deformation equations (3.7) are simplified as

$$\begin{cases} y_{1k} - \chi_k y_{1(k-1)} = c_{10} \left[y_{1(k-1)} - \left(1 - \chi_k\right) \frac{c_1}{a_1} - \int_0^1 G_1(x, s) p_1(s) H_{1(k-1)} ds \right], \\ y_{2k} - \chi_k y_{2(k-1)} = c_{20} \left[y_{2(k-1)} - \left(1 - \chi_k\right) \frac{c_2}{a_2} - \int_0^1 G_2(x, s) p_2(s) H_{2(k-1)} ds \right]. \end{cases} \quad (3.10)$$

By taking the initial approximations $y_{10} = \frac{c_1}{a_1}$ and $y_{20} = \frac{c_2}{a_2}$, the solutions components are

computed as

$$\left\{ \begin{array}{l} y_{11} = c_{10} y_{10} - c_{10} \frac{c_1}{a_1} - c_{10} \int_0^1 G_1(x, s) p_1(s) H_{10} ds, \\ y_{21} = c_{20} y_{20} - c_{20} \frac{c_2}{a_2} - c_{20} \int_0^1 G_2(x, s) p_2(s) H_{20} ds, \\ \vdots \\ y_{1k} = (1 + c_{10}) y_{1(k-1)} - c_{10} \int_0^1 G_1(x, s) p_1(s) H_{1(k-1)} ds, \\ y_{2k} = (1 + c_{20}) y_{2(k-1)} - c_{20} \int_0^1 G_2(x, s) p_2(s) H_{2(k-1)} ds, \quad k \geq 2. \end{array} \right. \quad (3.11)$$

The n th-order approximations to solutions are obtained as

$$\left\{ \begin{array}{l} \phi_{1n}(x, c_{10}, c_{20}) = \sum_{k=0}^n y_{1k}(x, c_{10}, c_{20}), \\ \phi_{2n}(x, c_{10}, c_{20}) = \sum_{k=0}^n y_{2k}(x, c_{10}, c_{20}). \end{array} \right. \quad (3.12)$$

The unknown parameters c_{10}, c_{20} have a significant influence on the convergence of the approximate solution. We next find the optimal values of c_{10}, c_{20} by solving

$$\frac{\partial E_{1n}(c_{10}, c_{20})}{\partial c_{10}} = 0, \quad \frac{\partial E_{2n}(c_{10}, c_{20})}{\partial c_{20}} = 0, \quad (3.13)$$

where E_{1n} and E_{2n} are given by

$$\left\{ \begin{array}{l} E_{1n}(c_{10}, c_{20}) = \frac{1}{N} \sum_{k=1}^N \left(N_1 \left[\phi_{1n}(x_k, c_{10}, c_{20}) \right] \right)^2, \\ E_{2n}(c_{10}, c_{20}) = \frac{1}{N} \sum_{k=1}^N \left(N_2 \left[\phi_{2n}(x_k, c_{10}, c_{20}) \right] \right)^2, \quad x_1, x_2, \dots, x_N \in [0, 1], \end{array} \right. \quad (3.14)$$

Then these optimal values of c_{10} and c_{20} will be substituting in (3.12) to obtain the optimal HAM-approximate solution

$$\left\{ \begin{array}{l} \phi_{1n}(x) = \sum_{k=0}^n y_{1k}(x), \\ \phi_{2n}(x) = \sum_{k=0}^n y_{2k}(x). \end{array} \right. \quad (3.15)$$

Remark 3.1. The ADM [1, 32] is a particular case of (3.11) when $c_{10} = c_{20} = -1$ and given by

$$\begin{cases} y_{10} = \frac{c_1}{a_1}, & y_{20} = \frac{c_2}{a_2}, \\ y_{1k} = \int_0^1 G_1(x, s) p_1(s) H_{1(k-1)} ds, \\ y_{2k} = \int_0^1 G_2(x, s) p_2(s) H_{2(k-1)} ds, & k \geq 1. \end{cases} \quad (3.16)$$

Hence, the ADM-approximate solutions are given by

$$\begin{cases} \psi_{1n}(x) = \sum_{k=0}^n y_{1k}(x), \\ \psi_{2n}(x) = \sum_{k=0}^n y_{2k}(x). \end{cases} \quad (3.17)$$

4 Convergence analysis

Let $E = (C[0, 1], \|y\|)$ be a Banach space with norm

$$\|y\| = \max\{\|y_1\|, \|y_2\|\}, \quad y \in E, \quad (4.1)$$

where, $\|y_1\| = \max_{x \in I=[0,1]} |y_1(x)|$ and $\|y_2\| = \max_{x \in I} |y_2(x)|$. We next discuss the convergence and error analysis of the proposed method. To do so, we first introduce the vector notations

$$\begin{aligned} y &= \begin{pmatrix} y_1 \\ y_2 \end{pmatrix}, \quad \mathbf{y}_k = \begin{pmatrix} y_{1k} \\ y_{2k} \end{pmatrix}, \quad c_0 = \begin{pmatrix} c_{10} \\ c_{20} \end{pmatrix}, \quad p(x) = \begin{pmatrix} p_1(x) \\ p_2(x) \end{pmatrix}, \\ G(x, s) &= \begin{pmatrix} G_1(x, s) \\ G_2(x, s) \end{pmatrix}, \quad H_k = \begin{pmatrix} H_{1k} \\ H_{2k} \end{pmatrix}, \quad f(x, y_1, y_2) = \begin{pmatrix} f_1(x, y_1, y_2) \\ f_2(x, y_1, y_2) \end{pmatrix}, \\ \phi_n &= \begin{pmatrix} \phi_{1n} \\ \phi_{2n} \end{pmatrix} \text{ and } M = \max \left(\begin{array}{c} \max_{x \in I} \left| \int_0^1 G_1(x, s) p_1(s) ds \right| \\ \max_{x \in I} \left| \int_0^1 G_2(x, s) p_2(s) ds \right| \end{array} \right). \end{aligned}$$

Theorem 4.1. *Suppose that the nonlinear function $f(x, z, w)$ satisfy Lipschitz condition: $|f(x, z, w) - f(x, z^*, w^*)| \leq l_1|z - z^*| + l_2|w - w^*|$, where l_1 and l_2 are Lipschitz constants. If the parameter c_0 is chosen such that there exists constant $\delta_{c_0} \in (0, 1)$, then the series ϕ_n defined by (3.15) is convergent in Banach space E .*

Proof. From (3.10) and (3.12), we have

$$\begin{aligned}\phi_n &= \sum_{k=0}^n \mathbf{y}_k(x) = (1 + c_0) \sum_{k=1}^n \mathbf{y}_{k-1} - c_0 \sum_{k=1}^n \left[\int_0^1 G(x, s) p(s) H_{k-1} ds \right] \\ &= (1 + c_0) \phi_{n-1} - c_0 \int_0^1 G(x, s) p(s) \sum_{k=1}^n H_{k-1} ds.\end{aligned}\tag{4.2}$$

From (4.2), we consider for $n > m$, for all $n, m \in \mathbb{N}$ as

$$\|\phi_n - \phi_m\| = \max_{x \in I} \left| (1 + c_0)(\phi_{n-1} - \phi_{m-1}) - c_0 \int_0^1 G(x, s) p(s) \left(\sum_{k=1}^n H_{k-1} - \sum_{k=1}^m H_{k-1} \right) ds \right|.$$

Using the inequality $\sum_{k=0}^n H_k \leq f(s, \phi_{1n}, \phi_{2n})$ from ([50]) we have

$$\begin{aligned}\|\phi_n - \phi_m\| &\leq |1 + c_0| \max_{x \in I} |\phi_{n-1} - \phi_{m-1}| + |c_0| \\ &\quad \max_{x \in I} \left| \int_0^1 G(x, s) p(s) \left(f(s, \phi_{1(n-1)}, \phi_{2(n-1)}) - f(s, \phi_{1(m-1)}, \phi_{2(m-1)}) \right) ds \right|.\end{aligned}$$

Applying the Lipschitz condition, we get

$$\begin{aligned}\|\phi_n - \phi_m\| &\leq |1 + c_0| \|\phi_{n-1} - \phi_{m-1}\| + |c_0| \max_{x \in I} \left| \int_0^1 G(x, s) p(s) ds \right| \times \sum_{i=1}^2 l_i \max_{t \in I} |\phi_{i(n-1)} - \phi_{i(m-1)}| \\ &\leq |1 + c_0| \|\phi_{n-1} - \phi_{m-1}\| + |c_0| \max_{x \in I} \left| \int_0^1 G(x, s) p(s) ds \right| \\ &\quad \times 2L \max \left\{ \left\| \phi_{1(n-1)} - \phi_{1(m-1)} \right\|, \left\| \phi_{2(n-1)} - \phi_{2(m-1)} \right\| \right\} \\ &\leq |1 + c_0| \left\| \phi_{n-1} - \phi_{m-1} \right\| + 2LM|c_0| \left\| \phi_{n-1} - \phi_{m-1} \right\| \\ &\leq \left(|1 + c_0| + 2LM|c_0| \right) \|\phi_{n-1} - \phi_{m-1}\| = \delta_{c_0} \|\phi_{n-1} - \phi_{m-1}\|,\end{aligned}$$

Thus, we have

$$\|\phi_n - \phi_m\| \leq \delta_{c_0} \|\phi_{n-1} - \phi_{m-1}\|.\tag{4.3}$$

where $L = \max\{l_1, l_2\}$, $\delta_{c_0} = (|1 + c_0| + 2LM|c_0|)$.

Setting $n = m + 1$ in (4.3), we have

$$\|\phi_{m+1} - \phi_m\| \leq \delta_{c_0} \|\phi_m - \phi_{m-1}\| \leq \delta_{c_0}^2 \|\phi_{m-1} - \phi_{m-2}\| \leq \dots \leq \delta_{c_0}^m \|\phi_1 - \phi_0\|.$$

For all $n, m \in \mathbb{N}$, with $n > m$, consider

$$\begin{aligned}
\|\phi_n - \phi_m\| &= \|(\phi_n - \phi_{n-1}) + (\phi_{n-1} - \phi_{n-2}) + \cdots + (\phi_{m+1} - \phi_m)\| \\
&\leq \|\phi_n - \phi_{n-1}\| + \|\phi_{n-1} - \phi_{n-2}\| + \cdots + \|\phi_{m+1} - \phi_m\| \\
&\leq [\delta_{c_0}^{n-1} + \delta_{c_0}^{n-2} + \cdots + \delta_{c_0}^m] \|\phi_1 - \phi_0\| \\
&= \delta_{c_0}^m [1 + \delta_{c_0} + \delta_{c_0}^2 + \cdots + \delta_{c_0}^{n-m-1}] \|\phi_1 - \phi_0\| = \delta_{c_0}^m \left(\frac{1 - \delta_{c_0}^{n-m}}{1 - \delta_{c_0}} \right) \|\mathbf{y}_1\|.
\end{aligned}$$

Since $\delta_{c_0} < 1$, we have $(1 - \delta_{c_0}^{n-m}) < 1$. It readily follows that

$$\|\phi_n - \phi_m\| \leq \frac{\delta_{c_0}^m}{1 - \delta_{c_0}} \|\mathbf{y}_1\|. \quad (4.4)$$

Letting $n, m \rightarrow \infty$, we obtain

$$\lim_{n, m \rightarrow \infty} \|\phi_n - \phi_m\| = 0.$$

Therefore, $\{\phi_n\}$ is a Cauchy sequences in the Banach space E . \square

Theorem 4.2. If the approximate solution ϕ_n converges to \mathbf{y} , then the maximum absolute truncated error is given by

$$\|\mathbf{y} - \phi_m\| \leq \frac{\delta_{c_0}^m M |c_0|}{1 - \delta_{c_0}} \max_{x \in I} |f(x, y_{10}, y_{20})|. \quad (4.5)$$

Proof. Following inequality (4.4), we have

$$\|\phi_n - \phi_m\| \leq \frac{\delta_{c_0}^m}{1 - \delta_{c_0}} \|\mathbf{y}_1\|.$$

For $n \geq m$, as $n \rightarrow \infty$ then $\phi_n \rightarrow \mathbf{y}(x)$, the above inequality becomes

$$\|\mathbf{y}(x) - \phi_m\| \leq \frac{\delta_{c_0}^m}{1 - \delta_{c_0}} \|\mathbf{y}_1\|. \quad (4.6)$$

From (3.11), we have $\mathbf{y}_1 = c_0 \left(\mathbf{y}_0 - \frac{c}{a} - \int_0^1 G(x, s) p(s) H_0 ds \right)$, where $\mathbf{y}_0 = \begin{pmatrix} y_{10} \\ y_{20} \end{pmatrix}$, $\frac{c}{a} = \begin{pmatrix} \frac{c_1}{a_1} \\ \frac{c_2}{a_2} \end{pmatrix}$, $H_0 = f(x, y_{10}, y_{20})$. Now using $\mathbf{y}_0 = \frac{c}{a}$, we find

$$\|\mathbf{y}_1\| = |c_0| \max_{x \in I} \left| \frac{c}{a} - \frac{c}{a} - \int_0^1 G(x, s) p(s) H_0 ds \right| \leq |c_0| M \max_{x \in I} |f(x, y_{10}, y_{20})|. \quad (4.7)$$

Combining (4.6) and (4.7), we obtain error estimate as

$$\|\mathbf{y} - \phi_m\| \leq \frac{\delta_{c_0}^m |c_0| M}{1 - \delta_{c_0}} \max_{x \in I} |f(x, y_{10}, y_{20})|. \quad (4.8)$$

which completes the proof. \square

Remark 4.1. Let us consider the parameter c_0 , with $\delta_{c_0} < 1$, so that

$$|1 + c_0| + 2LM|c_0| < 1 \Rightarrow 2LM < \frac{1 - |1 + c_0|}{|c_0|}, \quad c_0 \neq 0,$$

From the right hand side of this inequality, we have

$$\frac{1 - |1 + c_0|}{|c_0|} = \begin{cases} -1 - \frac{2}{c_0}, & c_0 < -1, \\ 1, & c_0 \in [-1, 0) \\ -1, & c_0 > 0 \end{cases}$$

We can choose value of parameter $c_0 \in [-1, 0)$.

5 Numerical results and discussion

In order to compare our numerical results with the results obtained by the ADM [32], we define the absolute residual error as

$$Res_{1n} = \left| \frac{1}{p_1(x)} \left(p_1(x) \phi'_{1n}(x) \right)' - f_1(x, \phi_{1n}(x), \phi_{2n}(x)) \right|, \quad (5.1)$$

$$Res_{2n} = \left| \frac{1}{p_2(x)} \left(p_2(x) \phi'_{2n}(x) \right)' - f_2(x, \phi_{1n}(x), \phi_{2n}(x)) \right|, \quad (5.2)$$

$$res_{1n} = \left| \frac{1}{p_1(x)} \left(p_1(x) \psi'_{1n}(x) \right)' - f_1(x, \psi_{1n}(x), \psi_{2n}(x)) \right|, \quad (5.3)$$

$$res_{2n} = \left| \frac{1}{p_2(x)} \left(p_2(x) \psi'_{2n}(x) \right)' - f_2(x, \psi_{1n}(x), \psi_{2n}(x)) \right|, \quad (5.4)$$

where $(\phi_{1n}(x), \phi_{2n}(x))$ are the HAM approximate solutions and $(\psi_{1n}(x), \psi_{2n}(x))$ are the ADM approximate solutions.

In Tables 1– 8, we compare the numerical results of $(\phi_{1n}(x), \phi_{2n}(x))$, $(\psi_{1n}(x), \psi_{2n}(x))$, (Res_{1n}, Res_{2n}) , and (res_{1n}, res_{2n}) , obtained by the HAM and the ADM [32]. From the comparison, we observe that the convergence of the proposed method is very fast and the only the first few terms of the series provide an excellent approximation to the exact solution. The optimal choice of the convergence control parameters permits to obtain an excellent approximation.

Example 5.1. Consider the system of Lane-Emden-Fowler equations that relates the concentration of the carbon substrate and the concentration of oxygen [1, 4, 5]

$$\begin{cases} (x^{k_1} y_1')' = x^{k_1} \left(-b + \frac{a y_1 y_2}{(l_1 + y_1)(m_1 + y_2)} + \frac{c y_1 y_2}{(l_2 + y_1)(m_2 + y_2)} \right), & x \in (0, 1), \\ (x^{k_2} y_2')' = x^{k_2} \left(\frac{d y_1 y_2}{(l_1 + y_1)(m_1 + y_2)} + \frac{e y_1 y_2}{(l_2 + y_1)(m_2 + y_2)} \right), \\ y_1'(0) = 0, \quad y_2'(0) = 0 \quad y_1(1) = 1, \quad y_2(1) = 1, \end{cases} \quad (5.5)$$

where the functions $y_1(x)$ and $y_2(x)$ are the concentration of the carbon substrate and the concentration of oxygen, respectively. We fix the parameters $l_1 = l_2 = m_1 = m_2 = \frac{1}{10000}$, $a = 5$, $b = 1$, $c = d = \frac{1}{10}$, $e = \frac{5}{100}$ and $a_1 = a_2 = 1$, $b_1 = b_2 = 0$, $c_1 = c_2 = 1$. Applying (3.11) with $y_{10} = y_{20} = 1$ and $k_1 = k_2 = 1$, we find the HAM-approximations as

$$\begin{aligned}\phi_{13}(x, c_{10}, c_{20}) &= 1 - 1.9995c_{10} - 0.9996c_{10}^2 + 3.51 \times 10^{-6}c_{10}c_{20} + (1.9995c_{10} + 0.999c_{10}^2 \\ &\quad - 4.69 \times 10^{-6}c_{10}c_{20})x^2 + (3.12 \times 10^{-5}c_{10}^2 + 1.17 \times 10^{-6}c_{10}c_{20})x^4. \\ \phi_{23}(x, c_{10}, c_{20}) &= 1 - 0.074985c_{20} + 2.81 \times 10^{-6}c_{10}c_{20} - 0.03749c_{20}^2 + (0.07498c_{20} \\ &\quad - 3.75 \times 10^{-6}c_{10}c_{20} + 0.0374c_{20}^2)x^2 + (9.37 \times 10^{-7}c_{10}c_{20} + 3.51 \times 10^{-8}c_{20}^2)x^4.\end{aligned}$$

Using the formula (3.13), we find $c_{10} = -1.00010501$, $c_{20} = -1.0000443$, and hence the HAM-approximations

$$\begin{cases} \phi_{13}(x) = 1.99985 - 0.99988x^2 + 3.24108 \times 10^{-5}x^4. \\ \phi_{23}(x) = 1.0375 - 0.0374964x^2 + 9.72266 \times 10^{-7}x^4. \end{cases}$$

Using (3.11) with $y_{10} = y_{20} = 1$ and $k_1 = k_2 = 2$, we find the HAM-approximations as

$$\begin{aligned}\phi_{13}(x, c_{10}, c_{20}) &= 1 - 1.333c_{10} - 0.666461c_{10}^2 + 1.45 \times 10^{-6}c_{10}c_{20} + (1.333c_{10} + 0.6664c_{10}^2 \\ &\quad - 2.08 \times 10^{-6}c_{10}c_{20})x^2 + (0.00001665c_{10}^2 + 6.24 \times 10^{-7}c_{10}c_{20})x^4. \\ \phi_{23}(x, c_{10}, c_{20}) &= 1 - 0.04999c_{20} + 1.16 \times 10^{-6}c_{10}c_{20} - 0.024995c_{20}^2 + (0.04999c_{20} \\ &\quad - 1.66 \times 10^{-6}c_{10}c_{20} + 0.02499c_{20}^2)x^2 + (4.99 \times 10^{-7}c_{10}c_{20} + 1.87 \times 10^{-8}c_{20}^2)x^4.\end{aligned}$$

Applying (3.13), we find $c_{10} = -0.995713$, $c_{20} = -0.996167$, and obtain the HAM-approximate solutions

$$\begin{cases} \phi_{13}(x) = 1.66653 - 0.666545x^2 + 1.713 \times 10^{-5}x^4. \\ \phi_{23}(x) = 1.025 - 0.0249963x^2 + 5.142 \times 10^{-7}x^4. \end{cases}$$

Table 1 Comparison of numerical results for $k_1 = k_2 = 1$ of Example 5.1

x	Solutions				Errors			
	ϕ_{13}	$\psi_{13}[1]$	ϕ_{23}	$\psi_{23}[1]$	Res_{13}	$res_{13}[1]$	Res_{23}	$res_{23}[1]$
0.1	1.9898484	1.9898484	1.0371204	1.0371204	2.46E-04	2.46E-04	7.40E-06	7.40E-06
0.3	1.9098583	1.9098583	1.0341207	1.0341207	2.17E-04	2.17E-04	6.51E-06	6.51E-06
0.5	1.7498793	1.7498793	1.0281213	1.0281213	1.61E-04	1.60E-04	4.83E-06	4.82E-06
0.7	1.5099140	1.5099140	1.0191224	1.0191224	8.62E-05	8.62E-05	2.58E-06	2.58E-06
0.9	1.1899659	1.1899659	1.0071239	1.0071239	1.51E-05	1.51E-05	4.55E-07	4.55E-07

Example 5.2. Consider the system of Lane-Emden-Fowler equation occurs in catalytic diffusion reactions [3, 4]

$$\begin{cases} (x^2 y_1')' = x^2(a y_1^2 + b y_1 y_2), & x \in (0, 1) \\ (x^2 y_2')' = x^2(c y_1^2 - d y_1 y_2), \\ y_1'(0) = 0, \quad y_2'(0) = 0, \quad y_1(1) = 1, \quad y_2(1) = 2. \end{cases} \quad (5.6)$$

Table 2 Comparison of numerical results for $k_1 = k_2 = 2$ of Example 5.1

x	Solutions				Errors			
	ϕ_{13}	$\psi_{13}[32]$	ϕ_{23}	$\psi_{23}[32]$	Res_{13}	$res_{13}[32]$	Res_{23}	$res_{23}[32]$
0.1	1.6598623	1.6598747	1.0247458	1.0247462	5.49E-05	1.31E-04	1.65E-06	3.94E-06
0.3	1.6065388	1.6065503	1.0227461	1.0227465	3.85E-05	1.14E-04	1.16E-06	3.44E-06
0.5	1.4998926	1.4999020	1.0187467	1.0187470	7.73E-06	8.34E-05	2.37E-07	2.50E-06
0.7	1.3399248	1.3399312	1.0127477	1.0127479	3.18E-05	4.31E-05	9.50E-07	1.29E-06
0.9	1.1266376	1.1266400	1.0047491	1.0047492	6.69E-05	7.12E-06	2.00E-06	2.13E-07

Here, $k_1 = k_2 = 2$, $a_1 = a_2 = 1$, $b_1 = b_2 = 0$, $c_1 = 1$ and $c_2 = 2$. Using (3.11) with $y_{10} = 1$, $y_{20} = 2$ and $a = 1$, $b = \frac{2}{5}$, $c = \frac{1}{2}$, $d = 1$, we find the HAM approximate solution as

$$\begin{aligned}
\phi_{14}(x, c_{10}, c_{20}) &= 1 + \frac{9c_{10}}{10} + \frac{597c_{10}^2}{500} + \frac{55973c_{10}^3}{105000} + \frac{7c_{10}c_{20}}{120} + \frac{1613c_{10}^2c_{20}}{47250} + \frac{65c_{10}c_{20}^2}{3024} \\
&+ \left(-\frac{9c_{10}}{10} - \frac{33c_{10}^2}{25} - \frac{9611c_{10}^3}{15000} - \frac{c_{10}c_{20}}{12} - \frac{1409c_{10}^2c_{20}}{27000} - \frac{67c_{10}c_{20}^2}{2160} \right) x^2 + \left(\frac{63c_{10}^2}{500} + \frac{563c_{10}^3}{5000} \right. \\
&+ \left. \frac{c_{10}c_{20}}{40} + \frac{91c_{10}^2c_{20}}{4500} + \frac{7c_{10}c_{20}^2}{720} \right) x^4 + \left(-\frac{173c_{10}^3}{35000} - \frac{137c_{10}^2c_{20}}{63000} - \frac{c_{10}c_{20}^2}{5040} \right) x^6. \\
\phi_{24}(x, c_{10}, c_{20}) &= 2 + \frac{5c_{20}}{4} + \frac{63c_{10}c_{20}}{200} + \frac{1961c_{10}^2c_{20}}{14000} + \frac{67c_{20}^2}{48} + \frac{673c_{10}c_{20}^2}{5040} + \frac{3139c_{20}^3}{6048} \\
&+ \left(-\frac{5c_{20}}{4} - \frac{9c_{10}c_{20}}{20} - \frac{413c_{10}^2c_{20}}{2000} - \frac{35c_{20}^2}{24} - \frac{713c_{10}c_{20}^2}{3600} - \frac{487c_{20}^3}{864} \right) x^2 + \left(\frac{27c_{10}c_{20}}{200} + \frac{141c_{10}^2c_{20}}{2000} \right. \\
&+ \left. \frac{c_{20}^2}{16} + \frac{83c_{10}c_{20}^2}{1200} + \frac{13c_{20}^3}{288} \right) x^4 + \left(-\frac{57c_{10}^2c_{20}}{14000} - \frac{13c_{10}c_{20}^2}{2800} - \frac{c_{20}^3}{2016} \right) x^6.
\end{aligned}$$

Using (3.13), we find $c_{10} = -0.767463$, $c_{20} = -0.789762$. Hence, the HAM approximate solution

$$\begin{cases} \phi_{14} = 0.780767 + 0.191485x^2 + 0.0244069x^4 + 0.00334088x^6. \\ \phi_{24} = 1.68960 + 0.273372x^2 + 0.0326694x^4 + 0.00436072x^6. \end{cases}$$

Using (3.11) with $a = b = c = d = 1$ and (3.13) with $c_{10} = -0.689796$, $c_{20} = -0.708697$, and obtain the HAM-approximations as

$$\begin{cases} \phi_{14} = 0.674423 + 0.271204x^2 + 0.0454739x^4 + 0.00889876x^6. \\ \phi_{24} = 1.67352 + 0.27178x^2 + 0.0455586x^4 + 0.00914259x^6. \end{cases}$$

Example 5.3. Consider the following system of Lane-Emden-Fowler with boundary conditions

$$\begin{cases} (x^3y_1')' = x^3(y_1y_2 + 7 + (y_1 - 1)^2), & x \in (0, 1), \\ (x^4y_2')' = x^4(y_1y_2 - 11 + (y_2 - 1)^2), \\ y_1'(0) = 0, \quad y_2'(0) = 0, \quad y_1(1) = 2, \quad y_2(1) = 0. \end{cases} \quad (5.7)$$

Table 3 Comparison of numerical results for $a = 1$, $b = \frac{2}{5}$, $c = \frac{1}{2}$, $d = 1$ of Example 5.2

x	Solutions				Errors			
	ϕ_{14}	$\psi_{14}[32]$	ϕ_{24}	$\psi_{24}[32]$	Res_{14}	$res_{14}[32]$	Res_{24}	$res_{24}[32]$
0.1	0.7826843	0.7658317	1.6923350	1.6713156	1.13E-02	2.26E-01	1.59E-02	7.63E-01
0.3	0.7982008	0.7835530	1.7144693	1.6962143	9.45E-03	1.95E-01	1.34E-02	7.54E-01
0.5	0.8302159	0.8194185	1.7600510	1.7466228	5.96E-03	1.41E-01	9.17E-03	7.21E-01
0.7	0.8808479	0.8746115	1.8319072	1.8241881	4.43E-04	7.71E-02	2.78E-03	6.85E-01
0.9	0.9536588	0.9517495	1.9347811	1.9324416	1.11E-02	2.06E-02	1.02E-02	6.77E-01

Table 4 Comparison of numerical results for $a = b = c = d = 1$ of Example 5.2

x	Solutions				Errors			
	ϕ_{14}	$\psi_{14}[32]$	ϕ_{24}	$\psi_{24}[32]$	Res_{14}	$res_{14}[32]$	Res_{24}	$res_{24}[32]$
0.1	0.6771397	0.5967530	1.6762408	1.5967530	4.27E-02	1.143631	4.62E-02	1.1436
0.3	0.6992063	0.6293170	1.6983544	1.6293170	3.57E-02	0.99354	3.94E-02	0.9935
0.5	0.7452053	0.6936848	1.7444538	1.6936848	2.26E-02	0.72673	2.71E-02	0.7267
0.7	0.8192784	0.7895710	1.8187051	1.7895710	1.36E-03	0.40619	8.10E-03	0.4061
0.9	0.9286631	0.9196325	1.9284103	1.9196325	4.41E-02	0.11286	3.25E-02	0.1128

The exact solutions are $y_1(x) = 3 - x^2$ and $y_2(x) = -1 + x^2$. Here, $a_1 = a_2 = 1$, $b_1 = b_2 = 0$ and $c_1 = 2$ and $c_2 = 0$. Using (3.11) with $y_{10} = 2$ and $y_{20} = 0$, we find the series solutions as

$$\begin{aligned}
\phi_{13}(x, c_{10}, c_{20}) &= 2 - 3c_{10} - \frac{5c_{10}^2}{2} - \frac{217c_{10}^3}{288} - \frac{c_{10}c_{20}}{2} - \frac{23c_{10}^2c_{20}}{288} - \frac{c_{10}c_{20}^2}{6} + \left(3c_{10} + \frac{9c_{10}^2}{4}\right. \\
&\quad \left. + \frac{2c_{10}^3}{3} + \frac{3c_{10}c_{20}}{4} + \frac{c_{10}^2c_{20}}{12} + \frac{c_{10}c_{20}^2}{4}\right)x^2 + \left(\frac{c_{10}^2}{4} + \frac{c_{10}^3}{16} - \frac{c_{10}c_{20}}{4} + \frac{c_{10}^2c_{20}}{48} - \frac{c_{10}c_{20}^2}{12}\right)x^4 \\
&\quad + \left(\frac{7c_{10}^3}{288} - \frac{7c_{10}^2c_{20}}{288}\right)x^6. \\
\phi_{23}(x, c_{10}, c_{20}) &= 3c_{20} + 3c_{20}^2 + \frac{89c_{10}c_{20}^2}{1890} + \frac{1801c_{20}^3}{1890} + \left(-3c_{20} - 3c_{20}^2 - \frac{c_{10}c_{20}^2}{10} - \frac{9c_{20}^3}{10}\right)x^2 \\
&\quad + \left(\frac{c_{10}c_{20}^2}{14} - \frac{c_{20}^3}{14}\right)x^4 + \left(-\frac{c_{10}c_{20}^2}{54} + \frac{c_{20}^3}{54}\right)x^6.
\end{aligned}$$

Applying (3.13), we find $c_{10} = c_{20} = -1$, and obtain the exact solutions

$$\phi_{13}(x) = 3 - x^2, \quad \phi_{23}(x) = -1 + x^2.$$

Example 5.4. Consider the following system of Lane-Emden-Fowler with boundary conditions

$$\begin{cases}
(x^5 y_1')' = x^5 (-8e^{y_1} - 16e^{-\frac{y_2}{2}}), & x \in (0, 1), \\
(x^3 y_1')' = x^3 (8e^{-y_2} + 8e^{\frac{y_1}{2}}), \\
y_1'(0) = 0, \quad y_2'(0) = 0, \quad y_1(1) = -2 \ln 2, \quad y_2(1) = 2 \ln 2.
\end{cases} \quad (5.8)$$

The exact solutions are $y_1(x) = -2\ln(1+x^2)$ and $y_2(x) = 2\ln(1+x^2)$. Here, $k_1 = 5$, $k_2 = 3$, $a_1 = a_2 = 1$, $b_1 = b_2 = 0$, $c_1 = -2\ln 2$ and $c_2 = 2\ln 2$. Making use of (3.11) and (3.13) with $y_{10} = -2\ln 2$, $y_{20} = 2\ln 2$, we obtain the HAM-approximations with ($c_{10} = -0.763735$, $c_{20} = -0.743226$) as

$$\left\{ \begin{array}{l} \phi_{15}(x) = -2.05166 + 0.587014x^2 + 0.0661967x^4 + 0.00974409x^6 + 0.00215741x^8 \\ \quad + 0.000103467x^{10} + 0.000153951x^{12}. \\ \phi_{25}(x) = 1.95556 - 0.499767x^2 - 0.0589602x^4 - 0.00851556x^6 - 0.00181135x^8 \\ \quad - 0.000100184x^{10} - 0.000115625x^{12}. \end{array} \right.$$

Table 5 Comparison of numerical results for $k_1 = 5$, $k_2 = 3$ of Example 5.4

x	Solutions				Errors			
	ϕ_{15}	$\psi_{15}[32]$	ϕ_{25}	$\psi_{25}[32]$	Res_{25}	$res_{25}[32]$	Res_{25}	$res_{25}[32]$
0.1	-2.0457870	-2.0358737	1.9505604	1.9379913	2.20E-03	0.398753	1.65E-03	0.290684
0.3	-1.9982891	-1.9904854	1.9101010	1.8999319	1.51E-03	0.302292	1.17E-03	0.219317
0.5	-1.9006122	-1.8958769	1.8267970	1.8202489	7.51E-04	0.168370	6.34E-04	0.120416
0.7	-1.7468574	-1.7447898	1.6954112	1.6922536	3.74E-04	6.38E-02	3.94E-04	4.39E-02
0.9	-1.5265642	-1.5261201	1.5066963	1.5059088	6.69E-04	1.18E-02	7.42E-04	7.51E-03

Example 5.5. Consider the following system of Lane-Emden-Fowler with boundary conditions

$$\left\{ \begin{array}{l} (x^2 y_1')' = x^2 (2(7 + e^{y_2}) e^{-2y_1}), \quad x \in (0, 1), \\ (x^2 y_2')' = x^2 (2(11 + e^{y_1}) e^{-2y_2}), \\ y_1'(0) = 0, \quad y_2'(0) = 0, \quad y_1(1) = \ln 4, \quad y_2(1) = \ln 5. \end{array} \right. \quad (5.9)$$

The exact solutions are $y_1(x) = \ln(4+x^2)$ and $y_2(x) = \ln(5+x^2)$. Here, $k_1 = k_2 = 2$, $a_1 = a_2 = 1$, $b_1 = b_2 = 0$, $c_1 = \ln 4$ and $c_2 = \ln 5$. Using (3.11) and (3.13) with $y_{10} = \ln 4$, $y_{20} = \ln 5$, we find the HAM-approximate series solutions with ($c_{10} = -0.766209$, $c_{20} = -0.800994$) as

$$\left\{ \begin{array}{l} \phi_{14} = 1.58464 - 0.18032x^2 - 0.0160564x^4 - 0.00163913x^6 - 0.000328184x^8. \\ \phi_{24} = 1.77423 - 0.152098x^2 - 0.0114857x^4 - 0.00103977x^6 - 0.00016868x^8. \end{array} \right.$$

Table 6 Comparison of numerical results for $k_1 = k_2 = 2$ of Example 5.5

x	Solutions				Errors			
	ϕ_{14}	$\psi_{14}[32]$	ϕ_{24}	$\psi_{24}[32]$	Res_{24}	$res_{24}[32]$	Res_{24}	$res_{24}[32]$
0.1	1.5828329	1.5769131	1.7727080	1.7709758	2.14E-03	8.30E-02	9.45E-04	2.33E-02
0.3	1.5682776	1.5632335	1.7604475	1.7589804	1.77E-03	6.88E-02	7.83E-04	1.90E-02
0.5	1.5385273	1.5349546	1.7354708	1.7344446	1.31E-03	4.57E-02	5.90E-04	1.21E-02
0.7	1.4922141	1.4902763	1.6968123	1.6962663	1.23E-03	2.21E-02	5.60E-04	5.48E-03
0.9	1.4270318	1.4264972	1.6428697	1.6427234	2.46E-03	5.17E-03	1.06E-03	1.12E-03

Example 5.6. Consider the following system of Lane-Emden-Fowler with boundary conditions

$$\begin{cases} (x^2 y_1')' = x^2 \left(-6 \left(e^{\frac{y_2}{3}} + 4 \right) e^{\frac{2y_1}{3}} \right), & x \in (0, 1), \\ (x^2 y_2')' = x^2 \left(6 \left(e^{\frac{-y_1}{3}} + 4 \right) e^{\frac{-2y_2}{3}} \right), \\ y_1'(0) = 0, \quad y_2'(0) = 0, \quad y_1(1) = -3 \ln 3, \quad y_2(1) = 3 \ln 3. \end{cases} \quad (5.10)$$

The exact solutions are $y_1 = -3 \ln(2 + x^2)$, $y_2 = 3 \ln(2 + x^2)$. Here, $k_1 = k_2 = 2$, $a_1 = a_2 = 1$, $b_1 = b_2 = 0$, $c_1 = -3 \ln 3$ and $c_2 = 3 \ln 3$. In view of (3.11) and (3.13) with $y_{10} = -3 \ln 3$, $y_{20} = 3 \ln 3$, we find the HAM-approximate series solutions with ($c_{10} = -0.764679$, $c_{20} = -0.764679$) as

$$\begin{cases} \phi_{14}(x) = -3.91525 + 0.564146x^2 + 0.049265x^4 + 0.00500793x^6 + 0.00099786x^8. \\ \phi_{24}(x) = 3.91525 - 0.564146x^2 - 0.049265x^4 - 0.00500793x^6 - 0.00099786x^8. \end{cases}$$

Table 7 Comparison of numerical results for $k_1 = 2$, $k_2 = 2$ of Example 5.6

t	Solutions				Errors			
	ϕ_{14}	$\psi_{14}[32]$	ϕ_{24}	$\psi_{24}[32]$	Res_{24}	$res_{24}[32]$	Res_{24}	$res_{24}[32]$
0.1	-3.9096075	-3.8933979	3.9096075	3.8933979	6.39E-03	0.225756	6.39E-03	2.25E-02
0.3	-3.8640780	-3.8502979	3.8640780	3.8502979	5.34E-03	0.186062	5.34E-03	5.58E-02
0.5	-3.7710561	-3.7613414	3.7710561	3.7613414	4.11E-03	0.122139	4.11E-03	6.10E-02
0.7	-3.6263470	-3.6211160	3.6263470	3.6211160	4.05E-03	5.78E-02	4.05E-03	4.04E-02
0.9	-3.4228817	-3.4214542	3.4228817	3.4214542	8.02E-03	1.30E-02	8.02E-03	1.17E-02

Example 5.7. Consider the following system of Lane-Emden-Fowler with boundary conditions

$$\begin{cases} (x^3 y_1')' = x^3 (- (3 + y_2^2) y_1^5), & x \in (0, 1), \\ (x^4 y_1')' = x^4 ((4y_1^{-2} + 1) y_2^{-3}), \\ y_1'(0) = 0, \quad y_2'(0) = 0, \quad y_1(1) = \frac{1}{\sqrt{2}}, \quad y_2(1) = \sqrt{2}. \end{cases} \quad (5.11)$$

The exact solutions are $y_1(x) = \frac{1}{\sqrt{1+x^2}}$ and $y_2(x) = \sqrt{1+x^2}$. Here, $a_1 = a_2 = 1$, $b_1 = b_2 = 0$, $c_1 = \frac{1}{\sqrt{2}}$ and $c_2 = \sqrt{2}$. In view of (3.11) and (3.13) with $y_{10} = \frac{1}{\sqrt{2}}$, $y_{20} = \sqrt{2}$, we find the HAM-approximate series solutions with ($c_{10} = -0.718977$, $c_{20} = -0.726659$), as

$$\begin{cases} \phi_{14}(x) = 0.626038 + 0.0695736x^2 + 0.0099351x^4 + 0.00129509x^6 + 0.000264696x^8. \\ \phi_{24}(x) = 1.67507 - 0.237503x^2 - 0.0198153x^4 - 0.00227264x^6 - 0.0012686x^8. \end{cases}$$

Table 8 Comparison of numerical results for $k_1 = 3$, $k_2 = 4$ of Example 5.7

t	Solutions				Errors			
	ϕ_{14}	ψ_{14} [32]	ϕ_{24}	ψ_{24} [32]	Res_{24}	res_{24} [32]	Res_{24}	res_{24} [32]
0.1	0.6267350	0.6326026	1.6726961	1.6660461	1.67E-03	0.123810	8.99E-03	0.195077
0.3	0.6323813	0.6373236	1.6535356	1.6483411	1.35E-03	0.103217	7.53E-03	0.146187
0.5	0.6440739	0.6474853	1.6144184	1.6113861	9.66E-04	0.069423	6.11E-03	7.46E-02
0.7	0.6626824	0.6644484	1.5535984	1.5524363	1.01E-03	0.034154	6.20E-03	1.78E-02
0.9	0.6897135	0.6901626	1.4679409	1.4677736	2.79E-03	0.008057	1.01E-02	1.44E-03

6 Concluding remarks

The theory of a system of singular differential equations finds its vital presence in many of the natural or physical processes such as catalytic diffusion reactions [4], and some system of Lane-Emden equations that relate the concentration of the carbon substrate and the concentration of oxygen [32]. In this paper, a general analytical approach has been presented for the approximate series solution of system of singular differential equations with Neumann and Robin type boundary conditions. Unlike the Adomian decomposition method, the proposed technique does not require the computation of unknown constants, and it contains convergence parameters which ensure the fast convergence of the series solution. Unlike, the other numerical methods, our approach does not require any linearization or discretization of variables. Numerical results obtained by the present method are better than the results obtained by the Adomian decomposition method reported in [32], as shown in Tables 1-8. Convergence and error estimation of the homotopy analysis method for a system of singular boundary value problems is provided under quite general conditions.

References

- [1] R. Singh, Solving coupled Lane-Emden equations by Green's function and decomposition technique, International Journal of Applied and Computational Mathematics 6 (2020) 80.
- [2] B. Muatjetjeja, C. M. Khalique, Noether, partial noether operators and first integrals for the coupled Lane-Emden system, Mathematical and Computational Applications 15 (3) (2010) 325–333.
- [3] D. Flockerzi, K. Sundmacher, On coupled Lane-Emden equations arising in dusty fluid models, Journal of Physics: Conference Series, Vol. 268, IOP Publishing, 2011, 012006.
- [4] R. Rach, J.S. Duan, A.M. Wazwaz, Solving coupled Lane-Emden boundary value problems in catalytic diffusion reactions by the Adomian decomposition method, Journal of Mathematical Chemistry 52 (1) (2014) 255–267.

- [5] A.M. Wazwaz, R. Rach, J.-S. Duan, A study on the systems of the volterra integral forms of the Lane–Emden equations by the Adomian decomposition method, *Mathematical Methods in the Applied Sciences* 37 (1) (2014) 10–19.
- [6] R. Singha, A.-M. Wazwazb, An efficient algorithm for solving coupled Lane–Emden boundary value problems in catalytic diffusion reactions: The homotopy analysis method, *Communications in mathematical and in computer chemistry* 81 (3) (2019) 785–800.
- [7] T.C. Hao, F.-Z. Cong, Y.-F. Shang, An efficient method for solving coupled Lane–Emden boundary value problems in catalytic diffusion reactions and error estimate, *Journal of Mathematical Chemistry* 56 (9) (2018) 2691–2706.
- [8] A.M. Wazwaz, R. Rach, Comparison of the Adomian decomposition method and the variational iteration method for solving the Lane-Emden equations of the first and second kinds, *Kybernetes* 40 (9/10) (2011) 1305–1318.
- [9] M. Inc, M. Ergut, Y. Cherruault, A different approach for solving singular two-point boundary value problems, *Kybernetes: The International Journal of Systems & Cybernetics* 34 (7) (2005) 934–940.
- [10] R. Mittal, R. Nigam, Solution of a class of singular boundary value problems, *Numerical Algorithms* 47 (2) (2008) 169–179.
- [11] A.M. Wazwaz, The variational iteration method for solving systems of equations of Emden–Fowler type, *International Journal of Computer Mathematics* 88 (16) (2011) 3406–3415.
- [12] S. Khuri, A. Sayfy, A novel approach for the solution of a class of singular boundary value problems arising in physiology, *Mathematical and Computer Modelling* 52 (3) (2010) 626–636.
- [13] A. Ebaid, A new analytical and numerical treatment for singular two-point boundary value problems via the Adomian decomposition method, *Journal of Computational and Applied Mathematics* 235 (8) (2011) 1914–1924.
- [14] A.-M. Wazwaz, R. Rach, J.-S. Duan, Adomian decomposition method for solving the volterra integral form of the Lane–Emden equations with initial values and boundary conditions, *Applied Mathematics and Computation* 219 (10) (2013) 5004–5019.
- [15] R. Singh, J. Kumar, G. Nelakanti, Numerical solution of singular boundary value problems using Green’s function and improved decomposition method, *Journal of Applied Mathematics and Computing* 43 (1-2) (2013) 409–425.
- [16] R. Singh, J. Kumar, The Adomian decomposition method with Green’s function for solving nonlinear singular boundary value problems, *Journal of Applied Mathematics and Computing* 44 (1-2) (2014) 397–416.

- [17] R. Singh, J. Kumar, An efficient numerical technique for the solution of nonlinear singular boundary value problems, *Computer Physics Communications* 185 (4) (2014) 1282–1289.
- [18] R. Singh, J. Kumar, G. Nelakanti, Approximate series solution of singular boundary value problems with derivative dependence using Green’s function technique, *Computational and Applied Mathematics* 33 (2) (2014) 451–467.
- [19] N. Das, R. Singh, A.-M. Wazwaz, J. Kumar, An algorithm based on the variational iteration technique for the Bratu-type and the Lane–Emden problems, *Journal of Mathematical Chemistry* 54 (2) (2016) 527–551.
- [20] R. Singh, A.-M. Wazwaz, J. Kumar, An efficient semi-numerical technique for solving nonlinear singular boundary value problems arising in various physical models, *International Journal of Computer Mathematics* 93 (8) (2016) 1330–1346.
- [21] M. Singh, A. K. Verma, An effective computational technique for a class of Lane–Emden equations, *Journal of Mathematical Chemistry* 54 (1) (2016) 231–251.
- [22] R. Singh, N. Das, J. Kumar, The optimal modified variational iteration method for the Lane–Emden equations with neumann and robin boundary conditions, *The European Physical Journal Plus* 132 (6) (2017) 251.
- [23] A. K. Verma, S. Kayenat, On the convergence of mickens type nonstandard finite difference schemes on Lane–Emden type equations, *Journal of Mathematical Chemistry* 56 (6) (2018) 1667–1706.
- [24] A. K. Verma, D. Tiwari, Higher resolution methods based on quasilinearization and haar wavelets on Lane–Emden equations, *International Journal of Wavelets, Multiresolution and Information Processing* 17 (03) (2019) 1950005.
- [25] R. Singh, H. Garg, V. Guleria, Haar wavelet collocation method for Lane–Emden equations with Dirichlet, Neumann and Neumann-Robin boundary conditions, *Journal of Computational and Applied Mathematics* 346 (2019) 150–161.
- [26] R. Singh, J. Shahni, H. Garg, A. Garg, Haar wavelet collocation approach for Lane–Emden equations arising in mathematical physics and astrophysics, *The European Physical Journal Plus* 134 (11) (2019) 548.
- [27] R. Singh, A modified homotopy perturbation method for nonlinear singular Lane–Emden equations arising in various physical models, *International Journal of Applied and Computational Mathematics* 5 (3) (2019) 64.
- [28] R. Singh, V. Guleria, M. Singh, Haar wavelet quasilinearization method for numerical solution of Emden–Fowler type equations, *Mathematics and Computers in Simulation* 174 (2020) 123–133.
- [29] J. Shahni, R. Singh, An efficient numerical technique for Lane–Emden–Fowler boundary value problems: Bernstein collocation method, *The European Physical Journal Plus* 135 (6) (2020) 475.

- [30] M. Kumar, et al., Numerical solution of singular boundary value problems using advanced Adomian decomposition method, *Engineering with Computers* (2020) 1–11.
- [31] S. Muthukumar, M. Veeramuni, R. Lakshmanan, Analytical expression of concentration of substrate and oxygen in excess sludge production using Adomian decomposition method, *Indian journal of applied research* 4 (2014) 387–391.
- [32] J.-S. Duan, R. Rach, A.M. Wazwaz, Oxygen and carbon substrate concentrations in microbial floc particles by the Adomian decomposition method, *Commun. Math. Comput. Chem* 73 (2015) 785–796.
- [33] H. Madduri, P. Roul, A fast-converging iterative scheme for solving a system of Lane–Emden equations arising in catalytic diffusion reactions, *Journal of Mathematical Chemistry* 57 (2) (2019) 570–582.
- [34] A. M. Wazwaz, R. Rach, J.S. Duan, Variational iteration method for solving oxygen and carbon substrate concentrations in microbial floc particles, *Match* 76 (2016) 511–523.
- [35] A. Saadatmandi, S. Fayyaz, Numerical study of oxygen and carbon substrate concentrations in excess sludge production using sinc-collocation method, *Communications in Mathematical and in Computer Chemistry* 80 (2018) 355–368.
- [36] T. Hao, S. Song, Solving coupled Lane-Emden equations arising in catalytic diffusion reactions by reproducing kernel hilbert space method, 2015 International conference on Applied Science and Engineering Innovation, Atlantis Press, 2015.
- [37] S. Liao, *Beyond perturbation: introduction to the homotopy analysis method*, CRC press, 2003.
- [38] S. Liao, On the homotopy analysis method for nonlinear problems, *Applied Mathematics and Computation* 147 (2) (2004) 499–513.
- [39] S. Liao, Y. Tan, A general approach to obtain series solutions of nonlinear differential equations, *Studies in Applied Mathematics* 119 (4) (2007) 297–354.
- [40] S. Liao, Series solution of nonlinear eigenvalue problems by means of the homotopy analysis method, *Nonlinear Analysis: Real World Applications* 10 (4) (2009) 2455–2470.
- [41] V. Marinca, N. Herişanu, Application of optimal homotopy asymptotic method for solving nonlinear equations arising in heat transfer, *International Communications in Heat and Mass Transfer* 35 (6) (2008) 710–715.
- [42] V. Marinca, N. Herişanu, I. Nemeş, Optimal homotopy asymptotic method with application to thin film flow, *Central European Journal of Physics* 6 (3) (2008) 648.
- [43] S. Liao, An optimal homotopy-analysis approach for strongly nonlinear differential equations, *Communications in Nonlinear Science and Numerical Simulation* 15 (8) (2010) 2003–2016.

- [44] S. Liao, Homotopy analysis method in nonlinear differential equations, Springer, 2012.
- [45] R. Singh, Optimal homotopy analysis method for the non-isothermal Reaction–Diffusion model equations in a spherical catalyst, *Journal of Mathematical Chemistry* 56 (2018) 2579–2590.
- [46] R. Singh, A.-M. Wazwaz, Optimal homotopy analysis method for oxygen diffusion in a spherical cell with nonlinear oxygen uptake kinetics, *Communications in Mathematical and in Computer Chemistry* 80 (2) (2018) 369–382.
- [47] R. Singha, A.-M. Wazwazb, An efficient algorithm for solving coupled Lane–Emden boundary value problems in catalytic diffusion reactions: The homotopy analysis method, *Communications in mathematical and in computer chemistry* 81 (3) (2019) 785–800.
- [48] R. Singh, Analytic solution of singular Emden-Fowler-type equations by Green’s function and homotopy analysis method, *The European Physical Journal Plus* 134 (11) (2019) 583.
- [49] R. Singh, Analytical approach for computation of exact and analytic approximate solutions to the system of Lane-Emden-fowler type equations arising in astrophysics, *The European Physical Journal Plus* 133 (8) (2018) 320.
- [50] R. Rach, A new definition of the Adomian polynomials, *Kybernetes* 37 (7) (2008) 910–955.