

# REVERSE SUPERPOSITION ESTIMATES IN SOBOLEV SPACES

JEAN VAN SCHAFTINGEN

**ABSTRACT.** We study when and how the norm of a function  $u$  in the homogeneous Sobolev spaces  $\dot{W}^{s,p}(\mathbb{R}^n, \mathbb{R}^m)$ , with  $p \geq 1$  and either  $s = 1$  or  $s > 1/p$ , is controlled by the norm of composite function  $f \circ u$  in the same space.

## 1. INTRODUCTION

The absolute value preserves weak differentiability despite its non-differentiability at 0 [5] (see also [3, lemma 7.6; 8, corollary 6.1.14; 9, corollary 2.1.8]). More precisely, if  $u$  belongs to the homogeneous first-order Sobolev space  $\dot{W}^{1,p}(\Omega, \mathbb{R})$  for some  $p \in [1, \infty)$ , that is, if the function  $u: \Omega \rightarrow \mathbb{R}$  is weakly differentiable on the open set  $\Omega \subseteq \mathbb{R}^n$  and its weak derivative  $Du$  satisfies the integrability condition  $\int_{\Omega} |Du|^p < +\infty$ , then  $|u| \in \dot{W}^{1,p}(\Omega, \mathbb{R})$ ; moreover, one has then

$$(1.1) \quad D|u| = \operatorname{sgn}(u)Du \quad \text{almost everywhere in } \Omega,$$

where the signum function  $\operatorname{sgn}$  is defined by  $\operatorname{sgn}(t) = -1$  when  $t < 0$ ,  $\operatorname{sgn}(0) = 0$  and  $\operatorname{sgn}(t) = 1$  when  $t > 0$ . A consequence of the identity (1.1) and of the fact that  $Du = 0$  almost everywhere on  $u^{-1}(\{0\})$  is the integral identity

$$(1.2) \quad \int_{\Omega} |D|u||^p = \int_{\Omega} |Du|^p,$$

which can be interpreted either as an estimate in the homogeneous Sobolev space  $\dot{W}^{1,p}(\Omega)$  for  $|u|$  in terms of  $u$ , or conversely as an a priori estimate for  $u$  in terms of  $|u|$ , *provided* it is known a priori that  $u \in \dot{W}^{1,p}(\Omega, \mathbb{R})$ . We will adopt the latter point of view.

This result about the absolute value is a particular case of reverse estimates for superposition operators  $u \mapsto f \circ u$ , for  $u \in \dot{W}^{1,p}(\mathbb{R}^n, \mathbb{R}^m)$  and  $f: \mathbb{R}^m \rightarrow \mathbb{R}^{\ell}$ . We state in theorem 2.1 below a wide condition on the function  $f$  which ensures that  $u \in \dot{W}^{1,p}(\mathbb{R}^n, \mathbb{R}^m)$  is controlled by  $f \circ u \in \dot{W}^{1,p}(\mathbb{R}^n, \mathbb{R}^{\ell})$ ; this condition does not require that  $f \circ u \in \dot{W}^{1,p}(\mathbb{R}^n, \mathbb{R}^{\ell})$  when  $u \in \dot{W}^{1,p}(\mathbb{R}^n, \mathbb{R}^m)$ .

We next consider the question whether such reverse superposition estimate extend to the *homogeneous fractional Sobolev space*

$$(1.3) \quad \dot{W}^{s,p}(\Omega, \mathbb{R}^m) := \left\{ u: \Omega \rightarrow \mathbb{R}^m \mid \iint_{\Omega \times \Omega} \frac{|u(y) - u(x)|^p}{|y - x|^{n+sp}} dy dx < +\infty \right\},$$

with  $0 < s < 1$  and  $1 \leq p < +\infty$ . Although there is no identity such as (1.1) for fractional Sobolev spaces, we prove that when  $sp > 1$  there exists a constant such that for every

---

2010 *Mathematics Subject Classification.* 46E35 (47H30).

*Key words and phrases.* Fractional Sobolev space; superposition operator; reverse inequality.

$u \in \dot{W}^{s,p}(\Omega, \mathbb{R})$ , the reverse estimate

$$(1.4) \quad \iint_{\Omega \times \Omega} \frac{|u(y) - u(x)|^p}{|y - x|^{n+sp}} dy dx \leq C \iint_{\Omega \times \Omega} \frac{||u(y)| - |u(x)||^p}{|y - x|^{n+sp}} dy dx$$

holds. The estimate (1.4) is a particular case of a class of reverse estimates for superposition operators (theorem 3.1). The proof of (1.4) is based on a reverse oscillation obtained by Petru Mironescu and the author in the lifting of fractional Sobolev mappings over a compact covering [6].

When  $sp \leq 1$ , the reverse estimate (1.4) fails. In fact there exists an unbounded sequence  $(u_j)_{j \in \mathbb{N}}$  in  $\dot{W}^{s,p}(\Omega, \mathbb{R})$  such that for each  $j \in \mathbb{N}$  the function  $|u_j|$  is constant on  $\Omega$  ( $sp < 1$ , proposition 4.1) or such that the sequence  $(|u_j|)_{j \in \mathbb{N}}$  remains bounded in  $\dot{W}^{s,p}(\Omega, \mathbb{R})$  ( $sp = 1$ , proposition 4.2).

When  $p = 2$ , an estimate of the form (1.4) still holds when  $1 < s < 3/2$  with a suitable definition of fractional Sobolev norm [7].

## 2. REVERSE ESTIMATES FOR FIRST-ORDER SOBOLEV SPACES

Our first result is a reverse estimate for weakly differentiable functions.

**Theorem 2.1.** *If the set  $\Omega \subseteq \mathbb{R}^n$  is open, if the function  $f : \mathbb{R}^m \rightarrow \mathbb{R}^\ell$  is Borel-measurable, if  $u \in \dot{W}_{\text{loc}}^{1,1}(\Omega, \mathbb{R}^m)$  and if  $f \circ u \in \dot{W}_{\text{loc}}^{1,1}(\Omega, \mathbb{R}^\ell)$ , then for almost every  $x \in \Omega$  and every  $h \in \mathbb{R}^n$ ,*

$$(2.1) \quad |Du(x)[h]| \leq |D(f \circ u)(x)[h]| \limsup_{y \rightarrow u(x)} \frac{|y - u(x)|}{|f(y) - f(u(x))|}$$

*Remark 2.2.* If the function  $f$  is classically differentiable at the point  $u(x)$ , then

$$(2.2) \quad \limsup_{y \rightarrow u(x)} \frac{|y - u(x)|}{|f(y) - f(u(x))|} = \frac{1}{\sup \{|Df(u(x))[k]|/|k| \mid k \in \mathbb{R}^m \setminus \{0\}\}}.$$

*Remark 2.3.* If for each  $y \in \mathbb{R}$  the function  $f$  is defined as  $f(y) := |y|$ , then we have for every  $z \in \mathbb{R}$ ,

$$(2.3) \quad \limsup_{y \rightarrow z} \frac{|y - z|}{||y| - |z||} = 1,$$

and (2.1) is then in this particular case a consequence of (1.1).

The proof of theorem 2.1 follows the strategy of the general chain rule for weakly differentiable functions [1].

*Proof of theorem 2.1 when  $n = 1$ .* By the characterisation of weakly differentiable functions on an interval (see for example [4, theorem 7.13]), for almost every  $x \in \Omega$  there exists a sequence  $(h_j)_{j \in \mathbb{N}}$  in  $\mathbb{R} \setminus \{0\}$  converging to 0 such that both

$$(2.4) \quad \lim_{j \rightarrow \infty} \frac{u(x + h_j) - u(x)}{h_j} = u'(x)$$

and

$$(2.5) \quad \lim_{j \rightarrow \infty} \frac{f(u(x + h_j)) - f(u(x))}{h_j} = (f \circ u)'(x).$$

Assuming without loss of generality that

$$\limsup_{y \rightarrow u(x)} \frac{|y - u(x)|}{|f(y) - f(u(x))|} < +\infty$$

we have for  $j \in \mathbb{N}$  large enough  $f(u(x) + h_j) \neq f(u(x))$ ; it then follows from the limits (2.4) and (2.5) that

$$\begin{aligned} |u'(x)| &= |(f \circ u)'(x)| \lim_{j \rightarrow \infty} \frac{|u(x + h_j) - u(x)|}{|f(u(x + h_j)) - f(u(x))|} \\ &\leq |(f \circ u)'(x)| \limsup_{y \rightarrow u(x)} \frac{|y - u(x)|}{|f(y) - f(u(x))|}. \end{aligned} \quad \square$$

*Proof of theorem 2.1 when  $n \geq 2$ .* The proof goes by noting that the restrictions of  $u$  and  $f \circ u$  to almost every one-dimensional line  $L$  are weakly differentiable (see for example [4, theorem 10.35]), applying the one-dimensional case and concluding by Fubini's theorem.  $\square$

### 3. FRACTIONAL SOBOLEV SPACES

In the fractional case, we have the following counterpart of theorem 2.1.

**Theorem 3.1.** *For every  $s \in (0, 1)$  and  $p \in [1, +\infty)$  satisfying  $sp > 1$ , there exists a constant  $C$  such that for every convex set  $\Omega \subseteq \mathbb{R}^n$  and every  $f : \mathbb{R}^m \rightarrow \mathbb{R}^\ell$ , if  $u \in \dot{W}^{s,p}(\Omega, \mathbb{R}^m)$  and if  $f \circ u \in \dot{W}^{s,p}(\Omega, \mathbb{R}^\ell)$ , then*

$$\begin{aligned} (3.1) \quad & \iint_{\Omega \times \Omega} \frac{|u(y) - u(x)|^p}{|y - x|^{n+sp}} dy dx \\ & \leq C \left( \sup \left\{ \frac{\text{diam}(K)}{\text{diam}(f(K))} \mid K \subset \text{ess rg } u \text{ compact, connected and } \text{diam}(K) > 0 \right\} \right)^p \\ & \quad \times \iint_{\Omega \times \Omega} \frac{|f(u(y)) - f(u(x))|^p}{|y - x|^{n+sp}} dy dx. \end{aligned}$$

Here  $\text{ess rg } u$  denotes the *essential range* of the function  $u : \Omega \rightarrow \mathbb{R}^m$  with respect to Lebesgue's  $n$ -dimensional measure  $\mathcal{L}^n$ , defined as

$$(3.2) \quad \text{ess rg } u := \left\{ y \in \mathbb{R}^m \mid \text{for each } \varepsilon > 0, \mathcal{L}^n(u^{-1}(B_\varepsilon(y))) > 0 \right\}.$$

Our main tool to prove theorem 3.1 is the following reverse oscillation inequality [6].

**Proposition 3.2.** *If the set  $\Omega \subseteq \mathbb{R}^n$  is convex and if  $sp > 1$ , then there exists a constant  $C$  such that for every  $u \in \dot{W}^{s,p}(\Omega, \mathbb{R}^m)$  one has*

$$(3.3) \quad \iint_{\Omega \times \Omega} \frac{(\text{ess osc}_{[x,y]} u)^p}{|y - x|^{n+sp}} dy dx \leq C \iint_{\Omega \times \Omega} \frac{|u(y) - u(x)|^p}{|y - x|^{n+sp}} dy dx.$$

Here we have defined the segment  $[x, y] = \{(1-t)x + ty \mid 0 \leq t \leq 1\}$  and the *essential oscillation*

$$(3.4) \quad \operatorname{ess\,osc}_{[x,y]} u := \operatorname{ess\,sup}_{t,r \in [0,1]} |u((1-r)x + ry) - u((1-t)x + ty)|$$

Proposition 3.2 is proved for  $n = 1$  and extended by Fubini-type arguments to higher dimension [6]; we give here a direct proof in all dimensions.

*Proof of proposition 3.2.* Since  $sp > 1$ , we can fix  $\sigma \in \mathbb{R}$  so that  $\frac{1}{p} < \sigma < s$ . There exists a constant  $C_1$  such that for every  $x, y \in \mathbb{R}^n$ , we have

$$(3.5) \quad (\operatorname{ess\,osc}_{[x,y]} u)^p \leq C_1 \iint_{[0,1] \times [0,1]} \frac{|u((1-t)x + ty) - u((1-r)x + ry)|^p}{|t-r|^{1+\sigma p}} dt dr.$$

Indeed, since  $\sigma p > 1$ , by the fractional Morrey–Sobolev embedding there exists a constant  $C_1$  such that (see [2, §8]) for almost every  $\rho, \tau \in [0, 1]$ ,

$$(3.6) \quad |u((1-\rho)x + \rho y) - u((1-\tau)x + \tau y)|^p \leq C_1 \iint_{[0,1] \times [0,1]} \frac{|u((1-t)x + ty) - u((1-r)x + ry)|^p}{|t-r|^{1+\sigma p}} dt dr,$$

and (3.5) follows from the definition of essential oscillation (3.4) and from the estimate (3.6).

Integrating (3.5) with respect to  $x, y \in \Omega$  we get

$$(3.7) \quad \iint_{\Omega \times \Omega} \frac{(\operatorname{ess\,osc}_{[x,y]} u)^p}{|y-x|^{n+sp}} dy dx \leq C_1 \iint_{\Omega \times \Omega} \iint_{[0,1] \times [0,1]} \frac{|u((1-t)x + ty) - u((1-r)x + ry)|^p}{|t-r|^{1+\sigma p} |y-x|^{n+sp}} dt dr dy dx.$$

Applying in the right-hand side of (3.5) the change of variable  $(x, y) \mapsto (w, z) = ((1-t)x + ty, (1-r)x + ry)$ , we get

$$(3.8) \quad \iint_{\Omega \times \Omega} \frac{(\operatorname{ess\,osc}_{[x,y]} u)^p}{|y-x|^{n+sp}} dy dx \leq C_1 \iint_{\Omega \times \Omega} \iint_{\Sigma_{z,w}} \frac{|u(z) - u(w)|^p}{|t-r|^{1-(s-\sigma)p} |z-w|^{n+sp}} dt dr dz dw.$$

where for each  $z, w \in \Omega$  we have defined the set

$$(3.9) \quad \Sigma_{z,w} := \left\{ (t, r) \in [0, 1]^2 \mid \frac{rz-tw}{r-t} \in \Omega \text{ and } \frac{(1-r)z-(1-t)w}{t-r} \in \Omega \right\}.$$

We conclude by estimating the innermost integral in the right-hand side of (3.9) by monotonicity of the integral as

$$(3.10) \quad \iint_{\Sigma_{z,w}} \frac{1}{|t-r|^{1-(s-\sigma)p}} dt dr \leq \iint_{[0,1] \times [0,1]} \frac{1}{|t-r|^{1-(s-\sigma)p}} dt dr = \frac{1}{(s-\sigma)p} \int_0^1 |1-r|^{(s-\sigma)p} + |r|^{(s-\sigma)p} dr < +\infty,$$

since  $\sigma > s$ . The conclusion follows from (3.8) and (3.10).  $\square$

*Proof of theorem 3.1.* Since  $sp > 1$ , for almost every  $[x, y]$ , by the fractional Morrey embedding, the closed set  $\text{ess rg}_{[x,y]} u \subset \text{ess rg}_\Omega u$  is compact and connected, and  $\text{ess rg}_{[x,y]} f \circ u = f(\text{ess rg}_{[x,y]} u)$ , we have thus for almost every  $x, y \in \Omega$ ,

$$(3.11) \quad \begin{aligned} |u(y) - u(x)| &\leq \text{ess osc}_{[x,y]} u = \text{diam}_{[x,y]}(\text{ess rg } u) \\ &\leq \lambda \text{diam}_{[x,y]}(\text{ess rg } f \circ u) = \lambda \text{ess osc}_{[x,y]} f \circ u. \end{aligned}$$

where

$$(3.12) \quad \lambda := \sup \left\{ \frac{\text{diam}(K)}{\text{diam}(f(K))} \mid K \subset \text{ess rg } u \text{ compact, connected and } \text{diam}(K) > 0 \right\}.$$

By (3.11) and the reverse oscillation inequality proposition 3.2, we conclude that there exists a constant  $C$  such that

$$(3.13) \quad \iint_{\Omega \times \Omega} \frac{|u(y) - u(x)|^p}{|y - x|^{n+sp}} dy dx \leq C \lambda^p \iint_{\Omega \times \Omega} \frac{|f(u(y)) - f(u(x))|^p}{|y - x|^{n+sp}} dy dx. \quad \square$$

#### 4. COUNTEREXAMPLES

The following example shows that in the rough case  $sp < 1$ , the fractional reverse estimate theorem 3.1 fails as soon as the function  $f$  is not injective.

**Proposition 4.1.** *Let  $\Omega \subseteq \mathbb{R}^n$ ,  $s \in (0, 1)$  and  $p \in [1, +\infty)$ . If  $sp < 1$  and if the function  $f : \mathbb{R}^m \rightarrow \mathbb{R}^\ell$  is not injective, then there exists a sequence  $(u_j)_{j \in \mathbb{N}}$  in  $\dot{W}^{s,p}(\Omega, \mathbb{R}^m)$  such that for every  $j \in \mathbb{N}$ , the function  $f \circ u_j$  is constant  $\Omega$  and such that*

$$(4.1) \quad \lim_{j \rightarrow \infty} \iint_{\Omega \times \Omega} \frac{|u_j(y) - u_j(x)|^p}{|y - x|^{n+sp}} dy dx = +\infty.$$

*Proof.* We consider the case  $\Omega = (0, 1) \subset \mathbb{R}$ ; the other cases are similar. By assumption, there exist two points  $b_0, b_1 \in \mathbb{R}^m$  such that  $f(b_0) = f(b_1)$ . For each  $j \in \mathbb{N}$  we define the function  $u_j : (0, 1) \rightarrow \mathbb{R}^m$  for every  $x \in (0, 1)$  by

$$(4.2) \quad u_j(x) := \begin{cases} b_0 & \text{if } jx \in [2k, 2k+1) \text{ for some } k \in \mathbb{Z}, \\ b_1 & \text{if } jx \in [2k+1, 2(k+1)) \text{ for some } k \in \mathbb{Z}. \end{cases}$$

By construction, for each  $j \in \mathbb{N}$ , we have  $f(u_j) = f(b_0) = f(b_1)$  everywhere in the interval  $(0, 1)$ . Estimating

$$\begin{aligned}
 \iint_{(0,1) \times (0,1)} \frac{|u_j(y) - u_j(x)|^p}{|y - x|^{1+sp}} dy dx &= \sum_{\ell=0}^{j-1} \iint_{(\frac{\ell}{j}, \frac{\ell+1}{j}) \times (0,1)} \frac{|u_j(y) - u_j(x)|^p}{|y - x|^{1+sp}} dy dx \\
 (4.3) \quad &\leq \sum_{\ell=0}^{j-1} \iint_{(\frac{\ell}{j}, \frac{\ell+1}{j}) \times \mathbb{R} \setminus (\frac{\ell}{j}, \frac{\ell+1}{j})} \frac{|b_0 - b_1|^p}{|y - x|^{1+sp}} dy dx \\
 &= j^{sp} \iint_{(0,1) \times \mathbb{R} \setminus (0,1)} \frac{|b_0 - b_1|^p}{|y - x|^{1+sp}} dy dx = \frac{2j^{sp}|b_0 - b_1|^p}{sp(1 - sp)},
 \end{aligned}$$

we infer that for each  $j \in \mathbb{N}$ , we have  $u_j \in \dot{W}^{s,p}((0, 1), \mathbb{R}^m)$ . Finally, we have if  $j \in \mathbb{N}_*$ ,

$$\begin{aligned}
 \iint_{(0,1) \times (0,1)} \frac{|u_j(y) - u_j(x)|^p}{|y - x|^{1+sp}} dy dx &\geq \sum_{\ell=1}^{j-1} \iint_{(\frac{\ell-1}{j}, \frac{\ell}{j}) \times (\frac{\ell}{j}, \frac{\ell+1}{j})} \frac{|b_0 - b_1|^p}{|y - x|^{1+sp}} dy dx \\
 (4.4) \quad &\geq (1 - \frac{1}{j})j^{sp} \iint_{(-1,0) \times (0,1)} \frac{|b_0 - b_1|^p}{|y - x|^{1+sp}} dy dx \\
 &= \frac{2(1 - \frac{1}{j})j^{sp}(1 - 2^{-sp})}{sp(1 - sp)} |b_1 - b_0|^p,
 \end{aligned}$$

which goes to  $+\infty$  as  $j \rightarrow \infty$ .  $\square$

Finally, in the critical case  $sp = 1$ , the fractional reverse estimate of theorem 3.1 fails when the function  $f$  is Lipschitz continuous and not injective.

**Proposition 4.2.** *Let  $\Omega \subseteq \mathbb{R}^n$ ,  $s \in (0, 1)$  and  $p \in [1, +\infty)$ . If  $sp = 1$ , if the function  $f : \mathbb{R}^m \rightarrow \mathbb{R}^\ell$  is Lipschitz-continuous and is not injective, then there exists a sequence  $(u_j)_{j \in \mathbb{N}}$  in  $\dot{W}^{s,p}(\Omega, \mathbb{R}^m)$  such that*

$$(4.5) \quad \lim_{j \rightarrow \infty} \iint_{\Omega \times \Omega} \frac{|u_j(y) - u_j(x)|^p}{|y - x|^{n+sp}} dy dx = +\infty$$

and

$$(4.6) \quad \sup_{j \in \mathbb{N}} \iint_{\Omega \times \Omega} \frac{|f \circ u_j(y) - f \circ u_j(x)|^p}{|y - x|^{n+sp}} dy dx < +\infty.$$

*Proof.* We concentrate on the case  $\Omega = (-1, 1) \subset \mathbb{R}$ , the other cases being similar. By our assumption, there are two points  $b_0, b_1 \in \mathbb{R}^m$  such that  $f(b_0) = f(b_1)$ . We define the function  $u_* : \mathbb{R} \rightarrow \mathbb{R}^n$  for each  $t \in \mathbb{R}$  by

$$(4.7) \quad u_*(t) := \begin{cases} b_0 & \text{if } t \leq -1, \\ \frac{1-t}{2}b_0 + \frac{1+t}{2}b_1 & \text{if } -1 < t < 1, \\ b_1 & \text{if } t \geq 1. \end{cases}$$

and we define for every  $j \in \mathbb{N}$  the function  $u_j : (-1, 1) \rightarrow \mathbb{R}^m$  by setting for each  $x \in (-1, 1)$ ,  $u_j(x) := u(jx)$ . Since the function  $u_*$  is Lipschitz-continuous, we have  $u_j \in \dot{W}^{s,p}((-1, 1), \mathbb{R}^m)$ . Since  $sp = 1$ , we have for every  $j \in \mathbb{N}$ ,

$$(4.8) \quad \begin{aligned} \iint_{(-1,1) \times (-1,1)} \frac{|f(u_j(y)) - f(u_j(x))|^p}{|y - x|^{1+sp}} dy dx &\leq \iint_{\mathbb{R} \times \mathbb{R}} \frac{|f(u_*(jy)) - f(u_*(jx))|^p}{|y - x|^2} dy dx \\ &= \iint_{\mathbb{R} \times \mathbb{R}} \frac{|f(u_*(y)) - f(u_*(x))|^p}{|y - x|^2} dy dx < +\infty. \end{aligned}$$

On the other hand, we have for every  $j \in \mathbb{N}$

$$(4.9) \quad \iint_{(-1,1) \times (-1,1)} \frac{|u_j(y) - u_j(x)|^p}{|y - x|^{1+sp}} dy dx \geq 2 \int_{-1}^{-\frac{1}{j}} \int_{\frac{1}{j}}^1 \frac{|b_1 - b_0|^p}{|y - x|^2} dy dx = 2|b_1 - b_0|^p \ln \frac{(j+1)^2}{4j},$$

which blows up as  $j \rightarrow \infty$ . □

## REFERENCES

- [1] L. Ambrosio and G. Dal Maso, *A general chain rule for distributional derivatives*, Proc. Amer. Math. Soc. **108** (1990), no. 3, 691–702, doi:10.2307/2047789.
- [2] E. Di Nezza, G. Palatucci, and E. Valdinoci, *Hitchhiker's guide to the fractional Sobolev spaces*, Bull. Sci. Math. **136** (2012), no. 5, 521–573, doi:10.1016/j.bulsci.2011.12.004. MR2944369
- [3] D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*, 2nd ed., Grundlehren der Mathematischen Wissenschaften, vol. 224, Springer, Berlin, 1983.
- [4] G. Leoni, *A first course in Sobolev spaces*, 2nd ed., Graduate Studies in Mathematics, vol. 181, American Mathematical Society, Providence, R.I., 2017.
- [5] M. Marcus and V. J. Mizel, *Absolute continuity on tracks and mappings of Sobolev spaces*, Arch. Rational Mech. Anal. **45** (1972), 294–320, doi:10.1007/BF00251378.
- [6] P. Mironescu and J. Van Schaftingen, *Lifting in compact covering spaces for fractional Sobolev mappings*, to appear in Anal. PDE, available at [arXiv:1907.01373](https://arxiv.org/abs/1907.01373).
- [7] R. Musina and A. I. Nazarov, *A note on truncations in fractional Sobolev spaces*, Bull. Math. Sci. **9**, no. 1, 1950001, 7, doi:10.1142/S1664360719500012.
- [8] M. Willem, *Functional analysis: Fundamentals and applications*, Birkhäuser/Springer, New York, 2013.
- [9] W. P. Ziemer, *Weakly differentiable functions: Sobolev spaces and functions of bounded variation*, Graduate Texts in Mathematics, vol. 120, Springer, New York, 1989.

UNIVERSITÉ CATHOLIQUE DE LOUVAIN, INSTITUT DE RECHERCHE EN MATHÉMATIQUE ET PHYSIQUE,  
CHEMIN DU CYCLOTRON 2 BTE L7.01.01, 1348 LOUVAIN-LA-NEUVE, BELGIUM  
Email address: [Jean.VanSchaftingen@UCLouvain.be](mailto:Jean.VanSchaftingen@UCLouvain.be)