

# The combined basic LP and affine IP relaxation for promise VCSPs on infinite domains\*

Caterina Viola  
University of Oxford, UK  
caterina.viola@cs.ox.ac.uk

Stanislav Živný  
University of Oxford, UK  
standa.zivny@cs.ox.ac.uk

## Abstract

Convex relaxations have been instrumental in solvability of constraint satisfaction problems (CSPs), as well as in the three different generalisations of CSPs: valued CSPs, infinite-domain CSPs, and most recently promise CSPs. In this work, we extend an existing tractability result to the three generalisations of CSPs combined: We give a sufficient condition for the combined basic linear programming and affine integer programming relaxation for exact solvability of promise valued CSPs over infinite-domains. This extends a result of Brakensiek and Guruswami [SODA’20] for promise (non-valued) CSPs (on finite domains).

## 1 Introduction

**Constraint satisfaction** Constraint satisfaction problems (CSPs) are a wide class of computational decision problems. An instance of a CSP is defined by finitely many relations (constraints) that must hold among finitely many given variables; the computational task is to decide whether it is possible to find an assignment of labels from a fixed set (the domain) to the variables so that all the constraints are satisfied. Many problems in computer science (e.g., from artificial intelligence, scheduling, computational linguistic, computational biology and verification) can be modelled as CSPs by choosing an appropriate set of constraints. However, there are many other problems in which some of the constraints may be violated at a cost or in which there are satisfying assignments which are preferable to others. These situations are captured by valued constraint satisfaction problems.

**Valued constraint satisfaction** An instance of a valued constraint satisfaction problem (VCSP) is defined by finitely many cost functions (valued constraints) depending on finitely many given variables and a (rational) threshold; the computational task is to decide whether it is possible to find an assignment of labels from the domain to the variables so that the value of the sum of the cost functions is at most the given threshold. In VCSP instances, cost functions

---

\*An extended abstract of this work appeared in the *Proceedings of the 45th International Symposium on Mathematical Foundations of Computer Science (MFCS’20)* [57]. Stanislav Živný was supported by a Royal Society University Research Fellowship. This project has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No 714532). The paper reflects only the authors’ views and not the views of the ERC or the European Commission. The European Union is not liable for any use that may be made of the information contained therein.

can take on rational or infinite values. VCSPs not only capture optimisation problems but are also a generalisation of CSPs: the non-feasibility of an assignment is modelled by allowing the cost functions to evaluate to  $+\infty$ . A CSP can thus be seen as a VCSP in which the cost functions take values in  $\{0, +\infty\}$ .

**Finite domains** In the case in which the domain (i.e., the fixed set of possible labels for the variables) is a *finite* set the computational complexity of both CSPs and VCSPs have been completely classified. Moreover, in both frameworks a dichotomy theorem holds: every CSP and VCSP is either in P or is NP-complete, depending on some algebraic condition of the underlying set of allowed relations and cost functions, respectively. A dichotomy theorem for CSPs was conjectured by Feder and Vardi [30]. The attempt to prove the conjecture motivated the introduction of the so-called universal algebraic approach [22] for CSPs, which was later extended to VCSPs in [27] and [43], where an analogue of the complexity dichotomy was conjectured for VCSPs. The dichotomy conjectures for finite-domain CSPs and VCSPs inspired an intensive line of research. A complexity classification of finite-domain VCSPs for sets of cost functions taking finite (rational-only) values was established in [52]. A complexity classification of VCSPs was consequently established in [39], assuming a dichotomy for CSPs, which was proved independently in [21] and [58].

**Infinite domains** Although most research on CSPs and VCSPs in the past two decades focused on finite-domain problems, the literature is full of problems (studied independently of CSPs and VCSPs) that can be modelled as CSPs or VCSPs *only* if infinite domains are allowed. For instance, solvability of linear Diophantine equations [26, 37] and the model-checking problem for Kozen’s modal  $\mu$ -calculus [42] are examples of problems that can be modelled as infinite-domain CSPs. Linear Programming, Linear Least Square Regression [16], and Minimum Correlation Clustering [4] are examples of problems that can be modelled as infinite-domain VCSPs. The classes of infinite-domain CSPs and infinite-domain VCSPs are huge! In fact, *every* computational problem over a finite alphabet is polynomial-time Turing-equivalent to an infinite-domain CSP [8]. Therefore, only by focussing on special classes of infinite-domain CSPs (and VCSPs) is it possible to obtain general complexity results. There is a rich literature on the computational complexity of special classes of infinite-domain CSPs, e.g., [11, 10, 15, 36, 9, 13, 14, 7].

**Promise constraint satisfaction** Both infinite-domains CSPs and VCSPs are extensions of the original (finite-domain) CSPs. Promise constraint satisfaction problems (PCSPs) are a third, recently introduced extension of CSPs [17, 23, 5, 31]. Informally, in a PCSP the goal is to find an approximately good solution to a problem under the assumption (the promise) that the problem has a solution. The difference between CSPs and PCSPs is that in a PCSP instance each constraint comes with two relations (not necessarily on the same domain), a “strict” and a “weak” relation. The computational task is then to distinguish between being able to satisfy all the strict constraints versus not being able to satisfy all the weak constraints. A CSP can be seen as a PCSP in which the strict and weak constraints coincide. Perhaps the most well-known example of a PCSP is the approximate graph colouring problem, in which the task is to distinguish  $k$ -colourable graphs from graphs that are not  $c$ -colourable, for some  $c > k$ . (For  $c = k$ , we get the standard  $k$ -colouring problem.) Kazda recently introduced the framework of promise VCSPs on finite domains [38], where he generalised some of the

algebraic reductions from (finite-domain) promise CSPs to (finite-domain) promise VCSPs. As far as we are aware, the only other related work on (finite-domain) promise VCSPs is [3].

**Convex relaxations** One of the most effective ways to design a polynomial-time algorithm for solving combinatorial and optimisation problems is to employ convex relaxations. The idea of convex relaxations is to transform the original problem to an integer program which is then relaxed to a polynomial-time solvable convex program [16], e.g. a linear program. In the context of CSPs, convex relaxations have been studied for robust solvability [44, 29, 6, 28]. Convex relaxations have been also successfully applied to the study of the three extensions of CSPs. For VCSPs, characterisations of the applicability of the basic linear programming relaxation [40], constant levels of the Sherali-Adams linear programming hierarchy [53], and a polynomial-size semidefinite programming relaxation [54] have been provided for exact solvability. In the PCSP framework, the polynomial-time tractability via a specific convex relaxation has been characterised for the basic linear programming relaxation [23], affine integer programming relaxation [23], and their combination [18, 19, 20]. For infinite-domain VCSPs, a sufficient condition has been identified for the solvability via a combination of the basic linear programming relaxation and an efficient sampling algorithm (that is, polynomial-time many-one reduction to a finite-domain VCSP) [12, 56].

## 1.1 Contributions

We initiate the study of convex relaxations for the three generalisations of CSPs *combined*; that is, convex relaxations for promise valued constraint satisfaction problems on infinite-domains. We focus on the combined basic linear programming (BLP) and affine integer programming (AIP) relaxation introduced by Brakensiek and Guruswami [19]. This relaxation is stronger than both the BLP and AIP relaxations individually in the sense that if a class of promise VCSPs is solved by, say, the BLP relaxation then it is also solved by the combined relaxation (and the same holds true for the AIP relaxation). The power of the combined relaxation for (finite-domain) promise CSPs was established in [20]. Rather surprisingly, the combined relaxation gives an algorithm that solves all tractable (non-promise) CSPs on Boolean domains, identified in Schaefer’s work [50], thus giving a unified algorithm.

By extending the argument from [19], we establish a sufficient algebraic condition on the combined relaxation for the solvability of promise VCSPs in which the domain of the “weak cost functions” is possibly infinite (Theorem 8). The proof of this result draws on ideas introduced in [19] but requires a non-trivial amount of technical machinery to make it work in the infinite-domain valued setting. While our relaxation is inspired by [19], it is appropriately modified to work in the optimisation setting (of valued (P)CSPs). We remark that the condition we give is known to be necessary already in special cases of our setting, namely for finite-domain non-valued PCSPs [20]. As an application of our main result, we derive an algebraic condition under which an *infinite-domain* promise VCSP admitting an efficient sampling algorithm can be solved in polynomial time using the combined relaxation (Theorem 16). We emphasise that our main results (Theorems 8 and 16) are appreciatively general, and in particular hold for various special cases of our framework; e.g., for finite-domain promise VCSPs and infinite-domain promise CSPs.

**Approximability of Max-CSPs** PCSPs are approximability problems in which we require that all constraints should be satisfied, although only in a weaker sense. Another very natural

and well-studied form of relaxation is to try to maximise the number of satisfied constraints. Convex relaxations have played a crucial role in this research direction on approximability of (finite-domain) Max-CSPs, going back to the work of Goemans and Williamson [33], e.g., [48, 55, 46, 25, 24, 41, 32].

## 2 Preliminaries

Throughout the paper, we denote by  $x_i$  the  $i$ -th component of a tuple  $x$ . We denote by  $\mathbb{N}$ ,  $\mathbb{Z}$ ,  $\mathbb{Q}$ , and  $\mathbb{Q}_{\geq 0}$  the set of whole numbers, integer numbers, rational numbers, and nonnegative rational numbers, respectively. For every  $m \in \mathbb{N}$ , we denote by  $[m]$  the set  $\{1, \dots, m\} \subset \mathbb{N}$ . Finally, for every  $k \in \mathbb{Q}$  we use the  $\lceil k \rceil$  and  $\lfloor k \rfloor$  to denote the minimum natural number that is at least  $k$  and the maximum natural number that is at most  $k$ , respectively.

### 2.1 Valued Constraint Satisfaction Problems

A *valued structure*  $\Gamma$  (over  $D$ ) consists of a signature  $\tau$  consisting of function symbols  $f$ , each equipped with an arity  $\text{ar}(f)$ ; a set  $D = \text{dom}(\Gamma)$  (the *domain*); and, for each  $f \in \tau$ , a *cost function*, i.e., a function  $f^\Gamma: D^{\text{ar}(f)} \rightarrow \mathbb{Q} \cup \{+\infty\}$ . Here,  $+\infty$  is an extra element with the expected properties that for all  $c \in \mathbb{Q} \cup \{+\infty\}$ , we have  $(+\infty) + c = c + (+\infty) = +\infty$  and  $c < +\infty$  for every  $c \in \mathbb{Q}$ . Given a valued structure  $\Gamma$  with signature  $\tau$ , for every  $f \in \tau$  we define  $\text{dom}(f) := \{t \in D^{\text{ar}(f)} \mid f^\Gamma(t) < +\infty\}$ .

Let  $\Gamma$  be a valued structure with domain  $D$  and signature  $\tau$ . The *valued constraint satisfaction problem* for  $\Gamma$ , denoted by  $\text{VCSP}(\Gamma)$ , is the following computational problem.

An *instance* of  $\text{VCSP}(\Gamma)$  is a triple  $I := (V, \phi, u)$  where  $V$  is a finite set of variables;  $\phi$  is an expression of the form  $\sum_{i=1}^m f_i(v_1^i, \dots, v_{\text{ar}(f_i)}^i)$ , where  $f_1, \dots, f_m \in \tau$  and all the  $v_j^i$  are variables from  $V$  (each summand is called a  $\tau$ -term); and  $u$  is a value from  $\mathbb{Q}$ . The task is to decide whether there exists an assignment  $s: V \rightarrow D$ , whose *cost*, defined as

$$\phi^\Gamma(s(v_1), \dots, s(v_{|V|})) := \sum_{i=1}^m f_i^\Gamma(s(v_1^i), \dots, s(v_{\text{ar}(f_i)}^i))$$

is finite, and if so, whether there is one whose cost is at most  $u$ .

We remark that, given a valued structure  $\Gamma$  over a *finite* signature, the representation of the structure  $\Gamma$  is inessential for computational complexity as  $\Gamma$  is not part of the input.

### 2.2 Fractional Homomorphisms and Fractional Polymorphisms

Let  $X$  be a set. A *discrete probability measure* on  $X$  is a map  $\mu: \mathcal{P}(X) \rightarrow [0, 1]$  such that  $\mu(X) = 1$  and  $\mu$  satisfies the countable additivity property; i.e., for every countable collection  $\{X_n\}_{n \in \mathbb{N}}$  of pairwise disjoint subsets  $X_n \subseteq X$ , it holds that  $\mu(\bigcup_{n \in \mathbb{N}} X_n) = \sum_{n \in \mathbb{N}} \mu(X_n)$ .

Given a probability measure  $\mu$  on a countable set  $X$ , we define its *support* as the set  $\text{Supp}(\mu) := \{x \in X \mid \mu(\{x\}) > 0\}$ . In the reminder, given a probability measure  $\mu$  on a set  $X$ , we use the notation  $\mu(x) := \mu(\{x\})$ . The following proposition is a well-known corollary of countable additivity (proved in Appendix A for completeness).

**Proposition 1.** *Let  $\mu$  be a discrete probability measure on a set  $X$ . Then  $\text{Supp}(\mu)$  is a countable set. Furthermore, if  $X$  is countable then  $\sum_{x \in \text{Supp}(\mu)} \mu(x) = 1$ ; that is,  $\text{Supp}(\mu)$  is non-empty.*

Proposition 1 will guarantee that all supports in this paper are countable sets.

Let  $\mu$  be a discrete probability measure on a countable set  $X$  and let  $Y$  be a random variable with countably many possible outcomes  $y_1, y_2, \dots$  occurring with probabilities  $\mu(x_1), \mu(x_2), \dots$ , respectively. The expectation of  $Y$  associated with  $\mu$  is  $\mathbb{E}_{x \sim \mu}[Y] = \sum_{n \in \mathbb{N}} \mu(x_n) y_n$ .

Let  $C$  and  $D$  be two sets. A map  $g: D^m \rightarrow C$  is called an  $m$ -ary operation. For any  $m \in \mathbb{N}$ , we denote by  $C^{D^m}$  the set of all maps  $g: D^m \rightarrow C$ .

Let  $\Gamma$  and  $\Delta$  be valued structures with the same signature  $\tau$  with domains  $C$  and  $D$ , respectively. A *fractional homomorphism* [51] from  $\Delta$  to  $\Gamma$  is a discrete probability measure  $\chi$  with a non-empty support on  $C^D$  such that for every function symbol  $\gamma \in \tau$  and tuple  $a \in D^{\text{ar}(\gamma)}$ , it holds that

$$\mathbb{E}_{h \sim \chi}[\gamma^\Gamma(h(a))] = \sum_{h \in \text{Supp}(\chi)} \chi(h) \gamma^\Gamma(h(a)) \leq \gamma^\Delta(a), \quad (1)$$

where the functions  $h$  are applied component-wise. We write  $\Delta \rightarrow_f \Gamma$  to indicate the existence of a fractional homomorphism from  $\Delta$  to  $\Gamma$ .

The following proposition, proved for completeness in Appendix A, is adapted from [51], where it was proved in the case of finite-domain valued structure, and appears in [12], where it was stated for valued structures with arbitrary domains and for fractional homomorphisms with finite supports.

**Proposition 2.** *Let  $\Gamma$  and  $\Delta$  be valued structures over the same signature  $\tau$  with domains  $C$  and  $D$ , respectively. Assume  $\Delta \rightarrow_f \Gamma$ . Let  $V = \{v_1, \dots, v_n\}$  be a set of variables and  $\phi$  a sum of finitely many  $\tau$ -terms with variables from  $V$ . For every  $u \in \mathbb{Q}$ , if there exists an assignment  $s: V \rightarrow D$  such that  $\phi^\Delta(s(v_1), \dots, s(v_n)) \leq u$ , then there exists an assignment  $s': V \rightarrow C$  such that  $\phi^\Gamma(s'(v_1), \dots, s'(v_n)) \leq u$ . In particular, it holds that  $\inf_C \phi^\Gamma \leq \inf_D \phi^\Delta$ .*

Let  $\Gamma$  be a valued structure with domain  $C$  and signature  $\tau$ . An  $m$ -ary *fractional polymorphism* of  $\Gamma$  is a discrete probability measure on  $C^{C^m}$  with a non-empty support such that for every  $f \in \tau$  and tuples  $a^1, \dots, a^m \in C^{\text{ar}(f)}$  it holds that

$$\mathbb{E}_{g \sim \omega}[f^\Gamma(g(a^1, \dots, a^m))] = \sum_{g \in C^{C^m}} \omega(g) f^\Gamma(g(a^1, \dots, a^m)) \leq \frac{1}{m} \sum_{i=1}^m f^\Gamma(a^i)$$

(where  $g$  is applied component-wise).

### 2.3 Promise VCSPs

Let  $\Gamma$  and  $\Delta$  be two valued structures over the same signature  $\tau$  with domains  $C$  and  $D$ , respectively. We say that  $(\Delta, \Gamma)$  is a *promise valued template* if there exists a fractional homomorphism from  $\Delta$  to  $\Gamma$ . Given a promise valued template  $(\Delta, \Gamma)$ , the *promise valued constraint satisfaction problem* [38] for  $(\Delta, \Gamma)$ , denoted by  $\text{PVCSP}(\Delta, \Gamma)$ , is the following computational problem.

An *instance*  $I$  of  $\text{PVCSP}(\Delta, \Gamma)$  is a triple  $I := (V, \phi, u)$  where  $V$  is a finite set of variables;  $\phi$  is an expression of the form  $\sum_{i=1}^m f_i(v_1^i, \dots, v_{\text{ar}(f_i)}^i)$ , where  $f_1, \dots, f_m \in \tau$  and all the  $v_j^i$  are variables from  $V$ ; and  $u$  is a value from  $\mathbb{Q}$ .

The task is to output YES if there exists an assignment  $s: V \rightarrow D$  with cost

$$\phi^\Delta(s(v_1), \dots, s(v_{|V|})) := \sum_{i=1}^m f_i^\Delta(s(v_1^i), \dots, s(v_{\text{ar}(f_i)}^i)) \leq u$$

and output NO if *every* assignment  $s': V \rightarrow C$  has cost

$$\phi^\Gamma(s'(v_1), \dots, s'(v_{|V|})) := \sum_{i=1}^m f_i^\Gamma(s'(v_1^i), \dots, s'(v_{\text{ar}(f_i)}^i)) \not\leq u.$$

Note that every valued structure  $\Gamma$  is fractionally homomorphic to itself and thus  $\text{VCSP}(\Gamma)$  is the same as  $\text{PVCSP}(\Gamma, \Gamma)$ .

Let  $(\Delta, \Gamma)$  be a promise valued template. We remark that if the common signature  $\tau$  is finite then the representation of the template is inessential for the computational complexity of  $\text{PVCSP}(\Delta, \Gamma)$  as  $(\Delta, \Gamma)$  is not part of the input.

Let  $e_i^{(m)}: D^m \rightarrow D$  denote the  $m$ -ary projection on  $D$  onto the  $i$ -th coordinate. Let  $\mathcal{J}_D^{(m)} := \{e_1^{(m)}, \dots, e_m^{(m)}\}$ , i.e., the set of all  $m$ -ary projections on  $D$ .

An  $m$ -ary *promise fractional polymorphism*<sup>1</sup> of a promise valued template  $(\Delta, \Gamma)$  is a pair  $\omega := (\omega_I, \omega_O)$  where  $\omega_O$  is a discrete probability measure on  $C^{D^m}$  with a non-empty support and  $\omega_I$  is a discrete probability measure with (finite) support  $\text{Supp}(\omega_I) = \mathcal{J}_D^{(m)}$  such that for every  $f \in \tau$  and tuples  $a^1, \dots, a^m \in D^{\text{ar}(f)}$  it holds that

$$\begin{aligned} \mathbb{E}_{g \sim \omega_O}[f^\Gamma(g(a^1, \dots, a^m))] &= \sum_{g \in \text{Supp}(\omega)} \omega_O(g) f^\Gamma(g(a^1, \dots, a^m)) \\ &\leq \sum_{i=1}^m \omega_I(e_i^{(m)}) f^\Delta(a^i) = \mathbb{E}_{e \sim \omega_I}[f^\Delta(e(a^1, \dots, a^m))]. \end{aligned} \quad (2)$$

**Remark 3.** An  $m$ -ary fractional polymorphism  $\omega$  of a valued structure  $\Gamma$  with domain  $C$  can be seen as an  $m$ -ary promise fractional polymorphism  $\mu = (\mu_I, \mu_O)$  of  $(\Gamma, \Gamma)$  such that  $\mu_O = \omega$  and  $\mu_I(e_i^{(m)}) = \frac{1}{m}$  for  $1 \leq i \leq m$ .

## 2.4 Block-Symmetric Maps

Let  $S_m$  be the symmetric group on  $\{1, \dots, m\}$ . An  $m$ -ary map  $g$  is *fully symmetric* if for every permutation  $\pi \in S_m$ , we have  $g(x_1, \dots, x_m) = g(x_{\pi(1)}, \dots, x_{\pi(m)})$ .

An  $m$ -ary map  $g$  is *block-symmetric* if there exists a partition of the coordinates of  $g$  into blocks  $B_1 \cup \dots \cup B_k = [m]$  such that  $g$  is permutation-invariant within each block  $B_i$ . Let  $\mathcal{P}_{\text{sym}}(g)$  be the set of all partitions into symmetric blocks of  $g$ . For  $B_1 \cup \dots \cup B_k \in \mathcal{P}_{\text{sym}}(m)$ , we define  $w(g, B_1 \cup \dots \cup B_k) := \min_{1 \leq i \leq k} |B_i|$  and we define the *width* of  $g$  to be

$$w(g) := \max\{w(g, B_1 \cup \dots \cup B_k) \mid B_1 \cup \dots \cup B_k \in \mathcal{P}_{\text{sym}}(g)\}.$$

Block-symmetric operation with width 1 are fully symmetric operations. The next example shows operations that are block-symmetric but not fully symmetric.

<sup>1</sup>These are called weighted polymorphisms in [38].



**Example 4.** We consider so-called “moving averages” (see, e.g., [35, 49]). For  $k \in \mathbb{N}$ , a  $k$ -ary 2-period weighted centred moving average is the operation  $\text{wMA}^{(k)}: \mathbb{Q}^k \rightarrow \mathbb{Q}$  defined by

$$\text{wMA}^{(k)}(x^1, \dots, x^k) := \frac{1}{3k} \left( \sum_{i=1}^{\lfloor \frac{k}{4} \rfloor} x^i + 2 \sum_{i=\lfloor \frac{k}{4} \rfloor + 1}^{\lfloor \frac{3k}{4} \rfloor} x^i + \sum_{i=\lfloor \frac{3k}{4} \rfloor + 1}^k x^i \right).$$

It can be verified that a  $k$ -ary 2-period weighted centred moving average, where  $k = 2m + 1$ , is a block-symmetric operation whose symmetric blocks are  $B_1 := \{1, \dots, \lfloor \frac{k}{4} \rfloor, \lfloor \frac{3k}{4} \rfloor + 1, \dots, k\}$  and  $B_2 := \{\lfloor \frac{k}{4} \rfloor + 1, \dots, \lfloor \frac{3k}{4} \rfloor\}$ . Observe that  $|B_1| = \lfloor \frac{k}{2} \rfloor$  and  $|B_2| = \lceil \frac{k}{2} \rceil$ .  $\triangle$

An  $m$ -ary fractional polymorphism  $\omega$  of a valued structure  $\Gamma$  is *block-symmetric* if there exists a partition of the coordinates of  $g$  into blocks  $B_1 \cup \dots \cup B_k = [m]$  such that every operation in  $\text{Supp}(\omega)$  is permutation-invariant within each coordinate block  $B_i$ .

**Example 5.** A  $k$ -ary *average* is a map  $A: \mathbb{Q}^k \rightarrow \mathbb{Q}$  such that, for every  $x \in \mathbb{Q}$ ,  $A(x, \dots, x) = x$  (i.e.,  $A$  is idempotent) and, for every  $x^1, \dots, x^k \in \mathbb{Q}$ , we have  $\min(x^1, \dots, x^k) \leq A(x^1, \dots, x^k) \leq \max(x^1, \dots, x^k)$ , and for every  $\lambda \in \mathbb{Q}$  it holds  $\lambda A(x^1, \dots, x^k) = A(\lambda x^1, \dots, \lambda x^k)$ . Examples of averages include the arithmetic average  $\text{avg}^{(k)}$  defined by  $\text{avg}^{(k)}(x^1, \dots, x^k) = \frac{1}{k}(x^1 + \dots + x^k)$  and the 2-period centred moving average (cf. Example 4).

Let  $A_1$  and  $A_2$  be two different averages. A function  $f: \mathbb{Q} \rightarrow \mathbb{Q} \cup \{+\infty\}$  is called  $(A_1, A_2)$ -convex<sup>2</sup> if for every  $k \in \mathbb{N}$  and every  $x^1, \dots, x^k \in \mathbb{Q}^n$  it holds  $f(A_1(x^1, \dots, x^k)) \leq A_2(f(x^1), \dots, f(x^k))$ .

A valued structure with domain  $\mathbb{Q}$  containing only  $(\text{wMA}, \text{avg})$ -convex functions has, for every  $m \in \mathbb{N}$ , a  $(2m + 1)$ -ary block-symmetric fractional polymorphism  $\omega^{(2m+1)}$  defined by  $\omega^{(2m+1)}(g) = 1$  if  $g = \text{wMA}^{(2m+1)}$  and 0 otherwise.  $\triangle$

Given a promise valued template  $(\Delta, \Gamma)$ , an  $m$ -ary promise fractional polymorphism  $\omega = (\omega_I, \omega_O)$  of  $(\Delta, \Gamma)$  is *block-symmetric* if

- there exists a partition of  $[m]$  into blocks  $B_1 \cup \dots \cup B_k$  such that every map in  $\text{Supp}(\omega_O)$  is a permutation-invariant within each coordinate block  $B_i$ , and
- $\sum_{i \in B_j} \omega_I(e_i^{(m)}) = \frac{|B_j|}{m}$  for every  $j \in \{1, \dots, k\}$ .

The proof of the following lemma can be found in Appendix A.

**Lemma 6.** Let  $(\Delta, \Gamma)$  be a promise valued template and let  $m \in \mathbb{N}$ . If  $\omega = (\omega_I, \omega_O)$  is an  $m$ -ary block symmetric promise fractional polymorphism of  $(\Delta, \Gamma)$ , then also  $\omega' = (\omega'_I, \omega_O)$ , where  $\omega'_I(e_i^{(m)}) = \frac{1}{m}$  for  $1 \leq i \leq m$ , is an  $m$ -ary block-symmetric promise fractional polymorphism of  $(\Delta, \Gamma)$ .

In view of Lemma 6, we will assume without loss of generality that any  $m$ -ary block-symmetric promise fractional polymorphism  $\omega = (\omega_I, \omega_O)$  is such that  $\omega_I$  assigns  $\frac{1}{m}$  to each  $m$ -ary projection on the domain of  $\Delta$  and we will identify  $\omega$  with  $\omega_O$ .

<sup>2</sup>In [1, 2], the notion of  $(A_1, A_2)$ -convexity is defined for  $A_1, A_2$  symmetric averages. An average  $A$  is symmetric if for every  $k \in \mathbb{N}$ , and every  $x^1, \dots, x^k \in \mathbb{Q}$ , it holds  $A(x^1, \dots, x^k) = A(x^{\pi(1)}, \dots, x^{\pi(k)})$  for every  $\pi \in S_k$ .

## 2.5 The Basic Linear Programming Relaxation

Every VCSP over a finite domain has a natural linear programming relaxation. Let  $\Delta$  be a valued structure with finite domain  $D$  and signature  $\tau$ . Let  $I$  be an instance of VCSP( $\Delta$ ) with set of variables  $V = \{x_1, \dots, x_d\}$ , objective function  $\phi(x_1, \dots, x_d) = \sum_{j \in J} f_j(x_1^j, \dots, x_{n_j}^j)$ , with  $f_j \in \tau$ ,  $x^j = (x_1^j, \dots, x_{n_j}^j) \in V^{n_j}$ , for all  $j \in J$  (the set  $J$  is finite and indexing the cost functions that are summands of  $\phi$ ), and a threshold  $u \in \mathbb{Q}$ .<sup>3</sup> Define the sets of variables as follows:  $W_1 := \{\lambda_j(t) \mid j \in J \text{ and } t \in D^{n_j}\}$ ,  $W_2 := \{\mu_{x_i}(a) \mid x_i \in V \text{ and } a \in D\}$ , and  $W := W_1 \cup W_2$ . Then the *basic linear programming* (BLP) relaxation associated to  $I$  (see [51], [40], and references therein) is a linear program with variables  $W$  and is defined in Figure 1.

subject to	$\text{BLP}(I, \Delta) := \min \sum_{j \in J} \sum_{t \in D^{n_j}} \lambda_j(t) f_j^\Delta(t)$
$\sum_{t \in D^{n_j} : t_\ell = a} \lambda_j(t) = \mu_{x_\ell^j}(a)$	for all $j \in J, \ell \in \{1, \dots, n_j\}, a \in D$ ,
$\sum_{a \in D} \mu_{x_i}(a) = 1$	for all $x_i \in V$ ,
$\lambda_j(t) = 0$	for all $j \in J, t \notin \text{dom}(f_j)$ ,
$0 \leq \lambda_j(t), \mu_{x_i}(a) \leq 1$	for all $\lambda_j(t) \in W_1, \mu_{x_i}(a) \in W_2$ .

Figure 1: BLP

We remark that a solution to the BLP also satisfies the constraints  $\sum_{t \in D^{n_j}} \lambda_j(t) = 1$  for all  $j \in J$ . If there is no feasible solution to the BLP then  $\text{BLP}(I, \Delta) = +\infty$ . For a finite-domain VCSP instance, the corresponding BLP relaxation can be computed in polynomial time.

We note that it is not difficult to lift the existing results characterising the power of BLP (in terms of fully symmetric operations) for CSPs [44], VCSPs [40], promise CSPs [23, 5], and infinite-domain VCSPs [12, 56] to our setting of promise VCSPs with infinite domains, cf. Theorem 20 in Appendix B. Our focus, however, is on the stronger combined relaxation presented in Section 3.

## 2.6 The Affine Integer Programming Relaxation

Let  $\Delta$  be a valued structure with finite domain  $D$  and signature  $\tau$ . Let  $I$  be an instance of VCSP( $\Delta$ ) with set of variables  $V = \{x_1, \dots, x_d\}$ , and objective function  $\phi(x_1, \dots, x_d) = \sum_{j \in J} f_j(x_1^j, \dots, x_{n_j}^j)$ , with  $f_j \in \tau$ ,  $x^j = (x_1^j, \dots, x_{n_j}^j) \in V^{n_j}$ , for all  $j \in J$  (the set  $J$  is finite and indexing the cost functions that are summands of  $\phi$ ), and a threshold  $u \in \mathbb{Q}$ .<sup>4</sup> Define the sets of variables as follows:  $R_1 := \{q_j(t) \mid j \in J \text{ and } t \in D^{n_j}\}$ ,  $R_2 := \{r_{x_i}(a) \mid x_i \in V \text{ and } a \in D\}$ , and  $R := R_1 \cup R_2$ . Then the *affine integer programming* (AIP) relaxation associated to  $I$  [18, 19] is an integer program with variables  $R$  and is defined in Figure 2.

<sup>3</sup>Note that the BLP relaxation does not depend on the threshold  $u$ .

<sup>4</sup>Note that the AIP relaxation does not depend on the threshold  $u$ .



$$\begin{array}{l}
\text{subject to} \quad \text{AIP}(I, \Delta) := \min \sum_{j \in J} \sum_{t \in D^{n_j}} q_j(t) f_j^\Delta(t) \\
\\
\sum_{t \in D^{n_j} : t_\ell = a} q_j(t) = r_{x_\ell^j}(a) \quad \text{for all } j \in J, \ell \in \{1, \dots, n_j\}, a \in D, \\
\\
\sum_{a \in D} r_{x_i}(a) = 1 \quad \text{for all } x_i \in V, \\
\\
q_j(t) = 0 \quad \text{for all } j \in J, t \notin \text{dom}(f_j), \\
q_j(t), r_{x_i}(a) \in \mathbb{Z} \quad \text{for all } q_j(t) \in R_1, r_{x_i}(a) \in R_2.
\end{array}$$

Figure 2: AIP

We remark that a solution to the AIP also satisfies the constraints  $\sum_{t \in D^{n_j}} \lambda_q(t) = 1$  for all  $j \in J$ . If there is no feasible solution to the AIP then  $\text{AIP}(I, \Delta) = +\infty$ . For a finite-domain VCSP instance, the corresponding AIP relaxation can be computed in polynomial time. Since the feasibility version of AIP can be solved in polynomial time [37, 18], (the optimisation version of) AIP can be solved in (oracle) polynomial time using an oracle for the feasibility version of the problem (see [34, Theorem 6.4.9]).

### 3 The Combined BLP and AIP Relaxation for PVCSPs

Let  $(\Delta, \Gamma)$  be a promise valued template such that the domain of  $\Delta$  is a *finite* set. We may solve  $\text{PVCSP}(\Delta, \Gamma)$  by using a *combination* of the BLP relaxation and the AIP relaxation of  $\Delta$ , as proposed (for finite-domain promise non-valued) CSPs in [19], appropriately modified to the valued setting.

To describe such an algorithm, we need the following definition.

**Definition 7.** Let  $\Delta$  be a valued structure with finite domain  $D$  and signature  $\tau$ . Let us consider an instance  $I := (V, \phi, u)$  of  $\text{VCSP}(\Delta)$  such that  $\phi(x_1, \dots, x_d) = \sum_{j \in J} f_j(x_1^j, \dots, x_{n_j}^j)$ . Assume that  $\text{BLP}(I, \Delta) \leq u$ . We define  $(\lambda^*, \mu^*)$  as follows.

- If there exists a relative interior point of the rational feasibility polytope of  $\text{BLP}(I, \Delta)$  with cost at most  $u$ ,<sup>5</sup> then  $(\lambda^*, \mu^*)$  is such a point;
- otherwise,  $(\lambda^*, \mu^*)$  is defined to be a point from the relative interior of the optimal polytope of  $\text{BLP}(I, \Delta)$ .<sup>6</sup>

The *refinement* of  $\text{AIP}(I, \Delta)$  with respect to  $(\lambda^*, \mu^*)$  is the integer program  $\text{AIP}^*(I, \Delta)$  obtained by adding to  $\text{AIP}(I, \Delta)$  the constraints

$$\begin{array}{ll}
q_j(t) = 0 & \text{for every } j \in J, t \in D^{n_j} \text{ such that } \lambda_j^*(t) = 0, \\
r_{x_i}(a) = 0 & \text{for every } x_i \in V, a \in D \text{ such that } \mu_{x_i}^*(a) = 0.
\end{array}$$

<sup>5</sup>There is a polynomial-time algorithm [34, 19] that decides the existence of a relative interior point in the rational feasibility polytope of  $\text{BLP}(I, \Delta)$  with cost at most  $u$  and, in the case it exists, finds it.

<sup>6</sup>Such a point can be found in polynomial time by applying the algorithm in [34, 19] to the feasibility linear program defined by adding to the constraints defining the feasibility polytope of  $\text{BLP}(I, \Delta)$  the additional constraint  $\sum_{j \in J} \sum_{t \in D^{n_j}} \lambda_j(t) f_j^\Delta(t) = u$ .

---

**ALGORITHM 1:** The combined BLP + AIP Relaxation Algorithm for PVCSP( $\Delta, \Gamma$ )

---

**Input:**
 $I := (V, \phi, u)$ , a valid instance of PVCSP( $\Delta, \Gamma$ )

**Output:**

YES if there exists an assignment  $s: V \rightarrow \text{dom}(\Delta)$  such that  $\phi^\Delta(s(x_1), \dots, s(x_{|V|})) \leq u$   
 NO if there is no assignment  $s: V \rightarrow \text{dom}(\Gamma)$  such that  $\phi^\Gamma(s(x_1), \dots, s(x_{|V|})) \leq u$

 BLP( $I, \Delta$ );

**if** BLP( $I, \Delta$ )  $\not\leq u$  **then**

output NO;

**else**

 ( $\lambda^*, \mu^*$ ), as in Definition 7;

 $\text{AIP}^*(I, \Delta) := \text{refinement of } \text{AIP}(I, \Delta) \text{ with respect to } (\lambda^*, \mu^*), \text{ as in Definition 7;}$ 
**if**  $\text{AIP}^*(I, \Delta) \not\leq u$  **then**

output NO;

**else**

output YES;

**end**
**end**


---

As our main result, we now present a sufficient condition under which Algorithm 1 correctly solves PVCSP( $\Delta, \Gamma$ ).

**Theorem 8.** *Let  $(\Delta, \Gamma)$  be a promise valued template such that  $\Delta$  has a finite domain. Assume that for all  $L \in \mathbb{N}$  there exists a block-symmetric promise fractional polymorphism of  $(\Delta, \Gamma)$  with arity  $2L + 1$  having two symmetric blocks of size  $L + 1$  and  $L$ , respectively. Then Algorithm 1 correctly solves PVCSP( $\Delta, \Gamma$ ) (in polynomial time).*

Note that in Theorem 8 the domain of the valued structure  $\Gamma$  can be finite or (countably) infinite.

To prove Theorem 8 we need to use a preliminary lemma and the notion of a *bimultiset-structures*. Let  $\Delta$  be a valued  $\tau$ -structure with domain  $D$ , let  $L \in \mathbb{N}$ , and let  $B_1 \cup B_2$  any partition of  $[2L + 1]$  such that  $|B_1| = L + 1$  and  $|B_2| = L$ . The *bimultiset-structure*  $\mathcal{B}_{B_1, B_2}^{2L+1}(\Delta)$  is the valued structure with domain  $\left(\binom{D}{L+1}\right) \times \left(\binom{D}{L}\right)$  i.e., the set whose elements  $(\alpha, \beta)$  are pairs of multisets of elements from  $D$  of size  $L + 1$  and of size  $L$ , respectively. For every  $k$ -ary function symbol  $f \in \tau$ , and  $(\alpha_1, \beta_1) \dots, (\alpha_k, \beta_k) \in \left(\binom{D}{L+1}\right) \times \left(\binom{D}{L}\right)$  the function  $f^{\mathcal{B}_{B_1, B_2}^{2L+1}(\Delta)}$  is defined as follows

$$f^{\mathcal{B}_{B_1, B_2}^{2L+1}(\Delta)}((\alpha_1, \beta_1) \dots, (\alpha_k, \beta_k)) := \frac{1}{2L + 1} \min_{\substack{t^1, \dots, t^k \in D^{2L+1}: \\ \{t^\ell\}_{B_1} = \alpha_\ell, \{t^\ell\}_{B_2} = \beta_\ell}} \sum_{i=1}^{2L+1} f^\Delta(t_i^1, \dots, t_i^k),$$

where  $\{t^\ell\}_{B_1}$  denotes the multiset  $\{t_i^\ell \mid i \in B_1\}$  and  $\{t^\ell\}_{B_2}$  denotes the multiset  $\{t_i^\ell \mid i \in B_2\}$ .

**Lemma 9.** *Let  $(\Delta, \Gamma)$  be a promise valued template such that  $\Delta$  has a finite domain. Let  $L \in \mathbb{N}$  and assume that  $(\Delta, \Gamma)$  has a block-symmetric promise fractional polymorphism of arity  $2L + 1$  with two symmetric blocks  $B_1$  and  $B_2$  of size  $L + 1$  and  $L$ , respectively. Then  $\mathcal{B}_{B_1, B_2}^{2L+1}(\Delta)$  is fractionally homomorphic to  $\Gamma$ .*

*Proof.* Let  $C$  be the (possibly infinite) domain of  $\Gamma$ , let  $D$  be the finite domain of  $\Delta$ , and let  $\tau$  be the common signature of  $\Gamma$  and  $\Delta$ . Let  $\omega$  be the  $(2L + 1)$ -ary block-symmetric promise

fractional polymorphism of  $(\Delta, \Gamma)$  with symmetric blocks  $B_1$  and  $B_2$  of size  $L+1$  and  $L$ , respectively. For every  $g \in \text{Supp}(\omega) \subseteq C^{D^{2L+1}}$  we define  $\tilde{g}: \left(\binom{D}{L+1}\right) \times \left(\binom{D}{L}\right) \rightarrow C$  by setting, for every  $\alpha = \{\xi^j \mid j \in B_1\} \in \left(\binom{D}{L+1}\right)$  and every  $\beta = \{\xi^j \mid j \in B_2\} \in \left(\binom{D}{L}\right)$ ,

$$\tilde{g}((\alpha, \beta)) = g(\xi^1, \xi^2, \dots, \xi^{2L+1}).$$

Observe that  $\tilde{g}$  is well defined as  $g$  is block-symmetric with symmetric blocks  $B_1$  and  $B_2$  (the order of the coordinates from a same block does not matter). We define the discrete probability measure  $\chi$  on  $C^{\left(\binom{D}{L+1}\right) \times \left(\binom{D}{L}\right)}$  as follows

$$\chi(Y) = \sum_{\substack{g \in \text{Supp}(\omega): \\ \tilde{g} \in Y}} \omega(g), \quad \text{for every } Y \subseteq \left(\binom{D}{L+1}\right) \times \left(\binom{D}{L}\right).$$

Observe that  $\chi$  satisfies the countable additivity property, since  $\omega$  does. Furthermore,  $\text{Supp}(\chi) = \{h \in C^{D^{2L+1}} \mid h = \tilde{g} \text{ for some } g \in \text{Supp}(\omega)\} = \{\tilde{g} \in C^{D^{2L+1}} \mid g \in \text{Supp}(\omega)\}$  is countable by Proposition 1 and it holds that

$$\sum_{h \in \text{Supp}(\chi)} \chi(h) = \sum_{g \in \text{Supp}(\omega)} \omega(g) = 1.$$

We claim that  $\chi$  is a fractional homomorphism from  $\mathcal{B}_{B_1, B_2}^{2L+1}(\Delta)$  to  $\Gamma$ . Indeed, for every  $f \in \tau$  and every tuple  $((\alpha_1, \beta_1), \dots, (\alpha_k, \beta_k)) \in \left(\left(\binom{D}{L+1}\right) \times \left(\binom{D}{L}\right)\right)^k$  with  $\alpha_i := \{\xi_i^j \mid j \in B_1\}$ ,  $\beta_i := \{\xi_i^j \mid j \in B_2\}$ , and  $k := \text{ar}(f)$ , it holds that

$$\begin{aligned} & \sum_{h \in C^{\left(\binom{D}{L+1}\right) \times \left(\binom{D}{L}\right)}} \chi(h) f^\Gamma(h((\alpha_1, \beta_1)), \dots, h((\alpha_k, \beta_k))) \\ &= \sum_{g \in \text{Supp}(\omega)} \omega(g) f^\Gamma(g(\xi_1^1, \dots, \xi_1^{2L+1}), \dots, g(\xi_k^1, \dots, \xi_k^{2L+1})) \\ &= \sum_{g \in \text{Supp}(\omega)} \omega(g) f^\Gamma(g(\xi_1^{\pi_1(1)}, \dots, \xi_1^{\pi_1(2L+1)}), \dots, g(\xi_k^{\pi_k(1)}, \dots, \xi_k^{\pi_k(2L+1)})) \end{aligned} \quad (3)$$

$$\leq \frac{1}{2L+1} \sum_{i=1}^{2L+1} f^\Delta(\xi_1^{\pi_1(i)}, \dots, \xi_k^{\pi_k(i)}) \quad (4)$$

for every  $\pi_1, \dots, \pi_k \in S_{2L+1}$  such that  $\pi_i$  is permutation-invariant within  $B_1$  and within  $B_2$  for  $1 \leq i \leq k$ . Equality (3) holds because the maps  $g \in \text{Supp}(\omega)$  are block-symmetric with symmetric blocks  $B_1$  and  $B_2$ . Inequality (4) holds because  $\omega$  is a promise fractional polymorphism of  $(\Delta, \Gamma)$ . Then, in particular, we obtain

$$\begin{aligned} & \sum_{h \in C^{\left(\binom{D}{L+1}\right) \times \left(\binom{D}{L}\right)}} \chi(h) f^\Gamma(h((\alpha_1, \beta_1), \dots, (\alpha_k, \beta_k))) \\ & \leq \frac{1}{2L+1} \min_{\substack{t^1, \dots, t^k \in D^{2L+1}: \\ \{t^\ell\}_{B_1} = \alpha_\ell, \{t^\ell\}_{B_2} = \beta_\ell}} \sum_{i=1}^{2L+1} f^\Delta(t_i^1, \dots, t_i^k) = f^{\mathcal{B}_{B_1, B_2}^{2L+1}(\Delta)}((\alpha_1, \beta_1), \dots, (\alpha_k, \beta_k)). \quad \square \end{aligned}$$

**Remark 10.** Although we do not need it for our main result, we note that the converse of Lemma 9 holds true. More precisely, let  $(\Delta, \Gamma)$  be a promise valued template such that  $\Delta$  has a finite domain. Let  $L \in \mathbb{N}$  and let  $B_1 \cup B_2$  be a partition of  $[2L+1]$  such that  $|B_1| = L+1$  and  $|B_2| = L$ . If  $\mathcal{B}_{B_1, B_2}^{2L+1}(\Delta)$  is fractionally homomorphic to  $\Gamma$ , then  $(\Delta, \Gamma)$  has a block-symmetric promise fractional polymorphism of arity  $2L+1$  with symmetric blocks  $B_1$  and  $B_2$ . To show this we reason as in the proof of [51, Lemma 2.2]. Let  $D$  be the domain of  $\Delta$ , and let  $\chi$  be a fractional homomorphism from  $\mathcal{B}_{B_1, B_2}^{2L+1}(\Delta)$  to  $\Gamma$ . Let us define  $h: D^{2L+1} \rightarrow \left(\binom{D}{L+1}\right) \times \left(\binom{D}{L}\right)$  such that it maps a tuple  $a = (a_1, \dots, a_m) \in D^{2L+1}$  to the bimultiset  $(\{a\}_{B_1}, \{a\}_{B_2})$ . Then, it is easily verified that the discrete probability measure  $\omega$  on  $C^{D^m}$  such that

$$\omega(g') = \sum_{\substack{g \in \text{Supp}(\chi): \\ g \circ h = g'}} \omega(g)$$

is the desired promise fractional polymorphism.

*Proof of Theorem 8.* Let  $C$  be the (possibly infinite) domain of  $\Gamma$  and let  $D$  be the finite domain of  $\Delta$ . Let  $\tau$  be the common signature of  $\Delta$  and  $\Gamma$ . Let  $I$  be an instance of  $\text{PVCSP}(\Delta, \Gamma)$  with variables  $V = \{x_1, \dots, x_n\}$ , objective function  $\phi(x_1, \dots, x_n) = \sum_{j \in J} \gamma_j(x^j)$  where  $J$  is a finite set of indices,  $\gamma_j \in \Gamma$ , and  $x^j \in V^{\text{ar}(j)}$ , and threshold  $u$ .

Assume that  $\min_D \phi^\Delta \leq u$ . Our goal is to show that Algorithm 1 outputs YES. Since  $\min_D \phi^\Delta \leq u$  we have  $\text{BLP}(I, \Delta) \leq u$  and in particular we have that either  $\text{BLP}(I, \Delta) < u$ , which by linearity implies the existence of a relative interior point in the feasibility polytope of  $\text{BLP}(I, \Delta)$  with value at most  $u$ ; or  $\text{BLP}(I, \Delta) = u = \min_D \phi^\Delta$ . In the first case, each coordinate of  $(\lambda^*, \mu^*)$  is positive if and only if the same coordinate is positive at some point in the feasibility polytope of the BLP. Therefore, the feasibility lattice of  $\text{AIP}^*(I, \Delta)$  includes every possible assignment which is in the support of some feasible solution to  $\text{BLP}(I, \Delta)$ , including integral solutions and as a consequence  $\text{AIP}^*(I, \Delta) \leq \min_D \phi^\Delta \leq u$ . In the second case, each coordinate of  $(\lambda^*, \mu^*)$  is positive if and only if the same coordinate is positive at some point in the optimal polytope of the BLP. Therefore, the feasibility lattice of  $\text{AIP}^*(I, \Delta)$  includes every possible assignment which is in the support of some optimal solution to  $\text{BLP}(I, \Delta)$ , including integral solutions and as a consequence  $\text{AIP}^*(I, \Delta) \leq \min_D \phi^\Delta = u$ . Thus, in both cases,  $\text{BLP}(I, \Delta) \leq u$  and  $\text{AIP}^*(I, \Delta) \leq u$  and hence Algorithm 1 indeed outputs YES, as required.

In the other direction, we want to show (by contrapositive) that if Algorithm 1 outputs YES then  $\inf_C \phi^\Gamma \leq u$ . Thus, assume that  $\text{BLP}(I, \Delta) \leq u$  and  $\text{AIP}^*(I, \Delta) \leq u$ . Let  $(\lambda^*, \mu^*)$  be as in Definition 7 and denote  $\text{BLP}^*(I, \Delta) := \sum_{j \in J} \sum_{t \in D^{n_j}} \lambda_j^*(t) f_j^\Delta(t)$ ; observe that  $\text{BLP}^*(I, \Delta) \leq u$  by the definition of  $(\lambda^*, \mu^*)$ . Let  $(q^*, r^*)$  be a solution to  $\text{AIP}^*(I, \Delta)$  with objective value at most  $u$ . Let  $\ell$  be a positive integer such that  $\ell \cdot \lambda^*$ , and  $\ell \cdot \mu^*$  are both integral, and let  $M$  be the maximum of the absolute values of the coordinates of both  $q^*$  and  $r^*$ . Let us set  $L := (M+1)\ell$ . From  $\text{BLP}^*(I, \Delta) \leq u$  and  $\text{AIP}^*(I, \Delta) \leq u$  it immediately follows that

$$\frac{2(M+1)\ell}{2(M+1)\ell+1} \text{BLP}^*(I, \Delta) + \frac{1}{2(M+1)\ell+1} \text{AIP}^*(I, \Delta) \leq u. \quad (5)$$

We claim that

$$\min_{\left(\binom{D}{L+1}\right) \times \left(\binom{D}{L}\right)} \phi^{\mathcal{B}_{B_1, B_2}^{2L+1}(\Delta)} \leq \frac{2(M+1)\ell}{2(M+1)\ell+1} \text{BLP}^*(I, \Delta) + \frac{1}{2(M+1)\ell+1} \text{AIP}^*(I, \Delta) \quad (6)$$

for all the partitions  $B_1 \cup B_2 = [2L + 1]$  such that  $|B_1| = L + 1$  and  $|B_2| = L$ . To prove the claim, let us define, for every  $i \in \{1, \dots, n\}$  and for every  $a \in D$ , the following nonnegative integers

$$\begin{aligned} W_{x_i, B_1}(a) &:= (M + 1)\ell\mu_{x_i}^*(a) + r_{x_i}^*(a), \\ W_{x_i, B_2}(a) &:= (M + 1)\ell\mu_{x_i}^*(a). \end{aligned}$$

(To check that  $W_{x_i, B_1}(a)$  and  $W_{x_i, B_2}(a)$  are nonnegative it is enough to observe that if  $\mu_{x_i}^*(a)$  is 0 then, by Definition 7,  $r_{x_i}^*(a)$  is also 0, otherwise  $\mu_{x_i}^*(a)$  is at least  $\frac{1}{\ell}$ , and the positivity of  $W_{x_i, B_1}(a)$ ,  $W_{x_i, B_2}(a)$  follows by the choice of  $M$ .) Observe that for every  $i \in \{1, \dots, n\}$  we have that

$$\begin{aligned} \sum_{a \in D} W_{x_i, B_1}(a) &= (M + 1)\ell \sum_{a \in D} \mu_{x_i}^*(a) + \sum_{a \in D} r_{x_i}^*(a) = (M + 1)\ell + 1 = L + 1, \\ \sum_{a \in D} W_{x_i, B_2}(a) &= (M + 1)\ell \sum_{a \in D} \mu_{x_i}^*(a) = (M + 1)\ell = L. \end{aligned}$$

Let  $\nu: V \rightarrow \left(\binom{D}{L+1}\right) \times \left(\binom{D}{L}\right)$  be the map defined, for every  $x_i \in V$ , by  $\nu(x_i) = (\alpha_i, \beta_i)$ , where  $\alpha_i$  is the multiset of  $\left(\binom{D}{L+1}\right)$  that contains  $W_{x_i, B_1}(a)$  many occurrences of  $a$ , for every  $a \in D$ , and  $\beta_i$  is the multiset of  $\left(\binom{D}{L}\right)$  that contains  $W_{x_i, B_2}(a)$  many occurrences of  $a$ , for every  $a \in D$ .

Let  $f_j$  be a  $k$ -ary function symbol appearing as a term of the objective function  $\phi$ . Let us define, for every  $t \in D^k$ , the following nonnegative integers

$$\begin{aligned} P_{j, B_1}(t) &:= (M + 1)\ell\lambda_j^*(t) + q_j^*(t), \\ P_{j, B_2}(t) &:= (M + 1)\ell\lambda_j^*(t). \end{aligned}$$

Observe that

$$\begin{aligned} \sum_{t \in D^k} P_{j, B_1}(t) &= (M + 1)\ell \sum_{t \in D^k} \lambda_j^*(t) + \sum_{t \in D^k} q_j^*(t) = (M + 1)\ell + 1 = L + 1, \\ \sum_{t \in D^k} P_{j, B_2}(t) &= (M + 1)\ell \sum_{t \in D^k} \lambda_j^*(t) = (M + 1)\ell = L. \end{aligned}$$

We write now

$$\sum_{t \in D^k} P_{j, B_1}(t) f_j^\Delta(t) = \sum_{h=1}^{(M+1)\ell+1} f_j^\Delta(\zeta_1^h, \dots, \zeta_k^h)$$

where  $\zeta^1, \dots, \zeta^{(M+1)\ell+1}$  are defined to be  $(M + 1)\ell + 1$  elements of  $D^k$  such that  $P_{j, B_1}(t)$  many of them are equal to  $t$ , for every  $t \in D^k$ ; and

$$\sum_{t \in D^k} P_{j, B_2}(t) f_j^\Delta(t) = \sum_{h=1}^{(M+1)\ell} f_j^\Delta(\xi_1^h, \dots, \xi_k^h)$$

where  $\xi^1, \dots, \xi^{(M+1)\ell}$  are defined to be  $(M+1)\ell$  elements of  $D^k$  such that  $P_{j,B_2}(t)$  many of them are equal to  $t$ , for every  $t \in D^k$ .

We obtain

$$\begin{aligned}
& \frac{2(M+1)\ell}{2(M+1)\ell+1} \lambda_j^*(t) f_j^\Delta(t) + \frac{1}{2(M+1)\ell+1} q_j^*(t) f_j^\Delta(t) \\
&= \frac{1}{2(M+1)\ell+1} \left( \sum_{t \in D^k} P_{j,B_1}(t) f_j^\Delta(t) + \sum_{t \in D^k} P_{j,B_2}(t) f_j^\Delta(t) \right) \\
&= \frac{1}{2L+1} \left( \sum_{h=1}^{L+1} f_j^\Delta(\zeta_1^h, \dots, \zeta_k^h) + \sum_{h=1}^L f_j^\Delta(\xi_1^h, \dots, \xi_k^h) \right) \\
&\geq \frac{1}{2L+1} \min_{t^1, \dots, t^k \in D^m: \{t^\ell\}_{B_1} = \zeta_\ell, \{t^\ell\}_{B_2} = \xi_\ell} \sum_{i=1}^{2L+1} f^\Delta(t_i^1, \dots, t_i^k) \\
&= f_j^{\mathcal{B}_{B_1, B_2}^{2L+1}(\Delta)}((\zeta_1, \xi_1), \dots, (\zeta_k, \xi_k)) = f_j^{\mathcal{B}_{B_1, B_2}^{2L+1}(\Delta)}(\nu(x_1^j), \dots, \nu(x_k^j)),
\end{aligned}$$

where the last equality follows because, for every  $a \in D$ , the number of  $a$ 's in  $\zeta_h$  is

$$\begin{aligned}
& \sum_{t \in D^k: t_h = a} P_{j,B_1}(t) = (M+1)\ell \sum_{t \in D^k: t_h = a} \lambda_j^*(t) + \sum_{t \in D^k: t_h = a} q_j^*(t) \\
&= (M+1)\ell \mu_{x_h^j}^*(a) + r_{x_h^j}^*(a) = W_{x_h^j, B_1}(a),
\end{aligned}$$

and, for every  $a \in D$ , the number of  $a$ 's in  $\xi_h$  is

$$\sum_{t \in D^k: t_h = a} P_{j,B_2}(t) = (M+1)\ell \sum_{t \in D^k: t_h = a} \lambda_j^*(t) = (M+1)\ell \mu_{x_h^j}^*(a) = W_{x_h^j, B_2}(a).$$

This proves the claim.

From Inequalities (5) and (6) it follows that for all partitions  $B_1 \cup B_2 = [2L+1]$  such that  $|B_1| = L+1$  and  $|B_2| = L$  it holds

$$\min_{\left(\binom{D}{L+1}\right) \times \left(\binom{D}{L}\right)} \phi^{\mathcal{B}_{B_1, B_2}^{2L+1}(\Delta)} \leq u.$$

Moreover, since there exists a block-symmetric promise fractional polymorphisms of  $(\Delta, \Gamma)$  of arity  $2L+1$  having two symmetric blocks  $B_1$  and  $B_2$  with respective size  $L+1$  and  $L$ , Lemma 9 implies the existence of a fractional homomorphism from  $\mathcal{B}_{B_1, B_2}^{2L+1}(\Delta)$  to  $\Gamma$ . From Proposition 2 it follows that

$$\inf_C \phi^\Gamma \leq \min_{\left(\binom{D}{L+1}\right) \times \left(\binom{D}{L}\right)} \phi^{\mathcal{B}_{B_1, B_2}^{2L+1}(\Delta)} \leq u$$

and this concludes the proof.  $\square$

We conclude the section with a couple of remarks showing that the number of symmetric blocks and their size do not play a crucial role in the proofs of Theorem 8 and Lemma 9. The notion of bimultiset-structure can be straightforwardly generalised to the notion of  $k$ -multiset-structure with blocks of size  $b_1, \dots, b_k$ .



**Remark 11.** The proof of Lemma 9 can be easily adapted to the case in which the promise template  $(\Delta, \Gamma)$  has a block-symmetric promise fractional polymorphism  $\omega$  of arity  $L \in \mathbb{N}$  with an arbitrary number  $k$  of symmetric blocks of size  $b_1, \dots, b_k$  (given that the blocks are the same for each map in  $\text{Supp}(\omega)$ ). In this case, the  $k$ -multiset structure  $\mathcal{B}_{B_1, \dots, B_k}(\Delta)$  is fractionally homomorphic to  $\Gamma$ . The converse (corresponding to Remark 10) is also true.

**Remark 12.** Let  $(\Delta, \Gamma)$  be a promise valued template such that  $\Delta$  has a *finite domain*. Assume that  $(\Delta, \Gamma)$  has block-symmetric promise fractional polymorphisms with arbitrarily many blocks of arbitrarily large size. Then Algorithm 1 correctly solves  $\text{PVCSP}(\Delta, \Gamma)$ . This statement can be proved by a slight modification of the proof of Theorem 8 employing a block-symmetric promise fractional polymorphism whose symmetric blocks  $B_1, \dots, B_k$  have each size at least  $M\ell^2$  and where the coefficients  $W_{x_i, B_b(a)}$ , and  $P_{j, B_b}(t)$  are defined as in the proof of [20, Theorem 4.1].

Finally, we point out that for finite-domain PCSPs, the condition of having a block-symmetric promise polymorphism of arity  $2L + 1$  is equivalent to the one of having a block-symmetric promise polymorphism of arity  $2L + 1$  with two symmetric blocks of arity  $L + 1$  and  $L$ , for every  $L \in \mathbb{N}$  [20].

## 4 The Combined Relaxation with Sampling for PVCSPs

We use the notion of a sampling algorithm for a valued structure from [12].

**Definition 13.** Let  $\Gamma$  be a valued structure with domain  $C$  and finite signature  $\tau$ . A *sampling algorithm* for  $\Gamma$  takes as input a positive integer  $d$  and computes a finite-domain valued structure  $\Delta$  fractionally homomorphic to  $\Gamma$  such that, for every finite sum  $\phi$  of  $\tau$ -terms having at most  $d$  distinct variables,  $V = \{x_1, \dots, x_d\}$ , and every  $u \in \mathbb{Q}$ , there exists a solution  $s: V \rightarrow C$  with  $\phi^\Gamma(s(x_1), \dots, s(x_d)) \leq u$  if and only if there exists a solution  $h': V \rightarrow D$  with  $\phi^\Delta(h'(x_1), \dots, h'(x_d)) \leq u$ . A sampling algorithm is called *efficient* if its running time is bounded by a polynomial in  $d$ . The finite-domain valued structure computed by a sampling algorithm is called a *sample*.

**Example 14.** A valued structure  $\Gamma$  with domain  $\mathbb{Q}$  and signature  $\tau$  is called *piecewise linear homogeneous (PLH)* if, for every  $\gamma \in \tau$ , the cost function  $\gamma^\Gamma$  is first-order definable over the structure  $\mathfrak{L} = (\mathbb{Q}; \leq, 1, \{c \cdot\}_{c \in \mathbb{Q}})$  where

- $<$  is a relation symbol (i.e., a  $\{0, \infty\}$ -valued function symbol) of arity 2 and  $<^\mathfrak{L}$  is the strict linear order of  $\mathbb{Q}$ ,
- 1 is a constant symbol and  $1^\mathfrak{L} := 1 \in \mathbb{Q}$ , and
- $c \cdot$  is a unary function symbol for every  $c \in \mathbb{Q}$  such that  $(c \cdot)^\mathfrak{L}$  is the function  $x \mapsto cx$ , i.e., the multiplication by  $c$ .

If  $\Gamma$  is a PLH valued structure with a finite signature, then it admits an efficient sampling algorithm [12].  $\triangle$

Let  $\Gamma$  be a valued structure with a finite signature that admits an efficient sampling algorithm. Observe that for every finite-domain valued structure  $\Delta_d$  computed by such an efficient sampling algorithm, the pair  $(\Delta_d, \Gamma)$  is a promise valued template. The following lemma follows from the definition of a sampling algorithm for a valued structure.

**Lemma 15.** *Let  $(\Gamma_1, \Gamma_2)$  be a promise valued template with a finite signature. Assume that  $\Gamma_1$  admits an efficient sampling algorithm. If  $\text{PVCSP}(\Delta_d, \Gamma_2)$  is polynomial-time solvable for every finite-domain valued structure  $\Delta_d$  computed by an efficient sampling algorithm for  $\Gamma_1$ , then  $\text{PVCSP}(\Gamma_1, \Gamma_2)$  is polynomial-time solvable.*

*Proof.* It can be verified that  $(\Delta_d, \Gamma_2)$  is a valid promise valued template, for every  $d \in \mathbb{N}$ ; that is, the existence of a fractional homomorphism from  $\Delta_d$  to  $\Gamma_1$  and a fractional homomorphism from  $\Gamma_1$  to  $\Gamma_2$  implies the existence of a fractional homomorphism from  $\Delta_d$  to  $\Gamma_2$ , which is obtained by a composition of the two fractional homomorphisms in the same fashion as in the proof of Theorem 16.

Let  $I := (V, \phi, u)$  be an instance of  $\text{PVCSP}(\Gamma_1, \Gamma_2)$  and let  $d := |V|$ . Let  $C_1, C_2$  and  $D_d$  be the domains of  $\Gamma_1, \Gamma_2$  and  $\Delta_d$ , respectively. Assume that there is a polynomial-time algorithm solving  $\text{PVCSP}(\Delta_d, \Gamma_2)$ . If the output of such an algorithm is NO then clearly the answer to  $\text{PVCSP}(\Gamma_1, \Gamma_2)$  is NO. Otherwise, if the output of such an algorithm is YES, then by the definition of sampling algorithm the answer to  $\text{PVCSP}(\Gamma_1, \Gamma_2)$  is YES.  $\square$

**Theorem 16.** *Let  $(\Gamma_1, \Gamma_2)$  be a promise valued template with a finite signature. Assume that  $\Gamma_1$  admits an efficient sampling algorithm. Moreover, assume that  $(\Gamma_1, \Gamma_2)$  has a block-symmetric promise fractional polymorphism of arity  $2L + 1$  with two symmetric blocks of size  $L + 1$  and  $L$ , respectively, for all  $L \in \mathbb{N}$ . Then  $\text{PVCSP}(\Gamma_1, \Gamma_2)$  is polynomial-time solvable.*

Note that in Theorem 16 (and in Lemma 15) both the respective domains of the valued structures  $\Gamma_1$  and  $\Gamma_2$  have arbitrary (countable) cardinality, that is, each of  $\Gamma_1$  and  $\Gamma_2$  can have a finite or an infinite domain.

*Proof.* For every positive integer  $d$  let  $\Delta_d$  be the finite-domain valued structure  $\Delta_d$  computed by an efficient sampling algorithm for  $\Gamma_1$  on input  $d$ . By Lemma 15, to prove that  $\text{PVCSP}(\Gamma_1, \Gamma_2)$  is polynomial-time solvable it is enough to prove that  $\text{PVCSP}(\Delta_d, \Gamma_2)$  is polynomial-time solvable for every positive integer  $d$ . We claim that for every  $d, L \in \mathbb{N}$  there exists a block-symmetric promise fractional polymorphism of  $(\Delta_d, \Gamma_2)$  with arity  $2L + 1$  having two symmetric blocks of size  $L + 1$  and  $L$ , respectively. By Theorem 8, proving our claim will imply that  $\text{PVCSP}(\Delta_d, \Gamma_2)$  is polynomial-time solvable and therefore, by Lemma 15, that  $\text{PVCSP}(\Gamma_1, \Gamma_2)$  is polynomial-time solvable.

We now prove the claim. Let  $C_1, C_2$  and  $D_d$  be the domains of  $\Gamma_1, \Gamma_2$  and  $\Delta_d$ , respectively, and let  $\tau$  be the signature of  $\Gamma_1$  and  $\Gamma_2$  (by the definition of sampling algorithm it is also the signature of  $\Delta_d$ ). By the definition of sampling algorithm, there exists a fractional homomorphism  $\chi$  from  $\Delta_d$  to  $\Gamma_1$  and, by assumption, there exists a block-symmetric promise fractional polymorphism  $\omega := (\omega_I, \omega_O)$  of  $(\Gamma_1, \Gamma_2)$  with two symmetric blocks  $B_1$  and  $B_2$  having size  $L + 1$  and  $L$ , respectively. For every  $h \in \text{Supp}(\chi) \subseteq C_1^{D_d}$  and every  $g \in \text{Supp}(\omega_O) \subseteq C_2^{C_1^{2L+1}}$  we define the map  $g \circ h: D_d^{2L+1} \rightarrow C_2$  by setting, for every  $a^1, \dots, a^{2L+1} \in D_d^{2L+1}$ ,

$$g \circ h(a^1, \dots, a^{2L+1}) = g(h(a^1), \dots, h(a^{2L+1})).$$

Observe that  $g \circ h$  is a block-symmetric map with symmetric block  $B_1$  and  $B_2$ , because so is  $g$ . We define the discrete probability measure  $\mu_O$  on  $C_2^{D_d^{2L+1}}$  as follows

$$\mu_O(Y) = \sum_{h \in \text{Supp}(\chi)} \sum_{\substack{g \in \text{Supp}(\omega_O): \\ g \circ h \in Y}} \chi(h) \omega_O(g), \quad \text{for every } Y \subseteq D_d^{2L+1}.$$

Since  $\sum_{h \in \text{Supp}(\chi)} \chi(h)$  and  $\sum_{g \in \text{Supp}(\omega_O)} \omega_O(g)$  are absolutely convergent series, it follows by Martens' Theorem (see, e.g., [45, Theorem 6.57]) that  $\mu_O$  is well defined. Observe that  $\mu_O$  satisfies the countable additivity property, since  $\omega_O$  and  $\chi$  do. Furthermore,  $\text{Supp}(\mu_O) = \{(g \circ h) \mid h \in \text{Supp}(\chi), g \in \text{Supp}(\omega_O)\}$  is countable and, again by Martens' Theorem, it holds

$$\sum_{g' \in \text{Supp}(\mu_O)} \mu_O(g') = \sum_{h \in \text{Supp}(\chi)} \chi(h) \sum_{g \in \text{Supp}(\omega_O)} \omega_O(g) = 1.$$

Let  $\mu_I$  be the discrete probability measure in  $\mathcal{J}_D^{(2L+1)}$  such that  $\mu_I(e_i^{(2L+1)}) = \frac{1}{2L+1}$ . We now show that  $\mu := (\mu_I, \mu_O)$  is a promise fractional polymorphism of  $(\Delta_d, \Gamma_1)$  with arity  $2L+1$ . By definition,  $\mu$  is block-symmetric with two symmetric blocks of respective size  $L+1$  and  $L$ .

For every  $f \in \tau$  and every  $a^1, \dots, a^{2L+1} \in D_d^{\text{ar}(f)}$  it holds that

$$\begin{aligned} & \sum_{g' \in \text{Supp}(\mu_O)} \mu_O(g') f^{\Gamma_2}(g(a^1, \dots, a^{2L+1})) \\ &= \sum_{h \in \text{Supp}(\chi)} \chi(h) \sum_{g \in \text{Supp}(\omega_O)} \omega_O(g) f^{\Gamma_2}(g(h(a^1), \dots, h(a^{2L+1}))) \\ &\leq \sum_{h \in \text{Supp}(\chi)} \chi(h) \frac{1}{2L+1} \sum_{i=1}^{2L+1} f^{\Gamma_1}(g(h(a^i))) \end{aligned} \tag{7}$$

$$\begin{aligned} &= \frac{1}{2L+1} \sum_{i=1}^{2L+1} \sum_{h \in \text{Supp}(\chi)} \chi(h) f^{\Gamma_1}(h(a^i)) \\ &\leq \frac{1}{2L+1} \sum_{i=1}^{2L+1} f^{\Delta_d}(a^i), \end{aligned} \tag{8}$$

where Inequality (7) holds because  $\omega$  is a promise fractional polymorphism of  $(\Gamma_1, \Gamma_2)$  and Inequality (8) holds because  $\chi$  is a fractional homomorphism from  $\Delta_d$  to  $\Gamma_1$ . We obtained that  $\mu$  is a promise fractional polymorphism of  $(\Delta_d, \Gamma_2)$  and this concludes the proof.  $\square$

In the particular case in which  $\Gamma_1 = \Gamma_2$ , from Theorem 16 we obtain the following result for infinite-domain (non-promise) VCSPs.

**Corollary 17.** *Let  $\Gamma$  be a valued structure with a finite signature that admits an efficient sampling algorithm. Assume that  $\Gamma$  has a block-symmetric fractional polymorphism of arity  $2L+1$  with two symmetric blocks of size  $L+1$  and  $L$ , respectively, for all  $L \in \mathbb{N}$ . Then  $\text{VCSP}(\Gamma)$  is polynomial-time solvable.*

Corollary 17 and the existence of an efficient sampling algorithm for PLH valued structures (see Example 14) imply the following tractability result for (wMA, avg)-convex PLH valued structures.

**Corollary 18.** *Let wMA be the 2-period centred weighted moving average. Let  $\Gamma$  be a PLH valued structure with a finite signature such that every cost function from  $\Gamma$  is a (wMA, avg)-convex cost function (see Example 5). Then  $\text{VCSP}(\Gamma)$  is polynomial-time solvable.*

## References

- [1] G.D. Anderson, M.K. Vamanamurthy, and M. Vuorinen. Generalized convexity and inequalities. *Journal of Mathematical Analysis and Applications*, 335(2):1294 – 1308, 2007. [doi:10.1016/j.jmaa.2007.02.016](https://doi.org/10.1016/j.jmaa.2007.02.016).
- [2] G Aumann. Konvexe Funktionen und Induktion bei Ungleichungen zwischen Mittelwerten. *Akad. Wiss. Math.-Natur. Kl. Abh., Math. Ann*, 109:405–413, 1933. URL: [https://www.zobodat.at/pdf/Sitz-Ber-Akad-Muenchen-math-Kl\\_1933\\_0403-0415.pdf](https://www.zobodat.at/pdf/Sitz-Ber-Akad-Muenchen-math-Kl_1933_0403-0415.pdf).
- [3] Per Austrin and Johan Håstad. On the usefulness of predicates. *ACM Transactions on Computation Theory*, 5(1):1:1–1:24, 2013. [arXiv:1204.5662](https://arxiv.org/abs/1204.5662), [doi:10.1145/2462896.2462897](https://doi.org/10.1145/2462896.2462897).
- [4] Nikhil Bansal, Avrim Blum, and Shuchi Chawla. Correlation clustering. *Machine Learning*, 56(1-3):89–113, 2004. [doi:10.1023/B:MACH.0000033116.57574.95](https://doi.org/10.1023/B:MACH.0000033116.57574.95).
- [5] Libor Barto, Jakub Bulín, Andrei Krokhin, and Jakub Opršal. Algebraic approach to promise constraint satisfaction. 2019. [arXiv:1811.00970v3](https://arxiv.org/abs/1811.00970v3).
- [6] Libor Barto and Marcin Kozik. Robustly solvable constraint satisfaction problems. *SIAM Journal on Computing*, 45(4):1646–1669, 2016. [arXiv:1512.01157](https://arxiv.org/abs/1512.01157), [doi:10.1137/130915479](https://doi.org/10.1137/130915479).
- [7] Libor Barto and Michael Pinsker. Topology Is Irrelevant (In a Dichotomy Conjecture for Infinite Domain Constraint Satisfaction Problems). *SIAM Journal on Computing*, 49(2):365–393, 2020. [arXiv:1909.06201](https://arxiv.org/abs/1909.06201), [doi:10.1137/18M1216213](https://doi.org/10.1137/18M1216213).
- [8] Manuel Bodirsky and Martin Grohe. Non-dichotomies in constraint satisfaction complexity. In *Proceedings of the 35th International Colloquium on Automata, Languages and Programming (ICALP)*, pages 184 –196. Springer Verlag, 2008. [doi:10.1007/978-3-540-70583-3\\_16](https://doi.org/10.1007/978-3-540-70583-3_16).
- [9] Manuel Bodirsky, Peter Jonsson, and Trung Van Pham. The complexity of phylogeny constraint satisfaction problems. *ACM Transactions on Computational Logic*, 18(3), 2017. [arXiv:1503.07310](https://arxiv.org/abs/1503.07310), [doi:10.1145/3105907](https://doi.org/10.1145/3105907).
- [10] Manuel Bodirsky, Peter Jonsson, and Timo Von Oertzen. Essential convexity and complexity of semi-algebraic constraints. *Logical Methods in Computer Science*, 8(4), 2012. [arXiv:1210.0420](https://arxiv.org/abs/1210.0420), [doi:10.2168/LMCS-8\(4:5\)2012](https://doi.org/10.2168/LMCS-8(4:5)2012).
- [11] Manuel Bodirsky and Jan Kára. The complexity of temporal constraint satisfaction problems. *Journal of the ACM*, 57(2), 2010. Article 9. [doi:10.1145/1667053.1667058](https://doi.org/10.1145/1667053.1667058).
- [12] Manuel Bodirsky, Marcello Mamino, and Caterina Viola. Piecewise linear valued CSPs solvable by linear programming relaxation. 2019. [arXiv:1912.09298](https://arxiv.org/abs/1912.09298).
- [13] Manuel Bodirsky, Barnaby Martin, and Antoine Mottet. Discrete temporal constraint satisfaction problems. *Journal of the ACM*, 65(2):9:1–9:41, 2018. [arXiv:1503.08572](https://arxiv.org/abs/1503.08572), [doi:10.1145/3154832](https://doi.org/10.1145/3154832).

- [14] Manuel Bodirsky, Barnaby Martin, Michael Pinsker, and András Pongrácz. Constraint satisfaction problems for reducts of homogeneous graphs. *SIAM Journal on Computing*, 48(4):1224–1264, 2019. [arXiv:1602.05819](#), [doi:10.1137/16M1082974](#).
- [15] Manuel Bodirsky and Michael Pinsker. Schaefer’s theorem for graphs. *Journal of the ACM*, 62(3):19:1–19:52, 2015. [arXiv:1011.2894](#), [doi:10.1145/2764899](#).
- [16] Stephen P. Boyd and Lieven Vandenbergh. *Convex Optimization*. Cambridge University Press, 2004. URL: [https://web.stanford.edu/~boyd/cvxbook/bv\\_cvxbook.pdf](https://web.stanford.edu/~boyd/cvxbook/bv_cvxbook.pdf).
- [17] Joshua Brakensiek and Venkatesan Guruswami. Promise constraint satisfaction: Structure theory and a symmetric boolean dichotomy. In *Proceedings of the 29th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA’18)*, pages 1782–1801. SIAM, 2018. [arXiv:1704.01937](#), [doi:10.1137/1.9781611975031.117](#).
- [18] Joshua Brakensiek and Venkatesan Guruswami. An Algorithmic Blend of LPs and Ring Equations for Promise CSPs. In *Proceedings of the 30th Annual ACM-SIAM Symposium on Discrete Algorithms (SODA’19)*, pages 436–455. SIAM, 2019. [arXiv:1807.05194](#), [doi:10.1137/1.9781611975482.28](#).
- [19] Joshua Brakensiek and Venkatesan Guruswami. Symmetric polymorphisms and efficient decidability of promise CSPs. In *Proceedings of the 31st ACM-SIAM Symposium on Discrete Algorithms (SODA’20)*, pages 297–304, 2020. [arXiv:1907.04383](#), [doi:10.1137/1.9781611975994.18](#).
- [20] Joshua Brakensiek, Venkatesan Guruswami, Marcin Wrochna, and Stanislav Živný. The power of the combined basic LP and affine relaxation for promise CSPs. *SIAM Journal on Computing*, 49:1232–1248, 2020. [arXiv:1907.04383](#), [doi:10.1137/20M1312745](#).
- [21] Andrei A. Bulatov. A dichotomy theorem for nonuniform CSPs. In *Proceedings of the 58th IEEE Annual Symposium on Foundations of Computer Science (FOCS)*, pages 319–330, 2017. [arXiv:1703.03021](#), [doi:10.1109/FOCS.2017.37](#).
- [22] Andrei A. Bulatov, Andrei A. Krokhin, and Peter G. Jeavons. Classifying the complexity of constraints using finite algebras. *SIAM Journal on Computing*, 34(3):720–742, 2005. [doi:10.1137/S0097539700376676](#).
- [23] Jakub Bulín, Andrei Krokhin, and Jakub Opršal. Algebraic approach to promise constraint satisfaction. In *Proceedings of the 51st ACM Annual Symposium on Theory of Computing (STOC’19)*, page 602–613. ACM, 2019. [doi:10.1145/3313276.3316300](#).
- [24] Siu On Chan. Approximation resistance from pairwise-independent subgroups. *Journal of the ACM*, 63(3), 2016. Article No. 27. [arXiv:2012/110](#), [doi:10.1145/2873054](#).
- [25] Siu On Chan, James R. Lee, Prasad Raghavendra, and David Steurer. Approximate constraint satisfaction requires large LP relaxations. *Journal of the ACM*, 63(4):34:1–34:22, 2016. [arXiv:1309.0563](#), [doi:10.1145/2811255](#).
- [26] Tsu-Wu J. Chou and George E. Collins. Algorithms for the solution of systems of linear Diophantine equations. *SIAM Journal on Computing*, 11:687–708, 1982. [doi:10.1137/0211057](#).

- [27] David A. Cohen, Martin C. Cooper, Páidí Creed, Peter Jeavons, and Stanislav Živný. An algebraic theory of complexity for discrete optimisation. *SIAM Journal on Computing*, 42(5):915–1939, 2013. [arXiv:1207.6692](#), [doi:10.1137/130906398](#).
- [28] Víctor Dalmau, Marcin Kozik, Andrei A. Krokhin, Konstantin Makarychev, Yury Makarychev, and Jakub Opršal. Robust algorithms with polynomial loss for near-unanimity csp. *SIAM Journal on Computing*, 48(6):1763–1795, 2019. [arXiv:1607.04787](#), [doi:10.1137/18M1163932](#).
- [29] Víctor Dalmau and Andrei A. Krokhin. Robust satisfiability for CSPs: Hardness and algorithmic results. *ACM Transactions on Computation Theory*, 5(4):15:1–15:25, 2013. [doi:10.1145/2540090](#).
- [30] Tomás Feder and Moshe Y. Vardi. The computational structure of monotone monadic SNP and constraint satisfaction: a study through Datalog and group theory. *SIAM Journal on Computing*, 28(1):57–104, 1999. [doi:10.1137/S0097539794266766](#).
- [31] Miron Ficak, Marcin Kozik, Miroslav Olšák, and Szymon Stankiewicz. Dichotomy for symmetric Boolean PCSPs. In *Proceedings of the 46th International Colloquium on Automata, Languages, and Programming (ICALP’19)*, volume 132 of *LIPIcs*, pages 57:1–57:12. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2019. [arXiv:1904.12424](#), [doi:10.4230/LIPIcs.ICALP.2019.57](#).
- [32] Mrinalkanti Ghosh and Madhur Tulsiani. From weak to strong linear programming gaps for all constraint satisfaction problems. *Theory of Computing*, 14(1):1–33, 2018. [arXiv:1608.00497](#), [doi:10.4086/toc.2018.v014a010](#).
- [33] Michel X. Goemans and David P. Williamson. Improved approximation algorithms for maximum cut and satisfiability problems using semidefinite programming. *Journal of the ACM*, 42(6):1115–1145, 1995. [doi:10.1145/227683.227684](#).
- [34] Martin Grötschel, László Lovász, and Alexander Schrijver. *Geometric Algorithms and Combinatorial Optimization*, volume 2. Springer-Verlag Berlin Heidelberg, 1993. [doi:10.1007/978-3-642-78240-4](#).
- [35] Xiao Hu, Zhong Xiao, and Ni Zhang. Removal of baseline wander from ecg signal based on a statistical weighted moving average filter. *Journal of Zhejiang University SCIENCE C*, 12(5):397–403, 2011. [doi:10.1631/jzus.C1010311](#).
- [36] Peter Jonsson and Johan Thapper. Constraint satisfaction and semilinear expansions of addition over the rationals and the reals. *Journal of Computer and System Sciences*, 82(5):912–928, 2016. [arXiv:1506.00479](#), [doi:10.1016/j.jcss.2016.03.002](#).
- [37] Ravindran Kannan and Achim Bachem. Polynomial algorithms for computing the Smith and Hermite normal forms of an integer matrix. *SIAM Journal on Computing*, 8(4):499–507, 1979. [doi:10.1137/0208040](#).
- [38] Alexander Kazda. Minion homomorphisms give reductions between promise valued CSPs. 2021. In preparation.



- [39] Vladimir Kolmogorov, Andrei A. Krokhin, and Michal Rolínek. The complexity of general-valued csps. *SIAM Journal on Computing*, 46(3):1087–1110, 2017. [arXiv:1502.07327](#), [doi:10.1137/16M1091836](#).
- [40] Vladimir Kolmogorov, Johan Thapper, and Stanislav Živný. The power of linear programming for general-valued CSPs. *SIAM Journal on Computing*, 44(1):1–36, 2015. [arXiv:1311.4219](#), [doi:10.1137/130945648](#).
- [41] Pravesh K. Kothari, Raghu Meka, and Prasad Raghavendra. Approximating rectangles by juntas and weakly-exponential lower bounds for LP relaxations of CSPs. In *Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing (STOC’17)*, pages 590–603, 2017. [arXiv:1610.02704](#), [doi:10.1145/3055399.3055438](#).
- [42] Dexter Kozen. Results on the propositional  $\mu$ -calculus. *Theoretical Computer Science*, 27:333–354, 1983. [doi:10.1016/0304-3975\(82\)90125-6](#).
- [43] Marcin Kozik and Joanna Ochremiak. Algebraic properties of valued constraint satisfaction problem. In *Proceedings of the 42nd International Colloquium on Automata, Languages and Programming (ICALP’15)*, volume 9134 of *Lecture Notes in Computer Science*, pages 846–858. Springer, 2015. [arXiv:1403.0476](#), [doi:10.1007/978-3-662-47672-7\\_69](#).
- [44] Gábor Kun, Ryan O’Donnell, Suguru Tamaki, Yuichi Yoshida, and Yuan Zhou. Linear programming, width-1 CSPs, and robust satisfaction. In *Proceedings of the 3rd Innovations in Theoretical Computer Science (ITCS’12)*, pages 484–495. ACM, 2012. [doi:10.1145/2090236.2090274](#).
- [45] Miklós Laczkovich and Vera T Sós. *Real Analysis: Series, Functions of Several Variables, and Applications*, volume 3. Springer, New York, NY, 2017. [doi:10.1007/978-1-4939-7369-9](#).
- [46] James R. Lee, Prasad Raghavendra, and David Steurer. Lower bounds on the size of semidefinite programming relaxations. In *Proceedings of the 47th Symposium on Theory of Computing (STOC’15)*, pages 567–576, 2015. [arXiv:1411.6317](#), [doi:10.1145/2746539.2746599](#).
- [47] M. Loève. *Elementary Probability Theory*, pages 1–52. Springer, New York, NY, 1977. [doi:10.1007/978-1-4684-9464-8\\_1](#).
- [48] Prasad Raghavendra. Optimal algorithms and inapproximability results for every CSP? In *Proceedings of the 40th Annual ACM Symposium on Theory of Computing (STOC’08)*, pages 245–254, 2008. [doi:10.1145/1374376.1374414](#).
- [49] Aistis Raudys, Vaidotas Lenčiauskas, and Edmundas Malčius. Moving averages for financial data smoothing. In *Proceedings of the 19th International Conference on Information and Software Technologies (ICIST’13)*, volume 403 of *Communications in Computer and Information Science*, pages 34–45. Springer, 2013. [doi:10.1007/978-3-642-41947-8\\_4](#).
- [50] Thomas J. Schaefer. The Complexity of Satisfiability Problems. In *Proceedings of the 10th Annual ACM Symposium on Theory of Computing (STOC’78)*, pages 216–226. ACM, 1978. [doi:10.1145/800133.804350](#).

- [51] Johan Thapper and Stanislav Živný. The power of linear programming for valued CSPs. In *Proceedings of the 53rd IEEE Annual Symposium on Foundations of Computer Science (FOCS'12)*, pages 669–678. IEEE Computer Society, 2012. [arXiv:1204.1079](#), [doi:10.1109/FOCS.2012.25](#).
- [52] Johan Thapper and Stanislav Živný. The complexity of finite-valued CSPs. *Journal of the ACM*, 63(4), 2016. Article No. 37. [arXiv:1210.2987](#), [doi:10.1145/2974019](#).
- [53] Johan Thapper and Stanislav Živný. The Power of Sherali-Adams Relaxations for General-Valued CSPs. *SIAM Journal on Computing*, 46(4):1241–1279, 2017. [arXiv:1606.02577](#), [doi:10.1137/16M1079245](#).
- [54] Johan Thapper and Stanislav Živný. The limits of SDP relaxations for general-valued csp. *ACM Transactions on Computation Theory*, 10(3):12:1–12:22, 2018. [arXiv:1612.01147](#), [doi:10.1145/3201777](#).
- [55] Madhur Tulsiani. CSP gaps and reductions in the Lasserre hierarchy. In *Proceedings of the 41st Annual ACM Symposium on Theory of Computing (STOC'09)*, pages 303–312. ACM, 2009. [doi:10.1145/1536414.1536457](#).
- [56] Caterina Viola. *Valued Constraint Satisfaction Problems over Infinite Domains*. PhD thesis, Technische Universität Dresden, Dresden, Germany, 2020.
- [57] Caterina Viola and Stanislav Živný. The combined basic LP and affine IP relaxation for promise VCSPs on infinite domains. In *Proceedings of the 45th International Symposium on Mathematical Foundations of Computer Science (MFCS'20)*, volume 170, pages 82:1–82:15, Dagstuhl, Germany, 2020. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik. [arXiv:2007.01779](#), [doi:10.4230/LIPIcs.MFCS.2020.85](#).
- [58] Dmitriy Zhuk. A proof of the CSP dichotomy conjecture. *J. ACM*, 67(5):30:1–30:78, 2020. [arXiv:1704.01914](#), [doi:10.1145/3402029](#).

## A Omitted Proofs from Section 2

Let  $\mu$  be a discrete probability measure as defined in Section 2. We note that countable additivity implies that  $\mu(\emptyset) = 0$  and that  $\mu(X_1) \leq \mu(X_2)$  for every  $X_1 \subseteq X_2$ , i.e.,  $\mu$  is monotone. We also remark that while the set  $X$  might be countable or uncountable, the terminology *discrete probability measure* refers to the fact that the  $\sigma$ -algebra is  $\mathcal{P}(X)$ , i.e., the discrete topology on  $X$  (see [47, Examples 2.8 and 2.9]).

**Proposition** (Proposition 1 restated). Let  $\mu$  be a discrete probability measure on a set  $X$ . Then  $\text{Supp}(\mu)$  is a countable set. Furthermore, if  $X$  is countable then  $\sum_{x \in \text{Supp}(\mu)} \mu(x) = 1$ ; that is,  $\text{Supp}(\mu)$  is non-empty.

*Proof.* Let us define, for all  $n \in \mathbb{N}$ , the subset  $A_n := \{x \in X \mid \mu(\{x\}) \geq \frac{1}{n}\}$ . We can write the support of  $\mu$  as  $\text{Supp}(\mu) = \bigcup_{n \in \mathbb{N}} A_n$ . Observe that, for every  $n \in \mathbb{N}$ , the set  $A_n$  is finite. If this was not the case, then there would exist a sequence  $(x_i)_{i \in \mathbb{N}} \in (A_n)^{\mathbb{N}}$  of pairwise distinct elements and, by the countable additivity property, we would obtain that  $\mu(\{x_i \mid i \in \mathbb{N}\}) = \sum_{i \in \mathbb{N}} \mu(\{x_i\}) \geq \sum_{i \in \mathbb{N}} \frac{1}{n} = +\infty$ , which contradicts the assumption that  $\mu(\{x_i \mid i \in \mathbb{N}\}) \leq \mu(X) = 1$ . Therefore,  $\text{Supp}(\mu)$  is countable, because it is the countable union of finite sets.  $\square$

**Proposition** (Proposition 2 restated). Let  $\Gamma$  and  $\Delta$  be valued structures over the same signature  $\tau$  with domains  $C$  and  $D$ , respectively. Assume  $\Delta \rightarrow_f \Gamma$ . Let  $V = \{v_1, \dots, v_n\}$  be a set of variables and  $\phi$  a sum of finitely many  $\tau$ -terms with variables from  $V$ . For every  $u \in \mathbb{Q}$ , if there exists an assignment  $s: V \rightarrow D$  such that  $\phi^\Delta(s(v_1), \dots, s(v_n)) \leq u$ , then there exists an assignment  $s': V \rightarrow C$  such that  $\phi^\Gamma(s'(v_1), \dots, s'(v_n)) \leq u$ . In particular, it holds that

$$\inf_C \phi^\Gamma \leq \inf_D \phi^\Delta.$$

*Proof.* Let  $\phi(v_1, \dots, v_n) := \sum_{j \in J} \gamma_j(v^j)$ , where  $\gamma_j \in \tau$  and  $v^j \in V^{ar(\gamma_j)}$ ; and let  $u \in \mathbb{Q}$ . Let  $\chi$  be a fractional homomorphism from  $\Delta$  to  $\Gamma$  and let  $s: V \rightarrow D$  be an assignment with cost  $\phi^\Delta(s(v_1), \dots, s(v_n)) \leq u$ . Then, by the definition of fractional homomorphism,

$$\sum_{h \in \text{Supp}(\chi)} \chi(h) \sum_{j \in J} \gamma_j^\Gamma(h(s(v^j))) = \sum_{j \in J} \sum_{h \in \text{Supp}(\chi)} \chi(h) \gamma_j^\Gamma(h(s(v^j))) \leq \sum_{j \in J} \gamma_j^\Delta(s(v^j)) \leq u.$$

Therefore, there exists at least one map  $h: D \rightarrow C$  from  $\text{Supp}(\chi)$ , such that  $\phi^\Gamma(h \circ s(v_1), \dots, h \circ s(v_n)) \leq \phi^\Delta(s(v_1), \dots, s(v_n)) \leq u$ , that is,  $h \circ s: V \rightarrow C$  is an assignment with cost at most  $u$ .  $\square$

**Lemma** (Lemma 6 restated). Let  $(\Delta, \Gamma)$  be a promise valued template and let  $m \in \mathbb{N}$ . If  $\omega = (\omega_I, \omega_O)$  is an  $m$ -ary block symmetric promise fractional polymorphism of  $(\Delta, \Gamma)$ , then also  $\omega' = (\omega'_I, \omega_O)$ , where  $\omega'_I(e_i^{(m)}) = \frac{1}{m}$  for  $1 \leq i \leq m$ , is an  $m$ -ary block-symmetric promise fractional polymorphism of  $(\Delta, \Gamma)$ .

*Proof.* Let  $C$  and  $D$  be the domains of  $\Gamma$  and  $\Delta$ , respectively, and let  $\tau$  be the common signature of the two valued structures. By the definition of block-symmetric promise fractional polymorphism, there exists a partition  $B_1 \cup \dots \cup B_k$  of  $[m]$  such that for every permutation  $\pi \in S_m$  that preserves the blocks  $B_1, \dots, B_k$ , every  $f \in \tau$  and every  $a^1, \dots, a^m \in D^{ar(f)}$  it holds that

$$\sum_{g \in \text{Supp}(\omega_O)} \omega_O(g) f^\Gamma(g(a^1, \dots, a^m)) \leq \sum_{i=1}^m \omega_I(e_i^{(m)}) f^\Delta(a^{\pi(i)}).$$

By summing the last inequality over all  $\pi \in S_m$  that preserve all the blocks  $B_1, \dots, B_k$ , i.e., over all  $\pi = \pi^1 \dots \pi^k$  such that  $\pi^j$  is a permutation of the elements in  $B_j$  for  $1 \leq j \leq k$ , we obtain

$$\begin{aligned} \prod_{j=1}^k |B_j|! \sum_{g \in \text{Supp}(\omega_O)} f^\Gamma(g(a^1, \dots, a^m)) &\leq \sum_{j=1}^k \prod_{h \neq j} |B_h|! \sum_{i \in B_j} \omega_I(e_i^{(m)}) \left( (|B_j| - 1)! \sum_{\ell \in B_j} f^\Delta(a^\ell) \right) \\ &= \prod_{j=1}^k |B_j|! \sum_{j=1}^k \frac{1}{|B_j|} \sum_{i \in B_j} \omega_I(e_i^{(m)}) \sum_{\ell \in B_j} f^\Delta(a^\ell). \end{aligned}$$

Since, by the definition of block-symmetric promise fractional polymorphism,  $\sum_{i \in B_j} \omega_I(e_i^{(m)}) = \frac{|B_j|}{m}$  for every  $j \in \{1, \dots, k\}$ , we obtain

$$\sum_{g \in \text{Supp}(\omega_O)} f^\Gamma(g(a^1, \dots, a^m)) \leq \frac{1}{m} \sum_{i=1}^m f^\Delta(a^i),$$

that is,  $\omega'$  is a promise fractional polymorphism of  $(\Delta, \Gamma)$ .  $\square$

## B The BLP Relaxation

A fractional polymorphism of a valued structure  $\Gamma$  is *fully symmetric* if every operation in its support is a fully symmetric map. Given a promise valued template  $(\Delta, \Gamma)$ , a promise fractional polymorphism  $\omega = (\omega_I, \omega_O)$  of  $(\Delta, \Gamma)$  is *fully symmetric* if every map in  $\text{Supp}(\omega_O)$  is fully symmetric.

The following lemma is a corollary of Lemma 6.

**Lemma 19.** *Let  $(\Delta, \Gamma)$  be a promise valued template and let  $m \in \mathbb{N}$ . If  $\omega = (\omega_I, \omega_O)$  is an  $m$ -ary fully symmetric promise fractional polymorphism of  $(\Delta, \Gamma)$ , then also  $\omega' = (\omega'_I, \omega_O)$ , where  $\omega'_I(e_i^{(m)}) = \frac{1}{m}$  for  $1 \leq i \leq m$ , is an  $m$ -ary fully symmetric promise fractional polymorphism of  $(\Delta, \Gamma)$ .*

In view of Lemma 19, we will assume without loss of generality that any  $m$ -ary fully symmetric promise fractional polymorphism  $\omega = (\omega_I, \omega_O)$  is such that  $\omega_I$  assign  $\frac{1}{m}$  to each  $m$ -ary projection on the domain of  $\Delta$  and we will identify  $\omega$  with  $\omega_O$ .

Recall the BLP relaxation from Section 2.5. Let  $(\Delta, \Gamma)$  be a promise valued template such that the domain of  $\Delta$  is a *finite* set. We may solve  $\text{PVCSP}(\Delta, \Gamma)$  by using the following algorithm that computes a BLP relaxation of  $\Delta$ .

---

### ALGORITHM 2: BLP Relaxation Algorithm for $\text{PVCSP}(\Delta, \Gamma)$

---

**Input:**

$I := (V, \phi, u)$ , a valid instance of  $\text{PVCSP}(\Delta, \Gamma)$

**Output:**

YES if there exists an assignment  $s: V \rightarrow \text{dom}(\Delta)$  such that  $\phi^\Delta(s(x_1), \dots, s(x_{|V|})) \leq u$   
 NO if there is no assignment  $s: V \rightarrow \text{dom}(\Gamma)$  such that  $\phi^\Gamma(s(x_1), \dots, s(x_{|V|})) \leq u$

BLP( $I, \Delta$ );

**if** BLP( $I, \Delta$ )  $\not\leq u$  **then**

    output NO;

**else**

    output YES;

**end**

---

Algorithm 2 runs in polynomial time in  $I$ , and that if it outputs NO, then indeed the answer to  $\text{PVCSP}(\Delta, \Gamma)$  is NO, without further assumptions.

We now present a sufficient condition under which Algorithm 2 correctly solves  $\text{PVCSP}(\Delta, \Gamma)$ .

**Theorem 20.** *Let  $(\Delta, \Gamma)$  be a promise valued template such that  $\Delta$  has a finite domain. Assume that for all  $m \in \mathbb{N}$  there exists a fully symmetric promise fractional polymorphisms of  $(\Delta, \Gamma)$  with arity  $m$ . Then Algorithm 2 correctly solves  $\text{PVCSP}(\Delta, \Gamma)$  (in polynomial time).*

To prove Theorem 20 we need to use a preliminary lemma and the notion of a *multiset-structures*.

Let  $\Delta$  be a valued  $\tau$ -structure with domain  $D$  and let  $m \in \mathbb{N}$ . The *multiset-structure*  $\mathcal{P}^m(\Delta)$  is the valued structure with domain  $\left(\binom{D}{m}\right)$  i.e., the set whose elements  $\alpha$  are multisets of elements from  $D$  of size  $m$ . For every  $k$ -ary function symbol  $f \in \tau$ , and  $\alpha_1, \dots, \alpha_k \in \left(\binom{D}{m}\right)$

the function  $f^{\mathcal{P}^m(\Delta)}$  is defined as follows

$$f^{\mathcal{P}^m(\Delta)}(\alpha_1, \dots, \alpha_k) := \frac{1}{m} \min_{\substack{t^1, \dots, t^k \in D^m: \\ \{t^\ell\} = \alpha_\ell}} \sum_{i=1}^m f^\Delta(t_i^1, \dots, t_i^k),$$

where  $\{t^\ell\}$  is the multiset of the coordinates of  $t^\ell$ , i.e., the multiset  $\{t_i^\ell \mid 1 \leq i \leq m\}$ .

**Lemma 21.** *Let  $(\Delta, \Gamma)$  be a promise valued template such that  $\Delta$  has a finite domain. Let  $m \in \mathbb{N}$  and let us assume that  $(\Delta, \Gamma)$  has a fully symmetric promise fractional polymorphism of arity  $m$ . Then  $\mathcal{P}^m(\Delta)$  is fractionally homomorphic to  $\Gamma$ .*

Lemma 21 is a generalisation of [51, Lemma 2.2] to promise valued templates  $(\Delta, \Gamma)$  where  $\Delta$  is a finite-domain valued structure. Furthermore, it already appeared in a similar form in [12, Lemma 6.9].<sup>7</sup>

*Proof of Lemma 21.* Let  $C$  be the (possibly infinite) domain of  $\Gamma$ , let  $D$  be the finite domain of  $\Delta$ , and let  $\tau$  be the common signature of  $\Gamma$  and  $\Delta$ . Let  $\omega$  be the  $m$ -ary fully symmetric promise fractional polymorphism of  $(\Delta, \Gamma)$ . For every  $g \in \text{Supp}(\omega) \subseteq C^{D^m}$  we define  $\tilde{g}: \binom{D}{m} \rightarrow C$  by setting, for every  $\alpha = \{\xi^1, \dots, \xi^m\} \in \binom{D}{m}$ ,

$$\tilde{g}(\alpha) = g(\xi^1, \xi^2, \dots, \xi^m).$$

Observe that  $\tilde{g}$  is well defined as  $g$  is fully symmetric. We define the discrete probability measure  $\chi$  on  $C^{\binom{D}{m}}$  as follows

$$\chi(Y) = \sum_{\substack{g \in \text{Supp}(\omega): \\ \tilde{g} \in Y}} \omega(g), \quad \text{for every } Y \subseteq \binom{D}{m}.$$

Observe that  $\chi$  satisfies the countable additivity property, since  $\omega$  does. Furthermore,  $\text{Supp}(\chi) = \{h \in C^{\binom{D}{m}} \mid h = \tilde{g} \text{ for some } g \in \text{Supp}(\omega)\} = \{\tilde{g} \in C^{\binom{D}{m}} \mid g \in \text{Supp}(\omega)\}$  is countable and it holds that

$$\sum_{h \in \text{Supp}(\chi)} \chi(h) = \sum_{g \in \text{Supp}(\omega)} \omega(g) = 1.$$

We claim that  $\chi$  is a fractional homomorphism from  $\mathcal{P}^m(\Delta)$  to  $\Gamma$ . Indeed, for every  $f \in \tau$  and every tuple  $(\alpha_1, \dots, \alpha_k) \in \binom{D}{m}^k$  with  $\alpha_i := \{\xi_i^1, \dots, \xi_i^m\}$  and  $k := \text{ar}(f)$ , it holds that

$$\begin{aligned} & \sum_{h \in C^{\binom{D}{m}}} \chi(h) f^\Gamma(h(\alpha_1), \dots, h(\alpha_k)) \\ &= \sum_{g \in \text{Supp}(\omega)} \omega(g) f^\Gamma(g(\xi_1^1, \dots, \xi_1^m), \dots, g(\xi_k^1, \dots, \xi_k^m)) \\ &= \sum_{g \in \text{Supp}(\omega)} \omega(g) f^\Gamma(g(\xi_1^{\pi_1(1)}, \dots, \xi_1^{\pi_1(m)}), \dots, g(\xi_k^{\pi_k(1)}, \dots, \xi_k^{\pi_k(m)})) \end{aligned} \quad (9)$$

$$\leq \frac{1}{m} \sum_{i=1}^m f^\Delta(\xi_1^{\pi_1(i)}, \dots, \xi_k^{\pi_k(i)}) \quad (10)$$

---

<sup>7</sup>In [12] the assumption of having an  $m$ -ary fully-symmetric promise fractional polymorphism of  $(\Delta, \Gamma)$  is replaced by the hypothesis of having an  $m$ -ary fully-symmetric fractional polymorphism of  $\Gamma$ . Another difference is that in [12] the notion of fractional polymorphism employed allows only finitely supported probability measures.

for every  $\pi_1, \dots, \pi_k \in S_m$ . Equality (9) holds because the maps  $g \in \text{Supp}(\omega)$  are fully symmetric and Inequality (10) holds because  $\omega$  is a promise fractional polymorphism of  $(\Delta, \Gamma)$ . Then, in particular, we obtain

$$\begin{aligned} \sum_{h \in C \left( \binom{D}{m} \right)} \chi(h) f^\Gamma(h(\alpha_1, \dots, \alpha_k)) &\leq \frac{1}{m} \min_{\substack{t^1, \dots, t^k \in D^m: \\ \{t^\ell\} = \alpha_\ell}} \sum_{i=1}^m f^\Delta(t_i^1, \dots, t_i^k) \\ &= f^{\mathcal{P}^m(\Delta)}(\alpha_1, \dots, \alpha_k). \end{aligned} \quad \square$$

With Lemma 21, the proof of Theorem 20 follows the same argument as in the finite-domain non-promise case [51, Theorem 3.2]. We include the proof here for completeness.

*Proof of Theorem 20.* Let  $C$  be the (possibly infinite) domain of  $\Gamma$  and let  $D$  be the finite domain of  $\Delta$ . Let  $\tau$  be the common signature of  $\Delta$  and  $\Gamma$ . Let  $I$  be an instance of  $\text{PVCSP}(\Delta, \Gamma)$  with variables  $V = \{x_1, \dots, x_n\}$ , objective function  $\phi(x_1, \dots, x_n) = \sum_{j \in J} \gamma_j(x^j)$  where  $J$  is a finite set of indices,  $\gamma_j \in \Gamma$ , and  $x^j \in V^{\text{ar}(j)}$ , and threshold  $u$ . Note that if  $\text{BLP}(I, \Delta) \not\leq u$  (this also includes the case  $\text{BLP}(I, \Delta) = +\infty$ , i.e., the case that  $I$  is not feasible) then  $\min_D \phi^\Delta \not\leq u$ . We may therefore safely output NO. Otherwise,  $\text{BLP}(I, \Delta) \leq u$  implies that  $\inf_C \phi^\Gamma \leq \text{BLP}(I, \Delta)$ . The proof of this last statement is contained in the first part of the proof of [51, Theorem 3.2]; we include it here for completeness. Let  $(\lambda^*, \mu^*)$  be an optimal solution to  $\text{BLP}(I, \Delta)$  and let  $M$  be a positive integer such that  $M \cdot \lambda^*$ , and  $M \cdot \mu^*$  are both integral. Let  $\nu: V \rightarrow \left( \binom{D}{M} \right)$  be defined by mapping the variable  $x_i$  to the multiset in which the elements are distributed accordingly to  $\mu_{x_i}^*$ , i.e., for every  $a \in D$  the number of occurrences of  $a$  in  $\nu(x_i)$  is equal to  $M \mu_{x_i}^*(a)$ . Let  $f_j$  be a  $k$ -ary function symbol in  $\tau$  that occurs in a term  $f_j(x^j)$  of the objective function  $\phi$ . Now we write

$$M \cdot \sum_{t \in D^k} \lambda_j^*(t) f_j^\Delta(t) = f_j^\Delta(\alpha^1) + \dots + f_j^\Delta(\alpha^M),$$

where the  $\alpha^i \in D^k$  are such that  $\lambda_j^*(t)$ -fractions are equal to  $t$ . Let us define  $\alpha'_\ell := (\alpha_\ell^1, \dots, \alpha_\ell^M)$  for  $1 \leq \ell \leq k$ . We get

$$\begin{aligned} \sum_{t \in D^k} \lambda_j^*(t) f_j^\Delta(t) &= \frac{1}{M} \sum_{i=1}^M f_j^\Delta(\alpha^i) = \frac{1}{M} \sum_{i=1}^M f_j^\Delta(\alpha_1^i, \dots, \alpha_k^i) \\ &\geq \frac{1}{M} \min_{\substack{t^1, \dots, t^k \in D^M: \\ \{t^\ell\} = \{\alpha'_\ell\}}} \sum_{i=1}^M f_j^\Delta(t_i^1, \dots, t_i^k) = f_j^{\mathcal{P}^M(\Delta)}(\alpha'_1, \dots, \alpha'_k) \\ &= f_j^{\mathcal{P}^M(\Delta)}(\nu(x)), \end{aligned}$$

where the last equality follows as the number of  $a$ 's in  $\alpha'_\ell$  is

$$M \cdot \sum_{t \in D^k: t^i = a} \lambda_j^*(t) = M \cdot \mu_{x_i}^*(a).$$



Then

$$\begin{aligned} \text{BLP}(I, \Delta) &= \sum_{j \in J} \sum_{t \in D^{\text{ar}(f_j)}} \lambda_j^*(t) f_j^\Delta(t) = \sum_{j \in J} \left( \sum_{t \in D^{\text{ar}(f_j)}} \lambda_j^*(t) f_j^\Delta(t) \right) \\ &\geq \sum_{j \in J} \left( f_j^{\mathcal{P}^M(\Delta)}(\nu(x)) \right) \geq \min_{\left( \binom{D}{m} \right)} \phi^{\mathcal{P}^M(\Delta)}. \end{aligned}$$

Since we assumed  $\text{BLP}(I, \Delta) \leq u$ , we obtain  $\min_{\left( \binom{D}{m} \right)} \phi^{\mathcal{P}^M(\Delta)} \leq u$ . Moreover, since  $\Gamma$  has fully symmetric fractional polymorphisms of all arities, Lemma 21 implies the existence of a fractional homomorphism  $\omega$  from  $\mathcal{P}^M(\Delta)$  to  $\Gamma$ . From Proposition 2 it follows that

$$\inf_C \phi^\Gamma \leq \min_{\left( \binom{D}{m} \right)} \phi^{\mathcal{P}^M(\Delta)} \leq \text{BLP}(I, \Delta) \leq u.$$

□