

Causal Mediation Analysis for Sparse and Irregular Longitudinal Data

Shuxi Zeng Stacy Rosenbaum Elizabeth Archie Susan Alberts Fan Li ¹

ABSTRACT

Causal mediation analysis aims to investigate how the treatment effect of an exposure on outcomes is mediated through intermediate variables. Although many applications involve longitudinal data, the existing methods are not directly applicable to the settings where the mediator and outcome are measured on sparse and irregular time grids. We extend the existing causal mediation framework from a functional data analysis perspective, viewing the sparse and irregular longitudinal data as realizations of underlying smooth stochastic processes. We define causal estimands of direct and indirect effects accordingly and provide corresponding identification assumptions. For estimation and inference, we employ a functional principal component analysis approach for dimension reduction and use the first few functional principal components instead of the whole trajectories in the structural equation models. We adopt the Bayesian paradigm to accurately quantify the uncertainties. The operating characteristics of the proposed methods are examined via simulations. We apply the proposed methods to a longitudinal data set from a wild baboon population in Kenya to estimate the causal effects between early adversity, the strength of social bonds, and adult glucocorticoid hormone concentrations. We find that early adversity has a significant direct effect (a 9-14% increase) on females glucocorticoid concentrations across adulthood, but find little evidence that these effects were mediated by weak social bonds.

KEY WORDS: Causal inference, functional principal component analysis, mediation, longitudinal data, sparse and irregular data

¹Shuxi Zeng is Ph.D. student, Department of Statistical Science, Duke University, Durham, NC, 27705 (email: shuxi.zeng@duke.edu); Stacy Rosenbaum is Assistant Professor, Department of Anthropology, University of Michigan, Ann Arbor, MI 48109 (email: rosenbas@umich.edu); Elizabeth Archie is Associate Professor, Department of Biological Sciences, University of Notre Dame, Notre Dame, IN, 46556 (email: earchie@nd.edu); Susan Alberts is Robert F. Durden Distinguished Professor, Department of Biology, Duke University, Durham, NC, 27705 (email: alberts@duke.edu); Fan Li is Associate Professor, Department of Statistical Science, Duke University, Durham, NC, 27705 (email: fl35@duke.edu).

1 Introduction

Mediation analysis seeks to understand the role of an intermediate variable (i.e. mediator) M that lies on the causal path between an exposure or treatment Z and an outcome Y . The most widely used mediation analysis method, proposed by Baron and Kenny (1986), fits two linear structural equation models (SEMs) between the three variables and interprets the model coefficients as causal effects. There is a vast literature on the Baron-Kenny framework across a variety of disciplines, including psychology, sociology, and epidemiology (see MacKinnon, 2012). A major advancement in recent years is the incorporation of the potential-outcome-based causal inference approach (Neyman, 1923; Rubin, 1974). This led to formal definition of relevant causal estimands, clarification of identification assumptions and new estimation strategies beyond linear SEMs (Robins and Greenland, 1992; Pearl, 2001; Sobel, 2008; Daniels et al., 2012; Tchetgen Tchetgen and Shpitser, 2012; VanderWeele, 2016). In particular, Imai et al. (2010) proved that the Baron-Kenny estimator can be interpreted as a special case of a causal mediation estimator given additional model assumptions. These methodological advancements also opened up new application areas including imaging, neuroscience and environmental health (Lindquist and Sobel, 2011; Lindquist, 2012; Zigler et al., 2012; Kim et al., 2019). Comprehensive reviews on causal mediation analysis are given in VanderWeele (2015); Nguyen et al. (2020).

In traditional settings of mediation analysis, exposure Z , mediation M and outcome Y are all univariate variables at a single time point. Recent work has extended to time-varying cases, where at least one of the triplet (Z, M, Y) is longitudinal. This line of research has primarily focused on the case with time varying mediators or outcomes that are observed on sparse and regular time grids (van der Laan and Petersen, 2008; Roth and MacKinnon, 2012; Lin et al., 2017). For example, VanderWeele and Tchetgen Tchetgen (2017) developed a method for identifying and estimating causal mediation effects with time-varying exposures and mediators based on marginal structural model (Robins et al., 2000). Another stream of research, motivated from applications in neuroimaging, focuses on the cases where mediators or outcomes are densely recorded continuous functions, e.g. the blood-oxygen-level-dependent (BOLD) signal collected in a functional magnetic resonance imaging (fMRI) session. In

particular, Lindquist (2012) introduced the concept of *functional mediation* in the presence of a functional mediator and extended causal SEMs to functional data analysis (Ramsay and Silverman, 2005). Zhao et al. (2018) further extended this approach to functional exposure, mediator and outcome.

Sparse and irregularly-spaced longitudinal data are increasingly available for causal studies. For example, in electronic health records (EHR) data, the number of observations usually varies between patients and the time grids are uneven. The same situation applies in animal behavior studies due to the inherent difficulties in observing wild animals. Such data structure pose challenges to the existing causal mediation methods. First, one cannot simply treat the trajectories of mediators and outcomes as functions as in Lindquist (2012) because the sparse observations render the trajectories volatile and non-smooth. Second, with irregular time grids the dependence between consecutive observations changes over time, making the methods based on sparse and regular longitudinal data such as VanderWeele and Tchetgen Tchetgen (2017) not applicable. A further complication arises when the mediator and outcome are measured with different frequencies even within the same individual.

In this paper, we propose a framework for causal mediation analysis with sparse and irregular longitudinal data that address the aforementioned challenges. Similar as Lindquist (2012) and Zhao et al. (2018), we adopt a functional data analysis perspective (Ramsay and Silverman, 2005), viewing the sparse and irregular longitudinal data as realizations of underlying smooth stochastic processes. We define causal estimands of direct and indirect effects accordingly and provide assumptions for nonparametric identification (Section 3). For estimation and inference, we proceed under the classical two-SEM mediation framework (Imai et al., 2010) but diverge from the existing methods in modeling (Section 4). Specifically, we employ the functional principal component analysis (FPCA) approach (Yao et al., 2005; Jiang et al., 2010, 2011) to project the mediator and outcome trajectories to a low-dimensional representation. We then use the first few functional principal components instead of the whole trajectories as predictors in the structural equation models. To accurately quantifying the uncertainties, we employ a Bayesian FPCA model (Kowal and Bourgeois, 2020) to simultaneously estimate the functional principal components and the structural equation models.

Our motivating application is the evaluation of the causal effects and mechanism between early

adversity, social bonds, and stress in wild baboons (Section 2). Here the exposure is early adversity (e.g. drought, maternal death), the mediators are the adult social bonds, and the outcomes are the adult glucocorticoid hormone concentrations, which is a measure of the stress level. The exposure is a binary variable measured at one time point, whereas both the mediators and outcomes are sparse and irregular longitudinal variables. We apply the proposed method to a prospective and longitudinal observational data set from the Amboseli Baboon Research Project (Alberts and Altmann, 2012) (Section 6). We find that experiencing one or more sources of early adversity leads to significant direct effects (a 9-14% increase) on females glucocorticoid concentrations across adulthood, but find little evidence that these effects were mediated by weak social bonds.

Though motivated from a specific application, the proposed method is readily applicable to other causal mediation studies with similar data structure, including the EHR and ecology studies. Note that our method is also applicable to regular longitudinal observations.

2 Motivating Application: Early Adversity, Social Bond and Stress

2.1 Biological Background

Conditions in early life can have profound consequences for individual development, behavior, and physiology across the life course (Lindström, 1999; Gluckman et al., 2008; Bateson et al., 2004). These early life effects are important, in part, because they have major implications for human health. One leading explanation for how early life environments affect adult health is provided by the biological embedding hypothesis, which posits that early life stress causes developmental changes that create a pro-inflammatory phenotype and elevated risk for several diseases of aging (Miller et al., 2011). The biological embedding hypothesis proposes at least two, non-exclusive causal pathways that connect early adversity to poor health in adulthood. In the first pathway, early adversity leads to altered hormonal profiles that contribute to inflammation and disease. Under this scenario, stress in early life leads to dysregulation of hormonal signals in the body's main stress response system, leading to the release of glucocorticoid hormones (GCs), which engage the body's fight-or-flight response. In turn, such acti-

vations is associated with inflammation and elevated disease risk (McEwen, 1998; Miller et al., 2002; McEwen, 2008). In the second causal pathway, early adversity hampers an individual's ability to form strong interpersonal relationships. Under this scenario, the social isolation contributes to both altered GCs profiles and inflammation.

Hence, the biological embedding hypothesis posits that early life adversity affects both GCs profiles and social conditions in adulthood, and that poor social relationships partly mediate the relationship between early adversity and GCs. Importantly, the second causal pathway mediated through adult social conditions suggests an opportunity to mitigate the negative health effect of early adversity. Specifically, strong and supportive social relationships may dampen the stress response or reduce individual exposure to stressful events, which in turn reduces GCs and inflammation. In support, strong and supportive social relationships have repeatedly been linked to reduced morbidity and mortality in humans and other social animals (Holt-Lunstad et al., 2010; Silk, 2007). In addition to the biological embedding hypothesis, this idea of social mitigation is central to several hypotheses that propose causal connections between adult social conditions and adult health, even independent of early life adversity; these hypotheses include the stress buffering and stress prevention hypotheses (Cohen and Wills, 1985; Landerman et al., 1989; Thorsteinsson and James, 1999) and the social causation hypothesis (Marmot et al., 1991; Anderson and Marmot, 2011).

Despite the aforementioned research, the causal relationships between early adversity, adult social conditions, and HPA dysregulation remain to be subject to considerable debate. While social relationships might exert direct effects on stress and health, it is also possible that poor health and high stress limit an individual's ability to form strong and supportive relationships. As such the causal arrow flows backwards, from stress to social relationships (Case and Paxson, 2011). In another causal scenario, early adversity exerts independent effects on social conditions and the HPA axis, and correlations between social relationships and GCs are spurious, arising solely as a result of their independent links to early adversity.

2.2 Data

In this paper, we test whether the links between early adversity, the strength of adult social bonds, and adult HPA axis activity are consistent with predictions of the biological embedding hypothesis in a wild primate population. Specifically, we use data from a well-studied population of savannah baboons in the Amboseli ecosystem in Kenya (Alberts and Altmann, 2012). Founded in 1971, the Amboseli Baboon Research Project (Alberts and Altmann, 2012) has prospective longitudinal data on early life experiences, and fine-grained longitudinal data on adult social bonds and GC hormones levels, which is a measure of HPA axis activation and the stress response.

Our study sample includes 192 female baboons. Each baboon was at least four years old, and we had complete information on her experience of six well-characterized sources of early adversity (i.e., exposure) (Tung et al., 2016; Zippel et al., 2019), as well as information on her adult social bonds (i.e. mediators) and fecal GC hormones concentrations (i.e. outcomes). Social bonds and GC hormones levels are measured repeatedly throughout the subjects lives on the same grid. For wild baboons, the observations are not on regular basis as the social bonds and GCs levels can be missing or measured multiple times within a year. We have 51.4 observations for each baboon on average for both social bonds and GCs levels, but the number of observations of a single baboon ranges from 3 to 113. Figure 1 shows the mediator and outcome trajectories of two randomly selected baboons in the sample. We can see that the frequency of the observations and time grids of the mediator or outcome trajectories vary significantly between baboons. More detailed information about the data, including covariates, is discussed in Section 6.1, and in Rosenbaum et al. (2020).

3 Causal Mediation Framework

3.1 The Setup and Estimands

Suppose we have a sample of N units; each unit i ($i = 1, 2, \dots, N$) is assigned to a treatment ($Z_i = 1$) or a control ($Z_i = 0$) group. For each unit i , we make observations at T_i different time points




Figure 1: Observed trajectories of social bond and GC hormones level of two randomly selected female baboons in the study sample

$\{t_{ij} \in [0, T], j = 1, 2, \dots, T_i\}$, and T_i can vary between units. At each time point t_{ij} , we measure an outcome Y_{ij} and a mediator M_{ij} prior to the outcome, and a vector of p time-varying covariates $\mathbf{X}_{ij} = (X_{ij,1}, \dots, X_{ij,p})'$. For each unit, the time points are sparse along the time span and irregularly spaced. Also, the observed time grids for the outcome and the mediator are not necessarily the same between units.

A key to our framework is to view the observed mediator and outcome values drawn from a smooth underlying process $M_i(t)$ and $Y_i(t)$, $t \in [0, T]$, with Normal measurement errors, respectively:

$$M_{ij} = M_i(t_{ij}) + \varepsilon_{ij}, \quad \varepsilon_{ij} \sim \mathcal{N}(0, \sigma_m^2), \quad (1)$$

$$Y_{ij} = Y_i(t_{ij}) + \nu_{ij}, \quad \nu_{ij} \sim \mathcal{N}(0, \sigma_y^2). \quad (2)$$

Hence, instead of directly exploring the relationship between the treatment Z_i , mediators M_{ij} and outcomes Y_{ij} , we investigate the relationship between Z_i and the stochastic processes $M_i(t_{ij})$ and $Y_i(t_{ij})$. In particular, we wish to answer two questions: (a) how much is the causal impact of the treatment on the outcome process, and (b) how much of that impact is mediated through the mediator process.

To be consistent with the standard notation of potential outcomes in causal inference, from now on we move the time index of the mediator and outcome process to the superscript: $M_i(t) = M_i^t, Y_i(t) =$

Y_i^t . Also, we use the following bold font notation to represent a process until time t : $\mathbf{M}_i^t \equiv \{M_i^s, s \leq t\} \in \mathcal{R}^{[0,t]}$, and $\mathbf{Y}_i^t \equiv \{Y_i^s, s \leq t\} \in \mathcal{R}^{[0,t]}$. Similarly, we denote the covariates up until the time between the j th and $j + 1$ th time point for unit i as $\mathbf{X}_i^t = \{X_{i1}, X_{i2}, \dots, X_{ij}\}$ for $t_{ij} \leq t < t_{ij+1}$.

We extend the definition of potential outcomes to define the causal estimands. Specifically, let $\mathbf{M}_i^t(z) \in \mathcal{R}^{[0,t]}$ for $z = 0, 1, t \in [0, T]$, denote the potential values of the underlying mediator process for unit i until time t under the treatment status z ; let $\mathbf{Y}_i^t(z, \mathbf{m}) \in \mathcal{R}^{[0,t]}$ be the potential outcome for unit i until time t under the treatment status z and the mediator process taking value of $\mathbf{M}_i^t = \mathbf{m}$ with $\mathbf{m} \in \mathcal{R}^{[0,t]}$. The above notation implicitly makes a standard assumption that the potential outcomes are determined solely by the treatment status z and the mediator values \mathbf{m} before time t , but not after t . For each unit, we can only observe one realization from the potential mediator or outcome process:

$$\mathbf{M}_i^t = \mathbf{M}_i^t(Z_i) = Z_i \mathbf{M}_i^t(1) + (1 - Z_i) \mathbf{M}_i^t(0), \quad (3)$$

$$\mathbf{Y}_i^t = \mathbf{Y}_i^t(z, \mathbf{M}_i^t(Z_i)) = Z_i \mathbf{Y}_i^t(1, \mathbf{M}_i^t(1)) + (1 - Z_i) \mathbf{Y}_i^t(0, \mathbf{M}_i^t(0)). \quad (4)$$

We define the total effect (TE) of the treatment Z_i on the outcome process at time t as:

$$\tau_{\text{TE}}^t = E\{Y_i^t(1, \mathbf{M}_i^t(1)) - Y_i^t(0, \mathbf{M}_i^t(0))\}. \quad (5)$$

In the presence of a mediator, the total effect can be decomposed into direct and indirect effects. Below we extend the framework of (Imai et al., 2010) to formally define these effects. First, we define the average causal mediation (or indirect) effect (ACME) under treatment z at time t by fixing the treatment status while altering the mediator process:

$$\tau_{\text{ACME}}^t(z) \equiv E\{Y_i^t(z, \mathbf{M}_i^t(1)) - Y_i^t(z, \mathbf{M}_i^t(0))\}, \quad z = 0, 1. \quad (6)$$

The ACME quantifies the difference between the potential outcomes, given a fixed treatment status z , corresponding to the potential mediator process under treatment $\mathbf{M}_i^t(1)$ and that under control $\mathbf{M}_i^t(0)$. In the previous literature, variants of the ACME are also called the *natural indirect effect* (Pearl, 2001), or the *pure indirect effect* for $\tau_{\text{ACME}}^t(0)$ and *total indirect effect* for $\tau_{\text{ACME}}^t(1)$ (Robins and Greenland, 1992)

Second, we define the average natural direct effect (ANDE) (Pearl, 2001; Imai et al., 2010) of treatment on the outcome at time t by fixing the mediator process while altering the treatment status:

$$\tau_{\text{ANDE}}^t(z) \equiv E\{Y_i^t(1, \mathbf{M}_i^t(z)) - Y_i^t(0, \mathbf{M}_i^t(z))\}, \quad (7)$$

The ANDE quantifies the portion in the total effects that does not pass through the mediators.

It is easy to verify that the total effect is the sum of ACME and ANDE:

$$\tau_{\text{TE}}^t = \tau_{\text{ACME}}^t(z) + \tau_{\text{ANDE}}^t(1 - z), \quad z = 0, 1. \quad (8)$$

This implies we only need to identify two of the three quantities τ_{TE}^t , $\tau_{\text{ACME}}^t(z)$, $\tau_{\text{ANDE}}^t(z)$. In this paper, we will focus on the estimation of τ_{TE}^t and $\tau_{\text{ACME}}^t(z)$. Also we make a common assumption that the ACME and ANDE are the same in the treatment and control groups: $\tau_{\text{ACME}}^t(0) = \tau_{\text{ACME}}^t(1)$, $\tau_{\text{ANDE}}^t(0) = \tau_{\text{ANDE}}^t(1)$. Because we only observe a portion of all the potential outcomes, we cannot directly identify these estimands from the observed data, which would require additional assumptions.

3.2 Identification assumptions

In this subsection, we list the causal assumptions necessary for identifying the ACME and ANDEs with sparse and irregular longitudinal data.

The first assumption extends the standard ignorability (or unconfoundedness) assumption and rules out the unmeasured treatment-outcome confounding.

Assumption 1 (Ignorability) *Conditional on the observed covariates, the treatment is unconfounded with respect to the potential mediator process and the potential outcomes process:*

$$\{\mathbf{Y}_i^t(1, \mathbf{m}), \mathbf{Y}_i^t(0, \mathbf{m}), \mathbf{M}_i^t(1), \mathbf{M}_i^t(0)\} \perp\!\!\!\perp Z_i \mid \mathbf{X}_i^t$$

for any t and $\mathbf{m} \in \mathcal{R}^{[0,t]}$.

The second assumption extends the sequential ignorability assumption in (Imai et al., 2010) to the functional data setting.

Assumption 2 (Sequential Ignorability) *There exists $\varepsilon > 0$, such that for any $0 < \Delta < \varepsilon$, the increment of mediator process is independent of the increment of potential outcomes process from time t to $t + \Delta$, conditional on the observed treatment status, covariates and the mediator process up to time t :*

$$\{Y_i^{t+\Delta}(z, \mathbf{m}) - Y_i^t(z, \mathbf{m})\} \perp\!\!\!\perp (M_i^{t+\Delta} - M_i^t) \mid \{Z_i, \mathbf{X}_i^t, \mathbf{M}_i^t\}$$

for any $z, 0 < \Delta < \varepsilon, t, t + \Delta \in [0, T], \mathbf{m} \in \mathcal{R}^{[0, T]}$.

Assumption 2 implies that conditioning on the observed treatment status, covariates, and the mediator process up to a given time point, the change in the mediator values within a sufficiently small time interval is randomized with respect to the change in the potential outcomes. Namely, there are no unobserved mediator-outcomes confounders in a sufficiently small time interval. Though differs in the specific form, Assumption 2 shares the essence with the previous sequential ignorability assumptions for the regularly spaced observations in Bind et al. (2015) and VanderWeele and Tchetgen Tchetgen (2017). This is a crucial assumption in mediation analysis, but is strong and generally untestable in practice because it is usually impossible to manipulate the mediator values, even in randomized trials.

Assumption 1 and 2 are illustrated by the directed acyclic graphs (DAG) in Figure 2a, which condition on the covariates \mathbf{X}_i^t and a window between two sufficiently closed time points t and $t + \Delta$. The arrows between Z_i, M_i^t, Y_i^t represent a causal relationship (i.e., nonparametric structural equation model), with solid and dashed lines representing measured and unmeasured relationships, respectively. Figure 2b and 2c depicts two possible scenarios where Assumption 1 and 2 is violated, respectively, where U_i represents an unmeasured confounder.

Assumptions 1 and 2 allow nonparametric identification of the total effect and ACME from the observed data, as summarized in the following theorem.

Theorem 1 *Under Assumption 1,2, and some regularity conditions (specified in the Appendix), the total effect, ACME and ANDE can be identified nonparametrically from the observed data: for $z = 0, 1$, we*




Figure 2: Directed acyclic graphs (DAG) of Assumption 1, 2 and examples of possible violation. The arrows between variables represent a causal relationship, with solid and dashed lines representing measured and unmeasured relationships, respectively.

have

$$\begin{aligned} \tau_{TE} &= \int_{\mathcal{X}} \{E(Y_i^t | Z_i = 1, \mathbf{X}_i^t = \mathbf{x}^t) - E(Y_i^t | Z_i = 0, \mathbf{X}_i^t = \mathbf{x}^t)\} dF_{\mathbf{X}_i^t}(\mathbf{x}^t), \\ \tau_{ACME}^t(z) &= \int_{\mathcal{X}} \int_{R^{[0,t]}} E(Y_i^t | Z_i = z, \mathbf{X}_i^t = \mathbf{x}^t, \mathbf{M}_i^t = \mathbf{m}) dF_{\mathbf{X}_i^t}(\mathbf{x}^t) \times \\ &\quad d\{F_{\mathbf{M}_i^t | Z_i=1, \mathbf{X}_i^t=\mathbf{x}^t}(\mathbf{m}) - F_{\mathbf{M}_i^t | Z_i=0, \mathbf{X}_i^t=\mathbf{x}^t}(\mathbf{m})\}, \end{aligned}$$

where $F_W(\cdot)$ and $F_{W|V}(\cdot)$ denotes the cumulative distribution of a random variable or vector W and

the conditional distribution given another random variable or vector V , respectively.

The proof of Theorem 1 is provided in the Supplementary A. Theorem 1 implies that estimating the causal effects requires modeling two components: (a) the distribution of observed outcome process given the treatment, covariates, and the observed mediator process, $E(Y_i^t|Z_i, \mathbf{X}_i^t, \mathbf{M}_i^t)$, (b) the distribution of the observed mediator process given the treatment and the covariates, $E(\mathbf{M}_i^t|Z_i, \mathbf{X}_i^t)$. These two components correspond to the two linear structural equations in the classic mediation framework of Baron and Kenny (1986). In the setting of functional data, we can employ more flexible models instead of linear regression models, and express the total effect and ACME as functions of the model parameters. Theorem 1 can be readily extended to more general scenarios such as discrete mediators and time-to-event outcomes.

4 Modeling mediator and outcome via functional principal component analysis

In this section, we propose to employ the functional principal component analysis (FPCA) approach to infer the mediator and outcome processes from sparse and irregular observations (Yao et al., 2005; Jiang et al., 2010, 2011). In order to take into account the uncertainty due to estimating the functional principal components (Goldsmith et al., 2013), we adopt a Bayesian model to jointly estimate the principal components and the structural equation models. Specifically, we impose a Bayesian FPCA model similar to that in Kowal and Bourgeois (2020) to project the observed mediator and outcome processes into lower-dimensional representations and then take the first few dominant principal components as the predictors in the structural equation models.

We assume the potential processes for mediators $\mathbf{M}_i^t(z)$ and outcomes $\mathbf{Y}_i^t(z, \mathbf{m})$ have the following

Karhunen-Loeve decomposition,

$$M_i^t(z) = \mu_M(\mathbf{X}_i^t) + \sum_{r=1}^{\infty} \zeta_{i,z}^r \psi_r(t), \quad (9)$$

$$Y_i^t(z, \mathbf{m}) = \mu_Y(\mathbf{X}_i^t) + \int_0^t \gamma(s, t) \mathbf{m}(s) ds + \sum_{s=1}^{\infty} \theta_{i,z}^s \eta_s(t). \quad (10)$$

where $\mu_M(\cdot)$ and $\mu_Y(\cdot)$ are the mean functions of the mediator process \mathbf{M}_i^t and outcome process \mathbf{Y}_i^t , respectively; $\psi_r(t)$ and $\eta_s(t)$ are the Normal orthogonal eigenfunctions for \mathbf{M}_i^t and \mathbf{Y}_i^t , respectively, and $\zeta_{i,z}^r$ and $\theta_{i,z}^s$ are the corresponding principal scores of unit i . The above model assumes that the treatment affects the mediation and the outcome processes only through the principal scores. We represent the mediator and outcome process of each unit with its principal score $\zeta_{i,z}^r$ and $\theta_{i,z}^s$. Given the principal scores, we can transform back to the smooth process with linear combination. As such, if we are interested in the differences on the process, it is equivalent to investigate the difference on the principal scores. Also, as we usually require only 3 or 4 components to explain the most variation, we reduce the dimensions of trajectories effectively by projecting the difference to principal scores.

The underlying processes \mathbf{M}_i^t and \mathbf{Y}_i^t are not directly observed. Instead, we assume the observations M_{ij} 's and Y_{ij} 's are randomly sampled from the respective underlying processes with errors. For the observed mediator trajectories, we posit the following model that truncates to the first R principal components of the mediator process:

$$M_{ij} = X'_{ij} \beta_M + \sum_{r=1}^R \zeta_i^r \psi_r(t_{ij}) + \varepsilon_{ij}, \quad \varepsilon_{ij} \sim \mathcal{N}(0, \sigma_m^2), \quad (11)$$

where $\psi_r(t)$ ($r = 1, \dots, R$) are the orthonormal principal components, ζ_i^r ($r = 1, \dots, R$) are the corresponding principal scores, and ε_{ij} is the measurement error. With similar parametrization in Kowal and Bourgeois (2020), we express the principal components as a linear combination of the spline basis $\mathbf{b}(t) = (1, t, b_1(t), \dots, b_L(t))'$ in $L + 2$ dimensions and choose the coefficients $\mathbf{p}_r \in \mathcal{R}^{L+2}$ to meet the normal orthogonality constraints of the r th principal component:

$$\psi_r(t) = \mathbf{b}(t)' \mathbf{p}_r, \quad \text{subject to } \int_0^T \psi_r^2(t) dt = 1, \quad \int_0^T \psi_{r'}(t) \psi_{r''}(t) dt = 0. \quad (12)$$

We assume the principal scores ζ_i^r are randomly drawn from normal distributions with different means in the treated and control groups, χ_1^r and χ_0^r , and diminishing variance as r increases:

$$\zeta_i^r \sim \mathcal{N}(\chi_{Z_i}^r, \lambda_r^2), \quad \lambda_1^2 \geq \lambda_2^2 \geq \dots \geq \lambda_R^2 \geq 0. \quad (13)$$

We select the truncation term R based on the fraction of explained variance (FEV), $\sum_{r=1}^R \lambda_r^2 / \sum_{r=1}^{\infty} \lambda_r^2$ being greater than 90%.

For the observed outcome trajectories, we posit a similar model that truncates to the first S principal components of the outcome process:

$$Y_{ij} = X_{ij}^T \beta_Y + \int_0^{t_{ij}} \gamma(u, t) M_i^u du + \sum_{s=1}^S \eta_s(t) \theta_i^s + \nu_{ij}, \quad \nu_{ij} \sim N(0, \sigma_y^2). \quad (14)$$

We express the principal components η_s as a linear combination of the spline basis $\mathbf{b}(t)$, with the normal orthogonality constraints:

$$\eta_s(t) = \mathbf{b}(t)' \mathbf{q}_s, \text{ subject to } \int_0^T \eta_s(t)^2 dt = 1, \int_0^T \eta_{s'}(t) \eta_{s''}(t) dt = 0. \quad (15)$$

Similarly, we assume that the principal scores of outcome process for each unit come from two different normal distributions in the treated and control group with means ξ_1^s and ξ_0^s respectively, and a shrinking variance ρ_s^2 :

$$\theta_i^s \sim \mathcal{N}(\xi_{Z_i}^s, \rho_s^2), \quad \rho_1^2 \geq \rho_2^2 \geq \dots \geq \rho_S^2 \geq 0. \quad (16)$$

We select the truncation term S based on the FEV being greater than 90%, namely $\sum_{s=1}^S \rho_s^2 / \sum_{s=1}^{\infty} \rho_s^2 \geq 90\%$.

We assume the effect of the mediation process on the outcome is concurrent, namely the outcome process at time t does not depend on the past value of the mediation process. As such, $\gamma(u, t)$ can be shrunk to γ instead of the integral in Model (14),

$$Y_{ij} = X_{ij}^T \beta_Y + \gamma M_{ij} + \sum_{s=1}^S \eta_s(t) \theta_i^s + \nu_{ij}, \quad \nu_{ij} \sim N(0, \sigma_y^2). \quad (17)$$

The causal estimands, the total effect and ACME, can be expressed as functions of the parameters in the above mediator and outcome models:

$$\tau_{\text{TE}}^t = \sum_{s=1}^S (\zeta_1^s - \zeta_0^s) \eta_s(t) + \gamma \sum_{r=1}^R (\chi_1^r - \chi_0^r) \psi_r(t), \quad (18)$$

$$\tau_{\text{ACME}}^t = \gamma (\chi_1^r - \chi_0^r) \psi_r(t). \quad (19)$$

To account for the uncertainty in estimating the above models, we adopt the Bayesian paradigm and impose prior distributions for the parameters Kowal and Bourgeois (2020). For the basis function $\mathbf{b}(t)$ to construct principal components, we choose the thin-plate spline which takes the form $\mathbf{b}(t) = (1, t, (|t - k_1|)^3, \dots, |t - k_L|^3)' \in \mathcal{R}^{L+2}$, where the k_l ($l = 1, 2, \dots, L$) is the pre-defined knots on the time span. We set the values of knots k_l with the quantiles of observation time grids. For the parameters of principal components, taking the mediator model for an example, we impose the following priors on parameters in (12):

$$\phi_r \sim N(0, h_r^{-1} \Omega^{-1}), h_r \sim \text{Uniform}(\lambda_r^2, 10^4),$$

where $\Omega \in \mathcal{R}^{(L+2) \times (L+2)}$ is the roughness penalty matrix and $h_r > 0$ is the smooth parameter. This implies a Gaussian Process prior on $\psi_r(t)$ with mean function zero and covariance function $\text{Cov}(\psi_r(t), \psi_r(s)) = h_r \mathbf{b}'(s) \Omega \mathbf{b}(t)$. We choose the Ω such that $[\Omega_r]_{l,l'} = (k_l - k_{l'})^2$, when $l, l' > 2$, and $[\Omega_r]_{l,l'} = 0$ when $l, l' \leq 2$. For the distribution of principal scores in (13), we specify a multiplicative Gamma prior (Bhattacharya and Dunson, 2011; Montagna et al., 2012) on the variance to encourage shrinkage as r increasing,

$$\chi_0^r, \chi_1^r \sim N(0, \sigma_{\chi_r}^2), \quad \sigma_{\chi_r}^{-2} = \prod_{l \leq r} \delta_{\chi_l}, \quad \delta_{\chi_1} \sim \text{Ga}(a_{\chi_1}, 1), \quad \delta_{\chi_l} \sim \text{Ga}(a_{\chi_2}, 1), \quad l \geq 2,$$

$$\lambda_r^{-2} = \prod_{l \leq r} \delta_l, \quad \delta_1 \sim \text{Ga}(a_1, 1), \quad \delta_l \sim \text{Ga}(a_2, 1), \quad l \geq 2,$$

$$a_1, a_{\chi_1} \sim \text{Ga}(2, 1), \quad a_2, a_{\chi_2} \sim \text{Ga}(3, 1).$$

Further details on the hyperparameters of the priors can be found in Bhattacharya and Dunson (2011) and Durante (2017). For the coefficients of covariates β_M , we specify a diffused normal prior $\beta_M \sim \mathcal{N}(0, 100^2 * \mathbf{I}_{\dim(X)})$. We impose similar prior distributions for the parameters in the outcome model.

Posterior inference can be obtained by Gibbs sampling. The credible intervals of the causal effects τ_{TE}^t and τ_{ACME}^t can be easily constructed using the posterior sample of the parameters in the model. Details of the Gibbs sampler are provided in the Supplementary B.

5 Simulations

In this section, we conduct simulations to evaluate the operating characteristics of the proposed method and compare with two standard methods.

5.1 Simulation design

We generate 200 units to approximate the sample size in our application. For each unit, we make T_i observations at the time grid $\{t_{ij} \in [0, 1], j = 1, 2, \dots, T_i\}$. We draw T_i from a Poisson distribution with mean T and randomly pick t_{ij} uniformly:

$$T_i \sim \text{Poisson}(T), \quad t_{ij} \sim \text{Uniform}(0, 1), \quad j = 1, 2, \dots, T_i.$$

For each unit i and time j , we generate three covariates from a tri-variate Normal distribution, $\mathbf{X}_{ij} = (X_{ij1}, X_{ij2}, X_{ij3}) \sim \mathcal{N}([0, 0, 0]^T, \sigma_X^2 \mathbf{I}_3)$. We simulate the binary treatment indicator from $Z_i = \mathbf{1}\{c_{i1} > 0\}$, where $c_{i1} \sim \mathcal{N}(0, 1)$. To simulate the sparse and irregular mediator trajectories, we first simulate a smooth underlying process $M_i^t(z)$ for the mediators:

$$M_i^t(z) = 0.2 + \{0.2 + 2t + \sin(2\pi t)\}(z + 1) - X_{ij1} + 0.5X_{ij2} + \varepsilon_i^m(t) + c_{i2},$$

where the error term $\varepsilon_i^m(t) \sim \text{GP}(0, \sigma_m^2 \exp\{-8(s - t)^2\})$ is drawn from a Gaussian Process (GP) with an exponential kernel and σ_m^2 controlling the volatility of the realized curves, and $c_{i2} \sim \mathcal{N}(0, \sigma_m^2)$ to represent the individual random intercepts. The mean value of the mediator process depends on the covariates and time. The polynomial term and the trigonometric function of t introduce the long term growth trend and periodic fluctuations, respectively. Also, the coefficient of z evolves as the time changes, implying a time varying treatment effect on the mediator. Similarly, we specify a GP model

for the outcome process,

$$Y_i^t(z, \mathbf{m}) = \mathbf{m}^t + \cos(2\pi t) + 0.1t^2 + 2t + \{\cos(2\pi t) + 0.2t^2 + 3t\}z - 0.5X_{ij2} + X_{ij3} + \varepsilon_i^y(t) + c_{i3},$$

where the error term $\varepsilon_i^y(t) \sim \text{GP}(0, \sigma_y^2 \exp\{-8(s-t)^2\})$ is drawn from a GP, and $c_{i3} \sim \mathcal{N}(0, \sigma_y^2)$ controls the individual random effect for the outcome process.

The above settings imply non-linear true causal effects (τ_{TE}^t and τ_{ACME}^t) in time, which are shown as the dashed lines in Figure 3. Upon simulating the processes, we evaluate the potential values of the mediators and outcomes at the sampled time point t_{ij} to obtain the observed trajectories with measurement error:

$$M_{ij} \sim \mathcal{N}(\mathbf{M}_i^{t_{ij}}(Z_i), 1), \quad Y_{ij} \sim \mathcal{N}(\mathbf{Y}_i^{t_{ij}}(Z_i, \mathbf{M}_i^{t_{ij}}(Z_i)), 1).$$

We control the sparsity of the mediator and outcome trajectories by varying the value of T in the grid of (15, 30, 50, 100), namely the average number of observations for each individual.

We compare the proposed method in Section 4 (referred to as MFPCA) with two standard methods in longitudinal data analysis: the random effects model (Laird and Ware, 1982) and the generalized estimating equations (GEE) (Liang and Zeger, 1986). To facilitate the comparisons, we aggregate the time-varying mediation effects into the following scalar values:

$$\tau_{\text{ACME}} = \int_0^T \tau_{\text{ACME}}^t dt, \quad \tau_{\text{TE}} = \int_0^T \tau_{\text{TE}}^t dt.$$

The true values for τ_{ACME} and τ_{TE} in the simulations are 1.20 and 2.77 respectively.

For the random effects approach, we fit the following two models:

$$M_{ij} = X_{ij}^T \beta_M + s_m(T_{ij}) + \tau_m Z_i + r_{ij}^m + \varepsilon_{ij}^m, \quad (20)$$

$$Y_{ij} = X_{ij}^T \beta_Y + s_y(T_{ij}) + \tau_y Z_i + \gamma M_{ij} + r_{ij}^y + \varepsilon_{ij}^y, \quad (21)$$

where r_{ij}^m and r_{ij}^y are normally distributed random effect with zero means, $s_m(T_{ij})$ and $s_y(T_{ij})$ are thin plate splines to capture the nonlinear effect of time. To model the time dependency, we specify an AR(1)

correlation structure for the random effects, thus $\text{Corr}(r_{ij}^m, r_{ij+1}^m) = p_1$, $\text{Corr}(r_{ij}^y, r_{ij+1}^y) = p_2$, namely the correlation decay exponentially within the observations of a given unit. Given the above random effect model, the mediation effect and total effect can be calculated as: $\hat{\tau}_{\text{ACME}}^{\text{RD}} = \hat{\gamma}\hat{\tau}_m$, $\hat{\tau}_{\text{TE}}^{\text{RD}} = \hat{\gamma}\hat{\tau}_m + \hat{\tau}_y$.

For the GEE approach, we specify the following estimation equations:

$$E(M_{ij}|X_{ij}, Z_i) = X_{ij}^T\beta_M + \tau_m Z_i, \quad (22)$$

$$E(Y_{ij}|M_{ij}, X_{ij}, Z_i) = X_{ij}^T\beta_M + \tau_y Z_i + \gamma M_{ij}. \quad (23)$$

For the working correlation structure, we consider the AR(1) correlation for both the mediators and outcomes. Similarly, we obtain the estimations through $\hat{\tau}_{\text{ACME}}^{\text{GEE}} = \hat{\gamma}\hat{\tau}_m$, $\hat{\tau}_{\text{TE}}^{\text{GEE}} = \hat{\gamma}\hat{\tau}_m + \hat{\tau}_y$ with two different correlation structures.

It is worth noting that both the random effects model and the GEE model generally lack the flexibility to accommodate irregularly-spaced longitudinal data, which renders specifying the correlation between consecutive observations difficult. For example, though the AR(1) correlation takes into account the temporal structure of the data, it still imposes the correlation between any two consecutive observations to be constant, which is unlikely the case in the cases with irregularly-spaced data. Nonetheless, we compare the proposed method with these two models as they are the standard methods in longitudinal data analysis.

5.2 Simulation results

We apply the proposed MFPCA method, the random effects model and the GEE model in Section 5.1 to the simulated data $\{Z_i, \mathbf{X}_{ij}, M_{ij}, Y_{ij}\}$, to estimate the causal effects τ_{TE} and τ_{ACME} .

Figure 3 shows the causal effects and associated 95% credible interval estimated from MFPCA in one randomly selected simulated dataset under each of the four levels of sparsity T . Regardless of T , MFPCA appears to estimate the time-varying causal effects satisfactorily, with the 95% credible interval covering the true effects at any time. As expected, the accuracy of the estimation increases as the frequency of the observations increases.

Table 1 presents the absolute bias, root mean squared error (RMSE) and coverage rate of the 95%




Figure 3: Posterior mean of $\tau_{TE}^t, \tau_{ACME}^t$ and 95% credible intervals in one simulated dataset under each level of sparsity. The solid line represents the true surface for τ_{TE}^t and τ_{ACME}^t

confidence interval of τ_{TE} and τ_{ACME} under the MFPCA, the random effect model and the GEE model based on 200 simulated datasets for each level of sparsity T in $[15, 25, 50, 100]$. The performance of all three methods improve as the frequency of observations increases. With low frequency ($T < 100$), i.e. sparse observations, MFPCA consistently outperforms random effect model, which in turn outperforms GEE in all measures. The advantage of MFPCA over the other two methods diminishes as the frequency increases. In particular, with dense observations ($T = 100$), MFPCA leads to similar results as random effects, both outperforming GEE. The simulation results validate the use of our method in sparse case.

6 Empirical Application

6.1 The data

As discussed in Section 2, the goal of this application is to investigate the causal mediation mechanisms between early adversity, adult social bonds, and adult GC hormones levels. We apply the method and models proposed in Section 3 and 4 to a longitudinal dataset on wild baboons collected in the Amboseli Baboon Research Project. Here we first provide more information about the data.

Table 1: Absolute bias, RMSE and coverage rate of the 95% confidence interval of MFPCA, the random effect model and the generalized estimating equation (GEE) model under different frequency of observations in the simulations.

Method	τ_{TE}			τ_{ACME}		
	Bias	RMSE	Coverage	Bias	RMSE	Coverage
<i>T=15</i>						
MFPCA	0.103	0.154	88.4%	0.134	0.273	86.4%
Random effect	0.165	0.208	78.2%	0.883	1.673	69.5%
GEE	0.183	0.304	77.6%	0.987	2.051	61.8%
<i>T=25</i>						
MFPCA	0.092	0.123	92.3%	0.102	0.246	90.6%
Random effect	0.124	0.165	81.2%	0.679	1.263	72.3%
GEE	0.152	0.273	80.3%	0.860	1.753	64.4%
<i>T=50</i>						
MFPCA	0.087	0.112	93.5%	0.094	0.195	92.3%
Random effect	0.109	0.134	90.3%	0.228	0.497	88.8%
GEE	0.121	0.175	83.5%	0.236	0.493	80.8%
<i>T=100</i>						
MFPCA	0.053	0.089	94.3%	0.064	0.163	93.1%
Random effect	0.046	0.093	93.1%	0.053	0.154	92.8%
GEE	0.093	0.124	90.5%	0.098	0.161	90.3%

Our sample includes 192 female baboons and 11658 observations in total. We retain 10626 observations after removing the observations with missing social bonds or GCs levels information. The longitudinal observations on the strength of social bonds (mediators) and GC hormones concentrations. The social bonds range from -1.47 to 3.31 with mean value at 1.04 and standard deviation 0.51 . The

fecal GCs levels range from 7.51 to 982.87 with mean 74.13 and standard deviation 38.25. Age is used to index within-individual observations on both social bond and GCs levels. All the baboons enter into the study after becoming mature at age 5. However, data on both GCs levels and social bond strength for females older than 18 years are extremely sparse and volatile (only about 20% baboons survive until age 18). Therefore, we truncated all trajectories at age 18, resulting in a final sample with 192 female baboons and 9878 observations.

The six adversity sources (exposure) are drought, maternal death, competing sibling, high group density, low maternal rank, maternal social isolation. Table 2 presents the number of baboons experienced early early adversity. Overall, while only a small proportion of subjects experienced any given source of early adversity, most subjects had experienced at least one source of early adversity. Therefore, in our analysis we also create a cumulative exposure variable that summarizes the total number of adversities.

Table 2: Sources of early adversity and the number of baboons experienced each type of early adversity. The last row summarizes the number of baboons had at least one of six individual adversity sources.

early adversity	no. subjects did not experience (control)	no. subjects did experience (exposure)
Drought	164	28
Competing Sibling	153	39
High group density	161	31
Maternal death	157	35
Low maternal rank	152	40
Maternal Social isolation	140	52
At least one	48	144

The time-varying covariates include reproductive state, density in the social group, max temperature in last 30 days, whether the sample is collected in wet or dry season, the amount of rainfall. More

information can be found in Rosenbaum et al. (2020).

6.2 Results of FPCA

We first summarize the results of FPCA of the observed trajectories. We posit model (11) for the social bonds and Model (17) for the GCs levels, with some modifications. First, we added two random effects, one for social group and one for hydrological year, in both models. Second, in the outcome model, we use the log transformed GCs level instead of the original scale as the outcome, which allows us to interpret the coefficient of early adversity as the effect on the percentage change of the GCs level. For both the mediator and outcome processes, the first three functional principal components explain more than 90% of the total variation and we will use them in the structural equation model for mediation analysis. Figure 4 shows the first two principal components extracted from the mediator (left panel) and outcome (right panel) processes. For the social bonds process, the first two principal components explain 53% and 31% of the total variation. The first component depicts a drastic change in the early stage of a baboon's life and stabilizes afterwards. For the GCs process, the first two functional principal components explain 54% and 34% of the total variation. The first component depicts a stable trend throughout the life span.

The left panel of Figure 5 displays the observed trajectory of GCs versus the posterior mean of its imputed smooth process of three baboons who experienced zero (EAG), one (OCT) and two (GUI) sources of early adversity, respectively. We can see that the imputed smooth process generally captures the overall time trend of each subject while reduce the noise in the observations. The pattern is similar for the social bonds, which is shown in Supplementary C with a few more randomly selected subjects. Recall that each subject's observed trajectory is fully captured by its vector of principal scores, and thus the principal scores of the first few dominant principal components adequately summarize the whole trajectory. The right panel of Figure 5 shows the principal scores of the first (X-axis) versus second (Y-axis) principal component for the GCs process of all subjects in the sample, color-coded based on the number of early adversities experienced. We can see that significant difference exists in the distributions of the first two principal scores between the group who experienced no early adversity and the groups




Figure 4: The first two functional principal components of the process of the mediator, i.e. social bonds (left panel) and the outcome, i.e., GCs level (right panel).

who experienced at least one source of adversity.

6.3 Results of causal mediation analysis

Before proceeding to the causal mediation analysis, we first assess the plausibility of the key causal assumptions in the application. The ignorability assumption states that there is no unmeasured confounding, besides the observed covariates, between early adversity and the social bond and GCs processes. This is plausible in our application because the six sources of early adversity (e.g. maternal death, drought, low maternal rank) are all largely randomized by the nature. The sequential ignorability assumption states that there is no unmeasured confounding, besides the observed covariates and the history of social bond strength, between the social bond process and the GCs process. One possible violation can be due to the ‘feedback’ between the social bond and GCs processes. We have performed a sensitivity analysis by adding (a) the most recent prior observed GCs value, or (b) the average of all past observed GCs values, as a predictor in the mediation model, which lead to little difference in the results and thus bolsters sequential ignorability. Though we are not aware of the existence of other sequential confounders, we also cannot rule them out.




Figure 5: Left panel: Observed trajectory of GCs versus the posterior mean of its imputed smooth process of three baboons who experienced zero (EAG), one (OCT) and two (GUI) sources of early adversity, respectively. Right panel: Principal scores of the first (X-axis) versus second (Y-axis) principal component for the GCs process of all subjects in the sample; color-coded based on the number of early adversities experienced.

We perform a separate causal mediation analysis for each source of early adversity. Table 3 presents the posterior mean and 95% credible interval of the total effect (TE), direct effect (ANDE) and indirect effect mediated through social bonds (ACME) of each source of early adversity on adult GCs level, as well as the effects of early adversity on the mediator (social bonds). First, from the first column of Table 3 we can see that experiencing any source of early adversity would reduce a baboon's strength of social bond with other baboons in the adulthood. The negative effect is particularly severe for those experiencing drought, high group density or maternal death in early life. For example, compared with the baboons who did not experience any early adversity, the baboons who experienced maternal death have a 0.221 unit decrease in social bonds, translating to approximately 0.4 standard deviation of social bonds in the population. Overall, experiencing at least one source of adversity have a decrease of 0.2 standard deviation of social bonds in adulthood.

Second, from the second column of Table 3 we can see a strong total effect of early adversity on

female baboon's GCs levels across adulthood. Baboons who experienced at least one source of adversity had GCs concentrations that were approximately 9% higher than their peers who did not experience any adversity. Although the range of total effect sizes across all individual adversity sources varies from 4% to 14%, the point estimates are consistent toward higher GCs levels, even for the early adversity sources of which the credible interval includes zero. Among the individual sources of adversity, females who were born during a drought, into a high-density group, or to a low-ranking mother had particularly elevated GCs concentrations (12-14%) in adulthood, although the credible interval of high group density includes zero.

Table 3: Total, direct and indirect causal effects of individual and cumulative sources of early adversity sources on social bonds and GC concentrations in adulthood in wild female baboons. 95% credible intervals are in the parenthesis.

Source of adversity	effect on mediator	τ_{TE}	τ_{ACME}	τ_{ANDE}
Drought	-0.164 (-0.314, -0.014)	0.124 (0.007, 0.241)	0.009 (0.000, 0.017)	0.114 (0.005, 0.222)
Competing sibling	-0.106 (-0.249, 0.030)	0.084 (-0.008, 0.172)	0.006 (0.003, 0.009)	0.078 (-0.012, 0.163)
High group density	-0.271 (-0.519, -0.023)	0.123 (-0.052, 0.281)	0.015 (0.000, 0.029)	0.108 (-0.053, 0.252)
Maternal death	-0.221 (-0.423, -0.019)	0.061 (-0.006, 0.129)	0.011 (0.005, 0.014)	0.049 (-0.014, 0.113)
Low maternal rank	-0.052 (-0.298, 0.001)	0.134 (0.011, 0.256)	0.008 (0.005, 0.011)	0.126 (0.008, 0.244)
Maternal social isolation	-0.040 (-0.159, 0.095)	0.035 (-0.045, 0.116)	0.002 (0.000, 0.005)	0.033 (-0.044, 0.111)
At least one	-0.102 (-0.195, -0.008)	0.092 (0.005, 0.178)	0.007 (0.002, 0.009)	0.084 (0.009, 0.159)

Third, while female baboons who experienced harsh conditions in early life show higher GCs levels in adulthood, we found no evidence that these effects were significantly mediated by the absence of strong social bonds. Specifically, the mediation effect τ_{ACME} (third column in Table 3) is consistently small; the strength of females social bonds with other females accounted for a difference in GCs of only 0.85% when averaged across the six individual adversity sources, even though it had credible intervals that did not include zero for five of the six individual sources of adversity. On the other hand, the direct effects τ_{ANDE} (fourth column in Table 3) are much stronger than the mediation effect. When averaged across the six adversity sources, the direct effect of early adversity on GCs level was 11.6 times stronger than the mediation effect running through social bonds. For example, for females who experienced at least one source of early adversity, the direct effect can explain 8.4% increase in GCs level, while the mediation effect only takes up 0.7% for the increase in GCs.

The above findings of the causal relationships between early adversity, social bonds, and GCs levels in wild baboons are compatible with observations in many other species that early adversity and weak relationships both give rise to poor health, and that early adversity predicts various forms of social dysfunction, including weaker relationships. However, they call into question the notion that social bonds play a major role in mediating the effect of early adversity on poor health. In wild female baboons, any such effect appears to be functionally biologically irrelevant, and what little exists is limited strictly to their relationships with other females.

7 Discussion

We proposed a framework for conducting causal mediation analysis with sparse and irregular longitudinal data. We defined several causal estimands (total, direct and indirect effects) in such settings and developed assumptions to nonparametrically identify these effects. For estimation and inference, we combine functional principal component analysis (FPCA) techniques and the standard two structural-equation-model system. In particular, we use a Bayesian FPCA model to reduce the dimensions of the observed trajectories of mediators and outcomes. We applied the proposed method to analyze the causal

effects of early adversity on adult social bonds and adult GC hormone level in a sample wild female baboons. We found that experiencing adversity before maturity generally hampers a baboon's ability to build social bond with other baboons and increase the GC hormones level in adulthood, which in turn lead to stress and diseases. However, the effects on the GC concentration from early adversity is not mediated by social bonds.

One limitation of our analysis is that the identification of mediation effects (ACME and ANDE) relies on strong assumptions, particularly sequential ignorability. Though not routinely performed in the literature of mediation analysis, a formal sensitivity analysis would shed light on how reliant the proposed method is on sequential ignorability. Given the complexity of mediation analysis, a model-based approach appears to be the most feasible for sensitivity analysis (Imai et al., 2010). Alternatively, one could consider the framework developed by Didelez et al. (2012); VanderWeele et al. (2014) and relax sequential ignorability to allow for observed treatment-induced mediator-outcome confounding. This framework targets at a different set of causal estimands, namely the interventional direct or indirect effect, instead of the natural direct or indirect effects considered in this paper.

We proposed a framework for conducting causal mediation analysis with sparse and irregular longitudinal data. We defined several causal estimands (total, direct and indirect effects) in such settings and developed assumptions to nonparametrically identify these effects. For estimation and inference, we combine functional principal component analysis (FPCA) techniques and the standard two structural-equation-model system. In particular, we use a Bayesian FPCA model to reduce the dimensions of the observed trajectories of mediators and outcomes. We applied the proposed method to analyze the causal effects of early adversity on adult social bonds and adult GC hormone level in a sample wild female baboons. We found that experiencing adversity before maturity generally hampers a baboon's ability to build social bond with other baboons and increase the GC hormones level in adulthood, which in turn lead to stress and diseases. However, the effects on the GC concentration from early adversity is not mediated by social bonds.

One limitation of our analysis is that the identification of mediation effects (ACME and ANDE) relies on strong assumptions, particularly sequential ignorability. Though not routinely performed in

the literature of mediation analysis, a formal sensitivity analysis would shed light on how reliant the proposed method is on sequential ignorability. Given the complexity of mediation analysis, a model-based approach appears to be the most feasible for sensitivity analysis (Imai et al., 2010). Alternatively, one could consider the framework developed by Didelez et al. (2012); VanderWeele et al. (2014) and relax sequential ignorability to allow for observed treatment-induced mediator-outcome confounding. This framework targets at a different set of causal estimands, namely the interventional direct or indirect effect, instead of the natural direct or indirect effects considered in this paper.

An important extension of our method is to incorporate time-to-event outcomes, a common setting in longitudinal studies (Lange et al., 2012; VanderWeele, 2011). For example, it is of much scientific interest to extend our application to investigate the causal mechanisms between early adversity, social bonds, GCs level and survival time. A typical complication in mediation analysis with time-to-event outcomes and time-varying mediators is that the mediators are undefined for the time period in which a unit was not observed (Didelez, 2019; Vansteelandt et al., 2019). Within our framework, we can bypass this problem by imputing the underlying process of the mediators in an identical range for every unit.

Acknowledgements

We thank Surya Tokdar, Stacy Rosenbaum, Fernando Campos, and Georgia Papadogeorgou for helpful discussions. This research is supported by NIH grants 1R01AG053330-01A1 and 1R01 AG053308-01A1.

Supplementary

A. Proof of Theorem 1

For the first part, identification of total effect, we have for any $z \in \{0, 1\}$,

$$E(Y_i^t | Z_i = z, \mathbf{X}_i^t) = E(Y_i^t(z, \mathbf{M}_i(z)) | Z_i = z, \mathbf{X}_i^t) = E(Y_i^t(z, \mathbf{M}_i(z)) | \mathbf{X}_i^t).$$

The second equality follows from Assumption 1. Therefore, we prove the identification of τ_{TE}^t ,

$$\begin{aligned}\tau_{TE}^t &= \int_X \{E(Y_i^t(1, \mathbf{M}_i(1)) | \mathbf{X}_i^t) - E(Y_i^t(0, \mathbf{M}_i(0)) | \mathbf{X}_i^t)\} dF_{\mathbf{X}_i^t}(\mathbf{x}^t), \\ &= \int_X \{E(Y_i^t | Z_i = 1, \mathbf{X}_i^t = \mathbf{x}^t) - E(Y_i^t | Z_i = 0, \mathbf{X}_i^t = \mathbf{x}^t)\} dF_{\mathbf{X}_i^t}(\mathbf{x}^t),\end{aligned}$$

For the second part, identification of τ_{ACME}^t , we make the following regularity assumptions. Suppose the potential outcomes $Y_i^t(z, \mathbf{m})$ as a function of \mathbf{m} is Lipschitz continuous on $[0, T]$ with probability one. There exists $A < \infty$ $|Y_i^t(z, \mathbf{m}) - Y_i^t(z, \mathbf{m}')| \leq A \|\mathbf{m} - \mathbf{m}'\|_2$, for any $z, t, \mathbf{m}, \mathbf{m}'$ almost surely.

For any $z, z' \in \{0, 1\}$, we have

$$\begin{aligned}& \int_{\mathcal{X}} \int_{R^{[0,t]}} E(Y_i^t | Z_i = z, \mathbf{X}_i^t = \mathbf{x}^t, \mathbf{M}_i^t = \mathbf{m}) dF_{\mathbf{X}_i^t}(\mathbf{x}^t) \times d\{F_{\mathbf{M}_i^t | Z_i = z', \mathbf{X}_i^t = \mathbf{x}^t}(\mathbf{m})\} \\ &= \int_{\mathcal{X}} \int_{R^{[0,t]}} E(Y_i^t(z, \mathbf{m}) | Z_i = z, \mathbf{X}_i^t = \mathbf{x}^t, \mathbf{M}_i^t = \mathbf{m}) \times d\{F_{\mathbf{M}_i^t | Z_i = z', \mathbf{X}_i^t = \mathbf{x}^t}(\mathbf{m})\}.\end{aligned}$$

For any path \mathbf{m} on the time span $[0, t]$, we make a finite partition into H pieces at points $\mathcal{M}_H = \{t_0 = 0, t_1 = t/H, t_2 = 2t/H, \dots, t_H = t\}$. Now we consider using a step functions with jumps at points \mathcal{M}_H . Denote the step function as \mathbf{m}_H , which is:

$$\mathbf{m}_H(x) = \begin{cases} \mathbf{m}(0) = m_0 & 0 \leq x < t/H, \\ \mathbf{m}(t/H) = m_1 & t/H \leq x < 2t/H, \\ \dots & \\ \mathbf{m}((H-1)t/H) = m_H & (H-1)t/H \leq x \leq t. \end{cases}$$

We wish to use this step function $\mathbf{m}_H(x)$ to approximate function \mathbf{m} . First, given \mathbf{m} is Lipschitz continuous, there exists $B > 0$ such that $|m(x_1) - m(x_2)| \leq B|x_1 - x_2|$. Therefore, the step functions \mathbf{m}_H approximates the original function \mathbf{m} well in the sense that,

$$\|\mathbf{m}_H - \mathbf{m}\|_2 \leq \sum_{i=1}^H \frac{t}{H} B^2 \frac{t^2}{H^2} \asymp O(H^{-2}).$$

As such we can approximate the expectation over a continuous process with expectation on a vector with values on the jumps, (m_0, m_1, \dots, m_H) . That is,

$$\begin{aligned} & \int_{\mathcal{X}} \int_{R^{[0,t]}} E(Y_i^t(z, \mathbf{m}) | Z_i = z, \mathbf{X}_i^t = \mathbf{x}^t, \mathbf{M}_i^t = \mathbf{m}) \times d\{\mathbf{F}_{\mathbf{M}_i^t | Z_i = z', \mathbf{X}_i^t = \mathbf{x}^t}(\mathbf{m})\} \\ & \asymp \int_{\mathcal{X}} \int_{R^{[0,t]}} E(Y_i^t(z, \mathbf{m}_H) | Z_i = z, \mathbf{X}_i^t = \mathbf{x}^t, \mathbf{M}_i^t = \mathbf{m}_H) \\ & \quad \times d\{\mathbf{F}_{\mathbf{M}_i^t | Z_i = z', \mathbf{X}_i^t = \mathbf{x}^t}(\mathbf{m}_H)\} + O(H^{-2}). \end{aligned}$$

This equivalence follows from the regularity condition that the potential outcome $Y_i^t(z, \mathbf{m})$ as a function of \mathbf{m} is continuous with the L_2 metrics of \mathbf{m} . As the values of steps function \mathbf{m}_H are completely determined by the values on finite jumps, we can further reduce to,

$$\begin{aligned} & \asymp \int_{\mathcal{X}} \int_{R^H} E(Y_i^t(z, \mathbf{m}_H) | Z_i = z, \mathbf{X}_i^t = \mathbf{x}^t, m_0, m_1, m_2, \dots, m_H) \\ & \quad \times d\{\mathbf{F}_{m_0, m_1, \dots, m_H | Z_i = z', \mathbf{X}_i^t = \mathbf{x}^t}(m_0, m_1, m_2, \dots, m_H)\} + O(H^{-2}). \end{aligned}$$

With Assumption 1, we can show that

$$\begin{aligned} & d\{\mathbf{F}_{m_0, m_1, \dots, m_H | Z_i = z', \mathbf{X}_i^t = \mathbf{x}^t}(m_0, m_1, m_2, \dots, m_H)\} \\ & = d\{\mathbf{F}_{m_0(z'), m_1(z'), \dots, m_H(z') | \mathbf{X}_i^t = \mathbf{x}^t}(m_0, m_1, m_2, \dots, m_H)\}, \\ & = d\{\mathbf{F}_{\mathbf{m}_H(z') | \mathbf{X}_i^t = \mathbf{x}^t}(\mathbf{m}_H)\}. \end{aligned}$$

With a slightly abuse of notations, we use $\mathbf{m}_H(z)$ to denote the potential process induced by the original potential process $\mathbf{M}_i^t(z)$ and $m_i(z)$ to denote potential values of $\mathbf{M}_i^t(z)$ evaluated at point $x_i = it/H$. Also, with the assumption 2, we can choose a large H such that $t/H \leq \varepsilon$. Then we have the following

conditional independence conditions,

$$\begin{aligned}
& Y_i^0(z, \mathbf{m}_H) \perp\!\!\!\perp m_0 | Z_i, \mathbf{X}_i^t, \\
& \{Y_i^{t/H}(z, \mathbf{m}_H) - Y_i^0(z, \mathbf{m}_H)\} \perp\!\!\!\perp (m_1 - m_0) | Z_i, \mathbf{X}_i^t, \mathbf{m}_H^0, \\
& \{Y_i^{2t/H}(z, \mathbf{m}_H) - Y_i^{t/H}(z, \mathbf{m}_H)\} \perp\!\!\!\perp (m_2 - m_1) | Z_i, \mathbf{X}_i^t, \mathbf{m}_H^{t/H}, \\
& \dots \\
& \{Y_i^t(z, \mathbf{m}_H) - Y_i^{t(H-1)/H}(z, \mathbf{m}_H)\} \perp\!\!\!\perp (m_H - m_{H-1}) | Z_i, \mathbf{X}_i^t, \mathbf{m}_H^{t(H-1)/H},
\end{aligned}$$

where are equivalent to,

$$\begin{aligned}
& Y_i^0(z, \mathbf{m}_H) \perp\!\!\!\perp m_0 | Z_i, \mathbf{X}_i^t, \\
& \{Y_i^{t/H}(z, \mathbf{m}_H) - Y_i^0(z, \mathbf{m}_H)\} \perp\!\!\!\perp (m_1 - m_0) | Z_i, \mathbf{X}_i^t, m_0, \\
& \{Y_i^{2t/H}(z, \mathbf{m}_H) - Y_i^{t/H}(z, \mathbf{m}_H)\} \perp\!\!\!\perp (m_2 - m_1) | Z_i, \mathbf{X}_i^t, m_0, m_1, \\
& \dots \\
& \{Y_i^t(z, \mathbf{m}_H) - Y_i^{t(H-1)/H}(z, \mathbf{m}_H)\} \perp\!\!\!\perp (m_H - m_{H-1}) | Z_i, \mathbf{X}_i^t, m_0, m_1, \dots, m_{H-1},
\end{aligned}$$

as the step function $m_H^{it/H}$ is completely determined by values m_0, \dots, m_i . With the above conditional independence, we have,

$$\begin{aligned}
& E(Y_i^t(z, \mathbf{m}_H) | Z_i = z, \mathbf{X}_i^t = \mathbf{x}^t, m_0, m_1, m_2, \dots, m_H) \\
& = E(Y_i^t(z, \mathbf{m}_H) | Z_i = z, \mathbf{X}_i^t = \mathbf{x}^t).
\end{aligned}$$

With similar arguments, it also equals:

$$\begin{aligned}
& E(Y_i^t(z, \mathbf{m}_H) | Z_i = z, \mathbf{X}_i^t = \mathbf{x}^t) = E(Y_i^t(z, \mathbf{m}_H) | Z_i = z', \mathbf{X}_i^t = \mathbf{x}^t), \\
& = E(Y_i^t(z, \mathbf{m}_H) | Z_i = z, \mathbf{X}_i^t = \mathbf{x}^t, m_0 = m_0(z'), \dots, m_H = m_H(z')), \\
& = E(Y_i^t(z, \mathbf{m}_H) | Z_i = z, \mathbf{X}_i^t = \mathbf{x}^t, \mathbf{m}_H(z') = \mathbf{m}_H), \\
& = E(Y_i^t(z, \mathbf{m}_H) | \mathbf{X}_i^t = \mathbf{x}^t, \mathbf{m}_H(z') = \mathbf{m}_H).
\end{aligned}$$

As a conclusion, we have shown that,

$$\begin{aligned}
& \int_{\mathcal{X}} \int_{R^{[0,t]}} E(Y_i^t(z, \mathbf{m}) | Z_i = z, \mathbf{X}_i^t = \mathbf{x}^t, \mathbf{M}_i^t = \mathbf{m}) \times d\{F_{\mathbf{M}_i^t | Z_i = z', \mathbf{X}_i^t = \mathbf{x}^t}(\mathbf{m})\}, \\
& \asymp \int_{\mathcal{X}} \int_{R^{[0,t]}} E(Y_i^t(z, \mathbf{m}_H) | \mathbf{X}_i^t = \mathbf{x}^t, \mathbf{m}_H(z') = \mathbf{m}_H) \\
& \quad \times d\{F_{\mathbf{m}_H(z') | \mathbf{X}_i^t = \mathbf{x}^t}(\mathbf{m}_H)\} + O(H^{-2}), \\
& \asymp \int_{\mathcal{X}} E(Y_i^t(z, \mathbf{m}_H(z')) | \mathbf{X}_i^t = \mathbf{x}^t) + O(H^{-2}), \\
& \asymp \int_{\mathcal{X}} E(Y_i^t(z, \mathbf{m}(z')) | \mathbf{X}_i^t = \mathbf{x}^t) + O(H^{-2}).
\end{aligned}$$

The last equivalence comes from the regularity condition of $Y_i^t(z, \mathbf{m}(z'))$ as a function of $\mathbf{m}(z')$. Let H goes to infinity, we have,

$$\begin{aligned}
& \int_{\mathcal{X}} \int_{R^{[0,t]}} E(Y_i^t | Z_i = z, \mathbf{X}_i^t = \mathbf{x}^t, \mathbf{M}_i^t = \mathbf{m}) dF_{\mathbf{X}_i^t}(\mathbf{x}^t) \times d\{F_{\mathbf{M}_i^t | Z_i = z', \mathbf{X}_i^t = \mathbf{x}^t}(\mathbf{m})\} \\
& \quad = \int_{\mathcal{X}} E(Y_i^t(z, \mathbf{m}(z')) | \mathbf{X}_i^t = \mathbf{x}^t) dF_{\mathbf{X}_i^t}(\mathbf{x}^t).
\end{aligned}$$

With this relationship established, it is straightforward to show that,

$$\begin{aligned}
\tau_{\text{ACME}}^t(z) &= \int_{\mathcal{X}} \{E(Y_i^t(z, \mathbf{m}(1)) | \mathbf{X}_i^t = \mathbf{x}^t) - E(Y_i^t(z, \mathbf{m}(0)) | \mathbf{X}_i^t = \mathbf{x}^t)\} dF_{\mathbf{X}_i^t}(\mathbf{x}^t), \\
&= \int_{\mathcal{X}} \int_{R^{[0,t]}} E(Y_i^t | Z_i = z, \mathbf{X}_i^t = \mathbf{x}^t, \mathbf{M}_i^t = \mathbf{m}) dF_{\mathbf{X}_i^t}(\mathbf{x}^t) \times \\
& \quad d\{F_{\mathbf{M}_i^t | Z_i = 1, \mathbf{X}_i^t = \mathbf{x}^t}(\mathbf{m}) - F_{\mathbf{M}_i^t | Z_i = 0, \mathbf{X}_i^t = \mathbf{x}^t}(\mathbf{m})\},
\end{aligned}$$

which completes the proof.

B. Gibbs Sampler

In this section, we provide detailed descriptions on the Gibbs sampler for the model in Section 4. We only include the sampler for mediator process as the sampling procedure is essentially identical for the outcome process. For simplicity, we introduce some notations to represent vector values, $M_i = (M_{i1}, M_{i2}, \dots, M_{in_i}) \in \mathcal{R}^{T_i}$, $X_i = [X_{i1}, X_{i2}, \dots, X_{in_i}]' \in \mathcal{R}^{T_i \times p}$, $\psi_r(\mathbf{t}_i) = [\psi_r(t_{i1}), \dots, \psi_r(t_{in_i})] \in \mathcal{R}^{T_i}$

1. **Sample the eigen function** $\psi_r(t), r = 1, 2 \dots, R$.

- (a) $\psi_r | \dots \sim N(Q_{\phi_{r,ss-1} l_{\phi_r}}, Q_{\phi_r}^{-1})$ conditional on $C_r \psi_r = 0, C_r = [\psi_1, \psi_2, \dots, \psi_{r-1}, \psi_{r+1}, \dots, \psi_R]' B_G' B_G$, B_G is the basis functions evaluated at a equal spaced grids on $[0,1], \{t_1, t_2, \dots, t_G\}$, $G = 50$ for example, $B_G = [\mathbf{b}(t_1), \dots, \mathbf{b}(t_G)]' \in R^{G \times (L+2)}$. The corresponding mean and covairance functions are,

$$Q_{\psi_r} = \frac{\sum_{i=1}^N B_i' B_i \zeta_{r,i}^2}{\sigma_m^2} + h_k \Omega,$$

$$l_{\psi_r} = \frac{\sum_{i=1}^N B_i' \zeta_{i,r} (M_i - X_i \beta_M^T - \sum_{r' \neq r}^R \psi_r(\mathbf{t}_i) \zeta_{r',i})}{\sigma_m^2}.$$

Update the $\psi_r \leftarrow \psi_r / \sqrt{\psi_r' B_G' B_G \psi_r} = \psi_r / \|\psi_r(t)\|_2$ to ensure $\|\psi_r(t)\|_2 = 1$ and $\psi_r(t) = \mathbf{b}(t) \psi_r$ and update $\zeta_{r,i} \Rightarrow \zeta_{r,i} * \|\psi_r(t)\|_2$ to maintain likelihood function.

- (b) $h_k | \dots \sim \text{Ga}((L+1)/2, \psi_r' \Omega \psi_r)$ truncated on $[\lambda_r^2, 10^4]$.

2. **Sample the principal score** $\zeta_{r,i}, \zeta_{r,i} | \dots \sim N(\mu_r / \lambda_r^2, \lambda_r^2)$

$$\sigma_r^2 = (\|\psi_r(\mathbf{t}_i)\|_2^2 / \sigma_m^2 + \xi_{i,r} / \lambda_r^2)^{-1}$$

$$\mu_r = \frac{(M_i - X_i \beta_M^T - (\sum_{r' \neq r} \phi_{r'}(\mathbf{t}_i) \zeta_{r',i}))' \psi_r(\mathbf{t}_i)}{\sigma_\varepsilon^2} + \frac{(\tau_{0,r}(1 - Z_i) + \tau_{1,r} Z_i) \xi_{i,r}}{\lambda_r^2}$$

3. **Sample the causal parameters** χ_0^r, χ_1^r . Let $\chi_z = (\chi_z^r, \dots, \chi_z^R), z = 0, 1, \chi_z^r | \dots \sim N(Q_{z,r}^{-1} l_{z,r}, Q_{z,r}^{-1})$.

$$Q_{z,r} = \left(\sum_{i=1}^N \xi_{r,i} \mathbf{1}_{Z_i=z} / \lambda_r^2 + 1 / \sigma_{\chi_r}^2 \right)^{-1}$$

$$l_{z,r} = \sum_{i=1}^N \zeta_{r,i} \xi_{r,i} \mathbf{1}_{Z_i=z} / \lambda_r^2$$

4. **Sample the coefficients** β_M . The coefficients for covariates are $\beta_M | \dots \sim N(Q_\beta^{-1} \mu_\beta, Q_\beta^{-1})$,

$$Q_\beta = X' X / \sigma_m^2 + 100^2 I_{\dim(X)}$$

$$\mu_\beta = \sum_{i=1}^N X_i' (M_i - \sum_{r=1}^R \psi_r(\mathbf{t}_i) \zeta_{i,r}) / \sigma_m^2$$

5. Sample the precision/variance parameters.

- (a) $\sigma_m^{-2} | \dots \sim \text{Ga}(\sum_{i=1}^N T_i/2, \sum_{i=1}^N \|M_i - X_i \beta'_M - \sum_{r=1}^R \psi_r(\mathbf{t}_i) \zeta_{i,r}\|_2^2/2)$
- (b) $\sigma_{\chi_r}^2 | \dots,$

$$\delta_{\chi_1} | \dots \sim \text{Ga}(a_{\chi_1} + R, 1 + \frac{1}{2} \sum_{r=1}^R \chi_1^{(r)} (\chi_0^{r2} + \chi_1^{r2})), \chi_l^{(r)} = \prod_{i=l+1}^r \delta_{\chi_i}$$

$$\delta_{\chi_r} | \dots \sim \text{Ga}(a_{\chi_2} + R + 1 - r, 1 + \frac{1}{2} \sum_{r'=r}^R \chi_{r'}^{(r)} (\tau_0^{r'2} + \chi_1^{r'2})), r \geq 2,$$

$$\sigma_{\chi_r}^{-2} = \prod_{r'=1}^r \delta_{\chi_{r'}}.$$

- (c) $\lambda_r^2 | \dots,$

$$\delta_1 | \dots \sim \text{Ga}(a_1 + RN/2, 1 + \frac{1}{2} \sum_{r=1}^R \chi_1^{(r)'} \xi_{i,r} (\zeta_{i,r} - (1 - Z_i) \chi_0^r - Z_i \chi_1^r)^2),$$

$$\chi_l^{(r)'} = \prod_{i=l+1}^r \delta_i$$

$$\delta_r | \dots \text{Ga}(a_2 + (R - r + 1)N/2,$$

$$1 + \frac{1}{2} \sum_{r'=r}^R \chi_{r'}^{(r)'} \xi_{i,r'} (\zeta_{i,r'} - (1 - Z_i) \chi_0^{r'} - Z_i \chi_1^{r'})^2), r \geq 2$$

$$\lambda_r^{-2} = \prod_{r'=1}^r \delta_{r'}.$$

- (d) $\xi_{i,r} | \dots \sim \text{Ga}(\frac{v+1}{2}, \frac{1}{2} (v + (\zeta_{i,r'} - (1 - Z_i) \chi_0^{r'} - Z_i \chi_1^{r'})^2 / \lambda_r^2))$.
- (e) $a_1, a_2, a_{\chi_1}, a_{\chi_2}$ can be sampled with Metropolis-Hasting algorithm.

The sampling for the outcomes model Y_{ij} is similar to that for the mediator model except that we added the imputed value of the mediator process $M(t_{ij})$ as a covariate.

C. Imputed processes of the mediators and outcomes of eight subjects

Figure 6 shows the posterior means of the imputed smooth processes of the mediators and the outcomes against their respective observed trajectories of eight randomly selected subjects in the sample. For

social bonds (left panel of Figure 6), the imputed smooth process adequately captures the overall time trend of each subject while reduce the noise in the observations, evident in the subjects with code name HOK, DUI and LOC.




Figure 6: The imputed underlying smooth process against the observed trajectories for social bonds (left panel) and GCs levels (right panel).

For the subjects with few observations or observations concentrating in a short time span, such as subject NEA, the imputed process matches the trend of the observations while extrapolating to the rest of the time span with little information. FPCA achieves this by borrowing information from other units when learning the principal component on the population level. Compared with social bonds, variation of the adult GCs levels across the lifespan is much smaller. In the right panel in Figure 6, we can see the

imputed processes for the GCs levels are much flatter than those for social bonds. It appears that most variation in the GCs trajectories is due to noise rather than intrinsic developmental trend.

References

- Alberts, S. C. and Altmann, J. (2012), “The Amboseli Baboon Research Project: 40 years of continuity and change,” in *Long-term Field Studies of Primates*, Springer, 261–287.
- Anderson, M. and Marmot, M. (2011), “The effects of promotions on heart disease: Evidence from Whitehall,” *The Economic Journal*, 122, 555–589.
- Baron, R. M. and Kenny, D. A. (1986), “The moderator–mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations.” *Journal of Personality and Social Psychology*, 51, 1173.
- Bateson, P., Barker, D., Clutton-Brock, T., Deb, D., D’udine, B., Foley, R. A., Gluckman, P., Godfrey, K., Kirkwood, T., Lahr, M. M., et al. (2004), “Developmental plasticity and human health,” *Nature*, 430, 419.
- Bhattacharya, A. and Dunson, D. B. (2011), “Sparse Bayesian infinite factor models,” *Biometrika*, 291–306.
- Bind, M.-A., Vanderweele, T., Coull, B., and Schwartz, J. (2015), “Causal mediation analysis for longitudinal data with exogenous exposure,” *Biostatistics*, 17, 122–134.
- Case, A. and Paxson, C. (2011), “The long reach of childhood health and circumstance: evidence from the Whitehall II Study,” *The Economic Journal*, 121, F183–F204.
- Cohen, S. and Wills, T. A. (1985), “Stress, social support, and the buffering hypothesis.” *Psychological Bulletin*, 98, 310.
- Daniels, M. J., Roy, J. A., Kim, C., Hogan, J. W., and Perri, M. G. (2012), “Bayesian inference for the causal effect of mediation,” *Biometrics*, 68, 1028–1036.

- Didelez, V. (2019), “Defining causal mediation with a longitudinal mediator and a survival outcome,” *Lifetime Data Analysis*, 25, 593–610.
- Didelez, V., Dawid, P., and Geneletti, S. (2012), “Direct and indirect effects of sequential treatments,” *arXiv preprint arXiv:1206.6840*.
- Durante, D. (2017), “A note on the multiplicative gamma process,” *Statistics & Probability Letters*, 122, 198–204.
- Gluckman, P. D., Hanson, M. A., Cooper, C., and Thornburg, K. L. (2008), “Effect of in utero and early-life conditions on adult health and disease,” *New England Journal of Medicine*, 359, 61–73.
- Goldsmith, J., Greven, S., and Crainiceanu, C. (2013), “Corrected confidence bands for functional data using principal components,” *Biometrics*, 69, 41–51.
- Holt-Lunstad, J., Smith, T. B., and Layton, J. B. (2010), “Social relationships and mortality risk: a meta-analytic review,” *PLoS Medicine*, 7, e1000316.
- Imai, K., Keele, L., and Yamamoto, T. (2010), “Identification, inference and sensitivity analysis for causal mediation effects,” *Statistical Science*, 51–71.
- Jiang, C.-R., Wang, J.-L., et al. (2010), “Covariate adjusted functional principal components analysis for longitudinal data,” *The Annals of Statistics*, 38, 1194–1226.
- (2011), “Functional single index models for longitudinal data,” *The Annals of Statistics*, 39, 362–388.
- Kim, C., Daniels, M. J., Hogan, J. W., Choirat, C., and Zigler, C. M. (2019), “Bayesian methods for multiple mediators: Relating principal stratification and causal mediation in the analysis of power plant emission controls,” *The Annals of Applied Statistics*, 13, 1927.
- Kowal, D. R. and Bourgeois, D. C. (2020), “Bayesian Function-on-Scalars Regression for High-Dimensional Data,” *Journal of Computational and Graphical Statistics*, 1–26.

- Laird, N. M. and Ware, J. H. (1982), “Random-effects models for longitudinal data,” *Biometrics*, 963–974.
- Landerman, R., George, L. K., Campbell, R. T., and Blazer, D. G. (1989), “Alternative models of the stress buffering hypothesis,” *American Journal of Community Psychology*, 17, 625–642.
- Lange, T., Vansteelandt, S., and Bekaert, M. (2012), “A simple unified approach for estimating natural direct and indirect effects,” *American Journal of Epidemiology*, 176, 190–195.
- Liang, K.-Y. and Zeger, S. L. (1986), “Longitudinal data analysis using generalized linear models,” *Biometrika*, 73, 13–22.
- Lin, S.-H., Young, J., Logan, R., Tchetgen Tchetgen, E. J., and VanderWeele, T. J. (2017), “Parametric mediational g-formula approach to mediation analysis with time-varying exposures, mediators, and confounders,” *Epidemiology (Cambridge, Mass.)*, 28, 266.
- Lindquist, M. A. (2012), “Functional causal mediation analysis with an application to brain connectivity,” *Journal of the American Statistical Association*, 107, 1297–1309.
- Lindquist, M. A. and Sobel, M. E. (2011), “Graphical models, potential outcomes and causal inference: Comment on Ramsey, Spirtes and Glymour,” *NeuroImage*, 57, 334–336.
- Lindström, J. (1999), “Early development and fitness in birds and mammals,” *Trends in Ecology & Evolution*, 14, 343–348.
- MacKinnon, D. (2012), *Introduction to statistical mediation analysis*, Routledge.
- Marmot, M. G., Stansfeld, S., Patel, C., North, F., Head, J., White, I., Brunner, E., Feeney, A., and Smith, G. D. (1991), “Health inequalities among British civil servants: the Whitehall II study,” *The Lancet*, 337, 1387–1393.
- McEwen, B. S. (1998), “Stress, adaptation, and disease: Allostasis and allostatic load,” *Annals of the New York Academy of Sciences*, 840, 33–44.

- (2008), “Central effects of stress hormones in health and disease: Understanding the protective and damaging effects of stress and stress mediators,” *European Journal of Pharmacology*, 583, 174–185.
- Miller, G. E., Chen, E., and Parker, K. J. (2011), “Psychological stress in childhood and susceptibility to the chronic diseases of aging: moving toward a model of behavioral and biological mechanisms.” *Psychological Bulletin*, 137, 959.
- Miller, G. E., Cohen, S., and Ritchey, A. K. (2002), “Chronic psychological stress and the regulation of pro-inflammatory cytokines: a glucocorticoid-resistance model.” *Health psychology*, 21, 531.
- Montagna, S., Tokdar, S. T., Neelon, B., and Dunson, D. B. (2012), “Bayesian latent factor regression for functional and longitudinal data,” *Biometrics*, 68, 1064–1073.
- Neyman, J. (1923), “On the application of probability theory to agricultural experiments. Essay on principles. Section 9 (translated).” *Statistical Science*, 465–480.
- Nguyen, T. Q., Schmid, I., and Stuart, E. A. (2020), “Clarifying causal mediation analysis for the applied researcher: Defining effects based on what we want to learn,” *Psychological Methods*, in press.
- Pearl, J. (2001), “Direct and indirect effects,” in *Proceedings of the Seventeenth Conference on Uncertainty in Artificial Intelligence*, Morgan Kaufmann Publishers Inc., 411–420.
- Ramsay, J. and Silverman, B. (2005), “Functional Data Analysis,” .
- Robins, J. M. and Greenland, S. (1992), “Identifiability and exchangeability for direct and indirect effects,” *Epidemiology*, 143–155.
- Robins, J. M., Hernan, M. A., and Brumback, B. (2000), “Marginal Structural Models and Causal Inference in Epidemiology,” *Epidemiology*, 11.
- Rosenbaum, S., Zeng, S., Campos, F., Gesquiere, L., Altmann, J., Alberts, S., Li, F., and Archie, E. (2020), “Social bonds do not mediate the relationship between early adversity and adult glucocorticoids in wild baboons,” *Proceedings of the National Academy of Sciences*, in press.

- Roth, D. L. and MacKinnon, D. P. (2012), “Mediation analysis with longitudinal data,” *Longitudinal data analysis: A practical guide for researchers in aging, health, and social sciences*, 181–216.
- Rubin, D. B. (1974), “Estimating causal effects of treatments in randomized and nonrandomized studies,” *Journal of Educational Psychology*, 66, 688.
- Silk, J. B. (2007), “The adaptive value of sociality in mammalian groups,” *Philosophical Transactions of the Royal Society B: Biological Sciences*, 362, 539–559.
- Sobel, M. E. (2008), “Identification of causal parameters in randomized studies with mediating variables,” *Journal of Educational and Behavioral Statistics*, 33, 230–251.
- Tchetgen Tchetgen, E. J. and Shpitser, I. (2012), “Semiparametric theory for causal mediation analysis: efficiency bounds, multiple robustness, and sensitivity analysis,” *Annals of Statistics*, 40, 1816.
- Thorsteinsson, E. B. and James, J. E. (1999), “A meta-analysis of the effects of experimental manipulations of social support during laboratory stress,” *Psychology and Health*, 14, 869–886.
- Tung, J., Archie, E. A., Altmann, J., and Alberts, S. C. (2016), “Cumulative early life adversity predicts longevity in wild baboons,” *Nature Communications*, 7, 1–7.
- van der Laan, M. J. and Petersen, M. L. (2008), “Direct effect models,” *The International Journal of Biostatistics*, 4.
- VanderWeele, T. (2015), *Explanation in causal inference: methods for mediation and interaction*, Oxford University Press.
- VanderWeele, T. J. (2011), “Causal mediation analysis with survival data,” *Epidemiology (Cambridge, Mass.)*, 22, 582.
- (2016), “Mediation analysis: a practitioner’s guide,” *Annual Review of Public Health*, 37, 17–32.

- VanderWeele, T. J. and Tchetgen Tchetgen, E. J. (2017), “Mediation analysis with time varying exposures and mediators,” *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 79, 917–938.
- VanderWeele, T. J., Vansteelandt, S., and Robins, J. M. (2014), “Effect decomposition in the presence of an exposure-induced mediator-outcome confounder,” *Epidemiology (Cambridge, Mass.)*, 25, 300.
- Vansteelandt, S., Linder, M., Vandenberghe, S., Steen, J., and Madsen, J. (2019), “Mediation analysis of time-to-event endpoints accounting for repeatedly measured mediators subject to time-varying confounding,” *Statistics in Medicine*, 38, 4828–4840.
- Yao, F., Müller, H.-G., and Wang, J.-L. (2005), “Functional data analysis for sparse longitudinal data,” *Journal of the American Statistical Association*, 100, 577–590.
- Zhao, Y., Luo, X., Lindquist, M., and Caffo, B. (2018), “Functional Mediation Analysis with an Application to Functional Magnetic Resonance Imaging Data,” *arXiv preprint arXiv:1805.06923*.
- Zigler, C. M., Dominici, F., and Wang, Y. (2012), “Estimating causal effects of air quality regulations using principal stratification for spatially correlated multivariate intermediate outcomes,” *Biostatistics*, 13, 289–302.
- Zipple, M. N., Archie, E. A., Tung, J., Altmann, J., and Alberts, S. C. (2019), “Intergenerational effects of early adversity on survival in wild baboons,” *Elife*, 8, e47433.