

# Constraints on Axion-Lepton coupling from Big Bang Nucleosynthesis

Diptimoy Ghosh <sup>a,1</sup>, Divya Sachdeva <sup>a,2</sup>

<sup>a</sup> Department of Physics, Indian Institute of Science Education and Research Pune, Pune 411008, India

## Abstract

In this article, we study the implications of the coupling between Axion-Like-Particles (ALPs) and Leptons to cosmology in particular, the Big Bang Nucleosynthesis (BBN). We show that the BBN, through the constraint on the effective number of relativistic neutrino species, provides the most stringent bound on the ALP-electron interaction strength for the mass of axion between 20 keV and 1 MeV. For other values of the mass, the BBN bound complements the stellar-evolution and laboratory bounds.

---

<sup>1</sup> diptimoy.ghosh@iiserpune.ac.in

<sup>2</sup> divya.sachdeva@students.iiserpune.ac.in

# 1 Introduction

The Standard Model of Particle Physics (SM) has been immensely successful; however, there remains a few unsolved puzzles, one of them being the smallness of the coefficient  $\bar{\theta}$  which is a sum of two apparently independent terms in the theory

$$\bar{\theta} = \theta + \arg \det(Y_u Y_d), \quad (1)$$

where  $\theta$  is the coefficient of the classically marginal<sup>1</sup> and CP odd operator  $-(1/32\pi^2)G_{\mu\nu}^a \tilde{G}^{a\mu\nu}$ , and  $Y_{u,d}$  are the quark Yukawa matrices. Interestingly, the coefficient  $\bar{\theta}$  contributes to Electric Dipole Moments (EDMs) of the leptons [3, 4] as well as the hadrons, e.g., the neutron [5, 6]. vacuum and the Yukawa coupling of Higgs boson to quarks, The strong experimental constraint on the neutron EDM gives an upper bound  $\bar{\theta} \leq 10^{-10}$  [5–7]. The smallness of this parameter is intriguing, and known as the “strong CP problem”.

The most popular solution of this problem, known as the Peccei Quinn (PQ) solution [8–11], predicts the existence of a light pseudo Nambu-Goldstone Boson (pNGB), called the axion, in the particle spectrum. The axion gets its mass from non-perturbative QCD dynamics, and in this case, the axion mass is completely determined by the axion decay constant denoted by  $f_a$ , see [12, 13] and the references therein for more details.

Light pNGBs, such as the QCD axion discussed above, appear in many extensions of the SM [14]. These particles, being pNGBs, share many properties with the QCD axion. However, since their mass, in general, does not arise from the QCD dynamics, the mass and the decay constant are independent parameters. These particles are traditionally called the Axion-Like-Particles (ALPs).

A lot of a theoretical and experimental efforts are underway to search for the ALPs, for example, they are actively being searched at Light-shining through wall experiments [15, 16], particle colliders [17–21], beam dump experiments [22–24], dark matter detection experiments [25–27] as well as various axion helioscopes and haloscopes [21, 28, 29]. There are also many new proposals such as [30–38] which are expected to cut into unprobed regions of parameter space, and hopefully yield a signal. Along with these, ALPs are also constrained by a variety of astrophysical and cosmological observations. For example, axions can be produced copiously inside Sun, Supernovae, Pulsars, White dwarfs etc., and affect the cooling time of these objects as well as the photon polarisation. This gives interesting constraints on axion properties [39–42] (note, however, that the astrophysical bounds often suffer from large uncertainties [43, 44]). Similarly, the Cosmic Microwave Background (CMB) power spectrum, and the Big Bang Nucleosynthesis (BBN) can also be sensitive to the properties of the ALPs, and are used to constrain significant part of the ALP parameter space [41, 45–51].

---

<sup>1</sup>The leading contribution to the beta function of  $\bar{\theta}$  in the SM appears at the 7-loop order [1, 2].

Most of the experiments mentioned above rely on the axion coupling to photons. The ALP-Lepton coupling has been relatively less studied, see however, [41, 50, 51]. In this work, we perform a detailed analysis of the cosmological implications of ALP-Lepton interaction, in particular, its impact on the BBN. The BBN occurred when the temperature of the Universe was between  $10 \text{ keV} < T < 1 \text{ MeV}$ , and therefore axions with mass  $< 1 \text{ MeV}$  can affect the physics in this era. The axions can be produced via

$$l^\pm \gamma \rightarrow l^\pm a \text{ and } l^+ l^- \rightarrow \gamma a$$

processes and can lead to non-negligible axion abundance during BBN and contribute to  $\Delta N_{\text{eff}}$ . In addition, the ALP-electron coupling  $c_e/f_a$  can also induce significantly large coupling to photons at one loop,  $g_{a\gamma\gamma} \sim 10^{-3} c_e/f_a$ , for  $m_a \gtrsim m_e$ . This extra interaction would allow ALPs to stay in equilibrium for longer providing stronger bounds.

The paper is organised as follows. In Sec. 2 we introduce the general effective Lagrangian describing the ALPs and their couplings to the photon and the SM fermions, and review the present constraints on ALP-photon and ALP-lepton couplings. Sec. 3 describes how the ALP-Lepton coupling can affect the thermal history of the universe, and impact the effective number of neutrinos,  $N_{\text{eff}}$ , measured during BBN. In Sec. 4, we numerically solve the Boltzmann equations to find the exact constraints on ALPs-Lepton couplings as a function of the ALP mass. Finally, in Sec. 5, we summarise our main results and conclude.

## 2 Axion properties

We use the following Lagrangian density describing the interactions of axions or ALPs to SM particles

$$\mathcal{L} = \frac{1}{2}(\partial_\mu a)^2 + \frac{\alpha_s}{8\pi} \frac{a}{f_a} G_{\mu\nu}^A \tilde{G}^{A\mu\nu} + \frac{g_{a\gamma\gamma}}{4} a F_{\mu\nu} \tilde{F}^{\mu\nu} + c_\psi \frac{\partial_\mu a}{2f_a} \bar{\psi} \gamma^\mu \gamma_5 \psi \quad (2)$$

where  $f_a$  is the effective axion decay constant and  $\alpha_s$  is the QCD fine structure constant. The axion-photon-photon coupling,  $g_{a\gamma\gamma}$ , is model dependent. The coupling  $g_{a\gamma\gamma}$  for the QCD axion can be written as

$$g_{a\gamma\gamma} = \frac{\alpha_{\text{em}}}{2\pi f_a} \left( \frac{E}{N} - 1.92 \right) \quad (3)$$

where the quantities  $E$  and  $N$  are the mixed anomaly coefficients of the PQ symmetry with Electro-Magnetism (EM) and QCD respectively, and the number 1.92 is the model-independent contribution from QCD [52]. The mass of the QCD axion is also determined by the axion decay constant in the following way [52]

$$m_a = 5.7 \mu\text{eV} \left( \frac{10^{12} \text{ GeV}}{f_a} \right). \quad (4)$$

This gives,

$$g_{a\gamma\gamma} = \left(\frac{E}{N} - 1.92\right) \left(\frac{m_a}{0.5 \text{ eV}}\right) (10^{-10} \text{ GeV}^{-1}) \quad (5)$$

$m_a$ (MeV)	$g_{a\gamma\gamma}$ (GeV $^{-1}$ )	Description
$\lesssim 10^{-7}$	$\lesssim 10^{-7}$	Light-shining-through-walls experiment [15]
$\lesssim 10^{-5}$	$\lesssim 10^{-10}$	Tokyo Axion Helioscope (SUMICO) and the CERN Axion Solar Telescope (CAST) [28, 53]
$\lesssim 10^{-2}$	$\lesssim 10^{-10}$	Horizontal Branch stars [39]
$\lesssim 10^2$	$\geq 10^{-2} \ \& \ \lesssim 10^{-9}$	Supernova Type II SN1987A [40]
$\sim 1 - 10^6$	$\lesssim 10^{-3}$	Mono-photon and tri-photon searches at LEP, CDF and LHC [20]

Table 1: *Model independent constraints on  $g_{a\gamma\gamma}$  for different values of  $m_a$ . Note that the ALP-photon coupling  $g_{a\gamma\gamma}$  can also give rise to interesting effects on the CMB properties which, in turn, can be used to put serious constraints on  $g_{a\gamma\gamma}$ , see for example, refs. [41, 47–51, 54].*

The current bounds on  $g_{a\gamma\gamma}$  and its variation with mass are shown in table 1. For example,  $g_{a\gamma\gamma} \lesssim 10^{-10} \text{ GeV}^{-1}$  for  $m_a \lesssim 10 \text{ keV}$  which translates to  $m_a \lesssim 0.5 \text{ eV}$  (and consequently,  $f_a \gtrsim 10^7 \text{ GeV}$ ) for QCD axion with  $(E/N - 1.92) \sim 1$ . The decay width of the ALP decaying to photons can be written as,

$$\Gamma_{a \rightarrow \gamma\gamma} = \left(\frac{g_{a\gamma\gamma}}{4}\right)^2 \frac{m_a^3}{4\pi} = \left(\frac{g_{a\gamma\gamma}}{10^{-10} \text{ GeV}^{-1}}\right)^2 \left(\frac{m_a}{350 \text{ eV}}\right)^3 \left(\frac{1}{3.3 \times 10^{17} \text{ sec.}}\right) \quad (6)$$

$$= \left(\frac{E}{N} - 1.92\right)^2 \left(\frac{m_a}{25.5 \text{ eV}}\right)^5 \left(\frac{1}{3.3 \times 10^{17} \text{ sec.}}\right) \text{ [for QCD axion]} \quad (7)$$

Thus, for the QCD axion, one necessary condition for it to have a lifetime more than the age of the universe is that the axion mass has to be less than about 25 eV (for  $(E/N - 1.92) \sim 1$ ). For a general ALP, its lifetime can be larger than the age of the universe even when the mass is much larger than 25 eV, and depends on  $g_{a\gamma\gamma}$  (and also couplings to matter) which is not determined by  $m_a$ .

Similar to  $g_{a\gamma\gamma}$ , the axion-fermion-fermion couplings,  $c_\psi$ , is also model dependent. In this work, we consider couplings only to the charged leptons,  $e$ ,  $\mu$  and  $\tau$  (couplings to the neutrinos are extremely suppressed due to smallness of the neutrino masses). In models such as the Dine-Fischler-Srednicki-Zhitnitsky (DFSZ) model [55, 56], axions have tree-level interactions with leptons and  $c_l$  can be as high as 1/3. In the Kim-Shifman-Vainshtein-Zakharov (KSVZ) model [57, 58], on the other hand, such direct interaction with leptons is absent. Even in models where there is no tree level coupling to leptons, it can be generated by the axion-photon-photon coupling at one loop. As the axion-photon-photon coupling already has one power of  $\alpha_{\text{em}}$ ,  $c_l$ , generated in this way, is  $\mathcal{O}(\alpha_{\text{em}}^2)$  suppressed. However, for QCD axions, large logarithms either of the size

$\text{Ln}(\Lambda_{\text{QCD}}/m_e) \approx 5$  or  $\text{Ln}(f_a/m_e) \approx 30$  may kill part of this suppression. In Fig. 1, we compile all the existing bounds on  $c_l/f_a$  for different values of  $m_a$ .




Figure 1: Existing model independent constraints on  $c_e/f_a$  (left) and  $c_\mu/f_a$  (right) for different values for  $m_a$  from observations of Red Giants [39, 59], SN1987A [41, 42, 59, 60], dark photon searches at Babar [19], Beam dump experiment at SLAC [24]. Note that collider bounds from LEP [21] assumes  $g_{a\gamma\gamma} \sim 10^{-3} \text{ GeV}^{-1}$ , so they are not model independent. Bounds from EDELWEISS [25] and LUX [26] are one order of magnitude stronger than shown here, if axion are assumed as cold dark matter. Similarly, BBN studies in [41] constrains  $(c_e/f_a, c_\mu/f_a, c_\tau/f_a) \sim (10^{-6}, 10^{-7}, 10^{-6})$  assuming relativistic axions during CMB decoupling i.e, with  $m_a < 10^{-1} \text{ eV}$ . The CAST experiment [61] constraints the product  $g_{a\gamma\gamma} c_e/f_a < 1.6 \times 10^{-19} \text{ GeV}^{-2}$  at 95% CL for  $m_a \lesssim 10^{-2} \text{ eV}$ . The solid black diagonal line corresponds to the DFSZ QCD axion model.

As we wish to extract constraints on the couplings  $c_l/f_a$  in a model independent way, we assume that the direct coupling to photons is zero (the bounds on  $c_l/f_a$  will only get stronger for non-zero  $g_{a\gamma\gamma}$ , thus our bounds are conservative). Note, however, that the ALP-Lepton coupling does generate the axion photon coupling,  $g_{a\gamma\gamma}$ , at one loop:

$$g_{a\gamma\gamma}^{\text{loop}} = \frac{\alpha_{\text{em}}}{4\pi} \frac{c_l}{f_a} 4 B_1(x_l) \quad (8)$$

where,  $B_1(x_l) = 1 - x_l f(x_l)^2$ ;  $x_l \equiv \frac{4m_l^2}{m_a^2}$

$$f(x) = \begin{cases} \arcsin \frac{1}{\sqrt{x}}; & x \geq 1, \\ \frac{\pi}{2} + \frac{i}{2} \ln \frac{1 + \sqrt{1-x}}{1 - \sqrt{1-x}}; & x < 1. \end{cases}$$

The fermion loop function  $B_1(x) \sim -\frac{m_a^2}{12m_l^2}$  for  $m_l \gg m_a$  and  $|B_1(x)| \sim 1$  for  $m_a \gg m_l$ . We plot  $g_{a\gamma\gamma}^{\text{loop}}$  as a function of the ALP mass for different leptons in Fig. 2(left).

It is evident that the value of  $g_{a\gamma\gamma}$  is non-negligible especially for  $m_a \gg m_l$  with




Figure 2: (left) The variation of ratio  $\left| \frac{g_{a\gamma\gamma}}{c_l/f_a} \right|$  with  $m_a$ . (right) Lifetime of axion decaying to photon as well as leptons as a function of mass for  $c_l/f_a = 10^{-4} \text{ GeV}^{-1}$ .

$g_{a\gamma\gamma} \stackrel{m_a \gg m_l}{\sim} \frac{\alpha_{\text{em}}}{\pi} \frac{c_l}{f_a}$ . To emphasize its significance, we plot lifetime of ALPs as function of their mass keeping  $c_l/f_a = 10^{-4} \text{ GeV}^{-1}$  in Fig. 2(right). To calculate the lifetime, we use eqn. 6 and the following expression for decay rate of the ALP to leptons,

$$\Gamma_{a \rightarrow l^+ l^-} = \frac{m_l^2}{8\pi} \left( \frac{c_l}{f_a} \right)^2 m_a \sqrt{1 - \frac{4m_l^2}{m_a^2}}. \quad (9)$$

### 3 Axions production in the early universe

Considering only the axion-lepton-lepton coupling,  $c_l$ , the axions can be generated in the early universe by the following processes

$$l^\pm(p_1) \gamma(p_2) \rightarrow l^\pm(p_3) a(p_4) \quad (10)$$

$$l^-(p_1) l^+(p_2) \rightarrow \gamma(p_3) a(p_4). \quad (11)$$

The corresponding Feynman diagrams are shown in Fig. 3. Based on the dimensional analysis, one expects  $\langle \sigma v \rangle$  to scale as  $\frac{c_l^2 m_l^2}{f_a^2 T^2}$  in the relativistic limit (for  $T \gg m_{l,a}$ ).

This implies that the ratio of interaction rate to Hubble expansion takes the form  $\frac{\Gamma}{H} \sim \frac{n_{l,\gamma} \langle \sigma v \rangle M_{\text{Pl}}}{T^2} \propto \frac{1}{T}$ . As Universe cools to temperatures below the lepton mass,  $\langle \sigma v \rangle$  goes as  $\frac{c_l^2 m_l^2}{f_a^2} \frac{1}{g(m_a^2, m_l^2)}$  where  $g(m_a^2, m_l^2)$  is a function having mass dimension 2.

Consequently,  $\Gamma/H$  would go as  $\frac{n_\gamma \langle \sigma v \rangle M_{\text{Pl}}}{T^2} \propto T$ . However, the detailed analysis discussed below suggests that the interaction rate in the non-relativistic limit drop more quickly than expected from the naive power counting. For that, we write below the




Figure 3: *Feynman diagrams of  $l^\pm(p_1)\gamma(p_2) \rightarrow l^\pm(p_3)a(p_4)$  and  $l^+(p_1)l^-(p_2) \rightarrow \gamma(p_3)a(p_4)$*

Boltzmann equation describing the evolution of the axion number density

$$\begin{aligned} \frac{dn_a}{dt} + 3Hn_a &= \left(2\Gamma^{l^\pm\gamma\rightarrow al^\pm} + \Gamma^{l^-l^+\rightarrow a\gamma}\right)(n_a - n_a^{\text{eq}}) \\ \frac{dY_a}{dx} &= -\frac{xs}{H(m)} \left(2\langle\sigma v\rangle^{l^\pm\gamma\rightarrow al^\pm} Y_l^{\text{eq}} \frac{Y_l^{\text{eq}}}{Y_a^{\text{eq}}} + \langle\sigma v\rangle^{l^-l^+\rightarrow a\gamma} Y_\gamma^{\text{eq}} \frac{Y_l^{\text{eq}}}{Y_a^{\text{eq}}}\right) (Y_a - Y_a^{\text{eq}}) \end{aligned} \quad (12)$$

where,  $Y_a = \frac{n_a}{s}$ , and  $x = \frac{m_l}{T}$ . The matrix element for process in eqn.10 gets two contributions associated to s- and u-channel exchange. The corresponding expressions are given as:

$$\begin{aligned} i\mathcal{M}^s &= \frac{ie c_l}{2f_a} \bar{u}(p_3) \gamma^\mu \gamma^5 \frac{i(\not{p}_1 - \not{p}_2 + m_l)}{s - m_l^2} \gamma^\nu u(p_1) p_{4\mu} \epsilon_\nu(p_2), \\ i\mathcal{M}^u &= \frac{ie c_l}{2f_a} \bar{u}(p_3) \gamma^\nu \frac{i(\not{p}_1 - \not{p}_4 + m_l)}{u - m_l^2} \gamma^\mu \gamma^5 u(p_1) p_{4\mu} \epsilon_\nu(p_2), \end{aligned}$$

Similar expression for the process in eqn.11 can be derived using crossing symmetry property. We see that the transformation of the Mandelstam variables from  $(s, t, u) \rightarrow (t, s, u)$  gives us the matrix element for pair-annihilation process.

Next, we calculate the cross-section in the limit  $m_a \rightarrow 0$ , useful in relativistic limit ( $T \gg m_{l,a}$ )

$$\begin{aligned} \sigma^{l^\pm\gamma\rightarrow l^\pm a} &= \frac{\alpha c_l^2 m_l^2}{8f_a^2 s^2} \frac{(-m_l^4 + 4m_l^2 s + s^2 \log\left(\frac{s^2}{m_l^4}\right) - 3s^2)}{(s - m_l^2)} \\ &\stackrel{s \gg m_l^2}{=} \frac{\alpha c_l^2 m_l^2}{8f_a^2 s} (\log(s^2/m_l^4) - 3) \end{aligned} \quad (13)$$

$$\begin{aligned}
\sigma^{l^+l^- \rightarrow \gamma a} &= \frac{\alpha c_l^2 m_l^2 \tanh^{-1} \left( \sqrt{1 - \frac{4m_l^2}{s}} \right)}{4f_a^2 (s - 4m_l^2)} \\
&\stackrel{s \gg m_l^2}{=} \frac{\alpha c_l^2 m_l^2}{4f_a^2 s} \tanh^{-1} \left( \sqrt{1 - \frac{4m_l^2}{s}} \right)
\end{aligned} \tag{14}$$

It should be noted that the logarithmic dependence leads to divergent cross-section in the  $m_l \rightarrow 0$  limit corroborating the t-channel singularity.

From eqn.12, 13 and 14, the reaction rate per axion particle in the limit  $T \gg m_{a,l}$  can be written as

$$\Gamma = \frac{n_{l/\gamma}^{\text{eq}} n_l^{\text{eq}}}{n_a^{\text{eq}}} \langle \sigma v \rangle \stackrel{\text{eqn.10,11}}{\sim} T^3 \left( \frac{\alpha_{\text{em}}}{16} \frac{c_l^2 m_l^2}{f_a^2} \frac{1}{T^2} \text{Log} \left[ \frac{4T^2}{m_l^2} \right] \right) = \frac{\alpha_{\text{em}} T}{16} \frac{c_l^2 m_l^2}{f_a^2} \text{Log} \left[ \frac{4T^2}{m_l^2} \right].$$

Since, the Hubble parameter,  $H \sim T^2/M_{\text{pl}}$ ,

$$\frac{\Gamma}{H} \propto \frac{1}{T} \text{Log} \left[ \frac{4T^2}{m_l^2} \right] \tag{15}$$

Thus, as the universe cools,  $\Gamma/H$  keeps growing allowing the possibility to bring the axions in thermal equilibrium with the plasma. However, when the universe cools further so that  $T \ll m_l$ , one expects

$$\Gamma \stackrel{\text{eqn.10}}{=} \frac{n_{\gamma}^{\text{eq}} n_l^{\text{eq}}}{n_a^{\text{eq}}} \langle \sigma v \rangle \sim T^3 \frac{n_l^{\text{eq}}}{n_a^{\text{eq}}} \left[ \frac{\alpha_{\text{em}}}{2m_l^2} \frac{c_l^2 m_l^2}{f_a^2} \left( a \frac{T}{m_l} + b \frac{T^2}{m_l^2} \right) \right] \tag{16}$$

$$\stackrel{\text{eqn.11}}{=} \frac{n_l^{\text{eq}} n_l^{\text{eq}}}{n_a^{\text{eq}}} \langle \sigma v \rangle \sim e^{-m_l/T} (m_l T)^{3/2} \frac{n_l^{\text{eq}}}{n_a^{\text{eq}}} \left[ \frac{\alpha_{\text{em}}}{2m_l^2} \frac{c_l^2 m_l^2}{f_a^2} \left( \tilde{a} + \tilde{b} \frac{T}{m_l} \right) \right]. \tag{17}$$

where  $a, \tilde{a}, b$  and  $\tilde{b}$  are functions of the axion and the lepton masses and are found numerically to be  $\mathcal{O}(1)$  numbers. It should be noted that the interaction in equation 10 is dominant at lower temperature as the other interaction has an extra Boltzmann suppression factor corresponding to the extra lepton in the initial state. Thus, the ratio  $\Gamma/H$  scales (other than the Boltzmann suppression) as  $T^n$  with  $n = 2$  in the non-relativistic regime i.e it falls quicker than expected within dimensional analysis. We also conclude that this ratio attains its maximum value around  $T \sim m_l$ . This feature can be seen in figure 4. Note that this feature arises only for axion-lepton-lepton couplings because it is effectively a marginal coupling. It can be seen from the figure that the axion produced in this way stays in thermal equilibrium (i.e.,  $\Gamma > H$ ) for some duration. Depending on the coupling and mass, it goes out of the equilibrium at lower temperatures. For ALPs interacting with only the muons or the tau-leptons, they go out of equilibrium before the neutrino decoupling; however, they can still contribute significantly to the total energy budget of the Universe. In fact, as we show in the next section, the energy density stored in the ALPs could be significant enough to affect the BBN. The non-negligible yield and hence, the energy density of axions can increase the Hubble parameter during the BBN. A larger Hubble parameter during the BBN






Figure 4: *Interaction rate and Hubble rate as function of temperature for  $m_a = 10$  keV and for various coupling strength.*

modifies the neutron-to-proton ratio, which in turn changes the abundance of Helium-4 and Deuterium. This effect can be captured by defining a quantity called  $\Delta N_{\text{eff}}^{\text{BBN}}$  in the following way. The total energy density during the BBN can be written as

$$\rho = \frac{\pi^2}{30} \left\{ g_\gamma + \frac{7}{8} (3g_\nu + 3g_{\bar{\nu}} + g_{e^+} + g_{e^-}) \right\} T_\gamma^4 + \rho_a \quad (18)$$

$$= \frac{\pi^2}{30} \frac{43}{4} T_\gamma^4 + \rho_a \quad (19)$$

$$= \frac{43}{8} \rho_\gamma + \rho_a \quad (20)$$

The quantity  $\Delta N_{\text{eff}}^{\text{BBN}}$  can now be defined by

$$\rho = \frac{\pi^2}{30} \left( \frac{43}{4} + \frac{7}{8} 2 \Delta N_{\text{eff}}^{\text{BBN}} \right) T_\gamma^4 \quad (21)$$

$$\Rightarrow \Delta N_{\text{eff}}^{\text{BBN}} = \frac{8}{7} \frac{\rho_a}{\rho_\gamma} \quad (22)$$

The latest measurement and analysis of Helium and Deuterium abundance constrain this parameter to  $N_{\text{eff}}^{\text{BBN}} = 2.878 \pm 0.278$  at 68.3% CL [62]. Using  $N_{\text{eff}} = 3.046$  for the SM [63, 64], this gives  $\Delta N_{\text{eff}}^{\text{BBN}} < 0.39$  at  $2\sigma$ . Since, some of the earlier works [65–69] had a slightly weaker upper bound, in order to be very conservative we take  $\Delta N_{\text{eff}}^{\text{BBN}} < 0.5$  in our work.

The maximum value of  $\Delta N_{\text{eff}}^{\text{BBN}}$  in the context of ALPs is obtained if the ALPs stay in thermal equilibrium during the BBN and are relativistic (i.e.,  $m_a \ll 1$  MeV). In this case,  $\Delta N_{\text{eff}}^{\text{BBN}} = 8/7 \times 1/2 = 0.57$  since the axion, being a pseudo-scalar, contributes only one extra degree of freedom. For larger axion mass,  $\Delta N_{\text{eff}}^{\text{BBN}}$  reduces from 0.57 owing to the Boltzmann suppression, as can be seen from Fig. 5 (left). In the right panel of Fig. 5, we also show  $\Delta N_{\text{eff}}^{\text{BBN}}$  as a function of the temperature for three different values of  $m_a$ . Note that Fig. 5 shows  $\Delta N_{\text{eff}}^{\text{BBN}}$  for particle at thermal equilibrium. In order to fully take into account various processes when ALPs are not in thermal equilibrium, we need to solve a complete set of Boltzmann equations for the distribution functions of ALPs as well as neutrinos. This will be discussed in the next section.




Figure 5:  $\Delta N_{\text{eff}}$  as function of  $m_a$  (left) and  $T$  (right) assuming that the ALPs are in thermal equilibrium. In the left panel, only the ALP-electron coupling has been considered.

We would like to stress here that the quantity  $\Delta N_{\text{eff}}$  is also measured from the CMB observations. However, in general,  $\Delta N_{\text{eff}}^{\text{CMB}}$  is not the same as  $\Delta N_{\text{eff}}^{\text{BBN}}$ ; depending on the details,  $\Delta N_{\text{eff}}^{\text{CMB}}$  can be more or less than  $\Delta N_{\text{eff}}^{\text{BBN}}$  [70]. In order for the axions to contribute to  $\Delta N_{\text{eff}}^{\text{CMB}}$ , they have to be relativistic at the time of CMB decoupling. This means that for the axion mass  $m_a \gtrsim \text{eV}$ , they would not contribute significantly to  $\Delta N_{\text{eff}}^{\text{CMB}}$ . This is why we have not used the stronger constraint on  $\Delta N_{\text{eff}}$  obtained from the CMB [62, 71]. The decay of the ALPs after BBN and close to the CMB decoupling can, however, lead to observable effects on radiation density and also on other observables like CMB spectral distortions which can in principle give rise to interesting bounds [72, 73].

## 4 Constraint on $c_l/f_a$

In the homogeneous and isotropic Universe with the FLRW metric, the relevant distribution functions fulfill the following Boltzmann equations:

$$\frac{\partial f_i(|\vec{p}|, t)}{\partial t} - H|\vec{p}| \frac{\partial f_i(|\vec{p}|, t)}{\partial |\vec{p}|} = C[f_i(|\vec{p}|, t)]$$

To solve these first-order partial differential equations, we adopt characteristics curves method [74] and introduce two dimensionless parameters  $z = m_l R(t)$  and  $Q = |\vec{p}| R(t)$ , where  $R(t)$  is scale factor in the FLRW metric. So, the Boltzmann equations is recast into the following form:

$$Hz \frac{\partial f_i(Q, z)}{\partial z} = C[f_i(Q, z)], \quad (23)$$

where the subscript  $i$  stands for  $e^\pm$ ,  $\nu$  and  $a$ . While photons and electrons initially follow the distributions in thermal equilibrium with zero chemical potentials, the initial abundance of axion is assumed to be negligible. The collision term for the ALPs can be written as:

$$C[f_{\text{ALPs}}] = \frac{1}{2E_a} \int dp_l dp'_l dp_\gamma (2\pi)^4 \delta^4(\sum p) [f_\gamma f_l (1 + f_a)(1 - f'_l) - f'_l f_a (1 - f_l)(1 + f_\gamma)] |\mathcal{M}|^2$$

where  $dp_i = g_i d^3 p_i / [(2\pi)^3 2E_i]$  with  $g_i$  being the internal degrees of freedom. The collision terms for the SM species are taken from references [75–77]. We use numerical techniques discussed in [75, 78] to solve multi-dimensional collision integration.

Along with the Boltzmann equation, the Friedmann equations,  $H^2 = \frac{8\pi G \rho^{\text{tot.}}}{3}$  and  $\frac{d\rho}{dt} = -3H(\rho + P)$  in terms of  $z$  and  $Q$  are also relevant to calculate the evolution of photon temperature:

$$z \frac{d\rho}{dz} = -3(\rho + P) \quad (24)$$

$$z \frac{dT_\gamma}{dz} \frac{d\rho_\gamma}{dT_\gamma} = -3(\rho + P) - z \frac{d\rho_e^\pm}{dz} - z \frac{d\rho_\nu}{dz} - z \frac{d\rho_a}{dz} \quad (25)$$

After solving the Boltzmann equation, energy density during BBN and hence  $\Delta N_{\text{eff}}$  are calculated using Eq. (22) at the time of neutrino decoupling. In Fig. 6(left) we show the variation of  $\Delta N_{\text{eff}}$  with respect to the coupling  $c_l/f_a$  for axions with mass  $m_a < 10$  keV (we have chosen 10 keV just as a benchmark. Also, for this mass range, the axions are relativistic and  $\Delta N_{\text{eff}}$  is independent of the mass). Fig. 6(right), on the other hand, shows the contours for  $\Delta N_{\text{eff}} = 0.5$  and  $0.4$  on the  $m_a - c_l/f_a$  plane. Note that we have restricted our analysis to  $m_a \leq 1$  MeV. However, in principle, the analysis can be extended to heavier axion masses, but the constraints are expected to be much weaker.

Let us now try to understand qualitatively the results in Fig. 6. First, note that the ALP-Lepton coupling is proportional to the ALP mass. Thus, for a given  $c_l/f_a$ , axions

will have stronger coupling to heavier leptons. This means that, in order to get the same axion yield, a relatively smaller value of  $c_\mu/f_a$  would be required than  $c_e/f_a$ . Consequently, the constraint on  $c_\mu/f_a$  will be stronger than that on  $c_e/f_a$  (for this argument, we have assumed the mass of the axion to be much less than the electron and the muon mass), which can be seen from the figure.

For the heavier leptons, however, another effect comes into play: lower number density of leptons for heavier leptons (due to Boltzmann suppression) leads to lower rate for the process  $l^-/\gamma + a \rightarrow l^-l^+$  reducing the yield. In order to maintain significant axion yield, this requires larger  $c_l/f_a$ . In this way, the competition between the number density of leptons and lepton mass dependent coupling strength plays a crucial role in obtaining the constraints given in Fig. 6. In the right panel of Fig. 6, the increase of required coupling strength with the axion mass in order to maintain a fixed  $\Delta N_{\text{eff}}$  is also easily understood in terms of the Boltzmann and phase space suppression of the process  $l^-l^+ \rightarrow l^-/\gamma + a$ .




Figure 6: (left)  $\Delta N_{\text{eff}}$  as a function of  $c_l/f_a$  for  $m_a < 10$  keV. (right) Contour satisfying  $\Delta N_{\text{eff}} = 0.5$  on  $m_a$  vs  $c_l/f$  plane.

As discussed above, the coupling of the ALPs with tau-lepton cannot be constrained by BBN as the contribution to  $\Delta N_{\text{eff}}$  is very small (Fig. 6(left)). In this case, the axions go out of equilibrium much before BBN (as can be seen from the lower panel of Fig. 4) and thus, their yield remains small. This result is consistent with the observation that, for a particle that decouples from the primordial thermal plasma before the QCD transition, one always gets  $\Delta N_{\text{eff}} \lesssim 0.3$  [41].

Finally, in Fig. 7 we compare our bounds with the existing constraints discussed in section. 2. It can be seen that, for the ALP-electron coupling, our bound is the strongest for axion mass between 20 keV and 1 MeV. We should also mention in passing that, indirect bound on the ALPs-lepton coupling can be obtained using Eq. (8) and the bounds on ALPs-photon coupling given in table 1 (shown as dashed and dotted black

lines in Fig. 7). However, unlike the BBN bound discussed in this work, this indirect bound is model dependent and relies on the exact value of the direct  $g_{a\gamma\gamma}$  coupling.




Figure 7: Constraints on  $c_e/f_a$  (upper panel) and  $c_\mu/f_a$  (lower panel) obtained in the present work along with the existing model independent constraints. We also show constraints derived using bounds on axion-photon couplings discussed in Tab. 1.

## 5 Summary and Conclusion

In this article, we have performed a detailed study of the constraints on ALP-lepton coupling,  $c_l/f_a$ , from BBN. In the presence of non-zero  $c_l/f_a$ , the ALPs can be produced in the early universe by the processes  $l^\pm\gamma \rightarrow l^\pm a$  and  $l^+l^- \rightarrow \gamma a$  (Fig. 3). When the temperature of the thermal plasma  $T \gg m_{a,l}$ , the quantity  $\Gamma/H$  keeps increasing as

the universe cools allowing the possibility for the axions to come into equilibrium with the rest of the plasma. However, as the universe cools further and the temperature goes below the mass of the lepton,  $\Gamma/H$  starts decreasing with the decreasing temperature. As a result, the ALPs, depending on their coupling strengths, first come into equilibrium, then stay in equilibrium for a while, and eventually goes out of the equilibrium (see Fig. 4). If the axions are relativistic and also in equilibrium during the BBN, their contribution to  $\Delta N_{\text{eff}}^{\text{BBN}}$  turns out to be as high as 0.57 which is clearly ruled out by observations. With the increasing mass of the axion, their contribution to  $\Delta N_{\text{eff}}^{\text{BBN}}$  keeps decreasing due to the Boltzmann suppression (shown in Fig. 5). However, they can still contribute to  $\Delta N_{\text{eff}}^{\text{BBN}}$  significantly which, in turn, provides interesting constraints on the ALP-Lepton couplings.

When ALPs are not in equilibrium with the thermal plasma, the full Boltzmann equations have to be solved in order to understand their effect on BBN quantitatively. We did this exercise in section 4. Based on our analysis, in Fig. 6, we showed contours for  $\Delta N_{\text{eff}}^{\text{BBN}} = 0.5$  on the  $c_{e,\mu}/f_a - m_a$  plane. We also discussed qualitatively why the constraints on  $c_\mu/f_a$  is stronger than the bound on  $c_e/f_a$ , and why the bound on  $c_\tau/f_a$  is extremely poor. Finally, in Fig. 7 we compare our bound (using a very conservative observational upper limit  $\Delta N_{\text{eff}}^{\text{BBN}} \leq 0.5$ ) with the other existing bounds and show that our bound is the most stringent one for the ALP-electron interaction strength for the mass of axion between 20 keV and 1 MeV.

## Acknowledgement

The authors acknowledge support through the Ramanujan Fellowship of the Department of Science and Technology, Government of India.

## References

- [1] J. R. Ellis and M. K. Gaillard, *Strong and Weak CP Violation*, *Nucl. Phys. B* **150** (1979) 141–162.
- [2] I. Khriplovich and A. Vainshtein, *Infinite renormalization of Theta term and Jarlskog invariant for CP violation*, *Nucl. Phys. B* **414** (1994) 27–32, [[hep-ph/9308334](#)].
- [3] K. Choi and J.-y. Hong, *Electron electric dipole moment and Theta (QCD)*, *Phys. Lett. B* **259** (1991) 340–344.
- [4] D. Ghosh and R. Sato, *Lepton Electric Dipole Moment and Strong CP Violation*, *Phys. Lett. B* **777** (2018) 335–339, [[1709.05866](#)].

- [5] R. Crewther, P. Di Vecchia, G. Veneziano and E. Witten, *Chiral Estimate of the Electric Dipole Moment of the Neutron in Quantum Chromodynamics*, *Phys. Lett. B* **88** (1979) 123. [Erratum: Phys.Lett.B 91, 487 (1980)].
- [6] V. Baluni, *CP Violating Effects in QCD*, *Phys. Rev. D* **19** (1979) 2227–2230.
- [7] J. M. Pendlebury et al., *Revised experimental upper limit on the electric dipole moment of the neutron*, *Phys. Rev. D* **92** (2015) 092003, [[1509.04411](#)].
- [8] R. D. Peccei and H. R. Quinn, *CP conservation in the presence of pseudoparticles*, *Phys. Rev. Lett.* **38** (Jun, 1977) 1440–1443.
- [9] R. D. Peccei and H. R. Quinn, *Constraints imposed by CP conservation in the presence of pseudoparticles*, *Phys. Rev. D* **16** (Sep, 1977) 1791–1797.
- [10] S. Weinberg, *A new light boson?*, *Phys. Rev. Lett.* **40** (Jan, 1978) 223–226.
- [11] F. Wilczek, *Problem of strong  $p$  and  $t$  invariance in the presence of instantons*, *Phys. Rev. Lett.* **40** (Jan, 1978) 279–282.
- [12] R. Peccei, *The Strong CP problem and axions*, *Lect. Notes Phys.* **741** (2008) 3–17, [[hep-ph/0607268](#)].
- [13] D. J. E. Marsh, *Axion Cosmology*, *Phys. Rept.* **643** (2016) 1–79, [[1510.07633](#)].
- [14] A. Ringwald, *Axions and Axion-Like Particles*, in *49th Rencontres de Moriond on Electroweak Interactions and Unified Theories*, pp. 223–230, 2014. [1407.0546](#).
- [15] J. Redondo and A. Ringwald, *Light shining through walls*, *Contemp. Phys.* **52** (2011) 211–236, [[1011.3741](#)].
- [16] K. E. et al, *New alps results on hidden-sector lightweights*, *Physics Letters B* **689** (2010) 149 – 155.
- [17] P. del Amo Sanchez, J. P. Lees, V. Poireau, E. Prencipe, V. Tisserand, J. Garra Tico et al., *Search for production of invisible final states in single-photon decays of  $\Upsilon(1s)$* , *Physical Review Letters* **107** (Jul, 2011) .
- [18] K. Mimasu and V. Sanz, *Alps at colliders*, 2014.
- [19] J. Lees, V. Poireau, V. Tisserand, E. Grauges, A. Palano, G. Eigen et al., *Search for a dark photon in  $e^+e^-$  collisions at babar*, *Physical Review Letters* **113** (Nov, 2014) .
- [20] J. Jaeckel and M. Spannowsky, *Probing mev to 90 gev axion-like particles with lep and lhc*, *Physics Letters B* **753** (Feb, 2016) 482–487.
- [21] M. Bauer, M. Neubert and A. Thamm, *Collider Probes of Axion-Like Particles*, *JHEP* **12** (2017) 044, [[1708.00443](#)].

- [22] E. M. Riordan, M. W. Krasny, K. Lang, P. de Barbaro, A. Bodek, S. Dasu et al., *Search for short-lived axions in an electron-beam-dump experiment*, *Phys. Rev. Lett.* **59** (Aug, 1987) 755–758.
- [23] J. D. Bjorken, S. Ecklund, W. R. Nelson, A. Abashian, C. Church, B. Lu et al., *Search for neutral metastable penetrating particles produced in the slac beam dump*, *Phys. Rev. D* **38** (Dec, 1988) 3375–3386.
- [24] R. Essig, R. Harnik, J. Kaplan and N. Toro, *Discovering new light states at neutrino experiments*, *Physical Review D* **82** (Dec, 2010) .
- [25] E. Armengaud, Q. Arnaud, C. Augier, A. Benoit, A. Benoit, L. Bergé et al., *Axion searches with the edelweiss-ii experiment*, *Journal of Cosmology and Astroparticle Physics* **2013** (Nov, 2013) 067–067.
- [26] D. Akerib, S. Alsum, C. Aquino, H. Araújo, X. Bai, A. Bailey et al., *First searches for axions and axionlike particles with the lux experiment*, *Physical Review Letters* **118** (Jun, 2017) .
- [27] XENON collaboration, E. Aprile et al., *Observation of Excess Electronic Recoil Events in XENON1T*, **2006.09721**.
- [28] P. W. Graham, I. G. Irastorza, S. K. Lamoreaux, A. Lindner and K. A. van Bibber, *Experimental Searches for the Axion and Axion-Like Particles*, *Ann. Rev. Nucl. Part. Sci.* **65** (2015) 485–514, [**1602.00039**].
- [29] P. Di Vecchia, M. Giannotti, M. Lattanzi and A. Lindner, *Round Table on Axions and Axion-like Particles*, *PoS Confinement2018* (2019) 034, [**1902.06567**].
- [30] B. Döbrich, J. Jaeckel, F. Kahlhoefer, A. Ringwald and K. Schmidt-Hoberg, *ALPtraum: ALP production in proton beam dump experiments*, *JHEP* **02** (2016) 018, [**1512.03069**].
- [31] S. Alekhin et al., *A facility to Search for Hidden Particles at the CERN SPS: the SHiP physics case*, *Rept. Prog. Phys.* **79** (2016) 124201, [**1504.04855**].
- [32] C. Abel et al., *Search for Axionlike Dark Matter through Nuclear Spin Precession in Electric and Magnetic Fields*, *Phys. Rev. X* **7** (2017) 041034, [**1708.06367**].
- [33] A. Berlin, R. T. D’Agnolo, S. A. Ellis, C. Nantista, J. Neilson, P. Schuster et al., *Axion Dark Matter Detection by Superconducting Resonant Frequency Conversion*, **1912.11048**.
- [34] T. Wu, J. W. Blanchard, G. P. Centers, N. L. Figueroa, A. Garcon, P. W. Graham et al., *Search for axionlike dark matter with a liquid-state nuclear spin comagnetometer*, *Physical Review Letters* **122** (May, 2019) .
- [35] R. Janish, V. Narayan, S. Rajendran and P. Riggins, *Axion production and detection with superconducting RF cavities*, *Phys. Rev. D* **100** (2019) 015036, [**1904.07245**].



- [36] C. Smorra et al., *Direct limits on the interaction of antiprotons with axion-like dark matter*, *Nature* **575** (2019) 310–314, [[2006.00255](#)].
- [37] A. Garcon, J. W. Blanchard, G. P. Centers, N. L. Figueroa, P. W. Graham, D. F. Jackson Kimball et al., *Constraints on bosonic dark matter from ultralow-field nuclear magnetic resonance*, *Science Advances* **5** (Oct, 2019) eaax4539.
- [38] J. B. Dent, B. Dutta, D. Kim, S. Liao, R. Mahapatra, K. Sinha et al., *New directions for axion searches via scattering at reactor neutrino experiments*, *Physical Review Letters* **124** (May, 2020) .
- [39] G. G. Raffelt, *Astrophysical axion bounds*, *Axions* 51–71.
- [40] J. Jaeckel and A. Ringwald, *The low-energy frontier of particle physics*, *Annual Review of Nuclear and Particle Science* **60** (Nov, 2010) 405–437.
- [41] C. Brust, D. E. Kaplan and M. T. Walters, *New Light Species and the CMB*, *JHEP* **12** (2013) 058, [[1303.5379](#)].
- [42] R. Bollig, W. DeRocco, P. W. Graham and H.-T. Janka, *Muons in supernovae: implications for the axion-muon coupling*, [2005.07141](#).
- [43] N. Bar, K. Blum and G. D’amico, *Is there a supernova bound on axions?*, *Phys. Rev. D* **101** (2020) 123025, [[1907.05020](#)].
- [44] J. Isern, *White Dwarfs as Advanced Physics Laboratories. The Axion case*, in *IAU Symposium 357: White Dwarfs as probes of fundamental physics and tracers of planetary, stellar & galactic evolution*, 2, 2020. [2002.08069](#).
- [45] M. Vysotsky, Y. Zeldovich, M. Khlopov and V. Chechetkin, *Some Astrophysical Limitations on Axion Mass*, *Pisma Zh. Eksp. Teor. Fiz.* **27** (1978) 533–536.
- [46] S. Chang and K. Choi, *Hadronic axion window and the big bang nucleosynthesis*, *Phys. Lett. B* **316** (1993) 51–56, [[hep-ph/9306216](#)].
- [47] D. Cadamuro, S. Hannestad, G. Raffelt and J. Redondo, *Cosmological bounds on sub-MeV mass axions*, *JCAP* **02** (2011) 003, [[1011.3694](#)].
- [48] D. Cadamuro and J. Redondo, *Cosmological bounds on pseudo Nambu-Goldstone bosons*, *JCAP* **02** (2012) 032, [[1110.2895](#)].
- [49] M. Millea, L. Knox and B. Fields, *New Bounds for Axions and Axion-Like Particles with keV-GeV Masses*, *Phys. Rev. D* **92** (2015) 023010, [[1501.04097](#)].
- [50] D. Baumann, D. Green and B. Wallisch, *New Target for Cosmic Axion Searches*, *Phys. Rev. Lett.* **117** (2016) 171301, [[1604.08614](#)].
- [51] F. D’Eramo, R. Z. Ferreira, A. Notari and J. L. Bernal, *Hot Axions and the  $H_0$  tension*, *JCAP* **11** (2018) 014, [[1808.07430](#)].

- [52] G. Grilli di Cortona, E. Hardy, J. Pardo Vega and G. Villadoro, *The QCD axion, precisely*, *JHEP* **01** (2016) 034, [[1511.02867](#)].
- [53] CAST collaboration, E. Arik et al., *Probing eV-scale axions with CAST*, *JCAP* **02** (2009) 008, [[0810.4482](#)].
- [54] P. F. Depta, M. Hufnagel and K. Schmidt-Hoberg, *Robust cosmological constraints on axion-like particles*, *JCAP* **05** (2020) 009, [[2002.08370](#)].
- [55] A. Zhitnitsky, *On Possible Suppression of the Axion Hadron Interactions. (In Russian)*, *Sov. J. Nucl. Phys.* **31** (1980) 260.
- [56] M. Dine, W. Fischler and M. Srednicki, *A simple solution to the strong cp problem with a harmless axion*, *Physics Letters B* **104** (1981) 199 – 202.
- [57] J. E. Kim, *Weak-interaction singlet and strong CP invariance*, *Phys. Rev. Lett.* **43** (Jul, 1979) 103–107.
- [58] M. Shifman, A. Vainshtein and V. Zakharov, *Can confinement ensure natural cp invariance of strong interactions?*, *Nuclear Physics B* **166** (1980) 493 – 506.
- [59] L. Calibbi, D. Redigolo, R. Ziegler and J. Zupan, *Looking forward to Lepton-flavor-violating ALPs*, [2006.04795](#).
- [60] D. Croon, G. Elor, R. K. Leane and S. D. McDermott, *Supernova Muons: New Constraints on Z' Bosons, Axions, and ALPs*, [2006.13942](#).
- [61] K. Barth and A. B. al., *CAST constraints on the axion-electron coupling*, *Journal of Cosmology and Astroparticle Physics* **2013** (may, 2013) 010–010.
- [62] B. D. Fields, K. A. Olive, T.-H. Yeh and C. Young, *Big-Bang Nucleosynthesis After Planck*, *Astrophys. J.* **894** (2020) 109, [[1912.01132](#)].
- [63] G. Mangano, G. Miele, S. Pastor, T. Pinto, O. Pisanti and P. D. Serpico, *Relic neutrino decoupling including flavor oscillations*, *Nucl. Phys. B* **729** (2005) 221–234, [[hep-ph/0506164](#)].
- [64] K. Akita and M. Yamaguchi, *A precision calculation of relic neutrino decoupling*, [2005.07047](#).
- [65] G. Mangano and P. D. Serpico, *A robust upper limit on  $N_{\text{eff}}$  from BBN, circa 2011*, *Phys. Lett. B* **701** (2011) 296–299, [[1103.1261](#)].
- [66] R. Cooke, M. Pettini, R. A. Jorgenson, M. T. Murphy and C. C. Steidel, *Precision measures of the primordial abundance of deuterium*, *Astrophys. J.* **781** (2014) 31, [[1308.3240](#)].
- [67] R. H. Cyburt, B. D. Fields, K. A. Olive and T.-H. Yeh, *Big Bang Nucleosynthesis: 2015*, *Rev. Mod. Phys.* **88** (2016) 015004, [[1505.01076](#)].

- [68] C. Pitrou, A. Coc, J.-P. Uzan and E. Vangioni, *Precision big bang nucleosynthesis with improved Helium-4 predictions*, *Phys. Rept.* **754** (2018) 1–66, [[1801.08023](#)].
- [69] A. Berlin, N. Blinov and S. W. Li, *Dark Sector Equilibration During Nucleosynthesis*, *Phys. Rev. D* **100** (2019) 015038, [[1904.04256](#)].
- [70] W. Fischler and J. Meyers, *Dark Radiation Emerging After Big Bang Nucleosynthesis?*, *Phys. Rev. D* **83** (2011) 063520, [[1011.3501](#)].
- [71] PLANCK collaboration, N. Aghanim et al., *Planck 2018 results. VI. Cosmological parameters*, [1807.06209](#).
- [72] Z. Hou, R. Keisler, L. Knox, M. Millea and C. Reichardt, *How massless neutrinos affect the cosmic microwave background damping tail*, *Physical Review D* **87** (Apr, 2013) .
- [73] S. K. Acharya and R. Khatri, *CMB anisotropy and BBN constraints on pre-recombination decay of dark matter to visible particles*, *JCAP* **12** (2019) 046, [[1910.06272](#)].
- [74] K. F. Riley, M. P. Hobson and S. J. Bence, *Mathematical Methods for Physics and Engineering: A Comprehensive Guide*. Cambridge University Press, 2 ed., 2002. 10.1017/CBO9781139164979.
- [75] S. Hannestad and J. Madsen, *Neutrino decoupling in the early universe*, *Phys. Rev. D* **52** (1995) 1764–1769, [[astro-ph/9506015](#)].
- [76] A. Dolgov, S. Hansen and D. Semikoz, *Nonequilibrium corrections to the spectra of massless neutrinos in the early universe*, *Nucl. Phys. B* **503** (1997) 426–444, [[hep-ph/9703315](#)].
- [77] S. Esposito, G. Miele, S. Pastor, M. Peloso and O. Pisanti, *Nonequilibrium spectra of degenerate relic neutrinos*, *Nucl. Phys. B* **590** (2000) 539–561, [[astro-ph/0005573](#)].
- [78] G.-y. Huang, T. Ohlsson and S. Zhou, *Observational Constraints on Secret Neutrino Interactions from Big Bang Nucleosynthesis*, *Phys. Rev. D* **97** (2018) 075009, [[1712.04792](#)].