

OPTIMAL CONTROL FOR THE INFINITY OBSTACLE PROBLEM

HENOK MAWI AND CHEIKH BIRAHIM NDIAYE

ABSTRACT. In this note, we show that a natural optimal control problem for the ∞ -obstacle problem admits an optimal control which is also an optimal state. Moreover, we show the convergence of the minimal value of an optimal control problem for the p -obstacle problem to the minimal value of our optimal control problem for the ∞ -obstacle problem, as $p \rightarrow \infty$.

1. INTRODUCTION

The obstacle problem corresponding to an obstacle f in

$$(1.1) \quad W_g^{1,2}(\Omega) = \{u \in W^{1,2}(\Omega) : u = g \text{ on } \partial\Omega\}$$

consists of minimizing the Dirichlet energy

$$\int_{\Omega} |Du(x)|^2 dx$$

over the set

$$(1.2) \quad \mathbb{K}_{f,g}^2 = \{u \in W_g^{1,2}(\Omega) : u(x) \geq f(x) \text{ in } \Omega\}$$

where $\Omega \subset \mathbb{R}^n$ is a bounded and smooth domain, Du is the gradient of u , and $g \in \text{tr}(W^{1,2}(\Omega))$ with tr the trace operator. In (1.1), the equality $u = g$ on $\partial\Omega$ is in the sense of trace. This problem is used to model the equilibrium position of an elastic membrane whose boundary is held fixed at g and is forced to remain above a given obstacle f . It is known that the obstacle problem admits a unique solution $v \in \mathbb{K}_{f,g}^2$. That is, there is a unique $v \in \mathbb{K}_{f,g}^2$ such that

$$\int_{\Omega} |Dv(x)|^2 dx \leq \int_{\Omega} |Du(x)|^2 dx, \quad \forall u \in \mathbb{K}_{f,g}^2.$$

In [3] Adams, Lenhart and Yong introduced an optimal control problem for the obstacle problem by studying the minimizer of the functional

$$J_2(\psi) = \frac{1}{2} \int_{\Omega} (|T_2(\psi) - z|^2 + |D\psi|^2) dx.$$

In the above variational problem, following the terminology in control theory [16], ψ is called the control variable and $T_2(\psi)$ is the corresponding state. The control ψ lies in the space $W_0^{1,2}(\Omega)$, the state $T_2(\psi)$ is the unique solution for the obstacle problem corresponding to the obstacle ψ and the profile z is in $L^2(\Omega)$. The authors proved that there exists a unique minimizer $\bar{\psi} \in W_0^{1,2}(\Omega)$ of the functional J_2 . Furthermore, they showed that $T_2(\bar{\psi}) = \bar{\psi}$.

February 26, 2022.

The first author was partially supported by NSF grant HRD-1700236.

Following suit, for $1 < p < \infty$, and $z \in L^p(\Omega)$, Lou in [17] considered the variational problem of minimizing the functional

$$(P_p) \quad \bar{J}_p(\psi) = \frac{1}{p} \int_{\Omega} |T_p(\psi) - z|^p + |D\psi|^p dx$$

for $\psi \in W_0^{1,p}(\Omega) := \{u \in W^{1,p}(\Omega) : u = 0 \text{ on } \partial\Omega\}$ and established that the problem admits a minimizer ψ . Here $T_p(\psi)$ is the unique solution for the p -obstacle problem with obstacle $\psi \in W_0^{1,p}(\Omega)$, see [6] and references therein for discussions about the p -obstacle problem. We remind the reader that the p -obstacle problem with obstacle $f \in W_g^{1,p}(\Omega)$ refers to the problem of minimizing the p -Dirichlet energy

$$\int_{\Omega} |Du(x)|^p dx$$

among all functions in the class

$$\mathbb{K}_{f,g}^p = \{u \in W^{1,p}(\Omega) : u \geq f \text{ in } \Omega \text{ and } u = g \text{ on } \partial\Omega\},$$

with $g \in \text{tr}(W^{1,p}(\Omega))$. It is further shown in [17] that, as in the case of $p = 2$, $T_p(\bar{\psi}) = \bar{\psi}$.

For the boundary data $g \in \text{Lip}(\partial\Omega)$, letting $p \rightarrow \infty$, one obtains a limiting variational problem of L^∞ -type which is referred in the literature as the infinity obstacle problem or ∞ -obstacle problem (see [20]). That is, given an obstacle $f \in W_g^{1,\infty}(\Omega)$ one considers the minimization problem:

$$(1.3) \quad \text{Finding } u_\infty \in \mathbb{K}_{f,g}^\infty : \|Du_\infty\|_\infty = \inf_{u \in \mathbb{K}_{f,g}^\infty} \|Du\|_\infty,$$

where

$$\mathbb{K}_{f,g}^\infty = \{u \in W^{1,\infty}(\Omega) : u \geq f \text{ in } \Omega \text{ and } u = g \text{ on } \partial\Omega\}, \text{ and } \|\cdot\|_\infty := \text{ess sup } |\cdot|.$$

It is established in [20] that the minimization problem (1.3) has a solution

$$(1.4) \quad u_\infty := u_\infty(f) \in \mathbb{K}_{f,g}^\infty$$

which verifies

$$(1.5) \quad -\Delta_\infty u_\infty \geq 0 \text{ in } \Omega \text{ in a weak sense.}$$

More importantly, the authors in [20] characterize u_∞ as the smallest infinity superharmonic function on Ω that is larger than the obstacle f and equals g on the boundary. Thus for a fixed $F \in \text{Lip}(\partial\Omega)$, this generates an obstacle to solution operator

$$T_\infty : W_F^{1,\infty}(\Omega) \longrightarrow W_F^{1,\infty}(\Omega)$$

defined by

$$(1.6) \quad T_\infty(f) := u_\infty(f) \in W_F^{1,\infty}(\Omega), \quad f \in W_F^{1,\infty}(\Omega),$$

where

$$W_F^{1,\infty}(\Omega) := \{u \in W^{1,\infty}(\Omega) : u = F \text{ on } \partial\Omega\}.$$

In this note, we consider a natural optimal control problem for the infinity obstacle problem. More precisely, for $F \in \text{Lip}(\partial\Omega)$ and for $z \in L^\infty(\Omega)$ fixed, we introduce the functional

$$J_\infty(\psi) = \max\{\|T_\infty(\psi) - z\|_\infty, \|D\psi\|_\infty\}, \quad \psi \in W_F^{1,\infty}(\Omega)$$

and study the problem of existence of $\psi_\infty \in W_F^{1,\infty}(\Omega)$ such that:

$$(P_\infty) \quad J_\infty(\psi_\infty) \leq J_\infty(\psi), \quad \forall \psi \in W_F^{1,\infty}(\Omega).$$

In deference to optimal control theory, a function ψ_∞ satisfying (P_∞) is called an *optimal control* and the state $T_\infty(\psi_\infty)$ is called an *optimal state*.

Several variants of control problems where the control variable is the obstacle have been studied by different authors since the first of such works appeared in [3]. The literature is vast, but to mention a few, in [2] the authors studied a generalization of [3] by adding a source term. In [1] a similar problem is studied when the state is a solution to a parabolic variational inequality. In [18] the author studied regularity of the optimal state obtained in [3]. When the state is governed by a bilateral variational inequality, results are obtained in [9], [10], [11] and [12]. Optimal control for higher order obstacle problems appears in [5] and [14]. Related works where the control variable is the obstacle are also studied in [13, 21] and the references therein.

In this note, we prove that the optimal control problem (P_∞) associated to J_∞ is solvable. Precisely we show the following result:

Theorem 1.1. *Assuming that $\Omega \subset \mathbb{R}^n$ is a bounded and smooth domain, $F \in Lip(\partial\Omega)$, and $z \in L^\infty(\Omega)$, J_∞ admits an optimal control $u_\infty \in W_F^{1,\infty}(\Omega)$ which is also an optimal state, i.e*

$$u_\infty = T_\infty(u_\infty).$$

Using also arguments similar to the ones used in the proof of Theorem 1.1, we show the convergence of the minimal value of an optimal control problem associated to \bar{J}_p to the minimal value of the optimal control problem corresponding to J_∞ as p tends to infinity. Indeed we prove the following result:

Theorem 1.2. *Let $\Omega \subset \mathbb{R}^n$ be a bounded and smooth domain, $F \in Lip(\partial\Omega)$, and $z \in L^\infty(\Omega)$. Then setting*

$$J_p = (p\bar{J}_p)^{\frac{1}{p}}, \quad C_p = \min_{\psi \in W_F^{1,p}(\Omega)} J_p(\psi) \text{ for } 1 < p < \infty, \text{ and } C_\infty = \min_{\psi \in W_F^{1,\infty}(\Omega)} J_\infty(\psi),$$

where \bar{J}_p is as in (P_p) , we have

$$\lim_{p \rightarrow \infty} C_p = C_\infty$$

In the proofs of the above results, we use the p -approximation technique as in the study of the ∞ -obstacle problem combined with the classical methods of weak convergence in Calculus of Variations. As in the study of the ∞ -obstacle problem, here also the key analytical ingredients are the L^q -characterization of L^∞ and Hölder's inequality. The difficulty arises from the fact that the unicity question for the ∞ -obstacle problem is still an open problem to the best of our knowledge. To overcome the latter issue, we make use of the characterization of the solution of the ∞ -obstacle problem by Rossi-Teixeira-Urbano [20].

2. PRELIMINARIES

One of the most popular way of approaching problems related to minimizing a functional of L^∞ -type is to follow the idea first introduced by Aronsson in [7] and which involves interpreting an L^∞ -type minimization problem as a limit when $p \rightarrow \infty$ of an L^p -type minimization problem. In this note, this p -approximation technique will be used to show existence of an optimal control for J_∞ . In order to prepare for our use of the p -approximation technique, we are going to start this section by discussing some related L^p -type variational problems.

Let $\Omega \subset \mathbb{R}^n$ be a bounded and smooth domain and $g \in Lip(\partial\Omega)$. Moreover let $\psi \in W_g^{1,\infty}(\Omega)$ be fixed and $1 < p < \infty$. Then as described earlier the p -obstacle problem with obstacle ψ corresponds to finding a minimizer of the functional

$$(2.1) \quad I_p(v) = \int_{\Omega} |Dv(x)|^p dx$$

over the space $\mathbb{K}_{\psi,g}^p = \{v \in W^{1,p}(\Omega) : v \geq \psi, \text{ and } v = g \text{ on } \partial\Omega\}$. The energy integral (2.1) admits a unique minimizer $u_p \in \mathbb{K}_{\psi,g}^p$. The minimizer u_p is not only p -superharmonic, i.e. $\Delta_p u_p \leq 0$, but is also a weak solution to the following system

$$(2.2) \quad \begin{cases} -\Delta_p u \geq 0 & \text{in } \Omega \\ -\Delta_p u (u - \psi) = 0 & \text{in } \Omega \\ u \geq \psi & \text{in } \Omega \end{cases}$$

where Δ_p is the p -Laplace operator given by

$$\Delta_p u := \operatorname{div}(|Du|^{p-2} Du).$$

Moreover, it is known that the p -obstacle problem is equivalent to the system (2.2) (see [16] or [19]) and hence we will refer to (2.2) as the p -obstacle problem as well. On the other hand, by the equivalence of weak and viscosity solutions established in [19] (and [15]) u_p is also a viscosity solution of (2.2) according to the following definition.

Definition 2.1. *A function $u \in C(\Omega)$ is said to be a viscosity subsolution (supersolution) to*

$$(2.3) \quad \begin{aligned} F(x, u, Du, D^2u) &= 0 & \text{in } \Omega \\ u &= 0 & \text{in } \partial\Omega \end{aligned}$$

if for every $\phi \in C^2(\Omega)$ and $x_0 \in \Omega$ whenever $\phi - u$ has a minimum (resp. maximum) in a neighborhood of x_0 in Ω we have:

$$F(x, u, D\phi, D^2\phi) \leq 0 \quad (\text{resp.} \quad \geq 0).$$

The function u is called a viscosity solution of (2.3) in Ω if u is both viscosity subsolution and viscosity supersolution of (2.3) in Ω .

The asymptotic behavior of the sequence of minimizers $(u_p)_{p>1}$ as p tends to infinity has been investigated in [20]. In fact, in [20], it is established that for a fixed $\psi \in W_g^{1,\infty}(\Omega)$, there exists $u_\infty = u_\infty(\psi) \in \mathbb{K}_{\psi,g}^\infty = \{v \in W_g^{1,\infty}(\Omega) : v \geq \psi\}$ such that $u_p \rightarrow u_\infty$ locally uniformly

in $\bar{\Omega}$, and that for every $q \geq 1$, u_p converges to u_∞ weakly in $W^{1,q}(\Omega)$. Furthermore, u_∞ is a solution to the ∞ -obstacle problem

$$(2.4) \quad \min_{v \in \mathbb{K}_{\psi,g}^\infty} \|Dv\|_\infty$$

For Ω convex (see [8]), the variational problem (2.4) is equivalent to the minimization problem

$$\min_{v \in \mathbb{K}_{\psi,g}^\infty} \mathcal{L}(v),$$

where

$$\mathcal{L}(v) = \inf_{(x,y) \in \Omega^2, x \neq y} \frac{|v(x) - v(y)|}{|x - y|}.$$

Moreover, in [20], it is shown that u_∞ is a viscosity solution to the following system.

$$\begin{cases} -\Delta_\infty u \geq 0 & \text{in } \Omega \\ -\Delta_\infty u (u - \psi) = 0 & \text{in } \Omega \\ u \geq \psi & \text{in } \Omega \end{cases}$$

where Δ_∞ is the ∞ -Laplacian and is defined by

$$\Delta_\infty u = \langle D^2 u Du, Du \rangle = \sum_{i=1}^n \sum_{j=1}^n u_{x_i} u_{x_j} u_{x_i x_j}.$$

Recalling that u is said to be *infinity superharmonic* or ∞ -*superharmonic*, if $-\Delta_\infty u \geq 0$ in the viscosity sense, we have the following characterization of u_∞ in terms of infinity superharmonic functions and it is proven in [20]. We would like to emphasize that this will play an important role in our arguments.

Lemma 2.2. *Setting*

$$\mathcal{F}^+ = \{v \in C(\Omega), -\Delta_\infty v \geq 0 \text{ in } \Omega \text{ in the viscosity sense}\}$$

and

$$\mathcal{F}_\psi^+ = \{v \in \mathcal{F}^+, v \geq \psi \text{ in } \Omega, \text{ and } v = \psi \text{ on } \partial\Omega\},$$

we have

$$(2.5) \quad T_\infty(\psi) = u_\infty = \inf_{v \in \mathcal{F}_\psi^+} v,$$

with T_∞ as defined earlier in (1.6).

Lemma 2.2 implies the following characterization of infinity superharmonic functions as fixed points of T_∞ . This characterization plays a key role in our p -approximation scheme for existence.

Lemma 2.3. *Assuming that $u \in W_g^{1,\infty}(\Omega)$, u being infinity superharmonic is equivalent to u being a fixed point of T_∞ , i.e*

$$T_\infty(u) = u.$$

Proof. Let $u \in W_g^{1,\infty}(\Omega)$ be an infinity superharmonic function and v be defined by $v = T_\infty(u)$. Then clearly the definition of v and lemma 2.2 imply $v \geq u$. On the other hand, since $u \in W_g^{1,\infty}(\Omega)$ and is an infinity superharmonic function, we deduce from lemma 2.2 that $u \geq T_\infty(u) = v$. Thus, we get $T_\infty(u) = u$. Now if $u = T_\infty(u)$, then using again lemma 2.2 or (1.4)-(1.6), we obtain u is an infinity superharmonic function. Hence the proof of the lemma is complete. \square

To run our p -approximation scheme for existence, another crucial ingredient that we will need is an appropriate characterization of the limit of sequence of solution w_p of the p -obstacle problem (2.2) with obstacle ψ_p under uniform convergence of both w_p and ψ_p . Precisely, we will need the following lemma.

Lemma 2.4. *If w_p is a solution to the p -obstacle problem (2.2) with obstacle ψ_p that is, w_p satisfies*

$$(2.6) \quad \begin{cases} -\Delta_p w_p \geq 0 & \text{in } \Omega \\ -\Delta_p w_p (w_p - \psi_p) = 0 & \text{in } \Omega \\ w_p \geq \psi_p & \text{in } \Omega \end{cases}$$

in the viscosity sense and if also that $w_p \rightarrow u_\infty$ and $\psi_p \rightarrow \psi_\infty$ locally uniformly in $\overline{\Omega}$, then u_∞ is a solution in the viscosity sense of the following system

$$(2.7) \quad \begin{cases} -\Delta_\infty w_\infty \geq 0 & \text{in } \Omega \\ -\Delta_\infty w_\infty (w_\infty - \psi_\infty) = 0 & \text{in } \Omega \\ w_\infty \geq \psi_\infty & \text{in } \Omega. \end{cases}$$

Proof. First of all, note that since $w_p \geq \psi_p$, $-\Delta_p w_p \geq 0$ in the viscosity sense in Ω for every p , $w_p \rightarrow u_\infty$, and $\psi_p \rightarrow \psi_\infty$ both locally uniformly in $\overline{\Omega}$, and $\overline{\Omega}$ is compact, we have $w_\infty \geq \psi_\infty$ and $-\Delta_\infty w_\infty \geq 0$ in the viscosity sense in Ω . It thus remains to prove that $-\Delta_\infty w_\infty (w_\infty - \psi_\infty) = 0$ in Ω which (because of $w_\infty \geq \psi_\infty$ in Ω) is equivalent to $-\Delta_\infty w_\infty = 0$ in $\{w_\infty > \psi_\infty\} := \{x \in \Omega : w_\infty(x) > \psi_\infty(x)\}$. Thus to conclude the proof, we are going to show $-\Delta_\infty w_\infty = 0$ in $\{w_\infty > \psi_\infty\}$. To that end, fix $y \in \{w_\infty > \psi_\infty\}$. Then, by continuity there exists an open neighborhood V of y in Ω such that \overline{V} is a compact subset of Ω , and a small real number $\delta > 0$ such that $w_\infty > \delta > \psi_\infty$ in \overline{V} . Thus, from $w_p \rightarrow w_\infty$, $\psi_p \rightarrow \psi_\infty$ locally uniformly in $\overline{\Omega}$, and \overline{V} compact subset of Ω , we infer that for sufficiently large p

$$(2.8) \quad w_p > \delta > \psi_p \quad \text{in } \overline{V}.$$

On the other hand, since w_p is a solution to the p obstacle problem (2.2) with obstacle ψ_p , then clearly $-\Delta_p w_p = 0$ in $\{w_p > \psi_p\} := \{x \in \Omega : w_p(x) > \psi_p(x)\}$. Thus, (2.8) imply $-\Delta_p w_p = 0$ in the sense of viscosity in V . Hence, recalling that $w_p \rightarrow w_\infty$ locally uniformly in $\overline{\Omega}$ and letting $p \rightarrow \infty$, we obtain

$$-\Delta_\infty w_\infty = 0 \quad \text{in the sense of viscosity in } V.$$

Thus, since $y \in V$ is arbitrary in $\{w_\infty > \psi_\infty\}$, then we arrive to

$$-\Delta_\infty w_\infty = 0 \quad \text{in the sense of viscosity in } \{w_\infty > \psi_\infty\},$$

thereby ending the proof of the lemma. \square

On the other hand, to show the convergence of the minimal values of J_p to that of J_∞ , we will make use of the following elementary results.

Lemma 2.5. *Suppose $\{a_p\}$ and $\{b_p\}$ are nonnegative sequences with*

$$\liminf_{p \rightarrow \infty} a_p = a \quad \text{and} \quad \liminf_{p \rightarrow \infty} b_p = b.$$

Then

$$\liminf_{p \rightarrow \infty} \max\{a_p, b_p\} = \max\{a, b\}.$$

Proof. Let $\{b_{p_k}\}$ be a subsequence converging to $b = \liminf_{p \rightarrow \infty} b_p$. Then

$$\lim_{k \rightarrow \infty} \max\{a_{p_k}, b_{p_k}\} = \max\{a, b\}.$$

Since the \liminf is the smallest limit point we have

$$(2.9) \quad \liminf_{p \rightarrow \infty} \max\{a_p, b_p\} \leq \max\{a, b\}.$$

On the other hand

$$a_p, b_p \leq \max\{a_p, b_p\}, \quad \text{for all } p.$$

Thus

$$b = \liminf_{p \rightarrow \infty} b_p \leq \liminf_{p \rightarrow \infty} \max\{a_p, b_p\},$$

and likewise

$$a \leq \liminf_{p \rightarrow \infty} \max\{a_p, b_p\}.$$

Consequently

$$(2.10) \quad \liminf_{p \rightarrow \infty} \max\{a_p, b_p\} \geq \max\{a, b\}.$$

Finally (2.9) and (2.10) conclude the proof of the lemma. \square

Lemma 2.6. *Suppose $\{a_p\}$ and $\{b_p\}$ are nonnegative sequences with*

$$\liminf_{p \rightarrow \infty} a_p = a \quad \text{and} \quad \liminf_{p \rightarrow \infty} b_p = b.$$

Then

$$\liminf_{p \rightarrow \infty} (a_p^p + b_p^p)^{1/p} = \max\{a, b\}.$$

Proof. It follows directly from the trivial inequality

$$2^{1/p} \max\{a_p, b_p\} \geq (a_p^p + b_p^p)^{1/p} \geq \max\{a_p, b_p\}, \quad \forall p \geq 1,$$

lemma 2.5 and the fact that $\liminf_n (a_n b_n) = (\lim_n a_n)(\liminf_n b_n)$ if $\lim_n a_n > 0$. \square

3. EXISTENCE OF OPTIMAL CONTROL FOR J_∞ AND LIMIT OF C_p

In this section, we show the existence of an optimal control for J_∞ and show that C_p converges to C_∞ as $p \rightarrow \infty$. We divide it in two subsections. In the first one we show existence of an optimal control for J_∞ via the p -approximation technique, and in the second one we show that C_p converges to C_∞ as p tends to infinity.

3.1. Existence of optimal control. In this subsection, we show the existence of a minimizer of J_∞ via the p -approximation technique using solutions of the optimal control for J_p . For this end, we start by recalling some optimality facts about J_p inherited from \bar{J}_p (see (P_p) for its definition) and mentioned in the introduction. For $\Omega \subset \mathbb{R}^n$ a bounded and smooth domain, $z \in L^\infty(\Omega)$, $F \in Lip(\partial\Omega)$, and $1 < p < \infty$, we recall that the functional J_p is defined by the formula

$$(3.1) \quad J_p(\psi) = \left[\int_{\Omega} |T_p(\psi) - z|^p + |D\psi|^p dx \right]^{1/p}, \quad \psi \in W_F^{1,p}(\Omega)$$

and that the optimal control problem for J_p is the variational problem of minimizing J_p , namely

$$(3.2) \quad \inf_{\psi \in W_F^{1,p}(\Omega)} J_p(\psi)$$

over $W_F^{1,p}(\Omega)$, where

$$W_F^{1,p}(\Omega) = \{\psi \in W^{1,p}(\Omega) : \psi = F \text{ on } \partial\Omega\},$$

and $T_p(\psi)$ is the solution to the p -obstacle problem with obstacle ψ . Moreover, as for the functional \bar{J}_p , J_p also admits a minimizer $\psi_p \in W_F^{1,p}(\Omega)$ verifying

$$(3.3) \quad T_p(\psi_p) = \psi_p.$$

As mentioned in the introduction, for more details about the latter results, see [3] for $p = 2$ and see [17] for $p > 2$.

To continue, let us pick $\eta \in W_F^{1,\infty}(\Omega)$. Since η competes in the minimization problem (3.2), we have

$$\int_{\Omega} |D\psi_p|^p dx \leq J_p(\eta) = \int_{\Omega} |T_p(\eta) - z|^p + |D\eta|^p dx.$$

Since $\bar{\Omega}$ is compact and $T_p(\eta) \rightarrow T_\infty(\eta)$ as $p \rightarrow \infty$ locally uniformly on $\bar{\Omega}$ (which follows from the definition of $T_\infty(\eta)$), we deduce that for p very large

$$(3.4) \quad \int_{\Omega} |D\psi_p|^p dx \leq M^p |\Omega|$$

for some M which depends only on $\|\eta\|_{W^{1,\infty}}$, $\|T_\infty(\eta)\|_{C^0}$ and $\|z\|_\infty$. Furthermore, let us fix $1 < q < p$. Then by using Holder's inequality, we can write

$$(3.5) \quad \int_{\Omega} |D\psi_p|^q dx \leq \left\{ \int_{\Omega} (|D\psi_p|^p)^{q/p} dx \right\}^{q/p} |\Omega|^{\frac{p-q}{p}}$$

and we obtain by using (3.4) that for p very large

$$\int_{\Omega} |D\psi_p|^q dx \leq M^q |\Omega|^{\frac{q}{p}} |\Omega|^{\frac{p-q}{p}}$$

and raising both sides to $1/q$, we derive that for p very large, there holds

$$\|D\psi_p\|_{L^q} \leq M |\Omega|^{1/q},$$

with $\|\cdot\|_{L^q}$ denoting the classical $L^q(\Omega)$ -norm. This shows, that the sequence $\{\psi_p\}$ is bounded in $W_F^{1,q}(\Omega)$ in the gradient norm for every q with a bound independent of q ,

and by Poincare's inequality, that for every $1 < q < \infty$, the sequence $\{\psi_p\}$ is bounded in $W_F^{1,q}(\Omega)$ in the standard $W^{1,q}(\Omega)$ -norm. Therefore, by classical weak compactness arguments, we have that, up to a subsequence,

(3.6)

$$\psi_p \longrightarrow \psi_\infty, \text{ as } p \rightarrow \infty \text{ locally uniformly in } \overline{\Omega} \text{ and weakly in } W^{1,q}(\Omega) \quad \forall \quad 1 < q < \infty.$$

Notice that consequently $\|D\psi_\infty\|_{L^q} \leq M|\Omega|^{1/q}$ for all $1 < q < \infty$. Thus, we deduce once again by Poincare's inequality that

$$(3.7) \quad \psi_\infty \in W_F^{1,\infty}(\Omega).$$

We want now to show that ψ_∞ is a minimizer of J_∞ . To that end, we make the following observation which is a consequence of lemma 2.4.

Lemma 3.1. *The function ψ_∞ is a fixed point of T_∞ , namely*

$$T_\infty(\psi_\infty) = \psi_\infty,$$

and the solutions $T_p(\psi_p)$ of the p -obstacle problem with obstacle ψ_p verify: as $p \rightarrow \infty$,

$$T_p(\psi_p) \longrightarrow T_\infty(\psi_\infty) \text{ locally uniformly in } \overline{\Omega} \text{ and weakly in } W^{1,q}(\Omega) \quad \forall \quad 1 < q < \infty.$$

Proof. We know that $T_p(\psi_p) = \psi_p$ (see (3.3)) Thus using (3.6) and Lemma 2.4 with $\phi_p = \psi_p$ and $w_p = T_p(\psi_p) = \psi_p$, we have $T_p(\psi_p) \rightarrow \psi_\infty$ locally uniformly in $\overline{\Omega}$, weakly in $W^{1,q}(\Omega)$ for every $1 < q < \infty$, and ψ_∞ is a infinity superharmonic. Thus, recalling (3.7), we have lemma 2.3 implies $T_\infty(\psi_\infty) = \psi_\infty$. Hence the proof of the lemma is complete. \square

Now, with all the ingredients at hand, we are ready to show that ψ_∞ is a minimizer of J_∞ . Indeed, we are going to show the following proposition:

Proposition 3.2. *Let $\Omega \subseteq \mathbb{R}^n$ be a bounded and smooth domain, $F \in Lip(\partial\Omega)$ and $z \in L^\infty(\Omega)$. Then ψ_∞ is a minimizer of J_∞ on $W_F^{1,\infty}(\Omega)$ That is:*

$$J_\infty(\psi_\infty) = \min_{\eta \in W_F^{1,\infty}(\Omega)} J_\infty(\eta)$$

Proof. We first introduce for $n < p < \infty$ and $\psi \in W_F^{1,p}(\Omega)$

$$H_p(\psi) = \max\{\|T_p(\psi) - z\|_\infty, \|D\psi\|_\infty\},$$

which is well defined by Sobolev Embedding Theorem. Then for any $\eta \in W_F^{1,\infty}(\Omega)$

$$\int_\Omega |D\psi_p|^p dx \leq J_p^p(\eta) = \int_\Omega (|T_p(\eta) - z|^p + |D\eta|^p) dx.$$

Therefore, using the trivial identity $(|a|^p + |b|^p)^{\frac{1}{p}} \leq 2^{\frac{1}{p}} \max\{|a|, |b|\}$, we get

$$\left(\int_\Omega |D\psi_p|^p dx \right)^{1/p} \leq 2^{1/p} |\Omega|^{1/p} H_p(\eta).$$

If we now set

$$(3.8) \quad I_p = \inf_{\eta \in W_F^{1,\infty}(\Omega)} H_p(\eta),$$

we deduce that

$$\left(\int_{\Omega} |D\psi_p|^p dx \right)^{1/p} \leq 2^{1/p} |\Omega|^{1/p} I_p.$$

Let us fix q such that $n < q < \infty$. Then for $q < p < \infty$, by proceeding as in (3.5), we obtain

$$\|D\psi_p\|_{L^q} \leq 2^{1/p} I_p |\Omega|^{1/q}.$$

Similarly,

$$\|T_p(\psi_p) - z\|_{L^q} \leq 2^{1/p} I_p |\Omega|^{1/q}.$$

Thus

$$(3.9) \quad \max\{\|T_p(\psi_p) - z\|_{L^q}, \|D\psi_p\|_{L^q}\} \leq 2^{1/p} I_p |\Omega|^{1/q}.$$

For any $\eta \in W_F^{1,\infty}(\Omega)$ we also have $I_p \leq H_p(\eta)$ and $\liminf_{p \rightarrow \infty} I_p \leq \liminf_{p \rightarrow \infty} H_p(\eta)$. Thus, since ψ_p converges weakly in $W^{1,q}(\Omega)$ to ψ_∞ as $p \rightarrow \infty$ and (3.9) holds, then by weak lower semicontinuity, we conclude that

$$\|D\psi_\infty\|_{L^q} \leq \liminf_{p \rightarrow \infty} \|D\psi_p\|_{L^q} \leq |\Omega|^{1/q} \liminf_{p \rightarrow \infty} H_p(\eta).$$

Moreover, since $T_p(\eta)$ converges locally uniformly on $\overline{\Omega}$ to $T_\infty(\eta)$ as $p \rightarrow \infty$ and $\overline{\Omega}$ is compact, then clearly

$$\lim_{p \rightarrow \infty} H_p(\eta) = J_\infty(\eta),$$

and hence

$$\|D\psi_\infty\|_{L^q} \leq J_\infty(\eta) |\Omega|^{1/q}.$$

Since this holds for any element η of $W_F^{1,\infty}(\Omega)$, we conclude that by taking the infimum over $W_F^{1,\infty}(\Omega)$ and letting $q \rightarrow \infty$

$$(3.10) \quad \|D\psi_\infty\|_\infty \leq \inf_{\eta \in W_F^{1,\infty}(\Omega)} J_\infty(\eta) \leq J_\infty(\psi_\infty).$$

Using lemma 3.1 and equation (3.9) combined with Rellich compactness Theorem or the continuous embedding of L^∞ into L^q , we conclude that

$$\|T_\infty(\psi_\infty) - z\|_{L^q} = \lim_{p \rightarrow \infty} \|T_p(\psi_p) - z\|_{L^q} \leq |\Omega|^{1/q} \liminf_{p \rightarrow \infty} H_p(\eta).$$

Thus, as above letting q goes to infinity and taking infimum in η over $W_F^{1,\infty}(\Omega)$, we also have

$$(3.11) \quad \|T_\infty(\psi_\infty) - z\|_\infty \leq \inf_{\eta \in W_F^{1,\infty}(\Omega)} J_\infty(\eta) \leq J_\infty(\psi_\infty).$$

Finally, from (3.7), (3.10) and (3.11) we deduce

$$J_\infty(\psi_\infty) = \min_{\eta \in W_F^{1,\infty}(\Omega)} J_\infty(\eta),$$

as desired. □

3.2. Convergence of Minimum Values. In this subsection, we show the convergence of the minimal value of the optimal control problem of J_p to the one of J_∞ as $p \rightarrow \infty$, namely Theorem 1.2 via the following proposition:

Proposition 3.3. *Let $\Omega \subset \mathbb{R}^n$ be a bounded and smooth domain, $F \in \text{Lip}(\partial\Omega)$ and $1 < p < \infty$. Then recalling that*

$$C_p = \min_{\psi \in W_F^{1,p}(\Omega)} J_p(\psi) \quad \text{and} \quad C_\infty = \min_{\psi \in W_F^{1,\infty}(\Omega)} J_\infty(\psi),$$

we have

$$\lim_{p \rightarrow \infty} C_p = C_\infty.$$

Proof. Let $\psi_p \in W_F^{1,p}(\Omega)$ and $\psi_\infty \in W_F^{1,\infty}(\Omega)$ be as in subsection 3.1. Then they satisfy $J_p(\psi_p) = C_p$ and $J_\infty(\psi_\infty) = C_\infty$. Moreover, up to a subsequence, we have ψ_p and ψ_∞ verify (3.6) and the conclusions of lemma 3.1. On the other hand, by minimality and Hölder's inequality, we have

$$J_p(\psi_p) \leq J_p(\psi_\infty) \leq 2^{1/p} |\Omega|^{1/p} \max\{\|T_p(\psi_\infty) - z\|_\infty, \|D\psi_\infty\|_\infty\}.$$

Thus

$$(3.12) \quad \limsup_{p \rightarrow \infty} J_p(\psi_p) \leq J_\infty(u_\infty).$$

Now we are going to show the following

$$(3.13) \quad J_\infty(\psi_\infty) \leq \liminf_{p \rightarrow \infty} J_p(\psi_p).$$

To that end observe that by definition of J_∞ , we have

$$(3.14) \quad J_\infty(\psi_\infty) = \max\{\|T_\infty(\psi_\infty) - z\|_\infty, \|D\psi_\infty\|_\infty\}.$$

Thus, using the L^q -characterization of L^∞ , we have that (3.14) imply

$$(3.15) \quad J_\infty(\psi_\infty) = \max\left\{\lim_{q \rightarrow \infty} \|T_\infty(\psi_\infty) - z\|_{L^q}, \lim_{q \rightarrow \infty} \|D\psi_\infty\|_{L^q}\right\},$$

and by using lemma 2.5, we get

$$(3.16) \quad J_\infty(\psi_\infty) = \lim_{q \rightarrow \infty} \max\{\|T_\infty(\psi_\infty) - z\|_{L^q}, \|D\psi_\infty\|_{L^q}\}.$$

On the other hand, by weak lower semicontinuity, and corollary 3.1, we have

$$(3.17) \quad \|D\psi_\infty\|_{L^q} \leq \liminf_{p \rightarrow \infty} \|D\psi_p\|_{L^q}.$$

Now, combining (3.16) and (3.17), we obtain

$$(3.18) \quad J_\infty(\psi_\infty) \leq \liminf_{q \rightarrow \infty} \max\{\|T_\infty(\psi_\infty) - z\|_{L^q}, \liminf_{p \rightarrow \infty} \|D\psi_p\|_{L^q}\}.$$

Next, using lemma 2.6, corollary 3.1, and (3.18), we get

$$(3.19) \quad J_\infty(\psi_\infty) \leq \liminf_{q \rightarrow \infty} \liminf_{p \rightarrow \infty} \{(\|T_p(\psi_p) - z\|_{L^q})^p + (\|D\psi_p\|_{L^q})^p\}^{1/p}.$$

To continue, we are going to estimate the right hand side of (3.19). Indeed, using Hölder's inequality, we have

$$\begin{aligned} (||T_p(\psi_p) - z||_{L^q})^p &= \left\{ \int_{\Omega} |T_p(\psi_p) - z|^q dx \right\}^{p/q} \\ &\leq \left\{ \int_{\Omega} |T_p(\psi_p) - z|^p dx \right\} |\Omega|^{(1-q/p)p/q} \\ &= \left\{ \int_{\Omega} |T_p(\psi_p) - z|^p dx \right\} |\Omega|^{(1-q/p)p/q}. \end{aligned}$$

Similarly, we obtain

$$(||D\psi_p||_{L^q})^p \leq \left\{ \int_{\Omega} |D\psi_p|^p dx \right\} |\Omega|^{(1-q/p)p/q}.$$

By using the latter two estimates in (3.19), we get

$$\begin{aligned} J_{\infty}(\psi_{\infty}) &\leq \liminf_{q \rightarrow \infty} \liminf_{p \rightarrow \infty} \left[\left\{ \int_{\Omega} (|T_p(\psi_p) - z|^p + |D\psi_p|^p) dx \right\}^{1/p} |\Omega|^{(1-q/p)p/q(1/p)} \right] \\ &= \liminf_{q \rightarrow \infty} \liminf_{p \rightarrow \infty} \left[\left\{ \int_{\Omega} (|T_p(\psi_p) - z|^p + |D\psi_p|^p) dx \right\}^{1/p} |\Omega|^{\frac{1}{q} - \frac{1}{p}} \right] \\ (3.20) \quad &= \liminf_{q \rightarrow \infty} \left[|\Omega|^{\frac{1}{q}} \liminf_{p \rightarrow \infty} J_p(\psi_p) \right] = \liminf_{p \rightarrow \infty} J_p(\psi_p) \end{aligned}$$

proving claim (3.13). Combining (3.12) with (3.20) we obtain

$$\lim_{p \rightarrow \infty} J_p(\psi_p) = J_{\infty}(u_{\infty}),$$

and recalling that we were working with a possible subsequence, then we have that up to a subsequence

$$\lim_{p \rightarrow \infty} C_p = C_{\infty}.$$

Hence, since the limit is independent of the subsequence, we have

$$\lim_{p \rightarrow \infty} C_p = C_{\infty}$$

as required. □

REFERENCES

- [1] David R. Adams and Suzanne Lenhart, *Optimal control of the obstacle for a parabolic variational inequality*, J. Math. Anal. Appl. **268** (2002), no. 2, 602–614.
- [2] ———, *An obstacle control problem with a source term*, Appl. Math. Optim. **47** (2003), no. 1, 79–95.
- [3] D. R. Adams, S. M. Lenhart, and J. Yong, *Optimal control of the obstacle for an elliptic variational inequality*, Appl. Math. Optim. **38** (1998), no. 2, 121–140.
- [4] David R. Adams and Suzanne Lenhart, *An obstacle control problem with a source term*, Appl. Math. Optim. **47** (2003), no. 1, 79–95.
- [5] David R. Adams, Volodymyr Hryniv, and Suzanne Lenhart, *Optimal control of a biharmonic obstacle problem*, Around the research of Vladimir Maz'ya. III, Int. Math. Ser. (N. Y.), vol. 13, Springer, New York, 2010, pp. 1–24.

- [6] John Andersson, Erik Lindgren, and Henrik Shahgholian, *Optimal regularity for the obstacle problem for the p -Laplacian*, J. Differential Equations **259** (2015), no. 6, 2167–2179.
- [7] Gunnar Aronsson, *Minimization problems for the functional $\sup_x F(x, f(x), f'(x))$* , Ark. Mat. **6** (1965), 33–53 (1965).
- [8] Gunnar Aronsson, Michael G. Crandall, and Petri Juutinen, *A tour of the theory of absolutely minimizing functions*, Bull. Amer. Math. Soc. (N.S.) **41** (2004), no. 4, 439–505.
- [9] Maïtine Bergounioux and Suzanne Lenhart, *Optimal control of bilateral obstacle problems*, SIAM J. Control Optim. **43** (2004), no. 1, 240–255.
- [10] Qihong Chen, *Optimal control of semilinear elliptic variational bilateral problem*, Acta Math. Sin. (Engl. Ser.) **16** (2000), no. 1, 123–140.
- [11] Qihong Chen and Yuquan Ye, *Bilateral obstacle optimal control for a quasilinear elliptic variational inequality*, Numer. Funct. Anal. Optim. **26** (2005), no. 3, 303–320.
- [12] Qihong Chen, Delin Chu, and Roger C. E. Tan, *Optimal control of obstacle for quasi-linear elliptic variational bilateral problems*, SIAM J. Control Optim. **44** (2005), no. 3, 1067–1080.
- [13] Daniela Di Donato and Dimitri Mugnai, *On a highly nonlinear self-obstacle optimal control problem*, Appl. Math. Optim. **72** (2015), no. 2, 261–290.
- [14] Radouen Ghanem and Ibtissam Nouri, *Optimal control of high-order elliptic obstacle problem*, Appl. Math. Optim. **76** (2017), no. 3, 465–500.
- [15] Vesa Julin and Petri Juutinen, *A new proof for the equivalence of weak and viscosity solutions for the p -Laplace equation*, Comm. Partial Differential Equations **37** (2012), no. 5, 934–946.
- [16] J. L. Lions, *Optimal Control of Systems Governed by Partial Differential Equations*, 1st ed., Vol. 170, 1971.
- [17] Hongwei Lou, *An optimal control problem governed by quasi-linear variational inequalities*, SIAM J. Control Optim. **41** (2002), no. 4, 1229–1253.
- [18] ———, *On the regularity of an obstacle control problem*, J. Math. Anal. Appl. **258** (2001), no. 1, 32–51.
- [19] P. Lindqvist, *Notes on the Infinity Laplace Equation*, . **17** (2015).
- [20] J. D. Rossi, E. V. Teixeira, and J. M. Urbano, *Optimal regularity at the free boundary for the infinity obstacle problem*, Interfaces Free Bound. **17** (2015), no. 3, 381–398.
- [21] Martin H. Strömqvist, *Optimal control of the obstacle problem in a perforated domain*, Appl. Math. Optim. **66** (2012), no. 2, 239–255.

DEPARTMENT OF MATHEMATICS, HOWARD UNIVERSITY, WASHINGTON, D.C. 20059
E-mail address: henok.mawi@howard.edu, cheikh.ndiaye@howard.edu