

# NORM-CONTROLLED COHOMOLOGY OF TRANSFORMATION GROUPS

MITSUAKI KIMURA

**ABSTRACT.** We generalize the result of Brandenbursky and Marcinkowski for the bounded cohomology of transformation groups to the infinite volume case. To state the result, we introduce the notion of norm-controlled cohomology as a generalization of bounded cohomology. This cohomology theory also provides a framework for relative quasimorphisms.

## 1. INTRODUCTION

Since Gromov's seminal paper [13], bounded cohomology has received considerable attention and has been studied by many authors. However, its computation is difficult in general. In particular, bounded cohomology of third degree or higher had rarely been calculated. Recently, Brandenbursky and Marcinkowski [4] proved that the third bounded cohomology is infinite-dimensional for certain transformation groups of non-positively curved manifolds. In this paper, we generalize the result of Brandenbursky and Marcinkowski by introducing a new cohomology theory.

Let  $M$  be a complete connected Riemannian manifold and  $\mu$  the measure on  $M$  induced by the Riemannian metric. Let  $\text{Homeo}_0(M, \mu)$  denote the identity component of the group of measure-preserving homeomorphisms of  $M$  with compact support. Here, we endow  $\text{Homeo}_0(M, \mu)$  with the compact-open topology. In [4], Brandenbursky and Marcinkowski gave a construction of bounded cohomology classes of  $\text{Homeo}_0(M, \mu)$  for the case when  $M$  is of finite volume. We consider the following problem: what happens if  $M$  is of infinite volume? We observe that their construction gives not necessarily a bounded cochain but a “norm-controlled” cochain (Proposition 4.1). In order to regard this cochain as a certain cohomology class, we introduce a notion of norm-controlled cohomology. Here we review the definition of a norm on a group. Let  $G$  be a group and  $1_G$  denote the identity element of  $G$ . A function  $\nu: G \rightarrow [0, \infty)$  is a *norm* if it satisfies

- (1)  $\nu(gh) \leq \nu(g) + \nu(h)$  for any  $g, h \in G$ ,
- (2)  $\nu(g^{-1}) = \nu(g)$  for any  $g \in G$ ,
- (3)  $\nu(1_G) = 0$ ,
- (4)  $\nu(g) > 0$  if  $g \neq 1_G$ .

If one drops condition (4),  $\nu$  is said to be a *pseudo norm*. If a norm  $\nu$  additionally satisfies the condition  $\nu(ghg^{-1}) = \nu(h)$  for every  $g, h \in G$ ,  $\nu$  is called a *conjugation-invariant norm*. The concept of conjugation-invariant norm was introduced and studied by Burago, Ivanov, and Polterovich in [5]. The definition (of the simplest version) of norm-controlled cohomology is the following.

---

2000 *Mathematics Subject Classification.* 57S05, 20J06.

*Key words and phrases.* bounded cohomology; relative quasimorphisms; diffeomorphism groups.

**Definition 1.1.** Let  $G$  be a group and  $\nu$  a (pseudo) norm on  $G$ . We define  $\bar{C}_\nu^n(G)$  to be the set of (inhomogeneous) cochains  $\bar{c} \in \bar{C}^n(G)$  satisfying the following: there exist  $C, D \geq 0$  such that for all  $g_1, \dots, g_n \in G$ ,

$$|\bar{c}(g_1, \dots, g_n)| \leq C(\nu(g_1) + \dots + \nu(g_n)) + D.$$

The *norm-controlled cohomology*, denoted by  $H_\nu^n(G)$ , is defined to be the cohomology of the cochain complex  $(\bar{C}_\nu^n(G), \bar{\delta})$ . The *exact norm-controlled cohomology*  $EH_\nu^n(G)$  is the kernel of the comparison map  $H_\nu^n(G) \rightarrow H^n(G)$ .

Here we only consider the cohomology with real coefficients. See Section 2 for more details on group cohomology. If  $\nu$  is a bounded norm, then  $H_\nu^n(G)$  is nothing but the bounded cohomology  $H_b^n(G)$  of  $G$ . Note that a similar generalization of bounded cohomology is studied for finitely generated groups and its word length, which is called the *polynomially bounded cohomology* (see [17] for example).

We mainly consider several subgroups of  $\text{Homeo}_0(M, \mu)$ . Let  $\text{Diff}_0(M, \text{vol})$  denote the identity component of the group of diffeomorphisms with compact support which preserve the volume form induced by the Riemannian metric. If  $(M, \omega)$  is a symplectic manifold, let  $\text{Symp}_0(M, \omega)$  denote the identity component of the group of symplectomorphisms with compact support. We regard  $\text{Symp}_0(M, \omega)$  as a subgroup of  $\text{Homeo}_0(M, \mu)$  by considering  $\mu$  induced by the volume form  $\omega^{(\dim M)/2}$ .

In this paper, we consider norm-controlled cohomology with respect to the fragmentation norm defined as follows.

**Definition 1.2.** Let  $U$  be an open subset of  $M$ . Let  $\mathcal{S}_U$  denote the set of elements  $h \in \mathcal{T}_M$  that satisfy the following condition: there exists an isotopy  $\{h_t\}_{0 \leq t \leq 1}$  of  $h$  such that  $\text{supp}(h_t) \subset U$  for every  $t \in [0, 1]$ . We define the *fragmentation norm*  $\nu_U$  with respect to  $U$  on  $\mathcal{T}_M$  by

$$\nu_U(g) = \min \left\{ k \left| \begin{array}{l} \exists f_i \in \mathcal{T}_M, \exists h_i \in \mathcal{S}_U, (i = 1, \dots, k) \\ g = (f_1^{-1} h_1 f_1) \cdots (f_k^{-1} h_k f_k) \end{array} \right. \right\}$$

for  $g \in \mathcal{T}_M$ . If no such decomposition of  $g$  exists, we define  $\nu_U(g) = +\infty$ . We say that  $\nu_U$  is *well-defined on  $\mathcal{T}_M$*  if  $\nu_U(g) < +\infty$  for all  $g \in \mathcal{T}_M$ .

**Example 1.3.** Let  $M$  be a manifold,  $U$  a non-empty open subset of  $M$ , and  $i: U \rightarrow M$  the inclusion.

- Let  $\mathcal{T}_M$  be  $\text{Symp}_0(M, \omega)$  and  $\text{Flux}_\omega: \mathcal{T}_M \rightarrow H_c^1(M; \mathbb{R})/\Gamma_\omega$  denote the symplectic flux homomorphism, where  $\Gamma_\omega$  is the symplectic flux group. Since  $\text{Ker}(\text{Flux}_\omega)$  has the fragmentation property [1] (see also [2]),  $\text{Ker}(\text{Flux}_\omega)$  is normally generated by  $\mathcal{S}_U$  and hence  $\nu_U$  is well-defined on  $\text{Ker}(\text{Flux}_\omega)$ .
- Let  $\mathcal{T}_M$  be  $\text{Diff}_0(M, \text{vol})$  and  $\text{Flux}: \mathcal{T}_M \rightarrow H_c^{n-1}(M; \mathbb{R})/\Gamma$  denote the volume flux homomorphism, where  $\Gamma$  is the volume flux group. Since  $\text{Ker}(\text{Flux})$  has the fragmentation property (an unpublished result of W. Thurston, see Banyaga's book [2]),  $\nu_U$  is well-defined on  $\text{Ker}(\text{Flux})$ .
- Let  $\mathcal{T}_M$  be  $\text{Homeo}_0(M, \mu)$  and  $\tilde{\mathcal{T}}_M$  its universal covering, where  $\mathcal{T}_M$  is endowed with the  $C^0$ -topology. In [9], Fathi defined the homomorphism  $\tilde{\theta}: \tilde{\mathcal{T}}_M \rightarrow H_1(M; \mathbb{R})$  and  $\tilde{\theta}$  induces the mass flow homomorphism  $\theta: \mathcal{T}_M \rightarrow H_1(M; \mathbb{R})/\Gamma$ , where  $\Gamma = \tilde{\theta}(\pi_1(\tilde{\mathcal{T}}_M))$  (see also [18]). Since  $\text{Ker}(\theta)$  has the fragmentation property [9],  $\nu_U$  is well-defined on  $\text{Ker}(\theta)$ .

Henceforth  $H_c^\bullet$  denotes the (de Rham) cohomology with compact support and  $H_c^\bullet$  defines a covariant functor.

**Example 1.4.** Let  $M$ ,  $U$ , and  $i: U \rightarrow M$  be as above. We can think of  $\mathcal{T}_U$  as a subgroup of  $\mathcal{T}_M$  by extending the diffeomorphism on  $U$  to  $M$ .

- Let  $\mathcal{T}_M = \text{Symp}_0(M, \omega)$ . If  $i^*: H_c^1(U; \mathbb{R}) \rightarrow H_c^1(M; \mathbb{R})$  is surjective, we can see that  $\nu_U$  is well-defined on  $\mathcal{T}_M$  as follows. For  $g \in \mathcal{T}_M$ , there exists  $h \in \mathcal{T}_U \subset \mathcal{T}_M$  such that  $\text{Flux}_\omega(g) = \text{Flux}_\omega(h)$  since the flux homomorphism is surjective and by the assumption of  $i^*$ . Thus  $g = (gh^{-1})h$  is written as a product of the conjugation of the elements of  $\mathcal{S}_U$  since  $h \in \mathcal{S}_U$  and  $gh^{-1} \in \text{Ker}(\text{Flux}_\omega)$ .

$$\begin{array}{ccccc} \pi_1(\mathcal{T}_M) & \longrightarrow & \widetilde{\mathcal{T}}_M & \longrightarrow & \mathcal{T}_M \\ \downarrow \widetilde{\text{Flux}}_\omega|_{\pi_1(\mathcal{T}_M)} & & \downarrow \widetilde{\text{Flux}}_\omega & & \downarrow \text{Flux}_\omega \\ \Gamma_\omega & \longrightarrow & H_c^1(M; \mathbb{R}) & \longrightarrow & H_c^1(M; \mathbb{R})/\Gamma_\omega \end{array}$$

- Let  $\mathcal{T}_M = \text{Diff}_0(M, \text{vol})$ . If  $i^*: H_c^{n-1}(U; \mathbb{R}) \rightarrow H_c^{n-1}(M; \mathbb{R})$  is surjective, we can see that  $\nu_U$  is well-defined on  $\mathcal{T}_M$  by the same argument.
- Let  $\mathcal{T}_M = \text{Homeo}_0(M, \mu)$ . If  $i_*: H_1(U; \mathbb{R}) \rightarrow H_1(M; \mathbb{R})$  is surjective, we can see that  $\nu_U$  is well-defined on  $\mathcal{T}_M$  by the same argument.

Let  $\pi_M$  denote the quotient group of the fundamental group of  $M$  by its center. The main theorem of this paper is the following.

**Theorem 1.5.** *Let  $\mathcal{T}_M$  be  $\text{Homeo}_0(M, \mu)$ ,  $\text{Diff}_0(M, \text{vol})$ , or  $\text{Symp}_0(M, \omega)$ . Assume that there exists an open subset  $U$  of  $M$  with finite volume such that  $\nu_U$  is well-defined on  $\mathcal{T}_M$ . If either*

- (1)  $\pi_M$  surjects onto  $F_2$  or
- (2)  $\pi_M$  is an acylindrically hyperbolic group,

then

$$\dim_{\mathbb{R}} EH_{\nu_U}^n(\mathcal{T}_M) \geq \dim_{\mathbb{R}} \overline{EH}_b^n(F_2)$$

for every degree  $n \geq 1$ .

Here,  $\overline{EH}_b^n(G)$  denotes the reduced exact bounded cohomology of  $G$  (see Section 2.2). If  $M$  is of finite volume and  $U = M$ , this implies (a weak version of) the result of [4, Theorem A and B]. It is known that most 3-manifold groups are acylindrically hyperbolic [16]. On the other hand, if  $M$  is 3-dimensional and  $\pi_1(M)$  is finitely generated, there exists a 3-dimensional compact submanifold  $C$  such that the inclusion  $C \rightarrow M$  is a homotopy equivalence by the Scott core theorem [21]. Thus we have many examples that satisfy the assumption of the above theorem. For more details about acylindrically hyperbolic groups, see [19].

We also define a variant of the norm-controlled cohomology  $H_{(d)}^n(G, \nu)$  for a non-negative integer  $d$  such that  $H_{(0)}^n(G, \nu) = H_\nu^n(G)$ . We will observe several basic properties of this cohomology, such as functoriality (Proposition 3.9). The cochain obtained by Burandenbursky–Marcinkowski’s construction for infinite volume manifolds is regarded as an element of this cohomology too.

As a corollary of Theorem 1.5, we obtain the following result.

**Corollary 1.6.** *Suppose  $M$  and  $U$  satisfy the assumption in Theorem 1.5. Then  $EH_{(d)}^3(\mathcal{T}_M, \nu_U)$  is uncountably infinite-dimensional for  $d = 0, 1, 2$ .*




FIGURE 1. Relation to other notions

Our cohomology is also motivated by relative quasimorphisms. Let  $G$  be a group and  $\nu: G \rightarrow [0, \infty)$  a conjugation-invariant norm on  $G$ . A function  $\phi: G \rightarrow \mathbb{R}$  is called a *relative quasimorphism* with respect to a conjugation-invariant norm  $\nu$  if there exists a constant  $C \geq 0$  such that

$$(1.1) \quad |\phi(gh) - \phi(g) - \phi(h)| \leq C \cdot \min\{\nu(g), \nu(h)\}$$

for all  $g, h \in G$ . Relative quasimorphisms first appeared in the paper of Entov and Polterovich [8]. They appear mainly in symplectic geometry, but they are also found in another context by the author [15]. For a quasimorphism  $\phi$ , its coboundary  $\bar{\delta}\phi$  is considered as an element of the second bounded cohomology class. Similarly, a relative quasimorphism  $\phi: G \rightarrow \mathbb{R}$  with respect to a conjugation-invariant norm  $\nu$  can be seen as an element of  $H_{(1)}^2(G, \nu)$  (see Figure 1). We hope that our cohomology has applications to the study of relative quasimorphisms.

We note that the idea of the technique in Proposition 4.1 comes from the construction of relative quasimorphisms by Kawasaki [14] and Brandenbursky and Kędra [3]. Moreover, our construction also can be seen as a generalization of [3]. We remark that both [3] and [4] can be seen as a generalization of the construction of Gambaudo–Ghys [12] and Polterovich [20] (Figure 2).




FIGURE 2. Relation to previous constructions

## 2. PRELIMINARY ON GROUP COHOMOLOGY

In this section, we review the definitions and terminology of (bounded) cohomology of groups. We refer to [6, 10] for more information on this topic. Throughout this paper, we only consider the cohomology with real coefficients.

**2.1. Group cohomology.** Let  $G$  be a group. We consider the space of (inhomogeneous)  $n$ -cochains

$$\bar{C}^n(G) = \{\bar{c}: G^n \rightarrow \mathbb{R}\},$$

and the coboundary map  $\bar{\delta} : \bar{C}^{n-1}(G) \rightarrow \bar{C}^n(G)$  defined by

$$\bar{\delta}\bar{c}(g_1, \dots, g_n) = \bar{c}(g_2, \dots, g_n) + \sum_{i=1}^{n-1} (-1)^i \bar{c}(g_1, \dots, g_i g_{i+1}, \dots, g_n) + (-1)^n \bar{c}(g_1, \dots, g_{n-1})$$

for  $\bar{c} \in \bar{C}^n(G)$  and  $g_1, \dots, g_n \in G$ . The cohomology of this cochain complex is called the *(group) cohomology* of  $G$  and denoted by  $H^n(G)$ .

There is another definition of this cohomology. A map  $c : G^{n+1} \rightarrow \mathbb{R}$  is said to be *homogeneous* if  $c(hg_0, \dots, hg_n) = c(g_0, \dots, g_n)$  for every  $g_0, \dots, g_n, h \in G$ . The space of (homogeneous)  $n$ -cochains is

$$C^n(G) = \{c : G^{n+1} \rightarrow \mathbb{R} \mid c \text{ is homogeneous}\},$$

and the coboundary map  $\delta : C^{n-1}(G) \rightarrow C^n(G)$  is defined by

$$\delta c(g_0, \dots, g_n) = \sum_{i=0}^n (-1)^i c(g_0, \dots, \widehat{g_i}, \dots, g_n)$$

for  $c \in C^n(G)$  and  $g_0, \dots, g_n \in G$ , where  $\widehat{g_i}$  means that we omit the entry  $g_i$ . The cohomology of  $(C^n(G), \delta)$  also defines  $H^n(G)$ .

The correspondence between an inhomogeneous cochain  $\bar{c} \in \bar{C}^n(G)$  and homogeneous one  $c \in C^n(G)$  is the following:

$$(2.1) \quad \bar{c}(g_1, g_2, \dots, g_n) = c(1, g_1, g_1 g_2, \dots, g_1 g_2 \dots g_n),$$

$$(2.2) \quad c(g_0, g_1, \dots, g_n) = \bar{c}(g_0^{-1} g_1, g_1^{-1} g_2, \dots, g_{n-1}^{-1} g_n).$$

We call a cochain  $c \in C^n(G)$  *alternating* if

$$c(g_{\sigma(0)}, \dots, g_{\sigma(n)}) = \text{sgn}(\sigma) c(g_0, \dots, g_n)$$

for any  $g_0, \dots, g_n \in G$  and  $\sigma \in \mathfrak{S}_{n+1}$ , where  $\text{sgn}(\sigma) \in \{\pm 1\}$  is the sign of  $\sigma$ . Let  $C_{\text{alt}}^n(G)$  denote the set of alternating  $n$ -cochains. Then  $(C_{\text{alt}}^n(G), \delta)$  is a subcomplex of  $(C^n(G), \delta)$ . It is known that the cohomology of  $(C_{\text{alt}}^n(G), \delta)$  also defines  $H^n(G)$ .

**2.2. Bounded cohomology.** We review the definition of bounded cohomology. We only mention the inhomogeneous case but the homogenous case is defined similarly. If we consider the subcomplex

$$\bar{C}_b^n(G) = \{\bar{c} : G^n \rightarrow \mathbb{R} \mid \bar{c} \text{ is bounded}\}$$

of  $\bar{C}^n$ , the homology of the complex  $(\bar{C}_b^n(G), \bar{\delta})$  is called the *bounded cohomology* of  $G$  and denoted by  $H_b^n(G)$ . The natural inclusion  $\bar{C}_b^n(G) \rightarrow \bar{C}^n(G)$  induces the homomorphism  $H_b^n(G) \rightarrow H^n(G)$  called the *comparison map*. The kernel of the comparison map  $H_b^n(G) \rightarrow H^n(G)$  is called the *exact bounded cohomology* and denoted by  $EH_b^n(G)$ .

For a cochain  $\bar{c} \in \bar{C}_b^n(G)$ , we define the *norm*  $\|\bar{c}\|$  of  $\bar{c}$  by

$$\|\bar{c}\| = \sup_{g_1, \dots, g_n \in G} |\bar{c}(g_1, \dots, g_n)|.$$

This norm induces a natural seminorm  $\|\cdot\|$  on  $H_b^n(G)$  defined by

$$\|\alpha\| = \inf \|\bar{c}\|$$

for  $\alpha \in H_b^n(G)$ , where the infimum is taken over all cochains  $\bar{c} \in \bar{C}_b^n(G)$  representing the cohomology class  $\alpha$ .

Let  $N^n(G)$  denote the norm zero subspace of  $H_b^n(G)$ , i.e.,

$$N^n(G) = \{\alpha \in H_b^n(G) \mid \|\alpha\| = 0\}.$$

The *reduced cohomology*  $\overline{H}_b^n(G)$  is defined by the quotient  $H_b^n(G)/N^n(G)$ . The *reduced exact cohomology*  $\overline{EH}_b^n(G)$  is defined by  $EH_b^n(G)/EN(G)$ , where  $EN^n(G) = N^n(G) \cap EH_b^n(G)$ .

We can consider the homogeneous complex  $C_b^\bullet(G)$ , alternating homogenous and inhomogeneous subcomplex  $C_{b,\text{alt}}^\bullet(G)$  and  $\bar{C}_{b,\text{alt}}^\bullet(G)$ , and they also define the cohomology  $H_b^\bullet(G)$  endowed with its natural seminorm.

### 3. NORM-CONTROLLED COHOMOLOGY

**3.1. Definition of the cohomology.** We refer to a group equipped with a pseudo norm as a *normed group*. We define the norm-controlled cohomology  $H_{(d)}^\bullet(G, \nu)$  for a normed group  $(G, \nu)$  and a non-negative integer  $d$ .

**Definition 3.1.** For a cochain  $\bar{c} \in \bar{C}^n(G)$  and a function  $\mu: G^n \rightarrow [0, \infty)$ , we say that  $\bar{c}$  is *Lipschitz with respect to  $\mu$*  if there exist constants  $C, D \geq 0$  such that for every  $g_1, \dots, g_n \in G$

$$|\bar{c}(g_1, \dots, g_n)| \leq C \cdot \mu(g_1, \dots, g_n) + D.$$

**Definition 3.2.** For a normed group  $(G, \nu)$  and non-negative integers  $n$  and  $d$ , we define  $\bar{C}_{(d)}^n(G, \nu)$  as follows.

- If  $n > d$ , we define  $\bar{C}_{(d)}^n(G, \nu)$  as the set of Lipschitz cochains  $\bar{c} \in \bar{C}^n(G)$  with respect to  $\nu_{(n,d)}$ , where  $\nu_{(n,d)}: G^n \rightarrow [0, \infty)$  is defined by

$$\begin{aligned} \nu_{(n,d)}(g_1, \dots, g_n) &= \min_{\substack{I \subset \{1, \dots, n\} \\ \#I = n-d}} \left\{ \sum_{i \in I} \nu(g_i) \right\} \\ &= \min_{1 \leq i_1 < \dots < i_d \leq n} \left\{ \nu(g_1) + \dots + \widehat{\nu(g_{i_1})} + \dots + \widehat{\nu(g_{i_d})} + \dots + \nu(g_n) \right\}. \end{aligned}$$

- If  $d \geq n$ , we define  $\bar{C}_{(d)}^n(G, \nu) = \bar{C}_b^n(G, \nu)$ .

Note that  $\bar{c} \in \bar{C}_{(0)}^n(G, \nu)$  implies

$$|\bar{c}(g_1, \dots, g_n)| \leq C \cdot \{\nu(g_1) + \dots + \nu(g_n)\} + D$$

and  $\bar{c} \in \bar{C}_{(n-1)}^n(G, \nu)$  implies

$$|\bar{c}(g_1, \dots, g_n)| \leq C \cdot \min\{\nu(g_1), \dots, \nu(g_n)\} + D.$$

**Lemma 3.3.** For any integer  $d \geq 0$ ,  $(\bar{C}_{(d)}^n(G, \nu), \bar{\delta})$  is a subcomplex of  $(\bar{C}^n(G), \bar{\delta})$ .

*Proof.* It is sufficient to prove that  $\bar{\delta}(\bar{C}_{(d)}^{n-1}(G, \nu)) \subset \bar{C}_{(d)}^n(G, \nu)$  for the case  $n-1 > d$ .

Let  $g_1, \dots, g_n$  be elements in  $G$ . It is easy to see that

$$\begin{aligned} \nu_{(n-1,d)}(g_2, \dots, g_n) &\leq \nu_{(n,d)}(g_1, \dots, g_n), \\ \nu_{(n-1,d)}(g_1, \dots, g_{n-1}) &\leq \nu_{(n,d)}(g_1, \dots, g_n). \end{aligned}$$

Since  $\nu(g_i g_{i+1}) \leq \nu(g_i) + \nu(g_{i+1})$ ,

$$\nu_{(n-1,d)}(g_1, \dots, g_i g_{i+1}, \dots, g_n) \leq \nu_{(n,d)}(g_1, \dots, g_i, g_{i+1}, \dots, g_n)$$

for  $i = 1, 2, \dots, n-1$ . Therefore, for  $\bar{c} \in \bar{C}_{(d)}^{n-1}(G, \nu)$ ,

$$\begin{aligned} & |\bar{\delta}\bar{c}(g_1, \dots, g_n)| \\ & \leq |\bar{c}(g_2, \dots, g_n)| + \sum_{i=1}^{n-1} |\bar{c}(g_1, \dots, g_i g_{i+1}, \dots, g_n)| + |\bar{c}(g_1, \dots, g_{n-1})| \\ & \leq (n+1)\{C \cdot \nu_{(n,d)}(g_1, \dots, g_n) + D\}. \end{aligned} \quad \square$$

**Definition 3.4.** For a normed group  $(G, \nu)$  and an integer  $d \geq 0$ , we define the *norm-controlled cohomology*  $H_{(d)}^\bullet(G, \nu)$  of level  $d$  to be the cohomology of the cochain complex  $(\bar{C}_{(d)}^n(G, \nu), \bar{\delta})$ .

Note that the complexes  $\{\bar{C}_{(d)}^n(G, \nu)\}_{n,d}$  can be seen as a filtered complex, i.e.,  $\bar{C}_{(d)}^n(G, \nu) \subset \bar{C}_{(d')}^n(G, \nu)$  if  $d \geq d'$ .

By the correspondence (2.2), we can define the homogeneous norm-controlled cochain complex  $C_{(d)}^n(G, \nu)$  as the set of cochains  $c \in C^n(G)$  which satisfies the following: there exist constants  $C, D \geq 0$  such that

$$|c(g_1, \dots, g_n)| \leq C \cdot \min_{\substack{I \subset \{1, \dots, n\} \\ \#I = n-d}} \left\{ \sum_{i \in I} \nu(g_{i-1}^{-1} g_i) \right\} + D.$$

We can also define the inhomogeneous (resp. homogeneous) alternating subcomplex  $\bar{C}_{(d), \text{alt}}^\bullet(G, \nu)$  (resp.  $C_{(d), \text{alt}}^\bullet(G, \nu)$ ) and they also define the cohomology  $H_{(d)}^\bullet(G, \nu)$ .

**Example 3.5.** Let  $\mathbb{Z}^n$  be the free abelian group of rank  $n$ . For a positive integer  $l \leq n$ , define a (pseudo) norm  $\nu_l$  on  $\mathbb{Z}^n$  by

$$\nu_l(m_1, \dots, m_l, \dots, m_n) = |m_1| + \dots + |m_l|$$

for  $m_1, \dots, m_n \in \mathbb{Z}$ . Now we compute  $H_{\nu_l}^1(\mathbb{Z}^n)$ . Note that  $H_\nu^1(G) = H_{(0)}^1(G, \nu) = \text{Ker}(\bar{\delta}: \bar{C}_{(d)}^1 \rightarrow \bar{C}_{(d)}^2)$  is the set of Lipschitz homomorphisms with respect to  $\nu$ .

We define a homomorphism  $\phi_i: \mathbb{Z}^n \rightarrow \mathbb{R}$  by  $\phi_i(m_1, \dots, m_l, \dots, m_n) = m_i$ .  $\text{Hom}(\mathbb{Z}^n, \mathbb{R}) \cong \mathbb{R}^n$  is generated by  $\phi_1, \dots, \phi_n$ . It is easy to see that  $\phi_i$  is Lipschitz with respect to  $\nu_l$  for  $i \leq l$  and not for  $i > l$ . Thus,  $H_{\nu_l}^1(\mathbb{Z}^n)$  is generated by  $\phi_1, \dots, \phi_l$  and isomorphic to  $\mathbb{R}^l$ .

**Example 3.6.** Let  $\phi$  be an unbounded relative quasimorphism (see (1.1)) on a group  $G$  with respect to a conjugation-invariant norm  $\nu$ . If  $\phi$  is at infinite distance from homomorphisms, then  $\bar{\delta}\phi$  defines a non-trivial cohomology class in  $EH_{(1)}^2(G, \nu)$ . Thus, for the following cases,  $EH_b^2(G)$  is trivial but  $EH_{(1)}^2(G, \nu)$  is non-trivial for a certain norm  $\nu$ .

- $G$  is the identity component of the group of symplectomorphisms  $\text{Symp}_0^c(\mathbb{R}^{2n}, \omega_0)$  of the standard symplectic space  $(\mathbb{R}^{2n}, \omega_0)$  with compact support [14].
- $G$  is the infinite braid group  $B_\infty$  [15].
- $G$  is the Hamiltonian diffeomorphism group  $\text{Ham}(T^*\Sigma_g \times \mathbb{R}^{2n})$  of  $T^*\Sigma_g \times \mathbb{R}^{2n}$ , where  $\Sigma_g$  is a closed surface of genus  $g > 1$  [3].

**3.2. Functoriality.** We show that our cohomology is a functor for a certain category. Recall that a normed group  $(G, \nu)$  is a pair of a group  $G$  and a (pseudo) norm  $\nu$  on  $G$ .

**Definition 3.7.** Let  $(G, \nu_G)$  and  $(H, \nu_H)$  be normed groups. A homomorphism  $\phi: G \rightarrow H$  is said to be *Lipschitz* if there exist  $C, D \geq 0$  such that for all  $g \in G$ ,

$$\nu_H(\phi(g)) \leq C \cdot \nu_G(g) + D.$$

**Definition 3.8.** We define the *category  $\mathbf{NGrp}$  of normed groups* as follows.

- The objects  $Ob(\mathbf{NGrp})$  are normed groups.
- The morphisms  $Mor(\mathbf{NGrp})$  are Lipschitz homomorphisms  $\phi: (G, \nu_G) \rightarrow (H, \nu_H)$  between normed groups  $(G, \nu_G)$  and  $(H, \nu_H)$ .

The composition of morphisms is the composition of group homomorphisms, and hence the associativity holds. For every  $(G, \nu) \in Ob(\mathbf{NGrp})$ , there exists the identity  $\text{id}_G: (G, \nu) \rightarrow (G, \nu)$  in  $Mor(\mathbf{NGrp})$ . Hence  $\mathbf{NGrp}$  is a category.

Let  $H_{(d)}^n$  denote the correspondence from a normed group  $(G, \nu)$  to  $H_{(d)}^n(G, \nu)$ .

**Proposition 3.9.** *The correspondence  $H_{(d)}^n$  is a contravariant functor from the category of normed groups  $\mathbf{NGrp}$  to the category of real vector spaces  $\mathbf{Vect}_{\mathbb{R}}$ .*

If two norms on a same group are bi-Lipschitz, then they define the same norm-controlled cohomology.

**Corollary 3.10.** *Let  $G$  be a group with (pseudo) norms  $\nu_1$  and  $\nu_2$ . If both  $\text{id}_G: (G, \nu_1) \rightarrow (G, \nu_2)$  and  $\text{id}_G: (G, \nu_2) \rightarrow (G, \nu_1)$  are Lipschitz, then  $\text{id}_G^*: H_{(d)}^n(G, \nu_2) \rightarrow H_{(d)}^n(G, \nu_1)$  is an isomorphism.*

The proof of Proposition 3.9 and Corollary 3.10 are elementary and will be omitted.

#### 4. COCYCLES ON TRANSFORMATION GROUPS

**4.1. Brandenbursky–Marcinkowski construction.** We briefly review the construction of Brandenbursky and Marcinkowski [4]. Let  $M$  be a finite volume complete Riemannian manifold and  $\mu$  the measure on  $M$  associated with the Riemannian structure. Fix a base point  $z \in M$ . Let  $\text{Homeo}_0(M, \mu)$  denote the identity component of the group of homeomorphisms of  $M$  with compact support that preserve the measure  $\mu$ . Recall that  $\pi_M$  denotes the quotient group  $\pi_1(M, z)/Z(\pi_1(M, z))$ , where  $Z(G)$  denotes the center of a group  $G$ .

For a subgroup  $\mathcal{T}_M$  of  $\text{Homeo}_0(M, \mu)$ , they constructed a map  $\Gamma_b^\bullet: H_b^\bullet(\pi_M) \rightarrow H_b^\bullet(\mathcal{T}_M)$  as follows. Let  $C$  denote the cut locus of  $z$ , i.e., the closure of the set of all points  $x \in M$  that have two or more distinct shortest paths in  $M$  from  $x$  to  $z$ . For  $x \in M$  and  $g \in \mathcal{T}_M$  such that  $x \notin C$  and  $g(x) \notin C$ , we define  $\gamma(g, x) \in \pi_1(M, z)$  by the concatenation of the minimizing geodesic between  $z$  and  $x$ , the path defined by  $\{g_t(x)\}_{0 \leq t \leq 1}$ , where  $\{g_t\}_{0 \leq t \leq 1}$  is an isotopy of  $g$  with  $g_0 = \text{id}$  and  $g_1 = g$ , and the minimizing geodesic between  $g(x)$  and  $z$ . Then  $\gamma(g, x)$  is defined uniquely up to center for any choice of isotopies. Thus it defines an element of  $\pi_M$ .

Since the measure  $\mu(C)$  of the cut locus  $C$  is zero and the map  $\gamma(f, \cdot): M \rightarrow \pi_M$  has essentially finite image for  $f \in \mathcal{T}_M$ , we can define the map  $\Phi_{BM}: C^n(\pi_M) \rightarrow C^n(\mathcal{T}_M)$  by

$$\Phi_{BM}(c)(g_0, \dots, g_n) = \int_M c(\gamma(g_0, x), \dots, \gamma(g_n, x)) d\mu(x)$$

for  $c \in C^n(\pi_M)$  and  $g_0, \dots, g_n \in \mathcal{T}_M$ . The map  $\Gamma^n: H^n(\pi_M) \rightarrow H^n(\mathcal{T}_M)$  is defined as the induced map from  $\Phi_{BM}$ . If  $c \in C_b^n(\pi_M)$  is a bounded cochain, then  $\Phi_{BM}(c)$



is also a bounded cochain since

$$|\Phi_{BM}(c)(g_0, \dots, g_n)| \leq \|c\| \cdot \text{vol}(M) < +\infty$$

for any  $g_0, \dots, g_n \in \pi_M$ . Hence  $\Phi_{BM}$  induces the map  $\Gamma_b^n: H_b^n(\pi_M) \rightarrow H_b^n(\mathcal{T}_M)$ . We also obtain the map of the exact part  $E\Gamma_b^n: EH_b^n(\pi_M) \rightarrow EH_b^n(\mathcal{T}_M)$ .

**4.2. Main result.** We consider the above construction for the case that  $M$  has infinite volume. In this case, the map  $\Phi_{BM}$  is well-defined on  $C_{b,\text{alt}}(\pi_M)$  since we consider homeomorphisms with compact support. On the other hand, the image  $\Phi_{BM}(c)$  of a bounded cochain  $c \in C_{b,\text{alt}}(\pi_M)$  might not be a bounded cochain. We prove that, however, the image is a norm-controlled cochain with respect to a fragmentation norm (Proposition 4.1).

**Proposition 4.1.** *For  $c \in C_{b,\text{alt}}^n(\pi_M)$ , there exists  $C \geq 0$  such that*

$$|\Phi_{BM}(c)(g_0, \dots, g_n)| \leq C \cdot \min_{0 \leq i < j \leq n} \{\nu_U(g_i^{-1}g_j)\}.$$

for all  $g_0, \dots, g_n \in \mathcal{T}_M$ . In particular,  $\Phi_{BM}(c) \in C_{(n-1),\text{alt}}^n(\mathcal{T}_M, \nu_U)$ .

*Proof.* We fix  $i$  and  $j$  ( $0 \leq i < j \leq n$ ). Assume that  $\nu_U(g_i^{-1}g_j) = m$ . Then we can write  $g_i^{-1}g_j = (f_1^{-1}h_1f_1) \dots (f_m^{-1}h_mf_m)$ , where  $h_k \in \mathcal{S}_U$  and  $f_k \in \mathcal{T}_M$  for  $k = 1, \dots, m$ . Take an isotopy  $\{(g_i)_t\}_t$  of  $g_i$  and isotopies  $\{(h_1)_t\}_t, \dots, \{(h_m)_t\}_t$  for  $h_1, \dots, h_m$  such that  $\text{supp}((h_k)_t) \subset U$  for every  $t \in [0, 1]$  and  $k = 1, \dots, m$ . We define  $(g_j)_t = (g_i)_t(f_1^{-1}(h_1)_tf_1) \dots (f_m^{-1}(h_m)_tf_m)$ . Then  $\{(g_j)_t\}_t$  is an isotopy of  $g_j$ . Set

$$U_{ij} = \bigcup_{0 \leq t \leq 1} \text{supp}((g_i)_t^{-1}(g_j)_t) = \bigcup_{0 \leq t \leq 1} \text{supp}((f_1^{-1}(h_1)_tf_1) \dots (f_m^{-1}(h_m)_tf_m)).$$

Note that  $U_{ij} \subset f_1(U) \cup \dots \cup f_m(U)$ . If  $x \notin U_{ij}$ ,  $(g_i)_t(x) = (g_j)_t(x)$  for every  $t \in [0, 1]$ . Thus  $\gamma(g_i, x) = \gamma(g_j, x) \in \pi_M$ . Since  $c$  is alternating,  $c(\gamma(g_0, x), \dots, \gamma(g_n, x)) = 0$ . Therefore,

$$|\Phi_{BM}(c)(g_0, \dots, g_n)| \leq \text{vol}(U_{ij}) \cdot \|c\| \leq m \cdot \text{vol}(U) \cdot \|c\|.$$

Since we can arbitrarily take  $i$  and  $j$ , the inequality holds for  $C = \text{vol}(U) \cdot \|c\|$ .  $\square$

*Remark 4.2.* For  $d \leq n-1$ , the map  $\Phi_{BM}: C_{b,\text{alt}}^n(\pi_M) \rightarrow C_{(d),\text{alt}}^n(\mathcal{T}_M, \nu_U)$  is well-defined. However, if  $d = n-1$ ,  $\Phi_{BM}$  does not induce the map  $H_b^n(\pi_M) \rightarrow H_{(n-1)}^n(\mathcal{T}_M, \nu_U)$  because the image of  $\bar{B}_{(n-1)}^n(\pi_M)$  might not be in  $\bar{B}_{(n-1)}^n(\mathcal{T}_M)$ . On the other hand, if  $d < n-1$ ,  $\Phi_{BM}$  induces  $H_b^n(\pi_M) \rightarrow H_{(d)}^n(\mathcal{T}_M, \nu_U)$ . Especially, if  $d = 0$ , then  $\Phi_{BM}: C_{b,\text{alt}}^n(\pi_M) \rightarrow C_{\nu_U,\text{alt}}^n(\mathcal{T}_M)$  induces  $H_b^n(\pi_M) \rightarrow H_{\nu_U}^n(\mathcal{T}_M)$  for any  $n \geq 2$ .

We prove the following key lemma which corresponds to [4, Lemma 3.3]. Let  $a$  and  $b$  be generators of  $F_2$ .

**Lemma 4.3.** *Let  $U \subset M$  be an open subset such that  $\nu_U$  is well-defined on  $\mathcal{T}_M$ . Assume that there exists an injection  $i: F_2 \rightarrow \pi_M$ . Let  $\alpha$  and  $\beta$  be two loops in  $M$  representing  $i(a)$  and  $i(b)$ . Suppose that  $\alpha$  and  $\beta$  are contained in  $U$ . Then there exist a constant  $\Lambda > 0$  and a family of Lipschitz homomorphisms  $\{\rho_\epsilon: (F_2, \nu_0) \rightarrow (\mathcal{T}_M, \nu_U)\}_{\epsilon \in (0,1)}$  such that*

$$\lim_{\epsilon \rightarrow +0} \|\rho_\epsilon^*(E\Gamma_{\nu_U}(c)) - \Lambda i^*(c)\| = 0$$

for every  $c \in EH_{\nu_U}^n(\pi_M)$ .

Here,  $\nu_0: F_2 \rightarrow [0, \infty)$  is the trivial norm defined by

$$\nu_0(w) = \begin{cases} 0 & (w = 1_{F_2}), \\ 1 & (w \neq 1_{F_2}). \end{cases}$$

The maps  $i^*$  and  $\rho_\epsilon^*$  represent the induced maps  $i^*: EH_b^n(\pi_M) \rightarrow EH_b^n(F_2)$  and  $\rho_\epsilon^*: EH_{\nu_U}^n(\pi_M) \rightarrow EH_{\nu_0}^n(F_2) = EH_b^n(F_2)$ .

*Proof.* Our proof is similar to [4, Lemma 3.3]. Let  $N(\alpha)$  denote a tubular neighborhood of  $\alpha$  in  $U$  and take a diffeomorphism  $n_\alpha: N(\alpha) \rightarrow S^1 \times B^{n-1}(1)$ . Here  $B^{n-1}(r)$  denotes the  $(n-1)$ -ball in  $\mathbb{R}^n$  with radius  $r$ . Let  $A_\epsilon(\alpha)$  denote  $n_\alpha^{-1}(S^1 \times B^{n-1}(1-\epsilon))$ . We define an element  $\rho_\epsilon(a) \in \mathcal{T}_M$  which “rotates” every point in  $A_\epsilon$  one lap in the direction of  $S^1$  and fixes outside of  $N(\alpha)$  (see [4] for more details). Similarly, we define  $N(\beta) \subset U$ ,  $B_\epsilon$  and  $\rho_\epsilon(b) \in \mathcal{T}_M$  (see Figure 3). Thus we obtain the representation  $\rho_\epsilon: F_2 \rightarrow \mathcal{T}_M$ . Since  $\text{supp}(\rho_\epsilon(w))$  is contained in  $U$  for any  $w \in F_2$ , the map  $\rho_\epsilon: (F_2, \nu_0) \rightarrow (\mathcal{T}_M, \nu_U)$  is a Lipschitz homomorphism. By the functoriality of the correspondence  $H_{(0)}^n$  (Proposition 3.9), the map  $\rho_\epsilon^*: EH_{\nu_U}^n(\mathcal{T}_M) \rightarrow EH_b^n(\pi_M)$  is well-defined.




FIGURE 3. loops  $\alpha$  and  $\beta$  and their neighborhood

For  $w_0, \dots, w_n \in F_2$ , we have


$$\rho_\epsilon^*(E\Gamma_{\nu_U}(c))(w_0, \dots, w_n) = \int_M c(\gamma(\rho_\epsilon(w_0), x), \dots, \gamma(\rho_\epsilon(w_n), x)) d\mu(x).$$

Let  $B_\epsilon(\alpha)$  and  $B_\epsilon(\beta)$  denote  $N(\alpha) - A_\epsilon(\alpha)$  and  $N(\beta) - A_\epsilon(\beta)$  respectively. We calculate this integral by decomposing  $M$  into 5 parts;  $A_\epsilon := A_\epsilon(\alpha) \cap A_\epsilon(\beta)$ ,  $A_\epsilon^a := A(\alpha) - N(\beta)$ ,  $A_\epsilon^b := A_\epsilon(\beta) - N(\alpha)$ ,  $B_\epsilon := B_\epsilon(\alpha) \cup B_\epsilon(\beta)$ , and their exterior  $M - (N(\alpha) \cup N(\beta))$  (see Figure 4).

The exterior part is 0 and it turns out that  $A_\epsilon^a$  and  $A_\epsilon^b$  part are also 0. The  $A_\epsilon$  part is calculated to be  $\mu(A_\epsilon)i^*(c)$  and the  $B_\epsilon$  is bounded by  $\mu(B_\epsilon)\|c\|$ . Hence the claim follows from  $\mu(A_\epsilon) \xrightarrow{\epsilon \rightarrow +0} \mu(N(\alpha) \cap N(\beta)) > 0$  and  $\mu(B_\epsilon) \xrightarrow{\epsilon \rightarrow +0} 0$ .  $\square$

We give proof of the main result. Our proof is similar to the proof of Theorem A and B in [4].

*Proof of Theorem 1.5.* First, we prove case (1). Let  $p: \pi_M \rightarrow F_2$  be a surjection. Assume that  $\dim(M) \geq 3$ . Then there exists an injection  $i: F_2 \rightarrow \pi_M$  such that  $p \circ i = \text{id}_{F_2}$ . If  $\dim(M) = 2$ , we can find an injection  $i: F_2 \rightarrow \pi_M$ , and there exists a retraction  $p: \pi_M \rightarrow F_2$  of  $i$ ; we use this  $p$  instead of the given  $p$ . If necessary, we retake  $U$  to be containing  $\alpha$  and  $\beta$  in Lemma 4.3.

FIGURE 4. the decomposition of  $M$  into 5 parts

$$\begin{array}{ccc}
 EH_b^n(\pi_M) & \xrightarrow{E\Gamma_{\nu_U}} & EH_{\nu_U}^n(\mathcal{T}_M) \\
 \uparrow p^* & \swarrow i^* & \searrow \rho_\epsilon^* \\
 EH_b^n(F_2) & & 
 \end{array}$$

Note that  $EH_{\nu_U}^n(\mathcal{T}_M) \supset \text{Im}(E\Gamma_{\nu_U} \circ p^*) \cong H_b^n(F_2) / \text{Ker}(E\Gamma_{\nu_U} \circ p^*)$ . For  $d \in \text{Ker}(E\Gamma_{\nu_U} \circ p^*)$ , set  $c = p^*(d) \in EH_b^n(\pi_M)$ . Since  $i^* \circ p^* = \text{id}$ ,  $i^*(c) = i^* \circ p^*(d) = d$ .

By Lemma 4.3, there exist  $\Lambda > 0$  and a family of representation  $\{\rho_\epsilon\}$  such that

$$\lim_{\epsilon \rightarrow +0} \|\rho_\epsilon^*(E\Gamma_{\nu_U}(c)) - \Lambda i^*(c)\| = 0.$$

Since  $E\Gamma_{\nu_U}(c) = E\Gamma_{\nu_U} \circ p^*(d) = 0$ ,  $\|i^*(c)\| = \|d\| = 0$ . Hence  $\text{Ker}(E\Gamma_{\nu_U} \circ p^*) \subset EN^n(F_2)$ . Therefore,

$$\dim_{\mathbb{R}} \left( H_b^n(F_2) / \text{Ker}(E\Gamma_{\nu_U} \circ i^*) \right) \geq \dim_{\mathbb{R}} \left( EH_b^n(F_2) / EN^n(F_2) \right) = \dim_{\mathbb{R}} \overline{EH}_b^n(F_2)$$

and we complete the proof for (1).

Next, we prove case (2). If  $\dim(M) = 2$ , we can use the argument in the proof of (1). Thus we can assume that  $\dim(M) \geq 3$ . Since  $\pi_M$  is an acylindrically hyperbolic group, then there exists a hyperbolic embedding  $j: F_2 \times K \rightarrow \pi_M$  [7]. We define  $s: F_2 \rightarrow F_2 \times K$  by  $r(x) = (x, \text{id})$  for  $x \in F_2$  and  $i: F_2 \rightarrow \pi_M$  by  $i = j \circ s$ . Since we assumed that  $\dim(M) \geq 3$ ,  $i$  is injective. If necessary, we retake  $U$  to be containing  $\alpha$  and  $\beta$  in Lemma 4.3.

The induced map  $j^*: EH_b^n(\pi_M) \rightarrow EH_b^n(F_2 \times K)$  is surjective [11]. Since  $s^*: EH_b^n(F_2 \times K) \rightarrow EH_b^n(F_2)$  induces an isomorphism,  $i^* = j^* \circ s^*$  is also surjective.

$$\begin{array}{ccc}
 EH_b^n(\pi_M) & \xrightarrow{E\Gamma_{\nu_U}} & EH_{\nu_U}^n(\mathcal{T}_M) \\
 \downarrow i^* & \swarrow \rho_\epsilon^* & \\
 EH_b^n(F_2) & & 
 \end{array}$$

Note that  $EH_{\nu_U}^n(\mathcal{T}_M) \supset \text{Im}(E\Gamma_{\nu_U}) \cong H_b^n(\pi_M)/\text{Ker}(E\Gamma_{\nu_U})$ . Let  $c \in EH_b^n(\pi_M)$ . If  $E\Gamma_{\nu_U}(c) = 0$ , then  $\|i^*(c)\| = 0$  by Lemma 4.3. Thus  $\text{Ker}(E\Gamma_{\nu_U}) \subset \text{Ker}(q \circ i^*)$ , where  $q: EH_b^n(F_2) \rightarrow \overline{EH}_b^n(F_2)$  is the quotient map. Therefore,

$$\dim_{\mathbb{R}} \left( H_b^n(\pi_M) / \text{Ker}(E\Gamma_{\nu_U}) \right) \geq \dim_{\mathbb{R}} \left( EH_b^n(F_2) / \text{Ker}(q \circ i^*) \right).$$

Since  $q \circ i^*$  is surjective,  $EH_b^n(F_2)/\text{Ker}(q \circ i^*) \cong \overline{EH}_b^n(F_2)$  and we complete the proof.  $\square$

*Proof of Corollary 1.6.* Since the dimension of  $\overline{EH}_b^3(F_2)$  is uncountably infinite [22], by Theorem 1.5,  $EH_{\nu_U}^3(\mathcal{T}_M) = EH_{(0)}^3(\mathcal{T}_M, \nu_U)$  is also uncountably infinite-dimensional. For  $d = 1, 2$ , There is the natural map  $EH_{(d)}^3(\mathcal{T}_M, \nu_U) \rightarrow EH_{(0)}^3(\mathcal{T}_M, \nu_U)$  induced by the inclusion  $C_{(d)}^3(\mathcal{T}_M, \nu_U) \rightarrow C_{(0)}^3(\mathcal{T}_M, \nu_U)$ . Since  $\Phi_{BM}(c) \in C_{(d)}^3(\mathcal{T}_M, \nu_U)$  for  $c \in C_b^3(\pi_M)$  by Proposition 4.1, this map surjects onto  $\text{Im}(E\Gamma_{\nu_U}) \subset EH_{\nu_U}^3(\mathcal{T}_M)$ . We can see that the dimension of  $\text{Im}(E\Gamma_{\nu_U})$  is uncountably infinite in the proof of Theorem 1.5, thus  $EH_{(d)}^3(\mathcal{T}_M, \nu_U)$  is also uncountably infinite-dimensional.  $\square$

#### ACKNOWLEDGMENT

The author would like to thank Morimichi Kawasaki for suggesting this work and for many helpful comments. He would also like to thank Michael Brandenbursky and Michał Marcinkowski for their comments on the webinar. He would like to thank Ryuma Orita for reading the manuscript and his advice on improving this paper. The author is supported by JSPS KAKENHI Grant Number JP20H00114 and JST-Mirai Program Grant Number JPMJMI22G1.

#### REFERENCES

- [1] A. Banyaga, *Sur la structure du groupe des difféomorphismes qui préservent une forme symplectique*, Comment. Math. Helv. **53** (1978), no. 2, 174–227.
- [2] ———, *The structure of classical diffeomorphism groups*, Mathematics and its Applications, vol. 400, Kluwer Academic Publishers Group, Dordrecht, 1997.
- [3] Michael Brandenbursky and Jarek Kędra, *Fragmentation norm and relative quasimorphisms*, Proc. Amer. Math. Soc. **150** (2022), no. 10, 4519–4531. MR 4470192
- [4] Michael Brandenbursky and Michał Marcinkowski, *Bounded cohomology of transformation groups*, Math. Ann. **382** (2022), no. 3–4, 1181–1197. MR 4403221
- [5] D. Burago, S. Ivanov, and L. Polterovich, *Conjugation-invariant norms on groups of geometric origin*, Groups of diffeomorphisms, Adv. Stud. Pure Math., vol. 52, Math. Soc. Japan, Tokyo, 2008, pp. 221–250.
- [6] D. Calegari, *scf*, MSJ Memoirs, vol. 20, Mathematical Society of Japan, Tokyo, 2009.
- [7] F. Dahmani, V. Guirardel, and D. Osin, *Hyperbolically embedded subgroups and rotating families in groups acting on hyperbolic spaces*, Mem. Amer. Math. Soc. **245** (2017), no. 1156, v+152.
- [8] M. Entov and L. Polterovich, *Quasi-states and symplectic intersections*, Comment. Math. Helv. **81** (2006), no. 1, 75–99.
- [9] A. Fathi, *Structure of the group of homeomorphisms preserving a good measure on a compact manifold*, Ann. Sci. École Norm. Sup. (4) **13** (1980), no. 1, 45–93.
- [10] R. Frigerio, *Bounded cohomology of discrete groups*, Mathematical Surveys and Monographs, vol. 227, American Mathematical Society, Providence, RI, 2017.
- [11] R. Frigerio, M. B. Pozzetti, and A. Sisto, *Extending higher-dimensional quasi-cocycles*, J. Topol. **8** (2015), no. 4, 1123–1155.
- [12] Jean-Marc Gambaudo and Étienne Ghys, *Commutators and diffeomorphisms of surfaces*, Ergodic Theory Dynam. Systems **24** (2004), no. 5, 1591–1617.
- [13] M. Gromov, *Volume and bounded cohomology*, Inst. Hautes Études Sci. Publ. Math. (1982), no. 56, 5–99 (1983).

- [14] M. Kawasaki, *Relative quasimorphisms and stably unbounded norms on the group of symplectomorphisms of the Euclidean spaces*, J. Symplectic Geom. **14** (2016), no. 1, 297–304.
- [15] M. Kimura, *Conjugation-invariant norms on the commutator subgroup of the infinite braid group*, J. Topol. Anal. **10** (2018), no. 2, 471–476.
- [16] A. Minasyan and D. Osin, *Acylindrical hyperbolicity of groups acting on trees*, Math. Ann. **362** (2015), no. 3-4, 1055–1105.
- [17] C. Ogle, *Polynomially bounded cohomology and discrete groups*, J. Pure Appl. Algebra **195** (2005), no. 2, 173–209.
- [18] Y. Oh and S. Müller, *The group of Hamiltonian homeomorphisms and  $C^0$ -symplectic topology*, J. Symplectic Geom. **5** (2007), no. 2, 167–219. MR 2377251
- [19] D. Osin, *Acylindrically hyperbolic groups*, Trans. Amer. Math. Soc. **368** (2016), no. 2, 851–888.
- [20] L. Polterovich, *Floer homology, dynamics and groups*, Morse theoretic methods in nonlinear analysis and in symplectic topology, NATO Sci. Ser. II Math. Phys. Chem., vol. 217, Springer, Dordrecht, 2006, pp. 417–438.
- [21] G. P. Scott, *Compact submanifolds of 3-manifolds*, J. London Math. Soc. (2) **7** (1973), 246–250.
- [22] T. Soma, *Bounded cohomology and topologically tame Kleinian groups*, Duke Math. J. **88** (1997), no. 2, 357–370.

DEPARTMENT OF MATHEMATICS, KYOTO UNIVERSITY, KITASHIRAKAWA OIWAKE-CHO, SAKYO-KU, KYOTO 606-8502, JAPAN

*Email address:* mkimura@math.kyoto-u.ac.jp