

ON THE EQUIDISTRIBUTION OF UNSTABLE CURVES FOR PSEUDO-ANOSOV DIFFEOMORPHISMS OF COMPACT SURFACES

GIOVANNI FORNI

ABSTRACT. We prove that the asymptotics of ergodic integrals along an invariant foliation of a toral Anosov diffeomorphism, or of a pseudo-Anosov diffeomorphism on a compact orientable surface of higher genus, are determined (up to a logarithmic error) by the action of the diffeomorphism on the cohomology of the surface. As a consequence of our argument and of the results of Giulietti and Liverani [GL] on horospherical averages, toral Anosov diffeomorphisms have no Ruelle resonances in the open interval $(1, e^{h_{top}})$.

1. INTRODUCTION

In this note we prove that the asymptotics of the equidistribution of unstable (or stable) curves for any C^r ($r > 1$) pseudo-Anosov diffeomorphism of a compact surface is entirely determined by the action of the diffeomorphism on the first cohomology (or homology) group up to a logarithmic error. This work was motivated by the question on whether non-trivial resonances, in the interval $(1, e^{h_{top}})$, do appear in the asymptotics of ergodic integrals of Giulietti and Liverani [GL] for Anosov diffeomorphisms of the torus and in the spectrum of the relevant transfer operator. By comparing our asymptotics with that of Giulietti and Liverani [GL], we conclude that no such non-trivial resonance exists. A direct, self-contained proof that there are no non-trivial resonances for the transfer operator has been given simultaneously and independently by V. Baladi [Ba]. Her proof was an additional motivation to write up the argument presented below.

The argument is inspired by the author's proof [F02] of deviation of ergodic averages for generic (almost all) translation flows on higher genus surfaces. Here we only deal with the special case of unstable foliations of diffeomorphisms, but we do not assume that the diffeomorphism is volume preserving. For this reason we work in Hölder spaces instead of Sobolev L^2 spaces (with respect to the invariant volume). A complete description of Ruelle resonances, as well as a complete asymptotics of ergodic averages and results on cohomological equations for linear pseudo-Anosov maps, has been given recently in the paper by Faure, Gouëzel and Lanneau [FGL]. Our argument gives a simplified proof of the part of their result concerning the Ruelle resonances in the interval $(1, e^{h_{top}})$ and the deviation

Date: March 17, 2024.

2010 Mathematics Subject Classification. 37D20, 37A25, 37C30.

Key words and phrases. Anosov diffeomorphisms, Deviation of ergodic averages for invariant foliations, Ruelle asymptotics.

of ergodic averages up to a logarithmic error. Our argument also extends to the “non-linear” case, which to the best of our knowledge has not been studied so far.

Let $A : M \rightarrow M$ be an orientation-preserving pseudo-Anosov diffeomorphism of class C^r for any $r > 1$ of a compact surface M , finite set of fixed points (singularities) at $\Sigma \subset M$, not necessarily linear (with respect to a translation structure on M), but topologically conjugated to a linear pseudo-Anosov.

Let $E^+ \subset H^1(M, \mathbb{C})$ and $E^- \subset H^1(M, \mathbb{C})$ denote respectively the unstable and the stable spaces of the finite dimensional linear map $A^\# : H^1(M, \mathbb{C}) \rightarrow H^1(M, \mathbb{C})$ induced by A on cohomology. Let $\{\mu_1, \dots, \mu_{2s}\} \subset \mathbb{C}$ denote its spectrum with

$$\mu_1 = \lambda > |\mu_2| \geq \dots \geq |\mu_k| > |\mu_{k+1}| = \dots |\mu_s| = 1.$$

Let $J_1(=1), \dots, J_k \in \mathbb{N} \setminus \{0\}$ denote the geometric multiplicities of the expanding eigenvalues $\mu_1 = \lambda > \mu_2 \geq \dots \geq \mu_k$ or, by symmetry, of the contracting eigenvalues $\mu_{2g} = \lambda^{-1} < \mu_{2g-1} \leq \dots \leq \mu_{2g-k+1}$ of the linear map $A^\# : H^1(M, \mathbb{C}) \rightarrow H^1(M, \mathbb{C})$ and let

$$\{\mathcal{C}_{i,j}^\pm | i \in \{1, \dots, k\}, j \in \{1, \dots, J_i\}\}$$

denote Jordan bases of the spaces $E^\pm \subset H^1(M, \mathbb{C})$ respectively.

Let J_A^0 denote the maximal geometric multiplicity of eigenvalues of A on the unit circle, that is, of eigenvalues $\mu_{k+1}, \dots, \mu_{2g-k}$ of $A|E^0$.

Let \mathcal{L}^\pm denote the conditional measures of the Margulis measure (measure of maximal entropy) along the leaves of the unstable and, respectively, stable foliations \mathcal{F}^+ and \mathcal{F}^- . For any $(x, \mathcal{L}) \in M \times \mathbb{R}^+$, let $\gamma_{\mathcal{L}}(x) \subset M \setminus \Sigma$ denote an unstable curve with initial point $x \in M$ and unstable Margulis “length” $\mathcal{L}^+(\gamma_{\mathcal{L}}(x)) = \mathcal{L}$ (whenever it exists). Since A is topologically conjugated by assumption to a linear pseudo-Anosov diffeomorphism, there exists a set of full measure (with respect to the Margulis measure) of $x \in M$ such that $\gamma_{\mathcal{L}}(x)$ is well-defined for all $\mathcal{L} > 0$.

1.1. Deviation of ergodic averages: invariant foliations. Let $\mathcal{Z}^{-1}(M)$ denote the subspace of closed currents of the space of currents of dimension 1 (and degree 1) dual to the space $\Omega^1(M)$ of 1-forms of class C^1 .

Theorem 1.1. *There exist injective maps $\mathcal{B}^\pm : E^\pm \rightarrow \mathcal{Z}^{-1}(M) \subset \Omega^1(M)^*$ into the subspace $\mathcal{Z}^{-1}(M)$ of closed currents (of degree and dimension 1) dual to the space $\Omega^1(M)$ of differential 1-forms of class C^1 on M such that*

$$A_* \circ \mathcal{B}^\pm = \mathcal{B}^\pm \circ A^\# \quad \text{on} \quad E^\pm \subset H^1(M, \mathbb{C}).$$

For $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, J_i\}$ let us adopt the notation

$$B_{i,j}^\pm := \mathcal{B}^\pm(\mathcal{C}_{i,j}^\pm) \in \Omega^1(M)^*.$$

The currents $B_1^+ := \mathcal{B}^+(\mathcal{C}_{1,1}^+)$ and $B_1^- := \mathcal{B}^-(\mathcal{C}_{1,1}^-)$ and can be explicitly written as follows: let \mathcal{M}^+ and \mathcal{M}^- denote the unstable and the stable Margulis measures. For any 1-form α of class C^1 on M we have

$$(1) \quad B_1^+(\alpha) = \int_M \mathcal{M}^- \otimes \alpha \quad \text{and} \quad B_1^-(\alpha) = \int_M \alpha \otimes \mathcal{M}^+.$$

There exists a constant $C > 0$ such that the following holds. For any unstable curve $\gamma_{\mathcal{L}}(x)$ of initial point $x \in M$ and unstable length $\mathcal{L} > 1$, there exists a set of uniformly bounded coefficients

$$\{c_{i,j}(x, \mathcal{L}) \mid i \in \{2, \dots, k\}, j \in \{1, \dots, J_i\}\}$$

such that, for any differential 1-form $\eta \in C^1(M)$, we have

$$(2) \quad \left| \int_{\gamma_{\mathcal{L}}(x)} \eta - \mathcal{L} B_1^+(\eta) - \sum_{i=2}^k \sum_{j=1}^{J_i} c_{i,j}(x, \mathcal{L}) B_{i,j}^+(\eta) (\log \mathcal{L})^{j-1} \mathcal{L}^{\frac{\log |\mu_j|}{h_{\text{top}}(A)}} \right| \\ \leq C \|\eta\|_{C^1(M)} [\log(1 + \mathcal{L})]^{\max(J_A^0, 1) + 1}.$$

In addition, there exists $c > 0$ such that for every $x \in M$ and for every $i \in \{2, \dots, k\}$ and $j \in \{1, \dots, J_i\}$, there exists a sequence $\mathcal{L}_n := \mathcal{L}_n^{(i,j)}(x)$ such that

$$(3) \quad \inf_{n \in \mathbb{N}} |c_{i,j}(x, \mathcal{L}_n)| \geq c.$$

From the above asymptotic result of formula (2), together with the lower bound of formula (3), we can derive *a posteriori* additional invariant properties of the closed currents $B_1^\pm, B_{i,j}^\pm$ which are not apparent from their construction. In order to state these invariance properties we recall the following:

Definition 1.2. A current B is called basic for a foliation \mathcal{F} on a smooth manifold if, for all vector fields Y tangent to \mathcal{F} ,

$$\mathcal{L}_Y B = \iota_Y B = 0.$$

If the current has dimension 1, by the identity $\mathcal{L}_Y B = \iota_Y dB + d\iota_Y B$ it follows that B is basic if and only

$$dB = \iota_Y B = 0.$$

Addendum 1.3. The currents B_1^\pm and $B_{i,j}^\pm$ for $i \in \{2, \dots, k\}$, $j \in \{1, \dots, J_i\}$, of Theorem 1.1 are basic for the unstable, respectively, stable, foliations \mathcal{F}^\pm on $M \setminus \Sigma$.

The asymptotic expansion of Theorem 1.1 can be refined by introducing finitely additive functionals on rectifiable arcs, following the work of A. Bufetov [Bu14] on translation flows (see also [BuFo14] on horocycle flows and [FoKa] on nilflows).

Theorem 1.4. Let Γ_r the set of all rectifiable paths (considered as a subset of the space of currents). There exists a map $\hat{\beta}^+ : \Gamma_r \rightarrow \mathcal{B}^+(E^+) \subset \mathcal{Z}^{-1}(M)$ into the space of closed currents (with image in the space of basic currents for the unstable foliation) such that the following holds. The map $\hat{\beta}^+$ has the following properties:

(1) (Additive property) For any decomposition $\gamma = \gamma_1 + \gamma_2$ into subarcs,

$$\hat{\beta}^+(\gamma) = \hat{\beta}^+(\gamma_1) + \hat{\beta}^+(\gamma_2);$$

(2) (Scaling) For any $\gamma \in \Gamma_r$, we have

$$\hat{\beta}^+(A\gamma) = A_* \hat{\beta}^+(\gamma),$$

(3) (Stable holonomy invariance) For all pair of arcs $\gamma_1, \gamma_2 \in \Gamma_r$ equivalent under the stable holonomy, we have

$$\hat{\beta}^+(\gamma_1) = \hat{\beta}^+(\gamma_2).$$

In addition, the functional $\hat{\beta}^+$ has the following asymptotic property: there exists a constant $C > 0$ such that, for every arc $\gamma \in \Gamma_r$ we have

$$\|\gamma - \hat{\beta}^+(\gamma)\|_{-1} \leq C(1 + \mathcal{L}^-(\gamma))[\log(1 + \mathcal{L}^+(\gamma))]^{\max(J_A^0, 1) + 1}.$$

1.2. Deviation of ergodic averages:unstable (horocyclic) flows. Let us then assume that the tangent map DA of the pseudo-Anosov diffeomorphism preserves the orientation of the unstable foliation of a pseudo-Anosov map A of class C^r with $r > 1 + \alpha$ and, as in [GL], let X denote a vector field of class $C^{1+\alpha}$, tangent to the unstable foliation and normalized to have constant norm with respect to a fixed Riemannian metric on M . By definition, for all $n \in \mathbb{N}$ there exists a function $v_n : M \setminus \Sigma \rightarrow \mathbb{R}$ such that (see [GL], formula (1.3))

$$D_x A^n(X_x) = v_n(x) X_{A^n(x)}, \quad \text{for all } x \in M \setminus \Sigma.$$

From Theorem 1.1 we derive a result on the asymptotics of ergodic integrals for the flow $h_{\mathbb{R}}^X$ generated by the unstable vector field X on $M \setminus \Sigma$.

For linear pseudo-Anosov maps on higher genus surfaces, a sharper asymptotics of ergodic integrals of the stable and unstable translation flows was obtained by F. Faure, S. Gouëzel and E. Laneeau [FGL], who also proved complete results on the existence and regularity of solutions of the cohomological equation.

For the case of toral Anosov diffeomorphisms the asymptotics of ergodic integrals of stable and unstable vector fields was studied in the pioneering work of P. Giulietti and C. Liverani [GL]. V. Baladi [Ba] has given a proof, independent of ours, that in the toral case there are no “deviation resonances” in the Giulietti-Liverani asymptotics (see Remark 1.6 below). Her argument also proves that for the stable or unstable vector fields of sufficiently regular Anosov diffeomorphisms every zero average function is a continuous coboundary, a result which is beyond the reach of our cohomological approach.

Let \hat{X} be a 1-form of class $C^{1+\alpha}$ dual to the vector field X , in the sense that

$$\hat{X}(X) \equiv 1 \quad \text{on } M \setminus \Sigma.$$

The unique invariant probability measure μ_X of the flow $h_{\mathbb{R}}^X$ is given by the condition

$$\mu_X \in \mathbb{R}^+(B_1^+ \wedge \hat{X}) \quad \left(\text{defined as } \int_M f d\mu_X = \frac{B_1^+(f\hat{X})}{B_1^+(\hat{X})}, \text{ for all } f \in C^1(M) \right).$$

Let $\{D_{i,j}^X | i \in \{2, \dots, k\}, j \in \{1, \dots, J_i\}\}$ the finite set of distributions

$$D_{i,j}^X := B_{i,j}^+ \wedge \hat{X}, \quad \text{for all } i \in \{2, \dots, k\}, j \in \{1, \dots, J_i\}.$$

It follows from Addendum 1.3 (see Section 4) that, for all $i \in \{2, \dots, k\}$ and $j \in \{1, \dots, J_i\}$, the distributions $D_{i,j}^X$ are X -invariant, in the sense that

$$X D_{i,j}^X = 0 \quad \text{in } \mathcal{D}'(M \setminus \Sigma).$$

The following asymptotic expansion of ergodic integrals holds.

Corollary 1.5. *There exist a constant $C_X > 0$ and, for all $(x, T) \in (M \setminus \Sigma) \times \mathbb{R}^+$, a finite set of uniformly bounded coefficients*

$$\{c_{i,j}^X(x, T) | i \in \{2, \dots, k\}, j \in \{1, \dots, J_i\}\},$$

such that, for all $f \in C^1(M)$ and for all $(x, T) \in (M \setminus \Sigma) \times \mathbb{R}^+$ with $h_{[0,T]}(x) \subset M \setminus \Sigma$, we have

$$(4) \quad \left| \int_0^T f \circ h_t^X(x) dt - T \int_M f d\mu_X - \sum_{i=2}^k \sum_{j=1}^{J_i} c_{i,j}^X(x, T) D_{i,j}^X(f) (\log T)^{j-1} T^{\frac{\log |\mu_i|}{h_{\text{top}}(A)}} \right| \leq C_X \|f\|_{C^1(M)} [\log(1+T)]^{\max(J_A^0, 1)+1}.$$

In addition, there exists $c > 0$ such that for every $x \in M$ and for every $i \in \{2, \dots, k\}$ and $j \in \{1, \dots, J_i\}$, there exists a sequence $T_n := T_n^{(i,j)}(x)$ such that

$$(5) \quad \inf_{n \in \mathbb{N}} |c_{i,j}(x, T_n)| \geq c.$$

Remark 1.6. (Comparison with a result of V. Baladi [Ba], Corollary 2.3) A similar, but more refined, asymptotics of ergodic averages is proved in [Ba] for the case of C^r Anosov diffeomorphisms of the 2-torus, a case for which the spectrum of $A^\# : H^1(\mathbb{T}^2) \rightarrow H^1(\mathbb{T}^2)$ has a unique expanding eigenvalue $\mu_1 = \lambda > 1$ (and no neutral eigenvalues).

In fact, she proves (see [Ba], Corollary 2.3) that there exists $r_0, r_1 > 1$ such that for any $r \geq \max\{r_0, r_1\}$ there exist constants $C > 0$ and $\theta_{\min} < 0$ such that for all $f \in C^{r-1}(\mathbb{T}^2)$, and for all $T > 0$,

$$\left| \int_0^T f \circ h_t^X(x) dt - T \int_{\mathbb{T}^2} f d\mu_X \right| \leq C(T^{\theta_{\min}} |f|_{C^{r-1}} + \sup |f|).$$

In particular, $\mu_X(f) = 0$ if and only if f is a continuous coboundary.

In comparison with the asymptotics of Corollary 1.5 above, Baladi's asymptotics has bounded, not logarithmic, error terms. However, such terms cannot be neglected for general functions of class C^1 . A refined asymptotics with bounded error terms may hold for general functions of class $C^{1+\alpha}$, for some $\alpha > 0$, and for such functions one can derive results on existence of solutions of the cohomological equation (see [F07] and [MY16] also in the higher genus case). Such refined results are beyond the purely cohomological approach presented here.

1.3. Ruelle-Pollicott asymptotics. From the above equidistribution results we derive a Ruelle-Pollicott asymptotics for pseudo-Anosov maps. For linear pseudo-Anosov maps on higher genus surfaces, a complete Ruelle-Pollicott asymptotics has been obtained by F. Faure, S. Gouëzel and E. Laneeau [FGL]. For toral Anosov diffeomorphisms the Ruelle-Pollicott asymptotics follows from the work of P. Giulietti and C. Liverani [GL]. V. Baladi [Ba] has given an independent proof that in the toral case there are no “deviation resonances” in the Giulietti-Liverani asymptotics (see Remark 1.8 below).

Corollary 1.7. *Let us assume that the diffeomorphism A is of class C^r with $r > 2$. Then A has a Ruelle-Pollicott asymptotics in the sense that, for any $f, g \in C^1(M)$, the correlations $\langle f \circ A^n, g \rangle$ with respect to the Margulis measure \mathcal{M} have an expansion*

$$(6) \quad \begin{aligned} \langle f \circ A^n, g \rangle &= \left(\int_M f d\mathcal{M} \right) \left(\int_M g d\mathcal{M} \right) \\ &+ \sum_{i=2}^k \sum_{j=1}^{J_i} \mathcal{C}_{i,j}(f, g, n) n^{j-1} \left(\frac{\mu_i}{e^{h_{\text{top}}(A)}} \right)^n + \mathcal{R}(f, g, n) \frac{n^{\max(J_A^0, 1)+1}}{e^{nh_{\text{top}}(A)}}, \end{aligned}$$

with $\mathcal{C}_{i,j}(f, g, n)$ and $\mathcal{R}(f, g, n)$ uniformly bounded: there exists a constant $C > 0$ such that, for all $n \in \mathbb{N}$ we have

$$\sum_{i=2}^k \sum_{j=1}^{J_i} |\mathcal{C}_{i,j}(f, g, n)| + |\mathcal{R}(f, g, n)| \leq C |f|_{C^1(M)} |g|_{C^1(M)}.$$

Remark 1.8. (Comparison with a result of V. Baladi [Ba], Corollary 2.5) A similar, but more refined, Ruelle-Pollicott asymptotics is proved in [Ba] for the case of C^r Anosov diffeomorphisms of the 2-torus, a case for which the spectrum of $A^\# : H^1(\mathbb{T}^2) \rightarrow H^1(\mathbb{T}^2)$ has a unique expanding eigenvalues $\mu_1 = \lambda > 1$ (and no neutral eigenvalues).

In fact, she proves (see [Ba], Corollary 2.5) that for any $r > 1$ there exist constants $C > 0$ and $\rho \in (0, 1)$ such that for all $f, g \in C^{r-1}(\mathbb{T}^2)$, and for all $n \in \mathbb{N}$,

$$\left| \langle f \circ A^n, g \rangle - \left(\int_{\mathbb{T}^2} f d\mathcal{M} \right) \left(\int_{\mathbb{T}^2} g d\mathcal{M} \right) \right| \leq C \rho^n |f|_{C^{r-1}} |g|_{C^{r-1}}.$$

The error term in the above estimate is refined as follows. Let $\lambda^+ > 1 > \lambda^-$ denote the Lyapunov exponents. There exists $\tilde{\rho}_A < e^{h_{\text{top}}(A)} \min(\lambda^+, (\lambda^-)^{-1})^{-\frac{r-1}{2}}$ such that for $\tilde{\rho}_A < 1$ the estimate holds for some $\rho < e^{-h_{\text{top}}(A)}$. In particular, there exists $r_1 > 1$ such that whenever A is of class C^r for $r > r_1$ then $\tilde{\rho}_A < 1$ and the estimates hold for some $\rho < e^{-h_{\text{top}}(A)}$. For $\tilde{\rho}_A \geq 1$ the estimate holds for any $\rho > e^{-h_{\text{top}}(A)} \tilde{\rho}_A$.

The paper is organized as follows. In Section 2 we prove a representation lemma for stable and unstable cohomology classes in terms, respectively, of stable and unstable closed (basic) currents in the dual space of 1-forms of class C^1 (Lemma 2.1). From the representation lemma and de Rham theorem we derive a result on the asymptotics of the action of the pseudo-Anosov diffeomorphism on the space of closed currents dual to the space of 1-forms of class C^1 . In Section 3 we apply the results of Section 2 first to a special sequence of return leaves of the unstable foliation, then after proving a standard decomposition lemma (see Lemma 3.1), to all leaves. We then complete the proof of the main result, Theorem 1.1. Finally, in Section 4 we derive the proof of Corollary 1.5 on the deviation of ergodic averages and Corollary 1.7 on the Ruelle-Pollicott asymptotics.

ACKNOWLEDGEMENTS

The author wishes to thank V. Baladi, P. Giulietti, S. Gouëzel and C. Liverani for their interest in the cohomological perspective presented here and their encouragement to write up this note. He is also grateful to the participants of the Oberwolfach Seminar on *Anisotropic Spaces and their Applications to Hyperbolic and Parabolic Systems* (June 9-15, 2019) for the stimulating environment and to MFO for making the Seminar possible. Part of the paper took shape while the author was preparing his lectures at the Seminar. This research was supported by the NSF grant DMS 1600687 and by a Research Chair of the Fondation Sciences Mathématiques de Paris (FSMP). The author is grateful to the Institut de Mathématiques de Jussieu (IMJ) for its hospitality during the preparation of the first draft of the paper.

2. GROWTH OF CLOSED CURRENTS

Lemma 2.1. *There exist injective maps $\mathcal{B}^\pm : E^\pm \rightarrow \mathcal{Z}^{-1}(M) \subset \Omega^1(M)^*$ into the subspace $\mathcal{Z}^{-1}(M)$ of closed currents (of degree and dimension 1) dual to the space $\Omega^1(M)$ of 1-forms of class C^1 on M such that*

$$A_* \circ \mathcal{B}^\pm = \mathcal{B}^\pm \circ A^\# \quad \text{on} \quad E^\pm \subset H^1(M, \mathbb{C}).$$

Proof. There exists a linear map $\alpha : H^1(M, \mathbb{C}) \rightarrow \mathcal{Z}^{-1}(M) \subset \Omega^1(M)^*$ of closed form of class C^1 on M such that, for any $C \in H^1(M, \mathbb{C})$ we have that $[\alpha(C)] = C \in H^1(M, \mathbb{C})$, hence there exists a (bounded) linear map $u : H^1(M, \mathbb{C}) \rightarrow C^{r+1}(M)$ such that

$$A^* \circ \alpha = \alpha \circ A^\# + du \quad \text{on} \quad H^1(M, \mathbb{C}).$$

We note that since the exterior derivative is elliptic, for every $C \in H^1(M, \mathbb{C})$, the function $u(C) \in C^{r+1}$ under the hypothesis that A is a C^r diffeomorphism.

By iterating the above identity, for any $C \in H^1(M, \mathbb{C})$, we have

$$(A^*)^n(\alpha(C)) = \alpha[(A^\#)^n(C)] + d[A^*(u((A^\#)^{n-1}(C)))] + \dots + d[(A^\#)^n \circ u(C)].$$

It follows that

$$(A^*)^n \circ \alpha \circ (A^\#)^{-n} = \alpha + d \circ A^* \circ u \circ (A^\#)^{-1} + \dots + d \circ (A^*)^n \circ u \circ (A^\#)^{-n}.$$

Let us give the argument for the unstable space $E^+ \subset H^1(M, \mathbb{C})$, otherwise we replace A with its inverse A^{-1} . We claim that, since the restriction $A^\#|_{E^+}$ is (strictly) expanding, it follows by completeness that for every $C \in E^+ \subset H^1(M, \mathbb{C})$, the following limit exists in $C^0(M)$ (and in $L^2(M)$ in the volume preserving case):

$$\begin{aligned} U(C) &:= (A^* \circ u)((A^\#)^{-1}C) + \dots + ((A^*)^n \circ u)((A^\#)^{-n}C) \\ &= \sum_{k=1}^{\infty} ((A^*)^k \circ u)((A^\#)^{-k}C) \in C^0(M). \end{aligned}$$

In fact, for every function $u \in C^0(M)$ we have that

$$\|(A^*)^n(u)\|_{C^0(M)} = \|u\|_{C^0(M)}$$

and, in the volume preserving case, also

$$\|(A^*)^n(u)\|_{L^2(M, \text{vol})} = \|u\|_{L^2(M, \text{vol})}.$$

Thus, since $A^\#|E^+$ is (strictly) expanding, there exists $\rho > 1$ such that, for all $k \in \mathbb{N}$,

$$\|((A^*)^k \circ u)((A^\#)^{-k}C)\|_{C^0(M)} \leq \left(\max_{|C|=1} \|u(C)\|_{C^0(M)} \right) \rho^{-k},$$

which implies the absolute convergence of the series, hence the claim.

It follows that the following limit exists in $(C^1(M))^*$:

$$\mathcal{B}^+(C) := \lim_{n \rightarrow +\infty} (A^*)^n \circ \alpha \circ (A^\#)^{-n}(C) = \alpha(C) + dU(C).$$

By construction we clearly have $[\mathcal{B}^+(C)] = [\alpha(C)] = [C] \in H^1(M, \mathbb{C})$ and

$$\begin{aligned} A_*(\mathcal{B}^+(C)) &= \lim_{n \rightarrow +\infty} (A^*)^{n+1} (\alpha((A^\#)^{-n}(C))) \\ &= \lim_{n \rightarrow +\infty} (A^*)^{n+1} \left(\alpha((A^\#)^{-(n+1)}(A^\#C)) \right) = \mathcal{B}^+(A^\#C). \end{aligned}$$

The map $\mathcal{B}^+ : E^+ \rightarrow \mathcal{Z}^{-1}(M)$ is therefore defined, it is linear by its definition and it is injective since $[\mathcal{B}^+(C)] = [C] \in H^1(M, \mathbb{C})$. \square

Remark 2.2. *By definition of the Margulis measure, the currents in formula are distributional eigenvectors for the diffeomorphism A for the eigenvalues $\lambda^{\pm 1}$ since*

$$\begin{aligned} A_*B_1^+(\alpha) &= B_1^+(A^*\alpha) = \int_M \mathcal{M}^- \otimes A^*\alpha = \int_M A_*^{-1} \mathcal{M}^- \otimes \alpha = \lambda B_1^+(\alpha), \\ A_*B_1^-(\alpha) &= B_1^-(A^*\alpha) = \int_M A^*\alpha \otimes \mathcal{M}^+ = \int_M \alpha \otimes A_*^{-1} \mathcal{M}^+ = \lambda^{-1} B_1^-(\alpha). \end{aligned}$$

Since the eigenvalues $\lambda^{\pm 1}$ for action A^* of the diffeomorphism A on $H^1(M, \mathbb{C})$ are simple, it follows that the Margulis currents B_1^\pm in formula (1) are (up to multiplicative constants) the unique distributional eigenvectors of eigenvalues $\lambda^{\pm 1}$ for the linear map A_* on the space of closed 1-currents.

Let us recall that $\mathcal{Z}^{-1}(M)$ denotes the space of closed currents of dimension and degree 1 on M dual to the space of 1-forms of class C^1 . Let E^+ , E^- and E^0 denote, respectively, the unstable, the stable and the central stable space of the linear map $A^\# : H^1(M, \mathbb{C}) \rightarrow H^1(M, \mathbb{C})$ induced by $A : M \rightarrow M$ on the first cohomology of M . There is a direct decomposition

$$(7) \quad H^1(M, \mathbb{C}) = E^+ \oplus E^- \oplus E^0.$$

By Lemma 2.1 there exist maps $\mathcal{B}^\pm : E^\pm \rightarrow \mathcal{Z}^{-1}(M)$ such that

$$A^* \circ \mathcal{B}^\pm = \mathcal{B}^\pm \circ A^\# \quad \text{on } E^\pm.$$

There is also a linear map $\mathcal{B}^0 : E^0 \rightarrow \mathcal{Z}^1(M)$, with values in the space $\mathcal{Z}^1(M)$ of closed smooth 1-forms, and a linear map $F : E^0 \rightarrow C^\infty(M)$ such that

$$A^* \circ \mathcal{B}^0 = \mathcal{B}^0 \circ A^\# + dF \quad \text{on } E^0.$$

The restriction $A^\#|E^0$ is by definition a unipotent linear operator. Let J_A^0 the dimension of the largest Jordan block of $A^\#|E^0$.

For every closed current $\gamma \in \mathcal{Z}^{-1}(M)$ let $[\gamma] \in H^1(M, \mathbb{R})$ denote its cohomology class and let $[\gamma]^+$, $[\gamma]^-$ and $[\gamma]^0$ denote, respectively, the projections of the cohomology class $[\gamma]$ on the subspaces E^+ , E^- and E^0 according to the decomposition in formula (7), that is, for all $\gamma \in \mathcal{Z}^{-1}(M)$ we have

$$[\gamma] = [\gamma]^+ + [\gamma]^- + [\gamma]^0, \quad \text{with } [\gamma]^\pm \in E^\pm \text{ and } [\gamma]^0 \in E^0.$$

Lemma 2.3. *There exists $C > 0$ such that, for any closed current $\gamma \in \mathcal{Z}^{-1}(M)$ of dimension 1 (and degree 1) and for any $n \in \mathbb{N}$, we have*

$$\|A_*^n(\gamma) - \mathcal{B}^+((A^\#)^n[\gamma]^+) \|_{-1} \leq C \|[\gamma]\| n^{\max(J_A^0, 1)}.$$

Proof. For every $n \in \mathbb{N}$, there exist $X^\pm(n) \in E^\pm$, $X^0(n) \in E^0$ and a current U_n of dimension 2 (and degree 0) in the dual space of the space of 2-forms with C^0 coefficients, such that we can write

$$A_*^n(\gamma) = \mathcal{B}^+(X^+(n)) + \mathcal{B}^-(X^-(n)) + \mathcal{B}^0(X^0(n)) + dU_n.$$

We therefore have the identities

$$\begin{aligned} A_*^{n+1}(\gamma) &= \mathcal{B}^+(X^+(n+1)) + \mathcal{B}^-(X^-(n+1)) + \mathcal{B}^0(X^0(n+1)) + dU_{n+1} \\ &= A_* (\mathcal{B}^+(X^+(n)) + \mathcal{B}^-(X^-(n)) + \mathcal{B}^0(X^0(n)) + dU_n) \\ &= \mathcal{B}^+(A^\#X^+(n)) + \mathcal{B}^-(A^\#X^-(n)) + \mathcal{B}^0(A^\#X^0(n)) + dF(X^0(n)) + dA^*U_n. \end{aligned}$$

By projecting the above identity on cohomology we have

$$\begin{aligned} X^\pm(n+1) &= A^\#X^\pm(n), \quad X^0(n+1) = A^\#X^0(n) \quad \text{and} \\ U_{n+1} &= F(X^0(n)) + A^*(U_n), \end{aligned}$$

from which we derive that

$$\begin{aligned} X_j^\pm(n) &= (A^\#)^n(x^\pm(0)) = (A^\#)^n([\gamma]^\pm), \\ X^0(n) &= (A^\#)^n(X^0(0)) = (A^\#)^n([\gamma]^0), \end{aligned}$$

and that there exists a constant $C > 0$ such that

$$\|U_n\|_{C^0(M)^*} \leq C \|[\gamma]^0\| n^{\max(J_A^0, 1)}.$$

In fact

$$\begin{aligned} \|U_{n+1}\|_{C^0(M)^*} &\leq \|A^*(U_n)\|_{C^0(M)^*} + C_F \|X^0(n)\| \\ &\leq \|U_n\|_{C^0(M)^*} + C'_F (n+1)^{\max(J_A^0, 1)-1} \|X^0(0)\|, \end{aligned}$$

hence

$$\|U_n\|_{C^0(M)^*} \leq C'_F \|[\gamma]^0\| \sum_{m=0}^n (m+1)^{\max(J_A^0, 1)-1},$$

thus the argument is concluded. \square

3. CLOSEST RETURN LEAVES

In this section we derive the asymptotics for closed currents given by closest return leaves of the unstable foliation. We measure distances on M along the unstable and stable foliations by the conditional measures \mathcal{L}^\pm of the Margulis measure.

For all $x \in M$ let $I^s(x)$ denote a stable curve centered at $x \in M$ of unit length and let $\tilde{\gamma}(x)$ be an unstable return curve with endpoints $x, y \in I^s(x)$ with no other intersections with $I^s(x)$. Let $\gamma(x) = \tilde{\gamma}(x) \cup J^s(x, y)$ denote the closed arc which is union of the unstable curve $\tilde{\gamma}(x)$ with the stable curve $J^s(x, y) \subset I^s(x)$ with endpoints $x, y \in I^s(x)$. By compactness there exist constants $C > c > 0$ such that

$$c \leq \mathcal{L}^+(\tilde{\gamma}(x)) \leq C.$$

For every $n \in \mathbb{N}$, let then $\tilde{\gamma}_n(x)$ and $\gamma_n(x)$ be the curves

$$\tilde{\gamma}_n(x) = A^n(\tilde{\gamma}(A^{-n}(x))) \quad \text{and} \quad \gamma_n(x) = A^n(\gamma(A^{-n}(x))).$$

Let $\lambda = e^{h_{\text{top}}(A)}$ denote the expansion rates of the diffeomorphism A on M with respect to the unstable Margulis measure. We have

$$c\lambda^n \leq \mathcal{L}^+(\tilde{\gamma}_n(x)) \leq C\lambda^n.$$

We have the following decomposition result:

Lemma 3.1. *Any unstable curve $\tilde{\gamma}$ has a decomposition into consecutive closest returns closed curves,*

$$(8) \quad \tilde{\gamma} = \sum_{\ell=1}^n \sum_{m=1}^{m_\ell} \tilde{\gamma}_\ell(x_{\ell,m}) + \gamma_0,$$

such that $c\lambda^n \leq \mathcal{L}^+(\tilde{\gamma}) \leq C\lambda^{n+1}$ and, for all $1 \leq \ell \leq n$,

$$(9) \quad m_\ell \leq C\lambda/c, \quad \mathcal{L}^+(\tilde{\gamma}_\ell(x_{\ell,m})) \in [c\lambda^\ell, C\lambda^\ell] \quad \text{and} \quad \mathcal{L}^+(\gamma_0) \leq C\lambda/c.$$

Proof. Let $x \in M$ denote the initial point of $\tilde{\gamma}$. Let $n \in \mathbb{N}$ denote the unique integer such that $\tilde{\gamma}_n(x) \subset \tilde{\gamma}$. We have

$$c\lambda^n \leq \mathcal{L}^+(\tilde{\gamma}_n(x)) \leq \mathcal{L}^+(\tilde{\gamma}) \leq \mathcal{L}^+(\tilde{\gamma}_{n+1}(x)) \leq C\lambda^{n+1}.$$

Let then $x_{n,1} = x$. Let y denote the endpoint of $\tilde{\gamma}_n(x)$. Let then n' denote the largest integer such that $\tilde{\gamma}_{n'}(y) \subset \tilde{\gamma} \setminus \tilde{\gamma}_n(x)$. If $n' = n$, we set $y = x_{n,2}$, otherwise if $n' < n$ we set $y = x_{n',1}$. We keep iterating this procedure. It is clear that

$$cm_\ell \lambda^\ell \leq C\lambda^{\ell+1}$$

otherwise the union of segments $\tilde{\gamma}_\ell(x_{\ell,1}) \cup \dots \tilde{\gamma}_\ell(x_{\ell,m_\ell})$ would be covered by a segment $\tilde{\gamma}_{\ell+1}(x_{\ell+1,m_{\ell+1}})$. In particular, we have that $\mathcal{L}^+(\gamma_0) \leq C\lambda/c$. \square

Theorem 3.2. *There exists a map $\mathcal{C}^+ : \Gamma^+ \rightarrow E^+$ on the set Γ^+ of unstable curves with bounded range in $E^+ \subset H^1(M, \mathbb{C})$, such that for any unstable curve $\tilde{\gamma}$ of unstable length $\mathcal{L}^+(\tilde{\gamma}) > 1$ we have*

$$(10) \quad \|\tilde{\gamma} - \mathcal{B}^+((A^\#)^{\lceil \frac{\log(\mathcal{L}^+(\tilde{\gamma}))}{h_{\text{top}}(A)} \rceil}(\mathcal{C}^+(\tilde{\gamma})))\|_{-1} \leq C[\log(1 + \mathcal{L}^+(\tilde{\gamma}))]^{\max(J_A^0, 1)+1}.$$

Proof. For any $x, y \in M$, let $I_{x,y}$ denote a curve of bounded length joining x to y . For any unstable curve $\tilde{\gamma}$ with endpoints x and $y \in M$, let $\gamma = \tilde{\gamma} \cup I_{x,y}$. There exists a current u_γ of degree 0 (and dimension 2) in the dual space of the space of 2-forms with C^0 coefficients

$$\gamma = \mathcal{B}^+([\gamma]^+) + \mathcal{B}^-([\gamma]^-) + \mathcal{B}^0([\gamma]^0) + du_\gamma.$$

By the decomposition lemma we also have

$$\tilde{\gamma} = \sum_{\ell=1}^n \sum_{m=1}^{m_\ell} \tilde{\gamma}_\ell(x_{\ell,m}) + \tilde{\gamma}_0,$$

with $\tilde{\gamma}_\ell(x_{\ell,m}) = A_*^\ell(\gamma(A^{-\ell}(x_{\ell,m})))$, hence there exists a constant $C > 0$ such that

$$\|\tilde{\gamma}_\ell(x_{\ell,m}) - \mathcal{B}^+((A^\#)^\ell[\gamma(A^{-\ell}(x_{\ell,m}))]^+)\|_{-1} \leq C\ell^{\max(J_A^0, 1)},$$

and by summation

$$\|\tilde{\gamma} - \sum_{\ell=1}^n \sum_{m=1}^{m_\ell} \mathcal{B}^+((A^\#)^\ell[\gamma(A^{-\ell}(x_{\ell,m}))]^+)\|_{-1} \leq Cn^{\max(J_A^0, 1)+1}.$$

Let $\mathcal{C}^+(\tilde{\gamma}) \in E^+$ denote the cohomology class

$$(11) \quad \mathcal{C}^+(\tilde{\gamma}) := \sum_{\ell=1}^n \sum_{m=1}^{m_\ell} (A^\#)^{-([\frac{\log(\mathcal{L}^+(\tilde{\gamma})}{h_{top}(A)})] - \ell)} [\gamma(A^{-\ell}(x_{\ell,m}))]^+.$$

Since by the decomposition lemma we have

$$n \leq \frac{\log(\mathcal{L}^+(\tilde{\gamma})/c)}{\log \lambda} = \frac{\log(\mathcal{L}^+(\tilde{\gamma})/c)}{h_{top}(A)},$$

and since $(A^\#)^{-1}$ is contracting on the unstable space E^+ , by the decomposition lemma, it follows that there exists a constant $C' > 0$ such that

$$\|\mathcal{C}^+(\tilde{\gamma})\| \leq C', \quad \text{for all } \tilde{\gamma} \in \Gamma^+.$$

The result follows. \square

Proof of Theorem 1.1. It follows from Theorem 3.2 by writing the asymptotics in formula (10) with respect to a Jordan basis of $A^\#|E^+$. The lower bound on the coefficients along subsequences in formula (3) holds by Lemma 2.3 along sequences of closest return leaves. In fact, it is known that for every projection given by a Jordan basis of $A^\#$ in $H^1(M, \mathbb{R})$ and for all $x \in M \setminus \Sigma$ with infinite forward orbit, there exists a close return orbit $\gamma(x)$ with non-zero projection (see for instance [Bu14], Prop. 2.9, or [DHL14], § 5.3). We recall below a proof of this statement for the convenience of the reader.

Let I denote any stable arc. The return map of the unstable flow $h_{\mathbb{R}}^X$ to I is an Interval Exchange Transformation and that the flow $h_{\mathbb{R}}^X$ is a suspension of that Interval Exchange Transformation under a piece-wise constant roof function.

It is well-known that the set of loops equal to the union of a first return orbit of $h_{\mathbb{R}}^X$ (to any given transverse stable arc I) with a transverse (stable) segment joining the endpoints spans the relative homology group $H_1(M \setminus \Sigma, \mathbb{Z})$ (see for instance [Yoc10], §4.5). By Poincaré duality, it follows that, for every given Jordan

projection (induced by a Jordan basis of the map $A^\#$ on $H^1(M, \mathbb{R})$), there exists a subinterval $J \subset I \subset M$ such that the cohomology class of the loop $\gamma(x)$, given by the union of the return orbit of any point $x \in J$ with a transverse (stable) arc $I_x \subset I$, has a non-zero projection.

Let now $x \in I \subset M \setminus \Sigma$ be an arbitrary point with infinite forward orbit. Let $\tilde{\gamma}'(x)$ denote the orbit arc until the first visit of the forward orbit $h_{\mathbb{R}^+}^X(x)$ to the subinterval J and let $\gamma'(x)$ denote the loop given by the union of $\tilde{\gamma}'(x)$ with a transverse (stable) interval $I' \subset I$. Let $\tilde{\gamma}''(x)$ denote the smallest orbit arc of the forward orbit $h_{\mathbb{R}^+}^X(x)$ which is strictly larger than $\tilde{\gamma}'(x)$ and let $\gamma''(x)$ denote the loop given by the union of $\tilde{\gamma}''(x)$ with a transverse (stable) interval $I'' \subset I$.

We claim that the given Jordan projection of either the cohomology class $[\gamma'(x)] \in H^1(M, \mathbb{Z})$ of the loop $\gamma'(x)$ or cohomology class $[\gamma''(x)] \in H^1(M, \mathbb{Z})$ of the loop $\gamma''(x)$ is non-zero.

In fact, let $x' \in J \subset I$ denote the endpoint of the orbit arc $\tilde{\gamma}'(x)$ and let $\gamma(x')$ denote the loop given by the union of the return orbit of $x' \in J$ with a transverse (stable) arc $I_{x'} \subset I$. On the one hand, by hypothesis we have that the given Jordan projection of the class $[\gamma(x')]$ is non-zero; on the other hand, by construction we have

$$[\gamma''(x)] = [\gamma'(x)] + [\gamma(x')] \quad \text{in } H^1(M, \mathbb{Z}),$$

hence the claim follows.

The lower bounds in the statement of Theorem 1.1 follow from the above claim on Jordan projections of orbit arcs and from Theorem 3.2. \square

From the statement of Theorem 1.1 we derive:

Proof of Addendum 1.3. Since the currents B_1^\pm and $B_{i,j}^\pm$ are closed, it is enough to prove that, for every vector field Y^\pm tangent to the unstable/stable foliation \mathcal{F}^\pm , we have that

$$\iota_{Y^\pm} B_1^\pm = \iota_{Y^\pm} B_{i,j}^\pm = 0, \quad \text{for all } i \in \{2, \dots, k\}, j \in \{1, \dots, J_i\}.$$

By Theorem 1.1 for any vector field $Y := Y^+$ tangent to the unstable foliation, and for every 2-form w of class C^1 we have, since $\int_{\gamma_{\mathcal{L}}(x)} \iota_Y w = 0$,

$$(12) \quad \begin{aligned} & |\mathcal{L} B_1^+(\iota_Y w) - \sum_{i=2}^k \sum_{j=1}^{J_i} c_{i,j}(x, \mathcal{L}) B_{i,j}^+(\iota_Y w) (\log \mathcal{L})^{j-1} \mathcal{L}^{\frac{\log |\mu_{\mathcal{L}}|}{h_{\text{top}}(A)}}| \\ & \leq C \|\iota_Y w\|_{C^1(M)} [\log(1 + \mathcal{L})]^{\max(J_A^0, 1) + 1}. \end{aligned}$$

The above inequality immediately implies that $(\iota_Y B_1^+)(w) = B_1^+(\iota_Y w) = 0$, then by the lower bound on coefficients $c_{i,j}(x, \mathcal{L})$ given in Theorem 1.1, it also follows by finite induction on (i, j) , with respect to the lexicographic order (such that $(i, j) < (i', j')$ iff $i < i'$ or $i = i'$ and $j > j'$) that

$$(\iota_Y B_{i,j}^+)(w) = B_{i,j}^+(\iota_Y w) = 0, \quad \text{for all } i \in \{2, \dots, k\}, j \in \{1, \dots, J_i\}.$$

Since w is an arbitrary 2-form of class C^1 , the Addendum is proved in the case of the unstable foliation. The statement for the stable foliation follows by considering the case of the unstable foliation for the inverse map A^{-1} .

□

Finally, we detail the construction of asymptotic functionals:

Proof of Theorem 1.4. Let $\tilde{\gamma}$ be any unstable arc. We claim that the limit

$$\beta^+(\tilde{\gamma}) := \lim_{k \rightarrow \infty} A_*^{-k} \mathcal{B}^+((A^\#)^{\lceil \frac{\log(\mathcal{L}^+(A^k \tilde{\gamma}))}{h_{top}(A)} \rceil} (\mathcal{C}^+(A^k \tilde{\gamma})))$$

exists. In fact, this claim is easily reduced to existence of the limit

$$\mathcal{C}_\infty(\tilde{\gamma}) := \lim_{n \rightarrow \infty} \mathcal{C}^+(A^n \tilde{\gamma}).$$

It follows from the definition of the map $\mathcal{C}^+ : \Gamma^+ \rightarrow E^+$ in formula (11) that there exists a sequence $\{x_{\ell,m} | \ell \in \mathbb{N}, 1 \leq m \leq m_\ell\} \subset \tilde{\gamma}$ such that, for all $k \in \mathbb{N}$,

$$\begin{aligned} \mathcal{C}^+(A^k \tilde{\gamma}) &= \sum_{\ell=1}^{n+k} \sum_{m=1}^{m_\ell} (A^\#)^{-\lceil \frac{\log(\mathcal{L}^+(A^k \tilde{\gamma}))}{h_{top}(A)} \rceil - \ell - k} [\gamma(A^{-\ell}(x_{\ell,m}))]^+ \\ &= \sum_{\ell=1}^{n+k} \sum_{m=1}^{m_\ell} (A^\#)^{-\lceil \frac{\log(\mathcal{L}^+(\tilde{\gamma}))}{h_{top}(A)} \rceil - \ell} [\gamma(A^{-\ell}(x_{\ell,m}))]^+. \end{aligned}$$

Since the cohomology classes $[\gamma(A^{-\ell}(x_{\ell,m}))]$ are uniformly bounded, the limit of the above sequence exists and it is equal to the sum of a convergent series, that is,

$$\mathcal{C}_\infty(\tilde{\gamma}) := \sum_{\ell=1}^{+\infty} \sum_{m=1}^{m_\ell} (A^\#)^{-\lceil \frac{\log(\mathcal{L}^+(\tilde{\gamma}))}{h_{top}(A)} \rceil - \ell} [\gamma(A^{-\ell}(x_{\ell,m}))]^+.$$

The scaling property follows since the definition is well-posed, in fact

$$\begin{aligned} \beta^+(A\gamma) &:= \lim_{k \rightarrow \infty} A_*^{-k} \mathcal{B}^+((A^\#)^{\lceil \frac{\log(\mathcal{L}^+(A^{k+1} \tilde{\gamma}))}{h_{top}(A)} \rceil} (\mathcal{C}^+(A^{k+1} \tilde{\gamma}))) \\ &= A_* \lim_{k \rightarrow \infty} A_*^{-(k+1)} \mathcal{B}^+((A^\#)^{\lceil \frac{\log(\mathcal{L}^+(A^{k+1} \tilde{\gamma}))}{h_{top}(A)} \rceil} (\mathcal{C}^+(A^{k+1} \tilde{\gamma}))) = A_* \beta^+(\gamma). \end{aligned}$$

It is clear by the definition that the map \mathcal{C}_∞ is invariant under (regular) stable holonomies of unstable arcs. As a consequence, it is possible to extend the functional \mathcal{B}^+ to any rectifiable arc by decomposition into finally many paths, which are equivalent to unstable paths under stable holonomies.

Finally, it follows immediately from Theorem 3.2 that there exists a constant $C > 0$ such that, for any unstable arc $\tilde{\gamma}$, we have

$$\|\tilde{\gamma} - \beta^+(\tilde{\gamma})\|_{-1} \leq C[\log(1 + \mathcal{L}^+(\tilde{\gamma}))]^{\max(J_A^0, 1) + 1}.$$

By the above-mentioned decomposition, we then derive that, for any $\gamma \in \Gamma_r$,

$$\|\gamma - \beta^+(\gamma)\|_{-1} \leq C(1 + \mathcal{L}^-(\gamma))[\log(1 + \mathcal{L}^+(\gamma))]^{\max(J_A^0, 1) + 1},$$

as stated. □

4. DEVIATIONS OF ERGODIC INTEGRALS AND RUELLE-POLLICOTT ASYMPTOTICS

In this final section, we complete the proofs of Corollary 1.5 on deviations of ergodic integrals for unstable vector fields and Corollary 1.7 on the Ruelle-Pollicott asymptotics.

We first verify our claim that, as a consequence of Addendum 1.3, the measure $D_1^X = B_1^+ \wedge \hat{X}$ and, for all $i \in \{2, \dots, k\}$ and all $j \in \{1, \dots, J_i\}$, the distributions $D_{i,j}^X = B_{i,j}^+ \wedge \hat{X}$ are X -invariant. Let B any basic current of degree and dimension 1 for the unstable foliation on $M \setminus \Sigma$. The current $B \wedge \hat{X}$ is a current of degree 2 and dimension 0. Since the contraction operator ι_X is surjective onto the space of functions, it is enough to prove that $\iota_X \mathcal{L}_X(B \wedge \hat{X}) = 0$, as the latter identity then implies $\mathcal{L}_X(B \wedge \hat{X}) = 0$. Indeed, since $\iota_X B = dB = 0$ and $\iota_X \hat{X} = 1$ we have

$$\iota_X \mathcal{L}_X(B \wedge \hat{X}) = \iota_X d\iota_X(B \wedge \hat{X}) = -\iota_X dB = 0.$$

Thus our claim is proved. We then proof our result on deviations of ergodic averages.

Proof of Corollary 1.5. For all $x \in M \setminus \Sigma$, let $\gamma_T^X(x)$ denote the orbit of the unstable vector flow $h_{\mathbb{R}}^X$ on $M \setminus \Sigma$ (which we assume defined). Let $\mathcal{L}_T(x)$ denote the Margulis unstable measure of the orbit $\gamma_T^X(x)$, that is, according to the notations in the statement of Theorem 1.1 we have

$$\gamma_T^X(x) = \gamma_{\mathcal{L}_T(x)}(x).$$

For any $f \in C^1(M)$, let $\eta_f := f\hat{X}$. By Theorem 1.1, formula (2) and our definitions, since

$$\int_{\gamma_{\mathcal{L}}(x)} \eta_f = \int_0^T f \circ h_t^X(x) dt,$$

we have the expansion (for some constant $C_X > 0$):

$$\begin{aligned} (13) \quad & \left| \int_0^T f \circ h_t^X(x) dt - \mathcal{L}_T(x) D_1^X(f) \right. \\ & \left. - \sum_{i=2}^k \sum_{j=1}^{J_i} c_{i,j}(x, \mathcal{L}_T(x)) D_{i,j}^X(f) (\log \mathcal{L}_T(x))^{j-1} \mathcal{L}_T(x)^{\frac{\log |\mu_i|}{h_{op}(A)}} \right| \\ & \leq C_X \|f\|_{C^1(M)} [\log(1 + \mathcal{L}_T(x))]^{\max(J_A^0, 1) + 1}. \end{aligned}$$

For the case of the constant function $f = 1$ we have that, for some constant $C'_X > 0$,

$$\begin{aligned} (14) \quad & \left| T - \mathcal{L}_T(x) D_1^X(1) - \sum_{i=2}^k \sum_{j=1}^{J_i} c_{i,j}(x, \mathcal{L}_T(x)) D_{i,j}^X(1) (\log \mathcal{L}_T(x))^{j-1} \mathcal{L}_T(x)^{\frac{\log |\mu_i|}{h_{op}(A)}} \right| \\ & \leq C'_X [\log(1 + \mathcal{L}_T(x))]^{\max(J_A^0, 1) + 1}. \end{aligned}$$

The statement then follows from formulas (13) and (14). Indeed, by formula (14) the ratio $\mathcal{L}_T(x)/T$ is uniformly bounded from above and below, so that the coefficients

$$c_{ij}^X(x, T) := c_{i,j}(x, \mathcal{L}_T(x)) \left(\frac{\log(\frac{\mathcal{L}_T(x)}{T})}{\log T} + 1 \right)^{j-1} \left(\frac{\mathcal{L}_T(x)}{T} \right)^{\frac{\log |\mu_i|}{h_{\text{top}}(A)}}$$

are uniformly bounded from above, and from formula (13), for any function f of class C^1 , of zero average with respect to the invariant measure μ_X , we have

$$(15) \quad \left| \int_0^T f \circ h_t^X(x) dt - \sum_{i=2}^k \sum_{j=1}^{J_i} c_{i,j}^X(x, T) D_{i,j}^X(f) (\log T)^{j-1} T^{\frac{\log |\mu_i|}{h_{\text{top}}(A)}} \right| \\ \leq C_X \|f\|_{C^1(M)} \left[\log \left(1 + \frac{\mathcal{L}_T(x)}{T} T \right) \right]^{\max(J_A^0, 1) + 1}.$$

For general functions of class C^1 the conclusion follows by considering their projections onto the subspace of functions of zero average. \square

Proof of Corollary 1.7. Since A is of class C^r with $r > 2$, its unstable foliation is of class $C^{1+\alpha}$ for $\alpha > 0$. Let $\phi^+ : M \times \mathbb{R} \rightarrow M$ the flow of class $C^{1+\alpha}$ defined by following the (one-dimensional) unstable leaves with the speed determined by the Margulis measure. In more precise terms, given an orientation of the unstable leaves, for every $(x, t) \in M \times \mathbb{R}$ the point $\phi_t^+(x)$ is the unique point which belongs to the unstable leaf $\mathcal{F}^+(x)$ of x , such that $\mathcal{L}^+(x, \phi_t^+(x)) = |t|$ with $[x, \phi_t^+(x)]$ oriented in positive or negative direction depending on the sign of $t \neq 0$.

By definition the flow ϕ^+ preserves the Margulis measure, so that we have

$$\langle f \circ A^n, g \rangle = \frac{1}{\sigma} \int_0^\sigma \langle f \circ A^n \circ \phi_t^+, g \circ \phi_t^+ \rangle dt.$$

By integration by parts we have the formula

$$(16) \quad \int_0^\sigma \langle f \circ A^n \circ \phi_t^+, g \circ \phi_t^+ \rangle dt = \left\langle \int_0^\sigma f \circ A^n \circ \phi_t^+ dt, g \circ \phi_\sigma^+ \right\rangle \\ - \int_0^t \left\langle \int_0^s f \circ A^n \circ \phi_\tau^+ d\tau, \frac{d}{ds}(g \circ \phi_s^+) \right\rangle ds$$

Since the unstable curves are parametrized by the Margulis measure, by change of variables we have

$$(17) \quad \int_0^s f \circ A^n \circ \phi_\tau^+ d\tau = \frac{1}{\lambda^n} \int_0^{\lambda^n s} f \circ \phi_\tau^+ \circ A^n d\tau.$$

Let then η^+ denote a 1-form of class $C^{1+\alpha}$ such that $\eta^+(\frac{d}{dt}\phi_t^+) \equiv 1$. The 1-form η^+ restricts to the oriented Margulis measure on unstable leaves and can be taken to have kernel tangent to the stable foliation. By this definition we have, for all $x \in M$, $n \in \mathbb{N}$ and $s \in \mathbb{R}$, and for all $f \in C^1(M)$,

$$\int_0^{\lambda^n s} f \circ \phi_\tau^+ \circ A^n(x) d\tau = \int_{\gamma_{\lambda^n s}(A^n(x))} f \eta^+.$$

By Theorem 1.1 and by formula (17) we have

$$\begin{aligned} \left| \int_0^{\lambda^n s} f \circ A^n \circ \phi_\tau^+(x) d\tau - \frac{1}{\lambda^n} \sum_{i=1}^k \sum_{j=1}^{J_i} c_{i,j}(x, \lambda^n s) \log(\lambda^n s)^{j-1} (\lambda^n s)^{\frac{\log|\mu_j|}{\text{htop}(A)}} B_{i,j}^+(f\eta^+) \right| \\ \leq C \|f\|_{C^1(M)} \frac{[\log(1 + \lambda^n s)]^{\max(J_A^0, 1)+1}}{\lambda^n}. \end{aligned}$$

We can assume that the function f has zero average, hence we have

$$B_1^+(f\eta^+) = \int_M f \mathcal{M}^- \otimes \eta^+ = \int_M f d\mathcal{M} = 0,$$

so that by inserting the above asymptotics in formula (16) we derive the desired expansion for correlations. □

REFERENCES

- [Ba] V. Baladi, There are no deviations for the ergodic averages of the Giulietti-Liverani horocycle flows on the two-torus. Preprint, arXiv:1907.03453.
- [Bu14] A. Bufetov, Limit theorems for translation flows, *Ann. of Math.* **179** (2) (2014), 431–499.
- [BuFo14] A. Bufetov & G. Forni, Limit theorems for horocycle flows, *Ann. Sci. ENS* **47** (5) (2014), 851–903.
- [DHL14] V. Delecroix, P. Hubert & S. Lelièvre, Diffusion for the periodic wind-tree model, *Ann. Sci. ENS* **47** (6) (2014), 1085–1110.
- [FGL] F. Faure, S. Gouëzel & E. Laneeau, Ruelle spectrum of linear pseudo-Anosov maps, *Journal de l'École polytechnique–Mathématiques* **6** (2019), 811–877.
- [F02] G. Forni, Deviation of Ergodic Averages for Area-Preserving Flows on Surfaces of Higher Genus, *Annals of Mathematics* **155** (1) (2002), 1–103.
- [F07] G. Forni, Sobolev regularity of solutions of the cohomological equation, *Ergodic Theory and Dynamical Systems*, 1–105. doi:10.1017/etds.2019.108.
- [FoKa] G. Forni & A. Kanigowski, Time-Changes of Heisenberg nilflows, in “Some aspects of the theory of dynamical systems: a tribute to Jean-Christophe Yoccoz (volume II)” (S. Crovisier, R. Krikorian, Carlos Matheus, S. Senti editors), *Astérisque* **416** (2020), 253–299.
- [GL] P. Giulietti & C. Liverani, Parabolic dynamics and Anisotropic Banach spaces, *Journal Europ. Math. Soc.* **21** (9) (2019), 2793–2858. Published online: 2019-05-20 DOI: 10.4171/JEMS/892.
- [MY16] S. Marmi & J.-C. Yoccoz, Hölder Regularity of the Solutions of the Cohomological Equation for Roth Type Interval Exchange Maps, *Commun. Math. Phys.* **344** (1) (2016) 117–139. <https://doi.org/10.1007/s00220-016-2624-9>.
- [Yoc10] J.-C. Yoccoz, Interval exchange maps and translation surfaces, in *Homogeneous Flows, Moduli Spaces and Arithmetic*, Clay Math. Proc. **10**, Amer. Math. Soc., Providence, RI, 2010, 1–69.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK, MD USA
E-mail address: gforni@math.umd.edu