

Information-theoretic convergence of extreme values to the Gumbel distribution

Oliver Johnson*

July 8, 2020

Abstract

We show how convergence to the Gumbel distribution in an extreme value setting can be understood in an information-theoretic sense. We introduce a new type of score function which behaves well under the maximum operation, and which implies simple expressions for entropy and relative entropy. We show that, assuming certain properties of the von Mises representation, convergence to the Gumbel can be proved in the strong sense of relative entropy.

1 Introduction and notation

It is well-known that convergence to the Gaussian distribution in the Central Limit Theorem regime can be understood in an information-theoretic sense, following the work of Stam [18], Blachman [4], Brown [5], and in particular Barron [1] who proved convergence in relative entropy (see [9] for an overview of this work). While a traditional characteristic function proof of the Central Limit Theorem may not give a particular insight into why the Gaussian is the limit, this information-theoretic argument (which can be understood to relate to Stein’s method [19]) offers an insight on this.

To be specific, we can understand this convergence through the (Fisher) score function with respect to location parameter $\rho_X(x) = f'_X(x)/f_X(x) = (\log f_X(x))'$ of a random variable X with density f_X , where $'$ represents the spatial derivative. Two key observations are (i) that a standard Gaussian random variable Z is characterized by having linear score $\rho_Z(x) = -x$ and (ii) there is a closed form expression for the score of the sum of independent random variables as a conditional expectation (projection) of the scores of the individual summands (see e.g. [4]). As a result of this, the score function becomes ‘more linear’ in the Central Limit Theorem regime (see [11, 10]). Similar arguments can be used to understand ‘law of small numbers’ convergence to the Poisson distribution [12].

*School of Mathematics, University of Bristol, Fry Building, Woodland Road, Bristol, BS8 1UG, UK
E-mail: O.Johnson@bristol.ac.uk

However, there exist other families of probabilistic limit theorems which we would like to understand in a similar framework. In this paper we will consider a standard extreme value theory setup [16]: that is, we take i.i.d. random variables $X_1, X_2, \dots \sim X$ and define $M_n = \max(X_1, \dots, X_n)$ and $N_n = (M_n - b_n)/a_n$ for some sequences a_n and b_n . We want to consider whether N_n converges in relative entropy to a standard extreme value distribution.

In this paper we show how to do this for the case of a Gumbel (Type I Extreme Value) limit, by introducing a different type of score function, which we refer to as the max-score Θ_X , which is designed for this problem. Corresponding properties to those described above hold for this new quantity: (i) a Gumbel random variable X can be characterized by having linear max-score Θ_X (see Example 1.2) (ii) there is a closed form expression (Lemma 1.3) for the max-score of the maximum of independent random variables.

In Section 2 we show that the entropy and relative entropy can be expressed in terms of the max-score, and in Section 3 we relate this to the standard von Mises representation (see [16, Chapter 1]) to deduce convergence in relative entropy in Theorem 3.4. Our aim is not to provide a larger class of random variables than papers such as [6, 7, 14] for which convergence to the Gumbel takes place, but rather to use ideas from information theory to understand why this convergence may be seen as natural, and to prove convergence in a strong (relative entropy) sense.

We briefly remark that entropy was studied in this regime by Saeb [17], though using a direct computation based on the density. The standard Fisher score was used in an extreme value context by Bartholmé and Swan [2] in a version of Stein's method. Extreme value distributions were considered in the context of Tsallis entropy by Bercher and Vignat [3]. However, this particular framework appears to be new, to the best of our knowledge.

Definition 1.1. *For continuous random variable $Z \in \mathbb{R}$ we write distribution function $F_Z(z) = \mathbb{P}(Z \leq z)$ and $f_Z(z) = F'_Z(z)$ for the corresponding density. We define the max-score function:*

$$\Theta_Z(z) = \log(f_Z(z)/F_Z(z)). \quad (1)$$

Note that we can write $F_X(x) = \exp(-w(x))$ for some decreasing function w with $w(\infty) = 0$. Then $f_X(x) = -w'(x)F_X(x)$ so that $\Theta_X(x) = \log(-w'(x))$. Equivalently, inverting this relationship gives

$$F_X(x) = \exp\left(-\int_x^\infty e^{\Theta_X(u)} du\right). \quad (2)$$

We now remark that under this definition the Gumbel distribution has linear max-score function:

Example 1.2. *A Gumbel random variable Y with parameters μ and β has distribution function $F_Y(y) = \exp(-e^{-(y-\mu)/\beta})$, so in the notation above $w(y) = e^{-(y-\mu)/\beta}$. Hence $w'(y) = -e^{-(y-\mu)/\beta}/\beta$ and a Gumbel random variable has max-score*

$$\Theta_Y(y) = -\log \beta - \frac{(y - \mu)}{\beta}.$$

Indeed, using (2), we can see the property of having a linear max-score Θ_Y characterizes the Gumbel.

For future reference in this paper, note that (see [13, Eq. (1.25)]) the $\mathbb{E}Y = \mu + \beta\gamma$, where γ is the Euler–Mascheroni constant, and $\mathcal{M}_Y(t) = e^{\mu t}\Gamma(1-\beta t)$ (see [13, Eq. (1.23)]).

We can further state how the max-score function behaves under the maximum and rescaling operations:

Lemma 1.3. *If we write $M_n = \max(X_1, \dots, X_n)$ and $N_n = (M_n - b_n)/a_n$ then*

$$\Theta_{N_n}(z) = \log(na_n) + \Theta_X(a_n z + b_n), \quad (3)$$

$$\Theta_{N_n}(N_n) = \log(na_n) + \Theta_X(M_n), \quad (4)$$

Proof. As usual (see for example [16, Chapter 0.3]), we know that by independence

$$F_{M_n}(x) = \mathbb{P}(\max(X_1, \dots, X_n) \leq x) = \mathbb{P}\left(\bigcap_{i=1}^n \{X_i \leq x\}\right) = F_X(x)^n, \quad (5)$$

so that $F_{N_n}(x) = F_{M_n}(a_n x + b_n) = F_X(a_n x + b_n)^n$. This means that $f_{N_n}(x) = na_n F_X(a_n x + b_n)^{n-1} f_X(a_n x + b_n)$, so $f_{N_n}(x)/F_{N_n}(x) = na_n f_X(a_n x + b_n)/F_X(a_n x + b_n)$ and (3) follows on taking logarithms. The second result, (4), follows by direct substitution using the fact that $M_n = a_n N_n + b_n$. \square

Example 1.4. *In particular, if X is exponential with parameter λ , so with $a_n = 1/\lambda$ and $b_n = \log n/\lambda$*

$$\Theta_X(z) = \log\left(\frac{\lambda e^{-\lambda z}}{1 - e^{-\lambda z}}\right),$$

this gives

$$\Theta_{N_n}(z) = \log(n/\lambda) + \log\left(\frac{\lambda e^{-z}/n}{1 - e^{-z}/n}\right) = -z - \log(1 - e^{-z}/n).$$

Hence, letting $n \rightarrow \infty$, we know that $\Theta_{N_n}(z)$ converges pointwise to $-z$, which is the max-score of the standard Gumbel (with parameters $\mu = 0$ and $\beta = 1$) – see Example 1.2 above.

However, while this gives us some intuition as to why the Gumbel is the limit in this case, pointwise convergence of the score function does not seem a particularly strong sense of convergence. We now discuss the question of convergence in relative entropy.

2 Max-score function and entropy

We next show that we can use the max-score function to give an alternative formulation for the entropy of a random variable, which allows us to quickly find the entropy of a Gumbel distribution. We first state a simple lemma:

Lemma 2.1. For any random variable X with distribution function F_X :

$$\mathbb{E} \log F_X(X) = \mathbb{E} \log (1 - F_X(X)) = -1.$$

Proof. By change of variables $y = F_X(x)$, since the $dy = f_X(x)dx$ the first expression becomes

$$\int_{-\infty}^{\infty} f_X(x) \log F_X(x) dx = \int_0^1 \log y dy = [y \log y - y]_0^1 = -1. \quad (6)$$

For the second expression the same change of variables gives

$$\int_{-\infty}^{\infty} f_X(x) \log (1 - F_X(x)) dx = \int_0^1 \log(1 - y) dy = [-(1 - y) \log(1 - y) - y]_0^1 = -1, \quad (7)$$

as stated. \square

We note a corollary which will be useful in due course:

Corollary 2.2. For M_n the maximum of n independent copies of X :

$$-\mathbb{E} \log F_X(M_n) = \frac{1}{n}$$

Proof. Recalling from (5) that $F_{M_n}(x) = F_X(x)^n$ we know from Lemma 2.1 that

$$-1 = \mathbb{E} \log F_{M_n}(M_n) = n \mathbb{E} \log F_X(M_n),$$

and the result follows on rearranging. \square

Proposition 2.3. For a random variable X with max-score function Θ_X , the entropy $H(X)$ satisfies

$$H(X) = 1 - \mathbb{E} \Theta_X(X). \quad (8)$$

Proof. The key observation is that $\log f_X(x) = \log F_X(x) + \Theta_X(x)$ so that:

$$\begin{aligned} H(X) &= - \int_{-\infty}^{\infty} f_X(x) \log f_X(x) dx \\ &= - \int_{-\infty}^{\infty} f_X(x) \log F_X(x) dx - \int_{-\infty}^{\infty} f_X(x) \Theta_X(x) dx \\ &= 1 - \mathbb{E} \Theta_X(X), \end{aligned} \quad (9)$$

where we apply Lemma 2.1 to find the first term of (9). \square

Example 2.4. In particular we recover the entropy of Y , a Gumbel distribution with parameters μ and β since $\mathbb{E}Y = \mu + \beta\gamma$ so that using Example 1.2:

$$H(Y) = 1 - \mathbb{E} \left(-\log \beta - \frac{Y - \mu}{\beta} \right) = 1 + \log \beta + \gamma.$$

(See for example [15, Theorem 1.6iii])

We can use similar arguments to give an expression for the relative entropy $D(X\|Y)$ where Y is Gumbel:

Proposition 2.5. *Given random variable X , we can write the relative entropy from X to Y , a Gumbel random variable with parameters μ and β , as*

$$D(X\|Y) = \left(\mathbb{E}\Theta_X(X) + \log \beta + \frac{\mathbb{E}X - \mu}{\beta} \right) + (\mathbb{E}e^{-(X-\mu)/\beta} - 1). \quad (10)$$

Proof.

$$\begin{aligned} D(X\|Y) &= \int f_X(x) \log \left(\frac{f_X(x)}{f_Y(x)} \right) dx \\ &= -H(X) - \int f_X(x) \log f_Y(x) dx \\ &= -(1 - \mathbb{E}\Theta_X(X)) - \int f_X(x) \log F_Y(x) dx - \int f_X(x) \Theta_Y(x) dx \\ &= \mathbb{E}\Theta_X(X) - 1 + \int f_X(x) \exp \left(-\frac{(x-\mu)}{\beta} \right) dx + \log \beta + \int f_X(x) \frac{(x-\mu)}{\beta} dx. \end{aligned}$$

substituting from Proposition 2.3 and using the value of Θ_Y from Example 1.2. \square

Observe that in the case of X itself Gumbel with the same parameters as Y , both bracketed terms in (10) vanish:

1. We can rewrite the first term as $\mathbb{E}(\Theta_X(X) - \Theta_Y(X))$, using the value of the max-score in the Gumbel case (Example 1.2). This suggests that (as in [9]) we may wish to consider this term as a standardized score function with the relevant linear term subtracted off.
2. We can rewrite the second term in (10) as $e^{\mu/\beta} \mathcal{M}_X(-1/\beta) - 1$, where $\mathcal{M}_X(t)$ is the moment generating function. Since (see Example 1.2) the moment generating function of a Gumbel random variable is $\mathcal{M}_Y(t) = e^{\mu t} \Gamma(1 - \beta t)$, we know that in the Gumbel case $e^{\mu/\beta} \mathcal{M}_X(-1/\beta) - 1 = e^{\mu/\beta} e^{-\mu/\beta} \Gamma(2) - 1 = 0$.

3 von Mises representation and convergence in relative entropy

We will demonstrate convergence of relative entropy in a restricted version of the domain of maximum attraction:

Definition 3.1. *Assume that the upper limit of the support of X is $x_0 := \sup\{x : F_X(x) < 1\}$ (which may be finite or infinite) and assume that the distribution function F_X has a von Mises representation [16, Eq. (1.3)]*

$$F_X(x) = 1 - c \exp \left(- \int_{z_0}^x \frac{1}{g(u)} du \right) = 1 - c \exp(-G(x)), \quad (11)$$

for some auxiliary function g such that $g'(u) \rightarrow 0$ as $u \rightarrow x_0$.

Example 3.2.

1. For the exponential we can take $c = 1$, $z_0 = 0$, $x_0 = \infty$, $g(u) = 1/\lambda$ and $b_n = \log n/\lambda$.
2. For the example of $F(x) = 1 - \exp(-x/(1-x))$ given by Gnedenko [8] (see also [16, P.39]) we can take $c = 1$, $z_0 = 0$, $x_0 = 1$, $g(u) = (1-u)^2$ and $b_n = \log n/(1 + \log n)$.

Note that (see [16, Proposition 1.4]) the normalized maximum N_n converges to the Gumbel if and only if the distribution function is of the form $F_X(x) = 1 - c(x) \exp(-G(x))$ where $\lim_{x \rightarrow x_0} c(x) = c$. However, the restricted class of distributions of Definition 3.1 is easier to deal with.

Assuming the von Mises representation (11) holds we can write the density

$$f_X(x) = cG'(x) \exp(-G(x)) = \frac{1 - F_X(x)}{g(x)}, \quad (12)$$

or note that $1/g(x)$ is the hazard function of X .

We now state a technical condition which we will use to prove convergence:

Condition 1.

1. Assume $\log g(x)$ is bounded and continuous, and that $g(x_0)$ is finite and non-zero.
2. Assume that $\int_{-\infty}^0 |x|^k dF_X(x) < \infty$ for all k .

Lemma 3.3. *Under Condition 1.2:*

1. The mean $\mathbb{E}N_n$ converges to the Euler–Mascheroni constant γ .
2. The moment generating function converges as follows:

$$\lim_{n \rightarrow \infty} \mathcal{M}_{N_n}(t) = \Gamma(1 - t), \quad (13)$$

by Taylor’s theorem.

Proof. Note that (see [16, Proposition 2.1]) under Condition 1.2 the k th moment of N_n converges:

$$\lim_{n \rightarrow \infty} \mathbb{E}(N_n)^k = (-1)^k \Gamma^{(k)}(1), \quad (14)$$

where $\Gamma^{(k)}(x)$ is the k th derivative of the Γ function at x .

We deduce convergence of the moment generating function

$$\lim_{n \rightarrow \infty} \mathcal{M}_{N_n}(t) = \lim_{n \rightarrow \infty} \sum_{r=0}^{\infty} \frac{t^r \mathbb{E}(N_n)^r}{r!} = \lim_{n \rightarrow \infty} \sum_{r=0}^{\infty} \frac{t^r \mathbb{E}(N_n)^r}{r!} = \sum_{r=0}^{\infty} \frac{(-t)^r \Gamma^{(k)}(1)}{r!} = \Gamma(1 - t),$$

by Taylor’s theorem. □

Theorem 3.4. *If the distribution function of X satisfies the von Mises representation (11) and Condition 1 holds, there exist norming constants a_n and b_n satisfying $\log(cn) = G(b_n)$ and $a_n = g(b_n)$ such that $N_n = (M_n - b_n)/a_n$ satisfies*

$$\lim_{n \rightarrow \infty} D(N_n \| Y) = 0,$$

where Y is a standard Gumbel distribution (with $\beta = 1$ and $\mu = 0$).

Proof. We use the norming constants from [16, Proposition 1.1(a)], that is $a_n = g(b_n)$ where b_n satisfies $1/n = 1 - F(b_n) = c \exp(-G(b_n))$. Taking logs and rearranging, this gives $\log(cn) = G(b_n)$.

We can give an alternative form of the max-score of F_X :

$$\begin{aligned} \Theta_X(x) &= \log(f_X(x)/F_X(x)) \\ &= \log(f_X(x)/(1 - F_X(x)) + \log(1 - F_X(x)) - \log F_X(x) \\ &= -\log g(x) + \log(1 - F_X(x)) - \log F_X(x) \end{aligned} \quad (15)$$

Then, using (4) from Lemma 1.3 and defining $H_n := 1 + 1/2 + 1/3 + \dots + 1/n$ to be the n th harmonic number we can write the first term in the relative entropy expression (10) in the case $\mu = 0$ and $\beta = 1$ as

$$\begin{aligned} \mathbb{E}\Theta_{N_n}(N_n) + \mathbb{E}N_n &= \log(na_n) + \mathbb{E}\Theta_X(M_n) + \mathbb{E}N_n \\ &= \log(na_n) - \mathbb{E}\log g(M_n) + \mathbb{E}\log(1 - F_X(M_n)) - \mathbb{E}\log F_X(M_n) + \mathbb{E}N_n \\ &= (\log g(b_n) - \mathbb{E}\log g(M_n)) + (\log n - H_n) + \frac{1}{n} + \mathbb{E}N_n \end{aligned} \quad (16)$$

where we use the expression $-\mathbb{E}\log F_X(M_n) = 1/n$ stated in Corollary 2.2 above, and use the expression $-\mathbb{E}\log(1 - F_X(M_n)) = H_n$ given in Lemma 3.5 below.

We can consider the behaviour as $n \rightarrow \infty$ of the four terms in (16) separately:

1. Using Condition 1.1 we know that $\log g(b_n) \rightarrow \log g(x_0)$ and since $M_n \rightarrow x_0$ in distribution we know that $\mathbb{E}\log g(M_n) \rightarrow \log g(x_0)$ by the portmanteau lemma.
2. It is a standard fact that $\log n - H_n$ is a monotonically increasing sequence which converges to $-\gamma$.
3. Clearly the first term converges to zero.
4. Lemma 3.3 tells us that the final term converges to γ .

Putting this all together, we deduce that (16) converges to

$$0 - \gamma + 0 + \gamma = 0.$$

In the case $\mu = 0$ and $\beta = 1$, the second term in the relative entropy expression (10) becomes

$$\mathbb{E}e^{-N_n} - 1 = \mathcal{M}_{N_n}(-1) - 1 \rightarrow \Gamma(2) - 1 = 0,$$

by (13). □

Lemma 3.5. *The expected value*

$$-\mathbb{E} \log(1 - F_X(M_n)) = H_n$$

is the same for all F_X .

Proof. By standard manipulations, we know that $-\log(1 - F_X(X))$ is exponential with parameter 1. This follows by direct substitution, since

$$\begin{aligned} \mathbb{P}(-\log(1 - F_X(X)) \leq z) &= \mathbb{P}(F_X(X) \leq 1 - \exp(-z)) \\ &= \mathbb{P}(X \leq F_X^{-1}(1 - \exp(-z))) = F_X(F_X^{-1}(1 - \exp(-z))) \\ &= 1 - \exp(-z), \end{aligned}$$

as required. Now, since $-\log(1 - F_X(t))$ is increasing in t , we can write

$$-\log(1 - F_X(M_n)) = -\log\left(1 - F_X\left(\max_{1 \leq i \leq n} X_i\right)\right) = \max_{1 \leq i \leq n} -\log(1 - F_X(X_i)) \sim \max_{1 \leq i \leq n} E_i,$$

where E_i are independent exponentials with parameter 1.

It is well-known that the expected value of $\max_{1 \leq i \leq n} E_i = H_n$, the n th harmonic number. The simplest proof of this is to write

$$\max_{1 \leq i \leq n} E_i = \sum_{i=1}^n U_i,$$

where U_i are independent exponentials with parameter $n - i + 1$. (This follows from the memoryless property of E_i , by thinking of U_1 as the time for the first exponential event to happen, U_2 as the time for the second, and so on). Since $\mathbb{E}U_i = 1/(n - i + 1)$, the result follows. \square

Corollary 3.6. *Assume that the distribution function F_X has a von Mises representation (11) whose auxiliary function g satisfies Condition 1. Then the entropy of the normalized maximum $N_n = (M_n - b_n)/a_n$ satisfies*

$$\lim_{n \rightarrow \infty} H(N_n) = 1 + \gamma,$$

which is the entropy of the corresponding Gumbel distribution.

Proof. By Proposition 2.3

$$\begin{aligned} H(N_n) &= 1 - \mathbb{E}\Theta_{N_n}(N_n) \\ &= (1 + \mathbb{E}N_n) - \mathbb{E}(\Theta_{N_n}(N_n) + N_n) \end{aligned}$$

The first term converges to $1 + \gamma$ by Lemma 3.3, the second term is precisely (16) and converges to zero as described in the proof of Theorem 3.4. \square

Remark 3.7. Note that for the exponential case of Example 3.2, Condition 1 is satisfied, so we can deduce convergence in relative entropy. Indeed, since g is constant in this case, the first term of (16) vanishes, meaning that we can deduce that $\mathbb{E}\Theta_{N_n}(N_n) = \log n - H_n + 1/n$ and the entropy is exactly

$$H(N_n) = 1 + H_n - \log n - 1/n, \quad (17)$$

which value may be of independent interest. Using the fact that $H_n = \log n + \gamma + 1/(2n) + O(1/n^2)$, we can deduce that $H(N_n) = 1 + \gamma - 1/(2n) + O(1/n^2)$. In the spirit of [11] and other papers, it may be of interest to ask under what conditions the convergence in Corollary 3.6 is at rate $O(1/n)$ in this way.

Theorem 3.4 shows that convergence in relative entropy occurs for a range of random variables that are ‘approximately exponential’ in some sense. However, observe that the Gnedenko example $g(u) = (1 - u)^2$ from Example 3.2 does not satisfy Condition 1.1, so Theorem 3.4 cannot be directly applied in this case. However, it is possible to deduce convergence in relative entropy in this example too, using a relatively simple adaption of the argument.

We use Condition 1.1 to control the term $(\log g(b_n) - \mathbb{E} \log g(M_n))$. We can do this under milder conditions as follows. Write $\psi(u) = \log g(u)$, then we can use the Taylor series around b_n to expand

$$\begin{aligned} \mathbb{E} \log g(M_n) - \log g(b_n) &= \sum_{k=1}^{\infty} \frac{\psi^{(k)}(b_n)}{k!} \mathbb{E}(M_n - b_n)^k \\ &= \sum_{k=1}^{\infty} \frac{1}{k!} (\psi^{(k)}(b_n) a_n^k) \mathbb{E} \left(\frac{M_n - b_n}{a_n} \right)^k \\ &= \sum_{k=1}^{\infty} \frac{1}{k!} (\psi^{(k)}(b_n) g(b_n)^k) \mathbb{E} N_n^k. \end{aligned} \quad (18)$$

In the specific case of $g(u) = (1 - u)^2$, notice that $\psi^{(k)}(u) = -2(k - 1)!/(1 - u)^k$, so that

$$(\psi^{(k)}(b_n) g(b_n)^k) = -2(k - 1)!(1 - b_n)^k,$$

and (18) becomes:

$$-2 \sum_{k=1}^{\infty} \frac{1}{k} (1 - b_n)^k \mathbb{E} N_n^k.$$

We can see that the $(1 - b_n)^k$ term converges to 0 since b_n converges to 1, and (14) gives that $\mathbb{E} N_n^k$ converges to a finite constant, and 18 (and hence relative entropy) converges to zero.

Clearly this argument will hold in greater generality, and we can deduce that convergence in relative entropy will hold for any random variable defined by an auxiliary function g such that $\psi^{(k)}(u)g(u)^k/k! \rightarrow 0$ as $u \rightarrow x_0$.

References

- [1] A. R. Barron. Entropy and the Central Limit Theorem. *Ann. Probab.*, 14(1):336–342, 1986.
- [2] C. Bartholmé and Y. Swan. Rates of convergence towards the Fréchet distribution. See [arXiv:1311.3896](https://arxiv.org/abs/1311.3896), 2013.
- [3] J.-F. Bercher and C. Vignat. An entropic view of Pickands’ theorem. In *2008 IEEE International Symposium on Information Theory (ISIT)*, pages 2625–2628. IEEE, 2008.
- [4] N. M. Blachman. The convolution inequality for entropy powers. *IEEE Trans. Information Theory*, 11:267–271, 1965.
- [5] L. D. Brown. A proof of the Central Limit Theorem motivated by the Cramér-Rao inequality. In G. Kallianpur, P. R. Krishnaiah, and J. K. Ghosh, editors, *Statistics and Probability: Essays in Honour of C.R. Rao*, pages 141–148. North-Holland, New York, 1982.
- [6] L. De Haan and A. Ferreira. *Extreme value theory: an introduction*. Springer Science & Business Media, 2007.
- [7] L. De Haan and S. I. Resnick. Local limit theorems for sample extremes. *The Annals of Probability*, pages 396–413, 1982.
- [8] B. Gnedenko. Sur la distribution limite du terme maximum d’une série aléatoire. *Annals of Mathematics*, pages 423–453, 1943.
- [9] O. T. Johnson. *Information theory and the Central Limit Theorem*. Imperial College Press, London, 2004.
- [10] O. T. Johnson. Maximal correlation and the rate of Fisher information convergence in the Central Limit Theorem. *IEEE Transactions on Information Theory (online first)*, 2020. doi:10.1109/TIT.2020.2985957.
- [11] O. T. Johnson and A. R. Barron. Fisher information inequalities and the Central Limit Theorem. *Probability Theory and Related Fields*, 129(3):391–409, 2004.
- [12] I. Kontoyiannis, P. Harremoës, and O. T. Johnson. Entropy and the law of small numbers. *IEEE Trans. Inform. Theory*, 51(2):466–472, 2005.
- [13] S. Kotz and S. Nadarajah. *Extreme value distributions: theory and applications*. World Scientific, 2000.
- [14] J. Pickands III. The continuous and differentiable domains of attraction of the extreme value distributions. *The Annals of Probability*, pages 996–1004, 1986.

- [15] S. Ravi and A. Saeb. A note on entropies of l -max stable, p -max stable, generalized Pareto and generalized log-Pareto distributions. In *ProbStat Forum*, volume 5, pages 62–79, 2012.
- [16] S. I. Resnick. *Extreme values, regular variation and point processes*. Springer, 2013.
- [17] A. Saeb. Rates of convergence for Rényi entropy in extreme value theory. See [arXiv:1402.4316](https://arxiv.org/abs/1402.4316), 2014.
- [18] A. J. Stam. Some inequalities satisfied by the quantities of information of Fisher and Shannon. *Information and Control*, 2:101–112, 1959.
- [19] C. Stein. A bound for the error in the normal approximation to the distribution of a sum of dependent random variables. In *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability, Volume 2: Probability Theory*. The Regents of the University of California, 1972.