

# STOCHASTIC FORMS OF BRUNN'S PRINCIPLE

PETER PIVOVAROV & JESÚS REBOLLO BUENO

ABSTRACT. A number of geometric inequalities for convex sets arising from Brunn's concavity principle have recently been shown to yield local stochastic formulations. Comparatively, there has been much less progress towards stochastic forms of related functional inequalities. We develop stochastic geometry of  $s$ -concave functions to establish local versions of dimensional forms of Brunn's principle à la Borell, Brascamp-Lieb, and Rinott. To do so, we define shadow systems of  $s$ -concave functions and revisit Rinott's approach in the context of multiple integral rearrangement inequalities.

## 1. INTRODUCTION

Brunn's concavity principle underpins a wealth of inequalities in geometry and analysis. One can formulate it as follows: for any convex body  $K \subseteq \mathbb{R}^n$  and any direction  $\theta$ , the  $(n-1)$ -volume of slices of  $K$  by parallel translates of  $\theta^\perp$  is  $1/(n-1)$ -concave on its support, i.e.,

$$A(t) = |K \cap (\theta^\perp + t\theta)|^{1/(n-1)} \quad (1.1)$$

is concave. A far-reaching extension of this principle in analysis is exemplified by a family of functional inequalities obtained by Borell [4] and Brascamp-Lieb [6], with an alternate approach put forth by Rinott [34]. These inequalities can be formulated in terms of certain means as follows: for  $a, b \geq 0$ ,  $s \geq -1/n$  and  $\lambda \in (0, 1)$ , set

$$\mathcal{M}_\lambda^s(a, b) = \begin{cases} (\lambda a^s + (1-\lambda)b^s)^{1/s} & \text{if } ab \neq 0 \\ 0 & \text{if } ab = 0, \end{cases}$$

where the cases  $s \in \{-1/n, 0, +\infty\}$  are defined as limits

$$\mathcal{M}_\lambda^{-1/n}(a, b) = \min\{a, b\}, \quad \mathcal{M}_\lambda^{+\infty}(a, b) = \max\{a, b\}, \quad \mathcal{M}_\lambda^0(a, b) = a^\lambda b^{1-\lambda}.$$

Then for measurable functions  $f, g, h : \mathbb{R}^n \rightarrow [0, \infty)$ ,  $0 < \lambda < 1$ , and  $s \geq -1/n$ , if

$$h(\lambda x_1 + (1-\lambda)x_2) \geq \mathcal{M}_\lambda^s(f(x_1), g(x_2)) \quad (1.2)$$

for all  $x_1, x_2 \in \mathbb{R}^n$ , one has

$$\int_{\mathbb{R}^n} h \geq \mathcal{M}_\lambda^{s/(1+ns)} \left( \int_{\mathbb{R}^n} f, \int_{\mathbb{R}^n} g \right). \quad (1.3)$$

The Prékopa-Leindler inequality [22, 31, 32] corresponds to the logarithmically concave case  $s = 0$ ; for earlier work on the real line, see Henstock and Macbeath [20]. These principles are now fundamental in analysis, geometry and probability, among other fields. For their considerable impact, we refer the reader to [16] and the references therein.

The inequalities (1.3) stem from principles rooted in convexity. Indeed, the standard approach to Brunn’s principle (1.1) connects concavity of the map  $t \mapsto A(t)$  to the convexity of  $K$  through suitable symmetrizations, see e.g., [2]. In [5], Brascamp and Lieb used symmetrization to prove certain cases of (1.3). Subsequently, they provided an alternate inductive approach, based on the Brunn-Minkowski inequality [6]. Rinott provided an alternate proof, starting with epigraphs of convex functions [34]. However, the inequalities ultimately do not require convexity as they hold for measurable functions. So convexity (or concavity) of the functions involved seems of no importance. In this paper, our focus is on what more can be said when the functions involved do possess some concavity.

Our motivation stems from recent work for convex sets in which a “local” stochastic dominance accompanies an isoperimetric principle. A concrete example, which can be derived via Brunn’s principle, is the Blaschke-Santaló [38] inequality. The latter says that the volume of the polar of an origin-symmetric convex body  $K$  is maximized by a Euclidean ball  $B$  under a constraint of equal volume. Proofs via symmetrization depend on variants of (1.3), e.g., Meyer-Pajor [26] and Campi-Gronchi [11]. In [14], the first-named author, together with Cordero-Erausquin, Fradelizi and Paouris, proved a stochastic version in which the dominance applies “locally” to random polytopes that naturally approximate  $K$  and  $B$  from within. By repeated sampling, this recovers the Blaschke-Santaló inequality by the law of large numbers. This example is indicative of recent research on isoperimetric inequalities: when an isoperimetric principle for convex sets can be proved by symmetrization, it is fruitful to instead carry out the symmetrization on product probability spaces. Multiple integral rearrangement inequalities of Rogers [35], Brascamp-Lieb-Luttinger [7], and Christ [13] then enter the picture and yield stronger stochastic formulations. This builds on principles in stochastic geometry e.g., [9, 10, 19]; see [14, 28, 29, 33] for further background.

While there is much work on stochastic isoperimetric inequalities for convex sets, there are far fewer results about random functions. In [30], we




FIGURE 1. Stochastic approximation of a log-concave function  $f$  by its least log-concave majorant  $[f]_N$  above a random sample under the graph of  $f$ .

initiated work on the Prékopa-Leindler inequality for random log-concave functions. Here we will show that the full family of inequalities (1.3) actually have “local” stochastic strengthenings for functions  $f$  that are  $s$ -concave, i.e.  $f^s$  is concave on its support; when  $s < 0$ , this means that  $f^s$  is convex. To formulate our main result, for each  $s$ -concave function  $f$ , we sample independent random vectors  $(X_1, Z_1), \dots, (X_N, Z_N)$  distributed uniformly under the graph of  $f$  according to Lebesgue measure. We associate random functions  $[f]_N$ , supported on the convex hull  $\text{conv}\{X_1, \dots, X_N\}$ , defined by

$$[f]_N(x) = \begin{cases} \inf\{z^{1/s} : (x, z) \in \mathcal{P}_{f,N}\}, & \text{if } s < 0 \\ \sup\{e^z : (x, z) \in \mathcal{P}_{f,N}\}, & \text{if } s = 0, \\ \sup\{z^{1/s} : (x, z) \in \mathcal{P}_{f,N}\}, & \text{if } s > 0, \end{cases}$$

where

$$\mathcal{P}_{f,N} = \begin{cases} \text{conv}\{(X_1, Z_1^s), \dots, (X_N, Z_N^s)\}, & \text{if } s \neq 0. \\ \text{conv}\{(X_1, \log Z_1), \dots, (X_N, \log Z_N)\}, & \text{if } s = 0. \end{cases}$$

In other words, when  $s = 0$  or  $s > 0$ ,  $[f]_N$  is the least log-concave or  $s$ -concave function, respectively, satisfying  $[f]_N(X_i) \geq Z_i$ ; similarly, when  $s < 0$ ,  $[f]_N$  is the greatest  $s$ -concave function with  $[f]_N(X_i) \leq Z_i$ . See Figure 1 for the case  $s = 0$ .

With this notation, we can state our main result, which we formulate in terms of the sup-convolution

$$(f \star_{\lambda,s} g)(v) = \sup\{\mathcal{M}_\lambda^s(f(x_1), g(x_2)) : v = \lambda x_1 + (1 - \lambda)x_2\}$$

and the symmetric decreasing rearrangements  $f^*$  and  $g^*$  of  $f$  and  $g$ , respectively (defined in (2.1)).

**Theorem 1.1.** *Let  $s \in (-1/n, \infty)$  and let  $f, g : \mathbb{R}^n \rightarrow [0, \infty)$  be integrable  $s$ -concave functions and  $N, M > n + 1$ . Then for  $\alpha > 0$ ,*

$$\mathbb{P} \left( \int_{\mathbb{R}^n} ([f]_N \star_{\lambda, s} [g]_M)(v) dv > \alpha \right) \geq \mathbb{P} \left( \int_{\mathbb{R}^n} ([f^*]_N \star_{\lambda, s} [g^*]_M)(v) dv > \alpha \right).$$

When  $N, M \rightarrow \infty$  one gets

$$\int_{\mathbb{R}^n} (f \star_{\lambda, s} g)(v) dv \geq \int_{\mathbb{R}^n} (f^* \star_{\lambda, s} g^*)(v) dv. \quad (1.5)$$

As mentioned, Brascamp and Lieb's first approach to cases of (1.3) used rearrangements. Recent interest in rearranged strengthenings for this and other means in (1.3) have been studied by Melbourne [25] for general functions. Roysdon and Xing have studied  $L_p$  variants of the Borell-Brascamp-Lieb inequality [37]. Our treatment will also allow for other means (see Remark 8.1). We focus on  $s$ -concave functions because in this case there is a stronger local stochastic dominance.

Special cases of Theorem 1.1, namely  $s = 1/q$  ( $q \in \mathbb{N}$ ) were treated in [30]. The approach used multiple integral rearrangement inequalities (as discussed above) and, additionally, built on ideas of Artstein-Avidan, Klartag and Milman [3] on moving from convex sets to log-concave functions. Here a new key step is inspired by Rinott's approach to (1.3) via epigraphs of convex functions [34]; the latter has recently been used in a dual setting by Artstein-Avidan, Florentin and Segal [1] for a new Prékopa-Leindler inequality.

Another new tool developed in this paper is that of linear parameter systems for  $s$ -concave functions. Linear parameter systems along a direction  $\theta \in \mathbb{S}^{n-1}$  are families of convex sets of the form

$$K_t = \text{conv}\{x_i + \lambda_i t \theta : i \in I\}, \quad t \in [a, b],$$

where  $I$  is an index set,  $\{x_i\}_{i \in I} \subseteq \mathbb{R}^n$  and  $\{\lambda_i\}_{i \in I} \subseteq \mathbb{R}$  are bounded sets. Rogers and Shephard [36] proved the fundamental fact that the volume of  $K_t$  is a convex function of  $t$ . This was extended to the more general notion of shadow systems by Shephard [41]. Shadow systems incorporate key features of Steiner symmetrization and have been successfully used in a variety of isoperimetric type inequalities, developed especially by Campi and Gronchi, e.g., [11, 12]; for other examples, see [15, 39, 27, 43] or [40] and the references therein.

The proof of Theorem 1.1 relies on extending linear parameter systems to  $s$ -concave functions. Let  $I$  be an index set, and  $\{(x_i + \lambda_i t \theta, z_i)\}_{i \in I} \subseteq \mathbb{R}^n \times [0, \infty)$  a collection of points lying under the graph of an integrable  $s$ -concave function and set  $w_i(t) = x_i + \lambda_i t \theta$ ,  $i \in I$ . Analogous to the definition of  $[f]_N$ , we define, for  $s \geq 0$ ,  $f_{t,s}$  to be the least  $s$ -concave function above

$\{w_i(t)\}$ ; similarly, for  $s < 0$ , we define  $f_{t,s}$  to be the greatest  $s$ -concave function beneath the points  $\{w_i(t)\}$  (see §5). In this setting, we show that

$$t \mapsto \int_{\mathbb{R}^n} f_{t,s}(v) dv$$

is convex. Just as linear parameter systems can be viewed as special shadow systems, the same applies to  $s$ -concave functions. We also give an interpretation of shadow systems of  $s$ -concave functions in terms of associated epigraphs and hypographs and establish related convexity properties in §5. We show in §7 how these interface with rearrangement inequalities and thus provide a path towards stochastic geometry of  $s$ -concave functions and associated extremal inequalities.

## 2. PRELIMINARIES

We will denote by  $\{e_1, \dots, e_n\}$  the standard basis in  $\mathbb{R}^n$ . Let  $K$  be a compact, convex set in  $\mathbb{R}^n$ ,  $\theta$  on the unit sphere  $\mathbb{S}^{n-1}$  and  $P := P_{\theta^\perp}$  the orthogonal projection onto  $\theta^\perp$ . We define  $u_K : PK \rightarrow \mathbb{R}$  by

$$u_K(x) := u(K, x) := \sup\{\lambda : x + \lambda\theta \in K\}$$

and  $\ell_K : PK \rightarrow \mathbb{R}$  by

$$\ell_K(x) := \ell(K, x) := \inf\{\lambda : x + \lambda\theta \in K\}.$$

Notice that  $u_K$  and  $\ell_K$  are concave and convex, respectively.

We recall that the Steiner symmetral of a non-empty compact set  $A \subseteq \mathbb{R}^n$  with respect to  $\theta^\perp$ ,  $S_{\theta^\perp}A$ , is the set with the property that for each line  $l$  orthogonal to  $\theta^\perp$  and meeting  $A$ , the set  $l \cap S_{\theta^\perp}A$  is a closed segment with midpoint on  $\theta^\perp$  and length equal to that of the set  $l \cap A$ . The mapping  $S_{\theta^\perp} : A \rightarrow S_{\theta^\perp}A$  is called the Steiner symmetrization of  $A$  with respect to  $\theta^\perp$ . In particular, if  $K$  is a convex body

$$S_{\theta^\perp}K = \{x + \lambda\theta : x \in PK, -\frac{u_K(x) - \ell_K(x)}{2} \leq \lambda \leq \frac{u_K(x) - \ell_K(x)}{2}\}.$$

This shows that  $S_{\theta^\perp}K$  is convex, since the function  $u_K - \ell_K$  is concave. Moreover,  $S_{\theta^\perp}K$  is symmetric with respect to  $\theta^\perp$ , it is closed, and by Fubini's theorem it has the same volume as  $K$ .

Let  $A \subseteq \mathbb{R}^n$  be a Borel set with finite Lebesgue measure. The symmetric rearrangement,  $A^*$ , of  $A$  is the open ball with center at the origin whose volume is equal to the measure of  $A$ . Since we choose  $A^*$  to be open,  $\mathbb{1}_{A^*}$  is lower semicontinuous. The symmetric decreasing rearrangement of  $\mathbb{1}_A$  is defined by  $\mathbb{1}_A^* = \mathbb{1}_{A^*}$ . We say a Borel measurable function  $f : \mathbb{R}^n \rightarrow [0, \infty)$  vanishes at infinity if for every  $t > 0$ , the set  $\{x \in \mathbb{R}^n : f(x) > t\}$  has finite

Lebesgue measure. In such a case, the symmetric decreasing rearrangement  $f^*$  is defined by

$$f^*(x) = \int_0^\infty \mathbb{1}_{\{f>t\}}^*(x) dt = \int_0^\infty \mathbb{1}_{\{f>t\}}(x) dt. \quad (2.1)$$

Observe that  $f^*$  is radially symmetric, radially decreasing, and equimeasurable with  $f$ , i.e.,  $\{f > t\}$  and  $\{f^* > t\}$  have the same measure for each  $t > 0$ . Let  $\{e_1, \dots, e_n\}$  be an orthonormal basis of  $\mathbb{R}^n$  such that  $e_1 = \theta$ . Then, for  $f$  vanishing at infinity, the Steiner symmetral  $f(\cdot|\theta)$  of  $f$  with respect to  $\theta^\perp$  is defined as follows: set  $f_{(x_2, \dots, x_n), \theta}(t) = f(t, x_2, \dots, x_n)$  and define  $f^*(t, x_2, \dots, x_n|\theta) := (f_{(x_2, \dots, x_n), \theta})^*(t)$ . In other words, we obtain  $f^*(\cdot|\theta)$  by rearranging  $f$  along every line parallel to  $\theta$ . We refer to the books [23, 42] or the introductory notes [8] for further background material on rearrangement of functions.

### 3. RINOTT'S APPROACH TO $s$ -CONCAVE FUNCTIONS VIA EPIGRAPHS AND HYPOGRAPHS

Rinott [34] provides a geometric proof of the Borell-Brascamp-Lieb inequalities (1.3) by deriving integral inequalities for functions using certain higher-dimensional measures. We start this section by recalling his approach.

A function  $f : \mathbb{R}^n \rightarrow [0, \infty)$  is called log-concave if  $\log f$  is concave on its support. We also use the notion of  $s$ -concavity as in [4] meaning that  $f$  is  $s$ -concave if  $f^s$  is concave on its support; this differs from other uses of this term [3, 21, 30]. Any  $s$ -concave function, for  $s > 0$ , is also log-concave. For  $A \subseteq \mathbb{R}^n$ , we define the epigraph of  $f$  on  $A$  in  $\mathbb{R}^n$  by

$$E_A(f) = \{(x, z) \in A \times \mathbb{R} : f(x) \leq z\}.$$

Analogously we define the hypograph of  $f$  on  $A$  by

$$H_A(f) = \{(x, z) \in A \times [0, \infty) : f(x) \geq z\}.$$

When we omit the subscript  $A$ , we assume that  $A$  is the support of  $f$ .

Let  $f : \mathbb{R}^n \rightarrow [0, \infty)$  be an  $s$ -concave function for  $s \in (-1/n, \infty)$ , and  $\nu$  a measure on  $\mathbb{R}^n \times \mathbb{R}$  such that

$$d\nu(x_1, \dots, x_{n+1}) = h(x_{n+1}) dx_1 \cdots dx_{n+1}, \quad (3.1)$$

for some continuous function  $h : \mathbb{R} \rightarrow \mathbb{R}$ . With this setup, we can express the integral of  $f$  in terms of the  $\nu$ -measure of the epigraph or hypograph of

a transformation of it. Moreover,

$$\int_A f(x)dx = \begin{cases} \nu(E_A(f^s)), & \text{for } h(x_{n+1}) = -\frac{1}{s}x_{n+1}^{\frac{1}{s}-1}, \text{ if } s < 0. \\ \nu(E_A(-\log f)), & \text{for } h(x_{n+1}) = e^{-x_{n+1}}, \text{ if } s = 0. \\ \nu(H_A(f^s)), & \text{for } h(x_{n+1}) = \frac{1}{s}x_{n+1}^{\frac{1}{s}-1}, \text{ if } s > 0. \end{cases} \quad (3.2)$$

Notice the  $s$ -concavity of  $f$  implies the convexity of  $H_A(f^s)$ ,  $E_A(-\log f)$ , and  $E_A(f^s)$  respectively.

For  $x_1, x_2 \in \mathbb{R}^n$  and  $\lambda \in [0, 1]$ , let  $x_1 +_\lambda x_2 = \lambda x_1 + (1 - \lambda)x_2$ ; similarly for  $K, L \subseteq \mathbb{R}^n$ , we write

$$K +_\lambda L := \lambda K + (1 - \lambda)L. \quad (3.3)$$

For convex functions  $\varphi, \psi : \mathbb{R}^n \rightarrow \mathbb{R}$  and  $\lambda \in [0, 1]$ , we define their infimal convolution by

$$(\varphi \square_\lambda \psi)(v) = \inf_{v=x_1+\lambda x_2} \{\mathcal{M}_\lambda^1(\varphi(x_1), \psi(x_2))\},$$

so that

$$E(\varphi \square_\lambda \psi) = E(\varphi) +_\lambda E(\psi).$$

Let  $f, g : \mathbb{R}^n \rightarrow [0, \infty)$  be  $s$ -concave functions. When  $s < 0$ ,  $f^s$  and  $g^s$  are convex and

$$\begin{aligned} (f \star_{\lambda, s} g)^s(v) &= \left( \sup_{v=x_1+\lambda x_2} \mathcal{M}_\lambda^1(f^s(x_1), g^s(x_2))^{1/s} \right)^s \\ &= \inf_{v=x_1+\lambda x_2} \mathcal{M}_\lambda^1(f^s(x_1), g^s(x_2)) \\ &= (f^s \square_\lambda g^s)(v), \end{aligned}$$

and we have

$$E((f \star_{\lambda, s} g)^s) = E(f^s) +_\lambda E(g^s). \quad (3.4)$$

For  $s = 0$ ,  $-\log f$  and  $-\log g$  are convex and

$$\begin{aligned} -\log(f \star_{\lambda, 0} g)(v) &= -\log \left( \sup_{v=x_1+\lambda x_2} \{\mathcal{M}_\lambda^0(f(x_1), g(x_2))\} \right) \\ &= \inf_{v=x_1+\lambda x_2} \{-\log \mathcal{M}_\lambda^0(f(x_1), g(x_2))\} \\ &= ((-\log f) \square_\lambda (-\log g))(v), \end{aligned}$$

which implies

$$E(-\log(f \star_{\lambda, 0} g)) = E(-\log f) +_\lambda E(-\log g). \quad (3.5)$$

Lastly, for  $s > 0$ ,  $-f^s$ ,  $-g^s$  are convex and

$$\begin{aligned} -(f \star_{\lambda, s} g)^s(v) &= - \sup_{v=x_1+\lambda x_2} \mathcal{M}_\lambda^1(f^s(x_1), g^s(x_2)) \\ &= \inf_{v=x_1+\lambda x_2} \mathcal{M}_\lambda^1(-f^s(x_1), -g^s(x_2)) \\ &= (-f^s \square_\lambda - g^s)(v), \end{aligned}$$

from which it follows that

$$H((f \star_{\lambda, s} g)^s) = H(f^s) +_\lambda H(g^s). \quad (3.6)$$

#### 4. CONVEX HULL AND $\mathcal{M}$ -ADDITION OPERATIONS

Let  $C \subseteq \mathbb{R}^N$  be a compact convex set; for  $x_1, \dots, x_N \in \mathbb{R}^n$ , we view the  $n \times N$  matrix  $[x_1, \dots, x_N]$  as an operator from  $\mathbb{R}^N$  to  $\mathbb{R}^n$ . Then

$$[x_1, \dots, x_N]C = \left\{ \sum_i c_i x_i : c = (c_i) \in C \right\} \quad (4.1)$$

produces a convex set in  $\mathbb{R}^n$ . This viewpoint was used by the first-named author and Paouris in [28] in randomized isoperimetric inequalities for convex sets; for the special case  $C = \text{conv}\{e_1, \dots, e_N\}$ , one has

$$[x_1, \dots, x_N]C = \text{conv}\{x_1, \dots, x_N\}.$$

Moreover, for vectors  $x_1, \dots, x_N, x_{N+1}, \dots, x_{N+M}$ , we have

$$\begin{aligned} [x_1, \dots, x_N]C_N + [x_{N+1}, \dots, x_{N+M}]C_M \\ &= [x_1, \dots, x_N]C_N + [x_{N+1}, \dots, x_{N+M}]C_M \\ &= [x_1, \dots, x_{N+M}](C_N + \widehat{C}_M), \end{aligned} \quad (4.2)$$

where  $C_k = \text{conv}\{e_1, \dots, e_k\}$  for  $k = N, M$  and  $\widehat{C}_M = \text{conv}\{e_{N+1}, \dots, e_{N+M}\}$ . The convex operations on points (4.1) can also be generalized to convex operations on sets by using the notion of  $\mathcal{M}$ -addition. This was introduced by Gardner, Hug, and Weil [17, 18] as a unifying framework for operations in Lutwak, Yang, and Zhang's  $L_p$  and Orlicz Brunn-Minkowski theory (see e.g., [24]). For  $\mathcal{M} \subseteq \mathbb{R}^N$  and subsets  $K_1, \dots, K_N$  in  $\mathbb{R}^n$ , their  $\mathcal{M}$ -combination is defined by

$$\oplus_{\mathcal{M}}(K_1, \dots, K_N) = \left\{ \sum_{i=1}^N m_i x_i : x_i \in K_i, (m_1, \dots, m_N) \in \mathcal{M} \right\}.$$

Thus, with this notation, for  $C = \mathcal{M}$ ,

$$\oplus_C(\{x_1\}, \dots, \{x_N\}) = [x_1, \dots, x_N]C.$$



Additionally, when  $K_1, \dots, K_N$  are convex and  $\mathcal{M}$  is compact, convex and contained in the positive orthant or origin-symmetric, then  $\oplus_{\mathcal{M}}(K_1, \dots, K_N)$  is convex [17, Theorem 6.1].

To connect with the epigraphs and hypographs defined in §3, we use  $\mathcal{M}$ -combinations of rays and line segments in  $\mathbb{R}^{n+1}$ . Let  $C \subseteq \mathbb{R}^N$  be a compact, convex set contained in the positive orthant,  $\rho_1, \dots, \rho_N \in \mathbb{R}$ , and  $x_1, \dots, x_N \in \mathbb{R}^n$ . We define the rays

$$R_{\rho_i}(x_i) = \{(x_i, r) \in \mathbb{R}^n \times \mathbb{R} : \rho_i \leq r\} \quad (4.3a)$$

and the line segments

$$\tilde{R}_{\rho_i}(x_i) = \{(x_i, r) \in \mathbb{R}^n \times [0, \infty) : \rho_i \geq r\}. \quad (4.3b)$$

Accordingly,

$$\oplus_C(R_{\rho_1}(x_1), \dots, R_{\rho_N}(x_N)) \quad \text{and} \quad \oplus_C(\tilde{R}_{\rho_1}(x_1), \dots, \tilde{R}_{\rho_N}(x_N))$$

form the epigraph of a convex function and the hypograph of a concave function, respectively. By choosing  $C = \text{conv}\{e_1, \dots, e_N\}$ , one simply takes the convex hull of the rays or line segments, respectively.

## 5. SHADOW SYSTEMS OF $s$ -CONCAVE FUNCTIONS

In this section, we recall the notion of linear parameter and shadow systems of convex sets and extend these to  $s$ -concave functions. We establish a corresponding convexity property in the functional setting.

Recall the notation for linear parameter systems from the introduction: for an index set  $I$ , bounded sets  $\{x_i\}_{i \in I} \subseteq \mathbb{R}^n$  and  $\{\lambda_i\}_{i \in I} \subseteq \mathbb{R}$  and  $\theta \in \mathbb{S}^{n-1}$ ,  $a \leq t \leq b$ , we set

$$K_t = \text{conv}\{x_i + \lambda_i t \theta : i \in I\}, \quad t \in [a, b].$$

In the notation of §2, we use  $u(K_t, \cdot)$ ,  $\ell(K_t, \cdot)$ ,  $P = P_{\theta^\perp}$ , and we set  $D = PK$ . Define  $L(x) = u(K_t, x) - \ell(K_t, x)$ . Rogers and Shephard proved the fundamental fact that for  $x \in D$ ,  $t \mapsto L(K_t, x)$  is convex. Consequently, the map

$$t \mapsto |K_t| = \int_D L(K_t, x) dx \quad (5.1)$$

is a convex function of  $t$ . For background on shadow systems, see e.g., [11, 27, 39], and [40, §10.4].

We will use certain linear parameter systems for a finite index set  $I = \{1, \dots, N\}$  associated to the  $\oplus_C$  operations of the previous section for epigraphs and hypographs.

**Proposition 5.1.** *Let  $C$  be a compact convex set in  $\mathbb{R}^N$  contained in the positive orthant. Let  $\{x_i\}_{i=1}^N \subseteq \mathbb{R}^n$ ,  $\{z_i\} \subseteq (0, \infty)$  and  $\{\lambda_i\}_{i=1}^N \subseteq \mathbb{R}$ . For  $s \neq 0$ , let  $\rho_i(s) = z_i^s$  and for  $s = 0$  let  $\rho_i(s) = -\log z_i$ . For  $a < b$  and  $t \in [a, b]$ , let*

$$E_t^s = \oplus_C(\{R_{\rho_i(s)}(x_i + \lambda_i t \theta)\}_{i=1}^N) \quad (s \geq 0)$$

and

$$H_t^s = \oplus_C(\{\tilde{R}_{\rho_i(s)}(x_i + \lambda_i t \theta)\}_{i=1}^N) \quad (s < 0).$$

Then for  $\nu$  as in (3.2),  $t \mapsto \nu(E_t^s)$  and  $t \mapsto \nu(H_t^s)$  are convex.

*Proof.* Let  $s \geq 0$ . For  $z \in \mathbb{R}$ , write  $\pi_z = e_{n+1}^\perp + z e_{n+1}$  so that

$$\nu(E_t^s) = \int_{\mathbb{R}} |E_t^s \cap \pi_z| h(z) dz.$$

Thus it suffices to show that for fixed  $z$ ,  $t \mapsto |E_t^s \cap \pi_z|$  is convex. We have

$$E_t^s = \left\{ \sum_{i=1}^N c_i (x_i + r_i e_{n+1}) + \left( \sum_{i=1}^N c_i \lambda_i \right) t \theta : c \in C, \rho_i(s) \leq r_i \right\}.$$

As noted in the previous section, since  $C$  is compact, convex and contained in the positive orthant,  $E_t^s$  is convex. For  $c \in C$ , we write  $x_c = \sum_{i=1}^N c_i x_i$ ,  $r_c = \sum_{i=1}^N c_i r_i$ , and  $\lambda_c = \sum_{i=1}^N c_i \lambda_i$ . For  $x_c + r_c e_{n+1} + \lambda_c t \theta \in E_t^s \cap \pi_z$ , we have  $r_c = z$  and the sets  $\{x_c + z e_{n+1}\}$ ,  $\{\lambda_c\}$  are bounded. Thus  $E_t^s \cap \pi_z$  is a linear parameter system of convex sets indexed by  $C$ . Then  $t \mapsto |E_t^s \cap \pi_z|$  is convex by (5.1). The argument for  $s < 0$  is analogous.  $\square$

Next, we define a linear parameter system of  $s$ -concave functions. Let  $I$  be an index set, and  $\{(x_i, z_i)\}_{i \in I} \subseteq \mathbb{R}^n \times [0, \infty)$  a collection of points under the graph of some integrable  $s$ -concave function. For  $s \geq 0$ , let  $T_{\{w_i\}, s}$  and be the least  $s$ -concave function above  $\{w_i\}$ . For  $s < 0$ , let  $T_{\{w_i\}, s}$  be the greatest  $s$ -concave function beneath  $\{w_i\}$ . More explicitly,  $T_{\{w_i\}, s}$  is supported on  $\text{conv}\{x_i : i \in I\}$  and given by

$$T_{\{w_i\}, s}(x) = \begin{cases} \inf\{z^{1/s} : (x, z) \in \mathcal{P}_{\{w_i\}}\}, & \text{if } s < 0 \\ \sup\{e^z : (x, z) \in \mathcal{P}_{\{w_i\}}\}, & \text{if } s = 0, \\ \sup\{z^{1/s} : (x, z) \in \mathcal{P}_{\{w_i\}}\}, & \text{if } s > 0, \end{cases}$$

where

$$\mathcal{P}_{\{w_i\}} = \begin{cases} \text{conv}\{(x_i, z_i^s)\}_{i \in I}, & \text{if } s \neq 0. \\ \text{conv}\{(x_i, \log z_i)\}_{i \in I}, & \text{if } s = 0. \end{cases}$$

With the above notation, we assume that  $w_i(t) = (x_i + \lambda_i t \theta, z_i)$ ,  $a \leq t \leq b$ . Then setting  $f_{t,s} = T_{\{w_i(t)\},s}$ , we call the family  $\{f_{t,s}\}$  a linear parameter system of  $s$ -concave functions. The convexity property corresponding to (5.1) reads as follows.

**Proposition 5.2.** *Let  $\{f_{t,s}\}$  be a linear parameter system of  $s$ -concave functions. Then*

$$t \mapsto \int_{\mathbb{R}^n} f_t(v) dv$$

*is a convex function.*

*Proof.* As for linear parameter systems of convex sets, we can assume without loss generality that  $I$  is finite, say  $I = \{1, \dots, N\}$ . We take  $C$  to be  $\text{conv}\{e_1, \dots, e_N\}$ . In the notation of the previous proposition, we have

$$E_t^s = \text{conv}\{R_{\rho_i(s)}(x_i + \lambda_i t \theta)\}_{i=1}^N$$

and  $E_t^s = E(-\log f_{t,s})$  for  $s = 0$ , while  $E_t^s = E(f_{t,s}^s)$  for  $s < 0$ . Similarly,

$$H_t^s = \text{conv}\{\tilde{R}_{\rho_i(s)}(x_i + \lambda_i t \theta)\}_{i=1}^N,$$

hence  $H_t^s = H(f_{t,s}^s)$  for  $s > 0$ . By (3.2), we have

$$\int f_{t,s}(v) dv = \begin{cases} \nu(E_t^s), & \text{for } h(x_{n+1}) = -\frac{1}{s} x_{n+1}^{\frac{1}{s}-1}, \text{ if } s < 0. \\ \nu(E_t^s), & \text{for } h(x_{n+1}) = e^{-x_{n+1}}, \text{ if } s = 0. \\ \nu(H_t^s), & \text{for } h(x_{n+1}) = \frac{1}{s} x_{n+1}^{\frac{1}{s}-1}, \text{ if } s > 0, \end{cases}$$

and we can conclude the proof by applying Proposition 5.1.  $\square$

Shephard [41] introduced shadow systems to extend linear parameter systems. Given a convex body  $K \subseteq \mathbb{R}^n$ , a bounded function  $\alpha : K \rightarrow \mathbb{R}$  and  $t \in [a, b]$ , a shadow system in direction  $\theta \in \mathbb{S}^{n-1}$  is a family of convex sets

$$K_t = \text{conv}\{x + \alpha(x)t\theta : x \in K\}. \quad (5.3)$$

Then  $K_t = P_t \overline{K}$ , where  $\overline{K} = \{(x, \alpha(x)) : x \in K\}$  and  $P_t : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^n$  is the projection parallel to  $e_{n+1} - t\theta$  given by  $P_t(x, y) = x + ty\theta$ . Conversely, for any convex body  $\overline{K} \subseteq \mathbb{R}^{n+1}$ ,  $\theta \in \mathbb{S}^{n-1}$ , and  $t \in [a, b]$ , the family  $\{K_t\} = \{P_t \overline{K}\} \subseteq \mathbb{R}^n$  is a shadow system of convex bodies.

The correspondence between linear parameter systems of  $s$ -concave functions and epigraphs/hypographs in the proof of Proposition 5.2 affords a similar extension to shadow systems. We can simply start with shadow systems of epigraphs or hypographs. Let  $\mathcal{E} \subseteq \mathbb{R}^{n+2}$  be the epigraph of a convex

function  $\Phi : \mathbb{R}^{n+1} \rightarrow [0, \infty)$ ,  $\theta \in \mathbb{S}^{n-1}$ ,  $t \in [a, b]$ , and the projection  $P_t$  from  $\mathbb{R}^n \times \mathbb{R} \times \mathbb{R}$  onto  $\mathbb{R}^n \times \mathbb{R}$  parallel to  $e_{n+2} - t\theta$  given by

$$P_t(x, z, y) = (x + ty\theta, z). \quad (5.4)$$

Then the family  $\{E_t\} = \{P_t\mathcal{E}\}$  is a shadow system of epigraphs of convex functions, where

$$E_t = \text{conv}\{(x + yt\theta, z) \in \mathbb{R}^n \times \mathbb{R} : (x, y, z) \in \mathcal{E}\}. \quad (5.5)$$

Consequently, let  $\phi : \mathbb{R}^n \rightarrow [0, \infty)$  be a convex function,  $\alpha : E(\phi) \rightarrow \mathbb{R}$  a function such that  $\alpha|_{E(\phi) \cap \pi_z} : \mathbb{R}^n \rightarrow \mathbb{R}$  is bounded for all  $z \in \mathbb{R}$ . Consider  $\Phi : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  described by its epigraph  $E(\Phi) = \text{conv}\{(x, z, \alpha(x, z)) : \phi(x) \leq z\}$  and set

$$(E(\phi))_t = P_tE(\Phi) = \text{conv}\{(x + \alpha(x, z)t\theta, z) : \phi(x) \leq z\}.$$

Then we define the shadow system of convex functions  $\phi_t : \mathbb{R}^n \rightarrow [0, \infty)$  in direction  $\theta \in \mathbb{S}^n$  by

$$\phi_t(x) = \inf\{z : (x, z) \in (E(\phi))_t\}, \quad (5.6)$$

Analogously, given the hypograph  $\mathcal{H} \subseteq \mathbb{R}^{n+2}$  of a concave function  $\Psi : \mathbb{R}^{n+1} \rightarrow [0, \infty)$  the family  $\{H_t\} = \{P_t\mathcal{H}\}$  is a shadow system of hypographs of concave functions where

$$H_t = \text{conv}\{(x + yt\theta, z) \in \mathbb{R}^n \times \mathbb{R} : (x, z, y) \in \mathcal{H}\}. \quad (5.7)$$

Similarly, for a concave function  $\psi : \mathbb{R}^n \rightarrow [0, \infty)$ ,  $\alpha : H(\psi) \rightarrow \mathbb{R}$  a function such that  $\alpha|_{H(\psi) \cap \pi_z} : \mathbb{R}^n \rightarrow \mathbb{R}$  is bounded for all  $z \in \mathbb{R}$ . Consider the function  $\Psi : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$  with hypograph  $H(\Psi) = \{(x, z, \alpha(x, z)) : 0 \leq z \leq \psi(x)\}$  and set

$$(H(\psi))_t = P_tH(\Psi) = \text{conv}\{(x + \alpha(x, z)t\theta, z) : 0 \leq z \leq \psi(x)\}.$$

Then we define the shadow system of concave functions  $\psi_t : \mathbb{R}^n \rightarrow \mathbb{R}$  by

$$\psi_t(x) = \sup\{z : (x, z) \in (H(\psi))_t\}. \quad (5.8)$$

**Proposition 5.3.** *For shadow systems of epigraphs  $E_t$  and hypographs  $H_t$  and  $\nu$  as in (3.2), we have that  $t \mapsto \nu(E_t)$  and  $t \mapsto \nu(H_t)$  are convex.*

*Proof.* Let  $\nu$  be as in (3.2). Then

$$\nu(E_t) = \int_{\mathbb{R}} |E_t \cap \pi_z| h(z) dz.$$

For each  $z$ , the restriction of the epigraph  $E_t$  to the parallel hyperplane  $\pi_z$  is a shadow systems of convex bodies. Thus the convexity of  $t \mapsto \nu(E_t)$  follows from the convexity of the function (5.1). The proof for  $H_t$  is analogous.  $\square$

## 6. RANDOM EPIGRAPHS AND HYPOGRAPHS

In this section, we take our stochastic model for  $[f]_N$ , as defined in the introduction, and reformulate it in terms of epigraphs and hypographs. Thus for an integrable  $s$ -concave function  $f : \mathbb{R}^n \rightarrow [0, \infty)$ , we sample independent random vectors  $(X_1, Z_1), \dots, (X_N, Z_N)$  according to Lebesgue measure on

$$G_f := \{(x, z) \in \mathbb{R}^n \times [0, \infty) : x \in \text{supp } f, z \leq f(x)\}. \quad (6.1)$$

For  $C_N = \text{conv}\{e_1, \dots, e_N\}$ , we set

$$[E^s]_N = \begin{cases} \oplus_{C_N}(R_{Z_1^s}(X_1), \dots, R_{Z_N^s}(X_N)), & \text{if } s < 0, \\ \oplus_{C_N}(R_{-\log Z_1}(X_1), \dots, R_{-\log Z_N}(X_N)), & \text{if } s = 0. \end{cases} \quad (6.2a)$$

$$[H^s]_N = \oplus_{C_N}(\tilde{R}_{Z_1^s}(X_1), \dots, \tilde{R}_{Z_N^s}(X_N)), \quad \text{if } s > 0. \quad (6.2b)$$

With this notation,

$$[E^s]_N = E([f]_N^s), \quad \text{if } s < 0.$$

$$[E^s]_N = E(-\log [f]_N), \quad \text{if } s = 0.$$

$$[H^s]_N = H([f]_N^s), \quad \text{if } s > 0$$

and, by (3.2),

$$\int [f]_N = \begin{cases} \nu([E^s]_N), & \text{for } h(x_{n+1}) = -\frac{1}{s}x_{n+1}^{\frac{1}{s}-1}, \text{ if } s < 0. \\ \nu([E^s]_N), & \text{for } h(x_{n+1}) = e^{-x_{n+1}}, \text{ if } s = 0. \\ \nu([H^s]_N), & \text{for } h(x_{n+1}) = \frac{1}{s}x_{n+1}^{\frac{1}{s}-1}, \text{ if } s > 0. \end{cases} \quad (6.3a)$$

## 7. MULTIPLE INTEGRAL REARRANGEMENT INEQUALITIES

**7.1. Rearrangements and Steiner convexity.** In this section, we show that the multiple integral rearrangement inequalities of Rogers [35], and Brascamp, Lieb, and Luttinger [7] interface well with our approach. In particular, Christ's version [13] of the latter inequalities is especially applicable; as in [29], the following formulation is convenient for our purpose.

**Theorem 7.1.** *Let  $f_1, \dots, f_N$  be non-negative integrable functions on  $\mathbb{R}^n$  and  $F : (\mathbb{R}^n)^N \rightarrow [0, \infty)$ . Then*

$$\begin{aligned} & \int_{(\mathbb{R}^n)^N} F(x_1, \dots, x_N) \prod_{i=1}^N f_i(x_i) dx_1 \dots dx_N \\ & \geq \int_{(\mathbb{R}^n)^N} F(x_1, \dots, x_N) \prod_{i=1}^N f_i^*(x_i) dx_1 \dots dx_N, \end{aligned}$$

*whenever  $F$  satisfies the following condition: for each  $\theta \in \mathbb{S}^{n-1}$  and all  $Y := \{y_1, \dots, y_N\} \subseteq \theta^\perp$ , the function  $F_Y : \mathbb{R}^N \rightarrow [0, \infty)$  defined by*

$$F_{Y,\theta}(t_1, \dots, t_N) := F(y_1 + t_1\theta, \dots, y_N + t_N\theta)$$

*is even and quasi-convex.*

The condition on  $F$ , namely *Steiner convexity*, allows the theorem to be proved via iterated Steiner symmetrization; notice this terminology differs from the one in [13]. Of special interest, this condition interfaces well with shadow systems, e.g., [10, 36]; see [29] for further background and references.

**Proposition 7.2.** *Let  $\rho_1, \dots, \rho_N \in \mathbb{R}$  and  $C$  a compact convex set contained in the positive orthant. Then the function  $F : (\mathbb{R}^n)^N \rightarrow [0, \infty)$  defined by*

$$F(x_1, \dots, x_N) = \nu(\oplus_C(R_{\rho_1}(x_1), \dots, R_{\rho_N}(x_N))) \quad (7.1)$$

*is Steiner convex.*

*Proof.* Let  $\theta \in \mathbb{S}^{n-1}$  and  $y_1, \dots, y_N \in \theta^\perp$ . Let  $(s_1, \dots, s_N), (t_1, \dots, t_N) \in \mathbb{R}^N$  and  $\tau \in (0, 1)$ . For  $i = 1, \dots, N$ , write

$$y_i + (\tau s_i + (1 - \tau)t_i)\theta = (y_i + t_i\theta) + \tau(t_i - s_i)\theta$$

and apply Proposition 5.1 with  $x_i = y_i + t_i\theta$ ,  $\lambda_i = t_i - s_i$  and  $t = \tau$  to obtain the convexity in  $\tau$ . Lastly, the sets

$$\oplus_C(R_{\rho_1}(y_1 + t_1\theta), \dots, R_{\rho_N}(y_N + t_N\theta))$$

and

$$\oplus_C(R_{\rho_1}(y_1 - t_1\theta), \dots, R_{\rho_N}(y_N - t_N\theta))$$

are reflections of one another and so the evenness condition required for Steiner convexity holds.  $\square$

Next, we state the analogous proposition involving the line segments (4.3b); the proof follows the same line.

**Proposition 7.3.** *Let  $\rho_1, \dots, \rho_N \in \mathbb{R}$  and  $C$  a compact convex set contained in the positive orthant. Then the function  $F : (\mathbb{R}^n)^N \rightarrow [0, \infty)$  defined by*

$$F(x_1, \dots, x_N) = \nu \left( \oplus_C (\tilde{R}_{\rho_1}(x_1), \dots, \tilde{R}_{\rho_N}(x_N)) \right) \quad (7.2)$$

*is Steiner convex.*

## 8. MAIN PROOF

*Proof of Theorem 1.1.* Let  $w_i = (x_i, z_i) \in \mathbb{R}^n \times \mathbb{R}$  for  $i = 1, \dots, M + N$ . For  $s \geq 0$ , let  $T_{\{w_i\},s}^{(N)}$  and  $T_{\{w_i\},s}^{(M)}$  be the least  $s$ -concave functions above the collections  $\{w_i\}_{i \leq N}$  and  $\{w_i\}_{N+1 \leq i \leq M+N}$ , respectively; similarly, when  $s < 0$ , let  $T_{\{w_i\}}^{(N)}$  and  $T_{\{w_i\}}^{(M)}$  be the greatest  $s$ -concave functions beneath the respective collections  $\{w_i\}_{i \leq N}$  and  $\{w_i\}_{N+1 \leq i \leq M+N}$ . With this notation, we set

$$F(w_1, \dots, w_{N+M}) = \int_{\mathbb{R}^n} T_{\{w_i\}}^{(N)} \star_{\lambda} T_{\{w_i\}}^{(M)}(v) dv.$$

Let  $f, g : \mathbb{R}^n \rightarrow [0, \infty)$  be integrable  $s$ -concave functions for  $s \in (-1/n, \infty)$ . Sample independent random vectors  $W_i = (X_i, Z_i)$ ,  $i = 1, \dots, N + M$  uniformly according to the Lebesgue measure on  $G_f$  for  $i = 1, \dots, N$  and  $G_g$  for  $i = N + 1, \dots, N + M$ . Then the random functions  $[f]_N, [g]_M$  satisfy

$$\begin{aligned} & \mathbb{P} \left( \int_{\mathbb{R}^n} [f]_N \star_{\lambda,s} [g]_M(v) dv > \alpha \right) \\ &= \frac{1}{\prod_{i=1}^{M+N} \|k_i\|_1} \int_{N+M} \mathbb{1}_{\{F > \alpha\}}(\bar{w}) \prod_{i=1}^{N+M} \mathbb{1}_{[0, k_i(x_i)]}(z_i) d\bar{w}, \end{aligned} \quad (8.1)$$

where  $\int_{N+M}$  is the integral on  $(\mathbb{R}^n \times [0, \infty))^{N+M}$ ,  $k_i = f$  for  $i = 1, \dots, N$ ,  $k_i = g$  for  $i = N + 1, \dots, N + M$ , and

$$\bar{w} = (w_1, \dots, w_{N+M}), \quad d\bar{w} = dw_1 \dots dw_{N+M}. \quad (8.2)$$

Also we write  $C_N = \text{conv}\{e_1, \dots, e_N\}$  and  $\hat{C}_M = \text{conv}\{e_{N+1}, \dots, e_{N+M}\}$ , and consider  $\nu$  as in (3.2) for each case.

**Case  $s > 0$ :** By (6.2b) and (3.6) it follows

$$\begin{aligned} H([f]_N \star_{\lambda,s} [g]_M)^s &= \oplus_{C_N} (\{\tilde{R}_{Z_i^s}(X_i)\}_{i=1}^N) +_{\lambda} \oplus_{\hat{C}_M} (\{\tilde{R}_{Z_i^s}(X_i)\}_{i=N+1}^{N+M}) \\ &= \oplus_{C_N +_{\lambda} \hat{C}_M} (\{\tilde{R}_{Z_i^s}(X_i)\}_{i=1}^{N+M}), \end{aligned}$$

hence

$$\int_{\mathbb{R}^n} [f]_N \star_{\lambda,s} [g]_M(v) dv = \nu \left( \oplus_{C_N +_{\lambda} \hat{C}_M} (\{\tilde{R}_{Z_i^s}(X_i)\}_{i=1}^{N+M}) \right). \quad (8.3)$$

By (8.1), Fubini, Proposition 7.2, and Theorem 7.1, we have

$$\begin{aligned}
& \mathbb{P} \left( \int_{\mathbb{R}^n} [f]_N \star_{\lambda,s} [g]_M(v) dv > \alpha \right) \\
&= \frac{1}{\prod_{i=1}^{M+N} \|k_i\|_1^{N+M}} \int_{(\mathbb{R}^n \times [0,\infty))^{N+M}} \mathbb{1}_{\{F>\alpha\}}(\bar{w}) \prod_{i=1}^{N+M} \mathbb{1}_{[0,k_i(x_i)]}(z_i) d\bar{w} \\
&= \frac{1}{\prod_{i=1}^{M+N} \|k_i\|_1^{N+M}} \int_{[0,\infty)^{N+M}} \left( \int_{(\mathbb{R}^n)^{N+M}} \mathbb{1}_{\{F>\alpha\}}(\bar{w}) \prod_{i=1}^{N+M} \mathbb{1}_{[0,k_i(x_i)]}(z_i) d\bar{x} \right) d\bar{z} \\
&\geq \frac{1}{\prod_{i=1}^{M+N} \|k_i^*\|_1^{N+M}} \int_{[0,\infty)^{N+M}} \left( \int_{(\mathbb{R}^n)^{N+M}} \mathbb{1}_{\{F>\alpha\}}(\bar{w}) \prod_{i=1}^{N+M} \mathbb{1}_{[0,k_i^*(x_i)]}(z_i) d\bar{x} \right) d\bar{z} \\
&= \frac{1}{\prod_{i=1}^{M+N} \|k_i^*\|_1^{N+M}} \int_{(\mathbb{R}^n \times [0,\infty))^N} \mathbb{1}_{\{F>\alpha\}}(\bar{w}) \prod_{i=1}^{N+M} \mathbb{1}_{[0,k_i^*(x_i)]}(z_i) d\bar{w} \\
&= \mathbb{P} \left( \int_{\mathbb{R}^n} [f^*]_N \star_{\lambda,s} [g^*]_M(v) dv > \alpha \right).
\end{aligned}$$

**Case  $s = 0$ :** Using (6.2a), (3.5), we have

$$E(-\log([f]_N \star_{\lambda,0} [g]_M)) = \oplus_{C_N + \lambda \hat{C}_M} (\{R_{-\log Z_i}(X_i)\}_{i=1}^{N+M}),$$

so

$$\int_{\mathbb{R}^n} [f]_N \star_{\lambda,0} [g]_M(v) dv = \nu \left( \oplus_{C_N + \lambda \hat{C}_M} (\{R_{-\log Z_i}(X_i)\}_{i=1}^{N+M}) \right). \quad (8.4)$$

It follows as before

$$\mathbb{P} \left( \int_{\mathbb{R}^n} [f]_N \star_{\lambda,0} [g]_M(v) dv > \alpha \right) \geq \mathbb{P} \left( \int_{\mathbb{R}^n} [f^*]_N \star_{\lambda,0} [g^*]_M(v) dv > \alpha \right).$$

**Case  $s < 0$ :** It follows by (6.2a) and (3.5) that

$$E([f]_N \star_{\lambda,s} [g]_M)^s = \oplus_{C_N + \lambda \hat{C}_M} (\{R_{Z_i^s}(X_i)\}_{i=1}^{N+M}),$$

hence

$$\int_{\mathbb{R}^n} [f]_N \star_{\lambda,s} [g]_M(v) dv = \nu \left( \oplus_{C_N + \lambda \hat{C}_M} (\{R_{Z_i^s}(X_i)\}_{i=1}^{N+M}) \right). \quad (8.5)$$

It follows as before

$$\mathbb{P} \left( \int_{\mathbb{R}^n} [f]_N \star_{\lambda,s} [g]_M(v) dv > \alpha \right) \geq \mathbb{P} \left( \int_{\mathbb{R}^n} [f^*]_N \star_{\lambda,s} [g^*]_M(v) dv > \alpha \right).$$

□



**Remark 8.1.** We have applied Propositions 7.2 and 7.3 only in the special case when  $C = C_N +_\lambda \widehat{C}_M$ . Since these propositions apply to more general convex sets  $C$ , they can be used to treat alternate means and different stochastic functions in Theorem 1.1. This direction and its geometric implications are outside of our present scope but will appear in a forthcoming work of the authors.

## REFERENCES

- [1] S. Artstein-Avidan, D. I. Florentin, and A. Segal. Functional Brunn-Minkowski inequalities induced by polarity. *Adv. Math.*, 364:107006, 2020.
- [2] S. Artstein-Avidan, A. Giannopoulos, and V. D. Milman. *Asymptotic geometric analysis. Part I*, volume 202 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2015.
- [3] S. Artstein-Avidan, B. Klartag, and V. Milman. The Santaló point of a function, and a functional form of the Santaló inequality. *Mathematika*, 51(1-2):33–48 (2005), 2004.
- [4] C. Borell. Convex set functions in  $d$ -space. *Periodica Mathematica Hungarica*, 6(2):111–136, 1975.
- [5] H. J. Brascamp and E. H. Lieb. Best constants in Young's inequality, its converse, and its generalization to more than three functions. *Advances in Math.*, 20(2):151–173, 1976.
- [6] H. J. Brascamp and E. H. Lieb. On extensions of the Brunn-Minkowski and Prékopa-Leindler theorems, including inequalities for log concave functions, and with an application to the diffusion equation. *J. Functional Analysis*, 22(4):366–389, 1976.
- [7] H. J. Brascamp, E. H. Lieb, and J.M. Luttinger. A general rearrangement inequality for multiple integrals. *Journal of functional analysis*, 17(2):227–237, 1974.
- [8] A. Burchard. A short course on rearrangement inequalities. Available at <http://www.math.utoronto.ca/almut/rearrange/pdf>.
- [9] H. Busemann. Volume in terms of concurrent cross-sections. *Pacific J. Math.*, 3:1–12, 1953.
- [10] S. Campi, A. Colesanti, and P. Gronchi. A note on Sylvester's problem for random polytopes in a convex body. *Rend. Istit. Mat. Univ. Trieste*, 31(1-2):79–94, 1999.
- [11] S. Campi and P. Gronchi. On volume product inequalities for convex sets. *Proc. Amer. Math. Soc.*, 134(8):2393–2402, 2006.
- [12] S. Campi and P. Gronchi. Volume inequalities for sets associated with convex bodies. In *Integral geometry and convexity*, pages 1–15. World Scientific, 2006.
- [13] M. Christ. Estimates for the  $k$ -plane transform. *Indiana Univ. Math. J.*, 33(6):891–910, 1984.
- [14] D. Cordero-Erausquin, M. Fradelizi, G. Paouris, and P. Pivovarov. Volume of the polar of random sets and shadow systems. *Math. Ann.*, 362(3-4):1305–1325, 2015.
- [15] M. Fradelizi, M. Meyer, and A. Zvavitch. An application of shadow systems to Mahler's conjecture. *Discrete & Computational Geometry*, 48(3):721–734, 2012.
- [16] R. J. Gardner. The Brunn-Minkowski inequality. *Bull. Amer. Math. Soc. (N.S.)*, 39(3):355–405, 2002.

- [17] R. J. Gardner, D. Hug, and W. Weil. Operations between sets in geometry. *J. Eur. Math. Soc. (JEMS)*, 15(6):2297–2352, 2013.
- [18] R. J. Gardner, D. Hug, and W. Weil. The Orlicz-Brunn-Minkowski theory: a general framework, additions, and inequalities. *J. of Differential Geom.*, 97(3):427–476, 2014.
- [19] H. Groemer. On the mean value of the volume of a random polytope in a convex set. *Arch. Math. (Basel)*, 25:86–90, 1974.
- [20] R. Henstock and A.M. Macbeath. On the measure of sum-sets.(i) the theorems of Brunn, Minkowski, and Lusternik. *Proceedings of the London Mathematical Society*, 3(1):182–194, 1953.
- [21] B. Klartag. Marginals of geometric inequalities. In *Geometric aspects of functional analysis*, volume 1910 of *Lecture Notes in Math.*, pages 133–166. Springer, Berlin, 2007.
- [22] L. Leindler. On a certain converse of Hölder’s inequality. II. *Acta Sci. Math. (Szeged)*, 33(3-4):217–223, 1972.
- [23] E. H. Lieb and M. Loss. *Analysis*, volume 14 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, second edition, 2001.
- [24] E. Lutwak, Yang D., and Zhang G. Orlicz centroid bodies. *Journal of Differential Geometry*, 84(2):365–387, 2010.
- [25] J. Melbourne. Rearrangement and Prékopa-Leindler type inequalities. Available at <https://arxiv.org/abs/1806.08837>.
- [26] M. Meyer and A. Pajor. On the Blaschke-Santaló inequality. *Arch. Math. (Basel)*, 55(1):82–93, 1990.
- [27] M. Meyer and S. Reisner. Shadow systems and volumes of polar convex bodies. *Mathematika*, 53(1):129–148 (2007), 2006.
- [28] G. Paouris and P. Pivovarov. A probabilistic take on isoperimetric-type inequalities. *Adv. Math.*, 230(3):1402–1422, 2012.
- [29] G. Paouris and P. Pivovarov. Randomized isoperimetric inequalities. In *Convexity and concentration*, volume 161 of *IMA Vol. Math. Appl.*, pages 391–425. Springer, New York, 2017.
- [30] P. Pivovarov and J. Rebollo Bueno. A stochastic Prékopa-Leindler inequality for log-concave functions. *Communications in Contemporary Mathematics*, 2020. Accepted for publication.
- [31] A. Prékopa. Logarithmic concave measures with application to stochastic programming. *Acta Sci. Math. (Szeged)*, 32:301–316, 1971.
- [32] A. Prékopa. On logarithmic concave measures and functions. *Acta Sci. Math. (Szeged)*, 34:335–343, 1973.
- [33] J. Rebollo Bueno. Stochastic reverse isoperimetric inequalities in the plane. Preprint. Available at [bit.ly/StochPlan](https://bit.ly/StochPlan).
- [34] Y. Rinott. On convexity of measures. *Ann. Probability*, 4(6):1020–1026, 1976.
- [35] C. A. Rogers. A single integral inequality. *Journal of the London Mathematical Society*, 1(1):102–108, 1957.
- [36] C. A. Rogers and G. C. Shephard. Some extremal problems for convex bodies. *Mathematika*, 5:93–102, 1958.
- [37] M. Roysdon and S. Xing. On  $L_p$ -Brunn-Minkowski type and  $L_p$ -isoperimetric type inequalities for general measures. <https://arxiv.org/abs/2004.09737>.
- [38] L. A. Santaló. Un invariante afin para los cuerpos convexos del espacio de  $n$  dimensiones. *Portugaliae Mathematica*, 8(4):155–161, 1949.

- [39] C. Saroglou. Shadow systems: remarks and extensions. *Archiv der Mathematik*, 100(4):389–399, 2013.
- [40] R. Schneider. *Convex bodies: the Brunn-Minkowski theory*, volume 151 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, expanded edition, 2014.
- [41] G. C. Shephard. Shadow systems of convex sets. *Israel Journal of Mathematics*, 2(4):229–236, 1964.
- [42] B. Simon. *Convexity*, volume 187 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2011. An analytic viewpoint.
- [43] M. Weberndorfer. Shadow systems of asymmetric  $L_p$  zonotopes. *Advances in Mathematics*, 240:613–635, 2013.

UNIVERSITY OF MISSOURI, DEPARTMENT OF MATHEMATICS, COLUMBIA MO 65201  
Email address: pivovarovp@missouri.edu & jrc65@mail.missouri.edu