

# ON HOMOGENEOUS QUASIPOSITIVE LINKS

TETSUYA ITO

ABSTRACT. We discuss when homogeneous quasipositive links are positive. In particular, we show that a homogeneous diagram of a quasipositive link whose number of Seifert circles is equal to the braid index is a positive diagram.

An  $n$ -braid is *quasipositive* if it is a product of conjugates of the standard generators  $\sigma_1, \sigma_2, \dots, \sigma_{n-1}$ . A link  $K$  in  $S^3$  is

- *quasipositive* if  $K$  can be represented by the closure of a quasipositive braid.
- *positive* if  $K$  can be represented by a positive diagram, a diagram without negative crossing.
- *homogeneous* if  $K$  can be represented by a homogenous diagram (see [Cr] for definition)

A homogenous diagram are common generalization of positive diagram and alternating diagram, sharing various nice properties of such diagrams.

In [Ba1, Question (4)] Baader asked whether an alternating quasipositive link is positive. He also posed a more general question, whether a homogenous quasipositive link is positive [Ba2, Problems (i)].

Recently, in [Or] Orevkov gave an affirmative answer to the first Baader's question for a class of alternating links which he call *DHL link* (Diao-Hertyi-Liu link); An alternating link is DHL link if it has an alternating diagram whose number of Seifert circles is equal to the braid index. In [DHL] Diao-Hertyi-Liu gave a simple and clear characterization of such an alternating diagram; for an alternating diagram  $D$  of a link  $K$ , the number of Seifert circles of  $D$  is equal to the braid index of  $K$  if and only if there is no pair of Seifert circles connected by a single crossing [DHL].

In this note we discuss Baader's second question, the positivity of homogenous quasipositive links.

For a link  $K$  let  $SL(K)$  be the maximal self-linking number and  $b(K)$  be the braid index. For a link diagram  $D$  of  $K$ , we denote by  $O(D)$  the number of Seifert circles of  $D$ , and denote by  $c_+(D), c_-(D)$  the number of positive and negative crossings of  $D$ . We define the *self-linking number* of diagram  $D$  by  $sl(D) = -O(D) + c_+(D) - c_-(D)$  (this is the self-linking number of closed braid obtained from  $D$  by Vogel-Yamada method [Vo, Ya]).

**Theorem 1.** *Let  $D$  be a homogenous diagram of a quasipositive link  $K$ . If  $SL(K) = sl(D)$ ,  $D$  is a positive diagram.*

This generalizes Orevkov's theorem for homogenous case;

**Corollary 1.** *Let  $D$  be a homogenous diagram of a quasipositive link  $K$ . If  $O(D) = b(K)$ <sup>1</sup>, then  $D$  is a positive diagram.*

<sup>1</sup>In [HIK] we call a homogeneous link having such a homogeneous diagram *strongly homogeneous*.

*Proof of Corollary 1.* Following Vogel-Yamada method [Vo, Ya], one can convert  $D$  into a closed  $O(D)$ -braid diagram  $\beta_D$  without changing the writhe  $c_+(D) - c_-(D)$ . Since  $\beta_D$  is a closed braid with minimum braid index, by generalized Jones' conjecture proven in [DP, LM], the closed braid  $\beta_D$  viewed as a transverse link attains the maximal self-linking number  $SL(K)$  hence  $SL(K) = sl(D)$ .  $\square$

*Proof of Theorem 1.* In the following we assume that  $D$  is non-split, since to prove the theorem it is sufficient to treat the case  $D$  is non-split.

Abe [Ab] showed that for a homogeneous non-split diagram  $D$  of  $K$ , the Rasmussen invariant  $s(K)$  of  $K$  is given by<sup>2</sup>

$$s(K) = -O(D) + c_+(D) - c_-(D) + 2O_+(D) - 1 = sl(D) + 2O_+(D) - 1$$

where  $O_+(D)$  is the number of connected components of  $D$  obtained by smoothing all the negative crossings.

Since  $K$  is quasipositive, slice Bennequin inequality gives an equality hence

$$SL(K) = s(K) - 1 = -\chi_4(K).$$

Here  $\chi_4(K)$  the maximum euler characteristic of smooth oriented surface in  $B^4$  whose boundary is  $K$  (see [Sh]). Therefore by our hypothesis  $sl(K) = sl(D)$ , we get  $O_+(D) = 1$ . For a homogeneous non-split diagram  $D$ ,  $O_+(D) = 1$  happens only if  $c_-(D) = 0$ . Therefore  $D$  is positive.  $\square$

Actually our argument shows that a homogeneous diagram of a quasipositive link should be close to a positive diagram in the following sense.

**Theorem 2.** *Let  $D$  be a non-split homogeneous diagram of a quasipositive link  $K$ . Then  $O(D) - b(K) \geq O_+(D) - 1$  holds.*

*Proof.* By generalized Jones' conjecture proven in [DP, LM]

$$sl(D) \geq SL(K) - 2(O(D) - b(K)).$$

Hence

$$sl(D) \geq SL(K) - 2(O(D) - b(K)) = sl(D) + 2(O_+(D) - 1) - 2(O(D) - b(K))$$

$\square$

Unfortunately, in general it is not true that a homogeneous (alternating) link  $K$  has a homogeneous (alternating) diagram  $D$  with  $SL(K) = sl(D)$  (see Remark 3 below).

To overcome the defect, we give a different class of homogeneous diagrams having an affirmative answer to Baader's question. In [MP] Murasugi-Prytycki gave an refinement of famous Motron-Franks-Williams inequalities.

$$(1) \quad \min \deg_v P_K(v, z) \geq -O(D) + c_+(D) - c_-(D) + 1 + 2 \text{ind}_-(D)$$

$$(2) \quad \max \deg_v P_K(v, z) \leq O(D) + c_+(D) - c_-(D) - 1 - 2 \text{ind}_+(D)$$

$$(3) \quad b(K) \leq O(D) - \text{ind}(D)$$

---

<sup>2</sup>We remark that the Rasmussen's invariant  $s(K)$  is originally defined for knots [Ra]. For link case we use Beliakova-Wherli's generalization of Rasmussen invariant for links [BW]. Although [Ab] proves the formula only for knot case, his proof can be applied to link cases, as he noted and mentioned in [AT] (if the diagram is non-split).

We call the last inequality (3) of the braid index the *MP inequality* (Murasugi-Prytycki inequality).

Here  $P_K(v, z)$  is the HOMFLY polynomial of  $K$ , and  $\text{ind}_\pm(D), \text{ind}(D)$  are the index of the Seifert graph of  $D$  (see [MP] for definitions). In particular, they proved for a given knot diagram  $D$  of  $K$  one can find a diagram  $D'$  of the same knot  $K$  so that

$$\begin{aligned} sl(D') &= sl(D) + 2\text{ind}_-(D), & O(D') &= O(D) - \text{ind}_-(D) \\ sl(D') &= sl(D), & O(D') &= O(D) - \text{ind}_+(D). \end{aligned}$$

For general link diagram  $D$ ,

$$\text{ind}_+(D) + \text{ind}_-(D) \geq \text{ind}(D).$$

When  $D$  is homogeneous,  $\text{ind}(D) = \text{ind}_-(D) + \text{ind}_+(D)$ , and there is a diagram  $D'$  such that

$$sl(D') = sl(D) + 2\text{ind}_-(D), \quad O(D') = O(D) - \text{ind}(D)$$

**Remark 3.** If  $D$  is a reduced alternating diagram of  $\text{ind}_-(D) > 1$ , by (1)  $sl(D) = SL(K)$  cannot happen. Since for a reduced alternating link  $K$ , the writhe of a reduced alternating diagram is invariant of an alternating link, such a link does not have an alternating diagram  $D$  with  $sl(D) = SL(K)$ .

Using these results we give another class of links having an affirmative answer to Baader's question.

**Theorem 4.** *Let  $D$  be an irreducible homogeneous diagram of a quasipositive link. If  $sl(D) + 2\text{ind}_-(D) = SL(K)$ , then  $D$  is positive.*

*Proof.* As in Theorem 1, it is sufficient to treat the case  $D$  is non-split. Then by quasipositivity  $SL(K) = sl(D) + 2(O_+(D) - 1)$ , we have  $\text{ind}_-(D) = O_+(D) - 1$ . However, since  $D$  is irreducible this is possible only if  $\text{ind}_-(D) = 0$ . This follows from the following definitions, results, and observations:

- the Seifert graph  $\Gamma(D)$  of homogeneous diagram  $D$  is written as the  $*$ -product  $\Gamma(D_1) * \Gamma(D_2) * \cdots * \Gamma(D_n)$ , where each diagram  $D_i$  is a positive or negative special diagram. Moreover, if  $D$  is reduced, no  $D_i$  contains a vertex of valence one (definition of homogeneous diagram [Cr]).
- $\text{ind}_\pm(D) = \sum_{i=1}^n \text{ind}_\pm(D_i)$ , since  $\Gamma(D_i)$  is bipartite [MP, Theorem 2.4].
- If  $D$  is connected,  $O_+(D) - 1 = \sum_{i=1}^n (O_+(D_i) - 1)$  (follows from the definition of  $*$ -product).
- For a negative special diagram  $D_i$ ,  $\text{ind}_-(D_i) > O_+(D_i) - 1$  whenever its Seifert graph  $\Gamma(D_i)$  has no vertices of valence one (follows from the definition of index).

Thus  $O_+(D) = 1$  and  $c_-(D) = 0$ , as desired.  $\square$

In particular, we have the following.

**Corollary 2.** *Let  $D$  be an irreducible homogeneous diagram of a quasipositive link. If the MP inequality of the braid index (3) is equality, then  $D$  is positive.*

In [MP] it is conjectured that for alternating diagram  $D$ , the MP inequality of the braid index is an equality. Our discussion so far shows that an affirmative answer to Murasugi-Prytycki's conjecture implies an affirmative answer to Baader's first question.

Finally, although it is irrelevant to our study of homogeneous quasipositive links, we give an improvement of the MP inequality (3) which is interesting in its own right.

**Theorem 5** (Refined MP inequality). *For any link diagram  $D$  of a link  $K$ ,*

$$b(K) \leq O(D) - (\text{ind}_+(D) + \text{ind}_-(D))$$

*Proof.* Let  $\overline{D}$  and  $\overline{K}$  be the mirror image of  $D$  and  $K$ , respectively. Since  $\text{ind}_-(\overline{D}) = \text{ind}_+(D)$  by definition,

$$sl(D) + 2\text{ind}_-(D) \leq SL(K), \quad sl(\overline{D}) + 2\text{ind}_+(D) \leq SL(\overline{K})$$

hence

$$(sl(D) + sl(\overline{D})) + 2(\text{ind}_+(D) + \text{ind}_-(D)) \leq SL(K) + SL(\overline{K})$$

On the other hand, (generalized) Jones' conjecture implies  $SL(K) + SL(\overline{K}) = -2b(K)$ . Since  $sl(D) + sl(\overline{D}) = -2O(D)$  by definition, we conclude

$$-2O(D) + 2(\text{ind}_+(D) + \text{ind}_-(D)) \leq -2b(K).$$

□

Unlike the proof of original MP inequality where they explicitly construct a link diagram  $D'$  with  $O(D') = O(D) - \text{ind}_-(D)$  from  $D$ , our proof of Theorem 5 does not give a link diagram  $D'$  with  $O(D') = O(D) - (\text{ind}_+(D) + \text{ind}_-(D))$ . Finding how to get such a link diagram  $D'$  from  $D$  would be an interesting question.

**Corollary 3.** *For a diagram  $D$  of a link  $K$ , If equalities hold in (1) and (2), then  $b(K) = O(D) - \text{ind}_-(D) - \text{ind}_+(D) = \frac{1}{2}\text{span}_v P_K(v, z) + 1$  (in particular, the Morton-Franks-Williams inequality is equality).*

*Proof.* If equalities hold in (1) and (2) then  $\frac{1}{2}\text{span}_v P_K(v, z) + 1 = O(D) - (\text{ind}_+(D) + \text{ind}_-(D))$ . By the Morton-Franks-Williams inequality and refined MP inequality

$$\frac{1}{2}\text{span}_v P_K(v, z) + 1 \leq b(K) \leq O(D) - (\text{ind}_+(D) + \text{ind}_-(D))$$

□

#### ACKNOWLEDGEMENT

The author has been partially supported by JSPS KAKENHI Grant Number 19K03490, 16H02145. He is grateful to Stepan Yu Orevkov for inspiring the study of positivity of homogeneous quasipositive links.

#### REFERENCES

- [Ab] T. Abe, *The Rasmussen invariant of a homogeneous knot*, Proc. Amer. Math. Soc. 139 (2011), no. 7, 2647–2656.
- [AT] T. Abe and K. Tagami, *Characterization of positive links and the  $s$ -invariant for links*. Canad. J. Math. 69 (2017), no. 6, 1201–1218.
- [Ba1] S. Baader, *Slice and Gordian numbers of track knots*. Osaka J. Math. 42 (2005), no. 1, 257–271.
- [Ba2] S. Baader, *Quasipositivity and homogeneity*, Math. Proc. Cambridge Philos. Soc. 139 (2005), no. 2, 287–290.
- [BW] A. Beliakova and S. Wherli *Categorification of the colored Jones polynomial and Rasmussen invariant of links*, Canad. J. Math. 60 (2008), no. 6, 1240–1266.
- [Cr] P. Cromwell, *Homogeneous links*. J. London Math. Soc. (2) 39 (1989), no. 3, 535–552.

- [DHL] Y. Diao, G. Heteyi, and P. Liu, *The braid index of reduced alternating links*. Math. Proc. Cambridge Philos. Soc. 168 (2020), no. 3, 415–434.
- [DP] I. Dynnikov, and M. Prasolov, *Bypasses for rectangular diagrams. A proof of the Jones conjecture and related questions* Trans. Moscow Math. Soc. 2013, 97–144
- [HIK] J. Hamer, T. Ito, and K. Kawamuro, *Positivities of knots and links and the defect of Bennequin inequality*, Experiment. Math. to appear.
- [LM] D. LaFountain, and W. Menasco, *Embedded annuli and Jones’ conjecture*. Algebr. Geom. Topol. 14 (2014), no. 6, 3589–3601.
- [MP] K. Murasugi, and J. Przytycki, *An index of a graph with applications to knot theory*, Mem. Amer. Math. Soc. 106 (1993), no. 508, x+101 pp.
- [Or] S. Yu. Orevkov, *On alternating quasipositive links*, arXiv:2007.00093v1
- [Ra] J. Rasmussen, *Khovanov homology and the slice genus*. Invent. Math. 182 (2010), no. 2, 419–447.
- [Sh] A. Shumakovitch, *Rasmussen invariant, slice-Bennequin inequality, and sliceness of knots*. J. Knot Theory Ramifications 16 (2007), no. 10, 1403–1412.
- [Vo] P. Vogel, *Representation of links by braids: a new algorithm*. Comment. Math. Helv. 65 (1990), no. 1, 104–113.
- [Ya] S. Yamada, *The minimal number of Seifert circles equals the braid index of a link*. Invent. Math. 89 (1987), no. 2, 347–356.

DEPARTMENT OF MATHEMATICS, KYOTO UNIVERSITY, KYOTO 606-8502, JAPAN  
*E-mail address*: tetitoh@math.kyoto-u.ac.jp