

Tightness of the Asymptotic Generalized Poor-Verdú Error Bound for the Memoryless Symmetric Channel

Ling-Hua Chang^{*}, Po-Ning Chen[†], Fady Alajaji[‡] and Yunghsiang S. Han[§]

Abstract

The generalized Poor-Verdú error lower bound for multihypothesis testing is revisited. Its asymptotic expression is established in closed-form as its tilting parameter grows to infinity. The asymptotic generalized lower bound is then studied in the classical channel coding context where it is proved that for any sequence of block codes sent over the memoryless binary symmetric channel (BSC), the minimum probability of decoding error has a relative deviation from the generalized bound that grows at most linearly in blocklength. A direct consequence of this result is that the asymptotic generalized bound achieves the error exponent (or reliability function) of the BSC at arbitrary coding rates. Finally, these tightness results are extended for the class of memoryless non-binary symmetric channels.

Index Terms

Binary symmetric channel, non-binary symmetric channel, error probability bounds, error exponent, channel reliability function, hypothesis testing, code rates.

I. INTRODUCTION

A well-known lower bound on the minimum probability of error P_e of multihypothesis testing is the so-called Poor-Verdú bound [2]. The bound was generalized in [3] by tilting, via a parameter $\theta \geq 1$, the posterior hypothesis distribution. The generalized bound was noted to progressively improve with θ except for some particular channels such as the memoryless binary erasure channel (BEC). However, its asymptotic formula as θ tends to infinity was not determined.

In this paper, we revisit this generalized bound and determine its asymptotic expression in closed-form. We then investigate the resulting asymptotic generalized bound in the classical context of the error probability of block codes \mathcal{C}_n with blocklength n and size $|\mathcal{C}_n| = M_n$ sent over the memoryless binary symmetric channel (BSC) with

^{*}Department of Electrical Engineering, Yuan Ze University, Taiwan, R.O.C. (iamjaung@gmail.com).

[†]Institute of Communications Engineering and Department of Electrical and Computer Engineering, National Chiao Tung University, Taiwan, R.O.C. (poning@faculty.nctu.edu.tw).

[‡]Department of Mathematics and Statistics, Queen's University, Kingston, ON, Canada (fa@queensu.ca).

[§]School of Electrical Engineering and Intelligentization, Dongguan University of Technology, China (yunghsiangh@gmail.com).

The work of Ling-Hua Chang is supported by the Ministry of Science and Technology, Taiwan, R.O.C. (Grant No. MOST 107-2218-E-155-012-MY2). The work of Po-Ning Chen is supported by the Ministry of Science and Technology, Taiwan, R.O.C. (Grant No. MOST 108-2221-E-009-024-MY2). The work of Fady Alajaji is supported by the Natural Sciences and Engineering Research Council of Canada. The work of Yunghsiang S. Han is supported by the National Natural Science Foundation of China (Grant No. 61671007) and Start Fund of Dongguan University of Technology (KCYXM2017025). Parts of this work were presented at the 2020 IEEE International Symposium on Information Theory [1].

crossover probability $p < 1/2$. More specifically, for any code \mathcal{C}_n , we establish the following upper bound on its minimum probability of decoding error, a_n , in terms of the asymptotic generalized Poor-Verdú lower bound denoted by b_n :

$$a_n \leq (1 + cn)b_n, \quad (1)$$

where $c = (1 - p)/p$ is the channel (likelihood ratio) constant. In other words, we prove that the deviation $a_n - b_n$ grows at most linearly in n as a factor of b_n . A direct by-product of inequality (1) is that $\limsup_{n \rightarrow \infty} \frac{1}{n} \log \frac{a_n}{b_n} = 0$, showing that the asymptotic generalized bound is exponentially tight for arbitrary sequences of codes and hence achieves the BSC error exponent or reliability function at any (non-negative) rate. We also extend the above results for the class of non-binary memoryless symmetric channels.

The exact characterization of the channel reliability function at low rates remains a long-standing open problem; in-depth studies on this focal information-theoretic function and related problems include the classical papers [4]–[7] and texts [8]–[11] and the more recent works [12]–[23] (see also the references therein). In previous related work, Poor and Verdú conjectured in [2] that their original error lower bound for multihypothesis testing, which yields an upper bound on the channel coding reliability function, is tight for all rates and arbitrary channels. The conjecture was disproved in [24], where the bound was shown to be loose for the BEC at low rates. In [15], Polyanskiy showed that the original Poor-Verdú bound [2] actually coincides with the sphere-packing error exponent bound for discrete memoryless channels (and is hence loose at low rates for this entire class of channels). Our results, while not explicitly determining the reliability function for the BSC and the non-binary symmetric channel, provide an alternative approach for studying it and answers in the positive a variant of the original conjecture in [2] by showing the exponential tightness of the asymptotic generalized Poor-Verdú bound for these channels.

In proving inequality (1) for any block code \mathcal{C}_n sent over the BSC, we first observe that the asymptotic generalized Poor-Verdú bound b_n exactly equals the probability of the set $\mathsf{N}(\mathcal{C}_n)$ of “no decoder ties,” which consists of all input-output n -tuple pairs $(x^n, y^n) \in \mathcal{C}_n \times \mathcal{Y}^n$ satisfying

$$d(x^n, y^n) > \min_{u^n \in \mathcal{C}_n \setminus \{x^n\}} d(u^n, y^n), \quad (2)$$

where $d(\cdot, \cdot)$ is the Hamming distance and \mathcal{Y} is the channel output alphabet. By adding the probability of all “decoder ties,” i.e., all $(x^n, y^n) \in \mathcal{C}_n \times \mathcal{Y}^n$ such that

$$d(x^n, y^n) = \min_{u^n \in \mathcal{C}_n \setminus \{x^n\}} d(u^n, y^n), \quad (3)$$

which are collected in the set $\mathsf{T}(\mathcal{C}_n)$, to $b_n = \Pr(\mathsf{N}(\mathcal{C}_n))$, an upper bound on the minimum probability of decoding error a_n is then obtained. The problem becomes equivalent to proving that $\Pr(\mathsf{T}(\mathcal{C}_n))/b_n$ is no larger than cn . This

is shown using an intricate sequence of constructing finite covers, partitions and disjoint subsets of the two key decoder sets $\mathbb{N}(\mathcal{C}_n)$ and $\mathbb{T}(\mathcal{C}_n)$, until we ultimately arrive at “atomic” subsets $\mathcal{B}_{i,j}(u^n|k)$ and $\Omega_{i,j}(u^n|k)$ of $\mathbb{N}(\mathcal{C}_n)$ and $\mathbb{T}(\mathcal{C}_n)$, respectively, whose probability ratio $\Pr(\mathcal{B}_{i,j}(u^n|k))/\Pr(\Omega_{i,j}(u^n|k))$ is diligently shown (via counting arguments) to be upper-bounded by cn . The essential ingredients of the proof are outlined in Steps 1-5 at the end of Section III and are illustrated in Fig. 1. With inequality (1) proven, the exponential tightness of b_n to a_n is directly confirmed. A similar proof technique, with slightly more elaborate modifications, is used to extend the results for the non-binary symmetric channel.

The rest of the paper is organized as follows. The exact expression of the asymptotic generalized Poor-Verdú lower bound on the error probability in multihypothesis testing is presented in Section II. The error analysis of this asymptotic bound is carried out in detail for the channel coding problem over the memoryless BSC in Sections III and IV. Extensions of these results to the memoryless q -ary symmetric channel (with $q > 2$) are given in Section V. Finally, conclusions are drawn in Section VI.

II. ASYMPTOTIC EXPRESSION OF THE GENERALIZED POOR-VERDÚ BOUND

In 1995, Poor and Verdú established a lower bound on the error probability of multihypothesis testing [2]. This bound was generalized in [3] in terms of a *tilted* posterior hypothesis distribution with tilting parameter $\theta \geq 1$ (with the original bound in [2] recovered when $\theta = 1$).

Proposition 1: (Generalized Poor-Verdú bound [3]): Consider random variables X and Y , governed by the joint distribution $P_{X,Y}$, and that take values in a discrete (i.e., finite or countably infinite) alphabet \mathcal{X} and an arbitrary alphabet \mathcal{Y} , respectively. The minimum probability of error P_e in estimating X from Y satisfies

$$P_e \geq (1 - \alpha) \cdot P_{X,Y} \left\{ (x, y) \in \mathcal{X} \times \mathcal{Y} : P_{X|Y}^{(\theta)}(x|y) \leq \alpha \right\} \quad (4)$$

for each $\alpha \in [0, 1]$ and arbitrary $\theta \geq 1$, where for $(x, y) \in \mathcal{X} \times \mathcal{Y}$,

$$P_{X|Y}^{(\theta)}(x|y) \triangleq \frac{(P_{X|Y}(x|y))^\theta}{\sum_{u \in \mathcal{X}} (P_{X|Y}(u|y))^\theta} \quad (5)$$

is the tilted distribution of $P_{X|Y}(x|y)$ with parameter θ .

In [3], it is illustrated via examples that the lower bound in (4) improves in general as θ grows. However, the asymptotic expression of (4), as θ goes to infinity, was not established in closed-form. This issue is resolved in what follows.

Lemma 1: Given random variables X and Y with joint distribution $P_{X,Y}$ such that the marginal distribution P_X have finite support $\mathcal{C} \subseteq \mathcal{X}$, we have that for $\alpha < 1/|\mathcal{C}|$,

$$\limsup_{\theta \rightarrow \infty} P_{X,Y} \left\{ (x, y) \in \mathcal{X} \times \mathcal{Y} : P_{X|Y}^{(\theta)}(x|y) \leq \alpha \right\} = P_{X,Y} \left\{ (x, y) \in \mathcal{X} \times \mathcal{Y} : P_{X|Y}(x|y) < \max_{u \in \mathcal{C}} P_{X|Y}(u|y) \right\}. \quad (6)$$

Proof: Setting $\alpha = e^{-\kappa}$ in the right-hand side (RHS) probability term in (4) yields

$$\begin{aligned} & P_{X,Y} \left\{ (x,y) \in \mathcal{X} \times \mathcal{Y} : P_{X|Y}^{(\theta)}(x|y) \leq e^{-\kappa} \right\} \\ &= P_{X,Y} \left\{ (x,y) \in \mathcal{X} \times \mathcal{Y} : \frac{(P_{X|Y}(x|y))^\theta}{\sum_{u \in \mathcal{C}} (P_{X|Y}(u|y))^\theta} \leq e^{-\kappa} \right\} \end{aligned} \quad (7)$$

$$= P_{X,Y} \left\{ (x,y) \in \mathcal{X} \times \mathcal{Y} : \log P_{X|Y}(x|y) \leq \frac{1}{\theta} \log \left(\sum_{u \in \mathcal{C}} (P_{X|Y}(u|y))^\theta \right) - \frac{\kappa}{\theta} \right\} \quad (8)$$

$$= P_{X,Y} \left\{ (x,y) \in \mathcal{X} \times \mathcal{Y} : \log P_{X|Y}(x|y) \leq \log \|P_{X|Y}(\cdot|y)\|_\theta - \frac{\kappa}{\theta} \right\}, \quad (9)$$

where $\|P_{X|Y}(\cdot|y)\|_\theta \triangleq (\sum_{u \in \mathcal{C}} (P_{X|Y}(u|y))^\theta)^{1/\theta}$ is the θ -norm of $P_{X|Y}(\cdot|y)$ for a fixed $y \in \mathcal{Y}$. Noting that

$$\begin{aligned} \log P_{X|Y}(x|y) &\leq \log \|P_{X|Y}(\cdot|y)\|_\theta - \frac{\kappa}{\theta} \\ &\iff \frac{\kappa}{\theta} \leq \log \|P_{X|Y}(\cdot|y)\|_\theta - \log P_{X|Y}(x|y), \end{aligned} \quad (10)$$

we separately consider the following two cases.

Case 1: For (x,y) with $P_{X|Y}(x|y) < \max_{u \in \mathcal{C}} P_{X|Y}(u|y)$, the RHS of (10) tends to

$$\log \max_{u \in \mathcal{C}} P_{X|Y}(u|y) - \log P_{X|Y}(x|y) > 0 \quad (11)$$

(which is strictly positive) as θ grows without bound (since the θ -norm of $P_{X|Y}(\cdot|y)$ approaches its infinity-norm given by $\max_{u \in \mathcal{C}} P_{X|Y}(u|y)$), while the left-hand side of (10) tends to zero. Hence, (10) holds for θ sufficiently large.

Case 2: For (x,y) with $P_{X|Y}(x|y) = \max_{u \in \mathcal{C}} P_{X|Y}(u|y)$, we have

$$\log \|P_{X|Y}(\cdot|y)\|_\theta - \log P_{X|Y}(x|y) = \log \left(\sum_{u \in \mathcal{C}} (P_{X|Y}(u|y))^\theta \right)^{1/\theta} - \log P_{X|Y}(x|y) \quad (12)$$

$$= \log \frac{\left(\sum_{u \in \mathcal{C}} (P_{X|Y}(u|y))^\theta \right)^{1/\theta}}{\left((P_{X|Y}(x|y))^\theta \right)^{1/\theta}} \quad (13)$$

$$= \frac{1}{\theta} \log \frac{\sum_{u \in \mathcal{C}} (P_{X|Y}(u|y))^\theta}{(P_{X|Y}(x|y))^\theta} \quad (14)$$

$$\leq \frac{1}{\theta} \log |\mathcal{C}|, \quad (15)$$

where the last inequality holds since $P_{X|Y}(u|y) \leq P_{X|Y}(x|y)$ for all $u \in \mathcal{C}$. Thus (10) is violated since $\kappa = -\log \alpha > \log |\mathcal{C}|$.

Verifying the above two cases completes the proof. ■

In light of Lemma 1, we can fix $\kappa = -\log \alpha > \log |\mathcal{C}|$, take θ to infinity, and obtain from (4) that

$$P_e \geq (1 - e^{-\kappa}) P_{X,Y} \left\{ (x,y) \in \mathcal{X} \times \mathcal{Y} : P_{X|Y}(x|y) < \max_{u \in \mathcal{C}} P_{X|Y}(u|y) \right\}. \quad (16)$$

Since (16) holds for $\kappa > \log |\mathcal{C}|$ arbitrarily large, the following asymptotic expression of the generalized Poor-Verdú bound is established.

Corollary 1: (Asymptotic Generalized Poor-Verdú bound): The minimum error probability P_e in estimating X from Y satisfies

$$P_e \geq P_{X,Y} \left\{ (x, y) \in \mathcal{X} \times \mathcal{Y} : P_{X|Y}(x|y) < \max_{u \in \mathcal{C}} P_{X|Y}(u|y) \right\}. \quad (17)$$

Remark 1: We make two remarks in light of Corollary 1.

- First, the optimal estimate of X from observing Y is known to be the maximum *a posteriori* estimate, given by

$$e(y) = \arg \max_{x \in \mathcal{C}} P_{X|Y}(x|y), \quad (18)$$

where (18) can in fact directly yield the lower bound in (17). This indicates that as the tilting parameter θ goes to infinity, the generalized Poor-Verdú bound can indeed approach¹

$$1 - P_{X,Y} \left\{ (x, y) \in \mathcal{X} \times \mathcal{Y} : P_{X|Y}(x|y) = P_{X|Y}(e(y)|y) \right\}. \quad (19)$$

As a consequence, the lower bound in (17) is tight if and only if the x that maximizes $P_{X|Y}(x|y)$ is unique for all $y \in \mathcal{Y}$. This elucidates why in the example of [3, Fig. 1] the generalized Poor-Verdú bound achieves the minimum probability of error P_e when θ grows unbounded.

- Second, an alternative lower bound for P_e is the Verdú-Han bound established in [25]. This bound was recently generalized in [26, Thm. 1]. We remark that the Verdú-Han bound is not tight even if $P_{X|Y}(\cdot|y)$ admits a unique maximizer for every $y \in \mathcal{Y}$. For example, we can obtain from the ternary hypothesis testing example in [3, Sec. III-A] and [26, Sec. III-A] that:

$$P_e = \frac{3}{5} > \max_{\gamma \geq 0} (\Pr [P_{X|Y}(X|Y) \leq \gamma] - \gamma) = \frac{27}{47}, \quad (20)$$

where the maximizer in (20) is $\gamma^* = \frac{20}{47}$. Noting the sub-optimality of the Verdú-Han bound, the authors in [26] generalized it by varying the output statistics. They also proved the tightness of the resulting generalized Verdú-Han bound:

$$P_e = \max_{Q_Y} \max_{\gamma \geq 0} \left(\Pr \left[\frac{P_{X,Y}(X, Y)}{Q_Y(Y)} \leq \gamma \right] - \gamma \right). \quad (21)$$

¹ Note that the set $\{(x, y) \in \mathcal{X} \times \mathcal{Y} : P_{X|Y}(x|y) = P_{X|Y}(e(y)|y)\}$ includes all ties. For example, for the 2-fold BSC (i.e., the BSC used twice to transmit 2-tuple inputs) with uniform P_X over $\mathcal{C} = \{00, 11\}$, both (00, 01) and (11, 01) will be in this set, i.e.,

$$\{(x, y) \in \mathcal{X} \times \mathcal{Y} : P_{X|Y}(x|y) = P_{X|Y}(e(y)|y)\} = \{(00, 00), (00, 01), (11, 01), (00, 10), (11, 10), (11, 11)\}.$$

It is pertinent to note that the maximizers of (21) are given by

$$\gamma^* = \int_{\mathcal{Y}} \max_{x \in \mathcal{X}} P_{X,Y}(x, y) dP_Y(y) = 1 - P_e \quad (22)$$

and

$$Q_Y^*(y) = \frac{\max_{x \in \mathcal{X}} P_{X,Y}(x, y)}{\int_{\mathcal{Y}} \max_{x \in \mathcal{X}} P_{X,Y}(x, y) dP_Y(y)} \quad (23)$$

$$= \frac{P_Y(y) P_{X|Y}(e(y)|y)}{1 - P_e}. \quad (24)$$

Hence, the determination of the maximizers of the above generalized Verdú-Han bound is equivalent to determining the minimum error probability P_e itself.

Similar to the generalized Poor-Verdú bound with parameter θ , any Q_Y and γ adopted for the generalized Verdú-Han bound yields a lower bound on P_e . However, an interesting difference between the generalized Poor-Verdú bound and the generalized Verdú-Han bound is that when P_X is uniformly distributed over its support \mathcal{C} , the former bound can be transformed into a function of the *information density*

$$i_{XW}(x, y) \triangleq \frac{P_{Y|X}(y|x)}{P_Y(y)} \quad (25)$$

of the channel $W = P_{Y|X}$ with input X and output Y , while the latter bound cannot. This transformation may facilitate the interpretation of the error exponent via the information density (or equivalently, the Hamming distance) for memoryless symmetric channels such as the BSC and the (non-binary) q -ary symmetric channel.

III. EXPONENTIAL TIGHTNESS OF THE ASYMPTOTIC GENERALIZED POOR-VERDÚ BOUND

In this section, we prove that the asymptotic expression of the generalized Poor-Verdú bound given in (17) exactly characterizes the error exponent, $E(R)$, for any rate $R \geq 0$, of the BSC with crossover probability $0 < p < \frac{1}{2}$. Note that while $E(R)$ for the BSC at zero-rate as well as at rates above the cutoff rate is already known in terms of a single-letter expression [8], its determination at low *positive* rates is still open. In this work, we do *not explicitly calculate* $E(R)$. Rather, we demonstrate that the bound in (17) is exponentially tight for arbitrary sequences of rate- R block codes transmitted over the BSC, hence indirectly achieving $E(R)$ for any $R \geq 0$. The precise determination (in single-letter form) of $E(R)$ at low rates indeed remains an unsolved challenging problem. However, proving the exponential tightness of the asymptotic generalized Poor-Verdú bound provides an alternative approach for studying $E(R)$ for the BSC (and non-binary symmetric channels as we prove a similar result in Section V), which may also be beneficial for a larger class of channels.

We show the exponential tightness result as a direct consequence of the following theorem, which is one the main contributions of the paper.

Theorem 1: For any sequence of codes $\{\mathcal{C}_n\}_{n=1}^{\infty}$ of blocklength n and size $|\mathcal{C}_n| = M_n$, with $\mathcal{C}_n \subseteq \mathcal{X}^n \triangleq \{0, 1\}^n$, let

$$a_n \triangleq P_e(\mathcal{C}_n) \quad (26)$$

denote the minimum probability of decoding error for transmitting \mathcal{C}_n over the BSC with crossover probability $0 < p < 1/2$, under a uniform distribution P_{X^n} over \mathcal{C}_n , where X^n is the n -tuple (X_1, \dots, X_n) . Then

$$b_n \leq a_n \leq \left(1 + \frac{(1-p)}{p}n\right) b_n, \quad (27)$$

where

$$b_n \triangleq P_{X^n, Y^n} \left\{ (x^n, y^n) \in \mathcal{X}^n \times \mathcal{Y}^n : P_{X^n|Y^n}(x^n|y^n) < \max_{u^n \in \mathcal{C}_n} P_{X^n|Y^n}(u^n|y^n) \right\} \quad (28)$$

denotes the asymptotic generalized Poor-Verdú bound (i.e., the RHS of (17) applied to this channel coding context).

Remark 2: Theorem 1 reveals that for any *arbitrary* sequence of block codes $\{\mathcal{C}_n\}_{n \geq 1}$ used over the BSC, the relative deviation of the minimum probability of decoding error a_n from the asymptotic generalized Poor-Verdú lower bound b_n is *at most linear* in the blocklength n : $(a_n - b_n)/b_n \leq cn$, where $c = (1-p)/p$. Indeed, (27) directly implies that for the BSC, b_n is exponentially tight with a_n (and hence achieves the BSC reliability function at any rate); this is summarized in the next corollary.

Corollary 2: (Exponential Tightness of (28) for the BSC): Under the same assumptions as in Theorem 1, the asymptotic generalized Poor-Verdú bound b_n achieves the exponent of the minimum decoding error probability a_n for all coding rates, i.e.,

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log \frac{a_n}{b_n} = 0. \quad (29)$$

Remark 3: It is worth emphasizing that Corollary 2 does *not* hold for the BEC; i.e., the asymptotic generalized Poor-Verdú bound in (28) is not exponentially tight for this channel [24]. Indeed for the BEC, the bound in (4) (when translated to the channel coding problem) is unchanged for every $\theta \geq 1$, including when $\theta \rightarrow \infty$, and is hence identical to the original Poor-Verdú bound. The latter bound was shown in [24] not to achieve the BEC's error exponent at low rates; see also [15]. One plausible factor for this difference in tightness of b_n between these two channels, is that the probability of decoder ties is significantly more dominant for the BEC than for the BSC.

Overview of the Proof of Theorem 1: Before providing the full proof of Theorem 1 in Section IV, we outline its steps and elucidate the underlying concepts behind it.

Unlike our earlier result in the companion conference paper [1], where we show (29) for zero coding rates by upper-bounding a_n via the union bound and by lower-bounding b_n via properly designed subsets of its set in (28) (see (32) below), the proof of (29) for *arbitrary* (i.e., zero or positive) coding rates requires us to considerably

tighten the upper and lower bounds on a_n and b_n , respectively.

We next introduce the necessary notation and highlight how we develop the aforementioned tightened bounds needed to prove (27). Fix a sequence of codes $\{\mathcal{C}_n\}_{n=1}^{\infty}$ of blocklength n and size M_n , with $\mathcal{C}_n \subseteq \{0, 1\}^n$, and let the channel input distribution P_{X^n} be uniform over \mathcal{C}_n . Then the code's minimal probability of error a_n is achieved under maximum likelihood (ML) decoding. For the BSC (with $p < 1/2$), the ML estimate based on any received n -tuple y^n at the channel output is obtained via the Hamming distances $\{d(x^n, y^n)\}_{x^n \in \mathcal{C}_n}$. Let $\mathbf{E}(\mathcal{C}_n)$ denote the set of input-output n -tuple pairs (x^n, y^n) that result in an *erroneous* decision for a non-randomized ML decoder of code \mathcal{C}_n ; then

$$a_n = P_e(\mathcal{C}_n) = P_{X^n, Y^n}(\mathbf{E}(\mathcal{C}_n)). \quad (30)$$

Defining the set of *ties* with respect to code \mathcal{C}_n as

$$\mathbf{T}(\mathcal{C}_n) \triangleq \left\{ (x^n, y^n) \in \mathcal{C}_n \times \mathcal{Y}^n : d(x^n, y^n) = \min_{u^n \in \mathcal{C}_n \setminus \{x^n\}} d(u^n, y^n) \right\} \quad (31)$$

and the set of *no-ties* with respect to \mathcal{C}_n

$$\mathbf{N}(\mathcal{C}_n) \triangleq \left\{ (x^n, y^n) \in \mathcal{C}_n \times \mathcal{Y}^n : d(x^n, y^n) > \min_{u^n \in \mathcal{C}_n \setminus \{x^n\}} d(u^n, y^n) \right\}, \quad (32)$$

we have the following set relationships

$$\mathbf{N}(\mathcal{C}_n) \subseteq \mathbf{E}(\mathcal{C}_n) \subseteq (\mathbf{N}(\mathcal{C}_n) \cup \mathbf{T}(\mathcal{C}_n)), \quad (33)$$

where the second inclusion occurs when we have decoding errors for $(x^n, y^n) \in \mathbf{T}(\mathcal{C}_n)$. Finally, examining the inequality condition in (28), we directly have that

$$b_n = P_{X^n, Y^n}(\mathbf{N}(\mathcal{C}_n)). \quad (34)$$

From (33), (30) and (34), we have that

$$b_n \leq a_n \leq b_n + \delta_n \iff 1 \leq \frac{a_n}{b_n} \leq 1 + \frac{\delta_n}{b_n}, \quad (35)$$

where²

$$\delta_n \triangleq P_{X^n, Y^n}(\mathbf{T}(\mathcal{C}_n)). \quad (36)$$

To establish (27), we next derive an upper bound on $\frac{\delta_n}{b_n}$ by upper-bounding $\delta_n = P_{X^n, Y^n}(\mathbf{T}(\mathcal{C}_n))$ and lower-bounding $b_n = P_{X^n, Y^n}(\mathbf{N}(\mathcal{C}_n))$. These two bounds are obtained via judicious choices of covers, partitions, and disjoint subsets of $\mathbf{T}(\mathcal{C}_n)$ and $\mathbf{N}(\mathcal{C}_n)$. The steps, which are illustrated in Fig. 1, are as follows:

- 1) Indexing the codewords as $\mathcal{C}_n = \{x_{(1)}^n, x_{(2)}^n, \dots, x_{(M_n)}^n\}$, we first partition $\mathbf{T}(\mathcal{C}_n)$ and $\mathbf{N}(\mathcal{C}_n)$ along the codewords $x_{(i)}^n \in \mathcal{C}_n$ via $\{\{x_{(i)}^n\} \times \mathcal{T}_i\}_{i=1}^{M_n}$ and $\{\{x_{(i)}^n\} \times \mathcal{N}_i\}_{i=1}^{M_n}$, respectively, where $\mathcal{T}_i \subset \mathcal{Y}^n$ and $\mathcal{N}_i \subset \mathcal{Y}^n$, $i = 1, \dots, M_n$.

²Note if $\delta_n = 0$, then (35) is tight. So without loss of generality, we assume that $\delta_n > 0$.




Fig. 1. Illustration of the concepts behind the proof of Theorem 1.

Note that subsets \mathcal{T}_i or \mathcal{N}_i might be empty for some i . Also, since we assume that $\delta_n = P_{X^n, Y^n}(\mathcal{T}(\mathcal{C}_n)) > 0$, there exists at least one index i such that $\mathcal{T}_i \neq \emptyset$.

- 2) For each \mathcal{T}_i , we construct a disjoint covering $\{\mathcal{B}_{i,j}\}_{j=1, j \neq i}^{M_n}$ for it; i.e., $\mathcal{T}_i \subseteq \cup_{j=1, j \neq i}^{M_n} \mathcal{B}_{i,j}$ and $\mathcal{B}_{i,j} \cap \mathcal{B}_{i,j'} = \emptyset$ for all $j \neq j'$ in $\{1, 2, \dots, M_n\} \setminus \{i\}$. Also, we find a collection of disjoint subsets of \mathcal{N}_i , denoted by $\{\Omega_{i,j}\}_{j=1, j \neq i}^{M_n}$.
- 3) For each $\mathcal{B}_{i,j}$, we partition it via $\{\mathcal{B}_{i,j}(k)\}_{k=0}^{\ell}$ for some finite ℓ . We also obtain a size- $(\ell + 1)$ collection of disjoint subsets of $\Omega_{i,j}$, denoted by $\{\Omega_{i,j}(k)\}_{k=0}^{\ell}$.
- 4) For each $\mathcal{B}_{i,j}(k)$, we further partition it via $\{\mathcal{B}_{i,j}(u^n|k)\}$, which is indexed by a set of properly chosen $u^n \in \mathcal{B}_{i,j}(k)$. Likewise, we obtain a collection of disjoint subsets of $\Omega_{i,j}(k)$, denoted by $\{\Omega_{i,j}(u^n|k)\}$.
- 5) Then, the desired upper bound on $\frac{P_{X^n, Y^n}(\mathcal{T}(\mathcal{C}_n))}{P_{X^n, Y^n}(\mathcal{N}(\mathcal{C}_n))}$ is derived by upper-bounding $\frac{P_{Y^n}(\mathcal{B}_{i,j}(u^n|k))}{P_{Y^n}(\Omega_{i,j}(u^n|k))}$ via systematic counting arguments.

We remark that the achievability of $E(0)$ in [1] was proven by adopting only Steps 1 and 2 above and the use of the union bound. However, in this work, different definitions of the $\mathcal{B}_{i,j}$ and $\Omega_{i,j}$ sets are required from the ones in [1] in order to arrive at a more refined estimate of the associated probabilities when the coding rate is positive. The additional partitions and collections of disjoint subsets in Steps 3 and 4 then allow the determination of the probabilities $\Pr(\mathcal{B}_{i,j})$ and $\Pr(\Omega_{i,j})$, leading in Step 5 to the desired result (27).

IV. THE PROOF OF THEOREM 1

In Section IV-A, we give the main body of the proof where Steps 1-3 described above are applied. For better readability, the verification of a key inequality (see (47) below), where Steps 4 and 5 are minutely carried out, is separately provided in Section IV-B.

A. Proof

Denote for convenience $\mathcal{C}_n = \{x_{(1)}^n, x_{(2)}^n, \dots, x_{(M_n)}^n\}$ and define

$$\mathcal{T}_i \triangleq \{y^n \in \mathcal{Y}^n : (x_{(i)}^n, y^n) \in \mathbb{T}(\mathcal{C}_n)\}, \quad i = 1, \dots, M_n. \quad (37)$$

Then $\{\{x_{(i)}^n\} \times \mathcal{T}_i\}_{i=1}^{M_n}$ forms a partition of $\mathbb{T}(\mathcal{C}_n)$,³ and we have

$$\delta_n = P_{X^n, Y^n}(\mathbb{T}(\mathcal{C}_n)) = \sum_{i=1}^{M_n} P_{X^n}(x_{(i)}^n) \Pr\left(Y^n \in \mathcal{T}_i \mid X^n = x_{(i)}^n\right). \quad (38)$$

Similarly, we define for $i = 1, \dots, M_n$,

$$\mathcal{N}_i \triangleq \{y^n \in \mathcal{Y}^n : (x_{(i)}^n, y^n) \in \mathbb{N}(\mathcal{C}_n)\}, \quad (39)$$

and obtain a partition of $\mathbb{N}(\mathcal{C}_n)$, denoted by $\{\{x_{(i)}^n\} \times \mathcal{N}_i\}_{i=1}^{M_n}$. We similarly have

$$b_n = P_{X^n, Y^n}(\mathbb{N}(\mathcal{C}_n)) = \sum_{i=1}^{M_n} P_{X^n}(x_{(i)}^n) \Pr\left(Y^n \in \mathcal{N}_i \mid X^n = x_{(i)}^n\right). \quad (40)$$

We next define $\{\mathcal{B}_{i,j}\}_{j=1, j \neq i}^{M_n}$ and $\{\Omega_{i,j}\}_{j=1, j \neq i}^{M_n}$, where the former is a disjoint covering of \mathcal{T}_i and the latter are disjoint subsets of \mathcal{N}_i . For $1 \leq j \leq M_n$ and $j \neq i$, let

$$\mathcal{B}_{i,j} \triangleq \left\{ y^n \in \mathcal{Y}^n : d(x_{(j)}^n, y^n) = d(x_{(i)}^n, y^n) < \min_{1 \leq k < j, k \neq i} d(x_{(k)}^n, y^n) \right\} \quad (41)$$

and

$$\Omega_{i,j} \triangleq \left\{ y^n \in \mathcal{Y}^n : d(x_{(j)}^n, y^n) + 2 = d(x_{(i)}^n, y^n) \leq \min_{1 \leq k < j, k \neq i} d(x_{(k)}^n, y^n) \right\}, \quad (42)$$

where we adopt the convention that the minimum of an empty set is infinity, e.g.,

$$\min_{1 \leq k < 2, k \neq 1} d(x_{(k)}^n, y^n) = \infty. \quad (43)$$

Note that every y^n in $\Omega_{i,j}$ is also in \mathcal{N}_i due to the constraint $d(x_{(j)}^n, y^n) + 2 = d(x_{(i)}^n, y^n)$.

In what follows, we consider the case of $i = 1$ first. The cases of $2 \leq i \leq M_n$ can be similarly treated. From (41) and (42), it readily follows that:

$$i) \mathcal{B}_{1,j_1} \cap \mathcal{B}_{1,j_2} = \Omega_{1,j_1} \cap \Omega_{1,j_2} = \emptyset \text{ for } 2 \leq j_1 < j_2 \leq M_n;$$

³In this paper, the definition of the partition of a set is different from the normal one where all subsets of the partition are not empty. In this work, subsets of a partition are allowed to be empty sets.

ii) $\{\mathcal{B}_{1,j}\}_{j=2}^{M_n}$ form a disjoint covering of \mathcal{T}_1 , i.e.,

$$\mathcal{T}_1 = \left\{ y^n \in \mathcal{Y}^n : d(x_{(1)}^n, y^n) = \min_{u^n \in \mathcal{C}_n \setminus \{x_{(1)}^n\}} d(u^n, y^n) \right\} \subset \bigcup_{j=2}^{M_n} \mathcal{B}_{1,j}; \quad (44)$$

iii) $\{\Omega_{1,j}\}_{j=2}^{M_n}$ are disjoint subsets of \mathcal{N}_1 and hence,

$$\bigcup_{j=2}^{M_n} \Omega_{1,j} \subset \mathcal{N}_1 = \left\{ y^n \in \mathcal{Y}^n : d(x_{(1)}^n, y^n) > \min_{u^n \in \mathcal{C}_n \setminus \{x_{(1)}^n\}} d(u^n, y^n) \right\}. \quad (45)$$

We then claim that either

$$\Pr \left(Y^n \in \mathcal{B}_{1,j} \mid X^n = x_{(1)}^n \right) = 0 \quad (46)$$

or

$$\frac{\Pr \left(Y^n \in \mathcal{B}_{1,j} \mid X^n = x_{(1)}^n \right)}{\Pr \left(Y^n \in \Omega_{1,j} \mid X^n = x_{(1)}^n \right)} \leq \frac{(1-p)}{p} n, \quad (47)$$

where the proof of the claim is given in Section IV-B.

Collecting the above results yields

$$\Pr \left(Y^n \in \mathcal{T}_1 \mid X^n = x_{(1)}^n \right) \leq \sum_{j=2, \mathcal{B}_{1,j} \neq \emptyset}^{M_n} \Pr \left(Y^n \in \mathcal{B}_{1,j} \mid X^n = x_{(1)}^n \right) \quad (48)$$

$$\leq \frac{(1-p)}{p} n \sum_{j=2, \mathcal{B}_{1,j} \neq \emptyset}^{M_n} \Pr \left(Y^n \in \Omega_{1,j} \mid X^n = x_{(1)}^n \right) \quad (49)$$

$$= \frac{(1-p)}{p} n \Pr \left(Y^n \in \bigcup_{j=2, \mathcal{B}_{1,j} \neq \emptyset}^{M_n} \Omega_{1,j} \mid X^n = x_{(1)}^n \right) \quad (50)$$

$$\leq \frac{(1-p)}{p} n \Pr \left(Y^n \in \mathcal{N}_1 \mid X^n = x_{(1)}^n \right), \quad (51)$$

where (48) follows from (44), (49) holds because of (46) and (47), (50) holds because $\{\Omega_{1,j}\}_{j=2}^{M_n}$ are mutually disjoint, and (51) is a consequence of (45). The same argument can be used to prove that for $2 \leq i \leq M_n$,

$$\Pr \left(Y^n \in \mathcal{T}_i \mid X^n = x_{(i)}^n \right) \leq \frac{(1-p)}{p} n \Pr \left(Y^n \in \mathcal{N}_i \mid X^n = x_{(i)}^n \right). \quad (52)$$

We conclude from (38), (40), (51), and (52) that

$$\frac{\delta_n}{b_n} = \frac{\sum_{i=1}^{M_n} P_{X^n}(x_{(i)}^n) \Pr \left(Y^n \in \mathcal{T}_i \mid X^n = x_{(i)}^n \right)}{\sum_{i=1}^{M_n} P_{X^n}(x_{(i)}^n) \Pr \left(Y^n \in \mathcal{N}_i \mid X^n = x_{(i)}^n \right)} \leq \frac{(1-p)}{p} n, \quad (53)$$

which together with (35) completes the proof of (27).

B. Substantiation of (47)

We next prove inequality (47). Let \mathcal{S}_j consist of the indices of the components where $x_{(j)}^n$ and $x_{(1)}^n$ differ, i.e., $d(x_{(1)}^n, x_{(j)}^n) = |\mathcal{S}_j|$. By this definition, we immediately have $|\mathcal{S}_j| \geq 1$ for $2 \leq j \leq M_n$. Since the situation of $\mathcal{B}_{1,j}$ being empty is covered in (46), and since an odd $|\mathcal{S}_j|$ implies that $\mathcal{B}_{1,j}$ is empty, it suffices to prove (47) for

$|\mathcal{S}_j| = 2\ell_j$ even. We then separately prove (47) for each of the following four cases: the specific cases of $j = 2$, $j = 3$ and $j = 4$ and the case for $5 \leq j \leq M_n$.

1) *Proof of (47) for $j = 2$ with $|\mathcal{S}_2| = 2\ell_2$:* Denote by $d(u^n, y^n | \mathcal{S})$ the Hamming distance between the sub-portions of u^n and y^n with indices in an index set \mathcal{S} . Define for $0 \leq k \leq 2\ell_2$,

$$\mathcal{B}_{1,2}(k) \triangleq \left\{ y^n \in \mathcal{B}_{1,2} : d(x_{(1)}^n, y^n | \mathcal{S}_2) = k \right\}. \quad (54)$$

Then, $\mathcal{B}_{1,2} = \cup_{k=0}^{2\ell_2} \mathcal{B}_{1,2}(k)$, and $\mathcal{B}_{1,2}(k_1)$ and $\mathcal{B}_{1,2}(k_2)$ are disjoint for $k_1 \neq k_2$.

In this particular case of $j = 2$, we actually have $\mathcal{B}_{1,2}(k)$ is an empty set for every $k \neq \ell_2$; hence, we have $\mathcal{B}_{1,2} = \mathcal{B}_{1,2}(\ell_2)$. We proceed to partition $\mathcal{B}_{1,2}(\ell_2)$ into $2^{n-2\ell_2}$ disjoint subsets, each of which is defined through an element u^n in $\mathcal{B}_{1,2}(\ell_2)$ as

$$\mathcal{B}_{1,2}(u^n | \ell_2) \triangleq \left\{ y^n \in \mathcal{B}_{1,2}(\ell_2) : d(u^n, y^n | \mathcal{S}_2^c) = 0 \right\}. \quad (55)$$

Note that we can pick $2^{n-2\ell_2}$ elements from $\mathcal{B}_{1,2}(\ell_2)$ and collect them in a set \mathcal{U} such that $\mathcal{B}_{1,2}(u^n | \ell_2)$ and $\mathcal{B}_{1,2}(\hat{u}^n | \ell_2)$ are disjoint for distinct u^n and \hat{u}^n in \mathcal{U} , and $\cup_{u^n \in \mathcal{U}} \mathcal{B}_{1,2}(u^n | \ell_2) = \mathcal{B}_{1,2}(\ell_2)$.

On the other hand, we can likewise define for $0 \leq k \leq 2\ell_2$,

$$\Omega_{1,2}(k) \triangleq \left\{ y^n \in \Omega_{1,2} : d(x_{(1)}^n, y^n | \mathcal{S}_2) = k + 1 \right\}, \quad (56)$$

and define for $u^n \in \mathcal{U}$,

$$\Omega_{1,2}(u^n | \ell_2) \triangleq \left\{ y^n \in \Omega_{1,2}(\ell_2) : d(u^n, y^n | \mathcal{S}_2^c) = 0 \right\}, \quad (57)$$

where \mathcal{S}_2^c denotes the complement of \mathcal{S}_2 . It can be seen from the above definitions that $\{\Omega_{1,2}(k)\}_{k=0}^{2\ell_2}$ are disjoint subsets of $\Omega_{1,2}$, and $\{\Omega_{1,2}(u^n | \ell_2)\}_{u^n \in \mathcal{U}}$ are disjoint subsets of $\Omega_{1,2}(\ell_2)$.

We then obtain that for each $u^n \in \mathcal{U}$,

$$|\mathcal{B}_{1,2}(u^n | \ell_2)| = \binom{2\ell_2}{\ell_2} \quad \text{and} \quad |\Omega_{1,2}(u^n | \ell_2)| = \binom{2\ell_2}{\ell_2 + 1}, \quad (58)$$

which implies

$$\frac{\Pr \left(Y^n \in \mathcal{B}_{1,2}(u^n | \ell_2) \mid X^n = x_{(1)}^n \right)}{\Pr \left(Y^n \in \Omega_{1,2}(u^n | \ell_2) \mid X^n = x_{(1)}^n \right)} = \frac{(1-p) \binom{2\ell_2}{\ell_2}}{p \binom{2\ell_2}{\ell_2+1}} \quad (59)$$

$$= \frac{(1-p) (\ell_2 + 1)}{p \ell_2} \quad (60)$$

$$\leq \frac{(1-p)}{p} 2 \quad (61)$$

$$\leq \frac{(1-p)}{p} n, \quad (62)$$

where we upper-bound 2 by n in the last inequality in order to be consistent with what is later shown in (81), (85)

and (99). As a result,

$$\Pr\left(Y^n \in \mathcal{B}_{1,2} \mid X^n = x_{(1)}^n\right) = \Pr\left(Y^n \in \mathcal{B}_{1,2}(\ell_2) \mid X^n = x_{(1)}^n\right) \quad (63)$$

$$= \sum_{u^n \in \mathcal{U}} \Pr\left(Y^n \in \mathcal{B}_{1,2}(u^n | \ell_2) \mid X^n = x_{(1)}^n\right) \quad (64)$$

$$\leq \frac{(1-p)}{p} n \sum_{u^n \in \mathcal{U}} \Pr\left(Y^n \in \Omega_{1,2}(u^n | \ell_2) \mid X^n = x_{(1)}^n\right) \quad (65)$$

$$= \frac{(1-p)}{p} n \Pr\left(Y^n \in \bigcup_{u^n \in \mathcal{U}} \Omega_{1,2}(u^n | \ell_2) \mid X^n = x_{(1)}^n\right) \quad (66)$$

$$\leq \frac{(1-p)}{p} n \Pr\left(Y^n \in \Omega_{1,2} \mid X^n = x_{(1)}^n\right), \quad (67)$$

where (66) follows from (62), and (66) and (67) hold because $\{\Omega_{1,2}(u^n | \ell_2)\}_{u^n \in \mathcal{U}}$ are disjoint subsets of $\Omega_{1,2}$. This completes the proof of (47) for $j = 2$ with $|\mathcal{S}_2| = 2\ell_2$.

2) *Proof of (47) for $j = 3$ with $|\mathcal{S}_3| = 2\ell_3$:* The proof for $j = 3$ is more involved than that for $j = 2$. We divide \mathcal{S}_3 into two disjoint sets:

$$\mathcal{S}_3^{(1)} \triangleq \mathcal{S}_2 \cap \mathcal{S}_3 \quad \text{and} \quad \mathcal{S}_3^{(2)} \triangleq \mathcal{S}_2^c \cap \mathcal{S}_3. \quad (68)$$

Here, the former set consists of those indices in \mathcal{S}_3 that have been included in \mathcal{S}_2 , while the latter set contains the remaining indices. Let $\ell_3^{(1)} \triangleq |\mathcal{S}_3^{(1)}|$ and $\ell_3^{(2)} \triangleq |\mathcal{S}_3^{(2)}|$. Note that $\mathcal{S}_3 = \mathcal{S}_3^{(1)} \cup \mathcal{S}_3^{(2)}$ and $|\mathcal{S}_3| = \ell_3^{(1)} + \ell_3^{(2)} = 2\ell_3 \geq 2$.

Define for $0 \leq k \leq \ell_3^{(1)}$,

$$\mathcal{B}_{1,3}^{(1)}(k) \triangleq \left\{ y^n \in \mathcal{B}_{1,3} : d(x_{(1)}^n, y^n | \mathcal{S}_3^{(1)}) = k \right\}, \quad (69)$$

and for $0 \leq k \leq \ell_3^{(2)}$,

$$\mathcal{B}_{1,3}^{(2)}(k) \triangleq \left\{ y^n \in \mathcal{B}_{1,3} : d(x_{(1)}^n, y^n | \mathcal{S}_3^{(1)}) = \ell_3^{(1)} \text{ and } d(x_{(1)}^n, y^n | \mathcal{S}_3^{(2)}) = k \right\}. \quad (70)$$

Since $y^n \in \mathcal{B}_{1,3}$ must satisfy $d(x_{(1)}^n, y^n | \mathcal{S}_3) = d(x_{(3)}^n, y^n | \mathcal{S}_3) = \ell_3 < 2\ell_3 = |\mathcal{S}_3| = \ell_3^{(1)} + \ell_3^{(2)}$, we immediately have

$$\mathcal{B}_{1,3}^{(2)}(\ell_3^{(2)}) = \emptyset. \quad (71)$$

Then, we have from the definitions above that⁴

- i) $\mathcal{B}_{1,3}^{(1)}(\ell_3^{(1)}) = \bigcup_{k=0}^{\ell_3^{(2)}-1} \mathcal{B}_{1,3}^{(2)}(k)$;
- ii) $\mathcal{B}_{1,3} = \bigcup_{k^{(1)}=0}^{\ell_3^{(1)}} \mathcal{B}_{1,3}^{(1)}(k^{(1)}) = \left(\bigcup_{k^{(1)}=0}^{\ell_3^{(1)}-1} \mathcal{B}_{1,3}^{(1)}(k^{(1)}) \right) \cup \left(\bigcup_{k^{(2)}=0}^{\ell_3^{(2)}-1} \mathcal{B}_{1,3}^{(2)}(k^{(2)}) \right)$;
- iii) $\{\mathcal{B}_{1,3}^{(1)}(k)\}_{k=0}^{\ell_3^{(1)}-1}$ and $\{\mathcal{B}_{1,3}^{(2)}(k)\}_{k=0}^{\ell_3^{(2)}-1}$ are mutually disjoint.

For $0 \leq k < \ell_3^{(1)}$, we further partition $\mathcal{B}_{1,3}^{(1)}(k)$ into disjoint subsets, each of which is indexed by a specifically

⁴ By convention, we set $\bigcup_{k=0}^{-1} \mathcal{A}_k = \emptyset$.

chosen $u^n \in \mathcal{B}_{1,3}^{(1)}(k)$ and is defined by

$$\mathcal{B}_{1,3}^{(1)}(u^n|k) \triangleq \left\{ y^n \in \mathcal{B}_{1,3}^{(1)}(k) : d(u^n, y^n | (\mathcal{S}_3^{(1)})^c) = 0 \right\}. \quad (72)$$

Collect those representative u^n in $\mathcal{U}_3^{(1)}(k)$ such that $\mathcal{B}_{1,3}^{(1)}(k) = \bigcup_{u^n \in \mathcal{U}_3^{(1)}(k)} \mathcal{B}_{1,3}^{(1)}(u^n|k)$. We can similarly partition $\mathcal{B}_{1,3}^{(2)}(k)$ into disjoint $\{\mathcal{B}_{1,3}^{(2)}(u^n|k)\}_{u^n \in \mathcal{U}_3^{(2)}(k)}$ with

$$\mathcal{B}_{1,3}^{(2)}(u^n|k) \triangleq \left\{ y^n \in \mathcal{B}_{1,3}^{(2)}(k) : d(u^n, y^n | (\mathcal{S}_3^{(2)})^c) = 0 \right\}, \quad (73)$$

and obtain $\mathcal{B}_{1,3}^{(2)}(k) = \bigcup_{u^n \in \mathcal{U}_3^{(2)}(k)} \mathcal{B}_{1,3}^{(2)}(u^n|k)$.

On the other hand, we can likewise define for $0 \leq k < \ell_3^{(1)}$,

$$\Omega_{1,3}^{(1)}(k) \triangleq \left\{ y^n \in \Omega_{1,3} : d(x_{(1)}^n, y^n | \mathcal{S}_3^{(1)}) = k + 1 \right\}, \quad (74)$$

and for $0 \leq k < \ell_3^{(2)}$,

$$\Omega_{1,3}^{(2)}(k) \triangleq \left\{ y^n \in \Omega_{1,3} : d(x_{(1)}^n, y^n | \mathcal{S}_3^{(1)}) = \ell_3^{(1)} \text{ and } d(x_{(1)}^n, y^n | \mathcal{S}_3^{(2)}) = k + 1 \right\}, \quad (75)$$

and for $0 \leq k < \ell_3^{(1)}$ and $u^n \in \mathcal{U}_3^{(1)}(k)$,

$$\Omega_{1,3}^{(1)}(u^n|k) \triangleq \left\{ y^n \in \Omega_{1,3}^{(1)}(k) : d(u^n, y^n | (\mathcal{S}_3^{(1)})^c) = 0 \right\}, \quad (76)$$

and for $0 \leq k < \ell_3^{(2)}$ and $u^n \in \mathcal{U}_3^{(2)}(k)$,

$$\Omega_{1,3}^{(2)}(u^n|k) \triangleq \left\{ y^n \in \Omega_{1,3}^{(2)}(k) : d(u^n, y^n | (\mathcal{S}_3^{(2)})^c) = 0 \right\}. \quad (77)$$

Thus, $\{\Omega_{1,3}^{(1)}(u^n|k)\}_{0 \leq k < \ell_3^{(1)}, u^n \in \mathcal{U}_3^{(1)}(k)}$ and $\{\Omega_{1,3}^{(2)}(u^n|k)\}_{0 \leq k < \ell_3^{(2)}, u^n \in \mathcal{U}_3^{(2)}(k)}$ are all mutually disjoint.

With the above notation, we are ready to derive the desired result (47). For $0 \leq k < \ell_3^{(1)}$ and $u^n \in \mathcal{U}_3^{(1)}(k)$, we establish $|\mathcal{B}_{1,3}^{(1)}(u^n|k)| = \binom{\ell_3^{(1)}}{k}$ and $|\Omega_{1,3}^{(1)}(u^n|k)| = \binom{\ell_3^{(1)}}{k+1}$, and

$$\frac{\Pr\left(Y^n \in \mathcal{B}_{1,3}^{(1)}(u^n|k) \mid X^n = x_{(1)}^n\right)}{\Pr\left(Y^n \in \Omega_{1,3}^{(1)}(u^n|k) \mid X^n = x_{(1)}^n\right)} = \frac{(1-p) \binom{\ell_3^{(1)}}{k}}{p \binom{\ell_3^{(1)}}{k+1}} \quad (78)$$

$$= \frac{(1-p) (k+1)}{p (\ell_3^{(1)} - k)} \quad (79)$$

$$\leq \frac{(1-p)}{p} \ell_3^{(1)} \quad (80)$$

$$\leq \frac{(1-p)}{p} n, \quad (81)$$

where the inequality in (80) follows from $\ell_3^{(1)} - k \geq 1$ and $k+1 \leq \ell_3^{(1)}$. Similarly, for $0 \leq k < \ell_3^{(2)}$ and $u^n \in \mathcal{U}_3^{(2)}(k)$,

we can establish that $|\mathcal{B}_{1,3}^{(2)}(u^n|k)| = \binom{\ell_3^{(2)}}{k}$ and $|\Omega_{1,3}^{(2)}(u^n|k)| = \binom{\ell_3^{(2)}}{k+1}$, and

$$\frac{\Pr\left(Y^n \in \mathcal{B}_{1,3}^{(2)}(u^n|k) \mid X^n = x_{(1)}^n\right)}{\Pr\left(Y^n \in \Omega_{1,3}^{(2)}(u^n|k) \mid X^n = x_{(1)}^n\right)} = \frac{(1-p) \binom{\ell_3^{(2)}}{k}}{p \binom{\ell_3^{(2)}}{k+1}} \quad (82)$$

$$= \frac{(1-p) (k+1)}{p (\ell_3^{(2)} - k)} \quad (83)$$

$$\leq \frac{(1-p)}{p} \ell_3^{(2)} \quad (84)$$

$$\leq \frac{(1-p)}{p} n, \quad (85)$$

where (84) holds since from $\ell_3^{(2)} - k \geq 1$ and $k + 1 \leq \ell_3^{(2)}$. We conclude that

$$\begin{aligned} \Pr\left(Y^n \in \mathcal{B}_{1,3} \mid X^n = x_{(1)}^n\right) &= \sum_{k^{(1)}=0}^{\ell_3^{(1)}-1} \sum_{u^n \in \mathcal{U}_3^{(1)}(k^{(1)})} \Pr\left(Y^n \in \mathcal{B}_{1,3}^{(1)}(u^n | k^{(1)}) \mid X^n = x_{(1)}^n\right) \\ &\quad + \sum_{k^{(2)}=0}^{\ell_3^{(2)}-1} \sum_{u^n \in \mathcal{U}_3^{(2)}(k^{(2)})} \Pr\left(Y^n \in \mathcal{B}_{1,3}^{(2)}(u^n | k^{(2)}) \mid X^n = x_{(1)}^n\right) \end{aligned} \quad (86)$$

$$\begin{aligned} &\leq \frac{(1-p)}{p} n \left[\sum_{k^{(1)}=0}^{\ell_3^{(1)}-1} \sum_{u^n \in \mathcal{U}_3^{(1)}(k^{(1)})} \Pr\left(Y^n \in \Omega_{1,3}^{(1)}(u^n | k^{(1)}) \mid X^n = x_{(1)}^n\right) \right. \\ &\quad \left. + \sum_{k^{(2)}=0}^{\ell_3^{(2)}-1} \sum_{u^n \in \mathcal{U}_3^{(2)}(k^{(2)})} \Pr\left(Y^n \in \Omega_{1,3}^{(2)}(u^n | k^{(2)}) \mid X^n = x_{(1)}^n\right) \right] \end{aligned} \quad (87)$$

$$\leq \frac{(1-p)}{p} n \Pr\left(Y^n \in \Omega_{1,3} \mid X^n = x_{(1)}^n\right), \quad (88)$$

where (87) follows from (81) and (85), and (88) holds because

$$\{\Omega_{1,3}^{(1)}(u^n | k)\}_{0 \leq k < \ell_3^{(1)}, u^n \in \mathcal{U}_3^{(1)}(k)} \quad \text{and} \quad \{\Omega_{1,3}^{(2)}(u^n | k)\}_{0 \leq k < \ell_3^{(2)}, u^n \in \mathcal{U}_3^{(2)}(k)} \quad (89)$$

are mutually disjoint subsets of $\Omega_{1,3}$. This completes the proof of (47) for $j = 3$ with $|\mathcal{S}_3| = 2\ell_3$.

3) *Proof of (47) for $j = 4$ with $|\mathcal{S}_4| = 2\ell_4$:* We partition \mathcal{S}_4 into four (disjoint) sets:

$$\mathcal{S}_4^{(1)} \triangleq \mathcal{S}_2 \cap \mathcal{S}_3 \cap \mathcal{S}_4, \quad \mathcal{S}_4^{(2)} \triangleq \mathcal{S}_2 \cap \mathcal{S}_3^c \cap \mathcal{S}_4, \quad \mathcal{S}_4^{(3)} \triangleq \mathcal{S}_2^c \cap \mathcal{S}_3 \cap \mathcal{S}_4 \quad \text{and} \quad \mathcal{S}_4^{(4)} \triangleq \mathcal{S}_2^c \cap \mathcal{S}_3^c \cap \mathcal{S}_4. \quad (90)$$

Letting $\ell_4^{(m)} \triangleq |\mathcal{S}_4^{(m)}|$ for $1 \leq m \leq 4$, we have $\mathcal{S}_4 = \bigcup_{m=1}^4 \mathcal{S}_4^{(m)}$ with $|\mathcal{S}_4| = \sum_{m=1}^4 \ell_4^{(m)} = 2\ell_4 \geq 2$.

We define for $0 \leq k \leq \ell_4^{(m)}$ and $1 \leq m \leq 4$,

$$\mathcal{B}_{1,4}^{(m)}(k) \triangleq \left\{ y^n \in \mathcal{B}_{1,4} : (\forall 1 \leq i < m) d(x_{(1)}^n, y^n | \mathcal{S}_4^{(i)}) = \ell_4^{(i)} \text{ and } d(x_{(1)}^n, y^n | \mathcal{S}_4^{(m)}) = k \right\}. \quad (91)$$

Since $y^n \in \mathcal{B}_{1,4}$ must satisfy $d(x_{(1)}^n, y^n | \mathcal{S}_4) = d(x_{(4)}^n, y^n | \mathcal{S}_4) = \ell_4 < 2\ell_4 = |\mathcal{S}_4| = \sum_{m=1}^4 \ell_4^{(m)}$, we immediately have

$$\mathcal{B}_{1,4}^{(4)}(\ell_4^{(4)}) = \emptyset. \quad (92)$$

The definitions above imply that

$$i) \mathcal{B}_{1,4} = \bigcup_{m=1}^4 \bigcup_{k=0}^{\ell_4^{(m)}-1} \mathcal{B}_{1,4}^{(m)}(k);$$

$$ii) \{\mathcal{B}_{1,4}^{(m)}(k)\}_{1 \leq m \leq 4, 0 \leq k < \ell_4^{(m)}} \text{ are mutually disjoint.}$$

For $1 \leq m \leq 4$ and $0 \leq k < \ell_4^{(m)}$, we further partition $\mathcal{B}_{1,4}^{(m)}(k)$ into disjoint subsets, each of which is indexed

by a specifically chosen $u^n \in \mathcal{B}_{1,4}^{(m)}(k)$ and is defined by

$$\mathcal{B}_{1,4}^{(m)}(u^n|k) \triangleq \left\{ y^n \in \mathcal{B}_{1,4}^{(m)}(k) : d(u^n, y^n | (\mathcal{S}_4^{(m)})^c) = 0 \right\}. \quad (93)$$

We collect those representative u^n into $\mathcal{U}_4^{(m)}(k)$ such that $\mathcal{B}_{1,4}^{(m)}(k) = \bigcup_{u^n \in \mathcal{U}_4^{(m)}(k)} \mathcal{B}_{1,4}^{(m)}(u^n|k)$.

On the other hand, we can likewise define for $0 \leq k < \ell_4^{(m)}$ and $1 \leq m \leq 4$,

$$\Omega_{1,4}^{(m)}(k) \triangleq \left\{ y^n \in \Omega_{1,4} : (\forall 1 \leq i < m) d(x_{(1)}^n, y^n | \mathcal{S}_4^{(i)}) = \ell_4^{(i)} \text{ and } d(x_{(1)}^n, y^n | \mathcal{S}_4^{(m)}) = k + 1 \right\}, \quad (94)$$

and for $1 \leq m \leq 4$, $0 \leq k < \ell_4^{(m)}$ and $u^n \in \mathcal{U}_4^{(m)}(k)$,

$$\Omega_{1,4}^{(m)}(u^n|k) \triangleq \left\{ y^n \in \Omega_{1,4}^{(m)}(k) : d(u^n, y^n | (\mathcal{S}_4^{(m)})^c) = 0 \right\}. \quad (95)$$

We therefore have $\{\Omega_{1,4}^{(m)}(u^n|k)\}_{1 \leq m \leq 4, 0 \leq k < \ell_4^{(m)}, u^n \in \mathcal{U}_4^{(m)}(k)}$ are all mutually disjoint.

With the above notation, we can now prove (47). For $1 \leq m \leq 4$, $0 \leq k < \ell_4^{(m)}$ and $u^n \in \mathcal{U}_4^{(m)}(k)$, we establish $|\mathcal{B}_{1,4}^{(m)}(u^n|k)| = \binom{\ell_4^{(m)}}{k}$ and $|\Omega_{1,4}^{(m)}(u^n|k)| = \binom{\ell_4^{(m)}}{k+1}$, and

$$\frac{\Pr\left(Y^n \in \mathcal{B}_{1,4}^{(m)}(u^n|k) \mid X^n = x_{(1)}^n\right)}{\Pr\left(Y^n \in \Omega_{1,4}^{(m)}(u^n|k) \mid X^n = x_{(1)}^n\right)} = \frac{(1-p) \binom{\ell_4^{(m)}}{k}}{p \binom{\ell_4^{(m)}}{k+1}} \quad (96)$$

$$= \frac{(1-p)}{p} \frac{(k+1)}{(\ell_4^{(m)} - k)} \quad (97)$$

$$\leq \frac{(1-p)}{p} \ell_4^{(m)} \quad (98)$$

$$\leq \frac{(1-p)}{p} n, \quad (99)$$

where the first inequality in (99) follows from $\ell_4^{(m)} - k \geq 1$ and $k + 1 \leq \ell_4^{(m)}$. Consequently, the same argument as for (88) can be used to prove that

$$\Pr\left(Y^n \in \mathcal{B}_{1,4} \mid X^n = x_{(1)}^n\right) \leq \frac{(1-p)}{p} n \Pr\left(Y^n \in \Omega_{1,4} \mid X^n = x_{(1)}^n\right). \quad (100)$$

This completes the proof of (47) for $j = 4$ with $|\mathcal{S}_4| = 2\ell_4$.

4) *Proof of (47) for $5 \leq j \leq M_n$ with $|\mathcal{S}_j| = 2\ell_j$:* The examination of \mathcal{S}_j for $5 \leq j \leq M_n$ to show that

$$\Pr\left(Y^n \in \mathcal{B}_{1,j} \mid X^n = x_{(1)}^n\right) \leq \frac{(1-p)}{p} n \Pr\left(Y^n \in \Omega_{1,j} \mid X^n = x_{(1)}^n\right) \quad (101)$$

can be similarly done as in the case for $j = 4$. More specifically, for each $5 \leq j \leq M_n$, the set \mathcal{S}_j is partitioned into 2^{j-2} (disjoint) subsets as follows.

Let $\mathcal{I}_j \triangleq \{2, 3, \dots, j-1\}$ and let $\mathcal{P}(\mathcal{I}_j)$ denote its power set, containing 2^{j-2} subsets $\mathcal{I}_j^{(m)}$ of \mathcal{I}_j , $1 \leq m \leq 2^{j-2}$.

For example, for $j = 5$, $\mathcal{I}_5 = \{2, 3, 4\}$ and

$$\mathcal{P}(\mathcal{I}_5) = \{\mathcal{I}_5^{(1)}, \dots, \mathcal{I}_5^{(8)}\} = \{\emptyset, \{2\}, \{3\}, \{4\}, \{2, 3\}, \{2, 4\}, \{3, 4\}, \{2, 3, 4\}\}. \quad (102)$$

Now \mathcal{S}_j is partitioned into the subsets $\mathcal{S}_j^{(m)}$, $m = 1, 2, \dots, 2^{j-2}$, with each $\mathcal{S}_j^{(m)}$ generated by $\mathcal{I}_j^{(m)} \in \mathcal{P}(\mathcal{I}_j)$ as

follows:

$$\mathcal{S}_j^{(m)} = \left(\bigcap_{i \in \mathcal{I}_j^{(m)}} \mathcal{S}_i^c \right) \cap \left(\bigcap_{i \in \mathcal{I}_j \setminus \mathcal{I}_j^{(m)}} \mathcal{S}_i \right) \cap \mathcal{S}_j. \quad (103)$$

Then via the same arguments⁵ as detailed for the case of $j = 4$, the “atomic” collections $\{\mathcal{B}_{1,j}^{(m)}(u^n|k)\}$ and $\{\Omega_{1,j}^{(m)}(u^n|k)\}$ can be constructed to deduce inequality (101) for $5 \leq j \leq M_n$.

The substantiation of (47) is hence completed. ■

V. EXTENSION TO THE MEMORYLESS q -ARY SYMMETRIC CHANNEL

Theorem 1 can be extended to the memoryless (non-binary) q -ary symmetric channel with transition probability

$$P_{Y|X}(y|x) = \begin{cases} 1 - (q-1)p, & y = x \\ p, & y \neq x \end{cases} \quad (104)$$

where $p < \frac{1}{q}$.

Theorem 2: For any sequence of non-binary codes $\{\mathcal{C}_n\}_{n=1}^\infty$ of blocklength n and code size $|\mathcal{C}_n| = M_n$, with $\mathcal{C}_n \subseteq \{0, 1, \dots, q-1\}^n$ and $q > 2$, let a_n denote the minimum probability of decoding error for transmitting \mathcal{C}_n over the memoryless q -ary symmetric channel given in (104) under a uniform distribution P_{X^n} over \mathcal{C}_n . Then

$$b_n \leq a_n \leq \left(1 + \frac{(1 - (q-1)p)}{p} (q-2)(n+1) \right) b_n \quad (105)$$

where b_n is the asymptotic generalized Poor-Verdú bound given in (28).

Theorem 2 directly implies that the asymptotic generalized Poor-Verdú bound achieves the channel reliability function of the q -ary symmetric channel, as summarized in the next corollary.

Corollary 3: (Exponential Tightness of (105) for the Non-Binary Symmetric Channel): Under the same assumptions as in Theorem 2, the asymptotic generalized Poor-Verdú bound b_n achieves the error exponent of a_n for all coding rates, i.e.,

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log \frac{a_n}{b_n} = 0. \quad (106)$$

The proof of Theorem 2 follows the same procedure as in Section IV-A except that the key step in (47) needs to be replaced with

$$\frac{\Pr \left(Y^n \in \mathcal{B}_{i,j} \mid X^n = x_{(1)}^n \right)}{\Pr \left(Y^n \in \Omega_{i,j} \mid X^n = x_{(1)}^n \right)} \leq \frac{(1 - (q-1)p)}{p} (q-2)(n+1) \quad \text{for } q > 2. \quad (107)$$

To see this, we consider $q = 3$ first in the following, where $\mathcal{X} = \mathcal{Y} = \{0, 1, 2\}$, and take $i = 1$ for simplicity.

⁵Indeed, the proof steps for $5 \leq j \leq M_n$ are identical to the ones starting from right after (90) and ending at (99) after replacing $\ell_4^{(m)}$ by $\ell_j^{(m)}$ with m ranging from 1 to 2^{j-2} (instead of 4), and replacing the sets \mathcal{S}_4 , $\mathcal{S}_4^{(m)}$, $\mathcal{B}_{1,4}^{(m)}(k)$, $\mathcal{U}_4^{(m)}(k)$ and $\Omega_{1,4}^{(m)}(k)$ by \mathcal{S}_j , $\mathcal{S}_j^{(m)}$, $\mathcal{B}_{1,j}^{(m)}(k)$, $\mathcal{U}_j^{(m)}(k)$ and $\Omega_{1,j}^{(m)}(k)$, respectively.

1) Examination of (107) for $j = 2$ with $\mathcal{B}_{1,j}$ non-empty: We define for $0 \leq k \leq |\mathcal{S}_2|$,

$$\mathcal{B}_{1,2}(k) \triangleq \left\{ y^n \in \mathcal{B}_{1,2} : d(x_{(1)}^n, y^n | \mathcal{S}_2) = k \right\}, \quad (108)$$

and

$$\Omega_{1,2}(k) \triangleq \left\{ y^n \in \Omega_{1,2} : d(x_{(1)}^n, y^n | \mathcal{S}_2) = k + 1 \right\}. \quad (109)$$

We then observe that *i)* $\mathcal{B}_{1,2} = \bigcup_{k=0}^{|\mathcal{S}_2|} \mathcal{B}_{1,2}(k)$, and $\{\mathcal{B}_{1,2}(k)\}_{k=0}^{|\mathcal{S}_2|}$ are mutually disjoint, and *ii)* $\{\Omega_{1,2}(k)\}_{k=0}^{|\mathcal{S}_2|}$ are disjoint subsets of $\Omega_{1,2}$.

As for (55), we further divide the non-empty $\mathcal{B}_{1,2}(k)$ sets into disjoint subsets indexed via selective elements, based on which a set of corresponding disjoint subsets of $\Omega_{1,2}(k)$ can be obtained. Note that different from $q = 2$, $\mathcal{B}_{1,2}$ can be non-empty even if $|\mathcal{S}_2|$ is odd. We then note by definition of $\mathcal{B}_{1,2}(k)$ that for $y^n \in \mathcal{B}_{1,2}(k)$, $d(x_{(1)}^n, y^n | \mathcal{S}_2) = d(x_{(2)}^n, y^n | \mathcal{S}_2) = k$ and hence unlike (54), where $k < |\mathcal{S}_2| = 2\ell_2$ must hold under the BSC, it is possible that $d(x_{(1)}^n, y^n | \mathcal{S}_2) + d(x_{(2)}^n, y^n | \mathcal{S}_2) = 2k = 2|\mathcal{S}_2|$ when the ternary symmetric channel is considered. We thus have to distinguish between two cases:

- $0 \leq k < |\mathcal{S}_2|$: Define for some $u^n \in \mathcal{B}_{1,2}(k)$,

$$\mathcal{B}_{1,2}(u^n | k) \triangleq \left\{ y^n \in \mathcal{B}_{1,2}(k) : d(u^n, y^n | \mathcal{S}_2^c) = 0 \text{ and } d(x_{(2)}^n, y^n | \mathcal{S}_2) = d(x_{(2)}^n, u^n | \mathcal{S}_2) \right\}. \quad (110)$$

We note that *i)* $\mathcal{B}_{1,2}(u^n | k)$ contains $\binom{|\mathcal{S}_2|}{2k-|\mathcal{S}_2|} \binom{|\mathcal{S}_2|-(2k-|\mathcal{S}_2|)}{|\mathcal{S}_2|-k}$ elements, and *ii)* $\mathcal{B}_{1,2}(k)$ can be decomposed into disjoint subsets indexed by $|\mathcal{X}|^{n-|\mathcal{S}_2|}$ specifically chosen representative u^n . Collecting these representatives in $\mathcal{U}_2(k)$, we have $\mathcal{B}_{1,2}(k) = \bigcup_{u^n \in \mathcal{U}_2(k)} \mathcal{B}_{1,2}(u^n | k)$. For example, suppose $x_{(1)}^n = 0000$ and $x_{(2)}^n = 2220$ with $n = 4$. Then, $|\mathcal{S}_2| = 3$. For $u^n = 2100 \in \mathcal{B}_{1,2}(2)$, we have

$$\mathcal{B}_{1,2}(u^n = 2100 | k = 2) = \{2100, 2010, 1200, 1020, 0120, 0210\}. \quad (111)$$

It can be verified that $\mathcal{B}_{1,2}(2)$ can be partitioned by using 2100, 2101 and 2102 as representatives in (110).

On the other hand, define for all $u^n \in \mathcal{U}_2(k)$,

$$\Omega_{1,2}(u^n | k) \triangleq \left\{ y^n \in \Omega_{1,2}(k) : d(u^n, y^n | \mathcal{S}_2^c) = 0 \text{ and } d(x_{(2)}^n, y^n | \mathcal{S}_2) = d(x_{(2)}^n, u^n | \mathcal{S}_2) - 1 = k - 1 \right\}. \quad (112)$$

Turning back to the previous example, we can see that

$$\Omega_{1,2}(u^n = 2100 | k = 2) = \{2120, 2210, 1220\}. \quad (113)$$

We note that *i)* $|\Omega_{1,2}(u^n | k)| = \binom{|\mathcal{S}_2|}{2k-|\mathcal{S}_2|} \binom{|\mathcal{S}_2|-(2k-|\mathcal{S}_2|)}{|\mathcal{S}_2|-k+1}$, and *ii)* $\{\Omega_{1,2}(u^n | k)\}_{u^n \in \mathcal{U}_2(k)}$ are disjoint subsets of $\Omega_{1,2}(k)$. As a result,

$$\frac{\Pr \left(Y^n \in \mathcal{B}_{1,2}(u^n | k) \mid X^n = x_{(1)}^n \right)}{\Pr \left(Y^n \in \Omega_{1,2}(u^n | k) \mid X^n = x_{(1)}^n \right)} \leq \frac{(1 - (q-1)p) \binom{|\mathcal{S}_2|}{2k-|\mathcal{S}_2|} \binom{|\mathcal{S}_2|-(2k-|\mathcal{S}_2|)}{|\mathcal{S}_2|-k}}{p \binom{|\mathcal{S}_2|}{2k-|\mathcal{S}_2|} \binom{|\mathcal{S}_2|-(2k-|\mathcal{S}_2|)}{|\mathcal{S}_2|-k+1}} \quad (114)$$

$$= \frac{(1 - (q-1)p)(|\mathcal{S}_2| - k + 1)}{p(|\mathcal{S}_2| - k)} \quad (115)$$

$$= \frac{(1 - (q-1)p)}{p} 2 \quad (116)$$

$$\leq \frac{(1 - (q-1)p)}{p} (n+1), \quad (117)$$

where we upper-bound 2 by $n+1$ in the last inequality in order to be consistent with what is shown in (139) and (145).

- $k = |\mathcal{S}_2|$: Define $\mathcal{B}_{1,2}(u^n | k = |\mathcal{S}_2|)$ by the same manner as in (110). In this case, it contains only $\binom{|\mathcal{S}_2|}{0} = 1$ element. We can still decompose $\mathcal{B}_{1,2}(k = |\mathcal{S}_2|)$ into disjoint subsets indexed by $|\mathcal{X}|^{n-|\mathcal{S}_2|}$ representative elements u^n . Collecting these representatives in $\mathcal{U}_2(k = |\mathcal{S}_2|)$, we have

$$\mathcal{B}_{1,2}(k = |\mathcal{S}_2|) = \bigcup_{u^n \in \mathcal{U}_2(k=|\mathcal{S}_2|)} \mathcal{B}_{1,2}(u^n | k = |\mathcal{S}_2|). \quad (118)$$

On the other hand, for any element $u^n \in \mathcal{U}_2(k = |\mathcal{S}_2|)$, we have $\binom{|\mathcal{S}_2|}{1}$ elements of y^n satisfying

$$d(u^n, y^n | \mathcal{S}_2^c) = 0 \text{ and } d(x_{(2)}^n, y^n | \mathcal{S}_2) = d(x_{(1)}^n, y^n | \mathcal{S}_2) - 1 = |\mathcal{S}_2| - 1, \quad (119)$$

where we particularly note that the conditions in (119) are different from those in (112) and thus the set of these y^n satisfying (119) is denoted differently as $\tilde{\Omega}_{1,2}(u^n | k = |\mathcal{S}_2|)$. By definition, we have $\tilde{\Omega}_{1,2}(u^n | k = |\mathcal{S}_2|) \subseteq \Omega_{1,2}(|\mathcal{S}_2| - 1)$. This can be demonstrated via the previous example. We obtain

$$\mathcal{B}_{1,2}(u^n = 1110 | k = |\mathcal{S}_2| = 3) = \{1110\}, \quad (120)$$

and

$$\tilde{\Omega}_{1,2}(u^n = 1110 | k = |\mathcal{S}_2| = 3) = \{1120, 1210, 2110\} \subset \Omega_{1,2}(2). \quad (121)$$

We thus have that

$$\frac{\Pr(Y^n \in \mathcal{B}_{1,2}(u^n | k) | X^n = x_{(1)}^n)}{\Pr(Y^n \in \tilde{\Omega}_{1,2}(u^n | k) | X^n = x_{(1)}^n)} \leq \frac{(1 - (q-1)p)}{p} \frac{1}{\binom{|\mathcal{S}_2|}{1}} \quad (122)$$

$$= \frac{(1 - (q-1)p)}{p} \frac{1}{|\mathcal{S}_2|} \quad (123)$$

$$\leq \frac{(1 - (q-1)p)}{p} (n+1), \quad (124)$$

where we again upper-bound $\frac{1}{|\mathcal{S}_2|}$ by $n+1$ in the last inequality in order to be consistent with what is derived in (139) and (145).

2) *Examination of (107) for $j = 3$ with $\mathcal{B}_{1,j}$ non-empty:* We partition \mathcal{S}_3 into two (disjoint) sets:

$$\mathcal{S}_3^{(1)} \triangleq \mathcal{S}_2 \cap \mathcal{S}_3 \quad \text{and} \quad \mathcal{S}_3^{(2)} \triangleq \mathcal{S}_2^c \cap \mathcal{S}_3. \quad (125)$$

Let $\ell_3^{(1)} \triangleq |\mathcal{S}_3^{(1)}|$ and $\ell_3^{(2)} \triangleq |\mathcal{S}_3^{(2)}|$.

Define for $0 \leq k^{(1)} \leq \ell_3^{(1)}$,

$$\mathcal{B}_{1,3}^{(1)}(k^{(1)}) \triangleq \left\{ y^n \in \mathcal{B}_{1,3} : d(x_{(1)}^n, y^n | \mathcal{S}_3^{(1)}) = k^{(1)} \right\}, \quad (126)$$

and for $0 \leq k^{(2)} \leq \ell_3^{(2)}$,

$$\mathcal{B}_{1,3}^{(2)}(k^{(2)}) \triangleq \left\{ y^n \in \mathcal{B}_{1,3} : d(x_{(1)}^n, y^n | \mathcal{S}_3^{(1)}) = \ell_3^{(1)} \text{ and } d(x_{(1)}^n, y^n | \mathcal{S}_3^{(2)}) = k^{(2)} \right\}. \quad (127)$$

Then, the definitions above yield that

- i) $\mathcal{B}_{1,3}^{(1)}(\ell_3^{(1)}) = \bigcup_{k^{(1)}=0}^{\ell_3^{(2)}} \mathcal{B}_{1,3}^{(2)}(k^{(1)})$;
- ii) $\mathcal{B}_{1,3} = \bigcup_{k^{(1)}=0}^{\ell_3^{(1)}} \mathcal{B}_{1,3}^{(1)}(k^{(1)}) = \left(\bigcup_{k^{(1)}=0}^{\ell_3^{(1)}-1} \mathcal{B}_{1,3}^{(1)}(k^{(1)}) \right) \cup \left(\bigcup_{k^{(2)}=0}^{\ell_3^{(2)}} \mathcal{B}_{1,3}^{(2)}(k^{(2)}) \right)$;
- iii) $\{\mathcal{B}_{1,3}^{(1)}(k^{(1)})\}_{k^{(1)}=0}^{\ell_3^{(1)}-1}$ and $\{\mathcal{B}_{1,3}^{(2)}(k^{(2)})\}_{k^{(2)}=0}^{\ell_3^{(2)}}$ are mutually disjoint.

For $0 \leq k^{(1)} < \ell_3^{(1)}$, we further partition $\mathcal{B}_{1,3}^{(1)}(k^{(1)})$ into disjoint subsets, each of which is indexed by a specifically chosen $u^n \in \mathcal{B}_{1,3}^{(1)}(k^{(1)})$ and is defined by

$$\mathcal{B}_{1,3}^{(1)}(u^n | k^{(1)}) \triangleq \left\{ y^n \in \mathcal{B}_{1,3}^{(1)}(k^{(1)}) : d(u^n, y^n | (\mathcal{S}_3^{(1)})^c) = 0 \text{ and } d(x_{(3)}^n, y^n | \mathcal{S}_3^{(1)}) = d(x_{(3)}^n, u^n | \mathcal{S}_3^{(1)}) \right\}. \quad (128)$$

Denote the set of these representative u^n by $\mathcal{U}_3^{(1)}(k^{(1)})$. Hence,

$$\mathcal{B}_{1,3}^{(1)}(k^{(1)}) = \bigcup_{u^n \in \mathcal{U}_3^{(1)}(k^{(1)})} \mathcal{B}_{1,3}^{(1)}(u^n | k^{(1)}).$$

We can similarly partition $\mathcal{B}_{1,3}^{(2)}(k^{(2)})$ into disjoint $\{\mathcal{B}_{1,3}^{(2)}(u^n | k^{(2)})\}_{u^n \in \mathcal{U}_3^{(2)}(k^{(2)})}$ with

$$\mathcal{B}_{1,3}^{(2)}(u^n | k^{(2)}) \triangleq \left\{ y^n \in \mathcal{B}_{1,3}^{(2)}(k^{(2)}) : d(u^n, y^n | (\mathcal{S}_3^{(2)})^c) = 0 \text{ and } d(x_{(3)}^n, y^n | \mathcal{S}_3^{(2)}) = d(x_{(3)}^n, u^n | \mathcal{S}_3^{(2)}) \right\}, \quad (129)$$

which implies $\mathcal{B}_{1,3}^{(2)}(k^{(2)}) = \bigcup_{u^n \in \mathcal{U}_3^{(2)}(k^{(2)})} \mathcal{B}_{1,3}^{(2)}(u^n | k^{(2)})$.

On the other hand, we can likewise define for $0 \leq k^{(1)} < \ell_3^{(1)}$,

$$\Omega_{1,3}^{(1)}(k^{(1)}) \triangleq \left\{ y^n \in \Omega_{1,3} : d(x_{(1)}^n, y^n | \mathcal{S}_3^{(1)}) = k^{(1)} + 1 \right\}, \quad (130)$$

and for $0 \leq k^{(2)} \leq \ell_3^{(2)}$,

$$\Omega_{1,3}^{(2)}(k^{(2)}) \triangleq \left\{ y^n \in \Omega_{1,3} : d(x_{(1)}^n, y^n | \mathcal{S}_3^{(1)}) = \ell_3^{(1)} \text{ and } d(x_{(1)}^n, y^n | \mathcal{S}_3^{(2)}) = k^{(2)} + 1 \right\}, \quad (131)$$

and for $0 \leq k^{(1)} < \ell_3^{(1)}$ and $u^n \in \mathcal{U}_3^{(1)}(k^{(1)})$,

$$\Omega_{1,3}^{(1)}(u^n | k^{(1)}) \triangleq \left\{ y^n \in \Omega_{1,3}^{(1)}(k^{(1)}) : d(u^n, y^n | (\mathcal{S}_3^{(1)})^c) = 0 \text{ and } d(x_{(3)}^n, y^n | \mathcal{S}_3^{(1)}) = k^{(1)} - 1 \right\}, \quad (132)$$

and for $0 \leq k^{(2)} \leq \ell_3^{(2)}$ and $u^n \in \mathcal{U}_3^{(2)}(k^{(2)})$,

$$\Omega_{1,3}^{(2)}(u^n | k^{(2)}) \triangleq \left\{ y^n \in \Omega_{1,3}^{(2)}(k^{(2)}) : d(u^n, y^n | (\mathcal{S}_3^{(2)})^c) = 0 \text{ and } d(x_{(3)}^n, y^n | \mathcal{S}_3^{(2)}) = k^{(2)} - 1 \right\}. \quad (133)$$

By the definitions above, we have $\{\Omega_{1,3}^{(1)}(u^n | k^{(1)})\}_{0 \leq k^{(1)} < \ell_3^{(1)}, u^n \in \mathcal{U}_3^{(1)}(k^{(1)})}$ and $\{\Omega_{1,3}^{(2)}(u^n | k^{(2)})\}_{0 \leq k^{(2)} < \ell_3^{(2)}, u^n \in \mathcal{U}_3^{(2)}(k^{(2)})}$ are all mutually disjoint.

With the above notation, we are ready to give the derivation of the desired result. For $0 \leq k^{(1)} < \ell_3^{(1)}$ and

$u^n \in \mathcal{U}_3^{(1)}(k^{(1)})$ with $d(x_{(3)}^n, u^n | \mathcal{S}_3^{(1)}) = m$, we establish that

$$|\mathcal{B}_{1,3}^{(1)}(u^n | k^{(1)})| = \binom{\ell_3^{(1)}}{k^{(1)}+m-\ell_3^{(1)}} \binom{2\ell_3^{(1)}-k^{(1)}-m}{\ell_3^{(1)}-m}, \quad (134)$$

$$|\Omega_{1,3}^{(1)}(u^n | k^{(1)})| = \binom{\ell_3^{(1)}}{k^{(1)}+m-\ell_3^{(1)}} \binom{2\ell_3^{(1)}-k^{(1)}-m}{\ell_3^{(1)}-m+1}, \quad (135)$$

and,

$$\frac{\Pr\left(Y^n \in \mathcal{B}_{1,3}^{(1)}(u^n | k^{(1)}) \mid X^n = x_{(1)}^n\right)}{\Pr\left(Y^n \in \Omega_{1,3}^{(1)}(u^n | k^{(1)}) \mid X^n = x_{(1)}^n\right)} = \frac{(1 - (q-1)p) \binom{2\ell_3^{(1)}-k^{(1)}-m}{\ell_3^{(1)}-m}}{p \binom{2\ell_3^{(1)}-k^{(1)}-m}{\ell_3^{(1)}-m+1}} \quad (136)$$

$$= \frac{(1 - (q-1)p) (\ell_3^{(1)} - m + 1)}{p (\ell_3^{(1)} - k^{(1)})} \quad (137)$$

$$\leq \frac{(1 - (q-1)p)}{p} (\ell_3^{(1)} + 1) \quad (138)$$

$$\leq \frac{(1 - (q-1)p)}{p} (n + 1), \quad (139)$$

where (138) follows from $\ell_3^{(1)} - k^{(1)} \geq 1$ and $m \geq 0$.

We proceed to take care of $k^{(2)}$ by considering separately the following two cases.

- $0 \leq k^{(2)} < \ell_3^{(2)}$: Similarly, we establish that for $u^n \in \mathcal{U}_3^{(2)}(k^{(2)})$ with $d(x_{(3)}^n, u^n | \mathcal{S}_3^{(2)}) = m$,

$$|\mathcal{B}_{1,3}^{(2)}(u^n | k^{(2)})| = \binom{\ell_3^{(2)}}{k^{(2)}+m-\ell_3^{(2)}} \binom{2\ell_3^{(2)}-k^{(2)}-m}{\ell_3^{(2)}-m}, \quad (140)$$

$$|\Omega_{1,3}^{(2)}(u^n | k^{(2)})| = \binom{\ell_3^{(2)}}{k^{(2)}+m-\ell_3^{(2)}} \binom{2\ell_3^{(2)}-k^{(2)}-m}{\ell_3^{(2)}-m+1}, \quad (141)$$

and

$$\frac{\Pr\left(Y^n \in \mathcal{B}_{1,3}^{(2)}(u^n | k^{(2)}) \mid X^n = x_{(1)}^n\right)}{\Pr\left(Y^n \in \Omega_{1,3}^{(2)}(u^n | k^{(2)}) \mid X^n = x_{(1)}^n\right)} = \frac{(1 - (q-1)p) \binom{2\ell_3^{(2)}-k^{(2)}-m}{\ell_3^{(2)}-m}}{p \binom{2\ell_3^{(2)}-k^{(2)}-m}{\ell_3^{(2)}-m+1}} \quad (142)$$

$$= \frac{(1 - (q-1)p) (\ell_3^{(2)} - m + 1)}{p (\ell_3^{(2)} - k^{(2)})} \quad (143)$$

$$\leq \frac{(1 - (q-1)p)}{p} (\ell_3^{(2)} + 1) \quad (144)$$

$$\leq \frac{(1 - (q-1)p)}{p} (n + 1), \quad (145)$$

where (144) follows from $\ell_3^{(2)} - k^{(2)} \geq 1$ and $m \geq 0$.

- $k^{(2)} = \ell_3^{(2)}$: Let $u^n \in \mathcal{B}_{1,3}^{(2)}(\ell_3^{(2)})$. Then $d(x_{(1)}^n, u^n | \mathcal{S}_3) = d(x_{(3)}^n, u^n | \mathcal{S}_3) = \ell_3^{(1)} + \ell_3^{(2)} = |\mathcal{S}_3|$. We note that

$$|\mathcal{B}_{1,3}^{(2)}(u^n | \ell_3^{(2)})| = \binom{|\mathcal{S}_3|}{0} = 1, \quad (146)$$

and $\mathcal{B}_{1,3}^{(2)}(\ell_3^{(2)})$ can be decomposed into distinct subsets $\mathcal{B}_{1,3}^{(2)}(u^n | \ell_3^{(2)})$ indexed by specifically chosen $|\mathcal{X}|^{n-|\mathcal{S}_3|}$ representative u^n .

On the other hand, for $u^n \in \mathcal{U}_3^{(2)}(k^{(2)})$, the number of y^n satisfying

$$d(u^n, y^n | \mathcal{S}_3^c) = 0 \text{ and } d(x_{(3)}^n, y^n | \mathcal{S}_3) = d(x_{(1)}^n, y^n | \mathcal{S}_3) - 1 = |\mathcal{S}_3| - 1 \quad (147)$$

is $\binom{|\mathcal{S}_3|}{1}$. Collect the elements satisfying (147) in $\tilde{\Omega}_{1,2}^{(2)}(u^n | \ell_3^{(2)})$. Note that $\tilde{\Omega}_{1,2}^{(2)}(u^n | \ell_3^{(2)}) \subset \Omega_{1,2}^{(2)}(\ell_3^{(2)} - 1)$, and

$$\frac{\Pr\left(Y^n \in \mathcal{B}_{1,3}^{(2)}(u^n | \ell_3^{(2)}) \mid X^n = x_{(1)}^n\right)}{\Pr\left(Y^n \in \tilde{\Omega}_{1,3}^{(2)}(u^n | \ell_3^{(2)}) \mid X^n = x_{(1)}^n\right)} \leq \frac{(1 - (q-1)p)}{p} \frac{1}{\binom{|\mathcal{S}_3|}{1}} \quad (148)$$

$$= \frac{(1 - (q-1)p)}{p} \frac{1}{|\mathcal{S}_3|} \quad (149)$$

$$\leq \frac{(1 - (q-1)p)}{p} (n+1). \quad (150)$$

A similar derivation can be applied for $j = 4, \dots, M_n$. The proof for $q = 3$ is thus completed.

The proof for $q > 3$ is analogous to the derivation of $q = 3$. Some exceptions are listed below:

1) In the case of $j = 2$:

- The number of elements in $\mathcal{B}_{1,2}(u^n | k)$, as defined in (110), is

$$(q-2)^{2k-|\mathcal{S}_2|} \binom{|\mathcal{S}_2|}{2k-|\mathcal{S}_2|} \binom{|\mathcal{S}_2| - (2k - |\mathcal{S}_2|)}{|\mathcal{S}_2| - k} \quad (151)$$

instead of $\binom{|\mathcal{S}_2|}{2k-|\mathcal{S}_2|} \binom{|\mathcal{S}_2| - (2k - |\mathcal{S}_2|)}{|\mathcal{S}_2| - k}$.⁶

- The number of elements in $\Omega_{1,2}(u^n | k)$, as defined in (112), is

$$(q-2)^{2k-|\mathcal{S}_2|} \binom{|\mathcal{S}_2| - (2k - |\mathcal{S}_2|)}{|\mathcal{S}_2 - k + 1|} \quad (152)$$

instead of $\binom{|\mathcal{S}_2|}{2k-|\mathcal{S}_2|} \binom{|\mathcal{S}_2| - (2k - |\mathcal{S}_2|)}{|\mathcal{S}_2 - k + 1|}$.

- The number of elements in $\mathcal{B}_{1,2}(u^n | k = |\mathcal{S}_2|)$, as defined in (110), is $(q-2)^{|\mathcal{S}_2|}$ instead of 1 .⁷
- The number of elements in $\tilde{\Omega}_{1,2}(u^n | k = |\mathcal{S}_2|)$, which satisfies (119), is $(q-2)^{|\mathcal{S}_2|-1} \binom{|\mathcal{S}_2|}{1}$ instead of $\binom{|\mathcal{S}_2|}{1}$.
- The upper bound in (122) is instead

$$\frac{\Pr\left(Y^n \in \mathcal{B}_{1,2}(u^n | k) \mid X^n = x_{(1)}^n\right)}{\Pr\left(Y^n \in \tilde{\Omega}_{1,2}(u^n | k) \mid X^n = x_{(1)}^n\right)} \leq \frac{(1 - (q-1)p)}{p} \frac{(q-2)^{|\mathcal{S}_2|}}{(q-2)^{|\mathcal{S}_2|-1} \binom{|\mathcal{S}_2|}{1}} \quad (153)$$

$$\leq \frac{(1 - (q-1)p)}{p} (q-2)(n+1). \quad (154)$$

2) In the case of $j = 3$:

- The number of elements in $\mathcal{B}_{1,2}(u^n | k)$, as defined in (134), is

$$(q-2)^{k^{(1)}+m-\ell_3^{(1)}} \binom{\ell_3^{(1)}}{k^{(1)}+m-\ell_3^{(1)}} \binom{2\ell_3^{(1)}-k^{(1)}-m}{\ell_3^{(1)}-m} \quad (155)$$

instead of $\binom{\ell_3^{(1)}}{k^{(1)}+m-\ell_3^{(1)}} \binom{2\ell_3^{(1)}-k^{(1)}-m}{\ell_3^{(1)}-m}$.

⁶Note that for the BSC, where $q = 2$, we only consider $k = \ell_2$ and $|\mathcal{S}_2| = 2\ell_2$, for which the formula $(q-2)^{2k-|\mathcal{S}_2|}$ is underdetermined and cannot be applied. Thus, $\mathcal{B}_{1,2}(u^n | k)$ and $\Omega_{1,2}(u^n | k)$ when $q = 2$ are treated separately.

⁷For the BSC, where $q = 2$, $\mathcal{B}_{1,2}(u^n | k = |\mathcal{S}_2|)$ is an empty set. Thus, it is unnecessary to define or consider $\tilde{\Omega}_{1,2}(u^n | k = |\mathcal{S}_2|)$ when $q = 2$.

- The number of elements in $\Omega_{1,2}(u^n|k)$, as defined in (135), is

$$(q-2)^{k^{(1)}+m-\ell_3^{(1)}} \binom{\ell_3^{(1)}}{k^{(1)}+m-\ell_3^{(1)}} \binom{2\ell_3^{(1)}-k^{(1)}-m}{\ell_3^{(1)}-m+1} \quad (156)$$

instead of $\binom{\ell_3^{(1)}}{k^{(1)}+m-\ell_3^{(1)}} \binom{2\ell_3^{(1)}-k^{(1)}-m}{\ell_3^{(1)}-m+1}$.

- The number of elements in $\mathcal{B}_{1,3}^{(2)}(u^n|\ell_3^{(2)})$, as defined in (146), is $(q-2)^{|\mathcal{S}_3|}$ instead of 1.
- The number of elements in $\tilde{\Omega}_{1,3}^{(2)}(u^n|\ell_3^{(2)})$, which satisfies (147), is $(q-2)^{|\mathcal{S}_3|-1} \binom{|\mathcal{S}_3|}{1}$ instead of $\binom{|\mathcal{S}_3|}{1}$.
- The upper bound in (148) is instead

$$\frac{\Pr\left(Y^n \in \mathcal{B}_{1,3}^{(2)}(u^n|\ell_3^{(2)}) \mid X^n = x_{(1)}^n\right)}{\Pr\left(Y^n \in \tilde{\Omega}_{1,3}^{(2)}(u^n|\ell_3^{(2)}) \mid X^n = x_{(1)}^n\right)} \leq \frac{(1-(q-1)p)}{p} \frac{(q-2)^{|\mathcal{S}_3|}}{(q-2)^{|\mathcal{S}_3|-1} \binom{|\mathcal{S}_3|}{1}} \quad (157)$$

$$\leq \frac{(1-(q-1)p)}{p} (q-2)(n+1). \quad (158)$$

■

VI. CONCLUSION

We derived a closed-form formula for the asymptotic generalized Poor-Verdú error bound to the multihypothesis testing error probability and proved that, unlike the case for the BEC [24], it achieves the error exponent of the BSC and the non-binary symmetric channel for arbitrary sequences of codes. This indicates that we can characterize the channel reliability function at any rate, without having to explicitly calculate it, via the asymptotic generalized Poor-Verdú bound. We established the exponent tightness results as a direct consequence of a key inequality, showing that for any block code used over these channels, the relative deviation of the code's minimum probability of error from the asymptotic generalized lower bound grows at most linearly in blocklength. Even though the exact determination of the reliability function of the BSC (and the non-binary symmetric channel) at low rates remains a daunting open problem, our results offer a new perspective or tool for investigating it via the asymptotic generalized Poor-Verdú bound. Future work includes studying the relationship between the generalized bound and the meta-converse channel coding bound [15], [27] and the further examination of tight bounds for codes with small-to-moderate blocklength (e.g., see [26]–[28]) used over channels with and without memory.

REFERENCES

- [1] L.-H. Chang, P.-N. Chen, F. Alajaji, and Y. S. Han, "The asymptotic generalized poor-verdu bound achieves the BSC error exponent at zero rate," in *IEEE Int. Symp. Inf. Theory (ISIT)*, Los Angeles, CA, USA, June 21–26, 2020.
- [2] H. V. Poor and S. Verdú, "A lower bound on the probability of error in multihypothesis testing," *IEEE Trans. Inf. Theory*, vol. 41, no. 6, pp. 1992–1994, November 1995.
- [3] P.-N. Chen and F. Alajaji, "A generalized Poor-Verdú error bound for multihypothesis testings," *IEEE Trans. Inf. Theory*, vol. 58, no. 1, pp. 311–316, January 2012.

- [4] R. G. Gallager, “A simple derivation of the coding theorem and some applications,” *IEEE Trans. Inf. Theory*, vol. 11, no. 1, pp. 3–18, January 1965.
- [5] C. E. Shannon, R. G. Gallager, and E. R. Berlekamp, “Lower bounds to error probability for coding on discrete memoryless channels - i,” *Inf. Contr.*, vol. 10, pp. 65–103, January 1967.
- [6] —, “Lower bounds to error probability for coding on discrete memoryless channels - ii,” *Inf. Contr.*, vol. 10, pp. 522–552, May 1967.
- [7] R. J. McEliece and J. K. Omura, “An improved upper bound on the block coding error exponent for binary-input discrete memoryless channels,” *IEEE Trans. Inf. Theory*, vol. 23, no. 5, pp. 611–613, September 1977.
- [8] R. G. Gallager, *Information Theory and Reliable Communication*. NY: Wiley, 1968.
- [9] A. J. Viterbi and J. K. Omura, *Principles of Digital Communication and Coding*. NY: McGraw-Hill, 1979.
- [10] I. Csiszár and J. Körner, *Information Theory: Coding Theorems for Discrete Memoryless Systems*. NY: Academic Press, 1981.
- [11] R. Blahut, *Principles and Practice of Information Theory*. A. Wesley, MA, 1988.
- [12] S. Litsyn, “New upper bounds on error exponents,” *IEEE Trans. Inf. Theory*, vol. 45, no. 2, pp. 385–398, March 1999.
- [13] A. Barg and A. McGregor, “Distance distribution of binary codes and the error probability of decoding,” *IEEE Trans. Inf. Theory*, vol. 51, no. 12, pp. 4237–4246, December 2005.
- [14] E. A. Haroutunian, M. E. Haroutunian, and A. N. Harutyunyan, “Reliability criteria in information theory and in statistical hypothesis testing,” *Foundations and Trends in Communications and Information Theory, Now Publishers Inc.*, vol. 4, pp. 97–263, January 2007.
- [15] Y. Polyanskiy, “Saddle point in the minimax converse for channel coding,” *IEEE Trans. Inf. Theory*, vol. 59, no. 5, pp. 2576–2595, May 2013.
- [16] M. Dalai, “Lower bounds on the probability of error for classical and classical-quantum channels,” *IEEE Trans. Inf. Theory*, vol. 59, no. 12, pp. 8027–8056, December 2013.
- [17] J. Scarlett, A. Martinez, and A. G. i Fábregas, “Mismatched decoding: Error exponents, second-order rates and saddlepoint approximations,” *IEEE Trans. Inf. Theory*, vol. 60, no. 5, pp. 2647–2666, May 2014.
- [18] J. Scarlett, I. Peng, N. Merhav, A. Martinez, and A. G. i Fábregas, “Expurgated random-coding ensembles: Exponents, refinements, and connections,” *IEEE Trans. Inf. Theory*, vol. 60, no. 8, pp. 4449–4462, August 2014.
- [19] Y. Altüg and A. B. Wagner, “Refinement of the random coding bound,” *IEEE Trans. Inf. Theory*, vol. 60, no. 10, pp. 6005–6023, October 2014.
- [20] M. V. Burnashev, “On the BSC reliability function: Expanding the region where it is known exactly,” *Probl. Inf. Trans.*, vol. 51, no. 4, pp. 307–325, January 2015.
- [21] A. Tandon, V. Y. F. Tan, and L. R. Varshney, “The bee-identification problem: Bounds on the error exponent,” *IEEE Trans. Commun.*, vol. 67, no. 11, pp. 7405–7416, November 2019.
- [22] —, “On the bee-identification error exponent with absentee bees,” *arXiv preprint arXiv:1910.10333*, October 2019.
- [23] N. Merhav, “A Lagrange-dual lower bound to the error exponent of the typical random code,” *IEEE Trans. Inf. Theory*, vol. 66, no. 6, pp. 3456–3464, June 2020.
- [24] F. Alajaji, P.-N. Chen, and Z. Rached, “A note on the Poor-Verdú conjecture for the channel reliability function,” *IEEE Trans. Inf. Theory*, vol. 48, no. 1, pp. 309–313, January 2002.
- [25] S. Verdú and T. S. Han, “A general formula for channel capacity,” *IEEE Trans. Inf. Theory*, vol. 40, no. 4, pp. 1147–1157, July 1994.
- [26] G. Vazquez-Vilar, A. T. Campo, A. G. i Fábregas, and A. Martinez, “Bayesian M -ary hypothesis testing: The meta-converse and Verdú-Han bounds are tight,” *IEEE Trans. Inf. Theory*, vol. 62, no. 5, pp. 2324–2333, May 2016.

- [27] Y. Polyanskiy, H. V. Poor, and S. Verdú, “Channel coding rate in the finite blocklength regime,” *IEEE Trans. Inf. Theory*, vol. 56, no. 5, pp. 2307–2359, May 2010.
- [28] P.-N. Chen, H.-Y. Lin, and S. M. Moser, “Optimal ultrasmall block-codes for binary discrete memoryless channels,” *IEEE Trans. Inf. Theory*, vol. 59, no. 11, pp. 7346–7378, November 2013.