

MANIFOLDS WITH CYLINDRICAL ENDS HAVING A FINITE AND POSITIVE NUMBER OF EMBEDDED EIGENVALUES

T. J. CHRISTIANSEN AND K. DATCHEV

ABSTRACT. We construct a surface with a cylindrical end which has a finite number of Laplace eigenvalues embedded in its continuous spectrum. The surface is obtained by attaching a cylindrical end to a hyperbolic torus with a hole. To our knowledge, this is the first example of a manifold with a cylindrical end whose number of eigenvalues is known to be finite and nonzero. The construction can be varied to give examples with arbitrary genus and with an arbitrarily large finite number of eigenvalues. The constructed surfaces also have resonance-free regions near the continuous spectrum and long-time asymptotic expansions of solutions to the wave equation.

Let (X, g) be a manifold with cylindrical ends and let $-\Delta \geq 0$ be its Laplacian. A basic question is: How many eigenvalues does $-\Delta$ have? This question is subtle because the essential spectrum of $-\Delta$ is $[0, \infty)$, and hence any eigenvalues must be embedded in it rather than isolated. In some examples, including the simplest example of the cylinder $\mathbb{R} \times (\mathbb{R}/\mathbb{Z})$, the number of eigenvalues is zero. In this note we prove that the number of eigenvalues can be nonzero but finite.

Theorem. *There exists a surface with a cylindrical end whose Laplacian has a nonzero but finite number of eigenvalues.*

In fact, we construct families of Riemannian surfaces (X, g) consisting of a compact hyperbolic piece with an infinite cylindrical end attached. More specifically, we start with a symmetric funneled torus of constant negative curvature, (X, g_H) , and then modify the metric in the funnel end to obtain a cylindrical end (a Riemannian product of a half-line with a circle) as in Figure 1.




FIGURE 1. The hyperbolic surface (X, g_H) , and the surface with a cylindrical end (X, g) .

In the proof below, we show that if the length ℓ of the geodesic at the boundary of the funnel is sufficiently small, then $-\Delta$ has an eigenvalue. For this, as in work of earlier authors including [EvLeVa, DaPa], we consider $-\Delta|_O$, a restriction of the Laplacian $-\Delta$ to a domain of functions with a suitable symmetry, and we show that $-\Delta|_O$ has an isolated eigenvalue.

The more difficult step is showing that $-\Delta$ does not have infinitely many eigenvalues. This is done in [ChDa1, Theorem 1.1] for a family of manifolds with cylindrical ends which includes (X, g) ,

and which also includes the variants of (X, g) described in the extensions after the proof below. This result is an application of two recent breakthroughs in the scattering theory of hyperbolic manifolds: Vasy's propagation of semiclassical singularities [Va], and the essential spectral gap of Dyatlov and Zahl [DyZa] and Bourgain and Dyatlov [BoDy].

Our results also imply the existence of resonance-free regions and asymptotic expansions for solutions to the wave equation $(\partial_t^2 - \Delta)u = 0$. See the discussion following the proof below for more.

Existence proofs for eigenvalues for manifolds with cylindrical ends (including the case of waveguides, or more generally manifolds with boundary, where we may have isolated eigenvalues below the essential spectrum as well as embedded ones within it) have been given by many authors under various assumptions. In some cases the number of eigenvalues is shown to be infinite [Wi, ChZw, Pa, IsKo], and in many others it is left open whether the number of embedded eigenvalues is finite or infinite: see [ExŠe, AvBeGiMa, GoJa, EvLeVa, BuGeReSi, DaPa] for some such results, [LoCaMu, KrKř, ExKo] for further results and surveys of the literature, and [CaDuNa, SaMe] for some recent results and more references. In general, the eigenvalue counting function is polynomially bounded [Do1, ChZw, Pa]. On the negative side, nonexistence results for eigenvalues have been obtained in many of the aforementioned papers, as well as in [MoWe, BrDiKr, ChDa2], and one for a resonance at the bottom of the spectrum in [GrJe]. To our knowledge, the examples in this note are the first manifolds with cylindrical ends whose total number of eigenvalues is known to be finite but not zero.

We discuss briefly other geometric situations in which finite numbers of embedded eigenvalues arise. Particularly important, and closely related to manifolds with cylindrical ends [Do1, Gu], are manifolds with cusps: see [Co], the surveys [Sa, Mü], and [Bo1, HiJu] for recent results and pointers to the substantial literature on this topic. Another direction of study is manifolds whose curvature converges to a constant at infinity. One asks: For what rates of convergence do embedded eigenvalues occur? See [Do2, JiLi] for some recent results and references, and [Ku] for an example with precisely one embedded eigenvalue.

Proof of Theorem. We construct the desired Riemannian surface (X, g) by modifying the metric of a certain hyperbolic surface (X, g_H) , which is the quotient of the hyperbolic plane \mathbb{H} by a Schottky group. We use the unit disk model $\mathbb{H} = \{z \in \mathbb{C} \mid |z| < 1\}$. For each $j \in \{1, 2, 3, 4\}$, let D_j be the open disk in \mathbb{C} , orthogonal to the unit disk, which is inscribed in the sector $\{z \in \mathbb{C} \mid |\arg z - (2j - 1)\pi/4| < \alpha\}$, where $\alpha < \pi/4$ will be chosen later (close to $\pi/4$). See Figure 2.

Let S_1 be the orientation-preserving isometry of \mathbb{H} which fixes $e^{\pi i/4}$ and $-e^{\pi i/4}$ and which maps D_1 to the exterior of D_3 : see [Bo2, Section 2.1] and [MuSeWr, Project 4.2]. Let

$$S_2(z) = \overline{S_1^{-1}(\bar{z})}. \quad (1)$$

Then S_2 maps D_2 to the exterior of D_4 .

Let Γ be the group generated by S_1 and S_2 , and let (X, g_H) be the quotient of \mathbb{H} by Γ . We recall some standard facts about (X, g_H) from [Bo2, Sections 2.4 and 15.1]. The surface (X, g_H)




FIGURE 2. On the left, the unit disk with the parts of the disks D_1 , D_2 , D_3 , and D_4 used to define the fundamental domain \mathcal{F} . On the right, the fundamental domain \mathcal{F} , with points having matching letters identified.

is obtained from the fundamental domain

$$\mathcal{F} := \mathbb{H} - \bigcup_{j=1}^4 D_j$$

by identifying ∂D_1 with ∂D_3 using S_1 and ∂D_2 with ∂D_4 using S_2 , so that points having matching letters in Figure 2 are identified. Let N be the convex core of X ; this is the part of X whose lift to \mathcal{F} is given by $\mathcal{F} - \bigcup_{j=1}^4 D'_j$, where each D'_j is the open disk in \mathbb{C} , orthogonal to the unit disk, to D_j , and to D_{j+1} (with the convention that $D_5 = D_1$), whose center has argument $j\pi/2$. Then N has genus 1 and ∂N is a geodesic of length $\ell = 4\rho(D_1, D_2)$, where ρ is the hyperbolic distance function. Furthermore, $X \setminus N$ is a hyperbolic funnel, i.e. there are coordinates $(r, y) \in (0, \infty) \times (\mathbb{R}/\ell\mathbb{Z})$ such that

$$X \setminus N = (0, \infty)_r \times (\mathbb{R}/\ell\mathbb{Z})_y, \quad g_H|_{X \setminus N} = dr^2 + \cosh^2 r dy^2. \quad (2)$$

By (1), the involution $z \mapsto \bar{z}$ on \mathcal{F} descends to an involution J on X , and $J|_{X \setminus N}$ is a reflection. For later convenience, we normalize the coordinate y in (2) so that $J(r, y) = (r, -y)$. For example, we may take $y = 0$ when $\arg z = 0$, $y = \ell/8$ at a , $y = 3\ell/8$ at b , $y = \pm\ell/2$ when $\arg z = \pm\pi$, $y = -3\ell/8$ at c , $y = -\ell/8$ at d .

Let $B = \{z \in \mathbb{C} \mid |z| < \sqrt{2} - 1\}$, and fix $\varphi \in C_c^\infty(B)$ such that $\varphi \not\equiv 0$ and $\varphi(\bar{z}) = -\varphi(z)$ for all z . Since $B \subset \mathcal{F}$, regardless of α , we may identify φ with a function on X , and $\varphi \circ J = -\varphi$. Moreover, this φ is supported in N , regardless of α .

Now choose $\alpha < \pi/4$ (sufficiently close to $\pi/4$) such that $\ell = \ell(\alpha)$ obeys

$$\frac{\|d\varphi\|_{L^2(X, g_H)}^2}{\|\varphi\|_{L^2(X, g_H)}^2} < \frac{1}{\ell^2}. \quad (3)$$

More precisely, $\tan(\alpha) \sinh \rho(0, D_1) = 1$ by the angle of parallelism formula [Be, Theorem 7.9.1], and $\cosh(\ell/4) = \sinh^2 \rho(0, D_1)$ by a property of right-angled pentagons [Be, (7.18.2)]. See Figure 3. Hence, $\tan^2(\alpha) \cosh(\ell/4) = 1$, and (3) is equivalent to $\tan^2(\alpha) \cosh(\|\varphi\|/4 \|d\varphi\|) > 1$.




FIGURE 3. The triangle and pentagon used to obtain (3).

Next, modify the metric in the funnel end so as to change it into a cylindrical end in the following way. Fix a metric g on X having the form

$$g|_N = g_H|_N, \quad g|_{X \setminus N} = dr^2 + F(r)dy^2,$$

where $F \in C^\infty((0, \infty))$ obeys $F(r) = \cosh^2 r$ near $r = 0$, $F'(r) > 0$ when $r \in (0, R)$ for some $R > 0$, and $F(r) = (2\pi)^2$ for $r \geq R$. Note that g , like g_H , is invariant under J .

Let $-\Delta|_O$ be the Laplacian on (X, g) , restricted to functions u obeying $u \circ J = -u$. Observe that we have, in the part of $X \setminus N$ where $r > R$, the Fourier series expansion

$$-\Delta u(r, y) = (-\partial_r^2 - \partial_y^2 / (2\pi)^2) \sum_{n=1}^{\infty} u_n(r) \sin(2\pi n y / \ell) \geq (-\partial_r^2 + 1/\ell^2) u(r, y),$$

for all u obeying $u \circ J = -u$, with equality if $u_n = 0$ for $n \geq 2$. This implies that the essential spectrum of $-\Delta|_O$ is $[1/\ell^2, \infty)$. But, by (3), φ is not in the range of the spectral projection of $-\Delta|_O$ onto $[1/\ell^2, \infty)$. Hence $-\Delta|_O$ has at least one eigenvalue, and in particular $-\Delta$ has at least one eigenvalue.

By [ChDa1, Theorem 1.1], $-\Delta$ does not have infinitely many eigenvalues. \square

Extensions.

- (i) To construct examples with at least m eigenvalues, replace (3) with

$$\max_{\varphi \in V} \frac{\|d\varphi\|_{L^2(X, g_H)}^2}{\|\varphi\|_{L^2(X, g_H)}^2} < \frac{1}{\ell^2},$$

where V is an m -dimensional space of functions in $C_c^\infty(B)$ satisfying $\varphi(\bar{z}) = -\varphi(z)$.

- (ii) The symmetry with respect to the vertical axis (but not that with respect to the horizontal axis) can be broken: after choosing α such that (3) holds, replace D_2 and D_3 by disks, orthogonal to the unit disk, inscribed in larger sectors $\{z \in \mathbb{C} \mid |\arg z - (2j-1)\pi/4| < \alpha'\}$ for some $\alpha < \alpha' < \pi/4$.
- (iii) To construct examples having genus g for any $g \geq 1$, start with disks D_1, \dots, D_{4g} , orthogonal to the unit disk, inscribed in sectors $\{z \in \mathbb{C} \mid |\arg z - (2j-1)\pi/4| < \alpha\}$, with $\alpha < \pi/4g$, and take S_j mapping D_j to the exterior of D_{j+2g} . Then the convex core of the quotient has genus g , and the boundary is a geodesic of length $\ell = 4g\rho(D_1, D_2)$, and we can choose $\alpha < \pi/4g$, close to $\pi/4g$, such that (3) holds and proceed as before.

- (iv) In all of these examples, we can also rule out resonances close to the continuous spectrum. We discuss this briefly here, referring the reader to [ChDa1, Theorem 5.6] for details. By [ChDa1, Theorem 1.1], the cut-off resolvent of $-\Delta$ is bounded at high energies, and by a resolvent identity of Vodev [Vo], this implies that there is a polynomial neighborhood of the continuous spectrum in a suitable Riemann surface (see [Gu, Me]) to which the cut-off resolvent continues holomorphically.
- (v) By [ChDa3, Theorem 1.2], we also have an asymptotic expansion for solutions to the wave equation on (X, g) . Let $u(t, x)$ solve

$$(\partial_t^2 - \Delta)u(t, x) = 0, \quad u(0, x) = f_1(x), \quad \partial_t u(0, x) = f_2(x),$$

where f_1 and f_2 are in $C_c^\infty(X)$. Then, for any $k_0 \geq 1$, uniformly as $t \rightarrow \infty$ and as x varies in a compact subset of X ,

$$u(t, x) = \int_X f_2 + \sum_{m=1}^M \left(\cos(E_m t) \int_X f_1 \eta_m + \frac{\sin(E_m t)}{E_m} \int_X f_2 \eta_m \right) \eta_m(x) + \\ \sum_{k=0}^{k_0-1} t^{-1/2-k} \sum_{n=1}^{\infty} \left(\cos(nt/\ell) a_{n,k}(x) + \sin(nt/\ell) b_{n,k}(x) \right) + \mathcal{O}(t^{-k_0}),$$

where $\{\eta_1, \dots, \eta_M\}$ is a maximal, real-valued, orthonormal set of eigenfunctions of $-\Delta$, $-\Delta \eta_m = E_m^2 \eta_m$, and $a_{n,k}, b_{n,k} \in C^\infty(X)$. By the Theorem, the sum over m is not empty.

Acknowledgments. The authors gratefully acknowledge the partial support of the Simons Foundation (TC, collaboration grant for mathematicians) and the National Science Foundation (KD, Grant DMS-1708511). TC is also grateful for partial support from a University of Missouri research leave. Thanks also to David Borthwick for helpful conversations about hyperbolic quotients.

REFERENCES

- [AvBeGiMa] Y. Avishai, D. Bessis, B. G. Giraud, and G. Mantica, *Quantum bound states in open geometries*. Phys. Rev. B 44:14 (1990), pp. 8028–8034.
- [Be] Alan F. Beardon, *The Geometry of Discrete Groups*. Grad. Texts in Math. 91, Springer-Verlag, 1983.
- [Bo1] Yannick Bonthouneau, *Weyl laws for manifolds with hyperbolic cusps*. Preprint available at arXiv:1512.05794.
- [Bo2] David Borthwick, *Spectral Theory of Infinite Area Hyperbolic Surfaces. Second Edition*. Progr. Math. 318, Birkhäuser, 2016.
- [BoDy] Jean Bourgain and Semyon Dyatlov, *Spectral gaps without the pressure condition*. Ann. of Math. 187:3 (2018), pp. 825–867.
- [BrDiKr] Ph. Briet, J. Dittrich, and D. Krejčířík, *Absolute continuity of the spectrum in a twisted Dirichlet-Neumann waveguide*. J. Math. Phys. 61:1 (2020), 013503.
- [BuGeReSi] W. Bulla, F. Gesztesy, W. Renger, and B. Simon, *Weakly coupled bound states in quantum waveguides*. Proc. Amer. Math. Soc. 125:5 (1997), pp. 1487–1495.
- [CaDuNa] G. Cardone, T. Durante, and S. A. Nazarov, *Embedded eigenvalues of the Neumann problem in a strip with a box-shaped perturbation*. J. Math. Pures Appl. 112 (2018), pp. 1–40.
- [ChDa1] T. J. Christiansen and K. Datchev, *Resolvent estimates on asymptotically cylindrical manifolds and on the half line*. To appear in Ann. Sci. Éc. Norm. Supér. Preprint available at arXiv:1705.08969.

- [ChDa2] T. J. Christiansen and K. Datchev, *Resolvent estimates, wave decay, and resonance-free regions for star-shaped waveguides*. To appear in Math. Res. Lett. Preprint available at arXiv:1910.01038.
- [ChDa3] T. J. Christiansen and K. Datchev, *Wave asymptotics for waveguides and manifolds with infinite cylindrical ends*. Preprint available at arXiv:1705.08972.
- [ChZw] T. Christiansen and M. Zworski, *Spectral asymptotics for manifolds with cylindrical ends*. Ann. Inst. Fourier, Grenoble. 45:1 (1995), pp. 251–263.
- [Co] Yves Colin de Verdière, *Pseudo-Laplaciens II*. Ann. Inst. Fourier, Grenoble. 33:2 (1983), pp. 87–113.
- [DaPa] E. B. Davies and L. Parnowski, *Trapped modes in acoustic waveguides*. Quart. J. Mech. Appl. Math. 51:3 (1998), pp. 477–492.
- [Do1] Harold Donnelly, *Eigenvalue estimates for certain noncompact manifolds*. Michigan Math. J. 31:3 (1984), pp. 349–357.
- [Do2] Harold Donnelly, *Spectral Theory of Complete Riemannian Manifolds*. Pure Appl. Math. Q. 6:2 (2010), pp. 439–456.
- [DyZa] Semyon Dyatlov and Joshua Zahl, *Spectral gaps, additive energy, and a fractal uncertainty principle*. Geom. Funct. Anal., 26:4 (2016), pp. 1011–1094.
- [EvLeVa] D. V. Evans, M. Levitin, and D. Vassiliev, *Existence theorems for trapped modes*. J. Fluid. Mech. 261 (1994), pp. 21–31.
- [ExKo] Pavel Exner and Hynek Kovařík, *Quantum Waveguides*. Springer International Publishing, 2015.
- [ExSe] P. Exner and P. Šeba, *Bound states in curved quantum waveguides*. J. Math. Phys. 30:11 (1989), pp. 2574–2580.
- [GoJa] J. Goldstone and R. L. Jaffe, *Bound states in twisting tubes*. Phys. Rev. B 45:24 (1992) 14100.
- [GrJe] D. Grieser and D. Jerison, *Asymptotics of eigenfunctions on plane domains*. Pacific J. Math. 240:1 (2009), pp. 109–133.
- [Gu] Laurent Guillopé, *Théorie spectrale de quelques variétés à bouts*. Ann. Sci. Éc. Norm. Supér. (4) 22:1 (1989), pp. 137–160.
- [HiJu] Luc Hillairet and Chris Judge, *Hyperbolic triangles without embedded eigenvalues*. Ann. of Math. 187:2 (2018), pp. 301–377.
- [IsKo] Hiroshi Isozaki and Evgeny Korotyaev, *Inverse resonance scattering on rotationally symmetric manifolds*. To appear in Asymptot. Anal. Preprint available at arXiv:1904.08908.
- [JiLi] Svetlana Jitomirskaya and Wencai Liu, *Noncompact complete Riemannian manifolds with dense eigenvalues embedded in the essential spectrum of the Laplacian*. Geom. Func. Anal. 29:1 (2019), pp. 238–257.
- [KrKř] David Krejčířík and Jan Kříž, *On the Spectrum of Curved Planar Waveguides*. Publ. Res. Inst. Math. Sci., Kyoto Univ. 41:3 (2005), pp. 757–791.
- [Ku] Hironori Kumura, *The radial curvature of an end that makes eigenvalues vanish in the essential spectrum I*. Math. Ann. 346:4 (2010), pp. 795–828.
- [LoCaMu] J. Timothy Londergan, John P. Carini, and David P. Murdock, *Binding and Scattering in Two-Dimensional Systems: Applications to Quantum Wires, Waveguides, and Photonic Crystals*. Springer-Verlag, 1999.
- [Me] Richard B. Melrose, *The Atiyah-Patodi-Singer Index Theorem*. Research Notes in Mathematics, 4. A K Peters, Ltd., Wellesley, MA, 1993.
- [MoWe] K. Morgenröther and P. Werner, *Resonances and standing waves*. Math. Methods Appl. Sci. 9:1 (1987), pp. 105–126.
- [Mü] Werner Müller, *Weyl’s law in the theory of automorphic forms*, Chapter 7 of *Groups and Analysis*. London Math. Soc. Lecture Note Ser. 354 (2010), pp. 133–163.
- [MuSeWr] David Mumford, Caroline Series, and David Wright. *Indra’s Pearls: The Vision of Felix Klein*. Cambridge University Press, 2006.
- [Pa] L. B. Parnowski, *Spectral asymptotics of the Laplace operator on manifolds with cylindrical ends*. Internat. J. Math. 6:6 (1995), pp. 911–920.
- [SaMe] Cristina V. Sargent and A. J. Mestel, *Trapped modes of the Helmholtz equation in infinite waveguides with wall indentations and circular obstacles*. IMA J. Appl. Math. 84:2 (2019), pp. 312–344.

- [Sa] Peter Sarnak, *Spectra of hyperbolic surfaces*. Bull. Amer. Math. Soc. (N.S.) 40:4 (2003), pp. 441–478.
- [Va] András Vasy, *Microlocal analysis of asymptotically hyperbolic and Kerr-de Sitter spaces (with an appendix by Semyon Dyatlov)*. Invent. Math. 194:2 (2013), pp. 381–513.
- [Vo] G. Vodev, *Semi-classical resolvent estimates and regions free of resonances*. Math. Nachr. 287:7 (2014), pp. 825–835.
- [Wi] Karl J. Witsch, *Examples of Embedded Eigenvalues for the Dirichlet-Laplacian in Domains with Infinite Boundaries*. Math. Methods Appl. Sci. 12:2 (1990), pp. 177–182.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211 USA

Email address: `christiansent@missouri.edu`

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, IN 47907 USA

Email address: `kdatchev@purdue.edu`