

TAME ALGEBRAS HAVE DENSE \mathbf{g} -VECTOR FANSPIERRE-GUY PLAMONDON AND TOSHIYA YURIKUSA,
WITH AN APPENDIX BY BERNHARD KELLER

ABSTRACT. The \mathbf{g} -vector fan of a finite-dimensional algebra is a fan whose rays are the \mathbf{g} -vectors of its 2-term presilting objects. We prove that the \mathbf{g} -vector fan of a tame algebra is dense. We then apply this result to obtain a near classification of quivers for which the closure of the cluster \mathbf{g} -vector fan is dense or is a half-space, using the additive categorification of cluster algebras by means of Jacobian algebras. As another application, we prove that for quivers with potentials arising from once-punctured closed surfaces, the stability and cluster scattering diagrams only differ by wall-crossing functions on the walls contained in a separating hyperplane. The appendix is devoted to the construction of truncated twist functors and their adjoints.

CONTENTS

| | |
|--|----|
| 1. Introduction | 1 |
| Notations and conventions | 4 |
| 2. Recollections on \mathbf{g} -vectors and varieties of representations | 4 |
| 3. \mathbf{g} -vector fans and tame algebras | 6 |
| 4. Density of the 2-term silting \mathbf{g} -vector fan | 10 |
| 5. Density of \mathbf{g} -vector fans for cluster algebras and Jacobian algebras | 18 |
| 6. Scattering diagrams | 23 |
| Acknowledgements | 28 |
| Appendix A. Truncated twist functors, by Bernhard Keller | 28 |
| References | 31 |

1. INTRODUCTION

In the theory of cluster algebras, \mathbf{g} -vectors (introduced in [FZ07]) play an important role: they endow the cluster algebra with principal coefficients with a grading, and are computationally lighter than cluster variables.

The \mathbf{g} -vectors of a cluster algebra are the rays of a fan $\mathcal{F}_{\text{cluster}}^{\mathbf{g}}$ whose cones are generated by compatible \mathbf{g} -vectors. As a consequence of results of [DWZ10], this fan is essential, rational, polyhedral and simplicial. The fan of \mathbf{g} -vectors plays an important role in the theory and appears in different guises in the literature: it appears in the tropical cluster \mathcal{X} -variety (or cluster Poisson variety) of [FG09]; it is a subfan of the underlying fan of the cluster scattering diagram of [GHKK18] and

Date: October 11, 2021.

2010 *Mathematics Subject Classification.*

The first author was supported by French ANR grant CHARMS (ANR-19-CE40-0017-02).

The second author was supported by JSPS KAKENHI Grant Number 20J00410.

of the stability scattering diagram of [Bri17]; for cluster algebras of finite type, it is a complete fan [FZ03], and is the normal fan of a polytope called the generalized associahedron [CFZ02, HPS18, AHBHY18, BDM⁺17, PPPP19].

A related notion in representation theory is the \mathbf{g} -vector fan $\mathcal{F}_{2\text{-silt}}^{\mathbf{g}}$ of 2-term silted complexes over a finite-dimensional algebra Λ . It follows from the work of [AIR14] that the fan $\mathcal{F}_{2\text{-silt}}^{\mathbf{g}}$ is an essential, rational, polyhedral, simplicial fan, and from [DIJ19] that it is complete if and only if the algebra Λ is τ -tilting finite.

In this paper, we are interested in the case where the \mathbf{g} -vector fans are not complete, but dense in the real space \mathbb{R}^n which contains them. An algebra such that $\mathcal{F}_{2\text{-silt}}^{\mathbf{g}}$ is dense was called \mathbf{g} -tame in [AY20]. Our main result is the following.

Theorem 1.1 (Theorem 4.1). *Let Λ be a finite-dimensional basic algebra over an algebraically closed field. If Λ is tame, then its fan $\mathcal{F}_{2\text{-silt}}^{\mathbf{g}}$ is dense in \mathbb{R}^n . In other words, if Λ is tame, then Λ is \mathbf{g} -tame.*

Theorem 1.1 was previously known for path algebras of extended Dynkin type by the results of [Hil06]; for representation-finite algebras by [DIJ19] (and in fact for any τ -tilting finite algebra, even if it is not tame); for Jacobian algebras associated to triangulated surfaces by [Yur20] (tame or not); for gentle algebras by [AY20]; and for special biserial algebras by [AY20] and [ADI20], both using different methods.

We note that the converse of Theorem 1.1 is false: the quiver associated with a triangulation of a torus with one boundary component and one marked point on it admits a wild potential by [GLFS16], even though its \mathbf{g} -vector fan is dense by [Yur20]. The class of \mathbf{g} -tame algebras is thus strictly larger than that of tame algebras.

The related notion of τ -tilting tameness was introduced in [BST19]: an algebra is τ -tilting tame if the closure $\overline{\text{Wall}}$ of the union of the walls containing non-trivial stability conditions in \mathbb{R}^n has measure zero. Using recent results, it is not hard to see that τ -tilting tameness implies \mathbf{g} -tameness: by [Asa19, Theorem 3.17], $\overline{\text{Wall}}$ is the complement of the union of the interior of the maximal cones of $\mathcal{F}_{2\text{-silt}}^{\mathbf{g}}$, so if the latter is not dense, $\overline{\text{Wall}}$ contains the cone generated by an open subset of \mathbb{R}^n , and so has infinite measure. We conjecture that τ -tilting tameness is equivalent to \mathbf{g} -tameness. This, combined with Theorem 1.1, would imply that tame algebras are τ -tilting tame, as conjectured in [BST19, Conjecture 3.22]. We do not discuss τ -tilting tameness further in this paper.

We provide two applications of Theorem 1.1. The first is a near classification of quivers Q whose cluster algebras have dense \mathbf{g} -vector fans. We say that a quiver Q is *cluster- \mathbf{g} -dense* in that case, and *half cluster- \mathbf{g} -dense* if the closure of both its cluster fan and that of Q^{op} are half-spaces.

Corollary 1.2 (Theorem 5.17). *Let Q be a quiver without loops and 2-cycles. Assume that Q is not mutation equivalent to X_6, X_7 (see Theorem 5.7) or K_m , with $m \geq 3$ (see Example 5.4). Then Q is mutation-finite if and only if it is cluster- \mathbf{g} -dense or half cluster- \mathbf{g} -dense. In this case,*

- Q is half cluster- \mathbf{g} -dense if it arises from a triangulation of a closed surface with exactly one puncture, and
- Q is cluster- \mathbf{g} -dense otherwise.

We conjecture that X_6 and X_7 behave no differently.

Conjecture 1.3. (1) *The \mathbf{g} -vector fan in type X_6 is dense.*

(2) *The closure of the \mathbf{g} -vector fan in type X_7 is a half-space.*

We prove Corollary 1.2 by applying Theorem 1.1 and a result of [GLFS16] that states that all mutation-finite quivers admit a tame non-degenerate potential, except for X_6 , X_7 and K_m , with $m \geq 3$. We note that Corollary 1.2 was already known for quivers of Dynkin type [FZ03], extended Dynkin type [Hil06] and of surface type [Yur20]; thus the novelty here is the inclusion of the quivers of type $E_6^{(1,1)}$, $E_7^{(1,1)}$ and $E_8^{(1,1)}$.

The second application is a consequence of an argument due to L. Mou [Mou19] to get information on a conjecture of [KS14].

Theorem 1.4 (Theorem 6.9). *Let Q be a quiver without loops and 2-cycles, and let W be a non-degenerate potential on it. Let \mathfrak{D}_Q be the cluster scattering diagram of Q , and let $\mathfrak{D}_{J(Q,W)}$ be the stability scattering diagram of the Jacobian algebra $J(Q,W)$ (see Section 6).*

- (1) *If Q is cluster- \mathbf{g} -dense, then the cluster scattering diagram \mathfrak{D}_Q is equal to the stability scattering diagram $\mathfrak{D}_{J(Q,W)}$.*
- (2) *If Q is half cluster- \mathbf{g} -dense, then \mathfrak{D}_Q and $\mathfrak{D}_{J(Q,W)}$ only differ by functions on walls in the separating hyperplane.*

This allows to obtain new information on the scattering diagrams of cluster algebras from once-punctured closed surfaces.

Corollary 1.5. *For a cluster algebra arising from a once punctured closed surface, the two scattering diagrams only differ by functions on the walls of the separating hyperplane.*

We note that Conjecture 1.3, together with Theorem 1.4, would imply similar results in types X_6 and X_7 . Cases where the two scattering diagrams are known to differ by central elements are Jacobian algebras admitting a green-to-red sequence and the Jacobian algebra of a once-punctured torus [Mou19], and examples arising from del Pezzo surfaces [BMP20]; the latter reference also contains example of scattering diagrams for which the property does not hold, as explained for example in the lecture [Bou20] by Pierrick Bousseau.

The proof of Theorem 1.1 relies on two main ingredients. On one hand, it uses in an essential way the results on generic decompositions of \mathbf{g} -vectors obtained in [DF15, Pla13] and their more precise formulation for tame algebras shown in [GLFS20] (see Theorem 3.8). On the other hand, it relies on an operation Cyl_X on the objects of the homotopy category $K^b(\text{proj } \Lambda)$. The operation Cyl_X is reminiscent of a spherical twist with respect to an object X [ST01], with the important caveat that Cyl_X is not an auto-equivalence; in fact, it is not even a functor. The main difficulty in our proof of Theorem 1.1 is to show that Cyl_X , despite not being as good as a twist functor, behaves nicely enough on a subcategory of $K^b(\text{proj } \Lambda)$.

Our strategy is similar to (and partly inspired by) the one used in [AY20], where Theorem 1.1 is proved for completed gentle algebras by using the geometric model of [OPS18]. The idea there is to notice that the boundary of the \mathbf{g} -vector fan is governed by the \mathbf{g} -vectors of band objects, which correspond to closed curves in the geometric model, and to approach this boundary by inflicting successive Dehn twists on arcs with respect to those closed curves. Our operators Cyl_X should be seen as an “algebraic counterpart” of these Dehn twists, and indeed the two notions coincide in the case of gentle algebras. Under suitable vanishing conditions, the

action of the operator Cyl_X coincides with that of a truncated dual twist functor as constructed in the appendix.

This paper is organized as follows. In Section 2, we collect definitions and results on generic decomposition of \mathbf{g} -vectors and varieties of representations of an algebra which will be needed in the paper. In Section 3, we define the \mathbf{g} -vector fan of 2-term silting objects of an algebra, and we state a result of [GLFS20] on the generic decomposition of \mathbf{g} -vectors for tame algebras (Theorem 3.8). In Section 4, we introduce the operator Cyl_X and prove Theorem 1.1. We then apply our results to the density of \mathbf{g} -vector fans of cluster algebras in Section 5 and to scattering diagrams in Section 6.

NOTATIONS AND CONVENTIONS

In this paper, k will always denote an algebraically closed field. All modules will be right modules unless stated otherwise. Arrows in a quiver are composed from right to left, as are morphisms in a category. The suspension functor of a triangulated category will always be denoted by Σ . In any category \mathcal{C} , the set of morphisms from an object A to an object B will be denoted by $\text{Hom}_{\mathcal{C}}(A, B)$, or sometimes (A, B) to save space. A general element of a (quasi-projective) variety will always mean an element of a dense open subset of the variety.

2. RECOLLECTIONS ON \mathbf{g} -VECTORS AND VARIETIES OF REPRESENTATIONS

In this section, we fix a finite-dimensional basic k -algebra Λ .

2.1. Generic decomposition of \mathbf{g} -vectors. Denote by $\Lambda = \bigoplus_{i=1}^n P_i$ a decomposition of Λ as direct sum of pairwise non-isomorphic indecomposable projective right Λ -modules. We will denote by $K^b(\text{proj } \Lambda)$ the homotopy category of bounded complexes of finitely generated projective right Λ -modules. We denote by $K^{[-1,0]}(\text{proj } \Lambda)$ the full subcategory whose objects are complexes concentrated in degrees -1 and 0 . An object P in $K^{[-1,0]}(\text{proj } \Lambda)$ will be denoted by

$$P = P^{-1} \xrightarrow{f} P^0;$$

by abuse of terminology, we will sometimes identify the object P with the morphism f .

The Grothendieck group of the triangulated category $K^b(\text{proj } \Lambda)$ will be denoted by $K_0(\text{proj } \Lambda)$; we note that it is a free abelian group with basis the images of the indecomposable projectives P_1, \dots, P_n . The image of an object X in the Grothendieck group will be denoted by $[X]$. The basis $[P_1], \dots, [P_n]$ gives a natural identification of $K_0(\text{proj } \Lambda)$ with \mathbb{Z}^n , which we will often use implicitly.

Definition 2.1. A \mathbf{g} -vector is an element of $K_0(\text{proj } \Lambda)$. The \mathbf{g} -vector of an object P of $K^{[-1,0]}(\text{proj } \Lambda)$ is the element $[P] \in K_0(\text{proj } \Lambda)$.

For any $\mathbf{g} \in K_0(\text{proj } \Lambda)$, we let $P^{\mathbf{g}+}$ and $P^{\mathbf{g}-}$ be the unique finitely generated projective modules without common non-zero direct summands (up to isomorphism) such that $\mathbf{g} = [P^{\mathbf{g}+}] - [P^{\mathbf{g}-}]$.

Definition 2.2. Let $\mathbf{g}, \mathbf{g}' \in K_0(\text{proj } \Lambda)$. We denote by $e(\mathbf{g}, \mathbf{g}')$ the minimal value of

$$\dim \text{Hom}_{K^b(\text{proj } \Lambda)}(P, \Sigma P'),$$

where P and P' are objects of $K^{[-1,0]}(\text{proj } \Lambda)$ such that $[P] = \mathbf{g}$ and $[P'] = \mathbf{g}'$.

The following result is due to H. Derksen and J. Fei [DF15, Theorem 4.4]. We use the formulation given in [Pla13, Theorem 2.7].

Theorem 2.3 (Generic decomposition of \mathbf{g} -vectors). *Any $\mathbf{g} \in K_0(\text{proj } \Lambda)$ can be written as*

$$\mathbf{g} = \mathbf{g}_1 + \dots + \mathbf{g}_r,$$

where

- (1) for each $i \in \{1, \dots, r\}$, a general element of $\text{Hom}_\Lambda(P^{(\mathbf{g}_i)-}, P^{(\mathbf{g}_i)+})$ is indecomposable;
- (2) for each $i, j \in \{1, \dots, r\}$ with $i \neq j$, we have that $e(\mathbf{g}_i, \mathbf{g}_j) = 0$.

Moreover, the elements $\mathbf{g}_1, \dots, \mathbf{g}_r$ are unique for these properties (up to re-ordering), and a general element of $\text{Hom}_\Lambda(P^{\mathbf{g}-}, P^{\mathbf{g}+})$ is a direct sum of elements in

$$\text{Hom}_\Lambda(P^{(\mathbf{g}_1)-}, P^{(\mathbf{g}_1)+}), \dots, \text{Hom}_\Lambda(P^{(\mathbf{g}_r)-}, P^{(\mathbf{g}_r)+}).$$

Definition 2.4. Let $\mathbf{g} \in K_0(\text{proj } \Lambda)$. The decomposition of \mathbf{g} given in Theorem 2.3 is the *generic decomposition* of \mathbf{g} . If $r = 1$, then \mathbf{g} is *generically indecomposable*.

2.2. Varieties of representations and generically τ -reduced components.

Since k is algebraically closed and Λ is basic, we can assume that Λ is the quotient of the path algebra of a finite quiver Q by an admissible ideal I (see, for instance, [ASS06]). Let e_1, \dots, e_n be the paths of length 0 corresponding to the vertices of Q ; we can assume that $P_i = e_i \Lambda$ for each $i \in \{1, \dots, n\}$. Let S_1, \dots, S_n be the simple tops of P_1, \dots, P_n , respectively.

We denote by $\text{mod } \Lambda$ the category of finitely generated right Λ -modules, and by $K_0(\text{mod } \Lambda)$ its Grothendieck group. Note that $K_0(\text{mod } \Lambda)$ is a free abelian group, and that $[S_1], \dots, [S_n]$ form a basis for it. Finally, let

$$K_0(\text{mod } \Lambda)^\oplus := \left\{ \sum_{i=1}^n a_i [S_i] \mid a_1, \dots, a_n \in \mathbb{Z}_{\geq 0} \right\}.$$

Definition 2.5. A *dimension vector* is an element of $K_0(\text{mod } \Lambda)^\oplus$. The *dimension vector of a module M* is the element $[M] \in K_0(\text{mod } \Lambda)^\oplus$.

For any $\mathbf{d} \in K_0(\text{mod } \Lambda)^\oplus$, we let

$$\text{rep}_{\mathbf{d}}(Q^{\text{op}}) := \bigoplus_{\alpha \in Q_1} \text{Hom}_k(k^{d_{t(\alpha)}}, k^{d_{s(\alpha)}}).$$

It is an affine space whose points correspond to representations of the quiver Q^{op} with dimension vector \mathbf{d} . Inside it is the Zariski-closed subset $\text{rep}_{\mathbf{d}}(\Lambda)$ of representations satisfying the relations in the ideal I ; points in $\text{rep}_{\mathbf{d}}(\Lambda)$ correspond to right Λ -modules M such that $[M] = \mathbf{d}$.

The affine algebraic group $\text{GL}_{\mathbf{d}} := \prod_{i \in Q_0} \text{GL}_{d_i}(k)$ acts on $\text{rep}_{\mathbf{d}}(\Lambda)$ by

$$(g_i)_{i \in Q_0} \cdot (f_\alpha)_{\alpha \in Q_1} = (g_{s(\alpha)} f_\alpha g_{t(\alpha)}^{-1})_{\alpha \in Q_1}.$$

The $\text{GL}_{\mathbf{d}}$ -orbits of $\text{rep}_{\mathbf{d}}(\Lambda)$ are in bijection with isomorphism classes of right Λ -modules with dimension vector \mathbf{d} . The orbit of a point M will be denoted by \mathcal{O}_M . By abuse of notation, we will identify a point M in $\text{rep}_{\mathbf{d}}(\Lambda)$ with the module that it represents.

Let \mathcal{Z} be an irreducible component of $\text{rep}_{\mathbf{d}}(\Lambda)$. It is known that \mathcal{Z} is stable under the action of $\text{GL}_{\mathbf{d}}$, and that for any point $M \in \mathcal{Z}$,

$$\text{codim}_{\mathcal{Z}} \mathcal{O}_M \leq \dim_k \text{Ext}_{\Lambda}^1(M, M) \leq \dim_k \text{Hom}_{\Lambda}(M, \tau M),$$

where τ is the Auslander–Reiten translation (here, the first inequality follows from Voigt’s lemma [Gab74, Proposition 1.1] and the second one from the Auslander–Reiten duality [ASS06, Theorem IV.2.13]). The following definition is a dualized version of the notion of *strongly reduced component* which was introduced by C. Geiss, B. Leclerc and J. Schröer in the context of additive categorification of cluster algebras and of G. Lusztig’s dual semicanonical bases.

Definition 2.6 (Section 7.1 of [GLS12]). Let \mathbf{d} be a dimension vector. An irreducible component \mathcal{Z} of $\text{rep}_{\mathbf{d}}(\Lambda)$ is *generically τ -reduced* if, for a general $M \in \mathcal{Z}$, we have that

$$\text{codim}_{\mathcal{Z}} \mathcal{O}_M = \dim_k \text{Hom}_{\Lambda}(M, \tau M).$$

The next result states that generically τ -reduced components are parametrized by \mathbf{g} -vectors (in the case of τ -rigid modules, this result is due to [DK08]; see also [DF15] for generic decompositions of projective presentations).

Theorem 2.7 (Theorem 1.2 of [Pla13]). *Let $\mathbf{g} \in K_0(\text{proj } \Lambda)$. There is an open dense subset \mathcal{U} of $\text{Hom}_{\Lambda}(P^{\mathbf{g}-}, P^{\mathbf{g}+})$ such that*

- (1) *there is a dimension vector $\mathbf{d} \in K_0(\text{mod } \Lambda)^{\oplus}$ such that for any $f \in \mathcal{U}$, we have that $[\text{Coker } f] = \mathbf{d}$;*
- (2) *the union of the orbits of the cokernels of all $f \in \mathcal{U}$ is a dense subset of a generically τ -reduced component \mathcal{Z} of $\text{rep}_{\mathbf{d}}(\Lambda)$.*

Moreover, all generically τ -reduced components arise in this way, and two vectors $\mathbf{g}, \mathbf{g}' \in K_0(\text{proj } \Lambda)$ give rise to the same generically τ -reduced component if and only if their generic decompositions are the same, up to non-positive generically indecomposable terms.

3. \mathbf{g} -VECTOR FANS AND TAME ALGEBRAS

3.1. \mathbf{g} -vector fans. The \mathbf{g} -vectors of presilting objects of $K^b(\text{proj } \Lambda)$ are arranged into a structure called a simplicial fan. We recall here the main definitions on fans, and refer to [Zie95] for a general treatment.

Definition 3.1. Let $d \in \mathbb{Z}_{>0}$.

- (1) For any non empty subset X of \mathbb{R}^d , a *supporting hyperplane* of X is a hyperplane H of \mathbb{R}^d such that $H \cap X$ is non-empty and such that X is contained in one of the two half-spaces defined by H .
- (2) A *polyhedral cone* in \mathbb{R}^d is a set of the form

$$C(c_1, \dots, c_r) = \left\{ \sum_{i=1}^r \lambda_i c_i \mid \lambda_i \in \mathbb{R}_{\geq 0} \text{ for all } i \right\},$$

where c_1, \dots, c_r are vectors in \mathbb{R}^d . Equivalently, a polyhedral cone is the intersection of finitely many linear half-spaces. The *dimension* of $C(c_1, \dots, c_r)$ is the dimension of the vector space spanned by c_1, \dots, c_r . A cone $C(c_1, \dots, c_r)$ is *simplicial* if c_1, \dots, c_r are linearly independent.

- (3) The *faces* of a polyhedral cone $C(c_1, \dots, c_r)$ are its intersections with its supporting hyperplanes. A *ray* of a cone is a face of dimension 1; a *facet* is a face of codimension 1.
- (4) A *polyhedral fan* in \mathbb{R}^d is a set \mathcal{F} of polyhedral cones in \mathbb{R}^d such that
 - if $C \in \mathcal{F}$, then any face of C is in \mathcal{F} ; and
 - the intersection of two cones in \mathcal{F} is a face of both.

- (5) A polyhedral fan is *simplicial* if all its cones are simplicial. It is *complete* if the union of its cones is \mathbb{R}^d . It is *essential* if it contains the cone $\{0\}$.

The following follows from the fundamental results of τ -tilting theory, see [AIR14].

Theorem 3.2 ([AIR14]). *Let Λ be any basic finite-dimensional k -algebra. Then the \mathbf{g} -vectors of indecomposable presilting objects of $K^{[-1,0]}(\text{proj } \Lambda)$ are the rays of an essential simplicial polyhedral fan whose maximal cones are $C(g_1, \dots, g_n)$, where g_1, \dots, g_n are the \mathbf{g} -vectors of pairwise non-isomorphic indecomposable objects S_1, \dots, S_n such that $S = \bigoplus_{i=1}^n S_i$ is a silting object.*

Definition 3.3. The simplicial polyhedral fan described in Theorem 3.2 is the *2-term silting \mathbf{g} -vector fan* of Λ , and it is denoted by $\mathcal{F}_{2\text{-silt}}^{\mathbf{g}}(\Lambda)$.

3.2. Tame algebras. The main class of algebras that we will study is the following.

Definition 3.4. The algebra Λ is *tame* if for any dimension vector \mathbf{d} , there are $k[t]$ - Λ -bimodules $M_1, \dots, M_{m(\mathbf{d})}$ such that

- (1) for all $i \in \{1, \dots, m(\mathbf{d})\}$, the module M_i is free of finite rank as a $k[t]$ -module;
- (2) all but finitely many indecomposable Λ -modules of dimension vector \mathbf{d} have the form

$$k[t]/(t - \lambda) \otimes_{k[t]} M_i$$

with $i \in \{1, \dots, m(\mathbf{d})\}$ and $\lambda \in k$.

For a given M_i , the family of indecomposable Λ -modules of the form $k[t]/(t - \lambda) \otimes_{k[t]} M_i$ will be called a *1-parameter family of indecomposable modules*.

We will be using an important result on tame algebras, due to W. Crawley-Boevey.

Theorem 3.5 (Theorem D of [CB88]). *Let Λ be a tame algebra. Then for any dimension vector \mathbf{d} , all but finitely many isomorphism classes of Λ -modules L of dimension vector \mathbf{d} satisfy $\tau L \cong L$.*

3.3. Generic decomposition of \mathbf{g} -vectors for tame algebras. In this section, we state a theorem due to C. Geiss, D. Labardini-Fragoso and J. Schröer [GLFS20]. We first recall some definitions.

Definition 3.6. An object X of $K^b(\text{proj } \Lambda)$ is *presilting* if $\text{Hom}_{K^b(\text{proj } \Lambda)}(X, X[i])$ vanishes for all $i > 0$. It is called *silting* if, moreover, it generates $K^b(\text{proj } \Lambda)$ as a triangulated category.

Definition 3.7. A Λ -module is a *brick* if its endomorphism algebra is isomorphic to the base field k .

Theorem 3.8 ([GLFS20]). *Let Λ be a tame algebra, and let $\mathbf{g} \in K_0(\text{proj } \Lambda)$ be a \mathbf{g} -vector. Then the generic decomposition of \mathbf{g} has the form*

$$\mathbf{g} = \mathbf{g}_1 + \dots + \mathbf{g}_r + \mathbf{h}_1 + \dots + \mathbf{h}_s,$$

where $r, s \geq 0$ and

- (1) for $i \in \{1, \dots, r\}$, the vector \mathbf{g}_i is generically indecomposable and such that for a general f in $\text{Hom}_{\Lambda}(P^{(\mathbf{g}_i)-}, P^{(\mathbf{g}_i)+})$, the object f of $K^{[-1,0]}(\text{proj } \Lambda)$ is presilting;

- (2) for $j \in \{1, \dots, s\}$, the vector \mathbf{h}_j is generically indecomposable and there is a dense open subset \mathcal{U} of $\text{Hom}_\Lambda(P^{(\mathbf{h}_j)^-}, P^{(\mathbf{h}_j)^+})$ such that the cokernels of morphisms in \mathcal{U} form a 1-parameter family of indecomposable Λ -modules which are bricks and are isomorphic to their own Auslander-Reiten translate. Moreover, $e(\mathbf{h}_j, \mathbf{h}_j) = 0$.

Theorem 3.8 was announced and its proof outlined in a lecture by J. Schröer at MFO in January 2020. It is contained in [GLFS20], where it is formulated in terms of generically τ -reduced components. The above formulation in terms of generic decomposition of \mathbf{g} -vectors can be obtained by combining Theorem 3.2 and Section 4.2 of [GLFS20] together with [DF15, Theorem 4.4] and [Pla13, Theorem 1.2]. We include a complete proof here for the convenience of the reader.

Proof. Assume first that \mathbf{g} is generically indecomposable. Let \mathcal{Z} be the irreducible component of $\text{rep}_{\mathbf{d}}(\Lambda)$ obtained by taking the closure of the union of the orbits of cokernels of generic elements in $\text{Hom}_\Lambda(P^{\mathbf{g}^-}, P^{\mathbf{g}^+})$. By Theorem 2.7, the component \mathcal{Z} is generically τ -reduced.

Since Λ is tame, there exist $k[t]$ - Λ -bimodules $M_1, \dots, M_{m(\mathbf{d})}$, free of rank 1 as $k[t]$ -modules, such that almost all indecomposable Λ -modules of dimension vector \mathbf{d} are isomorphic to

$$k[t]/(t - \lambda) \otimes_{k[t]} M_i$$

for some $\lambda \in k$ and some $i \in \{1, \dots, m(\mathbf{d})\}$. Each M_i induces a morphism of varieties

$$\phi_i : \mathbb{A}^1 \rightarrow \text{rep}_{\mathbf{d}}(\Lambda)$$

sending $\lambda \in \mathbb{A}^1$ to a point corresponding to the module $k[t]/(t - \lambda) \otimes_{k[t]} M_i$. Since \mathbb{A}^1 is irreducible, the image of each ϕ_i either is contained in \mathcal{Z} or has empty intersection with \mathcal{Z} . Assume that the indices for which the image of ϕ_i is contained in \mathcal{Z} are $1, \dots, m'$.

For each $i \in \{1, \dots, m'\}$, consider the morphism

$$\begin{aligned} \psi_i : \text{GL}_{\mathbf{d}} \times \mathbb{A}^1 &\longrightarrow \mathcal{Z} \\ (g, \lambda) &\longmapsto g \cdot \phi_i(\lambda). \end{aligned}$$

By our assumptions on \mathcal{Z} , the union of the images of the ψ_i is dense in \mathcal{Z} . Therefore, there is an $\ell \in \{1, \dots, m'\}$ such that the image of ψ_ℓ is dense in \mathcal{Z} .

Let L be a generic point in the image of ψ_ℓ ; we shall make a small abuse of notation and denote by L the corresponding Λ -module. Assume that $L = \psi_\ell(g, \lambda)$. Then $\psi_\ell(\text{GL}_{\mathbf{d}} \times \{\lambda\})$ is the orbit \mathcal{O}_L of L in \mathcal{Z} . Since $\psi_\ell(\text{GL}_{\mathbf{d}} \times \mathbb{A}^1)$ is dense in \mathcal{Z} , it is of codimension 0. Thus, \mathcal{O}_L is of codimension 0 or 1 in \mathcal{Z} .

Case 1: $\text{codim}_{\mathcal{Z}} \mathcal{O}_L = 0$. Since \mathcal{Z} is generically τ -reduced, we have the equalities $\dim \text{Hom}_\Lambda(L, \tau L) = \text{codim}_{\mathcal{Z}} \mathcal{O}_L = 0$. Therefore, L is τ -rigid. Since the projective presentations of τ -rigid modules are 2-term presilting objects, this shows that generic elements in the space $\text{Hom}_{K^b(\text{proj } \Lambda)}(P^{\mathbf{g}^-}, P^{\mathbf{g}^+})$ is presilting. Thus \mathbf{g} satisfies condition (1) of the Theorem.

Case 2: $\text{codim}_{\mathcal{Z}} \mathcal{O}_L = 1$. In this case, since \mathcal{Z} is generically τ -reduced, we get that $\dim \text{Hom}_\Lambda(L, \tau L) = 1$. In particular, \mathcal{Z} contains infinitely many orbits. By [CB88, Theorem D], since L is generic, we get that τL is isomorphic to L . Therefore $\dim \text{Hom}_\Lambda(L, L) = 1$, and L is a brick. Thus to show that \mathbf{g} satisfies condition (2) of the Theorem, all that remains is to see that $e(\mathbf{g}, \mathbf{g}) = 0$.

Let (f_1, f_2) be a generic element of $\text{Hom}_\Lambda(P^{\mathbf{g}^-}, P^{\mathbf{g}^+}) \times \text{Hom}_\Lambda(P^{\mathbf{g}^-}, P^{\mathbf{g}^+})$. Then

$$\begin{aligned} e(\mathbf{g}, \mathbf{g}) &= \dim \text{Hom}_{K^b(\text{proj } \Lambda)}(f_1, \Sigma f_2) \\ &= \dim \text{Hom}_\Lambda(\text{Coker}(f_2), \tau \text{Coker}(f_1)) \\ &= \dim \text{Hom}_\Lambda(\text{Coker}(f_2), \text{Coker}(f_1)), \end{aligned}$$

where the second equality is by [Pla13, Lemma 2.6] and the third one by [CB88, Theorem D]. Letting $L_i = \text{Coker}(f_i)$ for $i \in \{1, 2\}$, it suffices to show that $\text{Hom}_\Lambda(L_2, L_1) = 0$.

The following argument was suggested to us by A. Skowroński and G. Zwara. Since L_1 and L_2 are bricks (as proved above), we have that

$$\text{rad End}_\Lambda(L_1) = 0 = \text{rad End}_\Lambda(L_2).$$

Since the function $(X, Y) \mapsto \dim \text{rad}_\Lambda(X, Y)$ is upper semicontinuous on \mathcal{Z} , this implies that for generic L_1 and L_2 , we have that $\text{rad}_\Lambda(L_2, L_1) = 0$. Since L_1 and L_2 are indecomposable and non isomorphic, this in turn implies that $\text{Hom}_\Lambda(L_2, L_1) = 0$.

This proves that $e(\mathbf{g}, \mathbf{g}) = 0$.

The theorem is now proved for a generically indecomposable \mathbf{g} . If \mathbf{g} is generically decomposable, simply apply the above to each term of its generic decomposition, and the theorem is proved. \square

We will need a technical corollary of Theorem 3.8.

Corollary 3.9. *Let Λ be any algebra satisfying conditions (1) and (2) of Theorem 3.8. Then*

(1) *for $j \in \{1, \dots, s\}$ and a general*

$$h \in \text{Hom}_\Lambda(P^{(\mathbf{h}_j)^-}, P^{(\mathbf{h}_j)^+}),$$

the space $\text{Hom}_{K^b(\text{proj } \Lambda)}(h, \Sigma h)$ is one-dimensional;

(2) *for $i \in \{1, \dots, r\}$, $j \in \{1, \dots, s\}$ and a general*

$$(g, h) \in \text{Hom}_\Lambda(P^{(\mathbf{g}_i)^-}, P^{(\mathbf{g}_i)^+}) \times \text{Hom}_\Lambda(P^{(\mathbf{h}_j)^-}, P^{(\mathbf{h}_j)^+}),$$

all morphisms in $\text{Hom}_{K^b(\text{proj } \Lambda)}(g, h)$ factor through an object of $\text{add}(\Sigma \Lambda)$;

(3) *for $j, \ell \in \{1, \dots, s\}$, and a general*

$$(h, h') \in \text{Hom}_\Lambda(P^{(\mathbf{h}_j)^-}, P^{(\mathbf{h}_j)^+}) \times \text{Hom}_\Lambda(P^{(\mathbf{h}_\ell)^-}, P^{(\mathbf{h}_\ell)^+}),$$

all morphisms in $\text{Hom}_{K^b(\text{proj } \Lambda)}(h, h')$ factor through an object of $\text{add}(\Sigma \Lambda)$.

Proof. To prove (1), we note that

$$\begin{aligned} \text{Hom}_{K^b(\text{proj } \Lambda)}(h, \Sigma h) &\cong \text{Hom}_\Lambda(H^0 h, \tau H^0 h) \quad (\text{by [Pla13, Lemma 2.6]}) \\ &\cong \text{Hom}_\Lambda(H^0 h, H^0 h) \quad (\text{since } H^0 h \cong \tau H^0 h \text{ by [CB88]}) \\ &\cong k \quad (\text{since } H^0 h \text{ is a brick}). \end{aligned}$$

The proof of (2) is similar:

$$\begin{aligned} \text{Hom}_{K^b(\text{proj } \Lambda)}(g, h)/(\Sigma \Lambda) &\cong \text{Hom}_\Lambda(H^0 g, H^0 h) \\ &\cong \text{Hom}_\Lambda(H^0 g, \tau H^0 h) \quad (\text{since } H^0 h \cong \tau H^0 h \text{ by [CB88]}) \\ &\cong \text{Hom}_{K^b(\text{proj } \Lambda)}(h, \Sigma g) \quad (\text{by [Pla13, Lemma 2.6]}) \\ &\cong 0, \end{aligned}$$

where $(\Sigma\Lambda)$ is the subspace of morphisms factoring through $\text{add}(\Sigma\Lambda)$. Finally, (3) is proved as follows:

$$\begin{aligned} \text{Hom}_{K^b(\text{proj } \Lambda)}(h, h')/(\Sigma\Lambda) &\cong \text{Hom}_{\Lambda}(H^0 h, H^0 h') \\ &\cong \text{Hom}_{\Lambda}(H^0 h, \tau H^0 h') \quad (\text{since } H^0 h' \cong \tau H^0 h' \text{ by [CB88]}) \\ &\cong \text{Hom}_{K^b(\text{proj } \Lambda)}(h', \Sigma h) \quad (\text{by [Pla13, Lemma 2.6]}) \\ &\cong 0. \end{aligned}$$

□

3.4. \mathbf{g} -tame algebras. The property that we wish to study is encapsulated in the following definition. Let Λ be a finite-dimensional algebra over a field k . We denote by $\bar{\mathcal{F}}_{2\text{-silt}}^{\mathbf{g}}(\Lambda)$ the closure of the union of the cones of the 2-term silting \mathbf{g} -vector fan $\mathcal{F}_{2\text{-silt}}^{\mathbf{g}}(\Lambda)$.

Definition 3.10 ([AY20]). The algebra Λ is \mathbf{g} -tame if $\bar{\mathcal{F}}_{2\text{-silt}}^{\mathbf{g}}(\Lambda) = \mathbb{R}^n$.

In Section 5, we will need the following property of \mathbf{g} -tame algebras.

Proposition 3.11. *Let e be an idempotent of Λ . If Λ is \mathbf{g} -tame, then so is $\Lambda/(e)$.*

Proof. Let $|e| = n - m$. Suppose that $\Lambda/(e)$ is not \mathbf{g} -tame. Then by [Asa19, Theorem 3.17], there is some m -dimensional cone C in $K_0(\text{proj } \Lambda/(e)) \simeq \mathbb{R}^m$ such that there is a non-zero θ -semistable $(\Lambda/(e))$ -module $M(\theta)$ for any $\theta \in C \cap \mathbb{Q}^m$. Let $\phi : K_0(\text{proj } \Lambda) \simeq \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a natural coordinate projection. Then $M(\theta)_{\Lambda}$ is a $\phi^{-1}(\theta)$ -semistable Λ -module. Thus there is a non-zero θ' -semistable Λ -module for any $\theta' \in \phi^{-1}(C) \cap \mathbb{Q}^n$, where $\phi^{-1}(C)$ is an n -dimensional cone in $K_0(\text{proj } \Lambda)$. By [Asa19, Theorem 3.17] again, Λ is not \mathbf{g} -tame. □

4. DENSITY OF THE 2-TERM SILTING \mathbf{g} -VECTOR FAN

Our main theorem is the following.

Theorem 4.1. *Let Λ be a tame basic finite-dimensional k -algebra, and let n be the rank of its Grothendieck group. Then any vector of \mathbb{Q}^n is in the closure $\bar{\mathcal{F}}_{2\text{-silt}}^{\mathbf{g}}$ of the union of the cones of the 2-term silting \mathbf{g} -vector fan $\mathcal{F}_{2\text{-silt}}^{\mathbf{g}}$. In particular, Λ is \mathbf{g} -tame.*

This section is devoted to the proof of Theorem 4.1.

4.1. Cylinders. The proof of Theorem 4.1 involves a variation on the notion of spherical twist.

Definition 4.2. Let \mathcal{T} be an essentially small, Hom-finite, Krull-Schmidt, triangulated category with suspension functor Σ . Let X be an object of \mathcal{T} . For any object U of \mathcal{T} , choose a basis (f_1, \dots, f_d) of the space $\text{Hom}_{\mathcal{T}}(U, X)$ and a triangle

$$\Sigma^{-1}X^d \rightarrow \text{Cyl}_X U \rightarrow U \xrightarrow{f} X^d,$$

where $f = \begin{bmatrix} f_1 \\ f_2 \\ \vdots \\ f_d \end{bmatrix}$. The object $\text{Cyl}_X U$ is the *cylinder of U with respect to X* .

Remark 4.3. (1) The object $\text{Cyl}_X U$ is only defined up to (non-unique) isomorphism. Moreover, we do not define an action of Cyl_X on morphisms, so it is not a functor.

(2) The morphism $U \xrightarrow{f} X^d$ is functorial in U . Indeed, the functor

$$\text{Hom}_{\mathcal{T}}(?, X) : \mathcal{T} \rightarrow (\text{mod } k)^{\text{op}}$$

admits the right adjoint $D(?) \otimes_k X \cong \text{Hom}_k(?, X)$, and the morphism $U \xrightarrow{f} X^d$ is then the unit of adjunction

$$\text{Id}_{\mathcal{T}} \rightarrow \text{Hom}_k(\text{Hom}_{\mathcal{T}}(?, X), X)$$

applied to U .

- (3) Our definition of Cyl_X is similar to, but different from, the (dual) definition of the twist Tw_X (or spherical twist if X is a spherical object) [ST01, Definition 2.7]. Indeed, the latter involves the morphisms in every degree from U to X , while the definition of $\text{Cyl}_X U$ only involves the morphisms in degree 0.
- (4) In the Appendix A, Bernhard Keller constructs a truncated version t_X^0 of the twist functor. Applying Example A.2(2) to the case where $\mathcal{A} = \Lambda$, we get that the truncated twist functor t_X^0 and the operator Cyl_X act in the same way on certain objects; this happens, for instance, in the setting of Lemma 4.5.

4.2. Commuting cylinders. The first key lemma in the proof of Theorem 4.1 is a condition for cylinders to commute.

Lemma 4.4 (Commuting cylinders). *Let \mathcal{T} be an essentially small, Hom-finite, Krull-Schmidt, triangulated category with suspension functor Σ . Let X and Y be non-isomorphic indecomposable objects of \mathcal{T} , and let U be any object of \mathcal{T} . Assume that the following hold:*

- (1) $\text{Hom}_{\mathcal{T}}(X, \Sigma Y) = \text{Hom}_{\mathcal{T}}(Y, \Sigma X) = 0$;
- (2) *for any morphism $\phi \in \text{Hom}_{\mathcal{T}}(U, X)$ and any morphism $\psi \in \text{Hom}_{\mathcal{T}}(X, Y)$, the composition $\psi\phi$ vanishes;*
- (3) *for any morphism $\phi' \in \text{Hom}_{\mathcal{T}}(U, Y)$ and any morphism $\psi' \in \text{Hom}_{\mathcal{T}}(Y, X)$, the composition $\psi'\phi'$ vanishes.*

Then $\text{Cyl}_X \text{Cyl}_Y U$ is isomorphic to $\text{Cyl}_Y \text{Cyl}_X U$.

Proof. Let

$$\Sigma^{-1}X^d \rightarrow \text{Cyl}_X U \xrightarrow{x} U \xrightarrow{f} X^d \quad \text{and} \quad \Sigma^{-1}Y^e \rightarrow \text{Cyl}_Y U \xrightarrow{y} U \xrightarrow{g} Y^e$$

be the triangles defining $\text{Cyl}_X U$ and $\text{Cyl}_Y U$. Applying $\text{Hom}_{\mathcal{T}}(-, Y)$ to the first triangle, we get the exact sequence

$$\text{Hom}_{\mathcal{T}}(X^d, Y) \rightarrow \text{Hom}_{\mathcal{T}}(U, Y) \xrightarrow{x^*} \text{Hom}_{\mathcal{T}}(\text{Cyl}_X U, Y) \rightarrow \text{Hom}_{\mathcal{T}}(\Sigma^{-1}X^d, Y).$$

By (2), the leftmost morphism vanishes. By (1), the space $\text{Hom}_{\mathcal{T}}(\Sigma^{-1}X^d, Y)$ vanishes. Therefore $\text{Hom}_{\mathcal{T}}(U, Y) \xrightarrow{x^*} \text{Hom}_{\mathcal{T}}(\text{Cyl}_X U, Y)$ is an isomorphism.

By applying $\text{Hom}_{\mathcal{T}}(-, X)$ to the second triangle above, and using (3) instead of (2), we get that $\text{Hom}_{\mathcal{T}}(U, X) \xrightarrow{y^*} \text{Hom}_{\mathcal{T}}(\text{Cyl}_Y U, X)$ is also an isomorphism.

Since x^* is an isomorphism, we get that (g_1x, \dots, g_ex) is a basis of $\text{Hom}_{\mathcal{T}}(\text{Cyl}_X U, Y)$, and so the defining triangle for $\text{Cyl}_Y \text{Cyl}_X U$ is isomorphic to

$$\Sigma^{-1}Y^e \rightarrow \text{Cyl}_Y \text{Cyl}_X U \rightarrow \text{Cyl}_X U \xrightarrow{gx} Y^e.$$

Similarly, since y^* is an isomorphism, we get that the defining triangle for $\text{Cyl}_X \text{Cyl}_Y U$ is isomorphic to

$$\Sigma^{-1}X^d \rightarrow \text{Cyl}_X \text{Cyl}_Y U \rightarrow \text{Cyl}_Y U \xrightarrow{fy} X^d.$$

Finally, applying the octahedral axiom to the composition gx yields an octahedron

Thus we have a triangle

$$\Sigma^{-1}X^d \rightarrow \text{Cyl}_Y \text{Cyl}_X U \rightarrow \text{Cyl}_Y U \xrightarrow{fy} X^d.$$

Comparing with the previous triangle (whose last morphism was also fy), we get that the objects $\text{Cyl}_X \text{Cyl}_Y U$ and $\text{Cyl}_Y \text{Cyl}_X U$ are isomorphic. \square

4.3. Rigid cylinders. The second key lemma in the proof of Theorem 4.1 is a condition for cylinders of rigid objects to be rigid in the category $K^{[-1,0]}(\text{proj } \Lambda)$.

Lemma 4.5 (Rigid cylinders). *Let H be an indecomposable object of $K^{[-1,0]}(\text{proj } \Lambda)$ such that $\text{Hom}_{K^b(\text{proj } \Lambda)}(H, \Sigma H)$ is one-dimensional, and let U be another object of $K^{[-1,0]}(\text{proj } \Lambda)$ satisfying the following:*

- (1) U is rigid, that is, $\text{Hom}_{K^b(\text{proj } \Lambda)}(U, \Sigma U) = 0$;
- (2) $\text{Hom}_{K^b(\text{proj } \Lambda)}(H, \Sigma U) = 0$;
- (3) for any non-zero $g \in \text{Hom}_{\mathcal{D}\Lambda}(\Sigma H, \nu H)$ the induced morphism

$$\text{Hom}_{K^b(\text{proj } \Lambda)}(U, \Sigma H) \xrightarrow{g*} \text{Hom}_{\mathcal{D}\Lambda}(U, \nu H)$$

is injective, where $\nu = - \otimes_{\Lambda}^L D\Lambda$ is the Nakayama functor.

Then $\text{Cyl}_{\Sigma H} U$ is in $K^{[-1,0]}(\text{proj } \Lambda)$ and also satisfies (1), (2) and (3).

Proof. The triangle $H^d \rightarrow \text{Cyl}_{\Sigma H} U \rightarrow U \xrightarrow{f} \Sigma H^d$ shows that the object $\text{Cyl}_{\Sigma H} U$ is in $K^{[-1,0]}(\text{proj } \Lambda)$, since this category is closed under extensions. Moreover, it induces the following commutative diagram with exact rows and columns (where we write (A, B) instead of $\text{Hom}_{K^b(\text{proj } \Lambda)}(A, B)$).

$$\begin{array}{ccccccccc}
& & & & & & (H^d, U) & \longrightarrow & (\Sigma^{-1}U, U) \\
& & & & & & \downarrow & & \downarrow \\
(\Sigma H^d, \Sigma H^d) & \longrightarrow & (U, \Sigma H^d) & \longrightarrow & (\text{Cyl}_{\Sigma H} U, \Sigma H^d) & \longrightarrow & (H^d, \Sigma H^d) & \longrightarrow & (\Sigma^{-1}U, \Sigma H^d) \\
\downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
(\Sigma H^d, \Sigma \text{Cyl}_{\Sigma H} U) & \longrightarrow & (U, \Sigma \text{Cyl}_{\Sigma H} U) & \longrightarrow & (\text{Cyl}_{\Sigma H} U, \Sigma \text{Cyl}_{\Sigma H} U) & \longrightarrow & (H^d, \Sigma \text{Cyl}_{\Sigma H} U) & \longrightarrow & (\Sigma^{-1}U, \Sigma \text{Cyl}_{\Sigma H} U) \\
\downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
(\Sigma H^d, \Sigma U) & \longrightarrow & (U, \Sigma U) & \longrightarrow & (\text{Cyl}_{\Sigma H} U, \Sigma U) & \longrightarrow & (H^d, \Sigma U) & \longrightarrow & (\Sigma^{-1}U, \Sigma U)
\end{array}$$

All spaces in the rightmost column vanish. Indeed, the top one vanishes by condition (1), and all others vanish since for any pair objects A and B in $K^{[-1,0]}(\text{proj } \Lambda)$, the space $(\Sigma^{-1}A, \Sigma B)$ vanishes. Moreover, the spaces $(U, \Sigma U)$ and $(H^d, \Sigma U)$ of the bottom row vanish by conditions (1) and (2). Therefore, the space $(\text{Cyl}_{\Sigma H} U, \Sigma U)$ vanishes as well.

By construction, any morphism from U to ΣH factors through f . Thus the map $(\Sigma H^d, \Sigma H^d) \rightarrow (U, \Sigma H^d)$ in the second row is surjective, and so the one immediately to its right vanishes. This implies that $(\text{Cyl}_{\Sigma H} U, \Sigma H^d) \rightarrow (H^d, \Sigma H^d)$ is an isomorphism.

Using the above information, the commutative diagrams simplifies considerably.

$$\begin{array}{ccccccccc}
& & & & & & (H^d, U) & \longrightarrow & 0 \\
& & & & & & \downarrow & & \\
(\Sigma H^d, \Sigma H^d) & \longrightarrow & \twoheadrightarrow & (U, \Sigma H^d) & \xrightarrow{0} & (\text{Cyl}_{\Sigma H} U, \Sigma H^d) & \xrightarrow{\sim} & (H^d, \Sigma H^d) & \longrightarrow & 0 \\
\downarrow & & & \downarrow & & \downarrow & & \downarrow & & \\
(\Sigma H^d, \Sigma \text{Cyl}_{\Sigma H} U) & \longrightarrow & & (U, \Sigma \text{Cyl}_{\Sigma H} U) & \longrightarrow & (\text{Cyl}_{\Sigma H} U, \Sigma \text{Cyl}_{\Sigma H} U) & \longrightarrow & (H^d, \Sigma \text{Cyl}_{\Sigma H} U) & \longrightarrow & 0 \\
\downarrow & & & \downarrow & & \downarrow & & \downarrow & & \\
(\Sigma H^d, \Sigma U) & & & 0 & & 0 & & 0 & &
\end{array}$$

We are now ready to prove that $\text{Cyl}_{\Sigma H} U$ satisfies properties (1) to (3).

Proof of (2). To prove that $(H, \Sigma \text{Cyl}_{\Sigma H} U)$ vanishes, it suffices to prove that the map $(H^d, U) \xrightarrow{f_*} (H^d, \Sigma H^d)$ in the rightmost non-zero column is surjective. Applying the duality D , this is equivalent to showing that the map $D(H^d, \Sigma H^d) \xrightarrow{Df_*} D(H^d, U)$ is injective. Applying the properties of the Nakayama functor ν , this is equivalent to proving that the map $(\Sigma H^d, \nu H^d) \xrightarrow{f^*} (U, \nu H^d)$ is injective.

Now, $(H, \Sigma H)$ is one dimensional, and therefore, so is $(\Sigma H, \nu H)$. Let g be a non-zero element in $(\Sigma H, \nu H)$; it is unique up to rescaling. An element ϕ of $(\Sigma H^d, \nu H^d)$

can be viewed as a $d \times d$ matrix with multiples of g as entries, say $\phi = (\phi_{i,j}g)_{i,j}$. Then

$$f^*(\phi) = \phi f = \begin{bmatrix} \phi_{1,1}g & \cdots & \phi_{1,d}g \\ \vdots & \ddots & \vdots \\ \phi_{d,1}g & \cdots & \phi_{d,d}g \end{bmatrix} \begin{bmatrix} f_1 \\ \vdots \\ f_d \end{bmatrix} = \begin{bmatrix} \sum_{i=1}^d \phi_{1,i}gf_i \\ \vdots \\ \sum_{i=1}^d \phi_{d,i}gf_i \end{bmatrix} = \begin{bmatrix} g(\sum_{i=1}^d \phi_{1,i}f_i) \\ \vdots \\ g(\sum_{i=1}^d \phi_{d,i}f_i) \end{bmatrix}.$$

Thus $f^*(\phi)$ vanishes if and only if $g(\sum_{i=1}^d \phi_{j,i}f_i)$ vanishes for all $j \in \{1, \dots, d\}$. By condition (3), this only happens if $(\sum_{i=1}^d \phi_{j,i}f_i)$ vanishes for all j . Since the f_i are linearly independent, this is only possible if all $\phi_{i,j}$ vanish. Thus $f^*(\phi) = 0$ implies that $\phi = 0$, and f^* is injective. By the above, this shows that the map $(H^d, U) \xrightarrow{f^*} (H^d, \Sigma H^d)$ is surjective, and so $(H, \Sigma \text{Cyl}_{\Sigma H} U)$ vanishes.

Proof of (3). As above, let g be a non-zero element of $(\Sigma H, \nu H)$ (it is unique up to rescaling). There is a bifunctorial non-degenerate bilinear form

$$(-, -) : (A, B) \times (B, \nu A) \rightarrow k,$$

where A, B can be any objects of $K^b(\text{proj } \Lambda)$ (see, for instance, [Hap88]). In particular, if h is a non-zero element of $(H, \Sigma H)$, then $(h, g) \neq 0$, since $(H, \Sigma H)$ is one-dimensional and the form is non-degenerate. But the bifunctoriality of the bilinear form implies that $(h, g) = (id_H, gh)$. Thus $gh \neq 0$. This implies that the map $(H, \Sigma H) \xrightarrow{g^*} (H, \nu H)$ is injective.

We wish to prove that $(\text{Cyl}_{\Sigma H} U, \Sigma H) \xrightarrow{g^*} (\text{Cyl}_{\Sigma H} U, \nu H)$ is injective. Consider the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} (\Sigma H^d, \Sigma H) & \xrightarrow{f^*} & (U, \Sigma H) & \xrightarrow{0} & (\text{Cyl}_{\Sigma H} U, \Sigma H) & \hookrightarrow & (H^d, \Sigma H) \\ \downarrow & & \downarrow & & \downarrow g_* & & \downarrow g_* \\ (\Sigma H^d, \nu H) & \xrightarrow{f^*} & (U, \nu H) & \longrightarrow & (\text{Cyl}_{\Sigma H} U, \nu H) & \longrightarrow & (H^d, \nu H) \end{array}$$

The uppermost, leftmost morphism is surjective since all morphisms from U to ΣH factor through f ; thus the upper middle map is zero and the upper, rightmost map is injective. The rightmost vertical morphism is injective by the above. Thus the map $(\text{Cyl}_{\Sigma H} U, \Sigma H) \xrightarrow{g^*} (\text{Cyl}_{\Sigma H} U, \nu H)$ is injective.

Proof of (1). As a consequence of (2) proved above, we get a commutative square

$$\begin{array}{ccc} (U, \Sigma H^d) & \xrightarrow{0} & (\text{Cyl}_{\Sigma H} U, \Sigma H^d) \\ \downarrow & & \downarrow \\ (U, \Sigma \text{Cyl}_{\Sigma H} U) & \longrightarrow & (\text{Cyl}_{\Sigma H} U, \Sigma \text{Cyl}_{\Sigma H} U) \end{array}$$

where the top map is zero and the left and bottom maps are surjective. Thus the composition of the top and right morphism is both zero and surjective. Thus the space $(\text{Cyl}_{\Sigma H} U, \Sigma \text{Cyl}_{\Sigma H} U)$ vanishes. \square

4.4. g-vectors of cylinders.

Lemma 4.6. *Let H be an indecomposable object of $K^{[-1,0]}(\text{proj } \Lambda)$ such that $(H, \Sigma H)$ is one-dimensional, and let U be another object of $K^{[-1,0]}(\text{proj } \Lambda)$. Then for any $m \in \mathbb{Z}_{>0}$, the object $\text{Cyl}_{\Sigma H}^m U$ is in $K^{[-1,0]}(\text{proj } \Lambda)$, and if $d = \dim(U, \Sigma H)$, then*

$$[\text{Cyl}_{\Sigma H}^m U] = [U] + md[H].$$

Proof. The triangle $H^d \rightarrow \text{Cyl}_{\Sigma H} U \rightarrow U \rightarrow \Sigma H^d$ implies that the object $\text{Cyl}_{\Sigma H} U$ is in $K^{[-1,0]}(\text{proj } \Lambda)$, since this category is closed under extensions. It also implies that $[\text{Cyl}_{\Sigma H} U] = [U] + d[H]$. Applying the functor $(-, \Sigma H)$, we get an exact sequence

$$(\Sigma H^d, \Sigma H) \twoheadrightarrow (U, \Sigma H) \xrightarrow{0} (\text{Cyl}_{\Sigma H} U, \Sigma H) \longrightarrow (H^d, \Sigma H) \longrightarrow (\Sigma^{-1}U, \Sigma H).$$

Since $(\Sigma^{-1}U, \Sigma H) = 0$, we deduce that $(\text{Cyl}_{\Sigma H} U, \Sigma H)$ and $(H^d, \Sigma H)$ are isomorphic. Thus $\dim(\text{Cyl}_{\Sigma H} U, \Sigma H) = d$. From there, we apply induction on m to get the desired equality. \square

Lemma 4.7. *Under the hypotheses of Lemma 4.4, if $d_X = \dim(U, X)$ and $d_Y = \dim(U, Y)$, then*

$$[\text{Cyl}_X \text{Cyl}_Y U] = [U] - d_X[X] - d_Y[Y] \in K_0(\mathcal{T}).$$

Proof. In the proof of Lemma 4.4, we have obtained triangle $\Sigma^{-1}X^{d_X} \rightarrow \text{Cyl}_X U \rightarrow U \rightarrow X^{d_X}$ and $\Sigma^{-1}Y^{d_Y} \rightarrow \text{Cyl}_Y \text{Cyl}_X U \rightarrow \text{Cyl}_X U \xrightarrow{g_X} Y^{d_Y}$. It follows from them that $[\text{Cyl}_X U] = [U] - d_X[X]$ and $[\text{Cyl}_Y \text{Cyl}_X U] = [\text{Cyl}_X U] - d_Y[Y]$. Since the object $\text{Cyl}_Y \text{Cyl}_X U$ is isomorphic to $\text{Cyl}_X \text{Cyl}_Y U$ by Lemma 4.4, we get the desired result. \square

Lemma 4.8. *Let Λ be any finite-dimensional algebra, and let H_1, \dots, H_s be objects of $K^{[-1,0]}(\text{proj } \Lambda)$ such that*

- for each $i \in \{1, \dots, s\}$, the object $H = H_i$ satisfies the hypotheses of Lemma 4.6;
- for each pair of distinct $i, j \in \{1, \dots, s\}$, the objects $X = \Sigma H_i$ and $Y = \Sigma H_j$ satisfy the hypotheses of Lemma 4.4.

Let U be an object of $K^{[-1,0]}(\text{proj } \Lambda)$, $d_i = \dim(U, \Sigma H_i)$, and $a_1, \dots, a_s \in \mathbb{Z}_{>0}$. Then $\text{Cyl}_{\Sigma H_s}^{a_s} \cdots \text{Cyl}_{\Sigma H_1}^{a_1} U$ is an object of $K^{[-1,0]}(\text{proj } \Lambda)$, and

$$[\text{Cyl}_{\Sigma H_s}^{a_s} \cdots \text{Cyl}_{\Sigma H_1}^{a_1} U] = [U] + \sum_{i=1}^s a_i d_i [H_i].$$

Proof. Apply Lemmas 4.6 and 4.7, and induction on s . \square

4.5. Proof of density. We now turn to the proof of Theorem 4.1. Assume that Λ is tame. Let $\mathbf{g} \in \mathbb{Q}^n$. To prove that \mathbf{g} is in the closure $\bar{\mathcal{F}}_{2\text{-silt}}^{\mathbf{g}}(\Lambda)$, it suffices to prove that it is true for a positive scalar multiple of \mathbf{g} . Thus, up to positive rescaling, we can assume that $\mathbf{g} \in \mathbb{Z}^n$.

We apply Theorem 3.8 to \mathbf{g} , and let

$$\mathbf{g} = a_1 \mathbf{g}_1 + \dots + a_r \mathbf{g}_r + b_1 \mathbf{h}_1 + \dots + b_s \mathbf{h}_s$$

be the generic decomposition of \mathbf{g} . Here, we have grouped the terms so that $i \neq j$ implies $\mathbf{g}_i \neq \mathbf{g}_j$ and $\mathbf{h}_i \neq \mathbf{h}_j$, with all a_i and b_i in $\mathbb{Z}_{>0}$.

If $s = 0$ (that is, if no term of the form \mathbf{h}_i appears in the generic decomposition of \mathbf{g}), then $\mathbf{g} \in \mathcal{F}_{2\text{-silt}}^{\mathbf{g}}(\Lambda)$ and there is nothing to prove. Assume then that $s > 0$. Note that, by [CILFS15, Theorem 6.1], this implies that $r < n$.

For any $i \in \{1, \dots, s\}$, let H_i be a generic object with \mathbf{g} -vector \mathbf{h}_i .

Lemma 4.9. *The objects H_1, \dots, H_s satisfy the hypotheses of Lemma 4.8.*

Proof. To prove that each H_i satisfies the hypotheses of Lemma 4.6, one simply notes that $\dim(H_i, \Sigma H_i)$ by Corollary 3.9(1).

Let us prove that the hypotheses of Lemma 4.4 are satisfied with $X = \Sigma H_i$ and $Y = \Sigma H_j$. Condition (1) is satisfied since \mathbf{h}_i and \mathbf{h}_j are two distinct terms in the generic decomposition of \mathbf{g} .

To prove condition (2), recall from Corollary 3.9(3) that any morphism from H_i to H_j factors through $\text{add}(\Sigma\Lambda)$. Thus any morphism from ΣH_i to ΣH_j factors through $\text{add}(\Sigma^2\Lambda)$, which implies that precomposing it with a morphism from U to ΣH_i will give zero. Thus (2) is satisfied.

The proof of condition (3) of Lemma 4.4 is similar. \square

Let G be a presilting object whose \mathbf{g} -vector is $a_1\mathbf{g}_1 + \dots + a_r\mathbf{g}_r$. Let G' be its *Bongartz co-completion*, defined by the triangle

$$\Lambda \rightarrow G'' \rightarrow G' \rightarrow \Sigma\Lambda$$

where the left-most morphism is a left $\text{add } G$ -approximation of Λ . By the dual of [Jas15, Definition-Proposition 4.9], the object $G' \oplus G$ is a silting object in the category $K^{[-1,0]}(\text{proj } \Lambda)$.

Lemma 4.10. *Taking $U = G'$ and $H = H_i$ (for $i \in \{1, \dots, s\}$), conditions (1) to (3) of Lemma 4.5 are satisfied. Moreover, the space $\text{Hom}_{K^b(\text{proj } \Lambda)}(G', \Sigma H_i)$ does not vanish.*

Proof. That (1) is true follows from the fact that G' is presilting. Applying the functor $\text{Hom}_{K^b(\text{proj } \Lambda)}(H_i, -)$ to the triangle defining G' , we get an exact sequence

$$\text{Hom}_{K^b(\text{proj } \Lambda)}(H_i, \Sigma G'') \rightarrow \text{Hom}_{K^b(\text{proj } \Lambda)}(H_i, \Sigma G') \rightarrow \text{Hom}_{K^b(\text{proj } \Lambda)}(H_i, \Sigma^2\Lambda).$$

The left term vanishes since $\text{Hom}_{K^b(\text{proj } \Lambda)}(H_i, \Sigma G)$ does, and the right term vanishes since Λ and H_i are in $K^{[-1,0]}(\text{proj } \Lambda)$. Thus $\text{Hom}_{K^b(\text{proj } \Lambda)}(H_i, \Sigma G') = 0$, and condition (2) is true.

To prove condition (3), let g be a non-zero morphism from ΣH_i to νH_i . Then the map $(\Sigma\Lambda, \Sigma H_i) \xrightarrow{g^*} (\Sigma\Lambda, \nu H_i)$ is injective. Indeed, note that taking the cohomology in degree -1 yields isomorphisms

$$\text{Hom}_{K^b(\text{proj } \Lambda)}(\Sigma\Lambda, \Sigma H_i) \cong \text{Hom}_{\Lambda}(\Lambda, H^0 H_i)$$

and

$$\text{Hom}_{K^b(\text{proj } \Lambda)}(\Sigma\Lambda, \nu H_i) \cong \text{Hom}_{\Lambda}(\Lambda, H^{-1}\nu H_i).$$

But by Theorem 3.8(2), $H^0 H_i$ and $H^{-1}\nu H_i \cong \tau H^0 H_i$ are isomorphic as Λ -modules. Moreover, the morphism g induces a non-zero morphism $H^{-1}g : H^0 H_i \rightarrow H^{-1}\nu H_i$, which must thus be an isomorphism since these two isomorphic modules are bricks.

Therefore, if $f : \Sigma\Lambda \rightarrow \Sigma H_i$ is a morphism such that $gf = 0$, then $H^{-1}g \circ H^{-1}f = 0$, and since $H^{-1}g$ is an isomorphism, then $H^{-1}f = 0$, and so $f = 0$ by the above.

Consider now the commutative diagram with exact rows

$$\begin{array}{ccccc}
(\Sigma\Lambda, \Sigma H_i) & \longrightarrow & (G', \Sigma H_i) & \longrightarrow & (G'', \Sigma H_i) \\
\downarrow g_* & & \downarrow g_* & & \\
(\Sigma G'', \nu H_i) & \longrightarrow & (\Sigma\Lambda, \nu H_i) & \longrightarrow & (G', \nu H_i)
\end{array}$$

Now, $(G'', \Sigma H_i)$ vanishes since $(G, \Sigma H_i)$ does, and $(\Sigma G'', \nu H_i)$ is isomorphic to the space $D(H_i, \Sigma G'')$, which also vanishes. Thus the bottom middle map is injective and the top middle map is surjective. The left vertical map is injective by the above. This implies that the right vertical map is injective, so condition (3) is proved. This also implies that the top middle map has to be injective, and is thus an isomorphism. Thus $(G', \Sigma H_i)$ is isomorphic to $(\Sigma\Lambda, \Sigma H_i)$, which is non-zero. \square

Lemma 4.11. *Let U be an object of $K^{[-1,0]}(\text{proj } \Lambda)$ and $i \in \{1, \dots, s\}$. If $(G, \Sigma U) = (U, \Sigma G) = 0$, then $(G, \Sigma \text{Cyl}_{\Sigma H_i} U) = (\text{Cyl}_{\Sigma H_i} U, \Sigma G) = 0$.*

Proof. Simply apply $(G, -)$ and $(-, \Sigma G)$ to the triangle

$$H_i^d \rightarrow \text{Cyl}_{\Sigma H_i} U \rightarrow U \rightarrow \Sigma H_i^d.$$

\square

We can now prove that the $\mathbf{g} \in \mathbb{Z}^n$ chosen at the begining of this section is in $\bar{\mathcal{F}}_{2\text{-silt}}^{\mathbf{g}}(\Lambda)$, thus finishing the proof of Theorem 4.1. For each $i \in \{1, \dots, s\}$, let $d_i = \dim(G', \Sigma H_i)$. By Lemma 4.10, the d_i are non-zero. Let $d = \prod_{i=1}^s d_i$, and let $e_i = \frac{d}{d_i}$ for each i .

The, combining Lemmas 4.10, 4.5, 4.9 and 4.4, we get that

$$\text{Cyl}_{\Sigma H_s}^{b_s e_s} \cdots \text{Cyl}_{\Sigma H_1}^{b_1 e_1} G'$$

is a presilting object in $K^{[-1,0]}(\text{proj } \Lambda)$. Moreover,

$$G^{\oplus d} \oplus \text{Cyl}_{\Sigma H_s}^{b_s e_s} \cdots \text{Cyl}_{\Sigma H_1}^{b_1 e_1} G'$$

is presilting by Lemma 4.11. Now, using Lemma 4.8, we get that

$$\begin{aligned}
[G^{\oplus d} \oplus \text{Cyl}_{\Sigma H_s}^{b_s e_s} \cdots \text{Cyl}_{\Sigma H_1}^{b_1 e_1} G'] &= d[G] + [G'] + \sum_{i=1}^s b_i e_i d_i [H_i] \\
&= d([G] + \sum_{i=1}^s b_i [H_i]) + [G'] \\
&= d\mathbf{g} + [G'].
\end{aligned}$$

Similarly, for any $m \in \mathbb{Z}_{>0}$, we have that

$$G^{\oplus dm} \oplus \text{Cyl}_{\Sigma H_s}^{mb_s e_s} \cdots \text{Cyl}_{\Sigma H_1}^{mb_1 e_1} G'$$

is a presilting object whose \mathbf{g} -vector is $m d\mathbf{g} + [G']$.

The vector \mathbf{g} is in the line at the limit of those generated by $m d\mathbf{g} + [G']$ as m goes to infinity. Since each $m d\mathbf{g} + [G']$ is the \mathbf{g} -vector of a presilting objects, these vectors are in the fan $\mathcal{F}_{2\text{-silt}}^{\mathbf{g}}(\Lambda)$. Thus $\mathbf{g} \in \bar{\mathcal{F}}_{2\text{-silt}}^{\mathbf{g}}(\Lambda)$. This finishes the proof of Theorem 4.1.

5. DENSITY OF \mathbf{g} -VECTOR FANS FOR CLUSTER ALGEBRAS AND JACOBIAN ALGEBRAS

In this section, we apply our results above to the \mathbf{g} -vector fans arising from the theory of cluster algebras.

5.1. \mathbf{g} -vectors for cluster algebras and Jacobian algebras. We first recall the definition of \mathbf{g} -vectors for cluster algebras [FZ07].

Definition 5.1 (Proposition 6.6 of [FZ07]). Let Q be a quiver without loops and 2-cycles, and let $Q_0 = \{1, \dots, n\}$ be the set of its vertices. Let Q^{prin} be the quiver obtained by adding a vertex i' and an arrow $i' \rightarrow i$ for every vertex i of Q . The \mathbf{g} -vectors for the cluster algebras of type Q are obtained by the following mutation rule:

- (1) $(Q^{\text{prin}}, (\mathbf{e}_1, \dots, \mathbf{e}_n))$ is a \mathbf{g} -vector seed, where \mathbf{e}_i is the elementary vector of \mathbb{Z}^n in coordinate i ;
- (2) if $(R, (\mathbf{g}_1, \dots, \mathbf{g}_n))$ is a \mathbf{g} -vector seed, then for any vertex $k \in \{1, \dots, n\}$, the mutation $\mu_k(R, (\mathbf{g}_1, \dots, \mathbf{g}_n)) = (R', (\mathbf{g}'_1, \dots, \mathbf{g}'_n))$ is also a \mathbf{g} -vector seed, where

$$\mathbf{g}'_\ell = \begin{cases} \mathbf{g}_\ell & \text{if } \ell \neq k; \\ -\mathbf{g}_k + \sum_{i=1}^n [b_{i,k}]_+ \mathbf{g}_i - \sum_{j=1}^n [b_{j,k}]_+ \begin{pmatrix} b_{1,j} \\ b_{2,j} \\ \vdots \\ b_{n,j} \end{pmatrix} & \text{if } \ell = k, \end{cases}$$

where $b_{i,j} = \#\{\text{arrows } i \rightarrow j \text{ in } R\} - \#\{\text{arrows } j \rightarrow i \text{ in } R\}$, and where $[z]_+ = \max(z, 0)$ for any real number z .

The vectors \mathbf{g}_i that appear in any \mathbf{g} -vector seed obtained by successive mutations of the initial seed $(Q^{\text{prin}}, (\mathbf{e}_1, \dots, \mathbf{e}_n))$ are the \mathbf{g} -vectors for the cluster algebra of type Q ; \mathbf{g} -vectors belonging to a common seed are *compatible*.

The compatibility relation for \mathbf{g} -vectors allows one to organize them in a fan; this follows from the proof of conjectures in [FZ07].


Theorem 5.2 (Consequence of Theorem 1.7 of [DWZ10]). *Let Q be a quiver without loops and 2-cycles. Then the \mathbf{g} -vectors of cluster variables of $\mathcal{A}(Q)$ are the rays of a simplicial polyhedral fan whose maximal cones are generated by sets of compatible \mathbf{g} -vectors.*

Definition 5.3. The simplicial polyhedral fan described in Theorem 5.2 is called the *cluster \mathbf{g} -vector fan* associated with Q , and it is denoted by $\mathcal{F}_{\text{cluster}}^{\mathbf{g}}(Q)$. We also denote by $\bar{\mathcal{F}}_{\text{cluster}}^{\mathbf{g}}(Q)$ the closure of the union of the cones of $\mathcal{F}_{\text{cluster}}^{\mathbf{g}}(Q)$.

Example 5.4. Let $m \in \mathbb{Z}_{\geq 1}$ and K_m be an m -Kronecker quiver, that is,

$$K_m := [1 \xrightarrow{\quad \vdots \quad} 2],$$

where there are m arrows between vertices 1 and 2. In particular, K_1 is of type A_2 and K_2 is a Kronecker quiver. The cluster \mathbf{g} -vector fan $\mathcal{F}_{\text{cluster}}^{\mathbf{g}}(K_m)$ is well known and given as in Figure 1. Here $\bar{\mathcal{F}}_{\text{cluster}}^{\mathbf{g}}(K_m)$ contains infinitely many rays converging to the rays r_{\pm} of slope $(-m \pm \sqrt{m^2 - 4})/2$ for $m \geq 2$. If $m = 2$, then

FIGURE 1. Cluster \mathbf{g} -vector fans $\mathcal{F}_{\text{cluster}}^{\mathbf{g}}(K_m)$

$r_+ = r_-$. If $m \geq 3$, then $r_+ \neq r_-$ and the interior of the cone spanned by r_+ and r_- is the complement of $\bar{\mathcal{F}}_{\text{cluster}}^{\mathbf{g}}(K_m)$.

Second, we recall Jacobian algebras [DWZ08]. A *potential* W of Q is a (possibly infinite) linear combination of cycles in Q . The pair (Q, W) is called a *quiver with potential*. For a cycle $\alpha_1 \cdots \alpha_m$ in Q and an arrow α of Q , we define

$$\partial_{\alpha}(\alpha_1 \cdots \alpha_m) := \sum_{i: \alpha_i = \alpha} \alpha_{i+1} \cdots \alpha_m \alpha_1 \cdots \alpha_{i-1}.$$

By linearity, this defines the cyclic derivative $\partial_{\alpha}(W)$ of a potential W of Q . For a quiver with potential (Q, W) , the *Jacobian algebra* $J(Q, W)$ is a quotient algebra of the complete path algebra of Q by the closure of the ideal generated by the set $\{\partial_{\alpha}W \mid \alpha \in Q_1\}$. We say that a potential W of Q is *Jacobi-finite* if $J(Q, W)$ is finite-dimensional.

The *mutation* $\mu_k(Q, W)$ in direction k is defined as an analogue of mutations of quivers. We refer to [DWZ08] for details. A potential W of Q is called *non-degenerate* if every quiver with potential obtained from (Q, W) by any sequence of mutations has no 2-cycles. In this case, we have $\mu_k(Q, W) = (\mu_k Q, W')$ for a non-degenerate potential W' of $\mu_k Q$. If W is Jacobi-finite, then so is W' [DWZ08]. In this section, we mainly study non-degenerate Jacobi-finite potentials.

On the other hand, mutations are defined for 2-term sifting complexes in [IY08]. Let $2\text{-silt}^+ J(Q, W)$ (resp., $2\text{-silt}^- J(Q, W)$) be a subset of $2\text{-silt} J(Q, W)$ consisting of objects obtained from $J(Q, W)$ (resp., $\Sigma J(Q, W)$) by all sequences of mutations. The Jacobian algebra $J(Q, W)$ associated with a quiver with non-degenerate Jacobi-finite potential (Q, W) gives a categorification of the associated cluster algebra $\mathcal{A}(Q)$. In particular, the following result is due to the authors [AIR14, Corollary 4.8], [FK10, Theorem 6.3] and [CIKLFP13, Corollary 3.5] (see also [Yur20, Theorem 4.4]).

Theorem 5.5 (Additive categorification of cluster algebras). *Let Q be a quiver without loops and 2-cycles. Let W be a non-degenerate Jacobi-finite potential of Q .*

- (1) *There is a bijection*

$$2\text{-silt}^+ J(Q, W) \leftrightarrow \{\mathbf{g}\text{-vector seeds of } Q\}$$

that sends $J(Q, W)$ to the initial \mathbf{g} -vector seed for Q and commutes with mutations. In particular, it preserves their cones of \mathbf{g} -vectors.

(2) *There is a bijection*

$$x_{(-)} : 2\text{-silt}^- J(Q, W) \leftrightarrow \{\mathbf{g}\text{-vector seeds of } Q^{\text{op}}\}$$

that sends $\Sigma J(Q, W)$ to the initial \mathbf{g} -vector seed for Q^{op} and commutes with mutations. In particular, the cone of \mathbf{g} -vectors associated to $U \in 2\text{-silt}^- J(Q, W)$ is the negative of the cone of \mathbf{g} -vectors associated to x_U .

Theorem 5.5 immediately implies that the set

$$2\text{-silt}^\pm J(Q, W) := 2\text{-silt}^+ J(Q, W) \cup 2\text{-silt}^- J(Q, W)$$

is independent of the choice of non-degenerate Jacobi-finite W .

Corollary 5.6. *Let Q be a quiver without loops and 2-cycles. Let W and W' be non-degenerate Jacobi-finite potentials of Q . Then there is a bijection*

$$2\text{-silt}^\pm J(Q, W) \leftrightarrow 2\text{-silt}^\pm J(Q, W')$$

that sends $J(Q, W)$ to $J(Q, W')$ (resp., $\Sigma J(Q, W)$ to $\Sigma J(Q, W')$) and commutes with mutations. In particular, it preserves their cones of \mathbf{g} -vectors.

5.2. Mutation-finite quivers. In the rest of this section, we fix a quiver Q without loops and 2-cycles. We say that Q is

- *mutation equivalent to Q'* if Q is obtained from Q' by a sequence of mutations;
- *mutation-finite* if there are only finitely many quivers mutation equivalent to Q .

Felikson, Shapiro and Tumarkin [FST12] classified mutation-finite quivers, see also [FST08, Section 12].

Theorem 5.7. [FST12, Theorem 6.1] *A mutation-finite quiver Q is one of the followings:*

- *an m -Kronecker quiver K_m with $m \geq 3$;*
- *a quiver defined from a triangulated surface (see [FST08]);*
- *a quiver mutation equivalent to one of the quivers E_i , \tilde{E}_i , $E_i^{(1,1)}$, X_6 and X_7 for $i \in \{6, 7, 8\}$ as in Figure 2.*

Remark 5.8. (1) The quivers K_m , E_i and \tilde{E}_i are acyclic.

- (2) The Jacobian algebras J and cluster categories associated with $E_i^{(1,1)}$, called *tubular quivers*, were studied in [BG12, BGJ13, GGS15]. In particular, it follows from [BKL10, Theorem 8.6] that any two silting objects of $K^{[-1,0]}(\text{proj } J)$ are connected by a sequence of mutations, see [BG12, Section 3].
- (3) Derksen and Owen [DO08] found the quivers X_6 and X_7 as new mutation-finite quivers.

Jacobian algebras associated with mutation-finite quivers satisfy some properties.

Theorem 5.9. [GLFS16] *A quiver Q is a mutation-finite one that is not mutation equivalent to one of the quivers X_6 , X_7 and K_m with $m \geq 3$ if and only if there is a non-degenerate Jacobi-finite potential W of Q such that $J(Q, W)$ is tame.*

Note that, for example, such potential W of acyclic quivers in Theorem 5.9 is zero. In other cases, such potentials W were given in [LF09, LF16] for quivers defined from triangulated surfaces; in [GGS15, Lad11] for tubular quivers.




FIGURE 2. Exceptional quivers

Proposition 5.10. *Suppose that Q is mutation-finite except for mutation equivalence classes of X_6 and X_7 . Let W be a non-degenerate Jacobi-finite potential of Q . If Q is not defined from a triangulated surface with exactly one puncture, then any two siltling objects in $K^{[-1,0]}(\text{proj } J(Q, W))$ are connected by a sequence of mutations, that is, $2\text{-silt } J(Q, W) = 2\text{-silt}^+ J(Q, W)$. Otherwise, $2\text{-silt } J(Q, W) = 2\text{-silt}^+ J(Q, W) \sqcup 2\text{-silt}^- J(Q, W)$.*

Proof. By the assumption and Theorem 5.7, Q is mutation equivalent to one of the following quivers: an acyclic quiver; a quiver defined from a triangulated surface; a tubular quiver. For each case, the assertions follow from [BMR⁺06, Proposition 3.5]; [Yur20, Corollary 1.4]; Remark 5.8(2). \square

Corollary 5.11. *Suppose that Q is mutation-finite except for mutation equivalence classes of X_6 and X_7 . Let W and W' be non-degenerate Jacobi-finite potentials of Q . Then there is a bijection*

$$2\text{-silt } J(Q, W) \leftrightarrow 2\text{-silt } J(Q, W')$$

that sends $J(Q, W)$ to $J(Q, W')$ and commutes with mutations. In particular, it preserves their cones of \mathbf{g} -vectors.

Proof. The assertion follows from Corollary 5.6 and Proposition 5.10. \square

5.3. \mathbf{g} -tame Jacobian algebras. In order to apply the categorification of cluster algebras to the study of the closure of their \mathbf{g} -vector fans, we need to study \mathbf{g} -vector fans for finite-dimensional Jacobian algebras.

Lemma 5.12. *If (Q, W) is a quiver with potential such that $J(Q, W)$ is \mathbf{g} -tame, then so is $\mu_k(Q, W)$ (if it is defined).*

Proof. The transformation of mutations for \mathbf{g} -vectors is piecewise linear. Thus it preserves \mathbf{g} -tameness. \square

We note that the proof of Lemma 5.12 applies even if the Jacobian algebra $J(Q, W)$ is infinite-dimensional.

We now give a near classification of quivers admitting a potential such that the corresponding Jacobian algebra is \mathbf{g} -tame.

Theorem 5.13. *Suppose that Q is not mutation equivalent to one of the quivers X_6 , X_7 and K_m with $m \geq 3$. Then Q admits a non-degenerate potential W such that $J(Q, W)$ is \mathbf{g} -tame if and only if it is mutation-finite. Moreover, a Jacobian algebra for the quiver K_m is not \mathbf{g} -tame for $m \geq 3$.*

Proof. For $Q = K_m$, the assertion follows from Example 5.4. Suppose that Q is not mutation equivalent to one of the quivers X_6 , X_7 and K_m . When Q is mutation-finite, then Q admits a potential W such that $J(Q, W)$ is \mathbf{g} -tame by Theorems 4.1 and 5.9. When Q is not mutation-finite, Q is mutation equivalent to a quiver Q' with full subquiver K_m for $m \geq 3$. Namely, there is an idempotent e of $J(Q', W)$ for any non-degenerate potential W of Q' such that $J(Q', W)/(e) \simeq J(K_m, 0) = kK_m$. Proposition 3.11 implies that $J(Q', W)$ is not \mathbf{g} -tame. Mutating back to Q , we obtain the result by Lemma 5.12. \square

Corollary 5.14. *Suppose that Q is not mutation equivalent to one of the quivers X_6 and X_7 . If $J(Q, W)$ is \mathbf{g} -tame for some non-degenerate potential W , then it is \mathbf{g} -tame for any non-degenerate Jacobi-finite potential.*

Proof. If $J(Q, W)$ is \mathbf{g} -tame, then by Theorem 5.13, Q is mutation-finite. The assertion then follows from Corollary 5.11. \square

5.4. Cluster algebras with dense \mathbf{g} -vector fans. We now wish to apply our knowledge of \mathbf{g} -vector fans for Jacobian algebras to those of cluster algebras. To this end, the following definition will be useful.

Definition 5.15. We say that Q is *cluster- \mathbf{g} -dense* if $\bar{\mathcal{F}}_{\text{cluster}}^{\mathbf{g}}(Q) = \mathbb{R}^n$. We say that Q is *half cluster- \mathbf{g} -dense* if $\bar{\mathcal{F}}_{\text{cluster}}^{\mathbf{g}}(Q)$ and $\bar{\mathcal{F}}_{\text{cluster}}^{\mathbf{g}}(Q^{op})$ are closed half-spaces in \mathbb{R}^n .

In the same as Lemme 5.12, (half) cluster- \mathbf{g} -denseness is mutation-invariant. In particular, it gives a new class of cluster algebras, called (half) \mathbf{g} -dense cluster algebras. It is clear that the associated finite-dimensional Jacobian algebras are \mathbf{g} -tame.

Lemma 5.16. *If Q is (half) cluster- \mathbf{g} -dense, then for any non-degenerate Jacobi-finite potential W , the algebra $J(Q, W)$ is \mathbf{g} -tame.*

Proof. The assertion follows from Theorem 5.5. \square

We give a near classification of (half) cluster- \mathbf{g} -dense quivers, the only open cases being the mutation equivalence classes of X_6 and X_7 . It is a stronger statement of Theorem 5.13.

Theorem 5.17. *Suppose that Q is not mutation equivalent to one of the quivers X_6 , X_7 and K_m with $m \geq 3$. Then Q is cluster- \mathbf{g} -dense or half cluster- \mathbf{g} -dense if and only if it is mutation-finite. In this case, Q is half cluster- \mathbf{g} -dense if and only if it is defined from a triangulated surface with exactly one puncture. On the other hand, K_m is not (half) cluster- \mathbf{g} -dense for $m \geq 3$.*

Proof. The assertions follow from Proposition 5.10, Theorems 5.5(1) and 5.13 if Q is mutation-finite except for one defined from a triangulated surface with exactly one puncture. In which case, it was proved in [Yur20, Theorem 1.2].

On the other hand, [Mul16, Theorem 33] implies that a quiver with full subquiver K_m for $m \geq 3$ is not (half) cluster- \mathbf{g} -dense. Thus it is given by the same way as the proof of Theorem 5.13 that non-mutation-finite quivers are not (half) cluster- \mathbf{g} -dense. \square

We conjecture that X_6 and X_7 should not be exceptions to Theorem 5.17. It is known that:

- (1) [Mil17] If Q is mutation equivalent to the quiver X_6 , then there is a cluster whose cone of \mathbf{g} -vectors is $\mathbb{R}_{\leq 0}^n$;
- (2) [Sev14] If Q is mutation equivalent to the quiver X_7 , then $\mathcal{F}_{\text{cluster}}^{\mathbf{g}}(Q)$ is contained in some open half-space in \mathbb{R}^n .

Therefore, the following seems natural.

Conjecture 5.18. (1) *The quiver X_6 is cluster- \mathbf{g} -dense.*
 (2) *The quiver X_7 is half cluster- \mathbf{g} -dense.*

Remark that the Jacobian algebras associated with X_6 and X_7 are wild [GLFS16]. Thus we cannot apply Theorem 4.1 for these classes.

The above allows us to give a partial converse to Lemma 5.16.

Corollary 5.19. *Suppose that Q is not mutation equivalent to one of the quivers X_6 , X_7 and K_m with $m \geq 3$. Then the following are equivalent:*

- (1) *Q is mutation-finite;*
- (2) *Q admits a non-degenerate potential W such that $J(Q, W)$ is \mathbf{g} -tame;*
- (3) *Q is cluster- \mathbf{g} -dense or half cluster- \mathbf{g} -dense.*

Moreover, K_m with $m \geq 3$ is mutation-finite, but not (half) cluster- \mathbf{g} -dense and the unique potential 0 on K_m is such that $J(K_m, 0)$ is not \mathbf{g} -dense.

Proof. The assertions follow from Theorems 5.13 and 5.17. \square

6. SCATTERING DIAGRAMS

Our work in this paper allows for a modest contribution to the question of the equivalence between the cluster and the stability scattering diagrams for (half) cluster- \mathbf{g} -dense quivers; our result will follow by using an argument due to L. Mou. The statement of the result, however, requires some recollections on scattering diagrams which will occupy us for the beginning of the section. We refer to [Bri17, GHKK18, Mou19] for scattering diagrams.

6.1. Consistent scattering diagrams. Let $N \simeq \mathbb{Z}^{\oplus n}$ be a free abelian group. Fix a basis (e_1, \dots, e_n) of N . We use the notations $M = \text{Hom}_{\mathbb{Z}}(N, \mathbb{Z})$, $M_{\mathbb{R}} = M \otimes_{\mathbb{Z}} \mathbb{R}$,

$$N^{\oplus} := \left\{ \sum_{i=1}^n a_i e_i \mid a_i \in \mathbb{Z}_{\geq 0} \right\} \quad \text{and} \quad N^+ := N^{\oplus} \setminus \{0\},$$

and consider an N^+ -graded Lie algebra

$$\mathfrak{g} = \bigoplus_{d \in N^+} \mathfrak{g}_d.$$

First, we define \mathfrak{g} -scattering diagrams for \mathfrak{g} with finite support, that is,

$$\#\text{Supp}(\mathfrak{g}) < \infty, \quad \text{where} \quad \text{Supp}(\mathfrak{g}) := \{d \in N^+ \mid \mathfrak{g}_d \neq 0\}.$$

In this case, \mathfrak{g} is nilpotent and there is a unipotent algebraic group G such that $\exp : \mathfrak{g} \rightarrow G$ is a bijection.

For a cone $\sigma \subset M_{\mathbb{R}}$, there is a Lie subalgebra

$$\mathfrak{g}_{\sigma} := \bigoplus_{d \in N \cap \sigma^{\perp}} \mathfrak{g}_d \subset \mathfrak{g},$$

where $\sigma^{\perp} := \{d \in N \mid m(d) = 0 \text{ for any } m \in \sigma\}$.

Let P be a finite subset of N^+ . For a partition $P = P_+ \sqcup P_0 \sqcup P_-$ with $P_0 \neq \emptyset$, there is a cone

$$\{m \in M_{\mathbb{R}} \mid m(P_+) > 0, m(P_0) = 0, m(P_-) < 0\} \subset M_{\mathbb{R}}.$$

The set of all such cones forms a polyhedral fan \mathfrak{S}_P in $M_{\mathbb{R}}$. Cones with codimension one are called *walls*. We denote by $\mathcal{W}(\mathfrak{S})$ the set of walls of \mathfrak{S} .

Definition 6.1. Suppose that \mathfrak{g} has a finite support $S = \text{Supp}(\mathfrak{g})$. A \mathfrak{g} -scattering diagram is a pair $\mathfrak{D} = (\mathfrak{S}_S, \phi_{\mathfrak{D}})$ with function $\phi_{\mathfrak{D}} : \mathcal{W}(\mathfrak{S}_S) \rightarrow G$ such that for $\sigma \in \mathcal{W}(\mathfrak{S}_S)$, $\phi_{\mathfrak{D}}(\sigma) \in \exp(\mathfrak{g}_{\sigma})$.

Second, we consider the consistency of a \mathfrak{g} -scattering diagram $\mathfrak{D} = (\mathfrak{S}_S, \phi_{\mathfrak{D}})$ for \mathfrak{g} with finite support $S = \text{Supp}(\mathfrak{g})$. A \mathfrak{D} -generic curve is a smooth curve $\gamma : [0, 1] \rightarrow M_{\mathbb{R}}$ such that

- (1) the endpoints $\gamma(0)$ and $\gamma(1)$ lie in cones of \mathfrak{S}_S with dimension n ,
- (2) γ does not intersect cones of \mathfrak{S}_S with codimension at least two,
- (3) γ and walls of \mathfrak{S}_S intersect transversally.

Then there are finitely many points $0 < t_1 < \dots < t_l < 1$ and walls $\sigma_1, \dots, \sigma_l \in \mathcal{W}(\mathfrak{S}_S)$ such that $\gamma(t_i) \in \sigma_i$ and for $t \in [0, 1] \setminus \{t_1, \dots, t_l\}$, $\gamma(t)$ lie in cones of \mathfrak{S}_S with dimension n . We define the *path-ordered product*

$$\Phi_{\mathfrak{D}}(\gamma) := \phi_{\mathfrak{D}}(\sigma_l)^{\epsilon_l} \cdots \phi_{\mathfrak{D}}(\sigma_1)^{\epsilon_1} \in G,$$

where $\epsilon_i \in \{1, -1\}$ is the negative of sign of the derivative of $\gamma(t)$ at $t = t_i$. We say that two \mathfrak{g} -scattering diagrams \mathfrak{D}_1 and \mathfrak{D}_2 are *equivalent* if any \mathfrak{D}_1 -generic and \mathfrak{D}_2 -generic curve γ satisfies $\Phi_{\mathfrak{D}_1}(\gamma) = \Phi_{\mathfrak{D}_2}(\gamma)$.

Definition 6.2. We say that a \mathfrak{g} -scattering diagram \mathfrak{D} is *consistent* if any \mathfrak{D} -generic curves γ_1 and γ_2 with same endpoints satisfy $\Phi_{\mathfrak{D}}(\gamma_1) = \Phi_{\mathfrak{D}}(\gamma_2)$.

Let \mathfrak{D} be a consistent \mathfrak{g} -scattering diagram. For cones σ_1 and σ_2 of \mathfrak{S}_S with dimension n , we define

$$\Phi_{\mathfrak{D}}(\sigma_1, \sigma_2) := \Phi_{\mathfrak{D}}(\gamma),$$

where γ is any \mathfrak{D} -generic curve with $\gamma(0) \in \sigma_1$ and $\gamma(1) \in \sigma_2$. It does not depend on γ .

Finally, we define (consistent) \mathfrak{g} -scattering diagrams for a general \mathfrak{g} . We define a map $\delta : N \rightarrow \mathbb{Z}$ by $(d_i) \mapsto \sum_{i=1}^n d_i$. Let $\mathfrak{g}^{>k} := \bigoplus_{\delta(d) > k} \mathfrak{g}_d$ be a Lie subalgebra of \mathfrak{g} . Then $\mathfrak{g}^{\leq k} := \mathfrak{g}/\mathfrak{g}^{>k}$ is a nilpotent Lie algebra with finite support and there is the corresponding unipotent algebraic group $G^{\leq k}$. There is a bijection as sets between

$$\hat{\mathfrak{g}} := \varprojlim_k \mathfrak{g}^{\leq k} \quad \text{and} \quad \hat{G} := \varprojlim_k G^{\leq k}.$$

For $i > j$ and a $\mathfrak{g}^{\leq i}$ -scattering diagram $\mathfrak{D}_{\mathfrak{g}^{\leq i}} = (\mathfrak{S}, \phi)$, we have a $\mathfrak{g}^{\leq j}$ -scattering diagram $\pi_*^{ij}(\mathfrak{D}_{\mathfrak{g}^{\leq i}}) = (\mathfrak{S}, \pi^{ij}\phi)$, where π^{ij} is a natural homomorphism from $G^{\leq i}$ to $G^{\leq j}$.

Definition 6.3. A \mathfrak{g} -scattering diagram is a sequence of $\mathfrak{g}^{\leq k}$ -scattering diagrams $(\mathfrak{D}^{\leq k})_{k \geq 1}$ such that $\pi_*^{ij}(\mathfrak{D}^{\leq i})$ is equivalent to $\mathfrak{D}^{\leq j}$ for $i > j$. We say that it is *consistent* if so is $\mathfrak{D}^{\leq k}$ for any $k \geq 1$.

Remark that a consistent \mathfrak{g} -scattering diagram is considered as the pair (\mathfrak{S}, ϕ) consisting of the collection \mathfrak{S} of certain cones, that is not rational polyhedral in general, and a function $\phi : \mathfrak{S} \rightarrow \hat{G}$ (see [Mou19, Remark 2.26]).

In this section, our main subjects of study are cluster scattering diagrams and stability scattering diagrams. To define them, the following result plays an important role.

Theorem 6.4. [Bri17, Proposition 3.4][KS14, Theorem 2.1.6] *There is a bijection between the set of equivalence classes of consistent \mathfrak{g} -scattering diagrams and \hat{G} as sets.*

We denote by $\mathfrak{D}_g = (\mathfrak{D}_g^{\leq k})_{k \geq 1}$ the consistent \mathfrak{g} -scattering diagram corresponding to $g \in \hat{G}$.

6.2. Cluster scattering diagrams. Let Q be a quiver without loops and 2-cycles and $|Q_0| = n$. Assume that N has a skew-symmetric form $\{-, -\} : N \times N \rightarrow \mathbb{Z}$ given by

$$\{e_i, e_j\} := \#\{\text{arrows from } j \text{ to } i \text{ in } Q\} - \#\{\text{arrows from } i \text{ to } j \text{ in } Q\}$$

and \mathfrak{g} is skew-symmetric, that is, if $\{d_1, d_2\} = 0$ for $d_1, d_2 \in N$, $[\mathfrak{g}_{d_1}, \mathfrak{g}_{d_2}] = 0$.

For $m \in M_{\mathbb{R}}$, there is a decomposition

$$\mathfrak{g} = \mathfrak{g}_+^m \oplus \mathfrak{g}_0^m \oplus \mathfrak{g}_-^m,$$

where

$$\mathfrak{g}_{\pm}^m := \bigoplus_{d \in N^+ : \pm m(d) > 0} \mathfrak{g}_d \quad \text{and} \quad \mathfrak{g}_0^m := \bigoplus_{d \in N^+ : m(d) = 0} \mathfrak{g}_d.$$

We denote by \hat{G}_{\bullet}^m the subgroup of \hat{G} induced by \mathfrak{g}_{\bullet}^m for $\bullet \in \{+, -, 0\}$. This gives a unique decomposition of $\hat{G} \ni g = g_+^m \cdot g_0^m \cdot g_-^m$, where $g_{\bullet}^m \in \hat{G}_{\bullet}^m$. Thus there is a projection map $\pi_m : G \rightarrow \hat{G}_0^m$ given by $g \mapsto g_0^m$. Moreover, for a map $p^* : N \rightarrow M$ given by $d \mapsto \{d, -\}$, there is a decomposition

$$\mathfrak{g}_0^{p^*(d)} = \mathfrak{g}_d^{\parallel} \oplus \mathfrak{g}_d^{\perp},$$

where $\mathfrak{g}_d^{\parallel} := \bigoplus_{k=1}^{\infty} \mathfrak{g}_{kd}$. This naturally gives group homomorphisms $r_d : \hat{G}_0^{p^*(d)} \rightarrow \hat{G}_d^{\parallel}$ and $\psi_d := r_d \pi_{p^*(d)} : \hat{G} \rightarrow \hat{G}_d^{\parallel}$.

Proposition 6.5. [KS14, Proposition 3.3.2] *The map*

$$\psi := (\psi_d)_{d \in N^+ : \text{primitive}} : \hat{G} \rightarrow \prod_{d \in N^+ : \text{primitive}} \hat{G}_d^{\parallel}$$

is a bijection as sets.

Set the skew-symmetric N^+ -graded Lie algebra

$$\mathfrak{g} = \mathfrak{g}_{\text{cl}} := \bigoplus_{d \in N^+} \mathbb{Q}x^d \quad \text{with} \quad [x^{d_1}, x^{d_2}] = \{d_1, d_2\}x^{d_1+d_2} \quad \text{for } d_1, d_2 \in N^+.$$

Definition 6.6. Let $g = (g_d) \in \prod_{d \in N^+ : \text{primitive}} \hat{G}_d^{\parallel}$ given by

$$g_d = \begin{cases} \exp\left(\sum_{k=1}^{\infty} \frac{(-1)^{k-1} x^{ks_i}}{k^2}\right) & \text{if } d = e_i, \\ \text{id} & \text{otherwise.} \end{cases}$$

Then the corresponding consistent \mathfrak{g}_{cl} -scattering diagram $\mathfrak{D}_Q := \mathfrak{D}_{\psi^{-1}(g)}$ is called the *cluster scattering diagram* associated with Q .

6.3. Stability scattering diagrams. Let J be a Jacobian algebra, and denote by $K_0(\text{mod } J) \simeq \mathbb{Z}^n = N$ its Grothendieck group. In this subsection, we recall the stability scattering diagram associated with J . We refer to [Bri17] for the details.

Let $\mathfrak{M}(J)$ be the moduli stack associated with $\text{mod } J$. There is a decomposition

$$\mathfrak{M}(J) = \coprod_{d \in N^{\oplus}} \mathfrak{M}(J)_d,$$

where $\mathfrak{M}(J)_d$ is the moduli stack of J -modules with dimension d . Let $K(\text{St}/\mathbb{C})$ be the Grothendieck ring of stacks over \mathbb{C} . The Grothendieck group $K(\text{St}/\mathfrak{M}(J))$ of stacks over $\mathfrak{M}(J)$ is an N^{\oplus} -graded $K(\text{St}/\mathbb{C})$ -algebra. Since there is a unique ring homomorphism from $K(\text{St}/\mathbb{C})$ to $\mathbb{Q}(t)$ sending the classes of smooth projective varieties to their Poincaré polynomials, we have an N^{\oplus} -graded $\mathbb{Q}(t)$ -algebra

$$H(J) := K(\text{St}/\mathfrak{M}(J)) \otimes_{K(\text{St}/\mathbb{C})} \mathbb{Q}(t),$$

called the *motivic Hall algebra* associated with J . For $d \in N^{\oplus}$, let $H(J)_d$ be the subspace of $H(J)$ generated by elements with form $[X \rightarrow \mathfrak{M}(J)]$ factoring through the inclusion from $\mathfrak{M}(J)_d$ to $\mathfrak{M}(J)$. Then

$$\mathfrak{g}_{\text{Hall}}(J) := \bigoplus_{d \in N^+} H(J)_d$$

is an N^+ -graded Lie algebra with commutator bracket. We have an element of the prounipotent algebraic group $\exp(\hat{\mathfrak{g}}_{\text{Hall}}(J))$

$$1_{\mathfrak{M}(J)} := [\mathfrak{M}(J) \xrightarrow{\text{id}} \mathfrak{M}(J)] \in 1 + \hat{\mathfrak{g}}_{\text{Hall}}(J) \simeq \exp(\hat{\mathfrak{g}}_{\text{Hall}}(J)).$$

The consistent $\mathfrak{g}_{\text{Hall}}$ -scattering diagram $\mathfrak{D}_{1_{\mathfrak{M}(J)}}$ is called the *Hall algebra scattering diagram* associated with J .

Define a subalgebra

$$\mathbb{C}_{\text{reg}}(t) := \mathbb{C}[t, t^{-1}][(1 + t^2 + \cdots + t^{2k})^{-1} \mid k \geq 1] \subset \mathbb{C}(t).$$

We denote by $\mathfrak{g}_{\text{reg}}(J) \subset \mathfrak{g}_{\text{Hall}}(J)$ the $\mathbb{C}_{\text{reg}}(t)$ -submodule generated by $[X \rightarrow \mathfrak{M}(J)]$ with algebraic variety X .

Theorem 6.7 ([Joy08, Bri17]). *Any Hall algebra scattering diagram is a $\mathfrak{g}_{\text{reg}}(J)$ -scattering diagram. Moreover, there is an N^+ -graded Lie algebra homomorphism*

$$\mathcal{I} : \mathfrak{g}_{\text{reg}}(J) \rightarrow \mathfrak{g}_{\text{cl}}.$$

Definition 6.8. The consistent \mathfrak{g}_{cl} -scattering diagram $\mathfrak{D}_J := \mathcal{I}(\mathfrak{D}_{1_{\mathfrak{M}(J)}})$ is called the *stability scattering diagram* associated with J .

6.4. Cluster/Stability scattering diagrams. Let Q be a quiver without loops and 2-cycles and W a non-degenerate potential of Q . We state the main result of this section.

Theorem 6.9. (1) *If Q is cluster- \mathfrak{g} -dense, then the cluster scattering diagram \mathfrak{D}_Q is equivalent to the stability scattering diagram $\mathfrak{D}_{J(Q,W)}$.*
 (2) *If Q is half cluster- \mathfrak{g} -dense, then \mathfrak{D}_Q and $\mathfrak{D}_{J(Q,W)}$ only differ by functions on walls in the separating hyperplane.*

To prove Theorem 6.9, we need some preparations. Set $J = J(Q, W)$ and $\mathfrak{g} = \mathfrak{g}_{\text{cl}}$. We only need to compare two consistent $\mathfrak{g}^{\leq k}$ -scattering diagrams $\mathfrak{D}_Q^{\leq k} = (\mathfrak{S}_Q^{\leq k}, \phi_Q^{\leq k})$ and $\mathfrak{D}_J^{\leq k} = (\mathfrak{S}_J^{\leq k}, \phi_J^{\leq k})$ for any $k \geq 1$. Note that $\mathfrak{g}^{\leq k}$ has finite support.

We can consider $\mathcal{F}_{\text{cluster}}^{\mathfrak{g}}(Q) \cup (-\mathcal{F}_{\text{cluster}}^{\mathfrak{g}}(Q^{\text{op}}))$ as a fan in $M_{\mathbb{R}} \simeq \mathbb{R}^n$. For a wall $\sigma \subset d_0^\perp$ of $\mathcal{F}_{\text{cluster}}^{\mathfrak{g}}(Q) \cup (-\mathcal{F}_{\text{cluster}}^{\mathfrak{g}}(Q^{\text{op}}))$, where $d_0 \in N^+$ is primitive, let

$$\phi(\sigma) = \exp\left(\sum_{k=1}^{\infty} \frac{(-1)^{k-1} x^{kd_0}}{k^2}\right) \in \hat{G}.$$

For a projection map $p_k : \hat{G} \rightarrow G^{\leq k}$, we define a fan $\mathcal{F}_k^{\mathfrak{g}}(Q)$ consisting of cones σ of $\mathcal{F}_{\text{cluster}}^{\mathfrak{g}}(Q) \cup (-\mathcal{F}_{\text{cluster}}^{\mathfrak{g}}(Q^{\text{op}}))$ such that $p_k \phi(\sigma)$ is not trivial. In particular, it has two cones

$$M_{\mathbb{R}}^{\pm} := \{m \in M_{\mathbb{R}} \mid \pm m(e_i) \geq 0 \text{ for any } i\} \subset M_{\mathbb{R}}.$$

Theorem 6.10. [Mou19, Theorem 4.25] *The fan $\mathcal{F}_k^{\mathfrak{g}}(Q)$ is a common subfan of $\mathfrak{S}_Q^{\leq k}$ and $\mathfrak{S}_J^{\leq k}$ such that $\phi_Q^{\leq k}(\sigma) = \phi_J^{\leq k}(\sigma) = p_k \phi(\sigma)$ for any $\sigma \in \mathcal{W}(\mathcal{F}_k^{\mathfrak{g}}(Q))$.*

We are ready to prove Theorem 6.9. For any wall $\sigma \in \mathcal{W}(\mathfrak{S}_Q^{\leq k})$, there are two cones $\sigma^+, \sigma^- \in G_Q^{\leq k}$ with dimension n such that $\sigma^+ \cap \sigma^- = \sigma$ since $\mathfrak{S}_Q^{\leq k}$ is a finite fan. If Q is cluster- \mathfrak{g} -dense, then $\Phi_{G_Q^{\leq k}}(M_{\mathbb{R}}^+, \sigma^+)$ and $\Phi_{G_Q^{\leq k}}(M_{\mathbb{R}}^+, \sigma^-)$ are products of some functions on $\mathcal{F}_k^{\mathfrak{g}}(Q)$. Thus so is $\Phi_{G_Q^{\leq k}}(\sigma^+, \sigma^-)$. On the other hand, there is a $\mathfrak{D}_Q^{\leq k}$ -generic curve from σ^+ to σ^- that intersects only one wall σ . Thus we have $\Phi_{G_Q^{\leq k}}(\sigma^+, \sigma^-) = \phi_{G_Q^{\leq k}}(\sigma)^\epsilon$ for some $\epsilon \in \{1, -1\}$. Therefore, for any wall $\sigma \in \mathcal{W}(\mathfrak{S}_Q^{\leq k})$, $\phi_{G_Q^{\leq k}}(\sigma)$ is a product of some functions on $\mathcal{F}_k^{\mathfrak{g}}(Q)$.

Similarly, we have that, for any wall $\sigma \in \mathcal{W}(\mathfrak{S}_J^{\leq k})$, $\phi_{G_J^{\leq k}}(\sigma)$ is a product of some functions on $\mathcal{F}_k^{\mathfrak{g}}(Q)$. Consequently, Theorem 6.10 implies Theorem 6.9(1).

Suppose that Q is half cluster- \mathfrak{g} -dense with the separating hyperplane d^\perp for $d \in N^+$. If σ does not lie on d^\perp , then we have $\phi_{G_Q^{\leq k}}(\sigma) = \phi_{G_J^{\leq k}}(\sigma)$ in the same way as above. Thus Theorem 6.9(2) holds.

ACKNOWLEDGEMENTS

We are grateful to A. Skowroński and G. Zwara for suggesting an argument in the proof of Theorem 3.8. We also thank J. Schröer for discussing the results of [GLFS20] in a meeting at MFO in January 2020. We thank B. Keller for discussions on twists, for his comments on a previous version, and for agreeing to write the appendix to this paper. Finally, we thank O. Iyama for discussing the results of [ADI20], and K. Mousavand for valuable discussions.

APPENDIX A. TRUNCATED TWIST FUNCTORS, BY BERNHARD KELLER

A.1. Construction. We refer to [Kel06] for the terminology and notation that we will use for dg (=differential graded) categories. Let k be a commutative ring and \mathcal{S}, \mathcal{T} k -linear triangulated categories with dg enhancements \mathcal{S}_{dg} and \mathcal{T}_{dg} . Let $s : \mathcal{S} \rightarrow \mathcal{T}$ be a triangle functor induced by a dg functor $s_{dg} : \mathcal{S}_{dg} \rightarrow \mathcal{T}_{dg}$. Assume that s admits a right adjoint $r : \mathcal{T} \rightarrow \mathcal{S}$. Let there be given a t -structure with truncation functor $\tau_{\leq 0}$ on \mathcal{S} .

Proposition A.1. a) *There is a triangle functor $t : \mathcal{T} \rightarrow \mathcal{T}$ admitting a dg enhancement and fitting into a functorial triangle*

$$sr \longrightarrow \mathbf{1}_{\mathcal{T}} \longrightarrow t \longrightarrow \Sigma sr.$$

b) *There is a k -linear functor $t^0 : \mathcal{T} \rightarrow \mathcal{T}$ fitting into a functorial triangle*

$$s\tau_{\leq 0}r \longrightarrow \mathbf{1}_{\mathcal{T}} \longrightarrow t^0 \longrightarrow \Sigma s\tau_{\leq 0}r.$$

The functor t is called the *twist functor* and t^0 the *truncated twist functor* associated with s .

Examples A.2. (1) Let \mathcal{A} be a dg category and X an object of its derived category \mathcal{DA} . We can take $\mathcal{S} = \mathcal{D}k$, $\mathcal{T} = \mathcal{DA}$ and $s = ? \overset{\mathrm{L}}{\otimes}_k X : \mathcal{D}k \rightarrow \mathcal{DA}$ with right adjoint

$$r = \mathrm{RHom}_{\mathcal{A}}(X, ?) : \mathcal{DA} \rightarrow \mathcal{D}k.$$

We get a twist functor t_X and a truncated twist functor t_X^0 fitting into functorial triangles

$$\mathrm{RHom}_{\mathcal{A}}(X, ?) \overset{\mathrm{L}}{\otimes}_k X \longrightarrow \mathbf{1}_{\mathcal{DA}} \longrightarrow t_X \longrightarrow$$

and

$$\tau_{\leq 0}(\mathrm{RHom}_{\mathcal{A}}(X, ?)) \overset{\mathrm{L}}{\otimes}_k X \longrightarrow \mathbf{1}_{\mathcal{DA}} \longrightarrow t_X^0 \longrightarrow .$$

(2) With \mathcal{A} , X and \mathcal{S} as in example (1), we can take $\mathcal{T} = (\mathcal{DA})^{op}$. We have an adjoint pair

$$\begin{array}{ccc} & (\mathcal{DA})^{op} & \\ s = \mathrm{RHom}_{\mathcal{A}}(?, X) \downarrow & \updownarrow & \mathrm{RHom}_k(?, X) = r \\ & \mathcal{D}k. & \end{array}$$

This leads to the following functorial triangles in \mathcal{DA} :

$$\longrightarrow t_X \longrightarrow \mathbf{1}_{\mathcal{DA}} \longrightarrow \mathrm{RHom}_k(\mathrm{RHom}_{\mathcal{A}}(? , X), X)$$

and

$$\longrightarrow t_X^0 \longrightarrow \mathbf{1}_{\mathcal{DA}} \longrightarrow \mathrm{RHom}_k(\tau_{\leq 0} \mathrm{RHom}_{\mathcal{A}}(? , X), X).$$

Proof of the Proposition. a) This is well-known, cf. for example [AL17]: One shows that one can replace \mathcal{T}_{dg} and \mathcal{S}_{dg} with quasi-equivalent dg categories such that s_{dg} admits a dg adjoint functor r_{dg} .

b) Let us fix s_{dg} and r_{dg} as in a). From now on, we suppress the subscript ‘dg’ on these functors. Let \mathcal{M}_{dg} be the dg category whose objects are the triples

$$(U, V, f : sV \rightarrow U)$$

where U is in \mathcal{T} , V in \mathcal{S} and $f : sV \rightarrow U$ is a closed morphism of degree 0 in \mathcal{S}_{dg} . By definition, the morphism complex between two objects (U, V, f) and (U', V', f') is the cylinder over the morphism

$$[f^*, -f' \circ s(?)] : \mathcal{T}_{dg}(U, U') \oplus \mathcal{S}_{dg}(V, V') \rightarrow \mathcal{T}_{dg}(sV, U').$$

Thus, closed morphisms of degree 0 are triples (u, v, h) such that $u : U \rightarrow U'$ is closed in \mathcal{T}_{dg} , $v : V \rightarrow V'$ is closed in \mathcal{S}_{dg} and we have

$$u \circ f - f' \circ s(v) = d(h)$$

so that we have a homotopy commutative square

$$\begin{array}{ccc} sV & \xrightarrow{f} & U \\ s(v) \downarrow & \searrow h & \downarrow u \\ sV' & \xrightarrow{f'} & U' \end{array}$$

Composition is defined in the natural way. The projection functor $P : \mathcal{M}_{dg} \rightarrow \mathcal{T}_{dg}$ takes a triple (U, V, f) to U . It is easy to see that it has a fully faithful right dg adjoint P_ρ taking U to $(U, rU, srU \rightarrow U)$, where the last arrow is the adjunction morphism, and a fully faithful left dg adjoint P_λ taking U to $(U, 0, 0 \rightarrow U)$. The functor I taking V to $(0, V, sV \rightarrow 0)$ identifies \mathcal{S}_{dg} with the kernel of P . We can construct the dg twist functor t as the composition

$$\mathcal{T}_{dg} \xrightarrow{P_\rho} \mathcal{M}_{dg} \xrightarrow{\text{cone}} \mathcal{T}_{dg}.$$

Now let $\mathcal{M} = H^0(\mathcal{M}_{dg})$. We have a recollement

$$\mathcal{S} \begin{array}{c} \xleftarrow{I_\lambda} \\ \xleftarrow{I} \\ \xleftarrow{I_\rho} \end{array} \mathcal{M} \begin{array}{c} \xleftarrow{P_\lambda} \\ \xleftarrow{P} \\ \xleftarrow{P_\rho} \end{array} \mathcal{T}.$$

Let $\mathcal{S}_{\leq 0}$ be the left aisle of the given t -structure on \mathcal{S} . Using Théorème 1.4.10 of [BBD82] or directly, we see that \mathcal{M} admits a t -structure whose left aisle $\mathcal{M}_{\leq 0}$ has the objects (U, V, f) such that $V \in \mathcal{S}_{\leq 0}$. The corresponding morphism $\tau_{\leq 0} \rightarrow \mathbf{1}_{\mathcal{M}}$ is given on an object (U, V, f) by $(U, \tau_{\leq 0} V, g)$, where g is the composition

$$s\tau_{\leq 0} V \xrightarrow{sc} sV \xrightarrow{f} U$$

for any closed morphism c lifting the adjunction morphism $\tau_{\leq 0} V \rightarrow V$. We obtain the truncated twist functor t^0 as the composition

$$\mathcal{T} \xrightarrow{P_\rho} \mathcal{M} \xrightarrow{\tau_{\leq 0}} \mathcal{M}_{\leq 0} \xrightarrow{\text{cone}} \mathcal{T}.$$

□

A.2. Adjoints. We keep the assumptions and notations of the preceding section. Suppose that $s : \mathcal{S} \rightarrow \mathcal{T}$ admits a left adjoint $l : \mathcal{T} \rightarrow \mathcal{S}$ and that the inclusion $\mathcal{S}_{\leq 0} \rightarrow \mathcal{S}$ also admits a left adjoint $\tau'_{\leq 0}$ (for example, this holds if k is a field and $\mathcal{S} = \mathcal{D}k$ is endowed with the canonical t -structure).

Proposition A.3. a) *The twist functor t admits a left adjoint t' fitting into a functorial triangle*

$$\Sigma^{-1} sl \rightarrow t' \rightarrow \mathbf{1}_{\mathcal{T}} \rightarrow sl.$$

b) *The truncated twist functor t^0 admits a left adjoint t'^0 fitting into a functorial triangle*

$$\Sigma^{-1} s \tau'_{\leq -1} l \rightarrow t'^0 \rightarrow \mathbf{1}_{\mathcal{T}} \rightarrow s \tau'_{\leq -1} l.$$

Examples A.4. We keep the notations of Examples A.2. We suppose that k is a field.

- (1) Suppose that $X \in \mathcal{DA}$ is an object such that the homology of $X(A)$ is of finite total dimension for all $A \in \mathcal{A}$. Put $D = \text{Hom}_k(?, k)$. Then for $U \in \mathcal{DA}$ and $V \in \mathcal{D}k$, we have canonical isomorphisms

$$\text{RHom}_{\mathcal{A}}(U, V \otimes X) = \text{RHom}_{\mathcal{A}}(U, \text{Hom}_k(DX, V)) = \text{Hom}_k(U \overset{\text{L}}{\otimes}_{\mathcal{A}} DX, X).$$

We get functorial triangles

$$\longrightarrow t' \longrightarrow \mathbf{1}_{\mathcal{DA}} \longrightarrow (? \overset{\text{L}}{\otimes}_{\mathcal{A}} DX) \otimes X$$

and

$$\longrightarrow t'^0 \longrightarrow \mathbf{1}_{\mathcal{DA}} \longrightarrow (\tau'_{\leq -1} (? \overset{\text{L}}{\otimes}_{\mathcal{A}} DX)) \otimes X.$$

- (2) Let $\mathcal{T} \subseteq \mathcal{DA}$ be a Hom-finite triangulated subcategory and X an object in \mathcal{T} such that the homology of $X(A)$ is of finite total dimension for each $A \in \mathcal{A}$. Let

$$s = \text{Hom}_k(?, X) : \mathcal{D}^b(k) \rightarrow \mathcal{T}^{op}.$$

Then for U in \mathcal{T} and V in $\mathcal{D}^b(k)$, we have

$$\begin{aligned} (\mathcal{DA})^{op}(U, \text{Hom}_k(V, X)) &= (\mathcal{DA})(\text{Hom}_k(V, X), U) \\ &= (\mathcal{DA})(X \otimes DV, U) \\ &= (\mathcal{DA})(DV \otimes X, U) \\ &= (\mathcal{D}k)(DV, \text{RHom}_{\mathcal{A}}(X, U)) \\ &= (\mathcal{D}k)(D \text{RHom}_{\mathcal{A}}(X, U), V). \end{aligned}$$

Thus, the functor $s : \mathcal{D}^b(k) \rightarrow \mathcal{T}$ admits $l = D\mathrm{RHom}_{\mathcal{A}}(X, ?)$ as a left adjoint and we get functorial triangles

$$\longrightarrow t' \longrightarrow \mathbf{1}_{\mathcal{T}} \longrightarrow D\mathrm{RHom}_{\mathcal{A}}(?, X) \otimes X$$

and

$$\longrightarrow t'^0 \longrightarrow \mathbf{1}_{\mathcal{T}} \longrightarrow \tau'_{\leq -1}(D\mathrm{RHom}_{\mathcal{A}}(?, X)) \otimes X.$$

Proof of the Proposition. Part a) is well-known, cf. for example [AL17]. Let us prove part b). For objects U, U' of \mathcal{T} , we abbreviate

$$(U, U') = \mathrm{RHom}_{\mathcal{T}_{dg}}(U, U')$$

and similarly

$$(V, V') = \mathrm{RHom}_{\mathcal{S}_{dg}}(V, V')$$

for objects V, V' of \mathcal{S} . Let U and U' be objects of \mathcal{T} . We have natural isomorphisms

$$\begin{aligned} \mathrm{Hom}_{\mathcal{T}}(t'^0 U, U') &\simeq H^0(\mathrm{Cyl}((s\tau'_{\leq -1} lU, \Sigma U') \rightarrow (U, \Sigma U'))) \\ &\simeq H^0(\mathrm{Cyl}((\tau'_{\leq -1} lU, \Sigma rU') \rightarrow (U, \Sigma U'))) \\ &\simeq H^0(\mathrm{Cyl}((\tau'_{\leq -1} lU, \tau_{\leq -1} \Sigma rU') \rightarrow (U, \Sigma U'))). \end{aligned}$$

Here the last isomorphism holds because we have isomorphisms

$$\mathrm{Hom}_{\mathcal{S}}(\tau'_{\leq -1} lU, \tau_{\leq -1} \Sigma rU) \simeq \mathrm{Hom}_{\mathcal{S}}(\tau'_{\leq -1} lU, \Sigma rU)$$

and

$$\mathrm{Hom}_{\mathcal{S}}(\Sigma \tau'_{\leq -1} lU, \tau_{\leq -1} \Sigma rU) \simeq \mathrm{Hom}_{\mathcal{S}}(\Sigma \tau'_{\leq -1} lU, \Sigma rU).$$

We continue the chain of isomorphisms with

$$\begin{aligned} H^0(\mathrm{Cyl}((\tau'_{\leq -1} lU, \tau_{\leq -1} \Sigma rU') \rightarrow (U, \Sigma U'))) &\simeq H^0(\mathrm{Cyl}((lU, \Sigma \tau_{\leq 0} rU') \rightarrow (U, \Sigma U'))) \\ &\simeq H^0(\mathrm{Cyl}(U, \Sigma s\tau_{\leq 0} rU') \rightarrow (U, \Sigma U'))) \\ &\simeq \mathrm{Hom}_{\mathcal{T}}(U, t'^0 U'). \end{aligned}$$

This shows the claim. \square

REFERENCES

- [ADI20] Sota Asai, Laurent Demonet, and Osamu Iyama. On algebras with dense g -vector cones. Work in progress, 2020.
- [AHBHY18] Nima Arkani-Hamed, Yuntao Bai, Song He, and Gongwang Yan. Scattering forms and the positive geometry of kinematics, color and the worldsheet. *J. High Energy Phys.*, (5):096, front matter+75, 2018.
- [AIR14] Takahide Adachi, Osamu Iyama, and Idun Reiten. τ -tilting theory. *Compos. Math.*, 150(3):415–452, 2014.
- [AL17] Rina Anno and Timothy Logvinenko. Spherical DG-functors. *J. Eur. Math. Soc. (JEMS)*, 19(9):2577–2656, 2017.
- [Asa19] Sota Asai. The wall-chamber structures of the real Grothendieck groups. arXiv:1905.02180, 2019.
- [ASS06] Ibrahim Assem, Daniel Simson, and Andrzej Skowroński. *Elements of the representation theory of associative algebras. Vol. 1*, volume 65 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 2006. Techniques of representation theory.
- [AY20] Toshitaka Aoki and Toshiya Yurikusa. Complete gentle algebras are \mathbf{g} -tame. arXiv:2003.09797 [math.RT], 2020.
- [BBD82] Alexander A. Beilinson, Joseph Bernstein, and Pierre Deligne. *Analyse et topologie sur les espaces singuliers*, volume 100 of *Astérisque*. Soc. Math. France, 1982.

- [BDM⁺17] Thomas Brüstle, Guillaume Douville, Kaveh Mousavand, Hugh Thomas, and Emine Yıldırım. On the combinatorics of gentle algebras. Preprint, [arXiv:1707.07665](#), 2017.
- [BG12] Michael Barot and Christof Geiss. Tubular cluster algebras I: categorification. *Mathematische Zeitschrift*, 271(3-4):1091–1115, 2012.
- [BGJ13] Michael Barot, Christof Geiss, and Gustavo Jasso. Tubular cluster algebras II: exponential growth. *Journal of Pure and Applied Algebra*, 217(10):1825–1837, 2013.
- [BKL10] Michael Barot, Dirk Kussin, and Helmut Lenzing. The cluster category of a canonical algebra. *Transactions of the American Mathematical Society*, 362(8):4313–4330, 2010.
- [BMP20] Guillaume Beaujard, Jan Manschot, and Boris Pioline. Vafa-Witten invariants from exceptional collections. [arXiv:2004.14466 \[hep-th\]](#), 2020.
- [BMR⁺06] Aslak Bakke Buan, Robert Marsh, Markus Reineke, Idun Reiten, and Gordana Todorov. Tilting theory and cluster combinatorics. *Advances in mathematics*, 204(2):572–618, 2006.
- [Bou20] Pierrick Bousseau. DT invariants of 3-gon quivers. Lecture notes available at <https://webusers.imj-prg.fr/~bernhard.keller/gdtcluster/>, June 11th, 2020.
- [Bri17] Tom Bridgeland. Scattering diagrams, Hall algebras and stability conditions. *Algebraic Geometry*, 4(5):523–561, 2017.
- [BST19] Thomas Brüstle, David Smith, and Hipolito Treffinger. Wall and chamber structure for finite-dimensional algebras. *Adv. Math.*, 354, 2019.
- [CB88] W. W. Crawley-Boevey. On tame algebras and bocses. *Proc. London Math. Soc. (3)*, 56(3):451–483, 1988.
- [CFZ02] Frédéric Chapoton, Sergey Fomin, and Andrei Zelevinsky. Polytopal realizations of generalized associahedra. volume 45, pages 537–566. 2002. Dedicated to Robert V. Moody.
- [CIKLFP13] Giovanni Cerulli Irelli, Bernhard Keller, Daniel Labardini-Fragoso, and Pierre-Guy Plamondon. Linear independence of cluster monomials for skew-symmetric cluster algebras. *Compositio Mathematica*, 149(10):1753–1764, 2013.
- [CILFS15] Giovanni Cerulli Irelli, Daniel Labardini-Fragoso, and Jan Schröer. Caldero-Chapoton algebras. *Trans. Amer. Math. Soc.*, 367(4):2787–2822, 2015.
- [DF15] Harm Derksen and Jiarui Fei. General presentations of algebras. *Adv. Math.*, 278:210–237, 2015.
- [DIJ19] Laurent Demonet, Osamu Iyama, and Gustavo Jasso. τ -tilting finite algebras, bricks, and g -vectors. *Int. Math. Res. Not. IMRN*, (3):852–892, 2019.
- [DK08] Raika Dehy and Bernhard Keller. On the combinatorics of rigid objects in 2-Calabi-Yau categories. *Int. Math. Res. Not. IMRN*, (11):Art. ID rnn029, 17, 2008.
- [DO08] Harm Derksen and Theodore Owen. New graphs of finite mutation type. *Electron. J. Combin.*, 15(1):Research Paper 139, 15, 2008.
- [DWZ08] Harm Derksen, Jerzy Weyman, and Andrei Zelevinsky. Quivers with potentials and their representations I: Mutations. *Selecta Mathematica*, 14(1):59–119, 2008.
- [DWZ10] Harm Derksen, Jerzy Weyman, and Andrei Zelevinsky. Quivers with potentials and their representations II: applications to cluster algebras. *J. Amer. Math. Soc.*, 23(3):749–790, 2010.
- [FG09] Vladimir V. Fock and Alexander B. Goncharov. Cluster ensembles, quantization and the dilogarithm. *Ann. Sci. Éc. Norm. Supér. (4)*, 42(6):865–930, 2009.
- [FK10] Changjian Fu and Bernhard Keller. On cluster algebras with coefficients and 2-Calabi-Yau categories. *Transactions of the American Mathematical Society*, 362(2):859–895, 2010.
- [FST08] Sergey Fomin, Michael Shapiro, and Dylan Thurston. Cluster algebras and triangulated surfaces. part I: Cluster complexes. *Acta Math.*, 201(1):83–146, 2008.
- [FST12] Anna Felikson, Michael Shapiro, and Pavel Tumarkin. Skew-symmetric cluster algebras of finite mutation type. *J. Eur. Math. Soc.*, 14(4):1135–1180, 2012.
- [FZ03] Sergey Fomin and Andrei Zelevinsky. Y -systems and generalized associahedra. *Ann. of Math. (2)*, 158(3):977–1018, 2003.
- [FZ07] Sergey Fomin and Andrei Zelevinsky. Cluster algebras IV: coefficients. *Compositio Mathematica*, 143(1):112–164, 2007.

- [Gab74] Peter Gabriel. Finite representation type is open. In *Proceedings of the International Conference on Representations of Algebras (Carleton Univ., Ottawa, Ont., 1974)*, Paper No. 10, pages 23 pp. Carleton Math. Lecture Notes, No. 9, 1974.
- [GGS15] Christof Geiss and Raúl González-Silva. Tubular Jacobian algebras. *Algebras and Representation Theory*, 18(1):161–181, 2015.
- [GHKK18] Mark Gross, Paul Hacking, Sean Keel, and Maxim Kontsevich. Canonical bases for cluster algebras. *Journal of the American Mathematical Society*, 31(2):497–608, 2018.
- [GLFS16] Christof Geiss, Daniel Labardini-Fragoso, and Jan Schröer. The representation type of Jacobian algebras. *Advances in Mathematics*, 290:364–452, 2016.
- [GLFS20] Christof Geiss, Daniel Labardini-Fragoso, and Jan Schröer. Schemes of modules over gentle algebras and laminations of surfaces. arXiv:2005.01073 [math.RT], 2020.
- [GLS12] Christof Geiss, Bernard Leclerc, and Jan Schröer. Generic bases for cluster algebras and the Chamber ansatz. *J. Amer. Math. Soc.*, 25(1):21–76, 2012.
- [Hap88] Dieter Happel. *Triangulated categories in the representation theory of finite-dimensional algebras*, volume 119 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1988.
- [Hil06] L. Hille. On the volume of a tilting module. *Abh. Math. Sem. Univ. Hamburg*, 76:261–277, 2006.
- [HPS18] Christophe Hohlweg, Vincent Pilaud, and Salvatore Stella. Polytopal realizations of finite type \mathbf{g} -vector fans. *Adv. Math.*, 328:713–749, 2018.
- [IY08] Osamu Iyama and Yuji Yoshino. Mutation in triangulated categories and rigid Cohen–Macaulay modules. *Inventiones mathematicae*, 172(1):117–168, 2008.
- [Jas15] Gustavo Jasso. Reduction of τ -tilting modules and torsion pairs. *Int. Math. Res. Not. IMRN*, (16):7190–7237, 2015.
- [Joy08] Dominic Joyce. Configurations in abelian categories. iv. invariants and changing stability conditions. *Advances in Mathematics*, 217(1):125–204, 2008.
- [Kel06] Bernhard Keller. On differential graded categories. In *International Congress of Mathematicians. Vol. II*, pages 151–190. Eur. Math. Soc., Zürich, 2006.
- [KS14] Maxim Kontsevich and Yan Soibelman. Wall-crossing structures in Donaldson–Thomas invariants, integrable systems and mirror symmetry. In *Homological mirror symmetry and tropical geometry*, pages 197–308. Springer, 2014.
- [Lad11] Sefi Ladkani. Mutation classes of certain quivers with potentials as derived equivalence classes. *arXiv preprint arXiv:1102.4108*, 2011.
- [LF09] Daniel Labardini-Fragoso. Quivers with potentials associated to triangulated surfaces. *Proceedings of the London Mathematical Society*, 98(3):797–839, 2009.
- [LF16] Daniel Labardini-Fragoso. Quivers with potentials associated to triangulated surfaces, part iv: Removing boundary assumptions. *Selecta Mathematica*, 22(1):145–189, 2016.
- [Mil17] Matthew R Mills. Maximal green sequences for quivers of finite mutation type. *Advances in Mathematics*, 319:182–210, 2017.
- [Mou19] Lang Mou. Scattering diagrams of quivers with potentials and mutations. *arXiv preprint arXiv:1910.13714*, 2019.
- [Mul16] Greg Muller. The existence of a maximal green sequence is not invariant under quiver mutation. *Electron. J. Combin.*, 23(2):P2.47, 2016.
- [OPS18] Sebastian Oppel, Pierre-Guy Plamondon, and Sibylle Schroll. A geometric model for the derived category of gentle algebras. arXiv:1801.09659 [math.RT], 2018.
- [Pla13] Pierre-Guy Plamondon. Generic bases for cluster algebras from the cluster category. *Int. Math. Res. Not. IMRN*, (10):2368–2420, 2013.
- [PPPP19] Arnau Padrol, Yann Palu, Vincent Pilaud, and Pierre-Guy Plamondon. Associahedra for finite type cluster algebras and minimal relations between \mathbf{g} -vectors. arXiv:1906.06861 [math.RT], 2019.
- [Sev14] Ahmet I Seven. Maximal green sequences of exceptional finite mutation type quivers. *SIGMA. Symmetry, Integrability and Geometry: Methods and Applications*, 10:089, 2014.
- [ST01] Paul Seidel and Richard Thomas. Braid group actions on derived categories of coherent sheaves. *Duke Math. J.*, 108(1):37–108, 2001.

- [Yur20] Toshiya Yurikusa. Density of g-Vector Cones From Triangulated Surfaces. *International Mathematics Research Notices*, 02 2020. rnaa008.
- [Zie95] Günter M. Ziegler. *Lectures on polytopes*, volume 152 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.

P.-G.P. : UNIVERSITÉ PARIS-SACLAY, CNRS, LABORATOIRE DE MATHÉMATIQUES D'ORSAY,
91405, ORSAY, FRANCE

E-mail address: pierre-guy.plamondon@universite-paris-saclay.fr

T.Y. : MATHEMATICAL INSTITUTE, TOHOKU UNIVERSITY, AOBA-KU, SENDAI, 980-8578, JAPAN

E-mail address: toshiya.yurikusa.d8@tohoku.ac.jp

B. K. : UNIVERSITÉ PARIS DE PARIS, UFR DE MATHÉMATIQUES, CNRS, INSTITUT DE MATHÉMATIQUES
DE JUSSIEU-PARIS RIVE GAUCHE, IMJ-PRG, BÂTIMENT SOPHIE GERMAIN, 75205 PARIS CEDEX
13, FRANCE

E-mail address: bernhard.keller@imj-prg.fr

URL: <https://webusers.imj-prg.fr/~bernhard.keller/>