

Heat exchange and fluctuation in Gaussian thermal states in the quantum realm

A. R. Usha Devi^{1,2}, Sudha^{3,2}, A. K. Rajagopal² and A. M. Jayannavar⁴

¹ Department of Physics, Jnanabharathi, Bangalore University, Bangalore-560056, India

² Inspire Institute Inc., Alexandria, Virginia, 22303, USA.

³ Department of Physics, Kuvempu University, Shankaraghatta-577 451, India

⁴ Institute of Physics, Bhubaneswar, India

E-mail: arutth@rediffmail.com

Abstract. The celebrated exchange fluctuation theorem – proposed by Jarzynski and Wójcik, (Phys Rev. Lett. **92**, 230602 (2004)) for heat exchange between two systems in thermal equilibrium at different temperatures – is explored here for quantum Gaussian states in thermal equilibrium. We employ Wigner distribution function formalism for quantum states, which exhibits close resemblance with the classical phase-space trajectory description, to arrive at this theorem. For two Gaussian states in thermal equilibrium at two different temperatures kept in contact with each other for a fixed duration of time we show that the quantum Jarzynski-Wójcik theorem agrees with the corresponding classical result in the *high temperature limit*.

Keywords: heat exchange statistics, exchange-fluctuation theorem, Gaussian states, variance matrix, Wigner function, symplectic transformation

1. Introduction

Fluctuation theorems [1, 2, 3, 4, 5] are of fundamental significance in non-equilibrium statistical physics. They correspond to a collection of exact relations, which remain valid even when the system is driven far away from equilibrium. Various exchange-fluctuation theorems (XFT) involving thermodynamic quantities like work, heat, entropy have been proposed during the last two decades [5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22]. They have offered significant insight in understanding thermodynamical processes – especially the emergence of irreversibility from reversible dynamics and the directionality of heat flow implied by second law of thermodynamics. Some of these relations are applicable for systems in non-equilibrium steady state [15], while others hold in the transient regime. Fluctuation relations with underlying Hamiltonian dynamics [6, 13] as well as stochastic dynamics [10, 12, 14, 16] have also been proposed. There are ongoing efforts to generalize and broaden the applicability of XFT in the quantum scenario [8, 10, 17, 19, 21, 22].

The fluctuation-exchange relations can be considered as generalizations of second law of thermodynamics for small systems and they connect the probabilities of appearance of physical quantities such as work, heat, number of particles, in an experimental set up, to those obtainable in a time-reversed set up. For instance, the Jarzynski-Wótcik fluctuation theorem (XFT) [5] given by,

$$\ln \left[\frac{p_\tau(+\mathcal{Q})}{p_\tau(-\mathcal{Q})} \right] = \Delta \beta \mathcal{Q}, \quad \Delta \beta = \frac{1}{k T_B} - \frac{1}{k T_A} \quad (1)$$

quantifies the ratio of probability $p_\tau(+\mathcal{Q})$ of heat exchange during interaction of A and B for a fixed time duration τ , to its time-reversed counterpart $p_\tau(-\mathcal{Q})$. Here k denotes Boltzmann constant and \mathcal{Q} denotes the amount of heat exchanged.

The Jarzynski-Wótcik theorem [5] is one among the important XFTs and has drawn much attention. Microreversibility and the strict directionality of thermodynamical heat flow form the foundational features of this XFT relation. Generalizations of the theorem to include processes involving a system coupled to reservoirs [20], a chain of interacting particles connecting two heat baths [15], correlated thermal quantum systems [19, 22] and the like have been carried out.

In the classical scenario Jarzynski and Wótcik [5] had employed phase space description, for the forward and reverse dynamical evolution between statistical systems A and B in thermal equilibriums at temperatures T_A and T_B respectively, so that one gains physical intuition underlying the relation between heat exchange and fluctuations. In the quantum regime, they considered systems with discrete energy levels to arrive at the relation (1).

In the present work, we retain the flavour of phase-space approach in the quantum scenario, by confining ourselves to continuous variable Gaussian thermal states. Following similar lines as that of the original work [5] we arrive at the heat exchange-fluctuation theorem in the quantum realm for two Gaussian states in thermal equilibrium at temperatures T_A, T_B kept in contact with each other. What comes to our aid here

is the fact that the Wigner distribution function, characterizing Gaussian states is non-negative [26] and hence, it serves as a *legitimate* quantum counterpart of phase-space probability distribution. This enables us to carry out explicit evaluations and arrive at the Jarzynski-Wórcik XFT in this case.

We have organized our paper as follows: In Section 2 a brief outline of Jarzynski-Wórcik derivation of heat transfer and fluctuation relation (1) is presented. Necessary mathematical preliminaries on infinite dimensional continuous variable Gaussian systems, their characterization in terms of variance matrix and obtaining Wigner distribution function in terms of the variance matrix are given in Section 3. In Section 4, we derive the Jarzynski-Wórcik heat exchange-fluctuation theorem for Gaussian systems A and B , in thermal equilibrium at temperatures T_A, T_B respectively using the Wigner distribution function approach. Time reversal symmetric canonical transformations of phase space observables is employed to identify explicit forms of forward and backward heat probability distributions using the Wigner distribution function associated with Gaussian thermal states (or equivalently, quantum harmonic oscillator system in thermal equilibrium). Discussions on heat-exchange statistics of quantum and classical harmonic oscillators in thermal equilibrium, possible connection of Jarzynski-Wórcik XFT with energy equipartition theorem in the quantum scenario are given in Section 5.

2. Classical phase-space description for Jarzynski-Wórcik heat exchange-fluctuation relation

Jarzynski and Wórcik considered two systems, phase space evolution of which is governed by Hamiltonians $H_A(\xi_A)$ and $H_B(\xi_B)$; ξ_A, ξ_B denoting phase-space variables (e.g., positions and momenta) of systems A and B respectively. The systems are kept in contact with each other for a time duration τ via an interaction characterized by $H_{\text{int}}(\xi_A, \xi_B)$, which is switched ‘on’ at time $t = 0$, and turned ‘off’ at $t = \tau$. The phase space trajectory of the two systems is denoted collectively by $\xi^t = (\xi_A^t, \xi_B^t)$. Both the systems are initially in thermal equilibrium, at temperatures T_A, T_B respectively, and their phase space probability distributions at time $t = 0$ is given by

$$p(\xi^0) = \frac{e^{-H_A(\xi_A^0)/kT_A} e^{-H_B(\xi_B^0)/kT_B}}{Z_A Z_B} \quad (2)$$

where Z_A, Z_B denote partition functions. Phase-space dynamics of the systems is assumed to be time-reversal symmetric i.e.,

$$\begin{aligned} H_A(\xi_A) &\longrightarrow H_A(\xi_A^*) = H_A(\xi_A) \\ H_B(\xi_B) &\longrightarrow H_B(\xi_B^*) = H_B(\xi_B) \\ H_{\text{int}}(\xi) &\longrightarrow H_{\text{int}}(\xi^*) = H_{\text{int}}(\xi) \end{aligned} \quad (3)$$

where time reversal operation is denoted by the superscript symbol (*). In other words, for every legitimate forward trajectory ξ^0 to ξ^τ , there exists a time-reversed trajectory

$\bar{\xi}^0 = \xi^{\tau*}$ to $\bar{\xi}^\tau = \xi^{0*}$. Their likelihood ratio is given by (see (2))

$$\begin{aligned} \frac{p(\xi^0)}{p(\bar{\xi}^0)} &= e^{(H_A(\bar{\xi}_A^0) - H_A(\xi_A^0))/kT_A} e^{(H_B(\bar{\xi}^0) - H_B(\xi_A^0))/kT_B} \\ &= e^{\Delta E_A/kT_A} e^{\Delta E_B/kT_B} \end{aligned} \quad (4)$$

where $\Delta E_A = H_A(\xi_A^{\tau*}) - H_A(\xi_A^0)$, $\Delta E_B = H_B(\xi_B^{\tau*}) - H_B(\xi_B^0)$ denote change of internal energies of systems A and B respectively. Assuming that the interaction term H_{int} is negligible, it is seen that $H_A(\xi_A^0) + H_B(\xi_B^0) \approx H_A(\xi_A^{\tau*}) + H_B(\xi_B^{\tau*})$ or $\Delta E_A \approx -\Delta E_B$. Net energy change during the interaction represents the amount of heat transferred i.e., $\mathcal{Q} = \Delta E_A \approx -\Delta E_B$. The heat transfer \mathcal{Q} from A to B during forward process gets compensated by that in the reverse process from B to A and is expressed by

$$\mathcal{Q}(\xi^0) = -\mathcal{Q}(\bar{\xi}^0). \quad (5)$$

Thus, it is seen that

$$\frac{p(\xi^0)}{p(\bar{\xi}^0)} = e^{\Delta\beta \mathcal{Q}(\xi^0)}. \quad (6)$$

From (5) and (6) it follows that

$$\begin{aligned} p_\tau(\mathcal{Q}) &= \int d\xi^0 p(\xi^0) \delta(\xi^0 - \mathcal{Q}) \\ p_\tau(-\mathcal{Q}) &= \int d\bar{\xi}^0 p(\bar{\xi}^0) \delta(\bar{\xi}^0 + \mathcal{Q}) \\ &= e^{-\Delta\beta \mathcal{Q}} p_\tau(\mathcal{Q}) \\ \implies \frac{p_\tau(\mathcal{Q})}{p_\tau(-\mathcal{Q})} &= e^{\Delta\beta \mathcal{Q}}. \end{aligned} \quad (7)$$

thus proving the Jarzynski-Wójcik heat exchange fluctuation theorem in the classical scenario.

In the quantum realm Jarzynski and Wójcik considered two discrete level systems prepared initially in thermal equilibrium at temperatures T_A , T_B and measure their energies E_i^A , E_i^B ; the systems are allowed to interact weakly for a time duration τ after interaction is turned off and energies of both the systems E_f^A , E_f^B measured. As the systems are allowed to interact weakly it is expected that the total energy of the system is conserved: $E_i^A + E_i^B \approx E_f^A + E_f^B$. Heat transfer is then interpreted as $\mathcal{Q}_{i \rightarrow f} = E_i^A - E_f^A \approx E_f^B - E_i^B$ resulting in the relation

$$\ln \left[\frac{p \left(|i\rangle \xrightarrow{\tau} |f\rangle \right)}{p \left(|f\rangle \xrightarrow{-\tau} |i\rangle \right)} \right] = \Delta\beta \mathcal{Q}_{i \rightarrow f}. \quad (8)$$

Our interest here is to derive the relation (7) describing heat transfer processes in the forward and time-reversed dynamics of quantum Gaussian system consisting of two subsystems A , B , prepared initially in thermal equilibrium at temperatures T_A , T_B respectively. Wigner distribution function formalism [26] is employed in this approach. To this end, we give necessary mathematical preliminaries on quantum phase space description and symplectic evolution, Gaussian thermal states and the associated Wigner distribution function in Section 3.

3. Gaussian states – description through variance matrix and Wigner function

Gaussian states, the most important among continuous variable states [23, 24], find diverse applications in several fields including quantum stochastic processes and open system dynamics [25]. They naturally occur as the thermal equilibrium states of any physical system in the small oscillations limit [25]. Being fully characterized by its first and second moments Gaussian states are simpler to handle among the continuous variable states.

Any arbitrary two-mode Gaussian state $\hat{\rho}_{AB}$ is characterized completely by its first and second order moments, written concisely in the form of a 4×4 covariance matrix \mathbf{V} (referred to as variance matrix from now on), which is defined in terms of its elements as [23, 24]

$$V_{ij} = \frac{1}{2} \left\langle \left\{ \hat{\xi}_i, \hat{\xi}_j \right\} \right\rangle - \langle \hat{\xi}_i \rangle \langle \hat{\xi}_j \rangle, \quad i, j = 1, 2, 3, 4. \quad (9)$$

Here $\hat{\xi}_i$ denotes a 4×1 column $\hat{\xi}$ with positions and momenta (dimensionless) as its components:

$$\hat{\xi} = (\hat{q}_A, \hat{p}_A, \hat{q}_B, \hat{p}_B)^T, \quad (10)$$

where ‘T’ stands for transpose operation; we have denoted $\left\{ \hat{\xi}_i, \hat{\xi}_j \right\} = \hat{\xi}_i \hat{\xi}_j + \hat{\xi}_j \hat{\xi}_i$ and we have denoted $\langle \dots \rangle = \text{Tr}(\hat{\rho} \dots)$ in (9). The canonical phase space variables $\hat{q}_\alpha, \hat{p}_\alpha$, $\alpha, \beta = A, B$ satisfy the Bosonic commutation relations,

$$\begin{aligned} [\hat{q}_\alpha, \hat{q}_\beta] &= 0, & [\hat{p}_\alpha, \hat{p}_\beta] &= 0, \quad \text{and} \\ [\hat{q}_\alpha, \hat{p}_\beta] &= i \delta_{\alpha\beta}; \end{aligned} \quad (11)$$

where $\delta_{\alpha\beta} = 1$ when $\alpha = \beta$ and zero when $\alpha \neq \beta$, is the Kronecker delta function. In terms of the components $\hat{\xi}_i$, $i = 1, 2, 3, 4$, the canonical commutation relations (11) assume the form

$$[\hat{\xi}_i, \hat{\xi}_j] = i \Omega_{ij}, \quad i, j = 1, 2 \quad (12)$$

where Ω_{ij} denote elements of the 4×4 matrix

$$\mathbf{\Omega} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}. \quad (13)$$

The commutation relations (see (11), (12)) remain invariant under a *symplectic* transformation [23, 24]:

$$\mathbf{S} \mathbf{\Omega} \mathbf{S}^T = \mathbf{\Omega}. \quad (14)$$

The set of all 4×4 real matrices \mathbf{S} satisfying the property (14) constitutes the symplectic group of real linear canonical transformations $\text{Sp}(4, \mathbb{R})$ [23, 24].

The variance matrix \mathbf{V} of a two-mode quantum system given explicitly by

$$\mathbf{V} = \begin{pmatrix} \langle q_A^2 \rangle & \frac{1}{2}\langle \{q_A, p_A\} \rangle & \langle q_A q_B \rangle & \langle q_A p_B \rangle \\ \frac{1}{2}\langle \{q_A, p_A\} \rangle & \langle p_A^2 \rangle & \langle p_A q_B \rangle & \langle p_A p_B \rangle \\ \langle q_A q_B \rangle & \langle q_B p_A \rangle & \langle q_B^2 \rangle & \frac{1}{2}\langle \{q_B, p_B\} \rangle \\ \langle q_A p_B \rangle & \langle p_A p_B \rangle & \frac{1}{2}\langle \{q_B, p_B\} \rangle & \langle p_B^2 \rangle \end{pmatrix} \quad (15)$$

is a real symmetric positive definite matrix and it completely characterizes a two mode Gaussian state $\hat{\rho}_{AB}$. Under the symplectic transformation $\hat{\xi}' = \mathbf{S}\hat{\xi}$, the variance matrix \mathbf{V} undergoes a congruent transformation $\mathbf{V}' = \mathbf{S}\mathbf{V}\mathbf{S}^T$, where \mathbf{V}' is the variance matrix associated with the new canonical variables $\hat{q}'_\alpha, \hat{p}'_\beta$, $\alpha, \beta = A, B$.

From the fundamental theorem due to Williamson [27] it follows that variance matrix \mathbf{V} attains a canonical form under symplectic transformation \mathbf{S}_W such that

$$\mathbf{V}_W = \mathbf{S}_W \mathbf{V} \mathbf{S}_W^T = \text{diag}(\nu_A, \nu_A; \nu_B, \nu_B) \quad (16)$$

and \mathbf{V}_W is referred to as the *Williamson normal form* of the variance matrix and ν_A, ν_B are called the symplectic eigenvalues of the variance matrix. The real positive diagonal elements ν_α , $\alpha = A, B$ of \mathbf{V}_W are the positive square roots of the doubly degenerate eigenvalues of the matrix $-(\mathbf{V}\boldsymbol{\Omega})^2$ as,

$$-\mathbf{S}_W (\mathbf{V}\boldsymbol{\Omega})^2 \mathbf{S}_W^{-1} = -(\mathbf{V}_W\boldsymbol{\Omega})^2 = \text{diag}(\nu_A^2, \nu_A^2, \nu_B^2, \nu_B^2). \quad (17)$$

Corresponding to the symplectic transformation \mathbf{S}_W , there exists a unitary operator $\mathbf{U}(\mathbf{S}_W)$ transforming the density matrix $\hat{\rho}_{AB}$ of a two mode Gaussian state as follows:

$$\hat{\rho}_{AB} = \mathbf{U}^\dagger(\mathbf{S}_W) (\hat{\rho}_{\nu_A} \otimes \hat{\rho}_{\nu_B}) \mathbf{U}(\mathbf{S}_W) \quad (18)$$

where the single mode density matrices $\hat{\rho}_{\nu_A}, \hat{\rho}_{\nu_B}$ are given by [25]

$$\hat{\rho}_\alpha = \frac{1}{\nu_\alpha + \frac{1}{2}} \sum_{n_\alpha=0}^{\infty} \left(\frac{\nu_\alpha - \frac{1}{2}}{\nu_\alpha + \frac{1}{2}} \right)^{n_\alpha} |n_\alpha\rangle \langle n_\alpha|, \quad \alpha = A, B. \quad (19)$$

Here $|n_A\rangle, |n_B\rangle$ are the eigenstates of the number operators $\hat{N}_A = \hat{a}_A^\dagger \hat{a}_A$, $\hat{N}_B = \hat{a}_B^\dagger \hat{a}_B$ of the modes A, B and $\hat{a}_\alpha, \hat{a}_\alpha^\dagger$ are related to the dimensionless canonical position and momentum observables as follows:

$$\hat{a}_\alpha = \frac{\hat{q}_\alpha + i\hat{p}_\alpha}{\sqrt{2}}, \quad \hat{a}_\alpha^\dagger = \frac{\hat{q}_\alpha - i\hat{p}_\alpha}{\sqrt{2}}. \quad (20)$$

Let us denote $\hat{Q}_\alpha = \left(\frac{\hbar}{m_\alpha \omega_\alpha} \right)^{1/2} \hat{q}_\alpha$, $\hat{P}_\alpha = (m_\alpha \omega_\alpha \hbar)^{1/2} \hat{p}_\alpha$. Given the Hamiltonian of a harmonic oscillator of mass m_α , frequency ω_α ,

$$\begin{aligned} \hat{H}_\alpha &= \frac{\hat{P}_\alpha^2}{2m_\alpha} + \frac{1}{2} m_\alpha \omega_\alpha^2 \hat{Q}_\alpha^2, \\ &= \left(\hat{N}_\alpha + \frac{1}{2} \right) \omega_\alpha, \end{aligned} \quad (21)$$

a canonical ensemble of Bosonic oscillators in thermal equilibrium at temperature T_α is described by

$$\hat{\rho}_{T_\alpha} = \frac{e^{-\hat{H}_\alpha/kT_\alpha}}{Z_\alpha} = \frac{e^{-\hbar\omega_\alpha/2kT_\alpha}}{Z_\alpha} \sum_{n_\alpha=0}^{\infty} e^{-n_\alpha\hbar\omega_\alpha/kT_\alpha} |n_\alpha\rangle \langle n_\alpha| \quad (22)$$

where

$$\begin{aligned} Z_\alpha &= \text{Tr}[e^{-\hat{H}/kT_\alpha}] = \text{Tr}\left[e^{-\frac{\hbar\omega_\alpha}{kT_\alpha}(\hat{N}_\alpha + \frac{1}{2})}\right] \\ &= \frac{e^{-\hbar\omega_\alpha/2kT_\alpha}}{1 - e^{-\hbar\omega_\alpha/kT_\alpha}} \end{aligned} \quad (23)$$

denotes the partition function. The single mode thermal state

$$\hat{\rho}_{T_\alpha} = (1 - e^{-\hbar\omega_\alpha/kT_\alpha}) \sum_{n_\alpha=0}^{\infty} e^{-n_\alpha\hbar\omega_\alpha/kT_\alpha} |n_\alpha\rangle\langle n_\alpha|$$

may be readily identified with the canonical single mode Gaussian state ρ_α of (19) appearing in the Williamson canonical decomposition (18).

The variance matrix \mathbf{V} of any two-mode thermal state $\hat{\rho}_{AB} = \hat{\rho}_{T_A} \otimes \hat{\rho}_{T_B}$ is given by [24]

$$\begin{aligned} \mathbf{V} &= \begin{pmatrix} \mathbf{V}_A & 0 \\ 0 & \mathbf{V}_B \end{pmatrix} \\ \mathbf{V}_A &= \frac{1}{2} \coth\left(\frac{\hbar\omega_A}{2kT_A}\right) \mathbb{1}_2, \quad \mathbf{V}_B = \frac{1}{2} \coth\left(\frac{\hbar\omega_B}{2kT_B}\right) \mathbb{1}_2 \end{aligned} \quad (24)$$

where $\mathbb{1}_2$ denotes 2×2 identity matrix. Note that \mathbf{V} is in the Williamson normal form, with its symplectic eigenvalues related to the frequency and temperature of the thermal state of oscillator systems A and B as,

$$\nu_{T_\alpha} = \frac{1}{2} \coth\left(\frac{\hbar\omega_\alpha}{2kT_\alpha}\right), \quad \alpha = A, B. \quad (25)$$

In the next subsection we give an outline of some preliminary notions on quantum phase space formalism in terms of Wigner distribution functions.

3.1. Wigner distribution function of a Gaussian state

Wigner distribution function plays a central role in developing quantum phase space formalism involving non-commuting canonical observables $\hat{q}_\alpha, \hat{p}_\alpha$. In this subsection we focus on some preliminary notions of Wigner function associated with single mode continuous variable quantum system.

Wigner function $W(q, p)$ of a single mode continuous variable quantum state $\hat{\rho}$ is a real function of phase space canonical variables q, p defined by [26]

$$W(q, p) = \frac{1}{\pi} \int_{-\infty}^{\infty} dx \langle q-x | \hat{\rho} | q+x \rangle e^{2i\pi px/\hbar} \quad (26)$$

where $|q\rangle$ denotes eigenvector of the operator \hat{q} ; it satisfies the normalization property [26]

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} W(q, p) dq dp = \text{Tr}(\hat{\rho}) = 1 \quad (27)$$

which is true of any probability distribution and gives correct marginal probability distributions:

$$\int_{-\infty}^{\infty} W(q, p) dp = \langle q | \hat{\rho} | q \rangle, \quad \int_{-\infty}^{\infty} W(q, p) dq = \langle p | \hat{\rho} | p \rangle \quad (28)$$

Quantum expectation value of any operator $\hat{f}(\hat{q}, \hat{p})$ in a state $\hat{\rho}$ can be replaced by a phase space integration using Wigner function $W(q, p)$ as,

$$\text{Tr} \left[\hat{\rho} \hat{f}(\hat{q}, \hat{p}) \right] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} W(q, p) f(q, p) dq dp. \quad (29)$$

where Weyl's correspondence rule

$$q^k p^l \longleftrightarrow \frac{1}{2^k} \sum_{r=0}^k \frac{k!}{r! (k-r)!} \hat{q}^r \hat{p}^l \hat{q}^{k-r} \quad (30)$$

to associate classical functions with quantum operators has been employed [26].

Wigner function formalism allows phase space description (see (27), (28), (29)) of quantum theory in a classical language. However the function $W(q, p)$ is not necessarily non-negative everywhere and hence it is termed as *quasi*-probability distribution function [26].

Interestingly, the Wigner function associated with single mode quantum Gaussian states given by [24]

$$W(\xi) = \frac{1}{2\pi \sqrt{\det \mathbf{V}}} e^{-\frac{1}{2} \xi^T \mathbf{V}^{-1} \xi} \quad (31)$$

is non-negative everywhere. Here, \mathbf{V} is the variance matrix of the Gaussian state and phase space canonical variables are expressed compactly in the form of a column $\xi = (q, p)^T$. Wigner function of a single mode Gaussian state in thermal equilibrium at temperature T [26] takes the following form (by substituting the variance matrix of a thermal state (see (24)) and after simplification):

$$\begin{aligned} W(q, p) &= \frac{1}{2\pi \nu_T} \exp \left[-\frac{1}{2\nu_T} (q^2 + p^2) \right] \\ &= \frac{1}{2\pi \nu_T} \exp \left[-\frac{H(Q, P)}{\hbar\omega \nu_T} \right] \end{aligned} \quad (32)$$

where $H(Q, P) = \frac{P^2}{2m} + \frac{1}{2} m\omega^2 Q^2$, $Q = (\hbar/m\omega)^{1/2} q$, $P = (m\omega\hbar)^{1/2} p$ and $\nu_T = \frac{1}{2} \coth \left(\frac{\hbar\omega}{2kT} \right)$ is the symplectic eigenvalue of the variance matrix (see (25)). It is evident that $W(q, p)$ of (32) is a Gaussian function of q, p and is non-negative.

Using the Wigner function description it is possible to develop a time-reversal symmetric phase space trajectory approach to derive heat exchange fluctuation relation analogous to the Jarzynski-Wójcik approach in the classical scenario (as outlined in Section 2).

We discuss heat exchange statistics between two Gaussian systems, in thermal equilibrium at different temperatures in the next section.

4. Heat exchange fluctuation theorem for Gaussian thermal states

Let us consider two quantum systems A and B characterized by their respective Hamiltonians

$$\hat{H}_A(\hat{\xi}_A) = \frac{\hat{P}_A^2}{2m_A} + \frac{1}{2} m_A \omega_A^2 \hat{Q}_A^2, \quad \hat{H}_B(\hat{\xi}_B) = \frac{\hat{P}_B^2}{2m_B} + \frac{1}{2} m_B \omega_B^2 \hat{Q}_B^2, \quad (33)$$

where $\hat{\xi}_\alpha = (\hat{q}_\alpha, \hat{p}_\alpha)^T$, $\hat{q}_\alpha = \sqrt{\frac{m_\alpha \omega_\alpha}{\hbar}} \hat{Q}_\alpha$, $\hat{p}_\alpha = \frac{1}{\sqrt{m_\alpha \omega_\alpha \hbar}} \hat{P}_\alpha$, $\alpha = A, B$.

Let the systems be prepared in a thermal state at temperatures T_A, T_B respectively i.e.,

$$\hat{\rho}_{T_A} = \frac{e^{-\hat{H}_A(\hat{\xi}_A)/kT_A}}{Z_A}, \quad \hat{\rho}_{T_B} = \frac{e^{-\hat{H}_B(\hat{\xi}_B)/kT_B}}{Z_B}. \quad (34)$$

The Wigner function $W(\xi^0)$ at time $t = 0$ corresponding to the two-mode Gaussian thermal state $\hat{\rho}_{AB}^0 = \hat{\rho}_{T_A}^0 \otimes \hat{\rho}_{T_B}^0$ is a product of Wigner functions $W(\xi_A^0), W(\xi_B^0)$, where $\xi^0 = (\xi_A^0, \xi_B^0)^T$ and $\xi_A^0 = (q_A^0, p_A^0)^T$, $\xi_B^0 = (q_B^0, p_B^0)^T$ are classical phase space columns at $t = 0$. Using (32) we obtain

$$W(\xi^0) = \frac{1}{(2\pi)^2 \nu_{T_A} \nu_{T_B}} \exp \left[- \left(\frac{\xi_A^{0T} \xi_A^0}{2\nu_{T_A}} + \frac{\xi_B^{0T} \xi_B^0}{2\nu_{T_B}} \right) \right] \quad (35)$$

$$= \frac{1}{(2\pi)^2 \nu_{T_A} \nu_{T_B}} \exp \left[- \left(\frac{H_A(\xi_A^0)}{\hbar \omega_A \nu_{T_A}} + \frac{H_B(\xi_B^0)}{\hbar \omega_B \nu_{T_B}} \right) \right] \quad (36)$$

where $H_\alpha(\xi_\alpha^0) = \frac{P_\alpha^2}{2m} + \frac{1}{2} m \omega_\alpha^2 Q_\alpha^2$, $Q_\alpha = (\hbar/m_\alpha \omega_\alpha)^{1/2} q_\alpha$, $P_\alpha = (m_\alpha \omega_\alpha \hbar)^{1/2} p_\alpha$, $\alpha = A, B$. The systems A and B are kept in contact with each other in terms of a *quadratic* interaction Hamiltonian (representing a canonical transformation in phase space) $\hat{H}_{\text{int}}(\hat{\xi}^t)$, which is turned on and off at time $t = 0, t = \tau$ respectively.

Corresponding to the unitary time evolution operator

$$\hat{U}(\hat{\xi}^\tau) = \exp \left[-\frac{i\tau}{\hbar} \left(\hat{H}_A(\hat{\xi}_A^\tau) + \hat{H}_B(\hat{\xi}_B^\tau) + \hat{H}_{\text{int}}(\hat{\xi}^\tau) \right) \right] \quad (37)$$

there exists a 4×4 real symplectic matrix [23, 24] $\mathbf{S}^\tau \in \text{Sp}(4, \mathbb{R})$ which acts on the column of canonical operators as $\hat{\xi}^0 = (\hat{q}_A^0, \hat{p}_A^0; \hat{q}_B^0, \hat{p}_B^0)$ resulting in

$$\hat{\xi}^0 \xrightarrow{\mathbf{S}^\tau} \hat{\xi}^\tau = \mathbf{S}^\tau \hat{\xi}^0 = \hat{\xi}^\tau. \quad (38)$$

Consequently, the Wigner function undergoes the transformation: $W(\xi^0) \rightarrow W(\xi^\tau)$, $\xi^\tau = (q_A^\tau, p_A^\tau; q_B^\tau, p_B^\tau)^T$. We thus obtain,

$$W(\xi^\tau) = \frac{1}{(2\pi)^2 \nu_{T_A} \nu_{T_B}} \exp \left[- \left(\frac{H_A(\xi_A^\tau)}{\hbar \omega_A \nu_{T_A}} + \frac{H_B(\xi_B^\tau)}{\hbar \omega_B \nu_{T_B}} \right) \right] \quad (39)$$

Under Time-reversal operation one finds that $\hat{\xi} \rightarrow \hat{\xi}^* = (\hat{q}_A, -\hat{p}_A; \hat{q}_B, -\hat{p}_B)^T$. The unitary dynamics is chosen to be invariant under time-reversal i.e.,

$$\begin{aligned} \hat{H}_A(\hat{\xi}_A) &\rightarrow \hat{H}_A(\hat{\xi}_A^*) = \hat{H}_A(\hat{\xi}_A) \\ \hat{H}_B(\hat{\xi}_B) &\rightarrow \hat{H}_B(\hat{\xi}_B^*) = \hat{H}_B(\hat{\xi}_B) \\ \hat{H}_{\text{int}}(\hat{\xi}) &\rightarrow \hat{H}_{\text{int}}(\hat{\xi}^*) = \hat{H}_{\text{int}}(\hat{\xi}). \end{aligned} \quad (40)$$

Consider dynamical evolution of the system under time-reversal, transforming the phase space column of observables $\hat{\xi}^{0*} = \hat{\xi}^{\tau}$ to $\hat{\xi}^{\tau*} = \hat{\xi}^0$. Ratio of the Wigner functions $W(\xi^0)/W(\bar{\xi}^0)$ is then given by (see (35) and (39)),

$$\begin{aligned} \frac{W(\xi^0)}{W(\bar{\xi}^0)} &= \frac{W(\xi^0)}{W(\xi^{\tau*})} \\ &= \exp \left[\left(\frac{H_A(\xi_A^{\tau*}) - H_B(\xi_A^0)}{\hbar\omega_A \nu_{T_A}} \right) \right] \exp \left[\left(\frac{H_B(\xi_B^{\tau*}) - H_B(\xi_B^0)}{\hbar\omega_B \nu_{T_B}} \right) \right] \\ &= \exp \left[\frac{\Delta E_A}{\hbar\omega_A \nu_{T_A}} \right] \exp \left[\frac{\Delta E_B}{\hbar\omega_B \nu_{T_B}} \right] \end{aligned} \quad (41)$$

where

$$\Delta E_A = H_A(\xi_A^{\tau*}) - H_A(\xi_A^0), \quad \Delta E_B = H_B(\xi_B^{\tau*}) - H_B(\xi_B^0). \quad (42)$$

Considering a weak interaction $\hat{H}_{\text{int}}(\hat{\xi}^t)$, it is deduced that

$$\begin{aligned} H_A(\xi_A^0) + H_B(\xi_B^0) &\approx H_A(\xi_A^{\tau*}) + H_B(\xi_B^{\tau*}) \\ \Rightarrow \Delta E_A &\approx -\Delta E_B. \end{aligned} \quad (43)$$

implying that the net change in internal energy of system A is compensated by an opposite change in the internal energy of system B , when the phase space trajectory ξ^t is sampled. In other words, heat transfer during forward realization $\mathcal{Q}(\xi^0) = \Delta E_A$ is opposite to that of the reverse realization i.e., $\mathcal{Q}(\bar{\xi}^0) = \Delta E_B = -\mathcal{Q}(\xi^0)$. Thus, we obtain

$$\frac{W(\xi^0)}{W(\bar{\xi}^0)} = e^{\Delta\beta_\omega \mathcal{Q}(\xi^0)}, \quad (44)$$

where

$$\begin{aligned} \Delta\beta_\omega &= \frac{1}{\hbar\omega_A \nu_{T_A}} - \frac{1}{\hbar\omega_B \nu_{T_B}} \\ &= \frac{2 \tanh\left(\frac{\hbar\omega_A}{2kT_A}\right)}{\hbar\omega_A} - \frac{2 \tanh\left(\frac{\hbar\omega_B}{2kT_B}\right)}{\hbar\omega_B} \end{aligned} \quad (45)$$

Heat distribution $p_\tau(\mathcal{Q})$ can then be expressed in terms of the Wigner function as

$$\begin{aligned} p_\tau(\mathcal{Q}) &= \int d\xi^0 W(\xi^0) \delta(\mathcal{Q} - \mathcal{Q}(\xi^0)) \\ &= e^{\Delta\beta_\omega \mathcal{Q}} \int d\bar{\xi}^0 W(\bar{\xi}^0) \delta(\mathcal{Q} + \mathcal{Q}(\bar{\xi}^0)) \\ &= e^{\Delta\beta_\omega \mathcal{Q}} p_\tau(-\mathcal{Q}). \end{aligned} \quad (46)$$

We obtain

$$\ln \left(\frac{p_\tau(\mathcal{Q})}{p_\tau(-\mathcal{Q})} \right) = \Delta\beta_\omega \mathcal{Q} \quad (47)$$

where $\Delta\beta_\omega$ is given by (45). We thus arrive at a more general quantum Jarzynski-Wózcik heat-exchange fluctuation relation (47) for Gaussian thermal states.

5. Discussions

Some relevant discussions on different forms (1), (8), (47) of the Jarzynski-Wózcik heat-exchange fluctuation relations are summarised in the following.

- (i) In the high temperature limit $\frac{\hbar\omega}{kT} \rightarrow 0$ we get $\Delta\beta_\omega \rightarrow \Delta\beta$. Thus the quantum heat-exchange fluctuation relation (8) reduces to its classical analogue (1) in this limit.
- (ii) Equipartition theorem plays a fundamental role in classical statistical physics. It states that for a system in thermal equilibrium at temperature T the average energy per degree of freedom is given by $\frac{1}{2}kT$. Equipartition theorem of energy holds universally in classical statistical physics as it neither depends on the number of particles in the ensemble nor on the nature of the potential acting on the particles. For a system of one dimensional classical harmonic oscillators, in thermal equilibrium at temperature T , contribution to the average energy comes from mean kinetic energy and mean potential energy i.e., $\langle E \rangle = kT$. It has been pointed out [28, 29, 30] recently that the classical energy equipartition theorem does not hold in the quantum realm. It is seen that the average energies in the state $\rho_{AB} = \rho_{T_A} \otimes \rho_{T_B}$ (see (34)) – characterizing a system of quantum harmonic oscillators A and B in thermal equilibrium at temperatures T_A , T_B respectively – are given by

$$\begin{aligned} \langle \hat{H}_A \rangle &= \text{Tr}[\rho_{T_A} \hat{H}_A] \\ &= \frac{\hbar\omega_A}{2} (\langle \hat{q}_A^2 \rangle + \langle \hat{p}_A^2 \rangle) \\ &= \frac{\hbar\omega_A}{2} \coth\left(\frac{\hbar\omega_A}{2kT_A}\right) = \hbar\omega_A \nu_{T_A} \end{aligned} \quad (48)$$

$$\begin{aligned} \langle \hat{H}_B \rangle &= \text{Tr}[\rho_{T_B} \hat{H}_B] \\ &= \frac{\hbar\omega_B}{2} (\langle \hat{q}_B^2 \rangle + \langle \hat{p}_B^2 \rangle) \\ &= \frac{\hbar\omega_B}{2} \coth\left(\frac{\hbar\omega_B}{2kT_B}\right) = \hbar\omega_B \nu_{T_B} \end{aligned} \quad (49)$$

where we have made use of (15), (24) and (33).

Here the average energies (48), (49) depend on frequencies ω_A, ω_B (indicative of the nature of the potential) besides temperatures T_A, T_B . The factor $\Delta\beta_\omega$ in the Jarzynski-Wózcik heat exchange fluctuation relation (8) approaches its classical analogue $\Delta\beta$ of (1) only in the high temperature limit (see (i) above). Deviations of (8) from (1) could be attributed to the fact that classical energy equipartition is no longer valid in the quantum scenario. A series of recent papers [28, 29, 30] have proposed quantum counterpart of energy equipartition theorem, which may shed more light on quantum heat exchange statistics of thermal harmonic oscillator system.

- (iii) The heat distribution associated with systems of thermal quantum harmonic

oscillators, weakly coupled to a heat reservoir at different temperatures, has been derived by Denzler and Lutz [31] by solving the quantum master equation. It has been shown that the heat distribution satisfies Jarzynski-Wózcik XFT (8). Authors of Ref. [31] implement measurements on the discrete energy levels of the quantum thermal oscillator systems at time $t = 0$ and $t = \tau$ to arrive at the corresponding XFT (8) - as prescribed originally by Jarzynski-Wózcik [5]. We have employed quantum phase space formalism using Wigner function for Gaussian thermal states to arrive at the XFT (47), classical limit of which can be readily seen.

We believe that quantum phase-space formalism to study Jarzynski-Wózcik XFT opens up new perspectives towards understanding the heat distribution statistics and its classical limit.

Acknowledgements

Sudha and ARU acknowledge financial support from the Department of Science and Technology(DST), India through Project No. DST/ICPS/QUST/2018/107. AMJ thanks DST, India for J.C.Bose National fellowship.

References

- [1] Jarzynski C 1997 Nonequilibrium Equality for Free Energy Differences *Phys. Rev. Lett* **78** 2690–2693
- [2] Jarzynski C 1997 Equilibrium free-energy differences from nonequilibrium measurements: A master-equation approach *Phys. Rev. E* **56** 5018–5035
- [3] Crooks G E 1998 Nonequilibrium Measurements of Free Energy Differences for Microscopically Reversible Markovian Systems, *J.Stat.Phys.* **90** 1481–1487
- [4] Crooks G E 1999 Entropy production fluctuation theorem and the nonequilibrium work relation for free energy differences *Phys. Rev. E* **60** 2721–2726
- [5] Jarzynski C and Wózcik D K 2004 Classical and Quantum Exchange Theorems for heat exchange *Phys. Rev. Lett.* **92** 230602
- [6] Jarzynski C 2007 Comparison of far-from-equilibrium work relations *C. R. Phys* **8** 495–506
- [7] Andrieux D, Gaspard P, Monnai T and Tasaki S 2009 The fluctuation theorem for currents in open quantum systems *New. J. Phys* **11** 043014
- [8] Esposito M, Harbola U and Mukamel S 2009 Nonequilibrium fluctuations, fluctuation theorems and counting statistics in quantum systems *Rev. Mod. Phys* **81** 1665-1702
- [9] Campisi M, Talkner P and Hänggi P 2010 Fluctuation Theorems for Continuously Monitored Quantum Fluxes *Phys. Rev. Lett* **105**, 140601
- [10] Campisi M, Hänggi P and Talkner P, 2011 Quantum fluctuation relations: Foundations and applications *Rev. Mod. Phys* **83** 771–791
- [11] Cohen D and Imry Y 2012 Straightforward quantum-mechanical derivation of the Crooks fluctuation theorem and the Jarzynski equality *Phys. Rev. E* **86** 011111
- [12] Rana S, Lahiri S, Jayannavar A M 2012 Quantum Jarzynski equality with multiple measurement and feedback for isolated systems *Pramana J. Phys* **79** 233–241
- [13] Deffner S and Jarzynski C 2013 Information processing and the second law of thermodynamics *Phys. Rev. X* **3** 041003

- [14] Rana S, Lahiri S, Jayannavar A M 2013 Generalized entropy production fluctuation theorems for quantum systems *Pramana J. Phys* **80** 207–222
- [15] Lahiri S, Jayannavar A M 2014 Exchange fluctuation theorems for a chain of interacting particles in presence of two heat baths *Eur. Phys. J. B* **87** 141
- [16] Lahiri S, Jayannavar A M 2015 Derivation of not-so-common fluctuation theorems *Indian J. Phys* **89** 515–523
- [17] Alberg J 2018 Fully quantum fluctuation theorems *Phys. Rev. X* **8**, 011019
- [18] Hänggi P and Talkner P 2015 The other QFT *Nature Physics* **11** 108–110
- [19] Jevtic S, Rudolph T, Jennings D, Hirano Y, Nakayama S and Murao M 2015 Exchange fluctuation theorems for correlated quantum systems *Phys. Rev. E* **92** 042113
- [20] Pal P S, Lahiri S and Jayannavar A M 2017 Transient Exchange Fluctuation Theorem for heat using a hamiltonian framework *Phys. Rev. E* **95** 042124
- [21] Hänggi P and Talkner P 2019 Statistical mechanics and thermodynamics at strong coupling: Quantum and classical arXiv:1911.11660v3 (Reviews of Modern Physics, In press)
- [22] Micadei K, Landi G T and Lutz E 2020 Quantum fluctuation theorems beyond two-point measurements *Phys. Rev. Lett* **124** 090602
- [23] Simon R, Mukunda N, Dutta B 1994 Quantum-noise matrix for multimode systems: $U(n)$ invariance, squeezing, and normal forms *Physical Review A* **49** 1567–1583
- [24] Arvind, Dutta B, Mukunda N, Simon R 1995 The real symplectic groups in quantum mechanics and optics *Pramana - J. Phys.* **45** 471–477
- [25] Gerardo Adesso, Sammy Ragy , Antony R Lee 2014 Continuous variable quantum information: Gaussian states and beyond, *Open. Syst. Inf. Dyn.* **21** 1440001
- [26] Hillery M, O’Connell R F, Scully M O and Wigner E P 1984 Distribution functions in Physics: Fundamentals *Phys. Rep.* **106** 121–167
- [27] Williamson J 1936 On the Algebraic Problem Concerning the Normal Forms of Linear Dynamical Systems *Am. J. Math.* **58** 141–163
- [28] Bialas P, Spiechowicz J, Luczka J 2018 Partition of energy for a dissipative quantum oscillator *Sci. Rep* **8** 16080(1–12)
- [29] Bialas P, Spiechowicz J, Luczka J 2019 Quantum analogue of energy equipartition theorem *J. Phys. A* **52** 15LT01
- [30] Luczka J 2020 Quantum counterpart of classical equipartition of energy *J. Stat. Phys.* **179** 839845
- [31] Denzler T, Lutz E, 2018 Heat distribution of quantum thermal oscillator *Phys. Rev. E* **98** 052106