

# THE GENERALIZED SPECTRA IN $C^*$ -ALGEBRAS OF OPERATORS OVER $C^*$ -ALGEBRAS

STEFAN IVKOVIĆ

**ABSTRACT.** In this paper we consider shift operators, self-adjoint, unitary and normal operators on the standard module over a unital  $C^*$ -algebra  $\mathcal{A}$ . We define various generalized spectra in  $\mathcal{A}$  of these operators, give description of such spectra of these operators and investigate their further properties.

The Mathematical Institute of the Serbian Academy of Sciences and Arts,  
p.p. 367, Kneza Mihaila 36, 11000 Beograd, Serbia,  
Tel.: +381-69-774237  
email: stefan.iv10@outlook.com

## 1. INTRODUCTION

Hilbert  $C^*$ -modules are natural generalizations of Hilbert spaces when the field of scalars is replaced by an arbitrary  $C^*$ -algebra. In this paper we let  $\mathcal{A}$  be a unital  $C^*$ -algebra and  $H_{\mathcal{A}}$  denote the standard module over  $\mathcal{A}$ , that is  $H_{\mathcal{A}} = l_2(\mathcal{A})$ . Moreover, we let  $B^a(H_{\mathcal{A}})$  be the  $C^*$ -algebra of all  $\mathcal{A}$ -linear, bounded, adjointable operators on  $H_{\mathcal{A}}$ .

Our starting question is the following: If  $\mathcal{A}$  is a  $C^*$ -algebra, then for  $\alpha \in \mathcal{A}$  could we define in a suitable way the operator  $\alpha I$  on  $H_{\mathcal{A}}$  and the generalized spectra in  $\mathcal{A}$  of operators in  $B^a(H_{\mathcal{A}})$  by setting for every  $F \in B^a(H_{\mathcal{A}})$   $\sigma^{\mathcal{A}}(F) = \{\alpha \in \mathcal{A} \mid F - \alpha I \text{ is not invertible in } B^a(H_{\mathcal{A}})\}$ ?

For  $a \in \mathcal{A}$  we may let  $\alpha I$  be the operator on  $H_{\mathcal{A}}$  given by  $\alpha I(x_1, x_2, \dots) = (\alpha x_1, \alpha x_2, \dots)$ . It is straightforward to check that  $\alpha I$  is an  $\mathcal{A}$ -linear operator on  $H_{\mathcal{A}}$ . Moreover,  $\alpha I$  is bounded and  $\|\alpha I\| = \|\alpha\|$ . Finally,  $\alpha I$  is adjointable and its adjoint is given by  $(\alpha I)^* = \alpha^* I$ .

We introduce then the following notion:

$$\sigma^{\mathcal{A}}(F) = \{\alpha \in \mathcal{A} \mid F - \alpha I \text{ is not invertible in } B^a(H_{\mathcal{A}})\};$$

$$\sigma_p^{\mathcal{A}}(F) = \{\alpha \in \mathcal{A} \mid \ker(F - \alpha I) \neq \{0\}\};$$

$$\sigma_{rl}^{\mathcal{A}}(F) = \{\alpha \in \mathcal{A} \mid F - \alpha I \text{ is bounded below, but not surjective on } H_{\mathcal{A}}\};$$

$$\sigma_{cl}^{\mathcal{A}}(F) = \{\alpha \in \mathcal{A} \mid \text{Im}(F - \alpha I) \text{ is not closed}\}. \text{ (where } F \in B^a(H_{\mathcal{A}})\text{)}.$$

Recall that not all closed submodules of  $H_{\mathcal{A}}$  are orthogonally complementable in  $H_{\mathcal{A}}$  which differs from the situation of Hilbert spaces. It may happen that  $\overline{\text{Im}(F - \alpha I)} \oplus \text{Im}(F - \alpha I)^{\perp} \subsetneq H_{\mathcal{A}}$ . However, if  $\text{Im}(F - \alpha I)$  is closed, then

$Im(F^* - \alpha^*I)$  is closed and we also have  $H_{\mathcal{A}} = Im(F - \alpha I) \oplus \ker(F^* - \alpha^*I) = \ker(F - \alpha I) \oplus Im(F^* - \alpha^*I)$  whenever  $F \in B^a(H_{\mathcal{A}})$ .

Therefore, it is more convinient in this setting to work with  $\sigma_{rl}^{\mathcal{A}}(F)$  and  $\sigma_{cl}^{\mathcal{A}}(F)$  for  $F \in B^a(H_{\mathcal{A}})$  instead of the residual and the continuous spectrum.

Note that we have obviously  $\sigma^{\mathcal{A}}(F) = \sigma_p^{\mathcal{A}}(F) \cup \sigma_{rl}^{\mathcal{A}}(F) \cup \sigma_{cl}^{\mathcal{A}}(F)$  and  $\sigma^{\mathcal{A}}(F^*) = (\sigma^{\mathcal{A}}(F))^*$ .

The challenges which arise are the following:

- 1)  $\mathcal{A}$  may be non commutative;
- 2) Not all non-zero elements in  $\mathcal{A}$  are invertible. Moreover, even if  $\alpha \in \mathcal{A} \cap G(\mathcal{A})$ , we do not have in general that  $\|\alpha^{-1}\| = \frac{1}{\|\alpha\|}$ . Therefore  $\sigma^{\mathcal{A}}(F)$  may be unbounded. ( However  $\sigma^{\mathcal{A}}(F)$  is always closed in  $\mathcal{A}$  ).

The notion of generalized spectra has originally been introduced in [1]. In this paper we give description of the generalized spectra of shift operators, unitary, self-adjoint and normal operators on  $H_{\mathcal{A}}$  and investigate some further properties of these spectra. Most of the results in this paper are generalizations of the results from [2].

## 2. MAIN RESULTS

We start with the following proposition.

**Proposition 2.1.** *Let  $\mathcal{A}$  be a unital  $C^*$ -algebra  $\{e_k\}_{k \in \mathbb{N}}$  denote the standard orthonormal basis of  $H_{\mathcal{A}}$  and  $S$  be the operator defined by  $Se_k = e_{k+1}, k \in \mathbb{N}$ , that is  $S$  is unilateral shift and  $S^*e_{k+1} = e_k$  for all  $k \in \mathbb{N}$ . If  $\mathcal{A} = L^\infty((0, 1))$  or if  $\mathcal{A} = C([0, 1])$ , then  $\sigma^{\mathcal{A}}(S) = \{\alpha \in \mathcal{A} \mid \inf |\alpha| \leq 1\}$ , (where in the case when  $\mathcal{A} = L^\infty((0, 1))$ , we set  $\inf |\alpha| = \inf\{C > 0 \mid \mu(|\alpha|^{-1}[0, C]) > 0\} = \sup\{K > 0 \mid |\alpha| > K\}$  a.e. on  $(0, 1)\}$ ). Moreover,  $\sigma_p^{\mathcal{A}}(S) = \emptyset$  in both cases.*

*Proof.* We have two cases. Case 1: In this case we consider  $\mathcal{A} = C([0, 1])$ . Let  $\alpha \in \mathcal{A}$  and suppose that  $\inf |\alpha| < 1$ . Since  $|\alpha|$  is continuous, we may find an open interval  $(t_1, t_2) \subseteq (0, 1)$  such that  $|\alpha(t)| < 1 - \epsilon$  for all  $t \in (t_1, t_2)$ , where  $0 < \epsilon < 1 - \inf |\alpha|$ . We may find some  $g \in \mathcal{A}$  since that  $\text{supp } g \subseteq (t_1, t_2)$  and  $0 \leq g \leq 1$ . Consider  $x_\alpha = (g, \overline{\alpha}g, \overline{\alpha}^2g, \dots)$ . Then  $\langle (\alpha I - S)e_k, x_\alpha \rangle = \overline{\alpha}^k g - \overline{\alpha}^{k+1} g = 0$ . Hence  $x_\alpha \in Im(\alpha I - S)^\perp$  and  $x_\alpha \neq 0$ , which gives that  $\alpha \in \sigma^{\mathcal{A}}(S)$ . Hence  $\{\alpha \in \mathcal{A} \mid \inf |\alpha| < 1\} \subseteq \sigma^{\mathcal{A}}(S)$ . Since  $\sigma^{\mathcal{A}}(S)$  is closed in the norm topolgy in  $\mathcal{A}$  it follows that  $\inf \{\alpha \in \mathcal{A} \mid \inf |\alpha| \leq 1\} \subseteq \sigma^{\mathcal{A}}(S)$ . On the other hand, if  $\alpha \in \mathcal{A}$  and  $\inf |\alpha| > 1$ , then  $\alpha$  is invertible and  $\sup |\alpha^{-1}| = \|\alpha^{-1}\| < 1$ . It follows that  $\|\alpha^{-1}S\| \leq \|\alpha^{-1}\| \|S\| < 1$ , so  $\alpha I - S = \alpha(I - \alpha^{-1}S)$  is invertible in  $B^a(H_{\mathcal{A}})$ .

Next, suppose that  $(\alpha I - S)(x) = 0$  for some  $\alpha \in \mathcal{A}$  and  $x \in H_{\mathcal{A}}$ . This gives the following system of equations coordinatewise:  $\alpha x_1 = 0, \alpha x_2 - x_1 = 0, \alpha x_3 - x_2 = 0, \dots$ . Since  $\alpha x_1 = 0$ , we deduce that  $x_1|_{\text{supp } \alpha} = 0$ . However, since  $\alpha x_2 - x_1 = 0$ , it follows that  $x_1|_{(\text{supp } \alpha)^c} = 0$  also. Hence  $x_1 = 0$ . But then  $\alpha x_2 = 0$  and  $\alpha x_3 - x_2 = 0$ . Using the same argument we obtain that  $x_2 = 0$ . Proceeding inductively we obtain that  $x_k = 0$  for all  $k$ , so  $x = 0$ . Since  $\alpha \in \mathcal{A}$  was arbitrary chosen, we conclude that  $\sigma_p^{\mathcal{A}}(S) = \emptyset$ .

Case 2: In this case we consider  $\mathcal{A} = L^\infty((0, 1))$ . Let  $\alpha \in \mathcal{A}$  and assume that  $\inf |\alpha| < 1$ . This means that  $\mu(|\alpha|^{-1}((0, 1 - \epsilon)) > 0)$ . Set  $M_\epsilon = |\alpha|^{-1}((0, 1 - \epsilon))$ , there exists on  $\epsilon \in (0, 1 - \inf |\alpha|)$  such that  $\chi_{M_\epsilon} \neq 0$ . Letting  $\chi_{M_\epsilon}$  play the role of the function  $g$  in the previous proof, we deduce by the same arguments that  $\sigma^{\mathcal{A}}(S) = \{\alpha \in \mathcal{A} \mid \inf |\alpha| \leq 1\}$ . Next, assume that  $(\alpha I - S)(x) = 0$  for some  $\alpha \in \mathcal{A}$  and  $x \in H_{\mathcal{A}}$ . As in the previous proof we get the system of equations  $\alpha x_1 = 0, \alpha x_2 - x_1 = 0, \alpha x_3 - x_2 = 0, \dots$ . The first equation gives that  $x_1 = 0$  a.e. on  $|\alpha|^{-1}(0, \infty)$  whereas the second equation gives  $x_1 = 0$  a.e. on  $\alpha^{-1}(\{0\})$ . Hence  $x_1 = 0$ . Proceeding inductively as in the previous proof we get  $x = 0$ , hence  $\sigma_p^{\mathcal{A}}(S)$  is empty also in this case.  $\square$

**Lemma 2.2.** *Let  $T \in B(H)$  and suppose that  $T$  is invertible. Then the equation  $(TI - S)x = y$  has a solution in  $H_{\mathcal{A}}$  for all  $e_k, k \in \mathbb{N}$  if and only if the sequence  $(T^{-1}, T^{-2}, \dots, T^{-k}, \dots)$  is in  $H_{\mathcal{A}}$ .*

*Proof.* For  $k = 1$ , if  $(TI - S)x = e_1$  we must have  $TB_1 = I$  where  $x = (B_1, B_2, \dots)$ . Hence  $B_1 = T^{-1}$ . Next  $TB_2 - B_1 = 0$ , so  $TB_2 = B_1 = T^{-1}$  which gives  $B_2 = T^{-2}$ . Proceeding inductively, we obtain that  $B_k = T^{-k}$  for all  $k$ . So the equation  $(TI - S)x = e_1$  has a solution in  $H_{\mathcal{A}}$  if and only if the sequence  $T^{-1}, T^{-2}, \dots$  belongs to  $H_{\mathcal{A}}$ .

Now, if  $T^{-1}, T^{-2}, \dots \in H_{\mathcal{A}}$  then the sequence  $x^{(k)}$  in  $H_{\mathcal{A}}$  given by

$$x_n^{(k)} = \begin{cases} 0 & \text{if } n \in \{1, \dots, k-1\} \\ T^{-(n-k+1)} & \text{for } n \in \{k, k+1, \dots\} \end{cases}$$

is the solution of the equation  $(TI - S)x = e_k$  for each  $k \in \mathbb{N}$ .  $\square$

For each  $T \in B(H)$  let  $N_T$  denote the set of all right annihilators of  $T$ . For each right invertible  $T$ , let  $B_T$  denote right inverse of  $T$  and  $W$  be the set of all right invertible  $T \in B(H)$  such that there exists a non zero sequence  $\{Y_n\} \subseteq N_T$  with the property that the sequence  $\{Y_1, B_T Y_1 + Y_2, B_T^2 Y_1 + B_T Y_2 + Y_3, \dots\}$  belongs to  $l_2(B(H))$ .

In addition, let  $Z$  be the set of all right invertible  $T$  such that there is no sequence  $\{Y_n\}$  in  $N_T$  with the property that the sequence  $\{B_T + Y_1, B_T^2 + B_T Y_1 + Y_2, B_T^3 + B_T^2 Y_1 + B_T Y_2 + Y_3, \dots\}$  belongs to  $l_2(B(H))$ . Then we have the following proposition.

**Proposition 2.3.** *Let  $\mathcal{A} = B(H)$  and  $S$  be the unilateral shift on  $H_{\mathcal{A}}$ . Then  $\sigma^{\mathcal{A}}(S) = Z \cup W \cup \{T \in B(H) \mid T \text{ is not right invertible}\}$ .*

*Proof.* If  $(TI - S)x = y$  should have a solution for each  $e_k$ , then, by the similar calculations as in the proof of Lemma 2.2 we must have that  $T$  is right invertible and belongs to  $B(H) \setminus Z$ . Now, suppose that  $T$  is right invertible and that  $(TS - I)y = 0$  for some  $y = \{Y_1, Y_2, \dots\} \in H_{\mathcal{A}}$ . We get coordinatewise the system of equations  $TY_1 = 0, TY_2 = Y_1, TY_3 = Y_2, \dots$ . It is easily seen that  $(TS - I)y = 0$  has a non trivial solution if and only if  $T$  belongs to  $W$ . The proposition follows.  $\square$

**Corollary 2.4.** *Let  $\mathcal{A}$  be a commutative unital  $C^*$ -algebra. Then  $\sigma^{\mathcal{A}}(S) = \mathcal{A} \setminus G(\mathcal{A}) \cup \{\alpha \in G(\mathcal{A}) \mid (\alpha^{-1}, \alpha^{-2}, \dots, \alpha^{-k}, \dots) \notin H_{\mathcal{A}}\}$ .*

*Proof.* Since  $\mathcal{A}$  is commutative, then the set of right invertible elements is  $G(\mathcal{A})$ . Hence we can apply the arguments from the proof of Lemma 2.2.  $\square$

**Corollary 2.5.** *Let  $\mathcal{A}$  be a unital  $C^*$ -algebra. If  $1_{\mathcal{A}}$  denotes the unit in  $\mathcal{A}$ , then  $1_{\mathcal{A}} \in \sigma^{\mathcal{A}}(S)$ .*

*Proof.* We have obviously that the sequence  $(1_{\mathcal{A}}, 1_{\mathcal{A}}, 1_{\mathcal{A}}, \dots) = (1_{\mathcal{A}}^{-1}, 1_{\mathcal{A}}^{-2}, 1_{\mathcal{A}}^{-3}, \dots)$  is not an element of  $H_{\mathcal{A}}$ . Then apply the arguments from the proof of Lemma 2.2.  $\square$

**Example 2.6.** We may also consider a weighted shift  $S_w$  on  $H_{\mathcal{A}}$  given by  $S_w(x)_{j+1} = w_j x_j$  where  $w = (w_1, w_2, \dots)$  is a bounded sequence in  $\mathcal{A}$ . In this case, if  $\alpha$  has a common right annihilator as  $w_j$  for some  $j \in \mathbb{N}$ , then the sequence having this right annihilator in its  $j$ -th coordinate and 0 elsewhere belongs to the kernel of  $\alpha I - S_w$ . Hence  $\alpha \in \sigma^{\mathcal{A}}(S_w)$  in this case.

*Remark 2.7.* Notice that Proposition 2.3 can be generalized to arbitrary unital  $C^*$ -algebras.

**Example 2.8.** Let  $\mathcal{A} = L^{\infty}((0, 1))$ . Set  $\tilde{S}(f_1, f_2, \dots) = (f_1 \chi_{(0, \frac{1}{2})}, f_2 \chi_{(0, \frac{1}{2})} + f_1 \chi_{(\frac{1}{2}, 1)}, f_3 \chi_{(0, \frac{1}{2})} + f_2 \chi_{(\frac{1}{2}, 1)}, \dots)$ . Then  $\tilde{S}$  has the matrix  $\begin{bmatrix} 1 & 0 \\ 0 & S \end{bmatrix}$  w.r.t. the decomposition  $(H_{\mathcal{A}} \cdot \chi_{(0, \frac{1}{2})}) \oplus (H_{\mathcal{A}} \cdot \chi_{(\frac{1}{2}, 1)})$ . It follows that  $\sigma^{\mathcal{A}}(\tilde{S}) = \{\alpha \in \mathcal{A} \mid \inf |\alpha| \cdot \chi_{(\frac{1}{2}, 1)} \leq 1\} \cup \{\alpha \in \mathcal{A} \mid \mu(\{t \mid t \in (0, \frac{1}{2}) \text{ and } \alpha(t) = 1\}) > 0\}$ .

**Proposition 2.9.** *Let  $\alpha \in \mathcal{A}$ . We have*

1. *If  $\alpha I - F$  is bounded below, and  $F \in B^a(H_{\mathcal{A}})$  then  $\alpha \in \sigma_{rl}^{\mathcal{A}}(F)$  if and only if  $\alpha^* \in \sigma_p^{\mathcal{A}}(F^*)$ .*
2. *If  $F, D \in B^a(H_{\mathcal{A}})$  and  $D = U^* F U$  for some unitary operator  $U$ , then  $\sigma^{\mathcal{A}}(F) = \sigma^{\mathcal{A}}(D)$ ,  $\sigma_p^{\mathcal{A}}(F) = \sigma_p^{\mathcal{A}}(D)$ ,  $\sigma_{cl}^{\mathcal{A}}(F) = \sigma_{cl}^{\mathcal{A}}(D)$  and  $\sigma_{rl}^{\mathcal{A}}(F) = \sigma_{rl}^{\mathcal{A}}(D)$ .*

*Proof.* 1) Suppose first that  $F - \alpha I$  is bounded below and  $\alpha \in \sigma_{rl}^{\mathcal{A}}(F)$ . Then  $\text{Im}(F - \alpha I)$  is closed and  $\text{Im}(F - \alpha I)^{\perp} \neq \{0\}$ . Since  $\text{Im } \text{Im}(F - \alpha I)^{\perp} = \ker(F^* - \alpha^* I)$ , it follows that  $\alpha^* \in \sigma_p^{\mathcal{A}}(F^*)$ . Conversely, suppose that  $\alpha^* \in \sigma_p^{\mathcal{A}}(F^*)$  and  $F - \alpha I$  is bounded below. Then, again  $\text{Im } \text{Im}(F - \alpha I)$  is closed and moreover  $\text{Im}(F - \alpha I)^{\perp} = \ker(F^* - \alpha^* I) \neq \{0\}$ . It follows that  $\alpha \in \sigma_{rl}^{\mathcal{A}}(F)$ . It is straightforward to prove the statement in 2.  $\square$

**Proposition 2.10.** *Let  $U \in B^a(H_{\mathcal{A}})$  be unitary. Then  $\sigma^{\mathcal{A}}(U) \subseteq \{\alpha \in \mathcal{A} \mid \|\alpha\| \geq 1\}$  and  $\sigma^{\mathcal{A}}(U) \cap G(\mathcal{A}) \subseteq \{\alpha \in G(\mathcal{A}) \mid \|\alpha^{-1}\|, \|\alpha\| \geq 1\}$ .*

*Proof.* We have  $\alpha I - V = (\alpha V^* - I)V$  and  $\|V^*\| = \|V\| = 1$ .  $\square$

Consider again the orthonormal basis  $\{e_k\}_{k \in \mathbb{N}}$  for  $H_{\mathcal{A}}$ . We may enumerate this basis by indexes in  $\mathbb{Z}$ . Then we get orthonormal basis  $\{e_j\}_{j \in \mathbb{Z}}$  for  $H_{\mathcal{A}}$  and we can consider bilateral shift operator  $V$  w.r.t. this basis i.e.  $V e_k = e_{k+1}$  all  $k \in \mathbb{Z}$ , which gives  $V^* e_k = e_{k-1}$  for all  $k \in \mathbb{Z}$ .

**Proposition 2.11.** *Let  $V$  be bilateral shift operator. Then the following holds*

- 1) *If  $\mathcal{A} = C([0, 1])$ , then  $\sigma^{\mathcal{A}}(V) = \{f \in \mathcal{A} \mid |f|([0, 1]) \cap \{1\} \neq \emptyset\}$*
  - 2) *If  $\mathcal{A} = L^{\infty}([0, 1])$ , then  $\sigma^{\mathcal{A}}(V) = \{f \in \mathcal{A} \mid \mu(|f|^{-1}((1 - \epsilon, 1 + \epsilon))) > 0 \ \forall \epsilon > 0\}$ .*
- In both cases  $\sigma_p^{\mathcal{A}}(V) = \emptyset$ .*

*Proof.* Case 1:

In this case we consider  $\mathcal{A} = C([0, 1])$ . Suppose that  $\alpha \in \mathcal{A}$  and  $|\alpha(\tilde{t})| = 1$  for some  $\tilde{t} \in [0, 1]$ . Choose a function  $y \in \mathcal{A}$  such that  $y(\tilde{t}) = 1$ . If  $\alpha I - V$  is surjective, then there exists an  $x \in H_{\mathcal{A}}$  such that  $(\alpha I - V)x = e_1 \cdot y$ . Now  $x(\tilde{t}) \in l_2$  since  $x \in H_{\mathcal{A}}$ . If we let  $\tilde{V}$  denote the ordinary bilateral shift on  $l_2$  we get that  $\alpha(\tilde{t})x(\tilde{t}) - \tilde{V}(x(\tilde{t})) = (1, 0, 0, \dots)$ , as  $y(\tilde{t}) = 1$ . But this is not possible since  $|\alpha(\tilde{t})| = 1$  (for more details, see [2, Propozicija 19]). We conclude that  $\alpha I - V$  can not be surjective, so  $\alpha \in \sigma^{\mathcal{A}}(V)$ . On the other hand, if  $\alpha \in \mathcal{A}$  and  $|\alpha|([0, 1]) \cap \{1\} = \emptyset$ , then either  $|\alpha(t)| \geq C > 1$  or  $|\alpha(t)| \leq K < 1$  for all  $t \in [0, 1]$  and some constants  $C$  and  $K$  (here we muse that  $|\alpha|$  is continuous, hence  $|\alpha|([0, 1])$  must be connected). If  $|\alpha(t)| \geq C > 1$  for all  $t \in [0, 1]$ , then  $\alpha$  is invertible in  $\mathcal{A}$  and  $\|\alpha^{-1}\| \leq \frac{1}{C} < 1$ . Since  $\|V\| = 1$  it follows that  $\alpha \notin \sigma^{\mathcal{A}}(V)$  then. If  $|\alpha(t)| \leq K < 1$   $t \in [0, 1]$ , then  $\|\alpha\| \leq K < 1$ , so again using that  $\alpha I - V = V(\alpha V^* - I)$  it follows that  $\alpha \notin \sigma^{\mathcal{A}}(V)$  then. Hence  $\sigma^{\mathcal{A}}(V) = \{\alpha \in \mathcal{A} \mid |\alpha|([0, 1]) \cap \{1\} \neq \emptyset\}$ . Next, if  $(\alpha I - V)(x) = 0$  for some  $x \in H_{\mathcal{A}}$ , then we must have  $\alpha(t)x(t) - \tilde{V}x(t) = 0$  for all  $t \in [0, 1]$ . This means that  $x(t) = 0$  for all  $t \in [0, 1]$ .  $\square$

Case 2:

Let now  $\mathcal{A} = L^\infty((0, 1))$  and  $\alpha \in \mathcal{A}$  be such that  $\mu(|\alpha|^{-1}((1 - \epsilon, 1 + \epsilon))) > 0$  for all  $\epsilon > 0$ . If  $(\alpha I - V)x = e_0$  for some  $x \in H_{\mathcal{A}}$ , then we must have  $\alpha x_k - x_{k-1} = 0$  for all  $k \neq 0$  and  $\alpha x_0 - x_{-1} = 1$ . For small  $\epsilon > 0$  set  $M_\epsilon = |\alpha|^{-1}((1 - \epsilon, 1 + \epsilon))$ ,  $M_\epsilon^- = |\alpha|^{-1}((1 - \epsilon, 1))$ ,  $M_\epsilon^+ = |\alpha|^{-1}((1, 1 + \epsilon))$ , so  $M_\epsilon = M_\epsilon^- \cup M_\epsilon^+$ ,  $\chi_{M_\epsilon} \neq 0$ . From the equations above we get  $x_k = \alpha^{-(k+1)}x_{-1}$  for all  $k \leq -1$  and  $x_k = \alpha^{-k}x_0$  for all  $k \geq 1$  on any subset of  $(0, 1)$  on which  $\alpha$  is bounded below, thus in particular on  $M_\epsilon$ . It follows that  $x_k \chi_{M_\epsilon^+} = 0$  for all  $k \leq -1$  and  $x_k \chi_{M_\epsilon^-} = 0$  for all  $k \geq 0$ . Setting this into the second equation above, we get  $\alpha x_0 \chi_{M_\epsilon^+} - x_1 \chi_{M_\epsilon^-} = \chi_{M_\epsilon}$  which gives  $x_0 = \alpha^{-1} \chi_{M_\epsilon^+}$  and  $x_1 = -\chi_{M_\epsilon^-}$ . Hence  $x_k = \alpha^{-(k+1)} \chi_{M_\epsilon^+}$  for all  $k \geq 0$  and  $x_k = -\alpha^{-(k+1)} \chi_{M_\epsilon^-}$  for all  $k \leq -1$ . This gives  $|x_k| \geq (1 + \epsilon)^{-(k+1)} \chi_{M_\epsilon^+}$  for all  $k \geq 0$  and  $|x_k| \geq (1 - \epsilon)^{-(k+1)} \chi_{M_\epsilon^-}$  for all  $k \leq -1$ . Since this holds for all  $\epsilon > 0$  and moreover, we have that either  $\chi_{M_\epsilon^-}$  or  $\chi_{M_\epsilon^+}$  is non zero (because  $\chi_{M_\epsilon}$  for all  $\epsilon > 0$ ), we get that the infinite sum  $\sum_{k \in \mathbb{Z}} x_k^* x_k$  diverge in  $\mathcal{A}$ , so  $x$  can not be an element of  $H_{\mathcal{A}}$ . We conclude that  $e_0 \notin \text{Im}(\alpha I - V)$ , so  $\alpha \in \sigma^{\mathcal{A}}(V)$  in this case.

On the other hand, if  $\mu(|\alpha|^{-1}((1 - \epsilon, 1 + \epsilon))) = 0$  for some  $\alpha \in \mathcal{A}$  and some  $\epsilon > 0$ , then  $(0, 1) = N_\epsilon^- \cup N_\epsilon^+$  where  $N_\epsilon^- = |\alpha|^{-1}((0, 1 - \epsilon))$  and  $N_\epsilon^+ = |\alpha|^{-1}((1 + \epsilon, \|\alpha\|))$ . Since the submodules  $H_{\mathcal{A}} \cdot \chi_{N_\epsilon^-}$  and  $H_{\mathcal{A}} \cdot \chi_{N_\epsilon^+}$  clearly reduce the operator  $V$  and the restrictions of  $\alpha I - V$  on both these submodules are invertible, it follows that  $\alpha I - V$  is invertible, so  $\alpha \notin \sigma^{\mathcal{A}}(V)$ .

**Example 2.12.** Let  $\{\alpha_1, \alpha_2, \dots\}$  be a sequence in a unital  $C^*$ -algebra  $\mathcal{A}$  s.t. each  $\alpha_k$  is a unitary element of  $\mathcal{A}$ . Then the operator  $V$  defined by  $V(x_1, x_2, \dots) = (\alpha_1 x_1, \alpha_2 x_2, \dots)$  is a unitary operator on  $H_{\mathcal{A}}$ . We have  $\sigma^{\mathcal{A}}(V) = \{\beta \in \mathcal{A} \mid \beta - \alpha_k \text{ is not right invertible in } \mathcal{A} \text{ or } (\beta - \alpha_k)\gamma_k = 0 \text{ for some } \gamma_k \in \mathcal{A}\}$ . If  $\mathcal{A} = C([0, 1])$  or if  $\mathcal{A} = L^\infty((0, 1))$  and  $J_1, J_2$  are two closed subintervals of  $(0, 1)$  such that  $J_1 \cap J_2 = \emptyset$ , then we may easily find a function  $\beta \in \mathcal{A}$  such that  $\beta = \alpha_1$  on

$J_1$  and  $|\beta(t)| > 1$  for all  $t \in J_2$ . Hence  $\|\beta\| > 1$ , but we also have  $\beta \in \sigma^A(V)$ . Similarly, if  $\mathcal{A} = B(H)$  where  $H$  is a Hilbert space, then we may easily find two closed subspaces  $H_1$  and  $H_2$  such that  $H_1 \perp H_2$  and  $T \in B(H)$  is such that  $T|_{H_1} = \alpha_{1|_{H_1}}$  and  $\|T|_{H_2}\| > 1$ . Hence again  $T \in \sigma^A(V)$  and  $\|T\| > 1$ . So, if  $V$  is a unitary operator on  $H_A$ , we do not have in general that  $\sigma^A(V) \subseteq \{\alpha \in \mathcal{A} \mid \|\alpha\| = 1\}$ .

**Lemma 2.13.** *If  $F$  is a self-adjoint operator on  $H_A$ , then  $\sigma_p^A(F)$  is a self-adjoint subset of  $\mathcal{A}$ , that is  $\alpha \in \sigma_p^A(F)$  if and only if  $\alpha^* \in \sigma_p^A(F)$  in the case when  $\mathcal{A}$  is a commutative  $C^*$ -algebra.*

*Proof.* Since  $F - \alpha I$  and  $F - \alpha^* I = F^* - \alpha^* I$  mutually commute, we can deduce that  $\|(F - \alpha I)x\| = \|(F - \alpha^* I)x\|$  for all  $x \in H_A$ .  $\square$

**Example 2.14.** Let  $\mathcal{A} = C([0, 1])$  or  $\mathcal{A} = L^\infty((0, 1))$ . If  $G$  is the operator on  $H_A$  given by  $G(f_1, f_2, \dots) = (g_1 f_1, g_2 f_2, \dots)$ , where  $\{g_1, g_2, \dots\}$  is a bounded sequence of real valued functions in  $\mathcal{A}$ , then  $G$  is a self-adjoint operator. Suppose that there are two mutually disjoint, closed subintervals  $J_1$  and  $J_2$  of  $(0, 1)$  such that  $g_1|_{J_1} \neq 0$  and  $g_1|_{J_2} = 0$ . Set  $\tilde{g} = ig_1$ . Then, if we choose a function  $f$  in  $\mathcal{A}$  such that  $\text{supp } f \subseteq J_2$ , we get that  $(\tilde{g}I - G)(f, 0, 0, \dots) = 0$ . However  $\tilde{g} \neq \bar{\tilde{g}}$ , so we do not have that  $\sigma_p^A(G)$  is included in the set of self-adjoint elements of  $\mathcal{A}$ .

**Example 2.15.** Let  $\mathcal{A} = B(H)$  where  $H$  is a Hilbert space and let  $\{e_j\}_{j \in \mathbb{N}}$  be an orthonormal basis for  $H$ . If  $P$  denotes the orthogonal projection onto  $\text{Span}\{e_1\}$ , then the operator  $P \cdot I$  is a self-adjoint operator on  $H_A$ . Now, if  $S$  is the unilateral shift operator on  $H$  w.r.t. to the orthonormal basis  $\{e_j\}$ , then  $S - P$  is injective whereas  $S^* - P$  is not injective because  $(S^* - P)(e_1 + e_2) = 0$ . It follows that  $(S - P) \cdot I$  is an injective operator whereas  $(S^* - P) \cdot I = ((S - P) \cdot I)^*$  is **not** an injective operator. Hence, if  $\mathcal{A} = B(H)$  where  $H$  is a Hilbert space, we do not have in general that  $\sigma_p^A(F)$  is a self-adjoint subset of  $\mathcal{A}$  when  $F = F^*$ .

**Lemma 2.16.** *Let  $\mathcal{A}$  be a commutative  $C^*$ -algebra. If  $F$  is a self-adjoint operator on  $H_A$  and  $\alpha \in \mathcal{A} \setminus \sigma_p^A(F)$ , then  $\overline{\text{Im}(F - \alpha I)}^\perp = \{0\}$ . Hence, if  $\alpha \in \mathcal{A} \setminus \sigma_p^A(F)$  and in addition  $F - \alpha I$  is bounded below, then  $\alpha \in \mathcal{A} \setminus \sigma^A(F)$ .*

*Proof.* Suppose now that  $\alpha \in \mathcal{A} \setminus \sigma_p^A(F)$ . If  $y \in \overline{\text{Im}(F - \alpha I)}^\perp$ , then for all  $x \in H_A$  we have  $\langle (F - \alpha I)x, y \rangle = 0$ . This gives  $\langle x, (F - \alpha^* I)y \rangle = 0$  for all  $x \in H_A$ . It follows that  $(F - \alpha^* I)y = 0$  in this case. By the arguments above we obtain the  $(F - \alpha I)y = 0$ . Since  $\alpha \notin \sigma_p^A(F)$  by the choice of  $\alpha$ , we get that  $y = 0$ . Thus  $\overline{\text{Im}(F - \alpha I)}^\perp = \{0\}$ , when  $\alpha \in \mathcal{A} \setminus \sigma_p^A(F)$ .

Suppose next that  $\alpha \in \mathcal{A}$  is such that  $F - \alpha I$  is bounded below. Then  $\alpha \in \mathcal{A} \setminus \sigma_p^A(F)$  so from the previous arguments we deduce that  $\text{Im}(F - \alpha I)^\perp = \{0\}$ . Moreover, since  $\text{Im}(F - \alpha I)$  is then closed and  $F - \alpha I \in B^a(H_A)$ , from [3, Theorem 2.3.3] it follows that  $\text{Im}(F - \alpha I)$  is orthogonally complementable in  $H_A$ . But, since  $\text{Im}(F - \alpha I)^\perp = \{0\}$ , we must have that  $\text{Im}(F - \alpha I) = H_A$ . Hence  $F - \alpha I$  is invertible in  $B^a(H_A)$ , so  $\alpha$  is in  $\mathcal{A} \setminus \sigma^A(F)$ .  $\square$

**Corollary 2.17.** *Let  $\mathcal{A}$  be a unital commutative  $C^*$ -algebra and  $F$  be a self-adjoint operator on  $H_A$ . If  $\alpha \in \mathcal{A}$  and  $\alpha - \alpha^* \in G(\mathcal{A})$ , then  $F - \alpha I$  is invertible. In this case  $\|(F - \alpha I)^{-1}\| \leq 2 \|(\alpha - \alpha^*)^{-1}\|$ .*



*Proof.* If  $\alpha \in \mathcal{A}$  and  $\alpha - \alpha^* \in G(\mathcal{A})$ , we have  $\langle Fx - \alpha Ix, x \rangle - \langle x, Fx - \alpha Ix \rangle = \langle x, x \rangle \alpha - \alpha^* \langle x, x \rangle = (\alpha - \alpha^*) \langle x, x \rangle$  (here we use that  $\mathcal{A}$  is commutative). From the triangle inequality and the Cauchy-Schwartz inequality for the inner product we obtain  $\|(\alpha - \alpha^*) \langle x, x \rangle\| \leq 2\|x\| \|Fx - \alpha Ix\|$ . Since  $(\alpha - \alpha^*)$  is invertible by assumption, we get that  $\|(\alpha - \alpha^*) \langle x, x \rangle\| \geq \frac{1}{\|(\alpha - \alpha^*)^{-1}\|} \|\langle x, x \rangle\|$ . It follows that  $\|(F - \alpha I)(x)\| \geq \frac{1}{2\|(\alpha - \alpha^*)^{-1}\|} \|\langle x, x \rangle\|$  for all  $x \in H_{\mathcal{A}}$ .  $\square$

*Remark 2.18.* Let  $\mathcal{A} = C([0, 1])$  or  $\mathcal{A} = L^\infty((0, 1))$ . As we have seen in Example 2.14 the operator  $\tilde{g}I - G$  is not invertible, whereas  $\tilde{g} - \bar{\tilde{g}} = i g_1 \neq 0$ . Therefore, the requirement that  $a - a^*$  is invertible is indeed necessary in Corollary 2.17.

**Example 2.19.** Let  $\mathcal{A} = M_2(\mathbb{C})$  and  $T_1, T_2 \in \mathcal{A}$  be given as  $T_1 = \begin{bmatrix} 2 & 1 \\ 1 & 0 \end{bmatrix}$ ,  $T_2 = \begin{bmatrix} 0 & i \\ i & i \end{bmatrix}$ . Then  $T_1$  is self-adjoint and  $T_2 - T_2^* = 2i \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}$ , so  $T_2 - T_2^*$  is invertible. Now  $T_1 - T_2 = \begin{bmatrix} 1 & 1-i \\ 1-i & -i \end{bmatrix}$ , so  $\det(T_1 - T_2) = 0$  which gives  $T_1 - T_2$  is not invertible. Hence the operator  $F := T_1 \cdot I$  is a self-adjoint operator on  $H_{\mathcal{A}}$ , but  $F - T_2 \cdot I = (T_1 - T_2) \cdot I$  is not invertible. This shows that the assumption that  $\mathcal{A}$  is commutative in Corollary 2.17 is indeed necessary.

For a self-adjoint operator  $F$  on  $H_{\mathcal{A}}$ , set  $M(F) = \sup\{\|\langle Fx, x \rangle\| \mid \|x\| = 1\}$  and  $m(F) = \inf\{\|\langle Fx, x \rangle\| \mid \|x\| = 1\}$ . We have the following corollary.

**Corollary 2.20.** *If  $\mathcal{A} = C([0, 1])$  and  $F$  is a self-adjoint operator on  $H_{\mathcal{A}}$ , then  $\sigma^{\mathcal{A}}(F) \subseteq \{f \in \mathcal{A} \mid |f|(0, 1) \cap [m, M] \neq \emptyset\}$ . If  $\mathcal{A} = L^\infty((0, 1))$  and  $F$  is a self-adjoint operator on  $H_{\mathcal{A}}$  then  $\sigma^{\mathcal{A}}(F) \subseteq \{f \in \mathcal{A} \mid \mu(|f|^{-1}[m - \epsilon, M + \epsilon]) > 0 \text{ for some } \epsilon = \epsilon(f) > 0\}$ .*

*Proof.* Let  $F$  be a self-adjoint operator on  $H_{\mathcal{A}}$  where  $\mathcal{A} = L^\infty((0, 1))$  and let  $\alpha \in \mathcal{A}$  be such that there exists an  $\epsilon = \epsilon(\alpha)$  with the property that  $\mu(|f|^{-1}((m - \epsilon, M + \epsilon))) = 0$ . Then  $(0, 1) = M_1 \cup M_2$  where  $M_1$  and  $M_2$  are Borel measurable, mutually disjoint subsets of  $(0, 1)$  satisfying  $|\alpha| \chi_{M_1} \geq M + \epsilon$  and  $|\alpha| \chi_{M_2} \leq m - \epsilon$  a.e. Then, for all  $x \in H_{\mathcal{A}}$  we have  $\langle (F - \alpha I)x, x \rangle = \langle (F - \alpha I)x, x \rangle \cdot \chi_{M_1} + \langle (F - \alpha I)x, x \rangle \cdot \chi_{M_2}$

Now, we have

$$\begin{aligned} & \|\langle (F - \alpha I)x, x \rangle\| \geq \\ & \|\langle (F - \alpha I)x, x \rangle \chi_{M_1}\| \geq \|\overline{\alpha} \langle x, x \rangle \chi_{M_1}\| - \|\langle Fx \cdot \chi_1, x \cdot \chi_{M_1} \rangle\| \geq \\ & \geq (M + \epsilon) \|\langle x, x \rangle \chi_{M_1}\| - \|\langle F(x \cdot \chi_{M_1}), x \cdot \chi_{M_1} \rangle\| \geq \\ & \geq (M + \epsilon) \|\langle x, x \rangle \chi_{M_1}\| - M \|\langle x \cdot \chi_{M_1}, x \cdot \chi_{M_1} \rangle\| = \\ & = (M + \epsilon) \|\langle x, x \rangle \chi_{M_1}\| - M \|\chi_{M_1} \langle x, x \rangle \chi_{M_1}\| = \epsilon \|\langle x, x \rangle \chi_{M_1}\| \end{aligned}$$

(where we have used that

$$\|\langle Fy, y \rangle\| = \|y\|^2 \|\langle F(\frac{y}{\|y\|}), \frac{y}{\|y\|} \rangle\| \leq \|y\| \|M\|.$$

Next

$$\|\langle (F - \alpha I)x, x \rangle\| \geq$$

$$\geq \| \langle (F - \alpha I)x, x \rangle_{\chi_{M_2}} \| \geq \| \langle Fx, x \rangle_{\chi_{M_2}} \| - \| \tilde{\alpha} \langle x, x \rangle_{\chi_{M_2}} \| =$$

$$= \| \langle F(x\chi_{M_2}), x\chi_{M_2} \rangle \| - \| \tilde{\alpha} \langle x, x \rangle_{\chi_{M_2}} \| \geq m \| \langle x \cdot \chi_{M_2}, x \cdot \chi_{M_2} \rangle \| - \\ - (m - \epsilon) \| \langle x, x \rangle_{\chi_{M_2}} \| = \epsilon \| \langle x, x \rangle_{\chi_{M_2}} \|.$$

Hence  $\| \langle (F - \alpha I)x, x \rangle \| \geq \epsilon \max\{ \| \langle x, x \rangle_{\chi_{M_2}} \|, \| \langle x, x \rangle_{M_2} \| \} = \epsilon \| \langle x, x \rangle \|$ .

Thus  $\| (F - \alpha I)x \| \| x \| \geq \| \langle (F - \alpha I)x, x \rangle \| \geq \epsilon \| x \|^2$  for a  $x \in H_{\mathcal{A}}$ . It follows that  $F - \alpha I$  is bounded below, hence from the previous result we deduce that  $F - \alpha I$  is invertible in  $B^a(H_{\mathcal{A}})$ .

The proof in the case when  $\mathcal{A} = C([0, 1])$  is similar, but more simple, because if  $\alpha \in \mathcal{A}$  and  $|\alpha|(0, 1) \cap [m, M] = \emptyset$ , then by continuity of  $|\alpha|$  we must either have that  $|\alpha| < m$  or  $|\alpha| > M$  that on the whole interval  $[0, 1]$ . Moreover, there exist then  $\epsilon > 0$  such that  $|\alpha|m - \epsilon$  or  $|\alpha| \geq M + \epsilon$  on the whole  $[0, 1]$  (since  $|\alpha|$  reaches its maximum and minimum on  $[0, 1]$ ). Then we may proceed in the same way as in the proof above.  $\square$

**Lemma 2.21.** *Let  $\mathcal{A}$  be a commutative unital  $C^*$ -algebra and  $F$  be a normal operator on  $H_{\mathcal{A}}$ , that is  $FF^* = F^*F$ . If  $\alpha_1, \alpha_2 \in \sigma_p^{\mathcal{A}}(F)$  and  $\alpha_1 - \alpha_2$  is invertible in  $\mathcal{A}$ , then  $\ker(F - \alpha_1 I) \perp \ker(F - \alpha_2 I)$ .*

*Proof.* Since  $F$  commutes with  $F^*$  and  $\mathcal{A}$  is a commutative unital  $C^*$ -algebra, then  $F - \alpha_2 I$  and  $F^* - \alpha_2^* I$  commute. Hence  $\ker(F - \alpha_2 I) = \ker(F^* - \alpha_2^* I)$ . For  $x_1 \in \ker(F - \alpha_2 I)$  and  $x_2 \in \ker(F - \alpha_2 I) = \ker(F^* - \alpha_2^* I)$  we get  $\langle x_2 - x_1, (\alpha_1 - \alpha_2) \rangle = \langle x_2 - x_1, \alpha_1 - \alpha_2 \rangle = \langle x_2, Fx_1 \rangle - \langle F^*x_2, x_1 \rangle = 0$ . Since  $(\alpha_1 - \alpha_2)$  is invertible by assumption, it follows that  $\langle x_2, x_1 \rangle = 0$ .  $\square$

**Example 2.22.** Let  $\mathcal{A} = C([0, 1])$  or  $\mathcal{A} = L^\infty((0, 1))$ , consider the self-adjoint operator  $G$  from the previous example. For any function  $f$  in  $\mathcal{A}$  with the support contained in  $J_2$ , we have  $(f, 0, 0, \dots) \in \ker G \cap \ker(\tilde{g}I - G)$ . However  $\tilde{g} = ig_1 \neq 0$  and  $f \neq 0$ , but  $\tilde{g}$  is not invertible in  $\mathcal{A}$ .

**Example 2.23.** Let  $\mathcal{A} = B(H)$  and  $T \in \mathcal{A}$  be a normal and invertible operator. If  $H_1$  and  $H_2$  are two closed subspaces of  $H$  such that  $H = H_1 \oplus H_2$  and  $H_1 \neq H_2^\perp$  (that is  $H_1$  and  $H_2$  are not mutually orthogonal), then  $T\Pi$  and  $T(1 - \Pi)$  are elements of  $\sigma_p^{\mathcal{A}}(T \cdot I)$ , where  $\Pi$  stands for the skew projection onto  $H_1$  along  $H_2$ . Moreover, the operator  $T \cdot I$  is normal operator on  $H_{\mathcal{A}}$  and  $T\Pi - T(1 - \Pi)$  is invertible in  $\mathcal{A}$ . However, if  $P_1$  and  $P_2$  denote the orthogonal projections onto  $H_1$  and  $H_2$ , respectively, then  $e_j \cdot P_1 \in \ker(T\Pi \cdot I - T \cdot I)$ ,  $e_j \cdot P_2 \in \ker(T(I - \Pi) \cdot I - T \cdot I)$  for all  $j$  and  $P_1 P_2 \neq 0$ . So the assumption that  $\mathcal{A}$  is commutative is indeed necessary in Lemma 2.21.

**Lemma 2.24.** *Let  $\mathcal{A}$  be a commutative  $C^*$ -algebra and  $F$  be a normal operator on  $H_{\mathcal{A}}$ . Then  $\sigma_{rl}^{\mathcal{A}}(F) = \emptyset$ , hence  $\sigma^{\mathcal{A}}(F) = \sigma_p^{\mathcal{A}}(F) \cup \sigma_d^{\mathcal{A}}(F)$ .*

*Proof.* Suppose that  $\alpha \in \sigma_{rl}^{\mathcal{A}}(F)$ . Then  $F - \alpha I$  is bounded below. Again, since  $F - \alpha I$  and  $F^* - \alpha^* I$  commute (because  $\mathcal{A}$  is commutative, so  $\alpha I$  commutes with  $\alpha^* I$ ), we get that  $\ker(F - \alpha I) = \ker(F^* - \alpha^* I) = \{0\}$ . Next, since  $\text{Im}(F - \alpha I)$  is closed,



by [3, Theorem 2.3.3], we have  $H_{\mathcal{A}} = \ker(F^* - \alpha^*I) \oplus \text{Im}(F - \alpha I) = \text{Im}(F - \alpha I)$ . So  $F - \alpha I$  is surjective, thus invertible, which gives that  $\sigma_{rl}^{\mathcal{A}}(F) = \emptyset$ .  $\square$

**Example 2.25.** Let  $\mathcal{A} = B(H)$  and  $S, P$  be as in Example 2.15. Then  $P \cdot I$  is a normal operator on  $H_{\mathcal{A}}$  being self-adjoint and  $(S - P) \cdot I$  is bounded below on  $H_{\mathcal{A}}$ . Indeed,  $\|(S - P)x\| \geq \|x\|$  for all  $x \in H$ , hence  $m(S - P) \geq 1$  so  $T^*(S - P)^*(S - P)T \geq m(S - P)^2 T^*T$  for all  $T \in B(H)$  which gives that  $(S - P) \cdot I$  is bounded below on  $H_{\mathcal{A}}$ . However,  $\text{Im}((S - P) \cdot I)^{\perp} = \ker((S^* - P) \cdot I)$ , and  $\ker((S^* - P) \cdot I) \neq \{0\}$  as we have seen in Example 2.15. Hence  $P \cdot I$  is a normal operator on  $H_{\mathcal{A}}$  and  $S \in \sigma_{rl}^{\mathcal{A}}(P \cdot I)$  which shows that the assumption that  $\mathcal{A}$  is commutative is indeed necessary in Lemma 2.24.

Next, for  $F \in B^a(H_{\mathcal{A}})$ , set  $\sigma_a^{\mathcal{A}}(F) = \{\alpha \in \mathcal{A} \mid F - \alpha I \text{ is not bounded below}\}$ . The next two propositions can be proved in exactly the same way same way as for operators on Hilbert spaces.

**Proposition 2.26.** For  $F \in B^a(H_{\mathcal{A}})$ , we have that  $\sigma_a^{\mathcal{A}}(F)$  is a closed subset of  $\mathcal{A}$  in the norm topology and  $\sigma^{\mathcal{A}}(F) = \sigma_a^{\mathcal{A}}(F) \cup \sigma_{rl}^{\mathcal{A}}(F)$ .

**Proposition 2.27.** If  $F \in B^a(H_{\mathcal{A}})$ , then  $\delta\sigma^{\mathcal{A}}(F) \subseteq \sigma_a^{\mathcal{A}}(F)$ . Moreover, if  $M$  is a closed submodule of  $H_{\mathcal{A}}$  and invariant with respect to  $F$ , and  $F_0 = F|_M$ , then we have  $\partial\sigma^{\mathcal{A}}(F_0) \subseteq \sigma_a^{\mathcal{A}}(F)$ ,  $\sigma^{\mathcal{A}}(F_0) \cap \sigma^{\mathcal{A}}(F) = \sigma_{rl}^{\mathcal{A}}(F_0)$ .

**Example 2.28.** We may also consider the operators on  $H_{\mathcal{A}}$  when  $\mathcal{A}$  is a unital  $C^*$ -algebra defined as  $W'(e_k) = e_{2k}$  and  $W''(e_k) = e_{2k-1} \forall k \in \mathbb{N}$ . Also for these operators we have  $\sigma^{\mathcal{A}}(W') = \sigma^{\mathcal{A}}(W'') = \{\alpha \in \mathcal{A} \mid \inf |\alpha| \leq 1\}$  in the case when  $\mathcal{A} = C([0, 1])$  or when  $\mathcal{A} = L^{\infty}((0, 1))$ . Suppose now that  $\mathcal{A} = L^{\infty}((0, 1))$  and consider the operator  $F$  on  $H_{\mathcal{A}}$  given by  $F(f_1, f_2, f_3, \dots) =$

$$(\chi_{(0, \frac{1}{2})}f_1, \chi_{(\frac{1}{2}, 1)}f_1, \chi_{(0, \frac{1}{2})}f_2, \chi_{(\frac{1}{2}, 1)}f_2, \dots). \text{ It follows that } F \text{ has the matrix } \begin{bmatrix} W'' & 0 \\ 0 & W' \end{bmatrix}$$

w.r.t the decomposition  $(H_{\mathcal{A}} \cdot \chi_{(0, \frac{1}{2})}) \oplus (H_{\mathcal{A}} \cdot \chi_{(\frac{1}{2}, 1)})$ . Therefore  $\sigma^{\mathcal{A}}(F) = \{\alpha \in \mathcal{A} \mid \inf |\alpha| \leq 1\}$ . Next  $\sigma_p^{\mathcal{A}}(W') = \emptyset$ ,  $\sigma_p^{\mathcal{A}}(W'') = \{\alpha \in \mathcal{A} \mid \alpha = 1 \text{ on some closed subinterval } J \subseteq [0, 1]\}$  in the case when  $\mathcal{A} = C([0, 1])$  and  $\sigma_p^{\mathcal{A}}(W'') = \{\alpha \in \mathcal{A} \mid \mu(\{t \in (0, 1) \mid \alpha(t) = 1\}) > 0\}$  in the case  $\mathcal{A} = L^{\infty}((0, 1))$ . Hence  $\sigma_p^{\mathcal{A}}(F) = \{\alpha \in \mathcal{A} \mid \mu(\{t \in (0, \frac{1}{2}) \mid \alpha(t) = 1\}) > 0\}$ .

Consider now the operators

$$Z(e_j) = \begin{cases} e_k & \text{when } j = 2k \\ 0 & \text{else} \end{cases}, k \in \mathbb{N}$$

$$Z'(e_j) = \begin{cases} e_k & \text{when } j = 2k - 1 \\ 0 & \text{else} \end{cases}, k \in \mathbb{N}$$

Then  $\sigma^{\mathcal{A}}(Z) = \sigma^{\mathcal{A}}(Z') = \{\alpha \in \mathcal{A} \mid \inf |\alpha| \leq 1\}$ . This follows since  $Z = W'^*$  and  $Z' = W''^*$ . Moreover, we have  $\sigma_p^{\mathcal{A}}(Z) = \{\alpha \in \mathcal{A} \mid \inf |\alpha| < 1\}$  in both cases. In the case  $\mathcal{A} = L^{\infty}((0, 1))$  we have  $\sigma_p^{\mathcal{A}}(Z') = \{\alpha \in \mathcal{A} \mid \inf |\alpha| < 1 \text{ or } \mu(\{t \in (0, 1) \mid \alpha(t) = 1\}) > 0\}$  and in the case when  $\mathcal{A} = C([0, 1])$  we have  $\sigma_p^{\mathcal{A}}(Z') = \{\alpha \in \mathcal{A} \mid \inf |\alpha| < 1 \text{ or } \alpha = 1 \text{ on some closed subinterval } J \subseteq [0, 1]\}$ . Let operator  $D$  on  $H_{\mathcal{A}}$  be given by  $D(g_1, g_2, g_3, \dots) = (g_1\chi_{(0, \frac{1}{2})} + g_2\chi_{(\frac{1}{2}, 1)}, g_3\chi_{(0, \frac{1}{2})} + g_4\chi_{(\frac{1}{2}, 1)}, \dots)$

when  $\mathcal{A} = L^\infty((0, 1))$ . Then  $D = F^*$  and  $D$  has the matrix  $\begin{bmatrix} Z' & 0 \\ 0 & Z \end{bmatrix}$  w.r.t. the decomposition  $H_{\mathcal{A}\chi_{(0, \frac{1}{2})}} \oplus H_{\mathcal{A}\chi_{(\frac{1}{2}, 1)}}$ . It follows that  $\sigma^{\mathcal{A}}(D) = \{\alpha \in \mathcal{A} \mid \inf |\alpha| \leq 1\}$  and  $\sigma_p^{\mathcal{A}}(D) = \{\alpha \in \mathcal{A} \mid \inf |\alpha| < 1 \text{ or } \mu(\{t \in (0, \frac{1}{2}) \mid \alpha(t) = 1\}) > 0\}$ .

Now we are going to prove the statements in this example. Again, we consider two cases.

Case 1:  $\mathcal{A} = L^\infty(0, 1)$ . If  $\inf |\alpha| < 1$  for  $\alpha \in \mathcal{A}$ , then there exist an  $\epsilon > 0$  such that  $\mu(|\alpha|^{-1}((0, 1 - \epsilon))) > 0$ . Set  $M_\epsilon = |\alpha|^{-1}((0, 1 - \epsilon))$  and put  $x_\alpha = \sum_{k=0}^\infty e_{2k} \cdot \chi_{M_\epsilon} \bar{\alpha}^k$ . Then it is easily seen that  $x_\alpha$  indeed is an element of  $H_{\mathcal{A}}$  and moreover  $\langle (\alpha I - W')e_k, x_\alpha \rangle = 0$  for all  $k$ , so  $x_\alpha \in \text{Im}(\alpha I - W')^\perp$ . Therefore,  $\alpha \in \sigma^{\mathcal{A}}(W')$ . On the other hand, we have  $\|W'\| = 1$ , hence if  $\inf |\alpha| > 1$ , it follows by the same arguments as before that  $\alpha I - W' = \alpha(I - \alpha^{-1})W'$  is invertible in  $B^a(H_{\mathcal{A}})$ . Hence, using that  $\sigma^{\mathcal{A}}(W')$  is closed in the norm topology of  $\mathcal{A}$ , we obtain that  $\sigma^{\mathcal{A}}(W') = \{\alpha \in \mathcal{A} \mid \inf |\alpha| \leq 1\}$ . Next, if  $(\alpha I - W')(x) = 0$ , then we must have  $\alpha x_{2k} - x_k = 0$  for all  $k \in \mathbb{N}$  which gives  $x_k = 0$  a.e. on  $\{t \in (0, 1) \mid \alpha(t) = 0\}$  for all  $k \in \mathbb{N}$ . Moreover,  $\alpha x_m = 0$  for all odd  $m$ . All this together implies that  $x_m = 0$  for all odd  $m$ . Now, if  $j = 2^k m$  where  $m$  is odd, we get  $x_m = \alpha^k x_j$ . Since  $x_m = 0$ , it follows that  $x_j = 0$  a.e. on  $\{t \in (0, 1) \mid \alpha(t) \neq 0\}$ . But since  $x_n = 0$  a.e. on  $\{t \in (0, 1) \mid \alpha(t) = 0\}$  for all  $n \in \mathbb{N}$ , we must have  $x_j = 0$ . In particular  $x_2 = 0$ . Then, from the arguments above it also follows that  $x_{2k} = x_{2k-1,2} = 0$  for all  $k$ . Thus  $x = 0$ , so  $\sigma^{\mathcal{A}}(W') = \emptyset$ . The proof for the case when  $\mathcal{A} = C([0, 1])$  is similar.

Next, we consider the operator  $W''$ . If  $\inf |\alpha| < 1$  for some  $\alpha \in \mathcal{A} = L^\infty((0, 1))$ , we let  $M_\epsilon \subseteq (0, 1)$  be the same set as before. Put  $y_\alpha = \sum e_{r_k} \chi_{M_\epsilon} \bar{\alpha}^{(k-1)}$ , where  $\{r_k\}$  is the strictly increasing sequence of positive integers recursively defined by  $r_{k+1} = 2r_k - 1$  for  $k \in \mathbb{N}$  and  $r_1 = 2$ . Then we have  $\langle (\alpha I - W'')e_k, y_\alpha \rangle = 0$  for all  $k \in \mathbb{N}$ , so  $y_\alpha \in \text{Im}(\alpha I - W'')^\perp$ . By the same arguments as before using that  $\|W''\| = 1$ , we obtain that  $\sigma^{\mathcal{A}}(W'') = \{\alpha \in \mathcal{A} \mid \inf |\alpha| \leq 1\}$ . Again, the proof for the case when  $\mathcal{A} = C([0, 1])$  is similar. Suppose next that  $(\alpha I - W'')x = 0$  for some  $x \in H_{\mathcal{A}}$ . Then for all  $k \in \mathbb{N}$  we must have  $\alpha x_{2k-1} - x_k = 0$  which gives  $x_k = 0$  a.e. on  $\{t \in (0, 1) \mid \alpha(t) = 0\}$  for all  $k \in \mathbb{N}$ . Moreover  $\alpha x_{2k} = 0$  for all  $k$  and all this together implies that  $x_{2k} = 0$  for all  $k$ . If  $\alpha \neq 1$  a.e. on  $(0, 1)$ , then the equation  $\alpha x_1 - x_1 = 0$  gives  $x_1 = 0$ . Assume now that  $x_{2j-1} = 0$  for  $j \in \{1, \dots, m\}$ . We have  $\alpha x_{2m+1} - x_{m+1} = 0$ . If  $m+1$  is even, then we already have that  $x_{m+1} = 0$ . If  $m+1$  is odd, then by induction hypothesis we must have  $x_{m+1} = 0$  since  $m+1 \leq 2m-1$  in this case (we assume that  $m \in \mathbb{N}$ , so if  $m+1$  is odd, then  $m \geq 2$ ). Hence  $\alpha x_{2m+1} = 0$ . Combining this together with the fact that  $x_{2m+1} = 0$  a.e. on  $\{t \in (0, 1) \mid \alpha(t) = 0\}$ , we deduce that  $x_{2m+1} = 0$ . By induction we obtain that  $x_{2k-1} = 0$  for all  $k \in \mathbb{N}$ . Hence  $x = 0$ . So, if  $\alpha \in \mathcal{A}$  is such that  $\alpha \neq 1$  a.e. then  $\alpha \notin \sigma_p^{\mathcal{A}}(W'')$ . On the other hand, if  $\alpha \in \mathcal{A}$  is such that  $\mu(\{t \in (0, 1) \mid \alpha(t) = 1\}) > 0$ , then if we let  $M = \alpha^{-1}(\{1\})$ , we have  $\chi_M \neq 0$ . Hence  $(\alpha I - W'')(\chi_M, 0, 0, 0, \dots) = 0$  so  $\alpha \in \sigma_p^{\mathcal{A}}(W'')$  in this case. The proof for the case when  $\mathcal{A} = C([0, 1])$  is similar.

Indeed, the equation  $\alpha x_1 - x_1$  gives that  $x_1 = 0$  on  $\text{supp } (\alpha - 1)$ . If  $\text{supp } (\alpha - 1) \neq [0, 1]$ , then there exists an open interval  $J_0 \subseteq \text{supp } (\alpha - 1)^c$ , since  $\text{supp } (\alpha - 1)^c$  is open. On  $J_0$  we have that  $\alpha - 1 = 0$ , hence by continuity of  $\alpha - 1$ , we must have that  $\alpha - 1 = 0$  on  $\overline{J_0}$ , which is a closed interval. Hence, if there is no closed subinterval of  $[0, 1]$  on which  $\alpha = 1$ , we must have that  $\text{supp } (\alpha - 1) = [0, 1]$ , which gives  $x_1 = 0$ . Then we can proceed in the same way as above. On the other hand, if  $\alpha = 1$  on some subinterval  $[t_1, t_2] \subseteq [0, 1]$ , then we may choose an  $x \in C([0, 1])$  such that  $\text{supp } x \subseteq [t_1, t_2]$ . Then,  $e_1 \cdot x \in \ker(\alpha I - W'')$ .

Now we consider the operators  $Z$  and  $Z'$ . Suppose that  $\alpha \in \mathcal{A} = L^\infty((0, 1))$  and  $\inf |\alpha| < 1$ .

Let  $\epsilon > 0$  be s.t.  $\chi_{M_\epsilon} \neq 0$ , where  $M_\epsilon = |\alpha|^{-1}((0, 1 - \epsilon))$ . Set  $x_\alpha = \sum_{k=1}^{\infty} e_{2k} \cdot \chi_{M_\epsilon} \alpha^{k-1}$ , then  $x_\alpha \in H_{\mathcal{A}}$ ,  $x_\alpha \neq 0$  and  $(\alpha I - Z)x_\alpha = 0$ . Hence  $\alpha \in \sigma_p^{\mathcal{A}}(Z)$ . Let now  $x \in H_{\mathcal{A}}$  be such that  $\inf |\alpha| = 1$ . Then, if  $(\alpha I - Z)x = 0$  for some  $x \in H_{\mathcal{A}}$ , we must have  $\alpha x_k - x_{2k} = 0$  for all  $k \in \mathbb{N}$ . If  $x_j \neq 0$  for some  $j \in \mathbb{N}$ , then  $x_{2^k j}$  for all  $k \in \mathbb{N}$ . Hence  $|x_{2^k j}| \geq |x_j|$  for all  $k \in \mathbb{N}$  (since  $\inf |\alpha| = 1$ ). This is impossible since  $x_j \neq 0$ , so we deduce that  $x_j = 0$  for all  $j \in \mathbb{N}$ , hence  $x = 0$ . Therefore  $\sigma_p^{\mathcal{A}}(Z) = \{\alpha \in \mathcal{A} \mid \inf |\alpha| < 1\}$ . Next, we consider the operator  $Z'$ . Suppose that  $\alpha \in \mathcal{A}$  and  $\inf \|\alpha\| < 1$ . Let  $M_\epsilon$  and  $\chi_{M_\epsilon}$  be as before. Set  $x_\alpha = \sum_{k=1}^{\infty} e_{r_k} \cdot \chi_{M_\epsilon} \alpha^{k-1}$ , where  $\{r_k\}$  is the strictly increasing sequence of positive integers recursively defined by  $r_1 = 2, r_{k+1} = 2r_k - 1$ . Then  $(\alpha I - Z')x_\alpha = 0$ . Hence  $\alpha \in \sigma_p^{\mathcal{A}}(Z')$ . Assume now that  $\alpha \in \mathcal{A}$  and  $\mu(\{t \in (0, 1) \mid \alpha(t) = 1\}) > 0$ . Set  $M = \alpha^{-1}(\{1\})$ , then  $\chi_M \neq 0$  and  $(\alpha I - Z')(\chi_M, 0, 0, 0, \dots) = 0$ , so  $\alpha \in \sigma_p^{\mathcal{A}}(Z')$  in this case. On the other hand, if  $\alpha \in \mathcal{A}$  is such that  $\alpha \neq 1$  a.e. on  $(0, 1)$  then  $x_1 = 0$  for all  $x \in H_{\mathcal{A}}$  satisfying  $(\alpha I - Z')x = 0$ . This is because  $\alpha x_1 - x_1 = 0$  then. Assume in addition that  $\inf |\alpha| = 1$ . If  $(\alpha I - Z')x = 0$  for some  $x \in H_{\mathcal{A}}$  and  $x_j \neq 0$  for some  $j \geq 2$  then we must have  $x_{\tilde{r}_k} = \alpha^{k-1} x_j$  where  $\{\tilde{r}_k\}$  is the strictly increasing sequence of positive integers recursively defined by  $\tilde{r}_1 = j, \tilde{r}_{k+1} = 2\tilde{r}_k - 1$ . This gives  $|x_{\tilde{r}_k}| \geq |x_j|$  for all  $k$ , which is impossible since  $x \in H_{\mathcal{A}}$ . Hence  $x = 0$ , so we deduce that  $\alpha \notin \sigma_p^{\mathcal{A}}(Z')$  if  $\inf |\alpha| \geq 1$  and  $\alpha \neq 1$  a.e. on  $(0, 1)$ . The proof for the case when  $\mathcal{A} = C([0, 1])$  is similar.

Indeed, if  $\inf |\alpha| < 1$ , then we can find a closed subinterval  $[t_1, t_2] \subseteq [0, 1]$  such that  $|\alpha(t)| \leq 1 - \epsilon$  for all  $t \in [t_1, t_2]$ , where  $\epsilon \in (0, 1 - \inf |\alpha|)$ . Let  $g \in \mathcal{A}$  be such that  $\text{supp } g \subseteq [t_1, t_2]$ . Letting  $g$  play the role of  $\chi_{M_\epsilon}$  in the arguments above, we can show that  $\{\alpha \in \mathcal{A} \mid \inf |\alpha| < 1\} \subseteq \sigma_p^{\mathcal{A}}(Z')$ .

Next, the equation  $\alpha x_1 - x_1 = 0$  has a nontrivial solution in  $\mathcal{A} = C([0, 1])$  if and only if  $\text{supp } (\alpha - 1) \neq [0, 1]$  which holds if and only if  $\alpha = 1$  on some closed subinterval  $J \subseteq [0, 1]$ . In this case we choose an  $h \in \mathcal{A}$  such that  $\text{supp } h \subseteq J$ , then  $(h, 0, 0, 0, \dots) \in \ker(\alpha I - Z')$ .

#### Acknowledgement

First of all I am very grateful to Professor Camillo Trapani and Professor Dragan Djordjevic for suggesting me to consider generalized spectra in  $C^*$ -algebras of operators on Hilbert  $C^*$ -modules.

Also, I am very grateful to Professor Ajit Iqbal Singh for inspiring comments, remarks and advises that led to the improved version of the paper.

## REFERENCES

- [1] [IS4] S. Ivković, *On compressions and generalized spectra of operators over  $C^*$ -algebras*, *Annals of Functional Analysis*, <http://link.springer.com/article/10.1007/s43034-019-00034-z>
- [2] [KU] S. Kurepa, *Funkcionalna analiza, Elementi teorije operatora*, Skolska knjiga, Zagreb, 1981
- [3] [MT] V. M. Manuilov, E. V. Troitsky, *Hilbert  $C^*$ -modules*, In: *Translations of Mathematical Monographs*. 226, American Mathematical Society, Providence, RI, 2005.

Stefan Ivković

The Mathematical Institute of the Serbian Academy of Sciences and Arts,

p.p. 367, Kneza Mihaila 36, 11000 Beograd, Serbia,

Tel.: +381-69-774237

email: stefan.iv10@outlook.com