

STEINER SYSTEMS AND CONFIGURATIONS OF POINTS

EDOARDO BALLICO, GIUSEPPE FAVACCHIO, ELENA GUARDO,
AND LORENZO MILAZZO†

ABSTRACT. The aim of this paper is to make a connection between design theory and algebraic geometry/commutative algebra. In particular, given any Steiner System $S(t, n, v)$ we associate two ideals, in a suitable polynomial ring, defining a Steiner configuration of points and its Complement. We focus on the latter, studying its homological invariants, such as Hilbert Function and Betti numbers. We also study symbolic and regular powers associated to the ideal defining a Complement of a Steiner configuration of points, finding its Waldschmidt constant, regularity, bounds on its resurgence and asymptotic resurgence. We also compute the parameters of linear codes associated to any Steiner configuration of points and its Complement.

1. INTRODUCTION

Combinatorial design theory is the study of arranging elements of a finite set into patterns (subsets, words, arrays) according to specified rules. It is a field of combinatorics connected to several other areas of mathematics including number theory and finite geometries. In the last years, the main techniques used in combinatorial and algebraic geometry allow design theory to grow up involving applications in other areas such as in coding theory, cryptography, and computer science.

A $t - (v, n, \lambda)$ -design $D = (V, B)$ is a pair consisting of a set V of v points and a collection B of n -subsets of V , called blocks, such that every t -subset (or t -tuple) of V is contained in exactly λ blocks in B .

The numbers $v = |V|$, $b = |B|$, n , λ , and t are called the *parameters* of the design.

A Steiner system (V, B) of type $S(t, n, v)$ is a $t - (v, n, 1)$ design, that is, a collection B of n -subsets (blocks) of a v -set V such that each t -tuple of V is contained in a unique block in B . The elements in V are called vertices or points and those of B are called blocks. In particular, a Steiner triple system of order v , $STS(v)$, is a collection of triples (3-subsets) of V , such that each unordered pair of elements is contained in precisely one block, and a Steiner quadruple system of order v , $SQS(v)$, is a collection of quadruples (4-subsets) of V such that each triple is found in precisely one block.

A geometric study of some particular classes of Steiner systems, with an eye on the automorphism groups, can be found in [61]. There exists a very extensive literature on STSs and SQS, e.g. [6, 7, 12, 13, 14, 29, 30, 31, 32, 40, 44, 45, 46, 47, 48, 49, 51] to cite some of them.

2010 *Mathematics Subject Classification.* 51E10, 13F55, 13F20, 14G50, 94B27.

Key words and phrases. Steiner systems, monomial ideals, symbolic powers, Stanley Reisner rings, linear codes.

Last updated: May 22, 2020.

† Deceased, March 4, 2019.

In this paper we study special configurations of reduced points in \mathbb{P}^n constructed on Steiner Systems, combining Combinatorial Algebraic Geometry and Commutative Algebra. We found many results concerning geometric and algebraic codes and Steiner systems (or designs in general) but we did not find in the literature any result concerning the study of combinatorial structures, as Steiner systems, using their homological invariants. Thus, we consider this paper as a starting point of the study of combinatorial designs, such as Steiner systems, using the homological invariants of their defining ideals. An introduction of coding theory with the use of Algebraic Geometry can be found in [60]. In particular, we can refer to [52] for some techniques used to study algebraic varieties with special combinatorial features. Recently, connections between commutative algebra and coding theory have gained much attention (see [20, 16, 17, 26, 33, 36, 41, 56, 57, 58] for more results in this direction). For a background in commutative algebra with a view toward algebraic geometry we suggest [22].

The paper is structured as follow. Section 2 provides the background on designs, ideals of points, symbolic and regular powers of ideals, containment problem and monomial ideals. We introduce two finite sets of reduced points in \mathbb{P}^n called *Steiner configuration* and *Complement of a Steiner configuration* (see Definitions 2.3 and 2.4). As pointed out in Remark 2.5, a Steiner configuration of points and its Complement constructed from a Steiner system of type $S(t, n, v)$ are subschemes of a so called *star configuration* of $\binom{v}{n}$ points in \mathbb{P}^n .

In Section 3, we will focus on the Complement of a Steiner configuration of points because it is a *proper hyperplanes section* of a monomial ideal that is the Stanley-Reisner ideal of a matroid (see Theorem 3.4). Thus, each m -th symbolic power of such a monomial ideal is arithmetically Cohen-Macaulay and, after a proper hyperplanes section, it agrees with the m -th symbolic power of the ideal defining a Complement of a Steiner configuration (see Proposition 3.6). This connection allows us to study the homological invariants of the ideal I_{X_C} of the Complement of a Steiner configuration. We start with the description of the initial degree I_{X_C} , i.e. we describe the minimum integer d such that $(I_{X_C})_d \neq (0)$ or, equivalently, the least degree d of a minimal generator of I_{X_C} . We also compute its Waldschmidt constant and give results on the containment problem of I_{X_C} , i.e. the problem of determining for which m and d the containment $I_{X_C}^{(m)} \subseteq I_{X_C}^d$ holds. Section 4 is devoted to the computation of the Hilbert Function, the regularity and graded Betti numbers of I_{X_C} . In particular, we give some bounds for the asymptotic resurgence and the resurgence of I_{X_C} .

Unfortunately, the monomial ideal associated to a Steiner configuration of points does not lead to a Cohen-Macaulay algebra and we cannot apply the same methods as for its Complement. From a combinatorial point of view, two Steiner systems having the same parameters could have very different properties. Computational experiments give us examples where such differences effect the homological invariants. We pose Questions 4.15 and 4.16 on the behaviour of Steiner configurations of points to lead us and the interested researcher towards future works.

We end the paper with an application to coding theory. Following [16, 26, 56, 57, 58, 59], we study special linear codes associated to a Steiner configuration of points and its Complement. In particular, their combinatorial structure allows us to compute the Hamming distance of the associated code (Proposition 5.2 and Theorem 5.4).

The computer softwares CoCoA [1] and Macaulay2 [34] were indispensable for all the computations we made.

Acknowledgement This project began in April 2018 when E. Guardo and L. Milazzo met together with the idea of making an interdisciplinary project between their own fields of research, such as Combinatorics and Commutative Algebra/Algebraic Geometry. This is one of the two projects where L. Milazzo gave his contribution with the idea of involving other mathematicians, such as E. Ballico and G. Favacchio. There is another one, still not published and in progress, that involves other people from University of Catania. However, L. Milazzo became quite ill in February 2019, and after many years battle with his illness fought with huge strength and courage, unexpectedly he passed away on March 4, 2019 in Catania. Lorenzo gave ideas to this paper, and turned his attention to this paper every time he could do it. Unfortunately, Lorenzo was not able to see the final version of this paper and we feel that his contributions warrant an authorship. Lorenzos interests and contributions to the topics in this paper should be recognized. Lorenzo is greatly missed to all of us.

The first author was partially supported by MIUR (Italy). G. Favacchio, E. Guardo and L. Milazzo's work has been supported by the Università degli Studi di Catania, "Piano della Ricerca 2016/2018 Linea di intervento 2". All the authors are members of GNSAGA of INdAM (Italy). We thank M. Gionfriddo, B. Harbourne, and S. Milici for their useful conversations on the topic and for their encouragement in finishing the project. We also thank Zs. Tuza for his useful comments in the revised version of the paper.

The authors thank the two anonymous referees for their useful comments that improved the previous version of the paper.

2. PRELIMINARIES AND NOTATION

Steiner systems play an important role in Design Theory. We use [12] and [14] as main references for all the background on this topic.

The existence of a Steiner system strongly depends on the parameters (t, n, v) . For instance if $t = 2$ and $n = 3$ then $v \equiv 1, 3 \pmod{6}$ must hold. There are known necessary conditions for the existence of a Steiner system of type $S(t, n, v)$ that are not in general sufficient. If a Steiner system (V, B) of type $S(t, n, v)$ exists, then $|B| = \frac{\binom{v}{t}}{\binom{n}{t}}$.

Example 2.1. One of the simplest and most known examples of Steiner system is the Fano Plane. It is unique up to isomorphism and it is a Steiner system of type $S(2, 3, 7)$ with block set

$$B := \{\{1, 2, 3\}, \{1, 4, 5\}, \{1, 6, 7\}, \{2, 4, 6\}, \{2, 5, 7\}, \{3, 4, 7\}, \{3, 5, 6\}\}.$$

2.1. Ideals of Steiner configuration of points and its Complement. Let V be a set of v points and $\mathcal{H} := \{H_1, \dots, H_v\}$ be a collection of distinct hyperplanes H_j of \mathbb{P}^n defined by the linear forms ℓ_j for $j = 1, \dots, v$ and $n \leq v$. Assume that any n hyperplanes in \mathcal{H} meet in a point. There is a natural way to associate a point in \mathbb{P}^n to a n -subset of V . Indeed, given a n -subset $\sigma := \{\sigma_1, \dots, \sigma_n\}$ of V , we denote by $P_{\mathcal{H}, \sigma}$ the point intersection of the hyperplanes $H_{\sigma_1}, \dots, H_{\sigma_n}$. Then the ideal $I_{P_{\mathcal{H}, \sigma}} = (\ell_{\sigma_1}, \dots, \ell_{\sigma_n}) \subseteq k[\mathbb{P}^n]$ is the vanishing ideal of the point $P_{\mathcal{H}, \sigma}$.

Definition 2.2. Given a set V of v points and a general collection Y of subsets of V , we define the following set of points in \mathbb{P}^n with respect to \mathcal{H}

$$X_{\mathcal{H},Y} := \cup_{\sigma \in Y} P_{\mathcal{H},\sigma}$$

and its defining ideal

$$I_{X_{\mathcal{H},Y}} := \cap_{\sigma \in Y} I_{P_{\mathcal{H},\sigma}}.$$

In particular, if $n \leq v$ are positive integers and V is a set of v points, we denote by $C_{(n,v)}$ the set containing all the n -subsets of V .

Definition 2.3. Let (V, B) be a Steiner system of type $S(t, n, v)$ with $t < v \leq n$. We associate to B the following set of points in \mathbb{P}^n

$$X_{\mathcal{H},B} := \cup_{\sigma \in B} P_{\mathcal{H},\sigma}$$

and its defining ideal

$$I_{X_{\mathcal{H},B}} := \cap_{\sigma \in B} I_{P_{\mathcal{H},\sigma}}.$$

We call $X_{\mathcal{H},B}$ the Steiner configuration of points associated to the Steiner system (V, B) of type $S(t, n, v)$ with respect to \mathcal{H} (or just X_B if there is no ambiguity).

Definition 2.4. Let (V, B) be a Steiner system of type $S(t, n, v)$ with $t < n \leq v$. We associate to $C_{(n,v)} \setminus B$ the following set of points in \mathbb{P}^n

$$X_{\mathcal{H},C_{(n,v)} \setminus B} := \cup_{\sigma \in C_{(n,v)} \setminus B} P_{\mathcal{H},\sigma}$$

and its defining ideal

$$I_{X_{\mathcal{H},C_{(n,v)} \setminus B}} := \cap_{\sigma \in C_{(n,v)} \setminus B} I_{P_{\mathcal{H},\sigma}}.$$

We call $X_{\mathcal{H},C_{(n,v)} \setminus B}$ the Complement of a Steiner configuration of points with respect to \mathcal{H} (or C-Steiner X_C if there is no ambiguity).

Remark 2.5. Note that from Definition 2.2, it follows that $X_{\mathcal{H},C_{(n,v)}}$ is a so called *star configuration* of $\binom{v}{n}$ points in \mathbb{P}^n . In particular, from definitions 2.3 and 2.4, $X_{\mathcal{H},C_{(n,v)} \setminus B}$ is the Complement of the Steiner configuration $X_{\mathcal{H},B}$ in the star configuration $X_{\mathcal{H},C_{(n,v)}}$. Thus, a Steiner configuration of points and its Complement are subschemes of a star configuration of $\binom{v}{n}$ points in \mathbb{P}^n . See for instance [8, 10, 9, 27, 28] for recent results on star configurations and see [59] for the connections between linear codes and ideals of star configurations.

Remark 2.6. Since the set $X_{\mathcal{H},B}$ contains $|B|$ points, we have that

$$\deg X_{\mathcal{H},C_{(n,v)} \setminus B} = \binom{v}{n} - |B| = \binom{v}{n} - \frac{\binom{v}{t}}{\binom{n}{t}}.$$

Example 2.7. Consider the Steiner configuration associated to (V, B) of type $S(2, 3, 7)$ as in Example 2.1. Take $\mathcal{H} := \{H_1, \dots, H_7\}$ a collection of 7 distinct hyperplanes H_i in \mathbb{P}^3 defined by a linear form ℓ_i for $i = 1, \dots, 7$, respectively, with the property that any 3 of them meet in a point $P_{\mathcal{H},\sigma} = H_{\sigma_1} \cap H_{\sigma_2} \cap H_{\sigma_3}$, where $\sigma = \{\sigma_1, \sigma_2, \sigma_3\} \in B$. We get that $X_{\mathcal{H},C_{(3,7)}}$ is a *star configuration* of $\binom{7}{3} = 35$ points in \mathbb{P}^3 , $X_{\mathcal{H},B} := \cup_{\sigma \in B} \{P_{\mathcal{H},\sigma}\}$ is a Steiner configuration consisting of 7 points in \mathbb{P}^3 and $X_{\mathcal{H},C_{(3,7)} \setminus B}$ is a C-Steiner configuration consisting of $\binom{7}{3} - 7 = 28$ points in \mathbb{P}^3 . Their defining ideals are respectively,

$$I_{X_{\mathcal{H},B}} := \cap_{\sigma \in B} I_{P_{\mathcal{H},\sigma}} \text{ and } I_{X_{\mathcal{H},C_{(3,7)} \setminus B}} := \cap_{\sigma \in C_{(3,7)} \setminus B} I_{P_{\mathcal{H},\sigma}}.$$

2.2. Symbolic and regular powers of an ideal and the containment problem. In this section, we recall some definitions and known results concerning the symbolic and regular powers of ideal of points.

Let I be a homogeneous ideal in the standard graded polynomial ring $R := k[x_0, \dots, x_n]$. Given an integer m , we denote by I^m the regular power of the ideal I . The m -th *symbolic power* of I is defined as

$$I^{(m)} = \bigcap_{\mathfrak{p} \in \text{Ass}(I)} (I^m R_{\mathfrak{p}} \cap R)$$

where $\text{Ass}(I)$ denotes the set of associated primes of I . If I is a radical ideal (this includes for instance squarefree monomial ideals and ideals of finite sets of points) then

$$I^{(m)} = \bigcap_{\mathfrak{p} \in \text{Ass}(I)} \mathfrak{p}^m.$$

It always holds that $I^m \subseteq I^{(m)}$. In particular, the *containment problem* is of interest, i.e. the problem of determining for which m and d the containment $I^{(m)} \subseteq I^d$ holds. We refer the reader to [2, 3, 5, 18, 19, 39, 54] for a partial list of papers on this topic.

The investigation of this problem led to the introduction of other invariants with the aim of comparison between symbolic and ordinary powers of an ideal.

Let I be a homogeneous ideal, the real number

$$\rho(I) = \sup \left\{ \frac{m}{r} : I^{(m)} \not\subseteq I^r \right\}$$

is called *resurgence* of I (see [3, 5]). For a homogeneous ideal $(0) \subset I \subset k[\mathbb{P}^n]$, in [35] the authors define an asymptotic resurgence as follows:

$$\rho_a(I) = \sup \left\{ \frac{m}{r} : I^{(mt)} \not\subseteq I^{rt} \text{ for all } t \gg 0 \right\}.$$

The resurgence is strictly related to the *Waldschmidt constant* of I , introduced in [64] in a completely different setting. If $\alpha(I)$ is the minimum integer d such that $I_d \neq (0)$, then the Waldschmidt constant of I , $\hat{\alpha}(I)$, is the real number

$$\hat{\alpha}(I) = \lim_{m \rightarrow \infty} \frac{\alpha(I^{(m)})}{m}.$$

Given r distinct points $P_i \in \mathbb{P}^n$ and non-negative integers m_i , we denote by $Z = m_1 P_1 + \dots + m_r P_r \subset \mathbb{P}^n$ the set of fat points defined by $I_Z = \bigcap_i I_{P_i}^{m_i}$, where I_{P_i} is the prime ideal generated by all forms which vanish at P_i . The set $X = \{P_1, \dots, P_r\}$ is called the support of Z . When $m_i = m$ for all i , $I_Z = \bigcap_i I_{P_i}^m = I_X^{(m)}$.

If I is the ideal of a set of (fat) points, we denote by I_i (resp. R_i) the vector space span in I (resp. R) of the forms of degree i in I (resp. R). The function $H : \mathbb{N} \rightarrow \mathbb{N}$ such that $H(i) = \dim_k(R/I)_i = \dim_k R_i - \dim_k I_i$ is called the Hilbert Function of R/I and it is denoted by $H_{R/I}(i)$. The Castelnuovo-Mumford regularity of I , denoted by $\text{reg}(I)$, is the least degree $t > 0$ such that $\dim(R/I)_t = \dim(R/I)_{t-1}$.

Theorem 1.2.1 in [5] shows that if I defines a 0-dimensional scheme, i.e., a finite set of points, then

$$\frac{\alpha(I)}{\hat{\alpha}(I)} \leq \rho(I) \leq \frac{\text{reg}(I)}{\hat{\alpha}(I)}.$$

In [35], Theorem 1.2. shows that (1) $1 \leq \frac{\alpha(I)}{\widehat{\alpha}(I)} \leq \rho_a(I) \leq \rho(I) \leq h$ where $h = \min(N, h_I)$ and h_I is the maximum of the heights of the associated primes of I ; (2) if I is the ideal of a (non-empty) smooth subscheme of \mathbb{P}^n , then $\rho_a(I) \leq \frac{\omega(I)}{\widehat{\alpha}(I)} \leq \frac{\text{reg}(I)}{\widehat{\alpha}(I)}$ where $\omega(I)$ is the largest degree in a minimal homogeneous set of generators of I .

2.3. Monomial ideals, simplicial complexes. In this section, we show how monomial ideals form an important link between commutative algebra and combinatorics. A monomial ideal is uniquely determined by the monomials it contains. Monomial ideals also arise in graph theory. Given a graph G with vertices $\{x_1, \dots, x_v\}$, we associate the ideal I_G in $k[x_1, \dots, x_v]$ generated by the quadratic monomials $x_i x_j$ such that x_i is adjacent to x_j . From known results in the literature, it is possible to determine many invariants for monomial ideals.

In particular, ideals generated by squarefree monomials have a beautiful combinatorial interpretation in terms of simplicial complexes. In the next sections, we will use these properties to completely describe the most important homological invariants of an ideal defining a Complement of a Steiner configuration of points. This is equivalent to say that we are describing special subsets of star configurations using tools from combinatorics and algebraic geometry. Unfortunately, the ideal defining a Steiner configuration of points cannot be described using monomial ideals, and we cannot determine its homological invariants. So, the problem is still open. At the end of the paper, we will be able to describe the parameters of the linear codes associated to both a Steiner configuration of points and its Complement.

We refer to [37] for notation and basic facts on monomial ideals and to [5] for an extensive coverage of the theory of Stanley-Reisner ideals.

Definition 2.8. An ideal I in a polynomial ring R is called a monomial ideal if there is a subset $\mathcal{A} \subset \mathbb{Z}_{\geq 0}^n$ (possibly infinite) such that I consists of all polynomials of the form $\sum_{\alpha \in \mathcal{A}} h_{\alpha} x^{\alpha}$, where $h_{\alpha} \in R$.

Definition 2.9. A simplicial complex Δ over a set of vertices $V = \{x_1, \dots, x_v\}$ is a collection of subsets of V satisfying the following two conditions:

- (1) $\{x_i\} \in \Delta$ for all $1 \leq i \leq v$
- (2) if $F \in \Delta$ and $G \subset F$, then $G \in \Delta$.

An element F of Δ is called a face, and the dimension of a face F of Δ is $|F| - 1$, where $|F|$ is the number of vertices of F . The faces of dimensions 0 and 1 are called vertices and edges, respectively, and $\dim \emptyset = -1$.

The maximal faces of Δ under inclusion are called facets of Δ . The dimension of the simplicial complex Δ is $\dim \Delta = \max\{\dim F \mid F \in \Delta\}$. We refer to i -dimensional faces as i -faces. We denote the simplicial complex Δ with facets F_1, \dots, F_q by $\Delta = \langle F_1, \dots, F_q \rangle$ and we call $\{F_1, \dots, F_q\}$ the facet set of Δ .

If Δ is a d -dimensional simplicial complex, the most important invariant is the f -vector (or *face vector*) of Δ and it is denoted by $(f_0, \dots, f_d) \in \mathbb{N}^{d+1}$, where f_i denotes the number of i -dimensional faces in Δ . From the monomial ideal point of view, the f -vector is encoded in the Hilbert series of the quotient ring R/I_{Δ} and it is related to the h -vector of a suitable set of points in \mathbb{P}^n (see Section 4). We will use Remark 2.11 to show the connections between Stanley-Reisner ideals, simplicial complexes, matroids and Steiner systems to determine Waldschmidt constant and

bounds for the resurgence (see Section 3), Betti numbers, Hilbert function, and regularity of the ideal of a Complement of a Steiner configurations of points (see Section 4).

The Alexander dual of a simplicial complex Δ on $V = \{x_1, \dots, x_v\}$ is the simplicial complex Δ^\vee on V with faces $V \setminus \sigma$, where $\sigma \notin \Delta$.

The Stanley-Reisner ideal of Δ is the ideal $I_\Delta := (x^\sigma \mid \sigma \notin \Delta)$ of $R = k[x_1, \dots, x_v]$, where $x^\sigma = \prod_{i \in \sigma} x_i$. It is well known that the Stanley-Reisner ideals are precisely the squarefree monomial ideals. The quotient ring $k[\Delta] := R/I_\Delta$ is the Stanley-Reisner ring of the simplicial complex Δ .

Each simplicial complex has a geometric realization as a certain subset of a finite dimensional affine space.

If V is a set v points, we denote by $k[V] := k[x_1, \dots, x_v]$ the standard graded polynomial ring in v variables. Given a n -subset of V , $\sigma := \{i_1, i_2, \dots, i_n\} \subseteq V$, we will write

$$\mathfrak{p}_\sigma := (x_{i_1}, x_{i_2}, \dots, x_{i_n}) \subseteq k[V]$$

for the prime ideal generated by the variables indexed by σ , and

$$M_\sigma := x_{i_1} x_{i_2} \cdots x_{i_n} \in k[V]$$

for the monomial given by the product of the variables indexed by σ .

Let $n \leq v$ be positive integers, and V a set of v points; recall that $C_{(n,v)}$ is the set containing all the n -subsets of V .

Definition 2.10. If $T \subset C_{(n,v)}$, we define two ideals

$$I_T := (M_\sigma \mid \sigma \in T) \subseteq k[V]$$

and

$$J_T := \bigcap_{\sigma \in T} \mathfrak{p}_\sigma \subseteq k[V]$$

called the *face ideal* of T and the *cover ideal* of T , respectively.

Remark 2.11. It is well known that J_T is the Stanley-Reisner ideal I_{Δ_T} of the simplicial complex

$$\Delta_T := \langle V \setminus \sigma \mid \sigma \in T \rangle.$$

Then J_T is generated by the monomials M_b with $b \notin \Delta_T$. We also recall that I_T and J_T are the Alexander duals of each other.

Since J_T is a squarefree monomial ideal, the m -th symbolic power of J_T (Theorem 3.7 in [15]) is

$$J_T^{(m)} := \bigcap_{\sigma \in T} \mathfrak{p}_\sigma^m.$$

3. MATROID AND CONFIGURATIONS OF POINTS FROM STEINER SYSTEMS: WALDSCHMIDT CONSTANT AND CONTAINMENT PROBLEM

This section is devoted to show how the results of the previous sections are related to our special configurations of points. In particular, we will show that the Complement of a Steiner configuration of points is connected with the theory of matroids.

Definition 3.1. A simplicial complex Δ is said to be a matroid if $F, G \in \Delta$ and if $|F| > |G|$ then there exists $i \in F \setminus G$ such that $G \cup \{i\} \in \Delta$.

We also recall that we say that a homogeneous ideal J in a polynomial ring R is *Cohen-Macaulay* (CM) if R/J is Cohen-Macaulay, i.e. $\text{depth}(R/J) = \text{Krull-dim}(R/J)$. Varbaro in [62] and Minh and Trung in [50] have independently shown the following

Theorem 3.2 (Varbaro [62], Minh and Trung [50]). *Let Δ be a simplicial complex. Then $k[V]/I_{\Delta}^{(m)}$ is Cohen-Macaulay for each $m \geq 1$ if and only if Δ is a matroid.*

Teraai and Trung in [55] proved if $I_{\Delta}^{(m)}$ is Cohen-Macaulay for some $m \geq 3$ then Δ is a matroid.

To shorten the notation, from now on we set $C := C_{(n,v)} \setminus B$. Let $X_{\mathcal{H},B}$ be a Steiner configuration of points and $X_{\mathcal{H},C}$ its Complement as previous defined with respect to a collection \mathcal{H} of hyperplanes in \mathbb{P}^n .

We claim that $I_{X_{\mathcal{H},C}}^{(m)}$ and $J_C^{(m)}$ share the same homological invariants for any positive integer m . In particular, the key point of our argument is that J_C is the Stanley-Reisner ideal of the simplicial complex Δ_C , and that Δ_C is a matroid (see Theorem 3.4).

Several times in the literature simplicial complexes have been associated to Steiner systems. See for instance [13, 53]. In Example 4.6 in [19] the well known Fano matroid is used to construct a Cohen-Macaulay ideal I such that $I^{(3)} \neq I^3$ and $I^{(2)} \neq I^2$. The quoted example is one of the cases described in Corollary 3.10.

We need the following auxiliary lemma.

Lemma 3.3. *Let (V, B) be a Steiner system of type $S(t, n, v)$. If $C_{(v-n-1,v)}$ is the set containing all the $(v-n+1)$ -subsets of V , then $C_{(v-n-1,v)} \subseteq \Delta_C$.*

Proof. Let F be a $(v-n-1)$ -set of V . In order to prove the lemma we need to find $G \in C$ such that $F \subseteq V \setminus G$, i.e., $G \subseteq V \setminus F$. First note that $|V \setminus F| = n+1$, so we can take two n -subsets G_1, G_2 of $V \setminus F$ sharing a $n-1$ -subset. Since $n-1 \geq t$, from the definition of Steiner Systems at least one of them, say G_1 , does not belong to G . Then $G_1 \in C$ and $G \subseteq V \setminus F$. \square

The next results are useful to describe the Complement of a Steiner configuration of points using the combinatorial properties of matroids.

Theorem 3.4. *Let (V, B) be a Steiner system of type $S(t, n, v)$. Then Δ_C is a matroid.*

Proof. Let F, G be two maximal elements in Δ_C , i.e., F, G belong to the facet set of Δ_C . Assume by contradiction $(F \setminus \{i\}) \cup \{j\} \notin \Delta_C$ for each $i \in F$ and for each $j \in G$. Then all the sets $\sigma_{i,j} := V \setminus ((F \setminus \{i\}) \cup \{j\})$ belong to B . Since $n \geq 2$, at least two blocks of B share a $(n-1)$ -set. That is a contradiction because $n-1 > t-1$ from Definition 2.3. \square

Proposition 3.5. *Let (V, B) be a Steiner system of type $S(t, n, v)$. Then $k[V]/J_C^{(m)}$ is Cohen-Macaulay for each $m \geq 1$.*

Proof. Since $J_C = I_{\Delta_C}$, it is a consequence of Theorem 3.2 and Theorem 3.4. \square

Now, given a Steiner system of type $S(t, n, v)$ we are able to describe some homological invariants of the ideal associated to a Complement of Steiner configuration of points. Unfortunately, we have no similar result as Proposition 3.5 that should hold for a Steiner configuration of points.

We now prove the main claim of this section.

Proposition 3.6. $I_{X_{\mathcal{H},C}}^{(m)} \subseteq k[\mathbb{P}^n]$ and $I_{\Delta_C}^{(m)} \subseteq k[V]$ share the same homological invariants.

Proof. It is an immediate consequence of Theorem 3.6. in [28]. Indeed, $k[V]/I_{\Delta_C}^{(m)}$ is Cohen-Macaulay, and any subset of at most n linear forms in $\{\ell_1, \dots, \ell_v\}$ is a $k[\mathbb{P}^n]$ -regular sequence. \square

The Cohen-Macaulay property of $k[V]/I_{\Delta_C}^{(m)}$ also allows us to look at $I_{X_{\mathcal{H},C}}^{(m)}$ as a proper hyperplane section of $I_{\Delta_C}^{(m)}$ (see [61]). This construction is quite standard but is very useful to describe combinatorial properties for arithmetically Cohen-Macaulay varieties X , i.e, $\text{depth}(R/I_X) = \text{Krull-dim}(R/I_X)$, not only in projective spaces, but also in multiprojective spaces, see for instance [23, 24, 25, 28]. The common idea is to relate, when possible, ideals of arithmetically Cohen-Macaulay varieties to monomial ideals in order to study their invariants.

We start with the description of the initial degree $\alpha(J_C)$ of $J_C = I_{\Delta_C}$, i.e. we describe the minimum integer d such that $(J_C)_d \neq (0)$ or, equivalently, the least degree d of a minimal generator of J_C . For the ease of the reader, we prove the following result that shows that $\{\alpha(J_C^{(m)})\}_m$ is a strictly increasing sequence.

Lemma 3.7. *Let $\mathfrak{p} \subseteq k[V]$ be a squarefree monomial ideal and $M \in \mathfrak{p}^m$. Then $\frac{\partial M}{\partial x_j} \in \mathfrak{p}^{m-1}$ for any $j \in V$.*

Proof. Set $\mathfrak{p} = \mathfrak{p}_\sigma$ where $\sigma \subseteq V$ and let $x_1^{a_1} \cdots x_v^{a_v} \in \mathfrak{p}^m$. Then $\sum_{i \in \sigma} a_i \geq m$. If $a_j = 0$ then trivially $0 \in \mathfrak{p}^{m-1}$. If $a_j > 0$ then $\frac{\partial M}{\partial x_j} = a_j x_1^{a_1} \cdots x_j^{a_j-1} \cdots x_v^{a_v}$ and $\sum_{i \in \sigma} a_i \geq m-1$. \square

Proposition 3.8. *Let (V, B) be a Steiner system of type $S(t, n, v)$. Then*

- i) $\alpha(J_C) = v - n$;
- ii) $\alpha(J_C^{(q)}) = v - n + q$ for $2 \leq q < n$;
- iii) $\alpha(J_C^{(m)}) = \alpha(J_C^{(q)}) + pv$, where $m = pn + q$ and $0 \leq q < n$.

Proof. (i) We have $J_C = I_{\Delta_C} = (M_\sigma \mid \sigma \notin \Delta_C)$. From Lemma 3.3 we have that the elements not in Δ_C have cardinality at least $v - n$. On the other hand, for any $\beta \in B$ we have $V \setminus \beta \notin \Delta_C$.

(ii) First we show $\alpha(J_C^{(q)}) \leq v - n + q$. Let $\beta \in B$ and $\sigma \subseteq \beta$ with $|\sigma| = q$, then we claim $M_{(V \setminus \beta) \cup \sigma} \in \alpha(J_C^{(q)})$. Assume by contradiction $M_{(V \setminus \beta) \cup \sigma} \notin \mathfrak{p}_\tau^q$ for some $\tau \in V$. Then $(V \setminus \beta) \cap \sigma \cap \tau = \emptyset$ but this is a contradiction since $v - n + q + n > n$. On the other hand, if $q > 2$ the statement follows by Lemma 3.7, since $\alpha(J_C^{(q)}) > \alpha(J_C^{(q-1)}) = v - n + q - 1$. If $q = 2$ we proceed by contradiction. Let $M \in J_C^{(2)}$ be a monomial such that $\deg M = v - n + 1$. Then there are at least $n - 1$ variables that do not divide M . Since $t \leq n - 1$, there is at most one block in B containing these variables. Therefore there are at least $v - n$ elements of C containing them. Since $M \in J_C^{(2)}$, M belongs to each of these ideals to the power of 2, i.e., there are $v - n$ variables x_j such that $x_j^2 \mid M$. So $\deg M \geq 2(v - n) > v - n + 1$.

(iii) We show that $J_C^{(m)} : M_V = J_C^{(m-n)}$.

- If $F \in J_C^{(m-n)}$ then $M_VF \in J_C^{(n)} \cdot J_C^{(m-n)} \subseteq J_C^{(m)}$.
- If $FM_V \in J_C^{(m)}$ then, by Lemma 3.7, $FM_{V \setminus \sigma} \in J_C^{(m-n)}$ for any $\sigma \in T$.
Thus $FM_{V \setminus \sigma} \in \mathfrak{p}_\sigma^{(m-n)}$. This implies $F \in \mathfrak{p}_\sigma^{(m-n)}$.

Then $\alpha(J_C^{(m)}) = \alpha(J_C^{(m-n)}) + v = \dots = \alpha(J_C^{(q)}) + pv$, where $m = pn + q$ and $0 \leq q < n$.

□

From Proposition 3.6 and Proposition 3.8, the initial degree of the ideal of a C-Steiner configuration of points only depends on the parameters (t, n, v) of the Steiner system.

Using the previous results, we have the following theorem:

Theorem 3.9. *Let (V, B) be a Steiner system of type $S(t, n, v)$. Then*

- i) $\alpha(I_{X_C}) = v - n$;
- ii) $\alpha(I_{X_C}^{(q)}) = v - n + q$, for $2 \leq q < n$;
- iii) $\alpha(I_{X_C}^{(m)}) = \alpha(I_{X_C}^{(q)}) + pv$, where $m = pn + q$ and $0 \leq q < n$ and $\alpha(I_{X_C}^{(n)}) = \alpha(I_{X_C}^{(0)}) + v = v$.

An important consequence of Theorem 3.9 is related to the containment problem.

Corollary 3.10. *Let (V, B) be a Steiner system of type $S(t, n, v)$. Then $I_{X_C}^{(m)} \not\subseteq I_{X_C}^d$ for any pair (m, d) such that*

$$m \equiv 1 \pmod{n} \quad \text{and} \quad d > 1 + \frac{(m-1)v}{n(v-n)} \quad \text{or}$$

$$m \not\equiv 1 \pmod{n} \quad \text{and} \quad d > 1 + \frac{m-n}{n} + \frac{m}{(v-n)}$$

In particular, if $v > 2n$ then

$$I_{X_C}^{(n)} \not\subseteq I_{X_C}^2$$

Proof. From item i) in Theorem 3.9 $\alpha(I_{X_C}^d) = d(v-n)$. Therefore, it is enough to take m such that $\alpha(I_{X_C}^{(m)}) < d(v-n)$. If $m \equiv 1 \pmod{n}$ then, from items i) and iii) in Theorem 3.9, we get $\alpha(I_{X_C}^{(m)}) = (v-n) + \frac{(m-1)v}{n}$. Then $(v-n) + \frac{(m-1)v}{n} < d(v-n)$ implies the statement. If $m \not\equiv 1 \pmod{n}$ then $m = pn + q$ and $2 \leq q \leq n$ and, from items ii) and iii) in Theorem 3.9, we get $\alpha(I_{X_C}^{(m)}) = v - n + m - pn + pv = (1+p)(v-n) + m$. Thus $(1+p)(v-n) < d(v-n) + m$ implies $d > 1 + p + \frac{m}{(v-n)} \geq 1 + \frac{m-n}{n} + \frac{m}{(v-n)}$.

In the case $m = n$, we get $1 + \frac{m-n}{n} + \frac{m}{(v-n)} = 1 + \frac{n}{(v-n)} < 2$. □

Example 3.11. If $X := X_{\mathcal{H}, C_{(3,7)} \setminus B} \subseteq \mathbb{P}^3$ is the C-Steiner configuration of points as in Example 2.7, then $I_X^{(3)} \not\subseteq I_X^2$.

Another immediate corollary allows us to compute the Waldschmidt constant of a C-Steiner configuration. The Waldschmidt constant of a uniform matroid was computed in Theorem 7.5 in [2]. Properties of a uniform matroid were also studied in [28].

Corollary 3.12. *If (V, B) is a Steiner system of type $S(t, n, v)$, then the Waldschmidt constant of I_{X_C} is*

$$\widehat{\alpha}(I_{X_C}) = \frac{v}{n}.$$

Proof. From [11], Lemma 1, the limit $\widehat{\alpha}(I_{X_C}) = \lim_{m \rightarrow +\infty} \frac{\alpha(I_{X_C}^{(m)})}{m}$ exists. Then from Theorem 3.9 we have

$$\widehat{\alpha}(I_{X_C}) = \lim_{p \rightarrow +\infty} \frac{\alpha(I_{X_C}^{(pn)})}{pn} = \lim_{p \rightarrow +\infty} \frac{pv}{pn} = \frac{v}{n}.$$

□

Remark 3.13. The homological invariants do not depend on the choice of the hyperplanes, provided that we take them meeting properly.

4. HOMOLOGICAL INVARIANTS OF C-STEINER CONFIGURATIONS OF POINTS: HILBERT FUNCTION AND GRADED BETTI NUMBERS

In this section we describe the Hilbert function of a C-Steiner configuration of points. We recall some known definitions. For a finite set of points $X \subset \mathbb{P}^n$ the Hilbert function of X is defined as the numerical function $H_X : \mathbb{N} \rightarrow \mathbb{N}$ such that

$$H_X(i) = \dim_k(R/I_X)_i = \dim_k R_i - \dim_k (I_X)_i$$

where $R = k[\mathbb{P}^n]$ and the first difference of the Hilbert function is defined by $\Delta H_X(i) := H_X(i) - H_X(i-1)$. The h -vector of X is denoted by

$$h_X = h = (1, h_1, \dots, h_p)$$

where $h_i = \Delta H_X(i)$ and p is the last index such that $\Delta H_X(i) > 0$. If I is the ideal of a set of points, analogously to the case of fat points, $\text{reg}(I)$ denotes the Castelnuovo-Mumford regularity of I .

The next result gives us informations on the h -vector of a C-Steiner configuration of points X_C . Recall that given a Steiner system (V, B) of type $S(t, n, v)$, the number of blocks is $|B| = \frac{\binom{v}{t}}{\binom{n}{t}}$ and that the last entry h_p of h_{X_C} is in degree $p = v - n$. With the previous notation, we have the following:

Proposition 4.1. *If (V, B) is a Steiner system $S(t, n, v)$, then the h -vector of X_C is*

$$h_{X_C} = \left(1, n, \binom{n+1}{n-1}, \dots, \binom{v-2}{n-1}, \binom{v-1}{n-1} - |B|\right).$$

Proof. From Theorem 3.9 we have $\alpha(I_{\Delta_C})$ is $v - n$. From Remark 2.11, a set of minimal generators of degree $v - n$ has $|B|$ elements. So we only need to show that $h_{X_C}(v - n + 1) = 0$. This follows since $\sum_{j=0}^{v-n} h_{X_C}(j) = \binom{v}{n} - |B| = \deg X_C$. □

The regularity of I_{X_C} is an easy consequence of Proposition 4.1:

Corollary 4.2. $\text{reg}(I_{X_C}) = \alpha(I_{X_C}) + 1 = v - n + 1$.

Let (V, B) be a Steiner system of type $S(t, n, v)$. A minimal graded free resolution of I_{X_C} will be written as

$$(1) \quad 0 \rightarrow \bigoplus_j R(-j)^{\beta_{n-1,j}(I_{X_C})} \rightarrow \dots \rightarrow \bigoplus_j R(-j)^{\beta_{1,j}(I_{X_C})} \rightarrow \bigoplus_j R(-j)^{\beta_{0,j}(I_{X_C})} \rightarrow R \rightarrow R/I_{X_C} \rightarrow 0$$

where $R(-j)$ is the free R -module obtained by shifting the degrees of R by j , i.e. so that $R(-j)_a = R_{a-j}$.

The number $\beta_{i,j}(I_{X_C})$ is called the (i, j) -th graded Betti numbers of I_{X_C} and equals the number of minimal generators of degree j in the i -th syzygy module of I_{X_C} .

Another consequence of Proposition 4.1 is that

$$\beta_{i,j}(I_{X_C}) = 0 \text{ for any } j - i > v - n + 1.$$

This means that the nonzero graded Betti numbers only occur in two rows of the Betti table $\beta(I_{X_C}) := (\beta_{i,i+j}(I_{X_C}))_{i,j}$ of I_{X_C} .

The next proposition excludes the existence of first syzygies in degree $v - n + 1$.

Proposition 4.3. *If (V, B) is a Steiner system of type $S(t, n, v)$, then*

$$\beta_{1,v-n+1}(I_{X_C}) = 0.$$

Proof. From Proposition 3.6 it is enough to show that $\beta_{1,v-n+1}(J_C) = 0$. By contradiction, let $\alpha_1, \alpha_2 \in B$ be two distinct blocks of the Steiner system (V, B) such that the monomials $M_1 := M_{V \setminus \alpha_1}$ and $M_2 := M_{V \setminus \alpha_2}$ give a linear syzygy. Then $\gcd(M_1, M_2)$ has degree $v - n - 1$. This implies $|\alpha_1 \cap \alpha_2| = n - 1 \geq t$, a contradiction to Definition 2.3 of Steiner system. \square

Proposition 4.4. *If (V, B) is a Steiner system of type $S(t, n, v)$, then*

$$\beta_{0,v-n+1}(I_{X_C}) = n|B| - \binom{v}{n-1} = n \frac{\binom{v}{t}}{\binom{n}{t}} - \binom{v}{n-1}.$$

Proof. It is matter of computation to show that

$$\Delta h_{X_C}(v - n + 1) = -\binom{v-1}{n-1} + |B|$$

and

$$\Delta h_{X_C}(v - n) = \binom{v-1}{n-1} - |B| - \binom{v-2}{n-1} = \binom{v-2}{n-2} - |B|$$

and for $i < v - n$

$$\Delta h_{X_C}(i) = \binom{i+n-2}{n-2}.$$

Then

$$\Delta^2 h_{X_C}(v - n + 1) = -\binom{v-2}{n-2} - \binom{v-1}{n-1} + 2|B|$$

and

$$\Delta^2 h_{X_C}(v - n) = \binom{v-3}{n-3} - |B|$$

and for $i < v - n$

$$\Delta^2 h_{X_C}(i) = \binom{i+n-3}{n-3}.$$

So we have

$$\begin{aligned}\Delta^n h_{X_C}(v-n+1) &= n|B| - \left[\binom{v-1}{n-1} + \binom{v-2}{n-2} + \cdots + \binom{v-n}{0} \right] = \\ &= n \frac{\binom{v}{n}}{\binom{n}{n}} - \binom{v}{n-1}.\end{aligned}$$

Then the statement follows from Proposition 4.3. \square

Corollary 4.5. *If (V, B) is a Steiner system of type $S(t, n, v)$, then $\beta_{0, v-n+1}(I_{X_C}) = 0$ if and only if $t = n - 1$.*

Proof. From Proposition 4.4 we have

$$\begin{aligned}\beta_{0, v-n+1}(I_{X_C}) &= \Delta^n h_{X_C}(v-n+1) = n|B| - \left[\binom{v-1}{n-1} + \binom{v-2}{n-2} + \cdots + \binom{v-n}{0} \right] = \\ &= n \frac{\binom{v}{n}}{\binom{n}{n}} - \binom{v}{n-1}.\end{aligned}$$

If $t = n - 1$ then we get

$$\beta_{0, v-n+1}(I_{X_C}) = n \frac{\binom{v}{n-1}}{\binom{n-1}{n-1}} - \binom{v}{n-1} = 0.$$

If $\beta_{0, v-n+1}(I_{X_C}) = 0$ then we have

$$\frac{(n-t)!}{(v-t)!} = \frac{1}{(v-n+1)!}$$

and then

$$v-n+1 = \binom{v-t}{n-t}.$$

This implies

$$\binom{v-n+1}{v-n} = \binom{v-t}{v-n}$$

and then $t = n - 1$. \square

Corollary 4.6. *If $t = n - 1$, we have*

$$\beta_{i,j}(I_{X_C}) = \begin{cases} |B| & \text{if } (i, j) = (0, v-n) \\ (-1)^i \Delta^n h_{X_C}(v-n+1+i) & \text{if } j = v-n+1+i \\ 0 & \text{otherwise.} \end{cases}$$

Corollary 4.7. *We have*

$$\omega(I_{X_C}) = \begin{cases} \alpha(I_{X_C}) = v-n & \text{if } t = n-1 \\ \alpha(I_{X_C}) + 1 = \text{reg}(I_{X_C}) = v-n+1 & \text{if } t < n-1 \end{cases}$$

where $\omega(I_{X_C})$ is the largest degree in a minimal homogeneous set of generators of I_{X_C} .

Recalling that the resurgence and the asymptotic resurgence of I are defined as $\rho(I) := \sup \left\{ \frac{m}{d} \mid I^{(m)} \not\subseteq I^d \right\}$ and $\rho_a(I) = \sup \left\{ \frac{m}{r} : I^{(mt)} \not\subseteq I^{rt} \text{ for all } t \gg 0 \right\}$, respectively, the following results give the bounds:

Corollary 4.8. *Let (V, B) be a Steiner system of type $S(t, n, v)$.*

(1) *If $t = n - 1$ then*

$$\frac{(v-n)n}{v} = \rho_a(I_{X_C}) \leq \rho(I_{X_C}) \leq \frac{(v-n+1)n}{v}.$$

(2) *If $t < n - 1$*

$$\frac{(v-n)n}{v} \leq \rho_a(I_{X_C}) \leq \rho(I_{X_C}) \leq \frac{(v-n+1)n}{v}.$$

Proof. By Theorem 1.2.1 in [5] and by Theorem 3.9 and Corollary 4.7 we have

$$\frac{(v-n)n}{v} = \frac{\alpha(I_{X_C})}{\widehat{\alpha}(I_{X_C})} \leq \rho(I_{X_C}) \leq \frac{\text{reg}(I_{X_C})}{\widehat{\alpha}(I_{X_C})} = \frac{(v-n+1)n}{v}$$

□

Remark 4.9. From Lemma 2.3.4 in [5], if $d \text{reg}(I_X) \leq \alpha(I_X^{(m)})$, then $I_X^{(m)} \subseteq I^d$. Moreover, if $d\alpha(I_X) > \alpha(I_X^{(m)})$ then $I_X^{(m)} \not\subseteq I_X^d$. The range of values not covered by these bounds has length

$$\left\lfloor \frac{\alpha(I_X^{(m)})}{\alpha(I_X)} - \frac{\alpha(I_X^{(m)})}{\text{reg}(I_X)} \right\rfloor = \left\lfloor \frac{\alpha(I_X^{(m)})(\text{reg}(I_X) - \alpha(I_X))}{\alpha(I_X) \text{reg}(I_X)} \right\rfloor$$

For a C-Steiner configuration X_C , from Corollary 4.2, this number is equal to

$$\left\lfloor \frac{\alpha(I_{X_C}^{(m)})}{\alpha(I_{X_C}) \text{reg}(I_{X_C})} \right\rfloor.$$

Example 4.10. We compute the h -vector and the graded Betti numbers of a C-Steiner System X_C of type $S(2, 3, v)$. We have $|B| = \frac{v(v-1)}{6}$ and then

$$\begin{aligned} h_{X_C} &= (1, 3, 6, \dots, \binom{v-2}{2}, \binom{v-1}{2} - |B|) = \\ \Delta^3 h_{X_C} &= (1, 0, 0, \dots, 0, -|B|, 0, 2\binom{v-1}{2} + (v-2) - 3|B|, -\binom{v-1}{2} + |B|) = \\ &= \left(1, 0, 0, \dots, 0, -\frac{v(v-1)}{6}, 0, \frac{v(v-3)}{2}, -\frac{(v-1)(v-3)}{3}\right). \end{aligned}$$

In this case, we have: $\beta_{0,v-3} = \frac{v(v-1)}{6}$, $\beta_{1,v-2} = 0$, $\beta_{1,v-1} = -\frac{v(v-3)}{2}$ and $\beta_{2,v} = -\frac{(v-1)(v-3)}{6}$. In particular, $\alpha(I_{X_C}) = \omega(I_{X_C}) = v - 3$ and $\text{reg}(I_{X_C}) = v - 2$.

In the following example we compute the graded Betti numbers of a C-Steiner configuration of type $S(t, n, v)$ where $t < n - 1$.

Example 4.11. Consider a Steiner system (V, B) of type $S(2, 4, 13)$ where $|V| = 13$ and

$$\begin{aligned} B := & \{\{2, 3, 5, 11\}, \{3, 4, 6, 12\}, \{4, 5, 7, 13\}, \{1, 5, 6, 8\}, \{2, 6, 7, 9\}, \{3, 7, 8, 10\}, \\ & \{4, 8, 9, 11\}, \{5, 9, 10, 12\}, \{6, 10, 11, 13\}, \{1, 7, 11, 12\}, \{2, 8, 12, 13\}, \\ & \{1, 3, 9, 13\}, \{1, 2, 4, 10\}\}. \end{aligned}$$

Setting $C := C_{(4,13)} \setminus V$, we construct a C-Steiner configuration X_C in \mathbb{P}^4 . We have

$$h_{X_C} = (1, 4, 10, 20, 35, 56, 84, 120, 165, 207).$$

We have $\beta_{0,9} = 13$, $\beta_{0,10} = 234$, $\beta_{1,11} = 702$, $\beta_{2,12} = 663$, $\beta_{3,13} = 207$. In particular, we have $\alpha(I_{X_C}) = 9$ and $\omega(I_{X_C}) = \text{reg}(I_{X_C}) = 10$.

In general, the following question is open.

Question 4.12. Let $X_C \subseteq \mathbb{P}^n$ be a C-Steiner configuration of points of type $S(t, n, v)$. Do the graded Betti numbers and the Hilbert Function of $I_{X_C}^{(m)}$ only depend on the parameters (t, n, v) ?

We don't have any formula to describe the Hilbert function of $I_{X_C}^{(m)}$. For $m = 1$ Proposition 4.1 implies that the h -vector of X_C is a pure O -sequence (see [43] for all the terminology and background on this topic). This is not always true if $m > 1$. Using [1] and [34], we can show examples where the h -vectors could be not unimodal or not differentiable.

Example 4.13. Consider a Steiner system (V_1, B_1) of type $S(2, 3, 7)$. Set $X_{C_1} := X_{C_{(3,7)} \setminus B_1}$. Using [34], we found that $I_{X_{C_1}}^{(7)}$ has the following h -vector

$$h_{I_{X_{C_1}}^{(7)}} = (1, 3, 6, \dots, 153_{16}, 171, \mathbf{183}, \mathbf{182}, \mathbf{189}, 175, 140, 119, 84, 63, 42, 21, 14_{28})$$

that is a non-unimodal sequence (the index denotes the degree in which the dimension occurs, for instance, above the entry 153 is in degree 16).

Example 4.14. Consider a Steiner configuration (V_2, B_2) of type $S(2, 3, 9)$ and set $X_{C_2} := X_{C_{(3,7)} \setminus B_2}$. Using [34], we found that the h -vector of $I_{X_{C_2}}^{(10)}$ is

$$h_{I_{X_{C_2}}^{(10)}} = (1, 3, 6, \dots, 528_{31}, 561, \mathbf{583}, \mathbf{585}, \mathbf{603}, 621, 639, 621, 567, 540, 540, 528, 468, 396, \\ 360, 360, 324, 252, 216, 216, 204, 144, 108, 108, 108, 72, 36, 36, 36, 24_{60})$$

that is unimodal but not differentiable (the positive part of the first difference is not an O -sequence).

We end this section with the following questions on $X_{\mathcal{H},B}$ suggested by experimental evidences using [1] and [34]. From a combinatorial point of view, two Steiner systems having the same parameters could have very different properties. We have examples where such differences effect the homological invariants.

Question 4.15. Let (V, B) be a Steiner system of type $S(t, n, v)$, and $X_{\mathcal{H},B}$ the associated Steiner configuration of points. Assume that the hyperplanes in \mathcal{H} are chosen generically. Do the Hilbert function and the graded Betti numbers of $X_{\mathcal{H},B}$ only depend on t, n, v ?

Question 4.16. Let (V, B) be a Steiner system of type $S(t, n, v)$, and $X_{\mathcal{H},B}$ the associated Steiner configuration of points. Assume that the hyperplanes in \mathcal{H} are chosen generically. Are the Hilbert function and the graded Betti numbers of $X_{\mathcal{H},B}$ generic with respect to the Hilbert function? (i.e. the same as a set of $|B|$ generic points in \mathbb{P}^n ?)

5. APPLICATION TO CODING THEORY

In this section we show an application of the previous results on Steiner and C-Steiner configurations of points to coding theory. We recall the basic notion on linear coding and we compute the parameters of a linear code associated to a Steiner configuration and a C-Steiner configuration of points in \mathbb{P}^n . We refer to [16, 26, 56, 57, 58, 59] for a detailed initial motivation to study the connections between the minimum distance and some invariants coming from commutative/homological algebra. There are several ways to compute the minimum distance. One of them comes from linear algebra.

Let k be any field and $X = \{P_1, \dots, P_r\} \subseteq \mathbb{P}^n$ a not degenerate finite set of reduced points. The linear code associated to Γ denoted by $\mathcal{C}(X)$ is the image of the injective linear map $\varphi : k^{n+1} \rightarrow k^r$.

We are interested in three parameters $[|X|, k_X, d_X]$ that we use to evaluate the goodness of a linear code. The first number $|X|$ is the cardinality of X . The number k_X is the dimension of the code as k -linear vector space, that is the rank of the matrix associated to φ . The number d_X denotes the minimal distance of $\mathcal{C}(X)$, that is the minimum of the Hamming distance of two elements in $\mathcal{C}(X)$. The *Singleton bound* gives always an upper bound for this number: $d_X \leq |X| - n$. When $d_X = |X| - n$ the code is called a maximal distance separable code (MDS code).

The linear code associated to X has generating matrix of type $(n+1) \times r$

$$A(X) = [c_1 \dots c_r]$$

where c_i are the coordinates of P_i . Then the linear code $\mathcal{C}(X)$ has parameters $[|X|, \text{Rank}(A(X)), d_X]$. Assuming that $A(X)$ has no proportional columns is equivalent to say that the points P_i are distinct points in \mathbb{P}^n . Then $|X| = r$, $\text{Rank}(A(X)) = n+1$ and $r - d_X$ is the maximum number of these points that fit in a hyperplane of \mathbb{P}^n . Remark 2.2 in [56, 57] says that the minimum distance d_X is also the minimum number such that $r - d_X$ columns in $A(X)$ span an n -dimensional space. The generating matrix $A(X)$ of an $[|X|, n+1, d]$ -linear code \mathcal{C} naturally determines a matroid $M(\mathcal{C})$.

Denoted by $\text{hyp}(X)$ the maximum number of points contained in some hyperplane, d_X has also geometrical interpretation, that is $d_X = |X| - \text{hyp}(X)$, see Section 2 in [56] and Remark 2.7 in [58]. In particular, the authors borrowed this terminology from coding theory since d_X is exactly the minimum distance of the (equivalence class of) linear codes with generating matrix having as columns the coordinates of the points of X .

Set $X_B := X_{\mathcal{H}, B}$. We apply the results of the previous sections to compute the parameters of linear codes associated to both a Steiner configuration X_B of points and its Complement X_C .

Proposition 5.1. *Let (V, B) be a Steiner system of type $S(t, n, v)$ with $|V| = v$. If X_B is the Steiner configuration of points and X_C its Complement, we have*

$$\text{hyp}(X_B) = \frac{\binom{v-1}{t-1}}{\binom{n-1}{t-1}}$$

and

$$\text{hyp}(X_C) = \binom{v-1}{n-1} - \frac{\binom{v-1}{t-1}}{\binom{n-1}{t-1}}.$$

Proof. Let $H_1 \in \mathcal{H}$ be one of the hyperplanes involved in the construction of a Steiner configuration. Then it is clear by definition that the number of blocks in B containing 1, i.e., the number of points in H_1 is

$$\text{hyp}(X_B) = \frac{\binom{v-1}{t-1}}{\binom{n-1}{t-1}}.$$

Since the star configuration $X_{\mathcal{H}, C(n,v)}$ has $\binom{v-1}{n-1}$ points lying on H_1 , we get

$$\text{hyp}(X_C) = \binom{v-1}{n-1} - \frac{\binom{v-1}{t-1}}{\binom{n-1}{t-1}}.$$

□

Proposition 5.2. *Let (V, B) be a Steiner system of type $S(t, n, v)$ with $|V| = v$. Then a Steiner configuration of points $X_B \subseteq \mathbb{P}^n$ defines a linear code with*

$$d_{X_B} = \frac{\binom{v}{t}}{\binom{n}{t}} - \frac{\binom{v-1}{t-1}}{\binom{n-1}{t-1}}.$$

A C -Steiner configuration of points $X_C \subseteq \mathbb{P}^n$ defines a linear code with

$$d_{X_C} = \binom{v}{n} - \frac{\binom{v}{t}}{\binom{n}{t}} - \binom{v-1}{n-1} + \frac{\binom{v-1}{t-1}}{\binom{n-1}{t-1}}.$$

Proof. The statement follows from Proposition 5.1 since, from Remark 2.7 in [58], for a set of points X it is $d_X = |X| - \text{hyp}(X)$. □

Remark 5.3. For a Steiner triple system $S(2, 3, v)$ with block set B , define $C := C_{(3,v)} \setminus B$. Then X_B and X_C have the following minimal distance

$$d_{X_B} = \frac{\binom{v}{2}}{3} - \frac{v-1}{2} = \frac{(v-1)(v-3)}{6}$$

and

$$d_{X_C} = \binom{v}{3} - \frac{\binom{v}{2}}{3} - \binom{v-1}{2} + \frac{v-1}{2} = \frac{(v-1)(v-3)^2}{6}.$$

With the above results, we have

Theorem 5.4. *Let (V, B) be a Steiner system $S(t, n, v)$ with $|V| = v$. Then*

- (1) *the parameters of the linear code defined by a Steiner configuration of points X_B are $[|B|, n+1, d_{X_B}]$;*
- (2) *the parameters of the linear code defined by a Complement of a Steiner configuration of points X_C are $[\binom{v}{n} - |B|, n+1, d_{X_C}]$.*

Theorem 5.5. *Let (V, B) be a Steiner system $S(t, n, v)$ with $|V| = v$.*

- (1) *If $n = \frac{\binom{v-1}{t-1}}{\binom{n-1}{t-1}}$ then $\mathcal{C}(X_B)$ is a MDS code;*
- (2) *if $n = \binom{v-1}{t-1} - \frac{\binom{v-1}{t-1}}{\binom{n-1}{t-1}}$ then $\mathcal{C}(X_C)$ is a MDS code;*

Example 5.6. Consider the Steiner system $S(2, 3, 7)$. The blocks are, up to isomorphism, $B := \{\{1, 2, 3\}, \{1, 4, 5\}, \{1, 6, 7\}, \{2, 4, 6\}, \{2, 5, 7\}, \{3, 4, 7\}, \{3, 5, 6\}\}$.

Let $\ell_i := x + 2^i y + 3^i z + 5^i w \in \mathbb{C}[x, y, z, w] = \mathbb{C}[\mathbb{P}^3]$ be linear forms and let $H_i \subseteq \mathbb{P}^3$ be the hyperplane defined by ℓ_i for $i = 1, \dots, 7$. Set $\mathcal{H} := \{H_1, \dots, H_7\}$. One can check that any three hyperplanes in \mathcal{H} meet in one point. Computing with Cocoa [1] the generating matrix of the linear code $\mathcal{C}(X_{\mathcal{H}, B})$ defined by the Steiner configuration on B we get

$$A(X_{\mathcal{H}, B}) := \begin{pmatrix} -15 & -1983 & -438045 & -350 & -639000 & 9315 & 104625 \\ 20 & 1576 & 269060 & 160 & 240075 & -2610 & -25875 \\ -10 & -418 & -34230 & -35 & -37550 & 470 & 4250 \\ 1 & 17 & 523 & 1 & 666 & -9 & -99 \end{pmatrix}$$

where the columns are the seven coordinates (among the thirty-five) of the intersection points of any three hyperplanes H_1, \dots, H_7 corresponding to the seven blocks B . The parameters of the code $\mathcal{C}(X_{\mathcal{H}, B})$ are $[7, 4, 4]$.

We note that only in this particular case, the linear code $\mathcal{C}(X_{\mathcal{H}, B})$ associated to a Steiner system of type $S(2, 3, 7)$ has $d_{X_B} = 4 = 7 - 3$ and $\mathcal{C}(X_{\mathcal{H}, B})$ is a maximal distance separable code (MDS).

We now compute linear code $\mathcal{C}(X_{\mathcal{H}, C_{(3,7)} \setminus B})$ associated to the Complement of the Steiner configuration. We get

$$A(X_{\mathcal{H}, C_{(3,7)} \setminus B}) := \begin{pmatrix} 75 & -207 & 615 & 4845 & -465 & -70 & 27585 & -190 & 21300 & -5985 & 36273 & -3610 \\ -90 & 232 & -660 & -5060 & 460 & 64 & -23980 & 160 & -16005 & 4340 & -24728 & 2368 \\ 40 & -94 & 250 & 1830 & -170 & -21 & 7190 & -45 & 3755 & -930 & 4586 & -387 \\ -3 & 5 & -9 & -43 & 11 & 1 & -239 & 1 & -111 & 19 & -115 & 7 \\ 225 & 1125 & -621 & 9225 & 6975 & 413775 & -29745 & -108819 & -3375 & -33750 & 5250 \\ -150 & -675 & 348 & -4950 & -3450 & -179850 & 11820 & 37092 & 1125 & 10125 & -1200 \\ 50 & 200 & -94 & 1250 & 850 & 35950 & -2090 & -4586 & -250 & -2000 & 175 \\ -3 & -9 & 3 & -27 & -33 & -717 & 51 & 69 & 9 & 54 & -3 \\ -446175 & -101250 & 1012500 & -3138750 & 3037500 & & & & & & \\ 88650 & 16875 & -151875 & 388125 & -253125 & & & & & & \\ -10450 & -2500 & 20000 & -42500 & 25000 & & & & & & \\ 153 & 54 & -324 & 594 & -324 & & & & & & \end{pmatrix}$$

where the columns are the twenty-eight coordinates of the intersection points (among the thirty-five) of any three hyperplanes H_1, \dots, H_7 corresponding to the twenty-eight blocks

$$C_{(3,7)} \setminus B = \{\{1, 2, 4\}, \{1, 2, 5\}, \{1, 2, 6\}, \{1, 2, 7\}, \{1, 3, 4\}, \{1, 3, 5\}, \{1, 3, 6\}, \{1, 3, 7\}, \{1, 4, 6\}, \\ \{1, 4, 7\}, \{1, 5, 6\}, \{1, 5, 7\}, \{2, 3, 4\}, \{2, 3, 5\}, \{2, 3, 6\}, \{2, 3, 7\}, \{2, 4, 5\}, \{2, 4, 7\}, \\ \{2, 5, 6\}, \{2, 6, 7\}, \{3, 4, 5\}, \{3, 4, 6\}, \{3, 5, 7\}, \{3, 6, 7\}, \{4, 5, 6\}, \{4, 5, 7\}, \{4, 6, 7\}, \{5, 6, 7\}\}.$$

The parameters of the code $\mathcal{C}(X_{\mathcal{H}, C_{(3,7)} \setminus B})$ are $[28, 4, 16]$.

REFERENCES

- [1] Abbott J., Bigatti A.M., Robbiano L., *CoCoA: a system for doing Computations in Commutative Algebra*. Available at <http://cocoa.dima.unige.it>
- [2] Bocci C., Cooper S.M., Guardo E., Harbourne B., Janssen M., Nagel U., Seceleanu A., Van Tuyl A., Vu T. *The Waldschmidt constant for squarefree monomial ideals*. Journal of Algebraic Combinatorics. 2016 Dec 1;Vol.44(4), 875-904.
- [3] Bocci C., Harbourne B., *The resurgence of ideals of points and the containment problem* Proceedings of the American Mathematical Society Vol. 138(4), (APRIL 2010), 1175-1190
- [4] Bocci C., Harbourne B., *Comparing powers and symbolic powers of ideals*. Journal of Algebraic Geometry, 19(3), (2010), 399-417.
- [5] Bruns, W., Herzog, J. *Cohen-Macaulay Rings*, vol. 39, Cambridge studies in advanced mathematics, revised edition, 1998.
- [6] Buratti M., Gionfriddo G., Milazzo L., Voloshin V. *Lower and upper chromatic numbers for $BSTS(2^h - 1)s$* . Comput. Sci. J. Moldova, Vol. 9 (2), (2001), 259-272.
- [7] Buratti M., Rinaldi G., Traetta T. *3-pyramidal Steiner triple systems*. Ars Math. Contemp. Vol 13, No 1 (2017)

- [8] Carlini E., Catalisano M. V., Guardo E., Van Tuyl A., *Hadamard Star Configurations*. Rocky Mount. J. Math, Vol. 49(2), (2019), 419–432.
- [9] Carlini E., Guardo E., Van Tuyl A., *Star configurations on generic hypersurfaces*, J. Algebra, Vol. 407, (2014), 1–20.
- [10] Catalisano M.V., Guardo E., Shin Y.S., *The Waldschmidt constant of special k -configurations in \mathbb{P}^n* , Journal of Pure and Applied Algebra, Available online 17 February 2020, <https://doi.org/10.1016/j.jpaa.2020.106341>
- [11] Chudnovsky G.V. *Singular points on complex hypersurfaces and multidimensional Schwarz Lemma*, Seminaire de Theorie des Nombres, Paris 1979–80, Seminaire Delange–Pisot–Poitou, Progress in Math vol. 12, M–J Bertin, editor, Birkhauser, Boston–Basel–Stuttgart 1981
- [12] Colbourn C.J., Dinitz J. H. *CRC handbook of combinatorial designs*. CRC press; 2010 Dec 12.
- [13] Colbourn C.J., Pulleyblank W.R. *Matroid Steiner problems, the Tutte polynomial and network reliability*. Journal of Combinatorial Theory, Series B. 1989 Aug 1; Vol 47(1), 20–31.
- [14] Colbourn C.J., Rosa A. *Triple systems*. Oxford University Press; 1999.
- [15] Cooper S.M., Embree R.J., Há H.T., Hoefel A.H. *Symbolic powers of monomial ideals*. Proceedings of the Edinburgh Mathematical Society. 2017 Feb; Vol.60(1), 39–55.
- [16] Cooper S.M., Guardo E. *Fat points, partial intersections and Hamming distance* Journal of Algebra and Its Applications, Vol.19 (4), (2020), 2050071 (25 pages), <https://doi.org/10.1142/S0219498820500711>
- [17] Cooper S.M., Seceleanu A., Tohaneanu S., Vaz Pinto M., Villarreal R. H. *GMD functions for scheme based linear codes and algebraic invariants of Geramita ideals* Advances in Applied Mathematics, Vol. 112, (2020), 101940.
- [18] Czapliński A., Glówka A., Malara G., Lampa-Baczyńska M., Luszcz-Swidecka P., Pokora P., Szpond J. *A counterexample to the containment $I^{(3)} \subseteq I^2$ over the reals*. Advances in Geometry. 2016 Jan 1; vol.16(1), 77–82.
- [19] Dao H., De Stefani A., Grifo E., Huneke C., Núñez-Betancourt L. *Symbolic powers of ideals*. In Singularities and Foliations. Geometry, Topology and Applications 2015 Jul 13 387–432). Springer, Cham.
- [20] De Boer M., Pellikaan R. *Groebner bases for codes*. Some tapas of computer algebra. Chap. 10, Springer, Berlin (1999), 237–259.
- [21] Dumnicki M., Harbourne B., Nagel U., Seceleanu A., Szemberg T., Tutaj-Gasińska H. *Resurgences for ideals of special point configurations in \mathbb{P}^N coming from hyperplane arrangements*. Journal of Algebra. 2015 Dec 1; Vol. 443, 383–394.
- [22] Eisenbud D., *Commutative Algebra: with a view toward algebraic geometry*, Springer-Verlag, New York (1995)
- [23] Favacchio G., Guardo E., Migliore J. *On the arithmetically Cohen-Macaulay property for sets of points in multiprojective spaces*. Proc. Amer. Math. Soc. Vol.146 (2018), 2811–2825. PDF DOI: <https://doi.org/10.1090/proc/13981>
- [24] Favacchio G., Guardo E., Picone B. *Special arrangements of lines: codimension two ACM varieties in $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$* . Journal of Algebra and its Applications, Vol. 18(4), 1 April 2019
- [25] Favacchio G., Migliore J. *Multiprojective spaces and the arithmetically Cohen-Macaulay property*. Math. Proc. of the Cambridge Philosophical Society. Vol.166 (3), (2019), 583–597.
- [26] Garrounian M., Tohaneanu S.O. *Minimal distance of linear codes and α -invariants*. Advances in Applied Mathematics, Vol. 71 (2015), 190–207.
- [27] Geramita A., Harbourne B., Migliore J. *Star configurations in \mathbb{P}^n* . Journal of Algebra, Vol. 376, (2013), 279–299.
- [28] Geramita A., Harbourne B., Migliore J., Nagel U. *Matroid configurations and symbolic powers of their ideals*. Transactions of the American Mathematical Society. 2017; Vol.369(10):7049–66.
- [29] Gionfriddo L. *A conjecture of Berge about linear hypergraphs and Steiner systems $S(2, 4, v)$* , Discrete Math. Vol.255 (2002), 125–133.
- [30] Gionfriddo M., Milazzo L., Rosa A., Voloshin V. *Bicoloring Steiner systems $S(2, 4, v)$* , Discrete Math. Vol.283 (2004), 249–253.
- [31] Gionfriddo M., Guardo E., Milazzo L. *Extending bicolorings for Steiner triple systems*, Appl. Anal. Discrete Math. Vol. 7 (2), (2013), 225–234.
- [32] Gionfriddo M., Guardo E., Milazzo L., Voloshin V. *Feasible sets of small bicolorable STSs*, Austr. J. of Math, Vol. 59(1) (2014), Pages 107–119.

- [33] Gold L., Little J., Schenck H. *Cayley-Bacharach and evaluation codes on complete intersections*. J. Pure Appl. Algebra, Vol. 196, (2005), 91–99.
- [34] Grayson D.R., Stillman M.E. *Macaulay 2, a software system for research in algebraic geometry*
- [35] Guardo E., Harbourne B., Van Tuyl A. *Asymptotic resurgences for ideals of positive dimensional subschemes of projective space*. Adv. in Math., Vol 246 (2013), 114–127, <http://dx.doi.org/10.1016/j.aim.2013.05.027>.
- [36] Hansen J., *Linkage and codes on complete intersections*. Appl. Algebra Engrg. Comm. Comput. Vol. 14 (3) (2003), 175–185.
- [37] Herzog J., Hibi T. *Monomial ideals*. 2011. Springer, London.
- [38] Jurrius R., Pellikaan R., *Codes, arrangements and matroids*, E. Matinez-Moro (Ed), Algebraic Geometry Modeling in Information Theory, Ser. Coding Theory Cryptol., Vol. 8, World Scientific (2013), 219–325
- [39] Lampa-Baczyńska M., Malara G. *On the containment hierarchy for simplicial ideals*. Journal of Pure and Applied Algebra. 2015 Dec 1;219(12):5, 402–12.
- [40] Lindner C.C., Rosa A. *Steiner quadruple systems - a survey*, Discrete Math. 21 (1978), 147–181.
- [41] Martinez-Bernal J., Pitones Y., Villarreal R.H. *Minimum Distance Functions of Complete Intersections*. Journal of Algebra and Its Applications, 2017 (<https://www.worldscientific.com/doi/abs/10.1142/S0219498818502043>).
- [42] Migliore J., Introduction to liaison theory and deficiency modules, Birkhauser, Progress in Mathematics vol. 165 (1998).
- [43] Migliore J., Nagel U., Zanello F. *Pure O-Sequences: Known Results, Applications, and Open Problems*. In: Peeva I. (eds) Commutative Algebra. Springer, New York, NY (2013) .
- [44] Milazzo L., Tuza Zs. *Upper chromatic number of Steiner triple systems and quadruple systems*, Discrete Math. 174 (1997), 247–259.
- [45] Milazzo L., Tuza Zs. *Strict colourings for classes of Steiner triple systems*, Discrete Math. 182 (1998), 233–243.
- [46] Milazzo L., Tuza Zs. *Logarithmic upper bound for the upper chromatic number of $S(t, t+1, v)$ systems*, Ars Combinatoria 92 (2009), 213–223.
- [47] Milazzo L., Tuza Zs., Voloshin V., *Strict colorings of Steiner triple and quadruple systems: a survey*, Discrete Math. 261 (2003), 399–411.
- [48] Milici S., *Colouring Steiner triple systems*, Rendiconti del Seminario Matematico di Messina Serie II Tomo XXV Supplemento al n.8 (2002), 113–118.
- [49] Milici S., Rosa A., Voloshin V., *Steiner systems with specified block colour patterns*, Discrete Mathematics, 240 (2001), 145–160.
- [50] Minh N.C, Trung N.V. *Cohen-Macaulayness of monomial ideals and symbolic powers of Stanley-Reisner ideals*. Advances in mathematics. 2011;226(2):1285–306.
- [51] Reid C., Rosa A.. *Steiner systems $S(2, 4, v)$ - a survey*.
- [52] Smith G.G., Sturmfels B., *Combinatorial Algebraic Geometry: Selected papers From 2016 Apprenticeship Program*, Springer (2017).
- [53] Swartz E. *Topological representations of matroids*. Journal of the American Mathematical Society. 2003;16(2):427–42.
- [54] Szemberg T., Szpond J. *On the containment problem*. Rendiconti del Circolo Matematico di Palermo Series 2. 2017 Aug 1;66(2):233–45.
- [55] Terai N., Trung N.V. *CohenMacaulayness of large powers of StanleyReisner ideals*. Advances in Mathematics. 2012 Jan 30;229(2):711–30.
- [56] Tohaneanu S.O.. *The minimum distance of sets of points and the minimum socle degree*. Journal of Pure and Applied Algebra. 2011 Nov 1;215(11):2645–51.
- [57] Tohaneanu S.O. *Lower bounds on minimal distance of evaluation codes*. Applicable Algebra in Engineering, Communication and Computing. 2009 Dec 1;20(5-6):351.
- [58] Tohaneanu S.O., Van Tuyl A. *Bounding invariants of fat points using a coding theory construction*. Journal of Pure and Applied Algebra. 2013 Feb 1;217(2):269–79.
- [59] Tohaneanu S.O., Xie Y.. *Homological properties of ideals generated by α -product of linear forms*, Preprint 2019, <https://128.84.21.199/pdf/1906.08346v3.pdf>
- [60] van Lint J., van der Geer G. *Introduction to Coding Theory and Algebraic Geometry*. Birkhäuser, 6 dic 2012 .

- [61] Van Maldeghem H. *Some remarks on Steiner Systems*. Design, Codes and Cryptography, 29, (2003) pp 199–213
- [62] Varbaro M. *Symbolic powers and matroids*. Proceedings of the American Mathematical Society. 2011;139(7):2357–66.
- [63] Villareal R., *Monomial Algebras*, Monographs and textbooks in Pure and Applied Mathematics, **238**, Marcel Dekker, Inc NY 2001.
- [64] Waldschmidt M. *Propriétés arithmétiques de fonctions de plusieurs variables II*. Séminaire P. Lelong (Analyse), 1975/76, pp. 108–135, Lecture Notes Math. 578, Springer, Heidelberg (1977)

DIPARTIMENTO DI MATEMATICA, VIA SOMMARIVE, 14 - 38123 POVO (TN), ITALY
E-mail address: `edoardo.ballico@unitn.it`

DIPARTIMENTO DI MATEMATICA E INFORMATICA, VIALE A. DORIA, 6 - 95100 - CATANIA, ITALY
E-mail address: `favacchio@dmf.unict.it`
URL: `www.dmf.unict.it/~gfavacchio`

DIPARTIMENTO DI MATEMATICA E INFORMATICA, VIALE A. DORIA, 6 - 95100 - CATANIA, ITALY
E-mail address: `guardo@dmf.unict.it`
URL: `www.dmf.unict.it/~guardo`

DIPARTIMENTO DI MATEMATICA E INFORMATICA, VIALE A. DORIA, 6 - 95100 - CATANIA, ITALY
E-mail address: `milazzo@dmf.unict.it`
URL: `www.dmf.unict.it/~milazzo`