

Exploring Consequences of Simulation Design for Apparent Performance of Statistical Methods. 2: Results from simulations with normally and uniformly distributed sample sizes

Elena Kulinskaya, David C. Hoaglin, and Ilyas Bakbergenuly

February 12, 2022

Abstract

This report continues our investigation of effects a simulation design may have on the conclusions on performance of statistical methods. In the context of meta-analysis of log-odds-ratios, we consider five generation mechanisms for control probabilities and log-odds-ratios. Our first report (Kulinskaya et al. [2020]) considered constant sample sizes. Here we report on the results for normally and uniformly distributed sample sizes.

1 Introduction

Our interest lies in effects that simulation design choices may have on conclusions on the comparative merits of various methods, taking as an example meta-analysis of odds ratios. The basic data from K studies involve $2K$ binomial variables, $X_{ij} \sim \text{Bin}(n_{ij}, p_{ij})$ for $i = 1, \dots, K$ and $j = C$ or T (for the Control or Treatment arm); those data underlie the odds-ratios for the meta-analysis.

A design specifies the number of studies, K ; the sample sizes, n_{ij} ; the nuisance parameters (control-arm probabilities, p_{iC} , or, equivalently, their logits, α_i); the overall log-odds-ratio, θ ; and the between-study variance, τ^2 . For each situation the simulation uses M replications, where M is typically large, say 10,000.

For simplicity, we consider equal arm-level sample sizes, $n_{iC} = n_{iT} = n_i$. The control probabilities p_{iC} or their logits α_i can be constant or generated from some distribution. Normal and uniform distributions are the typical choices. As in our previous report Kulinskaya et al. [2020], we consider five possible generation mechanisms for control-arm probabilities and log-odds-ratios under the random-effects model of meta-analysis.

We consider two fixed-intercept random-effects models (FIM1 and FIM2) and two random-intercept random-effects models (RIM1 and RIM2), as in Bakbergenuly and Kulinskaya [2018]. These models are equivalent to Models 2 and 4 (for FIM) and Models 3 and 5 (for RIM), respectively, of Jackson et al. [2018]. Briefly, the FIMs include fixed control-arm effects (log-odds of the control-arm probabilities), and the RIMs replace these fixed effects with random effects. We also consider a model with uniformly distributed control-arm probabilities (URIM1).

Studies also vary in how they specify the sample sizes n_i . In our previous report (Kulinskaya et al. [2020]) we set $n_1 = \dots = n_K$ in all M replications. Here we investigate the use of normal and uniform distributions to generate a new set of n_i in each replication.

2 Generation of sample sizes

Several authors Cheng et al. [2016], Bakbergenuly and Kulinskaya [2018] use constant study-level sample sizes, either equal or unequal, in all replications. More often, however, authors generate sample sizes from a uniform or normal distribution. Jackson et al. [2018] use (mostly with $n_{iC} = n_{iT}$) sample sizes from discrete $U(50, 500)$. Langan et al. [2018] use either constant and equal sample sizes within and across studies, or sample sizes from $U(40, 400)$ and $U(2000, 4000)$; Sidik and Jonkman [2007] use $U(20, 200)$; and Abo-Zaid et al. [2013] use $U(30, 100)$ and $U(30, 1000)$. Viechtbauer [2007] generates study-level sample sizes ($n_i = n_{iC} = n_{iT}$) from $N(n, n/4)$ ($n/4$ is the variance) with $n = 10, 20, 40, 80, 160$. In an extensive simulation study for sparse data, Kuss [2015] uses FIM1 and the corresponding model with $\tau^2 = 0$, along with a large number of fitting methods; he generates both the number of studies K and their sample sizes n from log-normal distributions: $\text{LN}(0.65, 1.2)$ and $\text{LN}(3.05, 0.97)$ for K and $\text{LN}(4.615, 1.1)$ for sample sizes.

In general, if mutually independent random variables Y_i have a common distri-

bution $F(\cdot)$, and $N \sim G_n(\cdot)$ is independent of the Y_i , the sum $Y_1 + \dots + Y_N$ has a compound distribution Grubbström and Tang [2006]. A binomial distribution with a random number of trials is a compound Bernoulli distribution. The first two moments of such a distribution are $E(X) = pE(N)$ and $\text{Var}(X) = p(1-p)E(N) + p^2\text{Var}(N)$. This variance is larger than the variance of the $\text{Bin}(E(N), p)$ distribution. Therefore, random generation of sample sizes produces an overdispersed Binomial (compound Bernoulli) distribution for the control arm, and may also inflate, though in a more complicated way, the variance in the treatment arm.

In particular, when $N \sim N(E(N), \sigma_n^2)$, the compound Bernoulli distribution has variance $\text{Var}(X) = p(1-p)E(N) + p^2\sigma_n^2$. And when $N \sim U(n_l, n_u)$, $\text{Var}(X) = p(1-p)E(N) + p^2(n_u - n_l)^2/12$.

3 Variances of estimated log-odds-ratios for random sample sizes

The (conditional, given p_{ij} and n_{ij}) variance of the estimated log-odds-ratio $\hat{\theta}_i$, derived by the delta method, is

$$v_i^2 = \text{Var}(\hat{\theta}_i) = \frac{1}{n_{iT}p_{iT}(1-p_{iT})} + \frac{1}{n_{iC}p_{iC}(1-p_{iC})}, \quad (3.1)$$

estimated by substituting \hat{p}_{ij} for p_{ij} . (We follow the particular method's procedure for calculating \hat{p}_{ij} .)

Under the binomial-normal random-effects model (REM), the true study-level effects, θ_i , follow a normal distribution: $\theta_i \sim N(\theta, \tau^2)$.

To calculate the variance of $\hat{\theta}_i$ when sample sizes n_i are random, we use the law of total variance:

$$\text{Var}(\hat{\theta}_i) = E(\text{Var}(\hat{\theta}_i|n_i)) + \text{Var}(E(\hat{\theta}_i|n_i)).$$

The second term is $\text{Var}(\theta) = 0$, and the first term is obtained by substituting $E(n_{iC}^{-1})$ and $E(n_{iT}^{-1})$ in an expression for the variance of $\hat{\theta}$ under fixed sample sizes.

For a random sample size N , using the delta method,

$$E(N^{-1}) = (E(N))^{-1}(1 + [\text{CV}(N)]^2), \quad (3.2)$$

where CV is the coefficient of variation (i.e., the ratio of the standard deviation of N to its mean). Therefore, to order $1/E(N)$, random generation of sample sizes inflates the variance of $\hat{\theta}$ if and only if the coefficient of variation of the distribution of sample

sizes is of order 1. In the simulations of Viechtbauer [2007], where $\text{Var}(N) = n/4$, $\text{CV}(N) = O(1/\sqrt{n})$, so the variance is not inflated. In contrast, generating sample sizes from $N(n, n^2/4)$ would result in $\text{CV} = 1/2$ and would inflate variance. (Use of such a combination of mean and variance, however, is unlikely to produce realistic sets of sample sizes, and the probability of generating a negative sample size exceeds 2%.)

The variance of a uniform distribution on an interval of width Δ centered at n_0 is $\Delta^2/12$, and its CV is $\Delta/(\sqrt{12}n_0)$. Therefore, $\text{CV}(N)$ is of order 1 whenever the width of the interval is of the same order as its center. Hence, variance is inflated in simulations by Jackson et al. [2018], Langan et al. [2018], Sidik and Jonkman [2007], and Abo-Zaid et al. [2013], who all use wide intervals for n .

4 Design of the simulations for randomly distributed sample sizes

Our simulations keep the arm-level sample sizes equal and the control-arm probabilities p_{iC} and the log-odds-ratios θ_i independent. Table 1 shows the components of the simulations for normally and uniformly distributed sample sizes: parameters, data-generation mechanisms, and estimation targets. Our first report Kulinskaya et al. [2020] provides more details. We included the DerSimonian-Laird (DL), restricted maximum-likelihood (REML), Mandel-Paule (MP), and Kulinskaya-Dollinger (KD) estimators of τ^2 with corresponding inverse-variance-weighted estimators of θ and confidence intervals with critical values from the normal distribution. Bakbergenuly et al. [2020] studied those inverse-variance-weighted estimators in detail. We also included the SSW point estimator of θ , whose weights depend only on the studies' arm-level sample sizes, and a corresponding confidence interval, which uses $\hat{\theta}_{SSW}$ as the midpoint, $\hat{\tau}_{KD}^2$ in the estimate of its variance, and critical values from the t distribution on $K - 1$ degrees of freedom. Among the estimators, FIM2 and RIM2 denote the estimators in the corresponding GLMMs.

We generated the arm-level sample sizes, n_i , from a normal or a uniform distribution centered at 40, 100, 250, and 1000.

In generating sample sizes from a normal distribution, we want negative sample sizes to have reasonably small probability. For our choice of $\sigma_n^2 = 1.21n^2$ this probability is 0.0008. Unfortunately, we were still getting a small number of values below

zero out of thousands of simulated values, so we additionally truncate the n values generated from a normal distribution at 10. Truncation happens with probability 0.009.

To make uniform distributions of sample sizes comparable to the normal distributions, we centered them at the same value, n , and equated their variances. If a normal distribution has variance σ_n^2 , a uniform distribution with the same variance has interval width $\Delta_n = \sqrt{12\sigma_n^2}$. We set $\Delta_n = 1.1n$, resulting in $CV = \Delta_n/(\sqrt{12}n) = 0.318$ and a squared CV of 0.101. Therefore, by Equation (3.2), our simulations with random n inflate variances and covariances by 10% in comparison with simulations with constant n . Wider intervals of n would inflate variances more, but in generating sample sizes from a corresponding normal distribution, we wanted negative sample sizes to have reasonably small probability. For our choice of Δ_n this probability is 0.0008.

5 Summary of the results

Our simulations explored two main components of design: the data-generation mechanism and the distribution of study-level sample sizes. Results of our simulations with normally distributed sample sizes are provided in Appendix A, and those with uniformly distributed sample sizes in Appendix B.

The five data-generation mechanisms (FIM1, FIM2, RIM1, RIM2, and URIM1) often produced different results for at least one of the measures of performance (bias of estimators of τ^2 , bias of estimators of θ , and coverage of confidence intervals for θ). In the most frequent pattern FIM2 and RIM2 yield similar results, and FIM1, RIM1, and URIM1 also yield results that are similar but different from those of FIM2 and RIM2. In some situations URIM1 stands apart.

We also expected the coverage of θ to suffer because random sample sizes increase the variance of generated log-odds-ratios. However, generation of sample sizes from normal and uniform distributions had essentially no impact, as can be seen by comparing the results from this report with those from our report Kulinskaya et al. [2020] on the simulations with constant sample sizes. The explanation may lie in our choice of variance σ_n^2 (not large enough) for the normal and uniform distributions of the sample sizes, causing an increase of just 10% in the variance of LORs, or in the rather low coverage, even under constant sample sizes, resulting from considerable

biases of estimators of θ .

References

- G. Abo-Zaid, B. Guo, J. J. Deeks, T. P. A. Debray, E. W. Steyerberg, K. G.M. Moons, and R. David. Riley. Individual participant data meta-analyses should not ignore clustering. *Journal of clinical epidemiology*, 66(8):865–873, 2013.
- Ilyas Bakbergenuly and Elena Kulinskaya. Meta-analysis of binary outcomes via generalized linear mixed models: a simulation study. *BMC Medical Research Methodology*, 18(70), 2018.
- Ilyas Bakbergenuly, David C. Hoaglin, and Elena Kulinskaya. Methods for estimating between-study variance and overall effect in meta-analysis of odds-ratios. *Research Synthesis Methods*, 11:426–442, 2020. doi: 10.1002/jrsm.1404.
- Ji Cheng, Eleanor Pullenayegum, John K Marshall, Alfonso Iorio, and Lehana Thabane. Impact of including or excluding both-armed zero-event studies on using standard meta-analysis methods for rare event outcome: a simulation study. *BMJ Open*, 6(8):e010983, 2016.
- Robert W. Grubbström and Ou Tang. The moments and central moments of a compound distribution. *European Journal of Operational Research*, 170:106–119, 2006.
- Dan Jackson, Martin Law, Theo Stijnen, Wolfgang Viechtbauer, and Ian R White. A comparison of seven random-effects models for meta-analyses that estimate the summary odds ratio. *Statistics in Medicine*, 37:1059–1085, 2018.
- Elena Kulinskaya, David C. Hoaglin, and Ilyas Bakbergenuly. Exploring consequences of simulation design for apparent performance of statistical methods. 1: Results from simulations with constant sample sizes. In *eprint arXiv:2006.16638v1 [stat.ME]*, June 2020.
- O. Kuss. Statistical methods for meta-analyses including information from studies without any events add nothing to nothing and succeed nevertheless. *Statistics in Medicine*, 34(7):1097–1116, 2015.

- Dean Langan, Julian P.T. Higgins, Dan Jackson, Jack Bowden, Areti Angeliki Veroniki, Evangelos Kontopantelis, Wolfgang Viechtbauer, and Mark Simmonds. A comparison of heterogeneity variance estimators in simulated random-effects meta-analyses. *Research Synthesis Methods*, 0(0), 2018. doi: 10.1002/jrsm.1316.
- K. Sidik and J. N. Jonkman. A comparison of heterogeneity variance estimators in combining results of studies. *Statistics in Medicine*, 26(21):1964–1981, 2007.
- Wolfgang Viechtbauer. Confidence intervals for the amount of heterogeneity in meta-analysis. *Statistics in Medicine*, 26(1):37–52, 2007.

Table 1: Components of the simulations for log-odds-ratio

Parameter	Values
K	5, 10, 30
n	40, 100, 250, 1000
θ	0, 0.5, 1, 1.5, 2
τ^2	0(0.1)1
p_C	.1, .4
σ^2	0.1, 0.4
Generation of n	
Normal($n, 1.21n^2/12$) Uniform($n \pm 0.55n$)	
Generation of p_{iC} and p_{iT}	
FIM1 FIM2 RIM1 RIM2 URIM1	Fixed intercept models: $p_{iC} \equiv p_C$ Random intercept models: $\text{logit}(p_{iC}) \sim N(\alpha, \sigma^2)$ $p_{iC} \sim U(p_C - \sigma\sqrt{3}p_C(1 - p_C), p_C + \sigma\sqrt{3}p_C(1 - p_C))$
Estimation targets	Estimators
bias in estimating τ^2	DL, REML, MP, KD, FIM2, RIM2
bias in estimating θ	DL, REML, MP, KD, FIM2, RIM2, SSW
coverage of θ	DL, REML, MP, KD, FIM2, RIM2, SSW (with $\hat{\tau}_{KD}^2$ and t_{K-1} critical values)