

CONVEX HULLS FOR GRAPHS OF QUADRATIC FUNCTIONS WITH UNIT COEFFICIENTS: EVEN WHEELS AND COMPLETE SPLIT GRAPHS

MITCHELL HARRIS^{1,2} AND THOMAS KALINOWSKI²

ABSTRACT. We study the convex hull of the graph of a quadratic function $f(\mathbf{x}) = \sum_{ij \in E} x_i x_j$, where the sum is over the edge set of a graph G with vertex set $\{1, \dots, n\}$. Using an approach proposed by Gupte et al. (Discrete Optimization **36**, 2020, 100569), we investigate minimal extended formulations using additional variables y_{ij} , $1 \leq i < j \leq n$, representing the products $x_i x_j$. The basic idea is to identify a set of facets of the Boolean Quadric Polytope which is sufficient for characterizing the convex hull for the given graph. Our main results are extended formulations for the cases that the underlying graph G is either an even wheel or a complete split graph.

1. INTRODUCTION

In global optimization, the construction of convex envelopes for nonconvex functions is a crucial ingredient in state-of-the-art solvers. As a consequence, there has been a lot of interest in characterizing the convex hulls of graphs of nonlinear functions; see [15] for a comprehensive treatment. The convex hull might be a polytope with a prohibitively large number of facets, and in [12] it has been proposed to use extended formulations to obtain more compact representations of such convex hulls. The basic idea is to represent the polytope as a projection of a higher dimensional polytope with significantly fewer facets, an approach that has been highly successful in combinatorial optimization [9]. In this paper, we focus on functions $f : [0, 1]^n \rightarrow \mathbb{R}$ of the form

$$f(\mathbf{x}) = \sum_{ij \in E} x_i x_j$$

where $G = (V, E)$ is a graph with vertex set $V = [n] := \{1, 2, \dots, n\}$. The convex hull of the graph of f is the set

$$X(f) := \text{conv}\{(\mathbf{x}, z) \in [0, 1]^n \times \mathbb{R} : z = f(\mathbf{x})\}.$$

The set $X(f)$ is a polytope, as it is the convex hull of the 2^n points (\mathbf{x}, z) with $\mathbf{x} \in \{0, 1\}^n$ (see [22, 23]). A natural setting for an extended formulation is to introduce additional variables y_{ij} representing the products $x_i x_j$ of two original variables. For $x_i, x_j \in \{0, 1\}$, the equation $y_{ij} = x_i x_j$ is equivalent to the classical McCormick inequalities [17]: $y_{ij} \geq 0$, $y_{ij} \leq x_i$, $y_{ij} \leq x_j$, and $x_i + x_j - y_{ij} \leq 1$. The *Boolean quadric polytope*

$$BQP_n = \text{conv}\{(\mathbf{x}, \mathbf{y}) \in \{0, 1\}^{n(n+1)/2} : \\ y_{ij} \geq 0, y_{ij} \leq x_i, y_{ij} \leq x_j, x_i + x_j - y_{ij} \leq 1 \text{ for all } 1 \leq i < j \leq n\}$$

was introduced by Padberg [21] in the context of quadratic 0-1-programming, and its structure has been extensively studied [6, 7, 8, 25]. In our setting, BQP_n provides an extended formulation for

¹SCHOOL OF MATHEMATICS AND PHYSICS, UNIVERSITY OF QUEENSLAND, BRISBANE, QLD 4072, AUSTRALIA

²SCHOOL OF SCIENCE AND TECHNOLOGY, UNIVERSITY OF NEW ENGLAND, ARMIDALE, NSW 2351, AUSTRALIA

E-mail addresses: m.g.harris@uq.edu.au, tkalinow@une.edu.au.

2010 *Mathematics Subject Classification.* 90C57, 90C26, 52B12.

Key words and phrases. extended formulation, convex hull, bilinear, quadratic.

$X(f)$ in the sense that $X(f) = \pi[f](BQP_n)$, where the map $\pi[f] : [0, 1]^{n(n+1)/2} \rightarrow [0, 1]^n \times \mathbb{R}$ is defined by

$$\pi[f](\mathbf{x}, \mathbf{y}) = \left(\mathbf{x}, \sum_{ij \in E} y_{ij} \right).$$

As the number of facets of BQP_n is exponential in n , the complete list of facets is known only for small values of n , and some of the known facets are NP-hard to separate [3, 10, 14, 21], the question arises if there exists a relaxation P of BQP_n , obtained by selecting certain families of facets, such that $\pi[f](P) = X(f)$. Let M_n be the relaxation obtained by just keeping the McCormick inequalities:

$$M_n = \left\{ (\mathbf{x}, \mathbf{y}) \in [0, 1]^{n(n+1)/2} : y_{ij} \geq 0, y_{ij} \leq x_i, y_{ij} \leq x_j, x_i + x_j - y_{ij} \leq 1 \text{ for all } 1 \leq i < j \leq n \right\}.$$

Then $\pi[f](M_n) \supseteq X(f)$ with equality if and only if G is bipartite [11, 16], which has been generalized to functions $\sum_{ij} a_{ij} x_i x_j$ with arbitrary coefficients $a_{ij} \in \mathbb{R}$ in [4, 20]. Padberg also studied variants of these polytopes where the variables y_{ij} are only introduced for $ij \in E$:

$$BQP_n(G) = \text{conv}\{(\mathbf{x}, \mathbf{y}) \in \{0, 1\}^{n+|E|} : y_{ij} \geq 0, y_{ij} \leq x_i, y_{ij} \leq x_j, x_i + x_j - y_{ij} \leq 1 \text{ for all } ij \in E\},$$

$$M_n(G) = \{(\mathbf{x}, \mathbf{y}) \in [0, 1]^{n+|E|} : y_{ij} \geq 0, y_{ij} \leq x_i, y_{ij} \leq x_j, x_i + x_j - y_{ij} \leq 1 \text{ for all } ij \in E\},$$

for which we also have $\pi[f](M_n(G)) \supseteq \pi[f](BQP_n(G)) = X(f)$. In general, the McCormick relaxation can be quite weak [4], and in [12] the problem was raised to find relaxations P with $\pi[f](P) = X(f)$. This can be viewed as a weak version of the problem of characterizing the facets of $BQP_n(G)$ for certain classes of graphs G which has been studied extensively [5, 6, 19, 21, 24]. As $\pi[f](P) = X(f)$ is a weaker condition than $P = BQP_n(G)$, we hope to need fewer facets to achieve this. It is easy to check (and has been explicitly proved in [18]) that for every $\mathbf{x} \in [0, 1]^n$,

$$\begin{aligned} \max\{z : (\mathbf{x}, z) \in X(f)\} &= \sum_{ij \in E} \min\{x_i, x_j\} \\ &= \max \left\{ \sum_{ij \in E} y_{ij} : y_{ij} \leq x_i, y_{ij} \leq x_j \text{ for all } 1 \leq i < j \leq n \right\}. \end{aligned}$$

In other words, the McCormick inequalities are sufficient for characterizing the upper boundary of $X(f)$, and we can focus on the lower boundary. For a polytope $P \subseteq [0, 1]^{n+|E|}$ or $P \subseteq [0, 1]^{n(n+1)/2}$, let

$$\text{LB}_P(\mathbf{x}) = \min \left\{ \sum_{ij \in E} y_{ij} : (\mathbf{x}, \mathbf{y}) \in P \right\}.$$

Then $\pi[f](P) = X(f)$ if and only if, for every $\mathbf{x} \in [0, 1]^n$,

$$\{z : (\mathbf{x}, z) \in X(f)\} = \left[\text{LB}_P(\mathbf{x}), \sum_{ij \in E} \min\{x_i, x_j\} \right].$$

Combining this observation with a technique from [26], a characterization of polytopes P with $\pi[f](P) = X(f)$ has been proved in [12], which is based on certain subsets of the half-open unit interval $[0, 1)$. To state this criterion, we define \mathcal{L} to be the collection of unions of finitely many half-open intervals and μ the Lebesgue measure (restricted to \mathcal{L}), that is,

$$\begin{aligned} \mathcal{L} &= \{[a_1, b_1) \cup \dots \cup [a_k, b_k) : 0 \leq a_1 < b_1 < a_2 < b_2 < \dots < a_k < b_k \leq 1, k \in \mathbb{N}\}, \\ \mu(X) &= (b_1 - a_1) + \dots + (b_k - a_k) \quad \text{for } X = [a_1, b_1) \cup \dots \cup [a_k, b_k) \in \mathcal{L}. \end{aligned}$$

Theorem 1 ([12]). *Let $P \subseteq [0, 1]^{n+|E|}$ or $P \subseteq [0, 1]^{n(n+1)/2}$ be a polytope satisfying the following two conditions:*

- (1) *For all $(\mathbf{x}, \mathbf{y}) \in P$ and all $ij \in E$, $y_{ij} \leq \min\{x_i, x_j\}$.*
- (2) *For all $\mathbf{x} \in [0, 1]^n$, there exists $(\mathbf{x}, \mathbf{y}) \in P$ with $y_{ij} = \min\{x_i, x_j\}$ for all $ij \in E$.*

Then $\pi[f](P) = X(f)$ if and only if, for every $\mathbf{x} \in [0, 1]^n$ there exist sets $X_1, \dots, X_n \in \mathcal{L}$ such that $\mu(X_i) = x_i$ for all $i \in [n]$, and $\text{LB}_P(\mathbf{x}) \geq \sum_{ij \in E} \mu(X_i \cap X_j)$.

We illustrate Theorem 1 in the following example.

Example 1. For $n = 3$ and $f(x_1, x_2, x_3) = x_1x_2 + x_2x_3 + x_1x_3$, the graph G is a triangle. Suppose $P = M_3$ is the polytope defined by the McCormick inequalities. Then $(\mathbf{x}, \mathbf{y}) \in P$ where $\mathbf{x} = (1/2, 1/2, 1/2)$ and $\mathbf{y} = \mathbf{0}$, hence $\text{LB}_P(\mathbf{x}) = 0$. By Theorem 1, if $\pi[f](P) = X(f)$, then there are three pairwise disjoint subsets of $[0, 1]$, each of them having measure $1/2$. This is impossible, and we conclude that P is too weak. To obtain a polytope P with $\pi[f](P) = X(f)$ it is sufficient to add the triangle inequality $y_{12} + y_{23} + y_{13} \geq x_1 + x_2 + x_3 - 1$. This can be easily shown using the characterization in Theorem 1: pick $\mathbf{x} \in [0, 1]^3$ and write down corresponding sets X_1, X_2 and X_3 . Without loss of generality, $x_1 \geq x_2 \geq x_3$.

- If $x_1 + x_2 + x_3 \leq 1$, then

$$X_1 = [0, x_1), \quad X_2 = [x_1, x_1 + x_2), \quad X_3 = [x_1 + x_2, x_1 + x_2 + x_3),$$

and for every \mathbf{y} with $(\mathbf{x}, \mathbf{y}) \in P$,

$$\mu(X_1 \cap X_2) + \mu(X_2 \cap X_3) + \mu(X_1 \cap X_3) = 0 \leq y_{12} + y_{23} + y_{13}.$$

- If $x_1 + x_2 \leq 1 < x_1 + x_2 + x_3$ then

$$X_1 = [0, x_1), \quad X_2 = [x_1, x_1 + x_2), \quad X_3 = [x_1 + x_2, 1) \cup [0, x_1 + x_2 + x_3 - 1),$$

and for every \mathbf{y} with $(\mathbf{x}, \mathbf{y}) \in P$,

$$\mu(X_1 \cap X_2) + \mu(X_2 \cap X_3) + \mu(X_1 \cap X_3) = x_1 + x_2 + x_3 - 1 \leq y_{12} + y_{23} + y_{13}$$

due to the triangle inequality.

- If $x_1 + x_2 > 1$ and $x_1 + x_2 + x_3 \leq 2$ then

$$X_1 = [0, x_1), \quad X_2 = [x_1, 1) \cup [0, x_1 + x_2 - 1), \quad X_3 = [x_1 + x_2 - 1, x_1 + x_2 + x_3 - 1),$$

and for every \mathbf{y} with $(\mathbf{x}, \mathbf{y}) \in P$,

$$\mu(X_1 \cap X_2) + \mu(X_2 \cap X_3) + \mu(X_1 \cap X_3) = x_1 + x_2 + x_3 - 1 \leq y_{12} + y_{23} + y_{13}$$

due to the triangle inequality.

- If $x_1 + x_2 + x_3 > 2$ then

$$X_1 = [0, x_1), \quad X_2 = [x_1, 1) \cup [0, x_1 + x_2 - 1), \quad X_3 = [x_1 + x_2 - 1, 1) \cup [0, x_1 + x_2 + x_3 - 2),$$

and for every \mathbf{y} with $(\mathbf{x}, \mathbf{y}) \in P$,

$$\begin{aligned} \mu(X_1 \cap X_2) + \mu(X_2 \cap X_3) + \mu(X_1 \cap X_3) &= 1 + 2(x_1 + x_2 + x_3 - 2) \\ &= (x_1 + x_2 - 1) + (x_2 + x_3 - 1) + (x_1 + x_3 - 1) \leq y_{12} + y_{23} + y_{13} \end{aligned}$$

due to the McCormick inequalities.

One of the classes of facets for BQP_n discussed in [21] is described by the *triangle inequalities*

$$\begin{aligned} x_i + x_j + x_k - y_{ij} - y_{ik} - y_{jk} &\leq 1, \\ -x_i + y_{ij} + y_{ik} - y_{jk} &\leq 0, \\ -x_j + y_{ij} - y_{ik} + y_{jk} &\leq 0, \\ -x_k - y_{ij} + y_{ik} + y_{jk} &\leq 0. \end{aligned}$$

The strength of the relaxation obtained from the McCormick inequalities together with the triangle inequalities has been studied both theoretically [6] and computationally [1, 5]. We focus on the first of the triangle inequalities.

Definition 1. For a graph $G = (V, E)$ let $T(G) \subseteq [0, 1]^{|V|+|E|}$ be the polytope determined by the McCormick inequalities for each $ij \in E$ together with the inequalities $y_{ij} + y_{jk} + y_{ik} \geq x_i + x_j + x_k - 1$ for every triangle ijk in G .

As mentioned above, the McCormick inequalities are sufficient for bipartite graphs, and it would be interesting to characterize the graphs G such that for the corresponding function f , $\pi[f](T(G)) = X(f)$. Our first result provides a first step in this direction by showing that even wheels belong to this class. Let W_{n-1} be the graph with vertex set $V = [n]$ and edge set $E = E_0 \cup E_1$ where

$$E_0 = \{ij : i \in [n-1], j \equiv i+1 \pmod{n-1}\}, \quad E_1 = \{in : i \in [n-1]\}.$$

Theorem 2. Suppose $n-1$ is even, and let $P = T(W_{n-1}) \subseteq [0, 1]^{3n-2}$. Then $\pi[f](P) = X(f)$.

Since the wheel graphs have tree-width 3, it is known that $BQP_n(W_{n-1})$ has an extended formulation with $O(n)$ variables and constraints [13], but it is still interesting to have an explicit construction in terms of the natural variables corresponding to vertices and edges.

Our second result is a generalization of Theorem 1 from [12], which dealt with the case that G is a complete graph with one edge removed. We extend this result to the class of complete split graphs. Let $n = n_1 + n_2$, and let $G = (V, E)$ be the graph with vertex set $V = [n]$, where $V_1 = [n_1]$ is a clique, $V_2 = V \setminus V_1$ is an independent set, and every vertex of V_1 is adjacent to every vertex of V_2 , that is,

$$E = \{ij : 1 \leq i < j \leq n_1\} \cup \{ij : 1 \leq i \leq n_1, n_1 + 1 \leq j \leq n\}.$$

For convenience, we denote by $E^*(W)$ for a vertex set $W \subseteq V$ the set of all pairs of vertices in W (edges and non-edges), that is,

$$E^*(W) = \{ij : i, j \in W, i < j\}.$$

We also use the notation $x(W) = \sum_{i \in W} x_i$ and $y(F) = \sum_{ij \in F} y_{ij}$ for sets $W \subseteq V$ and $F \subseteq E^*(V)$. The clique inequalities

$$y(E^*(W)) \geq \alpha x(W) - \binom{\alpha + 1}{2}$$

were also introduced in [21]. They are valid inequalities for every $W \subseteq V$, $|W| \geq 2$ and every $\alpha \in \{1, 2, \dots, |W| - 1\}$, and facet-defining for $|W| \geq 3$ and $1 \leq \alpha \leq |W| - 2$. For the complete split graphs we need a certain subset of these split inequalities.

Theorem 3. If f corresponds to a complete split graph with clique V_1 and independent set V_2 , then $\pi[f](P) = X(f)$ where P is the polytope described by

$$\begin{aligned} y_{ij} &\geq x_i + x_j - 1 & i \in V_1, j \in V_2, \\ y_{ij} &\leq \min\{x_i, x_j\} & 1 \leq i < j \leq n, \\ y(E^*(V_1 \cup S)) &\geq \alpha x(V_1 \cup S) - \binom{\alpha + 1}{2} & S \subseteq V_2, 0 \leq |S| \leq n_1 - 1, 1 \leq \alpha \leq n_1 - 1. \end{aligned}$$

The proofs of Theorems 2 and 3 are contained in Sections 2 and 3, respectively. In both cases, the sets X_i are constructed greedily: The sets are specified one-by-one, and when fixing X_i we aim to minimize its intersection with the sets X_j , $ij \in E$, which are already fixed. This is a natural strategy, but proving $\text{LB}_P(\mathbf{x}) \geq \sum_{ij \in E} \mu(X_i \cap X_j)$ requires a surprising amount of work. In Section 4, we conclude the paper by stating a couple of open problems.

2. EVEN WHEELS: PROOF OF THEOREM 2

In this section, indices i and j are from the set $[n-1]$, and indices like $i+1$ or $j+1$ are from $[n-1]$ as well, and have to be interpreted modulo $n-1$. Fix an arbitrary $\mathbf{x} \in [0, 1]^n$. In view of Theorem 1, all we need to do is to find sets $X_i \subseteq [0, 1]$ with $\mu(X_i) = x_i$ and

$$\text{LB}_P(\mathbf{x}) \geq \sum_{i=1}^{n-1} (\mu(X_i \cap X_{i+1}) + \mu(X_i \cap X_n)).$$

As a first step we write down a lower bound for $\text{LB}_P(\mathbf{x})$. For a subset $T \subseteq [n-1]$, we define

$$\begin{aligned} \phi(T) = \sum_{i \in [n-1] \setminus T} \max\{0, x_i + x_{i+1} - 1\} + \sum_{i \in [n-1] \setminus (T \cup (T+1))} \max\{0, x_i + x_n - 1\} \\ + \sum_{i \in T} \max\{0, x_i + x_{i+1} + x_n - 1\}, \end{aligned}$$

and we set $\Phi^* = \max \{\phi(T) : T \subseteq [n-1], T \cap (T+1) = \emptyset\}$. Subsets $T \subseteq [n-1]$ with $T \cap (T+1) = \emptyset$ can be identified with certain feasible solutions for the dual to the linear program defining $\text{LB}_P(\mathbf{x})$, such that $\phi(T)$ is the dual objective value. As a consequence, Φ^* is a lower bound for $\text{LB}_P(\mathbf{x})$, and this is the content of the following lemma.

Lemma 1. $\text{LB}_P(\mathbf{x}) \geq \Phi^*$.

Proof. Let $T \subseteq [n-1]$ with $\phi(T) = \Phi^*$ and $T \cap (T+1) = \emptyset$. Using the partition

$$\begin{aligned} E = \{\{i, i+1\}, \{i, n\}, \{i+1, n\} : i \in T\} \cup \{\{i, i+1\} : i \in [n-1] \setminus T\} \\ \cup \{\{i, n\} : i \in [n-1] \setminus (T \cup (T+1))\} \end{aligned}$$

we have that for every \mathbf{y} with $(\mathbf{x}, \mathbf{y}) \in P$,

$$\begin{aligned} \sum_{ij \in E} y_{ij} &= \sum_{i \in T} (y_{i, i+1} + y_{i, n} + y_{i+1, n}) + \sum_{i \in [n-1] \setminus T} y_{i, i+1} + \sum_{i \in [n-1] \setminus (T \cup (T+1))} y_{i, n} \\ &\geq \sum_{i \in T} \max\{0, x_i + x_{i+1} + x_n - 1\} + \sum_{i \in [n-1] \setminus T} \max\{0, x_i + x_{i+1} - 1\} \\ &\quad + \sum_{i \in [n-1] \setminus (T \cup (T+1))} \max\{0, x_i + x_n - 1\} = \phi(T). \quad \square \end{aligned}$$

In the next lemma, we state a property of optimal sets T which will be useful in the subsequent arguments.

Lemma 2. *Let $T \subseteq [n-1]$ with $T \cap (T+1) = \emptyset$ and $\phi(T) = \Phi^*$. Then $x_i + x_{i+1} + x_n \leq 2$ for all $i \in T$.*

Proof. Suppose $i \in T$ with $x_i + x_{i+1} + x_n > 2$. Then $\min\{x_i + x_{i+1}, x_i + x_n, x_{i+1} + x_n\} > 1$, and therefore

$$\phi(T \setminus \{i\}) - \phi(T) = (2x_1 + 2x_2 + 2x_n - 3) - (x_1 + x_2 + x_n - 1) = x_1 + x_2 + x_n - 2 > 0,$$

which contradicts $\phi(T) = \Phi^*$. \square

It will be convenient to assume that T is a subset $\{i \in [n-1] : 1 \leq x_i + x_{i+1} + x_n \leq 2\}$, and that it is a maximal subset subject to $T \cap (T+1) = \emptyset$. The next lemma says that there exists such a T which also maximizes $\phi(T)$.

Lemma 3. *There exists a set $T \subseteq [n-1]$ with $T \cap (T+1) = \emptyset$ and $\phi(T) = \Phi^*$ such that the following two conditions are satisfied:*

- (i) $x_i + x_{i+1} + x_n \geq 1$ for all $i \in T$, and
- (ii) $1 \leq x_i + x_{i+1} + x_n \leq 2 \implies \{i-1, i, i+1\} \cap T^* \neq \emptyset$ for all $i \in [n-1]$.

Proof. If $x_i + x_{i+1} + x_n < 1$ then $\max\{x_i + x_{i+1}, x_i + x_n, x_{i+1} + x_n\} < 1$, and therefore $\phi(T \setminus \{i\}) = \phi(T)$. If $1 \leq x_i + x_{i+1} + x_n \leq 2$ and $\{i-1, i, i+1\} \cap T = \emptyset$, then $T \cup \{i\}$ is a feasible set with $\phi(T \cup \{i\}) \geq \phi(T)$. Therefore we can satisfy the conditions in the lemma by removing the elements $i \in T$ which violate the first condition, and adding the elements $i \in [n-1] \setminus T$ which violate the second condition. \square

From now on, we fix a set $T^* \subseteq [n-1]$ with

- (1) $T^* \cap (T^* + 1) = \emptyset$, and
- (2) $\phi(T^*) = \Phi^*$, and
- (3) $1 \leq x_i + x_{i+1} + x_n \leq 2$ for all $i \in T^*$, and
- (4) $T^* \cap \{i-1, i, i+1\} \neq \emptyset$ for all $i \in [n-1]$ with $1 \leq x_i + x_{i+1} + x_n \leq 2$.

In view of Lemma 1, it is sufficient to find sets $X_i \in \mathcal{L}$ with $\mu(X_i) = x_i$ and

$$\sum_{i=1}^{n-1} (\mu(X_i \cap X_{i+1}) + \mu(X_i \cap X_n)) = \phi(T^*). \quad (1)$$

Lemma 4. *Let $X_i \subseteq [0, 1]$, $i \in [n-1]$, with $\mu(X_i) = x_i$. Then (1) is true if and only if the following three conditions are satisfied:*

- (i) $\mu(X_i \cap X_{i+1}) + \mu(X_i \cap X_n) + \mu(X_{i+1} \cap X_n) = \max\{0, x_i + x_{i+1} + x_n - 1\}$ for every $i \in T^*$,
- (ii) $\mu(X_i \cap X_{i+1}) = \max\{0, x_i + x_{i+1} - 1\}$ for all $i \in [n-1] \setminus T^*$, and
- (iii) $\mu(X_i \cap X_n) = \max\{0, x_i + x_n - 1\}$ for all $i \in [n-1] \setminus (T^* \cup (T^* + 1))$.

Proof. The left-hand side of (1) can be expanded as follows:

$$\begin{aligned} \sum_{i=1}^{n-1} (\mu(X_i \cap X_{i+1}) + \mu(X_i \cap X_n)) &= \sum_{i \in T^*} (\mu(X_i \cap X_{i+1}) + \mu(X_i \cap X_n) + \mu(X_{i+1} \cap X_n)) \\ &\quad + \sum_{i \in [n-1] \setminus T^*} \mu(X_i \cap X_{i+1}) + \sum_{i \in [n-1] \setminus (T^* \cup (T^* + 1))} \mu(X_i \cap X_n). \end{aligned}$$

This implies immediately that the three conditions in the lemma are sufficient. To see that they are also necessary, we observe that by the inclusion-exclusion principle, for every $i \in [n-1]$ and every $j \in \{i+1, n\}$,

$$\mu(X_i \cap X_j) = \mu(X_i) + \mu(X_j) - \mu(X_i \cup X_j) \geq x_i + x_j - 1,$$

and for every $i \in T^*$,

$$\begin{aligned} \mu(X_i \cap X_{i+1}) + \mu(X_i \cap X_n) + \mu(X_{i+1} \cap X_n) \\ = \mu(X_i) + \mu(X_{i+1}) + \mu(X_n) + \mu(X_i \cap X_{i+1} \cap X_n) - \mu(X_i \cup X_{i+1} \cup X_n) \geq x_i + x_{i+1} + x_n - 1. \end{aligned}$$

As a consequence,

- $\mu(X_i \cap X_{i+1}) + \mu(X_i \cap X_n) + \mu(X_{i+1} \cap X_n) \geq \max\{0, x_i + x_{i+1} + x_n - 1\}$ for every $i \in T^*$,
- $\mu(X_i \cap X_{i+1}) \geq \max\{0, x_i + x_{i+1} - 1\}$ for all $i \in [n-1] \setminus T^*$, and

- $\mu(X_i \cap X_n) \geq \max\{0, x_i + x_n - 1\}$ for all $i \in [n-1] \setminus (T^* \cup (T^* + 1))$,

and if any of these inequalities is strict then the left-hand side of (1) is strictly larger than the right-hand side. \square

For the arguments in the proof of the next lemma it is sometimes convenient to use the following equivalent statements for the conditions in Lemma 4:

- (i) $X_i \cup X_{i+1} \cup X_n = [0, 1)$ and $X_i \cap X_{i+1} \cap X_n = \emptyset$ for every $i \in T^*$,
- (ii) $X_i \cap X_{i+1} = \emptyset$ or $X_i \cup X_{i+1} = [0, 1)$ for all $i \in [n-1] \setminus T^*$, and
- (iii) $X_i \cap X_n = \emptyset$ or $X_i \cup X_n = [0, 1)$ for all $i \in [n-1] \setminus (T^* \cup (T^* + 1))$.

In order to state another equivalent condition for the existence of the sets X_i , we introduce the notation

$$\begin{aligned} m_i &= \max\{0, x_i - x_n\}, & M_i &= \min\{x_i, 1 - x_n\}, \\ m'_i &= \min\{1, x_i + x_{i+1}\} - x_n, & M'_i &= \max\{1, x_i + x_{i+1}\} - x_n, \end{aligned}$$

for $i \in [n-1]$.

Lemma 5. *The following two statements are equivalent.*

- (i) *There exist $X_i \subseteq [0, 1]$ with $\mu(X_i) = x_i$ satisfying (1).*
- (ii) *There exists a vector $\mathbf{z} = (z_1, \dots, z_{n-1})$ which satisfies the system*

$$-z_i \leq -m_i \quad i \in T^* \cup (T^* + 1), \quad (2)$$

$$-z_i \leq -M_i \quad i \in [n-1] \setminus (T^* \cup (T^* + 1)), \quad (3)$$

$$z_i \leq M_i \quad i \in [n-1], \quad (4)$$

$$-z_i - z_{i+1} \leq -M'_i \quad i \in T^*, \quad (5)$$

$$-z_i - z_{i+1} \leq -m'_i \quad i \in [n-1] \setminus T^*, \quad (6)$$

$$z_i + z_{i+1} \leq M'_i \quad i \in [n-1] \setminus T^*. \quad (7)$$

Proof.

“(i) \Rightarrow (ii)” : For $i \in [n-1]$, set $z_i = \mu(X_i \setminus X_n)$. Then $z_i \geq \mu(X_i) - \mu(X_n) = x_i - x_n$, and together with $z_i \geq 0$, this implies (2). From $X_i \setminus X_n \subseteq X_i$ and $X_i \setminus X_n \subseteq [0, 1) \setminus X_n$, we obtain $z_i \leq x_i$ and $z_i \leq 1 - x_n$, hence (4). The remaining inequalities are obtained by expressing the conditions from Lemma 4 in terms of the variables z_i in the following way:

- For $i \in T^*$, $X_i \cup X_{i+1} \cup X_n = [0, 1)$ implies $(X_i \setminus X_{n-1}) \cup (X_{i+1} \setminus X_n) \supset [0, 1) \setminus X_n$, hence $z_i + z_{i+1} \geq 1 - x_n$. On the other hand, $X_i \cap X_{i+1} \cap X_n = \emptyset$ implies $\mu(X_i \cap X_n) + \mu(X_{i+1} \cap X_n) \leq \mu(X_n)$, hence $(x_i - z_i) + (x_{i+1} - z_{i+1}) \leq x_n$. Rearranging the latter inequality gives $z_i + z_{i+1} \geq x_i + x_{i+1} - x_n$. Combining these inequalities yields (5).
- For $i \in [n-1] \setminus T^*$ with $x_i + x_{i+1} \geq 1$, $X_i \cup X_{i+1} = [0, 1)$ implies

$$z_i + z_{i+1} = \mu(X_i \setminus X_n) + \mu(X_{i+1} \setminus X_n) \geq \mu((X_i \cup X_{i+1}) \setminus X_n) = \mu([0, 1) \setminus X_n) = 1 - x_n,$$

and

$$(x_i - z_i) + (x_{i+1} - z_{i+1}) = \mu(X_i \cap X_n) + \mu(X_{i+1} \cap X_n) \geq \mu((X_i \cup X_{i+1}) \cap X_n) = \mu(X_n) = x_n.$$

Rearranging the latter inequality gives $z_i + z_{i+1} \leq x_i + x_{i+1} - x_n$, and combining the two inequalities, we obtain $1 - x_n \leq z_i + z_{i+1} \leq x_i + x_{i+1} - x_n$, which is (6) and (7) for $i \in [n-1] \setminus T^*$ with $x_i + x_{i+1} \geq 1$.

- For $i \in [n-1] \setminus T^*$ with $x_i + x_{i+1} \leq 1$, $X_i \cap X_{i+1} = \emptyset$ implies

$$z_i + z_{i+1} = \mu(X_i \setminus X_n) + \mu(X_{i+1} \setminus X_n) \leq \mu([0, 1) \setminus X_n) = 1 - x_n,$$

and

$$(x_i - z_i) + (x_{i-1} - z_{i-1}) = \mu(X_i \cap X_n) + \mu(X_{i+1} \cap X_n) \leq \mu(X_n) = x_n.$$

Rearranging the latter inequality gives $z_i + z_{i+1} \geq x_i + x_{i+1} - x_n$, and combining the two inequalities, we obtain $x_i + x_{i+1} - x_n \leq z_i + z_{i+1} \leq 1 - x_n$, which is (6) and (7) for $i \in [n-1] \setminus T^*$ with $x_i + x_{i+1} \leq 1$.

- For $i \in [n-1] \setminus (T^* \cup (T^* + 1))$, $X_i \cup X_n = [0, 1]$ if $x_i + x_n \geq 1$ and $X_i \cap X_n = \emptyset$ if $x_i + x_n \leq 1$. This implies $z_i = \min\{x_i, 1 - x_n\}$, and in particular, (3).

“(ii) \Rightarrow (i)” : We set $X_n = [0, x_n]$, and then

$$X_i = \begin{cases} [x_n, x_n + z_i) \cup [0, x_i - z_i) & \text{for odd } i \in [n-1], \\ [1 - z_i, 1) \cup [x_n - x_i + z_i, x_n) & \text{for even } i \in [n-1]. \end{cases} \quad \square$$

We complete the proof of Theorem 2 by showing that the system (2)–(7) is feasible. Suppose it isn't. By Farkas' lemma this implies the existence of non-negative numbers π_i^- , π_i^+ and σ_i^- for $i \in [n-1]$, and σ_i^+ for $i \in [n-1] \setminus T^*$, such that

$$\begin{aligned} \pi_i^+ + \sigma_{i-1}^+ &= \pi_i^- + \sigma_i^- + \sigma_{i-1}^- & i \in T^*, \\ \pi_i^+ + \sigma_i^+ &= \pi_i^- + \sigma_i^- + \sigma_{i-1}^- & i \in T^* + 1, \\ \pi_i^+ + \sigma_i^+ + \sigma_{i-1}^+ &= \pi_i^- + \sigma_i^- + \sigma_{i-1}^- & i \in [n-1] \setminus (T^* \cup (T^* + 1)), \end{aligned}$$

and $h(\pi^-, \pi^+, \sigma^-, \sigma^+) < 0$, where

$$\begin{aligned} h(\pi^-, \pi^+, \sigma^-, \sigma^+) &= \sum_{i \in T^*} (-m_i \pi_i^- + M_i \pi_i^+ - M'_i \sigma_i^-) + \sum_{i \in T^* + 1} (-m_i \pi_i^- + M_i \pi_i^+ - m'_i \sigma_i^- + M'_i \sigma_i^+) \\ &\quad + \sum_{i \in [n-1] \setminus (T^* \cup (T^* + 1))} (M_i (\pi_i^+ - \pi_i^-) - m'_i \sigma_i^- + M'_i \sigma_i^+). \end{aligned}$$

This is a negative cost circulation in the network $N(\mathbf{x}, T^*)$ with node set $\{O\} \cup [n-1]$, and the arc set described as follows:

- There are arcs (O, i) and (i, O) for every $i \in [n-1]$. For $i \in T^* \cup (T^* + 1)$, the costs are

$$\text{cost}(O, i) = \begin{cases} M_i & \text{if } i \text{ is odd,} \\ -m_i & \text{if } i \text{ is even,} \end{cases} \quad \text{cost}(i, O) = \begin{cases} -m_i & \text{if } i \text{ is odd,} \\ M_i & \text{if } i \text{ is even,} \end{cases}$$

and for $i \notin T^* \cup (T^* + 1)$,




$$\text{cost}(O, i) = \begin{cases} M_i & \text{if } i \text{ is odd,} \\ -M_i & \text{if } i \text{ is even,} \end{cases} \quad \text{cost}(i, O) = \begin{cases} -M_i & \text{if } i \text{ is odd,} \\ M_i & \text{if } i \text{ is even.} \end{cases}$$

The flows on these arcs correspond to the variables π_i^+ and π_i^- .

- For every $i \in T^*$ there is an arc with cost $-M'_i$ corresponding to the variable σ_i^- . This arc is $(i, i+1)$ for odd i and $(i+1, i)$ for even i .
- For every $i \in [n-1] \setminus T^*$ there are two arcs with costs $-m'_i$ and M'_i , respectively, corresponding to the variables σ_i^- and σ_i^+ . For odd i , the arc with cost $-m'_i$ is $(i, i+1)$ and the arc with cost M'_i is $(i+1, i)$, and for even i it is the other way around.

The construction is illustrated in Figures 1 to 3. Figure 4 shows the complete network for $n = 13$ and $T^* = \{1, 3, 6, 8, 10\}$. We conclude the proof of Theorem 2 by proving the following lemma.


Lemma 6. *The network $N(\mathbf{x}, T^*)$ does not contain a directed cycle of negative cost.*


 FIGURE 1. The arcs incident with a node $i \in T^*$.

 FIGURE 2. The arcs incident with a node $i \in T^* + 1$.

 FIGURE 3. The arcs incident with a node $i \in [n-1] \setminus (T^* \cup (T^* + 1))$.

Proof. Suppose there is a negative cost cycle and let C be a negative cost cycle with the minimum number of arcs. We will argue that we can use C to modify T^* to obtain a set T with $T \cap (T+1) = \emptyset$ and $\phi(T) > \phi(T^*)$, which is the required contradiction. If C is a “backward” cycle, that is, $C = (1, n-1, n-2, \dots, 2, 1)$ or $C = (O, i+1, i, i-1, \dots, j, O)$, then in the network $N((x_2, x_3, \dots, x_{n-1}, x_1, x_n), T^* - 1)$, the cycle $C' = (1, 2, \dots, n-1, 1)$ or $C' = (O, i-1, i, \dots, j)$ is a cycle of negative cost. As a consequence, we can assume that C is a forward cycle, that is, we are in one of the following two cases, where $[i, j]$ denotes the set $\{i, i+1, \dots, j\}$ and the elements of $[n-1]$ are arranged cyclically so that $n-1$ is followed by 1.

Case 1: $C = (1, 2, \dots, n-1)$. Then

$$T^* = \{i \in [n-1] : i \text{ odd and } 1 \leq x_i + x_{i+1} + x_n \leq 2\},$$

FIGURE 4. The flow network for $n = 13$ and $T^* = \{1, 3, 6, 8, 10\}$.

and we can take $T = \{i \in [n-1] : i \text{ even and } x_i + x_{i+1} + x_n \leq 2\}$.

Case 2: $C = (O, i, i+1, \dots, j, j+1, O)$. Then

$$T^* \cap [i, j] = \{k \in [i, j] : k \text{ odd and } 1 \leq x_k + x_{k+1} + x_n \leq 2\},$$

and we can take $T = (T^* \setminus [i, j]) \cup \{k \in [i, j] : k \text{ even and } 1 \leq x_k + x_{k+1} + x_n \leq 2\}$.

The basic idea is that $-\text{cost}(C)$ is a lower bound for $\phi(T) - \phi(T^*)$. The verification is straightforward but rather tedious, and we have decided to provide the details in Appendix A. \square

3. COMPLETE SPLIT GRAPHS: PROOF OF THEOREM 3

In this section, we prove that for a complete split graph with clique V_1 and independent set V_2 , $\pi[f](P) = X(f)$ where P is the polytope described by the following inequalities:

$$y_{ij} \geq x_i + x_j - 1 \quad i \in V_1, j \in V_2, \quad (8)$$

$$y_{ij} \leq \min\{x_i, x_j\} \quad 1 \leq i < j \leq n, \quad (9)$$

$$y(E^*(V_1 \cup S)) \geq \alpha x(V_1 \cup S) - \binom{\alpha+1}{2} \quad S \subseteq V_2, 0 \leq |S| \leq n_1 - 1, 1 \leq \alpha \leq n_1 - 1. \quad (10)$$

We will prove this by providing a construction for sets X_i as required by Theorem 1. To avoid certain case distinctions, we start by showing that we can assume that \mathbf{x} is in the interior of $[0, 1]^n$.

Lemma 7. *Suppose there exists $\mathbf{x} \in [0, 1]^n$ such that whenever $X_1, \dots, X_n \subseteq [0, 1]$ satisfy $\mu(X_i) = x_i$ for all $i \in [n]$ then $\text{LB}_P(\mathbf{x}) < \sum_{ij \in E} \mu(X_i \cap X_j)$. Then there is such a vector \mathbf{x} with $0 < x_i < 1$ for all $i \in [n]$.*

Proof. For $\mathbf{x} \in [0, 1]^n$ set

$$\bar{\mu}(\mathbf{x}) = \min \left\{ \sum_{ij \in E} \mu(X_i \cap X_j) : \mu(X_i) = x_i \text{ for all } i \in V \right\}.$$

Suppose the statement of the lemma is wrong, and let $\mathbf{x} \in [0, 1]^n$ be a vector such that $\text{LB}_P(\mathbf{x}) = \bar{\mu}(\mathbf{x}) - \delta$ with $\delta > 0$. Let \mathbf{x}' be a vector obtained from \mathbf{x} by setting

$$x'_i = \begin{cases} x_i & \text{if } 0 < x_i < 1, \\ \varepsilon & \text{if } x_i = 0, \\ 1 - \varepsilon & \text{if } x_i = 1. \end{cases}$$

By assumption $\bar{\mu}(\mathbf{x}') = \text{LB}_P(\mathbf{x}')$. Let $X'_1, \dots, X'_n \subseteq [0, 1]$ with $\mu(X'_i) = x'_i$ for all $i \in V$ and

$$\sum_{ij \in E} \mu(X'_i \cap X'_j) = \bar{\mu}(\mathbf{x}') = \text{LB}_P(\mathbf{x}').$$

We define sets X_1, \dots, X_n by setting $X_i = X'_i$ if $0 < x_i < 1$, $X_i = \emptyset$ if $x_i = 0$ and $X_i = [0, 1]$ if $x_i = 1$. For $\varepsilon > 0$ sufficiently small,

$$\bar{\mu}(\mathbf{x}) \leq \sum_{ij \in E} \mu(X_i \cap X_j) \leq \sum_{ij \in E} \mu(X'_i \cap X'_j) + \varepsilon n(n-1) = \text{LB}_P(\mathbf{x}') + \varepsilon n(n-1) < \text{LB}_P(\mathbf{x}') + \delta/2.$$

The linear programs defining $\text{LB}_P(\mathbf{x})$ and $\text{LB}_P(\mathbf{x}')$ differ only in the right-hand sides of their constraints, hence by choosing ε sufficiently small, we can assume that $\text{LB}_P(\mathbf{x}') < \text{LB}_P(\mathbf{x}) + \delta/2$. This implies $\bar{\mu}(\mathbf{x}) < \text{LB}_P(\mathbf{x}) + \delta = \bar{\mu}(\mathbf{x})$, which is the required contradiction. \square

Using Lemma 7 we can assume without loss of generality that $1 > x_{n_1+1} \geq x_{n_1+2} \geq \dots \geq x_n > 0$ and $1 > x_1 \geq x_2 \geq \dots \geq x_{n_1} > 0$. We set $X_i = [0, x_i]$ for $i \in V_2$, and construct the sets X_i for $i \in V_1$ greedily. Suppose the sets X_j for $j < i$ have been specified. Then X_i is chosen to minimize

$$\sum_{j=n_1+1}^n \mu(X_i \cap X_j) + \sum_{j=1}^{i-1} \mu(X_i \cap X_j).$$

More precisely, this can be done as described in Algorithm 1.

Example 2. Let $n_1 = 4$, $n_2 = 5$ and $\mathbf{x} = (0.85, 0.8, 0.7, 0.5, 0.8, 0.6, 0.5, 0.3, 0.1)$. The steps of Algorithm 1 are illustrated in Figures 5 to 9. From Figure 9 we see that




FIGURE 5. $X_5 = [0, 0.8)$, $X_6 = [0, 0.6)$, $X_7 = [0, 0.5)$, $X_8 = [0, 0.3)$, $X_9 = [0, 0.1)$.

Algorithm 1 Construction of the sets X_i .

Initialize $(a_0, a_1, \dots, a_{n_2}, a_{n_2+1}) = (1, x_{n_1+1}, x_{n_1+2}, \dots, x_n, 0)$

for $i = 1, 2, \dots, n_1$ **do**

$p \leftarrow \min\{k : a_k + x_i < 1\}$

$X_i \leftarrow [a_{p-1}, 1) \cup [a_p, a_p + a_{p-1} + x_i - 1)$

if $p = 1$ **then**

$a_1 \leftarrow a_1 + x_i$

else

$(a_{p-1}, a_p, \dots, a_{n_2}) \leftarrow (a_p + a_{p-1} + x_i - 1, a_{p+1}, a_{p+2}, \dots, a_{n_2+1})$

return X_1, \dots, X_{n_1}




FIGURE 6. $X_1 = [0.3, 1) \cup [0.1, 0.25)$.




FIGURE 7. $X_2 = [0.25, 1) \cup [0, 0.05)$.





FIGURE 8. $X_3 = [0.5, 1) \cup [0.05, 0.25)$.

$$\sum_{ij \in E} \mu(X_i \cap X_j) = 0.35 \binom{6}{2} + 0.45 \binom{5}{2} + 0.2 \binom{4}{2} - 0.6 - 2 \times 0.5 - 3 \times 0.3 - 4 \times 0.1 = 8.05.$$


 FIGURE 9. $X_4 = [0.6, 1) \cup [0.25, 0.35)$.

In order to see that this is a lower bound for $y(E)$, whenever (\mathbf{x}, \mathbf{y}) satisfies (8), (9) and (10), we start with (10) for $S = \{7, 8, 9\}$ and $t = 3$:

$$y(E^*(V \setminus \{5, 6\})) \geq 3 \times 3.75 - \binom{4}{2} = 5.25.$$

With (9) for the pairs (i, j) with $i, j \in \{7, 8, 9\}$ and $i < j$, we obtain

$$y(E(V \setminus \{5, 6\})) \geq 5.25 - 0.3 - 2 \times 0.1 = 4.75.$$

Finally, we add the edges incident with nodes 5 or 6, using (8) for the lower bound:

$$y(E(V)) \geq 4.75 + 0.65 + 0.6 + 0.5 + 0.3 + 0.45 + 0.4 + 0.3 + 0.1 = 8.05.$$

To prove that the sets X_i constructed by Algorithm 1 have the required properties we will extend the algorithm so that it also produces a set $S \subseteq V_2$ with the following properties:

- (1) $|S| \leq n_1 - 1$.
- (2) $\mu(X_i \cap X_j) = \max\{0, x_i + x_j - 1\}$ for all $i \in V_1, j \in V_2 \setminus S$,
- (3) There exists $\alpha \in \{0, 1, \dots, n_1 - 1\}$ such that for every $t \in [0, 1)$,

$$|\{i \in V_1 \cup S : t \in X_i\}| \in \{\alpha, \alpha + 1\}.$$

Actually, the number α in the third condition is $\alpha = \lfloor x(V_1 \cup S) \rfloor$. Assuming the existence of such a set S , the theorem is proved as follows.

Proof of Theorem 3. Let $\mathbf{y} \in [0, 1]^{n(n-1)/2}$ be such that $(\mathbf{x}, \mathbf{y}) \in P$. We need to check that $y(E) \geq \sum_{i,j \in E} \mu(X_i \cap X_j)$. Let the elements of S be j_1, j_2, \dots, j_k such that $j_1 < j_2 < \dots < j_k$. By the third condition on S , there is a set $Y \subseteq [0, 1)$ with

$$|\{i \in V_1 \cup S : t \in X_i\}| = \begin{cases} \alpha & \text{for } t \in Y, \\ \alpha + 1 & \text{for } t \in [0, 1) \setminus Y. \end{cases}$$

Then $x(V_1 \cup S) = \alpha + \mu([0, 1) \setminus Y) = \alpha + 1 - \mu(Y)$, and

$$\begin{aligned} \sum_{i,j \in V_1 \cup S, i < j} \mu(X_i \cap X_j) &= \mu(Y) \binom{\alpha}{2} + (1 - \mu(Y)) \binom{\alpha + 1}{2} = \binom{\alpha + 1}{2} - \mu(Y) \alpha \\ &= \binom{\alpha + 1}{2} - (\alpha + 1 - x(V_1 \cup S)) \alpha = \alpha x(V_1 \cup S) - \binom{\alpha + 1}{2}. \end{aligned}$$

Together with the second condition on S , we obtain

$$\begin{aligned} \sum_{i,j \in E} \mu(X_i \cap X_j) &= \alpha x(V_1 \cup S) - \binom{\alpha + 1}{2} - x_{j_2} - 2x_{j_3} - \dots - (k-1)x_{j_k} \\ &\quad + \sum_{i \in V_1} \sum_{j \in V_2 \setminus S} \max\{0, x_i + x_j - 1\}. \end{aligned}$$

Now

$$y(E) = y(E^*(V_1 \cup S)) - y(E^*(S)) + \sum_{i \in V_1} \sum_{j \in V_2 \setminus S} y_{ij},$$

and the required inequality follows from

$$\begin{aligned} y(E^*(V_1 \cup S)) &\geq \alpha x(V_1 \cup S) - \binom{\alpha + 1}{2} && \text{by (10),} \\ y(E^*(S)) &\leq x_{j_2} + 2x_{j_3} + \cdots + (k-1)x_{j_k} && \text{by (9), and} \\ \sum_{i \in V_1} \sum_{j \in V_2 \setminus S} y_{ij} &\geq \sum_{i \in V_1} \sum_{j \in V_2 \setminus S} \max\{0, x_i + x_j - 1\} && \text{by (8).} \quad \square \end{aligned}$$

What remains to be done is to construct the set S . Algorithm 2 does not exactly do that, but it returns sets A_p from which we can read off the set S easily. Let k be the number of A_p with

Algorithm 2 Construction of the sets X_i and A_p .

Initialize $(a_0, a_1, \dots, a_{n_2}, a_{n_2+1}) = (1, x_{n_1+1}, x_{n_1+2}, \dots, x_n, 0)$

Initialize $(A_1, \dots, A_{n_2}, A_{n_2+1}) = (\{n_1 + 1\}, \{n_1 + 2\}, \dots, \{n\}, \{n + 1\})$

for $i = 1, 2, \dots, n_1$ **do**

$p \leftarrow \min\{k : a_k + x_i < 1\}$

$X_i \leftarrow [a_{p-1}, 1) \cup [a_p, a_p + a_{p-1} + x_i - 1)$

if $p = 1$ **then**

$a_1 \leftarrow a_1 + x_i$

$A_1 \leftarrow A_1 \cup \{i\}$

else

$a_{p-1} \leftarrow a_p + a_{p-1} + x_i - 1$

$A_{p-1} \leftarrow A_p \cup A_{p-1} \cup \{i\}$

for $q = p, p + 1, \dots, n_2$ **do**

$a_q \leftarrow a_{q+1}$

$A_q \leftarrow A_{q+1}$

return X_1, \dots, X_{n_1} and A_1, \dots, A_{n_2}

$A_p \cap V_2 \neq \emptyset$ when the algorithm terminates, that is, $0 < a_k \leq a_{k-1} \leq \cdots \leq a_1 < 1$. Set $j_p = \max A_p$ for all $p \in [k]$, and let $j^* = \min(A_{p_0} \cap V_2)$ where $p_0 = \min\{p : |A_p| > 1\}$ is the smallest index such that A_p has been updated by the algorithm. Note that $j_p \in V_2$ for $p < k$, and $j_k \in \{n, n + 1\}$. An important observation for our arguments below is that x_{j_p} lies between a_{p+1} and a_p , and x_{j^*} between a_{p_0} and $x_{j_{p_0-1}} = a_{p_0-1}$, that is, setting $x_{n+1} = 0$ for convenience, we have

$$\begin{aligned} 0 = a_{k+1} &\leq x_{j_k} \leq a_k \leq x_{j_{k-1}} \leq a_{k-1} \leq \cdots \leq a_{p_0+1} \leq x_{j_{p_0}} \leq a_{p_0} \leq x_{j^*} \leq x_{j_{p_0-1}} = a_{p_0-1} \\ &\leq x_{j_{p_0-2}} = a_{p_0-2} \leq \cdots \leq x_{j_2} = a_2 \leq x_{j_1} = a_1 < 1. \end{aligned}$$

We are now prepared to specify the set S :

$$S = \begin{cases} V_2 \setminus \{j_1, j_2, \dots, j_k, j^*\} & \text{if } p_0 \geq 2 \text{ or } |A_1 \cap V_2| = |A_1 \cap V_1| + 1, \\ V_2 \setminus \{j_1, j_2, \dots, j_k\} & \text{if } p_0 = 1 \text{ and } |A_1 \cap V_2| \leq |A_1 \cap V_1|. \end{cases}$$

Before proving that the set S has the required properties we illustrate the construction in an example.

Example 3. Let $n_1 = 8$, $n_2 = 13$, and

$$\mathbf{x} = (0.9, 0.85, 0.53, 0.49, 0.44, 0.23, 0.16, 0.1, \\ 0.96, 0.89, 0.82, 0.75, 0.67, 0.59, 0.55, 0.45, 0.38, 0.29, 0.22, 0.15, 0.07)$$

The algorithm proceeds as follows.

Step 0:

$$(a_0, \dots, a_{14}) = (1, 0.96, 0.89, 0.82, 0.75, 0.67, 0.59, 0.55, 0.45, 0.38, 0.29, 0.22, 0.15, 0.07, 0), \\ (A_1, \dots, A_{14}) = (\{9\}, \{10\}, \{11\}, \{12\}, \{13\}, \{14\}, \{15\}, \{16\}, \{17\}, \{18\}, \{19\}, \{20\}, \{21\}, \{22\}).$$

Step 1:

$$X_1 = [0.07, 0.12) \cup [0.15, 1), \\ (a_0, \dots, a_{13}) = (1, 0.96, 0.89, 0.82, 0.75, 0.67, 0.59, 0.55, 0.45, 0.38, 0.29, 0.22, 0.12, 0), \\ (A_1, \dots, A_{13}) = (\{9\}, \{10\}, \{11\}, \{12\}, \{13\}, \{14\}, \{15\}, \{16\}, \{17\}, \{18\}, \{19\}, \{1, 20, 21\}, \{22\}).$$

Step 2:

$$X_2 = [0.12, 0.19) \cup [0.22, 1), \\ (a_0, \dots, a_{12}) = (1, 0.96, 0.89, 0.82, 0.75, 0.67, 0.59, 0.55, 0.45, 0.38, 0.29, 0.19, 0), \\ (A_1, \dots, A_{12}) = (\{9\}, \{10\}, \{11\}, \{12\}, \{13\}, \{14\}, \{15\}, \{16\}, \{17\}, \{18\}, \{1, 2, 19, 20, 21\}, \{22\}).$$

Step 3:

$$X_3 = [0.45, 0.53) \cup [0.55, 1), \\ (a_0, \dots, a_{11}) = (1, 0.96, 0.89, 0.82, 0.75, 0.67, 0.59, 0.53, 0.38, 0.29, 0.19, 0), \\ (A_1, \dots, A_{11}) = (\{9\}, \{10\}, \{11\}, \{12\}, \{13\}, \{14\}, \{3, 15, 16\}, \{17\}, \{18\}, \{1, 2, 19, 20, 21\}, \{22\}).$$

Step 4:

$$X_4 = [0.38, 0.4) \cup [0.53, 1), \\ (a_0, \dots, a_{10}) = (1, 0.96, 0.89, 0.82, 0.75, 0.67, 0.59, 0.4, 0.29, 0.19, 0), \\ (A_1, \dots, A_{10}) = (\{9\}, \{10\}, \{11\}, \{12\}, \{13\}, \{14\}, \{3, 4, 15, 16, 17\}, \{18\}, \{1, 2, 19, 20, 21\}, \{22\}).$$

Step 5:

$$X_5 = [0.4, 0.43) \cup [0.59, 1), \\ (a_0, \dots, a_9) = (1, 0.96, 0.89, 0.82, 0.75, 0.67, 0.43, 0.29, 0.19, 0), \\ (A_1, \dots, A_9) = (\{9\}, \{10\}, \{11\}, \{12\}, \{13\}, \{3, 4, 5, 14, 15, 16, 17\}, \{18\}, \{1, 2, 19, 20, 21\}, \{22\}).$$

Step 6:

$$X_6 = [0.75, 0.8) \cup [0.82, 1), \\ (a_0, \dots, a_8) = (1, 0.96, 0.89, 0.8, 0.67, 0.43, 0.29, 0.19, 0), \\ (A_1, \dots, A_8) = (\{9\}, \{10\}, \{6, 11, 12\}, \{13\}, \{3, 4, 5, 14, 15, 16, 17\}, \{18\}, \{1, 2, 19, 20, 21\}, \{22\}).$$

Step 7:

$$X_7 = [0.8, 0.85) \cup [0.89, 1),$$

$$(a_0, \dots, a_7) = (1, 0.96, 0.85, 0.67, 0.43, 0.29, 0.19, 0),$$

$$(A_1, \dots, A_7) = (\{9\}, \{6, 7, 10, 11, 12\}, \{13\}, \{3, 4, 5, 14, 15, 16, 17\}, \{18\}, \{1, 2, 19, 20, 21\}, \{22\}).$$

Step 8:

$$X_8 = [0.85, 0.91) \cup [0.96, 1),$$

$$(a_0, \dots, a_6) = (1, 0.91, 0.67, 0.43, 0.29, 0.19, 0),$$

$$(A_1, \dots, A_6) = (\{6, 7, 8, 9, 10, 11, 12\}, \{13\}, \{3, 4, 5, 14, 15, 16, 17\}, \{18\}, \{1, 2, 19, 20, 21\}, \{22\}).$$

As $p_0 = 1$, $|A_1 \cap V_2| = 4 = |A_1 \cap V_2| + 1$, we obtain $S = \{10, 11, 14, 15, 16, 19, 20\}$. The outcome is illustrated in Figure 10.




FIGURE 10. The outcome of Algorithm 2 for the vector \mathbf{x} in Example 3.

The next three lemmas establish the three required properties of the set S .

Lemma 8. $|S| \leq n_1 - 1$.

Proof. The number of sets A_p with $A_p \cap V_2 \neq \emptyset$ is initially equal to n_2 and in each iteration it drops by 1, except when $a_1 + x_i < 1$ or $a_k + x_i \geq 1$. Therefore, $k \geq n_2 - n_1$, and we have the following cases:

Case 1: $k = n_2 - n_1$. Then $p_0 \geq 2$ or $|A_1 \cap V_2| = |A_1 \cap V_1| + 1$, hence $|S| = n_2 - (k + 1) = n_1 - 1$.


Case 2: $k = n_2 - n_1 + 1$. Then $j_k = n$ or $p_0 \geq 2$ or $|A_1 \cap V_2| = |A_1 \cap V_1| + 1$, hence $|S| \leq n_2 - k = n_1 - 1$.

Case 3: $k \geq n_2 - n_1 + 2$. Then $|S| \leq n_2 - (k - 1) \leq n_1 - 1$. \square

Lemma 9. $\mu(X_i \cap X_j) = \max\{0, x_i + x_j - 1\}$ for all $i \in V_1, j \in V_2 \setminus S$.

Proof. Fix $j \in V_2 \setminus S, i \in V_1$. Then $j = j_p$ for some p or $j = j^*$, and $i \in A_q$ for some q . First assume $j = j_p$. If $q \leq p$ then $X_i \subseteq [x_j, 1)$, hence $X_i \cap X_j = \emptyset$, and if $q > p$, then $X_i \supseteq [x_j, 1)$, hence $\mu(X_i \cap X_j) = x_i + x_j - 1$. Now assume $j = j^*$. If $q < p_0$ then $X_i \subseteq [x_j, 1)$, hence $X_i \cap X_j = \emptyset$, and if $q \geq p_0$, then $X_i \supseteq [x_j, 1)$, hence $\mu(X_i \cap X_j) = x_i + x_j - 1$. \square

Define a function $h : [0, 1) \rightarrow \mathbb{Z}$ by $h(t) = |\{i \in V_1 \cup S : t \in X_i\}|$. Our third condition on S says that the function h takes only two consecutive values.


 FIGURE 11. The function h for Example 3.

Example 4. The graph of the function h for the instance in Example 3 is shown in Figure 11.

Lemma 10. *There exists an integer $\alpha \in \{0, 1, \dots, n_1 - 1\}$ such that $h(t) \in \{\alpha, \alpha + 1\}$ for all $t \in [0, 1)$.*

Proof. We start with the function h_0 defined by $h_0(t) = |\{i \in V_1 \cup V_2 : t \in X_i\}|$. Denoting the value of h_0 on the interval $[a_1, 1)$ by γ , we have $h_0(t) = \gamma + p - 1$ for $t \in [a_p, a_{p-1})$, $p = 1, 2, \dots, k + 1$.

Case 1: $S = V_2 \setminus \{j_1, j_2, \dots, j_k, j^*\}$. Then

$$h(t) = \begin{cases} h_0(t) - (p - 1) & \text{for } t \in [a_p, a_{p-1}), p = 1, 2, \dots, p_0 - 1, \\ h_0(t) - (p_0 - 1) & \text{for } t \in [x_{j^*}, a_{p_0-1}), \\ h_0(t) - p_0 & \text{for } t \in [a_{p_0}, x_{j^*}), \\ h_0(t) - (p - 1) & \text{for } t \in [x_{j_{p-1}}, a_{p-1}), p = p_0 + 1, \dots, k, \\ h_0(t) - p & \text{for } t \in [a_p, x_{j_{p-1}}), p = p_0 + 1, \dots, k + 1. \end{cases}$$

Substituting $h_0(t) = \gamma + p - 1$ for $t \in [a_p, a_{p-1})$, $p = 1, 2, \dots, k + 1$, we obtain

$$h(t) = \begin{cases} \gamma & \text{for } t \in [x_{j^*}, 1) \\ \gamma - 1 & \text{for } t \in [a_{p_0}, x_{j^*}), \\ \gamma & \text{for } t \in [x_{j_p}, a_p), p = p_0, \dots, k, \\ \gamma - 1 & \text{for } t \in [a_{p+1}, x_{j_p}), p = p_0, \dots, k, \end{cases}$$

and this concludes the proof with $\alpha = \gamma - 1$.

Case 2: $S = V_2 \setminus \{j_1, j_2, \dots, j_k\}$. Then $p_0 = 1$ and $h(t) = h_0(t) - (p - 1)$ for $t \in [x_{j_p}, x_{j_{p-1}})$, $p = 1, 2, \dots, k + 1$ (with $x_{j_{k+1}} = 0$ and $x_{j_0} = 1$). Substituting $h_0(t) = \gamma + p - 1$ for $t \in [a_p, a_{p-1})$, $p = 1, 2, \dots, k + 1$, we obtain

$$h(t) = \begin{cases} \gamma & \text{for } t \in [a_p, x_{j_{p-1}}), p = 1, \dots, k, \\ \gamma + 1 & \text{for } t \in [x_{j_p}, a_p), p = 1, \dots, k + 1, \end{cases}$$

and this concludes the proof with $\alpha = \gamma$. \square

4. CONCLUSION

In this paper, we have extended the results from [12] to derive extended formulations for the graphs of quadratic functions corresponding to even wheel graphs and complete split graphs. It is a natural question what happens for odd wheels. For odd wheels, the triangle inequalities

are not sufficient. In fact, even adding in all the facets from [21] is not enough. In Figure 12 this is illustrated for the 5-wheel. For the two points shown in the picture we have $\pi[f](\mathbf{x}, \mathbf{y}) = (1/3, 1/3, 1/3, 1/3, 1/3, 2/3, 5/6)$ and $\pi[f](\mathbf{x}, \mathbf{y}) = (2/3, 2/3, 2/3, 2/3, 2/3, 1/3, 5/2)$, respectively, hence $\pi[f](P) \neq X(f)$, because $(\mathbf{x}, \mathbf{y}) \in BQP_6(W_5)$ implies

$$y(E) \geq 2x_6 + x_1 + x_2 + x_3 + x_4 + x_5 - 2, \quad (11)$$

$$y(E) \geq 3x_6 + 2(x_1 + x_2 + x_3 + x_4 + x_5) - 5. \quad (12)$$




FIGURE 12. For the 5-wheel the relaxation of BQP_6 obtained by using all the facets from [21] is not sufficient to describe $X(f)$. The vector (\mathbf{x}, \mathbf{y}) corresponding to the left picture satisfies all these inequalities and (12), but not (11), while vector corresponding to the right picture satisfies the inequalities from [21] and (11), but not (12).

For the 5-wheel it can be checked using polymake [2], that $\pi[f](P) = X(f)$ where P is the polytope described by the McCormick inequalities, the inequalities $y_{ij} + y_{jk} + y_{ik} \geq x_i + x_j + x_k - 1$ for all triangles in W_5 and the inequalities (11) and (12). This corrects the observation about the 5-wheel in [12], where the triangle inequalities were missing.

Problem 1. Let $n - 1$ be odd, let $f : [0, 1]^n \rightarrow \mathbb{R}$ be the function corresponding to the wheel graph $G = W_{n-1}$, and let $P \subseteq [0, 1]^{3n-2}$ be the polytope described by the McCormick inequalities, the inequalities $y_{ij} + y_{jk} + y_{ik} \geq x_i + x_j + x_k - 1$ for all triangles in G , and the two inequalities

$$y(E) \geq \frac{n-2}{2}x_n + \sum_{i=1}^{n-1} x_i - \frac{n-2}{2}, \quad y(E) \geq \frac{n}{2}x_n + 2 \sum_{i=1}^{n-1} x_i - (n-1).$$

Is it true that $X(f) = \pi[f](P)$?

As mentioned in the introduction, it would be interesting to characterize the graphs G for which the polytope $T(G)$ (see Definition 1) is sufficient.

Problem 2. Characterise the graphs G such that $\pi[f](T(G)) = X(f)$.

As a first step, one can try to generalise Theorem 2 in various directions.

Problem 3. Let $n = n_1 + n_2$ with n_1 even, and let G be the graph obtained by joining every vertex of a cycle of length n_1 to all vertices of an independent set of size n_2 . Is it true that $\pi[f](T(G)) = X(f)$?

Problem 4. Let G be a graph on n vertices, where the graph induced on $\{1, \dots, n-1\}$ is bipartite, and vertex n is adjacent to all vertices in $\{1, \dots, n-1\}$. Is it true that $\pi[f](T(G)) = X(f)$?

Theorem 2 is the special case for Problem 3 with $n_2 = 1$, and for Problem 4 with the induced bipartite graph being a single even cycle. Another class of graphs for which it seems plausible that $\pi[f](T(G)) = X(f)$ are the 2-trees, that is, the graphs obtained by starting with a triangle, and repeatedly adding a vertex and joining to two adjacent vertices (maximal series-parallel graphs).

Problem 5. Is it true that $\pi[f](T(G)) = X(f)$ whenever G is a 2-tree?

REFERENCES

- [1] Kurt M. Anstreicher. “On convex relaxations for quadratically constrained quadratic programming”. In: *Mathematical Programming* 136.2 (2012), pp. 233–251.
- [2] Benjamin Assarf, Ewgenij Gawrilow, Katrin Herr, Michael Joswig, Benjamin Lorenz, Andreas Paffenholz, and Thomas Rehn. “Computing convex hulls and counting integer points with polymake”. In: *Mathematical Programming Computation* 9.1 (2017), pp. 1–38.
- [3] Francisco Barahona and Ali Ridha Mahjoub. “On the cut polytope”. In: *Mathematical Programming* 36.2 (1986), pp. 157–173.
- [4] Natashia Boland, Santanu S. Dey, Thomas Kalinowski, Marco Molinaro, and Fabian Rigterink. “Bounding the gap between the McCormick relaxation and the convex hull for bilinear functions”. In: *Mathematical Programming* 162 (2017), pp. 523–535.
- [5] Pierre Bonami, Oktay Günlük, and Jeff Linderoth. “Globally solving nonconvex quadratic programming problems with box constraints via integer programming methods”. In: *Mathematical Programming Computation* 10.3 (2018), pp. 333–382.
- [6] Endre Boros, Yves Crama, and Peter L. Hammer. “Chvátal cuts and odd cycle inequalities in quadratic 0–1 optimization”. In: *SIAM Journal on Discrete Mathematics* 5.2 (1992), pp. 163–177.
- [7] Endre Boros and Peter L. Hammer. “Cut-polytopes, Boolean quadric polytopes and nonnegative quadratic pseudo-Boolean functions”. In: *Mathematics of Operations Research* 18.1 (1993), pp. 245–253.
- [8] Samuel Burer and Adam N. Letchford. “On nonconvex quadratic programming with box constraints”. In: *SIAM Journal on Optimization* 20.2 (2009), pp. 1073–1089.
- [9] Michele Conforti, Gérard Cornuéjols, and Giacomo Zambelli. “Extended formulations in combinatorial optimization”. In: *4OR* 8.1 (2010), pp. 1–48.
- [10] Michel Marie Deza and Monique Laurent. *Geometry of cuts and metrics*. Vol. 15. Algorithms and Combinatorics. Springer, 1997.
- [11] Oktay Günlük, Jon Lee, and Janny Leung. “A polytope for a product of real linear functions in 0/1 variables”. In: *Mixed Integer Nonlinear Programming*. Ed. by Jon Lee and Sven Leyffer. Vol. 154. The IMA Volumes in Mathematics and its Applications. Springer New York, 2012, pp. 513–529.
- [12] Akshay Gupte, Thomas Kalinowski, Fabian Rigterink, and Hamish Waterer. “Extended formulations for convex hulls of some bilinear functions”. In: *Discrete Optimization* 36 (2020), p. 100569.
- [13] Monique Laurent. “Sums of squares, moment matrices and optimization over polynomials”. In: *Emerging Applications of Algebraic Geometry*. Springer New York, 2009, pp. 157–270.

- [14] Adam N. Letchford and Michael M. Sørensen. “A new separation algorithm for the Boolean quadric and cut polytopes”. In: *Discrete Optimization* 14 (2014), pp. 61–71.
- [15] Marco Locatelli and Fabio Schoen. *Global optimization: theory, algorithms, and applications*. SIAM, 2013.
- [16] James Luedtke, Mahdi Namazifar, and Jeffrey T. Linderoth. “Some results on the strength of relaxations of multilinear functions”. In: *Mathematical Programming* 136.2 (2012), pp. 325–351.
- [17] Garth P. McCormick. “Computability of global solutions to factorable nonconvex programs: Part I – Convex underestimating problems”. In: *Mathematical Programming* 10.1 (1976), pp. 147–175.
- [18] Clifford A. Meyer and Christodoulos A. Floudas. “Convex envelopes for edge-concave functions”. In: *Mathematical Programming* 103.2 (2005), pp. 207–224.
- [19] Carla Michini. “Forbidden minors for tight cycle relaxations”. Optimization Online Preprint 5483. optimization-online:5483. 2018.
- [20] Ruth Misener, James B. Smadbeck, and Christodoulos A. Floudas. “Dynamically generated cutting planes for mixed-integer quadratically constrained quadratic programs and their incorporation into GloMIQO 2”. In: *Optimization Methods and Software* 30.1 (2015), pp. 215–249.
- [21] Manfred W. Padberg. “The boolean quadric polytope: Some characteristics, facets and relatives”. In: *Mathematical Programming* 45.1–3 (1989), pp. 139–172.
- [22] Anatoliy D. Rikun. “A Convex Envelope Formula for Multilinear Functions”. In: *Journal of Global Optimization* 10.4 (1997), pp. 425–437.
- [23] Hanif D. Sherali. “Convex envelopes of multilinear functions over a unit hypercube and over special discrete sets”. In: *Acta Mathematica Vietnamica* 22.1 (1997), pp. 245–270.
- [24] Caterina De Simone. “The cut polytope and the Boolean quadric polytope”. In: *Discrete Mathematics* 79.1 (1990), pp. 71–75.
- [25] Yasutoshi Yajima and Tetsuya Fujie. “A polyhedral approach for nonconvex quadratic programming problems with box constraints”. In: *Journal of Global Optimization* 13.2 (1998), pp. 151–170.
- [26] Mark Zuckerberg. “Geometric proofs for convex hull defining formulations”. In: *Operations Research Letters* 44.5 (2016), pp. 625–629.

APPENDIX A. PROOF OF LEMMA 6

In this appendix we provide the details for the proof of Lemma 6. Recall that we have fixed a set $T^* \subseteq [n - 1]$ satisfying the following conditions:

- (1) $T^* \cap (T^* + 1) = \emptyset$, and
- (2) $\phi(T^*) = \Phi^*$, and
- (3) $1 \leq x_i + x_{i+1} + x_n \leq 2$ for all $i \in T$, and
- (4) $T^* \cap \{i - 1, i, i + 1\} \neq \emptyset$ for all $i \in [n - 1]$ with $1 \leq x_i + x_{i+1} + x_n \leq 2$.

We have then defined a network $N = N(\mathbf{x}, T^*)$, and we have to verify that the network does not contain negative cost cycles. Let C be a negative cost cycle with the minimum number of arcs. We discuss the cases indicated in the proof outline separately.

A.1. Case 1. $C = (1, 2, \dots, n - 1)$. By the definition of the arc set of N , all elements of T^* are odd because the arc $(i, i + 1)$ does not exist for even $i \in T^*$. The last two properties of T^* listed above imply

$$T^* = \{i \in [n - 1] : i \text{ is odd and } 1 \leq x_i + x_{i+1} + x_n \leq 2\}.$$

We set $T = \{i \in [n-1] : i \text{ is even and } 1 \leq x_i + x_{i+1} + x_n \leq 2\}$, and want to argue that $\phi(T) > \phi(T^*)$. We start by writing down the cost of C in a convenient form:

$$\begin{aligned} \text{cost}(C) = & - \sum_{i \in T^*} M'_i - \sum_{i \in [n-1] \setminus T^* \text{ odd}} m'_i + \sum_{i \in [n-1] \text{ even}} M'_i \\ & \sum_{i \in T^*} (x_n - \max\{1, x_i + x_{i+1}\}) + \sum_{\text{odd } i \in [n-1] \setminus T^*} (x_n - \min\{1, x_i + x_{i+1}\}) + \sum_{\text{even } i \in [n-1]} (\max\{1, x_i + x_{i+1}\} - x_n). \end{aligned}$$

We observe that the x_n cancel, and we split the second sum into the sum over $i \in S_0$ and $i \in S_2$, where

$$S_0 = \{i \in [n-1] : i \text{ odd and } x_i + x_{i+1} + x_n < 1\}, \quad S_2 = \{i \in [n-1] : i \text{ odd and } x_i + x_{i+1} + x_n > 2\}.$$

In addition, we use

$$\min\{1, x_i + x_{i+1}\} = \max\{1, x_i + x_{i+1}\} - \begin{cases} 1 - x_i - x_{i+1} & \text{for } i \in S_0, \\ x_i + x_{i+1} - 1 & \text{for } i \in S_2. \end{cases}$$

Putting this together, we obtain

$$\begin{aligned} \text{cost}(C) = & \sum_{\text{even } i \in [n-1]} \max\{1, x_i + x_{i+1}\} - \sum_{\text{odd } i \in [n-1]} \max\{1, x_i + x_{i+1}\} \\ & + \sum_{i \in S_0} (1 - x_i - x_{i+1}) + \sum_{i \in S_2} (x_i + x_{i+1} - 1). \quad (13) \end{aligned}$$

Now let's write down $\phi(T^*)$ and $\phi(T)$, and then $\phi(T) - \phi(T^*)$. We are done once we are convinced that $\phi(T) - \phi(T^*) > 0$. The contributions to $\phi(T^*)$ are as follows:

- An $i \in T^*$ contributes $x_i + x_{i+1} + x_n - 1$.
- An $i \in S_0$ contributes $0 = x_i + x_{i+1} + x_n - 1 + (1 - x_i - x_{i+1} - x_n)$.
- An $i \in S_2$ contributes $2x_i + 2x_{i+1} + 2x_n - 3 = x_i + x_{i+1} + x_n - 1 + (x_i + x_{i+1} + x_n - 2)$.
- An even $i \in [n-1]$ contributes $\max\{0, x_i + x_i - 1\} = \max\{1, x_i + x_{i+1}\} - 1$.

We obtain

$$\begin{aligned} \phi(T^*) = & \sum_{\text{odd } i \in [n-1]} (x_i + x_{i+1} + x_n - 1) + \sum_{\text{even } i \in [n-1]} (\max\{1, x_i + x_{i+1}\} - 1) \\ & + \sum_{i \in S_0} (1 - x_i - x_{i+1} - x_n) + \sum_{i \in S_2} (x_i + x_{i+1} + x_n - 2), \end{aligned}$$

which simplifies to

$$\begin{aligned} \phi(T^*) = & \sum_{i \in [n-1]} x_i + \frac{n-1}{2}(x_n - 2) + \sum_{\text{even } i \in [n-1]} \max\{1, x_i + x_{i+1}\} \\ & + \sum_{i \in S_0} (1 - x_i - x_{i+1} - x_n) + \sum_{i \in S_2} (x_i + x_{i+1} + x_n - 2). \end{aligned}$$

Similarly,

$$\begin{aligned} \phi(T) = & \sum_{i \in [n-1]} x_i + \frac{n-1}{2}(x_n - 2) + \sum_{\text{odd } i \in [n-1]} \max\{1, x_i + x_{i+1}\} \\ & + \sum_{i \in S'_0} (1 - x_i - x_{i+1} - x_n) + \sum_{i \in S'_2} (x_i + x_{i+1} + x_n - 2), \end{aligned}$$

where

$$S'_0 = \{i \in [n-1] : i \text{ even and } x_i + x_{i+1} + x_n < 1\}, \quad S_2 = \{i \in [n-1] : i \text{ even and } x_i + x_{i+1} + x_n > 2\}.$$

As a consequence,

$$\begin{aligned} \phi(T) - \phi(T^*) &= \sum_{\text{odd } i \in [n-1]} \max\{1, x_i + x_{i+1}\} - \sum_{\text{even } i \in [n-1]} \max\{1, x_i + x_{i+1}\} \\ &+ \sum_{i \in S'_0} (1 - x_i - x_{i+1} - x_n) + \sum_{i \in S'_2} (x_i + x_{i+1} + x_n - 2) - \sum_{i \in S_0} (1 - x_i - x_{i+1} - x_n) - \sum_{i \in S_2} (x_i + x_{i+1} + x_n - 2). \end{aligned}$$

Using (13) this can be rewritten as

$$\phi(T) - \phi(T^*) = -\text{cost}(C) + \sum_{i \in S'_0} (1 - x_i - x_{i+1} - x_n) + \sum_{i \in S'_2} (x_i + x_{i+1} + x_n - 2) + |S_0|x_n + |S_2|(1 - x_n).$$

This implies the required inequality $\phi(T) - \phi(T^*) > 0$ because $-\text{cost}(C) > 0$ and the remaining terms on the right hand side are non-negative.

A.2. Case 2. $C = (O, i, i+1, \dots, j, j+1, O)$. The assumption that C is a shortest cycle of negative cost leads to some restrictions on i and j as it implies $\text{cost}(O, i) + \text{cost}(i, i+1) < \text{cost}(O, i+1)$ and $\text{cost}(j, j+1) + \text{cost}(j+1, O) < \text{cost}(j, O)$.

Lemma 11. $\{i, i+1\} \cap T^* \neq \emptyset$.

Proof. Suppose $\{i, i+1\} \cap T^* = \emptyset$. We distinguish the three cases shown in Figure 13.




FIGURE 13. The three cases in the proof of Lemma 11.

Case (a): i is odd. We need to show that

$$\min\{x_i, 1 - x_n\} - \min\{1, x_i + x_{i+1}\} + x_n \geq -\min\{x_{i+1}, 1 - x_n\} = \max\{-x_{i+1}, x_n - 1\}.$$

- If $x_i + x_n \leq 1$ then the left hand side is $x_i - \min\{1, x_i + x_{i+1}\} + x_n \geq x_n - x_{i+1}$.
- If $x_i + x_n > 1$ then the left hand side is $1 - \min\{1, x_i + x_{i+1}\} \geq 0$.

Case (b): i is even and $i \in T^* + 1$. We need to show that

$$-\max\{0, x_i - x_n\} + \max\{1, x_i + x_{i+1}\} - x_n \geq \min\{x_{i+1}, 1 - x_n\}.$$

- If $x_i \leq x_n$ then the left hand side is $\max\{1, x_i + x_{i+1}\} - x_n \geq 1 - x_n$.
- If $x_i > x_n$ then the left hand side is $\max\{1, x_i + x_{i+1}\} - x_i \geq x_{i+1}$.

Case (c): i is even and $i \notin T^* + 1$. We need to show that

$$-\min\{x_i, 1 - x_n\} + \max\{1, x_i + x_{i+1}\} - x_n \geq \min\{x_{i+1}, 1 - x_n\}.$$

From $T^* \cap \{i-1, i, i+1\} = \emptyset$ it follows that $x_i + x_{i+1} + x_n < 1$ or $x_i + x_{i+1} + x_n > 2$.

- If $x_i + x_{i+1} + x_n < 1$ then the left hand side is $1 - x_i - x_n \geq x_{i+1}$.
- If $x_i + x_{i+1} + x_n > 2$ then the left hand side is $x_i + x_{i+1} - 1 \geq 1 - x_n$. □

Lemma 12.

- (1) If $i \in T^*$ then i is odd and either $x_i + \max\{x_{i+1}, x_n\} < 1$ or $x_i + \min\{x_{i+1}, x_n\} > 1$.
(2) If $i + 1 \in T^*$ then i is even and $i - 1 \notin T^*$.




FIGURE 14. Illustrations for the proof of Lemma 12

Proof. (1) If $i \in T^*$ then i has to be odd, since otherwise the arc $(i, i + 1)$ does not exist. Assume that $x_i + \max\{x_{i+1}, x_n\} \geq 1$ and $x_i + \min\{x_{i+1}, x_n\} \leq 1$. We have to show that

$$\min\{x_i, 1 - x_n\} - \max\{1, x_i + x_{i+1}\} + x_n \geq -\max\{0, x_{i+1} - x_n\} = \min\{0, x_n - x_{i+1}\}.$$

- If $x_i + x_{i+1} \geq 1$ and $x_i + x_n \leq 1$, then the left hand side is $x_n - x_{i+1}$.
- If $x_i + x_{i+1} \leq 1$ and $x_i + x_n \geq 1$, then the left hand side is 0.

- (2) Suppose $i + 1 \in T^*$. If i is odd, then $i + 1$ is even and there is no arc $(i + 1, i + 2)$. This forces $j = i$, so that the cycle C is the triangle shown in Figure 14(b). Then

$$\text{cost}(C) = \min\{x_i, 1 - x_n\} - \min\{1, x_i + x_{i+1}\} + x_n + \min\{x_{i+1}, 1 - x_n\}.$$

To see that this cannot be negative we verify

$$\min\{x_i, 1 - x_n\} + \min\{x_{i+1}, 1 - x_n\} \geq \min\{1, x_i + x_{i+1}\} - x_n.$$

Consequently, we can assume that i is even. In order to rule out $i \in T^* + 1$, see Figure 14(c), we verify

$$-\max\{0, x_i - x_n\} + \max\{1, x_i + x_{i+1}\} - x_n \geq \min\{x_{i+1}, 1 - x_n\}.$$

- If $x_i \leq x_n$ then the left hand side is $\max\{1, x_i + x_{i+1}\} - x_n \geq 1 - x_n$.
- If $x_i > x_n$ then the left hand side is $\max\{1, x_i + x_{i+1}\} - x_i \geq x_{i+1}$. □

Lemma 13. $j + 1 \notin T^*$

Proof. Suppose $j + 1 \in T^*$. We distinguish the three cases indicated in Figure 15.




FIGURE 15. The three cases in the proof of Lemma 13.

Case (a): j is even. We need to show that

$$\max\{1, x_j + x_{j+1}\} - x_n - \max\{0, x_{j+1} - x_n\} \geq \min\{x_j, 1 - x_n\}.$$

- If $x_{j+1} \leq x_n$ then the left hand side is $\max\{1, x_j + x_{j+1}\} - x_n \geq 1 - x_n$.
- If $x_{j+1} > x_n$ then the left hand side is $\max\{1, x_j + x_{j+1}\} - x_{j+1} \geq x_j$.

Case (b): j is odd and $j \in T^* + 1$. We need to show that

$$-\min\{1, x_j + x_{j+1}\} + x_n + \min\{x_{j+1}, 1 - x_n\} \geq -\max\{0, x_j - x_n\} = \min\{0, x_n - x_j\}.$$

- If $x_{j+1} + x_n \leq 1$ then the left hand side is $x_n + x_{j+1} - \min\{1, x_j + x_{j+1}\} \geq x_n - x_j$.
- If $x_{j+1} + x_n > 1$ then the left hand side is $1 - \min\{1, x_j + x_{j+1}\} \geq 0$.

Case (c): j is odd and $j \notin T^* + 1$. We need to show that

$$-\min\{1, x_j + x_{j+1}\} + x_n + \min\{x_{j+1}, 1 - x_n\} \geq -\min\{x_j, 1 - x_n\} = \max\{-x_j, x_n - 1\}.$$

- If $x_{j+1} + x_n \leq 1$ then the left hand side is $x_n + x_{j+1} - \min\{1, x_j + x_{j+1}\} \geq x_n - x_j$.
- If $x_{j+1} + x_n > 1$ then the left hand side is $1 - \min\{1, x_j + x_{j+1}\} \geq 0$. \square

Lemma 14.

- (1) If $j \in T^*$ then j is odd and either $x_{j+1} + \max\{x_j, x_n\} < 1$ or $x_{j+1} + \min\{x_j, x_n\} > 1$.
- (2) If $j \notin T^*$ then j is even.




FIGURE 16. Illustrations for the proof of Lemma 14

Proof. (1) If $j \in T^*$ then j must be odd for the arc $(j, j+1)$ to exist. Now suppose $x_{j+1} + \max\{x_j, x_n\} \geq 1$ and $x_{j+1} + \min\{x_j, x_n\} \leq 1$. We need to show that (see Figure 16(a))

$$-\max\{1, x_j + x_{j+1}\} + x_n + \min\{x_{j+1}, 1 - x_n\} \geq -\max\{0, x_j - x_n\} = \min\{0, x_n - x_j\}.$$

- If $x_{j+1} + x_j \geq 1$ and $x_{j+1} + x_n \leq 1$ then the left hand side is $x_n - x_j$.
- If $x_{j+1} + x_j \leq 1$ and $x_{j+1} + x_n \geq 1$ then the left hand side is 0.

(2) Suppose $j \notin T^*$ and j is odd (see Figure 16(b) and (c)). Since $-m_j \geq -M_j$ it is sufficient to verify

$$-\min\{1, x_j + x_{j+1}\} + x_n + \min\{x_{j+1}, 1 - x_n\} \geq -\max\{0, x_j - x_n\} = \min\{0, x_n - x_j\}.$$

- If $x_{j+1} + x_n \leq 1$ then the left hand side is $x_n + x_{j+1} - \min\{1, x_j + x_{j+1}\} \geq x_n - x_j$.
- If $x_{j+1} + x_n > 1$ then the left hand side is $1 - \min\{1, x_j + x_{j+1}\} \geq 0$. \square

By Lemmas 11, 12, 13 and 14 we are left with the following possibilities (illustrated in Figure):

- (1) i odd and j odd. Then $i \in T^*$ and $j \in T^*$.
- (2) i odd and j even. Then $i \in T^*$ and $j \notin T^*$, $j+1 \notin T^*$.
- (3) i even and j odd. Then $i+1 \in T^*$, $i-1 \notin T^*$ and $j \in T^*$.
- (4) i even and j even. Then $i+1 \in T^*$, $i-1 \notin T^*$ and $j \notin T^*$, $j+1 \notin T^*$.

In every case, the elements of $T^* \cap [i, j]$ are the odd $k \in [i, j]$ with $1 \leq x_k + x_{k+1} + x_n \leq 2$, and the set

$$T = (T^* \setminus [i, j]) \cup \{k \in [i, j] : k \text{ even and } 1 \leq x_k + x_{k+1} + x_n \leq 2\}$$




FIGURE 17. The four types of cycles in Case 3.

is feasible, that is, $T \cap (T+1) = \emptyset$. We will write down $\text{cost}(C)$ and $\phi(T) - \phi(T^*)$, and then see that $\text{cost}(C) < 0$ implies $\phi(T) > \phi(T^*)$, as required. Similar to what we did in the discussion of Case 1, we introduce the following notation:

$$S_0 = \{k \in [i, j] : k \text{ odd and } x_k + x_{k+1} + x_n < 1\}, \quad S_2 = \{k \in [i, j] : k \text{ odd and } x_k + x_{k+1} + x_n > 2\},$$

$$S'_0 = \{k \in [i, j] : k \text{ even and } x_k + x_{k+1} + x_n < 1\}, \quad S'_2 = \{k \in [i, j] : k \text{ even and } x_k + x_{k+1} + x_n > 2\}.$$

In all cases the cost is a slight modification of

$$\text{cost}_0(C) = \sum_{\text{even } k \in [i, j]} \max\{1, x_k + x_{k+1}\} - \sum_{\text{odd } k \in [i, j]} \max\{1, x_k + x_{k+1}\} + \sum_{k \in S_0} (1 - x_k - x_{k+1}) + \sum_{k \in S_2} (x_k + x_{k+1} - 1).$$

More precisely,

$$\text{cost}(C) = \text{cost}_0(C) + \min\{x_i, 1 - x_n\} + \min\{x_{j+1}, 1 - x_n\} + x_n \quad \text{if both } i \text{ and } j \text{ are odd,} \quad (14)$$

$$\text{cost}(C) = \text{cost}_0(C) + \min\{x_i, 1 - x_n\} - \min\{x_{j+1}, 1 - x_n\} \quad \text{if } i \text{ is odd and } j \text{ is even,} \quad (15)$$

$$\text{cost}(C) = \text{cost}_0(C) - \min\{x_i, 1 - x_n\} + \min\{x_{j+1}, 1 - x_n\} \quad \text{if } i \text{ is even and } j \text{ is odd,} \quad (16)$$

$$\text{cost}(C) = \text{cost}_0(C) - \min\{x_i, 1 - x_n\} - \min\{x_{j+1}, 1 - x_n\} - x_n \quad \text{if both } i \text{ and } j \text{ are even.} \quad (17)$$

Next we will show that

$$\begin{aligned} \phi(T) - \phi(T^*) = & -\text{cost}(C) + \sum_{k \in S'_0} (1 - x_k - x_{k+1} - x_n) + \sum_{k \in S'_2} (x_k - x_{k+1} - x_n - 2) \\ & + |S_0|x_n + |S_2|(1 - x_n), \end{aligned} \quad (18)$$

and this will conclude the proof in Case 3. Let Φ_0 be the common part of $\phi(T^*)$ and $\phi(T)$, that is everything beyond the interval $[i, j+1]$. If i and j are both odd, then

$$\begin{aligned}\phi(T^*) = \Phi_0 + \sum_{k \in [i, j+1]} x_k + x_n(j-i+2)/2 - (j-i+1) + \sum_{\text{even } k \in [i, j]} \max\{1, x_k + x_{k+1}\} \\ + \sum_{k \in S_0} (1 - x_k - x_{k+1} - x_n) + \sum_{k \in S_2} (x_k + x_{k+1} + x_n - 2),\end{aligned}$$

and

$$\begin{aligned}\phi(T) = \Phi_0 + \max\{0, x_i + x_n - 1\} + \max\{0, x_{j+1} + x_n - 1\} + \sum_{k \in [i+1, j]} x_k + x_n(j-i)/2 - (j-i+1) \\ + \sum_{\text{odd } k \in [i, j]} \max\{1, x_k + x_{k+1}\} + \sum_{k \in S'_0} (1 - x_k - x_{k+1} - x_n) + \sum_{k \in S'_2} (x_k + x_{k+1} + x_n - 2).\end{aligned}$$

Using $\max\{0, x_k + x_n - 1\} - x_k = -\min\{x_k, 1 - x_n\}$ for $k = i$ and $k = j+1$, we obtain

$$\begin{aligned}\phi(T) - \phi(T^*) = \sum_{\text{odd } k \in [i, j]} \max\{1, x_k + x_{k+1}\} - \sum_{\text{even } k \in [i, j]} \max\{1, x_k + x_{k+1}\} \\ - \min\{x_i, 1 - x_n\} - \min\{x_{j+1}, 1 - x_n\} - x_n + \sum_{k \in S'_0} (1 - x_k - x_{k+1} - x_n) + \sum_{k \in S'_2} (x_k - x_{k+1} - x_n - 2) \\ - \sum_{k \in S_0} (1 - x_k - x_{k+1} - x_n) - \sum_{k \in S_2} (x_k - x_{k+1} - x_n - 2),\end{aligned}$$

and with (14) we obtain (18). If i is even and j is odd, then the same calculations lead to

$$\begin{aligned}\phi(T) - \phi(T^*) = \sum_{\text{odd } k \in [i, j]} \max\{1, x_k + x_{k+1}\} - \sum_{\text{even } k \in [i, j]} \max\{1, x_k + x_{k+1}\} \\ - \min\{x_i, 1 - x_n\} + \min\{x_{j+1}, 1 - x_n\} + \sum_{k \in S'_0} (1 - x_k - x_{k+1} - x_n) + \sum_{k \in S'_2} (x_k - x_{k+1} - x_n - 2) \\ - \sum_{k \in S_0} (1 - x_k - x_{k+1} - x_n) - \sum_{k \in S_2} (x_k - x_{k+1} - x_n - 2),\end{aligned}$$

and with (15) we obtain (18). If i is even and j is odd, then

$$\begin{aligned}\phi(T) - \phi(T^*) = \sum_{\text{odd } k \in [i, j]} \max\{1, x_k + x_{k+1}\} - \sum_{\text{even } k \in [i, j]} \max\{1, x_k + x_{k+1}\} \\ + \min\{x_i, 1 - x_n\} - \min\{x_{j+1}, 1 - x_n\} - x_n + \sum_{k \in S'_0} (1 - x_k - x_{k+1} - x_n) + \sum_{k \in S'_2} (x_k - x_{k+1} - x_n - 2) \\ - \sum_{k \in S_0} (1 - x_k - x_{k+1} - x_n) - \sum_{k \in S_2} (x_k - x_{k+1} - x_n - 2),\end{aligned}$$

and with (16) we obtain (18). Finally, if both i and j are even, then

$$\begin{aligned} \phi(T) - \phi(T^*) = & \sum_{\text{odd } k \in [i, j]} \max\{1, x_k + x_{k+1}\} - \sum_{\text{even } k \in [i, j]} \max\{1, x_k + x_{k+1}\} \\ & + \min\{x_i, 1 - x_n\} + \min\{x_{j+1}, 1 - x_n\} + x_n + \sum_{k \in S'_0} (1 - x_k - x_{k+1} - x_n) + \sum_{k \in S'_2} (x_k - x_{k+1} - x_n - 2) \\ & - \sum_{k \in S_0} (1 - x_k - x_{k+1} - x_n) - \sum_{k \in S_2} (x_k - x_{k+1} - x_n - 2), \end{aligned}$$

and with (17) we obtain (18).