

Exact first-passage time distributions for three random diffusivity models

Denis S. Grebenkov,^{1,2} Vittoria Sposini,^{2,3} Ralf Metzler,² Gleb Oshanin,⁴ and Flavio Seno⁵

¹*Laboratoire de Physique de la Matière Condensée (UMR 7643),
CNRS – Ecole Polytechnique, IP Paris, 91128 Palaiseau, France*

²*Institute for Physics and Astronomy, University of Potsdam, 14476 Potsdam-Golm, Germany*

³*Basque Center for Applied Mathematics, 48009 Bilbao, Spain*

⁴*Sorbonne Université, CNRS, Laboratoire de Physique Théorique de la Matière
Condensée (UMR 7600), 4 Place Jussieu, 75252 Paris Cedex 05, France*

⁵*INFN, Padova Section and Department of Physics and Astronomy "Galileo Galilei", University of Padova, 35131 Padova, Italy*

We study the extremal properties of a stochastic process x_t defined by a Langevin equation $\dot{x}_t = \sqrt{2D_0 V(B_t)} \xi_t$, where ξ_t is a Gaussian white noise with zero mean, D_0 is a constant scale factor, and $V(B_t)$ is a stochastic "diffusivity" (noise strength), which itself is a functional of independent Brownian motion B_t . We derive exact, compact expressions for the probability density functions (PDFs) of the first passage time (FPT) t from a fixed location x_0 to the origin for three different realisations of the stochastic diffusivity: a cut-off case $V(B_t) = \Theta(B_t)$ (Model I), where $\Theta(x)$ is the Heaviside theta function; a Geometric Brownian Motion $V(B_t) = \exp(B_t)$ (Model II); and a case with $V(B_t) = B_t^2$ (Model III). We realise that, rather surprisingly, the FPT PDF has exactly the Lévy-Smirnov form (specific for standard Brownian motion) for Model II, which concurrently exhibits a strongly anomalous diffusion. For Models I and III either the left or right tails (or both) have a different functional dependence on time as compared to the Lévy-Smirnov density. In all cases, the PDFs are broad such that already the first moment does not exist. Similar results are obtained in three dimensions for the FPT PDF to an absorbing spherical target.

Keywords: diffusing diffusivity, first passage time, Lévy-Smirnov density

PACS numbers: 02.50.-r, 05.40.-a, 02.70.Rr, 05.10.Gg

I. INTRODUCTION

There is strong experimental evidence that in some complex environments the observation of a "diffusive" behaviour, i.e., of a mean-squared displacement growing linearly with time t in the form $\overline{x^2(t)} \sim t$ does not necessarily imply that the position probability density function (PDF) $P(x, t)$ of finding a particle at position x at time t is Gaussian. In fact, significant departures from a Gaussian form have been reported, with $P(x, t)$ having cusp-like shapes in the vicinity of $x = 0$, and/or exhibiting non-Gaussian tails. Such a behaviour was observed, e.g., for the motion of micron-sized beads along nanotubes or in entangled polymer networks [1, 2], in colloidal suspensions [3] or suspensions of swimming microorganisms [4], dynamics of tracers in arrays of nanoposts [5], transport at fluid interfaces [6–8], as well as for the motion of nematodes [9]. Even more complicated non-Gaussian distributions were observed in *Dictyostelium discoideum* cell motion [10, 11] and protein-crowded lipid bilayer membranes [12, 13]. An apparent deviation from Gaussian forms was evidenced in numerical simulations of particles undergoing a polymerisation process [14].

One increasingly popular line of thought concerning the origins of such non-Gaussian diffusion advocates a picture based on the Langevin equation

$$\frac{dx_t}{dt} = \sqrt{2D_t} \xi_t, \quad (1)$$

in which ξ_t is a usual white noise with zero mean and covariance $\xi_t \xi_{t'} = \delta(t - t')$, while the diffusivity D_t is

an independent stochastic process which captures in a heuristic fashion all possible dynamical constraints, local stimuli and interactions that a particle may experience while moving in a heterogeneous complex environment.

In the pioneering work [15] Chubinsky and Slater put forth such a random diffusivity concept for dynamics in heterogeneous systems for which they coined the notion "diffusing diffusivity". Concretely, they modelled the diffusivity as a Brownian particle in a gravitational field limited by a reflecting boundary condition at $D_t = 0$ in order to guarantee positivity and stationarity of the $D(t)$ dynamics. In subsequent analyses elucidating various aspects of the diffusing diffusivity model it was assumed that D_t is a squared Ornstein-Uhlenbeck process [16–18]. Finally, [19] use a formulation directly including an Ornstein-Uhlenbeck process for D_t . All these models feature a stochastic diffusivity with bounded fluctuations around a mean value, with a finite correlation time. When the process is started with an equilibrated diffusivity distribution, the mean squared displacement has a *constant* effective amplitude at all times, in contrast to non-equilibrium initial conditions [20]. The probability density function (PDF) $P(x, t)$ of such a process is not a Gaussian function at intermediate times.¹ Instead, $P(x, t)$ exhibits a transient cusp-like behaviour

¹ The crossover from non-Gaussian to Gaussian forms distinguishes the diffusing diffusivity models here from the superstatistical approach [21] employed originally in [1, 2]. In the latter case the shape of the position PDF is permanently non-Gaussian.

in the vicinity of the origin and has exponential tails.² Further extensions of this basic model were discussed in [20, 22, 23]. Different facets of extremal and first-passage properties were scrutinised in [24–26]. We also mention recent models for non-Gaussian diffusion with Brownian scaling $\overline{x^2(t)} \sim t$ based on extreme value statistics [27] and multimerisation of the diffusing molecule [14, 28]. Generalisations to anomalous diffusion of the form $\overline{x^2(t)} \sim t^\alpha$ with $\alpha \in (0, 2)$ in terms of long-range correlated, fractional Gaussian noise was recently discussed [29, 30], as well as a fractional Brownian motion generalisation [31] of the Kärger switching-diffusivity model [32]. Finally, the role of quenched disorder is analysed in [33].

We note that the diffusing diffusivity process appeared earlier in the mathematical finance literature, where it is used for the modelling of stock price dynamics. Indeed, if we redefine dx_t/dt in (1) as $d\ln(S_t)/dt$, we recover the celebrated Black-Scholes equation [34] for the dynamics of an asset price S_t with zero-constant trend and stochastic volatility $\sqrt{2D_t}$. In this context, the choice of a squared Ornstein-Uhlenbeck process for D_t corresponds to the Heston model of stochastic volatility [35]. The process x_t in (1) thus has a wider appeal beyond the field of transport in complex heterogeneous media.

Several ad hoc diffusing diffusivity models in which the PDFs exhibit a non-Gaussian behaviour for all times have been analysed recently [36] from a more general perspective, i.e., not constraining the analysis to Brownian motion with $\overline{x^2(t)} \sim t$ only, but also extending it to anomalous diffusion. In reference [36], which focused mostly on power spectral densities of individual trajectories x_t of such processes—a topic which attracted recent interest [37–40]—the corresponding position PDFs were also obtained explicitly [36]. It was demonstrated that their functional form is very sensitive to the precise choice of D_t . Indeed, depending on the choice of D_t , one encounters a very distinct behaviour in the long time limit: the central part of the PDF may be Gaussian or non-Gaussian, diverge as $|x| \rightarrow 0$, or remain bounded in this limit, and also the tails may assume Gaussian, exponential, log-normal, or even power-law forms.

In what follows we focus on the extremal properties of three models of generalised diffusing diffusivity introduced in [36]. In these models particle dynamics obeys the Langevin equation (1) with $D_t = D_0 V(B_t)$, where D_0 is a proportionality factor—the diffusion coefficient—while $V(B_t)$ is a (dimensionless) functional of independent Brownian motion B_t , such that $B_{t=0} = 0$, $\langle B_t \rangle = 0$ and

$$\langle B_t B_{t'} \rangle = 2D_B \min(t, t'). \quad (2)$$

Here and henceforth, the angular brackets denote av-

eraging with respect to all possible realisations of the Brownian motion B_t , while the bar corresponds to averaging over realisations of the white noise process. We note parenthetically that the extremal properties of the Langevin dynamics subordinated to another Brownian motion have been actively studied in the last years within the context of the so-called run-and-tumble dynamics. In this experimentally-relevant situation, the force acting on the particle is a functional of the rotational Brownian motion (see e.g., reference [41]).

Specifically we concentrate on the first-passage times (FPTs) from a fixed position $x_0 > 0$ to a target placed at the origin and determine the full FPT PDF $H(t|x_0)$ for the following three choices for the functional V :

- (I) $V(B_t) = \Theta(B_t)$, where $\Theta(x)$ is the Heaviside theta function such that $\Theta(x) = 1$ for $x \geq 0$, and zero, otherwise;
- (II) $V(B_t) = \exp(-B_t/a)$ with a scalar parameter a ; and
- (III) $V(B_t) = B_t^2/a^2$.

In Model I, which we call "cut-off Brownian motion" the process x_t undergoes a standard Brownian motion with diffusion coefficient D_0 once $B_t > 0$, and pauses for a random time at its current location when B_t remains at negative values. Albeit the mean-squared displacement of x_t grows linearly in time in this model (see reference [36]), this is indeed a rather intricate process, in which a duration of the diffusive tours and of the pausing times have the same *broad* distribution. We note that this model represents an alternative to other standard processes describing waiting times and/or trapping events. One could think of, for instance, the comb model, in which a particle, while performing standard Brownian motion along one direction, gets stuck for a random time in branches perpendicular to the direction of the relevant diffusive motion [42–44].

In Model II the diffusivity D_t follows so-called Geometric Brownian Motion, as does an asset price in the Black-Scholes model [34]. Note that here the dynamics of x_t is not diffusive—the process progressively freezes when B_t goes in the positive direction and accelerates when B_t performs excursions in the negative direction. Overall the latter dominate and the mean-squared displacement exhibits a very fast (exponential) growth with time. Lastly, in Model III the process x_t accelerates when B_t goes away from the origin in either direction, and we are thus facing again a super-diffusive behaviour: the process x_t in (1) shows a random ballistic growth with time. In a way, such a behaviour resembles the so-called "scaled" diffusion because for typical realisations of the process B_t one has $|B_t| \sim \sqrt{t}$ and, hence, x_t evolves in the presence of a random force whose magnitude grows with time in proportion to \sqrt{t} . As shown in [36] the position PDF of this process is Gaussian around the origin and exponential in the tails. This can be compared to scaled Brownian motion, a Markovian process with time dependent diffusion coefficient $\mathcal{K}(t) \sim t^{\alpha-1}$ in the ballistic limit $\alpha \rightarrow 2$, whose position PDF stays Gaussian at all times [45]. Conversely, heterogenous diffu-

² Depending on the specific model and the spatial dimension the exponential may have a sub-dominant power-law prefactor [16–19].

sion processes with position dependent diffusion coefficient $\overline{\mathcal{K}}(x) \sim |x|^\beta$ in the limit $\beta = 1$ are also ballistic but have an exponential position PDF (with subdominant power-law correction) at all times [46].

For all three models we derive exact compact expressions for the FPT PDF $H(t|x_0)$ in one dimension and also evaluate their forms for the three-dimensional case. We remark that, in general, the exact FPT PDFs are known in closed-form only for a very limited number of situations (see, e.g., [47–52], compare [53, 54] for a "simple" spherical system). Thus, our results provide novel and non-trivial examples of stochastic processes for which the full FPT PDF can be calculated exactly.

The paper is outlined as follows. In section II we present some general arguments relating the FPT PDF and the position PDF $P(x, t)$ for the processes governed by (1). In section III we present our results for the three models under study. Finally, in section IV we conclude with a brief summary of our results and an outlook.

II. GENERAL SETUP

A general approach for evaluating the FPT PDF for the diffusing diffusivity models in (1) was developed in [24] (see also [25, 26]). In this approach, one takes the advantage of the statistical independence of thermal noise and of the stochastic diffusivity D_t . Qualitatively speaking, the thermal noise determines the statistics of the stochastic trajectories of the process x_t , whereas the diffusing diffusivity controls the "speed" at which the process runs along these trajectories. As a consequence, for a particle starting from x_0 at time 0 the FPT PDF to a target, $H(t|x_0)$, can be obtained via subordination [24] (see also [18]),

$$H(t|x_0) = \int_0^\infty dT q(t; T) H_0(T|x_0), \quad (3)$$

where $H_0(T|x_0)$ is the FPT PDF to the same target for ordinary Brownian motion with a constant diffusivity D_0 , and $q(t; T)$ is the PDF of the first-crossing time τ of a level T by the integrated diffusivity,

$$\tau = \inf\{t > 0 : T_t > T\}, \quad T_t = \int_0^t dt' D_{t'}. \quad (4)$$

In other words, T_t/D_0 plays the role of a "stochastic internal time" of the process x_t , which relates it to ordinary diffusion. The PDF $q(t; T)$ can be formally determined by inverting the identity [24]

$$\int_0^\infty dT e^{-\lambda T} q(t; T) = -\frac{\partial \Upsilon(t; \lambda)/\partial t}{\lambda}, \quad (5)$$

where $\Upsilon(t; \lambda)$ is the generating function of the integrated diffusivity,

$$\begin{aligned} \Upsilon(t; \lambda) &= \left\langle \exp \left(-\lambda \int_0^t dt' D_{t'} \right) \right\rangle \\ &= \left\langle \exp \left(-D_0 \lambda \int_0^t dt' V(B_{t'}) \right) \right\rangle. \end{aligned} \quad (6)$$

In the case of a constant diffusivity, $V(z) = 1$, one simply gets $\Upsilon(t; \lambda) = \exp(-D_0 \lambda t)$.

When the process x_t is confined to a bounded Euclidean domain $\Omega \subset \mathbb{R}^d$, the FPT PDF $H(t|x_0)$ can be obtained via a spectral expansion over the eigenvalues λ_n and eigenfunctions u_n of the Laplace operator in which the conventional time-dependence via $e^{-D_0 t \lambda_n}$ is replaced by $(-\partial \Upsilon(t; \lambda_n)/\partial t)$ [24]. In fact, substituting the spectral expansion for ordinary diffusion [48],

$$H_0(T|x_0) = \sum_n \lambda_n e^{-T \lambda_n} u_n(x_0) \int_\Omega dx u_n(x), \quad (7)$$

into equation (3), one gets with the aid of (5) that

$$H(t|x_0) = \sum_n (-\partial_t \Upsilon(t; \lambda_n)) u_n(x_0) \int_\Omega dx u_n(x). \quad (8)$$

In turn, the analysis is more subtle for unbounded domains as one can no longer rely on spectral expansions.

In what follows we focus on two emblematic unbounded domains, for which the FPT probability density $H_0(T|x_0)$ for ordinary diffusion is known:

(i) x_t evolving on a half-line $(0, \infty)$ with the starting point $x_0 > 0$ and a target placed at the origin, for which

$$H_0(T|x_0) = \frac{x_0 \exp(-x_0^2/(4T))}{\sqrt{4\pi D_0(T/D_0)^3}} \quad (9)$$

is the Lévy-Smirnov distribution (with $T = D_0 t$). Substituting this function into equation (3), one gets [24]

$$H(t|x_0) = \frac{2}{\pi} \int_0^\infty \frac{dk}{k} \sin(kx_0) (-\partial_t \Upsilon(t; k^2)). \quad (10)$$

Note that the position PDF reads

$$P(x, t|x_0) = \int_0^\infty \frac{dk}{\pi} \cos(k(x - x_0)) \Upsilon(t; k^2), \quad (11)$$

such that the two PDFs are related via $\partial P(0, t|x_0)/\partial t = 2\partial H(t|x_0)/\partial x_0$. This is specific to the half-line problem.

(ii) In the second case we consider the dynamics in a three-dimensional region outside of an absorbing sphere of radius R . In this case one has for ordinary diffusion

$$H_0^{3d}(T|x_0) = \frac{R \exp(-(|x_0| - R)^2/(4T))}{\sqrt{4\pi D_0(T/D_0)^3}}, \quad (12)$$

for any starting point $x_0 \in \mathbb{R}^3$ outside the target, i.e., with $|x_0| > R$. Comparing equations (10) and (12) one gets for *any* diffusing diffusivity process D_t :

$$H^{3d}(t|x_0) = \frac{R}{|x_0|} H(t||x_0| - R), \quad (13)$$

with $H(t|x_0)$ given by equation (10). Note that the position PDF in this case

$$P(x, t|x_0) = \frac{1}{2\pi^2|x-x_0|} \int_0^\infty dk k \sin(k|x-x_0|) \Upsilon(t; k^2). \quad (14)$$

We highlight that in an unbounded three-dimensional space some trajectories travel to infinity and never reach the target, such that the target survival probability reaches a non-zero value when time tends to infinity. This implies that the FPT PDF is not normalised with respect to the set of all possible trajectories x_t . In standard fashion, the PDF in equation (13) can be renormalised over the set of such trajectories which do reach the target up to time moment t .

Using these general results, we now obtain closed-formed expressions for the FPT PDF of models I, II, and III.

III. RESULTS

A. Model I

We first consider the functional form $V(B_t) = \Theta(B_t)$, for which the generating function of the integrated diffusivity can be straightforwardly determined by taking advantage of the celebrated results due to Kac [55] and Kac and Erdős [56]. In our notations, we have

$$\Upsilon(t; q^2) = \exp\left(-\frac{D_0 q^2 t}{2}\right) I_0\left(\frac{D_0 q^2 t}{2}\right), \quad (15)$$

where $I_\nu(z)$ is the modified Bessel function of the first kind. Note that the inverse Laplace transform of this expression produces the celebrated Lévy arcsine law [57]. Curiously, this expression does *not* depend on the diffusion coefficient D_B of Brownian motion B_t driving the diffusing diffusivity.

Substituting expression (15) into (10) we get

$$H(t|x_0) = \frac{x_0}{\sqrt{4\pi^3 D_0 t^3}} \exp\left(-\frac{x_0^2}{8D_0 t}\right) K_0\left(\frac{x_0^2}{8D_0 t}\right), \quad (16)$$

where $K_\nu(z)$ is the modified Bessel function of the second kind. For completeness we also provide the moment-generating function of the FPT \mathcal{T} ,

$$\overline{\langle \exp(-\lambda \mathcal{T}) \rangle} = \frac{2}{\pi} \int_{x_0 \sqrt{\lambda/D_0}}^\infty dz K_0(z), \quad (17)$$

where the integral can also be represented in terms of modified Struve functions.

Due to the presence of $K_\nu(z)$, the FPT PDF $H(t|x_0)$ is functionally different from the conventional Lévy-Smirnov probability density (9)—we denote it as $H^{(\text{LS})}(t|x_0)$ —and this difference manifests itself both in the left and right tails of the FPT PDF. At short times $t \ll x_0^2/(8D_0)$, (i.e., for the left tail of the FPT PDF), one gets

$$H(t|x_0) \simeq \frac{\exp(-x_0^2/(4D_0 t))}{\pi t} \quad (t \rightarrow 0), \quad (18)$$

meaning that the PDF acquires, due to the presence of $K_\nu(z)$, an additional factor $1/\sqrt{t}$. As a consequence, $H_0^{(\text{LS})}(t|x_0)/H(t|x_0) \simeq x_0/\sqrt{D_0 t} \rightarrow \infty$ in this limit, implying that $H(t|x_0)$ vanishes faster than the Lévy-Smirnov density.

Conversely, at long times $t \gg x_0^2/(8D_0)$ (i.e., for the right tail of the PDF), one has

$$H(t|x_0) \simeq \frac{x_0}{\sqrt{4\pi^3 D_0 t^3}} \left(\ln\left(\frac{16D_0 t}{x_0^2}\right) - \gamma \right), \quad (19)$$

where $\gamma = 0.5772..$ is the Euler-Mascheroni constant. Hence, in the long- t limit the FPT PDF of Model I due to the additional logarithmic factor has a *heavier* tail than the Lévy-Smirnov density. Figure 1 illustrates the FPT PDF and its asymptotic behaviour.

We also note that the FPT PDF $H(t|x_0)$ resembles the free propagator of this diffusing diffusivity motion [36],

$$\begin{aligned} P(x, t|x_0) &= \frac{e^{-(x-x_0)^2/(8D_0 t)}}{\sqrt{4\pi^3 D_0 t^3}} K_0((x-x_0)^2/(8D_0 t)) \\ &= \frac{x_0}{t} H(t||x-x_0|). \end{aligned} \quad (20)$$

This curious follows equations (10, 11) and from the fact that $\Upsilon(t; \lambda)$ for this model is only a function of $D_0 t \lambda$.

B. Model II

For Model II we have $V(B_t) = \exp(-B_t/a)$ and the corresponding function $\Upsilon(t; q^2)$ was evaluated within a different context in [58]—in fact, $\Upsilon(t; q^2)$ is related to the moment-generating function of the probability current in finite Sinai chains. Explicitly, $\Upsilon(t; q^2)$ is defined by the Kontorovich-Lebedev transform

$$\begin{aligned} \Upsilon(t; q^2) &= \frac{2}{\pi} \int_0^\infty dx \exp\left(-\frac{D_B t}{4a^2} x^2\right) \cosh\left(\frac{\pi x}{2}\right) \\ &\quad \times K_{ix}\left(2aq\sqrt{\frac{D_0}{D_B}}\right), \end{aligned} \quad (21)$$

where $K_{ix}(z)$ is the modified Bessel function of the second kind with purely imaginary index. We note that the exact forms of $\Upsilon(t; \lambda)$ are also known for the case




Figure 1: FPT PDF $H(t|x_0)$ of the diffusing diffusivity dynamics to the absorbing endpoint of the half-line $(0, \infty)$ for Model I. The thick solid line shows the exact form (16), while the thin dashed and dash-dotted lines present the short-time and long-time asymptotic relations (18) and (19), respectively. Here we set $x_0 = 1$ and $D_0 = 1$.

when B_t experiences a constant drift [59, 60]. Inserting this expression into (10) and performing the integrations we find that the FPT PDF of Model II is given explicitly by

$$H(t|x_0) = \frac{a \operatorname{arcsinh}\left(x_0 / \left(2a\sqrt{D_0/D_B}\right)\right)}{\sqrt{\pi D_B t^3}} \times \exp\left(-\frac{a^2 \operatorname{arcsinh}^2\left(\frac{x_0}{2a\sqrt{D_0/D_B}}\right)}{D_B t}\right). \quad (22)$$

Remarkably, this is exactly the Lévy-Smirnov density of the form

$$H(t|x_0) = \frac{X_0}{\sqrt{4\pi D_0 t^3}} \exp\left(-\frac{X_0^2}{4D_0 t}\right), \quad (23)$$

with an *effective* starting point

$$X_0 = 2a \operatorname{arcsinh}\left(x_0 / \left(2a\sqrt{D_0/D_B}\right)\right), \quad (24)$$

dependent not only on x_0 but also on the diffusion coefficients D_0 and D_B in a non-trivial way.

Expectedly, the moment-generating function for the FPT is simply given by a one-sided stable law of the form

$$\langle \exp(-\lambda \mathcal{T}) \rangle = \exp(-X_0 \sqrt{\lambda/D_0}), \quad (25)$$

as for Brownian motion.

C. Model III

For Model III we set $V(B_t) = B_t^2/a^2$,

the function $\Upsilon(t; q^2)$ can be calculated exactly by using the results of Cameron and Martin [61, 62] (see also [63])

$$\Upsilon(t; q^2) = \frac{1}{\sqrt{\cosh(cqt)}}, \quad (26)$$

where $c = 2\sqrt{D_B D_0}/a^2$. Inserting this expression into (10) and performing the integral, we arrive at the rather unusual form of the FPT PDF

$$H(t|x_0) = \frac{x_0}{\sqrt{2\pi^3} ct^2} \Gamma\left(\frac{1}{4} + \frac{ix_0}{2ct}\right) \Gamma\left(\frac{1}{4} - \frac{ix_0}{2ct}\right), \quad (27)$$

where $\Gamma(x)$ is the Gamma function. At short times, using the asymptotic formula $|\Gamma(a + ib)|^2 \simeq 2\pi e^{-\pi b}/\sqrt{b}$ as $b \rightarrow \infty$ for $a = 1/4$, we get

$$H(t|x_0) \simeq \frac{2\sqrt{x_0/c}}{\sqrt{\pi t^3}} \exp(-\pi x_0/(2ct)). \quad (28)$$

While the t -dependence of expression (28) is exactly the same as in the Lévy-Smirnov density, the dependence on x_0 is rather different, and also the PDF depends, through the constant c , on the diffusion coefficient D_B . In fact, setting

$$X_0^2 = \frac{2\pi x_0 D_0}{c} = \pi x_0 a \sqrt{D_0/D_B}, \quad (29)$$

we can rewrite the short-time behaviour as

$$H(t|x_0) \simeq \sqrt{8/\pi} \frac{X_0}{\sqrt{4\pi D_0 t^3}} \exp(-X_0^2/(4D_0 t)), \quad (30)$$

which is the Lévy-Smirnov distribution, except for the additional numerical factor $\sqrt{8/\pi}$. At long times, the PDF exhibits the heavy tail

$$H(t|x_0) \simeq \frac{x_0 [\Gamma(1/4)]^2}{\sqrt{2\pi^3} c} t^{-2}, \quad (31)$$

i.e., it decays *faster* than the Lévy-Smirnov distribution, but not fast enough to insure the existence of even the first moment. Figure 2 illustrates the FPT PDF $H(t|x_0)$ and its asymptotic behaviour.

Finally, the FPT PDF $H(t|x_0)$ is plotted for the three considered models in figure 3(a). In addition, we present the empirical histograms of the FPT generated by Monte Carlo simulations, observing excellent agreement. Moreover, we depict in figure 3(b) the FPT PDF $H(t|x_0)$ to an absorbing sphere of radius R along with Monte Carlo simulations results.

IV. CONCLUSION

To conclude, we studied the extremal properties of a stochastic process x_t generated by the Langevin equation (1) with a stochastic diffusivity $V(B_t)$. The latter is taken to be a functional of an independent Brownian




Figure 2: FPT PDF $H(t|x_0)$ to the absorbing endpoint of the half-line $(0, \infty)$ for Model III. The thick solid line shows the exact form (27) while the thin dashed and dash-dotted lines represent the short-time and long-time asymptotics (28) and (31), respectively. Here we set $x_0 = 1$, $D_0 = 1$, $D_B = 1$, and $a = 1$.

motion B_t . For three choices of the functional form of $V(B_t)$ we derived exact, compact expressions for the FPT PDF from a fixed initial location to the origin. Such distributions are known only for a very limited number of stochastic processes, and hence, our work provides novel examples of non-trivial processes for which this type of analysis can be carried out exactly. Similar results were obtained for the first passage time to an absorbing spherical target in three dimensions.

Following the recent reference [36], which revealed a universal large-frequency behaviour of spectral densities of individual trajectories x_t for the three models studied here, one could expect the *same* short-time asymptotic behaviour of the FPT PDF for all these models, with a generic Lévy-Smirnov form. Indeed, in [36] it was shown that the spectral densities of individual realisations of x_t decay as $1/f^2$ when $f \rightarrow \infty$, i.e., exactly as the spectral density of standard Brownian motion. However, we realised here that the FPT PDF is of Lévy-Smirnov form (with an effective starting point, dependent on the diffusion coefficients D_B and D_0) only for Model II, in which $V(B_t)$ is exponentially dependent on B_t , such that the process x_t is strongly anomalous and its mean-squared displacement grows exponentially with time. In turn, for Model I with the cut-off Brownian motion $V(B_t) = \Theta(B_t)$, which exhibits a diffusive behaviour $\langle x_t^2 \rangle \propto t$, we observed essential departures from the Lévy-Smirnov form. We saw that the corresponding FPT PDF decays faster than the Lévy-Smirnov law in the limit $t \rightarrow 0$, and slower than the Lévy-Smirnov law in the limit $t \rightarrow \infty$. For Model III with the squared Brownian motion $V(B_t) = B_t^2/a^2$, the left tail of the FPT PDF has the Lévy-Smirnov form with a renormalised starting point, while the right tail decays faster. In all models the distributions are broad such that even the first moment does not exist.

We conclude that the universal $1/f^2$ decay of the spec-




Figure 3: **(a)** Comparison of the FPT PDF $H(t|x_0)$ to the absorbing endpoint of the half-line $(0, \infty)$ for the three considered models. The solid lines represent the analytical results, while the symbols show the empirical histograms obtained from Monte Carlo simulations with 10^4 runs. We set $x_0 = 1$, $D_0 = 1$, $D_B = 1$, and $a = 1$ (for Models II and III). **(b)** Similar comparison for the FPT PDF $H(t|x_0)$ to an absorbing sphere of radius $R = 1$. The starting point is $|x_0| = \sqrt{3}$, while the other parameters are the same.

tral density does not distinguish between different diffusing diffusivity models. Indeed, as we discussed earlier, the white noise ξ_t in the Langevin equation (1) determines the statistics of trajectories in space, whereas its amplitude, $\sqrt{2D_t}$, can speed up or slow down the motion along each trajectory [24]. The spectral density is thus more sensitive to the spatial aspect of the dynamics, and its universal decay simply reflects that the statistics of the trajectories governed by the white noise is the same for all considered models. In turn, the FPT is also sensitive to the temporal aspect of the dynamics, i.e., to the "speed", at which the particle moves along the trajectory. This feature makes the spectral density and the first-passage time analysis techniques complementary.

It will be interesting to extend this analysis to other models for diffusion in heterogeneous media, in particu-

lar, when the driving noise is Gaussian but correlated, for which the shapes of the mean squared displacement and the position PDF were recently considered in a superstatistical approach [64, 65] as well as in the diffusing diffusivity picture [29, 30].

Acknowledgments

D.S.G. acknowledges a partial financial support from the Alexander von Humboldt Foundation through a

Bessel Research Award. RM acknowledges the German Science Foundation (DFG grant ME 1535/7-1) and an Alexander von Humboldt Honorary Polish Research Scholarship from the Foundation for Polish Science (Fundacja na rzecz Nauki Polskiej, FNP). FS acknowledges financial support of the 191017 BIRD-PRD project of the Department of Physics and Astronomy of Padua University.

-
- [1] B. Wang, J. Kuo, S. C. Bae, and S. Granick, *When Brownian diffusion is not Gaussian*, Nat. Mater. **11**, 481 (2012).
 - [2] B. Wang, S. M. Anthony, S. C. Bae, and S. Granick, *Anomalous yet Brownian*, Proc. Natl. Acad. Sci. USA **106**, 15160 (2009).
 - [3] J. Guan, B. Wang, and S. Granick, *Single-molecule observation of long jumps in polymer adsorption*, ACS Nano **8**, 3331 (2014).
 - [4] K. C. Leptos, J. S. Guasto, J. P. Gollub, A. I. Pesci, and R. E. Goldstein, *Dynamics of enhanced tracer diffusion in suspensions of swimming eukaryotic microorganisms*, Phys. Rev. Lett. **103**, 198103 (2009).
 - [5] K. He, F. B. Khorasani, S. T. Retterer, D. K. Tjomasn, J. C. Conrad, and R. Krishnamoorti, *Diffusive dynamics of nanoparticles in arrays of nanoposts*, ACS Nano **7**, 5122 (2013).
 - [6] C. Xue, X. Zheng, K. Chen, Y. Tian, and G. Hu, *Probing non-Gaussianity in confined diffusion of nanoparticles*, J. Phys. Chem. Lett. **7**, 514 (2016).
 - [7] D. Wang, R. Hu, M. J. Skaug, and D. Schwartz, *Temporally anticorrelated motion of nanoparticles at a liquid interface*, J. Phys. Chem. Lett. **6**, 54 (2015).
 - [8] S. Dutta and J. Chakrabarti, *Anomalous dynamical responses in a driven system*, EPL **116**, 38001 (2016).
 - [9] S. Hapca, J. W. Crawford, and I. M. Young, *Anomalous diffusion of heterogeneous populations characterized by normal diffusion at the individual level*, J. R. Soc. Interface **6**, 111 (2009).
 - [10] P. Witzel, M. Götz, Y. Lanoiselée, T. Franosch, D. S. Grebenkov, and D. Heinrich, *Heterogeneities Shape Passive Intracellular Transport*, Biophys. J. **117**, 203 (2019).
 - [11] A. G. Cherstvy, O. Nagel, C. Beta, and R. Metzler, *Non-Gaussianity, population heterogeneity, and transient superdiffusion in the spreading dynamics of amoeboid cells*, Phys. Chem. Chem. Phys. **20**, 23034 (2018).
 - [12] J.-H. Jeon, M. Javanainen, H. Martinez-Seara, R. Metzler, and I. Vattulainen, *Protein crowding in lipid bilayers gives rise to non-Gaussian anomalous lateral diffusion of phospholipids and proteins*, Phys. Rev. X **6**, 021006 (2016).
 - [13] W. He, H. Song, Y. Su, L. Geng, B. J. Ackerson, H. B. Peng, and P. Tong, *Dynamic heterogeneity and non-Gaussian statistics for acetylcholine receptors on live cell membrane*, Nat. Comm. **7**, 11701 (2016).
 - [14] F. Baldovin, E. Orlandini, and F. Seno, *Polymerization Induces Non-Gaussian Diffusion*, Front. Phys., 24 September 2019; <https://doi.org/10.3389/fphy.2019.00124>
 - [15] M. V. Chubynsky and G. W. Slater, *Diffusing Diffusivity: A Model for Anomalous, yet Brownian, Diffusion*, Phys. Rev. Lett. **113**, 098302 (2014).
 - [16] R. Jain and K. L. Sebastian, *Diffusion in a Crowded, Rearranging Environment*, J. Phys. Chem. B **120**, 3988 (2016).
 - [17] R. Jain and K. L. Sebastian, *Diffusing diffusivity: survival in a crowded rearranging and bounded domain*, J. Phys. Chem. B **120**, 9215 (2016).
 - [18] A. V. Chechkin, F. Seno, R. Metzler, and I. M. Sokolov, *Brownian yet Non-Gaussian Diffusion: From Superstatistics to Subordination of Diffusing Diffusivities*, Phys. Rev. X **7**, 021002 (2017).
 - [19] N. Tyagi and B. J. Cherayil, *Non-Gaussian Brownian diffusion in dynamically disordered thermal environments*, J. Phys. Chem. B **121**, 7204 (2017).
 - [20] V. Sposini, A. V. Chechkin, F. Seno, G. Pagnini, and R. Metzler, *Random diffusivity from stochastic equations: comparison of two models for Brownian yet non-Gaussian diffusion*, New. J. Phys. **20**, 043044 (2018).
 - [21] C. Beck and E. Cohen, *Superstatistics*, Physica A **322**, 267 (2003).
 - [22] Y. Lanoiselée and D. S. Grebenkov, *A model of non-Gaussian diffusion in heterogeneous media*, J. Phys. A. **51**, 145602 (2018).
 - [23] Y. Lanoiselée and D. S. Grebenkov, *Non-Gaussian diffusion of mixed origins*, J. Phys. A **52**, 304001 (2019).
 - [24] Y. Lanoiselée, N. Moutal, and D. S. Grebenkov, *Diffusion-limited reactions in dynamic heterogeneous media*, Nat. Commun. **9**, 4398 (2018).
 - [25] V. Sposini, A. V. Chechkin, and R. Metzler, *First passage statistics for diffusing diffusivity*, J. Phys. A **52**, 04LT01 (2019).
 - [26] D. S. Grebenkov, *A unifying approach to first-passage time distributions in diffusing diffusivity and switching diffusion models*, J. Phys. A **52**, 174001 (2019).
 - [27] E. Barkai and S. Burov, *Packets of Diffusing Particles Exhibit Universal Exponential Tails*, Phys. Rev. Lett. **124**, 060603 (2020).
 - [28] M. Hidalgo-Soria and E. Barkai, *Hitchhiker model for Laplace diffusion processes*, Phys. Rev. E **102**, 012109 (2020).
 - [29] W. Wang, F. Seno, I. M. Sokolov, A. V. Chechkin, and R. Metzler, *Unexpected crossovers in correlated random-diffusivity processes*, New J. Phys., at press.

- [30] W. Wang, A. G. Cherstvy, A. V. Chechkin, S. Thapa, F. Seno, X. Liu, and R. Metzler, *Fractional Brownian motion with random diffusivity: emerging residual nonergodicity below the correlation time*, J. Phys. A., at press.
- [31] A. Sabri, X. Xu, D. Krapf, and M. Weiss, *Elucidating the Origin of Heterogeneous Anomalous Diffusion in the Cytoplasm of Mammalian Cells*, E-print arXiv:1910.00102.
- [32] J. Kärger, *NMR self diffusion studies in heterogeneous systems*, Adv. Coll. Interf. Sci. **23**, 129 (1985).
- [33] E. B. Postnikov, A. V. Chechkin, and I. M. Sokolov, *Brownian yet non-Gaussian diffusion in heterogeneous media: from superstatistics to homogenization*, New J. Phys. **22**, 063046 (2020).
- [34] F. Black and M. Scholes, *The pricing of options and corporate liabilities*, J. Polit. Econ. **81**, 637 (1973).
- [35] S. L. Heston, *A Closed-Form Solution for Options with Stochastic Volatility with Applications to Bond and Currency Options*, Rev. Financial Studies **6**, 327 (1993).
- [36] V. Sposini, D. S. Grebenkov, R. Metzler, G. Oshanin, and F. Seno, *Universal spectral features of different classes of random diffusivity processes*, New J. Phys. **22**, 063056 (2020).
- [37] D. Krapf, E. Marinari, R. Metzler, G. Oshanin, X. Xu and A. Squarcini, *Power spectral density of a single Brownian trajectory: what one can and cannot learn from it*, New J. Phys. **20**, 023029 (2018).
- [38] D. Krapf, N. Lukat, E. Marinari, R. Metzler, G. Oshanin, C. Selhuber-Unkel, A. Squarcini, L. Stadler, M. Weiss, and X. Xu, *Spectral Content of a Single Non-Brownian Trajectory*, Phys. Rev. X **9**, 011019 (2019).
- [39] V. Sposini, R. Metzler and G. Oshanin, *Single-trajectory spectral analysis of scaled Brownian motion*, New J. Phys. **21**, 073043 (2019).
- [40] S. N. Majumdar and G. Oshanin, *Spectral content of fractional Brownian motion with stochastic reset*, J. Phys. A: Math. Theor. **51**, 435001 (2018).
- [41] F. Mori, P. Le Doussal, S. N. Majumdar and G. Schehr, *Universal survival probability for a d-dimensional run-and-tumble particle*, Phys. Rev. Lett. **124**, 090603 (2020).
- [42] V. E. Arkhincheev and E. M. Baskin, *Anomalous diffusion and drift in a comb model of percolation clusters*, Sov. Phys. JETP **73**, 161 (1991).
- [43] T. Sandev, A. Iomin, H. Kantz, R. Metzler and A. Chechkin, *Comb model with slow and Ultraslow diffusion*, Math. Model. Nat. Phenom. **11**, 18 (2016).
- [44] O. Bénichou, P. Illien, G. Oshanin, A. Sarracino and R. Voituriez, *Diffusion and subdiffusion of interacting particles on comblike structures*, Phys. Rev. Lett. **115**, 220601 (2015).
- [45] J.-H. Jeon, A. V. Chechkin, and R. Metzler, *Scaled Brownian motion: a paradoxical process with a time dependent diffusivity for the description of anomalous diffusion*, Phys. Chem. Chem. Phys. **16**, 15811 (2014).
- [46] A. G. Cherstvy, A. V. Chechkin, and R. Metzler, *Anomalous diffusion and ergodicity breaking in heterogeneous diffusion processes*, New J. Phys. **15**, 083039 (2013).
- [47] A. N. Borodin and P. Salminen, *Handbook of Brownian Motion: Facts and Formulae* (Birkhäuser Verlag, Basel-Boston-Berlin, 1996).
- [48] S. Redner, *A Guide to First Passage Processes* (Cambridge: Cambridge University Press, 2001).
- [49] C. Mejia-Monasterio, G. Oshanin and G. Schehr, *First passages for a search by a swarm of independent random searchers*, Journal of Statistical Mechanics: Theory and Experiment 2011 (06), P06022.
- [50] R. Metzler, G. Oshanin, and S. Redner (Eds.) *First-Passage Phenomena and Their Applications* (Singapore: World Scientific, 2014).
- [51] A. J. Bray, S. Majumdar, and G. Schehr, *Persistence and First-Passage Properties in Non-equilibrium Systems*, Adv. Phys. **62**, 225 (2013).
- [52] O. Bénichou and R. Voituriez, *From first-passage times of random walks in confinement to geometry-controlled kinetics*, Phys. Rep. **539**, 225 (2014).
- [53] A. Godec and R. Metzler, *Universal proximity effect in target search kinetics in the few encounter limit*, Phys. Rev. X **6**, 041037 (2016).
- [54] D. Grebenkov, R. Metzler, and G. Oshanin, *Strong defocusing of molecular reaction times: geometry and reaction control*, Comm. Chem. **1**, 96 (2018).
- [55] M. Kac, *On distributions of certain Wiener functionals*, Trans. Amer. Math. Soc. **65**, 1 (1949).
- [56] P. Erdős and M. Kac, *On the number of positive sums of independent random variables*, Bull. Amer. Math. Soc. **53**, 1011 (1947).
- [57] P. Lévy, *Sur certains processus stochastiques homogènes (On certain homogeneous stochastic processes)*, Compositio Mathematica **7**, 283 (1940).
- [58] G. Oshanin, A. Mogutov and M. Moreau, *Steady flux in a continuous-space Sinai chain*, J. Stat. Phys. **73**, 379 (1993).
- [59] C. Monthus and A. Comtet, *On the flux distribution in a one-dimensional disordered system*, J. Phys. I. France **4**, 635 (1994).
- [60] G. Oshanin and G. Schehr, *Two stock options at the races: Black-Scholes forecasts*, Quant. Fin. **12**, 1325 (2012).
- [61] R. H. Cameron and W. T. Martin, *Transformations of Wiener integrals under a general class of linear transformation*, Trans. Amer. Math. Soc. **58**, 184 (1945).
- [62] R. H. Cameron and W. T. Martin, *Evaluation of various Wiener integrals by use of certain Sturm-Liouville differential equations*, Bull. Amer. Math. Soc. **51**, 73 (1945).
- [63] M. Kac, *On distributions of certain Wiener functionals*, Trans. Amer. Math. Soc. **65**, 1 (1949).
- [64] J. Ślęzak, R. Metzler, and M. Magdziarz, *Superstatistical generalised Langevin equation: non-Gaussian viscoelastic anomalous diffusion*, New J. Phys. **20**, 023026 (2018).
- [65] T. J. Lampo, S. Stylianidou, M. P. Backlund, P. A. Wiggins, and A. J. Spakowitz, *Cytoplasmic RNA-Protein Particles Exhibit Non-Gaussian Subdiffusive Behavior*, Biophys. J. **112**, 532 (2017).