

Comment on “Chern-Simons theory and atypical Hall conductivity in the Varma phase”

Angel E. Obispo,^{1,*} Francisco A. Cruz Neto,¹ Andrés G. Jirón Vicente,^{2,1} and L. B. Castro^{1,3,†}

¹UFMA Universidade Federal do Maranhão - Campus do Bacanga - CCET, 65080-805, São Luís-MA, Brazil

²UNAC Universidad Nacional del Callao - Campus Central - FCNM. 07001, Bellavista - Callao, Perú

³UNESP Universidade Estadual Paulista - Campus de Guaratinguetá - DFQ, 12516-410, Guaratinguetá-SP, Brazil

In a recent paper published in this Journal [Phys. Rev. B **97**, 075135 (2018)], Menezes *et al.* analyze the topological behavior of a effective bosonic model defined on the Lieb lattice in presence of an electromagnetic field. In this context, the authors claim to have found an atypical quantum Hall effect for the quasiparticles. However, some inconsistencies related to the treatment of the propagator jeopardizes the main result in this system.

PACS numbers:

In an interesting Letter, Menezes *et al.* [1] analyzed the topological response of a effective bosonic theory defined on the Lieb lattice which is minimally coupled to an external $U(1)$ gauge field in $2 + 1$ dimensions. To this purpose, the authors consider a tight-binding hamiltonian with three different species of (pseudo-) gapped fermions (see equations (1)-(3) in Ref.[1]), similar to the one proposed in Ref.[2]. Such pseudo-gap behavior arises from the so-called Varma phase [3] which break time-reversal symmetry spontaneously (preserving the translational symmetry of the lattice) and whose realization would be possible in the copper-oxygen planes of high temperature cuprate superconductors [3, 4].

As a first result, in the Section II in Ref.[1], the authors showed that the dynamics of the charge carriers on that Lieb lattice in low-energy regime present a relativistic-like behavior, correctly described by Duffin-Kemmer-Petiau-like hamiltonian (DKP). Disregarding irrelevant constants, this effective hamiltonian is expressed in a simplified version as (see equation (4) in Ref.[1])

$$H_{DKP}\Psi = E\Psi, \quad (1)$$

$$H_{DKP} = [\beta^0, \beta^1] k^1 + [\beta^0, \beta^2] k^2 + m\beta^0, \quad (2)$$

where Ψ is the three-component spinor (see figure (1) in Ref.[1]),

$$\Psi(\mathbf{k}) = \begin{pmatrix} b(\mathbf{k}) \\ a(\mathbf{k}) \\ c(\mathbf{k}) \end{pmatrix}, \quad (3)$$

$\mathbf{k} = (k^1, k^2)$ is the momentum or wave vector, and β^i are two 3×3 anti-hermitian matrices which, together with another 3×3 hermitian matrix β^0 , satisfy the so-called DKP algebra

$$\beta^\mu \beta^\nu \beta^\sigma + \beta^\sigma \beta^\nu \beta^\mu = \beta^\mu \eta^{\nu\sigma} + \beta^\sigma \eta^{\nu\mu}, \quad (4)$$

with the metric tensor $\eta^{\mu\nu} = \text{diag}(1, -1, -1)$. This is the basis on which [1] develops.

In simple terms, the DKP equation is a first-order wave equation that defines spin 0 (scalar sector) and spin 1 (vectorial sector) fields and particles with a rich algebraic structure not capable of being expressed in the traditional Klein-Gordon (KG) and Proca theories [5]. Related to this, the authors in Ref.[1] state that the charge carriers on the Lieb lattice are described by *relativistic pseudospin-0 quasiparticles in two spatial dimensions*, i.e, these would exist in the scalar sector of the theory. We rebut this statement via the following argument. In $3 + 1$ spacetime dimensions, the algebra (4) generates a set of 126 independent matrices whose irreducible representations are a trivial representation, a five-dimensional representation for the scalar sector, and a ten-dimensional representation for the vectorial sector [5, 6]. Whatever the sector, it is clear that the DKP spinor will have an excess of components. In this case, the theory needs to be complemented by a constraint equation that allows to eliminate the redundant components, which is given by

$$\beta^i \beta^0 \beta^0 k_i \Psi = m(1 - \beta^0 \beta^0) \Psi, \quad i = 1, 2, 3. \quad (5)$$

With this constraint equation we can express the three (four) components of the spinor by the other two (six) components and their space derivatives in the scalar (vector) sector - for more detail see Ref.[5]. Thereby, we can exclude the redundant components and reexpress our system of equations to another that depends only on physical components (1 for the scalar sector and 4 for the vectorial sector) of the DKP theory. If we performed this same analysis in $2 + 1$ dimensions, we will see that the algebra (4) now generates a set of 35 independent matrices whose irreducible representations are a trivial representation, a four-dimensional representation, and two different three-dimensional representations. Note that only the three-dimensional representations adapt to the structure of the Hamiltonian (2), reducing Ψ to a three-component spinor, similar to (3). As the Lieb lattice has three bands at low-energy and we have three components to the spinor in (3), these components come from the three-component pseudo-spin 1 quasi-particles. The constraint

*Electronic address: angel.obispo@ufma.br, ae.obispo@gmail.com

†Electronic address: luis.castro@ufma.br

equation (5) (modified to two space dimensions, $i = 1, 2$) eliminates the redundant pseudo-spin 0 quasi-particles. Therefore, (3) describes pseudospin-1 quasiparticles. It is important to highlight that constraint equation also allows one to demonstrate the equivalence between the hamiltonian form (2) and the DKP equation of motion, which is not a trivial matter [5, 7]. In fact, a DKP quantum field theory is possible only if this equivalence is established, as can be verified in Refs.[8–10].

The section III in Ref.[1] represents the crucial point of this Comment. In that section, the authors analyzed the topological response generated by one-loop radiative corrections to the two-point function of the gauge and DKP fields in $2 + 1$ dimensions. As it is known from usual QED_{2+1} , such topological term - called Chern-Simons term - comes from the first-order in external momentum contribution of the vacuum polarization diagram [11]. In Physics of Condensed Matter (perhaps its most notable application), this emergent Chern-Simons theory naturally leads to the transverse conductivity observed from the Hall effect. In this context, the authors claim to have found an *atypical quantum Hall effect for the DKP quasiparticles* (eq. (15) in Ref.[1]), given by

$$\sigma^{xy} = \text{sign}(m) \frac{q^2}{4h}. \quad (6)$$

The above expression is their main result and represents a truly atypical result (one should expect to obtain an integer quantum hall effect), which shows for the first time in both condensed matter and higher-energy physics literature the derivation of an Abelian Chern-Simons theory from a non-Dirac system (namely DKP system). Unfortunately, we found some inconsistencies related to the treatment of the DKP propagator, in fact, the expression (13) in Ref.[1] is incorrect [8–10], therefore, the final result (6) is invalid. Below we justify our statement.

We start considering that the interaction of the electromagnetic field with the long wavelength (low-frequency) excitations of charge carriers on the Lieb lattice can be described by relativistic quantum electrodynamics for integer spin particles [8–10]. The action (8) in Ref.[1] is built upon the motion equation

$$(i\hbar\partial - q\mathcal{A} - m)\Psi = 0, \quad (7)$$

where m is the mass-gap parameter, q is coupling parameter, A_μ is the vector gauge potential,

$$\partial = \beta^\mu \frac{\partial}{\partial x^\mu}, \quad \mathcal{A} = \beta^\mu A_\mu.$$

The resulting polarization tensor in the momentum representation is given by

$$i\Pi^{\mu\nu}(p) = +\frac{q^2}{\hbar} \int \frac{d^3k}{(2\pi)^3} \text{Tr} [\beta^\mu G_\Psi(k-p) \beta^\nu G_\Psi(k)], \quad (8)$$




FIG. 1: Vacuum polarization diagram

where

$$G_\Psi(k) = i \frac{1}{\beta^\mu k_\mu - m}, \quad (9)$$

is the DKP free (Feynman) propagator. The tensor polarization (8) is equivalent to equation (12) in Ref.[1], by changing $\mu \leftrightarrow \nu$. The vacuum polarization diagram we have considered is the one shown in Fig.(1), which allowed us built $i\Pi^{\mu\nu}(p)$ following the same Feynman rules of the usual QED_{2+1} , except for the plus sign (+) in front, which is a reminiscent of the bosonic nature of the DKP theory [8]. The standard procedure to calculate $i\Pi^{\mu\nu}(p)$ says that we must first evaluate the trace of β matrices, which implies that $G_\Psi(k)$ in (9) must be rewritten in such a way that these matrices appear in the numerator. Nevertheless, this process in DKP theory is more complicated (as compared with Dirac theory) due mainly to its algebra and because the β matrices are singular ($\det[\beta] = 0$). As β^{-1} does not exist, it implies that some common identities are not valid anymore, for instance, $(\beta^\mu p_\mu)(\beta^\nu p_\nu)^{-1} = \mathbb{I}$. This did not take into account by the authors in Ref.[1], and as result, their propagator was incorrectly constructed (see equation (13) in Ref.[1]). The correct form for the DKP propagator is also performed in Refs.[12–14], and is expressed as follows

$$G_\Psi(k) = i \frac{1}{\beta^\mu k_\mu - m} = \frac{i}{m} \left[\frac{k(k+m)}{k^2 - m^2} - 1 \right]. \quad (10)$$

It is straightforward to see that the propagator in (10) is strictly defined for massive particles, as required for the DKP theory. In fact, the equation (2) with $m = 0$ represent a different relativistic equation, the so-called Harish-Chandra equation [15], whose analysis we will leave aside here. So, we go to focus on the propagator in (10), alternatively rewritten as

$$\begin{aligned} G_\Psi(k) &= G_1(k) + G_2(k) \\ &= i \frac{k(k+m)}{k^2 - m^2} + \frac{i}{m} \frac{(k^2 - k^2)}{k^2 - m^2}. \end{aligned} \quad (11)$$

Note that the first term on the right $G_1(k)$ coincides exactly with the propagator proposed by Menezes *et al.* (see equation (13) in Ref.[1]), which is used to determine the value of the Hall conductivity by considering the contribution from antisymmetric part of the polarization tensor $i\Pi^{\mu\nu} \sim \int d^3k \text{Tr}[\beta^\mu G_1 \beta^\nu G_1]$ and the second term $G_2(k)$ absent. The point of this Comment is to demonstrate that, when $G_2(k)$ is inserted into (8), the

term $\sim \int d^3k \text{Tr}[\beta^\mu G_1 \beta^\nu G_2 + \beta^\mu G_2 \beta^\nu G_1 + \beta^\mu G_2 \beta^\nu G_2]$ contributes a non-negligible value to the value of the Hall conductivity found in Ref.[1].

For this purpose, we rewrite the polarization tensor according to decomposition (11)

$$i\Pi^{\mu\nu}(p) = i\Pi_{(1,1)}^{\mu\nu} + i\Pi_{(1,2)}^{\mu\nu} + i\Pi_{(2,1)}^{\mu\nu} + i\Pi_{(2,2)}^{\mu\nu}, \quad (12)$$

where $i\Pi_{(i,j)}^{\mu\nu}$ are functions of momentum p , conveniently defined as

$$i\Pi_{(1,1)}^{\mu\nu} = \frac{q^2}{\hbar} \int \frac{d^3k}{(2\pi)^3} \text{Tr}[\beta^\mu G_1(k-p) \beta^\nu G_1(k)],$$

$$i\Pi_{(1,2)}^{\mu\nu} = \frac{q^2}{\hbar} \int \frac{d^3k}{(2\pi)^3} \text{Tr}[\beta^\mu G_1(k-p) \beta^\nu G_2(k)],$$

$$i\Pi_{(2,1)}^{\mu\nu} = \frac{q^2}{\hbar} \int \frac{d^3k}{(2\pi)^3} \text{Tr}[\beta^\mu G_2(k-p) \beta^\nu G_1(k)],$$

$$i\Pi_{(2,2)}^{\mu\nu} = \frac{q^2}{\hbar} \int \frac{d^3k}{(2\pi)^3} \text{Tr}[\beta^\mu G_2(k-p) \beta^\nu G_2(k)].$$

As previously mentioned, the result obtained in Ref.[1] can be reproduced by computing simply the antisymmetric part of $i\Pi_{(1,1)}^{\mu\nu}$, so, in that sense, it is convenient to divide the Eq. (12) into two contributions: $i\Pi_{(1,1)}^{\mu\nu}$ and $i\Pi_{(1,2)}^{\mu\nu} + i\Pi_{(2,1)}^{\mu\nu} + i\Pi_{(2,2)}^{\mu\nu}$. As we are only interested in the topological part of each contribution, we focus our attention on the terms $\sim \epsilon^{\mu\nu\alpha} p_\alpha$, which come from combining an odd number of β matrices,

$$\text{Tr}[\beta^\rho \beta^\sigma \beta^\theta] = i\epsilon^{\rho\sigma\theta}, \quad (13)$$

$$\begin{aligned} 2\text{Tr}[\beta^\rho \beta^\alpha \beta^\sigma \beta^\omega \beta^\theta] &= ig^{\rho\alpha} \epsilon^{\sigma\omega\theta} + ig^{\alpha\sigma} \epsilon^{\omega\theta\rho} \\ &\quad + ig^{\sigma\omega} \epsilon^{\theta\rho\alpha} + ig^{\omega\theta} \epsilon^{\rho\alpha\sigma} \\ &\quad + ig^{\rho\omega} \epsilon^{\alpha\sigma\theta} + ig^{\alpha\theta} \epsilon^{\rho\sigma\omega}. \end{aligned} \quad (14)$$

Thus, following the standard methods for QED calculations [8], we will compute the antisymmetric part of each contribution separately.

• Computing $i\Pi_{AS(1,1)}^{\mu\nu}$:

To determinate the contribution from this first term (which is used in Ref.[1] to get its main result), we start applying the trace property (13). After a few calculations we obtain

$$i\Pi_{AS(1,1)}^{\mu\nu} = -\frac{imq^2}{\hbar} \int_0^1 dx \int \frac{d^3k}{(2\pi)^3} \frac{p_\alpha}{[k^2 - \Delta^2]^2} \epsilon^{\mu\nu\alpha},$$

where we have used the Feynman parametrization procedure

$$\frac{1}{[(k-p)^2 - m^2][k^2 - m^2]} = \int_0^1 dx \frac{1}{[k^2 - \Delta^2]^2}, \quad (15)$$

together with the change $k \rightarrow k + xp$ and $\Delta = m^2 - p^2 x(1-x)$. These integrals are the massive one-loop Feynman integrals, widely studied in QED, and whose result we will use directly. Thereby, we get

$$\begin{aligned} i\Pi_{AS(1,1)}^{\mu\nu} &= \frac{m}{|m|} p_\alpha \epsilon^{\mu\nu\alpha} \frac{q^2}{4h} \int_0^1 dx \frac{1}{\sqrt{1-x(1-x)p^2/m^2}}, \\ &= \text{sgn}(m) \frac{1}{4} \frac{q^2}{h} p_\alpha \epsilon^{\mu\nu\alpha}, \end{aligned} \quad (16)$$

where in the last line we have considered the Chern-Simons regime ($m \gg p$). This is the result obtained by Menezes in their manuscript, as expected.

• Computing $i\Pi_{AS(1,2)}^{\mu\nu} + i\Pi_{AS(2,1)}^{\mu\nu} + i\Pi_{AS(2,2)}^{\mu\nu}$:

In this case, both (13) and (14) are required. A quick inspection allows us to demonstrate that $i\Pi_{AS(2,2)}^{\mu\nu} = 0$, i.e., this term does not have an antisymmetric part. By other hand, the sum of the cross terms provides

$$\begin{aligned} i\Pi_{AS(1,2)}^{\mu\nu} + i\Pi_{AS(2,1)}^{\mu\nu} &= -\frac{iq^2}{2m\hbar} \int_0^1 dx \int \frac{d^3k}{(2\pi)^3} \times \\ &\quad \frac{1}{[(k-px)^2 - \Delta^2]^2} \times \text{Tr}[p], \end{aligned}$$

where we use the parametrization (15), with $\Delta = m^2 - p^2 x(1-x)$ and

$$\begin{aligned} \text{Tr}[p] &= 2\epsilon^{\nu\alpha\omega} p_\alpha k^\mu k_\omega - 2\epsilon^{\mu\alpha\omega} p_\alpha k^\nu k_\omega \\ &\quad + 2\epsilon^{\mu\nu\alpha} k_\alpha (pk) - \epsilon^{\nu\alpha\omega} p_\alpha p^\mu k_\omega \\ &\quad + \epsilon^{\mu\alpha\omega} p_\alpha p^\nu k_\omega - \epsilon^{\mu\nu\alpha} p_\alpha (pk) - \epsilon^{\mu\nu\alpha} k_\alpha p^2. \end{aligned}$$

At this point, a regularization scheme is required. We use the Pauli-Villars regularization, which allows us to make the change $k \rightarrow k + px$, to then exclude the linear terms in k associated to odd-integrals, and replace $k^\mu k^\nu$ by $g^{\mu\nu} k^2/3$ in numerator. Thereby we obtain:

$$\begin{aligned} i\Pi_{AS(1,2)}^{\mu\nu} + i\Pi_{AS(2,1)}^{\mu\nu} &= \text{sgn}(m) \frac{3}{4} \frac{q^2}{h} \epsilon^{\mu\nu\alpha} p_\alpha \times \\ &\quad \int_0^1 dx \frac{3 - 2x(1-x)(p^2/m^2)}{3\sqrt{1-x(1-x)(p^2/m^2)}}. \end{aligned}$$

In the Chern-Simons regime ($m \gg p$)

$$i\Pi_{AS(1,2)}^{\mu\nu}(p) + i\Pi_{AS(2,1)}^{\mu\nu}(p) = \text{sgn}(m) \frac{3}{4} \frac{q^2}{h} \epsilon^{\mu\nu\alpha} p_\alpha. \quad (17)$$

The above expression represents the main result of this Comment. Note that it is three times larger than the one found in (16), which implies an important modification to the result found by Menezes *et al.* [1]. Thus, if we combine the eqs. (16) and (17) to find the full expression of the dynamically generated Chern-Simons term, we get

$$\begin{aligned} i\Pi_{AS}^{\mu\nu}(p) &= i\Pi_{AS(1,1)}^{\mu\nu} + i\Pi_{AS(1,2)}^{\mu\nu} + i\Pi_{AS(2,1)}^{\mu\nu}, \\ &= \text{sign}(m) \frac{q^2}{h} \epsilon^{\mu\nu\theta} p_\theta. \end{aligned} \quad (18)$$

The Hall conductivity can be obtained via the Kubo's formula

$$\sigma^{xy} = \lim_{\mathbf{p} \rightarrow 0, p_0 \rightarrow 0} \frac{i\Pi_{AS}^{xy}}{p_0} = \text{sign}(m) \frac{q^2}{h}, \quad (19)$$

which is the result expected according to the literature [3, 16, 17]. Therefore, there is no such atypical Hall conductivity, as reported by the authors.

We conclude by emphasizing that the results and conclusions presented in this Comment do not alter the others results shown in [1], concerning the obtaining of Landau levels in DKP theory and to the extension of Jackiw-Rebbi approach for the DKP quasiparticles.

ACKNOWLEDGMENTS

Angel E. Obispo thanks to CNPq (grant 312838/2016-6) and Secti/FAPEMA (grant FAPEMA DCR-02853/16), for financial support. L. B. Castro also thanks to CNPq, Brazil, Grant No. 307932/2017-6 (PQ) and No. 422755/2018-4 (UNIVERSAL), São Paulo Research Foundation (FAPESP), Grant No. 2018/20577-4, FAPEMA, Brazil, Grant No. UNIVERSAL-01220/18 and CAPES, Brazil. The authors would like to thank R. Casana for useful discussions.

-
- [1] N. Menezes, C.M. Smith, G. Palumbo, Phys. Rev. B **97**, 075135 (2018). DOI 10.1103/PhysRevB.97.075135. URL <https://link.aps.org/doi/10.1103/PhysRevB.97.075135>
 - [2] Y. He, J. Moore, C.M. Varma, Phys. Rev. B **85**, 155106 (2012). DOI 10.1103/PhysRevB.85.155106. URL <https://link.aps.org/doi/10.1103/PhysRevB.85.155106>
 - [3] C.M. Varma, Phys. Rev. Lett. **83**, 3538 (1999). DOI 10.1103/PhysRevLett.83.3538. URL <https://link.aps.org/doi/10.1103/PhysRevLett.83.3538>
 - [4] C.M. Varma, Phys. Rev. B **73**, 155113 (2006). DOI 10.1103/PhysRevB.73.155113. URL <https://link.aps.org/doi/10.1103/PhysRevB.73.155113>
 - [5] L.B. Castro, A.S. de Castro, Phys. Rev. A **90**, 022101 (2014). DOI 10.1103/PhysRevA.90.022101. URL <http://link.aps.org/doi/10.1103/PhysRevA.90.022101>
 - [6] E. Corson, *Introduction to Tensors, Spinors, and Relativistic Wave-equations (relation Structure)* (Hafner Publishing Company, 1953). URL <https://books.google.com.br/books?id=1IMuAAAAIAAJ>
 - [7] J.T. Lunardi, B.M. Pimentel, R.G. Teixeira, J.S. Valverde, Phys. Lett. A **268**, 165 (2000). DOI [http://dx.doi.org/10.1016/S0375-9601\(00\)00163-8](http://dx.doi.org/10.1016/S0375-9601(00)00163-8). URL <http://www.sciencedirect.com/science/article/pii/S0375960100001638>
 - [8] A.I. Akhiezer, V.B. Berestetskii, *Quantum Electrodynamics* (Jonh Wiley & Sons, New York, 1965)
 - [9] J.S. Valverde, A teoria de duffin-kemmer-petiau: Aspectos clássicos e quânticos. Master's thesis, IFT-UNESP (2000)
 - [10] S.I. Kruglov, Int. J. Mod. Phys. A **26**(15), 2487 (2011). DOI 10.1142/S0217751X11053523. URL <https://doi.org/10.1142/S0217751X11053523>
 - [11] A.E. Obispo, M. Hott, Phys. Rev. B **89**, 165405 (2014). DOI 10.1103/PhysRevB.89.165405. URL <http://link.aps.org/doi/10.1103/PhysRevB.89.165405>
 - [12] J.T. Lunardi, B.M. Pimentel, J.S. Valverde, L.A. Manzoni, Int. J. Mod. Phys. A **17**(02), 205 (2002). DOI 10.1142/S0217751X02005682. URL <https://doi.org/10.1142/S0217751X02005682>
 - [13] R. Bufalo, T.R. Cardoso, A.A. Nogueira, B.M. Pimentel, J. Phys.: Conference Series **706**, 052002 (2016). DOI 10.1088/1742-6596/706/5/052002. URL <https://doi.org/10.1088/1742-6596/706/5/052002>
 - [14] J. Kowalski-Glikman, G. Rosati, Phys. Rev. D **101**, 065003 (2020). DOI 10.1103/PhysRevD.101.065003. URL <https://link.aps.org/doi/10.1103/PhysRevD.101.065003>
 - [15] Harish-Chandra, Proc. Roy. Soc. Lon. A **186**(1007), 502 (1946). DOI 10.1098/rspa.1946.0061. URL <https://royalsocietypublishing.org/doi/abs/10.1098/rspa.1946.0061>
 - [16] M. Nita, B. Ostahie, A. Aldea, Phys. Rev. B **87**, 025428 (2013). DOI 10.1103/PhysRevB.87.025428. URL <https://link.aps.org/doi/10.1103/PhysRevB.87.025428>
 - [17] N. Goldman, D.F. Urban, D. Bercioux, Phys. Rev. A **83**, 063601 (2011). DOI 10.1103/PhysRevA.83.063601. URL <https://link.aps.org/doi/10.1103/PhysRevA.83.063601>