

Reply to Comment on “Chaotic-Integrable Transition in the Sachdev-Ye-Kitaev Model”

Antonio M. García-García,¹ Bruno Loureiro,² Aurelio Romero-Bermúdez,³ and Masaki Tezuka⁴

¹Shanghai Center for Complex Physics, Department of Physics and Astronomy,
Shanghai Jiao Tong University, Shanghai 200240, China*

²Université Paris-Saclay, CNRS, CEA, Institut de physique théorique, 91191, Gif-sur-Yvette, France†

³Institute Lorentz for Theoretical Physics, Leiden University, Leiden, The Netherlands‡

⁴Department of Physics, Kyoto University, Kyoto 606-8502, Japan§

In a recent comment [1] to the paper *Chaotic Integrable transition in the SYK model* [2], it was claimed that, in a certain region of parameters, the Lyapunov exponent of the N Majoranas Sachdev-Ye-Kitaev model with a quadratic perturbation studied in [2] is always positive. This implies that the model is quantum chaotic. In this reply, we show that the perturbative formalism of Ref. [1] breaks down precisely in the range of parameters investigated in Ref. [1] due to a lack of separation of time scales. Moreover, based on recent analytical results, we show that for any large and fixed N , the model has indeed a chaotic-integrable transition that invalidate the results of the comment.

The Lyapunov exponent λ_L is obtained by carrying out a semiclassical $1/N$ expansion of certain out of time correlation functions. To leading order, these correlation functions grow exponentially in time $\sim e^{\lambda_L t}/N^2$. Physically, this exponential growth defines a minimum time scale, called the Ehrenfest or scrambling time $t_E \sim \log N/\lambda_L$, for quantum effects to become important in an initially quasi-classical wavepacket. This exponential growth is restricted to quantum chaotic systems while for integrable systems the growth is typically power-law with a non-universal exponent. Therefore, only for quantum chaotic systems, the Lyapunov exponent is finite. For the deformed SYK model of interest [2], where the typical random quartic interaction of strength J is perturbed by a quadratic perturbation of strength κ , the Lyapunov exponent λ_L is a function of T, λ, κ where T is the temperature of the system. The main claim of [1], is that for N large and fixed, and κ sufficiently large, so that $T/\kappa \ll 1$ and $J/\kappa \ll 1$,

$$\frac{\lambda_L}{\kappa} = \frac{3T^2 J^2}{\kappa^4}. \quad (1)$$

so $\lambda_L > 0$, no transition occurs and the model is always quantum chaotic. This is in tension with the results of Ref.[2] where we show that, in this range of parameters, a chaotic-integrable transition takes place for sufficiently large κ . In the following, we show that the semiclassical formalism employed to obtain Eq. (1) breaks down in the range of parameters above. Moreover, we provide evidence that in any case, in that range of parameters, the system is no longer quantum chaotic. We note the perturbation theory leading to Eq. (1) in [1] is only valid for $t_E \leq t \ll t_H$ where $t_H \sim \frac{1}{\Delta}$ is the Heisenberg time and Δ is the mean level spacing. Physically, this is the time scale for non-perturbative quantum effects, related to the discreteness of the spectrum, to become dominant. The analytical treatment of Ref. [1] is semiclassical and therefore it breaks down when t_E becomes smaller but of the order of t_H . In our model, we can estimate $t_H \sim 2^{-N/2}$. Therefore, the semiclassical formalism leading to Eq. (1) is not applicable for:

$$t_E \geq t_H \quad \Leftrightarrow \quad \lambda_L(\kappa, T) \leq 2^{N/2} \log N \quad (2)$$




FIG. 1. Ensemble average of the adjacent gap ratio $\langle r_i \rangle$ for the lowest eigenvalues, labeled by the index i with respect to the ground state, that correspond with the limit of low temperatures. Agreement with Poisson statistics is excellent for sufficiently large, but finite, κ which confirms that the system is no longer quantum chaotic.

Using Eq. (1), it is straightforward to estimate the maximum $\kappa = \kappa_*$ for which the formalism of [1] applies:

$$\kappa_* \left(\frac{T}{\kappa}, N \right) \sim \left(\frac{T}{\kappa} \right)^{-2} 2^{N/2} \log N \quad (3)$$

This is in contradiction with the claim of Ref.[1] that, for any fixed value of $N \gg 1$, T, J , Eq. (1) applies for a sufficiently large κ . This is only true for $\kappa < \kappa_*$ which in principle invalidates the main claim of Ref. [1]. We say in principle, because a natural criticism to this argument is that, even though the semiclassical formalism does not apply for sufficiently large κ , it may be the system is still quantum chaotic and therefore no transition to integrability occurs. We show below that this is not the case and that the system undergoes a transition from quantum chaos to localization not only in this range of parameters but, for sufficiently large κ , at any temperature.

The fact that the system is not quantum chaotic in the large κ limit is rather evident from the level statistics analysis we performed in Ref. [2]. Spectral correlators such as the level




FIG. 2. Distribution function $P(r)$ of the adjacent gap ratio for $\kappa = 50$ (see definition of r around Eq. 35 of Ref. [3]), for different temperatures (T is defined in Ref. [2]). Agreement with Poisson statistics contradicts the claim of Ref [1] that the system is quantum chaotic in this region




FIG. 3. Lyapunov exponent as a function of temperature for various κ , circles are the results of Ref. [2] and the squares are the recalculated results. Only few points show a substantial difference with the results of Ref. [2]. However, the critical temperature is substantially reduced.

spacing distribution or the average adjacent gap ratio $\langle r \rangle$ are very close to the prediction for Poisson statistics typical of an insulator/integrable system. Indeed, correlations of eigenvalues closer to the edge of the spectrum, corresponding to the limit of low temperatures, tend to exhibit Poisson statistics, typical of localized states, at even relatively small κ , see the adjacent gap ratio in Fig. 11 of the supplemental information of Ref. [2] that for convenience we reprint in Fig. 1, or the lower panels of Fig. 1 and Fig. 2. in the main paper. As a further confirmation of this point, in Fig. 2, we depict results of the distribution function of the adjacent spacing ratio $P(r)$ for $\kappa = 50$ and different temperatures. In contradiction with the claims of Ref. [1] about the persistence of quantum chaos for $\kappa/T \gg 1$, we find that, in this region, $P(r)$ is already very close to Poisson statistics typical of an integrable or many-body localized phase. For larger values of κ , the agreement to Poisson statistics (not shown) is even better. Further evidence that the system is not quantum chaotic at finite N , for sufficiently large κ , has recently been reported in Ref. [3], coauthored

by one of us (M.T.). By an exact matrix integral representation, spectral and eigenfunction correlation functions of the SYK model with a quadratic perturbation were mapped onto a supersymmetric sigma model which was solved by standard field theory techniques. The system has a rather rich phase diagram with chaotic, localized/integrable, and intermediate chaotic but non-ergodic phases. For the purpose of this reply, the main result of Ref. [3] is that for $\kappa > \kappa_c \sim N^2 \log N$, the system is no longer quantum chaotic as *all* states are many-body localized, and spectral correlations are well described by Poisson statistics. In the language used in Ref. [1], this means the the Lyapunov exponent is zero for any temperature. We expect a much weaker N dependence, if any, of κ_c in the tail of the spectrum but this result refutes already the main claim of Ref. [1] that λ_L is finite. This analytical result for κ_c was supported by numerical simulations for $N = 22 - 30$ Majorana fermions, involving both level and eigenfunction statistics. In all cases, good agreement between analytical and numerical results was found.

We also mention that while we were checking the calculation of λ_L in [2], we noticed that for a few κ 's in the low temperature limit, our code did not pick up the largest λ_L but other sub-leading solutions of the eigenvalue problem that are not relevant for sufficiently long times. In Fig. 3, we compare Fig. 3 of [2] with the recalculated λ_L . As is observed, the corrections are only appreciable for a few points. A fitting of $\lambda_L(T)$ still suggests the possible existence of a critical temperature $T_c(\kappa)$ below which $\lambda_L = 0$. However, the values of T_c resulting from the fitting of the curve are relatively low. For instance, $T_c(2) = 0.003$ $(-0.01, 0.02)$ and $T_c(0.2) = 0.0003$ $(0.00005, 0.0005)$. The value of the 95% confidence interval, in parenthesis, is not conclusive about the vanishing of T_c at finite temperature. In any case, the conclusions of this reply are not modified because, as argued earlier, the theoretical formalism employed in Ref. [1] breaks down for a sufficiently large $\kappa/T \gg 1$ and, more importantly, in stark contrast with the claim of Ref. [1], level statistics results together with the analytical findings of Ref. [3], confirm that the model is not quantum chaotic but many-body localized/integrable for sufficiently large κ .

* amgg@sju.edu.cn

† brloureiro@gmail.com

‡ aurelio.romero.bermudez@gmail.com

§ tezuka@scphys.kyoto-u.ac.jp

- [1] J. Kim and X. Cao, Comment on chaotic-integrable transition in the Sachdev-Ye-Kitaev model (2020), arXiv:2004.05313 [cond-mat.stat-mech].
- [2] A. M. García-García, B. Loureiro, A. Romero-Bermúdez, and M. Tezuka, Chaotic-integrable transition in the Sachdev-Ye-Kitaev model, Phys. Rev. Lett. **120**, 241603 (2018).
- [3] F. Monteiro, T. Micklitz, M. Tezuka, and A. Altland, Fock space localization in the Sachdev-Ye-Kitaev model (2020), arXiv:2005.12809 [cond-mat.str-el].