

Nichols algebras over classical Weyl groups

Zhengtang Tan ^a, Weicai Wu ^b, Shouchuan Zhang ^c

a. College of Engineering and Design, Hunan Normal University, Changsha 410081, P.R. China

b. College of Mathematics, Hunan Institute of Science and Technology, Yueyang 414006, P.R. China

c. Department of Mathematics, Hunan University, Changsha 410082, P.R. China

Emails: weicaiwu@hnu.edu.cn (WCW)

Abstract

We show that except in several cases conjugacy classes of classical Weyl groups $W(B_n)$ and $W(D_n)$ are of type D. We prove that except in three cases Nichols algebras of irreducible Yetter-Drinfeld (YD in short) modules over the classical Weyl groups are infinite dimensional.

2010 Mathematics Subject Classification: 16W30, 22E60, 11F23

Keywords: Rack, classical Weyl groups, Nichols algebras.

1 Introduction

Nichols algebras play a fundamental role in the classification of finite-dimensional complex pointed Hopf algebras by means of the lifting method developed by Andruskiewitsch and Schneider [AS02]. In this context, given a group G , an important step to classify all finite-dimensional complex pointed Hopf algebras H with group-like $G(H) = G$ is to determine all the pairs (\mathcal{O}, ρ) such that the associated Nichols algebra $\mathfrak{B}(\mathcal{O}, \rho)$ is finite-dimensional. Here the pairs (\mathcal{O}, ρ) are such that \mathcal{O} runs over all conjugacy classes of G and ρ runs over all irreducible representations of the centralizer of g in G , with $g \in \mathcal{O}$ fixed. In general, this is a difficult task since Nichols algebra is defined by generators and relations. In practice, it is often useful to discard those pairs such that $\dim \mathfrak{B}(\mathcal{O}, \rho) = \infty$. There are properties of the conjugacy class \mathcal{O} that imply that $\dim \mathfrak{B}(\mathcal{O}, \rho) = \infty$ for any ρ , one of which is the property of being of type D. This is useful since it reduces the computations to operations inside the group and avoids hard calculations of generators and relations of the corresponding Nichols algebra.

This work contributes to the classification of finite-dimensional Hopf algebras over an algebraically closed field of characteristic 0. This problem was posed by Kaplansky

in 1975. The lifting method of Andruskiewitsch and Schneider describes a way to classify finite-dimensional complex Hopf algebras. In [AS10] they obtained the classification over the finite abelian groups whose order is relative prime with 210. Nichols algebras of braided vector spaces $(\mathbb{C}X, c_q)$, where X is a rack and q is a 2-cocycle in X , were studied in [AG03]. It was shown [AFGV08, AFZ09, AZ07] that Nichols algebras $\mathfrak{B}(\mathcal{O}_\sigma, \rho)$ over symmetry groups are infinite dimensional, except in a number of remarkable cases corresponding to \mathcal{O}_σ . Two of the present authors [ZZ12] showed that except in three cases Nichols algebras of irreducible Yetter-Drinfeld (YD) modules over classical Weyl groups $A \rtimes \mathbb{S}_n$ supported by \mathbb{S}_n are infinite dimensional. However, the classification has not been completed for Nichols algebras over general classic Weyl groups $W(B_n)$ and $W(D_n)$.

Note that $\mathbb{Z}_2^n \rtimes \mathbb{S}_n$ is isomorphic to Weyl groups $W(B_n)$ and $W(C_n)$ of B_n and C_n for $n > 2$. If $A = \{a \in \mathbb{Z}_2^n \mid a = (a_1, a_2, \dots, a_n) \text{ with } a_1 + a_2 + \dots + a_n \equiv 0 \pmod{2}\}$, then $A \rtimes \mathbb{S}_n$ is isomorphic to Weyl group $W(D_n)$ of D_n for $n > 3$. Obviously, when $A = \{a \in \mathbb{Z}_2^n \mid a = (a_1, a_2, \dots, a_n) \text{ with all } a_i \equiv 0 \pmod{2}\}$, $A \rtimes \mathbb{S}_n$ is isomorphic to Weyl group of A_{n-1} for $n > 1$. Note that \mathbb{S}_n acts on A as follows: for any $a \in A$ with $a = (a_1, a_2, \dots, a_n)$ and $\sigma, \tau \in \mathbb{S}_n$, $\sigma \cdot a := (a_{\sigma^{-1}(1)}, a_{\sigma^{-1}(2)}, \dots, a_{\sigma^{-1}(n)})$.

It is clear that

$$\begin{aligned} (a, \sigma)^{-1} &= (-(a_{\sigma(1)}, a_{\sigma(2)}, \dots, a_{\sigma(n)}), \sigma^{-1}) = (-\sigma^{-1}(a), \sigma^{-1}), \\ (b, \tau)(a, \sigma)(b, \tau)^{-1} &= (b + \tau(a) - \tau\sigma\tau^{-1}(b), \tau\sigma\tau^{-1}). \\ a\sigma &= (a, \sigma). \end{aligned}$$

Without specification, $K_n := \{a \in \mathbb{Z}_n \mid a_1 + a_2 + \dots + a_n = 0\}$ and $K_n \rtimes \mathbb{S}_n$ is a subgroup of $\mathbb{Z}_2^n \rtimes \mathbb{S}_n$. Let \mathbb{W}_n denote $K_n \rtimes \mathbb{S}_n$ or $\mathbb{Z}_2^n \rtimes \mathbb{S}_n$ throughout this paper. When we say type D we mean a rack or conjugacy class of type D and not a Weyl group of type D_n .

In this paper we prove that except in several cases conjugacy classes of classical Weyl groups \mathbb{W}_n are of type D, and except in three cases Nichols algebras of irreducible YD modules over the classical Weyl groups are infinite dimensional.

The work is organized as follows. In Section 1 we provide some preliminaries and set our notations. In Section 2 we determine when the conjugacy classes of juxtapositions of two elements are of type D. In Section 3 we examine the rack and conjugacy classes of \mathbb{W}_n . In Section 4 we classify Nichols algebras of irreducible YD modules over the classical Weyl groups.

Preliminaries and conventions

Let \mathbf{k} be an algebraically closed field of characteristic zero and (V, C) a braided vector space, i.e. V is a vector space over \mathbf{k} and $C \in \text{Aut}(V \otimes V)$ is a solution of the braid equation $(\text{id} \otimes C_{V,V})(C_{V,V} \otimes \text{id})(\text{id} \otimes C_{V,V}) = (C_{V,V} \otimes \text{id})(\text{id} \otimes C_{V,V})(C_{V,V} \otimes \text{id})$. $C_{i,i+1} :=$

$\text{id}^{\otimes i-1} \otimes C_{V,V} \otimes \text{id}^{\otimes n-i-1}$. Let $S_m \in \text{End}_{\mathbf{k}}(V^{\otimes m})$ and $S_{1,j} \in \text{End}_{\mathbf{k}}(V^{\otimes(j+1)})$ denote the maps $S_m = \prod_{j=1}^{m-1} (\text{id}^{\otimes m-j-1} \otimes S_{1,j})$, $S_{1,j} = \text{id} + C_{12}^{-1} + C_{12}^{-1}C_{23}^{-1} + \cdots + C_{12}^{-1}C_{23}^{-1} \cdots C_{j,j+1}^{-1}$

(in leg notation) for $m \geq 2$ and $j \in \mathbb{N}$. Then the subspace $S = \bigoplus_{m=2}^{\infty} \ker S_m$ of the tensor

$T(V) = \bigoplus_{m=0}^{\infty} V^{\otimes m}$ is a two-sided ideal, and algebra $\mathfrak{B}(V) = T(V)/S$ is termed the Nichols algebra associated to (V, C) .

For $s \in G$ and $(\rho, V) \in \widehat{G^s}$, here is a precise description of the YD module $M(\mathcal{O}_s, \rho)$, introduced in [Gr00]. Let $t_1 = s, t_2, \dots, t_m$ be a numeration of \mathcal{O}_s , which is a conjugacy class containing s , and let $g_i \in G$ such that $g_i \triangleright s := g_i s g_i^{-1} = t_i$ for all $1 \leq i \leq m$. Then $M(\mathcal{O}_s, \rho) = \bigoplus_{1 \leq i \leq m} g_i \otimes V$. Let $g_i v := g_i \otimes v \in M(\mathcal{O}_s, \rho)$, $1 \leq i \leq m$, $v \in V$. If $v \in V$ and $1 \leq i \leq m$, then the action of $h \in G$ and the coaction are given by

$$\delta(g_i v) = t_i \otimes g_i v, \quad h \cdot (g_i v) = g_j (\nu_i(h) \cdot v), \quad (1)$$

where $h g_i = g_j \nu_i(h)$, for unique $1 \leq j \leq m$ and $\nu_i(h) \in G^s$. The explicit formula for the braiding is then given by

$$C(g_i v \otimes g_j w) = t_i \cdot (g_j w) \otimes g_i v = g_{j'} (\nu_j(t_i) \cdot w) \otimes g_i v \quad (2)$$

for any $1 \leq i, j \leq m$, $v, w \in V$, where $t_i g_j = g_{j'} \nu_j(t_i)$ for unique j' , $1 \leq j' \leq m$ and $\nu_j(t_i) \in G^s$. Let $\mathfrak{B}(\mathcal{O}_s, \rho)$ denote $\mathfrak{B}(M(\mathcal{O}_s, \rho))$. $M(\mathcal{O}_s, \rho)$ is a simple YD module (see [DPR91, Ci97, AZ07]).

We briefly recall the definition and main properties of racks; see [AG03] for details, more information and bibliographical references. A rack is a pair (X, \triangleright) , where X is a non-empty set and $\triangleright : X \times X \rightarrow X$ is an operation such that $x \triangleright x = x$, $x \triangleright (y \triangleright z) = (x \triangleright y) \triangleright (x \triangleright z)$ and ϕ_x is invertible for any $x, y, z \in X$, where ϕ_x is a map from X to X sending y to $x \triangleright y$ for any $x, y \in X$. For example, $(\mathcal{O}_s^G, \triangleright)$ is a rack with $x \triangleright y := xyx^{-1}$.

If R and S are two subracks of X with $R \cup S = X$, $R \cap S = \emptyset$, $x \triangleright y \in S$, $y \triangleright x \in R$, for any $x \in R, y \in S$, then $R \cup S$ is called a decomposition of subracks of X . Furthermore, if there exist $a \in R, b \in S$ such that $\text{sq}(a, b) := a \triangleright (b \triangleright (a \triangleright b)) \neq b$, then X is called to be of type D. Notice that if a rack Y contains a subrack X of type D, then Y is also of type D (see [AFGV08]).

2 Conjugacy classes of juxtapositions

In this section we determine when the conjugacy classes of juxtapositions of two elements are of type D.

Lemma 2.1. *Let $G = \mathbb{W}_n$ and $(a, \tau), (b, \mu) \in G$. Let (c, λ) denote $(a, \tau) \triangleright ((b, \mu) \triangleright ((a, \tau) \triangleright (b, \mu)))$.*

(i) Then

$$\begin{aligned} c &= (a + \tau \cdot [b + \mu \cdot (a + \tau \cdot b + (\tau \triangleright \mu) \cdot a) + (\mu \triangleright (\tau \triangleright \mu)) \cdot b] \\ &\quad + (\tau \triangleright (\mu \triangleright (\tau \triangleright \mu))) \cdot a. \end{aligned} \quad (3)$$

(ii) If τ and μ are commutative, then

$$c = a + \tau\mu \cdot a + \tau\mu^2 \cdot a + \mu \cdot a + \tau \cdot b + \tau^2\mu \cdot b + \tau\mu \cdot b. \quad (4)$$

(iii) If τ and μ are commutative, then $\text{sq}((a, \tau), (b, \mu)) = (b, \mu)$ if and only if

$$a + \tau\mu \cdot a + \tau\mu^2 \cdot a + \mu \cdot a = b + \tau \cdot b + \tau^2\mu \cdot b + \tau\mu \cdot b. \quad (5)$$

(iv) If τ and μ are commutative with $(b, \mu) = \xi \triangleright (a, \tau)$ and $\xi \cdot a = a$, then $c = a + \tau\mu^2 \cdot a + \mu \cdot a + \tau \cdot a + \tau^2\mu \cdot a$.

(v) If τ and ξ are commutative with $\tau^2 = 1$ and $(b, \mu) = \xi \triangleright (a, \tau)$ and $\xi \cdot a = a$, then $c = a$.

Proof. It is clear. \square

If the lengths of independent cycles of π and τ are different, then they are called mutually orthogonal, written as $\pi \perp \tau$. If the lengths of independent sign cycles of (a, π) and (b, τ) are different, then they are called mutually orthogonal, written as $a\pi \perp b\tau$. Obviously, $a\pi \perp b\tau$ if and only if $\pi \perp \tau$. If $\text{ord}(\pi)$ and $\text{ord}(\tau)$ are coprime and one of the two elements does not have any fixed point, then $\pi \perp \tau$. Notice that if $\pi \perp \tau$, then $\mathbb{S}_{m+n}^{\pi\#\tau} = \mathbb{S}_n^\pi \# \mathbb{S}_m^\tau$.

For any $a\pi \in \mathbb{W}_n$ and $b\tau \in \mathbb{W}_m$, define $a\pi \# b\tau \in \mathbb{W}_{m+n}$ as follows:

$$(a\#b)_i := \begin{cases} a_i & \text{when } i \leq n \\ b_{i-n} & \text{when } i > n \end{cases}, \quad (\pi\#\tau)(i) := \begin{cases} \pi(i) & \text{when } i \leq n \\ \tau(i-n) + n & \text{when } i > n \end{cases} \quad \text{and}$$

$a\pi \# b\tau := (a\#b, \pi\#\tau)$. Obviously $a\pi \# b\tau \in \mathbb{W}_{m+n}$ and it is called a juxtaposition of $a\pi$ and $b\tau$ (see [AZ07]). Let $\overrightarrow{\nu_{n,m}}$ be a map from \mathbb{W}_n to \mathbb{W}_{m+n} by sending $a\pi$ to $\overrightarrow{\nu_{n,m}}(a\pi) := a\pi \# 1_{\mathbb{W}_m}$; let $\overleftarrow{\nu_{n,m}}$ be a map from \mathbb{W}_m to \mathbb{W}_{m+n} by sending $b\tau$ to $\overleftarrow{\nu_{n,m}}(b\tau) := 1_{\mathbb{W}_n} \# b\tau$.

Lemma 2.2. Assume $a\pi \perp b\tau$ with $a\pi, a'\pi' \in \mathbb{W}_n$ and $b\tau, b'\tau' \in \mathbb{W}_m$. Then

- (i) $(a\pi \# b\tau)(a'\pi' \# b'\tau') = (a\pi a'\pi' \# b\tau b'\tau')$.
- (ii) $a\pi \# b\tau = \overrightarrow{\nu_{n,m}}(a\pi) \overleftarrow{\nu_{n,m}}(b\tau) = \overleftarrow{\nu_{n,m}}(b\tau) \overrightarrow{\nu_{n,m}}(a\pi)$.
- (iii) $\mathbb{W}_{m+n}^{a\pi \# b\tau} = \mathbb{W}_n^{a\pi} \# \mathbb{W}_m^{b\tau} = \overrightarrow{\nu_{n,m}}(\mathbb{W}_n^{a\pi}) \overleftarrow{\nu_{n,m}}(\mathbb{W}_m^{b\tau})$ as directed products.
- (iv) For any $\rho \in \widehat{\mathbb{W}_{m+n}^{a\pi \# b\tau}}$, there exist $\rho_1 \in \widehat{\mathbb{W}_n^{a\pi}}$, $\rho_2 \in \widehat{\mathbb{W}_m^{b\tau}}$ such that $\rho = \rho_1 \otimes \rho_2$.
- (v) $(a\pi \# b\tau) \triangleright (a'\pi' \# b'\tau') = (a\pi \triangleright a'\pi') \# (b\tau \triangleright b'\tau')$.
- (vi) $\mathcal{O}_{a\pi \# b\tau}^{\mathbb{W}_{m+n}} = \mathcal{O}_{a\pi}^{\mathbb{W}_n} \# \mathcal{O}_{b\tau}^{\mathbb{W}_m}$.

Proof. (i), (ii) and (v) are clear.

(iii) By [AZ07, Section 2.2], $\mathbb{S}_{m+n}^{\pi\#\tau} = \mathbb{S}_n^\pi \# \mathbb{S}_m^\tau$. Obviously, $\mathbb{W}_n^{a\pi} \# \mathbb{W}_m^{b\tau} \subseteq \mathbb{W}_{m+n}^{a\pi \# b\tau}$. For any $c\xi \in \mathbb{W}_{m+n}^{a\pi \# b\tau}$, then $\xi \in \mathbb{S}_{m+n}^{\pi\#\tau}$ and there exist $\mu \in \mathbb{S}_n^\pi$ and $\lambda \in \mathbb{S}_m^\tau$ such that $\xi = \mu \# \lambda$.

Consequently, $c\xi = d\mu \# f\lambda$. Considering $c\xi(a\pi \# b\tau) = (a\pi \# b\tau)c\xi$, we have $d\mu \in \mathbb{W}_n^{a\pi}$ and $f\lambda \in \mathbb{W}_m^{b\tau}$. This completes the proof.

(iv) It follows from (iii).

(vi) By (v), $\mathcal{O}_{a\pi}^{\mathbb{W}_n} \# \mathcal{O}_{b\tau}^{\mathbb{W}_m} \subseteq \mathcal{O}_{a\pi \# b\tau}^{\mathbb{W}_{m+n}}$. Consequently (vi) follows from (iii). \square

Remark 2.3. (i), (ii) and (v) above still hold when $a\pi$ and $b\tau$ are not mutually orthogonal.

Theorem 2.4. If $\mathcal{O}_{a\tau}$ is of type D, then $\mathcal{O}_{a\tau \# b\mu}$ is also of type D.

Proof. Let $X = R \cup S$ be a subrack decomposition of $\mathcal{O}_{a\tau}$ and of type D. It is clear that $X \# b\mu = R \# b\mu \cup S \# b\mu$ is a subrack decomposition of $\mathcal{O}_{a\tau \# b\mu}$ and of type D. \square

Lemma 2.5. Assume $a\pi \perp b\tau$. Let $\rho = \rho_1 \otimes \rho_2 \in \widehat{\mathbb{W}_{m+n}^{a\pi \# b\tau}}$, $\rho_1 \in \widehat{\mathbb{W}_n^{a\pi}}$, $\rho_2 \in \widehat{\mathbb{W}_m^{b\tau}}$, $a\pi \in \mathbb{W}_n$ and $b\tau \in \mathbb{W}_m$ with $q_{a\pi, a\pi} \text{id} = \rho_1(a\pi)$ and $q_{b\tau, b\tau} \text{id} = \rho_2(b\tau)$. If $\dim \mathfrak{B}(\mathcal{O}_{a\pi \# b\tau}^{\mathbb{W}_{n+m}}, \rho) < \infty$, then

(i) $M(\mathcal{O}_{a\pi}^{\mathbb{W}_n}, \rho_1)$ is isomorphic to a YD submodule of $M(\mathcal{O}_{a\pi \# b\tau}^{\mathbb{W}_{n+m}}, \rho)$ over \mathbb{W}_n when $q_{b\tau, b\tau} = 1$; hence $\dim \mathfrak{B}(\mathcal{O}_{a\pi}^{\mathbb{W}_n}, \rho_1) < \infty$.

(ii) $q_{a\pi, a\pi} q_{b\tau, b\tau} = -1$.

(iii) $q_{b\tau, b\tau} = 1$ and $q_{a\pi, a\pi} = -1$ when $\text{ord}(b\tau) \leq 2$ and $\text{ord}(q_{a\pi, a\pi}) \neq 1$.

(iv) $q_{b\tau, b\tau} = 1$ and $q_{a\pi, a\pi} = -1$ when $\text{ord}(a\pi)$ and $\text{ord}(b\tau)$ are coprime and $\text{ord}(b\tau)$ is odd.

Proof. (i) The proof is similar to the one in [AZ07, Section 2.2]. Indeed, Let $t_1 = \pi, t_2, \dots, t_{|\mathcal{O}_\pi|}$ be a numeration of \mathcal{O}_π , and let $g_i \in G$ such that $g_i \triangleright \pi := g_i \pi g_i^{-1} = t_i$ for all $1 \leq i \leq |\mathcal{O}_\pi|$. Let $s_1 = \tau, s_2, \dots, s_{|\mathcal{O}_\tau|}$ be a numeration of \mathcal{O}_τ , and let $h_i \in G$ such that $h_i \triangleright \tau := h_i \tau h_i^{-1} = s_i$ for all $1 \leq i \leq |\mathcal{O}_\tau|$. Thus $\{t_i \# s_j \mid 1 \leq i \leq |\mathcal{O}_\pi|, 1 \leq j \leq |\mathcal{O}_\tau|\} = \mathcal{O}_{\pi \# \tau}$. Let V and W be representation spaces of ρ_1 and ρ_2 , respectively. Let $0 \neq w_0 \in W$. Define a map ψ from $M(\mathcal{O}_\pi^{\mathbb{W}_n}, \rho_1)$ to $M(\mathcal{O}_{\pi \# \tau}^{\mathbb{W}_m \# \mathbb{W}_n}, \rho_1 \otimes \rho_2)$ by sending $g_i v$ to $(g_i \# h_1)v \otimes w_0$ for any $v \in V$, $1 \leq i \leq |\mathcal{O}_\pi|$. It is clear that ψ is injective. Now we show that ψ is a homomorphism of braided vector spaces. For any $v, v' \in V$, we have that

$$\begin{aligned} (\psi \otimes \psi)(C(g_i v \otimes g_j v')) &= (\psi \otimes \psi)(g_{i \triangleright j} \rho_1(\nu_j(t_i))(v') \otimes g_i v) \\ &= (g_{i \triangleright j} \# h_1 \rho_1(\nu_j(t_i))(v') \otimes w_0) \otimes (g_i \# h_1 v \otimes w_0) \quad \text{and} \end{aligned}$$

$$\begin{aligned} C(\psi(g_i v) \otimes \psi(g_j v')) &= C((g_i \# h_1 v \otimes w_0) \otimes (g_j \# h_1 v' \otimes w_0)) \\ &= (g_{i \triangleright j} \# h_1 (\rho_1 \otimes \rho_2)(\nu_{j,1}(t_i \# s_1))(v' \otimes w_0)) \otimes (g_i \# h_1 v \otimes w_0). \end{aligned}$$

It is clear that $\nu_{j,1}(t_i \# h_1) = \nu_j(t_i) \# h_1$. Therefore ψ is a homomorphism of braided vector spaces.

(ii) and (iii) are clear.

(iv) Obviously, $\text{ord}(q_{a\pi, a\pi}) \mid \text{ord}(a\pi)$ and $\text{ord}(q_{b\tau, b\tau}) \mid \text{ord}(b\tau)$. Therefore, $\text{ord}(q_{b\tau, b\tau})$ is odd. The least common factor $(\text{ord}(q_{a\pi, a\pi}), \text{ord}(q_{b\tau, b\tau})) = 1$ since $(\text{ord}(a\pi), \text{ord}(b\tau)) = 1$. By Part (ii), $\text{ord}(q_{a\pi, a\pi})\text{ord}(q_{b\tau, b\tau}) = 2$. Consequently, $\text{ord}(q_{a\pi, a\pi}) = 2$ and $\text{ord}(q_{b\tau, b\tau}) = 1$. \square

3 Conjugacy classes of \mathbb{W}_n

In this section we prove that except in several cases conjugacy classes of classical Weyl groups \mathbb{W}_n are of type D.

Lemma 3.1. *Let p be odd with $p \geq 5$ and $a\tau \in \mathbb{W}_n$ with $\tau = (1\ 2 \cdots p)$. Then $\mathcal{O}_{a\tau}^{\mathbb{W}_n}$ is of type D.*

Proof. (i) Assume that $a\tau$ is a negative cycle (defined in [ZZ12, Appendix]) and $b = (1, 0, \dots, 0, a_{p+1}, a_{p+2}, \dots, a_n)$. Thus $a\tau$ and $b\tau$ are conjugate. We assume $a = (1, 1, \dots, 1, a_{p+1}, a_{p+2}, \dots, a_n)$ without lost generality. Obviously, the right hand side of (5) is non-vanishing for $\mu = \tau^2$, i.e. $\text{sq}(a\tau, b\tau^2) \neq b\tau^2$. Let $R := \mathbb{Z}_2^n \rtimes \tau \cap \mathcal{O}_{a\tau}^{\mathbb{W}_n}$ and $S := \mathbb{Z}_2^n \rtimes \tau^2 \cap \mathcal{O}_{a\tau}^{\mathbb{W}_n}$. It is clear that $R \cup S$ is a subrack decomposition of $\mathcal{O}_{a\tau}^{\mathbb{W}_n}$, hence it is of type D; notice that τ and τ^2 are conjugate in \mathbb{S}_n .

(ii) Assume that $a\tau$ is a positive cycle. Let $b = (1, 0, 0, 1, 0, a_{p+1}, a_{p+2}, \dots, a_n)$ when $p = 5$; $b = (1, 1, 0, \dots, 0, a_{p+1}, a_{p+2}, \dots, a_n)$ when $p > 5$. Thus 0τ , $a\tau$ and $b\tau$ are conjugate. We assume $a = 0$ without lost generality. It is clear that the right hand side of (5) is not equal to 0. Consequently, $R \cup S$ is a subrack decomposition and is of type D as Part (i). \square

Lemma 3.2. *If σ is of type (3^2) , then $\mathcal{O}_{a\sigma}^{\mathbb{W}_6}$ is of type D for all $a \in \mathbb{Z}_2^6$.*

Proof. Let $\pi = (1\ 2\ 3)$, $\xi = (4\ 5\ 6)$, $\tau = (1\ 2\ 3)(4\ 5\ 6)$ and $\mu = (1\ 2\ 3)^2(4\ 5\ 6)$. We have that the 4-th, 5-th and 6-th components of (5) are

$$(a_6 + a_5, a_4 + a_6, a_5 + a_4) = (b_6 + b_5, b_4 + b_6, b_5 + b_4). \quad (6)$$

By (5), (i) If $a = (1, 1, \dots, 1)$ and $b = (1, 0, 0, 1, 0, 0)$, then (6) does not hold.

(ii) If $a = 0$ and $b = (0, 0, 0, 1, 1, 0)$, then (6) does not hold.

(iii) If $a = (1, 0, 0, 0, 0, 0)$ and $b = (1, 0, 0, 1, 1, 0)$, then (6) does not hold.

(iv) If $a = (0, 0, 0, 1, 0, 0)$ and $b = (0, 0, 0, 0, 1, 0)$, then (6) does not hold. Let $R := \mathbb{Z}_2^6 \rtimes \tau \cap \mathcal{O}_{a\sigma}^{\mathbb{W}_6}$ and $S := \mathbb{Z}_2^6 \rtimes \mu \cap \mathcal{O}_{a\sigma}^{\mathbb{W}_6}$. It is clear that $R \cup S$ is a subrack decomposition of $\mathcal{O}_{a\sigma}^{\mathbb{W}_6}$. Consequently $\mathcal{O}_{a\sigma}^{\mathbb{W}_6}$ is of type D. \square

Lemma 3.3. *If σ is of type $(2^2, 3^1)$, then $\mathcal{O}_{a\sigma}^{\mathbb{W}_7}$ is of type D for all $a \in \mathbb{Z}_2^7$.*

Proof. Let $\pi = (5\ 6\ 7)$, $\xi = (1\ 2)(3\ 4)$, $\lambda = (1\ 3)(2\ 4)$, $\tau = \pi\xi$ and $\mu = \pi\lambda$. If $a = (a_1, a_2, a_3, a_4, 0, 0, 0)$, let $b = (b_1, b_2, b_3, b_4, 1, 1, 0)$. If $a = (a_1, a_2, a_3, a_4, 1, 1, 1)$, let $b = (b_1, b_2, b_3, b_4, 1, 0, 0)$. Then the 5-th, 6-th and 7-th components of (5) are $(a_6 + a_7, a_7 + a_5, a_5 + a_6) = (b_6 + b_7, b_7 + b_5, b_5 + b_6)$, respectively. Consequently, (5) does not hold.

Let $R := \mathbb{Z}_2^7 \rtimes \tau \cap \mathcal{O}_{a\sigma}^{\mathbb{W}_7}$ and $S := \mathbb{Z}_2^7 \rtimes \mu \cap \mathcal{O}_{a\sigma}^{\mathbb{W}_7}$. It is clear that $R \cup S$ is a subrack decomposition of $\mathcal{O}_{a\sigma}^{\mathbb{W}_7}$. Consequently $\mathcal{O}_{a\sigma}^{\mathbb{W}_7}$ is of type D. \square

Example 3.4. If τ and μ are of type (2^2) , then $a\tau$ and $b\mu$ are square commutative when they are conjugate to each other or $(-1)^{\sum_i^4 a_i} = (-1)^{\sum_i^4 b_i}$.

Proof. Let $\tau = (1\ 2)(3\ 4)$. Then $\mu = (1\ 3)(2\ 4)$ or $\mu = \tau$. It is clear that equation (5) becomes $a_1 + a_2 + a_3 + a_4 = b_1 + b_2 + b_3 + b_4$. \square

Lemma 3.5. (i) Assume that τ and μ are conjugate with $\text{sq}(\tau, \mu) \neq \mu$ in \mathbb{S}_n and $\tau(n) = \mu(n) = n$. If $a \in \mathbb{Z}_2^n$ and there exists i such that $a_i \neq a_n$ with $\tau(i) = \mu(i) = i$, then $\mathcal{O}_{a\tau}$ is of type D.

(ii) Assume that $n > 4$ and $a\tau \in \mathbb{W}_n$ with type $(1^{n-2}, 2)$ of τ . If there exist i, j such that $\tau(i) = i$ and $\tau(j) = j$ with $a_i \neq a_j$, then $\mathcal{O}_{a\tau}$ is of type D.

(iii) Assume that $n > 5$ and $a\tau \in \mathbb{W}_n$ with type $(1^{n-3}, 3)$ of τ . If there exist i, j such that $\tau(i) = i$ and $\tau(j) = j$ with $a_i \neq a_j$, then $\mathcal{O}_{a\tau}$ is of type D.

Proof. (i) Let $R := \{d\xi \in \mathcal{O}_{a\tau}^{\mathbb{W}_n} \mid \xi(n) = n; d_n = 0\}$ and $S := \{d\xi \in \mathcal{O}_{a\tau}^{\mathbb{W}_n} \mid \xi(n) = n; d_n = 1\}$. Obviously, $R \cup S$ is a subrack decomposition. Let $\xi \in \mathbb{S}_n$ with $\xi(i) = i$ and $\xi(n) = n$ such that $\xi \triangleright \tau = \mu$. By simple computation we have $(i, n)\xi \triangleright (a\tau) = b\mu$ with $b_n = a_i$. Consequently, $a\tau$ and $b\mu$ are in the same set of R and S , which implies that $R \cup S$ is of type D.

(ii) It is clear that $\text{sq}(\tau, \mu) \neq \mu$ with $\tau := (1, 2)$ and $\mu := (2, 3)$. Applying Part (i) we complete the proof.

(iii) It is clear that $\text{sq}(\tau, \mu) \neq \mu$ with $\tau = (1, 2, 3)$ and $\mu = (2, 4, 3)$. Applying Part (i) we complete the proof. \square

Lemma 3.6. Let $\sigma = (a, \tau) \in \mathbb{W}_n$. If $\mathcal{O}_\tau^{\mathbb{S}_n}$ is of type D, then so is $\mathcal{O}_{(a, \tau)}^{\mathbb{W}_n}$.

Proof. Let $X = S \cup T$ be a subrack decomposition of $\mathcal{O}_\tau^{\mathbb{S}_n}$ and $s \in S, t \in T$ such that

$$s \triangleright (t \triangleright (s \triangleright t)) \neq t. \quad (7)$$

Let $h \triangleright \tau = s$ and $g \triangleright \tau = t$ with $h, g \in \mathbb{S}_n$. It is clear $\text{sq}((h \cdot a, s), (g \cdot a, t)) \neq (g \cdot a, t)$ since (7); $(h \cdot a, s) = h \triangleright (a, \tau)$ and $(g \cdot a, t) = g \triangleright (a, \tau)$.

$\langle \mathbb{S}_n \cdot a \rangle, X$ is a subrack, where $\langle \mathbb{S}_n \cdot a \rangle$ is the subgroup generated by subset $\mathbb{S}_n \cdot a$ of \mathbb{Z}_2^n . In fact, for any $h, g \in \mathbb{S}_n, \xi, \mu \in X$, we have

$$(h \cdot a, \xi) \triangleright (g \cdot a, \mu) = ((h + \xi g + \xi \mu \xi^{-1} h) \cdot a, \xi \triangleright \mu) \in \langle \mathbb{S}_n \cdot a \rangle, X.$$

Thus $(\langle \mathbb{S}_n \cdot a \rangle, X)$ is a subrack. Consequently $(\langle \mathbb{S}_n \cdot a \rangle, X) \cap \mathcal{O}_{(a,\tau)}^{\mathbb{W}_n} = (\langle \mathbb{S}_n \cdot a \rangle, S) \cap \mathcal{O}_{(a,\tau)}^{\mathbb{W}_n} \cup (\langle \mathbb{S}_n \cdot a \rangle, T) \cap \mathcal{O}_{(a,\tau)}^{\mathbb{W}_n}$ is a subrack decomposition of $\mathcal{O}_{(a,\tau)}^{\mathbb{W}_n}$ and is of type D. \square

Theorem 3.7. *Let $G = \mathbb{W}_n$ with $n > 4$. Let $\tau \in \mathbb{S}_n$ be of type $(1^{\lambda_1}, 2^{\lambda_2}, \dots, n^{\lambda_n})$ and $a \in \mathbb{Z}_2^n$ with $\sigma = (a, \tau) \in G$ and $\tau \neq 1$. If \mathcal{O}_σ^G is not of type D, then the type of τ belongs to one in the following list.*

- (i) $(2, 3); (2^3);$
- (ii) $(2^4); (1, 2^2), (1^2, 3), (1^2, 2^2);$
- (iii) $(1^{n-2}, 2)$ and $(1^{n-3}, 3)$ ($n > 5$) with $a_i = a_j$ when $\tau(i) = i$ and $\tau(j) = j$.

Proof. It follows from Lemma 3.3, Lemma 3.5, Lemma 3.6, Lemma 3.1, Lemma 3.2 and [AFGV08, Theorem 4.1]. \square

4 Nichols algebras of irreducible YD module over \mathbb{W}_n

In this section we show that except in three cases Nichols algebras of irreducible YD modules over classical Weyl groups \mathbb{W}_n are infinite dimensional.

Let $\text{supp}M := \{g \in G \mid M_g \neq 0\}$ for G -comodule (M, δ) , where $M_g := \{x \in M \mid \delta(x) = g \otimes x\}$.

We shall use the following facts:

Theorem 4.1. ([HS08, Cor. 8.4]) *Let $n \in \mathbb{N}$, $n \geq 3$, and assume that $G = \mathbb{S}_n$ is the symmetric group. Let U be a YD module over G . If $\mathfrak{B}(U)$ is finite dimensional, then U is an irreducible YD module over G .*

Theorem 4.2. ([AFGV08, Th. 1.1]) *Let $m \geq 5$. Let $\sigma \in \mathbb{S}_m$ be of type $(1^{n_1}, 2^{n_2}, \dots, m^{n_m})$, let \mathcal{O}_σ be the conjugacy class of σ and let $\rho = (\rho, V) \in \widehat{S_m^\sigma}$. If $\dim \mathfrak{B}(\mathcal{O}_\sigma, \rho) < \infty$, then the type of σ and ρ are in the following list:*

- (i) $(1^{n_1}, 2)$, $\rho_1 = \text{sgn}$ or ϵ , $\rho_2 = \text{sgn}$.
- (ii) $(2, 3)$ in \mathbb{S}_5 , $\rho_2 = \text{sgn}$, $\rho_3 = \vec{\chi}_0$.
- (iii) (2^3) in \mathbb{S}_6 , $\rho_2 = \vec{\chi}_1 \otimes \epsilon$ or $\vec{\chi}_1 \otimes \text{sgn}$.

Theorem 4.3. *If G is a finite group and \mathcal{O}_σ^G is of type D, then $\dim \mathfrak{B}(\mathcal{O}_\sigma^G, \rho) = \infty$ for any $\rho \in \widehat{G^\sigma}$.*

Proof. It follows from [AFGV08, Theorem 3.6]. \square

Theorem 4.4. Assume $n > 4$. Let $\tau \in \mathbb{S}_n$ be of type $(1^{\lambda_1}, 2^{\lambda_2}, \dots, n^{\lambda_n})$ and $a \in \mathbb{Z}_2^n$ with $\sigma = (a, \tau) \in \mathbb{W}_n$ and $\tau \neq 1$. If $\dim \mathfrak{B}(\mathcal{O}_\sigma^{\mathbb{W}_n}, \rho) < \infty$, then the type of τ belongs to one in the following list:

- (i) $(2, 3); (2^3);$
- (ii) $(2^4); (1, 2^2), (1^2, 3), (1^2, 2^2);$
- (iii) $(1^{n-2}, 2)$ and $(1^{n-3}, 3)$ ($n > 5$) with $a_i = a_j$ when $\tau(i) = i$ and $\tau(j) = j$.

Proof. It follows from Theorem 3.7 and Theorem 4.3. \square

Let $a\mu = c\tau \# d\xi \in \mathbb{W}_n$ with $c\tau \in \mathbb{W}_m$, $d\xi \in \mathbb{B}_{n-m}$ and $c\tau \perp d\xi$. Let $\rho = \rho_1 \otimes \rho_2 \in \widehat{\mathbb{W}_n^{a\mu}} = \widehat{\mathbb{W}_m^{c\tau}} \times \widehat{\mathbb{W}_{n-m}^{d\xi}}$. If $c = (1, 1, \dots, 1)$, then $\rho_1 = (\chi_1 \otimes \mu_1) \uparrow_{G_{\chi_1}^{a\mu}}^{G_{\chi_1}^{a\mu}}$ with $\chi_1 \in \widehat{(Z_2^m)^\tau}$, $\mu_1 \in \widehat{(\mathbb{S}_m^\tau)_{\chi_1}}$ (see [ZZ12, previous Proposition 2.9 in Section 2.5]), where $G = \mathbb{W}_n$. Case $a = 0$ and $a = (1, 1, \dots, 1)$ were studied in paper [ZZ12, Theorem 1.1 and Table 1]. Other cases are listed as follows:

Corollary 4.5. Under notation above assume $\dim \mathfrak{B}(\mathcal{O}_{a\mu}^{\mathbb{W}_n}, \rho) < \infty$ with $d_1 = d_2 = \dots = d_{n-m}$ and $\xi = \text{id}^{\otimes(n-m)}$.

- (i) Then $\rho_1(c\tau) = -\text{id}$ when $\rho_2(d\xi) = \text{id}$ and $\rho_1(c\tau) = \text{id}$ when $\rho_2(d\xi) = -\text{id}$.
- (ii) Case $\tau = (1\ 2)$, $c = 0$ and $d = (1, 1, \dots, 1)$. Then $\rho_1(c\tau) = \pm \text{id}$, $\rho_2(d\xi) = \mp \text{id}$, $\chi_1(c) = 1$ and $\mu_1(\tau) = \pm 1$.
- (iii) Case $\tau = (1\ 2)$, $c = (1, 1)$ and $d = 0$. Then $\rho_1(c\tau) = -\text{id}$, $\rho_2(d\xi) = \text{id}$.
- (iv) Case $\tau = (1\ 2\ 3)$, $c = 0$ and $d = (1, 1, \dots, 1)$. Then $\rho_1(c\tau) = \text{id}$, $\rho_2(d\xi) = -\text{id}$, $\chi_1(c) = 1$ and $\mu_1(\tau) = 1$.
- (v) Case $\tau = (1\ 2\ 3)$, $c = (1, 1, 1)$ and $d = 0$. Then $\rho_1(c\tau) = -\text{id}$, $\rho_2(d\xi) = \text{id}$, $\chi_1(c) = -1$ and $\mu_1(\tau) = 1$.
- (vi) Case $\tau = (1\ 2)(3\ 4)$, $c = 0$ and $d = (1, 1)$. Then $\rho_1(c\tau) = \text{id}$, $\rho_2(d\xi) = -\text{id}$.
- (vii) Case $\tau = (1\ 2)(3\ 4)$, $c = (1, 0, 1, 0)$ and $d = (1, 1)$. Then $\rho_1(c\tau) = \pm \text{id}$, $\rho_2(d\xi) = \mp \text{id}$.
- (viii) Case $\tau = (1\ 2)(3\ 4)$, $c = (1, 0, 1, 0)$ and $d = 0$. Then $\rho_1(c\tau) = -\text{id}$, $\rho_2(d\xi) = \text{id}$.
- (ix) Case $\tau = (1\ 2)(3\ 4)$, $c = (1, 0, 0, 0)$ and $d = (1, 1)$. Then $\rho_1(c\tau) = \pm \text{id}$, $\rho_2(d\xi) = \mp \text{id}$.
- (x) Case $\tau = (1\ 2)(3\ 4)$, $c = (1, 0, 0, 0)$ and $d = (0, 0)$. Then $\rho_1(c\tau) = -\text{id}$, $\rho_2(d\xi) = \text{id}$.

Proof. (iv) If $\rho_2(d\xi) = \text{id}$, then $\dim \mathfrak{B}(\mathcal{O}_{c\tau}^{\mathbb{W}_3}, \rho_1) < \infty$ by Lemma 2.5, which constructs to [ZZ12, Theorem 1.1].

(v) If $\chi_1(c) = 1$, then there exists a contradiction by [ZZ12, Proposition 2.4, Theorem 1.1].

The others follow from Lemma 2.5 and [ZZ12, Theorem 1.1]. \square

Appendix

In this section we give the general FK' Conjecture and another proof of Theorem 4.3.

A General FK' Conjecture.

$\mathfrak{B}(\mathcal{O}_\sigma, \rho)$ is finite dimensional when $n \leq 5$ according to [MS, FK99, AZ07]. However, it is open whether $\mathfrak{B}(\mathcal{O}_\sigma, \rho)$ is finite dimensional when $n > 5$. We know the relationship between FK' Conjecture and Nichols algebra $\mathfrak{B}(\mathcal{O}_{(1,2)}, \epsilon \otimes \text{sgn})$ of transposition over symmetry group (see [MS, Theorem 5.7]). That is, if $\dim \mathfrak{B}(\mathcal{O}_{(1,2)}, \epsilon \otimes \text{sgn}) = \infty$, then so is $\dim \mathcal{E}_n$.

Definition A.1. (See [FK99, Def. 2.1]) algebra \mathcal{E}_n is generated by $\{x_{ij} \mid 1 \leq i < j \leq n\}$ with definition relations:

- (i) $x_{ij}^2 = 0$ for $i < j$.
- (ii) $x_{ij}x_{jk} = x_{jk}x_{ik} + x_{ik}x_{ij}$ and $x_{jk}x_{ij} = x_{ik}x_{jk} + x_{ij}x_{ik}$, for $i < j < k$.
- (iii) $x_{ij}x_{kl} = x_{kl}x_{ij}$ for any distinct i, j, k, l and $l, i < j, k < l$. Equivalently, algebra \mathcal{E}_n is generated by $\{x_{ij} \mid i \neq j, 1 \leq i, j \leq n\}$ with definition relations:

- (i) $x_{ij}^2 = 0, x_{ij} = -x_{ji}$;
 - (ii) $x_{ij}x_{jk} + x_{jk}x_{ki} + x_{ki}x_{ij} = 0$;
 - (iii) $x_{ij}x_{kl} = x_{kl}x_{ij}$;
- for any distinct i, j, k and l .

S. Fomin and A.N. Kirillov [FK99, Conjecture 2.2] point out a conjecture “ \mathcal{E}_n is finite dimensional”. Consequently, we have

Problem A.2. Let $\alpha_{ijk}, \beta_{ijk}, \gamma_{ij}, \lambda_{ijkl} \in \{1, -1\}$ for any distinct i, j, k and l with $1 \leq i, j, k, l \leq n$. Assume that algebra $A(\alpha, \beta, \gamma, \lambda)$ is generated by $\{x_{ij} \mid i \neq j, 1 \leq i, j \leq n\}$ with definition relations:

- (i) $x_{ij}^2 = 0, x_{ij} = \gamma_{ij}x_{ji}$;
 - (ii) $x_{ij}x_{jk} + \alpha_{ijk}x_{jk}x_{ki} + \beta_{ijk}x_{ki}x_{ij} = 0$;
 - (iii) $x_{ij}x_{kl} = \lambda_{ijkl}x_{kl}x_{ij}$;
- for any distinct i, j, k and l .

Then $A(\alpha, \beta, \gamma, \lambda)$ is finite dimensional when $n > 5$.

Obviously, $\mathcal{E}_n = A(1, 1, -1, 1)$, i.e. $\mathcal{E}_n = A(\alpha, \beta, \gamma, \lambda)$ with $\alpha_{i,j,k} = 1, \beta_{i,j,k} = 1, \gamma_{ij} = -1, \lambda_{i,j,k,l} = 1$ for any distinct i, j, k and l .

B Proof of Theorem 4.3.

Proof. (i) There exists subrack $X \subseteq \mathcal{O}_\sigma^G$. $R \cup S = X$ is a subrack decomposition of X there exists $a \in R$, $b \in S$ such that $\text{sq}(a, b) := a \triangleright (b \triangleright (a \triangleright b)) \neq b$. Let H be the subgroup generated by X in G . It is clear $\mathcal{O}_{\sigma_1}^H \neq \mathcal{O}_{\sigma_2}^H$ for any $\sigma_1 \in R, \sigma_2 \in S$. In fact, if $\mathcal{O}_{\sigma_1}^H = \mathcal{O}_{\sigma_2}^H$, then there exists $t \in H$, such that $\sigma_2 = t\sigma_1 t^{-1}$. $t = \prod_{i=1}^n t_i$ whit $t_i \in X$ or $t_i^{-1} \in X$. Therefore $\sigma_2 \in R$, which is a contradiction.

(ii) By (i), we have $\mathcal{O}_a^H \neq \mathcal{O}_b^H$ and they are not square commutative. By [HS08, Th. 8.6], $\dim \mathfrak{B}(M(\mathcal{O}_a^H, \lambda) + M(\mathcal{O}_b^H, \xi)) = \infty$, for any $\lambda \in \widehat{H}^a$, $\xi \in \widehat{H}^b$. Consequently, $\dim \mathfrak{B}(\mathcal{O}_\sigma^G, \rho) = \infty$ for any $\rho \in \widehat{G}^\sigma$ by [AFGV08, Lemma 3.2 ii]. \square

References

- [AFGV08] N. Andruskiewitsch, F. Fantino, M. Graña, L.Vendramin, Finite-dimensional pointed Hopf algebras with alternating groups are trivial, *Ann. Mat. Pura Appl.*, **190** (2011)2, 225-245.
- [AFZ09] N. Andruskiewitsch, F. Fantino, S. Zhang, On pointed Hopf algebras associated with the symmetric groups, *Manuscripta Math.* **128** (2009), 359-371.
- [AZ07] N. Andruskiewitsch, S. Zhang, On pointed Hopf algebras associated to some conjugacy classes in \mathbb{S}_n , *Proc. Amer. Math. Soc.*, **135** (2007), 2723-2731.
- [AG03] N. Andruskiewitsch, M. Grana, From racks to pointed Hopf algebras, *Adv. in Math.*, **178** (2003)2, 177-243.
- [AS02] N. Andruskiewitsch, H.-J. Schneider, Pointed Hopf algebras, in *New Directions in Hopf Algebras*, MSRI Series, pp.1-68, Cambridge Univ. Press, 2002.
- [AS10] N. Andruskiewitsch, H.-J. Schneider, On the classification of finite-dimensional pointed Hopf algebras, *Ann. Math.*, **171** (2010) 1, 375-417.
- [Ci97] C. Cibils, Tensor products of Hopf bimodules over a group algebra. *Proc. A.M.S.* **125** (1997), 1315-1321.
- [DPR91] R Dijkgraaf, V Pasquier, P Roche, Quasi Hopf algebras, group cohomology and orbifold models. *Nucl. Phys. B Proc. Suppl.* **18** (1991), 60-72.
- [FK99] S. Fomin, A.N. Kirillov, Quadratic algebras, Dunkl elements, and Schubert calculus, *Advances in geometry*, 147-182, *Progr. Math.*, 172, Birkhäuser Boston, Boston, MA, 1999.

- [Gr00] M. Graña, On Nichols algebras of low dimension, *Contemp. Math.*, **267** (2000), 111-134.
- [MS] A. Milinski, H.-J. Schneider, *Pointed Indecomposable Hopf Algebras over Coxeter Groups*, *Contemp. Math.*, **267** (2000), 1-22.
- [Se77] J.-P. Serre, *Linear representations of finite groups*, Springer-Verlag, New York 1977.
- [ZZ12] Shouchuan Zhang, Yao-Zhong Zhang, Pointed Hopf Algebras with classical Weyl Groups, *International Journal of Mathematics*, **23**(2012), 1250066.
- [HS08] I. Heckenberger, H.-J. Schneider, Root systems and Weyl groupoids for Nichols algebras. *Proceedings of the London Mathematical Society*, 2010, **101**(3):623-654.