

DISTRIBUTION OF THE VOLUME OF WEIGHTED GAUSSIAN SIMPLEX

TATIANA MOSEEEVA

ABSTRACT. Let $X_0, \dots, X_l$ be independent standard Gaussian vectors in $\mathbb{R}^d$ such that $l \leq d$. We derive an explicit formula for the distribution of the volume of weighted Gaussian simplex without the origin — l -dimensional simplex $\text{conv}(\sigma_0 X_0, \dots, \sigma_l X_l)$ ($\sigma_0, \dots, \sigma_l > 0$).

1. INTRODUCTION AND MAIN RESULT

Let X be a standard Gaussian vector in $\mathbb{R}^k$. By definition, its length is chi-distributed with k degrees of freedom:

$$|X| \stackrel{d}{=} \chi_k.$$

This distributional identity has a nice well-known generalization to several vectors. Namely, let $X_0, X_1, \dots, X_l$ be random points sampled independently from the standard normal distribution on $\mathbb{R}^d$ ($l \leq d$). Then using some geometrical reasoning based on the «height $\times$ base» formula for the volume of a simplex and the symmetry property of the multidimensional Gaussian distribution (for details, see, e.g., [1, Chapter 7] or [4]) it can be shown that the volume of the l -dimensional Gaussian simplex with the origin is distributed as follows:

$$(1.1) \quad |\text{conv}(0, X_1, \dots, X_l)| \stackrel{d}{=} \frac{1}{l!} \chi_{d-l+1} \dots \chi_d,$$

where $|\cdot|$ denotes the l -dimensional volume. It follows immediately that for any $\sigma_1, \dots, \sigma_l > 0$,

$$|\text{conv}(0, \sigma_1 X_1, \dots, \sigma_l X_l)| \stackrel{d}{=} \sigma_1 \dots \sigma_l \frac{1}{l!} \chi_{d-l+1} \dots \chi_d.$$

If we remove the origin from the set of the vertices, the situation becomes more subtle. Using the Blaschke–Petkantchin formula (see Statement 2.1 below) Miles showed [3] that

$$(1.2) \quad \mathbb{E}|\text{conv}(X_0, \dots, X_l)|^p = \left[\frac{2^{l/2} \sqrt{l+1}}{l!} \right]^p \prod_{i=d-l+1}^d \frac{\Gamma((i+p)/2)}{\Gamma(i/2)}.$$

From this, in view of

$$\mathbb{E}\chi_k^p = 2^{p/2} \frac{\Gamma((k+p)/2)}{\Gamma(k/2)},$$

he concluded (without a proof) that it follows from the method of moments that

$$|\text{conv}(X_0, \dots, X_l)| \stackrel{d}{=} \frac{\sqrt{l+1}}{l!} \chi_{d-l+1} \dots \chi_d,$$

Key words and phrases. Random simplex, Gaussian simplex, convex hull, volume, Blaschke–Petkantschin formula.

The work was supported by the Foundation for the Advancement of Theoretical Physics and Mathematics “BASIS”.

The work was supported by Ministry of Science and Higher Education of the Russian Federation, agreement № 075-15-2019-1619.

see also [2, Theorem 2.5].

Note that now the weighted case does not follow from the unweighted one. Specifically, in their study of the convex hulls of several Gaussian random walks, Randon-Furling and Zaporozhets [6, Theorem 6.1] derived as an auxiliary result the following formula:

$$(1.3) \quad \mathbb{E}|\text{conv}(\sigma_0 X_0, \dots, \sigma_l X_l)|^p = \left[\frac{2^{l/2} \sigma_0 \dots \sigma_l}{l!} \sqrt{\frac{1}{\sigma_0^2} + \dots + \frac{1}{\sigma_l^2}} \right]^p \prod_{i=d-l+1}^d \frac{\Gamma((i+p)/2)}{\Gamma(i/2)}.$$

Although (1.2) was already known, it took them three pages to prove it by using a generalization of the Sudakov–Tsirelson theorem obtained by Paouris and Pivovarov [5, Proposition 4.1]. Again, it was suggested [6, Remark 6.2] that from the method of moments the distribution of the volume of the weighted Gaussian simplex could be obtained. Since for their main result the authors needed only the moments of the volume, they didn't provide a detailed proof. Also they suggested that it is possible to find a direct derivation avoiding the method of moments.

In this paper, we provide such a derivation, which is intuitive and geometric. Our main result is the following theorem.

Theorem 1.1. *Fix some $l = 1, \dots, d$. Let $X_0, \dots, X_l$ be independent d -dimensional standard Gaussian vectors. Then for any $\sigma_0, \dots, \sigma_l > 0$*

$$(1.4) \quad |\text{conv}(\sigma_0 X_0, \dots, \sigma_l X_l)| \stackrel{d}{=} \frac{1}{l!} \sigma_0 \dots \sigma_l \sqrt{\frac{1}{\sigma_0^2} + \dots + \frac{1}{\sigma_l^2}} \chi_{d-l+1} \dots \chi_d,$$

where $\chi_{d-l+1}, \dots, \chi_d$ are independent random variables such that for any $k = d-l+1, \dots, d$ the random variable χ_k has the chi distribution with k degrees of freedom.

The paper is organized as follows. First we recall some necessary facts from stochastic geometry in Section 2 and then present the proof of Theorem 1.1 in Section 3.

2. AUXILIARY RESULTS

To prove the main result we will need the following statement ([7], Theorem 7.2.7):

Statement 2.1 (Blaschke-Petkantschin formula). *Let $h : (\mathbb{R}^d)^{l+1} \rightarrow \mathbb{R}$ be non-negative measurable function, $l \in \{1, \dots, d\}$. Then*

$$\begin{aligned} & \int_{(\mathbb{R}^d)^{l+1}} f(x_0, \dots, x_l) dx_0 \dots dx_l = \\ & = (l!)^{d-l} b_{d,l} \int_{A_{d,l}} \int_{E^{l+1}} f(x_0, \dots, x_l) |\text{conv}(x_0, \dots, x_l)|^{d-l} \lambda_E(dx_0) \dots \lambda_E(dx_l) \mu_{d,l}(dE), \end{aligned}$$

where $b_{d,k} = \frac{\omega_{d-k+1} \dots \omega_d}{\omega_1 \dots \omega_k}$.

The following result of this section is probably well-known, however, for the reader's convenience we present it with proof.

Lemma 2.1. *Let $\eta_1, \eta_2, \theta_1, \theta_2, \xi_1, \xi_2$ be positive with probability one random variables, such that*

$$\begin{aligned}
(2.1) \quad & \eta_1 \cdot \theta_1 = \xi_1, \\
(2.2) \quad & \eta_2 \cdot \theta_2 = \xi_2,
\end{aligned}$$

and η_i u θ_i are independent for any $i \in \{1, 2\}$. Let us assume that $\theta_1 \stackrel{d}{=} \theta_2$ and $\xi_1 \stackrel{d}{=} \xi_2$. Then $\eta_1 \stackrel{d}{=} \eta_2$.

Proof. Taking the logarithm of equations (2.1) and (2.2) we obtain

$$\begin{aligned}
\log \eta_1 + \log \theta_1 &= \log \xi_1, \\
\log \eta_2 + \log \theta_2 &= \log \xi_2.
\end{aligned}$$

Random variables in the left-hand side of both of these equations are independent thus the characteristic function of their sum is equal to the product of their characteristic functions. Therefore,

$$\phi_{\log \eta_1}(t) \cdot \phi_{\log \theta_1}(t) = \phi_{\log \xi_1}(t) = \phi_{\log \xi_2}(t) = \phi_{\log \eta_2}(t) \cdot \phi_{\log \theta_2}(t),$$

where in the second step we used the fact that $\xi_1 \stackrel{d}{=} \xi_2$. Taking into account equation $\theta_1 \stackrel{d}{=} \theta_2$, we obtain $\phi_{\log \eta_1}(t) = \phi_{\log \eta_2}(t)$, and the lemma follows. $\square$

3. PROOF OF THEOREM 1.1

Let us consider the case $l = d$ first.

By the simplex volume formula,

$$\begin{aligned}
|\text{conv}(\sigma_0 X_0, \dots, \sigma_d X_d)| &= |\text{conv}(0, \sigma_1 X_1 - \sigma_0 X_0, \dots, \sigma_d X_d - \sigma_0 X_0)| \\
&= \frac{1}{d!} \det[\sigma_1 X_1 - \sigma_0 X_0, \dots, \sigma_d X_d - \sigma_0 X_0] \\
&= \frac{1}{d!} \det \begin{pmatrix} \sigma_1 X_1^{(1)} - \sigma_0 X_0^{(1)} & \dots & \sigma_d X_d^{(1)} - \sigma_0 X_0^{(1)} \\ \vdots & \ddots & \vdots \\ \sigma_1 X_1^{(d)} - \sigma_0 X_0^{(d)} & \dots & \sigma_d X_d^{(d)} - \sigma_0 X_0^{(d)} \end{pmatrix} \\
&= \frac{1}{d!} \det \begin{pmatrix} \sigma_1 X_1^{(1)} - \sigma_0 X_0^{(1)} & \dots & \sigma_1 X_1^{(d)} - \sigma_0 X_0^{(d)} \\ \vdots & \ddots & \vdots \\ \sigma_d X_d^{(1)} - \sigma_0 X_0^{(1)} & \dots & \sigma_d X_d^{(d)} - \sigma_0 X_0^{(d)} \end{pmatrix},
\end{aligned}$$

where in the last step we used the fact that the determinant does not change after transposing.

Note that columns of the obtained matrix are independent Gaussian vectors. Let us denote the k -th column of this matrix by Y_k . $\mathbb{E}Y_k = 0$.

$$\begin{aligned}\text{cov}(Y_k^i, Y_k^j) &= \text{cov}(\sigma_i X_i^{(k)} - \sigma_0 X_0^{(k)}, \sigma_j X_j^{(k)} - \sigma_0 X_0^{(k)}) = \sigma_i \sigma_j \text{cov}(X_i^{(k)}, X_j^{(k)}) - \\ &\quad - \sigma_i \sigma_0 \underbrace{\text{cov}(X_i^{(k)}, X_0^{(k)})}_{=0} - \sigma_j \sigma_0 \underbrace{\text{cov}(X_0^{(k)}, X_j^{(k)})}_{=0} + \sigma_0^2 \text{cov}(X_0^{(k)}, X_0^{(k)}) = \\ &= \sigma_i \sigma_j \text{cov}(X_i^{(k)}, X_j^{(k)}) + \sigma_0^2 = \sigma_i \sigma_j \delta_{ij} + \sigma_0^2,\end{aligned}$$

where δ_{ij} is the Kronecker delta.

Therefore,

$$\text{cov}(Y_k) = \begin{pmatrix} \sigma_1^2 + \sigma_0^2 & \sigma_0^2 & \dots & \sigma_0^2 \\ \sigma_0^2 & \sigma_2^2 + \sigma_0^2 & \dots & \sigma_0^2 \\ \vdots & \vdots & \ddots & \vdots \\ \sigma_0^2 & \sigma_0^2 & \dots & \sigma_d^2 + \sigma_0^2 \end{pmatrix}.$$

Note that the covariance matrix of Y_k does not depend on k . Let us denote this matrix by $\mathbf{M}$. Then we can say, that $Y_k = \mathbf{A}\overline{X}_k$, where $\overline{X}_k$ are independent standard Gaussian vectors and $\mathbf{A}\mathbf{A}^T = \mathbf{M}$.

It follows that

$$(3.1) \quad |\text{conv}(\sigma_0 X_0, \dots, \sigma_d X_d)| = \frac{1}{d!} \det[\mathbf{A}\overline{X}_1, \dots, \mathbf{A}\overline{X}_d] = \frac{1}{d!} \det \mathbf{A} \det[\overline{X}_1, \dots, \overline{X}_d].$$

First let us calculate the determinant of $\mathbf{A}$. Since $\mathbf{A}\mathbf{A}^T = \mathbf{M}$, we obtain $\det \mathbf{A} = (\det \mathbf{M})^{\frac{1}{2}}$.

$$\begin{aligned}\det \mathbf{M} &= \det \left(\begin{pmatrix} \sigma_1^2 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & \sigma_d^2 \end{pmatrix} + \sigma_0^2 \cdot I \right) = \\ &= \sigma_1^2 \dots \sigma_d^2 + \sum_{k=1}^d \frac{\sigma_1^2 \dots \sigma_d^2}{\sigma_k^2} \cdot \sigma_0^2 = \sigma_0^2 \sigma_1^2 \dots \sigma_d^2 \sum_{k=0}^d \frac{1}{\sigma_k^2}.\end{aligned}$$

Consequently, $\det \mathbf{A} = \sqrt{\det \mathbf{M}} = \sigma_0 \dots \sigma_d \sqrt{\frac{1}{\sigma_0^2} + \dots + \frac{1}{\sigma_d^2}}$.

It remains to find the distribution of $\det[\overline{X}_1, \dots, \overline{X}_d]$, that is, the volume of the parallelotope spanned by vectors $\overline{X}_1, \dots, \overline{X}_d$. Let us denote that parallelotope by P .

$$|P| = d! \cdot |\text{conv}(0, \overline{X}_1, \dots, \overline{X}_d)|.$$

By (1.1), $|\text{conv}(0, \overline{X}_1, \dots, \overline{X}_d)| \stackrel{d}{=} \frac{1}{d!} \chi_1 \dots \chi_d$, thus

$$(3.2) \quad \det[\overline{X}_1, \dots, \overline{X}_d] \stackrel{d}{=} \chi_1 \dots \chi_d.$$

Substituting (3.2) in (3.1) we obtain

$$(3.3) \quad |\text{conv}(\sigma_0 X_0, \dots, \sigma_d X_d)| \stackrel{d}{=} \frac{1}{d!} \sigma_0 \dots \sigma_d \sqrt{\frac{1}{\sigma_0^2} + \dots + \frac{1}{\sigma_d^2}} \chi_1 \dots \chi_d,$$

where random variables $\chi_1, \dots, \chi_d$ are independent, χ_k has the chi distribution with k degrees of freedom for any $k = 1, \dots, d$.

Now suppose that $l < d$. Consider the affine hull

$$V_l := \text{aff}(\sigma_0 X_0, \dots, \sigma_l X_l).$$

With probability one, V_l is an affine l -plane. Denote by O_{V_l} the orthogonal projection of the origin onto V_l . Consider the linear subspace

$$W_l := V_l - O_{V_l}.$$

Lemma. W_l is uniformly distributed over the l -dimensional linear Grassmanian with respect to Haar measure and independently of $|\text{conv}(\sigma_0 X_0, \dots, \sigma_l X_l)|$.

Proof. Let $f : G_{d,l} \rightarrow \mathbb{R}^1$, $g : \mathbb{R}^1 \rightarrow \mathbb{R}^1$ be non-negative bounded measurable functions.

$$\begin{aligned} & \mathbb{E}[f(W_l)g(|\text{conv}(\sigma_0 X_0, \dots, \sigma_l X_l)|)] \\ &= \int_{(\mathbb{R}^d)^{l+1}} f(\text{aff}(\sigma_0 x_0, \dots, \sigma_l x_l) - O_{V_l}) g(|\text{conv}(\sigma_0 x_0, \dots, \sigma_l x_l)|) ((2\pi)^{-d/2})^{l+1} e^{-\frac{|x_0|^2}{2}} \dots e^{-\frac{|x_l|^2}{2}} dx_0 \dots dx_l \\ &= \frac{1}{\prod_{i=0}^l \sigma_i} ((2\pi)^{-d/2})^{l+1} \int_{(\mathbb{R}^d)^{l+1}} f(\text{aff}(y_0, \dots, y_l) - O_{V_l}) g(|\text{conv}(y_0, \dots, y_l)|) e^{-\frac{|y_0|^2}{2\sigma_0^2}} \dots e^{-\frac{|y_l|^2}{2\sigma_l^2}} dy_0 \dots dy_l. \end{aligned}$$

Using Statement 2.1 yields

$$\begin{aligned} \mathbb{E}[f(W_l)g(|\text{conv}(\sigma_0 X_0, \dots, \sigma_l X_l)|)] &= \underbrace{(l!)^{d-l} b_{d,l} \frac{1}{\prod_{i=0}^l \sigma_i}}_{=C} ((2\pi)^{-d/2})^{l+1} \cdot \\ & \int_{A_{d,l}} \int_{E^{l+1}} f(\text{aff}(y_0, \dots, y_l) - O_{V_l}) g(|\text{conv}(y_0, \dots, y_l)|) e^{-\left(\frac{|y_0|^2}{2\sigma_0^2} + \dots + \frac{|y_l|^2}{2\sigma_l^2}\right)} |\text{conv}(y_0, \dots, y_l)|^{d-l} \prod_{i=0}^l \lambda_E(dy_i) \mu_{d,l}(dE) \\ &= C \int_{G_{d,l}} f(L) \int_{L^\perp} \int_{(L+a)^{l+1}} g(|\text{conv}(y_0, \dots, y_l)|) e^{-\left(\frac{|y_0|^2}{2\sigma_0^2} + \dots + \frac{|y_l|^2}{2\sigma_l^2}\right)} |\text{conv}(y_0, \dots, y_l)|^{d-l} \prod_{i=0}^l \lambda_{L+a}(dy_i) da \nu_{d,l}(dL). \end{aligned}$$

Consider the linear l -plane

$$U := \text{lin}(e_1, \dots, e_l).$$

Fix some $L \in G_{d,l}$. Let $\mathbf{Q}$ be the orthogonal matrix such that $L = \mathbf{Q}U$. Then we have

$$\begin{aligned} & \int_{L^\perp} \int_{(L+a)^{l+1}} g(|\text{conv}(y_0, \dots, y_l)|) e^{-\left(\frac{|y_0|^2}{2\sigma_0^2} + \dots + \frac{|y_l|^2}{2\sigma_l^2}\right)} |\text{conv}(y_0, \dots, y_l)|^{d-l} \prod_{i=0}^l \lambda_{L+a}(dy_i) da \\ &= \int_{U^\perp} \int_{(U+a)^{l+1}} g(|\text{conv}(\mathbf{Q}y_0, \dots, \mathbf{Q}y_l)|) e^{-\left(\frac{|\mathbf{Q}y_0|^2}{2\sigma_0^2} + \dots + \frac{|\mathbf{Q}y_l|^2}{2\sigma_l^2}\right)} |\text{conv}(\mathbf{Q}y_0, \dots, \mathbf{Q}y_l)|^{d-l} \prod_{i=0}^l \lambda_{U+a}(dy_i) da \\ &= \int_{U^\perp} \int_{(U+a)^{l+1}} g(|\text{conv}(y_0, \dots, y_l)|) e^{-\left(\frac{|y_0|^2}{2\sigma_0^2} + \dots + \frac{|y_l|^2}{2\sigma_l^2}\right)} |\text{conv}(y_0, \dots, y_l)|^{d-l} \prod_{i=0}^l \lambda_{U+a}(dy_i) da. \end{aligned}$$

Note that the obtained expression does not depend on L , denote it by $\mathcal{J}(g)$.

Therefore,

$$\mathbb{E}[f(W_l)g(|\text{conv}(\sigma_0 X_0, \dots, \sigma_l X_l)|)] = C\mathcal{J}(g) \int_{G_{d,l}} f(L)\nu_{d,l}(dL).$$

Substituting $f = 1$ and $g = 1$ yields

$$1 = C\mathcal{J}(1).$$

Let A be some measurable subset of $G_{d,l}$. Consider $f = \mathbb{1}_A, g = 1$. We have

$$\mathbb{P}[W_l \in A] = C\mathcal{J}(1) \int_A \nu_{d,l}(dL) = \int_A \nu_{d,l}(dL),$$

therefore W is uniformly distributed over $G_{d,l}$.

Taking $f = 1, g = \mathbb{1}_B$ for some measurable $B \in \mathbb{R}$ we obtain

$$\mathbb{P}[|\text{conv}(\sigma_0 X_0, \dots, \sigma_l X_l)| \in B] = C\mathcal{J}(\mathbb{1}_B).$$

Substituting $f = \mathbb{1}_A$ and $g = \mathbb{1}_B$ finishes the proof. $\square$

Let P_l be the orthogonal projection from $\mathbb{R}^d$ onto the first l coordinates, denote by P_l^W its restriction to W_l .

We have

$$\begin{aligned} |\text{conv}(\sigma_0 P_l X_0, \dots, \sigma_l P_l X_l)| &= |\text{conv}(\sigma_0 P_l X_0 - P_l O_{V_l}, \dots, \sigma_l P_l X_l - P_l O_{V_l})| \\ &= |\text{conv}(P_l^W(\sigma_0 X_0 - O_{V_l}), \dots, P_l^W(\sigma_l X_l - O_{V_l}))| \\ (3.4) \quad &= |\text{conv}(\sigma_0 X_0 - O_{V_l}, \dots, \sigma_l X_l - O_{V_l})| \cdot |\det(P_l^W)| \\ &= |\text{conv}(\sigma_0 X_0, \dots, \sigma_l X_l)| \cdot |\det(P_l^W)|. \end{aligned}$$

As noted above, the orthogonal projection of standard Gaussian vector onto the l -dimensional subspace has the standard Gaussian distribution in that subspace, therefore, using (3.3), we have:

$$(3.5) \quad |\text{conv}(\sigma_0 P_l X_0, \dots, \sigma_l P_l X_l)| \stackrel{d}{=} \frac{1}{l!} \sigma_0 \dots \sigma_l \sqrt{\frac{1}{\sigma_0^2} + \dots + \frac{1}{\sigma_l^2}} \chi_1 \dots \chi_l,$$

where $\chi_1, \dots, \chi_l$ are independent chi-distributed random variables.

Thus we have

$$(3.6) \quad |\text{conv}(\sigma_0 X_0, \dots, \sigma_l X_l)| \cdot |\det(P_l^W)| \stackrel{d}{=} \frac{1}{l!} \sigma_0 \dots \sigma_l \sqrt{\frac{1}{\sigma_0^2} + \dots + \frac{1}{\sigma_l^2}} \chi_1 \dots \chi_l,$$

wherein $|\text{conv}(\sigma_0 X_0, \dots, \sigma_l X_l)|$ and $|\det(P_l^W)|$ are independent.

Consider the linear subspace $W := \text{lin}(0, Y_1, \dots, Y_l)$, where Y_i are independent standard Gaussian vectors in $\mathbb{R}^d$. With probability one, W is a linear l -plane. Since Y_i are independent and spherically invariant, W is uniformly distributed over the l -dimensional Grassmanian.

Thus,

$$(3.7) \quad |\text{conv}(0, Y_1, \dots, Y_l) \cdot |\det(P_l^W)| = |\text{conv}(0, P_l Y_1, \dots, P_l Y_l)|,$$

where $|\det(P_l^W)|$ does not depend on $|\text{conv}(0, Y_1, \dots, Y_l)|$.

Since (1.1),

$$(3.8) \quad |\text{conv}(0, Y_1, \dots, Y_l)| \stackrel{d}{=} \frac{1}{l!} \chi_{d-l+1} \dots \chi_d,$$

where χ_i are independent chi-distributed random variables.

Similarly, using the fact, that $P_l Y_i$ are independent standard Gaussian vectors in $\mathbb{R}^l$, we obtain

$$(3.9) \quad |\text{conv}(0, P_l Y_1, \dots, P_l Y_l)| \stackrel{d}{=} \frac{1}{l!} \chi_1 \dots \chi_l.$$

Combining (3.7), (3.8) and (3.9), we have

$$(3.10) \quad \frac{1}{l!} \chi_{d-l+1} \dots \chi_d \cdot |\det(P_l^W)| \stackrel{d}{=} \frac{1}{l!} \chi_1 \dots \chi_l.$$

Multiplying (3.10) by $\sigma_0 \dots \sigma_l \sqrt{\frac{1}{\sigma_0^2} + \dots + \frac{1}{\sigma_l^2}}$ we obtain

$$(3.11) \quad \frac{1}{l!} \sigma_0 \dots \sigma_l \sqrt{\frac{1}{\sigma_0^2} + \dots + \frac{1}{\sigma_l^2}} \chi_{d-l+1} \dots \chi_d \cdot |\det(P_l^W)| \stackrel{d}{=} \frac{1}{l!} \sigma_0 \dots \sigma_l \sqrt{\frac{1}{\sigma_0^2} + \dots + \frac{1}{\sigma_l^2}} \chi_1 \dots \chi_l.$$

Applying Lemma 2.1 to (3.6) and (3.11) we conclude

$$|\text{conv}(\sigma_0 X_0, \dots, \sigma_l X_l)| \stackrel{d}{=} \frac{1}{l!} \sigma_0 \dots \sigma_l \sqrt{\frac{1}{\sigma_0^2} + \dots + \frac{1}{\sigma_l^2}} \chi_{d-l+1} \dots \chi_d,$$

which finishes the proof.

REFERENCES

- [1] T. W. Anderson. *An introduction to multivariate statistical analysis*. Wiley Series in Probability and Statistics. Wiley-Interscience [John Wiley & Sons], Hoboken, NJ, third edition, 2003.
- [2] J. Grote, Z. Kabluchko, and Ch. Thäle. Limit theorems for random simplices in high dimensions. *ALEA Lat. Am. J. Probab. Math. Stat.*, 16(1):141–177, 2019.
- [3] R. E. Miles. Isotropic random simplices. *Advances in Applied Probability*, 3(2):353–382, 1971.
- [4] H. H. Nguyen and V. Vu. Random matrices: law of the determinant. *Ann. Probab.*, 42(1):146–167, 2014.
- [5] G. Paouris and P. Pivovarov. Small-ball probabilities for the volume of random convex sets. *Discrete Comput. Geom.*, 49(3):601–646, 2013.
- [6] Julien Randon-Furling and Dmitry Zaporozhets. Convex hulls of several multidimensional gaussian random walks. *arXiv preprint arXiv:2007.02768*, 2020.
- [7] R. Schneider and W. Weil. *Stochastic and integral geometry*. Probability and its Applications (New York). Springer-Verlag, Berlin, 2008.

TATIANA MOSEVA, LEONHARD EULER INTERNATIONAL MATHEMATICAL INSTITUTE, RUSSIA
E-mail address: polezina@yandex.ru