

# Separation of congruence intervals and implications

Andrei A. Bulatov

## Abstract

The Constraint Satisfaction Problem (CSP) has been intensively studied in many areas of computer science and mathematics. The approach to the CSP based on tools from universal algebra turned out to be the most successful one to study the complexity and algorithms for this problem. Several techniques have been developed over two decades. One of them is through associating edge-colored graphs with algebras and studying how the properties of algebras are related with the structure of the associated graphs. This approach has been introduced in our previous two papers (A.Bulatov, Local structure of idempotent algebras I,II. CoRR abs/2006.09599, CoRR abs/2006.10239, 2020). In this paper we further advance it by introducing new structural properties of finite idempotent algebras omitting type **1** such as separation congruences, collapsing polynomials, and their implications for the structure of subdirect products of finite algebras. This paper also provides the algebraic background for our proof of Feder-Vardi Dichotomy Conjecture (A. Bulatov, A Dichotomy Theorem for Nonuniform CSPs. FOCS 2017: 319-330).

## 1 Introduction

Over the last two decades methods from universal algebra found strong applications in computer science, specifically in the study of the Constraint Satisfaction Problem (CSP) and related combinatorial problems. The original research problem where the algebraic approach was used is the complexity of so-called nonuniform CSPs, and more specifically the Dichotomy Conjecture posed by Feder and Vardi in [15, 16] and refined in [12]. The Dichotomy Conjecture states that every nonuniform CSP is either solvable in polynomial time or is NP-complete, and also delineates the precise borderline between the two cases. Every nonuniform CSP can be associated with a finite algebra, and the complexity of the CSP is completely determined by this algebra [19, 12]. The Dichotomy Conjecture was confirmed independently by the author [6, 7] and by Zhuk [26, 27], and the algebraic approach played a key role in both proofs.

The specific version of the algebraic approach used in [6, 7] was developed in [2, 13, 3, 4, 8, 11]. In this paper we further advance this approach preparing

the ground for a proof of the Dichotomy Conjecture. We will introduce two structural features of finite algebras and demonstrate how they influence the structure of subdirect products of finite idempotent algebras omitting type **1**.

First we introduce the notion of separability of prime intervals in the congruence lattice by a unary polynomial. More precisely, we say that a prime interval  $\alpha \prec \beta$  in the congruence lattice of an algebra  $\mathbb{A}$  can be separated from interval  $\gamma \prec \delta$  if there is a unary polynomial  $f$  of  $\mathbb{A}$  such that  $f(\delta) \subseteq \gamma$ , but  $f(\beta) \not\subseteq \alpha$ . This concept can be extended to subdirect products of algebras, say,  $R \subseteq \mathbb{A} \times \mathbb{B}$ , in which case intervals  $\alpha \prec \beta$  and  $\gamma \prec \delta$  may be in the congruence lattices of different factors, say  $\alpha, \beta \in \text{Con}(\mathbb{A}_1)$ ,  $\gamma, \delta \in \text{Con}(\mathbb{A}_2)$ , and  $f$  is a polynomial of  $R$ . The relation ‘cannot be separated from’ on the set of prime intervals is clearly reflexive and transitive. Our first result, Theorem 38 shows that it is also to some extent symmetric.

The property proved in Theorem 38 is used to prove the existence of the second structural feature of subdirect products, collapsing polynomials, see Theorem 53. A unary polynomial of a subdirect product  $R \subseteq \mathbb{A}_1 \times \cdots \times \mathbb{A}_n$  for a prime interval  $\alpha \prec \beta$  in  $\text{Con}(\mathbb{A}_i)$  for some  $i$  is collapsing if for any  $j$  and any prime interval  $\gamma \prec \delta$  in  $\text{Con}(\mathbb{A}_j)$  it holds that  $f(\delta) \not\subseteq \gamma$  if and only if  $\alpha \prec \beta$  cannot be separated from  $\gamma \prec \delta$ . Collapsing polynomials are one of the main tools in the proof of the Dichotomy Conjecture [6, 7], as they are very useful in the study of the structure of subdirect products. One example of such results is the Congruence Lemma 56, which provides much information about the fine structure of a subdirect product of algebras when one of its factors is restricted on its congruence block. The Congruence Lemma is another important tool in the proof of the Dichotomy Conjecture.

Besides congruence separation and collapsing polynomials we also introduce an alternative definition of the centralizer and use it to derive certain properties of subdirect products. In addition, we introduce two more technical properties of subdirect products, chaining and polynomial closure, and study their properties that again are instrumental in the proof of the Dichotomy Conjecture.

## 2 Preliminaries

Here we introduce all the notation and terminology used in this paper. It mainly follows the standard books [14, 22].

### 2.1 Notation and agreements

By  $[n]$  we denote the set  $\{1, \dots, n\}$ . For sets  $A_1, \dots, A_n$  tuples from  $A_1 \times \cdots \times A_n$  are denoted in boldface, say,  $\mathbf{a}$ ; the  $i$ th component of  $\mathbf{a}$  is referred to

as  $\mathbf{a}[i]$ . An  $n$ -ary relation  $R$  over sets  $A_1, \dots, A_n$  is any subset of  $A_1 \times \dots \times A_n$ . For  $I = \{i_1, \dots, i_k\} \subseteq [n]$  by  $\text{pr}_I \mathbf{a}, \text{pr}_I R$  we denote the *projections*  $\text{pr}_I \mathbf{a} = (\mathbf{a}[i_1], \dots, \mathbf{a}[i_k])$ ,  $\text{pr}_I R = \{\text{pr}_I \mathbf{a} \mid \mathbf{a} \in R\}$  of tuple  $\mathbf{a}$  and relation  $R$ . If  $\text{pr}_i R = A_i$  for each  $i \in [n]$ , relation  $R$  is said to be a *subdirect product* of  $A_1 \times \dots \times A_n$ . It will be convenient to use  $\overline{A}$  for  $A_1 \times \dots \times A_n$  if the sets  $A_i$  are clear from the context. For  $I \subseteq [n]$  we will use  $\overline{A}_I$ , for  $\prod_{i \in I} A_i$ , or if  $I$  is clear from the context just  $\overline{A}$ .

Algebras will be denoted by  $\mathbb{A}, \mathbb{B}$  etc.; we often do not distinguish between subuniverses and subalgebras. For  $B \subseteq \mathbb{A}$  the subalgebra generated by  $B$  is denoted  $\text{Sg}_{\mathbb{A}}(B)$  or just  $\text{Sg}(B)$ . For  $C \subseteq \mathbb{A}^2$  the congruence generated by  $C$  is denoted  $\text{Cg}_{\mathbb{A}}(C)$  or just  $\text{Cg}(C)$ . The equality relation and the full congruence of algebra  $\mathbb{A}$  are denoted  $\underline{0}_{\mathbb{A}}$  and  $\underline{1}_{\mathbb{A}}$ , respectively. Often when we need to use one of these trivial congruences of an algebra indexed in some way, say,  $\mathbb{A}_i$ , we write  $\underline{0}_i, \underline{1}_i$  for  $\underline{0}_{\mathbb{A}_i}, \underline{1}_{\mathbb{A}_i}$ . The set of all polynomials (unary, binary polynomials) of  $\mathbb{A}$  is denoted by  $\text{Pol}(\mathbb{A})$  and  $\text{Pol}_1(\mathbb{A}), \text{Pol}_2(\mathbb{A})$ , respectively. A unary polynomial  $f$  is *idempotent* if  $f \circ f = f$ . We frequently use operations on subalgebras of direct products of algebras, say,  $R \subseteq \mathbb{A}_1 \times \dots \times \mathbb{A}_n$ . If  $f$  is such an operation (say,  $k$ -ary) then we denote its component-wise action also by  $f$ , e.g.  $f(a_1, \dots, a_k)$  for  $a_1, \dots, a_k \in \mathbb{A}_i$ . In the same way we denote the action of  $f$  on projections of  $R$ , e.g.  $f(\mathbf{a}_1, \dots, \mathbf{a}_k)$  for  $I \subseteq [n]$  and  $\mathbf{a}_1, \dots, \mathbf{a}_k \in \text{pr}_I R$ . What we mean will always be clear from the context. We use similar agreements for collections of congruences. If  $\alpha_i \in \text{Con}(\mathbb{A}_i)$ ,  $i \in [n]$ , then  $\overline{\alpha}$  denotes the congruence  $\alpha_1 \times \dots \times \alpha_n$  of  $R$ . If  $I \subseteq [n]$  we use  $\overline{\alpha}_I$  to denote  $\prod_{i \in I} \alpha_i$ . If it does not lead to a confusion we write  $\overline{\alpha}$  for  $\overline{\alpha}_I$ . Sometimes  $\alpha_i$  are specified for  $i$  from a certain set  $I \subseteq [n]$ , then by  $\overline{\alpha}$  we mean the congruence  $\prod_{i \in [n]} \alpha'_i$  where  $\alpha'_i = \alpha_i$  if  $i \in I$  and  $\alpha'_i$  is the equality relation otherwise. For example, if  $\alpha \in \text{Con}(\mathbb{A}_1)$  then  $R/\overline{\alpha}$  means the factor of  $R$  modulo  $\alpha \times \underline{0}_2 \times \dots \times \underline{0}_n$ . For  $\alpha \in \text{Con}(\mathbb{A})$  and a polynomial  $f$  of  $\mathbb{A}$ , we will often abuse notation and denote the action of  $f$  on  $\mathbb{A}/\alpha$  by the same symbol  $f$ . For  $\alpha, \beta \in \text{Con}(\mathbb{A})$  we write  $\alpha \prec \beta$  if  $\alpha < \beta$  and  $\alpha \leq \gamma \leq \beta$  in  $\text{Con}(\mathbb{A})$  implies  $\gamma = \alpha$  or  $\gamma = \beta$ . In this paper all algebras are finite, idempotent and omit type **1**, except the definition of edges and Theorem 5 in the beginning of Section 2.3.

## 2.2 Minimal sets and polynomials

We will use the following basic facts from the tame congruence theory [18], often without further notice.

Let  $\mathbb{A}$  be a finite algebra and  $\alpha, \beta \in \text{Con}(\mathbb{A})$  with  $\alpha \prec \beta$ . An  $(\alpha, \beta)$ -*minimal set* is a set minimal with respect to inclusion among the sets of the form  $f(\mathbb{A})$ , where  $f \in \text{Pol}_1(\mathbb{A})$  is such that  $f(\beta) \not\subseteq \alpha$ . Sets  $B, C \subseteq \mathbb{A}$  are said to be *polynomially isomorphic* in  $\mathbb{A}$  if there are  $f, g \in \text{Pol}_1(\mathbb{A})$  such that  $f(B) = C$ ,  $g(C) = B$ ,

and  $f \circ g, g \circ f$  are identity mappings on  $C$  and  $B$ , respectively.

**Lemma 1** (Theorem 2.8, [18]). *Let  $\alpha, \beta \in \text{Con}(\mathbb{A})$ ,  $\alpha \prec \beta$ . Then the following hold.*

- (1) *Any  $(\alpha, \beta)$ -minimal sets  $U, V$  are polynomially isomorphic.*
- (2) *For any  $(\alpha, \beta)$ -minimal set  $U$  and any  $f \in \text{Pol}_1(\mathbb{A})$ , if  $f(\beta|_U) \not\subseteq \alpha$  then  $f(U)$  is an  $(\alpha, \beta)$ -minimal set,  $U$  and  $f(U)$  are polynomially isomorphic, and  $f$  witnesses this fact.*
- (3) *For any  $(\alpha, \beta)$ -minimal set  $U$  there is  $f \in \text{Pol}_1(\mathbb{A})$  such that  $f(\mathbb{A}) = U$ ,  $f(\beta) \not\subseteq \alpha$ , and  $f$  is idempotent, in particular,  $f$  is the identity mapping on  $U$ .*
- (4) *For any  $(a, b) \in \beta - \alpha$  and an  $(\alpha, \beta)$ -minimal set  $U$  there is  $f \in \text{Pol}_1(\mathbb{A})$  such that  $f(\mathbb{A}) = U$  and  $(f(a), f(b)) \in \beta|_U - \alpha|_U$ .*
- (5) *For any  $(\alpha, \beta)$ -minimal set  $U$ ,  $\beta$  is the transitive closure of*

$$\alpha \cup \{(f(a), f(b)) \mid (a, b) \in \beta|_U, f \in \text{Pol}_1(\mathbb{A})\}.$$

*In fact, as  $\alpha \prec \beta$ , this claim can be strengthened as follows. For any  $(a, b) \in \beta - \alpha$ ,  $\beta$  is the symmetric and transitive closure of*

$$\alpha \cup \{(f(a), f(b)) \mid f \in \text{Pol}_1(\mathbb{A})\}.$$

- (6) *For any  $f \in \text{Pol}_1(\mathbb{A})$  such that  $f(\beta) \not\subseteq \alpha$  there is an  $(\alpha, \beta)$ -minimal set  $U$  such that  $f$  witnesses that  $U$  and  $f(U)$  are polynomially isomorphic.*

For an  $(\alpha, \beta)$ -minimal set  $U$  and a  $\beta$ -block  $B$  such that  $\beta|_{U \cap B} \neq \alpha|_{U \cap B}$ , the set  $U \cap B$  is said to be an  $(\alpha, \beta)$ -trace. A 2-element set  $\{a, b\} \subseteq U \cap B$  such that  $(a, b) \in \beta - \alpha$ , is called an  $(\alpha, \beta)$ -subtrace. Depending on the structure of its minimal sets the interval  $(\alpha, \beta)$  can be of one of the five types, **1–5**. Since we assume the tractability conditions of the Dichotomy Conjecture, type **1** does not occur in algebras we deal with.

**Lemma 2** (Section 4 of [18]). *Let  $\alpha, \beta \in \text{Con}(\mathbb{A})$  and  $\alpha \prec \beta$ . Then the following hold.*

- (1) *If  $\text{typ}(\alpha, \beta) = \mathbf{2}$  then every  $(\alpha, \beta)$ -trace is polynomially equivalent to a 1-dimensional vector space.*

- (2) If  $\text{typ}(\alpha, \beta) \in \{3, 4, 5\}$  then every  $(\alpha, \beta)$ -minimal set  $U$  contains exactly one trace  $T$ , and if  $\text{typ}(\alpha, \beta) \in \{3, 4\}$ ,  $T$  contains only 2 elements. Also,  $T/\alpha$  is polynomially equivalent to a Boolean algebra, 2-element lattice, or 2-element semilattice, respectively.

Intervals  $(\alpha, \beta), (\gamma, \delta)$ ,  $\alpha, \beta, \gamma, \delta \in \text{Con}(\mathbb{A})$  and  $\alpha \prec \beta, \gamma \prec \delta$  are said to be *perspective* if  $\beta = \alpha \vee \delta, \gamma = \alpha \wedge \delta$ , or  $\delta = \beta \vee \gamma, \alpha = \beta \wedge \gamma$ .

**Lemma 3** (Lemma 6.2, [18]). *Let  $\alpha, \beta, \gamma, \delta \in \text{Con}(\mathbb{A})$  be such that  $\alpha \prec \beta, \gamma \prec \delta$  and intervals  $(\alpha, \beta), (\gamma, \delta)$  are perspective. Then  $\text{typ}(\alpha, \beta) = \text{typ}(\gamma, \delta)$  and a set  $U$  is  $(\alpha, \beta)$ -minimal if and only if it is  $(\gamma, \delta)$ -minimal.*

We will also use polynomials that behave on a minimal set in a particular way.

**Lemma 4** (Lemmas 4.15, 4.17, [18]). *Let  $\alpha, \beta \in \text{Con}(\mathbb{A})$ ,  $\alpha \prec \beta$ , and  $\text{typ}(\alpha, \beta) \in \{3, 4, 5\}$ . Let  $U$  be an  $(\alpha, \beta)$ -minimal set and  $T$  its only trace. Then there is an element  $1 \in T$  and a binary polynomial  $p$  of  $\mathbb{A}$  such that*

- (1)  $(1, a) \notin \alpha$  for any  $a \in U - \{1\}$ ;
- (2) for all  $a \in U - \{1\}$ , the algebra  $(\{a, 1\}, p)$  is a semilattice with neutral element 1, that is,  $p(1, 1) = 1$  and  $p(1, a) = p(a, 1) = p(a, a) = a$ .

*Polynomial  $p$  is said to be a pseudo-meet operation on  $U$ .*

*If  $\text{typ}(\alpha, \beta) \in \{3, 4\}$  then  $|T| = 2$ , say,  $T = \{0, 1\}$ , and there is a binary polynomial  $q$  of  $\mathbb{A}$ , a pseudo-join operation, that satisfies the conditions of item (2) except  $q(1, 0) = q(0, 1) = 1$ .*

## 2.3 Coloured graphs

In [2, 13, 3, 10, 11, 9] we introduced a local approach to the structure of finite algebras. As we use this approach throughout the paper, we present it here in some detail, see also [10, 11]. For the sake of the definitions below we slightly abuse terminology and by a module mean an algebra term equivalent to the full idempotent reduct of a module.

For an algebra  $\mathbb{A}$  the (undirected) graph  $\mathcal{G}(\mathbb{A})$  is defined as follows. The vertex set is the universe  $A$  of  $\mathbb{A}$ . A pair  $ab$  of vertices is an *edge* if and only if there exists a congruence  $\theta$  of  $\text{Sg}(a, b)$ , and either  $\text{Sg}(a, b)/\theta$  is a set (that is an algebra all of whose term operations are projections), or there is a term operation  $f$  of  $\mathbb{A}$  such that either  $\text{Sg}(a, b)/\theta$  is a module and  $f$  is an affine operation  $x - y + z$  on it ('affine operation' will always refer to  $x - y + z$ ), or  $f$  is a semilattice operation on  $\{a/\theta, b/\theta\}$ , or  $f$  is a majority operation on  $\{a/\theta, b/\theta\}$ . (Note that we use the same operation symbol in these cases.)

If there are a proper congruence  $\theta$  and a term operation  $f$  of  $\mathbb{A}$  such that  $f$  is a semilattice operation on  $\{a/\theta, b/\theta\}$  then  $ab$  is said to have the *semilattice type*. An edge  $ab$  is of *majority type* if there are a proper congruence  $\theta$  and a term operation  $f$  such that  $f$  is a majority operation on  $\{a/\theta, b/\theta\}$  and there is no semilattice term operation on  $\{a/\theta, b/\theta\}$ . Also,  $ab$  has the *affine type* if there are proper  $\theta$  and  $f$  such that  $f$  is an affine operation on  $\text{Sg}(a, b)/\theta$  and  $\text{Sg}(a, b)/\theta$  is a module; in particular it implies that there is no semilattice or majority operation on  $\{a/\theta, b/\theta\}$ . Finally, if  $\{a/\theta, b/\theta\}$  is a set,  $ab$  is said to have the *unary type*. In all cases we say that congruence  $\theta$  *witnesses* the type of edge  $ab$ . For an edge  $ab$  the set  $\{a/\theta, b/\theta\}$  is said to be a *thick edge*. Observe that a pair  $ab$  can be an edge of more than one type as witnessed by different congruences, although this has no consequences in this paper.

Omitting type **1** is characterized in [10, Theorem 5]. The second part of the next statement easily follows from [10, Theorem 5].

**Theorem 5** (Theorem 5, [10]). *An idempotent algebra  $\mathbb{A}$  omits type **1** (that is, the variety generated by  $\mathbb{A}$  omits type **1**) if and only if  $\mathcal{G}(\mathbb{A})$  contains no edges of the unary type.*

*Moreover, a finite class  $\mathcal{K}$  of similar idempotent algebras closed under subalgebras and quotient algebras omits type **1** if and only if  $\mathcal{G}(\mathbb{A})$  contains no edges of the unary type for any  $\mathbb{A} \in \mathcal{K}$ .*

For the sake of the Dichotomy Conjecture, it suffices to consider *reducts* of an algebra  $\mathbb{A}$  omitting type **1**, that is, algebras with the same universe but reduced set of term operations, as long as reducts also omit type **1**. In particular, we are interested in reducts of  $\mathbb{A}$ , in which semilattice and majority edges are subalgebras. An algebra  $\mathbb{A}$  such that  $a/\theta \cup b/\theta$  is a subuniverse of  $\mathbb{A}$  for every semilattice or majority edge  $ab$  of  $\mathbb{A}$  is called *smooth*. It is easy to see that that every subalgebra and every quotient of a smooth algebra is smooth. By [10, Theorem 12] if  $\mathcal{G}(\mathbb{A})$  contains no unary edges, there exists a reduct  $\mathbb{A}'$  of  $\mathbb{A}$  such that  $\mathbb{A}'$  is smooth and  $\mathcal{G}(\mathbb{A}')$  contains no edges of the unary type. From this point on all algebras occurring in the paper are idempotent and omit type **1**, unless stated otherwise.

Many concepts and results in the paper involve a class of algebras rather than a single algebra. Such a class, usually denoted by  $\mathcal{K}$ , is finite, consists of smooth algebras, and is closed under taking subalgebras and quotient algebras. A class of similar algebras satisfying these conditions will be called a *smooth class*. For a smooth class  $\mathcal{K}$  let  $\mathcal{V}$  be the class of finite algebras from the variety it generates, that is, the pseudovariety generated by  $\mathcal{K}$ . We will slightly abuse the terminology and call  $\mathcal{V}$  the variety generated by  $\mathcal{K}$ .

Observe that as the following example shows even though  $\mathcal{K}$  consists of smooth algebras, algebras in  $\mathcal{V}$  are not necessarily smooth.

**Example 6.** Let  $\mathbb{A}$  be an algebra with universe  $A = \{a, b, c\}$  and two basic operations  $f$  and  $g$ . The operation  $f$  is majority on  $A$ , and  $g$  is minority on  $\{a, b\}$  and  $\{a, c\}$ , and a 2/3-minority on  $\{b, c\}$ , that is,  $g(x, y, y) = g(x, y, x) = g(y, y, x) = x$  on  $\{b, c\}$ . If  $\{x, y, z\} = \{a, b, c\}$  then  $f(x, y, z) = g(x, y, z) = x$ . As is easily seen, all three 2-element subsets of  $A$  are subuniverses and there is no term operation of  $\mathbb{A}$  that is semilattice on any of the three 2-element subsets. Therefore, each of the pairs  $ab, bc, ac$  is a majority edge as witnessed by the equality relation, and  $\mathbb{A}$  is smooth. Consider  $\mathbb{A}^2$  and  $(a, b), (b, c) \in \mathbb{A}^2$ . Since  $\mathbb{A}^2$  does not have a binary term operation acting as a semilattice operation on these two pairs,  $(a, b)(b, c)$  is a majority edge in  $\mathbb{A}^2$  witnessed by the equality relation. However,  $\{(a, b), (b, c)\}$  is not a subalgebra, because

$$g\left(\begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} b \\ c \end{pmatrix}, \begin{pmatrix} a \\ b \end{pmatrix}\right) = \begin{pmatrix} g(a, b, a) \\ g(b, c, b) \end{pmatrix} = \begin{pmatrix} b \\ b \end{pmatrix}.$$

The next statement uniformizes the operations witnessing the type of edges in smooth algebras.

**Theorem 7** (Theorem 21 and Corollary 22, [10]). *Let  $\mathcal{K}$  be a smooth class. There are term operations  $f, g, h$  of  $\mathcal{K}$  such that for any  $\mathbb{A} \in \mathcal{K}$  and any  $a, b \in \mathbb{A}$ , the operation  $f$  is a semilattice operation on  $\{a/\theta, b/\theta\}$  if  $ab$  is a semilattice edge;  $g$  is a majority operation on  $\{a/\theta, b/\theta\}$  if  $ab$  is a majority edge;  $h$  is the affine operation  $x - y + z$  on  $\text{Sg}(a, b)/\theta$  if  $ab$  is an affine edge, where  $\theta$  witnesses the type of the edge.*

Operations  $f, g, h$  from Theorem 7 above can be chosen to satisfy certain identities.

**Lemma 8** (Lemma 23 of [10]). *Operations  $f, g, h$  identified in Theorem 7 can be chosen such that*

- (1)  $f(x, f(x, y)) = f(x, y)$  for all  $x, y \in \mathbb{A} \in \mathcal{K}$ ;
- (2)  $g(x, g(x, y, y), g(x, y, y)) = g(x, y, y)$  for all  $x, y \in \mathbb{A} \in \mathcal{K}$ ;
- (3)  $h(h(x, y, y), y, y) = h(x, y, y)$  for all  $x, y \in \mathbb{A} \in \mathcal{K}$ .

We will assume that for a class  $\mathcal{K}$  operations  $f, g, h$  satisfying the conditions of Theorem 7 and Lemma 8 are chosen and fixed.

Thin edges also introduced in [10] offer a better technical tool. Note that thin edges are defined for any algebra from the variety generated by  $\mathcal{K}$ .

For a smooth class  $\mathcal{K}$ ,  $\mathbb{A} \in \mathcal{K}$ , and  $a, b \in \mathbb{A}$ , the pair  $ab$  is called a *thin semilattice edge* if the equality relation witnesses that it is a semilattice edge; or in other words if  $f(a, b) = f(b, a) = b$ . The binary operation  $f$  from Theorem 7 can be chosen to satisfy a special property.



**Proposition 9** (Proposition 24, [10]). *Let  $\mathcal{K}$  be a smooth class. There is a binary term operation  $f$  of  $\mathcal{K}$  such that  $f$  is a semilattice operation on every thick semilattice edge of every  $\mathbb{A} \in \mathcal{K}$  and for any  $a, b \in \mathbb{A}$ ,  $\mathbb{A} \in \mathcal{K}$ , either  $a = f(a, b)$  or the pair  $(a, f(a, b))$  is a thin semilattice edge.*

We assume that operation  $f$  satisfying the conditions of Proposition 9 is fixed, and use  $\cdot$  to denote it (think multiplication). If  $ab$  is a thin semilattice edge, that is,  $a \cdot b = b \cdot a = b$ , we write  $a \leq b$ .

Let again  $\mathcal{K}$  be a smooth class and  $\mathcal{V}$  the variety it generates. Let  $\mathbb{A} \in \mathcal{V}$ ,  $a, b \in \mathbb{A}$ ,  $\mathbb{B} = \text{Sg}(a, b)$ , and let  $\theta$  be a congruence of  $\mathbb{B}$ . Pair  $ab$  is said to be *minimal* with respect to  $\theta$  if for any  $b' \in b/\theta$ ,  $b \in \text{Sg}(a, b')$ . A ternary term  $g'$  is said to satisfy the *majority condition* (with respect to  $\mathcal{K}$ ) if it satisfies the conditions of Lemma 8(2) and  $g'$  is a majority operation on every thick majority edge of every algebra from  $\mathcal{K}$ . A ternary term  $h'$  is said to satisfy the *minority condition* if it satisfies the conditions of Lemma 8(3) and for any  $\mathbb{B} \in \mathcal{K}$  and every affine edge  $ab$  of  $\mathbb{B}$  witnessed by a congruence  $\theta$  of  $\text{Sg}_{\mathbb{B}}(a, b)$ , the operation  $h'$  is a Mal'tsev operation on  $\text{Sg}_{\mathbb{B}}(a, b)/\theta$ . By Theorem 7 operations satisfying the majority and minority conditions exist.

Let  $\mathbb{A} \in \mathcal{V}$  and  $a, b \in \mathbb{A}$ . The (ordered) pair  $ab$  is a *thin semilattice edge* if the term  $\cdot$  of  $\mathcal{V}$  is a semilattice operation on  $\{a, b\}$  and  $ab = b$ .

A pair  $ab$  is called a *thin majority edge* if

- (\*) for any term operation  $g'$  satisfying the majority condition the subalgebras  $\text{Sg}(a, g'(a, b, b))$ ,  $\text{Sg}(a, g'(b, a, b))$ ,  $\text{Sg}(a, g'(b, b, a))$  contain  $b$ .

A pair  $ab$  is called a *thin affine edge* if

- (\*\*)  $h(b, a, a) = b$  (where  $h$  is the fixed operation satisfying the conditions of Theorem 7), and for any term operation  $h'$  satisfying the minority condition  $b \in \text{Sg}(a, h'(a, a, b))$ .

The operations  $g, h$  from Theorem 7 do not have to satisfy any specific conditions on the set  $\{a, b\}$ , when  $ab$  is a thin majority or affine edge, except what follows from their definition. Also, both thin majority and thin affine edges are directed, since  $a, b$  in the definition occur asymmetrically. Note also, that what pairs of an algebra  $\mathbb{A}$  are thin majority and affine edges depend not only on the algebra itself, but also on the underlying class  $\mathcal{K}$ . If we are not interested in any particular class, just the algebra itself, set  $\mathcal{K} = \text{HS}(\mathbb{A})$ . We now fix a smooth class  $\mathcal{K}$  and the variety  $\mathcal{V}$  it generates along with operations  $f, g, h$  satisfying the conditions of Theorem 7, Lemma 8, and Proposition 9. All algebras in the rest of the paper are from  $\mathcal{V}$  and smooth ones are from  $\mathcal{K}$ , unless otherwise stated.



**Lemma 10** (Corollary 25, Lemmas 28,32, [10]). *Let  $\mathbb{A} \in \mathcal{K}$ , and let  $ab$  be a semilattice (majority, affine) edge,  $\theta$  a congruence of  $\text{Sg}(a, b)$  that witnesses this, and  $c \in a/\theta$ . Then, if  $ab$  is a semilattice [majority] edge, then for any  $d \in b/\theta$  such that  $cd$  is a minimal pair with respect to  $\theta|_{\text{Sg}(c, d)}$  the pair  $cd$  is a thin semilattice [respectively, thin majority] edge. If  $ab$  is affine then for any  $d \in b/\theta$  such that  $ad$  is a minimal pair with respect to  $\theta|_{\text{Sg}(c, d)}$  and  $h(d, a, a) = d$  the pair  $ad$  is a thin affine edge. Moreover,  $d \in b/\theta$  satisfying these conditions exists.*

The following simple properties of thin edges will be useful. Note that a subdirect product of algebras (a relation) is also an algebra, and so edges and thin edges can be defined for relations as well.

**Proposition 11** (Proposition 8, [11]). *For every  $\mathbb{A} \in \mathcal{V}$  and for any  $a, b \in \mathbb{A}$  either  $a = a \cdot b$  or the pair  $(a, a \cdot b)$  is a thin semilattice edge.*

Items (1) and (2) of the following lemma are Lemma 11 from [11], and item (3) follows from the definitions.

**Lemma 12** (Lemma 11, [11]). *(1) Let  $\mathbb{A} \in \mathcal{V}$ ,  $\overline{ab}$  be a thin edge in  $\mathbb{A}/\theta$ , and  $a \in \overline{a}$ . Then there is  $b \in \overline{b}$  such that  $ab$  is a thin edge in  $\mathbb{A}$  of the same type.*

*(2) Let  $\mathbb{A} \in \mathcal{V}$  and  $ab$  be a thin edge. Then  $ab$  is a thin edge in any subalgebra of  $\mathbb{A}$  containing  $a, b$ , and  $a/\theta b/\theta$  is a thin edge in  $\mathbb{A}/\theta$  for any congruence  $\theta$ .*

*(3) Let  $\mathbb{A} \in \mathcal{V}$  and  $\mathbb{B}$  its subalgebra. Then every thin edge of  $\mathbb{B}$  is a thin edge of  $\mathbb{A}$  of the same type.*

We will need operations that act in a specific way on pairs of thin edges.

**Lemma 13** (Lemma 36, [10], Lemma 9, [11]). *(1) Let  $ab$  be a thin majority edge of an algebra  $\mathbb{A} \in \mathcal{V}$ . There is a term operation  $t_{ab}$  such that  $t_{ab}(a, b) = b$  and  $t_{ab}(c, d) \stackrel{\eta}{=} c$  for all affine edges  $cd$  of all  $\mathbb{B} \in \mathcal{V}$ , where the type of  $cd$  is witnessed by the congruence  $\eta$ .*

*(2) Let  $ab$  be a thin affine edge of an algebra  $\mathbb{A} \in \mathcal{V}$ . There is a term operation  $h_{ab}$  such that  $h_{ab}(a, a, b) = b$  and  $h_{ab}(c, d, d) \stackrel{\eta}{=} c$  for all affine edges  $cd$  of all  $\mathbb{B} \in \mathcal{V}$ , where the type of  $cd$  is witnessed by the congruence  $\eta$ . Moreover,  $h_{ab}(x, c', d')$  is a permutation of  $\text{Sg}(c, d)/\eta$  for any  $c', d' \in \text{Sg}(c, d)$ .*

*(3) Let  $ab$  and  $cd$  be thin edges in  $\mathbb{A}_1, \mathbb{A}_2 \in \mathcal{V}$ . If they have different types there is a binary term operation  $p$  such that  $p(b, a) = b$ ,  $p(c, d) = d$ . If both edges are affine then there is a term operation  $h'$  such that  $h'(a, a, b) = b$  and  $h'(d, c, c) = d$ .*

*Proof.* Item (3) is items (1) and (2) of Lemma 9 from [11]. Lemma 36 from [10] proves items (1) and (2) for smooth algebras from  $\mathcal{K}$ . We generalize this statement for non-smooth algebras from  $\mathcal{V}$ . The proof is similar to that of Lemma 36 from [10], except we need to prove an additional statement that is common for both items (1) and (2).

CLAIM. For any  $\mathbb{B} \in \mathcal{V}$  and any affine edge  $cd$  of  $\mathbb{B}$  (note that  $\mathbb{B}$  does not have to be smooth to have affine edges), where  $\eta$  is the congruence of  $\mathbb{C} = \text{Sg}_{\mathbb{B}}(a, b)$  witnessing that  $cd$  is an affine edge, it holds that  $g(c, d, d) \stackrel{\eta}{\equiv} c$  and  $h(c, d, d) \stackrel{\eta}{\equiv} c$ .

Let  $\mathbb{B}$  be a quotient algebra modulo a congruence  $\theta$  of a subdirect product  $\mathbb{B}'$  of  $\mathbb{A}_1, \dots, \mathbb{A}_k \in \mathcal{K}$ . Pick elements  $c', d'$  from the  $\theta$ -blocks  $c, d$  — elements of  $\mathbb{B}$ , — respectively. Then  $c'd'$  is an affine edge of  $\mathbb{B}'$  as witnessed by the congruence  $\eta' = \{(e_1, e_2) \in \mathbb{B}'^2 \mid e_1/\theta \stackrel{\eta}{\equiv} e_2/\theta\}$ . If we prove that the statement of the Claim is true for  $c'd'$  and  $\eta'$ , the result for  $cd$  and  $\eta$  follows. Therefore, it suffices to assume that  $\mathbb{B}$  is a subdirect product of  $\mathbb{A}_1, \dots, \mathbb{A}_k$ .

Suppose now that  $\mathbb{B}, c, d, \mathbb{C} = \text{Sg}_{\mathbb{B}}(c, d)$ , and  $\eta$  is a minimal counterexample to the statement of the Claim in terms of the number  $k$  and the size of  $\mathbb{C}' = \mathbb{C}/\eta$ . By Lemma 31 from [10] the Claim is true for all affine edges of algebras from  $\mathcal{K}$ . Hence, if  $k = 1$ , we have the result. Otherwise  $\mathbb{C}'$  is a module over some ring  $\mathbb{R}$ , and the operation  $g$  satisfies the identity from Lemma 8(2). It is easy to see that  $g$  on  $\mathbb{C}'$  satisfies the following conditions:  $g(x, y, z) = \alpha x + \beta y + \gamma z$ , where  $\alpha$  and  $\beta + \gamma$  are idempotents of  $\mathbb{R}$  such that  $\alpha(\beta + \gamma) = 0$  and  $\alpha + \beta + \gamma = 1$ . A similar statement holds for  $h$ . Therefore  $\mathbb{C}' = \alpha\mathbb{C}' \oplus (\beta + \gamma)\mathbb{C}'$ . If  $\alpha \neq 0$  (for  $g$  or  $h$  alike) then we can replace  $\mathbb{C}'$  with its submodule  $(\beta + \gamma)\mathbb{C}'$  and  $c, d$  with  $(\beta + \gamma)c, (\beta + \gamma)d$ , obtaining a smaller counterexample. Indeed, if, say,  $g((\beta + \gamma)c, (\beta + \gamma)d, (\beta + \gamma)d) = (\beta + \gamma)c$ , then  $(\beta + \gamma)d = (\beta + \gamma)c$ , and therefore  $g(c, d, d) = \alpha c + (\beta + \gamma)d = \alpha c + (\beta + \gamma)c = c$ , a contradiction. Thus, we can assume that  $\alpha = 0$  and that  $g(x, y, z) = \beta y + \gamma z$  ( $h(x, y, z) = \beta y + \gamma z$ ) on  $\mathbb{C}'$  and  $\beta + \gamma = 1$ .

Consider the projection congruence  $\theta$  of  $\mathbb{C}$ :  $(a_1, \dots, a_k) \stackrel{\theta}{\equiv} (b_1, \dots, b_k)$  if and only if  $a_1 = b_1, \dots, a_{k-1} = b_{k-1}$ . If  $\theta \leq \eta$  then we can replace  $\mathbb{C}$  with  $\text{pr}_{[k-1]}\mathbb{C}$ , as  $g$  (or  $h$ ) does not satisfy the statement of the Claim for any affine edge of  $\text{pr}_{[k-1]}\mathbb{C}$ . Let  $\eta < \eta \vee \theta$ , then  $\eta|_{a/\theta}$  is not the full congruence of the subalgebra  $a/\theta$  for some  $a = (a_1, \dots, a_k) \in \mathbb{C}$ . Note that  $a/\theta$  is isomorphic to the subalgebra  $\mathbb{D} = \{e \mid (a_1, \dots, a_{k-1}, e) \in \mathbb{C}\}$  of  $\mathbb{A}_k$ . Let  $\eta'$  be the congruence of  $\mathbb{D}$  that is the isomorphic image of  $\eta$ . Then  $\mathbb{D}/\eta'$  is a module isomorphic to a subalgebra of  $\mathbb{C}/\eta$ , and therefore every pair of elements of  $\mathbb{D}/\eta'$  is an affine edge. Since  $g(c', d', d') = h(c', d', d') = d'$  for any  $c', d' \in \mathbb{D}/\eta'$  and  $\mathbb{D}/\eta' \in \mathcal{K}$ , we get a contradiction with Lemma 36 from [10]. The Claim is proved.

(1) Let  $b' = g(a, b, b)$ . By the definition of thin majority edges  $b \in \text{Sg}_{\mathbb{A}}(a, b')$  and there is a binary term operation  $r$  such that  $b = r(a, b')$ . By Claim  $g(x, y, y)$  is the first projection on  $\text{Sg}(c, d)/\eta$  for any affine edge  $cd$  of any algebra  $\mathbb{B} \in \mathcal{V}$ , where the congruence  $\eta$  witnesses this fact. Let  $t_{ab}(x, y) = r(x, g(x, y, y))$ . Then

$$\begin{aligned} t_{ab}(a, b) &= r(a, g(a, b, b)) = b, \\ t_{ab}(c, d) &= r(c, g(c, d, d)) \stackrel{\eta}{\equiv} c. \end{aligned}$$

This means that  $t_{ab}$  satisfies the required conditions.

(2) Let  $b' = h(a, a, b)$ . By the definition of thin affine edges  $b \in \text{Sg}_{\mathbb{A}}(a, b')$  and there is a binary term operation  $r$  such that  $b = r(a, b')$ . By Claim  $h(x, y, y)$  is the first projection on  $\text{Sg}(c, d)/\eta$  for any affine edge  $cd$  of any algebra  $\mathbb{B} \in \mathcal{V}$ , where the congruence  $\eta$  witnesses this fact. Let  $h_{ab}(x, y, z) = r(x, h(x, y, z))$ . Then

$$\begin{aligned} h_{ab}(a, a, b) &= r(a, h(a, a, b)) = b, \\ h_{ab}(c, d, d) &= r(c, h(c, d, d)) \stackrel{\eta}{\equiv} c. \end{aligned}$$

This means that  $h_{ab}$  satisfies the required conditions. To prove the second statement let  $c', d' \in \text{Sg}(c, d)$ . Since  $\mathbb{C} = \text{Sg}(c, d)/\eta$  is a module, in particular, it is an Abelian algebra and  $h_{ab}(x, c^*, c^*) = x$  for all  $c^* \in \mathbb{C}$ , the second result follows.  $\square$

## 2.4 Maximality

Let  $\mathbb{A} \in \mathcal{V}$ . A *path* in  $\mathbb{A}$  is a sequence  $a_0, a_1, \dots, a_k$  such that  $a_{i-1} = a_i$  or  $a_{i-1}a_i$  is a thin edge for all  $i \in [k]$  (note that thin edges are always assumed to be directed). We will distinguish paths of several types depending on what types of edges are allowed. If  $a_{i-1} \leq a_i$  for all  $i \in [k]$  then the path is called a *semilattice* or *s-path*. If for every  $i \in [k]$  either  $a_{i-1} \leq a_i$  or  $a_{i-1}a_i$  is a thin affine edge then the path is called *affine-semilattice* or *as-path*. The path is called *asm-path* when all types of edges are allowed. If there is a path  $a = a_0, a_1, \dots, a_k = b$  which is arbitrary (semilattice, affine-semilattice) then  $a$  is said to be *asm-connected* (or *s-connected*, or *as-connected*) to  $b$ . We will also say that  $a$  is *connected* to  $b$  if it is asm-connected. We denote this by  $a \sqsubseteq^{asm} b$  (for asm-connectivity),  $a \sqsubseteq b$ , and  $a \sqsubseteq^{as} b$  for s-, and as-connectivity, respectively.

Let  $\mathcal{G}_s(\mathbb{A})$  ( $\mathcal{G}_{as}(\mathbb{A})$ ,  $\mathcal{G}_{asm}(\mathbb{A})$ ) denote the digraph whose nodes are the elements of  $\mathbb{A}$ , and the edges are the thin semilattice edges (thin semilattice and affine edges, arbitrary thin edges, respectively). The strongly connected component of  $\mathcal{G}_s(\mathbb{A})$  containing  $a \in \mathbb{A}$  will be denoted by  $s(a)$ . The set of strongly connected components of  $\mathcal{G}_s(\mathbb{A})$  are ordered in the natural way (if  $a \leq b$  then  $s(a) \leq s(b)$ ),

the elements belonging to maximal ones will be called *maximal*, and the set of all maximal elements from  $\mathbb{A}$  will be denoted by  $\max(\mathbb{A})$ .

The strongly connected component of  $\mathcal{G}_{as}(\mathbb{A})$  containing  $a \in \mathbb{A}$  will be denoted by  $as(a)$ . A maximal strongly connected component of this graph is called an *as-component*, an element from an as-component is called *as-maximal*, and the set of all as-maximal elements is denoted by  $\max(\mathbb{A})$ .

Finally, the strongly connected component of  $\mathcal{G}_{asm}(\mathbb{A})$  containing  $a \in \mathbb{A}$  will be denoted by  $asm(a)$ . A maximal strongly connected component of  $\mathcal{G}_{asm}(\mathbb{A})$  is called a *universally maximal component* (or *u-maximal component* for short), an element from a u-maximal component is called *u-maximal*, and the set of all u-maximal elements is denoted by  $\max(\mathbb{A})$ .

Sometimes we use the notation  $\mathcal{G}_s(B)$ ,  $\mathcal{G}_{as}(B)$ ,  $\mathcal{G}_{asm}(B)$  and  $\max(B)$ ,  $\max(B)$ ,  $\max(B)$  for a subset  $B$  of  $\mathbb{A}$  that is not necessarily a subalgebra. In this case  $\mathcal{G}_s(B)$ ,  $\mathcal{G}_{as}(B)$ ,  $\mathcal{G}_{asm}(B)$  denote the subgraphs of  $\mathcal{G}_s(\mathbb{A})$ ,  $\mathcal{G}_{as}(\mathbb{A})$ ,  $\mathcal{G}_{asm}(\mathbb{A})$ , respectively, induced by  $B$ , and  $\max(B)$ ,  $\max(B)$ ,  $\max(B)$  refer to elements of the maximal strongly connected components of those subgraphs.

Alternatively, maximal, as-maximal, and u-maximal elements can be characterized as follows: an element  $a \in \mathbb{A}$  is maximal (as-maximal, u-maximal) if for every  $b \in \mathbb{A}$  such that  $a \sqsubseteq b$  ( $a \sqsubseteq^{as} b$ ,  $a \sqsubseteq^{asm} b$ ) it also holds that  $b \sqsubseteq a$  ( $b \sqsubseteq^{as} a$ ,  $b \sqsubseteq^{asm} a$ ). Sometimes it will be convenient to specify what the algebra is, in which we consider maximal components, as-components, or u-maximal components, and the corresponding connectivity. In such cases we will specify it by writing  $s_{\mathbb{A}}(a)$ ,  $as_{\mathbb{A}}(a)$ , or  $asm_{\mathbb{A}}(a)$ . For connectivity we will use  $a \sqsubseteq_{\mathbb{A}} b$ ,  $a \sqsubseteq_{\mathbb{A}}^{as} b$ , and  $a \sqsubseteq_{\mathbb{A}}^{asm} b$ .

As a straightforward implication of Lemma 12(3) we have the following statement.

**Corollary 14.** *Let  $\mathbb{A} \in \mathcal{V}$ ,  $\mathbb{B}$  its subalgebra, and  $a, b \in \mathbb{B}$ . Then if  $a \sqsubseteq_{\mathbb{B}}$  (or  $a \sqsubseteq_{\mathbb{B}}^{as} b$ ,  $a \sqsubseteq_{\mathbb{B}}^{asm} b$ ) then  $a \sqsubseteq_{\mathbb{A}}$  (respectively,  $a \sqsubseteq_{\mathbb{A}}^{as} b$ ,  $a \sqsubseteq_{\mathbb{A}}^{asm} b$ ). Moreover, if  $\mathbb{B}$  contains a maximal component (an as-component)  $C$  of  $\mathbb{A}$ , then  $C$  is a maximal component (as-component) of  $\mathbb{B}$ .*

As the following result shows, the graph  $\mathcal{G}_{asm}(\mathbb{A})$  is well connected.

**Proposition 15** (Corollary 22, [11]). *Let  $\mathbb{A} \in \mathcal{V}$ . Then any  $a, b \in \max(\mathbb{A})$  (or  $a, b \in \max(\mathbb{A})$ , or  $a, b \in \max(\mathbb{A})$ ) are connected in  $\mathcal{G}_{asm}(\mathbb{A})$  with a directed path.*

Since for every  $a \in \mathbb{A}$  there is a maximal  $a' \in \mathbb{A}$  such that  $a \sqsubseteq a'$ , Proposition 15 implies that there is only one u-maximal component. Moreover, Proposition 15 implies the following connection between maximal, as-maximal, and u-maximal elements.

**Corollary 16.** *Let  $\mathbb{A} \in \mathcal{V}$ . Then  $\max(\mathbb{A}), \text{amax}(\mathbb{A}) \subseteq \text{umax}(\mathbb{A})$ .*

U-maximality has an additional useful property, it is somewhat hereditary, as it is made precise in the following

**Lemma 17.** *Let  $\mathbb{B}$  be a subalgebra of  $\mathbb{A} \in \mathcal{V}$ , containing a u-maximal element of  $\mathbb{A}$ . Then every element u-maximal in  $\mathbb{B}$  is also u-maximal in  $\mathbb{A}$ . In particular, if  $\alpha$  is a congruence of  $\mathbb{A}$  and  $\mathbb{B}$  is a u-maximal  $\alpha$ -block, that is,  $\mathbb{B}$  is a u-maximal element in  $\mathbb{A}/\alpha$ , then  $\text{umax}(\mathbb{B}) \subseteq \text{umax}(\mathbb{A})$ .*

*Proof.* Let  $a \in \mathbb{B}$  be an element u-maximal in  $\mathbb{A}$ , let  $b \in \text{umax}(\mathbb{B})$ . For any  $c \in \mathbb{A}$  with  $b \sqsubseteq_{\mathbb{A}}^{asm} c$  we also have  $c \sqsubseteq_{\mathbb{A}}^{asm} a$ . Finally, since  $b \in \text{umax}(\mathbb{B})$  and  $a \in \mathbb{B}$ , we have  $a \sqsubseteq_{\mathbb{A}}^{asm} b$ , and therefore  $c \sqsubseteq_{\mathbb{A}}^{as} b$ . For the second part of the lemma we need to find a u-maximal element in  $\mathbb{B}$ . Let  $b \in \text{umax}(\mathbb{A})$ . Then as  $\mathbb{B}$  is u-maximal in  $\mathbb{A}/\alpha$  applying Lemma 12(1) we get that there is  $a' \in \mathbb{B}$  such that  $b \sqsubseteq_{\mathbb{A}}^{asm} a'$ . Clearly,  $a' \in \text{umax}(\mathbb{A})$ .  $\square$

**Lemma 18** (The Maximality Lemma, Lemma 15, Corollaries 16,17, [11]). *Let  $R$  be a subdirect product of  $\mathbb{A}_1, \dots, \mathbb{A}_n \in \mathcal{V}$ , and  $I \subseteq [n]$ .*

- (1) *For any  $\mathbf{a} \in R$ ,  $\mathbf{a}^* = \text{pr}_I \mathbf{a}$ ,  $\mathbf{b} \in \text{pr}_I R$  such that  $\mathbf{a}^* \mathbf{b}$  is a thin edge, there is  $\mathbf{b}' \in R$ ,  $\text{pr}_I \mathbf{b}' = \mathbf{b}$ , such that  $\mathbf{a} \mathbf{b}'$  is a thin edge of the same type.*
- (2) *If  $\mathbf{a} \mathbf{b}$  is a thin edge in  $R$  then  $\text{pr}_I \mathbf{a} \text{pr}_I \mathbf{b}$  is a thin edge in  $\text{pr}_I R$  of the same type (including the possibility that  $\text{pr}_I \mathbf{a} = \text{pr}_I \mathbf{b}$ ).*
- (3) *For any  $\mathbf{a} \in R$ , and an s- (as-, asm-) path  $\mathbf{b}_1, \dots, \mathbf{b}_k \in \text{pr}_I R$  with  $\text{pr}_I \mathbf{a} = \mathbf{b}_1$ , there is an s- (as-, asm-) path  $\mathbf{b}'_1, \dots, \mathbf{b}'_k \in R$  such that  $\mathbf{b}'_1 = \mathbf{a}$  and  $\text{pr}_I \mathbf{b}'_k = \mathbf{b}_k$ .*
- (4) *If  $\mathbf{a}_1, \dots, \mathbf{a}_k$  is an s- (as-, asm-) path in  $R$ , then  $\text{pr}_I \mathbf{a}_1, \dots, \text{pr}_I \mathbf{a}_k$  is an s- (as-, asm-) path in  $\text{pr}_I R$ .*
- (5) *For any maximal (as-maximal, u-maximal) (in  $\text{pr}_I R$ ) element  $\mathbf{b} \in \text{pr}_I R$ , there is  $\mathbf{b}' \in R$  which is maximal (as-maximal, u-maximal) in  $R$  and such that  $\text{pr}_I \mathbf{b}' = \mathbf{b}$ . In particular,  $\text{pr}_{[n]-I} \mathbf{b}'$  is a maximal (as-maximal, u-maximal) in  $\text{pr}_{[n]-I} R$ .*
- (6) *If  $\mathbf{a}$  is a maximal (as-maximal, u-maximal) in  $R$ , then  $\text{pr}_I \mathbf{a}$  is maximal (as-maximal, u-maximal) in  $\text{pr}_I R$ .*

We will need the following statement that easily follows from Lemma 18(3).

**Corollary 19.** *Let  $R$  be a subdirect product of  $\mathbb{A}_1, \dots, \mathbb{A}_n \in \mathcal{V}$ ,  $I \subseteq [n]$ ,  $B, C$  as-components of  $\text{pr}_I R$  and  $\text{pr}_{[n]-I} R$ , respectively, such that  $R' = R \cap (B \times C) \neq \emptyset$ . Then  $R'$  is a subdirect product of  $B$  and  $C$ .*

The following lemma considers a special case of maximal components (as well as as- and u-components) in subdirect products, and is straightforward.

**Lemma 20** (Lemma 18, [11]). *Let  $R$  be a subdirect product of  $\mathbb{A}_1, \mathbb{A}_2 \in \mathcal{V}$ ,  $B, C$  maximal components (as-components, u-components) of  $\mathbb{A}_1, \mathbb{A}_2$ , respectively, and  $B \times C \subseteq R$ . Then  $B \times C$  is a maximal component (as-component, u-component) of  $R$ .*

Lemma 12 applied to paths implies

**Lemma 21.** *Let  $\mathbb{A} \in \mathcal{V}$ , let  $\theta$  be a congruence of  $\mathbb{A}$ .*

- (1) *If  $C$  is a maximal component (as-component, u-component) of  $\mathbb{A}$ , then  $C/\theta$  is a maximal component (as-component, u-component) of  $\mathbb{A}/\theta$ .*
- (2) *If  $C$  is a maximal component (as-component, u-component) of  $\mathbb{A}/\theta$ , then every  $\bar{a} \in C$  contains a maximal (as-maximal, u-maximal) element of  $\mathbb{A}$ .*

We complete this section with an auxiliary statement that will be needed later.

**Lemma 22.** *Let  $\mathbb{A} \in \mathcal{V}$ ,  $\alpha \prec \beta$ ,  $\alpha, \beta \in \text{Con}(\mathbb{A})$ , let  $B$  be a  $\beta$ -block and  $\text{tp}(\alpha, \beta) = \mathbf{2}$ . Then  $B/\alpha$  is term equivalent to a module. In particular, every pair of elements of  $B/\alpha$  is a thin affine edge in  $\mathbb{A}/\alpha$ .*

*Proof.* As  $\mathbb{A}$  is an idempotent algebra that generates a variety omitting type **1**, and  $(\alpha, \beta)$  is a prime interval in  $\text{Con}(\mathbb{A})$  of type **2**, by [18, Theorem 7.12] there is a term operation of  $\mathbb{A}$  that is Mal'tsev on  $B/\alpha$ . Since  $\beta$  is Abelian on  $B/\alpha$ , we get the result.  $\square$

## 2.5 Quasi-decomposition and quasi-majority

We make use of the property of quasi-2-decomposability proved in [11].

**Theorem 23** (The Quasi-2-Decomposition Theorem, Corollary 31, [11]). *Let  $\mathbb{A}_1, \dots, \mathbb{A}_n \in \mathcal{V}$ . If  $R$  is a subdirect product of  $\mathbb{A}_1, \dots, \mathbb{A}_n$  and tuple  $\mathbf{a}$  is such that  $\text{pr}_J \mathbf{a} \in \text{pr}_J R$  for any  $J \subseteq [n]$ ,  $|J| = 2$ , then there is a tuple  $\mathbf{b} \in R$  with  $\text{pr}_J \mathbf{a} \sqsubseteq^{as} \text{pr}_J \mathbf{b}$  for any  $J \subseteq [n]$ ,  $|J| = 2$ .*

One useful implication of the Quasi-2-Decomposition Theorem 23 is the existence of a term operation resembling a majority operation.

**Theorem 24** (Theorem 32, [11]). *There is a term operation  $\text{maj}$  of  $\mathcal{V}$  such that for any  $\mathbb{A} \in \mathcal{V}$  and any  $a, b \in \mathbb{A}$ , it holds  $a \sqsubseteq^{as} \text{maj}(a, a, b), \text{maj}(a, b, a), \text{maj}(b, a, a)$ .*

*In particular, if  $a$  is as-maximal, then  $\text{maj}(a, a, b), \text{maj}(a, b, a), \text{maj}(b, a, a)$  belong to  $\text{as}(a)$ .*

A function  $\text{maj}$  satisfying the properties from Theorem 24 is called a *quasi-majority operation*.

## 2.6 Rectangularity

Let  $R$  be a subdirect product of (arbitrary) algebras  $\mathbb{A}_1, \mathbb{A}_2$ . By  $R[c], R^{-1}[c']$  for  $c \in \mathbb{A}_1, c' \in \mathbb{A}_2$  we denote the sets  $\{b \mid (c, b) \in R\}, \{a \mid (a, c') \in R\}$ , respectively, and for  $C \subseteq \mathbb{A}_1, C' \subseteq \mathbb{A}_2$  we use  $R[C] = \bigcup_{c \in C} R[c], R^{-1}[C'] = \bigcup_{c' \in C'} R^{-1}[c']$ , respectively. Binary relations  $\text{tol}_1, \text{tol}_2$  on  $\mathbb{A}_1, \mathbb{A}_2$  given by  $\text{tol}_1(R) = \{(a, b) \mid R[a] \cap R[b] \neq \emptyset\}$  and  $\text{tol}_2(R) = \{(a, b) \mid R^{-1}[a] \cap R^{-1}[b] \neq \emptyset\}$ , respectively, are called *link tolerances* of  $R$ . They are tolerances of  $\mathbb{A}_1, \mathbb{A}_2$ , respectively, that is, invariant reflexive and symmetric relations. The transitive closures  $\text{lk}_1, \text{lk}_2$  of  $\text{tol}_1(R), \text{tol}_2(R)$  are called *link congruences*, and they are, indeed, congruences. Relation  $R$  is said to be *linked* if the link congruences are full congruences.

**Lemma 25** (Lemma 24, [11]). *Let  $R$  be a subalgebra of  $\mathbb{A}_1 \times \mathbb{A}_2$ ,  $\mathbb{A}_1, \mathbb{A}_2 \in \mathcal{V}$ , and let  $a \in \mathbb{A}_1$  and  $B = R[a]$ . For any  $b \in \mathbb{A}_1$  such that  $ab$  is thin edge, and any  $c \in R[b] \cap B, d \in R[b]$  whenever  $c \sqsubseteq^{as} d$  in  $B$ .*

Any subalgebra  $R$  of a direct product of Mal'tsev algebras satisfy the *rectangularity* property: if  $(a, c), (a, d), (b, d) \in R$  then  $(b, c) \in R$ . This implies in particular that for any  $\text{lk}_1$ -block  $B_1$  and any  $\text{lk}_2$ -block  $B_2$ , it holds  $B_1 \times B_2 \subseteq R$  whenever  $R \cap (B_1 \times B_2) \neq \emptyset$ . The following two statements proved in [11] make this property more general.

**Proposition 26** (Corollary 27, [11]). *Let  $R$  be a subdirect product of  $\mathbb{A}_1, \mathbb{A}_2 \in \mathcal{V}$ ,  $\text{lk}_1, \text{lk}_2$  the link congruences, and let  $B_1, B_2$  be as-components of an  $\text{lk}_1$ -block and an  $\text{lk}_2$ -block, respectively, such that  $R \cap (B_1 \times B_2) \neq \emptyset$ . Then  $B_1 \times B_2 \subseteq R$ .*

**Proposition 27** (Proposition 28, [11]). *Let  $R$  be a subdirect product of  $\mathbb{A}_1, \mathbb{A}_2 \in \mathcal{V}$ ,  $\text{lk}_1, \text{lk}_2$  the link congruences, and let  $B_1$  be an as-component of an  $\text{lk}_1$ -block and  $B_2 = R[B_1]$ . Then  $B_1 \times \text{umax}(B_2) \subseteq R$ .*

We complete this section with a technical lemma that will be useful later.

**Lemma 28.** *Let  $\mathbb{A} \in \mathcal{V}$  be an algebra and  $C$  its as-component such that  $\mathbb{A} = \text{Sg}(C)$ , let  $R = \mathbb{A} \times \mathbb{A} = \text{Sg}(C \times C)$ , and let  $\alpha$  be a congruence of  $R$ . Then for some  $a, b \in C, a \neq b$ , the pair  $(a, b)$  is as-maximal in an  $\alpha$ -block.*



*Proof.* We start with a general claim.

CLAIM. If  $\beta, \gamma \in \text{Con}(R)$  are such that  $\beta \vee \gamma = \underline{1}_R$ , then  $\beta|_{C^2} \circ \gamma|_{C^2} = \gamma|_{C^2} \circ \beta|_{C^2} = C^2 \times C^2$ .

Let  $R_1 \subseteq R/\beta \times R$ ,  $R_2 \subseteq R/\gamma \times R$  be given by

$$R_1 = \{(a/\beta, a) \mid a \in R\}, \quad R_2 = \{(a/\gamma, a) \mid a \in R\}.$$

Consider a subdirect product  $Q$  of  $R/\beta \times R/\gamma$  defined as follows

$$S(x, y, z) = R_1(x, z) \wedge R_2(y, z),$$

and  $Q = \text{pr}_{12}S$ . As is easily seen, for a  $\beta$ -block  $B_1$  and a  $\gamma$ -block  $B_2$ ,  $(B_1, B_2) \in Q$  if and only if  $B_1 \cap B_2 \neq \emptyset$ . As  $\beta \vee \gamma = \underline{1}_R$ , the relation  $Q$  is linked. By  $C^2/\beta, C^2/\gamma$  we denote the sets of  $\beta$ - and  $\gamma$ -blocks of  $R$  that intersect  $C^2$ . By Lemma 21  $C^2/\beta$  is an as-component of  $R/\beta$  and  $C^2/\gamma$  is an as-component of  $R/\gamma$ . Therefore, Proposition 26 implies that  $C^2/\beta \times C^2/\gamma \subseteq Q$ . Therefore for any  $\beta$ - and  $\gamma$ -blocks  $B_1, B_2$  such that  $B_1 \cap C^2 \neq \emptyset, B_2 \cap C^2 \neq \emptyset$  we have  $B_1 \cap B_2 \neq \emptyset$ .

Now, consider the relation  $S$  defined above. For any  $a \in C^2$  we have  $(a/\beta, a/\gamma, a) \in S$ . By Lemma 20  $C^2$  is an as-component of  $R$ , and therefore, again by Lemma 20,  $C^2 \times C^2$  is an as-component of  $R^2$ . By Lemma 21  $C^2/\beta \times C^2/\gamma$  is an as-component of  $R/\beta \times R/\gamma$ . Since  $C^2/\beta \times C^2/\gamma \subseteq Q$ , by Corollary 14  $C^2/\beta \times C^2/\gamma$  is an as-component of  $Q$ . Thus, for any  $(B_1, B_2) \in C^2/\beta \times C^2/\gamma$ , we have  $(a/\beta, a/\gamma) \sqsubseteq^{as} (B_1, B_2)$  in  $Q$ . By the Maximality Lemma 18(3) there is  $b \in C^2$ ,  $a \sqsubseteq^{as} b$ , such that  $(B_1, B_2, b) \in S$ . The element  $b$  belongs to  $B_1 \cap B_2 \cap C^2$ , as required.

Let  $\beta$  be a maximal congruence of  $\mathbb{A}$  and  $\gamma_1 = \beta \times \underline{1}_{\mathbb{A}}, \gamma_2 = \underline{1}_{\mathbb{A}} \times \beta$ . As is easily seen,  $\gamma_1, \gamma_2$  are maximal congruences of  $R$ , and  $\gamma_1 \wedge \gamma_2 = \beta \times \beta$ . Indeed, if  $\delta \in \text{Con}(R)$  is such that  $\gamma_1 \subsetneq \delta$ , there are  $(a_1, b_1) \stackrel{\delta}{\equiv} (a_2, b_2)$  such that  $(a_1, a_2) \notin \beta$ . Therefore for any  $c_1, c_2 \in \mathbb{A}$  there are  $d_1, d_2 \in \mathbb{A}$  with  $(c_1, d_1) \stackrel{\delta}{\equiv} (c_2, d_2)$ . Since  $\gamma_1 \subseteq \delta$ , this holds for any  $d_1, d_2 \in \mathbb{A}$ . The statement for  $\gamma_2$  is similar and  $\gamma_1 \wedge \gamma_2 = \beta \times \beta$  is straightforward. There are two cases.

CASE 1.  $\alpha \vee (\beta \times \beta) = \underline{1}_R$ .

By the Claim for any  $\beta$ -blocks  $B_1, B_2$  such that  $B_1 \cap C, B_2 \cap C \neq \emptyset$  and an  $\alpha$ -block  $B$  with  $B \cap C^2 \neq \emptyset$ , we also have  $B \cap (B_1 \times B_2) \cap C^2 \neq \emptyset$ . As  $C$  generates  $\mathbb{A}$  and  $\beta$  is a proper congruence,  $C$  intersects with at least two distinct  $\beta$ -blocks, and therefore one can assume  $B_1 \neq B_2$ . Fix such  $B_1, B_2$  and let  $(a, b) \in B \cap (B_1 \times B_2) \cap C^2$ . Let  $(a', b') \in \text{amax}(B)$  be such that  $(a, b) \sqsubseteq_B^{as} (a', b')$ . Since  $a, b \in C$  and  $a \sqsubseteq_{\mathbb{A}}^{as} a', b \sqsubseteq_{\mathbb{A}}^{as} b'$ , we have  $a', b' \in C$ . Let  $(a', b') \in B'_1 \times B'_2$ , where  $B'_1, B'_2$  are  $\beta$ -blocks. By Lemma 12(2)  $(B_1, B_2) \sqsubseteq^{as} (B'_1, B'_2)$  in  $R/\beta \times \beta$ . As  $(B_1, B_2)$

is as-maximal in  $R/\beta \times \beta$ , there is also an as-path from  $(B'_1, B'_2)$  to  $(B_1, B_2)$  in  $R/\beta \times \beta$ . Let  $(B'_1, B'_2) = (B_1^1, B_2^1), (B_1^2, B_2^2), \dots, (B_1^\ell, B_2^\ell) = (B_1, B_2)$ . Since  $a', b' \in C$ , by Lemma 12(1) every block  $B_1^i \times B_2^i$  contains a pair from  $C^2$ . By the Claim it means that  $B \cap (B_1^i \times B_2^i) \cap C^2 \neq \emptyset$  for every  $i \in [\ell]$ . Consider the sequence  $B \cap (B_1^1 \times B_2^1), \dots, B \cap (B_1^\ell \times B_2^\ell)$  of blocks of the congruence  $\beta' = (\beta \times \beta)|_B$ . Since  $B/\beta'$  is a subalgebra of  $R/\beta \times \beta$ <sup>1</sup>, by Lemma 12(2) this sequence is an as-path in  $B/\beta'$ . By Lemma 12(1) there are  $(a', b') = (a^1, b^1), \dots, (a^\ell, b^\ell)$  such that  $(a^i, b^i) \in B \cap (B_1^i \times B_2^i)$ , which form an as-path in  $B$ . Since  $(a', b') \in \text{amax}(B)$ , we also have  $(a^\ell, b^\ell) \in \text{amax}(B)$ , and as  $a', b' \in C$ , we have  $a^\ell, b^\ell \in C$ . Finally, as  $(a^\ell, b^\ell) \in B_1 \times B_2$  and  $B_1 \neq B_2$ , we have  $a^\ell \neq b^\ell$ , as required.

CASE 2.  $\alpha \vee (\beta \times \beta) \neq \underline{1}_R$ .

In this case consider  $\mathbb{A}' = \mathbb{A}/\beta$ ,  $R' = R/\beta \times \beta$ ,  $\alpha' = (\alpha \vee (\beta \times \beta))/\beta \times \beta$ ; note that  $\mathbb{A}'$  is a simple idempotent algebra, and as  $R = \mathbb{A} \times \mathbb{A}$ , we have  $R' = \mathbb{A}' \times \mathbb{A}'$ . The idea here is to find an  $\alpha'$ -block  $D$  that does not contain pairs of the form  $(a, a)$  and find a pair  $(a, b) \in D \cap C^2$ . Then, if  $B$  denotes the  $\alpha$ -block containing  $(a, b)$ , then any  $(a', b') \in \text{amax}(B)$  such that  $(a, b) \sqsubseteq_B^{as} (a', b')$  satisfies the required conditions. Indeed,  $(a', b') \in C^2$ , because  $a \sqsubseteq_{\mathbb{A}}^{as} a', b \sqsubseteq_{\mathbb{A}}^{as} b'$ , and  $(a, b) \in C^2$ .

By [20] and [24, 23] (see also [10, Proposition 3]) either  $\mathbb{A}'$  has an *absorbing element*  $a$ , that is,  $f(a_1, \dots, a_k) = a$  for any term operation  $f$  of  $\mathbb{A}'$ , whenever  $a_i = a$  for some essential variable  $x_i$  of  $f$ , or  $\mathbb{A}'$  is a module, or the only nontrivial congruences of  $\mathbb{A}'^2$  are  $\gamma'_1 = \gamma_1/\beta \times \beta$ ,  $\gamma'_2 = \gamma_2/\beta \times \beta$ . Since  $C$  is a nontrivial as-component, the first option is impossible. Indeed, if  $a$  is an absorbing element, for any  $b \in \mathbb{A}'$  the pair  $ba$  is a thin semilattice edge, and  $\mathbb{A}'$  has only one as-component,  $\{a\}$ . By Lemma 21  $C/\beta = \{a\}$ , which contradicts the assumptions that  $\beta$  is proper and  $C$  generates  $\mathbb{A}$ . If  $\mathbb{A}'$  is a simple module, the only proper nonzero congruence that is different from  $\gamma'_1, \gamma'_2$  is the skew congruence with  $\Delta = \{(a, a) \mid a \in \mathbb{A}'\}$  as a congruence block. If  $\alpha'$  is the skew congruence, let  $D$  be any  $\alpha \vee (\beta \times \beta)$ -block different from  $\Delta$ . As  $C$  is not contained in a  $\beta$ -block,  $D$  can be chosen such that  $D \cap C^2 \neq \emptyset$  (in fact, as  $\mathbb{A}'$  is a module, it is not difficult to see that any  $\alpha \vee (\beta \times \beta)$ -block satisfies this condition). Then we complete the proof as indicated in the beginning of Case 2.

So, suppose  $\alpha \leq \gamma_1$ . If  $\alpha \leq \gamma_1 \wedge \gamma_2 = \beta \times \beta$ , choose a  $\beta \times \beta$ -block  $B_1 \times B_2$  such that  $B_1 \neq B_2$  and  $B_1 \cap C, B_2 \cap C \neq \emptyset$ ; clearly  $B_1, B_2$  are  $\beta$ -blocks. Then for any  $\alpha$ -block  $B \subseteq B_1 \times B_2$  such that  $B \cap (B_1 \times B_2) \cap C^2 \neq \emptyset$  we can find an element  $(a, b) \in B \cap C^2$  and  $a \neq b$  as required.

Finally, suppose  $\alpha \not\leq \gamma_2$ , then  $\alpha \vee \gamma_2 = \underline{1}_R$ . Take an  $\alpha$ -block  $B$ ,  $B \cap C^2 \neq \emptyset$ , we have  $B \subseteq B_1 \times \mathbb{A}$  for some  $\beta$ -block  $B_1$ . Moreover, by the Claim for any  $\beta$ -

<sup>1</sup>Strictly speaking instead of  $B/\beta'$  we need to use  $B^{\beta \times \beta}/\beta \times \beta$ , where  $B^{\beta \times \beta}$  is the union of all  $\beta \times \beta$ -blocks of  $R$  intersecting with  $B$ , see [14, Definition 6.16].

block  $B_2$  with  $B_2 \cap C \neq \emptyset$  there is  $(a, b) \in B \cap C^2$  such that  $b \in B_2$ . Choose  $B_2 \neq B_1$ . Then we complete the proof as in Case 1.  $\square$

### 3 Separating congruences

In this section we introduce and study the relationship between prime intervals in the congruence lattice of an algebra, or in the congruence lattices of factors in subdirect products. It was first introduced in [1] and used in the CSP research in [5, 8].

#### 3.1 Special polynomials, mapping pairs

We start with several technical results. They demonstrate the connection between minimal sets of an algebra  $\mathbb{A} \in \mathcal{V}$  and the structure of its graph  $\mathcal{G}_{asm}(\mathbb{A})$ . Let  $\mathbb{A} \in \mathcal{V}$  be an algebra and let  $Q_{ab}^{\mathbb{A}}$ ,  $a, b \in \mathbb{A}$ , denote the subdirect product of  $\mathbb{A}^2$  generated by  $\{(x, x) \mid x \in \mathbb{A}\} \cup \{(a, b)\}$ .

**Lemma 29.** *Let  $\mathbb{A} \in \mathcal{V}$  and  $a, b \in \mathbb{A}$ .*

- (1)  $Q_{ab}^{\mathbb{A}} = \{(f(a), f(b)) \mid f \in \text{Pol}_1(\mathbb{A})\}$ .
- (2) *For any  $f \in \text{Pol}_1(\mathbb{A})$ ,  $(f(a), f(b)) \in \text{tol}_1(Q_{ab}^{\mathbb{A}})$ . In particular,  $\text{lk}_1(Q_{ab}^{\mathbb{A}}) = \text{Cg}(a, b)$ ; denote this congruence by  $\alpha$ .*
- (3)  $Q_{ab}^{\mathbb{A}} \subseteq \alpha$ .
- (4) *Let  $B$  be an  $\alpha$ -block, and  $C_1, C_2$  as-components of  $B$  such that  $f(a) \in C_1$  and  $f(b) \in C_2$  for a polynomial  $f \in \text{Pol}_1(\mathbb{A})$ . Then  $C_1 \times C_2 \subseteq Q_{ab}^{\mathbb{A}}$ .*

*Proof.* (1) follows directly from the definitions.

(2) Take  $f \in \text{Pol}_1(\mathbb{A})$  and let  $f(x) = g(x, a_1, \dots, a_k)$  for a term operation  $g$  of  $\mathbb{A}$ . Then  $\begin{pmatrix} f(a) \\ f(b) \end{pmatrix} = g\left(\begin{pmatrix} a \\ b \end{pmatrix}, \begin{pmatrix} a_1 \\ a_1 \end{pmatrix}, \dots, \begin{pmatrix} a_k \\ a_k \end{pmatrix}\right) \in Q_{ab}^{\mathbb{A}}$  and  $(f(b), f(b)) \in Q_{ab}^{\mathbb{A}}$  since  $\{(x, x) \mid x \in \mathbb{A}\} \subseteq Q_{ab}^{\mathbb{A}}$  by definition. Thus  $(f(a), f(b)) \in \text{tol}_1(Q_{ab}^{\mathbb{A}})$ . This implies  $\text{Cg}(a, b) \subseteq \text{lk}_1(Q_{ab}^{\mathbb{A}})$ . Then (1) implies that  $\text{lk}_1(Q_{ab}^{\mathbb{A}}) \subseteq \text{Cg}(a, b)$ .

(3) follows from (1), and (4) follows from (2), (3), and Proposition 26.  $\square$

Lemma 29(4) immediately implies

**Corollary 30.** *Let  $\mathbb{A} \in \mathcal{V}$  and  $\alpha \in \text{Con}(\mathbb{A})$  such that  $\underline{0} \prec \alpha$ . Then for any  $a, b \in \mathbb{A}$  with  $a \stackrel{\alpha}{\equiv} b$ ,  $a \neq b$ , and any  $c, d \in \mathbb{A}$  such that  $c \stackrel{\alpha}{\equiv} d$  and  $c, d$  belong to the same as-component of  $c/\alpha$ , there is  $f \in \text{Pol}_1(\mathbb{A})$  such that  $c = f(a)$ ,  $d = f(b)$ .*

**Corollary 31.** *Let  $\mathbb{A} \in \mathcal{V}$ ,  $\alpha \in \text{Con}(\mathbb{A})$  such that  $\underline{0} \prec \alpha$ , and let  $c, d \in \mathbb{A}$ ,  $c \stackrel{\alpha}{\equiv} d$ ,  $c \neq d$ , be as-maximal in  $B = c/\alpha$ .*

- (1) *If  $c, d$  belong to the same as-component of  $B$ , then  $\{c, d\}$  is a  $(\underline{0}, \alpha)$ -subtrace.*
- (2) *If there is a  $(\underline{0}, \alpha)$ -subtrace  $\{c', d'\}$  such that  $c' \in \text{as}(c)$  and  $d' \in \text{as}(d)$  then  $\{c, d\}$  is a  $(\underline{0}, \alpha)$ -subtrace as well.*

*Proof.* (1) Take any  $(\underline{0}, \alpha)$ -minimal set  $U$ , and  $a, b \in U$  with  $a \stackrel{\alpha}{\equiv} b$ ,  $a \neq b$ . By Corollary 30 there is  $f \in \text{Pol}_1(\mathbb{A})$  with  $c = f(a)$ ,  $d = f(b)$ . By Lemma 1(2)  $U' = f(U)$  is a  $(\underline{0}, \alpha)$ -minimal set.

(2) Since  $(c', d') \in Q_{c'd'}^{\mathbb{A}}$ , we have  $Q_{c'd'}^{\mathbb{A}} \cap (\text{as}_B(c) \times \text{as}_B(d)) \neq \emptyset$ . By Lemma 29(4),  $\text{as}_B(c) \times \text{as}_B(d) \subseteq Q_{c'd'}^{\mathbb{A}}$ , in particular,  $(c, d) \in Q_{c'd'}^{\mathbb{A}}$ , hence there is a polynomial  $f$  such that  $f(c') = c$ ,  $f(d') = d$ . Let  $U$  be a  $(\underline{0}, \alpha)$ -minimal set containing  $c', d'$ . Then  $f(U)$  is a  $(\underline{0}, \alpha)$ -minimal set containing  $c, d$ .  $\square$

**Lemma 32.** *Let  $\mathbb{A} \in \mathcal{V}$ . For any  $\alpha \in \text{Con}(\mathbb{A})$  with  $\underline{0} \prec \alpha$  such that  $|D| > 1$  for some as-component  $D$  of an  $\alpha$ -block, the prime interval  $(\underline{0}, \alpha)$  has type **2** or **3**.*

*Proof.* Let  $a, b \in D$  for an as-component  $D$  of an  $\alpha$ -block. Then by Corollary 30 there is a polynomial  $f$  such that  $f(a) = b$  and  $f(b) = a$ . Also,  $a, b$  belong to some  $(\underline{0}, \alpha)$ -minimal set. This rules out types **4** and **5**. Since  $\mathbb{A}$  omits type **1**, this only leaves types **2** and **3**.  $\square$

**Lemma 33.** *Let  $\mathbb{A} \in \mathcal{V}$ , and let  $\alpha \in \text{Con}(\mathbb{A})$  with  $\underline{0} \prec \alpha$  be such that some  $\alpha$ -block contains a semilattice or majority edge. Then the prime interval  $(\underline{0}, \alpha)$  has type **3**, **4** or **5**.*

*Proof.* We need to show that  $(\underline{0}, \alpha)$  does not have type **2**. Let  $B$  be the  $\alpha$ -block containing a semilattice or majority edge. Then  $B$  contains a non-Abelian subalgebra. By Lemma 22 if  $(\underline{0}, \alpha)$  has type **2**, then  $B$  is term equivalent to a module. Since every subalgebra of a module is also term equivalent to a module,  $(\underline{0}, \alpha)$  cannot have type **2**.  $\square$

### 3.2 Separation

The first several definitions and results of this section are valid for arbitrary algebras, not only for algebras from  $\mathcal{V}$ .

Let  $\mathbb{A}$  be an algebra, and let  $\alpha \prec \beta$  and  $\gamma \prec \delta$  be prime intervals in  $\text{Con}(\mathbb{A})$ . We say that  $(\alpha, \beta)$  can be *separated* from  $(\gamma, \delta)$  if there is a unary polynomial  $f \in \text{Pol}_1(\mathbb{A})$  such that  $f(\beta) \not\subseteq \alpha$ , but  $f(\delta) \subseteq \gamma$ . The polynomial  $f$  in this case is said to *separate*  $(\alpha, \beta)$  from  $(\gamma, \delta)$ .

Since we often consider relations rather than single algebras, we also introduce separability in a slightly different way. Let  $R$  be a subdirect product of  $\mathbb{A}_1, \dots, \mathbb{A}_n$ . Let  $I, J \subseteq [n]$  and let  $\alpha_I \prec \beta_I, \alpha_J \prec \beta_J$  be prime intervals in  $\text{Con}(\text{pr}_I R)$  and  $\text{Con}(\text{pr}_J R)$ , respectively. Interval  $(\alpha_I, \beta_I)$  can be separated from  $(\alpha_J, \beta_J)$  if there is a unary polynomial  $f$  of  $R$  such that  $f(\beta_I) \not\subseteq \alpha_I$  but  $f(\beta_J) \subseteq \alpha_J$ . Similarly, the polynomial  $f$  in this case is said to *separate*  $(\alpha_I, \beta_I)$  from  $(\alpha_J, \beta_J)$ .

First, we observe a connection between separation in a single algebra and in relations. Clearly  $\alpha_I, \alpha_J, \beta_I, \beta_J$  give rise to congruences of  $R$  and therefore separation as defined above can be expressed as separation of intervals in  $\text{Con}(R)$ , although one needs to be careful to preserve the primality of congruence intervals. But we can go a bit further.

**Lemma 34.** (1) *Let  $R$  be the binary equality relation on an arbitrary idempotent algebra  $\mathbb{A}$ . Let  $\alpha_1 = \alpha, \beta_1 = \beta$  be viewed as congruences of the first factor of  $R$ , and  $\alpha_2 = \gamma, \beta_2 = \delta$  as congruences of the second factor of  $R$ . The prime interval  $(\alpha, \beta)$  can be separated from  $(\gamma, \delta)$  as intervals in  $\text{Con}(\mathbb{A})$  if and only if  $(\alpha_1, \beta_1)$  can be separated from  $(\alpha_2, \beta_2)$  in  $R$ .*

(2) *Let  $R$  be a subdirect product of arbitrary idempotent algebras  $\mathbb{A}_1, \dots, \mathbb{A}_n$ ,  $I, J \subseteq [n]$ , and  $R^*$  constructed as follows:  $K = I \cap J$ , say,  $K = [k]$  and  $K^* = \{n+1, \dots, n+k\}$ ,*

$$R^* = \{(a_1, \dots, a_n, a_1, \dots, a_k) \mid (a_1, \dots, a_n) \in R\},$$

*$I^* = I, J^* = (J-K) \cup K^*$ . Then  $\text{pr}_I R, \text{pr}_J R$  are isomorphic to  $\text{pr}_{I^*} R^*, \text{pr}_{J^*} R^*$ , let  $\varphi_I, \varphi_J$  be the isomorphisms.*

- *For every polynomial  $f$  of  $R$  there is a polynomial  $f^*$  of  $R^*$  such that  $f^*$  acts on  $\text{pr}_{I^*} R^* \text{pr}_{J^*} R^*$  as  $\varphi_I(f), \varphi_J(f)$ , respectively, and for every polynomial  $f$  of  $R^*$  there is a polynomial  $f'$  of  $R$  such that  $f'^* = f$ .*
- *Let  $\alpha_I, \beta_I \in \text{Con}(\text{pr}_I R)$ ,  $\alpha_J, \beta_J \in \text{Con}(\text{pr}_J R)$  be such that  $\alpha_I \prec \beta_I$ ,  $\alpha_J \prec \beta_J$  and  $\alpha_{I^*}^* = \varphi_I(\alpha_I)$ ,  $\beta_{I^*}^* = \varphi_I(\beta_I)$ ,  $\alpha_{J^*}^* = \varphi_J(\alpha_J)$ ,  $\beta_{J^*}^* = \varphi_J(\beta_J)$  be congruences of  $\text{pr}_{I^*} R^*, \text{pr}_{J^*} R^*$ . Then  $\alpha_{I^*}^* \prec \beta_{I^*}^*$ ,  $\alpha_{J^*}^* \prec \beta_{J^*}^*$ , and  $(\alpha_{I^*}^*, \beta_{I^*}^*)$  can be separated from  $(\alpha_{J^*}^*, \beta_{J^*}^*)$  if and only if  $(\alpha_I, \beta_I)$  can be separated from  $(\alpha_J, \beta_J)$ .*

*Proof.* (1) Note that for any polynomial  $f$  of  $R$  its action on the first and second projections of  $R$  is the same polynomial of  $\mathbb{A}$ . Therefore  $(\alpha, \beta)$  can be separated from  $(\gamma, \delta)$  in  $\text{Con}(\mathbb{A})$  if and only if there is  $f \in \text{Pol}_1(\mathbb{A})$ ,  $f(\beta) \not\subseteq \alpha$  while  $f(\delta) \subseteq \gamma$ . This condition can be expressed as follows: there is  $f \in \text{Pol}_1(R)$ ,  $f(\beta_1) \not\subseteq \alpha_1$  while  $f(\beta_2) \subseteq \alpha_2$ , which precisely means that  $(\alpha_1, \beta_1)$  can be separated from  $(\alpha_2, \beta_2)$  in  $R$ .

(2) The isomorphism part of the lemma is straightforward, while the separation part can be proved in a way similar to item (1).  $\square$

In what follows when proving results about separation we will always assume that we deal with a relation — a subdirect product — and that the prime intervals in question are from congruence lattices of projections of the subdirect product that do not overlap. If this is not the case, one can duplicate some of the factors and apply Lemma 34(2).

Let  $R$  be a subdirect product of  $\mathbb{A}_1, \dots, \mathbb{A}_n$ ,  $I \subseteq [n]$ , and let  $f$  be a polynomial of  $\text{pr}_I R$ , that is, there are a term operation  $g$  of  $R$  and  $\mathbf{a}_1, \dots, \mathbf{a}_k \in \text{pr}_I R$  such that  $f(x_1, \dots, x_\ell) = g(x_1, \dots, x_\ell, \mathbf{a}_1, \dots, \mathbf{a}_k)$ . The tuples  $\mathbf{a}_i$  can be extended to tuples  $\mathbf{a}'_i \in R$ . Then the polynomial of  $R$  given by  $f(x_1, \dots, x_\ell) = g(x_1, \dots, x_\ell, \mathbf{a}'_1, \dots, \mathbf{a}'_k)$  is said to be an *extension* of  $f$  to a polynomial of  $R$ .

**Lemma 35.** *Let  $\mathbb{A}_1, \dots, \mathbb{A}_n$  be arbitrary algebras. Let  $R$  be a subdirect product of  $\mathbb{A}_1, \dots, \mathbb{A}_n$ ,  $I, J \subseteq [n]$ , and  $\alpha_I \prec \beta_I$ ,  $\alpha_J \prec \beta_J$  for  $\alpha_I, \beta_I \in \text{Con}(\text{pr}_I R)$ ,  $\alpha_J, \beta_J \in \text{Con}(\text{pr}_J R)$ .*

- (1) *Let a unary polynomial  $f$  of  $R$  separate  $(\alpha_I, \beta_I)$  from  $(\alpha_J, \beta_J)$ . Then  $f$  can be chosen to be idempotent and such that  $f(\text{pr}_I R)$  is an  $(\alpha_I, \beta_I)$ -minimal set.*
- (2) *Let  $Q \subseteq R$  also be a subdirect product of  $\mathbb{A}_1, \dots, \mathbb{A}_n$  with  $\text{pr}_I Q = \text{pr}_I R$ ,  $\text{pr}_J Q = \text{pr}_J R$ . If  $(\alpha_I, \beta_I)$  can be separated from  $(\alpha_J, \beta_J)$  in  $Q$ ,  $(\alpha_I, \beta_I)$  can also be separated from  $(\alpha_J, \beta_J)$  in  $R$ .*

*Proof.* (1) Let  $g$  be a polynomial of  $R$  separating  $(\alpha_I, \beta_I)$  from  $(\alpha_J, \beta_J)$ . Since  $g(\beta_I) \not\subseteq \alpha_I$ , by Lemma 1(6) there is an  $(\alpha_I, \beta_I)$ -minimal set  $U$  such that  $g(\beta_I|_U) \not\subseteq \alpha_I$ . Let  $V = g(U)$ , by Lemma 1(2)  $V$  is a  $(\alpha_I, \beta_I)$ -minimal set. Let  $h$  be a unary polynomial such that  $h$  maps  $V$  onto  $U$  and  $h \circ g|_U$  is the identity mapping. Let also  $h'$  be an extension of  $h$  to a polynomial of  $R$ . Let  $p$  be an idempotent unary polynomial of  $\text{pr}_I R$  such that  $p(\text{pr}_I R) = U$  and let  $p'$  be an extension of  $p$  to a polynomial of  $R$ . If  $(h' \circ g)^m$  is an idempotent polynomial of  $R$  (the idempotent power of  $h' \circ g$ ), then let  $q = p' \circ (h' \circ g)^m$ . Now we can choose  $f$  as the idempotent power of  $q$ . By the choice of  $q$  we have  $q(\text{pr}_I R) = U$ , while on the projection on  $J$  the polynomial  $q$  collapses  $\beta_J$  to  $\alpha_J$ , so the idempotent power of  $q$  has all the required properties.

(2) Let a polynomial  $g$  of  $Q$  separate  $(\alpha_I, \beta_I)$  from  $(\alpha_J, \beta_J)$  in  $Q$ . Then, as  $Q \subseteq R$ ,  $g$  gives rise to a polynomial of  $R$  and, as  $\text{pr}_I Q = \text{pr}_I R$ ,  $\text{pr}_J Q = \text{pr}_J R$ , its action on  $\text{pr}_I R$  and  $\text{pr}_J R$  is the same. The result follows.  $\square$

### 3.3 Chaining

For a subdirect product  $R \subseteq \mathbb{A}_1 \times \cdots \times \mathbb{A}_n$  the relation ‘cannot be separated’ on prime intervals of the  $\mathbb{A}_i$ ’s is clearly reflexive and transitive. If the algebras  $\mathbb{A}_i$  are Mal’tsev, it is also symmetric (for partial results see [1, 5]). Moreover, it can be shown that it remains ‘almost’ symmetric when the  $\mathbb{A}_i$ ’s contain no majority edges. In the general case however the situation is more complicated. Next we introduce conditions that make the ‘cannot be separated’ relation to some extent symmetric, at least in what concerns our needs, as it will be demonstrated in Theorem 38. From now on we again assume that algebras we work with belong to  $\mathcal{V}$ .

For an algebra  $\mathbb{A} \in \mathcal{V}$ ,  $\alpha \in \text{Con}(\mathbb{A})$ , a set  $\mathcal{U}$  of unary polynomials of  $\mathbb{A}$ , and  $B \subseteq \mathbb{A}^2$ , we denote by  $\text{Cg}_{\mathbb{A}, \alpha, \mathcal{U}}(B)$  the transitive-symmetric closure of the set  $T(B, \mathcal{U}) = \{(f(a), f(b)) \mid (a, b) \in B, f \in \mathcal{U}\} \cup \alpha$ . We will write  $\{c, d\} \in T(B, \mathcal{U})$  if  $(c, d) \in T(B, \mathcal{U})$  or  $(d, c) \in T(B, \mathcal{U})$ . Let  $\alpha, \beta \in \text{Con}(\mathbb{A})$ ,  $\alpha \leq \beta$ , and  $D$  a subuniverse of  $\mathbb{A}$ . We say that  $\alpha$  and  $\beta$  are  $\mathcal{U}$ -chained with respect to  $D$  if for any  $a, b \in D$ ,  $a \stackrel{\beta}{=} b$ , such that  $\beta = \text{Cg}_{\mathbb{A}}(\{(a, b)\}) \vee \alpha$  and any  $\beta$ -block  $B$  such that  $B' = B \cap \text{umax}(D) \neq \emptyset$ , we have  $(\text{umax}(B'))^2 \subseteq \text{Cg}_{\mathbb{A}, \alpha, \mathcal{U}}(\{(a, b)\})$ . Note that if  $\beta = \text{Cg}_{\mathbb{A}}(\{(a, b)\}) \vee \alpha$  for no  $a, b \in D$ , then  $\alpha, \beta$  are trivially  $\mathcal{U}$ -chained.

Let  $R$  be a subdirect product of  $\mathbb{A}_1, \dots, \mathbb{A}_n \in \mathcal{V}$ ,  $\beta_i \in \text{Con}(\mathbb{A}_i)$ , let  $B_i$  be a  $\beta_i$ -block for  $i \in [n]$ , and let  $R' = R \cap \overline{B}$ ,  $B'_i = \text{pr}_i R'$ . A unary polynomial  $f$  of  $R$  is said to be  $\overline{B}$ -preserving if  $f(R') \subseteq R'$ . We call an  $n$ -ary relation  $R$  chained with respect to  $\overline{\beta}, \overline{B}$  if

(Q1) for any  $I \subseteq [n]$  and  $\alpha, \beta \in \text{Con}(\text{pr}_I R)$  such that  $\alpha \leq \beta \leq \overline{\beta}_I$ ,  $\alpha, \beta$  are  $\mathcal{U}_{\overline{B}}$ -chained with respect to  $\text{pr}_I R'$ , where  $\mathcal{U}_{\overline{B}}$  is the set of all  $\overline{B}$ -preserving polynomials of  $R$ ;

(Q2) for any  $I, J \subseteq [n]$  and  $\alpha, \beta \in \text{Con}(\text{pr}_I R)$ ,  $\gamma, \delta \in \text{Con}(\text{pr}_J R)$  (note that it may happen that  $I \cap J \neq \emptyset$ ), such that  $\alpha \prec \beta \leq \overline{\beta}_I$ ,  $\gamma \prec \delta \leq \overline{\beta}_J$ , and  $(\alpha, \beta)$  can be separated from  $(\gamma, \delta)$ , the congruences  $\alpha$  and  $\beta$  are  $\mathcal{U}(\gamma, \delta, \overline{B})$ -chained with respect to  $\text{pr}_I R'$ , where  $\mathcal{U}(\gamma, \delta, \overline{B})$  is the set of all  $\overline{B}$ -preserving polynomials  $g$  of  $R$  such that  $g(\delta) \subseteq \gamma$ .

Next we make an observation about a link between the property of chaining, projections, and factor algebras.

**Lemma 36.** *Let  $\mathbb{A}_1, \dots, \mathbb{A}_n, R, \overline{\beta}, \overline{B}$ , and  $R'$  be as in the definition of chaining.*

- (1) *Let  $K \subseteq [n]$  and consider  $R$  as a subdirect product  $R^*$  of  $\text{pr}_K R$  and  $\mathbb{A}_i$ ,  $i \in [n] - K$ . Let  $\beta_i^* = \beta_i$  and  $B_i^* = B_i$  for  $i \in [n] - K$  and  $\beta_K^* = \overline{\beta}_K$ ,  $B_K^* = \overline{B}_K$ . Then relation  $R^*$  is chained with respect  $\overline{\beta}^*, \overline{B}^*$  if and only if  $R$  is chained with respect to  $\overline{\beta}, \overline{B}$ .*



(2) Let  $R$  be chained with respect to  $\overline{\beta}, \overline{B}$  and let  $\alpha_i \in \text{Con}(\mathbb{A}_i)$  and  $\alpha_i \leq \beta_i$ ,  $i \in [n]$ . Then  $R/\overline{\alpha}$  is chained with respect to  $\overline{\beta}/\overline{\alpha}, \overline{B}/\overline{\alpha}$ .

*Proof.* (1) Suppose  $K = [k]$ . Let  $R$  be chained with respect to  $\overline{\beta}, \overline{B}$ . Let  $I, J \subseteq \{K, k+1, \dots, n\}$  and  $\alpha, \beta \in \text{Con}(\text{pr}_I R^*)$ ,  $\gamma, \delta \in \text{Con}(\text{pr}_J R^*)$  as in the definition of chaining. Also, if  $K \in I$  [if  $K \in J$ ] set  $I' = (I - \{K\}) \cup K$  [respectively,  $J' = (J - \{K\}) \cup K$ ] and  $I' = I$  [respectively,  $J' = J$ ] otherwise. Then the congruences  $\alpha, \beta, \gamma, \delta$  viewed as congruences of  $\text{pr}_I R^*, \text{pr}_J R^*$  satisfy the same conditions when viewed as congruences of  $\text{pr}_{I'} R, \text{pr}_{J'} R$ .

Now, let  $R^*$  be chained,  $I, J \subseteq [n]$ , and let  $\alpha, \beta, \gamma, \delta$  be as in the definition of chaining. We set  $I^* = I' = I$  [ $J^* = J' = J$ ] if  $I \cap K = \emptyset$  [if  $J \cap K = \emptyset$ ], and  $I' = I \cup K$ ,  $I^* = (I - K) \cup \{K\}$  [respectively,  $J' = J \cup K$ ,  $J^* = (J - K) \cup \{K\}$ ] otherwise. There are one-to-one correspondences between tuples from  $\text{pr}_{I'} R, \text{pr}_{J'} R$  and tuples from  $\text{pr}_{I^*} R^*, \text{pr}_{J^*} R^*$ , denote then  $\varphi_I, \varphi_J$ . Next, we need to define congruences of  $\text{pr}_{I^*} R^*, \text{pr}_{J^*} R^*$  in an appropriate way. Note that if  $\beta = \text{Cg}(\{(\mathbf{a}, \mathbf{b})\}) \vee \alpha$  for no  $\mathbf{a}, \mathbf{b} \in \text{pr}_I R'$ , there is nothing to prove. So, suppose there are  $\mathbf{a}, \mathbf{b} \in \text{pr}_I R'$  such that  $\beta = \text{Cg}(\{(\mathbf{a}, \mathbf{b})\}) \vee \alpha$ . If  $I \cap K = \emptyset$  or  $J \cap K = \emptyset$ , set  $\alpha^* = \alpha, \beta^* = \beta$  and  $\gamma^* = \gamma, \delta^* = \delta$ , respectively. If  $I \cap K \neq \emptyset$  then let  $\alpha' = \alpha \times \prod_{i \in K-I} \beta_i$ . As is easily seen, it is the maximal congruence of  $\text{pr}_{I'} R$  such that  $\alpha' \leq \overline{\beta}_{I'}$  and  $\{(\text{pr}_I \mathbf{c}, \text{pr}_I \mathbf{d}) \mid (\mathbf{c}, \mathbf{d}) \in \alpha'_{|\text{pr}_{I'} R'}\} = \alpha'_{|\text{pr}_I R'}$ . Then we set  $\alpha^* = \varphi_I(\alpha')$ . Also let  $\mathbf{a}', \mathbf{b}'$  be arbitrary tuples from  $\text{pr}_{I'} R'$  such that  $\text{pr}_I \mathbf{a}' = \mathbf{a}$  and  $\text{pr}_I \mathbf{b}' = \mathbf{b}$  and let  $\beta = \text{Cg}_{\text{pr}_{I'} R'}(\{(\mathbf{a}', \mathbf{b}')\}) \vee \alpha$ . We set  $\beta^* = \text{Cg}_{\text{pr}_{I^*} R^*}(\{(\varphi_I(\mathbf{a}'), \varphi_I(\mathbf{b}'))\}) \vee \alpha^*$ . Observe that if  $\alpha \prec \beta$  then  $\alpha^* \prec \beta^*$ . Indeed, for any  $\beta'$  with  $\alpha^* \leq \beta' \leq \beta^*$ , if  $\alpha^* \neq \beta'$ , there are  $\mathbf{c}, \mathbf{d} \in \text{pr}_{I^*} R^*$ ,  $\mathbf{c} \equiv^{\beta^*} \mathbf{d}$ , such that  $(\varphi_I^{-1}(\mathbf{c}), \varphi_I^{-1}(\mathbf{d})) \notin \alpha'$ . Therefore there are  $\mathbf{c}', \mathbf{d}' \in \text{pr}_{I'} R'$  such that  $(\text{pr}_I \mathbf{c}', \text{pr}_I \mathbf{d}') \in \beta - \alpha$ . Thus,  $(\mathbf{a}, \mathbf{b}) \in \text{Cg}_{\text{pr}_I R}(\{(\text{pr}_I \mathbf{c}', \text{pr}_I \mathbf{d}')\}) \vee \alpha$  implying  $(\mathbf{a}', \mathbf{b}') \in \text{Cg}_{\text{pr}_{I'} R'}(\{(\text{pr}_{I'} \mathbf{c}', \text{pr}_{I'} \mathbf{d}')\}) \vee \alpha'$ . The congruences  $\gamma^*, \delta^*$  are constructed in the same way. Since conditions (Q1), (Q2) hold for  $R^*, \alpha^*, \beta^*, \gamma^*, \delta^*$ , it is straightforward to see that they also hold for  $R, \alpha, \beta, \gamma, \delta$ .

(2) is straightforward from the definitions.  $\square$

In many cases Lemma 36 allows us to replace large sets  $I, J$  in the definition of chaining with singletons.

We conclude this subsection with an auxiliary statement. For an algebra  $\mathbb{A} \in \mathcal{V}$ ,  $\alpha \in \text{Con}(\mathbb{A})$ ,  $a', b' \in \mathbb{A}$ , and a set  $\mathcal{U}$  of unary polynomials of  $\mathbb{A}$  we use  $T_\alpha(a, b, \mathcal{U})$ , where  $a = a'/\alpha, b = b'/\alpha$ , to denote  $T(\{(a', b')\}, \mathcal{U})/\alpha$ .

**Lemma 37.** Let  $\mathbb{A}_1, \dots, \mathbb{A}_n, R, \overline{\beta}, \overline{B}$ , and  $R'$  be as in the definition of chaining, and  $R$  is chained with respect to  $\overline{\beta}, \overline{B}$ . Let  $I, J \subseteq [n]$ ,  $\alpha \leq \overline{\beta}_I$ , and  $\gamma \prec \delta \leq \overline{\beta}_J$ . Let also  $\mathcal{U} \in \{\mathcal{U}_{\overline{B}}, \mathcal{U}(\gamma, \delta, \overline{B})\}$ .

- (1) Any constant polynomial from  $R \cap \overline{B}$  belongs to  $\mathcal{U}$ .
- (2) If  $f$  is a  $k$ -ary term operation of  $R$  and  $g_1, \dots, g_k \in \mathcal{U}$ , then  $f(g_1(x), \dots, g_k(x)) \in \mathcal{U}$ .
- (3) For any  $\mathbf{a}, \mathbf{b} \in \text{pr}_I R' / \alpha$ , a congruence  $\beta \in \text{Con}(\text{pr}_I R)$ ,  $\beta \leq \overline{\beta}_I$ , such that  $\text{Cg}_{\text{pr}_I R / \alpha}(\{\mathbf{a}, \mathbf{b}\}) = \beta / \alpha$  and any  $\beta / \alpha$ -block  $E$  with  $E \cap \text{umax}(\text{pr}_I R' / \alpha) \neq \emptyset$ , it holds that  $(\mathbf{c}, \mathbf{d}) \in T_\alpha(\mathbf{a}, \mathbf{b}, \mathcal{U})$  for any  $\mathbf{c}, \mathbf{d}$  from the same as-component of  $E \cap \text{pr}_I R' / \alpha$ .
- (4) For any  $\mathbf{a}, \mathbf{b} \in \text{pr}_I R' / \alpha$ , a congruence  $\beta \in \text{Con}(\text{pr}_I R)$ ,  $\beta \leq \overline{\beta}_I$ , such that  $\text{Cg}_{\text{pr}_I R / \alpha}(\{\mathbf{a}, \mathbf{b}\}) = \beta / \alpha$ , any  $\beta / \alpha$ -block  $E$  with  $E \cap \text{umax}(\text{pr}_I R' / \alpha) \neq \emptyset$ , and any  $\mathbf{c}, \mathbf{d} \in \text{umax}(E')$ , where  $E' = E \cap \text{pr}_I R' / \alpha$ , there is a sequence  $\mathbf{c} = \mathbf{c}_1, \dots, \mathbf{c}_k = \mathbf{d}$  in  $E'$  such that  $\{\mathbf{c}_i, \mathbf{c}_{i+1}\} \in T_\alpha(\mathbf{a}, \mathbf{b}, \mathcal{U})$  for  $i \in [k-1]$ . In particular, if  $|\text{umax}(E')| > 1$ , for any  $\mathbf{c} \in \text{umax}(E')$  there exists  $\mathbf{c}' \in E'$ ,  $\mathbf{c} \neq \mathbf{c}'$ , such that  $\{\mathbf{c}, \mathbf{c}'\} \in T_\alpha(\mathbf{a}, \mathbf{b}, \mathcal{U})$ .
- (5) Let  $\mathbf{a}, \mathbf{b} \in \text{pr}_I R' / \alpha$ . If  $(\mathbf{c}, \mathbf{d}) \in T_\alpha(\mathbf{a}, \mathbf{b}, \mathcal{U})$  then  $T_\alpha(\mathbf{c}, \mathbf{d}, \mathcal{U}) \subseteq T_\alpha(\mathbf{a}, \mathbf{b}, \mathcal{U})$ .
- (6) Let  $\beta \in \text{Con}(\text{pr}_I R)$ ,  $\alpha \prec \beta \leq \overline{\beta}_I$ , and  $(\alpha, \beta)$  can be separated from  $(\gamma, \delta)$ . Let  $E$  be a  $\beta / \alpha$ -block such that  $E \cap \text{umax}(\text{pr}_I R' / \alpha) \neq \emptyset$ , and let  $E' = E \cap \text{pr}_I R' / \alpha$ . For any  $\mathbf{a}, \mathbf{b} \in E'$ ,  $\mathbf{a} \neq \mathbf{b}$ , there are  $\mathbf{a}', \mathbf{b}' \in E'$ ,  $\mathbf{a}' \neq \mathbf{b}'$ , such that  $(\mathbf{a}', \mathbf{b}') \in T_\alpha(\mathbf{a}, \mathbf{b}, \mathcal{U})$  and  $(\mathbf{a}', \mathbf{b}') \in T_\alpha(\mathbf{a}', \mathbf{b}', \mathcal{U})$ .<sup>2</sup> If  $|\text{umax}(E')| > 1$  and  $\mathbf{a} \in \text{umax}(E')$ ,  $\mathbf{a}'$  can be chosen to be  $\mathbf{a}$ . Moreover, if  $E'$  contains a nontrivial as-component, then there is a set  $T \subseteq \beta / \alpha$  such that  $T \subseteq T_\alpha(\mathbf{c}, \mathbf{d}, \mathcal{U})$  for any  $\mathbf{c}, \mathbf{d} \in \text{pr}_I R' / \alpha$ ,  $\mathbf{c} \neq \mathbf{d}$ ,  $\mathbf{c} \stackrel{\beta / \alpha}{\equiv} \mathbf{d}$ , and  $T = T_\alpha(\mathbf{c}', \mathbf{d}', \mathcal{U})$  for some  $\mathbf{c}', \mathbf{d}'$ .

*Proof.* Items (1),(2) are straightforward, and (4) follows from the definitions and the assumption that  $R$  is chained with respect to  $\overline{\beta}, \overline{B}$ . Let  $T(\mathbf{a}, \mathbf{b})$  denote  $T_\alpha(\mathbf{a}, \mathbf{b}, \mathcal{U})$ .

(3) By item (2)  $Q = T(\mathbf{a}, \mathbf{b}) \cap (\text{pr}_I R' / \alpha)^2$  is a subalgebra of  $(\text{pr}_I R / \alpha)^2$ . Since  $R$  is chained,  $\text{umax}(E \cap \text{pr}_I R' / \alpha)$  is a subset of a block of the link congruence of  $Q$ . Therefore, as by Corollary 16  $\text{amax}(E \cap \text{pr}_I R' / \alpha) \subseteq \text{umax}(E \cap \text{pr}_I R' / \alpha)$ , by Proposition 26 for any as-component  $D$  of  $E \cap \text{pr}_I R' / \alpha$  we have  $D^2 \subseteq Q$ .

(5) Let  $\{\mathbf{a}', \mathbf{b}'\} \in T(\mathbf{c}, \mathbf{d})$ . Then there are polynomials  $f, g \in \mathcal{U}$  with  $\{\mathbf{c}, \mathbf{d}\} = f(\{\mathbf{a}, \mathbf{b}\})$  and  $\{\mathbf{a}', \mathbf{b}'\} = g(\{\mathbf{c}, \mathbf{d}\})$ . Then  $g \circ f \in \mathcal{U}$  by item (2) of the lemma and the definition, and  $g \circ f(\{\mathbf{a}, \mathbf{b}\}) = \{\mathbf{a}', \mathbf{b}'\}$ .

(6) Consider  $T(\mathbf{a}, \mathbf{b})$  and let  $(\mathbf{a}', \mathbf{b}') \in E'^2 \cap T(\mathbf{a}, \mathbf{b})$  be such that  $T(\mathbf{a}', \mathbf{b}')$  is minimal possible with respect to inclusion. Since  $R$  is chained, it is easy to see

---

<sup>2</sup>If  $\mathcal{U} = \mathcal{U}_{\overline{B}}$  this claim is trivial for any  $\mathbf{a}', \mathbf{b}'$ , because the identity operation witnesses that  $(\mathbf{a}', \mathbf{b}') \in T_\alpha(\mathbf{a}', \mathbf{b}', \mathcal{U}_{\overline{B}})$ . However if  $\mathcal{U} = \mathcal{U}(\gamma, \delta, \overline{B})$ , it does not have to be the case.

that  $(\mathbf{a}', \mathbf{b}') \in T(\mathbf{a}', \mathbf{b}')$ . If  $\mathbf{a} \in \text{umax}(E')$  by item (4)  $(\mathbf{a}, \mathbf{b}'') \in T(\mathbf{a}', \mathbf{b}')$  for some  $\mathbf{b}'' \in E'$ . By the minimality of  $T(\mathbf{a}', \mathbf{b}')$  and item (5) we have  $T(\mathbf{a}', \mathbf{b}') = T(\mathbf{a}, \mathbf{b}'')$ . For the second part of the claim take  $\mathbf{a}, \mathbf{b} \in C$  where  $C$  is a nontrivial as-component in  $E'$ . By item (3)  $\{\mathbf{a}, \mathbf{b}\} \in T(\mathbf{c}, \mathbf{d})$  for any appropriate  $\mathbf{c}, \mathbf{d}$ . Therefore by item (5)  $T = T(\mathbf{a}, \mathbf{b}) \subseteq T(\mathbf{c}, \mathbf{d})$ .  $\square$

### 3.4 Symmetricity of separation

The following theorem establishes the weak symmetricity of separability relation mentioned before.

**Theorem 38.** *Let  $R$  be a subdirect product of  $\mathbb{A}_1, \dots, \mathbb{A}_n \in \mathcal{V}$ , for each  $i \in [n]$ ,  $\beta_i \in \text{Con}(\mathbb{A}_i)$ ,  $B_i$  a  $\beta_i$ -block such that  $R$  is chained with respect to  $\overline{\beta}, \overline{B}$ ;  $R' = R \cap \overline{B}$ ,  $B'_i = \text{pr}_i R'$ . Let also  $i, j \in [n]$ ,  $\alpha \prec \beta \leq \beta_i$ ,  $\gamma \prec \delta = \beta_j$ , where  $\alpha, \beta \in \text{Con}(\mathbb{A}_i)$ ,  $\gamma, \delta \in \text{Con}(\mathbb{A}_j)$ . If  $B'_j/\gamma$  has a nontrivial as-component  $C_j$  and  $(\alpha, \beta)$  can be separated from  $(\gamma, \delta)$ , then there is a  $\overline{B}$ -preserving polynomial  $g$  of  $R$  such that  $g(\beta|_{B'_i}) \subseteq \alpha$  and  $g(\delta|_{B'_j}) \not\subseteq \gamma$ . Moreover, for any  $c, d \in C_j$ ,  $c \neq d$ , the polynomial  $g$  can be chosen such that  $g(c) = c, g(d) = d$ .*

Without loss of generality assume  $i = 1, j = 2$ . Before proving Theorem 38 we describe two auxiliary constructions and prove several intermediate results. Fix a relation  $R$  as specified in the statement of Theorem 38. In the rest of Section 3.4 we use the notation from Theorem 38. As is easily seen, by Lemma 36(2) we can assume that  $\alpha, \gamma$  are equality relations. So, let  $\alpha = \underline{0}_1, \gamma = \underline{0}_2$ .

We use the following two constructions. By  $Q^*(a, b, c, d, \mathbf{a}) \subseteq \mathbb{A}_1^2 \times \mathbb{A}_2^2 \times R$  we denote the relation generated by  $\{(a, b, c, d, \mathbf{a})\} \cup \{(x, x, y, y, \mathbf{z}) \mid \mathbf{z} \in R, \mathbf{z}[1] = x, \mathbf{z}[2] = y\}$ , where  $a, b \in B'_1, c, d \in B'_2$  and  $\mathbf{a} \in R'$ . Let  $Q(a, b, c, d, \mathbf{a}) = \text{pr}_{1234} Q^*(a, b, c, d, \mathbf{a})$  and  $Q'(a, b, c, d, \mathbf{a}) = \text{pr}_{1234}(Q^*(a, b, c, d, \mathbf{a}) \cap (B'_1 \times B'_1 \times B'_2 \times B'_2 \times \overline{B}))$ .

Also, by  $P^*(c_1, c_2, c_3, \mathbf{a}) \subseteq \mathbb{A}_2^3 \times R$  we denote the relation generated by  $\{(c_1, c_2, c_3, \mathbf{a})\} \cup \{(x, x, x, \mathbf{z}) \mid \mathbf{z} \in R, \mathbf{z}[2] = x\}$ , where  $c_1, c_2, c_3 \in B'_2$  and  $\mathbf{a} \in R'$ . Let  $P(c_1, c_2, c_3, \mathbf{a}) = \text{pr}_{123} P^*(c_1, c_2, c_3, \mathbf{a})$  and  $P'(c_1, c_2, c_3, \mathbf{a}) = \text{pr}_{123}(P^*(c_1, c_2, c_3, \mathbf{a}) \cap (B'_2 \times B'_2 \times B'_2 \times \overline{B}))$ .

**Lemma 39.** *For any  $a, b \in B'_1$   $a \stackrel{\beta}{=} b$ , any  $c, d, c_1, c_2, c_3 \in B'_2$ , and any  $\mathbf{a} \in R'$ ,*

- (1)  $Q = Q(a, b, c, d, \mathbf{a})$  is the set of quadruples  $(f(a), f(b), f(c), f(d))$  for unary polynomials  $f$  of  $R$ , and  $Q' = Q'(a, b, c, d, \mathbf{a})$  is the set of quadruples  $(f(a), f(b), f(c), f(d))$  for  $\overline{B}$ -preserving unary polynomials  $f$  of  $R$ .<sup>3</sup>

<sup>3</sup>In fact, that we are looking for  $\overline{B}$ -preserving polynomials is the reason we consider the big relation  $Q^*$  rather than starting directly with the 4-ary  $Q(a, b, c, d, \mathbf{a})$ .

- (2)  $\text{pr}_1 Q' = \text{pr}_2 Q' = B'_1$ ,  $\text{pr}_3 Q' = \text{pr}_4 Q' = B'_2$ , and  $\text{pr}_{12} Q = Q_{ab}^{\mathbb{A}_1}$ ,  $\text{pr}_{34} Q = Q_{cd}^{\mathbb{A}_2}$ .
- (3)  $P = P(c_1, c_2, c_3, \mathbf{a})$  is the set of triples  $(f(c_1), f(c_2), f(c_3))$  for unary polynomials  $f$  of  $R$ , and  $P' = P'(c_1, c_2, c_3, \mathbf{a})$  is the set of triples  $(f(c_1), f(c_2), f(c_3))$  for  $\overline{B}$ -preserving unary polynomials  $f$  of  $R$ .
- (4)  $\text{pr}_1 P' = \text{pr}_2 P' = \text{pr}_3 P' = B'_2$ , and  $\text{pr}_{12} P = Q_{c_1 c_2}^{\mathbb{A}_2}$ ,  $\text{pr}_{23} P = Q_{c_2 c_3}^{\mathbb{A}_2}$ ,  $\text{pr}_{13} P = Q_{c_1 c_3}^{\mathbb{A}_2}$ .
- (5) Let  $\text{lk}_{12}, \text{lk}_{34}$  denote the link congruences of  $Q'$  viewed as a subdirect product of  $Q'_{12} = \text{pr}_{12} Q'$  and  $Q'_{34} = \text{pr}_{34} Q'$ .<sup>4</sup> There are  $a', b' \in B'_1$  and  $c', d' \in C_2$ ,  $c' \neq d'$ , such that  $(a', b', c', d') \in Q'$  and  $(a', b') \in \text{amax}(Q'_{12} \cap E^2)$  for a  $\beta$ -block  $E$  such that  $E \cap \text{umax}(B'_1) \neq \emptyset$ . Moreover, if  $(c', d')$  can be chosen from  $\text{amax}(D_{34})$ , where  $D_{34}$  is the  $\text{lk}_{34}$ -block containing  $(c', d')$ , then  $(a', b')$  can also be chosen from  $\text{amax}(D_{12})$ , where  $D_{12}$  is the  $\text{lk}_{12}$ -block containing  $(a', b')$ , and  $\{(a', b')\} \times C_{34} \subseteq Q'$ , where  $C_{34}$  is the as-component of  $D_{34}$  containing  $(c', d')$ .

**Remark 40.** Observe that by Lemma 39(1,3) the choice of  $\mathbf{a} \in R'$  is immaterial for  $Q(a, b, c, d, \mathbf{a})$ ,  $Q'(a, b, c, d, \mathbf{a})$ ,  $P(c_1, c_2, c_3, \mathbf{a})$ ,  $P'(c_1, c_2, c_3, \mathbf{a})$ . We therefore will often omit  $\mathbf{a}$  from this notation in the future.

*Proof.* (1) To prove the second part of (1) observe that  $(a', b', c', d') \in Q'$  if and only if there are a term operation  $g(x, y_1, \dots, y_k)$  and  $\mathbf{b}_1 = (a_1, a_1, c_1, c_1, \mathbf{a}_1), \dots, \mathbf{b}_k = (a_k, a_k, c_k, c_k, \mathbf{a}_k)$  with  $\mathbf{a}_i \in R$ ,  $\mathbf{a}_i[1] = a_i, \mathbf{a}_i[2] = c_i$  for  $i \in [k]$ , such that  $(a', b', c', d', \mathbf{b}') = g((a, b, c, d, \mathbf{a}), \mathbf{b}_1, \dots, \mathbf{b}_k)$  and  $\mathbf{b}' \in \overline{B}$ . Consider the unary polynomial  $f(x) = g(x, \mathbf{a}_1, \dots, \mathbf{a}_k)$ ; clearly  $\mathbf{b}' = f(\mathbf{a})$ . As  $R \cap \overline{B}$  is a congruence block, the latter implies that  $f$  is  $\overline{B}$ -preserving. Finally, since  $a_i = \mathbf{a}_i[1], c_i = \mathbf{a}_i[2]$ , we have  $a' = f(a), b' = f(b)$  in  $\text{pr}_1 R$  and  $c' = f(c), d' = f(d)$  in  $\text{pr}_2 R$ . The first part of (1) is proved in a similar way, except we do not need to care about polynomials being  $\overline{B}$ -preserving.

(2) follows from the definitions.

(3) and (4) are proved in a way similar to (1) and (2).

(5) Consider the relation  $S = Q' \cap (B'_1 \times B'_1 \times \text{Sg}(C_2) \times \text{Sg}(C_2))$ . Since  $R$  is chained, by item (2) and Lemma 37(3)  $C_2^2 \subseteq \text{pr}_{34} S$  and  $\text{pr}_{34} S$  is generated by  $C_2^2$ . Let

$$S/\beta = \{(a'/\beta, b'/\beta, c', d') \mid (a', b', c', d') \in S\} \subseteq B'_1/\beta \times B'_1/\beta \times \text{Sg}(C_2) \times \text{Sg}(C_2).$$

<sup>4</sup>Note that these congruences may be different from the link congruences of  $Q$  restricted to  $\text{pr}_{12} Q \cap (B'_1 \times B'_1)$ ,  $\text{pr}_{34} Q \cap (B'_2 \times B'_2)$ , respectively.

Let  $\text{lk}_{12}^\beta, \text{lk}_{34}^\beta$  be the link congruences of  $S/\beta$  treated as a subdirect product of  $\text{pr}_{12}S/\beta, \text{pr}_{34}S/\beta$ . By Lemma 28 there is a  $\text{lk}_{34}^\beta$ -block  $D_{34}^\beta$  such that for some  $c', d' \in C_2, c' \neq d'$ , the pair  $(c', d')$  is as-maximal in  $D_{34}^\beta$ . Let  $C_{34}^\beta$  be the as-component of  $D_{34}^\beta$  containing  $(c', d')$ . Note that, as  $a \stackrel{\beta}{\equiv} b$ ,  $\text{pr}_{12}S/\beta$  is the equality relation. Let  $E$  be a  $\beta$ -block such that  $(E, E)$  is as-maximal in the  $\text{lk}_{12}^\beta$ -block  $D_{12}^\beta$  corresponding to  $D_{34}^\beta$  (i.e.  $S/\beta \cap (D_{12}^\beta \times D_{34}^\beta) \neq \emptyset$ ) and such that  $(E, E, c', d') \in S/\beta$ . Such a  $\beta$ -block exists by the Maximality Lemma 18(5). Note that  $E \cap \text{umax}(B'_1) \neq \emptyset$ . Indeed, as  $D_{34}^\beta$  contains as-maximal elements, so does  $D_{12}^\beta$ . By Lemma 17 every element u-maximal in  $D_{12}^\beta$  is u-maximal in  $\text{pr}_{12}S/\beta$ . This means that  $E$  is u-maximal in  $B'_1/\beta$ , and again by Lemma 17  $E$  contains elements u-maximal in  $B'_1$ . We need several observations concerning  $S$  and the block  $E$ .

(i) By Proposition 26  $\{(E, E)\} \times C_{34}^\beta \subseteq S/\beta$ . This means that for any  $(c'', d'') \in C_{34}^\beta$  there is  $(a', b') \in S_{12}^E$ , where  $S_{12}^E = \text{pr}_{12}S \cap E^2$ , such that  $(a', b', c'', d'') \in S$ . Also, by construction for any  $(a', b') \in S_{12}^E$  there is  $(c'', d'') \in D_{34}^\beta$  such that  $(a', b', c'', d'') \in S$ .

(ii) Let  $D'_{12} = \{(a', b') \in \text{pr}_{12}S \mid (a'/\beta, b'/\beta) \in D_{12}^\beta\}$ . We claim that for any  $(a', b', c'', d'') \in S$  with  $(a', b') \in D'_{12}$  and  $(c'', d'') \in C_{34}^\beta$ , and any  $(a'', b'') \in Q'_{12}$  with  $(a', b') \sqsubseteq_{Q'_{12}}^{as} (a'', b'')$ , there is  $(c^*, d^*) \in C_2^2$  such that  $(a'', b'', c^*, d^*) \in S$ . Moreover, if  $(a'', b'') \in D'_{12}$  and  $(a', b') \sqsubseteq_{D'_{12}}^{as} (a'', b'')$  then  $(c^*, d^*)$  can be chosen from  $C_{34}^\beta$ . Indeed, the first claim follows from the Maximality Lemma 18(3) applied to  $Q'$ : there is  $(c^*, d^*) \in Q'_{34}$  such that  $(a'', b'', c^*, d^*) \in Q'$  and  $(c'', d'') \sqsubseteq_{Q'_{34}}^{as} (c^*, d^*)$ . Then  $(c^*, d^*) \in C_2^2$ , and so  $(a'', b'', c^*, d^*) \in S$ . Assuming  $(a', b') \sqsubseteq_{D'_{12}}^{as} (a'', b'')$ , again by the Maximality Lemma 18(3) the as-path from  $(c'', d'')$  to  $(c^*, d^*)$  lies in  $D_{34}^\beta$ . Therefore  $(c^*, d^*) \in C_{34}^\beta$ .

(iii) In particular, (ii) implies that if  $(a', b') \in \text{amax}(S_{12}^E)$ ,  $(a', b', c'', d'') \in S$  for some  $(c'', d'') \in C_{34}^\beta$ , and  $(a', b') \sqsubseteq_{Q'_{12} \cap E^2}^{as} (a'', b'')$ , then  $(a'', b'') \in \text{pr}_{12}S$ . This also implies that  $(a', b') \in \text{amax}(Q'_{12} \cap E^2)$ .

(iv) The last observation we need is that by Lemma 21(2) and the choice of  $E$ ,  $S_{12}^E$  contains a pair  $(a', b') \in \text{amax}(D'_{12})$ . Moreover, for any  $(a', b') \in S_{12}^E$  there is  $(a'', b'') \in \text{amax}(S_{12}^E) \cap \text{amax}(D'_{12})$  and such that  $(a', b') \sqsubseteq_{D'_{12}}^{as} (a'', b'')$ .

Hence, starting from any  $(a', b', c', d') \in S$ , where  $(a', b') \in S_{12}^E$  and  $c', d'$  are as identified in the beginning of the proof, we first find  $(a'', b'', c^*, d^*) \in S$  such that  $(a'', b'') \in \text{amax}(S_{12}^E)$  and  $(c^*, d^*) \in C_{34}^\beta$ . Since  $\{(E, E)\} \times C_{34}^\beta \subseteq S/\beta$  there is an as-path in  $S \cap (E^2 \times C_{34}^\beta)$  from  $(a'', b'', c^*, d^*)$  to  $(a^\dagger, b^\dagger, c', d')$  for some  $(a^\dagger, b^\dagger) \in \text{amax}(S_{12}^E)$ . Note that by (iii)  $(a^\dagger, b^\dagger)$  can also be assumed to be in

$\text{amax}(Q'_{12} \cap E^2)$ . The tuple  $(a^\dagger, b^\dagger, c', d')$  is as required.

For the second claim of item (5) suppose that  $(c', d') \in \text{amax}(D_{34})$ , where  $D_{34}$  is the  $\text{lk}_{34}$ -block containing  $(c', d')$ . Recall that  $C_{34}$  denotes the as-component of  $D_{34}$  containing  $(c', d')$ . Let  $D_{12}$  be the  $\text{lk}_{12}$ -block containing  $(a^\dagger, b^\dagger)$ . Then similar to (ii), (iii) there is  $(a^*, b^*) \in \text{amax}(D_{12}) \cap \text{amax}(Q'_{12} \cap E'^2)$  for some  $\beta$ -block  $E'$  such that  $(a^\dagger, b^\dagger) \sqsubseteq_{D_{12}}^{as} (a^*, b^*)$ . By Proposition 26  $\{(a^*, b^*)\} \times C_{34} \subseteq S$  and hence the pairs  $(a^*, b^*)$  and  $(c', d')$  satisfy the desired conditions. Part (5) is proved.  $\square$

**Lemma 41.** *Let  $c, d \in C_2$ ,  $c \neq d$ , and let  $a, b \in B'_1$ ,  $a \stackrel{\beta}{\equiv} b$  be such that  $(a, b) \in T_0 = T(a, b, \mathcal{U}(\underline{0}_2, \delta, \overline{B}))$  and such that  $(a, c) \in \text{pr}_{12}R'$ . Let  $\text{lk}_{12}, \text{lk}_{34}$  denote the link congruences of  $Q' = Q'(a, b, c, d)$  viewed as a subdirect product of  $Q'_{12} = \text{pr}_{12}Q'$  and  $Q'_{34} = \text{pr}_{34}Q'$ . Then*

- (1)  $(\underline{0}_1 \times \beta)|_{Q'_{12}} \subseteq \text{lk}_{12}$  and  $(\underline{0}_2 \times \delta)|_{Q'_{34}} \subseteq \text{lk}_{34}$ .
- (2) Let  $E = B \cap B'_1$ , where  $B$  is the  $\beta$ -block containing  $a, b$ . Then  $(\beta \times \beta)|_{\text{umax}(E) \times \text{umax}(E)} \subseteq \text{lk}_{12}$ , or equivalently,  $(\text{umax}(E) \times \text{umax}(E))^2 \subseteq \text{lk}_{12}$ .
- (3) for any  $c', d' \in C_2$  the pair  $(c', d')$  is as-maximal in a  $\text{lk}_{34}$ -block  $D'_{34}$ , and the as-component of  $D'_{34}$  containing  $(c', d')$  is either  $\{c'\} \times C_2$ , or  $C_2 \times C_2$ .
- (4) There is  $a' \in B'_1$  such that  $(a', a', c, d) \in Q'$ . In other words, there is a  $\overline{B}$ -preserving polynomial  $g$  of  $R$  such that  $g(a) = g(b)$  and  $g(c) = c, g(d) = d$ .

*Proof.* (1) The relation  $Q'$  contains tuples  $(a, b, c, d)$ ,  $(a, b, c', c')$ ,  $(a, a, c', c')$ ,  $(a, a, c, c)$  for some  $c' \in B'_2$ . Indeed,  $(a, b, c, d) \in Q'$  by definition,  $(a, a, c, c) \in Q'$  because  $(a, c) \in \text{pr}_{12}R'$ , and  $(a, b, c', c')$ ,  $(a, a, c', c')$  can be chosen to be the images of  $(a, b, c, d)$  and  $(a, a, c, c)$ , respectively, under a  $\overline{B}$ -preserving polynomial  $g$  such that  $g(a) = a, g(b) = b$  and  $g(\delta) \subseteq \underline{0}_2$ . Such a polynomial exists by the choice of  $a, b$  such that  $(a, b) \in T_0$  and because  $(\underline{0}_1, \beta)$  can be separated from  $(\underline{0}_2, \delta)$ . This implies that  $(a, b) \stackrel{\text{lk}_{12}}{\equiv} (a, a), (c, d) \stackrel{\text{lk}_{34}}{\equiv} (c, c)$ . Let  $\eta_{12}, \eta_{34}$  be congruences of  $\text{pr}_{12}Q(a, b, c, d), \text{pr}_{34}Q(a, b, c, d)$  generated by  $((a, b), (a, a))$  and  $((c, d), (c, c))$ , respectively. Then

$$\eta_{12}|_{Q'_{12}} = (\underline{0}_1 \times \beta)|_{Q'_{12}}, \quad \text{and} \quad \eta_{34}|_{Q'_{34}} = (\underline{0}_2 \times \delta)|_{Q'_{34}}.$$

Indeed, in the case of, say,  $\underline{0}_1 \times \beta$ , since  $(a, b) \stackrel{\underline{0}_1 \times \beta}{\equiv} (a, a)$ ,  $\eta_{12} \leq \underline{0}_1 \times \beta$ . On the other hand, the relation  $Q'_{12}$  consists of pairs  $(g(a), g(b))$  for  $\overline{B}$ -preserving unary polynomials  $g$  of  $\mathbb{A}_1$ . Since  $(a, b) \stackrel{\eta_{12}}{\equiv} (a, a)$ , for any  $(a', b') = (g(a), g(b)) \in Q'_{12}$  it holds that

$$(a', b') = (g(a), g(b)) \stackrel{\eta_{12}}{\equiv} (g(a), g(a)) = (a', a'),$$



showing that  $(\underline{Q}_1 \times \beta)|_{Q'_{12}} \subseteq \eta_{12}$ . For  $Q'_{34}$  and  $\underline{Q}_2 \times \delta$  the argument is similar.

Finally, as  $(a, b), (a, a)$  are in the same  $\text{lk}_{12}$ -block,  $\eta_{12}|_{Q'_{12}} \subseteq \text{lk}_{12}$ ; and, as  $(c, d), (c, c)$  are in the same  $\text{lk}_{34}$ -block,  $\eta_{34}|_{Q'_{34}} \subseteq \text{lk}_{34}$ . Item (1) is proved.

(2) By the definition of  $T_0$ , for any pair  $(a', b') \in T_0 \cap E^2$  there is a  $\overline{B}$ -preserving polynomial  $g$  satisfying  $g(a) = a', g(b) = b'$ , and  $g(B'_2) = \{c'\} \subseteq B'_2$ . Applying  $g$  to tuples  $(a, a, c, c)$ , and  $(b, b, d', d')$  for any  $d'$  such that  $(b, d') \in \text{pr}_{12}R'$ , we obtain  $(a', a', c', c'), (b', b', c', c') \in Q'$ . Therefore,  $(a', a') \stackrel{\text{lk}_{12}}{\equiv} (b', b')$ . Since  $R$  is chained with respect to  $\overline{\beta}, \overline{B}$ , by (Q1) and Lemma 37(4)  $(a'', a'') \stackrel{\text{lk}_{12}}{\equiv} (b'', b'')$  for any  $a'', b'' \in \text{umax}(E)$ . Together with part (1) this proves the result.

(3) Suppose first that for some  $e, e' \in B'_2$ ,  $e \neq e'$ , we have  $(e, e) \stackrel{\text{lk}_{34}}{\equiv} (e', e')$ . Then for any pair  $(e'', e''') \in T(\{(e, e'), \mathcal{U}_{\overline{B}}\})$  there is a  $\overline{B}$ -preserving polynomial  $g$  with  $g(e) = e'', g(e') = e'''$ . Applying this polynomial to the tuples witnessing that  $(e, e) \stackrel{\text{lk}_{34}}{\equiv} (e', e')$  we get  $(e'', e'') \stackrel{\text{lk}_{34}}{\equiv} (e''', e''')$ . Therefore by condition (Q1) all tuples of the form  $(x, x)$ ,  $x \in \text{umax}(B'_2)$ , are  $\text{lk}_{34}$ -related. Since by Lemma 37(3)  $(c', d')$  is a pair from  $T(\{(e, e'), \mathcal{U}_{\overline{B}}\})$ , using part (1) of Lemma 41 this implies that  $\text{lk}_{34}|_{Q'_{34}{}^u} = (\delta \times \delta)|_{Q'_{34}{}^u} = Q'_{34}{}^u \times Q'_{34}{}^u$ , where  $Q'_{34}{}^u = Q'_{34} \cap (\text{umax}(B'_2) \times \text{umax}(B'_2))$ . In particular,  $C_2^2$  is contained in  $Q'_{34}$ , and is contained in a  $\text{lk}_{34}$ -block  $D'_{34}$ . All elements of  $C_2^2$  are as-maximal in  $D'_{34}$ .

Suppose  $(e, e) \stackrel{\text{lk}_{34}}{\equiv} (e', e')$  for no  $e, e' \in B'_2$ ,  $e \neq e'$ . The inclusion  $(\underline{Q}_2 \times \delta)|_{Q'_{34}} \subseteq \text{lk}_{34}$  implies that if  $(c_1, d_1) \stackrel{\text{lk}_{34}}{\equiv} (c_2, d_2)$  then  $(c_1, c_1) \stackrel{\text{lk}_{34}}{\equiv} (c_2, c_2)$ . Therefore, by part (1) we have  $\text{lk}_{34}|_{Q'_{34}{}^u} = (\underline{Q}_2 \times \delta)|_{Q'_{34}{}^u}$ . In particular,  $\{c'\} \times C_2$  is contained in a  $\text{lk}_{34}$ -block. Since  $c', d'$  are as-maximal,  $(c', d')$  is as-maximal in this  $\text{lk}_{34}$ -block. Part (3) is proved.

(4) By Lemma 39(5) there is  $(a', b', c', d') \in Q'$  such that  $(a', b') \in \text{amax}(Q'^E_{12})$ ,  $Q'^E_{12} = Q'_{12} \cap E^2$ , for some  $\beta$ -block  $E$  and  $c', d' \in C_2$ ,  $c' \neq d'$ . We start by proving that  $(a', a')$  is as-maximal in  $Q'^E_{12}$ . Let  $E' = E \cap B'_1$  and let  $\text{lk}_1^{Q'}, \text{lk}_2^{Q'}$  be the link congruences of the two copies of  $B'_1$  with respect to  $Q'_{12}$ . As by Lemma 39(2)  $Q'_{12} \subseteq Q_{ab}^{\mathbb{A}_1}$ , we have  $\text{lk}_1^{Q'}, \text{lk}_2^{Q'} \leq \beta|_{B'_1}$ . On the other hand,  $Q'_{12}$  consists of pairs of the form  $(x, x)$  and pairs from  $T_1 = T(\{(a, b)\}, \mathcal{U}_{\overline{B}})$  and, as  $R$  is chained,  $\text{umax}(E')$  belongs to a block of the transitive closure of  $T_1$ . Therefore, it is easy to see that  $\text{umax}(E')$  is a subset of both a  $\text{lk}_1^{Q'}$ - and a  $\text{lk}_2^{Q'}$ -block. Indeed, let  $e, e' \in \text{umax}(E')$  and  $e = e_1, \dots, e_k = e'$  be such that  $\{e_i, e_{i+1}\} \in T_1$ . This means that either  $(e_i, e_{i+1}) \in Q'_{12}$  or  $(e_{i+1}, e_i) \in Q'_{12}$ . Since  $(e_i, e_i), (e_{i+1}, e_{i+1}) \in Q'_{12}$  by construction, in either case we have  $(e_i, e_{i+1}) \in \text{lk}_1^{Q'}, \text{lk}_2^{Q'}$ .

Let  $C_1$  be the as-component of  $E'$  containing  $a'$ ; such an as-component exists by the choice of  $a'$ . As  $(a', a') \in Q'_{12} \cap (C_1 \times C_1) \neq \emptyset$  and  $C_1 \subseteq \text{umax}(E')$ , by



Proposition 26  $C_1 \times C_1 \subseteq Q'_{12}$ . Since  $C_1$  is an as-component in  $E'$ , by Lemma 20  $C_1 \times C_1$  is an as-component in  $Q'^E_{12}$ . In particular  $(a', a')$  is as-maximal in  $Q'^E_{12}$ .

By Lemma 41(3)  $(c', d')$  is as-maximal in a  $\text{lk}_{34}$ -block, which is equal to  $\{c'\} \times C_2$  or to  $C_2^2$ . Therefore, by Lemma 39(5)  $\{(a', b')\} \times \{c'\} \times C_2 \subseteq Q'$ . Since by Lemma 41(2)  $(a', b')$  and  $(a', a')$  are as-maximal in a  $\text{lk}_{12}$ -block and  $(a', a', c', c') \in Q'$ , by Proposition 26  $\{(a', a')\} \times \{c'\} \times C_2 \subseteq Q'$ . In particular,  $(a', a', c', d') \in Q'$ . Therefore, there is a  $\overline{B}$ -preserving polynomial  $f$  such that  $f(a) = f(b)$  and  $f(c) = c', f(d) = d'$ .

Finally, as  $R$  is chained, by Lemma 37(3) there is a  $\overline{B}$ -preserving polynomial  $f'$  such that  $f'(c') = c, f'(d') = d$ . As is easily seen,  $g = f' \circ f$  satisfies the desired conditions.  $\square$

**Corollary 42.** *Let  $c, d \in C_2$  and a  $\beta$ -block  $E$ ,  $E \cap \text{umax}(B'_1) \neq \emptyset$ , be such that  $(a, c) \in \text{pr}_{12}R'$  for some  $a \in \text{umax}(E')$ ,  $E' = E \cap B'_1$ . Then there is a  $\overline{B}$ -preserving polynomial  $h$  such that  $|h(\text{umax}(E'))| = 1$  and  $h(c) = c, h(d) = d$ .*

*Proof.* If  $|\text{umax}(E')| = 1$  then  $h$  can be chosen to be the identity mapping, so we assume  $|\text{umax}(E')| > 1$ . Since  $a \in \text{umax}(E')$ , by Lemma 37(6)  $(a, b) \in T_0 = T(\{(a, b)\}, \mathcal{U}(\mathbb{Q}_2, \delta, \overline{B}))$  for some  $b \in E'$ . Let us consider  $T_0$  as a directed graph with all the loops present. By Lemma 37(1,5) any  $\overline{B}$ -preserving polynomial  $g$  is a homomorphism of  $T_0$ . Moreover, let  $T_{0c}$  denote the subgraph of  $T_0$  induced by  $(\text{pr}_{12}R')^{-1}[c]$ , that is,  $a'$  is a vertex of  $T_{0c}$  if and only if  $(a', c) \in \text{pr}_{12}R'$ . Note that any  $\overline{B}$ -preserving polynomial  $g$  such that  $g(c) = c$  maps  $T_{0c}$  to itself. Since  $R$  is chained, the set  $\text{umax}(E')$  is contained in a connected component (not necessarily strongly connected) of  $T_0$ .

By Lemma 41(4) for any  $(a', b') \in T_0$ ,  $a' \in V(T_{0c})$ , there is a polynomial  $h_{cd}^{a'b'}$  such that  $h_{cd}^{a'b'}(a') = h_{cd}^{a'b'}(b')$  and  $h_{cd}^{a'b'}(c) = c, h_{cd}^{a'b'}(d) = d$ . We construct a sequence of induced subgraphs  $T_0^i$  of  $T_0$  as follows. Let  $T_0^1 = T_0$ . Then if  $T_0^i$  contains an edge  $(a', b')$  or  $(b', a')$  such that  $a' \in V(T_{0c})$ , set  $T_0^{i+1} = h_{cd}^{a'b'}(T_0^i)$ . Observe that a connected component of  $T_0^i$  is mapped to a connected component of  $T_0^{i+1}$ . This means that when the sequence stops at, say,  $T_0^k$ , every connected component of  $T_0^k$  containing a vertex from  $T_{0c}$  is a singleton. Therefore, if  $g_i$  is the polynomial that is used to map  $T_0^i$  to  $T_0^{i+1}$ , then  $h = g_{k-1} \circ \dots \circ g_1$  is such that  $h(c) = c, h(d) = d$ , and  $|h(\text{umax}(E'))| = 1$ . The polynomial  $h$  is as required.  $\square$

Next, we show the existence of polynomials of  $R$  that satisfy certain conditions on  $B'_2$ .

**Lemma 43.** *Let  $c, d \in C_2$ ,  $c \neq d$ , and  $c' \in B'_2$ ,  $c' \notin \{c, d\}$ . Either there is a  $\overline{B}$ -preserving polynomial  $f$  of  $R$  such that  $f(c) = f(c') = c$  and  $f(d) = d$ , or for any  $\overline{B}$ -preserving polynomial  $f$  such that  $f(c') = c, f(d) = d$ , it holds that  $f(c) \neq d$ .*

*Proof.* We consider  $P(c', c, d)$ , and  $P'(c', c, d)$  introduced before Lemma 39. Let  $P' = P'(c', c, d)$ . As  $R$  is chained, by Lemma 39(4) and Lemma 37(3)  $C_2^2 \subseteq \text{pr}_{12}P', \text{pr}_{13}P', \text{pr}_{23}P'$ . Moreover, as  $(c, c, c) \in P'$ , by the Maximality Lemma 18(3)  $P' \cap C_2^3$  is subdirect. Suppose that there is a  $\overline{B}$ -preserving polynomial  $f$  of  $R$  such that  $f(c') = c$  and  $f(c) = f(d) = d$ . This means that  $(c, d, d) \in P'$ . Since  $(d, d, d) \in P'$ , and since  $R$  is chained, for any  $x, y \in C_2$  there is  $z \in B'_2$  such that  $(x, z, z), (y, z, z) \in P'$ . Therefore  $C_2$  is a subset of a class of the link congruence of  $\text{pr}_1P'$ , when  $P'$  is considered a subdirect product of  $\text{pr}_1P'$  and  $\text{pr}_{23}P'$ . Since  $C_2$  and  $C_2^2$  are as-components in  $\text{pr}_1P'$  and  $\text{pr}_{23}P'$ , respectively, by Proposition 26  $C_2^3 \subseteq P'$ . In particular,  $(c, c, d) \in P'$  witnessing that there is a  $\overline{B}$ -preserving polynomial  $f$  of  $R$  such that  $f(c) = f(c') = c$  and  $f(d) = d$ .  $\square$

*Proof of Theorem 38.* We need to show that there is a  $\overline{B}$ -preserving polynomial  $g$  such that  $g$  collapses  $\beta|_{B'_1}$  but does not collapse  $\beta_2 = \delta$ .

First we show that for any  $c, d \in C_2$  and any  $(a, b) \in \beta|_{B'_1}$  there is a polynomial  $h_{cd}^{ab}$  of  $R$  such that

- (1)  $h_{cd}^{ab}$  is idempotent;
- (2)  $h_{cd}^{ab}(a) = h_{cd}^{ab}(b)$ ;
- (3)  $h_{cd}^{ab}(c) = c, h_{cd}^{ab}(d) = d$ .

Fix  $c, d \in C_2$ . By Corollary 42 such a polynomial exists whenever  $a, b \in \text{umax}(E \cap B'_1)$  for a  $\beta$ -block  $E$ ,  $E \cap \text{umax}(B'_1) \neq \emptyset$ , and  $(a', c) \in \text{pr}_{12}R'$  for some  $a' \in \text{umax}(E \cap B'_1)$ . We need to prove this for arbitrary  $a, b \in B'_1$  with  $a \stackrel{\beta}{=} b$ .

Take any  $c' \in B'_2$  such that  $(a, c') \in \text{pr}_{12}R'$ . If  $c' = d$  we swap the roles of  $c$  and  $d$ . If there exists  $h_{dc}^{ab}$  with the required properties, this polynomial can also be chosen as  $h_{cd}^{ab}$ . Otherwise by Lemma 37(3) there is a  $\overline{B}$ -preserving polynomial  $g$  such that  $g(c') = c, g(d) = d$ . If, see Lemma 43, there is a polynomial that maps  $c', c$  to  $c$  and  $d$  to  $d$ , we set  $g$  to be that polynomial. Let  $Q' = Q'(g(a), g(b), c, d)$  be as defined before Lemma 39. We use the notation from Lemma 39. By Lemma 39(1) it suffices to prove that a tuple of the form  $(a^*, a^*, c'', d'')$ ,  $c'', d'' \in C_2, c'' \neq d''$  belongs to  $Q'$ . Indeed, if this is the case, there is a  $\overline{B}$ -preserving polynomial  $h$  of  $R$  such that  $h(g(a)) = h(g(b))$  and  $h(c) = c'', h(d) = d''$ . Since  $R$  is chained, there is also a  $\overline{B}$ -preserving polynomial  $h'$  such that  $h'(c'') = c, h'(d'') = d$ . Set  $g' = h' \circ h \circ g$ . If  $g$  is such that  $g(c) = c$ , then  $g'(a) = g'(b)$  and  $g'(c) = c, g'(d) = d$ , and we can set  $h_{cd}^{ab} = g'$ . Otherwise by Lemma 43  $g'(c) \neq d$  and there is a  $\overline{B}$ -preserving polynomial  $h''$  such that  $h''(g'(c)) = c$  and  $h''(d) = d$ . As is easily seen, the polynomial  $h_{cd}^{ab} = h'' \circ g'$  is as required.

Finally, we use polynomials  $h_{cd}^{ab}$  to construct a single polynomial that collapses  $\beta$  on  $E' = E \cap B'_1$  for every  $\beta$ -block  $E$ . Fix  $c, d$  and  $h_{cd}^{ab}$  for every pair  $a, b \in B'_1$ ,  $a \stackrel{\beta}{\equiv} b$ . Let  $V_1, \dots, V_k$  be the list of all such pairs, and if  $V_\ell = \{a, b\}$  is the pair number  $\ell$ ,  $h^\ell$  denotes  $h_{cd}^{ab}$ . Take a sequence  $1 = \ell_1, \ell_2, \dots$  such that  $h^{(1)} = h^{\ell_1}$ ,  $V_{\ell_2}$  is a subset of  $h^{(1)}(\mathbb{A}_1)$ , and, for  $s > 2$ ,  $V_{\ell_s}$  is a subset of the range of  $h^{(s-1)} = h^{\ell_{s-1}} \circ \dots \circ h^{\ell_1}$ . Since  $|\text{Im}(h^{(s)})| < |\text{Im}(h^{(s-1)})|$ , there is  $r$  such that  $\text{Im}(h^{(r)})$  contains no pair  $V_\ell$  for any  $\ell$ . Therefore setting  $h(x) = h_{\ell_r} \circ \dots \circ h_{\ell_1}(x)$  we have that  $h$  collapses all the pairs  $V_\ell$ , and  $h$  acts identically on  $\{c, d\}$ . The result follows.  $\square$

### 3.5 Separation and minimal sets

The results of this section hold for arbitrary algebras, not only for algebras from  $\mathcal{V}$ .

We say that prime intervals  $(\alpha, \beta)$  and  $(\gamma, \delta)$  *cannot be separated* if  $(\alpha, \beta)$  cannot be separated from  $(\gamma, \delta)$  and  $(\gamma, \delta)$  cannot be separated from  $(\alpha, \beta)$ . In this section we show a connection between the fact that two prime intervals cannot be separated, their types, and link congruences.

**Lemma 44.** *Let  $\mathbb{A}$  be an algebra.*

- (1) *If prime intervals  $(\alpha, \beta)$  and  $(\gamma, \delta)$  in  $\text{Con}(\mathbb{A})$  are perspective, they cannot be separated.*
- (2) *If  $\alpha \prec \beta$  and  $\gamma \prec \delta$  from  $\text{Con}(\mathbb{A})$  cannot be separated, then a set  $U$  is a  $(\alpha, \beta)$ -minimal set if and only if it is a  $(\gamma, \delta)$ -minimal set.*
- (3) *Let  $R$  be a subdirect product of  $\mathbb{A}$  and  $\mathbb{B}$ ,  $\alpha, \beta \in \text{Con}(\mathbb{A})$ ,  $\gamma, \delta \in \text{Con}(\mathbb{B})$  such that  $\alpha \prec \beta$ ,  $\gamma \prec \delta$ , and let  $\alpha \prec \beta$  and  $\gamma \prec \delta$  cannot be separated. Then for any  $(\alpha, \beta)$ -minimal set  $U$  there is a unary idempotent polynomial  $f$  such that  $f(\mathbb{A}) = U$  and  $f(\mathbb{B})$  is a  $(\gamma, \delta)$ -minimal set.*

*Proof.* (1) Follows from Lemma 3.

(2) Let  $f$  be a polynomial of  $\mathbb{A}$  such that  $f(\mathbb{A}) = U$  and  $f(\beta) \not\subseteq \alpha$ . Since  $(\alpha, \beta)$  cannot be separated from  $(\gamma, \delta)$ , we have  $f(\delta) \not\subseteq \gamma$  and therefore by Lemma 1(6)  $U$  contains a  $(\gamma, \delta)$ -minimal set  $U'$ . If  $U' \neq U$ , there is a polynomial  $g$  with  $g \circ f(\delta) \not\subseteq \gamma$  and  $g \circ f(\mathbb{A}) = U'$ . In particular,  $|g(U)| < |U|$ , and so  $g \circ f(\beta) \subseteq \alpha$ ; a contradiction with the assumption that  $(\gamma, \delta)$  cannot be separated from  $(\alpha, \beta)$ .

(3) Take an idempotent polynomial  $g$  of  $R$  such that  $g(\mathbb{B})$  is a  $(\gamma, \delta)$ -minimal set. Then, as  $(\gamma, \delta)$  cannot be separated from  $(\alpha, \beta)$ ,  $g(\beta) \not\subseteq \alpha$ . By Lemma 1(6) there is an  $(\alpha, \beta)$ -minimal set  $U' \subseteq g(\mathbb{A})$ . Let  $g', h$  be polynomials of  $R$  such that  $g'(U) = U'$ ,  $h(U') = U$  and  $h(\mathbb{A}) = U$ , which exist by Lemma 1(1). Then

$h' = h \circ g \circ g'$  is such that  $h'(\mathbb{A}) = h'(U) = U$ ,  $h'(\beta) \not\subseteq \alpha$  and therefore  $h'(\delta) \not\subseteq \gamma$ . Then iterating  $h'$  sufficiently many times we get an idempotent polynomial  $f$  satisfying the same properties.  $\square$

**Lemma 45.** *Let  $R$  be a subdirect product of  $\mathbb{A}$  and  $\mathbb{B}$  and let  $\alpha, \beta \in \text{Con}(\mathbb{A})$ ,  $\gamma, \delta \in \text{Con}(\mathbb{B})$  be such that  $\alpha \prec \beta$ ,  $\gamma \prec \delta$ , and  $(\alpha, \beta), (\gamma, \delta)$  cannot be separated. Then  $\text{typ}(\alpha, \beta) = \text{typ}(\gamma, \delta)$ .*

*Proof.* Let  $f$  be a unary idempotent polynomial of  $R$  such that  $f(\mathbb{A}) = U$  is an  $(\alpha, \beta)$ -minimal set and  $f(\mathbb{B}) = V$  is a  $(\gamma, \delta)$ -minimal set. Observe that, as  $R$  is subdirect,  $R \cap (U \times V)$  is subdirect on  $U$  and  $V$ . Let also  $N$  be an  $(\alpha, \beta)$ -trace in  $U$ . Let first  $\text{typ}(\alpha, \beta), \text{typ}(\gamma, \delta) \in \{1, 2\}$ , say,  $\text{typ}(\alpha, \beta) = 1$  and  $\text{typ}(\gamma, \delta) = 2$ . Then, every polynomial of  $R$  is essentially unary on  $N/\alpha$ . Also, for any  $(\gamma, \delta)$ -trace  $N' \subseteq V$ ,  $N'/\gamma$  is polynomially equivalent to a one-dimensional vector space over  $\text{GF}(q)$  where  $q$  is a prime power. There is a polynomial  $g(x, y)$  of  $R$  such that  $g$  is the first projection on  $U/\alpha$  and  $x + y$  on  $N'/\gamma$ . Then the polynomial  $h(x) = g\left(\begin{pmatrix} a \\ b \end{pmatrix}, x\right)$  for any  $(a, b) \in R \cap (U \times V)$ ,  $b \in N'$ , witnesses that  $(\gamma, \delta)$  can be separated from  $(\alpha, \beta)$ .

Suppose now that  $\text{typ}(\gamma, \delta) \neq 1, 2$  and  $N'$  is the only  $(\gamma, \delta)$ -trace in  $V$  and  $N' = \{1'\} \cup O'$ , where  $\{1'\}$  and  $O'$  are the two  $\gamma|_V$ -blocks contained in  $N'$ , and  $p(x, y)$  is the pseudo-meet operation on  $V$  that exists by Lemma 4. In particular  $p(x, x)$  is idempotent.

**CLAIM.** If there exists a binary polynomial  $h \in \text{Pol}_2(R)$  such that  $h(R) \subseteq U \times V$ ,  $h$  acts as  $p$  on  $V/\gamma$ , and  $h$  is the first projection on  $N/\alpha$ , then  $(\alpha, \beta)$  and  $(\gamma, \delta)$  can be separated.

Suppose  $(a, b) \in R \cap (U \times V)$  for  $a \in N$  and  $b \in V - \{1'\}$ . Then the polynomial  $g(x) = h\left(x, \begin{pmatrix} a \\ b \end{pmatrix}\right)$  is the first projection on  $N/\alpha$  but is  $p(x, b) \stackrel{\gamma}{\equiv} b$  on  $N'$ . Since  $N'$  is the only  $(\gamma, \delta)$ -trace in  $V$ , we also get  $p(a_1, b) \stackrel{\gamma}{\equiv} p(a_2, b)$  whenever  $a_1 \stackrel{\delta}{\equiv} a_2$  for  $a_1, a_2 \in V$ . Therefore,  $g(\beta) \not\subseteq \alpha$  and  $g(\delta) \subseteq \gamma$ . Otherwise there is  $(a, 1') \in R$ ,  $a \in N$ . Set  $g(x) = h\left(\begin{pmatrix} a \\ 1' \end{pmatrix}, x\right)$ . We have  $g(x) \stackrel{\alpha}{\equiv} a$  on  $N$  and  $g(x) \stackrel{\gamma}{\equiv} x$  on  $N'$ . The result follows.

If  $\text{typ}(\alpha, \beta) = 1$ , then  $p$  is a projection on  $N/\alpha$  and  $(\gamma, \delta)$  can be separated from  $(\alpha, \beta)$  by the Claim.

If  $\text{typ}(\alpha, \beta) = 2$ , then the  $(\alpha, \beta)$ -trace  $N/\alpha$  is polynomially equivalent to a one-dimensional vector space over  $\text{GF}(q)$  where  $q$  is a prime power. Then  $p$  on  $N/\alpha$  is a linear operation of the form  $\varrho x + (1 - \varrho)y$ ,  $\varrho \in \text{GF}(q)$ . We may assume

that  $\varrho = 1$ . Indeed, if  $\varrho = 0$  then consider  $p(y, x)$  instead of  $p(x, y)$ . Otherwise the operation

$$\underbrace{p(p \dots p(x, y), y \dots, y)}_{q-1 \text{ times}}$$

satisfies the required conditions. Then  $p$  satisfies the conditions of the Claim above.

Next assume  $\text{typ}(\alpha, \beta) \in \{4, 5\}$  and  $\text{typ}(\gamma, \delta) = 3$ . Let  $N'$  be the only  $(\gamma, \delta)$ -trace in  $V$ , and let  $\{1\}, O$  be the two  $\gamma|_V$ -blocks contained in  $V$ . In this case  $|O'| = 1$ , say,  $O' = \{o'\}$ , and there is a unary polynomial  $h$ ,  $h(R) \subseteq U \times V$ , such that  $h(1') = o'$  and  $h(o') = 1'$ . Consider  $g(x) = p(x, h(x))$ ; we have  $g(x) = o'$  on  $N'$ . Since  $(\alpha, \beta), (\gamma, \delta)$  cannot be separated  $h(x)$  is a permutation on  $N/\alpha$ , that is,  $h(x) = x$  on  $N/\alpha$ . This implies that  $g(\beta) \not\subseteq \alpha$  and  $g(\delta) \subseteq \gamma$ , a contradiction again.

Finally, let  $\text{typ}(\alpha, \beta) = 5$  and  $\text{typ}(\gamma, \delta) = 4$ . Then in addition to the pseudo-meet operation there is also a pseudo-join operation  $q \in \text{Pol}_2(R)$  having the properties listed in Lemma 4. If any of  $p, q$  act as a projection on  $N/\alpha$  the result follows by the Claim. Suppose that both  $p$  and  $q$  are semilattice operations on  $N/\alpha$ . Since  $\text{typ}(\alpha, \beta) = 5$ , they are the same semilattice operation on  $N/\alpha$ . Then  $g(x) = p(x, q(x, y))$  satisfies the conditions of the Claim with the roles of  $\mathbb{A}$  and  $\mathbb{B}$  swapped.  $\square$

## 4 Centralizers

### 4.1 Centralizers and quasi-centralizers

The results of Section 4 hold for arbitrary algebras. If  $\mathbb{A}$  is an algebra,  $g \in \text{Pol}_2(\mathbb{A})$  and  $a \in \mathbb{A}$ , then  $g^a$  denotes the unary polynomial  $g^a(x) = g(a, x)$ . Recall that for  $\alpha, \beta \in \text{Con}(\mathbb{A})$ , the *centralizer* (see, eg. [17])  $(\alpha : \beta)$  is the largest congruence  $\theta \in \text{Con}(\mathbb{A})$  which centralizes  $\beta$  modulo  $\alpha$ , i.e., satisfies the condition  $C(\theta, \beta; \alpha)$  given by

$$\text{for any } f \in \text{Pol}_2(\mathbb{A}), \text{ any } (a, b) \in \theta \text{ and any } (c, d) \in \beta \text{ it holds that } f^a(c) \stackrel{\alpha}{\equiv} f^a(d) \text{ if and only if } f^b(c) \stackrel{\alpha}{\equiv} f^b(d).$$

In [7] we also introduced a somewhat related notion of *quasi-centralizer*  $\zeta(\alpha, \beta)$ :

$$(a, b) \in \zeta(\alpha, \beta) \text{ if for any } g \in \text{Pol}_2(D), g^a(\beta) \subseteq \alpha \text{ if and only if } g^b(\beta) \subseteq \alpha.$$

A relation basically identical to quasi-centralizer also appeared in [18], but completely inconsequentially, they did not study it at all. Kearnes observed that  $(\alpha :$

$\beta) \subseteq \zeta(\alpha, \beta)$ , and later Willard [25] demonstrated that the notions of the centralizer and the quasi-centralizer are equivalent at least in the case important for the purpose of this paper. With Willard's permission we reproduce his proof here and will use the usual centralizer in the sequel.

**Proposition 46.** *Let  $\mathbb{A}$  be a finite algebra,  $\alpha, \beta \in \text{Con}(\mathbb{A})$ ,  $\alpha \prec \beta$ . If  $\text{typ}(\alpha, \beta) \neq 1$ , then  $\zeta(\alpha, \beta) = (\alpha : \beta)$ .*

*Proof.* Let  $\theta = (\alpha : \beta)$ . We first show  $\theta \subseteq \zeta(\alpha, \beta)$ . Suppose  $(a, b) \in \theta$ ,  $g \in \text{Pol}_2(\mathbb{A})$ , and  $g^a(\beta) \subseteq \alpha$ . Pick  $(c, d) \in \beta$ . Then  $g(a, c) = g^a(c) \stackrel{\alpha}{=} g^a(d) = g(a, d)$ . As  $C(\theta, \beta; \alpha)$ ,  $(a, b) \in \theta$  and  $(c, d) \in \beta$ , we can replace the  $a$ 's with  $b$ 's to get  $g(b, c) \stackrel{\alpha}{=} g(b, d)$ , i.e.,  $(g^b(c), g^b(d)) \in \alpha$ , proving  $g^b(\beta) \subseteq \alpha$ . The reverse implication is proved similarly, so  $(a, b) \in \zeta(\alpha, \beta)$ . This proves  $\theta \subseteq \zeta(\alpha, \beta)$ .

Next we prove that  $\zeta(\alpha, \beta) \subseteq \theta$ . Fix an  $(\alpha, \beta)$ -minimal set  $U$  and a unary idempotent polynomial  $e \in \text{Pol}_1(\mathbb{A})$  satisfying  $e(\mathbb{A}) = U$ . Now consider cases.

CASE 1.  $\text{typ}(\alpha, \beta) \neq 2$ .

We will show  $\zeta(\alpha, \beta) \cap \beta = \alpha$ , as  $C(\zeta(\alpha, \beta), \beta; \alpha)$  will then follow ([18, Proposition 3.4(4)]), which will then imply  $\zeta(\alpha, \beta) \leq \theta$ . Clearly  $\alpha \subseteq \zeta(\alpha, \beta) \cap \beta$ , so it remains to prove the opposite inclusion. Let  $N$  be the unique  $(\alpha, \beta)$ -trace in  $U$ . Write  $N = \{1\} \cup O$ , where  $N^2 \subseteq \beta$  and  $\{1\}, O$  are the two  $\alpha|_U$ -classes contained in  $N$ , and for which there exists a pseudo-meet operation  $p$ , see Lemma 4, of  $\mathbb{A}|_U$  for  $N$  with neutral element 1. Thus  $p$  is a binary polynomial of  $\mathbb{A}|_U$  satisfying (among other things)  $p(1, x) = x$  and  $p(o, x) \neq 1$  for all  $x \in U$  and  $o \in O$ . Suppose  $(a, b) \in (\zeta(\alpha, \beta) \cap \beta) - \alpha$ . By Lemma 1(4) there exists  $f \in \text{Pol}_1(\mathbb{A})$  such that  $(f(a), f(b)) \in \beta|_U - \alpha|_U$ ; thus without loss of generality we can assume  $a, b \in U$ . Because  $N$  is the unique  $(\alpha, \beta)$ -trace of  $U$ , one of  $a, b$  must equal 1 and the other must be in  $O$ . Assume for concreteness that  $b = 1$  and  $a \in O$ . Define  $g(x, y) = p(e(x), e(y)) \in \text{Pol}_2(\mathbb{A})$ . Then  $g^1(1) = p(1, 1) = 1$  while  $g^1(a) = p(1, a) = a$ , so  $g^1(\beta|_U) \not\subseteq \alpha$  which certainly implies  $g^1(\beta) \not\subseteq \alpha$ . Suppose there exists  $(c, d) \in \beta$  such that  $(g^a(c), g^a(d)) \notin \alpha$ . Let  $c' = e(c)$  and  $d' = e(d)$ , so  $(c', d') \in \beta|_U$  and  $(p(a, c'), p(a, d')) \in \beta|_U - \alpha|_U$ . Since  $N$  is the unique  $\beta|_U$ -class containing more than one  $\alpha|_U$ -class, one of  $p(a, c')$  or  $p(a, d')$  must equal 1, which is impossible as  $a \in O$ . Thus such  $(c, d)$  does not exist, which proves  $g^a(\beta) \subseteq \alpha$ . This contradicts the assumption that  $(a, 1) = (a, b) \in \zeta(\alpha, \beta)$ .

CASE 2.  $\text{typ}(\alpha, \beta) = 2$ .

Suppose  $(a, b) \in \zeta(\alpha, \beta)$ . To prove  $(a, b) \in \theta$ , it suffices by the proof of [21, Lemma 2.6] to show that for all  $f \in \text{Pol}_2(\mathbb{A})$ ,  $e \circ f^a|_U$  is a permutation if and only if  $e \circ f^b|_U$  is a permutation. Suppose for concreteness that  $e \circ f^a|_U$  is a permutation but  $e \circ f^b|_U$  is not. Let  $g(x, y) = e(f(x, e(y))) \in \text{Pol}_2(\mathbb{A})$ . Because  $e \circ f^a|_U$  is

a permutation,  $e|_U$  is the identity map, and  $\beta|_U \not\subseteq \alpha$ , we get  $g^a(\beta) \not\subseteq \alpha$ . On the other hand, because  $e \circ f^b|_U$  is not the identity map, we get  $e \circ f^b(\beta|_U) \subseteq \alpha$  by Lemma 1(2) and hence  $g^b(\beta) \subseteq \alpha$ , contradicting  $(a, b) \in \zeta(\alpha, \beta)$ .  $\square$

## 4.2 Alignment

In this and the next sections we prove several properties of the centralizer. The first one concerns properties of a relation with respect to centralizer blocks.

Let  $R$  be a subdirect product of  $\mathbb{A}_1 \times \cdots \times \mathbb{A}_n$ ,  $i, j \in [n]$ , and  $\alpha_i \in \text{Con}(\mathbb{A}_i)$ ,  $\alpha_j \in \text{Con}(\mathbb{A}_j)$ . The coordinate positions  $i, j$  are said to be  $\alpha_i \alpha_j$ -aligned in  $R$  if, for any  $(a, c), (b, d) \in \text{pr}_{ij} R$ ,  $(a, b) \in \alpha_i$  if and only if  $(c, d) \in \alpha_j$ . Or in other words, the link congruences of  $\mathbb{A}_i, \mathbb{A}_j$  with respect to  $\text{pr}_{ij} R$  are no greater than  $\alpha_i, \alpha_j$ , respectively.

**Lemma 47.** *Let  $R$  be a subdirect product of  $\mathbb{A}_1 \times \mathbb{A}_2$ ,  $\alpha_i, \beta_i \in \text{Con}(\mathbb{A}_i)$ ,  $\alpha_i \prec \beta_i$ , for  $i = 1, 2$ . If  $(\alpha_1, \beta_1)$  and  $(\alpha_2, \beta_2)$  cannot be separated from each other, then the coordinate positions 1,2 are  $\zeta(\alpha_1, \beta_1)\zeta(\alpha_2, \beta_2)$ -aligned in  $R$ . If, in addition,  $\text{typ}(\alpha_1, \beta_1), \text{typ}(\alpha_2, \beta_2) \neq \mathbf{1}$ , then the coordinate positions 1,2 are  $(\alpha_1 : \beta_1)(\alpha_2 : \beta_2)$ -aligned in  $R$ .*

*Proof.* Let us assume the contrary, that is, without loss of generality there are  $a, b \in \mathbb{A}_1$  and  $c, d \in \mathbb{A}_2$  with  $(a, c), (b, d) \in R$ ,  $(a, b) \in \zeta(\alpha_1, \beta_1)$ , but  $(c, d) \notin \zeta(\alpha_2, \beta_2)$ . Therefore there is  $g(x, y) \in \text{Pol}_2(\mathbb{A}_2)$  such that  $g^c(\beta_2) \subseteq \alpha_2$  but  $g^d(\beta_2) \not\subseteq \alpha_2$ , or the other way round. Extend  $g$  to a polynomial of  $R$ . We have  $g^a(\beta_1) \subseteq \alpha_1$  if and only if  $g^b(\beta_1) \subseteq \alpha_1$ . Therefore, there is a polynomial of  $R$  that separates  $(\alpha_1, \beta_1)$  from  $(\alpha_2, \beta_2)$  or the other way round, a contradiction. The second statement of the lemma follows from Proposition 46.  $\square$

By [18, Theorem 5.5] if  $\text{typ}(\alpha, \beta) \neq \mathbf{1}, \mathbf{2}$  then  $(\alpha : \beta) \wedge \beta \leq \alpha$ . This and Lemmas 45,47 imply the following

**Lemma 48.** *Let  $R$  be a subdirect product of  $\mathbb{A}$  and  $\mathbb{B}$  and let  $\alpha, \beta \in \text{Con}(\mathbb{A})$ ,  $\gamma, \delta \in \text{Con}(\mathbb{B})$  be such that  $\alpha \prec \beta$ ,  $\gamma \prec \delta$ , and  $(\alpha, \beta), (\gamma, \delta)$  cannot be separated. Let also  $\text{lk}_1, \text{lk}_2$  be the link congruences of  $\mathbb{A}, \mathbb{B}$ , respectively. If  $\text{typ}(\alpha, \beta) \neq \mathbf{1}, \mathbf{2}$  then  $\text{lk}_1 \wedge \beta \leq \alpha$ ,  $\text{lk}_2 \wedge \delta \leq \gamma$ .*

## 4.3 Centralizer and congruence blocks

In this section we prove several properties indicating relationship between congruence blocks inside a centralizer block. All the algebras are assumed to be from  $\mathcal{V}$ .



**Lemma 49.** *Let  $\mathbb{A} \in \mathcal{V}$ ,  $\alpha, \beta \in \text{Con}(\mathbb{A})$  be such that  $\alpha \prec \beta$  and  $\text{typ}(\alpha, \beta) = \mathbf{2}$ , and so  $\beta \leq \zeta = (\alpha : \beta)$ , and let  $B, C$  be  $\beta$ -blocks from the same  $\zeta$ -block such that  $BC$  is a thin edge in  $\mathbb{A}/\beta$  or  $B = C$ . For any  $b \in B, c \in C$  such that  $bc$  is a thin edge of the same type (such  $b, c$  exist by Lemma 12(1)), or any type if  $B = C$ , the polynomial  $f(x) = x \cdot c$  if  $b \leq c$ ,  $f(x) = t_{bc}(x, c)$  if  $bc$  is majority, and  $f(x) = h_{bc}(x, b, c)$  if  $bc$  is affine, where  $t_{bc}, h_{bc}$  are the operations from Lemma 13, is an injective mapping from  $B/\alpha$  to  $C/\alpha$  and  $f(b) = c$ .*

*Moreover, if  $BC$  is a semilattice edge then for any  $a \in B/\alpha$ ,  $a \leq f(a)$  and  $a \not\leq b$  for any other  $b \in C/\alpha$ .*

*Proof.* We can assume that  $\alpha$  is the equality relation  $\underline{0}_1$ . Suppose  $f(a_1) = f(a_2)$  for some  $a_1, a_2 \in B$ . Since  $\text{typ}(\underline{0}_1, \beta) = \mathbf{2}$ , by Corollary 31(1) every pair of elements of  $B$  is a  $(\underline{0}_1, \beta)$ -subtrace. Let  $f'$  be an idempotent unary polynomial such that  $f'(a_1) = a_1, f'(a_2) = a_2$ , and  $f'(\mathbb{A})$  is a  $(\underline{0}_1, \beta)$ -minimal set. Note that if  $B = C$  and  $b \neq c$  then  $bc$  has to be of the affine type.

If  $b \leq c$ , let  $g(x, y) = f'(y) \cdot x$ . Then  $g^c(x) = g(c, x) = f(x)$  on  $\{a_1, a_2\}$ , that is,  $g^c(a_1) = g^c(a_2)$  implying  $g^c(\beta) \subseteq \underline{0}_1$ . On the other hand,  $g^b(x) = f'(x)$  on  $\{a_1, a_2\}$  implying  $g^b(\beta) \not\subseteq \underline{0}_1$ , a contradiction with the assumption  $b \stackrel{\zeta}{\equiv} c$ .

If  $bc$  is a thin majority edge, set  $g(x, y) = t_{bc}(f'(y), x)$ . Then  $g^c(a_1) = f(a_1) = f(a_2) = g^c(a_2)$ , and so  $g^c(\beta) \subseteq \underline{0}_1$ . On the other hand, since  $B$  is a module,  $a_1b, a_2b$  are affine edges and  $\underline{0}_1$  witnesses that. Therefore  $g^b(a_1) = a_1$  and  $g^b(a_2) = a_2$ , implying  $g^b(\beta) \not\subseteq \underline{0}_1$ , and we have a contradiction again.

If  $bc$  is a thin affine edge, we consider the polynomials  $g(x, y, z) = h_{bc}(f'(x), y, z)$  and  $g^{bc}(x) = g(x, b, c)$ ,  $g^{a_1a_1}(x) = g(x, a_1, a_1)$ . Again,  $g^{bc}(a_1) = f(a_1) = f(a_2) = g^{bc}(a_2)$ , while

$$g^{a_1a_1}(a_1) = h_{bc}(f'(a_1), a_1, a_1) = a_1 \neq a_2 = h_{bc}(f'(a_2), a_1, a_1) = g^{a_1a_1}(a_2),$$

since  $a_1a_2$  is an affine edge as witnessed by the equality congruence. Here  $b, c$  and  $a_1, a_2$  play the role  $a, b$  and  $c, d$ , respectively, in Lemma 13. This implies that  $g^{bc}(\beta) \subseteq \underline{0}_1$  and  $g^{a_1a_1}(\beta) \not\subseteq \underline{0}_1$ , a contradiction.

Finally, to prove the last statement of the lemma, suppose that there are  $c, d \in C$ ,  $c \neq d$ , such that  $a \leq c$  and  $a \leq d$ . Consider the term operation  $t(x, y) = x \cdot y$ . Then  $t^a(c) = c$  and  $t^a(d) = d$ . However, since  $C$  is a module, it contains no semilattice edges, and hence  $t^c(c) = c = t^c(d)$ . This contradicts the assumption  $(a, c) \in \zeta$ .  $\square$

**Corollary 50.** *Let  $\mathbb{A} \in \mathcal{V}$ ,  $\alpha, \beta \in \text{Con}(\mathbb{A})$  be such that  $\alpha \prec \beta$  and  $\text{typ}(\alpha, \beta) = \mathbf{2}$ , and so  $\beta \leq \zeta = (\alpha : \beta)$ . Let  $a, b, c \in \mathbb{A}$  be such that  $a \stackrel{(\alpha:\beta)}{\equiv} b \stackrel{\beta}{\equiv} c$ . Then  $ab \stackrel{\alpha}{\equiv} ac$ .*

*Proof.* We have  $ab \stackrel{\beta}{\equiv} ac$  and  $a \leq ab, a \leq ac$ . By the last statement of Lemma 49  $ab \stackrel{\alpha}{\equiv} ac$ .  $\square$

Another straightforward application of Lemma 49 is the following

**Corollary 51.** *Let  $\mathbb{A} \in \mathcal{V}$ ,  $\alpha, \beta \in \text{Con}(\mathbb{A})$  be such that  $\alpha \prec \beta$  and  $\text{typ}(\alpha, \beta) = 2$ , and let  $\zeta = (\alpha : \beta)$ . Then for any  $\beta$ -blocks  $B_1, B_2$  that belong to the same  $\zeta$ -block  $C$  and such that  $B_1 \sqsubseteq^{asm} B_2$  and  $B_2 \sqsubseteq^{asm} B_1$  in  $C/\beta$ ,  $|B_1/\alpha| = |B_2/\alpha|$ .*

## 5 Collapsing polynomials

In this section we introduce and prove the existence of polynomials that collapse all prime intervals in congruence lattices of factors of a subproduct, except for a set of factors that cannot be separated from each other.

We start with an auxiliary statement.

**Lemma 52.** *Let  $\mathbb{A} \in \mathcal{V}$  and  $\alpha, \beta \in \text{Con}(\mathbb{A})$  be such that  $\alpha \prec \beta$  and  $\text{typ}(\alpha, \beta) = 2$ ; let  $B$  be a  $\beta$ -block containing more than one  $\alpha$ -block, and  $a, b \in B$  with  $a \sqsubseteq_B^{asm} b$ .*

- (1) *There exists a polynomial  $f$  such that  $f(a) = b$  and  $f(\beta|_B) \not\subseteq \alpha$ .*
- (2) *If  $a$  belongs to an  $(\alpha, \beta)$ -trace, so does  $b$ . In particular, every element from  $\text{umax}(B)$  belongs to an  $(\alpha, \beta)$ -trace.*
- (3) *Let  $a \stackrel{\alpha}{\equiv} b$ ,  $a \sqsubseteq^{asm} b$  in  $a/\alpha$ , and  $N$  an  $(\alpha, \beta)$ -trace with  $a \in N$ . Then there is a polynomial  $f$  such that  $f(a) = b$ ,  $N' = f(N)$  is an  $(\alpha, \beta)$ -trace containing  $b$ , and  $N'/\alpha = N/\alpha$ .*

*Proof.* (1) follows from Lemma 49.

(2) Suppose  $a \in N$ , an  $(\alpha, \beta)$ -trace. Then the polynomial  $f$  constructed in item (1) satisfies  $f(a) = b$  and for any  $a' \in N$  with  $a'/\alpha \neq a/\alpha$  it holds that  $f(a') \not\stackrel{\alpha}{\equiv} f(a)$ . Therefore  $f(N)$  is a trace and  $b \in f(N)$ . The second claim of item (2) follows from the observation that by Lemma 1(5)  $B$  contains an  $(\alpha, \beta)$ -trace.

(3) Let  $f$  be the polynomial constructed in the proof of item (1). It suffices to notice that if  $a \stackrel{\alpha}{\equiv} b$  then  $f(x) \stackrel{\alpha}{\equiv} f(x)$  for  $x \in B$ .  $\square$

Let  $R$  be a subdirect product of  $\mathbb{A}_1, \dots, \mathbb{A}_n \in \mathcal{V}$ ,  $\beta_j \in \text{Con}(\mathbb{A}_j)$ , and  $B_j$  a  $\beta_j$ -block,  $j \in [n]$ . Let also  $i \in [n]$ , and  $\alpha, \beta \in \text{Con}(\mathbb{A}_i)$  be such that  $\alpha \prec \beta \leq \beta_i$ . We call an idempotent unary polynomial  $f$  of  $R$   $\alpha\beta$ -collapsing for  $\overline{\beta}, \overline{B}$  if

- (a)  $f$  is  $\overline{B}$ -preserving;

- (b)  $f(\mathbb{A}_i)$  contains an  $(\alpha, \beta)$ -minimal set, in particular,  $f(\beta) \not\subseteq \alpha$ ;
- (c)  $f(\delta|_{B_j}) \subseteq \gamma|_{B_j}$  for every  $\gamma, \delta \in \text{Con}(\mathbb{A}_j)$ ,  $j \in [n]$ , with  $\gamma \prec \delta \leq \beta_j$ , and such that  $(\alpha, \beta)$  can be separated from  $(\gamma, \delta)$  or  $(\gamma, \delta)$  can be separated from  $(\alpha, \beta)$ .

**Theorem 53.** *Let  $R$  be a subdirect product of  $\mathbb{A}_1, \dots, \mathbb{A}_n \in \mathcal{V}$  and choose  $\beta_j \in \text{Con}(\mathbb{A}_j)$  and a  $\beta_j$ -block  $B_j$  for each  $j \in [n]$ ; let  $R$  be chained with respect to  $\overline{\beta}, \overline{B}$  and  $R' = R \cap \overline{B}$ ,  $B'_j = \text{pr}_j R'$ ,  $j \in [n]$ . Let also  $i \in [n]$ , and  $\alpha \in \text{Con}(\mathbb{A}_i)$  be such that  $\alpha \prec \beta_i$ . Then if  $B'_i/\alpha$  contains a nontrivial as-component, then there exists an  $\alpha\beta_i$ -collapsing polynomial  $f$  for  $\overline{\beta}, \overline{B}$ . Moreover,  $f$  can be chosen to satisfy any one of the following conditions:*

- (d) *for any  $(\alpha, \beta_i)$ -subtrace  $\{a, b\} \subseteq \text{amax}(\text{pr}_i R')$  with  $b \in \text{as}(a)$  and any  $j \in [n]$ , the polynomial  $f$  can be chosen such that  $a/\alpha, b/\alpha \in f(\mathbb{A}_i)/\alpha$  and  $f(B_j) \cap \text{umax}(B'_j) \neq \emptyset$ ;*
- (e) *if  $\text{typ}(\alpha, \beta_i) = \mathbf{2}$ , for any  $\mathbf{a} \in \text{umax}(R')$  the polynomial  $f$  can be chosen such that  $f(\mathbf{a}) = \mathbf{a}$ ;*
- (f) *if  $\text{typ}(\alpha, \beta_i) = \mathbf{2}$ , a tuple  $\mathbf{a} \in R'$  is such that  $\mathbf{a} \in \text{umax}(R'')$ , where  $R'' = \{\mathbf{b} \in R \mid \mathbf{b}[i] \stackrel{\alpha}{\equiv} \mathbf{a}[i]\}$  and  $\{a, b\} \subseteq \text{amax}(\text{pr}_i R')$  is an  $(\alpha, \beta_i)$ -subtrace such that  $\mathbf{a}[i] = a$  and  $b \in \text{as}(a)$ , then the polynomial  $f$  can be chosen such that  $f(\mathbf{a}) = \mathbf{a}$  and  $b' \in f(\mathbb{A}_i)$  for some  $b' \stackrel{\alpha}{\equiv} b$ .*

*Proof.* First, we prove that an  $\alpha\beta_i$ -collapsing polynomial exists. Suppose  $i = 1$ , let  $C$  be a nontrivial as-component of  $B'_1/\alpha$ . Take  $a, b \in B'_1$  such that  $a/\alpha, b/\alpha \in C$  and  $a/\alpha \neq b/\alpha$ . Let  $f$  be a  $\overline{B}$ -preserving polynomial of  $R$  such that  $f(a/\alpha) = a/\alpha, f(b/\alpha) = b/\alpha$ . Such a polynomial exists, as the identity mapping satisfies these conditions. Let  $M(f)$  denote the set of triples  $(j, \gamma, \delta)$  such that  $j \in [n]$ ,  $\gamma, \delta \in \text{Con}(\mathbb{A}_j)$ ,  $\gamma \prec \delta \leq \beta_j$ , and  $f(\delta|_{B_j}) \subseteq \gamma$ . Choose  $f$  for which  $M(f)$  is maximal (under inclusion). Note that  $f$  can be replaced with its idempotent power, so it can be assumed idempotent. We show that  $f$  is  $\alpha\beta_1$ -collapsing.

Suppose there are  $j \in [n]$  and  $\gamma, \delta \in \text{Con}(\mathbb{A}_j)$  such that  $\gamma \prec \delta \leq \beta_j$ ,  $(\alpha, \beta_1), (\gamma, \delta)$  can be separated, and  $(j, \gamma, \delta) \notin M(f)$ . Then, since  $R$  is chained, there is a unary  $\overline{B}$ -preserving polynomial  $f_{j\gamma\delta}$  of  $R$  such that  $f_{j\gamma\delta}(a/\alpha) = a/\alpha, f_{j\gamma\delta}(b/\alpha) = b/\alpha$  (this implies  $f_{j\gamma\delta}(\beta_1) \not\subseteq \alpha$ ) and  $f_{j\gamma\delta}(\delta|_{B'_j}) \subseteq \gamma|_{B'_j}$ . Indeed, if  $(\alpha, \beta_1)$  can be separated from  $(\gamma, \delta)$ , by Lemma 37(3)  $(a/\alpha, b/\alpha) \in T_\alpha(a/\alpha, b/\alpha, \mathcal{U}(\gamma, \delta, \overline{B}))$ , and  $f_{j\gamma\delta}$  can be chosen to be the polynomial witnessing this. If  $(\gamma, \delta)$  can be separated from  $(\alpha, \beta_1)$ , then  $f_{j\gamma\delta}$  exists by Theorem 38. Let  $g$  be the idempotent power of  $f_{j\gamma\delta} \circ f$ . We have  $g(a/\alpha) = a/\alpha, g(b/\alpha) = b/\alpha, M(f) \subseteq M(g)$ , but  $g(\delta|_{B_j}) \subseteq \gamma$ , a contradiction with the choice of  $f$ .

Next we prove that any one of conditions (d)–(f) can be satisfied. For condition (d), that for any  $(\alpha, \beta_1)$ -subtrace  $\{a, b\} \subseteq \text{amax}(\text{pr}_i R')$  with  $b \in \text{as}(a)$  the polynomial  $f$  can be chosen such that  $a/\alpha, b/\alpha \in f(\mathbb{A}_i)/\alpha$  follows from what is proved above, since if  $\{a, b\}$  is an  $(\alpha, \beta_1)$ -subtrace such that  $a, b \in \text{amax}(\text{pr}_i R')$  and  $b \in \text{as}(a)$ , then  $a/\alpha, b/\alpha$  are members of a nontrivial as-component of  $B'_1/\alpha$  and the result follows from the chained condition and Lemma 37(3). It remains to show that, for any  $j \in [n]$ ,  $f$  can also be chosen to satisfy  $f(B_j) \cap \text{umax}(\text{pr}_j R') \neq \emptyset$ .

Let  $j = 2$  and  $c \in f(B'_2)$ . Let also  $C = \text{as}(a)/\alpha$ . Similar to the proof of Theorem 38 by  $Q^* \subseteq (\mathbb{A}_1/\alpha)^2 \times \mathbb{A}_2 \times R$  we denote the relation generated by  $\{(a/\alpha, b/\alpha, c, \mathbf{a})\} \cup \{(x/\alpha, x/\alpha, y, \mathbf{z}) \mid \mathbf{z} \in R, \mathbf{z}[1] = x, \mathbf{z}[2] = y\}$ , where  $\mathbf{a}$  is an arbitrary element from  $R'$ . Let  $Q = \text{pr}_{123}(Q^* \cap (B'_1/\alpha \times B'_1/\alpha \times B'_2 \times \overline{B}))$ . Observe that  $Q$  is exactly the set of triples  $(g(a)/\alpha, g(b)/\alpha, g(c))$  for  $\overline{B}$ -preserving unary polynomials  $g$  of  $R$ . By what has been proved,  $C^2 \subseteq \text{pr}_{12}Q$  and therefore  $(a/\alpha, b/\alpha)$  is an as-maximal element of  $\text{pr}_{12}Q$ . By the Maximality Lemma 18(5) there is  $d \in \text{umax}(\text{pr}_3 Q) = \text{umax}(B'_2)$  such that  $(a/\alpha, b/\alpha, d) \in Q$ . Thus, for any  $\overline{B}$ -preserving polynomial  $g$  such that  $g(a/\alpha) = a/\alpha, g(b/\alpha) = b/\alpha$ , and  $c \in g(B'_2)$ , there is a  $\overline{B}$ -preserving polynomial  $g_c$  with  $g_c(a/\alpha) = a/\alpha, g_c(b/\alpha) = b/\alpha$ , and  $g_c(c) \in \text{umax}(B'_2)$ . Let  $c_0 \in \text{umax}(B'_2)$  be such that there is a  $\overline{B}$ -preserving polynomial  $h$  with  $h(a/\alpha) = a/\alpha, h(b/\alpha) = b/\alpha$ , and  $c_0 \in h(B'_2)$ . Construct a sequence  $c_0, c_1, \dots$  by setting  $c_{\ell+1} = g_{c_\ell}(c_\ell)$ . Clearly, all the  $c_\ell$  are u-maximal in  $B'_2$ . There are  $r < s$  such that  $c_r = c_s$ . Then for the polynomial  $h = g_{c_{s-1}} \circ \dots \circ g_{c_r}$  it holds that  $h(c_r) = c_r$ . The idempotent power of  $h$  satisfies the desired conditions.

Now, suppose that  $\text{typ}(\alpha, \beta_1) = 2$ . We will use Lemma 52, so we need to identify some congruences of  $R$  related to  $\alpha\beta_1$ -collapsing polynomials. Consider congruences  $\alpha' = \alpha \times \beta_2 \times \dots \times \beta_n$  and  $\overline{\beta} = \beta_1 \times \beta_2 \times \dots \times \beta_n$ , and a maximal congruence  $\alpha^*$  of  $R$ ,  $\alpha' \leq \alpha^* \leq \overline{\beta}$ , such that  $\alpha^*|_{R'} = \alpha'|_{R'}$ . Let  $b_1^1, b_1^2 \in B'_1$  be such that  $b_1^1/\alpha \neq b_1^2/\alpha$ , and  $b_1^1, b_1^2$  belong to an as-component of  $B'_1$ . Such elements exist by Lemma 21(2). By the Maximality Lemma 18(3,5) these elements can be extended to  $\mathbf{b}_1, \mathbf{b}_2 \in \text{amax}(R')$  such that  $\mathbf{b}_1[1] = b_1^1, \mathbf{b}_2[1] = b_1^2$  and  $\mathbf{b}_1, \mathbf{b}_2$  belong to the same as-component of  $R'$ . Let  $\beta^* = \text{Cg}_R(\alpha^* \cup \{(\mathbf{b}_1, \mathbf{b}_2)\})$ . Since  $R$  is chained with respect to  $\overline{\beta}, \overline{B}$ , applying (Q1) for  $I = [n]$ , by Lemma 37(3) for any  $\mathbf{c}_1, \mathbf{c}_2 \in R'$  with  $\mathbf{c}_1[1] \not\stackrel{\alpha}{\sim} \mathbf{c}_2[1]$ , it holds  $\beta^* \leq \text{Cg}_R(\alpha^* \cup \{\mathbf{c}_1, \mathbf{c}_2\})$ . Therefore  $\alpha^* \prec \beta^*$ .

As  $f(\beta^*) \not\subseteq \alpha^*$  for the idempotent polynomial  $f$  found in the first part of the proof of Theorem 53 the set  $f(R)$  contains an  $(\alpha^*, \beta^*)$ -minimal set. Observe that  $\beta^*|_{R'} = \overline{\beta}|_{R'}$ . Indeed, as  $\text{typ}(\alpha, \beta_1) = 2$ ,  $B'_1/\alpha$  is a module, in particular,  $B'_1/\alpha$  is an as-component. Since  $R$  is chained, for any  $c/\alpha, d/\alpha \in B'_1/\alpha$  there is a  $\overline{B}$ -preserving polynomial  $g$  such that  $g(b_1^1/\alpha) = c/\alpha, g(b_1^2/\alpha) = d/\alpha$ , and so  $\mathbf{c} \stackrel{\beta^*}{\equiv} \mathbf{d}$  for any  $\mathbf{c}, \mathbf{d} \in R'$  such that  $\mathbf{c}[1] = c, \mathbf{d}[1] = d$ . Therefore, by Lemma 52(1) for

any  $\mathbf{a} \in \text{umax}(R')$  there exists a  $\overline{B}$ -preserving polynomial  $h$  such that  $hf(\mathbf{a}) = \mathbf{a}$  and  $hf(R)$  contains an  $(\alpha^*, \beta^*)$ -minimal set. The polynomial  $hf$  is as required for condition (e).

To prove condition (f) first notice that  $\text{umax}(R'') \subseteq \text{umax}(R')$ . Indeed, as  $\text{typ}(\alpha, \beta_1) = 2$ , every  $\alpha'$ -block, where  $\alpha'$  is as defined above, contains a u-maximal element from  $R'$  and Lemma 17 applies. Now, choose an  $\alpha\beta_1$ -collapsing polynomial  $f$  such that  $a, b' \in f(\mathbb{A}_1)$  for some  $(\alpha, \beta_1)$ -subtrace  $\{a, b'\}$ . Such a subtrace exists by part (d) and Lemma 52(1). Then, since  $\mathbf{a}$  is u-maximal in an  $\alpha^*$ -block, by Lemma 52(3) similar to the previous case  $f$  can be chosen such that  $f(\mathbf{a}) = \mathbf{a}$ .  $\square$

## 6 The Congruence Lemma

The main result of this section is the Congruence Lemma 56. We start with introducing three closure properties of algebras and their subdirect products.

Let  $\mathbb{C}$  be a subalgebra of  $\mathbb{A} \in \mathcal{V}$ . A subset  $B \subseteq \mathbb{C}$  is *as-closed* in  $\mathbb{C}$  if for any  $a, b \in \mathbb{C}$  with  $a \in \text{umax}(B)$  and  $a \sqsubseteq_{\mathbb{C}}^{as} b$ , it holds that  $b \in B$ . Similarly, the set  $B$  is *s-closed* in  $\mathbb{C}$  if for any  $a, b \in \mathbb{C}$  with  $a \in \text{umax}(B)$  and  $a \leq b$ , it holds that  $b \in B$ . Thus, an as-closed (s-closed) set is just a set of elements closed under thin semilattice and affine edges (respectively, thin semilattice edges). Note that the subalgebra  $\mathbb{C}$  is very important here, as we normally want to ‘contain’ as-closed (s-closed) sets within some subalgebra, and thin edges do not respect subalgebras.

Let  $R$  be a subdirect product of  $\mathbb{A}_1, \dots, \mathbb{A}_n \in \mathcal{V}$  and  $Q$  a subalgebra of  $R$ . By  $\text{Cg}(Q)$  we denote the congruence of  $R$  generated by pairs of elements from  $Q$ . That is,  $\text{Cg}(Q)$  is the smallest congruence such that  $Q$  is a subset of a  $\text{Cg}(Q)$ -block, denote it  $\text{Block}(Q)$ . The subalgebra  $Q$  is said to be *polynomially closed* in  $R$  if  $Q$  is as-closed in  $\text{Block}(Q)$ .

**Remark 54.** *Polynomially closed subalgebras of Mal'tsev algebras are blocks of congruences. In the general case the structure of polynomially closed subalgebras is more intricate. The intuition (although not entirely correct) is that if some block  $B$  of a congruence  $\beta$  contains several as-components, a polynomially closed subalgebra contains some of them and has empty intersection with the rest. However, since this is true only for factor sets, and we do not even consider non-as-maximal elements, the actual structure is more ‘fractal’.*

The next lemma follows from the definitions and Lemma 12, and the fact that congruences are invariant under polynomials.

**Lemma 55.** *Let  $\mathbb{A} \in \mathcal{V}$  and let  $R$  be a subdirect product of  $\mathbb{A}_1, \dots, \mathbb{A}_n \in \mathcal{V}$ .*

- (1)  *$R$  is polynomially closed in  $R$  and  $\mathbb{A}$  is as-closed and s-closed in  $\mathbb{A}$ .*

(2) For any congruence  $\beta \in \text{Con}(R)$  and a  $\beta$ -block  $Q$ , the subalgebra  $Q$  is polynomially closed in  $R$ .

(3) Let  $Q_1, Q_2$  be subalgebras of  $R$ ,  $Q_1, Q_2$  polynomially closed in  $R$ , and  $\text{umax}(Q_1) \cap \text{umax}(Q_2) \neq \emptyset$ . Then  $Q_1 \cap Q_2$  is polynomially closed in  $R$ .

In particular, let  $\beta \in \text{Con}(R)$  and  $T$  a  $\beta$ -block such that  $\text{umax}(Q_1) \cap \text{umax}(T) \neq \emptyset$ . Then  $Q_1 \cap T$  is polynomially closed in  $R$ .

Let  $\mathbb{B}$  be a subalgebra of  $\mathbb{A}$ , and let  $\mathbb{C}_1, \mathbb{C}_2$  be subalgebras of  $\mathbb{B}$  as-closed (s-closed) in  $\mathbb{B}$  and such that  $\text{umax}(\mathbb{C}_1) \cap \text{umax}(\mathbb{C}_2) \neq \emptyset$ . Then  $\mathbb{C}_1 \cap \mathbb{C}_2$  is as-closed (respectively, s-closed) in  $\mathbb{B}$ .

(4) Let  $\mathbb{B}$  be a subalgebra of  $\mathbb{A}$  and a subalgebra  $\mathbb{C} \subseteq \mathbb{B}$  as-closed (s-closed) in  $\mathbb{B}$ . Then for any  $\beta \in \text{Con}(\mathbb{A})$ , the algebra  $\mathbb{C}/\beta$  is as-closed (respectively, s-closed) in  $\mathbb{B}/\beta$ .

(5) Let  $R_i, i \in [k]$ , be a subdirect product of some algebras from  $\mathcal{V}$ , and  $Q_i$  polynomially closed in  $R_i, i \in [k]$ . Let  $R, Q$  be conjunctive-defined through  $R_1, \dots, R_k$  and  $Q_1, \dots, Q_k$ , respectively, by the same conjunctive formula  $\Phi$ ; that is,  $R = \Phi(R_1, \dots, R_k)$  and  $Q = \Phi(Q_1, \dots, Q_k)$ . In other words  $R$  consists of all tuples that are satisfying assignments of  $\Phi$  which uses the  $R_i$ 's as atoms, while  $Q$  is obtained in the same way only replacing the  $R_i$ 's with the  $Q_i$ 's. Suppose that for every atom  $Q_i(x_{j_1}, \dots, x_{j_\ell})$  it holds that  $\text{umax}(\text{pr}_{\{x_{j_1}, \dots, x_{j_\ell}\}} Q) \subseteq \text{umax}(Q_i)$ . Then  $Q$  is polynomially closed in  $R$ .

(6) Let  $R_i$  be a subdirect product of some algebras from  $\mathcal{V}$ ,  $Q_i$  subalgebra of  $R_i$ , and  $T_i$  a subalgebra of  $Q_i$  as-closed (s-closed) in  $Q_i, i \in [k]$ . Let  $R, Q$ , and  $T$  be pp-defined through  $R_1, \dots, R_k, Q_1, \dots, Q_k$ , and  $T_1, \dots, T_k$ , respectively, by the same pp-formula  $\exists \bar{x} \Phi$ ; that is,  $R = \exists \bar{x} \Phi(R_1, \dots, R_k)$ ,  $Q = \exists \bar{x} \Phi(Q_1, \dots, Q_k)$ , and  $T = \exists \bar{x} \Phi(T_1, \dots, T_k)$ . Let also  $Q' = \Phi(Q_1, \dots, Q_k)$  and  $T' = \Phi(T_1, \dots, T_k)$  and, and suppose that for every atom  $T_i(x_{j_1}, \dots, x_{j_\ell})$  it holds that  $\text{umax}(\text{pr}_{\{x_{j_1}, \dots, x_{j_\ell}\}} T') \subseteq \text{umax}(T_i)$ . Then  $T$  is as-closed (respectively, s-closed) in  $Q$ .

*Proof.* (1) is straightforward. (2) holds because if  $Q$  is a congruence block of  $\beta$ , then clearly  $\text{Block}(Q) = Q$ .

For (3) it suffices to observe that, as  $\text{umax}(Q_1) \cap \text{umax}(Q_2) \neq \emptyset$ , by Lemma 17 it holds that  $\text{umax}(Q_1 \cap Q_2) \subseteq \text{umax}(Q_1) \cap \text{umax}(Q_2)$ . We have  $\text{Cg}(Q_1 \cap Q_2) \leq \text{Cg}(Q_1) \wedge \text{Cg}(Q_2)$  and  $\text{Block}(Q_1 \cap Q_2) \subseteq \text{Block}(Q_1) \cap \text{Block}(Q_2)$ . Therefore for any  $\mathbf{a} \in \text{umax}(Q_1 \cap Q_2)$  and  $\mathbf{b} \in \text{Block}(Q_1 \cap Q_2)$  such that  $\mathbf{a} \sqsubseteq^{\text{as}} \mathbf{b}$  in  $\text{Block}(Q_1 \cap Q_2)$ , it holds that  $\mathbf{b} \in Q_1$  and  $\mathbf{b} \in Q_2$ .



The statement about as-closed (s-closed) subalgebras is straightforward because by Lemma 17  $\text{umax}(\mathbb{C}_1 \cap \mathbb{C}_2) \subseteq \text{umax}(\mathbb{C}_1) \cap \text{umax}(\mathbb{C}_2)$ .

Part (4) follows from Lemmas 12 and 17.

(5) Suppose that  $\Phi$  involves  $n$  variables  $x_1, \dots, x_n$ . Take  $\mathbf{a} \in \text{umax}(Q)$  and  $\mathbf{b} \in \text{Block}(Q)$  such that  $\mathbf{a} \sqsubseteq^{as} \mathbf{b}$  in  $\text{Block}(Q)$ . Consider some atoms  $R_i, Q_i$  of  $\Phi$ ; let  $X$  be the set of variables involved in  $R_i, Q_i$ , and set  $\mathbf{a}_i = \text{pr}_X \mathbf{a}, \mathbf{b}_i = \text{pr}_X \mathbf{b}$ . By the Maximality Lemma 18(4,6)  $\mathbf{a}_i \in \text{umax}(Q_i)$  and  $\mathbf{a}_i \sqsubseteq_{R_i}^{as} \mathbf{b}_i$ . As is easily seen,  $\text{pr}_X \text{Block}(Q) \subseteq \text{Block}(Q_i)$ , and since  $Q_i$  is polynomially closed,  $\mathbf{b}_i \in Q_i$ . As this holds for every  $i \in [k]$ ,  $\mathbf{b} \in Q$ .

(6) Suppose again that  $\Phi$  involves  $n$  variables  $x_1, \dots, x_n$  and  $x_{m+1}, \dots, x_n$  are quantified. Take  $\mathbf{a} \in \text{umax}(T)$ , and  $\mathbf{b} \in Q$  such that  $\mathbf{a} \sqsubseteq^{as} \mathbf{b}$ . By the assumptions of the lemma and the Maximality Lemma 18(3,5) there are  $\mathbf{a}' \in \text{umax}(T')$  and  $\mathbf{b}' \in Q'$  such that  $\mathbf{a} = \text{pr}_{[m]} \mathbf{a}', \mathbf{b} = \text{pr}_{[m]} \mathbf{b}'$ , and  $\mathbf{a}' \sqsubseteq_{Q'}^{as} \mathbf{b}'$ . Consider an atom  $Q_i$  (respectively,  $T_i$ ) of  $\Phi$ ; let  $X$  be the set of variables involved in  $Q_i, T_i$ , and set  $\mathbf{a}_i = \text{pr}_X \mathbf{a}', \mathbf{b}_i = \text{pr}_X \mathbf{b}'$ . By the Maximality Lemma 18(4,6)  $\mathbf{a}_i \in \text{umax}(T_i)$  and  $\mathbf{a}_i \sqsubseteq_{Q_i}^{as} \mathbf{b}_i$ . Since  $T_i$  is as-closed,  $\mathbf{b}_i \in T_i$ . As this holds for every  $i \in [k]$ ,  $\mathbf{b}' \in T'$ , and so  $\mathbf{b} \in T$ . For s-closeness the proof is identical replacing  $\sqsubseteq^{as}$  with  $\leq$ .  $\square$

To explain what Lemma 56 (the Congruence Lemma) amounts to saying consider this: let  $Q \subseteq \mathbb{A}' \times \mathbb{B}'$  be a subdirect product and the link congruence of  $\mathbb{A}'$  is the equality relation. Then, clearly,  $Q$  is the graph of a mapping  $\sigma : \mathbb{B}' \rightarrow \mathbb{A}'$ , and the kernel of this mapping is the link congruence  $\eta$  of  $\mathbb{B}'$  with respect to  $Q$ . Suppose now that  $Q$  is a subalgebra of  $R$ , a subdirect product of  $\mathbb{A} \times \mathbb{B}$  such that  $\mathbb{A}'$  is a subalgebra of  $\mathbb{A}$  and  $\mathbb{B}'$  is a subalgebra of  $\mathbb{B}$ . Then the restriction of the link congruence of  $\mathbb{A}$  with respect to  $R$  to  $\mathbb{A}'$  does not have to be the equality relation, and similarly the restriction of the link congruence of  $\mathbb{B}$  to  $\mathbb{B}'$  does not have to be  $\eta$ , even if  $Q = R \cap (\mathbb{A}' \times \mathbb{B}')$ . Most importantly, the restriction of  $\text{Cg}_{\mathbb{B}}(\eta)$ , the congruence of  $\mathbb{B}$  generated by  $\eta$ , to  $\mathbb{B}'$  does not have to be  $\eta$ . The Congruence Lemma 56 shows, however, that this is exactly what happens when  $R, Q$  and  $\mathbb{A}', \mathbb{B}'$  satisfy some additional conditions, such as being chained and polynomially closed.

**Lemma 56** (The Congruence Lemma). *Let  $R$  be a subdirect product of  $\mathbb{A}_1, \dots, \mathbb{A}_n \in \mathcal{V}$ ,  $\beta_i$  a congruence of  $\mathbb{A}_i$  and let  $B_i$  be a  $\beta_i$ -block,  $i \in [n]$ . Also, let  $R$  be chained with respect to  $\bar{\beta}, \bar{B}$  and  $R' = R \cap \bar{B}$ ,  $B'_i = \text{pr}_i R'$ . Let  $\alpha \in \text{Con}(\mathbb{A}_1)$  be such that  $\alpha \prec \beta_1$ , let  $I \subseteq [n] - \{1\}$ ,  $I' = I \cup \{1\}$ ,  $J = [n] - I'$ , and let  $Q$  be a subalgebra of  $R'$  polynomially closed in  $R$  and such that  $E_1 = \text{pr}_1 Q$  contains an as-component  $C$  of  $B'_1$  and  $Q \cap \text{umax}(R') \neq \emptyset$ . We consider  $R, R'$ , and  $Q$  as subdirect products of  $\text{pr}_1 R \times \text{pr}_I R \times \text{pr}_J R$ ,  $\text{pr}_1 R' \times \text{pr}_I R' \times \text{pr}_J R'$ , and  $\text{pr}_1 Q \times \text{pr}_I Q \times \text{pr}_J Q$ , respectively. Let  $R'/\alpha = R'/\alpha \times \underline{0}_2 \times \dots \times \underline{0}_n$ ,  $Q/\alpha = Q/\alpha \times \underline{0}_2 \times \dots \times \underline{0}_n$ , and let  $\text{lk}'_1, \text{lk}'_2$  and  $\text{lk}_1^Q, \text{lk}_2^Q$  be the link congruences of  $\text{pr}_{I'} R'/\alpha$  and  $\text{pr}_{I'} Q/\alpha$ , respectively, and*



$E_2 = \text{pr}_I Q$ . Finally, let  $\eta \leq \bar{\beta}_I = \prod_{i \in I} \beta_i$  be a maximal (under inclusion) congruence of  $\text{pr}_I R$  with  $\eta|_{\text{umax}(E_2^C)} \subseteq \text{lk}_2^Q$ , and  $\gamma = \text{Cg}_{\text{pr}_I R}(\text{umax}(E_2^C)) \vee \eta$ , where  $E_2^C = Q[C/\alpha] = \{\text{pr}_I \mathbf{a} \mid \mathbf{a} \in Q, \mathbf{a}[1]/\alpha \in C/\alpha\}$ . Then either

- (1)  $C/\alpha \times \text{umax}(E_2^C) \subseteq \text{pr}_{I'} Q/\alpha$ , or
- (2)  $\eta \prec \gamma \leq \beta_I$  and the intervals  $(\alpha, \beta_1)$  and  $(\eta, \gamma)$  cannot be separated.

Moreover, in case (2)  $\text{pr}_{I'} Q/\alpha$  is the graph of a mapping  $\varphi : E_2 \rightarrow E_1/\alpha$  such that the kernel  $\text{lk}_2^Q$  of  $\varphi$  is the restriction of  $\eta$  on  $E_2$ , and for any  $\bar{B}$ -preserving polynomial  $f$  such that  $f(\beta_1|_{B_1}) \not\subseteq \alpha$  it holds that  $f(\gamma|_{\bar{B}_I}) \not\subseteq \eta$ .<sup>5</sup>

We prove the lemma in three steps. First, we assume that  $Q = R'$  (Lemma 57), then we assume that  $Q$  is a congruence block of  $R$  (Lemma 58), and finally we will use polynomial closeness to prove Lemma 56 itself. In all of these intermediate lemmas we use the notation from Lemma 56 and assume that the conditions of Lemma 56 hold. Throughout the proof we will assume that  $R$  is viewed as a ternary relation, a subdirect product of  $\mathbb{A}_1$ ,  $\text{pr}_I R$ , and  $\text{pr}_J R$  ( $R'$  and  $Q$  will also be assumed ternary). We will also assume that  $\alpha = \underline{0}_1$  everywhere except the first part of the proof of Lemma 56 itself. The first assumption does not restrict generality, because by Lemma 36(1) and because it does not affect the congruences of  $\text{pr}_I R$  and  $\text{Cg}(Q)$ , relations  $R$  and  $Q$  viewed as ternary relations satisfy the conditions of the lemma, and the conclusion of the lemma is true for  $R, Q$  regardless of a partitioning of the subdirect product. The second assumption requires caution, because while the chaining condition still holds by Lemma 36(2),  $Q/\alpha$  is not necessarily polynomially closed. However, in the proof we will use the polynomial closeness of  $Q$  only once, and can assume  $\alpha = \underline{0}_1$  elsewhere. So, from now on  $n = 3$ ,  $I = \{2\}$ ,  $J = \{3\}$ , and  $\eta = \underline{0}_2$  unless stated otherwise.

**Lemma 57.** *If  $Q = R'$ , then the conclusion of Lemma 56 holds.*

*Proof.* Let  $R'' = \text{pr}_{12} R'$ . Since  $R$  is chained, if  $\text{lk}_1'$  is not the equality relation or  $|C| = 1$ , by Lemma 37(3)  $C^2 \subseteq \text{lk}_1'$ . In this case  $C \times \text{umax}(R''[C]) \subseteq R''$  by Proposition 27 and we obtain option (1) of Lemma 56 for  $R'$ . So, suppose that  $\text{lk}_1'$  is the equality relation and  $|C| > 1$ . Then  $R''$  is the graph of a mapping  $\varphi : B_2' \rightarrow B_1'$ , note that  $\text{lk}_2'$  is the kernel of  $\varphi$ . Let  $\gamma$  denote the congruence constructed in Lemma 56, that is,  $\gamma = \text{Cg}_{\mathbb{A}_2}(\text{umax}(B_2^C)) \vee \eta$ , where  $B_2^C = R''[C]$ . We prove that  $\text{lk}_2'|_{\text{umax}(B_2')} = \eta|_{\text{umax}(B_2')}$ , that  $\eta \prec \gamma$ , and that  $(\underline{0}_1, \beta_1), (\eta, \gamma)$  cannot be separated. This clearly implies option (2) of Lemma 56 for  $R'$ .

<sup>5</sup>This property is somewhat stronger than non-separability. The non-separability of  $(\alpha, \beta_1)$  and  $(\eta, \gamma)$  only implies that  $f(\gamma) \not\subseteq \eta$  without any restrictions on  $f(\gamma|_{\bar{B}_I})$ .

CLAIM 1.  $\text{lk}'_2|_{\text{umax}(B'_2)} = \eta|_{\text{umax}(B'_2)}$

Suppose that  $\text{lk}'_2|_{\text{umax}(B'_2)} \neq \eta|_{\text{umax}(B'_2)}$ . We show that in this case there is a congruence  $\theta$  of  $\mathbb{A}_2$  that is strictly greater than  $\eta$  and still satisfies the condition  $\theta|_{\text{umax}(B'_2)} \subseteq \text{lk}'_2|_{\text{umax}(B'_2)}$ . There are  $c, d \in \text{umax}(B'_2)$ ,  $(c, d) \notin \eta$  and  $a \in B'_1$  such that  $(a, c), (a, d) \in R''$  (as  $(c, d) \in \text{lk}'_2$  and  $\text{lk}'_2$  is the kernel of  $\varphi$ ). Let  $\theta = \text{Cg}_{\mathbb{A}_2}(\{(c, d)\}) \vee \eta$ . We claim that  $\theta|_{\text{umax}(B'_2)} \subseteq \text{lk}'_2|_{\text{umax}(B'_2)}$ , and, as  $\theta$  is strictly greater than  $\eta$ , it contradicts the choice of  $\eta$ . Firstly, notice that if  $e, e'$  are u-maximal in the same  $\theta$ -block then  $(e, e') \in \text{lk}'_2$ . Indeed, since  $R$  is chained, for any  $\theta$ -block  $D \subseteq B_2$  and any  $e, e' \in \text{umax}(D \cap B'_2)$  there are  $e = e_1, \dots, e_{k+1} = e' \in B'_2$ ,  $e'_1, \dots, e'_k \in B'_2$ ,  $e_i \stackrel{\eta}{\equiv} e'_i$  for  $i \in [k]$ , and  $\overline{B}$ -preserving polynomials  $f_1, \dots, f_k$  such that  $f_i(\{c, d\}) = \{e'_i, e_{i+1}\}$ . As  $(a, c), (a, d) \in R''$ , for every  $i \in [k-1]$ ,  $(e_i, e_{i+1}) \in \text{lk}'_2$ , and therefore  $(e, e') \in \text{lk}'_2$ . Note that we are not done yet, because there may be elements from  $\text{umax}(B'_2)$  that are not u-maximal in any  $\theta$ -block.

Suppose now that  $(e, e') \in \theta|_{B'_2} - \text{lk}'_2$ . Then  $(a, e), (b, e') \in R''$  for some  $a, b \in B'_1$  and  $a \neq b$ . Since  $R$  is chained with respect to  $\overline{\beta}, \overline{B}$ , and by Lemma 37(3) for any  $a', b' \in C$  there is a  $\overline{B}$ -preserving polynomial  $f$  such that  $f(a) = a', f(b) = b'$ . On the other hand,  $f(e) \stackrel{\theta}{\equiv} f(e')$ . Let  $R^\dagger$  be the algebra generated by  $(R'' \cap (C \times B'_2))/\underline{0}_1 \times \theta$  in  $B_1 \times B_2/\theta$ . By what is proved  $R^\dagger$  is linked and by Proposition 27 applied to  $R^\dagger$  there exists a  $\theta$ -block  $D$  such that for any  $a' \in C$  there is a  $c' \in D$  with  $(a', c') \in R''$ . By the Maximality Lemma 18(3)  $c'$  can be chosen to be u-maximal in  $D$ . Together with what is proved above this contradicts the assumption that  $\text{lk}'_1$  is the equality relation.

CLAIM 2. For all  $\theta \in \text{Con}(\mathbb{A}_2)$  with  $\eta \prec \theta \leq \beta_2$  the intervals  $(\underline{0}_1, \beta_1), (\eta, \theta)$  cannot be separated, and for any  $\overline{B}$ -preserving polynomial  $f$  such that  $f(\beta_1|_{B_1}) \not\subseteq \alpha$  it also holds that  $f(\theta|_{B_2}) \not\subseteq \eta$ .

Suppose that for some  $\theta \in \text{Con}(\mathbb{A}_2)$  with  $\eta \prec \theta \leq \beta_2$  Claim 2 is not true.

Let  $c, d \in \text{umax}(R''[C])$  with  $(c, d) \in \theta - \eta$  and  $a, b \in B'_1$  such that  $(a, c), (b, d) \in R''$ . We can assume that  $a, b \in C$ . By Corollary 31  $\{a, b\}$  is a  $(\underline{0}_1, \beta_1)$ -subtrace. Since  $R$  is chained, if  $(\underline{0}_1, \beta_1), (\eta, \theta)$  cannot be separated, by Theorem 53(d) there is a  $\overline{B}$ -preserving polynomial  $f$  such that  $f(a) = a, f(b) = b$ , and  $f(\theta|_{B'_2}) \subseteq \eta$ . If there is a  $\overline{B}$ -preserving polynomial  $g$  such that  $g(\beta_1|_{B_1}) \not\subseteq \alpha$  but  $g(\theta|_{B_2}) \subseteq \eta$ , we obtain  $g(a) = a', g(b) = b'$ , and then by chaining and Lemma 37(3) there is a  $\overline{B}$ -preserving polynomial  $g'$  with  $g'(a') = a, g'(b') = b$ . Then we set  $f = g \circ g'$ . Apply  $f$  to the pairs  $(a, c), (b, d)$  to obtain  $(a, f(c)), (b, f(d)) \in R''$ . Since  $f(\theta|_{B'_2}) \subseteq \eta|_{B'_2} \subseteq \text{lk}'_2$ , we obtain  $a \stackrel{\text{lk}'_1}{\equiv} b$ , a contradiction with the assumption that  $(c, d) \notin \text{lk}'_2$ . Claim 2 is proved.

Now we use  $\text{lk}'_2|_{\text{umax}(B'_2)} = \eta|_{\text{umax}(B'_2)}$  to show that  $\eta \prec \gamma$ . Clearly,  $\eta \leq \gamma$  and, by the choice of  $\eta, \gamma$ ,  $\eta \neq \gamma$ . Again, by the choice of  $\eta$  for any  $\theta \in \text{Con}(\mathbb{A}_2)$  with  $\eta < \theta \leq \beta_2$  there is a pair  $(c, d) \in \theta|_{\text{umax}(B'_2)} - \eta$ . Let  $(a, c), (b, d) \in R''$  for some  $a, b \in B'_1$ . Note that  $a \neq b$ , because  $(c, d) \notin \text{lk}'_2$ . We argue similarly to the proof of Claim 1. Since  $R$  is chained with respect to  $\bar{\beta}, \bar{B}$  and by Lemma 37(3), for any  $a', b' \in C$  there is a  $\bar{B}$ -preserving polynomial  $f$  such that  $f(a) = a', f(b) = b'$ . Let  $R^\dagger$  be the algebra generated by  $R'' \cap (C \times B'_2))/_{\underline{0}_1 \times \theta}$  in  $B_1 \times B_2/\theta$ . By what is proved  $R^\dagger$  is linked and by Proposition 27  $\text{umax}(R''[C])$  is a subset of a  $\text{lk}'_2 \vee \theta|_{B'_2}$ -block. Since  $\text{lk}'_2|_{\text{umax}(B'_2)} = \eta|_{\text{umax}(B'_2)} \subseteq \theta$ , it follows that  $\text{umax}(R''[C])$  is contained in a  $\theta$ -block, and therefore  $\gamma \subseteq \theta$ .  $\square$

Next, we assume that  $Q$  is a congruence block. Note that in this case polynomial closeness is trivial and the remaining conditions of Lemma 56 are assumed to be true.

**Lemma 58.** *Let in the notation of Lemma 56  $Q \subseteq R'$  be a block of  $\xi \in \text{Con}(R)$  with  $\xi \leq \bar{\beta}$ ,  $Q \in \text{umax}(R'/\xi)$ . Then the conclusion of Lemma 56 holds.*

*Proof.* By Lemma 57 Lemma 56 holds for  $R'$  in place of  $Q$ . Suppose first that option (2) of the lemma holds for  $R'$ . Then  $R'' = \text{pr}_{12}R'$ , and therefore  $Q'' = \text{pr}_{12}Q$ , as well, are the graphs of mappings, and we only need to verify that the congruences  $\eta, \gamma$  satisfy the desired conditions. Let  $\eta'$  be a maximal congruence such that  $\eta'|_{\text{umax}(B'_2)} \subseteq \text{lk}'_2$  and  $\eta' \leq \beta_2$ . Let  $B_2^C = \text{umax}(R''[C])$  and  $\gamma' = \text{Cg}_{\mathbb{A}_2}(\text{umax}(B_2^C)) \vee \eta'$ . As  $\text{umax}(E_2) \subseteq \text{umax}(B'_2)$ ,  $\text{umax}(E_2^C) \subseteq \text{umax}(B_2^C)$ , and  $\text{lk}_2^Q = \text{lk}'_2 \cap E_2^2$ , we have  $\eta'|_{\text{umax}(E_2^C)} \subseteq \text{lk}_2^Q$ , and therefore  $\eta'$  can be chosen such that  $\eta' \leq \eta$ . Since option (2) holds for  $R'$ ,  $\eta' \prec \gamma'$ . Let  $\theta \in \text{Con}(\mathbb{A}_2)$  be such that  $\eta \prec \theta \leq \gamma$ . As  $\eta' \prec \gamma'$ , it also holds that  $\gamma' = \text{Cg}_{\mathbb{A}_2}((c, d)) \vee \eta'$  for any  $(c, d) \in (\text{umax}(E_2^C))^2 \cap (\theta - \eta)$  (recall that  $\text{umax}(E_2^C) \subseteq \text{umax}(B_2^C)$  and therefore  $(c, d) \in \gamma'$ ). Therefore  $\gamma' \leq \theta$ . By the definition of  $\gamma'$  it holds that  $\text{umax}(B_2^C)$  is in a single  $\theta$ -block, hence,  $\gamma \leq \theta$ , implying  $\eta \prec \gamma$ . Also,  $\gamma' \wedge \eta = \eta'$ , because  $\eta' \leq \gamma' \wedge \eta$  and  $\eta' \prec \gamma'$ , and  $\gamma' \vee \eta = \gamma$ , because  $\text{umax}(E_2^C) \subseteq \text{umax}(B_2^C)$  and  $\gamma' \leq \theta$  for any  $\theta$  with  $\eta \prec \theta \leq \gamma$ , as observed above. Therefore, the intervals  $(\eta', \gamma')$  and  $(\eta, \gamma)$  are perspective. By Lemma 44 these intervals cannot be separated. Since  $(\underline{0}_1, \beta_1)$  and  $(\eta', \gamma')$  cannot be separated, it implies that  $(\underline{0}_1, \beta_1)$  and  $(\eta, \gamma)$  cannot be separated as well.

So, suppose that  $C \times \text{umax}(R''[C]) \subseteq R''$ . If  $\eta = \beta_2$ , we are done. Indeed, in this case  $\text{lk}_2^Q|_{\text{umax}(E_2^C)} = (\text{umax}(E_2^C))^2$ , and, since  $C \subseteq E_1$  and  $C$  is an as-component of  $E_1$ , we have  $C \times \text{umax}(E_2^C) \subseteq Q''$  by Proposition 27. Also, if, more generally,  $C$  is a subset of a  $\text{lk}_1^Q$ -block, then we are done for the same reason. Thus, assume  $\eta \neq \beta_2$  and  $C$  is not a subset of a  $\text{lk}_1^Q$ -block.

CLAIM 1. For any  $\theta \in \text{Con}(\mathbb{A}_2)$ ,  $\eta \prec \theta \leq \beta_2$  the intervals  $(\underline{0}_1, \beta_1)$  and  $(\eta, \theta)$  cannot be separated, and for any  $\overline{B}$ -preserving polynomial  $f$  such that  $f(\beta_1|_{B_1}) \not\subseteq \alpha$  it also holds that  $f(\theta|_{B_2}) \not\subseteq \eta$ .

Suppose that for some  $\theta \in \text{Con}(\mathbb{A}_2)$  with  $\eta \prec \theta \leq \beta_2$  Claim 1 is not true. We prove that in this case  $C \times \text{umax}(E_2^C) \subseteq Q''$ , a contradiction with our assumptions.

By the choice of  $\eta$  we have  $\theta|_{\text{umax}(E_2^C)} \not\subseteq \text{lk}_2^Q$ . Therefore, there are  $c, d \in \text{umax}(E_2^C)$  with  $(c, d) \in \theta - \text{lk}_2^Q$ . Let  $a, b \in E_1$  be such that  $a, b \in C$  and  $(a, c, e), (b, d, e') \in Q$  for some  $e, e' \in B_3'$ . By Corollary 31  $\{a, b\}$  is a  $(\underline{0}_1, \beta_1)$ -subtrace. If  $(\underline{0}_1, \beta_1)$  can be separated from  $(\eta, \theta)$ , by Theorem 53(d), there is a  $\overline{B}$ -preserving idempotent polynomial  $g$  of  $R$  with  $g(a) = a, g(b) = b, g(c) \stackrel{\eta}{=} g(d)$ . If there is a  $\overline{B}$ -preserving polynomial  $f$  such that  $g(\beta_1|_{B_1}) \not\subseteq \alpha$  but  $g(\theta|_{B_2}) \subseteq \eta$ , we obtain  $f(a) = a', f(b) = b'$ , and then by chaining and Lemma 37(3) there is a  $\overline{B}$ -preserving polynomial  $f'$  with  $f'(a') = a, f'(b') = b$ . Then we set  $g = f \circ f'$ . Although this looks like an indication that  $\text{lk}_1^Q \neq \underline{0}_1$ , the tuples  $(a, g(c), g(e)), (b, g(d), g(e'))$  do not necessarily belong to  $Q$ , and we have to make further steps.

As  $R$  is chained, for any  $a', b' \in C$  there is a  $\overline{B}$ -preserving polynomial  $h$  with  $h(a) = a', h(b) = b'$ . Note also  $(hg(a), hg(c), hg(e)) \stackrel{\xi}{=} (hg(b), hg(d), hg(e'))$ . Consider the subalgebra  $S$  of the product  $R'/\xi \times B_2'/\eta \times B_1'$  given by

$$S = \{((x, y, z)/\xi, y/\eta, x) \mid (x, y, z) \in R'\},$$

as a subdirect product of  $\text{pr}_{12}S$  and  $B_1'$ . By what is shown above, for any  $a', b' \in C$  there is  $(P, c') \in \text{pr}_{12}S$  such that  $(P, c', a'), (P, c', b') \in S$ . Therefore  $C$  is a subset of a block of the link congruence of  $S$ , and by Proposition 27 for any  $(P, c') \in \text{umax}(S^{-1}(C))$  it holds that  $\{(P, c')\} \times C \subseteq S$ . This means that for any  $a', b' \in C$  there are  $c', d' \in E_2$  and  $e_1, e_1' \in B_3'$  such that  $(a', c', e_1), (b', d', e_1') \in Q$  and  $c' \stackrel{\eta}{=} d'$ , a contradiction with the assumption that  $C$  is not contained in a class of  $\text{lk}_1^Q$ . Claim 1 is proved.

Since  $C \times \text{umax}(R''[C]) \subseteq R''$  and  $(\underline{0}_1, \beta_1), (\eta, \theta)$  cannot be separated, by Lemmas 48, 45  $\text{typ}(\underline{0}_1, \beta_1) = \text{typ}(\eta, \theta) = \mathbf{2}$ , for any  $\eta \prec \theta \leq \beta_2$ . Therefore, the  $\beta_1$ -block  $B_1$  is a module, and so is  $B_1'$ . Hence,  $B_1' = C = E_1$ . Also, this implies that  $E_2^C/\text{lk}_2^Q = E_2/\text{lk}_2^Q$ , and every  $\theta$ -block is a module modulo  $\eta$ .

CLAIM 2.  $\text{lk}_2^Q = \eta|_{E_2}$ .

Suppose  $\text{lk}_2^Q \neq \eta|_{E_2}$  and  $(c_1, d_1) \in \text{lk}_2^Q - \eta$ . The elements  $c_1, d_1$  can be chosen such that there is  $a_0 \in E_1$  and  $e_1, e_1' \in B_3'$  with  $(a_0, c_1, e_1), (a_0, d_1, e_1') \in Q$ . Let  $\varrho = \text{Cg}_{\mathbb{A}_2}((c_1, d_1)) \vee \eta$ .

By the choice of  $\eta$ ,  $\varrho|_{\text{umax}(E_2)} \not\subseteq \text{lk}_2^Q$ . Let  $D$  be a  $\varrho$ -block that intersects more than one  $\text{lk}_2^Q$ -blocks on  $\text{umax}(E_2)$ . Since  $R$  is chained, for any  $\bar{c}_2, \bar{d}_2 \in \text{umax}(D/\eta)$  there are  $\bar{c}_2 = \bar{c}_1^*, \dots, \bar{c}_k^* = \bar{d}_2$  in  $D/\eta$  and polynomials  $f_1, \dots, f_{k-1}$  with  $\{f_i(c_1), f_i(d_1)\} = \{\bar{c}_i^*, \bar{c}_{i+1}^*\}$ . Note that the triples  $(f_i(a_0), f_i(c_1), f_i(e_1))$ ,  $(f_i(a_0), f_i(d_1), f_i(e_1'))$  are related in  $\xi$ . Fix  $\bar{c}_2 \in \text{umax}(D/\eta) \subseteq \text{umax}(B'_2/\eta)$ .

Let  $\tau = \xi \wedge (\underline{0}_1 \times \eta \times \underline{1}_3)$ . We consider  $R'/\tau$  as a subdirect product  $R^\dagger$  of  $R'/\xi \times B'_1 \times B'_2/\eta$ . Let  $\text{lk}_3^\eta$  be the link congruence of the third coordinate of  $R^\dagger$ . By what is proved above  $\text{umax}(D/\eta)$  is contained in a  $\text{lk}_3^\eta$ -block. By Proposition 27 for every as-component  $C^\dagger$  of  $D/\eta$  there is  $b_0 \in B'_1$  and a  $\xi$ -block  $T$  such that for any  $\eta$ -block  $D' \in C^\dagger$  there is  $d \in D'$  and  $e'_2 \in B'_3$  with  $(b_0, d, e'_2) \in T$ . Choose an as-maximal  $\bar{c}_2 \in C/\eta$ ,  $c_2 \in \bar{c}_2$ ,  $b_0 \in B'_1$ , and  $e_2 \in B'_3$  such that  $(b_0, c_2, e_2) \in T$ .

It seems that  $b_0$  is related to multiple  $\eta$ -blocks across  $\text{lk}_2^Q$  in contradiction with the choice of  $\eta$ . However, two obstacles remain. First, while  $D$  intersects multiple  $\text{lk}_2^Q$ -blocks, the connection between  $C^\dagger$  and  $E_2$  is unknown, yielding no contradiction. It may even be the case that  $|C^\dagger| = 1$ . Second,  $T$  is not necessarily equal to  $Q$ , again avoiding a contradiction.

To overcome these obstacles, choose  $\theta \in \text{Con}(\mathbb{A}_2)$  such that  $\eta \prec \theta \leq \varrho$ . By Claim 1  $(\underline{0}_1, \beta_1)$  and  $(\eta, \theta)$  cannot be separated and by Lemma 45  $\text{typ}(\eta, \theta) = \mathbf{2}$ . Moreover, since  $B'_1 \times \text{umax}(B'_2) \subseteq R''$ , by Lemma 48,  $(\eta : \theta) \geq \beta_2$ . By the choice of  $\eta$  there is a  $\theta$ -block  $F$  and  $c_3, d_3 \in F \cap \text{umax}(E_2)$  with  $(c_3, d_3) \notin \text{lk}_2^Q$ , that is,  $|(F \cap \text{umax}(E_2))/\eta| > 1$ .

There exist  $b \in B'_1$  and  $e_3 \in B'_3$  such that  $(b, c_3, e_3) \in \text{umax}(Q) \subseteq \text{umax}(R')$ . Take an asm-path  $(b_0, c_2, e_2) = \mathbf{a}_1, \dots, \mathbf{a}_\ell = (b, c_3, e_3)$ , where  $\mathbf{a}_i = (b_i, c_i^\dagger, e_i^\dagger)$  in  $R'$ . For  $i \in [\ell - 1]$  let  $f_i$  be a polynomial of  $R$  given in Lemma 49 and defined as follows: If  $\mathbf{a}_i \mathbf{a}_{i+1}$  is semilattice (majority, affine), then

$$f_i(x) = x \cdot \mathbf{a}_{i+1}, \quad f_i(x) = t_{\mathbf{a}_i \mathbf{a}_{i+1}}(x, \mathbf{a}_{i+1}), \quad f_i(x) = h_{\mathbf{a}_i \mathbf{a}_{i+1}}(x, \mathbf{a}_i, \mathbf{a}_{i+1}),$$

respectively. As is easily seen,  $f_i(\mathbf{a}_i) = \mathbf{a}_{i+1}$ ,  $i \in [\ell - 1]$ . Let  $G \subseteq D$  be the  $\theta$ -block containing  $c_2$ ; note that  $G/\eta \subseteq C^\dagger$ . Composing all the polynomials  $f_i$  by Lemma 49 we obtain a polynomial  $f$  such that  $f(b_0, c_2, e_2) = (b, c_3, e_3)$  and, as  $(\eta : \theta) \geq \beta_2$ ,  $f$  maps  $G/\eta$  injectively into  $F/\eta$ . Since  $\{b_0\} \times G/\eta \subseteq (\text{pr}_{12}T)/\underline{0}_1 \times \eta$ , we obtain  $\{b\} \times f(G/\eta) \subseteq Q''/\underline{0}_1 \times \eta$ . It remains to show that  $f(G/\eta)$  is sufficiently large.

As  $\bar{c}_2 \in \text{umax}(B'_2/\eta)$  (recall that  $D$  contains u-maximal elements of  $B'_2$  and therefore every its u-maximal element belongs to  $\text{umax}(B'_2)$ ), there is  $c'_2 \in G \cap \text{umax}(B'_2)$ . Using an asm-path from  $c_3$  to  $c'_2$  by Lemma 49 we can show that there is also an injective mapping from  $F/\eta$  to  $G/\eta$ . Therefore,  $|G/\eta| = |F/\eta|$ , and so  $f(G/\eta) = F/\eta$ . As we proved above, it means that for any  $\eta$ -block  $F' \subseteq F$  there

is  $d \in F'$  with  $(b, d) \in Q''$ , a contradiction with the assumption that  $F$  contains elements from several  $\text{lk}_2^Q$ -blocks. Claim 2 is proved.

Let  $\eta^* = \text{Cg}_{\mathbb{A}_2}(\eta|_{E_2})$ . Replacing  $R$  with  $R/\underline{0}_1 \times \eta^* \times \underline{0}_3$  we may assume that  $\eta|_{E_2}$  is the equality relation. Then making use of Claim 2 assume that  $\text{lk}_2^Q = \underline{0}_2|_{E_2}$ .

CLAIM 3.  $\text{lk}_1^Q$  is either the equality relation or the full relation.

By what has been proved so far  $B'_1 = E_1$  is a module. Suppose that  $\text{lk}_1^Q \neq \underline{0}_{E_1}$ . Then there are  $a, b \in E_1$ ,  $a \neq b$ , and  $c \in E_2$ ,  $e_1, e_2 \in B'_3$  such that  $(a, c, e_1), (b, c, e_2) \in Q$  and  $\mathbf{a} = (a, c, e_1) \in \text{umax}(R')$ . Since  $R$  is chained, for any  $b' \in E_1$  there exists a  $\overline{B}$ -preserving polynomial  $f$  such that  $f(a) = a$  and

$$f(b) = b'. \text{ Then } ((a, d, e'_1), (b', d, e'_2)) \in \xi, \text{ where } \begin{pmatrix} a \\ d \\ e'_1 \end{pmatrix} = f \begin{pmatrix} a \\ c \\ e_1 \end{pmatrix}, \begin{pmatrix} b' \\ d \\ e'_2 \end{pmatrix} = f \begin{pmatrix} b \\ c \\ e_2 \end{pmatrix}. \text{ Since } a, b' \in \text{umax}(B'_1), d \text{ can be chosen from } \text{umax}(B'_2). \text{ We argue that } (a, b') \in \text{lk}_1^Q, \text{ implying that } \text{lk}_1^Q \text{ is the full relation.}$$

Let  $\mathbf{b} = (a, d, e'_1)$ . As  $\mathbf{a} \in \text{umax}(R')$  and  $B'_1 \times \text{umax}(B'_2) \subseteq R''$ , there is an asm-path  $\mathbf{b} = \mathbf{a}_1, \dots, \mathbf{a}_\ell = \mathbf{a}$  in  $R'$ , where  $\mathbf{a}_i = (a, c_i, e_i^*)$ . We now use polynomials similar to those constructed in Lemma 49. For  $i \in [\ell - 1]$  let  $g_i$  be a polynomial of  $R$  defined as follows: If  $\mathbf{a}_i \mathbf{a}_{i+1}$  is semilattice (majority, affine), then

$$g_i(x) = x \cdot \mathbf{a}_{i+1}, \quad g_i(x) = t_{\mathbf{a}_i \mathbf{a}_{i+1}}(x, \mathbf{a}_{i+1}), \quad g_i(x) = h_{\mathbf{a}_i \mathbf{a}_{i+1}}(x, \mathbf{a}_i, \mathbf{a}_{i+1}),$$

respectively. As is easily seen,  $g_i(\mathbf{a}_i) = \mathbf{a}_{i+1}$ . Also, if  $(b', d, e'_2) = \mathbf{a}'_1, \dots, \mathbf{a}'_\ell$ , where  $\mathbf{a}'_{i+1} = g_i(\mathbf{a}'_i)$  for  $i \in [\ell - 1]$ , then  $(\mathbf{a}_i, \mathbf{a}'_i) \in \xi$  for all  $i \in [\ell]$ , because  $(\mathbf{a}_1, \mathbf{a}'_1) \in \xi$  by construction. Moreover,  $\mathbf{a}'_i = (b', c_i, e_i^*)$  for some  $e_i^* \in B'_3$ . Therefore  $(b', c) \in Q''$ , implying  $(a, b') \in \text{lk}_1^Q$ . Claim 3 is proved.

Recall that we assume  $\eta|_{E_2} = \text{lk}_2^Q$  to be the equality relation. This however does not imply that  $\eta$  itself or even its restriction to  $B'_2$  are the equality relations. Since  $(\underline{0}_1, \beta_1)$  and  $(\eta, \theta)$  cannot be separated for any  $\theta \in \text{Con}(\mathbb{A}_2)$  with  $\eta \prec \theta \leq \beta_2$ , by Claim 2 it remains to prove that  $\eta \prec \gamma$ . By Claim 2  $E_2/\eta$  is a module. Take any  $c, d, d' \in E_2^C = E_2$ ,  $c \neq d$ . We will show that  $(c, d') \in \theta = \text{Cg}_{\mathbb{A}_2}((c, d)) \vee \eta$ , which, as  $c, d, d'$  are arbitrary, proves that  $\eta \prec \gamma$ . Since  $E_2$  is a module,  $\text{umax}(E_2) = E_2$ . Let  $a, b, b' \in E_1$  and  $e_1, e_2, e_3 \in B'_3$  be such that  $(a, c, e_1), (b, d, e_2), (b', d', e_3) \in Q$ . As  $c \in \text{umax}(E_2) \subseteq \text{umax}(B'_2)$  and  $C \times \text{umax}(B'_2) \subseteq R''$ , the triple  $(a, c, e_1)$  can be chosen u-maximal in  $R'$ .

Since  $c \neq d$ , it also holds that  $a \neq b$ . As  $R$  is chained, there is a  $\overline{B}$ -preserving polynomial  $f$  with  $f(a) = a, f(b) = b'$ . Let  $f(c) = c'', f(d) = d'', f(e_1) =$



$e'_1, f(e_2) = e'_2, f(e_3) = e'_3$ . By the choice of  $(a, c, e_1)$  we have  $(a, c'', e'_1) \sqsubseteq_{R'}^{asm} (a, c, e_1)$ . As in the proof of Claim 3 using again the polynomials introduced in Lemma 49 there is a  $\overline{B}$ -preserving polynomial  $g$  such that  $g(a) = a, g(c'') = c, g(e'_1) = e_1$ , and  $g(b') = b'$ . Since  $(a, c, e_1) \stackrel{\xi}{\equiv} (g(b'), g(d''), g(e'_1))$ , it holds that  $(b', g(d'')) \in Q''$  and so  $g(d'') = d'$ . Therefore  $c = gf(c) \stackrel{\theta}{\equiv} gf(d) = d'$ .  $\square$

*Proof of Lemma 56.* Recall that unlike the two previous lemmas we cannot assume  $\alpha = \underline{0}_1$  at the moment. Let again  $R'' = \text{pr}_{I'} R'$  and  $Q'' = \text{pr}_{I'} Q$ . Note that if  $|C/\alpha| = 1$ , the lemma is trivially true, because  $E_2^C = Q''[C/\alpha]$ , so (1) holds. By Lemma 32 this happens in particular when  $\text{typ}(\underline{0}_1, \beta_1) \in \{4, 5\}$ .

By Lemma 58 the statement of Lemma 56 holds for  $\text{Block}(Q)$ . Suppose first that option (1) of the lemma holds. Then, as  $Q$  is polynomially closed, for any  $(a, c, e) \in \text{umax}(Q)$ ,  $a \in C$ , and any  $(b, d, e') \in \text{Block}(Q)$  with  $(a, c, e) \sqsubseteq^{as} (b, d, e')$  in  $\text{Block}(Q)$ , we have  $(b, d, e') \in Q$ . Thus, option (1) of the lemma holds for  $Q$ , as well.

Suppose now that option (2) of the lemma holds for  $S = \text{Block}(Q)$ . Since we are not using polynomial closeness anymore, we again assume that  $\alpha = \underline{0}_1$ . The proof in this case is almost verbatim the argument in the beginning of the proof of Lemma 58. The algebra  $S' = \text{pr}_{12} S$ , and therefore  $Q''$  is the graph of a mapping, and we only need to verify that the congruences  $\eta, \gamma$  satisfy the desired conditions. Let  $E_i^S = \text{pr}_i S$ , and  $\text{lk}_i^S$  be the link congruence of  $E_i^S$  with respect to  $S$ ,  $i = 1, 2$ . Let  $\eta^S$  be a maximal congruence such that  $\eta^S|_{\text{umax}(E_2^S)} \subseteq \text{lk}_2^S$  and  $\eta^S \leq \beta_2$ . Let  $D_2^C = \text{umax}(S'[C])$  and  $\gamma^S = \text{Cg}_{\mathbb{A}_2}(D_2^C) \vee \eta^S$ . As  $\text{umax}(E_2) \subseteq \text{umax}(E_2^S)$ ,  $\text{umax}(E_2^C) \subseteq D_2^C$ , and  $\text{lk}_2^Q = \text{lk}_2^S \cap E_2^Q$ , we have  $\eta^S|_{\text{umax}(E_2^C)} \subseteq \text{lk}_2^Q$ , and therefore  $\eta^S$  can be chosen such that  $\eta^S \leq \eta$ . As option (2) holds for  $S$ ,  $\eta^S \prec \gamma^S$ . Let  $\theta \in \text{Con}(\mathbb{A}_2)$  be such that  $\eta \prec \theta \leq \gamma$ . As  $\eta^S \prec \gamma^S$ , it also holds that  $\gamma^S = \text{Cg}_{\mathbb{A}_2}((c, d)) \vee \eta^S$  for any  $(c, d) \in (\text{umax}(E_2^C))^2 \cap (\theta - \eta)$  (recall that  $\text{umax}(E_2^C) \subseteq \text{umax}(D_2^C)$  and therefore  $(c, d) \in \gamma^S$ ). Therefore  $\gamma^S \leq \theta$ . By the definition of  $\gamma^S$  it holds that  $E_2^S$  is in a single  $\theta$ -block, hence,  $\gamma \leq \theta$ , implying  $\eta \prec \gamma$ . Also,  $\gamma^S \wedge \eta = \eta^S$ , because  $\eta^S \leq \gamma^S \wedge \eta$  and  $\eta^S \prec \gamma^S$ , and  $\gamma^S \vee \eta = \gamma$ , because  $\text{umax}(E_2^C) \subseteq D_2^C$ . Therefore, the intervals  $(\eta^S, \gamma^S)$  and  $(\eta, \gamma)$  are perspective. By Lemma 44 these intervals cannot be separated. Since  $(\alpha, \beta_1)$  and  $(\eta^S, \gamma^S)$  cannot be separated, it implies that  $(\alpha, \beta_1)$  and  $(\eta, \gamma)$  cannot be separated as well.  $\square$



## 7 Chaining and maximality

In this section we show that the chaining condition holds in a fairly broad range of circumstances. In particular, it is preserved under certain transformations of the relation.

**Lemma 59.** *Let  $R$  be a subdirect product of  $\mathbb{A}_1, \dots, \mathbb{A}_n \in \mathcal{V}$ ,  $\beta_i = \underline{1}_i$ ,  $B_i = \mathbb{A}_i$ . Then  $R$  is chained with respect to  $\overline{\beta}, \overline{B}$ .*

*Proof.* Let  $I, J \subseteq [n]$ . By Lemmas 34(2), 36(1) we may assume that  $I \cap J = \emptyset$  and that  $|I| = |J| = 1$ . Let  $i, j \in [n]$ ,  $\alpha \leq \beta$ ,  $\alpha, \beta \in \text{Con}(\mathbb{A}_i)$ , and  $\gamma \prec \delta$ ,  $\gamma, \delta \in \text{Con}(\mathbb{A}_j)$ , be such that if  $\alpha \prec \beta$ , then  $(\alpha, \beta)$  can be separated from  $(\gamma, \delta)$ . Consider first condition (Q1). Since any polynomial of  $R$  is  $\overline{B}$ -preserving, (Q1) follows from the definitions.

For condition (Q2) let  $(a, b) \in \beta - \alpha$ , and let  $f$  be a unary idempotent polynomial such that  $f(\mathbb{A}_i)$  is an  $(\alpha, \beta)$ -minimal set and  $f(a) \not\stackrel{\alpha}{\equiv} f(b)$ . By Lemma 35(1)  $f$  can be chosen such that  $f(\delta) \subseteq \gamma$ . Since  $\beta = \text{Cg}_{\mathbb{A}_i}(\{(f(a), f(b))\}) \vee \alpha$ , for any  $(c, d) \in \beta$  there are polynomials  $f_1, \dots, f_k$  such that for some  $c_1, \dots, c_k, c'_1, \dots, c'_k$  with  $\{f_i(f(a)), f_i(f(b))\} = \{c_i, c'_i\}$ ,  $c'_i \stackrel{\alpha}{\equiv} c_{i+1}$  for  $i \in [k]$ ,  $f_1(f(a)) = c$ , and  $c'_k \stackrel{\alpha}{\equiv} d$ . Then the polynomials  $f_i \circ f$ ,  $i \in [k]$ , witness that  $(c, d) \in \text{Cg}_{\mathbb{A}_i, \alpha, \mathcal{U}(\gamma, \delta, \overline{B})}(\{(a, b)\})$ , and, as  $(c, d)$  is arbitrary, that  $\beta = \text{Cg}_{\mathbb{A}_i, \alpha, \mathcal{U}(\gamma, \delta, \overline{B})}(\{(a, b)\})$ .  $\square$

**Lemma 60.** *Let  $R$  be a subdirect product of  $\mathbb{A}_1, \dots, \mathbb{A}_n \in \mathcal{V}$ ,  $\beta_i \in \text{Con}(\mathbb{A}_i)$  and  $B_i$  a  $\beta_i$ -block,  $i \in [n]$ , such that  $R$  is chained with respect to  $\overline{\beta}, \overline{B}$ . Let  $R' = R \cap \overline{B}$  and  $B'_i = \text{pr}_i R'$ . Fix  $i \in [n]$ ,  $\beta'_i \prec \beta_i$ , and let  $D_i$  be a  $\beta'_i$ -block that is as-maximal in  $B'_i / \beta'_i$ . Let also  $\beta'_j = \beta_j$  and  $D_j = B_j$  for  $j \neq i$ . Then  $R$  is chained with respect to  $\overline{\beta'}, \overline{D}$ .*

*Proof.* Let  $R'' = R \cap \overline{D}$  and  $D'_j = \text{pr}_j R''$ ,  $j \in [n]$ . Take  $I, J$  from the definition of chaining. Let  $I = [\ell]$ ; if  $|I| > 1$ , by Lemma 36(1) we may consider  $R$  as a subdirect product of  $\text{pr}_I R$  and  $\mathbb{A}_{\ell+1}, \dots, \mathbb{A}_n$ , so we assume  $I = \{1\}$  and, for the same reason in condition (Q2), we assume  $|J| = 1$  and  $J = \{n\}$ . Let  $\alpha, \beta \in \text{Con}(\mathbb{A}_1)$ ,  $\gamma, \delta \in \text{Con}(\mathbb{A}_n)$  be such that  $\alpha \leq \beta \leq \beta'_1$ ,  $\gamma \prec \delta \leq \beta'_n$  (we assume  $\alpha \prec \beta$  when considering condition (Q2)). Clearly, we may assume  $\alpha = \underline{0}_1$ ,  $\gamma = \underline{0}_n$ , and  $\beta'_i = \underline{0}_i$ . Note that again by Lemma 36(1) replacing  $R$  with the  $n+1$ -ary relation  $\{(\mathbf{a}, \mathbf{a}[i]) \mid \mathbf{a} \in R\}$  we may assume that  $i \notin I \cup J$ . Without loss of generality assume  $i = 2$ . By the assumption  $\beta'_2 = \underline{0}_2$ , the classes of  $\beta'_2$  are just elements of  $\mathbb{A}_2$ , so let  $D'_2$  be denoted by  $c$ . Let  $C$  be the as-component of  $B'_2$  containing  $c$ .

To prove the lemma let  $\mathcal{U}^* \in \{\mathcal{U}_{\overline{D}}, \mathcal{U}(\gamma, \delta, \overline{D})\}$ . We assume that  $\beta$  is a principal congruence of  $\mathbb{A}_1$  such that  $\underline{0}_1 < \beta \leq \beta_1 = \beta'_1$ . Let  $a, b \in D'_1$  be any elements such that  $\text{Cg}_{\mathbb{A}_1}(\{(a, b)\}) = \beta$ . We need to prove that for any  $\beta$ -block  $B$  such that  $B \cap \text{umax}(D'_1) \neq \emptyset$ , it holds that  $(B \cap \text{umax}(D'_1))^2 \subseteq \text{Cg}_{\mathbb{A}_1, \underline{0}_1, \mathcal{U}^*}(a, b)$ . Note that it would be true if  $\mathcal{U}^*$  were one of  $\mathcal{U}_{\overline{B}}, \mathcal{U}(\gamma, \delta, \overline{B})$  by the assumption that  $R$  is chained with respect to  $\overline{\beta}, \overline{B}$ , which is witnessed by polynomials from  $\mathcal{U}_{\overline{B}}$  and  $\mathcal{U}(\gamma, \delta, \overline{B})$ . We need to change these polynomials so that they work for  $\overline{D}$ . Let  $Q' = \text{pr}_{12}R'$ ,  $B_1^C = Q'^{-1}[C]$ , and  $Q^C = Q' \cap (B_1^C \times C)$ . We divide the proof into two cases, depending on whether or not  $Q^C$  is linked.

First, we consider the case when  $Q^C$  is not linked, this case is relatively easy. Let  $\text{lk}_1^C, \text{lk}_2^C$  be defined for  $Q^C$  in the same way as link congruences are defined for a subdirect product. Note that  $Q^C$  is not necessarily a subalgebra, and  $\text{lk}_1^C, \text{lk}_2^C$  are not necessarily congruences.

CLAIM 1. Let  $Q^C$  be not linked and let  $\text{lk}_1^C, \text{lk}_2^C$  be as above. Then  $\text{lk}_2^C = \underline{0}_2|_C$  and either  $\beta|_{B_1^C} \subseteq \text{lk}_1^C$  or  $(\beta \cap \text{lk}_1^C)|_{B_1^C} = \theta|_{B_1^C}$  for some  $\theta \in \text{Con}(\mathbb{A}_1)$ ,  $\underline{0}_1 \leq \theta < \beta$ .

The relation  $Q'$  is a (trivial) subalgebra of  $\text{pr}_{12}R'$  and is polynomially closed in  $\text{pr}_{12}R$  by Lemma 55(2,5), and the second statement of Lemma 55(3). By the Congruence Lemma 56 if  $Q^C$  is not linked then  $Q^C$  is the graph of a mapping  $\varphi : B_1^C \rightarrow C$ . This means  $\text{lk}_2^C = \underline{0}_2|_C$  and  $\text{lk}_1^C$  is the restriction of a congruence  $\eta$  of  $\mathbb{A}_1$  defined in the Congruence Lemma 56 onto  $B_1^C$ . If  $\beta \leq \eta$ , we obtain the first option of the conclusion of the claim, otherwise  $\theta = \beta \wedge \eta$  and we have the second option.

Note that if  $\beta|_{B_1^C} \leq \text{lk}_1^C$  then any  $\overline{B}$ -preserving polynomial that maps a pair of  $\beta$ -related elements from  $D'_1$  on a pair from  $D'_1$  is also  $\overline{D}$ -preserving, because  $\text{lk}_2 = \underline{0}_2|_C$ . The result follows, as  $\text{umax}(D'_1) \subseteq \text{umax}(B'_1)$  by Lemma 17. If  $(\beta \wedge \text{lk}_1)|_{B_1^C} = \theta|_{B_1^C}$  for some  $\theta < \beta$ , there is nothing to prove, because no pair  $(a, b) \in \beta \cap (D'_1)^2$  generates  $\beta$ . Therefore we may assume  $Q^C$  is linked.

We start by choosing a  $\beta$ -block containing elements u-maximal in  $D'_1$  and studying some of its properties. Observe that since  $c$  is as-maximal in  $B'_2$ , the set  $D'_1$  also contains as-maximal elements of  $B'_1$ . Therefore by Lemma 17  $\text{umax}(D'_1) \subseteq \text{umax}(B'_1)$ . Let  $E$  be a  $\beta$ -block such that  $E \cap \text{umax}(D'_1) \cap B_1^C \neq \emptyset$ , and let  $E' = E \cap B'_1$ ,  $E^C = E \cap B_1^C$ , and  $E^* = E \cap D'_1$ . By the Maximality Lemma 18(4)  $\text{amax}(E^C)$  is a union of as-components of  $E'$ . Indeed, let  $a \in E^C$  and let  $\mathbf{a} \in R'$  be such that  $\mathbf{a}[1] = a$  and  $\mathbf{a}[2] \in C$ ; let also  $b \in E'$  with  $a \sqsubseteq^{as} b$  in  $E'$ . Then by the Maximality Lemma 18(4) there is  $\mathbf{b} \in R'$  such that  $\mathbf{b}[1] = b$  and  $\mathbf{a} \sqsubseteq^{as} \mathbf{b}$  in  $R'$ . In particular,  $\mathbf{a}[2] \sqsubseteq^{as} \mathbf{b}[2]$  implying  $\mathbf{b}[2] \in C$ . Also, by Proposition 27 applied to the subalgebra generated by  $Q^C$ , since  $Q^C$  is linked and  $\text{umax}(E^C) \subseteq \text{umax}(B_1^C)$ , we have  $\text{umax}(E^C) \times C \subseteq Q^C$ , and therefore  $\text{umax}(E^C) = \text{umax}(E^*) \subseteq \text{umax}(E')$ .

In particular,  $\text{amax}(E^*)$  is a union of as-components of  $E'$ . The last inclusion here is because  $E^C$  contains some as-maximal elements of  $E'$ .

First we prove that there are polynomials with images from an as-component of  $E^*$  for both conditions (Q1),(Q2) for  $\overline{\beta}', \overline{D}$ .

CLAIM 2. For any  $a, b \in E^*$ ,  $a', b' \in D'_1$  such that  $a, b$  belong to the same as-component of  $E^*$  and  $\text{Cg}_{\mathbb{A}_1}(\{(a', b')\}) = \beta$ , there exists a polynomial  $f$  from  $\mathcal{U}_{\overline{D}}$  with  $f(\{a', b'\}) = \{a, b\}$ . Moreover, if  $\alpha \prec \beta$  polynomial  $f$  can be chosen from  $\mathcal{U}(\gamma, \delta, \overline{D})$ .

Consider the relation  $S$ , a subdirect product of  $\mathbb{A}_1 \times \mathbb{A}_1 \times \mathbb{A}_2 \times \cdots \times \mathbb{A}_n$ , produced from  $(a', b', \mathbf{a})$ , where  $\mathbf{a}$  is a fixed tuple from  $\text{pr}_{\{2, \dots, n\}} R''$ , as follows:

$$S = \{(f(a'), f(b'), f(\mathbf{a})) \mid f \text{ is a unary polynomial of } R\},$$

and for (Q2) we also assume that polynomials in the definition of  $S$  satisfy  $f(\delta) \subseteq \gamma$ . By Lemma 37(1,2)  $S$  is a subalgebra, and, in particular it contains all the tuples of the form  $(\mathbf{b}[1], \mathbf{b}[1], \mathbf{b}[2], \dots, \mathbf{b}[n])$  for  $\mathbf{b} \in R$ . Let  $S' = S \cap \overline{B}$ , and  $S'' = S \cap \overline{D}$ . Observe that  $S'' = \{\mathbf{b} \in S' \mid \mathbf{b}[3] = c\}$ . Every tuple from  $S'$  or from  $S''$  corresponds to a  $\overline{B}$ - or  $\overline{D}$ -preserving polynomial. Therefore it suffices to prove that  $(a, b) \in \text{pr}_{12} S''$  or equivalently, that  $(a, b, c) \in P = \text{pr}_{123} S'$ .

Let  $F$  be the as-component of  $E^*$  containing  $a, b$ ; as observed above  $F$  is also an as-component of  $E'$ . As  $R$  is chained with respect of  $\overline{\beta}, \overline{B}$ , by Lemma 37(3) it holds that  $F^2 \subseteq \text{pr}_{12} P$ . Also,  $(e, e) \in \text{pr}_{12} S''$  and  $(e, e, d) \in P$  for any  $e \in F$  and  $d \in C$ , since  $F \times C \subseteq Q^C$ . Therefore considering  $P$  as a subdirect product of  $\text{pr}_{12} P$  and  $B'_2$ , the as-component  $C$  belongs to a block of the link congruence. As  $F^2 \subseteq \text{pr}_{12} P$ , every pair from  $F^2$  is as-maximal in  $\text{pr}_{12} P$ . Moreover, by the Maximality Lemma 18(4)  $P \cap (F^2 \times C)$  is subdirect on  $F^2$  and  $C$ . Hence  $F^2$  also belongs to a block of the link congruence. By Proposition 26  $F^2 \times C \subseteq P$ , in particular  $(a, b, c) \in P$ , as required. Claim 2 is proved.

Now we extend the result of Claim 2 to pairs from  $\text{umax}(E^*)$ . We prove the result in two steps. First, we show that for any  $a', b' \in D'_1$  such that  $\text{Cg}_{\mathbb{A}_1}(\{(a', b')\}) = \beta$  and any  $a, b \in \text{umax}(E^*)$  there is a sequence of  $\overline{B}$ -preserving polynomials  $f_1, \dots, f_k$  such that  $f_1(\{a', b'\}), \dots, f_k(\{a', b'\}) \subseteq E^*$  form a chain connecting  $a$  and  $b$ , and  $f_i(c) \in C$  for  $i \in [k]$ . Then we prove that  $f_1, \dots, f_k$  can be chosen in such a way that  $f_1(\{a', b'\}), \dots, f_k(\{a', b'\}) \subseteq E^*$  and  $f_1(c) = \dots = f_k(c) = c$ . Clearly, it suffices to prove in the case when  $b$  is as-maximal in  $E^*$ , since we can link  $a$  and  $b$  to an as-maximal element  $b'$  and then concatenate the two chains. We will also observe that in both cases the polynomials  $f_1, \dots, f_k$  can be chosen from  $\mathcal{U}(\gamma, \delta, \overline{B})$  when necessary.

By the assumption there are  $a = a_1, a_2, \dots, a_k = b$ ,  $a_1, \dots, a_k \in E'$  and polynomials  $f_1, \dots, f_{k-1}$  from  $\mathcal{U}_{\overline{B}}$  (or from  $\mathcal{U}(\gamma, \delta, \overline{B})$  when  $\alpha \prec \beta$ ) such that

$f_i(\{a', b'\}) = \{a_i, a_{i+1}\}$ , and also  $f_i(c) \in B'_2$ . We need to show that  $a_1, \dots, a_{k-1}$  and  $f_1, \dots, f_{k-1}$  can be chosen such that  $f_i(c) \in C$ . Choose  $\mathbf{a}, \mathbf{b} \in R''$  such that  $\mathbf{a}[1] = a, \mathbf{b}[1] = b$  and  $\mathbf{a}[2] = \mathbf{b}[2] = c$ . Now let  $g_i(x) = \text{maj}(\mathbf{a}, f_i(x), \mathbf{a})$  and  $h_i(x) = \text{maj}(\mathbf{a}, \mathbf{b}, f_i(x))$ , where  $\text{maj}$  is a quasi-majority operation, see Theorem 24. By Lemma 37(2)  $g_i, h_i$  are from  $\mathcal{U}_{\overline{B}}$  and from  $\mathcal{U}(\gamma, \delta, \overline{B})$  whenever  $f_1, \dots, f_k$  are from  $\mathcal{U}(\gamma, \delta, \overline{B})$ . Also, for each of them either  $\{b_i, b_{i+1}\} = g_i(\{a', b'\})$  ( $\{c_i, c_{i+1}\} = h_i(\{a', b'\})$ ) is from  $T(\{(a', b')\}, \mathcal{U}^*)$ ,  $\mathcal{U}^* \in \{\mathcal{U}_{\overline{B}}, \mathcal{U}(\gamma, \delta, \overline{B})\}$ , or  $g_i(a') = g_i(b')$  (respectively,  $h_i(a') = h_i(b')$ ). The polynomials  $g_i, h_i$  satisfying the first option form a sequence of pairs connecting  $a$  with  $\text{maj}(a, b, a)$  — by pairs of the form  $\{b_i, b_{i+1}\}$ , — and  $\text{maj}(a, b, a)$  with  $\text{maj}(a, b, b)$  — by pairs of the form  $\{c_i, c_{i+1}\}$ . Also, by Theorem 24  $\text{maj}(a, b, b)$  belongs to the as-component of  $E^*$  (and therefore of  $E'$ ) containing  $b$ . Therefore by Claim 2 this sequence of polynomials and pairs can be continued to connect  $\text{maj}(a, b, b)$  to  $b$ . Finally, by the same theorem  $g_i(c) = \text{maj}(c, f_i(c), c) \in C$  and  $h_i(c) = \text{maj}(c, c, f_i(c)) \in C$ .

For the second step we assume that  $a$  and  $b$  are connected with pairs  $\{a_i, a_{i+1}\}$ ,  $i \in [k-1]$  witnessed by polynomials  $f_i$  from  $\mathcal{U}_{\overline{B}}$  (or from  $\mathcal{U}(\gamma, \delta, \overline{B})$  when needed) such that  $c_i = f_i(c) \in C$ . We need to show that  $f_i$  can be chosen such that  $f_i(c) = c$ . Suppose that  $c_i \neq c$  for some  $i \in [k-1]$ . Since  $c_i$  and  $c$  belong to the same as-component, there is an as-path  $c_i = d_1, \dots, d_\ell = c$  in  $C$ . Suppose that there is a sequence of pairs  $\{b_j, b_{j+1}\} = g_j(\{a', b'\})$ ,  $b_1 = a$  and  $b_k = b$ , for some polynomials  $g_j \in \mathcal{U}_{\overline{B}}$  (or from  $\mathcal{U}(\gamma, \delta, \overline{B})$  when needed),  $j \in [k-1]$ , such that  $g_j(c) = c$  whenever  $f_j(c) = c$ , and  $g_i(c) = d_t$ . We will show that there are also pairs  $\{b'_j, b'_{j+1}\} = g'_j(\{a', b'\})$  for some polynomials  $g'_j \in \mathcal{U}_{\overline{B}}$  (or from  $\mathcal{U}(\gamma, \delta, \overline{B})$  when needed) such that  $b'_1 = a$  and  $b'_k$  is in the as-component containing  $b$ ,  $g'_i(c) = d_{t+1}$  and  $g'_j(c) = c$  whenever  $g_j(c) = c$ .

As is easily seen, it suffices to find a ternary term operation  $p$  such that  $p(a, a, b)$  belongs to the as-component containing  $b$ , and  $p(d_{t+1}, d_t, d_t) = d_{t+1}$ . Indeed, if such a term operation exists, then we set  $g'_j(x) = p(\mathbf{a}, \mathbf{a}, g_j(x))$ , where  $\mathbf{a}$  is as in the first step above, for  $j \in [k-1] - \{i\}$ , and  $\{b'_j, b'_{j+1}\} = g'_j(\{a', b'\})$ . We have  $g'_1(a') = p(a, a, g_1(a')) = a$  and  $g'_j(c) = p(c, c, g_j(c)) = c$  whenever  $g_j(c) = c$ . Finally, since  $g'_{k-1}(b) = p(a, a, b)$  belongs to the as-component containing  $b$ , we can use Claim 2 as before to connect  $p(a, a, b)$  to  $b$ . For  $g'_i$  we set  $g'_i(x) = p(\mathbf{a}', \mathbf{a}'', g_i(x))$  where  $\mathbf{a}', \mathbf{a}'' \in R''$  are such that  $\mathbf{a}'[1] = \mathbf{a}''[1] = a$  and  $\mathbf{a}'[2] = d_{t+1}, \mathbf{a}''[2] = d_t$ . Note that such  $\mathbf{a}', \mathbf{a}''$  exist, because  $\text{umax}(E^C) \times C \subseteq Q'$ . It follows from the assumption about  $p$  that  $g'_i$  is as required.

If  $d_t \leq d_{t+1}$ , then  $p(x, y, z) = z \cdot x$  fits the requirements. If  $d_t d_{t+1}$  is an affine edge, consider the relation  $Q^\dagger \subseteq \mathbb{A}_1 \times \mathbb{A}_2$  generated by  $\{(a, d_{t+1}), (a, d_t), (b, d_t)\}$ . Let  $\mathbb{B} = \text{Sg}_{\mathbb{A}_1}(a, b)$  and  $\mathbb{C} = \text{Sg}_{\mathbb{A}_2}(d_t, d_{t+1})$ ; then  $\mathbb{B} \times \{d_t\}, \{a\} \times \mathbb{C} \subseteq Q^\dagger$ . By Lemma 25, as  $d_t d_{t+1}$  is a thin affine edge,  $\text{umax}(\mathbb{B}) \times \{d_{t+1}\} \subseteq Q^\dagger$ . There is  $b'$

with  $b \sqsubseteq^{as} b'$  in  $\mathbb{B}$  such that  $b' \in \text{umax}(\mathbb{B})$ . Therefore there is a term operation  $p$  with  $p(a, a, b) = b'$  and  $p(d_{t+1}, d_t, d_t) = d_{t+1}$ , as required.  $\square$

## Declarations

### Data availability

Data sharing not applicable to this article as datasets were neither generated nor analyzed.

### Compliance with ethical standards

The author is a member of the Editorial Board of Algebra Universalis. Apart from this the author declares that he has no conflict of interest.

## References

- [1] A. Bulatov. Three-element Mal'tsev algebras. *Acta Sci. Math (Szeged)*, 71(3-4):469–500, 2002.
- [2] Andrei A. Bulatov. A graph of a relational structure and constraint satisfaction problems. In *LICS*, pages 448–457, 2004.
- [3] Andrei A. Bulatov. Graphs of finite algebras, edges, and connectivity. available at <https://www.cs.sfu.ca/~abulatov/papers/graph-connect-1.pdf>, 2016.
- [4] Andrei A. Bulatov. Graphs of relational structures: restricted types. In *LICS*, 2016.
- [5] Andrei A. Bulatov. Constraint satisfaction problems over semilattice block Mal'tsev algebras. *CoRR*, abs/1701.02623, 2017.
- [6] Andrei A. Bulatov. A dichotomy theorem for nonuniform CSPs. In *FOCS*, pages 319–330, 2017.
- [7] Andrei A. Bulatov. A dichotomy theorem for nonuniform CSPs. *CoRR*, abs/1703.03021, 2017.
- [8] Andrei A. Bulatov. Constraint satisfaction problems over semilattice block Mal'tsev algebras. *Inf. Comput.*, 268, 2019.
- [9] Andrei A. Bulatov. Graphs of relational structures: restricted types. *CoRR*, abs/2006.11713, 2020.

- [10] Andrei A. Bulatov. Local structure of idempotent algebras I. *CoRR*, abs/2006.09599, 2020.
- [11] Andrei A. Bulatov. Local structure of idempotent algebras II. *CoRR*, abs/2006.10239, 2020.
- [12] Andrei A. Bulatov, Peter Jeavons, and Andrei A. Krokhin. Classifying the complexity of constraints using finite algebras. *SIAM J. Comput.*, 34(3):720–742, 2005.
- [13] Andrei A. Bulatov and Matthew Valeriote. Recent results on the algebraic approach to the CSP. In *Complexity of Constraints - An Overview of Current Research Themes [Result of a Dagstuhl Seminar]*., pages 68–92, 2008.
- [14] Stanley Burris and H.P. Sankappanavar. *A course in universal algebra*, volume 78 of *Graduate Texts in Mathematics*. Springer-Verlag, 1981.
- [15] Tomas Feder and Moshe Vardi. Monotone monadic SNP and constraint satisfaction. In *STOC*, pages 612–622, 1993.
- [16] Tomas Feder and Moshe Vardi. The computational structure of monotone monadic SNP and constraint satisfaction: A study through datalog and group theory. *SIAM Journal of Computing*, 28:57–104, 1998.
- [17] Ralph Freese and Ralph McKenzie. *Commutator theory for congruence modular varieties*, volume 125 of *London Math. Soc. Lec. Notes*. London, 1987.
- [18] David Hobby and Ralph McKenzie. *The Structure of Finite Algebras*, volume 76 of *Contemporary Mathematics*. American Mathematical Society, Providence, R.I., 1988.
- [19] Peter Jeavons, David A. Cohen, and Marc Gyssens. Closure properties of constraints. *J. ACM*, 44(4):527–548, 1997.
- [20] Keith Kearnes. Idempotent simple algebras. In *Logic and algebra*, volume 180 of *Lecture Notes in Pure and Appl. Math.*, pages 529–572, 1996.
- [21] Ágnes Szendrei Keith Kearnes and Ross Willard. A finite basis theorem for difference-term varieties with a finite residual bound. *Trans. Amer. Math. Soc.*, 368:2115–2143, 2016.
- [22] Ralph McKenzie, George McNulty, and Walter Taylor. *Algebras, Lattices, Varieties, I*. Wadsworth–Brooks/Cole, Monterey, California, 1987.

- [23] Ágnes Szendrei. Idempotent algebras with restrictions on subalgebras. *Acta Sci. Math. (Szeged)*, 51:251–268, 1987.
- [24] Matthew Valeriote. Finite Simple Abelian Algebras are Strictly Simple. *Proceedings of the American Mathematical Society*, 108(1):49–57, 1990.
- [25] Ross Willard. Personal communication, 2019.
- [26] Dmitriy Zhuk. A proof of CSP dichotomy conjecture. In *FOCS*, pages 331–342, 2017.
- [27] Dmitriy Zhuk. The proof of CSP dichotomy conjecture. *CoRR*, abs/1704.01914, 2017.