

Fermionic model of unitary transport of qubits from a black hole

Bogusław Broda*

*Department of Theoretical Physics, Faculty of Physics and Applied Informatics,
University of Łódź, 90-236 Łódź, Pomorska 149/153, Poland[†]*

Inspired by a recent model of Osuga and Page, we propose an explicitly unitary fermionic toy model for transferring information from a black hole to the outgoing radiation. The model treats the unitary evolution as a composition of the Hawking pair creation outside the black hole and of pair annihilation inside the black hole.

I. INTRODUCTION

The black hole (BH) information (loss) paradox concerns difficulties around the issue of unitarity of BH evaporation (for recent reviews see e.g. [4, 8, 11, 17]). There are a lot of approaches proposed to date to analyze and resolve the paradox — some of them suggest to study simplified situations embodied in various qubit models (see e.g. [2, 5–7, 12–14, 16]). A successful model of BH evolution should include description of particle pair production according to the Hawking prescription, following gradual evaporation (“vanishing”) of the BH, and it should be unitary. A model strictly motivated by actual physical phenomena would certainly be greatly appreciated, but in fact any model respecting at least general physical laws, even without any real physical mechanism built in, would be welcome as a “proof of concept”.

Recently Osuga and Page [16] (inspired by [1]) have proposed an explicitly unitary toy qubit transport model for BH evaporation (without firewalls). Another version of the model (with additional features) has been presented in [3]. In the present paper, building on the both models, we propose yet another toy qubit transport model for BH evaporation, which is explicitly unitary. Since the model, by assumption, operates on qubits and particle pair production scheme exactly follows the Hawking mechanism for fermions, we shall work in terms of fermionic modes rather than bosonic ones. A new and important feature of our present proposal is explicit incorporation of the (fermionic) Hawking pair creation mechanism into the chain of unitary processes. In other words, the global unitary evolution considered is given by the composition $U = U'' \cdot U'$, where U' corresponds to creation of fermionic pairs outside a BH according to the Hawking prescription, whereas U'' corresponds to annihilation of fermion pairs inside the BH (as described in [3]).

For reader’s convenience, we will follow notation of [16] (and [3]) as closely as possible.

II. THE TOY MODEL

An initial total quantum state describing a newly formed (fermionic) BH and “fermionic radiation” in the vacuum state is assumed in the following (partially product) form [3] (cf.

* boguslaw.broda@uni.lodz.pl

† <http://merlin.phys.uni.lodz.pl/BBroda>

[16])

$$|\Psi\rangle = \sum_{q_1, q_2, \dots, q_N=0}^1 A_{q_1 q_2 \dots q_N} \bigotimes_{k=1}^N |q_k\rangle_{a_k} \otimes |O\rangle_{b_k c_k}. \quad (1)$$

Here $A_{q_1 q_2 \dots q_N}$ are amplitudes for inner BH modes a_k , which encode a quantum state of the BH, and the vacuum state for ‘‘fermionic radiation’’ is

$$|O\rangle_{b_k c_k} \equiv |0\rangle_{b_k} \otimes |0\rangle_{c_k}, \quad (2)$$

where the Hawking (fermionic) modes b_k and c_k are infalling and outgoing modes, respectively. The same range of indices ($k = 1, 2, \dots, N$) postulated for BH modes a_k and b_k, c_k pairs is not only a convenient computational simplification in our model but also it is a physically justified assumption, at least approximately (see e.g. [3]).

In the language of k -mode blocks, the first step of (unitary) evolution denoted by U'_k yields the Hawking (fermion) pair for a single ‘‘ k ’’ mode, i.e.

$$U'_k (|q_k\rangle_{a_k} \otimes |O\rangle_{b_k c_k}) = |q_k\rangle_{a_k} \otimes |H_1\rangle_{b_k c_k}, \quad (3)$$

where the fermionic Hawking state can be chosen in the form (cf. Eq.(116) in [10])

$$|H_1\rangle_{b_k c_k} \equiv \cos \omega_k |0\rangle_{b_k} \otimes |0\rangle_{c_k} + \sin \omega_k |1\rangle_{b_k} \otimes |1\rangle_{c_k}, \quad (4)$$

with ω_k determined by BH parameter(s).

The total (i.e. for all modes k) U' -evolution yields by virtue of (3) the total (intermediate) state

$$|\Psi'\rangle = \sum_{q_1, q_2, \dots, q_N=0}^1 A_{q_1 q_2 \dots q_N} \bigotimes_{i=1}^N |q_i\rangle_{a_i} \otimes |H_1\rangle_{b_i c_i}. \quad (5)$$

We could possibly consider a slight generalization of the unitary evolution (3) allowing some unitary transformation $\mathcal{U}'_k |q_k\rangle_{a_k}$ of the internal BH mode a_k on the RHS of (3), but we ignore this option, because it would merely give rise to a redefinition of A -amplitudes ($A_{q_1 q_2 \dots q_N} \mapsto A'_{q_1 q_2 \dots q_N}$) in the final state.

The second step of (unitary) evolution denoted by U''_k yields particle pair annihilation inside the BH (see [3] and cf. Eq. (3.3) in [1, 16]), i.e.

$$U''_k (|q_k\rangle_{a_k} \otimes |H_1\rangle_{b_k c_k}) = |O\rangle_{a_k b_k} \otimes |q_k\rangle_{c_k}, \quad (6)$$

where the vacuum state $|O\rangle_{a_k b_k}$ is defined analogously to (2) (with appropriate replacements of modes).

Consequently, for the entire evolution $U_k \equiv U_k'' \cdot U_k'$ we have

$$U_k (|q_k\rangle_{a_k} \otimes |O\rangle_{b_k c_k}) = |O\rangle_{a_k b_k} \otimes |q_k\rangle_{c_k}, \quad (7)$$

and the final total state assumes the form

$$|\Psi''\rangle = \sum_{q_1, q_2, \dots, q_N=0}^1 A_{q_1 q_2 \dots q_N} \bigotimes_{k=1}^N |O\rangle_{a_k b_k} \otimes |q_k\rangle_{c_k}. \quad (8)$$

Eq. (8) means that all information has been transferred from a BH to the outgoing radiation, and the BH is in the vacuum state.

Obviously, the total operators U' , U'' , U are the following tensor products of the above-defined k -mode operators

$$U' = \bigotimes_{k=1}^N U_k', \quad U'' = \bigotimes_{k=1}^N U_k'', \quad U = \bigotimes_{k=1}^N U_k'' \cdot U_k' \equiv \bigotimes_{k=1}^N U_k, \quad (9)$$

respectively, and

$$U' |\Psi\rangle = |\Psi'\rangle, \quad U'' |\Psi'\rangle = |\Psi''\rangle, \quad U |\Psi\rangle \equiv U'' \cdot U' |\Psi\rangle = |\Psi''\rangle. \quad (10)$$

III. UNITARY OPERATORS

We shall now explicitly derive the implicitly defined unitary operators U_k' , U_k'' and U_k . To this end let us first observe the following elementary fact from linear algebra: namely, for each pair of orthonormal bases $\{|E_\Lambda\rangle\}$, $\{|E'_{\Lambda'}\rangle\}$ ($\langle E_\Lambda | E_{\Lambda'} \rangle = \langle E'_{\Lambda'} | E'_{\Lambda'} \rangle = \delta_{\Lambda\Lambda'}$) in a finite dimensional Hilbert space \mathcal{H} we can construct an operator

$$U = \sum_{\Lambda} |E'_{\Lambda}\rangle \langle E_{\Lambda}|, \quad (11)$$

which is explicitly unitary. Really, we can easily check that e.g.

$$U^\dagger \cdot U = \sum_{\Lambda, \Lambda'} |E_\Lambda\rangle \langle E'_{\Lambda'} | E'_{\Lambda'} \rangle \langle E_{\Lambda'}| = \sum_{\Lambda, \Lambda'} \delta_{\Lambda\Lambda'} |E_\Lambda\rangle \langle E_{\Lambda'}| = \mathbb{I}. \quad (12)$$

Since the total Hilbert space \mathcal{H} is a tensor product of N k -mode Hilbert spaces \mathcal{H}_k , i.e. $\mathcal{H} = \bigotimes_{k=1}^N \mathcal{H}_k$, we can confine our construction to the single k -mode space $\mathcal{H}_k = \mathcal{H}_{a_k} \otimes \mathcal{H}_{b_k} \otimes \mathcal{H}_{c_k}$, where $\dim_{\mathbb{C}} \mathcal{H}_k = 2 \cdot 2 \cdot 2 = 8$. Then, our unitary operators will be defined by three 8-dimensional orthonormal (k -dependent) bases in \mathcal{H}_k .

The first base, $\{|E_\Lambda\rangle_k\}_{\Lambda=0}^7$, assumes the following standard form:

$$\begin{aligned}
|E_0\rangle_k &= |0\rangle_{a_k} \otimes |0\rangle_{b_k} \otimes |0\rangle_{c_k} \\
|E_1\rangle_k &= |0\rangle_{a_k} \otimes |0\rangle_{b_k} \otimes |1\rangle_{c_k} \\
&\vdots \\
|E_7\rangle_k &= |1\rangle_{a_k} \otimes |1\rangle_{b_k} \otimes |1\rangle_{c_k}
\end{aligned} \tag{13}$$

(or $|E_\Lambda\rangle_k = |\Lambda\rangle_{a_k b_k c_k}$, in short).

The second (τ -dependent) base, $\{|E'_\Lambda(\tau)\rangle_k\}_{\Lambda=0}^7$, is given by (for further convenience, we have also included expansions in terms of τ):

$$\begin{aligned}
|E'_0(\tau)\rangle_k &= |0\rangle_{a_k} \otimes [\cos(\omega_k \tau) |0\rangle_{b_k} \otimes |0\rangle_{c_k} + \sin(\omega_k \tau) |1\rangle_{b_k} \otimes |1\rangle_{c_k}] \\
&\equiv |0\rangle_{a_k} \otimes |H_\tau\rangle_{b_k c_k} \\
&\equiv \cos(\omega_k \tau) |E_0\rangle_k + \sin(\omega_k \tau) |E_3\rangle_k \\
&= |E_0\rangle_k + \omega_k \tau |E_3\rangle_k + \mathcal{O}(\tau^2) \\
|E'_1\rangle_k &= |0\rangle_{a_k} \otimes |0\rangle_{b_k} \otimes |1\rangle_{c_k} \equiv |E_1\rangle_k \\
|E'_2\rangle_k &= |0\rangle_{a_k} \otimes |1\rangle_{b_k} \otimes |0\rangle_{c_k} \equiv |E_2\rangle_k \\
|E'_3(\tau)\rangle_k &= |0\rangle_{a_k} \otimes [-\sin(\omega_k \tau) |0\rangle_{b_k} \otimes |0\rangle_{c_k} + \cos(\omega_k \tau) |1\rangle_{b_k} \otimes |1\rangle_{c_k}] \\
&\equiv |0\rangle_{a_k} \otimes |H_\tau^\perp\rangle_{b_k c_k} \\
&\equiv \cos(\omega_k \tau) |E_3\rangle_k - \sin(\omega_k \tau) |E_0\rangle_k \\
&= |E_3\rangle_k - \omega_k \tau |E_0\rangle_k + \mathcal{O}(\tau^2) \\
|E'_4(\tau)\rangle_k &= |1\rangle_{a_k} \otimes [\cos(\omega_k \tau) |0\rangle_{b_k} \otimes |0\rangle_{c_k} + \sin(\omega_k \tau) |1\rangle_{b_k} \otimes |1\rangle_{c_k}] \\
&\equiv |1\rangle_{a_k} \otimes |H_\tau\rangle_{b_k c_k} \\
&\equiv \cos(\omega_k \tau) |E_4\rangle_k + \sin(\omega_k \tau) |E_7\rangle_k \\
&= |E_4\rangle_k + \omega_k \tau |E_7\rangle_k + \mathcal{O}(\tau^2) \\
|E'_5\rangle_k &= |0\rangle_{a_k} \otimes |0\rangle_{b_k} \otimes |1\rangle_{c_k} \equiv |E_5\rangle_k \\
|E'_6\rangle_k &= |0\rangle_{a_k} \otimes |0\rangle_{b_k} \otimes |1\rangle_{c_k} \equiv |E_6\rangle_k \\
|E'_7(\tau)\rangle_k &= |1\rangle_{a_k} \otimes [-\sin(\omega_k \tau) |0\rangle_{b_k} \otimes |0\rangle_{c_k} + \cos(\omega_k \tau) |1\rangle_{b_k} \otimes |1\rangle_{c_k}] \\
&\equiv |1\rangle_{a_k} \otimes |H_\tau^\perp\rangle_{b_k c_k} \\
&\equiv \cos(\omega_k \tau) |E_7\rangle_k - \sin(\omega_k \tau) |E_4\rangle_k \\
&= |E_7\rangle_k - \omega_k \tau |E_4\rangle_k + \mathcal{O}(\tau^2),
\end{aligned} \tag{14}$$

where $|H_\tau^\perp\rangle$ denotes a Hawking state orthogonal to the Hawking state $|H_\tau\rangle$. Since, from a geometrical point of view, ω_k is an angle of rotation in the 8-dimensional Hilbert space \mathcal{H}_k for the k -mode sector, the auxiliary ‘‘time’’ parameter τ is dimensionless. More precisely, τ

is a scale parameter governing the evolution, and $\tau \in [0, 1]$. In particular, for $\tau = 1$ the evolution is supposed to be completed.

The third base, $\{|E''_{\Lambda}(\tau)\rangle_k\}_{\Lambda=0}^7$, is given by:

$$\begin{aligned}
|E''_0\rangle_k &= |0\rangle_{a_k} \otimes |0\rangle_{b_k} \otimes |0\rangle_{c_k} \equiv |E_0\rangle_k \\
|E''_1(\tau)\rangle_k &= \cos\left(\frac{\pi\tau}{2}\right) |0\rangle_{a_k} \otimes |0\rangle_{b_k} \otimes |1\rangle_{c_k} - \sin\left(\frac{\pi\tau}{2}\right) |1\rangle_{a_k} \otimes |0\rangle_{b_k} \otimes |0\rangle_{c_k} \\
&\equiv \cos\left(\frac{\pi\tau}{2}\right) |E_1\rangle_k - \sin\left(\frac{\pi\tau}{2}\right) |E_4\rangle_k \\
&= |E_1\rangle_k - \frac{\pi\tau}{2} |E_4\rangle_k + \mathcal{O}(\tau^2) \\
|E''_2\rangle_k &= |0\rangle_{a_k} \otimes |1\rangle_{b_k} \otimes |0\rangle_{c_k} \equiv |E_2\rangle_k \\
|E''_3\rangle_k &= |0\rangle_{a_k} \otimes |1\rangle_{b_k} \otimes |1\rangle_{c_k} \equiv |E_3\rangle_k \\
|E''_4(\tau)\rangle_k &= \sin\left(\frac{\pi\tau}{2}\right) |0\rangle_{a_k} \otimes |0\rangle_{b_k} \otimes |1\rangle_{c_k} + \cos\left(\frac{\pi\tau}{2}\right) |1\rangle_{a_k} \otimes |0\rangle_{b_k} \otimes |0\rangle_{c_k} \\
&\equiv \cos\left(\frac{\pi\tau}{2}\right) |E_4\rangle_k + \sin\left(\frac{\pi\tau}{2}\right) |E_1\rangle_k \\
&= |E_4\rangle_k + \frac{\pi\tau}{2} |E_1\rangle_k + \mathcal{O}(\tau^2) \\
|E''_5\rangle_k &= |1\rangle_{a_k} \otimes |0\rangle_{b_k} \otimes |1\rangle_{c_k} \equiv |E_5\rangle_k \\
|E''_6\rangle_k &= |1\rangle_{a_k} \otimes |1\rangle_{b_k} \otimes |0\rangle_{c_k} \equiv |E_6\rangle_k \\
|E''_7\rangle_k &= |1\rangle_{a_k} \otimes |1\rangle_{b_k} \otimes |1\rangle_{c_k} \equiv |E_7\rangle_k.
\end{aligned} \tag{15}$$

Now we define (τ -dependent) unitary operators according to the recipe (11) as follows:

$$U'_k(\tau) = \sum_{\Lambda=0}^7 |E'_{\Lambda}(\tau)\rangle_k \langle E_{\Lambda}|_k, \tag{16}$$

$$U''_k(\tau) = \sum_{\Lambda=0}^7 |E''_{\Lambda}(\tau)\rangle_k \langle E'_{\Lambda}(\tau)|_k, \tag{17}$$

and

$$\begin{aligned}
U_k(\tau) &\equiv U''_k(\tau) \cdot U'_k(\tau) = \sum_{\Lambda, \Lambda'=0}^7 |E''_{\Lambda}(\tau)\rangle \langle E'_{\Lambda}(\tau) | E'_{\Lambda'}(\tau) \rangle \langle E_{\Lambda'}| \\
&= \sum_{\Lambda=0}^7 |E''_{\Lambda}(\tau)\rangle \langle E_{\Lambda}|.
\end{aligned} \tag{18}$$

We can easily confirm that for $\tau = 1$ the (explicitly) unitary operators (16), (17) and (18) act according to the rules (3), (6) and (7), respectively. For example, for $U'_k(\equiv U'_k(1))$ we

confirm that

$$\begin{aligned}
U'_k (|q_k\rangle_{a_k} \otimes |O\rangle_{b_k c_k}) &= U'_k \{ [(1 - q_k) |0\rangle_{a_k} + q_k |1\rangle_{a_k}] \otimes |O\rangle_{b_k c_k} \} \\
&= \sum_{\Lambda=0}^7 |E'_\Lambda(1)\rangle_k \langle E_\Lambda|_k [(1 - q_k) |E_0\rangle_k + q_k |E_4\rangle_k] \\
&= (1 - q_k) |E'_0(1)\rangle_k + q_k |E'_4(1)\rangle_k = [(1 - q_k) |0\rangle_{a_k} + q_k |1\rangle_{a_k}] \otimes |H_1\rangle_{b_k c_k} \\
&= |q_k\rangle_{a_k} \otimes |H_1\rangle_{b_k c_k},
\end{aligned} \tag{19}$$

as expected (see (3)).

IV. PHYSICAL PICTURE

Now, let us determine a corresponding (infinitesimal) generator \mathbb{H}_k (“Hamiltonian”) for the unitary evolution operator $U_k(\tau)$, i.e.

$$U_k(\tau) = \mathbb{I}_k - i\tau\mathbb{H}_k + \mathcal{O}(\tau^2). \tag{20}$$

To this end we will utilize expansions (in terms of the “time” parameter τ) of the bases $\{|E'_\Lambda(\tau)\rangle\}$ and $\{|E''_\Lambda(\tau)\rangle\}$ given in (14) and (15), respectively, around the standard base $\{|E_\Lambda\rangle\}$.

To make a comparison to a known case (i.e. to the fermion squeezing operator [9, 18]), let us derive the form of the generator \mathbb{H}'_k for the unitary transformation $U'_k(\tau)$ (16). By virtue of (16), (14) and (20) (with unprimed quantities replaced by primed ones) we get

$$\begin{aligned}
U'_k(\tau) &= \mathbb{I}_k - i\tau i\omega_k (|E_3\rangle_k \langle E_0|_k - |E_0\rangle_k \langle E_3|_k \\
&\quad + |E_7\rangle_k \langle E_4|_k - |E_4\rangle_k \langle E_7|_k) + \mathcal{O}(\tau^2).
\end{aligned} \tag{21}$$

Introducing the identification:

$$\begin{aligned}
|0\rangle_{x_k} \langle 0|_{x_k} &= \hat{x}_k \hat{x}_k^\dagger \\
|0\rangle_{x_k} \langle 1|_{x_k} &= \hat{x}_k \\
|1\rangle_{x_k} \langle 0|_{x_k} &= \hat{x}_k^\dagger \\
|1\rangle_{x_k} \langle 1|_{x_k} &= \hat{x}_k^\dagger \hat{x}_k,
\end{aligned} \tag{22}$$

for the modes $x_k = a_k, b_k, c_k$ ($k = 1, 2, \dots, N$), from (21) we obtain a representation of the operator \mathbb{H}'_k in the Fock space, i.e.

$$\hat{\mathbb{H}}'_k = i\omega_k \hat{b}_k^\dagger \hat{c}_k^\dagger + \text{H.c.}, \tag{23}$$




Figure 1. A Feynman-like diagram/line (the S-shaped thick line with an arrow) depicts the entire qubit transport process U_k , i.e. a composition of the Hawking particle pair creation U'_k outside the BH and of particle pair annihilation U''_k inside the BH, in the spirit of a tunneling phenomenon.

where “H.c.” means Hermitian conjugate. The operator (23) is known as a two-mode fermion squeezing operator, responsible for creation of fermionic pairs in the framework of the Hawking effect (see e.g. Sect. 5.2 in [10]).

Let us now derive the generator \mathbb{H}_k , and its Fock space counterpart $\hat{\mathbb{H}}_k$, for the entire evolution $U_k(\tau)$. By virtue of (18), (15) and (20) we obtain

$$U_k(\tau) = \mathbb{I}_k - i\tau \frac{i\pi}{2} (|E_1\rangle_k \langle E_4|_k - |E_4\rangle_k \langle E_1|_k) + \mathcal{O}(\tau^2). \quad (24)$$

Implementing the identification (22) we obtain the corresponding Fock space generator (“Hamiltonian”)

$$\hat{\mathbb{H}}_k = \frac{i\pi}{2} \hat{a}_k \hat{b}_k \hat{b}_k^\dagger \hat{c}_k^\dagger + \text{H.c.} \quad (25)$$

One should note that the operator $\hat{\mathbb{H}}_k$ is 4-linear, which should be contrasted with a bilinear structure of the squeezing operator (23) and a trilinear structure of the operator discussed in the context of the Hawking effect in [15]. According to (9) the total “Hamiltonian” is $\hat{\mathbb{H}} = \sum_{k=1}^N \hat{\mathbb{H}}_k$.

Coming back to the global description of the BH unitary evolution, we would like to draw the reader’s attention to a possible interpretation, depicted in Fig. 1. Namely, Fig. 1, in an intuitive way, presents the entire unitary process U_k of qubit transfer from a BH to the outgoing radiation as a composition of the two processes, U'_k and U''_k , i.e. the Hawking particle pair creation outside the BH and later particle pair annihilation inside the BH, respectively. A Feynman-like diagram/line depicts the transfer of a qubit in the spirit of a tunneling phenomenon.

Taking into account “reversibility” of unitary processes, the Reader could rightfully expect that at some moment of the evolution the “reverse” process of a given one should also occur, and a “reverse diagram” to the one presented in Fig. 1 should also appear. The unitary transformations introduced in Section III are actually rotations in 8-dimensional complex vector spaces with rotation angles proportional to the “time” parameter τ . Then, if we start from a vector corresponding to an a_k -mode and “rotate” it onto a c_k -mode, and next we stop the evolution, we effectively swap one particle for another. If we instead further continued the evolution we would return to the a_k -mode back. Therefore, we have to assume that the proposed evolution is only valid within a limited period of “time” τ , namely $\tau \in [0, 1]$. That limitation could be justified by the assumption that the evolution is “effective” rather than “fundamental” in the sense that its temporary form is determined by the current structure of spacetime and the distribution of matter. Then, when a_k -modes become transformed into c_k -modes, the situation changes and the evolution in the proposed form switches off. Moreover, one should also note that the whole result is influenced by the initial state of the system, i.e. we start with modes only inside the black hole, while all other modes are in the vacuum state. Such an assumption corresponds to the idealized (model) situation assuming that in the beginning of the evolution we only deal with a (single) black hole (represented by a_k -modes occupied) and empty spacetime outside the black hole (c_k -modes empty).

V. FINAL REMARKS

Primarily inspired by a recent paper of Osuga and Page [16], in particular by their Eq. (3.3), essentially the same as Eq. (3.3) in [1] (accidental coincidence of the numbers of the equations!), we have proposed a unitary toy model of BH evaporation, which is an extension of the model introduced in [3]. By virtue of the construction the model is explicitly unitary and it describes transport of qubits from a BH to the outgoing radiation. For interpretational simplicity and a more direct relation to particle language, we have decided to formulate our qubit model in terms of fermions. As a byproduct of our construction, for possible reference to other models of the Hawking effect and BH evaporation, besides the global version of the evolution operator, we have determined its infinitesimal form (“Hamiltonian”). In turn the global evolution, involving the Hawking creation as well as latter annihilation inside the BH,

can intuitively be interpreted as a tunneling phenomenon as depicted in Fig. 1.

-
- [1] Ahmed Almheiri, Donald Marolf, Joseph Polchinski, Douglas Stanford, and James Sully. An Apologia for Firewalls. *JHEP*, 09:018, 2013.
 - [2] Steven G. Avery. Qubit Models of Black Hole Evaporation. *JHEP*, 01:176, 2013.
 - [3] Bogusław Broda. Unitary toy qubit transport model for black hole evaporation. *Eur. Phys. J. C*, 80(5):418, may 2020.
 - [4] Sumanta Chakraborty and Kinjalk Lochan. Black Holes: Eliminating Information or Illuminating New Physics? *Universe*, 3(3):55, 2017.
 - [5] Steven B. Giddings. Black holes, quantum information, and unitary evolution. *Phys. Rev. D*, 85:124063, Jun 2012.
 - [6] Steven B. Giddings. Models for unitary black hole disintegration. *Phys. Rev. D*, 85:044038, Feb 2012.
 - [7] Steven B. Giddings and Yinbo Shi. Quantum information transfer and models for black hole mechanics. *Phys. Rev. D*, 87:064031, Mar 2013.
 - [8] D. Harlow. Jerusalem lectures on black holes and quantum information. *Rev. Mod. Phys.*, 88:015002, Feb 2016.
 - [9] Faqir C. Khanna, Adolfo P. C. Malbouisson, Jorge M. C. Malbouisson, and Ademir E. Santana. *Thermal quantum field theory. Algebraic aspects and applications*. Hackensack, NJ: World Scientific, 2009.
 - [10] Robert B. Mann. *Black Holes: Thermodynamics, Information, and Firewalls*. SpringerBriefs in Physics. Springer, 2015.
 - [11] Donald Marolf. The Black Hole information problem: past, present, and future. *Rept. Prog. Phys.*, 80(9):092001, 2017.
 - [12] Samir D. Mathur. The Information paradox: A Pedagogical introduction. *Class. Quant. Grav.*, 26:224001, 2009.
 - [13] Samir D. Mathur. What Exactly is the Information Paradox? *Lect. Notes Phys.*, 769:3–48, 2009.
 - [14] Samir D. Mathur and Christopher J. Plumberg. Correlations in Hawking radiation and the infall problem. *JHEP*, 09:093, 2011.

- [15] Paul D. Nation and Miles P. Blencowe. The Trilinear Hamiltonian: A Zero Dimensional Model of Hawking Radiation from a Quantized Source. *New Journal of Physics*, 12:095013, 2010.
- [16] Kento Osuga and Don N. Page. Qubit transport model for unitary black hole evaporation without firewalls. *Phys. Rev. D*, 97:066023, Mar 2018.
- [17] Joseph Polchinski. The Black Hole Information Problem. In *Proceedings, Theoretical Advanced Study Institute in Elementary Particle Physics: New Frontiers in Fields and Strings (TASI 2015): Boulder, CO, USA, June 1–26, 2015*, pages 353–397, 2017.
- [18] K. Svozil. Squeezed fermion states. *Phys. Rev. Lett.*, 65:3341–3343, Dec 1990.