

AS-REGULAR ALGEBRAS FROM ACYCLIC SPHERICAL HELICES

SHINNOSUKE OKAWA AND KAZUSHI UEDA

ABSTRACT. We discuss a construction of AS-regular algebras from acyclic spherical helices, generalizing works of Bondal and Polishchuk [BP93] and Van den Bergh [VdB11].

Let $\mathcal{C}$ be an idempotent-complete pretriangulated dg category over a field $\mathbf{k}$. See e.g. [Kel06] and references therein for basic definitions on dg categories. For $X, Y \in \mathfrak{Ob}(\mathcal{C})$, we write the dg vector space of morphisms in $\mathcal{C}$ and its i -th cohomology as $\mathrm{hom}(X, Y)$ and $\mathrm{Hom}^i(X, Y)$ respectively.

A $\mathcal{C}$ -module is a dg functor $\mathcal{C}^{\mathrm{op}} \rightarrow \mathrm{Mod} \mathbf{k}$ to the dg category of dg vector spaces. A dg functor $F: \mathcal{C} \rightarrow \mathcal{C}'$ will be identified with its *graph bimodule* $\Gamma_F \in \mathrm{Mod}(\mathcal{C}^{\mathrm{op}} \otimes \mathcal{C}')$.

Assume that $\mathcal{C}$ is *proper* in the sense that $\mathrm{hom}(X, Y) \in \mathrm{per} \mathbf{k}$ for any $X, Y \in \mathfrak{Ob}(\mathcal{C})$. Assume further that $\mathcal{C}$ has a Serre functor $\mathbb{S}$ in the sense of [BK89]. The graph of the Serre functor is given by

$$(1) \quad \mathrm{hom}(Y, \mathbb{S}(X)) = \mathrm{hom}(X, Y)^*,$$

where $(-)^* := \mathrm{hom}_{\mathbf{k}}(-, \mathbf{k}): (\mathrm{Mod} \mathbf{k})^{\mathrm{op}} \rightarrow \mathrm{Mod} \mathbf{k}$.

A prototypical example of an idempotent-complete proper pretriangulated dg category admitting a Serre functor is the category of perfect complexes on a proper Gorenstein scheme X . The Serre functor is the tensor product with the canonical bundle followed by the shift by the dimension of X ; $\mathbb{S}(-) = (-) \otimes \omega_X[\dim X]$.

Let d be a natural number. An object S of $\mathcal{C}$ is said to be *spherical of dimension d* if $\mathbb{S}(S) \simeq S[d]$ and

$$(2) \quad \mathrm{Hom}^i(S, S) \cong \begin{cases} \mathbf{k} & i = 0, d, \\ 0 & \text{otherwise.} \end{cases}$$

The *spherical twist* along S is the functorial cone T_S over the evaluation morphism $\mathrm{ev}: \mathrm{hom}(S, -) \otimes S \rightarrow \mathrm{id}_{\mathcal{C}}$. It is an autoequivalence of the dg category $\mathrm{per} \mathcal{C}$ of perfect $\mathcal{C}$ -modules, whose inverse is given by the *dual twist* $T_S^{\vee}$ defined as the cone over the coevaluation morphism $\mathrm{coev}: \mathrm{id}_{\mathcal{C}} \rightarrow \mathrm{hom}(-, S)^{\vee} \otimes S$ shifted by -1 [ST01].

Given a sequence $\mathbf{S} = (S_i)_{i=1}^{\ell}$ of spherical objects, let $\mathcal{B} = \mathcal{B}_{\mathbf{S}}$ be the full subcategory of $\mathcal{C}$ consisting of $\{S_i\}_{i=1}^{\ell}$. We identify $\mathrm{per} \mathcal{B}$ with the full idempotent-complete pretriangulated subcategory of $\mathcal{C}$ generated by $\mathcal{B}$.

Definition 1. A sequence $\mathbf{S} = (S_i)_{i=1}^{\ell}$ of spherical objects is a *spherical collection* if the Serre functor of $\mathcal{C}$ preserves $\mathrm{per} \mathcal{B}$ and acts as the shift functor there; $\mathbb{S}_{\mathcal{C}}|_{\mathrm{per} \mathcal{B}} \simeq [d]$.

The number ℓ of objects in the collection is called the *length* of the collection.

The *directed subcategory* $\mathcal{A} = \mathcal{A}_{\mathbf{S}}$ of $\mathcal{B}$ is the dg category whose set of objects is $\{E_i\}_{i=1}^{\ell}$ and whose dg vector spaces of morphisms are given by

$$(3) \quad \mathrm{hom}_{\mathcal{A}}(E_i, E_j) := \begin{cases} \mathrm{hom}_{\mathcal{B}}(S_i, S_j) & i < j, \\ \mathbf{k} \cdot \mathrm{id}_{E_i} & i = j, \\ 0 & i > j. \end{cases}$$

The differentials and the compositions of morphisms in $\mathcal{A}$ are inherited from that of $\mathcal{B}$ in the obvious way.

Since the collection $\mathbf{S}$ is spherical, the $\mathcal{A}$ -bimodule $\mathcal{B}/\mathcal{A}$, defined as the cone of the natural morphism $\mathcal{A} \rightarrow \mathcal{B}$ of $\mathcal{A}$ -bimodules, is isomorphic to the shift $\mathcal{A}^*[-d]$ of the graph $\mathcal{A}^* := \mathrm{hom}_{\mathbf{k}}(\mathcal{A}, \mathbf{k})$ of the Serre functor. In other words, the category $\mathcal{B}$ is a *noncommutative anticanonical divisor* of $\mathcal{A}$ in the sense of [Sei17].

The sequence $(E_i)_{i=1}^\ell$ is a full exceptional collection of per $\mathcal{A}$. The existence of a full exceptional collection implies that per $\mathcal{A}$ is proper and *smooth* (in the sense that the diagonal bimodule is perfect), so that per $\mathcal{A}$ has a Serre functor.

Let $\iota_*: \text{per } \mathcal{B} \rightarrow \text{per } \mathcal{A}$ be the dg functor sending a $\mathcal{B}$ -module to the same module considered as an $\mathcal{A}$ -module by the embedding $\mathcal{A} \hookrightarrow \mathcal{B}$. The graph of the left adjoint functor $\iota^*: \text{per } \mathcal{A} \rightarrow \text{per } \mathcal{B}$ is $\mathcal{B}$ considered as an $\mathcal{A}^{\text{op}} \otimes \mathcal{B}$ -module. The right adjoint functor is given by $\iota^! = \mathbb{S}_{\mathcal{B}} \circ \iota^* \circ \mathbb{S}_{\mathcal{A}}^{-1} \simeq \iota^* \circ \mathbb{S}_{\mathcal{A}}^{-1}[d]$.

The *dual twist* $T_{\iota_*}^\vee$ along ι_* is defined as the shift by -1 of the cone over the unit $\text{id}_{\mathcal{A}} \rightarrow \iota_* \circ \iota^*$ of the adjunction. The graph of the functor $\iota_* \circ \iota^*$ is $\mathcal{B}$ considered as an $\mathcal{A}$ -bimodule, so that the graph of the dual twist is the $\mathcal{A}$ -bimodule $\mathcal{B}/\mathcal{A}[-1]$, which is the graph of $\mathbb{S}[-d-1]$. The *twist* T_{ι_*} along ι_* , defined as the cone over the counit $\iota_* \circ \iota^! \rightarrow \text{id}_{\mathcal{A}}$ of the adjunction, is right adjoint to $T_{\iota_*}^\vee$ by [AL17, Proposition 5.3], and hence is quasi-isomorphic to $\mathbb{S}^{-1}[d+1]$.

The *dual cotwist* $C_{\iota_*}^\vee$ along ι_* , defined as the cone over the counit $\iota^* \circ \iota_* \rightarrow \text{id}_{\mathcal{B}}$, is described explicitly as the (iterated) cone of

$$(4) \quad \text{hom}(S_\ell, -) \otimes \text{hom}(S_{\ell-1}, S_\ell) \otimes \cdots \otimes \text{hom}(S_1, S_2) \otimes S_1 \rightarrow \cdots \\ \rightarrow \bigoplus_{1 \leq i_1 < i_2 \leq \ell} \text{hom}(S_{i_2}, -) \otimes \text{hom}(S_{i_1}, S_{i_2}) \otimes S_{i_1} \rightarrow \bigoplus_{i=1}^\ell \text{hom}(S_i, -) \otimes S_i \rightarrow \text{id}_{\mathcal{B}},$$

which is quasi-isomorphic to the composition of spherical twists;

$$(5) \quad C_{\iota_*}^\vee \simeq T_{S_1} \circ T_{S_2} \circ \cdots \circ T_{S_\ell}.$$

Since both T_{ι_*} and $C_{\iota_*}^\vee$ are equivalences, the functor ι_* is spherical in the sense of [AL17, Theorem 1.1]. We expect that ι_* has a *relative Calabi–Yau structure* in the sense of [BD19], which is a slight strengthening of an identification of the spherical twist with a shift of the inverse Serre functor.

For $1 \leq i \leq \ell - 1$, the *right mutation* of a spherical collection $\mathbf{S} = (S_j)_{j=1}^\ell$ at position i is defined by

$$(6) \quad R_i(\mathbf{S}) := (S_1, \dots, S_{i-1}, S_{i+1}, T_{S_{i+1}}^\vee S_i, S_{i+2}, \dots, S_\ell).$$

The inverse operation is the *left mutation* defined by

$$(7) \quad L_i(\mathbf{S}) := (S_1, \dots, S_{i-1}, T_{S_i} S_{i+1}, S_i, S_{i+2}, \dots, S_\ell).$$

The *left mutation* $L_i(\mathcal{A})$ of $\mathcal{A}$ at position i is the full subcategory of per $\mathcal{A}$ consisting of E_j for $j \neq i+1$ and

$$(8) \quad L_{E_i} E_{i+1} := \text{Cone} \left(\text{hom}(E_i, E_{i+1}) \otimes E_i \xrightarrow{\text{ev}} E_{i+1} \right).$$

The passage to the directed subcategory commutes with mutations; $\mathcal{A}_{L_i(\mathbf{S})} \simeq L_i(\mathcal{A}_{\mathbf{S}})$.

The *helix* generated by an exceptional collection $(E_i)_{i=1}^\ell$ is the sequence $(E_i)_{i \in \mathbb{Z}}$ of exceptional objects of per $\mathcal{A}$ satisfying

$$(9) \quad E_{i-\ell} = L_{E_{i-\ell+1}} \circ L_{E_{i-\ell+2}} \circ \cdots \circ L_{E_{i-1}}(E_i)[-d-1]$$

for any $i \in \mathbb{Z}$. The shift is chosen in such a way that $E_{i-\ell} \cong \mathbb{S}(E_i)[-d-1]$, so that if per $\mathcal{A} \simeq D^b \text{coh } X$ for a smooth projective variety X of dimension $d+1$, then one has $E_{i-\ell} \cong E_i \otimes \omega_X$. The length ℓ of the exceptional collection generating the helix is called the *period* of the helix. For any $i \in \mathbb{Z}$, the exceptional collection $(E_j)_{j=i}^{i+\ell-1}$ consisting of ℓ consecutive objects in the helix is called a *foundation* of the helix. We say that a helix $(E_i)_{i \in \mathbb{Z}}$ is *acyclic* if $\text{Hom}^{\neq 0}(E_i, E_j) = 0$ for any $-\infty < i < j < \infty$.

Given an invertible $\mathcal{A}$ -bimodule θ , the *tensor algebra* over $\mathcal{A}$ is defined by

$$(10) \quad T_{\mathcal{A}}(\theta) := \bigoplus_{i=0}^{\infty} \theta^{\otimes_{\mathcal{A}} i} \\ (11) \quad = \mathcal{A} \oplus \theta \oplus (\theta \otimes_{\mathcal{A}} \theta) \oplus \cdots \\ (12) \quad \simeq \bigoplus_{i=0}^{\infty} \bigoplus_{j,k=1}^{\ell} \text{hom}(E_j, \theta^i(E_k)).$$

In addition to the cohomological grading coming from that of $\mathcal{A}$ and θ , the tensor algebra $T_{\mathcal{A}}(\theta)$ has an additional grading such that $\theta^{\otimes_{\mathcal{A}} i}$ has degree i . The positively graded part $T_{\mathcal{A}}(\theta)^+ := \bigoplus_{i=1}^{\infty} \theta^{\otimes_{\mathcal{A}} i} \simeq \theta \otimes_{\mathcal{A}} T_{\mathcal{A}}(\theta)$ is the kernel (the cone shifted by -1) of the natural morphism $\sigma^{\sharp}: T_{\mathcal{A}}(\theta) \rightarrow \mathcal{A}$.

The category $\text{per } T_{\mathcal{A}}(\theta)$ is a generalization of the derived category of coherent sheaves on the total space of a line bundle, and the pull-back $\sigma_*: \text{per } \mathcal{A} \rightarrow \text{per } T_{\mathcal{A}}(\theta)$ along $\sigma^{\sharp}$ is a generalization of the push-forward along the zero section. Its left adjoint $\sigma^*: \text{per } T_{\mathcal{A}}(\theta) \rightarrow \text{per } \mathcal{A}$ is given by tensoring $\mathcal{A}$ over $T_{\mathcal{A}}(\theta)$. Since the graph of $\sigma^* \circ \sigma_*$ is

$$(13) \quad \text{Cone}(\theta \otimes_{\mathcal{A}} T_{\mathcal{A}}(\theta) \rightarrow T_{\mathcal{A}}(\theta)) \otimes_{T_{\mathcal{A}}(\theta)} \mathcal{A} \simeq \text{Cone}(\theta \rightarrow \mathcal{A}),$$

the dual cotwist along σ_* is the autoequivalence $\theta[1]$ of $\text{per } \mathcal{A}$. See also [Seg18] for a related construction.

The push-forward $\pi^* := (-) \otimes_{\mathcal{A}} T_{\mathcal{A}}(\theta): \text{per } \mathcal{A} \rightarrow \text{per } T_{\mathcal{A}}(\theta)$ along the structure morphism $\pi^{\sharp}: \mathcal{A} \rightarrow T_{\mathcal{A}}(\theta)$ of the tensor algebra is a generalization of the pull-back along the projection to the base space. For any pair (E, F) of objects of $\text{per } \mathcal{A}$, one has

$$(14) \quad \text{hom}(\sigma_* E, \pi^* F) \simeq \text{hom}(\pi^* E \otimes_{T_{\mathcal{A}}(\theta)} \text{Cone}(\theta \otimes_{\mathcal{A}} T_{\mathcal{A}}(\theta) \rightarrow T_{\mathcal{A}}(\theta)), \pi^* F)$$

$$(15) \quad \simeq \text{hom}(\pi^* E, \pi^* F \otimes_{T_{\mathcal{A}}(\theta)} \text{Cone}(T_{\mathcal{A}}(\theta) \rightarrow \theta^{-1} \otimes_{\mathcal{A}} T_{\mathcal{A}}(\theta))[-1])$$

$$(16) \quad \simeq \text{hom}(\pi^* E, \sigma_*(\theta^{-1}(F)))[-1]$$

$$(17) \quad \simeq \text{hom}(E, \theta^{-1}(F))[-1].$$

The dg category $\text{gr } T_{\mathcal{A}}(\theta)$ of graded perfect $T_{\mathcal{A}}(\theta)$ -modules is a generalization of the derived category of equivariant coherent sheaves on the total space of a line bundle with respect to the $\mathbb{G}_m$ -action by fiberwise dilation. The dg $\mathbb{Z}$ -algebra $A = \bigoplus_{i,j=-\infty}^{\infty} A_{ij}$ defined by

$$(18) \quad A_{i+k\ell, j+m\ell} = \text{hom}(\theta^m(E_j), \theta^k(E_i))$$

for $1 \leq i, j \leq \ell$ and $-\infty < m \leq k < \infty$ is ℓ -periodic, and the 1-periodic dg $\mathbb{Z}$ -algebra $A' = \bigoplus_{i,j=-\infty}^{\infty} A'_{ij}$ defined by

$$(19) \quad A'_{km} = \bigoplus_{i=(k-1)\ell+1}^{k\ell} \bigoplus_{j=(m-1)\ell+1}^{m\ell} A_{ij}$$

is the dg $\mathbb{Z}$ -algebra associated to the $\mathbb{Z}$ -graded algebra $T_{\mathcal{A}}(\theta)$. One has quasi-equivalences

$$(20) \quad \text{gr } T_{\mathcal{A}}(\theta) \simeq \text{mod } A' \simeq \text{mod } A$$

of dg categories.

Let $(F_i)_{i=\ell}^1$ be the exceptional collection in $\text{per } \mathcal{A}$ right dual to $(E_i)_{i=1}^{\ell}$ so that $\text{hom}(E_i, F_j) \simeq \mathbf{k}$ if $i = j$ and 0 otherwise [Bon89, Section 7]. Then E_i (resp. F_i) represents the i -th projective (resp. simple) $\mathcal{A}$ -module, and $\pi^* E_i$ (resp. $\sigma_* F_i$) represents the i -th projective (resp. simple) $T_{\mathcal{A}}(\theta)$ -module.

The dg algebra $T_{\mathcal{A}}(\theta)$ in the case $\theta = \mathbb{S}^{-1}[d+1]$ is known as the *derived $(d+1)$ -preprojective algebra*, which is a model of the *trivial Calabi–Yau completion* [Kel11];

$$(21) \quad \Pi_{d+1}(\mathcal{A}) := T_{\mathcal{A}}(\mathbb{S}^{-1}[d+1]).$$

It is given explicitly as

$$(22) \quad \bigoplus_{i=0}^{\infty} \bigoplus_{j,k=1}^{\ell} \text{hom}(E_j, \mathbb{S}^{-i}(E_k)[i(d+1)]) \simeq \bigoplus_{j=1}^{\ell} \bigoplus_{k=1}^{\ell} \text{hom}(E_j, E_k),$$

and also known as the *rolled-up helix dg algebra*. The dg $\mathbb{Z}$ -algebra A in this case will be called the *helix dg algebra*.

For $1 \leq i, j \leq \ell$, one has

$$(23) \quad \text{hom}(\sigma_* F_i, \pi^* E_j) \simeq \text{hom}(F_i, \mathbb{S}(E_j)[-d-1])[-1]$$

$$(24) \quad \simeq \text{hom}(E_j, F_i)[-d-2]$$

$$(25) \quad \simeq \begin{cases} \mathbf{k}[-d-2] & i = j, \\ 0 & \text{otherwise.} \end{cases}$$

An object of $\text{gr } T_{\mathcal{A}}(\theta)$ is said to be a *torsion* if the grading is bounded. Let $\text{tor } T_{\mathcal{A}}(\theta)$ be the full subcategory of $\text{gr } T_{\mathcal{A}}(\theta)$ consisting of torsion modules, and $\text{qgr } T_{\mathcal{A}}(\theta) := \text{gr } T_{\mathcal{A}}(\theta) / \text{tor } T_{\mathcal{A}}(\theta)$ be the dg quotient.

Proposition 2. *One has an equivalence $\text{qgr } T_{\mathcal{A}}(\theta) \simeq \text{per } \mathcal{A}$.*

Proof. One has a semiorthogonal decomposition

$$(26) \quad \text{tor } T_{\mathcal{A}}(\theta) = \langle \mathcal{S}_i \rangle_{i \in \mathbb{Z}},$$

where $\mathcal{S}_i \simeq \text{per } \mathcal{A}$ is the admissible subcategory consisting of graded modules concentrated in degree i . One has

$$(27) \quad \text{gr } T_{\mathcal{A}}(\theta) = \langle \dots, \mathcal{S}_{-2}, \mathcal{S}_{-1}, (\text{gr } T_{\mathcal{A}}(\theta))_{\geq 0} \rangle$$

just as in [Orl09, Lemma 14], where $(\text{gr } T_{\mathcal{A}}(\theta))_{\geq 0}$ is the full subcategory of $\text{gr } T_{\mathcal{A}}(\theta)$ consisting of non-negatively graded modules. It follows from the quasi-isomorphism

$$(28) \quad \mathcal{A} \simeq \text{Cone}(T_{\mathcal{A}}(\theta)^+ \rightarrow T_{\mathcal{A}}(\theta))$$

of graded $T_{\mathcal{A}}(\theta)$ -modules that the free module $T_{\mathcal{A}}(\theta)$ is right orthogonal to $\langle \mathcal{S}_i \rangle_{i \geq 0}$, and that its shift $T_{\mathcal{A}}(\theta)(i)$ for any $i < 0$ is in the full pretriangulated subcategory generated by $T_{\mathcal{A}}(\theta)$ and $\langle \mathcal{S}_i \rangle_{i \geq 0}$. Hence $T_{\mathcal{A}}(\theta)$ generates the right orthogonal to $\langle \mathcal{S}_i \rangle_{i \geq 0}$ in $(\text{gr } T_{\mathcal{A}}(\theta))_{\geq 0}$, so that $\text{qgr } T_{\mathcal{A}}(\theta)$ is equivalent to the full subcategory $\langle T_{\mathcal{A}}(\theta) \rangle$ of $\text{gr } T_{\mathcal{A}}(\theta)$, which in turn is equivalent to $\text{per } \mathcal{A}$. $\square$

Definition 3. A *spherical helix* generated by a spherical collection $(S_i)_{i=1}^{\ell}$ is a sequence $(S_i)_{i \in \mathbb{Z}}$ of spherical objects of $\mathcal{C}$ satisfying

$$(29) \quad S_{i-\ell} = T_{S_{i-\ell+1}} \circ T_{S_{i-\ell+2}} \circ \dots \circ T_{S_{i-1}}(S_i)[-d-1]$$

for any $i \in \mathbb{Z}$.

The length ℓ of the spherical collection generating the spherical helix is called the *period* of the spherical helix. For any $i \in \mathbb{Z}$, the spherical collection $(S_j)_{j=i}^{i+\ell-1}$ consisting of ℓ consecutive objects in the spherical helix is called a *foundation* of the spherical helix $\mathcal{H}$.

Definition 4. A spherical helix $(S_i)_{i \in \mathbb{Z}}$ is *acyclic* if $\text{Hom}^{\neq 0}(S_i, S_j) = 0$ for any $-\infty < i < j < \infty$.

One has

$$(30) \quad \text{hom}(S_i, S_j) \simeq \text{hom}(\iota^* E_i, \iota^* E_j)$$

$$(31) \quad \simeq \text{hom}(E_i, \iota_* \iota^* E_j)$$

$$(32) \quad \simeq \text{hom}(E_i, \text{Cone}(E_{j-\ell} \rightarrow E_j)),$$

which immediately yields the following:

Theorem 5. *If the spherical helix $(S_i)_{i \in \mathbb{Z}}$ is acyclic, then the helix $(E_i)_{i \in \mathbb{Z}}$ is acyclic.*

Recall that a connected $\mathbb{Z}$ -algebra $A = \bigoplus_{i,j=-\infty}^{\infty} A_{ij}$ over $\mathbf{k}$ is said to be *AS-Gorenstein* if

$$(33) \quad \sum_{j,k=-\infty}^{\infty} \dim \text{Hom}^k(S_{i,A}, P_{j,A}) = 1$$

for any $i \in \mathbb{Z}$, where $S_{i,A}$ and $P_{i,A}$ are the i -th simple and projective A -modules respectively. It follows from (25) that the acyclicity of the helix $(E_i)_{i \in \mathbb{Z}}$ implies the AS-Gorenstein property of the helix $\mathbb{Z}$ -algebra A . Since $\mathcal{A}$ is smooth and θ is perfect, the tensor algebra $T_{\mathcal{A}}(\theta)$ is smooth, so

that A has finite global dimension (see e.g. [Kel11, Theorem 4.8]). The Hilbert polynomial of A is determined by the projective resolutions of the simples.

Acknowledgement: S. O. was partially supported by JSPS Grant-in-Aid for Scientific Research No.16H05994, 16H02141, 16H06337, 18H01120, 20H01797. K. U. was partially supported by JSPS Grant-in-Aid for Scientific Research No.16H03930.

REFERENCES

- [AL17] Rina Anno and Timothy Logvinenko, *Spherical DG-functors*, J. Eur. Math. Soc. (JEMS) **19** (2017), no. 9, 2577–2656. MR 3692883
- [BD19] Christopher Brav and Tobias Dyckerhoff, *Relative Calabi-Yau structures*, Compos. Math. **155** (2019), no. 2, 372–412. MR 3911626
- [BK89] A. I. Bondal and M. M. Kapranov, *Representable functors, Serre functors, and reconstructions*, Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), no. 6, 1183–1205, 1337. MR MR1039961 (91b:14013)
- [Bon89] A. I. Bondal, *Representations of associative algebras and coherent sheaves*, Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), no. 1, 25–44. MR MR992977 (90i:14017)
- [BP93] A. I. Bondal and A. E. Polishchuk, *Homological properties of associative algebras: the method of helices*, Izv. Ross. Akad. Nauk Ser. Mat. **57** (1993), no. 2, 3–50. MR 1230966 (94m:16011)
- [Kel06] Bernhard Keller, *On differential graded categories*, International Congress of Mathematicians. Vol. II, Eur. Math. Soc., Zürich, 2006, pp. 151–190. MR MR2275593 (2008g:18015)
- [Kel11] ———, *Deformed Calabi-Yau completions*, J. Reine Angew. Math. **654** (2011), 125–180, With an appendix by Michel Van den Bergh. MR 2795754
- [Orl09] Dmitri Orlov, *Derived categories of coherent sheaves and triangulated categories of singularities*, Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II, Progr. Math., vol. 270, Birkhäuser Boston Inc., Boston, MA, 2009, pp. 503–531. MR 2641200 (2011c:14050)
- [Seg18] Ed Segal, *All autoequivalences are spherical twists*, Int. Math. Res. Not. IMRN (2018), no. 10, 3137–3154. MR 3805198
- [Sei17] Paul Seidel, *Fukaya A_∞ -structures associated to Lefschetz fibrations. II*, Algebra, geometry, and physics in the 21st century, Progr. Math., vol. 324, Birkhäuser/Springer, Cham, 2017, pp. 295–364. MR 3727564
- [ST01] Paul Seidel and Richard Thomas, *Braid group actions on derived categories of coherent sheaves*, Duke Math. J. **108** (2001), no. 1, 37–108. MR MR1831820 (2002e:14030)
- [VdB11] Michel Van den Bergh, *Noncommutative quadrics*, International Mathematics Research Notices. IMRN (2011), no. 17, 3983–4026. MR MR2836401 (2012m:14004)

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY, MACHIKANNEYAMA 1-1, TOYONAKA, OSAKA, 560-0043, JAPAN.

E-mail address: okawa@math.sci.osaka-u.ac.jp

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, 3-8-1 KOMABA, MEGURO-KU, TOKYO, 153-8914, JAPAN.

E-mail address: kazushi@ms.u-tokyo.ac.jp