

Expressions for joint moments of elliptical distributions

Baishuai Zuo^a, Chuancun Yin^{a,*}, Narayanaswamy Balakrishnan^{b,*}

^a *School of Statistics, Qufu Normal University, Qufu, Shandong 273165, China*

^b *Department of Mathematics and Statistics, McMaster University, Hamilton, Ontario, Canada*

Abstract

Inspired by Stein's lemma, we derive two expressions for the joint moments of elliptical distributions. We use two different methods to derive $E[X_1^2 f(\mathbf{X})]$ for any measurable function f satisfying some regularity conditions. Then, by applying this result, we obtain new formulae for expectations of product of normally distributed random variables, and also present simplified expressions of $E[X_1^2 f(\mathbf{X})]$ for multivariate Student- t , logistic and Laplace distributions.

Keywords:

Multivariate elliptical distributions; Multivariate normal distribution; Joint moments; Stein's lemma

1. Introduction and motivation

Stein's (1981) lemma discusses the determination of $Cov(X, h(Y))$ for a bivariate normal random vector (X, Y) , where h is any differentiable function with finite extensions $E|h'(Y)|$. Inspired by the original work of Stein, several and generalizations have appeared in the literature. Liu (1994) generalized the lemma to the multivariate normal case, while Landsman (2006) showed that Stein's type lemma also holds when (X, Y) is distributed as bivariate elliptical. This result got extended to multivariate elliptical vectors

*Corresponding author.

Email addresses: ccyin@qfnu.edu.cn (Chuancun Yin), bala@mcmaster.ca (Narayanaswamy Balakrishnan)

by Landsman and Neslehova (2008); see also Landsman et al. (2013) for a simple proof. Recently, Shushi (2018) derived the multivariate Stein's lemma for truncated elliptical random vectors.

In this work, we derive expressions of the joint moments $E[X_1^2 f(\mathbf{X})]$ for any measurable function f satisfying some regularity conditions. In particular, we obtain new formulae for the expectation of product of normally distributed random variables, and also present expressions of $E[X_1^2 f(\mathbf{X})]$ for multivariate Student- t , logistic and Laplace distributions.

The rest of the paper is organized as follows. Section 2 reviews some definitions and properties of the family of elliptical distributions. Section 3 presents explicit expressions of joint moments of elliptical distributions by a direct method. Section 4 derives these joint moments by the use of Stein's lemma. Section 5 shows the equivalence of the two expressions under the condition that the scale matrix is positive definite. Section 6 presents expression for expectation of products of correlated normal variables. Section 7 presents simplified results for the special cases of multivariate Student- t , logistic and Laplace distributions as illustrative examples of the general result established here. Finally, Section 8 gives some concluding remarks.

2. Family of elliptical distributions

Elliptical distributions are generalizations of multivariate normal distribution and possess many novel tractable properties, in addition to allowing fat tails with suitably chosen kernels. This class of distributions was first introduced by Kelker (1970), and has been widely discussed in detail by Fang et al. (1990), and Kotz et al. (2000). A $n \times 1$ random vector $X = (X_1, \dots, X_n)^T$ is said to have an elliptically symmetric distribution if

its characteristic function has the form

$$E[\exp(it^T \mathbf{X})] = e^{it^T \boldsymbol{\mu}} \phi\left(\frac{1}{2} \mathbf{t}^T \boldsymbol{\Sigma} \mathbf{t}\right)$$

for all $\mathbf{t} \in \mathbb{R}^n$, denoted $\mathbf{X} \sim E_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \phi)$, where ϕ is called the characteristic generator with $\phi(0) = 1$, $\boldsymbol{\mu}$ (n -dimensional vector) is the location parameter, and $\boldsymbol{\Sigma}$ ($n \times n$ matrix with $\boldsymbol{\Sigma} \geq \mathbf{0}$) is the dispersion matrix (or scale matrix). The mean vector $E(\mathbf{X})$ (if it exists) coincides with the location vector and the covariance matrix $\text{Cov}(\mathbf{X})$ (if it exists) is $-\phi'(0)\boldsymbol{\Sigma}$. The generator of the multivariate normal distribution, for example, is given by $\phi(u) = \exp(-u)$.

In general, the elliptical vector $\mathbf{X} \sim E_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \phi)$ may not have a density function. However, if the density $f_{\mathbf{X}}(\mathbf{x})$ exists, then it is of the form

$$f_{\mathbf{X}}(\mathbf{x}) = \frac{c_n}{\sqrt{|\boldsymbol{\Sigma}|}} g_n\left(\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu})\right), \quad \mathbf{x} \in \mathbb{R}^n, \quad (1)$$

where $\boldsymbol{\mu}$ is an $n \times 1$ location vector, $\boldsymbol{\Sigma}$ is an $n \times n$ positive definite scale matrix, and $g_n(u)$, $u \geq 0$, is the density generator of \mathbf{X} . This density generator satisfies the condition

$$\int_0^\infty t^{(n/2)-1} g_n(t) dt < \infty,$$

and the normalizing constant c_n is given by

$$c_n = \frac{\Gamma(n/2)}{(2\pi)^{n/2}} \left[\int_0^\infty t^{(n/2)-1} g_n(t) dt \right]^{-1}. \quad (2)$$

Two important special cases are multivariate normal family with $g_n(u) = e^{-u}$, and multivariate generalized Student- t family with $g_n(u) = (1 + \frac{u}{k_{n,p}})^{-p}$, where the parameter $p > \frac{n}{2}$ and $k_{n,p}$ is some constant that may depend on n and p .

To derive the mixed moments of elliptical distributions, we use the cumulative generators $\overline{G}_n(u)$ and $\overline{\mathcal{G}}_n(u)$, which are given by

$$\overline{G}_n(u) = \int_u^\infty g_n(v) dv \quad (3)$$

and

$$\overline{\mathcal{G}}_n(u) = \int_u^\infty \overline{\mathcal{G}}_n(v) dv, \quad (4)$$

respectively (see Landsman et al. (2018)), and the corresponding normalizing constants are

$$c_n^* = \frac{\Gamma(n/2)}{(2\pi)^{n/2}} \left[\int_0^\infty t^{n/2-1} \overline{\mathcal{G}}_n(t) dt \right]^{-1} \quad (5)$$

and

$$c_n^{**} = \frac{\Gamma(n/2)}{(2\pi)^{n/2}} \left[\int_0^\infty t^{n/2-1} \overline{\mathcal{G}}_n(t) dt \right]^{-1}. \quad (6)$$

Throughout this paper, $\mathbf{x} \in \mathbb{R}^n$ will denote an n -dimensional vector and $\mathbf{x}^T = (x_1, \dots, x_n)$ its transpose. For an $n \times n$ matrix $\mathbf{\Sigma} \in \mathbb{R}^{n \times n}$, $|\mathbf{\Sigma}|$ is the determinant of $\mathbf{\Sigma}$. If $\mathbf{\Sigma}$ is positive definite, then its Cholesky decomposition is known to be unique.

3. Direct method of derivation

Consider a random vector $\mathbf{X} \sim E_n(\boldsymbol{\mu}, \mathbf{\Sigma}, g_n)$ with mean vector $\boldsymbol{\mu} = (\mu_1, \dots, \mu_n)^T$ and positive definite matrix $\mathbf{\Sigma} = (\sigma_{ij})_{i,j=1}^n$. Partition $\mathbf{y} \in \mathbb{R}^n$ into two parts as $\mathbf{y} = (y_1, \mathbf{y}_{(2)})^T$ each with 1 and $n-1$ components, respectively. By Cholesky decomposition (see Golub and Van Loan (2012), for example), there exists a unique lower triangular matrix $\mathbf{A} = (a_{ij})_{i,j=1}^n$ such that $\mathbf{A}\mathbf{A}^T = \mathbf{\Sigma}$. In terms of components, we have

$$a_{11} = \sqrt{\sigma_{11}}, \quad a_{i1} = \frac{\sigma_{i1}}{a_{11}}, \quad i = 2, \dots, n, \quad a_{ik} = 0, \quad i < k, \quad (7)$$

$$a_{kk} = \sqrt{\sigma_{kk} - \sum_{i=1}^{k-1} a_{ki}^2}, \quad a_{ik} = \frac{\sigma_{ik} - \sum_{j=1}^{k-1} a_{ij}a_{kj}}{a_{kk}}, \quad i = k+1, \dots, n. \quad (8)$$

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a twice continuously differentiable function, and shall use $(\nabla_{i,j}f(\mathbf{x}))_{i,j=1}^n$ to denote the Hessian matrix of f . In addition, we denote

$$\nabla_{i,j}f(\mathbf{x}) = \frac{\partial^2 f(\mathbf{x})}{\partial x_i \partial x_j}, \quad i, j = 1, 2, \dots, n,$$

$$\nabla_i f(\mathbf{x}) = \frac{\partial f(\mathbf{x})}{\partial y_i}, \quad i = 1, 2, \dots, n$$

and

$$\nabla f(\mathbf{x}) = \left(\frac{\partial f(\mathbf{x})}{\partial x_1}, \frac{\partial f(\mathbf{x})}{\partial x_2}, \dots, \frac{\partial f(\mathbf{x})}{\partial x_n} \right)^T.$$

Let $\mathbf{X}^* \sim E_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \overline{G}_n)$ and $\mathbf{X}^{**} \sim E_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \overline{\mathcal{G}}_n)$ be two elliptical random vectors with generators $\overline{G}_n(u)$ and $\overline{\mathcal{G}}_n(u)$, respectively.

The following theorem gives an expression for joint moments of elliptical distributions.

Theorem 1. *Let $\mathbf{X} \sim E_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, g_n)$ be an n -dimensional elliptical random vector with density generator g_n , positive definite matrix $\boldsymbol{\Sigma} = (\sigma_{i,j})_{i,j=1}^n$, and finite expectation $\boldsymbol{\mu}$. Further, let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a twice continuously differentiable function satisfying $E[\nabla_{i,j}f(\mathbf{X}^{**})] < \infty$ and $E[\nabla_i f(\mathbf{X}^*)] < \infty$. Let, in addition,*

$$\lim_{|\mathbf{x}| \rightarrow \infty} \mathbf{x}_1 f(\mathbf{A}\mathbf{x} + \boldsymbol{\mu}) \overline{G}_n\left(\frac{1}{2}\mathbf{x}^T \mathbf{x}\right) = 0 \quad (9)$$

and

$$\lim_{|\mathbf{x}| \rightarrow \infty} [\nabla_1 f(\mathbf{A}\mathbf{x} + \boldsymbol{\mu})] \overline{\mathcal{G}}_n\left(\frac{1}{2}\mathbf{x}^T \mathbf{x}\right) = 0. \quad (10)$$

Then,

$$\begin{aligned} E[X_1^2 f(\mathbf{X})] &= \sigma_{11} b_n^* E[f(\mathbf{X}^*)] + b_n^{**} \sum_{i=1}^n \sum_{j=1}^n \sigma_{i1} \sigma_{j1} E[\nabla_{i,j} f(\mathbf{X}^{**})] \\ &\quad + 2\mu_1 b_n^* \sum_{i=1}^n \sigma_{i1} E[\nabla_i f(\mathbf{X}^*)] + \mu_1^2 E[f(\mathbf{X})], \end{aligned} \quad (11)$$

where $b_n^* = \frac{c_n}{c_n^*}$ and $b_n^{**} = \frac{c_n}{c_n^{**}}$.

Proof. By definition,

$$\begin{aligned} E[X_1^2 f(\mathbf{X})] &= \frac{c_n}{\sqrt{|\Sigma|}} \int_{\mathbb{R}^n} x_1^2 f(\mathbf{x}) g_n \left\{ \frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^T \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right\} d\mathbf{x} \\ &= \frac{c_n}{\sqrt{|\Sigma|}} \int_{\mathbb{R}^n} x_1^2 f(\mathbf{x}) g_n \left\{ \frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^T (\mathbf{A} \mathbf{A}^T)^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right\} d\mathbf{x}. \end{aligned}$$

Now, setting $\mathbf{y} = \mathbf{A}^{-1}(\mathbf{x} - \boldsymbol{\mu})$, we obtain

$$\begin{aligned} E[X_1^2 f(\mathbf{X})] &= \frac{c_n |\mathbf{A}|}{\sqrt{|\Sigma|}} \int_{\mathbb{R}^n} (a_{1,1} y_1 + \mu_1)^2 f(\mathbf{A} \mathbf{y} + \boldsymbol{\mu}) g_n \left\{ \frac{1}{2} \mathbf{y}^T \mathbf{y} \right\} d\mathbf{y} \\ &= c_n \int_{\mathbb{R}^n} (a_{1,1} y_1 + \mu_1)^2 f(\mathbf{A} \mathbf{y} + \boldsymbol{\mu}) g_n \left\{ \frac{1}{2} \mathbf{y}^T \mathbf{y} \right\} d\mathbf{y} \\ &= a_{1,1}^2 c_n \int_{\mathbb{R}^n} y_1^2 f(\mathbf{A} \mathbf{y} + \boldsymbol{\mu}) g_n \left\{ \frac{1}{2} \mathbf{y}^T \mathbf{y} \right\} d\mathbf{y} \\ &\quad + 2a_{1,1} \mu_1 c_n \int_{\mathbb{R}^n} y_1 f(\mathbf{A} \mathbf{y} + \boldsymbol{\mu}) g_n \left\{ \frac{1}{2} \mathbf{y}^T \mathbf{y} \right\} d\mathbf{y} \\ &\quad + \mu_1^2 c_n \int_{\mathbb{R}^n} f(\mathbf{A} \mathbf{y} + \boldsymbol{\mu}) g_n \left\{ \frac{1}{2} \mathbf{y}^T \mathbf{y} \right\} d\mathbf{y} \\ &= a_{1,1}^2 c_n \int_{\mathbb{R}^{n-1}} I_1 d\mathbf{y}_{(2)} + 2a_{1,1} \mu_1 c_n \int_{\mathbb{R}^{n-1}} I_2 d\mathbf{y}_{(2)} \\ &\quad + \mu_1^2 c_n \int_{\mathbb{R}^n} f(\mathbf{A} \mathbf{y} + \boldsymbol{\mu}) g_n \left\{ \frac{1}{2} \mathbf{y}^T \mathbf{y} \right\} d\mathbf{y}, \end{aligned}$$

where

$$\begin{aligned} I_1 &= \int_{\mathbb{R}} y_1^2 f(\mathbf{A} \mathbf{y} + \boldsymbol{\mu}) g_n \left\{ \frac{1}{2} \mathbf{y}^T \mathbf{y} \right\} dy_1 \\ &= - \int_{\mathbb{R}} y_1 f(\mathbf{A} \mathbf{y} + \boldsymbol{\mu}) \frac{\partial}{\partial y_1} \overline{G}_n \left\{ \frac{1}{2} \mathbf{y}^T \mathbf{y} \right\} dy_1 \\ &= \int_{\mathbb{R}} [f(\mathbf{A} \mathbf{y} + \boldsymbol{\mu}) + y_1 \nabla_1 f(\mathbf{A} \mathbf{y} + \boldsymbol{\mu})] \overline{G}_n \left\{ \frac{1}{2} \mathbf{y}^T \mathbf{y} \right\} dy_1 \\ &= \int_{\mathbb{R}} f(\mathbf{A} \mathbf{y} + \boldsymbol{\mu}) \overline{G}_n \left\{ \frac{1}{2} \mathbf{y}^T \mathbf{y} \right\} dy_1 - \int_{\mathbb{R}} \nabla_1 f(\mathbf{A} \mathbf{y} + \boldsymbol{\mu}) \frac{\partial}{\partial y_1} \overline{G}_n \left\{ \frac{1}{2} \mathbf{y}^T \mathbf{y} \right\} dy_1 \\ &= \int_{\mathbb{R}} f(\mathbf{A} \mathbf{y} + \boldsymbol{\mu}) \overline{G}_n \left\{ \frac{1}{2} \mathbf{y}^T \mathbf{y} \right\} dy_1 + \int_{\mathbb{R}} \nabla_{1,1} f(\mathbf{A} \mathbf{y} + \boldsymbol{\mu}) \overline{G}_n \left\{ \frac{1}{2} \mathbf{y}^T \mathbf{y} \right\} dy_1, \end{aligned}$$

and

$$\begin{aligned}
I_2 &= \int_{\mathbb{R}} y_1 f(\mathbf{A}\mathbf{y} + \boldsymbol{\mu}) g_n \left\{ \frac{1}{2} \mathbf{y}^T \mathbf{y} \right\} dy_1 \\
&= - \int_{\mathbb{R}} f(\mathbf{A}\mathbf{y} + \boldsymbol{\mu}) \frac{\partial}{\partial y_1} \overline{G}_n \left\{ \frac{1}{2} \mathbf{y}^T \mathbf{y} \right\} dy_1 \\
&= \int_{\mathbb{R}} \nabla_1 f(\mathbf{A}\mathbf{y} + \boldsymbol{\mu}) \overline{G}_n \left\{ \frac{1}{2} \mathbf{y}^T \mathbf{y} \right\} dy_1.
\end{aligned}$$

We then obtain

$$\begin{aligned}
E[X_1^2 f(\mathbf{X})] &= a_{11}^2 \{ b_n^* E[f(\mathbf{X}^*)] + b_n^{**} \mathbf{A}_1^T E(\nabla_{i,j} f(\mathbf{X}^{**}))_{i,j=1}^n \mathbf{A}_1 \} \\
&\quad + 2a_{11}\mu_1 b_n^* \mathbf{A}_1^T E[\nabla f(\mathbf{X}^*)] + \mu_1^2 E[f(\mathbf{X})],
\end{aligned}$$

where $\mathbf{A} = (a_{ij})_{i,j=1}^n = (\mathbf{A}_1, \dots, \mathbf{A}_n)$. Now, upon using (7) and (8), we obtain (11), completing the proof of the theorem.

Kan (2008) and Song and Lee (2015) have presented explicit formulae for product moments of multivariate Gaussian random variables. The following corollary presents an explicit expression for product moments of multivariate elliptical random variables, in general.

Corollary 1. Suppose $\mathbf{X} \sim E_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, g_n)$, $\mathbf{X}^* \sim E_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \overline{G}_n)$ and $\mathbf{X}^{**} \sim E_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \overline{\mathcal{G}}_n)$. Let p_1, \dots, p_n be nonnegative integers with $p_1 \geq 2$. Then, we have

$$\begin{aligned}
E \left[\prod_{i=1}^n X_i^{p_i} \right] &= b_n^* \sigma_{11} E \left[(X_1^*)^{p_1-2} \prod_{k=2}^n (X_k^*)^{p_k} \right] \\
&\quad + b_n^{**} \left\{ \sigma_{11}^2 (p_1 - 2)(p_1 - 3) E \left[(X_1^{**})^{p_1-4} \prod_{k=2}^n (X_k^{**})^{p_k} \right] \right. \\
&\quad \left. + 2 \sum_{j=2}^n \sigma_{11} \sigma_{j1} (p_1 - 2) p_j E \left[(X_1^{**})^{p_1-3} (X_j^{**})^{p_j-1} \prod_{k=2, k \neq j}^n (X_k^{**})^{p_k} \right] \right. \\
&\quad \left. + \sum_{j=2}^n \sigma_{j1}^2 p_j (p_j - 1) E \left[(X_1^{**})^{p_1-2} (X_j^{**})^{p_j-2} \prod_{k=2, k \neq j}^n (X_k^{**})^{p_k} \right] \right\}
\end{aligned}$$

$$\begin{aligned}
& + \sum_{j=2}^n \sum_{i=2, i \neq j}^n \sigma_{j1} \sigma_{i1} p_j p_i E \left[(X_i^{**})^{p_i-1} (X_j^{**})^{p_j-1} (X_1^{**})^{p_1-2} \prod_{k=2, k \neq i, j}^n (X_k^{**})^{p_k} \right] \Big\} \\
& + 2b_n^* \mu_1 \left\{ \sigma_{11} (p_1 - 2) E \left[(X_1^{**})^{p_1-3} \prod_{k=2}^n (X_k^{**})^{p_k} \right] \right. \\
& \left. + \sum_{j=2}^n \sigma_{j1} p_j E \left[(X_j^*)^{p_j-1} (X_1^*)^{p_1-2} \prod_{k=2, k \neq j}^n (X_k^*)^{p_k} \right] \right\} + \mu_1^2 E \left[X_1^{p_1-2} \prod_{k=2}^n X_k^{p_k} \right].
\end{aligned} \tag{12}$$

From Corollary 1, we readily deduce the following relations for the special case of $p_1 = 3$, for example:

$$\begin{aligned}
E \left[X_1^3 \prod_{i=2}^n X_i^{p_i} \right] &= b_n^* \sigma_{11} E \left[X_1^* \prod_{k=2}^n (X_k^*)^{p_k} \right] + \mu_1^2 E \left[X_1 \prod_{k=2}^n X_k^{p_k} \right] \\
&+ b_n^{**} \left\{ 2 \sum_{j=2}^n \sigma_{11} \sigma_{j1} p_j E \left[(X_j^{**})^{p_j-1} \prod_{k=2, k \neq j}^n (X_k^{**})^{p_k} \right] \right. \\
&+ \sum_{j=2}^n \sigma_{j1}^2 p_j (p_j - 1) E \left[X_1^{**} (X_j^{**})^{p_j-2} \prod_{k=2, k \neq j}^n (X_k^{**})^{p_k} \right] \\
&+ \sum_{j=2}^n \sum_{i=2, i \neq j}^n \sigma_{j1} \sigma_{i1} p_j p_i E \left[(X_i^{**})^{p_i-1} (X_j^{**})^{p_j-1} X_1^{**} \prod_{k=2, k \neq i, j}^n (X_k^{**})^{p_k} \right] \Big\} \\
&+ 2b_n^* \mu_1 \left\{ \sigma_{11} E \left[\prod_{k=2}^n (X_k^{**})^{p_k} \right] + \sum_{j=2}^n \sigma_{j1} p_j E \left[(X_j^*)^{p_j-1} X_1^* \prod_{k=2, k \neq j}^n (X_k^*)^{p_k} \right] \right\}.
\end{aligned}$$

4. Derivation by the use of Stein's lemma

We now derive an expression for $E[X_1^2 f(\mathbf{X})]$ by using Stein's Lemma. It should be noted that the positive definiteness of Σ is not necessary in this case.

Theorem 2. *Suppose $\mathbf{X} \sim E_n(\boldsymbol{\mu}, \Sigma, \phi)$, and all the conditions of Theorem 1 hold. Then,*

$$\begin{aligned}
E[X_1^2 f(\mathbf{X})] &= \sum_{i=1}^n \sum_{j=1}^n Cov(X_1, X_i) Cov(X_1^*, X_j^*) E[\nabla_{i,j} f(\mathbf{X}^{**})] \\
&\quad + 2\mu_1 \sum_{i=1}^n Cov(X_1, X_i) E[\nabla_i f(\mathbf{X}^*)] \\
&\quad + Cov(X_1, X_1) E[f(\mathbf{X}^*)] + \mu_1^2 E[f(\mathbf{X})]. \tag{13}
\end{aligned}$$

Proof. By Lemma 2 of Landsman et al. (2013), we have

$$Cov(X_1, f(\mathbf{X})) = \sum_{i=1}^n Cov(X_1, X_i) E[\nabla_i f(\mathbf{X}^*)],$$

and so

$$\begin{aligned}
E[X_1 f(\mathbf{X})] &= Cov(X_1, f(\mathbf{X})) + E[X_1] E[f(\mathbf{X})] \\
&= \sum_{i=1}^n Cov(X_1, X_i) E[\nabla_i f(\mathbf{X}^*)] + E[X_1] E[f(\mathbf{X})]. \tag{14}
\end{aligned}$$

Upon replacing $f(\mathbf{X})$ by $X_1 f(\mathbf{X})$ in (14), we obtain

$$\begin{aligned}
E[X_1^2 f(\mathbf{X})] &= \sum_{i=1}^n Cov(X_1, X_i) E[\nabla_i (X_1^* f(\mathbf{X}^*))] + E[X_1] E[X_1 f(\mathbf{X})] \\
&= \sum_{i=1}^n Cov(X_1, X_i) E[X_1^* \nabla_i f(\mathbf{X}^*)] + Cov(X_1, X_1) E[f(\mathbf{X}^*)] + E[X_1] E[X_1 f(\mathbf{X})] \\
&= \sum_{i=1}^n Cov(X_1, X_i) \left\{ \sum_{j=1}^n Cov(X_1^*, X_j^*) E[\nabla_{i,j} f(\mathbf{X}^{**})] + E[X_1^*] E[\nabla_i f(\mathbf{X}^*)] \right\} \\
&\quad + Cov(X_1, X_1) E[f(\mathbf{X}^*)] + E[X_1] \left\{ \sum_{i=1}^n Cov(X_1, X_i) E[\nabla_i f(\mathbf{X}^*)] + E[X_1] E[f(\mathbf{X})] \right\} \\
&= \sum_{i=1}^n \sum_{j=1}^n Cov(X_1, X_i) Cov(X_1^*, X_j^*) E[\nabla_{i,j} f(\mathbf{X}^{**})] + (E[X_1])^2 E[f(\mathbf{X})] \\
&\quad + 2E[X_1] \sum_{i=1}^n Cov(X_1, X_i) E[\nabla_i f(\mathbf{X}^*)] + Cov(X_1, X_1) E[f(\mathbf{X}^*)],
\end{aligned}$$

as required.

5. Equivalence of the two expressions

We shall now establish the equivalence of the two expressions in (11) and (13) under the condition that Σ is positive definite. For this purpose, we will use the following lemma; see, for example, Fang et al. (1990).

Lemma 1. *For any non-negative measurable function $f : \mathbb{R} \rightarrow \mathbb{R}^+$, we have*

$$\int_{\mathbb{R}^n} f\left(\frac{1}{2}\mathbf{y}^T\mathbf{y}\right) d\mathbf{y} = \frac{(2\pi)^{n/2}}{\Gamma(n/2)} \int_0^\infty u^{n/2-1} f(u) du.$$

Proposition 1. *Under the conditions in Theorem 1, we have*

$$\frac{c_n}{c_n^*} = -\phi'(0), \quad \frac{c_n^*}{c_n^{**}} = -\phi^{*'}(0),$$

where ϕ and ϕ^* are the characteristic generating functions corresponding to the density generators g_n and \overline{G}_n , respectively.

Proof. Let $\mathbf{Y} = \sqrt{-\phi'(0)\Sigma^{-1}}(\mathbf{X} - E(\mathbf{X}))$. Then, $\mathbf{X} \sim E_n(\boldsymbol{\mu}, \Sigma, g_n)$ implies $\mathbf{Y} \sim E_n(\mathbf{0}, \mathbf{I}_n, g_n)$ and $\text{Cov}(\mathbf{Y}) = -\phi'(0)\mathbf{I}_n$. It then follows that

$$\begin{aligned} E(\mathbf{Y}^T\mathbf{Y}) &= -n\phi'(0) \\ &= c_n \int_{\mathbb{R}^n} \mathbf{y}^T\mathbf{y} g_n\left(\frac{1}{2}\mathbf{y}^T\mathbf{y}\right) d\mathbf{y} \\ &= 2c_n \frac{(2\pi)^{n/2}}{\Gamma(n/2)} \int_0^\infty t^{n/2} g_n(t) dt, \end{aligned}$$

by Lemma 1. Thus,

$$c_n = \frac{-n\phi'(0)}{2 \frac{(2\pi)^{n/2}}{\Gamma(n/2)} \int_0^\infty t^{n/2} g_n(t) dt}. \quad (15)$$

Now, by using Eq.(9) of Landsman et al. (2013), we have

$$1 = c_n^* \int_{\mathbb{R}^n} \overline{G}_n\left(\frac{1}{2}\mathbf{y}^T\mathbf{y}\right) d\mathbf{y} = c_n^* \frac{(2\pi)^{n/2}}{\Gamma(n/2+1)} \int_0^\infty t^{n/2} g_n(t) dt,$$

and hence

$$c_n^* = \frac{1}{\frac{(2\pi)^{n/2}}{\Gamma(n/2+1)} \int_0^\infty t^{n/2} g_n(t) dt}. \quad (16)$$

The result that $\frac{c_n}{c_n^*} = -\phi'(0)$ readily follows from (15) and (16).

For $\mathbf{X}^* \sim E_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \overline{G}_n)$, let $\mathbf{Z} = \sqrt{-\phi^{*'}(0)\boldsymbol{\Sigma}^{-1}}(\mathbf{X}^* - E(\mathbf{X}^*))$. Then, $\mathbf{Z} \sim E_n(\mathbf{0}, \mathbf{I}_n, \overline{G}_n)$ and $Cov(\mathbf{Z}) = -\phi^{*'}(0)\mathbf{I}_n$. By using the same arguments as above, we have

$$\begin{aligned} E(\mathbf{Z}^T \mathbf{Z}) &= -n\phi^{*'}(0) \\ &= c_n^* \int_{\mathbb{R}^n} \mathbf{z}^T \mathbf{z} \overline{G}_n \left(\frac{1}{2} \mathbf{z}^T \mathbf{z} \right) d\mathbf{z} \\ &= 2c_n^* \frac{(2\pi)^{n/2}}{\Gamma(n/2)} \int_0^\infty t^{n/2} \overline{G}_n(t) dt \\ &= nc_n^* \frac{(2\pi)^{n/2}}{\Gamma(n/2)} \int_0^\infty t^{n/2-1} \overline{G}_n(t) dt \\ &= \frac{nc_n^*}{c_n^{**}}, \end{aligned}$$

and hence

$$\frac{c_n^*}{c_n^{**}} = -\phi^{*'}(0),$$

as required.

Remark 1. From Proposition 1, we find that, when $\boldsymbol{\Sigma}$ is a positive definite, the expressions in (11) and (13) are indeed equivalent. Moreover, for a positive semidefinite matrix $\boldsymbol{\Sigma}$, we can rewrite (13) in terms of characteristic generators as follows:

$$\begin{aligned} E[X_1^2 f(\mathbf{X})] &= \phi'(0)\phi^{*'}(0) \sum_{i=1}^n \sum_{j=1}^n \sigma_{1i}\sigma_{1j} E[\nabla_{i,j} f(\mathbf{X}^{**})] \\ &\quad - 2\mu_1\phi'(0) \sum_{i=1}^n \sigma_{1i} E[\nabla_i f(\mathbf{X}^*)] \\ &\quad - \phi'(0)\sigma_{11} E[f(\mathbf{X}^*)] + \mu_1^2 E[f(\mathbf{X})], \end{aligned} \quad (17)$$

where ϕ and ϕ^* are the characteristic generators corresponding to the density generators g_n and \overline{G}_n , respectively.

6. Product moments of correlated normal random variables

We derive now expressions for $E[X_1^2 f(\mathbf{X})]$ of multivariate normal distribution and moments of products of correlated normal distribution.

Corollary 2. Suppose $\mathbf{X} \sim E_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \phi)$ with density generator $g(u) = \exp\{-u\}$ and characteristic generator $\phi(t) = \exp\{-t\}$. Note in this case that $g(u) = \overline{G}(u) = \overline{\mathcal{G}}(u) = \exp\{-u\}$, $\phi(t) = \phi^*(t)$ and $b_n^* = b_n^{**} = 1$. Then, we have

$$\begin{aligned} E[X_1^2 f(\mathbf{X})] &= \sum_{i=1}^n \sum_{j=1}^n \sigma_{1i} \sigma_{1j} E[\nabla_{i,j} f(\mathbf{X})] + 2\mu_1 \sum_{i=1}^n \sigma_{1i} E[\nabla_i f(\mathbf{X})] \\ &\quad + \sigma_{11} E[f(\mathbf{X})] + \mu_1^2 E[f(\mathbf{X})]. \end{aligned}$$

In general, for any $p_1 \geq 2$, we have the following recursive relation:

$$\begin{aligned} E[X_1^{p_1} f(\mathbf{X})] &= \sum_{i=1}^n \sum_{j=1}^n \sigma_{1i} \sigma_{1j} E[\nabla_{i,j} (X_1^{p_1-2} f(\mathbf{X}))] \\ &\quad + 2\mu_1 \sum_{i=1}^n \sigma_{1i} E[\nabla_i (X_1^{p_1-2} f(\mathbf{X}))] + \sigma_{11} E[X_1^{p_1-2} f(\mathbf{X})] + \mu_1^2 E[X_1^{p_1-2} f(\mathbf{X})] \\ &= \sum_{i=1}^n \sum_{j=1}^n \sigma_{1i} \sigma_{1j} E[X_1^{p_1-2} \nabla_{i,j} f(\mathbf{X})] + \sigma_{11}^2 (p_1 - 2)(p_1 - 3) E[X_1^{p_1-4} f(\mathbf{X})] \\ &\quad + 2\mu_1 \sum_{i=1}^n \sigma_{1i} E[X_1^{p_1-2} \nabla_i f(\mathbf{X})] + 2\mu_1 \sigma_{11} (p_1 - 2) E[X_1^{p_1-3} f(\mathbf{X})] \\ &\quad + \sigma_{11} E[X_1^{p_1-2} f(\mathbf{X})] + \mu_1^2 E[X_1^{p_1-2} f(\mathbf{X})]. \end{aligned}$$

This formula can be seen as a supplement to the following multivariate version of Stein's identity (see Stein (1981) and Liu (1994)):

$$\text{Cov}(X_1, f(\mathbf{X})) = \sum_{i=1}^n \text{Cov}(X_1, X_i) E(\nabla_i f(\mathbf{X})),$$

or equivalently,

$$E(X_1 f(\mathbf{X})) = \sum_{i=1}^n \sigma_{1i} E(\nabla_i f(\mathbf{X})) + E(X_1) E(f(\mathbf{X})).$$

Stein's identity for multivariate elliptical distributions also has a similar result, that can be found in Landsman et al. (2013).

In particular, when $\phi(t) = \exp\{-t\}$, then $\mathbf{X} \sim N_n(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, $\mathbf{X}^* \sim N_n(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ and $\mathbf{X}^{**} \sim N_n(\boldsymbol{\mu}, \boldsymbol{\Sigma})$. In this case, we obtain the following recursion formula:

$$\begin{aligned} E \left[\prod_{i=1}^n X_i^{p_i} \right] &= \sigma_{11} E \left[(X_1)^{p_1-2} \prod_{i=2}^n (X_i)^{p_i} \right] + \mu_1^2 E \left[X_1^{p_1-2} \prod_{k=2}^n X_k^{p_k} \right] \\ &+ \sigma_{11}^2 (p_1 - 2)(p_1 - 3) E \left[(X_1)^{p_1-4} \prod_{k=2}^n (X_k)^{p_k} \right] \\ &+ 2 \sum_{j=2}^n \sigma_{11} \sigma_{j1} (p_1 - 2) p_j E \left[(X_1)^{p_1-3} (X_j)^{p_j-1} \prod_{k=2, k \neq j}^n (X_k)^{p_k} \right] \\ &+ \sum_{j=2}^n \sigma_{j1}^2 p_j (p_j - 1) E \left[(X_1)^{p_1-2} (X_j)^{p_j-2} \prod_{k=2, k \neq j}^n (X_k)^{p_k} \right] \\ &+ \sum_{j=2}^n \sum_{i=2, i \neq j}^n \sigma_{j1} \sigma_{i1} p_j p_i E \left[(X_i)^{p_i-1} (X_j)^{p_j-1} (X_1)^{p_1-2} \prod_{k=2, k \neq i, j}^n (X_k)^{p_k} \right] \\ &+ 2\mu_1 \left\{ \sigma_{11} (p_1 - 2) E \left[(X_1)^{p_1-3} \prod_{k=2}^n (X_k)^{p_k} \right] \right. \\ &\left. + \sum_{j=2}^n \sigma_{j1} p_j E \left[(X_j)^{p_j-1} (X_1)^{p_1-2} \prod_{k=2, k \neq j}^n (X_k)^{p_k} \right] \right\}. \end{aligned}$$

7. Results for some special cases

In this section, we present the results for the special case of multivariate Student- t , logistic and Laplace distributions.

Example 7.1 Multivariate Student- t distribution. A n -dimensional Student- t random vector \mathbf{X} , with location parameter $\boldsymbol{\mu}$, scale matrix $\boldsymbol{\Sigma}$ and $p > 0$

degrees of freedom, has its density function as

$$f_{\mathbf{X}}(\mathbf{x}) = \frac{c_n}{\sqrt{|\Sigma|}} \left[1 + \frac{(\mathbf{x} - \boldsymbol{\mu})^T \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu})}{p} \right]^{-\frac{p+n}{2}}, \quad \mathbf{x} \in \mathbb{R}^n,$$

where $c_n = \frac{\Gamma(\frac{p+n}{2})}{\Gamma(p/2)(p\pi)^{\frac{n}{2}}}$. We denote it by $\mathbf{X} \sim St_n(\boldsymbol{\mu}, \Sigma, p)$. In this case, the density generator is

$$g_n(t) = \left(1 + \frac{2t}{p} \right)^{-(p+n)/2},$$

and so $\overline{G}_n(t)$ and $\overline{\mathcal{G}}_n(t)$ can be expressed, respectively, as

$$\overline{G}_n(t) = \frac{p}{p+n-2} \left(1 + \frac{2t}{p} \right)^{-(p+n-2)/2}$$

and

$$\overline{\mathcal{G}}_n(t) = \frac{p}{p+n-2} \frac{p}{p+n-4} \left(1 + \frac{2t}{p} \right)^{-(p+n-4)/2}.$$

In addition,

$$\begin{aligned} c_n^* &= \frac{(p+n-2)\Gamma(n/2)}{(2\pi)^{n/2}p} \left[\int_0^\infty t^{n/2-1} \left(1 + \frac{2t}{p} \right)^{-(p+n-2)/2} dt \right]^{-1} \\ &= \frac{(p+n-2)\Gamma(n/2)}{(p\pi)^{n/2}pB(\frac{n}{2}, \frac{p-2}{2})}, \quad \text{if } p > 2 \end{aligned}$$

and

$$\begin{aligned} c_n^{**} &= \frac{(p+n-2)(p+n-4)\Gamma(n/2)}{(2\pi)^{n/2}p^2} \left[\int_0^\infty t^{n/2-1} \left(1 + \frac{2t}{p} \right)^{-(p+n-4)/2} dt \right]^{-1} \\ &= \frac{(p+n-2)(p+n-4)\Gamma(n/2)}{(p\pi)^{n/2}p^2B(\frac{n}{2}, \frac{p-4}{2})}, \quad \text{if } p > 4, \end{aligned}$$

where $\Gamma(\cdot)$ and $B(\cdot, \cdot)$ are Gamma function and Beta function, respectively.

Then, we have

$$\begin{aligned} E[X_1^2 f(\mathbf{X})] &= \sigma_{11} b_n^* E[f(\mathbf{X}^*)] + b_n^{**} \sum_{i=1}^n \sum_{j=1}^n \sigma_{i1} \sigma_{j1} E[\nabla_{i,j} f(\mathbf{X}^{**})] \\ &\quad + 2\mu_1 b_n^* \sum_{i=1}^n \sigma_{i1} E[\nabla_i f(\mathbf{X}^*)] + \mu_1^2 E[f(\mathbf{X})], \end{aligned} \quad (18)$$

where $\mathbf{X} \sim E_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, g_n)$, $\mathbf{X}^* \sim E_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \overline{G}_n)$, $\mathbf{X}^{**} \sim E_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \overline{\mathcal{G}}_n)$,

$$b_n^* = \frac{p^2 \Gamma(\frac{p+n}{2}) B(\frac{n}{2}, \frac{p-2}{2})}{(p+n-2) \Gamma(\frac{p}{2}) \Gamma(\frac{n}{2})} = \frac{p^2}{p-2}, \text{ if } p > 2$$

and

$$b_n^{**} = \frac{p \Gamma(\frac{p+n}{2}) B(\frac{n}{2}, \frac{p-4}{2})}{(p+n-2)(p+n-4) \Gamma(\frac{p}{2}) \Gamma(\frac{n}{2})} = \frac{p}{(p-2)(p-4)}, \text{ if } p > 4.$$

Example 7.2 Multivariate logistic distribution. The density function of a n -dimension logistic random vector \mathbf{X} , with location parameter $\boldsymbol{\mu}$ and scale matrix $\boldsymbol{\Sigma}$, is given by

$$f_{\mathbf{X}}(\mathbf{x}) = \frac{c_n}{\sqrt{|\boldsymbol{\Sigma}|}} \frac{\exp\left\{-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu})\right\}}{\left[1 + \exp\left\{-\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\mathbf{x} - \boldsymbol{\mu})\right\}\right]^2}, \quad \mathbf{x} \in \mathbb{R}^n,$$

where

$$\begin{aligned} c_n &= \frac{\Gamma(n/2)}{(2\pi)^{n/2}} \left[\int_0^\infty t^{n/2-1} \frac{\exp(-t)}{[1 + \exp(-t)]^2} dt \right]^{-1} \\ &= \frac{1}{(2\pi)^{n/2} \Psi_2^*(-1, \frac{n}{2}, 1)}. \end{aligned}$$

Here $\Psi_\mu^*(z, s, a)$ is the generalized Hurwitz-Lerch zeta function defined by (cf. Lin et al.(2006))

$$\Psi_\mu^*(z, s, a) = \frac{1}{\Gamma(\mu)} \sum_{n=0}^{\infty} \frac{\Gamma(\mu+n)}{n!} \frac{z^n}{(n+a)^s},$$

which has an integral representation

$$\Psi_\mu^*(z, s, a) = \frac{1}{\Gamma(s)} \int_0^\infty \frac{t^{s-1} e^{-at}}{(1 - ze^{-t})^\mu} dt,$$

where $\mathcal{R}(a) > 0$; $\mathcal{R}(s) > 0$ when $|z| \leq 1$ ($z \neq 1$); $\mathcal{R}(s) > 1$ when $z = 1$. We denote it by $\mathbf{X} \sim Lo_n(\boldsymbol{\mu}, \boldsymbol{\Sigma})$. The density generator in this case is

$$g_n(t) = \frac{\exp(-t)}{[1 + \exp(-t)]^2},$$

and $\overline{G}_n(t)$ and $\overline{\mathcal{G}}_n(t)$ are given by

$$\overline{G}_n(t) = \frac{\exp(-t)}{1 + \exp(-t)}, \quad \overline{\mathcal{G}}_n(t) = \ln[1 + \exp(-t)].$$

In addition,

$$\begin{aligned} c_n^* &= \frac{\Gamma(n/2)}{(2\pi)^{n/2}} \left[\int_0^\infty t^{n/2-1} \frac{\exp(-t)}{1 + \exp(-t)} dt \right]^{-1} \\ &= \frac{1}{(2\pi)^{n/2} \Psi_1^*(-1, \frac{n}{2}, 1)} \end{aligned}$$

and

$$\begin{aligned} c_n^{**} &= \frac{\Gamma(n/2)}{(2\pi)^{n/2}} \left\{ \int_0^\infty t^{n/2-1} \ln[1 + \exp(-t)] dt \right\}^{-1} \\ &= \frac{\Gamma(n/2)}{(2\pi)^{n/2}} \left[\frac{2}{n} \int_0^\infty t^{n/2} \frac{e^{-t}}{1 + e^{-t}} dt \right]^{-1} \\ &= \frac{1}{(2\pi)^{n/2} \Psi_1^*(-1, \frac{n}{2} + 1, 1)}. \end{aligned}$$

Then, we have

$$\begin{aligned} E[X_1^2 f(\mathbf{X})] &= \sigma_{11} b_n^* E[f(\mathbf{X}^*)] + b_n^{**} \sum_{i=1}^n \sum_{j=1}^n \sigma_{i1} \sigma_{j1} E[\nabla_{i,j} f(\mathbf{X}^{**})] \\ &\quad + 2\mu_1 b_n^* \sum_{i=1}^n \sigma_{i1} E[\nabla_i f(\mathbf{X}^*)] + \mu_1^2 E[f(\mathbf{X})], \end{aligned} \tag{19}$$

where $\mathbf{X} \sim E_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, g_n)$, $\mathbf{X}^* \sim E_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \overline{G}_n)$, $\mathbf{X}^{**} \sim E_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \overline{\mathcal{G}}_n)$,

$$b_n^* = \frac{\Psi_1^*(-1, \frac{n}{2}, 1)}{\Psi_2^*(-1, \frac{n}{2}, 1)}$$

and

$$b_n^{**} = \frac{\Psi_1^*(-1, \frac{n}{2} + 1, 1)}{\Psi_2^*(-1, \frac{n}{2}, 1)}.$$

Remark 2 A simplification of c_n can be found in Yin et al. (2018):

$$c_n = \begin{cases} \frac{1}{\sqrt{2\pi}} [\Psi_2^*(-1, \frac{n}{2}, 1)]^{-1}, & \text{if } n = 1, \\ \frac{1}{\pi}, & \text{if } n = 2, \\ \frac{1}{4\pi^2 \ln 2}, & \text{if } n = 4, \\ \frac{1}{\pi^{n/2} (2^{n/2} - 4) \zeta(\frac{n}{2} - 1)}, & \text{if } n \geq 3, n \neq 4, \end{cases}$$

where ζ is Riemann zeta function, and it is defined as

$$\zeta(s) = \begin{cases} \sum_{n=1}^{\infty} \frac{1}{n^s} = \frac{1}{1-2^{-s}} \sum_{n=1}^{\infty} \frac{1}{(2n-1)^s}, & \text{if } \mathcal{R}(s) > 1, \\ \frac{1}{1-2^{1-s}} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^s}, & \text{if } \mathcal{R}(s) > 0, s \neq 1, \end{cases}$$

which can, except for a simple pole at $s = 1$ with its residue 1, be continued meromorphically to the whole complex s -plane (see Srivastava (2003) and Choi et al. (2004) for details).

Example 7.3 Multivariate Laplace distribution. The density function of a Laplace random vector \mathbf{X} , with location parameter $\boldsymbol{\mu}$ and scale matrix $\boldsymbol{\Sigma}$, is given by

$$f_{\mathbf{X}}(\mathbf{x}) = \frac{c_n}{\sqrt{|\boldsymbol{\Sigma}|}} \exp \left\{ -[(\mathbf{x} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu})]^{1/2} \right\}, \quad \mathbf{x} \in \mathbb{R}^n,$$

where $c_n = \frac{\Gamma(n/2)}{2\pi^{n/2}\Gamma(n)}$. We denote it by $\mathbf{X} \sim La_n(\boldsymbol{\mu}, \boldsymbol{\Sigma})$. The density generator in this case is $g_n(t) = \exp(-\sqrt{2t})$, and so

$$\overline{G}_n(t) = (1 + \sqrt{2t}) \exp(-\sqrt{2t}),$$

$$\overline{\mathcal{G}}_n(t) = (3 + 2t + 3\sqrt{2t}) \exp(-\sqrt{2t}).$$

In addition,

$$c_n^* = \frac{n\Gamma(n/2)}{2\pi^{n/2}\Gamma(n+2)}, \quad c_n^{**} = \frac{n(n+2)\Gamma(n/2)}{2\pi^{n/2}\Gamma(n+4)}.$$

Then, we have

$$\begin{aligned}
E[X_1^2 f(\mathbf{X})] &= \sigma_{11} b_n^* E[f(\mathbf{X}^*)] + b_n^{**} \sum_{i=1}^n \sum_{j=1}^n \sigma_{i1} \sigma_{j1} E[\nabla_{i,j} f(\mathbf{X}^{**})] \\
&\quad + 2\mu_1 b_n^* \sum_{i=1}^n \sigma_{i1} E[\nabla_i f(\mathbf{X}^*)] + \mu_1^2 E[f(\mathbf{X})], \tag{20}
\end{aligned}$$

where $\mathbf{X} \sim E_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, g_n)$, $\mathbf{X}^* \sim E_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \overline{G}_n)$, $\mathbf{X}^{**} \sim E_n(\boldsymbol{\mu}, \boldsymbol{\Sigma}, \overline{\mathcal{G}}_n)$,

$$b_n^* = n + 1 \quad \text{and} \quad b_n^{**} = (n + 3)(n + 1).$$

8. Concluding remarks

In this work, we have derived expressions for the joint moments of elliptical distributions by two methods and have shown their equivalence when the dispersion matrix $\boldsymbol{\Sigma}$ is positive definite. We have used the result to derive expectations of products of correlated normal random variables and have also presented simplified expressions for the joint moments of the special case of multivariate Student- t , logistic and Laplace distributions. It will be of interest to extend the results established here to truncated elliptical distributions along the lines of Shushi (2018), and to the family of skew-elliptical distributions proposed by Branco and Dey (2001). Work on these problems is currently under progress and hope to report the findings in a future paper.

Acknowledgments

The research was supported by the National Natural Science Foundation of China (No. 11571198, 11701319).

References

References

- [1] Branco, M.D., Dey, D.K., 2001. A general class of multivariate skew-elliptical distributions. *Journal of Multivariate Analysis* 79 (1), 99-113.

- [2] Choi, J., Cho, Y.J., Srivastava, H.M., 2004. Series involving the zeta function and multiple Gamma functions. *Applied Mathematics and Computation* 159, 509-537.
- [3] Fang, K.T., Kotz, S., Ng, K.W., 1990. *Symmetric Multivariate and Related Distributions*. Chapman and Hall, New York.
- [4] Golub, G.H., Van Loan, C.F., 2012. *Matrix Computation* (4th Edition). Johns Hopkins University Press, Baltimore, Maryland.
- [5] Kan, R., 2008. From moments of sum to moments of product. *Journal of Multivariate Analysis* 99 (3), 542-554.
- [6] Kelker, D., 1970. Distribution theory of spherical distributions and location-scale parameter generalization. *Sankhya* 32, 419-430.
- [7] Kotz, S., Balakrishnan, N., Johnson, N.L., 2000. *Continuous Multivariate Distributions-Vol.1*, Second edition. John Wiley and Sons, New York.
- [8] Landsman, Z., 2006. On the generalization of Stein's lemma for elliptical class of distributions. *Statistics and Probability Letters* 76, 1012-1016.
- [9] Landsman, Z., Makov, U., Shushi, T., 2018. A multivariate tail covariance measure for elliptical distributions. *Insurance: Mathematics and Economics* 81, 27-35.
- [10] Landsman, Z., Nešlehová, J., 2008. Stein's Lemma for elliptical random vectors. *Journal of Multivariate Analysis* 99, 912-927.
- [11] Landsman, Z., Vanduffel, S., Yao, J., 2013. A note on Stein's lemma for multivariate elliptical distributions. *Journal of Statistical Planning and Inference* 143, 2016-2022.

- [12] Lin, S.D., Srivastava, H. M., Wang, P. Y., 2006. Some expansion formulas for a class of generalized Hurwitz-Lerch Zeta functions, *Integral Transforms and Special Functions* 17 (11), 817-827.
- [13] Liu, J.S., 1994. Siegel's formula via Stein's identities. *Statistics and Probability Letters* 21, 247-251.
- [14] Shushi, T., 2018. Stein's lemma for truncated elliptical random vectors. *Statistics and Probability Letters* 137, 297-303.
- [15] Song, I., Lee, S., 2015. Explicit formulae for product moments of multivariate Gaussian random variables. *Statistics and Probability Letters* 100, 27-34.
- [16] Srivastava, H.M., 2003. Certain classes of series associated with the zeta and related functions. *Applied Mathematics and Computation* 141, 13-49.
- [17] Stein, C.M., 1981. Estimation of the mean of a multivariate normal distribution. *The Annals of Statistics* 9, 1135-1151.
- [18] Yin, C.C., Wang, Y., Sha, X.Y., 2018. A new class of symmetric distributions including the elliptically symmetric logistic. Submitted for publication.